

# Possible origin of the KM3-230213A neutrino event from dark matter decay

Debasish Borah,<sup>1,2,\*</sup> Nayan Das,<sup>1,†</sup> Nobuchika Okada,<sup>3,‡</sup> and Prantik Sarmah<sup>4,§</sup>

<sup>1</sup>*Department of Physics, Indian Institute of Technology Guwahati, Assam 781039, India*

<sup>2</sup>*Pittsburgh Particle Physics, Astrophysics, and Cosmology Center,*

*Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, PA 15260, USA*

<sup>3</sup>*Department of Physics, University of Alabama, Tuscaloosa, Alabama 35487, USA*

<sup>4</sup>*Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, 100049, People's Republic of China*

## Abstract

We study the possibility of the highest energy neutrino event with 220 PeV energy, detected recently by the KM3NeT experiment to be originating from heavy dark matter (DM) decay. Considering a heavy right handed neutrino (RHN) DM for illustrative purpose, we show that DM mass of 440 PeV, can explain the observed flux. The required DM lifetime to produce the best-fit value of the neutrino flux saturates the existing gamma-ray bounds. Due to the large uncertainty in the flux, it is possible to explain the KM3NeT event from RHN DM decay at  $3\sigma$  confidence level (CL) while being in agreement with gamma-ray bounds and non-observation of similar events at IceCube. While we consider a gauged  $B - L$  scenario where DM relic can be generated due to other interactions, we also briefly discuss some alternate DM possibilities where the gamma-ray bounds can be alleviated compared to the minimal RHN DM discussed here.

## I. INTRODUCTION

Recently, the KM3NeT collaboration has reported the detection of a neutrino event named as KM3-230213A having  $\mathcal{O}(100\text{PeV})$  energy [1]. The deep-sea neutrino telescope has detected an ultra-high-energy (UHE) muon with energy  $120_{-60}^{+110}$  PeV arriving from a near-horizontal direction (RA:  $94.3^\circ$ , Dec:  $-7.8^\circ$ ) which is most likely to be created by a neutrino of even higher energy in the vicinity of the detector. The required neutrino energy was found to be in the range 110-790 PeV with median energy being 220 PeV. This is the highest energy neutrino event ever detected. Creation of such energetic neutrinos through cosmic ray interactions ( $pp$  or  $p\gamma$ ) in conventional astrophysical sources would require protons to be accelerated to around EeV energies. However, such sources are not available in our Milky Way or any nearby Galaxies. The KM3NeT collaboration has analyzed several classes of source candidates such as blazars, gamma-ray bursts using different electromagnetic telescope catalogs and found no conclusive evidence [1]. The absence of any conclusive astrophysical source [2, 3] points towards a cosmic origin of the highest energy neutrino event detected so far [4]. Cosmogenic UHE neutrinos together with UHE gamma-rays are expected to be created in the interactions of UHE cosmic rays with the cosmic microwave background (CMB) photons, the so called Greisen-Zatsepin-Kuzmin (GZK) process [5, 6]. Several model estimates of these UHE neutrinos and gamma-rays flux can be found in the literature [e.g., see 7–10]. The UHE neutrino flux required to produce the KM3-230213A event was found to overshoot not only these model estimates, but also the sensitivi-

ties of other presently running neutrino telescopes such as IceCube and Pierre Auger Observatory (PAO). This could be due to the large uncertainty (about 3 orders of magnitude) in the KM3-230213A flux, possibly created by an upward fluctuation [1]. On the other hand, non-observation of similar events at other experiments like IceCube and PAO leads to a  $2.5\sigma - 3\sigma$  tension<sup>1</sup> with the cosmogenic origin hypothesis of the KM3-230213A event [12]. Several follow-up studies on implications of this neutrino event for UHE cosmic rays [13, 14], Lorentz invariance violation [15–17], possible astrophysical origin [18, 19], possible origin from primordial black hole evaporation [20], in-vacuo dispersion of neutrinos [21] have also appeared in the literature.

In this work, we consider a dark matter (DM) origin of the KM3-230213A event flux at  $3\sigma$  CL while also addressing the the mild tension with IceCube and PAO for such diffuse, isotropic origin of the event. DM decay origin of the PeV neutrino events at IceCube [22, 23] more than a decade ago was also explored in several works including [24, 25]. We consider a heavy right handed neutrino (RHN) to be the DM candidate whose late decay into neutrinos is responsible for the highest energy neutrino event mentioned above. While the neutrino event can, in principle, originate from DM annihilation as well, the flux will be proportional to DM density squared and hence suppressed for such superheavy DM. Such a RHN appears in type-I seesaw scenario which explains the origin of light neutrino masses. While two RHNs are sufficient to generate light neutrino mass and mixing, the third RHN can be DM if sufficiently long-lived due to feeble Yukawa coupling with the standard model (SM) leptons. Since RHN can decay into other SM particles due to the same Yukawa interactions, we also have secondary gamma-ray productions, tightly constrained from

\*Electronic address: [dborah@iitg.ac.in](mailto:dborah@iitg.ac.in)

†Electronic address: [nayan.das@iitg.ac.in](mailto:nayan.das@iitg.ac.in)

‡Electronic address: [okadan@ua.edu](mailto:okadan@ua.edu)

§Electronic address: [prantiksarmah@ihep.ac.cn](mailto:prantiksarmah@ihep.ac.cn)

<sup>1</sup> An independent analysis [11] finds a slightly larger tension.

astrophysical observations. We show that the required flux and energy of the KM3-230213A neutrino event can be explained within the uncertainties from decay of a RHN DM having mass  $M_{\text{DM}} \sim 440$  PeV while being in agreement with the gamma-ray bounds. While the required relic of such superheavy DM can not be generated in the minimal type-I seesaw model, we consider a gauged  $U(1)_{B-L}$  extension of the model where relic can be generated non-thermally. We also comment on possible alternative DM realisations in which the gamma-ray bounds can be weaker due to phase space suppression. Depending upon the specific realisation and DM mass, gauged  $U(1)_{B-L}$  offers very interesting multi-messenger detection prospects at energy, intensity as well as cosmic frontiers [26–30].

This paper is organised as follows. In section II, we discuss our model followed by the details of the flux calculation in section III. We discuss our results in section IV and briefly discuss some other DM realisations to alleviate gamma-ray bounds in section V. We finally conclude in section VI.

## II. THE MODEL

We consider heavy right handed neutrino DM in a gauged  $B-L$  extension of the SM [31–36]. In its minimal realisation, the SM particle content is extended with three RHNs  $N_i$  ( $i = 1, 2, 3$ ) and one complex singlet scalar ( $\Phi$ ) all of which are singlet under the SM gauge symmetry. The requirement of triangle anomaly cancellation in this minimal model requires each of the RHNs to have  $B-L$  charge  $-1$ . The complex singlet scalar having  $B-L$  charge 2 not only leads to spontaneous breaking of gauge symmetry but also generate RHN masses or the seesaw scale dynamically. The Lagrangian involving RHN can be written as

$$\begin{aligned} \mathcal{L}_{\text{fermion}} = & i \sum_{\kappa=1}^3 \overline{N_{R_\kappa}} \not{D} N_{R_\kappa} - \sum_{\substack{j=1 \\ \alpha=e,\mu,\tau}}^3 Y_D^{\alpha j} \overline{l_L^\alpha} \tilde{H} N_R^j \\ & - \sum_{i,j=1}^3 Y_{N_{ij}} \Phi \overline{N_{R_i}^C} N_{R_j} + \text{h.c.}, \end{aligned} \quad (1)$$

where  $l_L, H$  denote the SM Lepton and Higgs doublets, respectively. The covariant derivative for  $N_{R_\kappa}$  is defined as

$$\not{D} N_{R_\kappa} = \gamma^\mu (\partial_\mu - i g_{\text{BL}} Z'_\mu) N_{R_\kappa}, \quad (2)$$

with  $g_{\text{BL}}, Z'$  being the coupling and gauge boson respectively for  $U(1)_{B-L}$ . After the singlet scalar  $\Phi$  acquires a non-zero vacuum expectation value (VEV) denoted by  $v_{\text{BL}}$ , the RHNs and  $Z'$  acquire masses as,

$$M_{Z'} = 2g_{\text{BL}}v_{\text{BL}}, \quad M_i = \sqrt{2}Y_{N_i}v_{\text{BL}}, \quad (3)$$

where we have assumed the Yukawa coupling  $Y_N$  to be diagonal. We consider the lightest RHN namely  $N_1$

to be the DM candidate such that the corresponding Yukawa coupling  $Y_D^{\alpha 1}$  to ensure a long lifetime. Due to such Yukawa interactions which also mixes  $N_1$  with the light neutrinos after electroweak symmetry breaking, we can have dominant tree-level decays like  $N_1 \rightarrow \nu, h; \nu, Z; l^\mp, W^\pm$  with  $h$  being the SM-like Higgs. Ignoring the masses of SM particles in comparison to  $N_1$  mass  $M_1$ , these decay widths are given as [37]

$$\begin{aligned} \Gamma(N_1 \rightarrow \nu_\alpha h(Z)) &= \frac{|Y_D^{\alpha 1}|^2 M_1}{32\pi}, \\ \Gamma(N_1 \rightarrow l_\alpha^- W^+) &= \frac{|Y_D^{\alpha 1}|^2 M_1}{16\pi}. \end{aligned} \quad (4)$$

We can also have a radiative decay  $N_1 \rightarrow \gamma \nu_\alpha$  as [38, 39]

$$\Gamma_{N_1 \rightarrow \gamma \nu_\alpha} \simeq \frac{9G_F^2 \alpha_{\text{EM}}}{1024\pi^4} \sin^2(\theta_{\alpha 1}) M_1^5, \quad (5)$$

where  $\alpha_{\text{EM}} = 1/137$  is the fine structure constant,  $G_F = 1.166 \times 10^{-5} \text{ GeV}^{-2}$  is the Fermi constant. The active-sterile mixing is given by  $\sin(\theta_{\alpha 1}) \sim \frac{Y_D^{\alpha 1} v}{\sqrt{2}M_1}$  with  $v$  being the VEV of the SM Higgs doublet. The lifetime of  $N_1$ , dictated by its tree-level two-body decays, can then be estimated as

$$\tau_{N_1} \approx 8 \times 10^{28} \text{ s} \left( \frac{440 \text{ PeV}}{M_1} \right) \left( \sum_\alpha \left| \frac{Y_D^{\alpha 1}}{4.7 \times 10^{-31}} \right|^2 \right)^{-1} \quad (6)$$

which is identified as the DM lifetime  $\tau_{\text{DM}}$  in our setup. The dominant tree level decay widths given in Eq. (4) have 2 : 1 : 1 ratio for  $l_L^\pm W^\mp, \nu Z, \nu h$  final states, respectively. Using the best-fit values of neutrino oscillation parameters for normal ordering [40], the branching ratio (BR) into different lepton flavours for a particular decay mode follows the ratio  $\text{BR}(e) : \text{BR}(\mu) : \text{BR}(\tau) = 0.68 : 0.08 : 0.24$ .

While DM of required mass can not be of thermal origin due to the violation of the unitarity bound [41], it can not be produced non-thermally via Yukawa portal interactions as well due to the suppressed Yukawa couplings in Eq. (6) required to ensure its longevity. However, due to additional interactions present in the  $B-L$  model which are unrelated to DM lifetime, it is still possible to produce the desired relic via the freeze-in mechanism [42–45].

Using the approach of [43, 44], we consider the production of superheavy  $N_1$  of mass 440 PeV from decay of heavier particle. The required non-thermal criteria forces the gauge coupling  $g_{\text{BL}}$  to be tiny keeping  $Z'$  out-of-equilibrium as well. We consider the singlet scalar to be in equilibrium sourcing  $Z'$  with the latter acting as source of DM. The relic can be calculated by solving the coupled Boltzmann equations written in terms of comov-

ing number densities  $Y_X = \frac{n_X}{s}$  as follows:

$$\begin{aligned} \frac{dY_{Z'}}{dx} &= \frac{\beta}{x\mathcal{H}} \Gamma_{\phi \rightarrow Z'Z'} \frac{K_1(M_\phi/T)}{K_2(M_\phi/T)} Y_\phi^{\text{eq}} \\ &\quad - \frac{\beta}{x\mathcal{H}} \Gamma_{Z' \rightarrow \text{all}} \frac{K_1(M_{Z'}/T)}{K_2(M_{Z'}/T)} Y_{Z'}, \\ \frac{dY_{N_1}}{dx} &= \frac{\beta}{x\mathcal{H}} \Gamma_{\phi \rightarrow N_1 N_1} \frac{K_1(M_\phi/T)}{K_2(M_\phi/T)} Y_\phi^{\text{eq}} \\ &\quad + \frac{\beta}{x\mathcal{H}} \Gamma_{Z' \rightarrow N_1 N_1} \frac{K_1(M_{Z'}/T)}{K_2(M_{Z'}/T)} Y_{Z'}. \end{aligned} \quad (7)$$

Here  $x = M_{Z'}/T$ ,  $\beta = 1 + \frac{Tdg_s/dT}{3g_s}$ , with  $g_s$  being the relativistic entropy degrees of freedom.  $\mathcal{H}$  is the Hubble expansion parameter,  $\Gamma$  denotes the decay width and  $K_i$ 's denote the modified Bessel functions of the second kind.

### III. NEUTRINO AND GAMMA-RAY FLUX FROM DM DECAY

In this section, we describe the method to compute the flux of high energy neutrinos and gamma-rays produced from DM decay. As discussed above, we consider the RHN  $N_1$  decay via all possible two-body decay modes namely,  $N_1 \rightarrow \nu, h(Z, \gamma)$  and  $N_1 \rightarrow \ell^\pm, W^\mp$ , emitting secondaries such as high energy neutrinos, gamma-rays and electrons. Note that we do not distinguish between particle and anti-particle final states here. The DM particle  $N_1$  is considered to be the constituent of the Milky Way DM halo. Thus, the decay of  $N_1$  in the Milky Way can give rise to backgrounds of high energy neutrinos and gamma-rays. The flux of these secondaries  $\phi_i(E_i)$  can be estimated as follows [46, 47]:

$$\frac{d^2\phi_i}{dE_i d\Omega} = \frac{\mathcal{D}}{4\pi M_{\text{DM}} \tau_{\text{DM}}} \frac{dN_i}{dE_i}, \quad (8)$$

where,  $E_i$  denotes the energy of the  $i$ th ( $\nu$  or  $\gamma$ ) particle and  $M_{\text{DM}} \equiv M_1$  is DM mass with lifetime denoted by  $\tau_{\text{DM}}$ . The dependence of the flux on DM density profile is given by [48],

$$\mathcal{D} = \frac{1}{\Delta\Omega} \int_{\Delta\Omega} d\Omega \int_0^{s_{\text{max}}} ds \rho_{\text{DM}}(s, b, l). \quad (9)$$

Here,  $\Delta\Omega$  is the angular region of observation,  $(l, b)$  are Galactic coordinates (longitude and latitude), and  $s$  is the line-of-sight coordinate. These coordinates are related as  $r(s, b, l) = \sqrt{s^2 + r_\odot^2 - 2sr_\odot \cos b \cos l}$ . The differential solid angle,  $d\Omega$  is given by  $d\Omega = (\sin b)dl$  and  $r_\odot = 8.3$  kpc is the distance to Milky Way centre from the Sun. The DM density,  $\rho_{\text{DM}}(s, b, l)$  can be obtained from various numerical models in the literature [49–53]. For this work, we consider the Navarro-Frenk-White (NFW) density profile [49] as a function of the Galactocentric radius,  $R_{\text{GC}}$  given by,

$$\rho_{\text{NFW}}(R_{\text{GC}}) = \frac{\rho_c}{(R_{\text{GC}}/R_c)(1 + R_{\text{GC}}/R_c)^2}, \quad (10)$$

where  $\rho_c$  is the DM density at the characteristic scale  $R_c = 11$  kpc, and its value is obtained by normalizing  $\rho_{\text{NFW}}$  with the DM density in the solar neighbourhood,  $\rho_\odot = 0.43$  GeV cm $^{-3}$ . Note that all the DM profiles in the literature nearly predict similar DM density except at the Galactic center. Therefore, the variation of the DM profile is only important for signals from the Galactic center direction. Since the arrival direction of the KM3-230213A event is far away from the Galactic center direction, we do not consider other DM profiles in this work.

The differential energy spectra,  $dN_i/dE_i$  in Eq. (8) is obtained from the publicly available code `HDMSpectra` [54]. These spectra depend on the DM mass and the inclusive decay rate of the DM particle.

### IV. RESULTS AND DISCUSSION

In this section, we discuss the results obtained by computing the fluxes of UHE neutrinos and gamma-rays from DM decay using Eq. (8) and analyze them in view of the KM3-230213A event. The mass of the DM particle  $N_1$  is chosen to be 440 PeV in order to match the energy requirement (220 PeV) of the KM3-230213A event. For this analysis, we take into account the constraints from three different gamma-ray observations. For energies below 1 PeV, we take the diffuse gamma-ray measurements from the inner Galactic plane by the Large High Altitude Air Shower Observatory (LHAASO) [55], whereas for energies above 40 PeV, we consider the upper limits on UHE photon flux from Moscow State University Extensive Air Shower (EAS-MSU) array [56] and PAO (PAO HECO +SD750) [57]. In addition to these gamma-ray constraints, we also consider the IceCube's High Energy Starting Events (HESE) [58].

We show the gamma-ray (left) and neutrino (right) fluxes together with these constraints in Fig. 1. The gamma-ray fluxes shown here are computed for the inner Galactic plane ( $15^\circ < l < 125^\circ$ ,  $-5^\circ < b < 5^\circ$ ) to remain consistent with the LHAASO data (shown by the purple data points). This flux is slightly larger (about a factor of  $\sim 1.25$ ) than the overall diffuse flux of the DM halo. For comparison, we also include the high energy diffuse gamma-ray limits, ESU-MSU and PAO HECO +SD750 shown by the black and magenta downward arrows, respectively. Given these constraints, the gamma-ray flux can be appropriately normalized by tuning the DM lifetime,  $\tau_{\text{DM}}$  (see Eq. (8)). To demonstrate this, the fluxes for two different values of  $\tau_{\text{DM}}$  are plotted in the left panel of Fig. 1. The flux corresponding to  $\tau_{\text{DM}} \approx 10^{29}$  s (red curve) remain consistent with the LHAASO data, however, it is in tension with the ESU-MSU and PAO HECO+SD750 limits. This tension can be relaxed by choosing a larger lifetime,  $\tau_{\text{DM}} \approx 10^{30}$  s as shown by the blue curve. Note that these gamma-ray fluxes can undergo absorption due to pair production losses on low energy background photons such as interstellar radiation




FIG. 1: Gamma-ray (left-panel) and single flavour neutrino (right-panel) fluxes with DM mass  $M_1 = 440$  PeV and two different values of DM lifetime  $\tau_{\text{DM}} = 10^{29}$  s (solid red),  $\tau_{\text{DM}} = 10^{30}$  s (solid blue). The fluxes are summed over the all possible decay modes. The data points on the left panel plot correspond to gamma-ray bounds from LHAASO [55], EAS-MSU [56] and PAO [57]. The data points on the right panel plot correspond to UHE neutrino events at IceCube [58] and KM3NeT [1]. The gray dotted contour (right panel) denotes IceCube’s sensitivity to UHE neutrino flux[59].

and CMB. However, this absorption is negligible above 10 PeV [60]. Apart from absorption, the fluxes can also be affected by inverse Compton radiation from secondary electrons. However, the inverse Compton gamma-ray flux has uncertainties as it depends on energy loss of electrons due to magnetic field [60]. Although we do not compute the inverse Compton flux explicitly, we note that this can enhance the gamma-ray flux by a numerical factor around 100 PeV energies.

For the two specific values of  $\tau_{\text{DM}}$ , we show the corresponding single flavour neutrino fluxes in the right panel of Fig. 1 following the same color coding as the left panel. The neutrino flavour ratio at Earth is assumed to be 1 : 1 : 1. We also show the flux of the KM3-230213A event by orange data point with  $3\sigma$  CL. The black data points and the gray dotted curve show the IceCube’s HESE events [58] and UHE neutrino flux sensitivity at 90% CL [59]. For the estimation of the neutrino fluxes, we consider the arrival direction (RA:  $94.3^\circ$ , Dec:  $-7.8^\circ$ ) of KM3-230213A with an angular uncertainty of  $3^\circ$  as reported by the KM3NeT collaboration [1]. The flux obtained with this conservative approach is about a factor of 2 smaller than the overall diffuse neutrino flux of the DM halo. Clearly, RHN DM with lifetime  $\tau_{\text{DM}} = 10^{29}$  s and mass  $M_1 = 440$  PeV can give rise to a flux close to the central value of the KM3-230213A event at 220 PeV energy. However, it also produces a large flux of secondary gamma-rays saturating the existing upper limits from LHAASO while being in tension with EAS-MSU and PAO data as discussed above. Lowering the decay width of DM or increasing the lifetime leads to agreement with gamma-ray bounds at the cost of reducing the neutrino flux while still being consistent at  $3\sigma$  CL, as depicted by the flux corresponding to  $\tau_{\text{DM}} \approx 10^{30}$  s. Interestingly, this neutrino flux can also alleviate the tension between KM3NeT and IceCube regarding the diffuse, isotropic origin of the KM3-230213A




FIG. 2: Evolution of  $Z'$  and DM  $N_1$  as a function of  $x = \frac{M_{Z'}}{T}$  for a benchmark point.

event. This tension arises as the central value of the KM3-230213A flux is larger than the IceCube’s sensitivity. While the present DM model can not simultaneously explain all UHE neutrino events including IceCube HESE and KM3-230213A, the former can be explained with astrophysical sources [58].

Finally, we show the viability of generating the observed relic of RHN DM of mass 440 PeV in this model by solving the coupled Boltzmann equations in Eq. (7). Fig. 2 shows the non-thermal production of  $Z'$  and DM  $N_1$  as a function of  $x = M_{Z'}/T$  for one particular benchmark point corresponding to  $M_{Z'} = 895$  PeV,  $M_\phi = 60000$  PeV, and DM mass  $M_{N_1} = 440$  PeV. We consider the  $B - L$  symmetry breaking scale to be  $v_{\text{BL}} = 4 \times 10^{19}$  GeV which implies  $g_{\text{BL}} \sim 10^{-11}$  and  $Y_{N_{11}} \sim 8 \times 10^{-12}$ . For this particular benchmark point, the production of DM from non-thermal  $Z'$  dominates over the production of DM from thermal  $\phi$ . The asymptotic comoving abundance of  $N_1$  gives rise to present DM relic  $\Omega h^2 = 0.12$  consistent with PLANCK data [61]. The required lifetime of DM can be obtained by appropriate choice of

Yukawa coupling  $Y_D^{\alpha 1}$  without affecting DM relic. We can also produce DM non-thermally from scattering of SM bath particles into  $N_1$  mediated by  $Z'$ , applicable when  $Z' \rightarrow N_1 N_1$  is kinematically forbidden. As shown in [45], we can get correct DM relic in such a case for  $v_{\text{BL}} \gtrsim 10^{13}$  GeV allowing sub-Planckian symmetry breaking scales.

## V. OTHER DARK MATTER SCENARIOS

Since RHN DM decays into neutrino and other SM particles via two-body decay with the same Yukawa coupling controlling all these partial decay widths, it is difficult to reduce the gamma-ray flux without reducing the neutrino flux. This is observed while comparing fluxes for two different  $\tau_{\text{DM}}$  values in Fig. 1 discussed in the previous section. One can make suitable changes to this model in order to decouple the gamma-ray and neutrino flux to some extent. Here we briefly outline two such possibilities.

### A. Singlet Scalar DM

We can consider another complex singlet scalar  $\Phi'$  added to the model such that  $M_{\Phi'} < M_i$ . It couples to RHNs as  $y_{\Phi'} \Phi' \bar{N}_i^c N_i$ . Unlike  $\Phi$ , this new singlet scalar does not acquire VEV and be a long-lived DM. Since RHNs mix with active neutrinos after electroweak symmetry breaking we can have scalar DM decay as  $\Phi' \rightarrow \nu\nu$ . In such a case DM decay to charged particles will arise only at three-body or higher final states, possibly giving a softer spectra for diffuse gamma-rays. Such charged final states arise via  $\phi \rightarrow \nu\ell^\pm W^\mp$ . One can also have production of other SM particles via similar three-body decay processes. Such a scalar singlet DM can be generated non-thermally via scalar portal or  $Z'$  portal interactions, similar to RHN DM discussed above.

### B. Doublet Scalar DM

Instead of considering Majorana neutrino, one can consider RHNs to be the right chiral parts  $\nu_R$  of sub-eV Dirac neutrinos. In this simplest version, they can acquire sub-eV neutrino masses from the SM Higgs itself with tiny Yukawa couplings. In such a case, the gauged  $B - L$  symmetry needs to be broken by a singlet scalar with  $B - L$  charge different from 2 such that Majorana mass terms are not dynamically generated.

Let us consider a neutrinophilic scalar doublet  $\eta$  which couples to neutrinos as  $y_\eta \bar{l}_L \tilde{\eta} \nu_R$ , where  $\nu_R$  is the right chiral part of light Dirac neutrinos. Assuming a  $Z_2$  symmetry in the scalar sector under which  $\eta \rightarrow -\eta$ , the scalar potential is similar to the inert doublet model given by

$$V = \mu_H^2 |H|^2 + \mu_\eta^2 |\eta|^2 + \lambda_1 |H|^4 + \lambda_2 |\eta|^4 + \lambda_3 |H|^2 |\eta|^2 + \lambda_4 |\eta^\dagger H|^2 + \lambda_5 [(\eta^\dagger H)^2 + \text{h.c.}] \quad (11)$$

After electroweak symmetry breaking, the components of  $\eta$  split and the lightest neutral real scalar  $\eta_R$  can be DM. Since the scalar sector has a  $Z_2$  symmetry and  $\mu_\eta^2 > 0$ , the neutral component of  $\eta$  does not acquire any VEV. Therefore, DM in this setup can decay only via the Yukawa interactions which break  $Z_2$  symmetry. We consider the corresponding Yukawa coupling  $y_\eta$  to be tuned such that DM is still long-lived on cosmological scales. Its decay  $\eta_R \rightarrow \nu_L \nu_R$  can give rise to the observed neutrino event while its decay into charged final states appear only via three-body decay like  $\eta_R \rightarrow \ell^\pm W^\mp \nu_R$  keeping the resulting secondary gamma-ray spectrum softer compared to the RHN DM studied earlier. While pure thermal scalar doublet DM will be overproduced for DM mass above the unitarity limit, incomplete thermalisation with suitable reheating temperature can generate the desired relic [62].

## VI. CONCLUSION

We have explored the possibility of generating the recently detected highest energy neutrino event of 220 PeV energy by the KM3NeT collaboration from decay of superheavy dark matter. Considering the simplest possibility of a right handed neutrino coupling with Yukawa coupling to leptons in type-I seesaw model to be the DM candidate, we calculate the flux of neutrino and gamma-ray by considering all possible two-body decay modes of DM. Considering the best-fit value of the neutrino flux reported by the KM3NeT collaboration, we find that the present gamma-ray constraints disfavour the DM decay origin of the event, in addition to the mild tension with non-observation of similar events at IceCube and PAO. Due to the large uncertainty associated with the KM3-230213A flux at present, it is still possible to explain the event at within  $3\sigma$  CL while being consistent with the above-mentioned constraints. As we consider two-body decay of DM, the preferred DM mass is 440 PeV, the relic of which can be generated non-thermally, if we embed the type-I seesaw scenario in a gauged  $U(1)_{B-L}$  setup where additional scalar and vector boson play the crucial role in DM production. We also outline the possibility of two more DM realisations where gamma-ray constraints can be made weaker. To conclude, the possibility of DM origin of the KM3-230213A event is marginal at present due to gamma-ray constraints and non observation of similar events at other experiments. More precise estimates of flux, multi-messenger observations or identification of astrophysical sources in future will be able to settle this hypothesis more concretely.

### Acknowledgments

The work of D.B. is supported by the Science and Engineering Research Board (SERB), Government of India grants MTR/2022/000575 and CRG/2022/000603. D.B.

also acknowledges the support from the Fulbright-Nehru Academic and Professional Excellence Award 2024-25. D.B. is thankful to the Department of Physics and Astronomy, University of Alabama, Tuscaloosa for kind hospitality during February 11-13, 2025 when this work was initiated. The work of N.D. is supported by the Min-

istry of Education, Government of India via the Prime Minister's Research Fellowship (PMRF) December 2021 scheme. The work of N.O. is supported in part by the United States Department of Energy Grant Nos. DE-SC0012447 and DE-SC0023713.

- 
- [1] **KM3NeT** Collaboration, S. Aiello *et al.*, *Observation of an ultra-high-energy cosmic neutrino with KM3NeT*, *Nature* **638** (2025), no. 8050 376–382.
- [2] **KM3NeT**, **MessMapp Group**, **Fermi-LAT**, **Owens Valley Radio Observatory 40-m Telescope Group**, **SVOM** Collaboration, O. Adriani *et al.*, *Characterising Candidate Blazar Counterparts of the Ultra-High-Energy Event KM3-230213A*, [arXiv:2502.08484](#).
- [3] **KM3NeT** Collaboration, O. Adriani *et al.*, *On the Potential Galactic Origin of the Ultra-High-Energy Event KM3-230213A*, [arXiv:2502.08387](#).
- [4] **KM3NeT** Collaboration, O. Adriani *et al.*, *On the potential cosmogenic origin of the ultra-high-energy event KM3-230213A*, [arXiv:2502.08508](#).
- [5] K. Greisen, *End to the cosmic-ray spectrum?*, *Phys. Rev. Lett.* **16** (Apr, 1966) 748–750.
- [6] G. T. Zatsepin and V. A. Kuz'min, *Upper Limit of the Spectrum of Cosmic Rays*, *Soviet Journal of Experimental and Theoretical Physics Letters* **4** (Aug., 1966) 78.
- [7] G. Gelmini, O. E. Kalashev, and D. V. Semikoz, *GZK photons as ultra high energy cosmic rays*, *J. Exp. Theor. Phys.* **106** (2008) 1061–1082, [[astro-ph/0506128](#)].
- [8] R. Aloisio, D. Boncioli, A. di Matteo, A. F. Grillo, S. Petreria, and F. Salamida, *Cosmogenic neutrinos and ultra-high energy cosmic ray models*, *JCAP* **10** (2015) 006, [[arXiv:1505.04020](#)].
- [9] G. B. Gelmini, O. Kalashev, and D. Semikoz, *Upper Limit on the Diffuse Radio Background from GZK Photon Observation*, *Universe* **8** (2022), no. 8 402, [[arXiv:2206.00408](#)].
- [10] S. Chakraborty, P. Mehta, and P. Sarmah, *A relook at the GZK neutrino-photon connection: impact of extra-galactic radio background & UHECR properties*, *JCAP* **01** (2024) 058, [[arXiv:2307.15667](#)].
- [11] S. W. Li, P. Machado, D. Naredo-Tuero, and T. Schwemberger, *Clash of the Titans: ultra-high energy KM3NeT event versus IceCube data*, [arXiv:2502.04508](#).
- [12] **KM3NeT** Collaboration, O. Adriani *et al.*, *The ultra-high-energy event KM3-230213A within the global neutrino landscape*, [arXiv:2502.08173](#).
- [13] M. S. Muzio, T. Yuan, and L. Lu, *Emergence of a neutrino flux above 5 PeV and implications for ultrahigh energy cosmic rays*, [arXiv:2502.06944](#).
- [14] K. Fang, F. Halzen, and D. Hooper, *Cascaded Gamma-ray Emission Associated with the KM3NeT Ultra-High-Energy Event KM3-230213A*, [arXiv:2502.09545](#).
- [15] **KM3NeT** Collaboration, O. Adriani *et al.*, *KM3NeT Constraint on Lorentz-Violating Superluminal Neutrino Velocity*, [arXiv:2502.12070](#).
- [16] P. Satunin, *Ultra-high-energy event KM3-230213A constraints on Lorentz Invariance Violation in neutrino sector*, [arXiv:2502.09548](#).
- [17] Y.-M. Yang, X.-J. Lv, X.-J. Bi, and P.-F. Yin, *Constraints on Lorentz invariance violation in neutrino sector from the ultra-high-energy event KM3-230213A*, [arXiv:2502.18256](#).
- [18] T. A. Dzhatdov, *The blazar PKS 0605-085 as the origin of the KM3-230213A ultra high energy neutrino event*, [arXiv:2502.11434](#).
- [19] A. Neronov, F. Oikonomou, and D. Semikoz, *KM3-230213A: An Ultra-High Energy Neutrino from a Year-Long Astrophysical Transient*, [arXiv:2502.12986](#).
- [20] A. Boccia and F. Iocco, *A strike of luck: could the KM3-230213A event be caused by an evaporating primordial black hole?*, [arXiv:2502.19245](#).
- [21] G. Amelino-Camelia, G. D'Amico, G. Fabiano, D. Frattulillo, G. Gubitosi, A. Moia, and G. Rosati, *On testing in-vacuo dispersion with the most energetic neutrinos: KM3-230213A case study*, [arXiv:2502.13093](#).
- [22] **IceCube** Collaboration, M. G. Aartsen *et al.*, *First observation of PeV-energy neutrinos with IceCube*, *Phys. Rev. Lett.* **111** (2013) 021103, [[arXiv:1304.5356](#)].
- [23] **IceCube** Collaboration, M. G. Aartsen *et al.*, *Evidence for High-Energy Extraterrestrial Neutrinos at the IceCube Detector*, *Science* **342** (2013) 1242856, [[arXiv:1311.5238](#)].
- [24] K. Murase, R. Laha, S. Ando, and M. Ahlers, *Testing the Dark Matter Scenario for PeV Neutrinos Observed in IceCube*, *Phys. Rev. Lett.* **115** (2015), no. 7 071301, [[arXiv:1503.04663](#)].
- [25] T. Cohen, K. Murase, N. L. Rodd, B. R. Safdi, and Y. Soreq,  *$\gamma$ -ray Constraints on Decaying Dark Matter and Implications for IceCube*, *Phys. Rev. Lett.* **119** (2017), no. 2 021102, [[arXiv:1612.05638](#)].
- [26] T. Bose *et al.*, *Report of the Topical Group on Physics Beyond the Standard Model at Energy Frontier for Snowmass 2021*, [arXiv:2209.13128](#).
- [27] N. Okada and O. Seto, *DAMPE excess from decaying right-handed neutrino dark matter*, *Mod. Phys. Lett. A* **33** (2018), no. 27 1850157, [[arXiv:1712.03652](#)].
- [28] B. Barman, A. Das, S. Jyoti Das, and M. Merchand, *Hunting for heavy  $Z'$  with IceCube neutrinos and gravitational waves*, [arXiv:2502.13217](#).
- [29] D. Borah, S. Jyoti Das, A. K. Saha, and R. Samanta, *Probing WIMP dark matter via gravitational waves' spectral shapes*, *Phys. Rev. D* **106** (2022), no. 1 L011701, [[arXiv:2202.10474](#)].
- [30] K. N. Abazajian and J. Heeck, *Observing Dirac neutrinos in the cosmic microwave background*, *Phys. Rev. D* **100** (2019) 075027, [[arXiv:1908.03286](#)].
- [31] A. Davidson,  *$B - L$  as the fourth color within an  $SU(2)_L \times U(1)_R \times U(1)$  model*, *Phys. Rev. D* **20** (1979)

- [32] R. N. Mohapatra and R. E. Marshak, *Local B-L Symmetry of Electroweak Interactions, Majorana Neutrinos and Neutron Oscillations*, *Phys. Rev. Lett.* **44** (1980) 1316–1319. [Erratum: *Phys.Rev.Lett.* 44, 1643 (1980)].
- [33] R. E. Marshak and R. N. Mohapatra, *Quark - Lepton Symmetry and B-L as the U(1) Generator of the Electroweak Symmetry Group*, *Phys. Lett. B* **91** (1980) 222–224.
- [34] A. Masiero, J. F. Nieves, and T. Yanagida, *B<sup>-</sup>l Violating Proton Decay and Late Cosmological Baryon Production*, *Phys. Lett. B* **116** (1982) 11–15.
- [35] R. N. Mohapatra and G. Senjanovic, *Spontaneous Breaking of Global B<sup>-</sup>l Symmetry and Matter - Antimatter Oscillations in Grand Unified Theories*, *Phys. Rev. D* **27** (1983) 254.
- [36] W. Buchmuller, C. Greub, and P. Minkowski, *Neutrino masses, neutral vector bosons and the scale of B-L breaking*, *Phys. Lett. B* **267** (1991) 395–399.
- [37] T. Higaki, R. Kitano, and R. Sato, *Neutrino in the Universe*, *JHEP* **07** (2014) 044, [[arXiv:1405.0013](#)].
- [38] P. B. Pal and L. Wolfenstein, *Radiative Decays of Massive Neutrinos*, *Phys. Rev. D* **25** (1982) 766.
- [39] R. E. Shrock, *Electromagnetic Properties and Decays of Dirac and Majorana Neutrinos in a General Class of Gauge Theories*, *Nucl. Phys. B* **206** (1982) 359–379.
- [40] I. Esteban, M. C. Gonzalez-Garcia, M. Maltoni, I. Martinez-Soler, J. a. P. Pinheiro, and T. Schwetz, *NuFit-6.0: updated global analysis of three-flavor neutrino oscillations*, *JHEP* **12** (2024) 216, [[arXiv:2410.05380](#)].
- [41] K. Griest and M. Kamionkowski, *Unitarity Limits on the Mass and Radius of Dark Matter Particles*, *Phys. Rev. Lett.* **64** (1990) 615.
- [42] L. J. Hall, K. Jedamzik, J. March-Russell, and S. M. West, *Freeze-In Production of FIMP Dark Matter*, *JHEP* **03** (2010) 080, [[arXiv:0911.1120](#)].
- [43] A. Biswas and A. Gupta, *Freeze-in Production of Sterile Neutrino Dark Matter in U(1)<sub>B-L</sub> Model*, *JCAP* **09** (2016) 044, [[arXiv:1607.01469](#)]. [Addendum: *JCAP* 05, A01 (2017)].
- [44] D. Borah, S. Jyoti Das, and A. K. Saha, *Cosmic inflation in minimal U(1)<sub>B-L</sub> model: implications for (non) thermal dark matter and leptogenesis*, *Eur. Phys. J. C* **81** (2021), no. 2 169, [[arXiv:2005.11328](#)].
- [45] N. Okada, S. Okada, and Q. Shafi, *Light Z' and dark matter from U(1)<sub>X</sub> gauge symmetry*, *Phys. Lett. B* **810** (2020) 135845, [[arXiv:2003.02667](#)].
- [46] K. Murase and J. F. Beacom, *Constraining Very Heavy Dark Matter Using Diffuse Backgrounds of Neutrinos and Cascaded Gamma Rays*, *JCAP* **10** (2012) 043, [[arXiv:1206.2595](#)].
- [47] P. Sarmah, N. Das, D. Borah, S. Chakraborty, and P. Mehta, *The Amaterasu particle: constraining the superheavy dark matter origin of UHECRs*, [[arXiv:2406.03174](#)].
- [48] T. N. Maity, A. K. Saha, A. Dubey, and R. Laha, *Search for dark matter using sub-PeV  $\gamma$ -rays observed by Tibet AS $\gamma$* , [[arXiv:2105.05680](#)].
- [49] J. F. Navarro, C. S. Frenk, and S. D. M. White, *A Universal density profile from hierarchical clustering*, *Astrophys. J.* **490** (1997) 493–508, [[astro-ph/9611107](#)].
- [50] J. F. Navarro, E. Hayashi, C. Power, A. Jenkins, C. S. Frenk, S. D. M. White, V. Springel, J. Stadel, and T. R. Quinn, *The Inner structure of Lambda-CDM halos 3: Universality and asymptotic slopes*, *Mon. Not. Roy. Astron. Soc.* **349** (2004) 1039, [[astro-ph/0311231](#)].
- [51] J. Einasto and U. Haud, *Galactic models with massive corona. I - Method. II - Galaxy*, *aap* **223** (Oct., 1989) 89–106.
- [52] A. W. Graham, D. Merritt, B. Moore, J. Diemand, and B. Terzic, *Empirical models for Dark Matter Halos. I. Nonparametric Construction of Density Profiles and Comparison with Parametric Models*, *Astron. J.* **132** (2006) 2685–2700, [[astro-ph/0509417](#)].
- [53] A. Burkert, *The Structure of Dark Matter Halos in Dwarf Galaxies*, *apjl* **447** (July, 1995) L25–L28, [[astro-ph/9504041](#)].
- [54] C. W. Bauer, N. L. Rodd, and B. R. Webber, *Dark matter spectra from the electroweak to the Planck scale*, *JHEP* **06** (2021) 121, [[arXiv:2007.15001](#)].
- [55] **LHAASO** Collaboration, Z. Cao *et al.*, *Measurement of Ultra-High-Energy Diffuse Gamma-Ray Emission of the Galactic Plane from 10 TeV to 1 PeV with LHAASO-KM2A*, *Phys. Rev. Lett.* **131** (2023), no. 15 151001, [[arXiv:2305.05372](#)].
- [56] Y. A. Fomin, N. N. Kalmykov, I. S. Karpikov, G. V. Kulikov, M. Y. Kuznetsov, G. I. Rubtsov, V. P. Sulakov, and S. V. Troitsky, *Constraints on the flux of  $\sim (10^{16} - 10^{17.5})$  eV cosmic photons from the EAS-MSU muon data*, *Phys. Rev. D* **95** (2017), no. 12 123011, [[arXiv:1702.08024](#)].
- [57] **Pierre Auger** Collaboration, A. Castellina, *Highlights from the Pierre Auger Observatory*, *PoS ICRC2019* (2021) 004, [[arXiv:1909.10791](#)].
- [58] **IceCube** Collaboration, R. Abbasi *et al.*, *The IceCube high-energy starting event sample: Description and flux characterization with 7.5 years of data*, *Phys. Rev. D* **104** (2021) 022002, [[arXiv:2011.03545](#)].
- [59] **IceCube** Collaboration, M. G. Aartsen *et al.*, *Differential limit on the extremely-high-energy cosmic neutrino flux in the presence of astrophysical background from nine years of IceCube data*, *Phys. Rev. D* **98** (2018), no. 6 062003, [[arXiv:1807.01820](#)].
- [60] A. Esmaili and P. D. Serpico, *Gamma-ray bounds from EAS detectors and heavy decaying dark matter constraints*, *JCAP* **10** (2015) 014, [[arXiv:1505.06486](#)].
- [61] **Planck** Collaboration, N. Aghanim *et al.*, *Planck 2018 results. VI. Cosmological parameters*, *Astron. Astrophys.* **641** (2020) A6, [[arXiv:1807.06209](#)]. [Erratum: *Astron.Astrophys.* 652, C4 (2021)].
- [62] N. Okada and O. Seto, *Superheavy WIMP dark matter from incomplete thermalization*, *Phys. Lett. B* **820** (2021) 136528, [[arXiv:2103.07832](#)].