

GROTHENDIECK RINGS OF LOCALIZATIONS OF THE INTEGERS AS ORDERED ABELIAN GROUPS

NEER BHARDWAJ[†] AND FRODO MOONEN

ABSTRACT. Let \mathbb{Z}_S be the ring generated by the inverses of all elements of a non-empty proper subset S of integer primes. We show that the ring generated by the values of unary definable sets in the model-theoretic Grothendieck ring of $(\mathbb{Z}_S; +, <)$ is a quotient of $(\mathbb{Z}/q\mathbb{Z})[T]/(T + T^2)$, where q is the largest odd integer that divides $p - 1$ for all $p \notin S$.

This implies that the Grothendieck ring of $(\mathbb{Z}_S; +, <)$ is trivial in various salient cases, for example when S is finite, or when S does not contain any prime of the form $2^n + 1$, $n \in \mathbb{N}$.

1. INTRODUCTION

The model-theoretic notion of a Grothendieck ring was introduced in [13, 9] and captures some interesting combinatorial and geometric properties of sets definable in a first-order structure. In this paper, we focus on computing the Grothendieck rings of subrings of \mathbb{Q} in the language of ordered abelian groups. The edge cases are known: the Grothendieck ring of $(\mathbb{Z}; +, <)$ is trivial [13], and for the o-minimal structure $(\mathbb{Q}; +, <)$ this ring is isomorphic to $\mathbb{Z}[T]/(T^2 + T)$ [12].

Grothendieck ring of a structure. For an \mathcal{L} -structure \mathcal{M} , we denote by $\text{Def}(\mathcal{M})$ the family of definable (with parameters) subsets of \mathcal{M} . The *Grothendieck group* of \mathcal{M} is the abelian group generated by symbols $[X]$, $X \in \text{Def}(\mathcal{M})$, with the following relations:

- $[X] = [Y]$ when X and Y are definably isomorphic,
- $[X \cup Y] = [X] + [Y]$ when $X \cap Y = \emptyset$.

Extending this group by multiplication defined by $[X][Y] = [X \times Y]$ we obtain the *Grothendieck ring* of \mathcal{M} ; denoted by $K_0(\mathcal{M})$. We will refer to $[X]$ as the *value* of the definable set X in $K_0(\mathcal{M})$.

Let S be a non-empty proper subset of the set of all primes in \mathbb{Z} . We study the first-order structures $(\mathbb{Z}_S; +, <)$, where \mathbb{Z}_S is the ring generated by inverting all elements of S . Note that every proper subring of \mathbb{Q} which properly contains \mathbb{Z} is given by such a localization of the ring of integers [1]. We establish the following in Section 2.

Theorem 1.1. *The Grothendieck ring of the structure $(\mathbb{Z}_S; +, <)$ is trivial if no odd prime divides $p - 1$ for every prime $p \notin S$.*

2020 *Mathematics Subject Classification.* Primary 03C07, 03C64, 06F20.

Key words and phrases. Archimedean ordered abelian groups, Grothendieck ring.

[†]Bhardwaj was partially funded by KU Leuven IF C16/23/010.

In Section 3, we show that the Grothendieck ring of $(\mathbb{Z}_S; +, <)$ is generated by values of sets definable by quantifier-free formulas in the language $(+, <)$. Also, we compute all possible values realized by unary definable sets in $(\mathbb{Z}_S; +, <)$ and generalize Theorem 1.1 as follows.

Theorem 1.2. *The ring generated by the Grothendieck ring values of unary definable sets in $(\mathbb{Z}_S; +, <)$ is a quotient of $(\mathbb{Z}/q\mathbb{Z})[T]/(T + T^2)$, where q is the largest odd integer which divides $p - 1$ for all $p \notin S$.*

Grothendieck rings often play a crucial role in motivic integration [11, 7, 15], and the structures of ordered abelian groups are especially relevant in this context. Apart from determining the Grothendieck ring, it is also an important theme to study definable sets up to definable bijection, sometimes even up to some special definable bijections, depending on the context. This leads to some finer classification results, see for example [2, 3, 6], but we do not address this finer challenge in this paper.

There has been much work to compute Grothendieck rings of various structures in the language of rings. It follows easily that the Grothendieck ring of real closed fields is given by \mathbb{Z} [13], and in contrast the same paper shows that the Grothendieck ring of the complex field embeds a polynomial ring over \mathbb{Z} generated by continuum many indeterminates. Cluckers and Haskell showed in [8] that the field of p -adic numbers is trivial. Moreover, the Grothendieck ring of formal Laurent series over p -adic numbers and formal Laurent series over local fields of strictly positive characteristic are also trivial by work of Cluckers [4]. See also [14] for another exact computation of the Grothendieck ring of a first-order structure.

Acknowledgements. Our foremost thanks goes to Pierre Touchard for suggesting this research direction for the second author's Masters dissertation topic. We also express our gratitude to Raf Cluckers for his support and guidance throughout this project. In addition, we are grateful to Pierre Touchard, Mathias Stout, and Floris Vermeulen for helpful discussions; indeed we attribute the idea for Lemma 2.5 to these discussions.

2. BASIC VALUES AND TORSION FROM NON-INVERTED PRIMES

In this section we compute Grothendieck ring values for certain \mathbb{Q} -intervals in \mathbb{Z}_S , which are sets of the form $(a, b) \cap \mathbb{Z}_S$, where $a, b \in \mathbb{Q}$ with $a < b$. When the context is clear, we use (a, b) to denote the \mathbb{Q} -interval $(a, b) \cap \mathbb{Z}_S$. These sets are clearly definable in $(\mathbb{Z}_S; +, <)$, and the main focus of this paper is to compute the values realized by \mathbb{Q} -intervals in the Grothendieck ring. We achieve this goal by adapting the results of [12, Claim 7] and using various number theoretic tricks.

Lemma 2.1. *The following hold for bounded \mathbb{Q} -intervals in \mathbb{Z}_S .*

- (i) $[(a, b)] = -1$ if $a, b \in \mathbb{Z}_S$.
- (ii) $[(a, b)] = -\frac{1}{2}$ for some $a \in \mathbb{Z}_S$ and $b \in \mathbb{Q} \setminus \mathbb{Z}_S$.

Proof. We first prove (i). For $a, b \in \mathbb{Z}_S$ we have that

$$[(a, \infty)] = [(a, b)] + [\{b\}] + [(b, \infty)].$$

Since $b - a \in \mathbb{Z}_S$, the map $x \rightarrow x + b - a$ gives a definable isomorphism from (a, ∞) to (b, ∞) . It follows that $[(a, b)] = -1$ as desired since the value of a singleton is 1.

For (ii), choose some n such that $1/n \in \mathbb{Z}_S$ and $1/(n+1) \notin \mathbb{Z}_S$. Such an n clearly exists since we assume that S is a non-empty proper subset of $\mathbb{N}^{\geq 2}$. We have that

$$\begin{aligned} [(0, 1)] &= [(0, \frac{1}{n+1})] + [(\frac{1}{n+1}, 1)] \\ &= [(0, \frac{1}{n+1})] + [(0, \frac{n}{n+1})] \\ &= 2 \cdot [(0, \frac{1}{n+1})] \end{aligned}$$

We use here that $x \rightarrow 1-x$ and $x \rightarrow x/n$ are definable isomorphisms. Now (i) implies that $[(0, \frac{1}{n+1})] = -\frac{1}{2}$. \square

Lemma 2.2. *Set $X := [(0, +\infty)]$. The following hold for unbounded \mathbb{Q} -intervals in \mathbb{Z}_S .*

- (i) $[(a, +\infty)] = [(-\infty, a)] = X$ for any $a \in \mathbb{Z}_S$.
- (ii) $[(-\infty, +\infty)] = 2X + 1$.
- (iii) $[(c, +\infty)] = [(-\infty, c)] = X + \frac{1}{2}$ for any $c \in \mathbb{Q} \setminus \mathbb{Z}_S$.

Proof. The assertions in (i), (ii) are immediate, and then (iii) follows by a direct application of Lemma 2.1(ii). \square

Lemma 2.3. *In the Grothendieck ring of $(\mathbb{Z}_S; +, <)$ we have that $X^2 + X = 0$.*

Proof. For $I := (0, \infty) \cap \mathbb{Z}_S$, we set $(f, g)_I = \{(x, y) \in I \times \mathbb{Z}_S \mid f(x) < y < g(x)\}$, where f and g functions in $(\mathbb{Z}_S; +, <)$, or $f, g \in \{-\infty, +\infty\}$, with $f(x) < g(x)$ for all $x \in I$. We split I^2 into three disjoint parts and obtain

$$[I^2] = [(0, \text{Id}_I)_I] + [\Gamma(\text{Id}_I)] + [(\text{Id}_I, \infty)_I], \quad (1)$$

where Id_I is the identity map on I and $\Gamma(\text{Id}_I)$ denotes the graph its graph. We work with three definable isomorphisms given as follows.

$$\begin{aligned} f : (0, \text{Id}_I)_I &\rightarrow (\text{Id}_I, \infty)_I : (x, y) \rightarrow (y, x) \\ g : I^2 &\rightarrow (\text{Id}_I, \infty)_I : (x, y) \rightarrow (x, x+y) \\ h : \Gamma(\text{Id}_I) &\rightarrow I : (x, x) \rightarrow x \end{aligned}$$

Hence we may rewrite (1) as

$$[I^2] = [I^2] + [I] + [I^2],$$

and it follows that $X^2 + X = 0$ as claimed. \square

We will use following simple fact in the proof of the lemma below.

Fact 2.4. *Suppose that $n \in \mathbb{N}$ with $n \geq 2$ is not divisible by any prime in S . Then for all $y \in \mathbb{Z}_S$, n divides exactly one of $y, y+1, \dots, y+n-1$ in \mathbb{Z}_S .*

Theorem 1.1 follows directly from the following result.

Lemma 2.5. *Consider a prime number p such that $p \notin S$. Then in the Grothendieck ring of $(\mathbb{Z}_S; +, <)$ we have that $q = 0$ where q is the largest odd factor of $p-1$.*

Proof. The idea is to partition $(-\infty, +\infty) \cap \mathbb{Z}_S$ into equivalence classes modulo p . For $i = 0, \dots, p-1$, set $N_i := \{x \in \mathbb{Z}_S : p \mid (x+i)\}$. Then we have

$$2X + 1 = [(-\infty, +\infty)] = \sum_{i=0}^{p-1} [N_i].$$

For each N_i we have a definable isomorphism with $(0, \infty)$ given by $x \rightarrow \frac{x+i}{p}$. This implies that

$$2X + 1 = p \cdot (2X + 1)$$

and hence $(p-1)(2X+1) = 0$. Multiplying by $2X+1$ and using Lemma 2.2 we get that $p-1 = 0$. Since 2 is invertible in the Grothendieck ring of \mathbb{Z}_S by Lemma 2.1(ii), the desired conclusion follows. \square

Corollary 2.6. *The Grothendieck ring of $(\mathbb{Z}_S; +, <)$ is trivial in the following cases:*

- *S contains only finitely many primes.*
- *S does not contain a prime of form $2^n + 1$, $n \in \mathbb{N}$. In other words, S does not contain 2 or a Fermat prime.*

Proof. The second assertion is immediate. For the first assertion, note that for any odd integer $q > 1$, $qm + 2$ contains infinitely many primes p by the Dirichlet prime number theorem and for any such prime p , we have that such that $q \nmid (p-1)$. \square

3. GROTHENDIECK RING VALUES OF UNARY DEFINABLE SETS

It is well-known that the ordered abelian group of integers admits quantifier elimination in the Presburger language $\langle (0, 1, +, <, (P_n)_{n \in \mathbb{N}^{\geq 1}}) \rangle$, where for every $n \in \mathbb{N}^{\geq 1}$, P_n is unary set given by $a \in P_n$ if and only if $\exists x(a = nx)$. In fact any ordered abelian group that does not have a convex definable subgroup also admits quantifier elimination in this language [5]; in this general setting 1 is defined to be the minimal positive element if such an element exists, and $1 = 0$ otherwise.

So we have for every S , that the structure $(\mathbb{Z}_S; 0, +, <, (P_n)_{n \in \mathbb{N}^{\geq 1}})$ has quantifier elimination; and we use this fact to transfer certain aspects of cell decomposition from $(\mathbb{Q}; +, <)$ to $(\mathbb{Z}_S; +, <)$. See [10, Chapter 3] for details about o-minimal cells and cell decomposition of definable cells.

Consider a definable subset $A \subset \mathbb{Z}_S^n$. Recall that Fact 2.4 gives that $\neg P_n(x)$ is equivalent to $\bigvee_{i=1}^{n-1} P_n(x+i)$ for all $n \in \mathbb{N}^{\geq 2}$ coprime to every prime in S . Using this together with quantifier elimination and usual arguments with the Presburger language we obtain $M \in \mathbb{N}^{\geq 1}$, which is coprime to every of prime in S , such that A is a disjoint union of sets defined by formulas $\phi_{(l_1, \dots, l_n)}^A$ of the following form.

$$\bigvee \left(\left(\bigwedge_i \left(\sum_j m_{i,j} x_j = g_i \right) \right) \wedge \left(\bigwedge_j P_M(x_j - l_j) \right) \wedge \left(\bigwedge_i \left(c_i < \sum_j e_{i,j} x_j < d_i \right) \right) \right),$$

where $l_1, \dots, l_n \in \{0, \dots, M-1\}$, $m_{i,j}, e_{i,j} \in \mathbb{Z}$, $g_i, c_i, d_i \in \mathbb{Z}_S$.

Clearly the map $x_i \rightarrow (Mz_j + l_j)$ gives a definable isomorphism set defined by $\phi_{(l_1, \dots, l_n)}^A$, to the set defined by the following formula

$$\bigvee \left(\left(\bigwedge_i \left(\sum_j m_{i,j} (Mz_j + l_j) = g_i \right) \right) \wedge \left(\bigwedge_i \left(c_i < \sum_j e_{i,j} (Mz_j + l_j) < d_i \right) \right) \right).$$

This formula has no divisibility conditions and hence the set it defines in \mathbb{Q}^n can be partitioned into cells. This implies that in the Grothendieck ring of $(\mathbb{Z}_S; +, <)$ we have that

$$[A] = \sum_{(l_1, \dots, l_n)} A_{(l_1, \dots, l_n)} = \sum_i [C_i \cap \mathbb{Z}_S^n],$$

where C_i are cells in $(\mathbb{Q}; +, <)$.

Definition. For C a cell in \mathbb{Q}^n , we say that $C \cap \mathbb{Z}_S^n$ is a *quasi-cell*. Clearly the quasi-cells are definable in $(\mathbb{Z}_S; +, <)$.

Proposition 3.1. *The Grothendieck ring of $(\mathbb{Z}_S; +, <)$ is generated by the values of quasi-cells.*

Note that the unary quasi-cells are finite subsets of \mathbb{Z}_S and the \mathbb{Q} -intervals. The next result is the final ingredient for the proof of 1.2.

Lemma 3.2. *Suppose there is an odd prime which divides $p - 1$ for all primes $p \notin S$. Then the following holds for \mathbb{Q} -intervals in \mathbb{Z}_S .*

- (i) $[(a, b)] = -\frac{1}{2}$ if $a \in \mathbb{Z}_S$ and $b \in \mathbb{Q} \setminus \mathbb{Z}_S$ or vice versa.
- (ii) $[(a, b)] = 0$ if $a, b \in \mathbb{Q} \setminus \mathbb{Z}_S$.
- (iii) $[(a, +\infty)] = [(-\infty, a)] = X + \frac{1}{2}$ if $a \in \mathbb{Q} \setminus \mathbb{Z}_S$.

Proof. We begin with the proof of (i). It suffices to consider the case of $a \in \mathbb{Z}_S$ and $b \in \mathbb{Q} \setminus \mathbb{Z}_S$.

By translating, we may assume that $a = 0$ and $b = m/n \notin \mathbb{Z}_S$ such that $b < 1$ and $m, n \in \mathbb{N}$ with $\gcd(m, n) = 1$. Since multiplying and dividing by elements of S is definable isomorphisms in $(\mathbb{Z}_S; +, <)$, we may assume without loss of generality that all prime factors of m and n are not in S . For $b < 1/2$, we have that

$$\begin{aligned} [(0, 1)] &= [(0, \frac{m}{n})] + [(\frac{m}{n}, \frac{n-m}{n})] + [(\frac{n-m}{n}, 1)] \\ &= 2[(0, \frac{m}{n})] + [(\frac{m}{n}, \frac{n-m}{n})]. \end{aligned}$$

We show for every $i, j \in \mathbb{N}$ with $0 < i < j < n$ and $\gcd(i, n) = \gcd(j, n) = 1$ that $[(\frac{i}{n}, \frac{j}{n})] = 0$. It then follows that $[(0, \frac{m}{n})] = -1/2$ as desired. We then also obtain the case $b > 1/2$ as follows

$$[(0, \frac{m}{n})] = [(0, \frac{n-m}{n})] + [(\frac{m-n}{n}, \frac{m}{n})] = [(0, \frac{n-m}{n})] = -1/2$$

also follows in an analogous manner.

Since \mathbb{Z}_S is dense in \mathbb{Q} , we have $c = c_1/c_2 \in \mathbb{Z}_S$, $\gcd(c_1, c_2) = 1$, such that $i/n < c < j/n$. Using definable isomorphisms we get that

$$\begin{aligned} [(\frac{i}{n}, \frac{j}{n})] &= [(\frac{i}{n}, c)] + [\{c\}] + [(c, \frac{j}{n})] \\ &= [(\frac{i}{n} - c, 0)] + [(0, \frac{j}{n} - c)] + 1 \\ &= [(-\frac{nc_1 - ic_2}{n}, 0)] + [(0, \frac{jc_2 - nc_1}{n})] + 1 \end{aligned} \tag{2}$$

Next we aim to find $Q \in S$ such that n divides $(Qj - i)$ in the integers. Since we can then translate by $T = -(Q - 1)c_1 + c_2(Qj - i)/n \in \mathbb{Z}$ and multiply by Q for

the first and second summand of equation (2) respectively to get our desired claim as follows.

$$\begin{aligned} \left[\left(\frac{i}{n}, \frac{j}{n}\right)\right] &= [(-Qc_1 + Q\frac{c_2j}{n}, T)] + [(0, Q\frac{jc_2 - nc_1}{n})] + 1 \\ &= [(0, T)] + 1 \\ &= 0 \end{aligned}$$

Here we used Lemma 2.1(ii). By our assumption, we have an odd prime q which divides $p - 1$ for all $s \notin S$, and we solve for prime Q satisfying

$$Qj - i \equiv 0 \pmod{n} \quad \text{and} \quad Q \equiv 2 \pmod{q}.$$

Note that $q \in S$ and hence in addition to i, j , we have that q is coprime to n as well. Then by the Chinese remainder theorem and Dirichlet's theorem on primes in arithmetic progressions we have that there is a prime Q satisfying the displayed divisibility conditions. Note such a Q necessarily belongs in S since q does not divide $Q - 1$. This finishes the proof for (i).

For (ii), we start with $c, d \in \mathbb{Q} \setminus \mathbb{Z}_S$, and take $a, b \in \mathbb{Z}_S$ with $a < c < d < b$. Then we have that

$$-1 = [(a, b)] = [(a, c)] + [(c, d)] + [(d, b)] = -\frac{1}{2} + [(c, d)] - \frac{1}{2}.$$

It follows that $[(c, d)] = 0$.

Assertion (iii) follows by a direct application of (i) and Lemma 2.2(i). \square

Proof of Theorem 1.2. By Proposition 3.1 and the comment just preceding that statement, we need to only consider the values realized by \mathbb{Q} -intervals.

If there is no odd prime which divides $p - 1$ for all $p \notin S$, then the Grothendieck ring of $(\mathbb{Z}_S; +, <)$ is trivial by Theorem 1.1. In the remaining case, the desired result follows immediately by Lemmas 2.1, 2.2, 2.3, 2.5, and 3.2.

REFERENCES

1. Ross A. Beaumont and H. S. Zuckerman, *A characterization of the subgroups of the additive rationals*, Pacific J. Math. **1** (1951), 169–177. MR 44522
2. R. Cluckers, *Classification of semi-algebraic p -adic sets up to semi-algebraic bijection*, J. Reine Angew. Math. **540** (2001), 105–114. MR 1868600
3. ———, *Presburger sets and p -minimal fields*, The Journal of Symbolic Logic **68** (2003), no. 1, 153–162.
4. ———, *Grothendieck rings of Laurent series fields*, J. Algebra **272** (2004), no. 2, 692–700. MR 2028076
5. R. Cluckers and I. Halupczok, *Quantifier elimination in ordered abelian groups*, Confluentes Math. **3** (2011), no. 4, 587–615. MR 2899905
6. ———, *Definable sets up to definable bijections in Presburger groups*, Trans. London Math. Soc. **5** (2018), no. 1, 47–70. MR 3747040
7. ———, *A p -adic variant of Kontsevich-Zagier integral operation rules and of Hrushovski-Kazhdan style motivic integration*, J. Reine Angew. Math. **779** (2021), 105–121. MR 4319066
8. R. Cluckers and D. Haskell, *Grothendieck rings of \mathbb{Z} -valued fields*, Bull. Symbolic Logic **7** (2001), no. 2, 262–269. MR 1839548
9. J. Denef and F. Loeser, *Definable sets, motives and p -adic integrals*, J. Amer. Math. Soc. **14** (2001), no. 2, 429–469. MR 1815218
10. L. van den Dries, *Tame topology and o -minimal structures*, London Mathematical Society Lecture Note Series, vol. 248, Cambridge University Press, Cambridge, 1998. MR 1633348
11. E. Hrushovski and D. Kazhdan, *Integration in valued fields*, Algebraic geometry and number theory, Progr. Math., vol. 253, Birkhäuser Boston, Boston, MA, 2006, pp. 261–405. MR 2263194

12. M. Kageyama and M. Fujita, *Grothendieck rings of o-minimal expansions of ordered abelian groups*, J. Algebra **299** (2006), no. 1, 8–20. MR 2225762
13. J. Krajíček and T. Scanlon, *Combinatorics with definable sets: Euler characteristics and Grothendieck rings*, Bull. Symbolic Logic **6** (2000), no. 3, 311–330. MR 1803636
14. E. Elbaz Saban, *Grothendieck ring of the pairing function without cycles*, ArXiv e-prints (2020).
15. M. Stout and F. Vermeulen, *Integration in Hensel minimal fields*, in preparation.

DEPARTMENT OF MATHEMATICS, KU LEUVEN, B-3001 LEUVEN, BELGIUM
Email address: nbhardwaj@msri.org

DEPARTMENT OF MATHEMATICS, KU LEUVEN, B-3001 LEUVEN, BELGIUM
Email address: frodo.moonen@student.kuleuven.be