THE L_p GAUSSIAN MINKOWSKI PROBLEM FOR *C*-PSEUDO-CONES

JUNJIE SHAN*, WENCHUAN HU

ABSTRACT. The L_p Gaussian Minkowski problem for *C*-pseudo-cones is studied in this paper, and the existence and uniqueness results are established. This extends our previous work on the Minkowski problem for *C*-pseudo-cones with respect to the Gaussian surface area measure (p = 1) and the Gaussian cone measure (p = 0).

1. INTRODUCTION

For convex bodies, the Minkowski problem aims to establish necessary and sufficient conditions for a Borel measure on the unit sphere to be the surface area measure. The L_p surface area measure is a generalization of the classical surface area measure. The L_p Minkowski problem, which characterizes the L_p surface area measure, was first introduced by Lutwak [22, 23], extending the classical Brunn-Minkowski theory into a broader framework. When p = 1, the L_p Minkowski problem reduces to the classical Minkowski problem. Consequently, many classical results in convex geometry become part of the more general L_p theory, such as the affine isoperimetric inequality [4, 12, 26], the Minkowski problem [3, 14, 18, 24, 25, 40]. Notably, for $p \ge 1$, many forms of the L_p Minkowski problem have been resolved. However, significant open questions remain for the case p < 1.

A special class of unbounded closed convex sets, known as C-pseudo-cones, has recently attracted considerable research interest. Within the copolarity framework, C-pseudo-cones are the counterparts of convex bodies with the origin in their interiors [2, 27, 31, 37]. Let $C \subset \mathbb{R}^n$ be a pointed (i.e., containing no lines), *n*-dimensional closed convex cone. A *pseudo-cone* is defined as a nonempty closed convex set K that does not contain the origin and satisfies $\lambda K \subseteq K$ for all $\lambda \geq 1$. The *recession cone* of K, denoted by rec K, is given by

$$\operatorname{rec} K = \{ z \in \mathbb{R}^n : K + z \subseteq K \}.$$

A *C*-pseudo-cone is a pseudo-cone *K* whose recession cone satisfies rec K = C. As $x \in \mathbb{R}^n$ tends to infinity, the shape of the *C*-pseudo-cone *K* is strongly governed by the cone *C*.

²⁰¹⁰ Math Subject Classifications: 52A40, 52A38.

Keywords: C-pseudo-cone, Gaussian measure, L_p Minkowski problem.

^{*} Corresponding author.

The *polar cone* of C, denoted by C° , is defined as

$$C^{\circ} = \{ x \in \mathbb{R}^n : \langle x, y \rangle \le 0 \text{ for all } y \in C \},\$$

we will write

$$\Omega_{C^{\circ}} = S^{n-1} \cap \operatorname{int} C^{\circ}.$$

Unlike convex bodies, the surface area measure of a C-pseudo-cone may be infinite. The Minkowski problem for the surface area measure and cone volume measure of C-pseudo-cones was first studied by Schneider [29, 30]. To generate finite measures, Schneider [33, 32] introduced the weighted Minkowski problem, where the weighted functions are homogeneous. Furthermore, Yang, Ye, and Zhu [38] later explored the L_p -Brunn-Minkowski theory for C-coconvex sets, while Li, Ye, and Zhu [19] developed dual versions. Further developments in the theory of pseudo-cones can be found in [1, 34, 36, 39].

In [35], the first author, the second author, and Xu introduced a class of nonhomogeneous measures—the Gaussian surface area measure and the Gaussian cone measure for *C*-pseudo-cones, and posed the corresponding Gaussian Minkowski problem. The Gaussian surface area measure of a *C*-pseudo-cone K in \mathbb{R}^n is defined on Ω_{C° as

$$S_{\gamma^n}(K,\eta) = \frac{1}{(\sqrt{2\pi})^n} \int_{\nu_K^{-1}(\eta)} e^{-\frac{|x|^2}{2}} d\mathcal{H}^{n-1}(x)$$
(1.1)

for every Borel set $\eta \subset \Omega_{C^{\circ}}$, where ν_K is the outer unit normal vector of K. The measure $S_{\gamma^n}(K, \cdot)$ is obviously finite.

Naturally, for any real number p, the L_p Gaussian surface area measure, as the L_p generalization of (1.1), can be defined as

$$S_{p,\gamma^n}(K,\eta) = \frac{1}{(\sqrt{2\pi})^n} \int_{\nu_K^{-1}(\eta)} |\langle x, \nu_K(x) \rangle|^{1-p} e^{-\frac{|x|^2}{2}} d\mathcal{H}^{n-1}(x).$$
(1.2)

In particular, the cases p = 1 and p = 0 correspond to the Gaussian surface area measure and the Gaussian cone measure, respectively, as introduced in our previous work [35].

It is worth noting that, in the case of convex bodies, the corresponding Gaussian Minkowski problem was studied in [15], while its L_p version was provided in [20]. Additional results on the Gaussian Minkowski problem for convex bodies can be found in [5, 8, 9, 17, 21].

In this paper, we consider the (normalized) L_p Gaussian Minkowski problem for C-pseudo-cones. For the existence theorem, we discuss two cases: p > 0 and p < 0. When p > 0, the result corresponds to a generalization of the case p = 1 in [35], but when p < 0, the situation is completely different. For p > 0, the corresponding result is **Theorem 1.1.** If p > 0, for any nonzero finite Borel measure μ on $\Omega_{C^{\circ}}$, there exists a C-pseudo-cone K such that

$$\mu = \frac{c}{p} S_{p,\gamma^n}(K, \cdot),$$

where $c = \frac{\int_{\Omega_{C^{\circ}}} \bar{h}_{K}^{p} d\mu}{\gamma^{n}(K)}$.

Here $\gamma^n(K)$ denotes the *n*-dimensional Gaussian measure of K and \bar{h}_K denotes the absolute support function of K. For the case p < 0, the existence of solutions to the (normalized) L_p Gaussian Minkowski problem can be stated as follows.

Theorem 1.2. If p < 0, for any nonzero finite Borel measure μ on $\Omega_{C^{\circ}}$, there exists a C-pseudo-cone K such that

$$\mu = -\frac{c}{p} S_{p,\gamma^n}(K, \cdot),$$

where $c = \frac{\int_{\Omega_{C^o}} \bar{h}_K^p d\mu}{\gamma^n(K)}$.

For the case p = 0, the limiting case of L_p theory, referred to as the Gaussian logarithmic Minkowski problem, has been studied in [35].

The class $\mathcal{K}(C, \omega)$ consists of C-pseudo-cones whose construction is determined by the compact set $\omega \subset \Omega_{C^{\circ}}$. Specifically, $K \in \mathcal{K}(C, \omega)$ implies

$$K = C \cap \bigcap_{u \in \omega} \{ x \in \mathbb{R}^n : \langle x, u \rangle \le h_K(u) \},\$$

where h_K denotes the support function of K. Under volume constraints, elements in $\mathcal{K}(C, \omega)$ satisfy the following uniqueness property:

Theorem 1.3. For $p \in (0,1]$, let $\omega \subset \Omega_{C^{\circ}}$ be a nonempty compact set. If $K, L \in \mathcal{K}(C, \omega)$ have the same L_p Gaussian surface area measure, i.e.,

$$S_{p,\gamma^n}(K,\cdot) = S_{p,\gamma^n}(L,\cdot),$$

and if $\gamma^n(K) = \gamma^n(L)$, then K = L.

For certain directional hyperplanes H_t , the Gaussian surface area density of $C \cap H_t$ tends to zero as $t \to 0^+$ and $t \to +\infty$, where t denotes the distance from the origin to the hyperplane. This asymptotic behavior implies that for any cone C and any compact set $\omega \subset \Omega_{C^\circ}$, one can always construct examples in $\mathcal{K}(C, \omega)$ such that the solution to the L_p Gaussian Minkowski problem loses uniqueness:

Theorem 1.4. If p < n, for any pointed, n-dimensional closed convex cone C in \mathbb{R}^n and any nonempty compact set $\omega \subset \Omega_{C^\circ}$, there exist distinct sets $K, L \in \mathcal{K}(C, \omega)$ such that their L_p Gaussian surface area measures coincide, i.e., $S_{p,\gamma^n}(K, \cdot) = S_{p,\gamma^n}(L, \cdot)$, but $K \neq L$.

2. Preliminaries

Let $(\mathbb{R}^n, \langle \cdot, \cdot \rangle)$ denote the *n*-dimensional Euclidean space with the standard inner product. A subset $C \subset \mathbb{R}^n$ is called a *cone* if it satisfies $\lambda x \in C$ for all $x \in C$ and $\lambda \geq 0$. A closed convex set $K \subset \mathbb{R}^n$ is termed a *pseudo-cone* if it satisfies $\lambda x \in K$ for all $x \in K$ and $\lambda \geq 1$. The recession cone of K, characterizing its asymptotic behavior, is defined as $\operatorname{rec} K = \{z \in \mathbb{R}^n : K + z \subseteq K\}$. For a pointed closed convex cone $C \subset \mathbb{R}^n$, a *C-pseudo-cone* refers to a pseudo-cone with $\operatorname{rec} K = C$. The dual cone is given by $C^\circ = \{x \in \mathbb{R}^n : \langle x, y \rangle \leq 0 \ \forall y \in C\}$. Throughout this paper, we assume that any cone C is pointed (i.e., it contains no lines) and *n*-dimensional.

Let S^{n-1} represent the unit sphere and B the open unit ball. For $u \in S^{n-1}$ and $t \in \mathbb{R}$, we define $H(u,t) = \{x \in \mathbb{R}^n : \langle x, u \rangle = t\}$ and its associated halfspace

$$H^{-}(u,t) = \{ x \in \mathbb{R}^{n} : \langle x, u \rangle \le t \}, \quad H^{+}(u,t) = \{ x \in \mathbb{R}^{n} : \langle x, u \rangle \ge t \}.$$

Fixing a unit vector $\mathfrak{v} \in \operatorname{int} C \cap \operatorname{int}(-C^\circ)$, we partition C via

$$C^{-}(t) = C \cap H^{-}(\mathfrak{v}, t), C^{+}(t) = C \cap H^{+}(\mathfrak{v}, t), \text{ and } C(t) = C \cap H(\mathfrak{v}, t)$$

for t > 0. Correspondingly, any C-pseudo-cone K is decomposed into $K(t) = K \cap C(t)$, $K^{-}(t) = K \cap C^{-}(t)$, and $K^{+}(t) = K \cap C^{+}(t)$.

For a C-pseudo-cone K, the support function $h_K : C^{\circ} \to \mathbb{R}$, defined as

$$h_K(x) = \sup\{\langle x, y \rangle : y \in K\},\$$

and it is bounded and non-positive. Its absolute support function is given by

$$\bar{h}_K = -h_K.$$

The radial function $\rho_K : \Omega_C \to \mathbb{R}$, where $\Omega_C = S^{n-1} \cap \operatorname{int} C$, given by

$$\varrho_K(v) = \min\{\lambda \in \mathbb{R} : \lambda v \in K\},\$$

generates the radial map $r_K = \rho_K(v)v$ and radial Gauss map $\alpha_K = \nu_K \circ r_K$, where ν_K denotes the outer unit normal on ∂K .

Convergence of C-pseudo-cones follows Schneider [31]:

Definition 2.1. A sequence K_i converges to K if there exists $t_0 > 0$ such that $K_i^-(t_0) \neq \emptyset$ for all i and $K_i^-(t) \rightarrow K^-(t)$ in Hausdorff metric for $t \ge t_0$.

The following selection theorem of C-pseudo-cones is essential:

Lemma 2.2. For a sequence K_i with uniform bounds $a < \text{dist}(o, K_i) < b$, there exists a convergent subsequence $K_{i_i} \to K$.

An integral transformation formula for C-pseudo-cones states that [33]:

Lemma 2.3. Let K be a C-pseudo-cone, for any nonnegative and Borel measurable or \mathcal{H}^{n-1} -integrable function $F : \partial_i K \to \mathbb{R}$:

$$\int_{\partial_i K} F(y) d\mathcal{H}^{n-1}(y) = \int_{\Omega_C} F(r_K(v)) \frac{\varrho_K^n(v)}{\bar{h}_K(\alpha_K(v))} dv$$
$$= \int_{\Omega_C} F(r_K(v)) \frac{\varrho_K^{n-1}(v)}{|\langle v, \alpha_K(v) \rangle|} dv,$$

where $\partial_i K := \partial K \cap \operatorname{int} C$.

In this paper, we define $\Omega := \Omega_{C^{\circ}} = S^{n-1} \cap \operatorname{int} C^{\circ}$. For any $u \in \Omega$, the support hyperplane of a C-pseudo-cone K is defined as

$$H_K(u) = \{x \in \mathbb{R}^n : \langle x, u \rangle = h_K(u)\}, \quad H_K^-(u) = \{x \in \mathbb{R}^n : \langle x, u \rangle \le h_K(u)\}.$$

A C-pseudo-cone K is C-determined by a compact set $\omega \subset \Omega = S^{n-1} \cap \operatorname{int} C^{\circ}$ if $K = C \cap \bigcap_{u \in \omega} H_{K}^{-}(u)$. The collection $\mathcal{K}(C, \omega)$ consists of such C-pseudo-cones that are C-determined by ω .

We denote

$$K^{(\omega)} = C \cap \bigcap_{u \in \omega} H_K^-(u)$$

for a C-pseudo-cone K. $K^{(\omega)} \in \mathcal{K}(C, \omega)$.

The Wulff shape construction provides critical examples: For compact $\omega \subset \Omega$ and positive continuous function $h : \omega \to \mathbb{R}_+$, the Wulff shape [h] associated with (C, ω, h) is defined as:

$$[h] = C \cap \bigcap_{u \in \omega} \{ y \in \mathbb{R}^n : \langle y, u \rangle \le -h(u) \}$$

belongs to $\mathcal{K}(C, \omega)$.

The following lemma, established in [33], will be used in our analysis:

Lemma 2.4. Let K_j be a sequence of *C*-pseudo-cones converging to *K*, *i.e.*, $K_j \to K$. For any nonempty compact set $\omega \subset \Omega$, then $K_i^{(\omega)}$ converge to $K^{(\omega)}$.

Let E be a convex set in \mathbb{R}^n . Its n-dimensional Gaussian measure $\gamma^n(E)$ is given by

$$\gamma^n(E) := \frac{1}{(2\pi)^{\frac{n}{2}}} \int_E e^{-\frac{|x|^2}{2}} dx$$

The Gaussian measure γ^n is log-concave with respect to convex sets, that is:

$$\gamma^n((1-t)K+tL) \ge \gamma^n(K)^{1-t}\gamma^n(L)^t$$

equality holds if and only if K = L. For further inequalities related to the Gaussian measure, we refer to [6, 7, 11, 16, 17, 22, 23]. For a comprehensive treatment of the theory of convex bodies, the classical texts [10] and [28] are recommended references.

3. The L_p Gaussian surface area measure

The primary aim of this section is to establish the L_p variational formula. For this purpose, we need the following lemma from Schneider [33]. Let $\Omega := \Omega_{C^{\circ}} = S^{n-1} \cap \operatorname{int} C^{\circ}$.

Lemma 3.1. Let $\omega \subset \Omega$ be a nonempty compact set, and a C-pseudo-cone $K \in \mathcal{K}(C, \omega)$. Given a continuous function $f : \omega \to \mathbb{R}$, then there exists a constant $\delta > 0$ which is small enough such that for any $|t| \leq \delta$, the function h_t defined as

$$\log h_t(u) = \log \bar{h}_K(u) + tf(u) + o(t, u), \quad u \in \omega,$$
(3.1)

where $o(t, \cdot) : \omega \to \mathbb{R}$ is a continuous function satisfying $\lim_{t\to 0} o(t, \cdot)/t = 0$ uniformly on ω . The Wulff shape $[h_t]$ associated with (C, ω, h_t) is termed a logarithmic family of Wulff shapes.

For almost all $v \in \Omega_C$, the derivative of the radial function is given by

$$\frac{\mathrm{d}\varrho_{[h_t]}(v)}{\mathrm{d}t}\bigg|_{t=0} = \lim_{t \to 0} \frac{\varrho_{[h_t]}(v) - \varrho_K(v)}{t} = f\left(\alpha_K(v)\right)\varrho_K(v).$$
(3.2)

The Gaussian covolume $V_G(K)$ of a C-pseudo-cone K is given in [35] by

Definition 3.2.

$$V_G(K) = \gamma^n(C \setminus K).$$

Recall the L_p Gaussian surface area measure $S_{p,\gamma^n}(K, \cdot)$ of a C-pseudo-cone K is defined as follows:

Definition 3.3. For every Borel set $\eta \subset \Omega$,

$$S_{p,\gamma^n}(K,\eta) := \frac{1}{(\sqrt{2\pi})^n} \int_{\nu_K^{-1}(\eta)} |\langle x, \nu_K(x) \rangle|^{1-p} e^{-\frac{|x|^2}{2}} d\mathcal{H}^{n-1}(x) = \int_{\eta} \bar{h}_K^{1-p}(u) dS_{\gamma^n}(K,u).$$

The following L_p variational formula serves as the foundation for solving the L_p Minkowski problem using variational methods. We begin by proving the variational formula for the Gaussian covolume. The variational approach originated in [13].

Lemma 3.4. If $K \in \mathcal{K}(C, \omega)$ for some nonempty compact set $\omega \subset \Omega$. For $p \neq 0$, let $f : \omega \to \mathbb{R}$ be a continuous function, and $\left[\left(\bar{h}_K(u)^p + tf(u)\right)^{\frac{1}{p}}\right]$ denotes the Wulff shapes associated with $\left(C, \omega, \left(\bar{h}_K(u)^p + tf(u)\right)^{\frac{1}{p}}\right)$, then

$$\lim_{t \to 0} \frac{V_G\left(\left[\left(\bar{h}_K(u)^p + tf(u)\right)^{\frac{1}{p}}\right]\right) - V_G(K)}{t} = \frac{1}{p} \int_{\omega} f(u) dS_{p,\gamma^n}(K, u).$$
(3.3)

Proof. For $p \neq 0$, let

$$h_t(u) = \left(\bar{h}_K(u)^p + tf(u)\right)^{\frac{1}{p}},$$

then

$$\log h_t = \log \bar{h}_K + \frac{tf}{p\bar{h}_K^p} + o(t, u), \qquad (3.4)$$

where $o(t, \cdot) : \omega \to \mathbb{R}$ is a continuous function satisfying $\lim_{t\to 0} o(t, \cdot)/t = 0$ uniformly on ω .

Using polar coordinates, we have

$$V_G([h_t]) = \frac{1}{(\sqrt{2\pi})^n} \int_{\Omega_C} \int_0^{\varrho_{[h_t]}(v)} e^{-\frac{r^2}{2}} r^{n-1} dr dv = \frac{1}{(\sqrt{2\pi})^n} \int_{\Omega_C} F_t(v) dv,$$

where we denote $F_t(v) = \int_0^{\varrho_{[h_t]}(v)} e^{-\frac{r^2}{2}} r^{n-1} dr$. Therefore,

$$\frac{F_t(v) - F_0(v)}{t} = \frac{1}{t} \int_{\varrho_K(v)}^{\varrho_{[h_t]}(v)} e^{-\frac{r^2}{2}} r^{n-1} dr$$
(3.5)

$$= \frac{\varrho_{[h_t]}(v) - \varrho_K(v)}{t} \cdot \frac{1}{\varrho_{[h_t]}(v) - \varrho_K(v)} \int_{\varrho_K(v)}^{\varrho_{[h_t]}(v)} e^{-\frac{r^2}{2}} r^{n-1} dr.$$
(3.6)

Moreover, the last term converges to $e^{-\frac{\varrho_K^2}{2}}\varrho_K^{n-1}$. By Lemma 3.1, we have

$$\frac{\mathrm{d}\varrho_{[h_t]}(v)}{\mathrm{d}t}\Big|_{t=0} = \lim_{t \to 0} \frac{\varrho_{[h_t]}(v) - \varrho_K(v)}{t} = \frac{f\left(\alpha_K(v)\right)\varrho_K(v)}{p\bar{h}_K^p(\alpha_K(v))},\tag{3.7}$$

it follows that

$$\lim_{t \to 0} \frac{F_t(v) - F_0(v)}{t} = e^{-\frac{\varrho_K^2}{2}} \frac{f(\alpha_K(v)) \,\varrho_K^n(v)}{p\bar{h}_K^p(\alpha_K(v))}$$
(3.8)

for almost $v \in \Omega_C$. Moreover, for sufficiently small |t|, there exists a constant M satisfying

$$\left|\frac{F_t(v) - F_0(v)}{t}\right| \le M.$$

Applying the dominated convergence theorem and Lemma 2.3, we obtain

$$\lim_{t \to 0} \frac{V_G\left([h_t]\right) - V_G(K)}{t} = \frac{1}{(\sqrt{2\pi})^n} \int_{\Omega_C} \lim_{t \to 0} \frac{F_t(v) - F_0(v)}{t} dv$$
$$= \frac{1}{(\sqrt{2\pi})^n} \int_{\Omega_C} e^{-\frac{\vartheta_K^2}{2}} \frac{f\left(\alpha_K(v)\right) \varrho_K^n(v)}{p\bar{h}_K^p(\alpha_K(v))} dv$$
$$= \int_{\Omega} f(u)\bar{h}_K(u)^{1-p} dS_{\gamma^n}(K, u)$$
$$= \int_{\omega} f(u) dS_{p,\gamma^n}(K, u).$$

Naturally, the L_p variational formula for the Gaussian volume is as follows.

Lemma 3.5. If $K \in \mathcal{K}(C, \omega)$ for some nonempty compact set $\omega \subset \Omega$. For $p \neq 0$, let $f : \omega \to \mathbb{R}$ be a continuous function, and $\left[\left(\bar{h}_K(u)^p + tf(u)\right)^{\frac{1}{p}}\right]$ denotes the Wulff shapes associated with $\left(C, \omega, \left(\bar{h}_K(u)^p + tf(u)\right)^{\frac{1}{p}}\right)$, then

$$\lim_{t \to 0} \frac{\gamma^n \left(\left[\left(\bar{h}_K(u)^p + tf(u) \right)^{\frac{1}{p}} \right] \right) - \gamma^n(K)}{t} = -\frac{1}{p} \int_{\omega} f(u) dS_{p,\gamma^n}(K, u).$$
(3.9)

Proof. This follows from Lemma 3.4 and the fact

$$\gamma^{n}\left(\left[\left(\bar{h}_{K}(u)^{p}+tf(u)\right)^{\frac{1}{p}}\right]\right)+V_{G}\left(\left[\left(\bar{h}_{K}(u)^{p}+tf(u)\right)^{\frac{1}{p}}\right]\right)=\gamma^{n}(K)+V_{G}(K)=\gamma^{n}(C).$$

By the definition of $S_{p,\gamma^n}K$ and the results established in our previous work [35] regarding the finiteness and weak convergence of the Gaussian surface area measure (p = 1), we conclude that the L_p Gaussian surface area measure is also a finite measure. Furthermore, it is weakly convergent in the sense of Definition 2.1 when restricted to the set $\mathcal{K}(C, \omega)$, for some nonempty compact set $\omega \subset \Omega$:

Lemma 3.6. For a nonempty compact set $\omega \subset \Omega$, and $K_j \in \mathcal{K}(C, \omega)$, $j \in \mathbb{N}$. If $K_j \to K$ as $j \to \infty$, then the measure $S_{p,\gamma^n}(K_j, \cdot)$ converge weakly to $S_{p,\gamma^n}(K, \cdot)$.

Proof. This follows from the support functions \bar{h}_{K_j} converge uniformly to \bar{h}_K and the weak convergence of the Gaussian surface area measure.

4. EXISTENCE OF SOLUTIONS WHEN p > 0

We first need to establish a uniform estimate for the decay properties of the Gaussian measure.

Lemma 4.1.

$$\gamma^n(\mathbb{R}^n - rB) \le \frac{2n\sqrt{n}}{(2\pi)^{\frac{1}{2}}r}e^{-\frac{r^2}{2n}}.$$

Proof. If t > 0,

$$\begin{split} \gamma^{1}(x > t) &= \frac{1}{(2\pi)^{\frac{1}{2}}} \int_{t}^{+\infty} e^{-\frac{r^{2}}{2}} dr \\ &= \frac{1}{(2\pi)^{\frac{1}{2}}} \int_{t}^{+\infty} \frac{1}{r} \cdot r e^{-\frac{r^{2}}{2}} dr \\ &= \frac{1}{(2\pi)^{\frac{1}{2}}} \int_{t}^{+\infty} -\frac{1}{r} d(e^{-\frac{r^{2}}{2}}) \\ &= \frac{1}{(2\pi)^{\frac{1}{2}}} \left(-\frac{1}{r} e^{-\frac{r^{2}}{2}} \Big|_{t}^{+\infty} -\int_{t}^{+\infty} \frac{1}{r^{2}} e^{-\frac{r^{2}}{2}} dr \right) \\ &= \frac{1}{(2\pi)^{\frac{1}{2}}} \left(\frac{1}{t} e^{-\frac{t^{2}}{2}} - \int_{t}^{+\infty} \frac{1}{r^{2}} e^{-\frac{r^{2}}{2}} dr \right) \\ &\leq \frac{1}{(2\pi)^{\frac{1}{2}}} \cdot \frac{1}{t} e^{-\frac{t^{2}}{2}}. \end{split}$$

For a ball rB of radius r in \mathbb{R}^n , let P_r be its inscribed cube. Then, the edge length of P_r is $\frac{2r}{\sqrt{n}}$. Since $P_r \subset rB$, then $\mathbb{R}^n - rB \subset \mathbb{R}^n - P_r$, we conclude

$$\gamma^{n}(\mathbb{R}^{n} - rB) \leq \gamma^{n}(\mathbb{R}^{n} - P_{r})$$

$$\leq \gamma^{n}(|x_{1}| > \frac{r}{\sqrt{n}}) + \gamma^{n}(|x_{2}| > \frac{r}{\sqrt{n}}) + \dots + \gamma^{n}(|x_{n}| > \frac{r}{\sqrt{n}})$$

$$= 2n\gamma^{1}(x > \frac{r}{\sqrt{n}})$$

$$\leq 2n \cdot \frac{1}{(2\pi)^{\frac{1}{2}}} \cdot \frac{1}{\frac{r}{\sqrt{n}}} e^{-\frac{\left(\frac{r}{\sqrt{n}}\right)^{2}}{2}}$$

$$= \frac{2n\sqrt{n}}{(2\pi)^{\frac{1}{2}}r} e^{-\frac{r^{2}}{2n}}.$$

Next, we hope to transform the L_p Gaussian Minkowski problem into an optimization problem. When p > 0, the method for solving this problem is similar to [35], but with modifications.

We assume $\omega \subset \Omega$ is a nonempty compact subset. For any nonzero finite Borel measure μ on ω , we define the set $C^+(\omega)$ as the collection of continuous functions $f: \omega \to (0, \infty)$. If p > 0, we introduce a functional $I_{\mu}: C^+(\omega) \to (0, \infty)$ given by

$$I_{\mu}(f) := \gamma^{n}([f]) \int_{\omega} f^{p} d\mu$$

for each $f \in C^+(\omega)$, where [f] represents the Wulff shape associated with (C, ω, f) .

Given p > 0, the following optimization problem is then considered:

$$\sup\{I_{\mu}(f) = \gamma^{n}([f]) \int_{\omega} f^{p} d\mu : f \in C^{+}(\omega)\}.$$

Obviously, for any $K \in \mathcal{K}(C, \omega)$, $I_{\mu}(\bar{h}_K) = \gamma^n(K) \int_{\omega} \bar{h}_K^p d\mu > 0$.

If I_{μ} attains a maximum at \bar{h}_{K} for some $K \in \mathcal{K}(C, \omega)$, for any continuous function $f : \omega \to \mathbb{R}$, the function $(\bar{h}_{K}^{p} + tf)^{\frac{1}{p}}$ belongs to $C^{+}(\omega)$ for sufficiently small |t|. By Lemma 3.5, we obtain

$$0 = \frac{d}{dt}\Big|_{t=0} I_{\mu}(\left(\bar{h}_{K}^{p} + tf\right)^{\frac{1}{p}}) = \gamma^{n}(K) \int_{\omega} f d\mu - \frac{1}{p} \int_{\omega} f dS_{p,\gamma^{n}}(K, \cdot) \int_{\omega} \bar{h}_{K}^{p} d\mu.$$

In other words,

$$\gamma^{n}(K) \int_{\omega} f d\mu = \frac{1}{p} \int_{\omega} f dS_{p,\gamma^{n}}(K, \cdot) \int_{\omega} \bar{h}_{K}^{p} d\mu$$

holds for all continuous functions f on ω . Then we conclude that

$$\mu = \frac{\int_{\omega} \bar{h}_K^p d\mu}{p\gamma^n(K)} S_{p,\gamma^n}(K, \cdot).$$
(4.1)

Therefore, in what follows, we will show that I_{μ} indeed attains its maximum at support function \bar{h}_K for some $K \in \mathcal{K}(C, \omega)$. From the preceding discussion, this is equivalent to K being a solution to the (normalized) L_p Gaussian Minkowski problem.

Lemma 4.2. If p > 0, suppose $\omega \subset \Omega$ is a nonempty compact subset. For any nonzero finite Borel measure μ on ω , there exists a C-pseudo-cone $K \in \mathcal{K}(C, \omega)$ such that

$$\mu = \frac{\int_{\omega} h_K^p d\mu}{p\gamma^n(K)} S_{p,\gamma^n}(K,\cdot).$$

Proof. For any $f \in C^+(\omega)$, we have $\bar{h}_{[f]} \geq f$. Consequently,

$$I_{\mu}(f) = \gamma^{n}([f]) \int_{\omega} f^{p} d\mu \leq \gamma^{n}([f]) \int_{\omega} \bar{h}^{p}_{[f]} d\mu = I_{\mu}(\bar{h}_{[f]}).$$

This shows that the supremum of $I_{\mu}(f)$ is attained by the support functions of $\mathcal{K}(C,\omega)$. Given a sequence $\{\bar{h}_{K_i}\}$, where each $K_i \in \mathcal{K}(C,\omega)$, satisfying

$$\lim_{i \to \infty} I_{\mu}(\bar{h}_{K_i}) = \sup \left\{ I_{\mu}(f) : f \in C^+(\omega) \right\}.$$

Define

 $r_i = \min\{r : rB \cap K_i \neq \emptyset\},\$

then we have

$$h_{K_i} \leq r_i$$

and

$$K_i \subset \mathbb{R}^n - r_i B.$$

If there exists a subsequence, still denoted by r_i , such that $r_i \to \infty$, this leads to the estimate by Lemma 4.1:

$$\begin{split} I_{\mu}(\bar{h}_{K_{i}}) &= \gamma^{n}(K_{i}) \int_{\omega} \bar{h}_{K_{i}}^{p} d\mu \\ &\leq \gamma^{n}(\mathbb{R}^{n} - r_{i}B)\mu(\omega)r_{i}^{p} \\ &\leq \frac{c}{r_{i}}e^{-\frac{r_{i}^{2}}{2n}} \cdot r_{i}^{p} \to 0, \end{split}$$

for some constant c. This contradicts the fact that $\lim_{i\to\infty} I_{\mu}(\bar{h}_{K_i}) = \sup I_{\mu}$, since $I_{\mu}(\bar{h}_K) > 0$ for each $K \in \mathcal{K}(C, \omega)$.

Moreover, assume that $r_i \to 0$, then we have

$$\gamma^n(\mathbb{R}^n - r_i B) \to 1,$$

and through the same estimate, we obtain

$$I_{\mu}(\bar{h}_{K_{i}}) = \gamma^{n}(K_{i}) \int_{\omega} \bar{h}_{K_{i}}^{p} d\mu$$

$$\leq \gamma^{n}(\mathbb{R}^{n} - r_{i}B)\mu(\omega)r_{i}^{p} \to 0.$$

Similarly, we arrive at a contradiction.

This implies the uniform estimate $0 < c_1 < \text{dist}(o, \partial K_i) < c_2$, for some constants c_1 and c_2 . From Lemma 2.2, we can find a subsequence K_{i_j} , such that $K_{i_j} \to K$ for some $K \in \mathcal{K}(C, \omega)$. Consequently, we get $\lim_{i\to\infty} I_{\mu}(\bar{h}_{K_{i_j}}) = I_{\mu}(\bar{h}_K) = \sup I_{\mu}(f)$. By the preceding discussion (4.1), the result is proved.

If μ is a nonzero Borel measure on Ω , p > 0, we may approximate Ω by an increasing sequence of compact sets $\{\omega_i\}$, i.e., $\omega_i \subset \operatorname{int} \omega_{i+1}$ and $\bigcup_{i \in \mathbb{N}} \omega_i = \Omega$. For each $i \in \mathbb{N}$, denote $\mu_i = \mu \sqcup \omega_i$ as

$$\mu\llcorner\omega_i(\sigma):=\mu(\sigma\cap\omega_i)$$

for any Borel subset $\sigma \subset \Omega$. We further require that $\mu_0 = \mu \llcorner \omega_0$ is a nonzero measure. By Lemma 4.2, for each $i \in \mathbb{N}$, there exists a *C*-pseudo-cone $K_i \in \mathcal{K}(C, \omega_i)$ satisfying

$$\mu_i = \frac{\int_{\omega_i} h_{K_i}^p d\mu_i}{p\gamma^n(K_i)} S_{p,\gamma^n}(K_i, \cdot).$$
(4.2)

To establish the existence of solutions to the L_p Gaussian Minkowski problem on Ω , it remains to show that the sets $\{K_i\}$ are uniformly bounded, i.e., there exist positive uniform lower and upper bounds for dist $(o, \partial K_i)$. This uniform boundedness will allow us to extract a convergent subsequence, ultimately yielding the desired solution on Ω via an approximation argument. The uniform estimate as follows.

Lemma 4.3. For the sequence $\{K_i\}$ in (4.2), there exist two constant m and M such that

$$0 < m < \operatorname{dist}(o, \partial K_i) < M$$

for all $i \in \mathbb{N}$.

Proof. Recall the optimization problems

$$I_{\mu_i}(\bar{h}_{K_i}) = \sup\left\{I_{\mu_i}(f) = \gamma^n([f])\int_{\omega_i} f^p d\mu_i : f \in C^+(\omega_i)\right\}.$$

Since $\mu_i \leq \mu_{i+1} \leq \mu$, $\omega_i \subset \operatorname{int} \omega_{i+1}$ and $\mathcal{K}(C, \omega_i) \subset \mathcal{K}(C, \omega_{i+1})$, we get

$$0 < I_{\mu_{i}}(\bar{h}_{K_{i}}) = \gamma^{n}([\bar{h}_{K_{i}}|_{\omega_{i}}]) \int_{\omega_{i}} \bar{h}_{K_{i}}^{p} d\mu_{i}$$

$$= \gamma^{n}([\bar{h}_{K_{i}}|_{\omega_{i+1}}]) \int_{\omega_{i}} \bar{h}_{K_{i}}^{p} d\mu_{i}$$

$$\leq \gamma^{n}([\bar{h}_{K_{i}}|_{\omega_{i+1}}]) \int_{\omega_{i+1}} \bar{h}_{K_{i}}^{p} d\mu_{i+1}$$

$$\leq I_{\mu_{i+1}}(\bar{h}_{K_{i+1}}).$$

If we set

 $r_i = \min\{r : rB \cap K_i \neq \emptyset\},\$

Then we also have

$$h_{K_i} \leq r_i$$
 and $K_i \subset \mathbb{R}^n - r_i B$

as Lemma 4.2. If $r_i \to \infty$, then

$$I_{\mu_i}(\bar{h}_{K_i}) = \gamma^n(K_i) \int_{\omega_i} \bar{h}_{K_i}^p d\mu_i$$

$$\leq \gamma^n(\mathbb{R}^n - r_i B) \mu_i(\omega_i) r_i^p$$

$$\leq \gamma^n(\mathbb{R}^n - r_i B) \mu(\Omega) r_i^p$$

$$\leq \frac{d}{r_i} e^{-\frac{r_i^2}{2n}} \cdot r_i^p \to 0,$$

for some constant d. This contradicts the monotonicity property of $I_{\mu_i}(\bar{h}_{K_i})$, namely $0 < I_{\mu_i}(\bar{h}_{K_i}) \leq I_{\mu_{i+1}}(\bar{h}_{K_{i+1}}) \leq \cdots$. Meanwhile, if $r_i \to 0$, then

$$I_{\mu_i}(\bar{h}_{K_i}) = \gamma^n(K_i) \int_{\omega_i} \bar{h}_{K_i}^p d\mu_i$$

$$\leq \gamma^n(\mathbb{R}^n - r_i B) \mu(\Omega) r_i^p \to 0,$$

this also arrive at a contradiction.

By the uniform estimate and standard approximation argument, the (normalized) L_p Gaussian Minkowski theorem can be established.

Theorem 4.4. If p > 0, for any nonzero finite Borel measure μ on Ω , there exists a C-pseudo-cone K such that

$$\mu = \frac{\int_{\Omega} h_K^p d\mu}{p\gamma^n(K)} S_{p,\gamma^n}(K,\cdot).$$

Proof. For the sequence $\{K_i\}$ in (4.2) and the uniform estimate of dist $(o, \partial K_i)$, we can extract a convergent subsequence $K_i \to K$.

Fix $i \in \mathbb{N}$, choose a nonempty compact set $\beta \subset \Omega$ with $\omega_i \subset \operatorname{int} \beta$. Recall the set $K^{(\beta)} \in \mathcal{K}(C,\beta)$ is given by

$$K^{(\beta)} := C \cap \bigcap_{u \in \beta} H_K^-(u).$$

By Lemma 13 in [33], we have

$$\nu_{K_j}^{-1}(\omega_i) = \nu_{K_j^{(\beta)}}^{-1}(\omega_i), \quad \nu_K^{-1}(\omega_i) = \nu_{K^{(\beta)}}^{-1}(\omega_i),$$

Moreover, by Lemma 2.4, it follows that

$$K_i^{(\beta)} \to K^{(\beta)}.$$

From Lemma 3.6, the measure $S_{p,\gamma^n}(K_j) \sqcup \omega_i$ converges weakly to $S_{p,\gamma^n}(K^{(\beta)}) \sqcup \omega_i$. Consequently, we obtain

$$c_j S_{p,\gamma^n}(K_j,\cdot) \llcorner \omega_i \xrightarrow{w} c S_{p,\gamma^n}(K,\cdot) \llcorner \omega_i,$$

where

$$c_i = \frac{\int_{\omega_i} h_{K_i}^p d\mu_i}{p\gamma^n(K_i)}$$
 and $c = \frac{\int_{\Omega} \bar{h}_K^p d\mu}{p\gamma^n(K)}$

Since $\mu \llcorner \omega_i = c_i S_{p,\gamma^n}(K_i, \cdot) \llcorner \omega_i$, the equality

$$cS_{p,\gamma^n}(K,\sigma) = cS_{p,\gamma^n}(K^{(\beta)},\sigma) = \mu_i(\sigma) = \mu(\sigma)$$

holds for every Borel subset $\sigma \subset \omega_i$. Furthermore, since $\bigcup_{j \in \mathbb{N}} \omega_j = \Omega$, we deduce that $cS_{p,\gamma^n}(K, \cdot) = \mu$.

Remark 4.5. For p < 0, the corresponding optimization problems

$$\sup\{I_{\mu}(f) = \gamma^{n}([f]) \int_{\omega} f^{p} d\mu : f \in C^{+}(\omega)\}$$

$$(4.3)$$

or

$$\inf\{I_{\mu}(f) = \gamma^{n}([f]) \int_{\omega} f^{p} d\mu : f \in C^{+}(\omega)\}$$

$$(4.4)$$

are both invalid for the following reasons:

(i) For (4.3), since $p < 0, f^p \ge \bar{h}_{[f]}^p$, in other words, $I_{\mu}(f) \ge I_{\mu}(\bar{h}_{[f]})$.

(ii) For (4.4), if the function f sufficiently big, $f \to +\infty$, then $\gamma^n([f]) \to 0$ and $\int_{\omega} f^p d\mu \to 0$. Therefore, $\inf I_{\mu}(f) = 0$, the variational argument fails.

5. EXISTENCE OF SOLUTIONS WHEN p < 0

If p < 0, for a nonempty compact subset $\omega \subset \Omega$ and any nonzero Borel measure μ on ω , unlike the case for p > 0 (see Remark 4.5), we consider a new variational functional $\phi_{\mu} : C^{+}(\omega) \to (0, \infty)$ given by

$$\phi_{\mu}(f) := \frac{\gamma^n([f])}{\int_{\omega} f^p d\mu}$$

and the following optimization problem:

$$\sup\{\phi_{\mu}(f) = \frac{\gamma^{n}([f])}{\int_{\omega} f^{p} d\mu} : f \in C^{+}(\omega)\}.$$

Obviously, for any $f \in C^+(\omega)$, we have $\bar{h}_{[f]}^p \leq f^p$. Then

$$\phi_{\mu}(f) = \frac{\gamma^{n}([f])}{\int_{\omega} f^{p} d\mu} \leq \frac{\gamma^{n}([f])}{\int_{\omega} \bar{h}^{p}_{[f]} d\mu} = \phi_{\mu}(\bar{h}_{[f]}).$$

And for each $K \in \mathcal{K}(C, \omega), \ \phi_{\mu}(\bar{h}_{K}) = \frac{\gamma^{n}(K)}{\int_{\omega} h_{K}^{p} d\mu} > 0.$

If ϕ_{μ} attains a maximum at \bar{h}_{K} for some $K \in \mathcal{K}(C, \omega)$, for any continuous function $f: \omega \to \mathbb{R}$, the function $(\bar{h}_{K}^{p} + tf)^{\frac{1}{p}}$ belongs to $C^{+}(\omega)$ for sufficiently small |t|. By Lemma 3.5, we obtain

$$0 = \left. \frac{d}{dt} \right|_{t=0} \phi_{\mu}(\left(\bar{h}_{K}^{p} + tf\right)^{\frac{1}{p}}) = \frac{-\frac{1}{p} \int_{\omega} f dS_{p,\gamma^{n}}(K, \cdot) \int_{\omega} \bar{h}_{K}^{p} d\mu - \gamma^{n}(K) \int_{\omega} f d\mu}{\left(\int_{\omega} \bar{h}_{K}^{p} d\mu\right)^{2}},$$

is equivalent to

$$\gamma^{n}(K) \int_{\omega} f d\mu = -\frac{1}{p} \int_{\omega} f dS_{p,\gamma^{n}}(K, \cdot) \int_{\omega} \bar{h}_{K}^{p} d\mu$$

holds for all continuous functions f on ω . Then we obtain

$$\mu = -\frac{1}{p} \frac{\int_{\omega} \bar{h}_K^p d\mu}{\gamma^n(K)} S_{p,\gamma^n}(K,\cdot).$$
(5.1)

As the case p > 0, we will also show that ϕ_{μ} indeed attains its maximum at support function \bar{h}_K for some $K \in \mathcal{K}(C, \omega)$.

Lemma 5.1. If p < 0, suppose $\omega \subset \Omega$ is a nonempty compact subset. For any nonzero finite Borel measure μ on ω , there exists a C-pseudo-cone $K \in \mathcal{K}(C, \omega)$ such that

$$\mu = -\frac{\int_{\omega} \bar{h}_{K}^{p} d\mu}{p\gamma^{n}(K)} S_{p,\gamma^{n}}(K,\cdot).$$

Proof. For a maximizing sequence $\{\bar{h}_{K_i}\}, K_i \in \mathcal{K}(C, \omega)$, such that

$$\lim_{i \to \infty} \phi_{\mu}(\bar{h}_{K_i}) = \sup \left\{ \phi_{\mu}(f) : f \in C^+(\omega) \right\}.$$

Denote

$$r_i = \min\{r : rB \cap K_i \neq \emptyset\},\$$

If $r_i \to \infty$, since $\bar{h}_{K_i}^p \ge r_i^p$, the following estimate can obtained by Lemma 4.1:

$$\phi_{\mu}(\bar{h}_{K_{i}}) = \frac{\gamma^{n}(K_{i})}{\int_{\omega} \bar{h}_{K_{i}}^{p} d\mu}$$
$$\leq \frac{\gamma^{n}(\mathbb{R}^{n} - r_{i}B)}{\mu(\omega)r_{i}^{p}}$$
$$\leq \frac{c}{r_{i}}e^{-\frac{r_{i}^{2}}{2n}} \cdot r_{i}^{-p} \to 0,$$

for some constant c. This contradicts the fact that $\lim_{i\to\infty} \phi_{\mu}(\bar{h}_{K_i}) = \sup \phi_{\mu} > 0$. Moreover, if $r_i \to 0$, since

$$\gamma^n(\mathbb{R}^n - r_i B) \to 1,$$

we get

$$\phi_{\mu}(\bar{h}_{K_{i}}) = \frac{\gamma^{n}(K_{i})}{\int_{\omega} \bar{h}_{K_{i}}^{p} d\mu} \\ \leq \frac{\gamma^{n}(\mathbb{R}^{n} - r_{i}B)}{\mu(\omega)r_{i}^{p}} \to 0.$$

We arrive at a contradiction again.

This implies the uniform estimate $0 < n_1 < \text{dist}(o, \partial K_i) < n_2$, for some constants n_1 and n_2 . From Lemma 2.2, we can get a subsequence K_{i_j} , such that $K_{i_j} \to K$ for some $K \in \mathcal{K}(C, \omega)$. Therefore, we have $\lim_{i \to \infty} \phi_{\mu}(\bar{h}_{K_{i_j}}) = \phi_{\mu}(\bar{h}_K) = \sup \phi_{\mu}(f)$. By (5.1), the result is proved.

Let μ be a nonzero Borel measure on Ω , p < 0, applying approximation argument again. Choose an increasing sequence of compact sets $\{\omega_i\}$, such that $\omega_i \subset \operatorname{int} \omega_{i+1}$ and $\bigcup_{i \in \mathbb{N}} \omega_i = \Omega$, where ω_0 satisfies $\mu_0 = \mu \llcorner \omega_0$ nonzero. By Lemma 5.1, for each $i \in \mathbb{N}$ and $\mu_i = \mu \llcorner \omega_i$, there exists a *C*-pseudo-cone $K_i \in \mathcal{K}(C, \omega_i)$ satisfying

$$\mu_i = -\frac{\int_{\omega_i} h_{K_i}^p d\mu_i}{p\gamma^n(K_i)} S_{p,\gamma^n}(K_i, \cdot).$$
(5.2)

The uniform estimate of K_i as follows.

Lemma 5.2. For the sequence $\{K_i\}$ in (5.2), there exist two constant m and M such that

$$0 < m < \operatorname{dist}(o, \partial K_i) < M$$

for all $i \in \mathbb{N}$.

Proof. The sequence $\{K_i\}$ in (5.2) satisfies

$$\phi_{\mu_i}(\bar{h}_{K_i}) = \sup\left\{\phi_{\mu_i}(f) = \frac{\gamma^n([f])}{\int_{\omega_i} f^p d\mu_i} : f \in C^+(\omega_i)\right\}.$$

Let $1|_{\omega_i} \in C^+(\omega_i)$ denote the unit function on ω_i . For any $i \in \mathbb{N}$, the corresponding Wulff shape satisfies the inclusion

$$[1|_{\omega_i}] = C \cap \bigcap_{u \in \omega_i} \{ x \in \mathbb{R}^n : \langle x, u \rangle \le -1 \}$$
$$\supset C \cap \bigcap_{u \in \overline{\Omega}} \{ x \in \mathbb{R}^n : \langle x, u \rangle \le -1 \}$$
$$= \bigcap_{u \in \overline{\Omega}} \{ x \in \mathbb{R}^n : \langle x, u \rangle \le -1 \}$$
$$\triangleq A.$$

Thus

$$\gamma^n([1|_{\omega_i}]) \ge \gamma^n(A) > 0$$

for all $i \in \mathbb{N}$. Substituting into ϕ_{μ_i} , since $\mu_i \leq \mu$, we have

$$\phi_{\mu_i}(1|_{\omega_i}) = \frac{\gamma^n([1|_{\omega_i}])}{\mu_i(\omega_i)}$$
$$\geq \frac{\gamma^n([1|_{\omega_i}])}{\mu(\Omega)}$$
$$\geq \frac{\gamma^n(A)}{\mu(\Omega)}.$$

Therefore, $\phi_{\mu_i}(\bar{h}_{K_i}) = \sup \{ \phi_{\mu_i}(f) : f \in C^+(\omega_i) \} \ge \phi_{\mu_i}([1|_{\omega_i}]) \ge \frac{\gamma^n(A)}{\mu(\Omega)} > 0.$ Denote again

$$r_i = \min\{r : rB \cap K_i \neq \emptyset\},\$$

Then we also have $\bar{h}_{K_i}^p \ge r_i^p$. If $r_i \to \infty$, then

$$\phi_{\mu_i}(\bar{h}_{K_i}) = \frac{\gamma^n(K_i)}{\int_{\omega_i} \bar{h}_{K_i}^p d\mu_i} \\ \leq \frac{\gamma^n(\mathbb{R}^n - r_i B)}{\mu_i(\omega_i)r_i^p} \\ \leq \frac{\gamma^n(\mathbb{R}^n - r_i B)}{\mu_0(\omega_0)r_i^p} \\ \leq \frac{l}{r_i} e^{-\frac{r_i^2}{2n}} \cdot r_i^{-p} \to 0,$$

for some constant *l*. This contradicts the uniform lower bound estimate $\phi_{\mu_i}(\bar{h}_{K_i}) \geq \frac{\gamma^n(A)}{\mu(\Omega)}$. If $r_i \to 0$, then

$$\phi_{\mu_i}(\bar{h}_{K_i}) = \frac{\gamma^n(K_i)}{\int_{\omega_i} \bar{h}_{K_i}^p d\mu_i} \\ \leq \frac{\gamma^n(\mathbb{R}^n - r_i B)}{\mu_i(\omega_i)r_i^p} \\ \leq \frac{\gamma^n(\mathbb{R}^n - r_i B)}{\mu_0(\omega_0)r_i^p} \to 0,$$

this also arrive at a contradiction.

Theorem 5.3. If p < 0, for any nonzero finite Borel measure μ on Ω , there exists a C-pseudo-cone K such that

$$\mu = -\frac{c}{p} S_{p,\gamma^n}(K, \cdot),$$

where $c = \frac{\int_{\Omega} \bar{h}_K^p d\mu}{\gamma^n(K)}$.

Proof. This follows from the uniform estimate of $\{K_i\}$ in Lemma 5.2 and applying approximation argument as Theorem 4.4 again.

6. UNIQUENESS OF SOLUTION

The Gaussian measure is log-concave with respect to convex sets (in this paper, we focus on a special class of convex sets—C-pseudo-cones), that is:

Lemma 6.1. For C-pseudo-cones K and L in \mathbb{R}^n , 0 < t < 1,

$$\gamma^n((1-t)K+tL) \ge \gamma^n(K)^{1-t}\gamma^n(L)^t \tag{6.1}$$

equality holds if and only if K = L.

Similar to the method for obtaining uniqueness in classical convex body theory, the following L_p "mixed volume" inequality is a necessary technique for establishing the uniqueness of the corresponding L_p Minkowski problem.

Lemma 6.2. Let $\omega \subset \Omega$ be a nonempty compact set and $p \in (0,1]$, for $K, L \in \mathcal{K}(C, \omega)$, we have

$$\frac{1}{p\gamma^n(K)} \int_{\omega} \bar{h}_K^p - \bar{h}_L^p dS_{p,\gamma^n} K \ge \log \frac{\gamma^n(L)}{\gamma^n(K)},\tag{6.2}$$

equality holds if and only if K = L.

Proof. If $p \in (0, 1]$, $f(t) = t^p$ is a concave function, consequently, for $t \in [0, 1]$,

$$\left((1-t)\bar{h}_K + t\bar{h}_L\right)^p \ge (1-t)\bar{h}_K^p + t\bar{h}_L^p$$

In other words,

$$(1-t)\bar{h}_K + t\bar{h}_L \ge \left((1-t)\bar{h}_K^p + t\bar{h}_L^p\right)^{\frac{1}{p}},$$

and

$$-((1-t)\bar{h}_{K}+t\bar{h}_{L}) \leq -((1-t)\bar{h}_{K}^{p}+t\bar{h}_{L}^{p})^{\frac{1}{p}}.$$

Therefore, we have the set inclusion

$$\begin{split} \left[\left((1-t)\bar{h}_{K}^{p} + t\bar{h}_{L}^{p} \right)^{\frac{1}{p}} \Big|_{\omega} \right] &= C \cap \bigcap_{u \in \omega} \left\{ y \in \mathbb{R}^{n} : \langle y, u \rangle \leq - \left((1-t)\bar{h}_{K}(u)^{p} + t\bar{h}_{L}(u)^{p} \right)^{\frac{1}{p}} \right\} \\ &\supset C \cap \bigcap_{u \in \omega} \left\{ y \in \mathbb{R}^{n} : \langle y, u \rangle \leq - \left((1-t)\bar{h}_{K}(u) + t\bar{h}_{L}(u) \right) \right\} \\ &= \left[\bar{h}_{K} \Big|_{\omega} + t(\bar{h}_{L} \Big|_{\omega} - \bar{h}_{K} \Big|_{\omega} \right) \right] \\ &\supset C \cap \bigcap_{u \in \Omega} \left\{ y \in \mathbb{R}^{n} : \langle y, u \rangle \leq - \left((1-t)\bar{h}_{K}(u) + t\bar{h}_{L}(u) \right) \right\} \\ &= (1-t)K + tL. \end{split}$$

By the log-concave property in Lemma 6.1 we get

$$\log \gamma^{n} \left(\left[\left((1-t)\bar{h}_{K}^{p} + t\bar{h}_{L}^{p} \right)^{\frac{1}{p}} \Big|_{\omega} \right] \right) \geq \log \gamma^{n} \left(\left[\bar{h}_{K} \Big|_{\omega} + t(\bar{h}_{L} \Big|_{\omega} - \bar{h}_{K} \Big|_{\omega} \right) \right] \right)$$
$$\geq \log \gamma^{n} ((1-t)K + tL)$$
$$\geq (1-t) \log \gamma^{n}(K) + t \log \gamma^{n}(L).$$

Using the L_p variational formula in Lemma 3.5, we differentiate both sides of the above inequality at $t = 0^+$:

$$-\frac{1}{p\gamma^n(K)}\int_{\omega}\bar{h}_L^p - \bar{h}_K^p dS_{p,\gamma^n}K \ge \log\gamma^n(L) - \log\gamma^n(K).$$

Assuming equality holds, it follows that

$$\lim_{t \to 0^+} \frac{\log \gamma^n \left(\left[\left((1-t)\bar{h}_K^p + t\bar{h}_L^p \right)^{\frac{1}{p}} \Big|_{\omega} \right] \right) - \log \gamma^n(K)}{t}$$

$$= \lim_{t \to 0^+} \frac{\log \gamma^n \left([\bar{h}_K |_{\omega} + t(\bar{h}_L |_{\omega} - \bar{h}_K |_{\omega})] \right) - \log \gamma^n(K)}{t}$$

$$= \lim_{t \to 0^+} \frac{\log \gamma^n((1-t)K + tL) - \log \gamma^n(K)}{t}$$

$$= \log \gamma^n(L) - \log \gamma^n(K).$$
(6.3)

From the log-concave inequality (6.1), the function $k(t) = \log (\gamma^n((1-t)K + tL))$ is concave on the interval [0, 1]. In the case where equality is achieved, k'(0) = k(1) - k(0)in (6.3), then k(t) must be a linear function. As a result, the equality condition in (6.1) leads to the conclusion that K = L.

Using the L_p mixed volume inequality, we can derive the L_p generalization of the uniqueness result for the Gaussian Minkowski problem for C-pseudo-cones in [35].

Theorem 6.3. For $p \in (0,1]$, let $\omega \subset \Omega$ be a nonempty compact set. If $K, L \in \mathcal{K}(C, \omega)$ have the same L_p Gaussian surface area measure, i.e.,

$$S_{p,\gamma^n}(K,\cdot) = S_{p,\gamma^n}(L,\cdot),$$

and if $\gamma^n(K) = \gamma^n(L)$, then K = L.

Proof. Since $\gamma^n(K) = \gamma^n(L)$, by Lemma 6.2, we get

$$\int_{\omega} \bar{h}_K^p - \bar{h}_L^p dS_{p,\gamma^n} K \ge 0.$$

Therefore,

$$\int_{\omega} \bar{h}_L^p dS_{p,\gamma^n} L = \int_{\omega} \bar{h}_L^p dS_{p,\gamma^n} K \le \int_{\omega} \bar{h}_K^p dS_{p,\gamma^n} K.$$

By substituting K for L and vice versa,

$$\int_{\omega} \bar{h}_{K}^{p} dS_{p,\gamma^{n}} K = \int_{\omega} \bar{h}_{K}^{p} dS_{p,\gamma^{n}} L \le \int_{\omega} \bar{h}_{L}^{p} dS_{p,\gamma^{n}} L.$$

Thus,

$$\int_{\omega} \bar{h}_{L}^{p} dS_{p,\gamma^{n}} L \leq \int_{\omega} \bar{h}_{K}^{p} dS_{p,\gamma^{n}} K \leq \int_{\omega} \bar{h}_{L}^{p} dS_{p,\gamma^{n}} L.$$

As a result, equality holds in both inequalities. By Lemma 6.2, we conclude that K = L.

In the absence of a volume constraint, the following lemma extends the nonuniqueness of the Gaussian Minkowski problem to the L_p setting.

Theorem 6.4. If p < n, for any pointed, n-dimensional closed convex cone C in \mathbb{R}^n and any nonempty compact set $\omega \subset \Omega$, there exist distinct sets $K, L \in \mathcal{K}(C, \omega)$ such that their L_p Gaussian surface area measures coincide, i.e., $S_{p,\gamma^n}(K, \cdot) = S_{p,\gamma^n}(L, \cdot)$, but $K \neq L$.

Proof. For each $v \in \omega$, due to the rotational invariance of the Gaussian measure, suppose that $v = -e_n \in \Omega$. Let H(t) denote the intersection $C \cap H(e_n, t)$, and define

$$A = \{ x \in \mathbb{R}^{n-1} : (x, 1) \in H(1) \}.$$

For t > 0, the Gaussian surface area function s(t) of H(t) (up to a constant) is given by $s(t) = \int_{H(t)} e^{-\frac{|y|^2}{2}} d\mathcal{H}^{n-1}(y)$. Let B_a^{n-1} be a ball in \mathbb{R}^{n-1} with radius a, the Gaussian volume estimate of tB_a^{n-1} , as $t \to 0$, we know

$$\int_{tB_a^{n-1}} e^{-\frac{|y|^2}{2}} d\mathcal{H}^{n-1}(y) \sim t^{n-1}, \qquad t \to 0.$$
(6.4)

Since

$$\int_{tB_a^{n-1}} e^{-\frac{|y|^2}{2}} d\mathcal{H}^{n-1}(y) = \int_{S^{n-2}} \int_0^{at} e^{-\frac{r^2}{2}} r^{n-2} dr du = \mathcal{H}^{n-2}(S^{n-2}) \int_0^{at} e^{-\frac{r^2}{2}} r^{n-2} dr,$$

and

$$\frac{\int_0^{at} e^{-\frac{r^2}{2}} r^{n-2} dr}{t^{n-1}} \to \frac{a^{n-1}}{n-1} \quad \text{as} \quad t \to 0.$$

Then we get the estimate (6.4). There exists a ball B_a^{n-1} such that $A \subset B_a^{n-1}$, therefore, for t > 0,

$$0 < t^{1-p}s(t) = t^{1-p} \int_{H(t)} e^{-\frac{|y|^2}{2}} d\mathcal{H}^{n-1}(y)$$

= $t^{1-p} \int_{(x,t)\in H(t)} e^{-\frac{|x|^2+t^2}{2}} d\mathcal{H}^{n-1}(x)$
= $t^{1-p} e^{-\frac{t^2}{2}} \int_{tA} e^{-\frac{|x|^2}{2}} d\mathcal{H}^{n-1}(x)$
 $\leq t^{1-p} e^{-\frac{t^2}{2}} \int_{B_a^{n-1}} e^{-\frac{|x|^2}{2}} d\mathcal{H}^{n-1}(x)$
 $\sim t^{1-p} \cdot t^{n-1} = t^{n-p} \text{ as } t \to 0.$

If p < n, we obtain $t^{1-p}s(t) \to 0$ as $t \to 0$. If $t \to \infty$, the convergence rate of s(t) is determined by $e^{-\frac{t^2}{2}}$, and thus $t^{1-p}s(t) \to 0$. Then there exist positive numbers $t_1, t_2 > 0$, with $t_1 \neq t_2$, such that $t_1^{1-p}s(t_1) = t_2^{1-p}s(t_2)$.

Define $K = H(t_1) + C$ and $L = H(t_2) + \overline{C}$. Both K and L belong to the class $\mathcal{K}(C, v) \subset \mathcal{K}(C, \omega)$. We then compute

$$S_{p,\gamma^n}(K,v) = \bar{h}_K^{1-p}(v)S_{\gamma^n}(K,v) = \frac{1}{(\sqrt{2\pi})^n} t_1^{1-p}s(t_1)$$

and similarly,

$$S_{p,\gamma^n}(L,v) = \bar{h}_L^{1-p}(v)S_{\gamma^n}(L,v) = \frac{1}{(\sqrt{2\pi})^n} t_2^{1-p} s(t_2).$$

Since the measures $S_{p,\gamma^n}(K,\cdot)$ and $S_{p,\gamma^n}(L,\cdot)$ are both supported on v, it follows that $S_{p,\gamma^n}(K,\cdot) = S_{p,\gamma^n}(L,\cdot)$. This completes the proof of the desired result.

References

- [1] W. Ai, Y. Yang, D. Ye, The L_p dual Minkowski problem for unbounded closed convex sets, (2024), arXiv:2404.09804v1.
- [2] S. Artstein-Avidan, S. Sadovsky, K. Wyczesany, A zoo of dualities, J. Geom. Anal. 33 (2023), 1–40.
- [3] K.J. Böröczky, E. Lutwak, D. Yang, G. Zhang, *The logarithmic Minkowski prob*lem, J. Amer. Math. Soc. 26 (2013), 831–852.
- [4] S. Campi, P. Gronchi, The L_p-Buseman-Petty centroid inequality, Adv. Math. 167 (2002), 128–141.
- [5] S. Chen, S. Hu, W. Liu, Y. Zhao, On the planar Gaussian-Minkowski problem, Adv. Math. 435 (2023), Paper No. 109351.

- [6] A. Cianchi, N. Fusco, F. Maggi, A. Pratelli, On the isoperimetric deficit in Gauss space, Amer. J. Math. 133 (2011), 131–186.
- [7] A. Eskenazis, G. Moschidis, The dimensional Brunn-Minkowski inequality in Gauss space, J. Funct. Anal. 280 (2021), 108914.
- [8] Y. Feng, S. Hu, L. Xu, On the L_p Gaussian Minkowski problem, J. Differ. Equa. 363 (2023), 350–390.
- [9] Y. Feng, W. Liu, L. Xu, Existence of non-symmetric solutions to the Gaussian Minkowski problem, J. Geom. Anal. 33 (2023), Paper No. 89.
- [10] R.J. Gardner, *Geometric Tomography*, Cambridge Univ. Press, Cambridge, 2nd edition, 2006.
- [11] R.J. Gardner, A. Zvavitch, Gaussian Brunn-Minkowski inequalities, Trans. Amer. Math. Soc. 362 (2010), 5333–5353.
- [12] C. Haberl, F. Schuster, General L_p affine isoperimetric inequalities, J. Differential Geom. 83 (2009), 1–26.
- [13] Y. Huang, E. Lutwak, D. Yang, G. Zhang, Geometric measures in the dual Brunn-Minkowski theory and their associated Minkowski problems, Acta Math. 216 (2016), 325–388.
- [14] Y. Huang, E. Lutwak, D. Yang, G. Zhang, The L_p-Aleksandrov problem for L_pintegral curvature, J. Differential Geom. **110** (2018), 1–29.
- [15] Y. Huang, D. Xi, Y. Zhao, The Minkowski problem in Gaussian probability space, Adv. Math. 385 (2021), 107769.
- [16] M. Klartag, V. Milman, Geometry of log-concave functions and measures, Geom. Dedicata 112 (2005), 169–182.
- [17] L. Kryvonos, D. Langharst, Weighted Minkowski's existence theorem and projection bodies, Trans. Amer. Math. Soc. 376 (2023), 8447–8493.
- [18] Q. Li, J. Liu, J. Lu, Nonuniqueness of Solutions to the L_p Dual Minkowski Problem, Int. Math. Res. Not. **12** (2022), 9114–9150.
- [19] N. Li, D. Ye, B. Zhu, The dual Minkowski problem for unbounded closed convex sets, Math. Ann. 388 (2024), 2001–2039.
- [20] J. Liu, The L_p Gaussian Minkowski problem, Calc. Var. Partial Differ. Equ. 61 (2022), 1–23.
- [21] G.V. Livshyts, An extension of Minkowski's theorem and its applications questions about projections for measures, Adv. Math. 356 (2019), Paper No.106803, 40pp.
- [22] E. Lutwak, The Brunn-Minkowski-Firey theory I: Mixed volumes and the Minkowski Problem, J. Differential Geom. 38 (1993), 131–150.
- [23] E. Lutwak, The Brunn-Minkowski-Firey theory II: Affine and geominimal surface areas, Adv. Math. 118 (1996), 244–294.
- [24] E. Lutwak, D. Yang, G. Zhang, On the L_p -Minkowski problem, Trans. Amer. Math. Soc. **356** (2004), 4359–4370.
- [25] E. Lutwak, D. Yang, G. Zhang, L_p dual curvature measures, Adv. Math. **329** (2018), 85–132.

- [26] E. Lutwak, D. Yang, G. Zhang, Sharp affine L_p Sobolev inequalities, J. Differential Geom. 62 (2002), 17–38.
- [27] A. Rashkovskii, *Copolar convexity*, Ann. Pol. Math. **120** (2017), 83–95.
- [28] R. Schneider, *Convex Bodies: The Brunn-Minkowski Theory*, Cambridge Univ. Press, Cambridge, 2014.
- [29] R. Schneider, A Brunn-Minkowski theory for coconvex sets of finite volume, Adv. Math. 332 (2018), 199–234.
- [30] R. Schneider, Minkowski type theorems for convex sets in cones, Acta Math. Hungar. 164 (2021), 282–295.
- [31] R. Schneider, *Pseudo-cones*, Adv. in Appl. Math. **155** (2024), Paper No. 102657, 22pp.
- [32] R. Schneider, A weighted Minkowski theorem for pseudo-cones, Adv. Math. 450 (2024), Paper No. 109760, 26pp.
- [33] R. Schneider, Weighted cone-volume measures of pseudo-cones, (2024), arXiv:2407.05095v1.
- [34] V. Semenov, Y. Zhao, The growth rate of surface area measure for noncompact convex sets with prescribed asymptotic cone, (2024), arXiv:2409.18699v1.
- [35] J. Shan, W. Hu, W. Xu, The Gaussian Minkowski-type problems for C-pseudocones, (2025), arXiv:2501.16449.
- [36] X. Wang, W. Xu, J. Zhou, B. Zhu, Asymptotic theory of C-pseudo-cones, (2024), arXiv:2410.14962v2.
- [37] Y. Xu, J. Li, G. Leng, Dualities and endomorphisms of pseudo-cones, Adv. in Appl. Math. 142 (2023), 1–31.
- [38] J. Yang, D. Ye, B. Zhu, On the L_p Brunn-Minkowski theory and the L_p Minkowski problem for C-coconvex sets, Int. Math. Res. Not. **2023** (2022), 6252–6290.
- [39] N. Zhang, The Minkowski problem for the non-compact convex set with an asymptotic boundary condition, (2024), arXiv:2402.12802v1.
- [40] G. Zhu, The L_p Minkowski problem for polytopes for 0 , J. Funct. Anal.**269**(2015), 1070–1094.

SCHOOL OF MATHEMATICS, SICHUAN UNIVERSITY, CHENGDU, SICHUAN, 610064, P. R. CHINA *Email address:* shanjjmath@163.com, wenchuan@scu.edu.cn