

An adaptive forward-backward-forward splitting algorithm for solving pseudo-monotone inclusions

Flavia Chorobura

Automatic Control and Systems Engineering Department, National University of Science and Technology Politehnica Bucharest, 060042 Bucharest, Romania and Federal Institute of Education, Science and Technology of Paraná, campus Foz do Iguçu, Brazil, flavia.chorobura@ifpr.edu.br

Ion Necoara

Automatic Control and Systems Engineering Department, National University of Science and Technology Politehnica Bucharest, 060042 Bucharest and Gheorghe Mihoc-Caius Iacob Institute of Mathematical Statistics and Applied Mathematics of the Romanian Academy, 050711 Bucharest, Romania, ion.necoara@upb.ro

Jean-Christophe Pesquet

University Paris-Saclay, CentraleSupélec, CVN, Inria, Gif-sur-Yvette, France, jean-christophe.pesquet@centralesupelec.fr

Authors are encouraged to submit new papers to INFORMS journals by means of a style file template, which includes the journal title. However, use of a template does not certify that the paper has been accepted for publication in the named journal. INFORMS journal templates are for the exclusive purpose of submitting to an INFORMS journal and are not intended to be a true representation of the article's final published form. Use of this template to distribute papers in print or online or to submit papers to another non-INFORM publication is prohibited.

Abstract. In this paper, we propose an adaptive forward-backward-forward splitting algorithm for finding a zero of a pseudo-monotone operator which is split as a sum of three operators: the first is continuous single-valued, the second is Lipschitzian, and the third is maximally monotone. This setting covers, in particular, constrained minimization scenarios, such as problems having smooth and convex functional constraints (e.g., quadratically constrained quadratic programs) or problems with a pseudo-convex objective function minimized over a simple closed convex set (e.g., quadratic over linear fractional programs). For the general problem, we design a forward-backward-forward splitting type method based on novel adaptive stepsize strategies. Under an additional generalized Lipschitz property of the first operator, sublinear convergence rate is derived for the sequence generated by our adaptive algorithm. Moreover, if the sum is uniformly pseudo-monotone, linear/sublinear rates are derived depending on the parameter of uniform pseudo-monotonicity. Preliminary numerical experiments demonstrate the good performance of our method when compared to some existing optimization methods and software.

Funding: The research leading to these results has received funding from:

ITN-ETN project TraDE-OPT funded by the European Union's Horizon 2020 Research and Innovation Programme under the Marie Skłodowska-Curie grant agreement No. 861137; UEFISCDI, Romania, PN-III-P4-PCE-2021-0720, under project L2O-MOC, nr. 70/2022.

Key words: Pseudo-monotone operators, forward-backward-forward splitting, adaptive stepsize, convergence analysis, nonconvex optimization.

1. Introduction Let \mathbb{H} be a finite-dimensional real vector space endowed with a scalar product $\langle \cdot, \cdot \rangle$ and the corresponding norm $\| \cdot \|$. Our goal is to find a zero of a sum of three operators $A: \mathbb{H} \rightarrow \mathbb{H}$, $B: \mathbb{H} \rightarrow \mathbb{H}$, and $C: \mathbb{H} \rightarrow 2^{\mathbb{H}}$, that is

$$\text{find } \bar{z} \in \mathbb{H} \text{ such that } 0 \in A\bar{z} + B\bar{z} + C\bar{z}, \quad (1)$$

where $A + B + C$ is pseudo-monotone, and A satisfies some generalized smoothness condition, B is smooth, and C is maximally monotone as detailed in Assumption 1 of Section 3. Finding a zero of a sum of operators is a very general problem and covers, in particular, constrained optimization, and minimax optimization problems frequently encountered in signal processing [25], triangulation in computer vision [2], semi-supervised learning [15], learning of kernel matrices [28], steering direction estimation for RADAR detection [20], generative adversarial networks [31] among others.

Previous work. The problem of finding a zero of a sum of operators is considered in many works. For example, [8, 10, 16, 17, 19, 26, 39] cover the monotone case, while [3, 11, 35] consider the nonmonotone case. In [19, 26] all three operators are assumed maximally monotone and, additionally, the first is Lipschitz continuous. Under these settings, algorithms based on resolvent and forward operators, activated one at a time successively, are proposed together with a detailed convergence analysis. Furthermore, finding a zero of a sum of two maximally monotone operators, $A + C$, such that A is a continuous single-valued operator, is investigated in [39] and a forward-backward-forward algorithm is proposed (also known as Tseng's algorithm), where the stepsize is chosen constant when A is Lipschitz or based on an Armijo-Goldstein-type rule, otherwise. Linear rate was derived for this method when $A + C$ is strongly monotone. In [3, 11, 35], A is assumed Lipschitz continuous, possibly nonmonotone, and C is maximally monotone, such that either $A + C$ satisfies the weak Minty condition or a cohyponotonicity assumption. In particular, [35] considers an extragradient algorithm with adaptive and constant stepsizes, which reduces, for a specific choice of stepsize, to the forward-backward-forward algorithm in the monotone case. Moreover, [11] analyzes an optimistic gradient algorithm, while in [3] algorithms based on classical Halpern and Krasnosel'skii-Mann iterations are analyzed. For all these methods, under suitable assumptions, sublinear rates are derived. Finally, finding a zero of a sum of three operators $A + B + C$ is considered in [17], where A, C are maximally monotone and B is Lipschitz continuous and monotone, and asymptotic convergence is proved for an error-tolerant forward-backward-forward algorithm.

The forward-backward-forward algorithm was also extended to solve variational inequalities. For example, [7, 38] consider a variational inequality, where the operator is Lipschitz continuous, and a (modified) Tseng algorithm is employed with a constant stepsize or an adaptive stepsize, so that it is not necessary to know the Lipschitz constant. Convergence is derived when the operator is pseudo-monotone. Moreover, the Lipschitz assumption on the operator involved in the variational inequality is relaxed in [37], the operator being assumed continuous. Then, Tseng's algorithm is considered with an Armijo-Goldstein rule for the stepsize. Under standard conditions, the weak and strong convergence of the method is obtained in the pseudo-monotone case. Our approach differs from [37], as we consider that the operator A satisfies a relaxed Lipschitz condition and we employ Tseng's algorithm with novel adaptive stepsize rules (e.g., based on the positive root of a polynomial equation). Other methods for solving variational inequalities with a Lipschitz operator in the monotone case were considered e.g., in [33] and in the nonmonotone case (under weak Minty condition) in [21].

Furthermore, specific algorithms were also developed for particular classes of variational inequalities, such as convex-concave minimax optimization problems [14, 18]. More specifically, these papers address problems of the form:

$$\min_{x \in \mathcal{X}} \max_{y \in \mathcal{Y}} \langle Lx, y \rangle + \varphi(x) - \psi(y), \quad (2)$$

where \mathcal{X} and \mathcal{Y} are Hilbert spaces, L is a linear operator and $\varphi : \mathcal{X} \rightarrow \bar{\mathbb{R}} := \mathbb{R} \cup \{+\infty\}$ and $\psi : \mathcal{Y} \rightarrow \bar{\mathbb{R}}$ are proper, convex, lower semicontinuous functions. For such problems, a primal-dual proximal algorithm is proposed in [14] for which a sublinear rate is derived in the optimality measure:

$$G(\bar{x}, \bar{y}) = \max_{y \in \mathcal{Y}} \langle L\bar{x}, y \rangle - \psi(y) + \varphi(\bar{x}) - \min_{x \in \mathcal{X}} \langle Lx, \bar{y} \rangle + \varphi(x) - \psi(\bar{y}),$$

for a given $(\bar{x}, \bar{y}) \in \mathcal{X} \times \mathcal{Y}$. An extension of the algorithm from [14] is given in [18], where φ is split as $\varphi_1 + \varphi_2$, with $\varphi_1 : \mathcal{X} \rightarrow \mathbb{R}$ convex, differentiable with a Lipschitz continuous gradient, while φ_2 is a proper, convex, lower semicontinuous function. It is proved that this algorithm converges weakly to a solution to problem (2) and, if $\varphi_1 = 0$, then [18] recovers the primal-dual algorithm in [14].

Contributions. In this paper, we propose a method for finding a zero of a sum of three operators, which are not necessarily monotone. For this general problem we design a forward-backward-forward splitting type method based on novel adaptive stepsize strategies and then perform a detailed convergence analysis. More specifically, our main contributions are the following.

(i) *Mathematical modelling:* We consider the general problem (1) of finding a zero of a sum of three operators, $A + B + C$, such that A is continuous, B is Lipschitz continuous, and C is maximally monotone. In contrast to other works that assume A to be Lipschitz continuous and the sum to be monotone, we relax these conditions, i.e., we require the operator A to satisfy a generalized Lipschitz condition and the sum to be pseudo-monotone. Our assumptions cover important classes of optimization problems such as problems minimizing smooth and convex functional constraints (e.g., quadratically constrained quadratic programs) or problems minimizing pseudo-convex objective functions over a simple closed convex set (e.g., quadratic over linear fractional programs).

(ii) *Algorithm:* For solving this general problem we propose a variant of the forward-backward-forward algorithm [39], based on *two novel adaptive stepsize strategies*. In contrast to previous works where computationally expensive Armijo-Goldstein stepsize rules are used when the operator is continuous, we propose two *adaptive* stepsize strategies that require finding the root of a certain nonlinear equation whose coefficients depend on the current iterate and on the parameters characterizing the operator properties. In particular, for quadratically constrained quadratic (resp. quadratic over linear fractional) programs the stepsize is computed solving a second-order (resp. third-order) polynomial equation.

(iii) *Convergence analysis:* Within the considered settings, we provide a detailed convergence analysis for the forward-backward-forward algorithm based on our adaptive stepsize rules. In particular, when the sum of the operators is pseudo-monotone, we prove the global asymptotic convergence for the whole sequence generated by the algorithm and, additionally, establish sublinear convergence rate. An improved linear rate is obtained when the sum is uniformly pseudo-monotone of order $q \in [1, 2]$.

(iv) *Experiments:* Detailed numerical experiments using synthetic and real data demonstrate the effectiveness of our method and allows us to evaluate its performance when compared to some existing state-of-the-art optimization methods from [37, 39], and existing software [22].

In conclusion, enhancing a forward-backward-forward splitting algorithm by introducing novel adaptive stepsize strategies and considering suitable assumptions on the generalized problem (covering a wide spectrum of applications) to allow the derivation of a complete convergence analysis, represent important contributions for solving inclusion problems.

2. Background We denote by $\text{zer}(A)$ the set of zeros of an operator A , by $\Gamma_0(\mathbb{H})$ the set of proper lower semicontinuous convex functions on \mathbb{H} with values in $] -\infty, +\infty]$, by $\mathbf{0}_{m \times m}$ the $m \times m$

null matrix and by $\mathbf{0}_m$ the m -dimensional null vector. Further, let us recall the definition of the subdifferential of a convex function.

DEFINITION 1. The subdifferential of a proper convex function $f : \mathbb{H} \rightarrow \bar{\mathbb{R}}$ is the set-valued operator $\partial f : \mathbb{H} \rightarrow 2^{\mathbb{H}}$ which maps every point $x \in \mathbb{H}$ to the set

$$\partial f(x) = \{u \in \mathbb{H} \mid (\forall y \in \mathbb{H}) \langle y - x, u \rangle + f(x) \leq f(y)\}.$$

Note that $\partial f(x) = \emptyset$ for $x \notin \text{dom} f$. For example, let D be a nonempty closed and convex subset of \mathbb{H} and let its indicator function ι_D be defined as

$$\iota_D : \mathbb{H} \rightarrow \bar{\mathbb{R}} : x \mapsto \begin{cases} 0 & \text{if } x \in D \\ +\infty, & \text{otherwise.} \end{cases} \quad (3)$$

Then, $\partial \iota_D = \mathcal{N}_D$, where \mathcal{N}_D is the normal cone to D , i.e.

$$\mathcal{N}_D(x) = \begin{cases} \{u \in \mathbb{H} \mid (\forall y \in D) \langle y - x, u \rangle \leq 0\} & \text{if } x \in D \\ \emptyset & \text{otherwise.} \end{cases} \quad (4)$$

Moreover, if f is differentiable at a point $x \in \text{dom} f$, its gradient is denoted by $\nabla f(x)$. Let us also recall the definition of functions with Hölder continuous gradient.

DEFINITION 2. Let $\nu \in]0, 1]$. Then, the differentiable function $g : \mathbb{H} \rightarrow \mathbb{R}$ has a ν -Hölder continuous gradient, if there exists $L_g > 0$ such that

$$(\forall (x, w) \in \mathbb{H}^2) \quad \|\nabla g(x) - \nabla g(w)\| \leq L_g \|x - w\|^\nu. \quad (5)$$

If g has ν -Hölder continuous gradient, then the following inequality holds, see [40, Lemma 1]:

$$(\forall (x, w) \in \mathbb{H}^2) \quad |g(w) - g(x) - \langle \nabla g(x), w - x \rangle| \leq \frac{L_g}{1 + \nu} \|w - x\|^{1+\nu}. \quad (6)$$

Next, we present the definitions of pseudo-convex functions and operators.

DEFINITION 3. Let $\mathcal{X} \subseteq \mathbb{H}$ be an open set, let $f : \mathcal{X} \rightarrow \mathbb{R}$ be a differentiable function and let \mathcal{Z} be a subset of \mathcal{X} . Then, f is said to be pseudo-convex on \mathcal{Z} if, for every $(x, w) \in \mathcal{Z}^2$, one has:

$$\langle \nabla f(x), w - x \rangle \geq 0 \implies f(x) \leq f(w).$$

Clearly, any convex function is pseudo-convex and any stationary point of a pseudo-convex function is a global minimum. However, there are also pseudo-convex functions that are not convex. For

example, consider an open convex set $\mathcal{X} \subset \mathbb{R}^n$ and differentiable functions $g : \mathcal{X} \rightarrow [0, +\infty[$ and $h : \mathcal{X} \rightarrow]0, +\infty[$ such that g is convex and h is concave. Then, the function $f : \mathcal{X} \rightarrow]0, +\infty[: x \mapsto g(x)/h(x)$, is pseudo-convex on any subset of \mathcal{X} [6]. Other examples of pseudo-convex functions are given in Example 3 below, see also [29]. The notion of pseudo-convexity was also extended to nondifferentiable functions, see for example [4].

DEFINITION 4. An operator $T : \mathbb{H} \rightarrow 2^{\mathbb{H}}$ is said to be pseudo-monotone if

$$(\forall (x, y) \in \mathbb{H}^2) (\exists \hat{x} \in Tx) \langle \hat{x}, y - x \rangle \geq 0 \implies (\forall \hat{y} \in Ty) \langle \hat{y}, y - x \rangle \geq 0.$$

For example, [27] shows that any differentiable pseudo-convex function has a pseudo-monotone gradient. In addition, [4] proves that a lower semicontinuous radially continuous function is pseudo-convex if and only if its subdifferential is pseudo-monotone. Moreover, note that every monotone operator is pseudo-monotone

Finally, let us recall the definition of the resolvent of an operator $C : \mathbb{H} \rightarrow 2^{\mathbb{H}}$. The resolvent of C is the operator $J_C = (\text{Id} + C)^{-1}$, that is

$$(\forall (x, p) \in \mathbb{H}^2) \quad p \in J_C x \iff x - p \in Cp.$$

If $C : \mathbb{H} \rightarrow 2^{\mathbb{H}}$ is maximally monotone, then J_C is single-valued, defined everywhere on \mathbb{H} , and firmly nonexpansive [5]. Moreover, if $C = \partial f$, the subdifferential operator of a convex function f , then its resolvent is the proximal mapping prox_f . If $f = \iota_D$, where ι_D is defined in (3) and D is a nonempty closed convex subset of \mathbb{H} , then $\text{prox}_{\gamma \iota_D} = \text{proj}_D$, where proj_D is the projection operator onto the set D .

3. Assumptions and examples In this section we provide our main assumptions and also several examples of problems that fit into our framework. First, let us present our standing assumptions for the operators A , B , and C .

ASSUMPTION 1. *The following assumptions hold for problem (1):*

- i) C has nonempty closed convex domain, $\text{dom } C$, and is maximally monotone.
- ii) A is a continuous single-valued operator on $\text{dom } C \subseteq \mathbb{H}$.
- iii) B is a single-valued operator and Lipschitz on $\text{dom } C$ with a Lipschitz constant $L_B > 0$ (when $B = 0$, we can take an arbitrarily small positive value for L_B).
- iv) $A + B + C$ is a pseudo-monotone operator.

v) There exist $(\zeta, \tau) \in]0, +\infty[^2$ such that for every $(u, w) \in \mathbb{H}^2$, $\gamma \in]0, +\infty[$, $q = \text{proj}_{\text{dom } C} w$, and $z = q - \gamma u$, the following holds:

$$\|q - J_{\gamma C} z\| \leq \gamma(\zeta \|u\| + \tau). \quad (7)$$

vi) A satisfies a generalized Lipschitz condition, that is, there exist $\mu \in]0, 2]$, $(\beta, \theta) \in [2, +\infty[^2$ and continuous functions a, b , and c from \mathbb{H} to $[0, +\infty[$ such that, for every $(z_1, z_2) \in (\text{dom } C)^2$,

$$\|Az_1 - Az_2\|^2 \leq a(z_1)\|z_1 - z_2\|^\mu + b(z_1)\|z_1 - z_2\|^\theta + c(z_1)\|z_1 - z_2\|^\beta. \quad (8)$$

REMARK 1. Assumption 1.vi can be generalized to more than three terms, i.e., A satisfies a generalized Lipschitz condition if there exist $\mu \in]0, +\infty[$, $(\theta_i)_{1 \leq i \leq m} \in [2, +\infty[^m$, and continuous functions a and $(b_i)_{1 \leq i \leq m}$ from \mathbb{H} to $[0, +\infty[$ such that

$$(\forall (z_1, z_2) \in (\text{dom } C)^2) \quad \|Az_1 - Az_2\|^2 \leq a(z_1)\|z_1 - z_2\|^\mu + \sum_{i=1}^m b_i(z_1)\|z_1 - z_2\|^{\theta_i}.$$

Although the convergence analysis from this paper can be derived under this more general condition, Assumption 1.vi turns out to be sufficient in most of the applications of interest. Additionally, note that we require $a(\cdot)$, $b(\cdot)$ and $c(\cdot)$ to depend only on z_1 in order to make our adaptive stepsize choices introduced in the following sections dependent only on the current iterate.

Note that our assumptions are quite general. Next, we present some important examples of optimization problems that can be recast as problem (1), showing the versatility of our settings.

EXAMPLE 1. (Minimizing the sum of three functions). The most straightforward example of inclusion (1) arises from the optimization problem:

$$\min_{x \in \mathbb{R}^n} F(x) := f(x) + g(x) + h_1(x) + h_2(Lx), \quad (9)$$

where $L \in \mathbb{R}^{m \times n}$, $f \in \Gamma_0(\mathbb{R}^n)$ has a ν -Hölder continuous gradient with constant $L_f > 0$, $g \in \Gamma_0(\mathbb{R}^n)$ has a Lipschitz continuous gradient with constant $L_g > 0$, and $h_2 \in \Gamma_0(\mathbb{R}^m)$ is finite at a point in the relative interior of the range of L . Moreover, we assume that h_1 is the indicator function of a nonempty closed convex set D and h_2 is Lipschitz on its domain with modulus $L_{h_2} > 0$. We assume that $L(D) \subseteq \text{dom } h_2$ and there is a point in the intersection of the relative interiors of $L(D)$ and $\text{dom } h_2$. The latter condition ensures that $\partial(h_1 + h_2 \circ L) = \partial h_1 + L^\top \circ \partial h_2 \circ L$. This formulation covers composite ($f = 0$), Hölder composite ($g = 0$), or hybrid composite problems, respectively.

The first-order optimality condition for (9) at \bar{x} reduces to $0 \in A\bar{x} + B\bar{x} + C\bar{x}$, where $A = \nabla f$, $B = \nabla g$, and $C = \partial h_1 + L^\top \circ \partial h_2 \circ L$. Next, we show that Assumption 1 holds for this example:

(i) - (iii) Note that, since $h_1 \in \Gamma_0(\mathbb{R}^n)$, $h_2 \in \Gamma_0(\mathbb{R}^m)$, the gradient of f is continuous and g has Lipschitz continuous gradient, operator C is maximally monotone, operator A is a single-valued continuous operator, and operator B is Lipschitz continuous with constant $L_B = L_g$.

(iv) Since the functions f and g are convex, then operators A and B are monotone. Moreover, since C is maximal monotone, we have that the operator $A + B + C$ is monotone, and consequently, pseudo-monotone.

(v) Let $(u, w) \in \mathbb{H}^2$, $q = \text{proj}_{\text{dom}C} w$, and $z = q - \gamma u$. Then, we have

$$\text{dom}C = D \quad \text{and} \quad J_{\gamma C} = \text{prox}_{\gamma(h_1+h_2 \circ L)}.$$

In [1] it was proved that

$$J_{\gamma C} = \text{proj}_{\text{dom}C} \circ \text{prox}_{\gamma h_2 \circ L}^{h_1}, \quad \text{where} \quad \text{prox}_{\gamma h_2 \circ L}^{h_1} = (\text{Id} + \gamma L^\top \circ \partial h_2 \circ L \circ \text{proj}_{\text{dom}C})^{-1}.$$

Define $p = \text{prox}_{\gamma h_2 \circ L}^{h_1}(z)$. Then, $p + \gamma \hat{p} = z$ for some $\hat{p} \in L^\top \circ \partial h_2 \circ L(\text{proj}_{\text{dom}C} p)$. Moreover, $\|\hat{p}\| \leq L_{h_2} \|L\|$ and

$$\begin{aligned} \|\text{proj}_{\text{dom}C} w - J_{\gamma C} z\| &\leq \|\text{proj}_{\text{dom}C} w - \text{prox}_{\gamma h_2 \circ L}^{h_1}(z)\| = \|\text{proj}_{\text{dom}C} w - z + \gamma \hat{p}\| \\ &\leq \|\text{proj}_{\text{dom}C} w - z\| + \gamma \|\hat{p}\| \leq \|\text{proj}_{\text{dom}C} w - z\| + \gamma L_{h_2} \|L\| = \gamma \|u\| + \gamma L_{h_2} \|L\|, \end{aligned}$$

where, in the first inequality, we have used the nonexpansiveness of the projection operator and $\text{proj}_{\text{dom}C}(\text{proj}_{\text{dom}C}(w)) = \text{proj}_{\text{dom}C}(w)$, and in the last one, we have used the linear relation between z , $\text{proj}_{\text{dom}C} w$, and u . Therefore, in this case, $\zeta = 1$ and $\tau = L_{h_2} \|L\|$.

(vi) From (5) and the definition of operator A , we have, for every $(z_1, z_2) \in D^2$,

$$\|Az_1 - Az_2\|^2 \leq L_f^2 \|z_1 - z_2\|^{2\nu}$$

Hence, $a(z_1) = L_f^2$, $\mu = 2\nu$, $b(z_1) = 0$, and $c(z_1) = 0$.

EXAMPLE 2. (Minimax problems). Let m and n be positive integers, and consider the following minimax problem:

$$\min_{x \in \mathbb{R}^n} \max_{y \in \mathbb{R}^m} F(x, y) + \varphi(x) - \psi(y), \quad (10)$$

where $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$ is a differentiable function, $\psi = \psi_1 + \psi_2$, and $\varphi = \varphi_1 + \varphi_2$, with ψ_1 and φ_1 having Lipschitz gradients, $\varphi_2 \in \Gamma_0(\mathbb{R}^n)$, and $\psi_2 \in \Gamma_0(\mathbb{R}^m)$. Note that the minimax problem (10) is

more general than problem (2) considered in previous works, as we allow more general expressions for F beyond bilinear terms. The first-order optimality conditions for this problem are equivalent to solving the inclusion:

$$0 \in A(\bar{x}, \bar{y}) + B(\bar{x}, \bar{y}) + C(\bar{x}, \bar{y}), \quad (11)$$

where $\mathbb{H} = \mathbb{R}^n \times \mathbb{R}^m$ and the three operators are

$$A : \mathbb{H} \rightarrow \mathbb{H} : (x, y) \mapsto (\nabla_x F(x, y), -\nabla_y F(x, y)) \quad (12)$$

$$B : \mathbb{H} \rightarrow \mathbb{H} : (x, y) \mapsto (\nabla \varphi_1(x), \nabla \psi_1(y)),$$

$$C : \mathbb{H} \rightarrow 2^{\mathbb{H}} : (x, y) \mapsto \partial \varphi_2(x) \times \partial \psi_2(y).$$

Next, we give two examples of this minimax problem.

a) For simplicity, let us consider $m = 1$ and

$$(\forall (x, y) \in \mathbb{H}) \quad F(x, y) = yg(x), \quad (13)$$

where g is a twice differentiable convex function, which has a ν -Hölder continuous gradient with constant L_g , φ_1 and ψ_1 are convex functions, ψ_2 is the indicator function of the interval $[0, +\infty[$ and $\varphi_2 = t_1 + t_2$, where t_1 is the indicator function of a nonempty closed convex set D , and t_2 is a proper lower-semicontinuous convex function which is Lipschitz continuous on its domain with modulus $L_{t_2} > 0$. We assume that $D \subseteq \text{dom } t_2$ and there is a point in the intersection of the relative interiors of D and $\text{dom } t_2$. Let us show that Assumption 1 holds for this example.

(i) - (iii) From the definitions of operators A , B and C , we can easily verify that C is maximally monotone, A is a continuous single-valued operator, and B is Lipschitz continuous.

(iv) Operator A is given by

$$A : \mathbb{H} \rightarrow \mathbb{H} : (x, y) \mapsto (y\nabla g(x), -g(x)). \quad (14)$$

The Jacobian \mathcal{J}_A of A at $(x, y) \in \mathbb{H}$ is

$$\mathcal{J}_A(x, y) = \begin{bmatrix} y\nabla^2 g(x) & \nabla g(x) \\ -\nabla g(x)^\top & 0 \end{bmatrix}.$$

Note that, for every $(x, y) \in \mathbb{R}^n \times [0, +\infty[$, $\mathcal{J}_A(x, y)$ is positive semidefinite matrix. Since A is continuous and monotone on $\mathbb{R}^n \times [0, +\infty[$, then A is maximally monotone on $\mathbb{R}^n \times [0, +\infty[$, see

[5]. Moreover, since φ and ψ_1 are proper lower semicontinuous convex functions, then $A + B + C$ is a monotone operator, which is a particular instance of a pseudo-monotone operator.

(v) Note that in this case $\text{dom } C = D \times [0, \infty[$ and we have:

$$J_{\gamma C}(x, y) = (\text{prox}_{\gamma(t_1+t_2)}(x), \text{proj}_{[0, +\infty[}(y)).$$

Following Example 1, the following relation holds:

$$\text{prox}_{\gamma(t_1+t_2)}(x) = \text{proj}_D \circ \text{prox}_{\gamma t_2}^{t_1}, \text{ where } \text{prox}_{\gamma t_2}^{t_1} = (\text{Id} + \gamma \partial t_2 \circ \text{proj}_D)^{-1}.$$

Let $(u, w) \in \mathbb{H}^2$, $q = \text{proj}_{\text{dom} C} w$, and $z = q - \gamma u$. Considering the decomposition $w = (w_1, w_2)$, $u = (u_1, u_2)$ and $z = (z_1, z_2)$. Define $p = \text{prox}_{\gamma t_2}^{t_1}(z_1)$. Then, $p + \gamma \hat{p} = z_1$, for some $\hat{p} \in \partial t_2(\text{proj}_D p)$, and $\|\hat{p}\| \leq L_{t_2}$. We further have

$$\begin{aligned} \|\text{proj}_{\text{dom} C} w - J_{\gamma C} z\|^2 &= \|\text{proj}_D w_1 - \text{proj}_D \circ \text{prox}_{\gamma t_2}^{t_1} z_1\|^2 + \|\text{proj}_{[0, +\infty[} w_2 - \text{proj}_{[0, +\infty[} z_2\|^2 \\ &\leq \|\text{proj}_D w_1 - \text{prox}_{\gamma t_2}^{t_1} z_1\|^2 + \|\text{proj}_{[0, +\infty[} w_2 - z_2\|^2 \\ &= \|\text{proj}_D w_1 - z_1 + \gamma \hat{p}\|^2 + \|\text{proj}_{[0, +\infty[} w_2 - z_2\|^2 \\ &\leq 2\|\text{proj}_D w_1 - z_1\|^2 + 2\gamma^2 \|\hat{p}\|^2 + \|\text{proj}_{[0, +\infty[} w_2 - z_2\|^2 \\ &= 2\gamma^2 \|u_1\|^2 + 2\gamma^2 \|\hat{p}\|^2 + \gamma^2 \|u_2\|^2 \leq 2\gamma^2 \|u\|^2 + 2\gamma^2 L_{t_2}^2. \end{aligned}$$

Thus, we get $\zeta = \sqrt{2}$ and $\tau = \sqrt{2} L_{t_2}$.

(vi) For every $z = (x, y) \in \mathbb{R}^n \times \mathbb{R}$ and $\bar{z} = (\bar{x}, \bar{y}) \in \mathbb{R}^n \times \mathbb{R}$, we have

$$\|Az - A\bar{z}\|^2 \leq 2\|\nabla g(x)\|^2 \|z - \bar{z}\|^2 + 4L_g^2 \|z - \bar{z}\|^{2+2\nu} + 4L_g^2 |y|^2 \|z - \bar{z}\|^{2\nu}.$$

Indeed, from the definition of A (see (14)),

$$\|A(x, y) - A(\bar{x}, \bar{y})\|^2 = \|\nabla g(x)y - \nabla g(\bar{x})\bar{y}\|^2 + |g(\bar{x}) - g(x)|^2. \quad (15)$$

Moreover,

$$\begin{aligned} \|\nabla g(x)y - \nabla g(\bar{x})\bar{y}\|^2 &= \|\nabla g(x)y - \nabla g(x)\bar{y} + \nabla g(x)\bar{y} - \nabla g(\bar{x})\bar{y}\|^2 \\ &\leq 2\|\nabla g(x)\|^2 |y - \bar{y}|^2 + 2|\bar{y}|^2 \|\nabla g(x) - \nabla g(\bar{x})\|^2 \\ &\stackrel{(5)}{\leq} 2\|\nabla g(x)\|^2 |y - \bar{y}|^2 + 2L_g^2 |\bar{y}|^2 \|x - \bar{x}\|^{2\nu} \\ &\leq 2\|\nabla g(x)\|^2 |y - \bar{y}|^2 + 4L_g^2 |y|^2 \|x - \bar{x}\|^{2\nu} + 4L_g^2 |y - \bar{y}|^2 \|x - \bar{x}\|^{2\nu}, \quad (16) \end{aligned}$$

where, in the first and last inequalities, we used the fact that $\|a + b\|^2 \leq 2\|a\|^2 + 2\|b\|^2$, for all $a, b \in \mathbb{R}^n$ and $n \in \mathbb{N}$. On other hand, from (6) we deduce that

$$\begin{aligned} |g(\bar{x}) - g(x)|^2 &\leq 2|\langle \nabla g(x), \bar{x} - x \rangle|^2 + \frac{2L_g^2}{(1+\nu)^2} \|\bar{x} - x\|^{2+2\nu} \\ &\leq 2\|\nabla g(x)\|^2 \|\bar{x} - x\|^2 + \frac{2L_g^2}{(1+\nu)^2} \|\bar{x} - x\|^{2+2\nu} \\ &\leq 2\|\nabla g(x)\|^2 \|\bar{x} - x\|^2 + 4L_g^2 \|\bar{x} - x\|^{2+2\nu}. \end{aligned} \quad (17)$$

Altogether, (15), (16), and (17) lead to

$$\begin{aligned} &\|A(x, y) - A(\bar{x}, \bar{y})\|^2 \\ &\leq 2\|\nabla g(x)\|^2 \|(x, y) - (\bar{x}, \bar{y})\|^2 + 4L_g^2 \|(x, y) - (\bar{x}, \bar{y})\|^2 \|\bar{x} - x\|^{2\nu} + 4L_g^2 |y|^2 \|x - \bar{x}\|^{2\nu} \\ &\leq 2\|\nabla g(x)\|^2 \|z - \bar{z}\|^2 + 4L_g^2 \|z - \bar{z}\|^{2+2\nu} + 4L_g^2 |y|^2 \|x - \bar{x}\|^{2\nu}. \end{aligned}$$

Hence, Assumption 1.vi holds with $\mu = 2\nu$, $\beta = 2 + 2\nu$ and $\theta = 2$. Finally, the inequality $\|x - \bar{x}\| \leq \|(x, y) - (\bar{x}, \bar{y})\|$ allows us to prove the statement.

b) Consider

$$(\forall (x, y) \in \mathbb{H}) \quad F(x, y) = \sum_{i=1}^m y_i g_i(x), \quad (18)$$

where $(g_i)_{1 \leq i \leq \bar{m}}$ are twice differentiable convex functions, which have Lipschitz continuous gradients with constants $L_{g_i} > 0$, $(g_i)_{\bar{m}+1 \leq i \leq m}$ are the following affine functions:

$$(\forall i \in \{\bar{m} + 1, \dots, m\}) \quad g_i: x \mapsto l_i^\top x - r_i$$

with $(l_i, r_i)_{\bar{m}+1 \leq i \leq m}$ vectors in $\mathbb{R}^n \times \mathbb{R}$. In addition, φ_1 and ψ_1 are convex functions, ψ_2 is the indicator function of the set $[0, +\infty[^{\bar{m}} \times \mathbb{R}^{m-\bar{m}}$, and $\varphi_2 = t_1 + t_2$, where t_1 is the indicator function of a nonempty closed convex set D and t_2 is a proper lower-semicontinuous convex function which is Lipschitz on its domain with modulus $L_{t_2} > 0$. We assume that $D \subseteq \text{dom } t_2$ and there is a point in the intersection of the relative interiors of D and $\text{dom } t_2$. Let us show that Assumption 1 holds for this example.

(i)-(iii) From the definitions of the operators A , B , and C , it is straightforward that C is maximally monotone, A is a continuous single-valued operator, and B is Lipschitz continuous.

(iv) First note that $A + \{\mathbf{0}_n\} \times \mathcal{N}_{[0,+\infty]^m}$ is maximally monotone. Indeed, if we consider the notation $A = A_1 \times A_2$, where, for every $(x, y) \in \mathbb{R}^n \times \mathbb{R}^m$, $A_1(x, y) \in \mathbb{R}^n$ and $A_2(x, y) \in \mathbb{R}^m$, then $(A + \{\mathbf{0}_n\} \times \mathcal{N}_{[0,+\infty]^m})(x, y) = A_1(x, y) \times [A_2(x, y) + \mathcal{N}_{[0,+\infty]^m}(y)]$. Moreover, we have

$$A : \mathbb{H} \rightarrow \mathbb{H} : (x, y) \mapsto \left(\underbrace{\sum_{i=1}^m y_i \nabla g_i(x)}_{A_1(x,y)}, \underbrace{-g(x)}_{A_2(x,y)} \right), \quad (19)$$

where $g(x) = [g_1(x), \dots, g_m(x)]^\top$. The Jacobian \mathcal{J}_A of A at (x, y) is

$$\mathcal{J}_A(x, y) = \begin{bmatrix} \sum_{i=1}^m y_i \nabla^2 g_i(x) & \nabla g(x) \\ -\nabla g(x)^\top & \mathbf{0}_{m \times m} \end{bmatrix}, \quad \text{with } \nabla g(x) = [\nabla g_1(x), \dots, \nabla g_m(x)].$$

Note that, for every $(x, y) \in \mathbb{R}^n \times ([0, +\infty]^{\bar{m}} \times \mathbb{R}^{m-\bar{m}})$, $\mathcal{J}_A(x, y)$ is a positive semidefinite matrix. Since A is continuous and monotone on $\mathbb{R}^n \times ([0, +\infty]^{\bar{m}} \times \mathbb{R}^{m-\bar{m}})$, then A is maximally monotone on $\mathbb{R}^n \times ([0, +\infty]^{\bar{m}} \times \mathbb{R}^{m-\bar{m}})$, see [5]. Moreover, since φ and ψ_1 are proper lower semicontinuous convex functions, then $A + B + C$ is a monotone operator.

(v) Following the same argument as in the case $m = 1$, we get $\zeta = \sqrt{2}$ and $\tau = \sqrt{2}L_{t_2}$.

(vi) For every $z = (x, y) \in \mathbb{R}^n \times \mathbb{R}^m$ and $\bar{z} = (\bar{x}, \bar{y}) \in \mathbb{R}^n \times \mathbb{R}^m$, we have

$$\|A(x, y) - A(\bar{x}, \bar{y})\|^2 \leq a(x, y) \|(x, y) - (\bar{x}, \bar{y})\|^2 + b \|(x, y) - (\bar{x}, \bar{y})\|^4, \quad (20)$$

with $b = \frac{5}{2} \sum_{i=1}^m L_{g_i}^2$, $a(x, y) = 2(\rho(x, y) + \sum_{i=1}^m \|\nabla g_i(x)\|^2)$, and

$$\rho(x, y) = 2 \max \left(m \max_{1 \leq i \leq m} \|\nabla g_i(x)\|^2, \left(\sum_{i=1}^m L_{g_i} |y_i| \right)^2 \right).$$

Indeed, similarly to the previous example,

$$\|A(x, y) - A(\bar{x}, \bar{y})\|^2 = \left\| \sum_{i=1}^m \nabla g_i(x) y_i - \nabla g_i(\bar{x}) \bar{y}_i \right\|^2 + \sum_{i=1}^m |g_i(\bar{x}) - g_i(x)|^2. \quad (21)$$

Moreover,

$$\begin{aligned}
 & \left\| \sum_{i=1}^m (\nabla g_i(x)y_i - \nabla g_i(\bar{x})\bar{y}_i) \right\|^2 = \left\| \sum_{i=1}^m (\nabla g_i(x)y_i - \nabla g_i(x)\bar{y}_i + \nabla g_i(x)\bar{y}_i - \nabla g_i(\bar{x})\bar{y}_i) \right\|^2 \\
 & \leq \left(\sum_{i=1}^m \|\nabla g_i(x)\| |y_i - \bar{y}_i| + \sum_{i=1}^m |\bar{y}_i| \|\nabla g_i(x) - \nabla g_i(\bar{x})\| \right)^2 \\
 & \leq \left(\max_{1 \leq i \leq m} \|\nabla g_i(x)\| \sum_{i=1}^m |y_i - \bar{y}_i| + \sum_{i=1}^m L_{g_i} |\bar{y}_i| \|x - \bar{x}\| \right)^2 \\
 & \leq 2 \left(\sqrt{m} \max_{1 \leq i \leq m} \|\nabla g_i(x)\| \|y - \bar{y}\| + \sum_{i=1}^m L_{g_i} |y_i| \|x - \bar{x}\| \right)^2 + 2 \left(\sum_{i=1}^m L_{g_i} |\bar{y}_i - y_i| \|x - \bar{x}\| \right)^2,
 \end{aligned}$$

where, in the last inequality, we used the Cauchy-Schwarz inequality. Hence,

$$\begin{aligned}
 \left\| \sum_{i=1}^m \nabla g_i(x)y_i - \nabla g_i(\bar{x})\bar{y}_i \right\|^2 & \leq \rho(x, y) (\|y - \bar{y}\| + \|x - \bar{x}\|)^2 + 2 \left(\sum_{i=1}^m L_{g_i}^2 \right) \|\bar{y} - y\|^2 \|x - \bar{x}\|^2 \\
 & \leq 2\rho(x, y) \|(x, y) - (\bar{x}, \bar{y})\|^2 + 2 \left(\sum_{i=1}^m L_{g_i}^2 \right) \|(x, y) - (\bar{x}, \bar{y})\|^4. \quad (22)
 \end{aligned}$$

On other hand, using (6) with $\nu = 1$,

$$\begin{aligned}
 \sum_{i=1}^m |g_i(\bar{x}) - g_i(x)|^2 & \leq \sum_{i=1}^m 2|\langle \nabla g_i(x), \bar{x} - x \rangle|^2 + \sum_{i=1}^m \frac{L_{g_i}^2}{2} \|x - \bar{x}\|^4 \\
 & \leq 2 \sum_{i=1}^m \|\nabla g_i(x)\|^2 \|x - \bar{x}\|^2 + \sum_{i=1}^m \frac{L_{g_i}^2}{2} \|x - \bar{x}\|^4. \quad (23)
 \end{aligned}$$

Hence, (21), (22), (23), and the fact $\|x - \bar{x}\| \leq \|(x, y) - (\bar{x}, \bar{y})\|$ yield (20). From (20) it follows that Assumption 1.vi holds with $\mu = 2$ and $\theta = 4$ (note that in this case $c(z_1) = 0$).

REMARK 2. One concrete application of the example above is the quadratically constrained quadratic program (QCQP) problem:

$$\begin{aligned}
 & \min_{x \in \mathbb{R}^n, x \geq 0} \frac{1}{2} x^\top Q_0 x + b^\top x + c \\
 & \text{s.t. } \frac{1}{2} x^\top Q_i x + l_i^\top x \leq r_i \quad \forall i \in \{1, \dots, \bar{m}\}, \quad l_i^\top x = r_i \quad \forall i \in \{\bar{m} + 1, \dots, m\}, \quad (24)
 \end{aligned}$$

where $(Q_i)_{0 \leq i \leq \bar{m}}$ are positive semidefinite matrices of dimension $n \times n$, and $(l_i)_{1 \leq i \leq m}$ and b are vectors in \mathbb{R}^n . Rewriting the QCQP into the Lagrange primal-dual form using the dual variables $y = (y_i)_{1 \leq i \leq m}$, we get

$$\begin{aligned} A : \mathbb{H} &\rightarrow \mathbb{H} : (x, y) \mapsto \left(\sum_{i=1}^m (Q_i x + l_i) y_i, \left(-\frac{1}{2} x^\top Q_i x - l_i^\top x + r_i \right)_{1 \leq i \leq m} \right) \\ B : \mathbb{H} &\rightarrow \mathbb{H} : (x, y) \mapsto (Q_0 x + b, \mathbf{0}_m) \\ C : \mathbb{H} &\rightarrow 2^{\mathbb{H}} : (x, y) \mapsto (\mathcal{N}_{[0, +\infty)^n}(x) \times (\mathcal{N}_{[0, +\infty)^{\bar{m}}}((y_i)_{1 \leq i \leq \bar{m}}) \times \{\mathbf{0}_{m-\bar{m}}\}), \end{aligned} \quad (25)$$

where we have set $\mathbb{H} = \mathbb{R}^n \times \mathbb{R}^m$ and $Q_i = 0$ for every $i \in \{\bar{m} + 1, \dots, m\}$. QCQP's have many applications, e.g., in signal processing [25], triangulation in computer vision [2], semi-supervised learning [15], learning of kernel matrices [28], or steering direction estimation for RADAR detection [20].

EXAMPLE 3. Consider the following problem:

$$\min_{x \in D} f(x), \quad (26)$$

where D is a nonempty closed convex subset of $\mathbb{H} = \mathbb{R}^n$ and $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is a continuously differentiable pseudo-convex function on D . Some examples of pseudo-convex functions are encountered in fractional programs, see below and also [23, 36]. If \bar{x} satisfies the first-order optimality condition for (26), then we have the following inclusion:

$$0 \in A\bar{x} + C\bar{x}, \quad (27)$$

where the operators are

$$Ax = \begin{cases} \nabla f(x) & \text{if } x \in D \\ \emptyset & \text{otherwise,} \end{cases} \quad B = 0, \text{ and } C = \mathcal{N}_D. \quad (28)$$

(Fractional programming) Consider the following quadratic fractional programming problem:

$$\min_{x \in D} f(x) := \frac{\frac{1}{2} x^\top Q x - h^\top x + h_0}{d^\top x + d_0} \quad \text{with } D = \{x \in \mathbb{R}^n \mid d^\top x \geq 0\}, \quad (29)$$

where $f : D \rightarrow \mathbb{R}$, $d_0 \in]0, +\infty[$, $h_0 \in \mathbb{R}$, $(h, d) \in (\mathbb{R}^n)^2$, and $Q \in \mathbb{R}^{n \times n}$ is a symmetric matrix. When the matrix Q is positive semidefinite, the function f is pseudo-convex since it is the ratio of convex

over concave functions, see [6]. Otherwise, [12, 13] present necessary and sufficient conditions for the function f to be pseudo-convex. Particular cases of (29) are problems whose objective is a sum of a linear and a linear fractional function, i.e., when $Q = (rd^\top + dr^\top)/2$, which yields the following formulation:

$$\min_{x \in D} f(x) := r^\top x + \frac{h^\top x + h_0}{d^\top x + d_0} \quad \text{with} \quad D = \{x \in \mathbb{R}^n \mid d^\top x \geq 0\}. \quad (30)$$

Indeed, setting $Q = (rd^\top + dr^\top)/2$ in (29) yields

$$\frac{\frac{1}{2}x^\top Qx - h^\top x + h_0}{d^\top x + d_0} = \frac{\frac{1}{2}x^\top rd^\top x - h^\top x + h_0}{d^\top x + d_0} = \frac{1}{2}r^\top x + \frac{h_0 - (h + \frac{d_0}{2}r)^\top x}{d^\top x + d_0}.$$

Then, by defining $\bar{r} = r/2$ and $\bar{h} = -h - \frac{d_0}{2}r$, we obtain a problem of the form (30). Reference [30] presents several cases when f is pseudo-convex over the polyhedral set $\{x \in \mathbb{R}^n \mid d^\top x + d_0 > 0\}$, namely, if $r = \eta d$, with $\eta \geq 0$, or $h = \zeta d$, with $h_0 - \zeta d_0 \geq 0$ (see [30, Theorem 1] for more details). Fractional programming arises e.g., in portfolio and transportation problems (see [30] for more details).

Below, we show that operators A , B , and C defined in (28) satisfy Assumption 1.

(i) - (iii) From the definitions of the operators A , C , we can easily see that C is maximally monotone and A is a continuous single-valued operator.

(iv) Let us show that $A + C$ is pseudo-monotone. Indeed, consider $(x, w) \in D^2$. Assume that

$$(\forall \hat{x} \in Cx) \quad \langle \nabla f(x) + \hat{x}, w - x \rangle \geq 0. \quad (31)$$

We need to show that

$$(\forall \hat{w} \in Cw) \quad \langle \nabla f(w) + \hat{w}, w - x \rangle \geq 0. \quad (32)$$

It follows from the definition of the normal cone in (4) that

$$(\forall \hat{x} \in Cx) \quad \langle \hat{x}, w - x \rangle \leq 0 \quad \text{and} \quad (\forall \hat{w} \in Cw) \quad \langle \hat{w}, x - w \rangle \leq 0. \quad (33)$$

Combining (31) and the first inequality in (33) yields

$$\langle \nabla f(x), w - x \rangle \geq 0.$$

Since f is pseudo-convex, then the above inequality implies that

$$\langle \nabla f(w), w - x \rangle \geq 0.$$

Hence, from the previous inequality and the second one in (33), we derive (32). Therefore, Assumption 1.iv holds.

(v) Note that $J_{\gamma C}z = \text{proj}_{\text{dom}C}(z)$. Using the nonexpensiveness of the projection operator, since $q = \text{proj}_{\text{dom}C}w$ and $z = q - \gamma u$, we get

$$\|\text{proj}_{\text{dom}C}(w) - \text{proj}_{\text{dom}C}(z)\| \leq \|q - z\| = \gamma \|u\|.$$

Hence, in this case the inequality holds with $\zeta = 1$ and any $\tau > 0$.

(vi) Before showing that this assumption is valid for (29), let us first derive a more focused inequality for (30).

1. Consider problem (30) and operator A defined in (28). The Hessian of f is

$$(\forall x \in D) \quad \nabla^2 f(x) = \frac{2(h^\top x + h_0)}{(d^\top x + d_0)^3} dd^\top - \frac{1}{(d^\top x + d_0)^2} (dh^\top + hd^\top).$$

Consider $(x, \bar{x}) \in D^2$. By the mean value inequality, there exists $w \in (x, \bar{x})$ s.t.

$$\begin{aligned} \|\nabla f(x) - \nabla f(\bar{x})\|^2 &\leq \|\nabla^2 f(w)\|^2 \|x - \bar{x}\|^2 \\ &\leq \left(\frac{2}{(d^\top w + d_0)^3} \|d\|^2 |h^\top w + h_0| + \frac{2}{(d^\top w + d_0)^2} |d^\top h| \right)^2 \|x - \bar{x}\|^2 \\ &\leq \left(\frac{2}{d_0^3} \|d\|^2 |h^\top(x - w)| + \frac{2}{d_0^3} \|d\|^2 |h^\top x + h_0| + \frac{2}{d_0^2} |d^\top h| \right)^2 \|x - \bar{x}\|^2 \\ &\leq 2 \left[\left(\frac{2}{d_0^3} \|d\|^2 |h^\top(x - w)| \right)^2 + \left(\frac{2}{d_0^3} \|d\|^2 |h^\top x + h_0| + \frac{2}{d_0^2} |d^\top h| \right)^2 \right] \|x - \bar{x}\|^2 \\ &\leq \frac{8}{d_0^6} \|d\|^4 \|h\|^2 \|x - \bar{x}\|^4 + \frac{8}{d_0^4} \left(\frac{\|d\|^2}{d_0} |h^\top x + h_0| + |d^\top h| \right)^2 \|x - \bar{x}\|^2, \end{aligned}$$

where in the third inequality we used the fact that, since D is convex, $w \in D$, hence $d^\top w \geq 0$,

in the fourth inequality, we used the convexity of $(\cdot)^2$, and in the last one we used that $w \in (x, \bar{x})$. Hence, Assumption 1.vi holds with $a(z_1) = \frac{8}{d_0^4} \left(\frac{\|d\|^2}{d_0} |h^\top z_1 + h_0| + |d^\top h| \right)^2$, $b(z_1) =$

$\frac{8}{d_0^6} \|d\|^4 \|h\|^2$, $c(z_1) = 0$, $\mu = 2$, and $\theta = 4$.

2. Consider problem (29) and operator A defined in (28). The Hessian of f is

$$(\forall x \in D) \quad \nabla^2 f(x) = \frac{Q}{d^\top x + d_0} + \frac{(2f(x)dd^\top - (Qx - h)d^\top - d(Qx - h)^\top)}{(d^\top x + d_0)^2}.$$

Consider $(x, \bar{x}) \in D^2$. By the mean value inequality, there exists $w \in (x, \bar{x})$ s.t.

$$\|\nabla f(x) - \nabla f(\bar{x})\| \leq \|\nabla^2 f(w)\| \|x - \bar{x}\|. \quad (34)$$

After some calculations similar to those in the previous example, we get

$$\begin{aligned} \|\nabla^2 f(w)\| &\leq \frac{1}{d^\top w + d_0} \|Q\| + \frac{|w^\top Qw - 2h^\top w + 2h_0|}{(d^\top w + d_0)^3} \|d\|^2 + \frac{2\|d\| \|Qw - h\|}{(d^\top w + d_0)^2} \\ &\leq \frac{\|Q\|}{d_0} + \frac{\|d\|^2}{d_0^3} (\|Q\| \|w\|^2 + 2|h^\top w - h_0|) + \frac{2\|d\| \|Qw - h\|}{d_0^2} \\ &\leq \frac{\|Q\|}{d_0} + \frac{\|d\|^2}{d_0^3} (2\|Q\| \|w - x\|^2 + 2\|Q\| \|x\|^2 + 2|h^\top(w - x)| + 2|h^\top x - h_0|) \\ &\quad + \frac{2\|d\|}{d_0^2} (\|Qx - h\| + \|Q\| \|w - x\|) \\ &\leq \frac{\|Q\|}{d_0} + \frac{2\|d\|^2}{d_0^3} (\|Q\| \|\bar{x} - x\|^2 + \|Q\| \|x\|^2 + \|h\| \|\bar{x} - x\| + |h^\top x - h_0|) \\ &\quad + \frac{2\|d\|}{d_0^2} (\|Qx - h\| + \|Q\| \|\bar{x} - x\|) \\ &\leq \frac{\|Q\|}{d_0} + \frac{2\|d\|^2}{d_0^3} (\|Q\| \|x\|^2 + |h^\top x - h_0|) + \frac{2\|d\|}{d_0^2} \|Qx - h\| \\ &\quad + \left(\frac{2\|d\|^2}{d_0^3} \|h\| + \frac{2\|d\|}{d_0^2} \|Q\| \right) \|\bar{x} - x\| + \frac{2\|d\|^2}{d_0^3} \|Q\| \|\bar{x} - x\|^2. \end{aligned} \quad (35)$$

We deduce from (34) and (35) that

$$\begin{aligned} &\|\nabla f(x) - \nabla f(\bar{x})\|^2 \\ &\leq \frac{12\|d\|^4}{d_0^6} \|Q\|^2 \|\bar{x} - x\|^6 + 3 \left(\frac{2\|d\|^2}{d_0^3} \|h\| + \frac{2\|d\|}{d_0^2} \|Q\| \right)^2 \|\bar{x} - x\|^4 \\ &\quad + 3 \left(\frac{\|Q\|}{d_0} + \frac{2\|d\|^2}{d_0^3} (\|Q\| \|x\|^2 + |h^\top x - h_0|) + \frac{2\|d\|}{d_0^2} \|Qx - h\| \right)^2 \|\bar{x} - x\|^2. \end{aligned}$$

Therefore, Assumption 1.vi is satisfied with $\mu = 2$, $\theta = 4$, $\beta = 6$, and

$$\begin{aligned} c(z_1) &= \frac{12\|d\|^4}{d_0^6} \|Q\|^2, & b(z_1) &= 12 \left(\frac{\|d\|^2}{d_0^3} \|h\| + \frac{\|d\|}{d_0^2} \|Q\| \right)^2, \\ a(z_1) &= 3 \left(\frac{\|Q\|}{d_0} + \frac{2\|d\|^2}{d_0^3} (\|Q\| \|z_1\|^2 + |h^\top z_1 - h_0|) + \frac{2\|d\|}{d_0^2} \|Qz_1 - h\| \right)^2. \end{aligned} \quad (36)$$

From the previous discussion, one can see that our assumptions cover a broad range of optimization problems arising in applications.

4. An adaptive forward-backward-forward algorithm Adaptive methods are widely used in optimization due to their ability to simplify stepsize tuning [7, 35, 38]. Unlike previous approaches to the forward-backward-forward algorithm, such as those in [39], we introduce two novel adaptive strategies that bypass the need for computationally expensive Armijo-Goldstein stepsize rules, typically employed when the operator is assumed to be continuous [37, 39]. In both strategies, the stepsize is determined using the current iterate and the parameters that define the operator's properties.

4.1. Investigated algorithm In this section, we introduce a new algorithm for solving problem (1). Our algorithm is similar to the forward-backward-forward splitting algorithm in [39] as it also involves two explicit (forward) steps using A and B , and one implicit (backward) step using C . However, the novelty of our iterative process lies in the adaptive way we choose the stepsize γ_k at each iteration k , which is adapted to the assumptions considered on the operators A , B , and C (see Assumption 1).

Adaptive Forward-Backward-Forward Algorithm (AFBF):

1. Choose the initial estimate $x_0 \in \text{dom } C$.
2. For $k \geq 0$ do:
 - (a) Compute the stepsize $\gamma_k > 0$.
 - (b) $z_k = x_k - \gamma_k (Ax_k + Bx_k)$
 - (c) $p_k = J_{\gamma_k C} z_k$
 - (d) $q_k = p_k - \gamma_k (Ap_k + Bp_k)$
 - (e) $\hat{x}_k = q_k - z_k + x_k$
 - (f) $x_{k+1} = \text{proj}_{\text{dom } C}(\hat{x}_k)$

Typically, to prove the convergence of a forward-backward-forward splitting algorithm, one needs the operators A and B to satisfy a Lipschitz type inequality [39]:

$$\gamma_k^2 \|Ax_k + Bx_k - Ap_k - Bp_k\|^2 \leq \alpha_k \|x_k - p_k\|^2, \quad (37)$$

where $\alpha_k \in]\alpha_{\min}, \alpha_{\max}[\subset]0, 1[$, $k \in \mathbb{N}$. In our case it appears difficult to find a positive stepsize γ_k satisfying (37) as the operator A is not assumed to be Lipschitz. However, imposing appropriate assumptions on the operator A (e.g., some generalized Lipschitz type inequality as in Assumption 1.vi), we will show that we can ensure (37). In the next sections we provide two adaptive choices for γ_k that enable us to prove the convergence of AFBF.

4.2. First adaptive choice for the stepsize In this section, we design a novel strategy to choose γ_k when the operator A satisfies Assumption 1.vi with $\mu = 2$. Recall that $(\zeta, \tau) \in]0, +\infty[^2$ are the parameters which satisfying Assumption 1.v.

Stepsize Choice 1:

1. Choose $0 < \alpha_{\min} \leq \alpha_{\max} < 1$ and $\sigma > 0$.
2. For $k \geq 0$ do:
 - (a) Compute $d(x_k) = \zeta \|Ax_k + Bx_k\| + \tau$ and choose $\alpha_k \in [\alpha_{\min}, \alpha_{\max}]$.
 - (b) Choose γ_k such that

$$\gamma_k \in \begin{cases} [\sigma, \bar{\gamma}_k] & \text{if } \sigma \leq \bar{\gamma}_k \\ \bar{\gamma}_k & \text{otherwise,} \end{cases} \quad (38)$$

where $\bar{\gamma}_k > 0$ is the root of the following equation in γ :

$$b(x_k)d(x_k)^{\theta-2}\gamma^\theta + c(x_k)d(x_k)^{\beta-2}\gamma^\beta + (L_B^2 + a(x_k))\gamma^2 = \frac{\alpha_k}{2}. \quad (39)$$

REMARK 3. Stepsize choice (38) could be just considered $\gamma_k = \bar{\gamma}_k$. However, the positive parameter σ allows us to choose a larger range of stepsizes.

Note that equation (39), which is a polynomial equation when μ , θ , and β are integers, is well defined, i.e., there exists $\bar{\gamma}_k > 0$ satisfying equation (39). Indeed, define

$$(\forall \gamma \in [0, +\infty)) \quad h(\gamma) = 2b(x_k)d(x_k)^{\theta-2}\gamma^\theta + 2c(x_k)d(x_k)^{\beta-2}\gamma^\beta + 2(a(x_k) + L_B^2)\gamma^2 - \alpha_k,$$

and $w_k = \sqrt{\alpha_k}/L_B$. Note that, we have $h(w_k) \geq \alpha_k > 0$ and $h(0) < 0$. Since h is continuous on $[0, w_k]$, there exists $\bar{\gamma}_k \in]0, w_k[$ such that $h(\bar{\gamma}_k) = 0$. Moreover, since $h'(\gamma) \geq 4L_B^2\gamma > 0$ for every $\gamma \in]0, +\infty[$, then h is strictly increasing on $]0, +\infty[$. Hence, there exists exactly one $\bar{\gamma}_k > 0$ such that the equality in (39) is satisfied and $h(\gamma_k) \leq 0 = h(\bar{\gamma}_k)$ for γ_k defined in (38).

LEMMA 1. *Let Assumption 1 hold with $\mu = 2$. Let $k \geq 0$ and let γ_k be given by (38). Then, inequality (37) is satisfied and*

$$\gamma_k < \eta := \sqrt{\frac{\alpha_{\max}}{2L_B^2}}. \quad (40)$$

Proof From basic properties of the norm,

$$\gamma_k^2 \|Ax_k + Bx_k - Ap_k - Bp_k\|^2 \leq 2\gamma_k^2 \|Bx_k - Bp_k\|^2 + 2\gamma_k^2 \|Ax_k - Ap_k\|^2.$$

Using the Lipschitz continuity of operator B on $\text{dom } C$ and Assumption 1.vi for $\mu \in]0, +\infty[$, we get

$$\begin{aligned} & \gamma_k^2 \|Ax_k + Bx_k - Ap_k - Bp_k\|^2 \\ & \leq 2\gamma_k^2 L_B^2 \|x_k - p_k\|^2 + 2\gamma_k^2 a(x_k) \|x_k - p_k\|^\mu + 2\gamma_k^2 b(x_k) \|x_k - p_k\|^\theta + 2\gamma_k^2 c(x_k) \|x_k - p_k\|^\beta \\ & = 2\gamma_k^2 \left(L_B^2 + a(x_k) \|x_k - p_k\|^{\mu-2} + b(x_k) \|x_k - p_k\|^{\theta-2} + c(x_k) \|x_k - p_k\|^{\beta-2} \right) \|x_k - p_k\|^2. \end{aligned} \quad (41)$$

Using (7) with $q = x_k$, $w = \hat{x}_{k-1}$, $u = Ax_k + Bx_k$, and $\gamma = \gamma_k$,

$$\|x_k - p_k\| = \|\text{proj}_{\text{dom } C}(\hat{x}_{k-1}) - J_{\gamma_k C} z_k\| \stackrel{(7)}{\leq} \gamma_k (\zeta \|Ax_k + Bx_k\| + \tau) = \gamma_k d(x_k). \quad (42)$$

From (39), (41), (42), and the fact that $h(\gamma_k) \leq h(\bar{\gamma}_k)$, for $\mu = 2$, we deduce that

$$\begin{aligned} & \gamma_k^2 \|Ax_k + Bx_k - Ap_k - Bp_k\|^2 \\ & \stackrel{(41),(42)}{\leq} 2(L_B^2 \gamma_k^2 + a(x_k) \gamma_k^2 + b(x_k) d(x_k)^{\theta-2} \gamma_k^\theta + c(x_k) d(x_k)^{\beta-2} \gamma_k^\beta) \|x_k - p_k\|^2 \\ & \leq 2 \left((L_B^2 + a(x_k)) \bar{\gamma}_k^2 + b(x_k) d(x_k)^{\theta-2} \bar{\gamma}_k^\theta + c(x_k) d(x_k)^{\beta-2} \bar{\gamma}_k^\beta \right) \|x_k - p_k\|^2 \\ & \stackrel{(39)}{=} \alpha_k \|x_k - p_k\|^2. \end{aligned} \quad (43)$$

From the above inequality, the first statement holds. Moreover, from (39), since $a(x_k)$, $b(x_k)$, $c(x_k)$, and $d(x_k)$ are nonnegative for every $k \geq 0$, we have $2L_B^2 \bar{\gamma}_k^2 - \alpha_k \leq 0$. Since $\alpha_k < \alpha_{\max}$, for every $k \geq 0$, inequality (40) holds. \square

From previous examples, one can see that Stepsize Choice 1 requires the computation of a positive root of a second-order polynomial equation for quadratically constrained quadratic programs in (24), while for quadratic over linear fractional programs (29), one needs to compute the positive root of a third-order equation. More explicitly:

(i) If we consider the quadratically constrained quadratic program (24), then the operator A defined

in (25) for problem (24) satisfies (20) where, for every $i \in \{1, \dots, m\}$ $g_i: x \mapsto \frac{1}{2}x^\top Q_i x + l_i^\top x - r_i$. Hence, equation (39) becomes:

$$b d(x_k)^2 \gamma^4 + (\|Q_0\|^2 + a(x_k)) \gamma^2 - \frac{\alpha_k}{2} = 0, \quad (44)$$

with the functions a and b given in (20). Defining, $\hat{\gamma} = \gamma^2$ we have

$$c d(x_k)^2 \hat{\gamma}^2 + (\|Q_0\|^2 + b(x_k)) \hat{\gamma} - \frac{\alpha_k}{2} = 0.$$

Note that, the positive root of the second order equation is

$$\hat{\gamma}_k = \frac{-(\|Q_0\|^2 + b(x_k)) + \sqrt{(\|Q_0\|^2 + b(x_k))^2 + 2\alpha_k c d(x_k)^2}}{2c d(x_k)^2}.$$

Then, using $\hat{\gamma}_k = \gamma_k^2$, we obtain that the positive root of (44) is

$$\bar{\gamma}_k = \left(\frac{\sqrt{(\|Q_0\|^2 + b(x_k))^2 + 2c d(x_k)^2 \alpha_k} - (\|Q_0\|^2 + b(x_k))}{2c d(x_k)^2} \right)^{1/2}.$$

(ii) For the quadratic fractional program (29), equation (39) becomes

$$c(x_k) d(x_k)^4 \gamma^6 + b(x_k) d(x_k)^2 \gamma^4 + (a(x_k) + L_B^2) \gamma^2 - \frac{\alpha_k}{2} = 0,$$

where functions a , b , and c are given in (36). Setting $\eta = \gamma^2$, we obtain a cubic equation with a positive root η_k , and then $\bar{\gamma}_k = \sqrt{\eta_k}$.

4.2.1. Convergence results under pseudo-monotonicity Next, we show the asymptotic convergence of the sequences $(x_k)_{k \in \mathbb{N}}$ and $(p_k)_{k \in \mathbb{N}}$ generated by AFBF Algorithm when $\mu = 2$ and the stepsize is computed according to (38). The following sequence will play a key role in our convergence analysis:

$$(\forall k \in \mathbb{N}) \quad u_k = \gamma_k^{-1} (z_k - p_k) + A p_k + B p_k \in A p_k + B p_k + C p_k. \quad (45)$$

THEOREM 1. *Suppose that $\text{zer}(A + B + C) \neq \emptyset$ and Assumption 1 holds with $\mu = 2$. Let $(x_k)_{k \in \mathbb{N}}$, $(z_k)_{k \in \mathbb{N}}$, $(p_k)_{k \in \mathbb{N}}$, and $(q_k)_{k \in \mathbb{N}}$ be sequences generated by AFBF algorithm with stepsizes $(\gamma_k)_{k \in \mathbb{N}}$ given by (38). Then, the following hold:*

- i) $(x_k)_{k \in \mathbb{N}}$ is a Fejèr monotone sequence with respect to $\text{zer}(A + B + C)$;
- ii) $\sum_{k=0}^{+\infty} \|x_k - p_k\|^2 < +\infty$ and $\sum_{k=0}^{+\infty} \|z_k - q_k\|^2 < +\infty$;
- iii) there exists $\bar{z} \in \text{zer}(A + B + C)$ such that $x_k \rightarrow \bar{z}$, $p_k \rightarrow \bar{z}$, and $u_k \rightarrow 0$.

Proof i) Let $k \in \mathbb{N}$ and let $\bar{z} \in \text{zer}(A + B + C)$. Then,

$$\begin{aligned}
\|x_k - \bar{z}\|^2 &= \|x_k - p_k + p_k - \bar{z}\|^2 \\
&= \|x_k - p_k\|^2 + 2\langle x_k - p_k, p_k - \bar{z} \rangle + \|p_k - \hat{x}_k + \hat{x}_k - \bar{z}\|^2 \\
&= \|x_k - p_k\|^2 + \|p_k - \hat{x}_k\|^2 + \|\hat{x}_k - \bar{z}\|^2 + 2\langle x_k - p_k, p_k - \bar{z} \rangle + 2\langle p_k - \hat{x}_k, \hat{x}_k - \bar{z} \rangle \\
&= \|x_k - p_k\|^2 - \|p_k - \hat{x}_k\|^2 + \|\hat{x}_k - \bar{z}\|^2 + 2\langle x_k - p_k, p_k - \bar{z} \rangle + 2\langle p_k - \hat{x}_k, p_k - \bar{z} \rangle \\
&= \|x_k - p_k\|^2 - \|p_k - \hat{x}_k\|^2 + \|\hat{x}_k - \bar{z}\|^2 + 2\langle x_k - \hat{x}_k, p_k - \bar{z} \rangle.
\end{aligned}$$

Moreover, using $p_k - \hat{x}_k = \gamma_k (Ap_k + Bp_k - Ax_k - Bx_k)$, we deduce that

$$\|x_k - \bar{z}\|^2 = \|x_k - p_k\|^2 - \gamma_k^2 \|Ap_k + Bp_k - Ax_k - Bx_k\|^2 + \|\hat{x}_k - \bar{z}\|^2 + 2\langle x_k - \hat{x}_k, p_k - \bar{z} \rangle. \quad (46)$$

Note that $\bar{z} \in \text{dom}C$. Using the nonexpansiveness of the projection, (46) yields

$$\begin{aligned}
\|x_{k+1} - \bar{z}\|^2 &\leq \|\hat{x}_k - \bar{z}\|^2 \\
&\stackrel{(46)}{=} \|x_k - \bar{z}\|^2 - \|x_k - p_k\|^2 + \gamma_k^2 \|Ap_k + Bp_k - Ax_k - Bx_k\|^2 - 2\langle x_k - \hat{x}_k, p_k - \bar{z} \rangle. \quad (47)
\end{aligned}$$

We deduce from Lemma 1 that

$$\|x_{k+1} - \bar{z}\|^2 \leq \|x_k - \bar{z}\|^2 - (1 - \alpha_k) \|x_k - p_k\|^2 - 2\langle x_k - \hat{x}_k, p_k - \bar{z} \rangle. \quad (48)$$

Since $z_k \in (\text{Id} + \gamma_k C)p_k$, the inclusion relation in (45) holds and

$$x_k - \hat{x}_k = \gamma_k u_k. \quad (49)$$

Since $A + B + C$ is pseudo-monotone and \bar{z} is a zero of $A + B + C$, we obtain:

$$\langle u_k, p_k - \bar{z} \rangle \geq 0. \quad (50)$$

Using the last inequality with (49) and $\alpha_k \leq \alpha_{\max} < 1$, it follows from (48) that

$$\|x_{k+1} - \bar{z}\|^2 \leq \|x_k - \bar{z}\|^2 - (1 - \alpha_{\max}) \|x_k - p_k\|^2 \leq \|x_k - \bar{z}\|^2. \quad (51)$$

This shows that $(x_k)_{k \in \mathbb{N}}$ is a Fejèr monotone sequence w.r.t. $\text{zer}(A + B + C)$.

ii) Since $(x_k)_{k \in \mathbb{N}}$ is a Fejèr monotone sequence, then it is bounded and (51) yields

$$(1 - \alpha_{\max}) \sum_{j=0}^k \|x_j - p_j\|^2 \leq \|x_0 - \bar{z}\|^2 < +\infty. \quad (52)$$

It follows that $(p_k)_{k \in \mathbb{N}}$ is also bounded. In addition, by using Steps 2.(b) and 2.(e) of AFBF algorithm, Lemma 1, Cauchy-Schwarz inequality, and the fact that $\alpha_k \leq \alpha_{\max}$, we further get

$$\begin{aligned} \|z_k - q_k\|^2 &\stackrel{2.(e),(49)}{=} \gamma_k^2 \|u_k\|^2 \stackrel{2.(b),(45)}{=} \|x_k - p_k + \gamma_k(Ap_k + Bp_k - Ax_k - Bx_k)\|^2 \\ &= \|x_k - p_k\|^2 + 2\gamma_k \langle x_k - p_k, Ap_k + Bp_k - Ax_k - Bx_k \rangle \\ &\quad + \gamma_k^2 \|Ap_k + Bp_k - Ax_k - Bx_k\|^2 \\ &\leq \|x_k - p_k\|^2 + 2\gamma_k \|x_k - p_k\| \|Ap_k + Bp_k - Ax_k - Bx_k\| \\ &\quad + \gamma_k^2 \|Ap_k + Bp_k - Ax_k - Bx_k\|^2 \\ &\leq (1 + \sqrt{\alpha_k})^2 \|x_k - p_k\|^2. \end{aligned} \quad (53)$$

As a consequence of (52) and the boundedness of $(\alpha_k)_{k \in \mathbb{N}}$, $\sum_{k=0}^{+\infty} \|z_k - q_k\|^2 < +\infty$.

iii) Let u_k be defined by (45). According to (53), since $\alpha_k \in]0, 1[$,

$$\|u_k\| \leq \left(\gamma_k^{-1}\right) (1 + \sqrt{\alpha_k}) \|x_k - p_k\| \leq 2\gamma_k^{-1} \|x_k - p_k\|. \quad (54)$$

On other hand, from the definition of γ_k and Lemma 1, it follows that

$$\begin{aligned} \alpha_{\min} &\leq \alpha_k \quad (55) \\ &= 2\gamma_k^2 \left(L_B^2 + a(x_k) + b(x_k)d(x_k)^{\theta-2} \bar{\gamma}_k^{\theta-2} + c(x_k)d(x_k)^{\beta-2} \bar{\gamma}_k^{\beta-2} \right) \\ &\stackrel{(40)}{\leq} 2\bar{\gamma}_k^2 \left(L_B^2 + a(x_k) + b(x_k)d(x_k)^{\theta-2} \eta^{\theta-2} + c(x_k)d(x_k)^{\beta-2} \eta^{\beta-2} \right). \end{aligned}$$

Since A, B, a, b , and c are continuous on $\text{dom } C$ and, $(x_k)_{k \in \mathbb{N}}$ and $(\gamma_k)_{k \in \mathbb{N}}$ are bounded, then $(d_k)_{k \in \mathbb{N}}$ is bounded and there exist $(R_1, R_2, R_3) \in]0, +\infty[^3$ such that

$$a(x_k) \leq R_1, \quad b(x_k)d(x_k)^{\theta-2} \leq R_2, \quad \text{and} \quad c(x_k)d(x_k)^{\beta-2} \leq R_3.$$

This allows us to lower-bound γ_k as follows:

$$\gamma_k \geq \gamma_{\min} := \min \left\{ \sigma, \sqrt{\frac{\alpha_{\min}}{2(L_B^2 + R_1 + R_2 \eta^{\theta-2} + R_3 \eta^{\beta-2})}} \right\}. \quad (56)$$

Hence, from (54), we deduce that

$$\|u_k\| \leq 2\gamma_{\min}^{-1}\|x_k - p_k\|. \quad (57)$$

As (52) implies that $x_k - p_k \rightarrow 0$, we have

$$u_k \rightarrow 0. \quad (58)$$

To prove the convergence of $(x_k)_{k \in \mathbb{N}}$, according to the Fejèr-monotone convergence theorem [9, Lemma 6], applied in our finite dimensional setting, where strong and weak convergences are equivalent, it is sufficient to show that every sequential cluster point of $(x_k)_{k \in \mathbb{N}}$ is a zero of $A + B + C$. Let w be such a sequential cluster point. There thus exists a subsequence $(x_{k_n})_{n \in \mathbb{N}}$ of $(x_k)_{k \in \mathbb{N}}$ such that $x_{k_n} \rightarrow w$. It follows from (52) and (58) that

$$p_{k_n} \rightarrow w \quad \text{and} \quad u_{k_n} \rightarrow 0$$

Since A and B are continuous operators on $\text{dom } C$, $u_{k_n} - Ap_{k_n} - Bp_{k_n} \rightarrow -Aw - Bw$. It follows from (45) that $(p_{k_n}, u_{k_n} - Ap_{k_n} - Bp_{k_n})$ lies in $\text{gra } C$. Maximally monotonicity of C implies that $(w, -Aw - Bw) \in \text{gra } C$ [5, Proposition 20.38(iii)]. Thus, $w \in \text{zer}(A + B + C)$. Hence $x_k \rightarrow w$ and, since $x_k - p_k \rightarrow 0$, $(p_k)_{k \in \mathbb{N}}$ has the same limit. This concludes our proof. \square

In [3, 11, 35], the problem of finding a zero of the sum of two operators B and C is considered when B is Lipschitz continuous, C is maximally monotone, and $B + C$ satisfies the weak Minty condition. Next, we analyze the case when we replace the pseudo-monotonicity assumption with the weak Minty condition. Let us first recall this condition.

DEFINITION 5. An operator $T : \mathbb{H} \rightarrow 2^{\mathbb{H}}$ satisfies the weak Minty condition on $\mathcal{Z} \subset \mathbb{H}$ if there exists some $\rho \geq 0$ such that the following holds:

$$\langle \hat{w}, w - z \rangle \geq -\rho \|\hat{w}\|^2 \quad \text{for every } z \in \mathcal{Z}, w \in \mathbb{H}, \text{ and } \hat{w} \in Tw. \quad (59)$$

Note that pseudo-monotone operators (see Definition 4) satisfy the weak Minty condition on their set of zeros \mathcal{Z} with $\rho = 0$. Weak Minty condition covers, in particular, minimization problems having star-convex or quasar-convex differentiable objective functions [24].

REMARK 4.

- i) First, one can notice that our proof works with a weak Minty type condition, where $\mathcal{Z} = \text{zer}(A + B + C)$ and $\rho = 0$, instead of Assumption 1.iv. Indeed, in the proof of Theorem 1, the pseudo-monotonicity of $A + B + C$ has been used to derive inequality (50), which can also be derived from the weak Minty condition with $\rho = 0$.

ii) Second, let us replace the pseudo-monotone condition in Assumption 1.iv with the assumption that $A+B+C$ satisfies the weak Minty condition on $\text{zer}(A+B+C)$ with $\rho > 0$ and, additionally, assume $\text{dom } C$ bounded. From the continuity of A , B , a , b , and c , and the boundedness of $\text{dom } C$, there exists $(R_a, R_b, R_c) \in]0, +\infty[^3$ such that, for every $z \in \text{dom } C$,

$$a(z) \leq R_a, \quad b(z)d(z) \leq R_b, \quad \text{and} \quad c(z)d(z) \leq R_c. \quad (60)$$

Then, the results from the last theorem hold as long as the following conditions are satisfied:

$$\rho < \frac{2^{-\frac{3}{2}} \sqrt{\alpha_{\min}} (1 - \sqrt{\alpha_{\max}})}{(1 + \sqrt{\alpha_{\max}}) \sqrt{L_B^2 + R_a + R_b \eta^{\theta-2} + R_c \eta^{\beta-2}}} := \rho_{\max}, \quad (61)$$

and either

$$\sigma \geq \sqrt{\frac{\alpha_{\min}}{2(L_B^2 + R_a + R_b \eta^{\theta-2} + R_c \eta^{\beta-2})}}, \quad \text{or} \quad (\forall k \geq 0) \gamma_k = \bar{\gamma}_k. \quad (62)$$

Indeed, from (48), (49) and (59),

$$(\forall k \in \mathbb{N}) \quad \|x_{k+1} - \bar{z}\|^2 \leq \|x_k - \bar{z}\|^2 - (1 - \alpha_k) \|x_k - p_k\|^2 + 2\gamma_k \rho \|u_k\|^2.$$

Using (54), we obtain

$$\|x_{k+1} - \bar{z}\|^2 \leq \|x_k - \bar{z}\|^2 - (1 - \alpha_k - 2\gamma_k^{-1} \rho (1 + \sqrt{\alpha_k})^2) \|x_k - p_k\|^2. \quad (63)$$

On other hand, (38), (55), (60), (61), and (62) yield

$$\begin{aligned} \gamma_k \geq \min \{ \sigma, \bar{\gamma}_k \} &\geq \sqrt{\frac{\alpha_{\min}}{2(L_B^2 + R_a + R_b \eta^{\theta-2} + R_c \eta^{\beta-2})}} \\ &\geq \frac{2\rho_{\max}(1 + \sqrt{\alpha_{\max}})}{1 - \sqrt{\alpha_{\max}}} \geq \frac{2\rho_{\max}(1 + \sqrt{\alpha_k})}{1 - \sqrt{\alpha_k}}. \end{aligned}$$

Hence, it follows that

$$1 - \alpha_k - 2\gamma_k^{-1} \rho (1 + \sqrt{\alpha_k})^2 \geq \left(1 - \frac{\rho}{\rho_{\max}}\right) (1 - \alpha_k) \geq \left(1 - \frac{\rho}{\rho_{\max}}\right) (1 - \alpha_{\min}).$$

The inequality above and (63) show that $(x_k)_{k \in \mathbb{N}}$ is a Fejèr monotone sequence with respect to $\text{zer}(A+B+C)$ and $\sum_{k=0}^{+\infty} \|x_k - p_k\|^2 < +\infty$. By proceeding similarly to the proof of Theorem 1, the convergence of $(x_k)_{k \in \mathbb{N}}$ to a zero of $A+B+C$ can be proved.

Now, we show a sublinear convergence rate result for the iterates of AFBF algorithm.

THEOREM 2. *Under the same assumptions as in Theorem 1, the following hold: for every $k_0 \in \mathbb{N}$ and $k \in \mathbb{N}^*$,*

$$\begin{aligned} \gamma_{\min} \min_{k_0 \leq j \leq k_0+k-1} \|u_j\| &\leq \min_{k_0 \leq j \leq k_0+k-1} \|x_j - \hat{x}_j\| \\ &\leq (1 + \sqrt{\alpha_{\max}}) \min_{k_0 \leq j \leq k_0+k-1} \|x_j - p_j\| \leq \frac{\varepsilon_{k_0}}{\sqrt{k}}, \end{aligned}$$

where u_k is defined in (45) and $\varepsilon_{k_0} \rightarrow 0$ as $k_0 \rightarrow +\infty$.

Proof According to (49), (54), and (56),

$$(\forall j \in \mathbb{N}) \quad \gamma_{\min} \|u_j\| \leq \|x_j - \hat{x}_j\| \leq (1 + \sqrt{\alpha_{\max}}) \|x_j - p_j\|. \quad (64)$$

Let \bar{z} be the limit of $(x_j)_{j \in \mathbb{N}}$. It follows from (51) that

$$(1 - \alpha_{\max}) \sum_{j=k_0}^{k_0+k-1} \|x_j - p_j\|^2 \leq \|x_{k_0} - \bar{z}\|^2,$$

which leads to

$$\min_{k_0 \leq j \leq k_0+k-1} \|x_j - p_j\|^2 \leq \frac{1}{(1 - \alpha_{\max})k} \|x_{k_0} - \bar{z}\|^2.$$

The result follows from the latter equation and (64), by setting

$$\varepsilon_{k_0} = \frac{1}{\sqrt{1 - \alpha_{\max}}} \|x_{k_0} - \bar{z}\|.$$

Hence, the statement follows. \square

Note that convergence results in Theorems 1 and 2 are consistent with those obtained in the literature on (non)monotone inclusion problems [8, 17, 19, 26, 39].

4.2.2. Convergence results under uniform pseudo-monotonicity In this section, we refine our convergence results when the operator $A + B + C$ is uniformly pseudo-monotone. Next, we present the definition of a uniformly monotone/pseudo-monotone operator.

DEFINITION 6. Let $T: \mathbb{H} \rightarrow 2^{\mathbb{H}}$.

- i) T is said to be uniformly monotone with modulus $q \geq 1$ if there exists a constant $\nu > 0$ such that, for every $(x, y) \in \mathbb{H}^2$ and $(\hat{x}, \hat{y}) \in Tx \times Ty$,

$$\langle \hat{x} - \hat{y}, x - y \rangle \geq \frac{\nu}{2} \|x - y\|^q.$$

- ii) T is said to be uniformly pseudo-monotone with modulus $q \geq 1$ if there exists a constant $\nu > 0$ such that, for every $(x, y) \in \mathbb{H}$ and $(\hat{x}, \hat{y}) \in Tx \times Ty$,

$$\langle \hat{x}, y - x \rangle \geq 0 \quad \implies \langle \hat{y}, y - x \rangle \geq \frac{\nu}{2} \|x - y\|^q.$$

When $q = 2$ in the definition above, we say that operator T is strongly monotone / pseudo-monotone. Note that, if T is uniformly monotone, then T is also uniformly pseudo-monotone.

EXAMPLE 4. Consider a proper uniformly convex function $f: \mathbb{R}^n \rightarrow]-\infty, +\infty[$. The subdifferential ∂f of f is uniformly monotone [5, Example 22.5]

Below we give an example of a strongly pseudo-monotone map that is not monotone.

EXAMPLE 5. Consider the unit ball $U = \{x \in \mathbb{R}^n \mid \|x\| \leq 1\}$ and the map $F: U \setminus \{0\} \rightarrow \mathbb{R}^n$ such that

$$(\forall x \in U \setminus \{0\}) \quad F(x) = \left(\frac{2}{\|x\|} - 1 \right) x.$$

Note that F is not monotone on $U \setminus \{0\}$. For example, setting $y = (1, 0, \dots, 0)$ and $w = (1/2, 0, \dots, 0)$ yields

$$\langle F(y) - F(w), y - w \rangle = -\frac{1}{4}.$$

However, F is strongly pseudo-monotone on $U \setminus \{0\}$. Indeed, for every $(x, y) \in (U \setminus \{0\})^2$, if $\langle F(x), y - x \rangle \geq 0$, then $\langle x, y - x \rangle \geq 0$, and consequently:

$$\langle F(y), y - x \rangle = (2\|y\|^{-1} - 1) \langle y, y - x \rangle \geq (2\|y\|^{-1} - 1) \langle y - x, y - x \rangle \geq \|y - x\|^2.$$

Next, considering operators A, B, C satisfying Assumption 1 with $\mu = 2$ and stepsizes $(\gamma_k)_{k \in \mathbb{N}}$ computed as in (38), we derive linear convergence rates when $A + B + C$ is uniformly pseudo-monotone with modulus $q \in [1, 2]$, and sublinear rates when $q > 2$.

THEOREM 3. *Suppose that Assumption 1 holds with $\mu = 2$. Let $(x_k)_{k \in \mathbb{N}}$, and $(p_k)_{k \in \mathbb{N}}$ be sequences generated by AFBF algorithm with stepsizes $(\gamma_k)_{k \in \mathbb{N}}$ given by (38). Assume that $A + B + C$ is uniformly pseudo-monotone with modulus $q \geq 1$ and constant $\nu > 0$. Then, for some $\bar{z} \in \text{zer}(A + B + C)$ and constants*

$$R = \sup_{k \in \mathbb{N}} \|p_k - \bar{z}\| < +\infty \quad \text{and} \quad r = \min\{1 - \alpha_{\max}, \gamma_{\min} \nu R^{q-2}\} < 1, \quad (65)$$

the following hold:

i) For $q \in [1, 2]$, x_k converges to \bar{z} linearly:

$$(\forall k \in \mathbb{N}) \quad \|x_k - \bar{z}\| \leq \left(1 - \frac{r}{2}\right)^{k/2} \|x_0 - \bar{z}\|. \quad (66)$$

ii) For $q > 2$ and $\bar{r} = \frac{r}{2^{q-1} R^{q-2}}$, x_k converges to \bar{z} sublinearly:

$$(\forall k \in \mathbb{N}) \quad \|x_k - \bar{z}\| \leq \frac{\|x_0 - \bar{z}\|}{\left(\frac{q-2}{2} \bar{r} \|x_0 - \bar{z}\|^{q-2} k + 1\right)^{\frac{1}{q-2}}}.$$

Proof From Theorem 1, the sequence $(p_k)_{k \in \mathbb{N}}$ generated by AFBF algorithm is convergent. Hence, for some $\bar{z} \in \text{zer}(A + B + C)$, we have $R = \sup_{k \in \mathbb{N}} \|p_k - \bar{z}\| < +\infty$. Since $A + B + C$ is uniformly pseudo-monotone with modulus q and constant $\nu > 0$,

$$(\forall k \in \mathbb{N}) \quad \langle u_k, p_k - \bar{z} \rangle \geq \frac{\nu}{2} \|p_k - \bar{z}\|^q.$$

It follows from (47) and (49) that

$$\|x_{k+1} - \bar{z}\|^2 \leq \|x_k - \bar{z}\|^2 - \|x_k - p_k\|^2 + \gamma_k^2 \|Ap_k + Bp_k - Ax_k - Bx_k\|^2 - \gamma_k \nu \|p_k - \bar{z}\|^q.$$

Since $\alpha_k \leq \alpha_{\max}$ and $\gamma_{\min} \leq \gamma_k$, we deduce from Lemma 1 that

$$\|x_{k+1} - \bar{z}\|^2 \leq \|x_k - \bar{z}\|^2 - (1 - \alpha_{\max}) \|x_k - p_k\|^2 - \gamma_{\min} \nu \|p_k - \bar{z}\|^q. \quad (67)$$

i) If $q \in [1, 2]$, using the definition of R , we get

$$\begin{aligned} & (1 - \alpha_{\max}) \|x_k - p_k\|^2 + \gamma_{\min} \nu \|p_k - \bar{z}\|^q \\ & \stackrel{(65)}{\geq} (1 - \alpha_{\max}) \|x_k - p_k\|^2 + \gamma_{\min} \nu R^{q-2} \|p_k - \bar{z}\|^2 \\ & \geq \min\{1 - \alpha_{\max}, \gamma_{\min} \nu R^{q-2}\} \left(\|x_k - p_k\|^2 + \|p_k - \bar{z}\|^2 \right) \stackrel{(65)}{\geq} \frac{r}{2} \|x_k - \bar{z}\|^2. \end{aligned}$$

Combining the two last inequalities we obtain

$$\|x_{k+1} - \bar{z}\|^2 \leq \left(1 - \frac{r}{2}\right) \|x_k - \bar{z}\|^2.$$

Therefore, unrolling the above inequality allows us to prove the first statement.

ii) If $q > 2$, it follows from (65) that

$$\begin{aligned} & (1 - \alpha_{\max}) \|x_k - p_k\|^2 + \gamma_{\min} \nu \|p_k - \bar{z}\|^q \\ & \stackrel{(65)}{\geq} \frac{(1 - \alpha_{\max})}{R^{q-2}} \|x_k - p_k\|^q + \gamma_{\min} \nu \|p_k - \bar{z}\|^q \\ & \geq \min \left\{ \frac{(1 - \alpha_{\max})}{R^{q-2}}, \gamma_{\min} \nu \right\} (\|x_k - p_k\|^q + \|p_k - \bar{z}\|^q) \geq \bar{r} \|x_k - \bar{z}\|^q. \end{aligned}$$

where in the last inequality we used the fact that $\|a + b\|^q \leq 2^{q-1} \|a\|^q + 2^{q-1} \|b\|^q$ for $q \geq 1$. Therefore, using (67), we obtain $\|x_{k+1} - \bar{z}\|^2 \leq \|x_k - \bar{z}\|^2 - \bar{r} \|x_k - \bar{z}\|^q$. Multiplying the inequality above by $\bar{r}^{\frac{2}{q-2}}$, we obtain

$$\bar{r}^{\frac{2}{q-2}} \|x_{k+1} - \bar{z}\|^2 \leq \bar{r}^{\frac{2}{q-2}} \|x_k - \bar{z}\|^2 - \left(\bar{r}^{\frac{2}{q-2}} \|x_k - \bar{z}\|^2\right)^{\frac{q}{2}}.$$

Applying [32, Lemma 8] (see Appendix) for $\zeta = \frac{q-2}{2} > 0$, we get

$$\|x_k - \bar{z}\| \leq \frac{\|x_0 - \bar{z}\|}{\left(\frac{q-2}{2}\bar{r}\|x_0 - \bar{z}\|^{q-2}k + 1\right)^{\frac{1}{q-2}}}.$$

This proves the second statement of the theorem. \square

REMARK 5. In Theorem 3, we can replace the assumption of uniform pseudo-monotonicity with the following one: there exists $\nu > 0$ and $q \geq 1$, such that, for every $w \in \mathbb{H}$, $\hat{w} \in (A + B + C)w$, and $\bar{z} \in \text{zer}(A + B + C)$, the following inequality holds:

$$\langle \hat{w}, w - \bar{z} \rangle \geq \nu \|w - \bar{z}\|^q. \quad (68)$$

Proceeding similarly to the proof of Theorem 3, linear and sublinear rates can be derived under this condition. Condition (68), with $q = 2$, covers, e.g., minimization problems with strongly star-convex differentiable objective function or strongly quasi-convex objective functions [24].

4.3. Second adaptive choice for the stepsize In this section, we present another possible *adaptive* choice for the stepsize when the operator A satisfies Assumption 1.vi with $\mu \in]0, 2[$. Let $\epsilon \in]0, 1[$ be the desired accuracy for solving problem (1), i.e., to obtain u in the range of $A + B + C$ such that $\|u\| \leq \epsilon$. The procedure is described below.

Stepsize Choice 2:

1. Choose $\epsilon \in]0, 1[$, $0 < \alpha_{\min} \leq \alpha_{\max} < 1$, and $\sigma > 0$.

2. For $k \geq 0$ do:

(a) Choose $\alpha_k \in [\alpha_{\min}, \alpha_{\max}]$ and compute $d(x_k) = \zeta \|Ax_k + Bx_k\| + \tau$.

(b) Compute $\bar{\gamma}_k^{(1)} > 0$ as the solution to the equation

$$L_B^2 \gamma^2 + b(x_k)d(x_k)^{\theta-2}\gamma^\theta + c(x_k)d(x_k)^{\beta-2}\gamma^\beta + 2^{2-\mu}a(x_k)\gamma^\mu \epsilon^{\mu-2} = \frac{\alpha_k}{2} \quad (69)$$

(c) Compute $\bar{\gamma}_k^{(2)} > 0$ as the solution to the equation

$$L_B^2 d(x_k)^{2-\mu}\gamma^2 + b(x_k)d(x_k)^{\theta-\mu}\gamma^\theta + c(x_k)d(x_k)^{\beta-\mu}\gamma^\beta + a(x_k)\gamma^\mu = \frac{\epsilon^{2-\mu}}{2^{3-\mu}}\alpha_k \quad (70)$$

(d) Update

$$\bar{\gamma}_k = \min \left\{ \bar{\gamma}_k^{(1)}, \bar{\gamma}_k^{(2)} \right\} \quad (71)$$

(e) Choose γ_k such that

$$\gamma_k \in \begin{cases} [\sigma, \bar{\gamma}_k] & \text{if } \sigma \leq \bar{\gamma}_k \\ \bar{\gamma}_k & \text{otherwise.} \end{cases} \quad (72)$$

Note that γ is well-defined in Steps 2.(b) and 2.(c) of this second procedure for the choice of the stepsize, i.e., there exist unique $\bar{\gamma}_k^{(1)}, \bar{\gamma}_k^{(2)}$ satisfying (69) and (70), respectively. Indeed, consider the functions

$$\begin{aligned} h(\gamma) &= \gamma^2 L_B^2 + b(x_k)d(x_k)^{\theta-2}\gamma^\theta + c(x_k)d(x_k)^{\beta-2}\gamma^\beta + 2^{2-\mu}a(x_k)\gamma^\mu \epsilon^{\mu-2} - \frac{\alpha_k}{2} \\ r(\gamma) &= L_B^2 d(x_k)^{2-\mu}\gamma^2 + b(x_k)d(x_k)^{\theta-\mu}\gamma^\theta + c(x_k)d(x_k)^{\beta-\mu}\gamma^\beta + a(x_k)\gamma^\mu - \frac{\epsilon^{2-\mu}}{2^{3-\mu}}\alpha_k, \end{aligned}$$

and variables $w_k = \frac{\sqrt{\alpha_k}}{L_B}$ and $\bar{w}_k = \frac{\sqrt{\alpha_k}}{L_B d(x_k)^{\frac{2-\mu}{2}}}$. Note that $h(0) < 0$ and $h(w_k) \geq \alpha_k/2 > 0$. Since h is continuous on $[0, w_k]$ there exist $\bar{\gamma}_k^{(1)} \in]0, w_k[$ such that $h(\bar{\gamma}_k^{(1)}) = 0$. Moreover, since $h'(\gamma) \geq 2\gamma L_B^2 > 0$ for every $\gamma \in]0, +\infty[$, then h is strictly increasing in $(0, +\infty)$. Hence, there exists exactly one $\bar{\gamma}_k^{(1)} > 0$ such that $h(\bar{\gamma}_k^{(1)}) = 0$. Using the same arguments, we can conclude that r is strictly increasing on $]0, +\infty[$ and there exists only one $\bar{\gamma}_k^{(2)} \in]0, \bar{w}_k[$ such that $r(\bar{\gamma}_k^{(2)}) = 0$. Since both functions h and r are strictly increasing in $(0, +\infty)$, $h(0) < 0$ and $r(0) < 0$, γ_k defined in (72) satisfies the following two inequalities:

$$h(\gamma_k) \leq 0 \quad \text{and} \quad r(\gamma_k) \leq 0. \quad (73)$$

Note that

$$\bar{\gamma}_k^{(1)} \leq \eta \quad \text{and} \quad \bar{\gamma}_k^{(2)} \leq \bar{\eta} := \left(\frac{\epsilon^{2-\mu} \alpha_{\max}}{2^{3-\mu} L_B^2 \tau^{2-\mu}} \right)^{\frac{1}{2}}, \quad (74)$$

with η defined in (40), where in the second inequality we used the fact that $d(x_k) \geq \tau$. The theorem below provides a bound on the number of iterations required, for a given $\epsilon > 0$, to generate $\|u_k\| \leq \epsilon$, with u_k defined in (45).

THEOREM 4. *Let $\epsilon \in]0, 1[$. Suppose that Assumption 1 holds with $\mu \in]0, 2[$. Let $(x_k)_{k \in \mathbb{N}}$ and $(p_k)_{k \in \mathbb{N}}$ be the sequences generated by AFBF algorithm with stepsizes $(\gamma_k)_{k \in \mathbb{N}}$ given by (72). Then, for $u_k = \gamma_k^{-1}(x_k - p_k) + Ap_k + Bp_k - Ax_k - Bx_k \in Ap_k + Bp_k + Cp_k$ and $\gamma_{\min}(\epsilon) = \mathcal{O}(\epsilon^{(2-\mu)/\mu})$, performing*

$$K \geq \frac{1}{\epsilon^2} \left(\frac{(1 + \sqrt{\alpha_{\max}})^2}{\gamma_{\min}^2(\epsilon)(1 - \alpha_{\max})} \right) \|x_0 - \bar{z}\|^2$$

iterations ensures that there exists $k \in \{0, \dots, K-1\}$ such that $\|u_k\| \leq \epsilon$.

Proof i) First, consider the case when, for every $k \in \{0, \dots, K-1\}$, $\gamma_k^{-1}\|x_k - p_k\| > \epsilon/2$. We deduce from (41), (42) and (73) that

$$\begin{aligned}
 & \gamma_k^2 \|Ax_k + Bx_k - Ap_k - Bp_k\|^2 \\
 & \stackrel{(41),(42)}{\leq} 2 \left(\gamma_k^2 L_B^2 + b(x_k)d(x_k)^{\theta-2} \gamma_k^\theta + c(x_k)d(x_k)^{\beta-2} \gamma_k^\beta + 2^{2-\mu} a(x_k) \gamma_k^\mu \epsilon^{\mu-2} \right) \|x_k - p_k\|^2 \\
 & \stackrel{(73)}{\leq} \alpha_k \|x_k - p_k\|^2.
 \end{aligned}$$

Let $\bar{z} \in \text{zer}(A + B + C)$. Since $\alpha_k \leq \alpha_{\max}$, using a similar reasoning as in (47), the inequality (51) also holds when $k \in \{0, \dots, K-1\}$, for the Stepsize Choice 2. This implies that

$$\begin{aligned}
 (1 - \alpha_{\max}) \sum_{k=0}^{K-1} \|x_k - p_k\|^2 & \leq \|x_0 - \bar{z}\|^2, \\
 (\forall k \in \{0, \dots, K\}) \quad \|x_k - \bar{z}\| & \leq \|x_0 - \bar{z}\|.
 \end{aligned} \tag{75}$$

Let D be the closed ball of center \bar{z} and radius $\|x_0 - \bar{z}\|$. Since A, B, a, b , and c are continuous on $\text{dom } C$, the quantities define below take finite values:

$$\begin{aligned}
 R_1 &= \sup_{x \in D} a(x), \quad R_2 = \sup_{x \in D} b(x)d(x)^{\theta-2}, \quad R_3 = \sup_{x \in D} c(x)d(x)^{\beta-2} \\
 R_4 &= \sup_{x \in D} d(x)^{2-\mu}, \quad R_5 = \sup_{x \in D} b(x)d(x)^{\theta-\mu}, \quad R_6 = \sup_{x \in D} c(x)d(x)^{\beta-\mu}.
 \end{aligned}$$

From (74), (69) and (70), one can lower-bound the stepsize as

$$(\forall k \in \{0, \dots, K-1\}) \quad \gamma_k \geq \gamma_{\min}(\epsilon) := \min\{\gamma_{\min}^{(1)}(\epsilon), \gamma_{\min}^{(2)}(\epsilon), \sigma\},$$

with

$$\gamma_{\min}^{(1)}(\epsilon) := \left(\frac{\alpha_{\min}}{2(L_B^2 \eta^{2-\mu} + 2^{2-\mu} R_1 \epsilon^{\mu-2} + R_2 \eta^{\theta-\mu} + R_3 \eta^{\beta-\mu})} \right)^{\frac{1}{\mu}} \tag{76}$$

and

$$\gamma_{\min}^{(2)}(\epsilon) := \left(\frac{\epsilon^{2-\mu} \alpha_{\min}}{2^{3-\mu} (R_1 + L_B^2 \bar{\eta}^{2-\mu} R_4 + R_5 \bar{\eta}^{\theta-\mu} + R_6 \bar{\eta}^{\beta-\mu})} \right)^{\frac{1}{\mu}}. \tag{77}$$

Note that, if ϵ is sufficiently small, then $\gamma_{\min}(\epsilon) = \mathcal{O}(\epsilon^{(2-\mu)/\mu})$. Using (54), we finally obtain

$$(\forall k \in \{0, \dots, K-1\}) \quad \|u_k\| \leq (\gamma_{\min}(\epsilon))^{-1} (1 + \sqrt{\alpha_{\max}}) \|x_k - p_k\|,$$

which, by virtue of (75), yields

$$\min_{0 \leq k \leq K-1} \|u_k\|^2 \leq \frac{1}{K} \left(\frac{(1 + \sqrt{\alpha_{\max}})^2}{\gamma_{\min}^2(\epsilon) (1 - \alpha_{\max})} \right) \|x_0 - \bar{z}\|^2.$$

ii) Second, consider the case when, there exists $k \in \{0, \dots, K-1\}$ such that $\gamma_k^{-1} \|x_k - p_k\| \leq \epsilon/2$.

Let us prove that $\|Ap_k + Bp_k - Ax_k - Bx_k\| \leq \epsilon/2$. Indeed, we deduce from (41) that

$$\begin{aligned} & \|Ap_k + Bp_k - Ax_k - Bx_k\|^2 \\ & \leq 2L_B^2 \|x_k - p_k\|^2 + 2a(x_k) \|x_k - p_k\|^\mu + 2b(x_k) \|x_k - p_k\|^\theta + 2c(x_k) \|x_k - p_k\|^\beta \\ & \stackrel{(42)}{\leq} 2(L_B^2 d(x_k)^{2-\mu} \gamma_k^{2-\mu} + b(x_k) d(x_k)^{\theta-\mu} \gamma_k^{\theta-\mu} + c(x_k) d(x_k)^{\beta-\mu} \gamma_k^{\beta-\mu} + a(x_k)) \|x_k - p_k\|^\mu \\ & \leq 2(L_B^2 d(x_k)^{2-\mu} \gamma_k^2 + b(x_k) d(x_k)^{\theta-\mu} \gamma_k^\theta + c(x_k) d(x_k)^{\beta-\mu} \gamma_k^\beta + a(x_k) \gamma_k^\mu) \frac{\epsilon^\mu}{2^\mu} \stackrel{(73)}{\leq} \frac{\epsilon^2 \alpha_k}{4} \leq \frac{\epsilon^2}{4}. \end{aligned}$$

Hence, from the definition of u_k , applying the triangle inequality leads to $\|u_k\| \leq \epsilon$. Hence, the statement of the theorem is proved. \square

It can be noticed that the literature on convergence rates for the general inclusion problem addressed in this section is scarce. Existing results predominantly focus on the composite problem outlined in Example 1, particularly when $g = 0$ and $L = I_n$, spanning both the convex case [34] and the nonconvex one [40].

5. Simulations In this section, we evaluate the performance of our algorithm on convex quadratically constrained quadratic programs (QCQPs), see (24), using synthetic and real data. Then, we also test our algorithm on a pseudo-convex problem using synthetic data. We compare our Adaptive Forward-Backward-Forward (AFBF) algorithm to Tseng's algorithm [39], and one dedicated commercial optimization software package, Gurobi [22] (which has a specialized solver for QCQPs). We implemented the algorithm AFBF as follows: at each iteration $k \in \mathbb{N}$, the stepsize $\gamma_k = \bar{\gamma}_k$, where $\bar{\gamma}_k$ is computed as in (39), $b(x_k)$ and $c(x_k)$ are computed as in (20), and $\alpha_k = 0.99$. The code was implemented using MATLAB R2020a on a computer equipped with an AMD Ryzen CPU operating at 3.4 GHz and 64 GB of RAM.

5.1. Solving convex QCQPs We consider the following convex QCQP

$$\begin{aligned} \min_{x \in \mathbb{R}^n} f(x) &= \frac{1}{2} x^\top Q_0 x + b^\top x \\ \text{s.t. } g_i(x) &= \frac{1}{2} x^\top Q_i x + l_i^\top x - r_i \leq 0, \quad \forall i \in \{1, \dots, m\}, \end{aligned} \quad (78)$$

where $(Q_i)_{0 \leq i \leq m}$ are symmetric positive semidefinite matrices in $\mathbb{R}^{n \times n}$, $(l_i)_{1 \leq i \leq m}$ and b are vectors in \mathbb{R}^n , and $(r_i)_{1 \leq i \leq m}$ are nonnegative reals. Note that the operator A defined in (25) for QCQPs fits

(20). For every $i \in \{0, \dots, m\}$, Q_i was generated as $Q_i = R_i^\top R_i$, where $R_i \in \mathbb{R}^{p \times n}$ is a sparse random matrix whose elements are drawn independently from a uniform distribution over $[0, 1]$. Moreover, the components of vectors b and $(l_i)_{1 \leq i \leq m}$ were generated from a standard normal distribution $\mathcal{N}(0, 1)$. Constants $(r_i)_{1 \leq i \leq m}$ and the components of the algorithm starting point were generated from a uniform distribution over $[0, 1]$. For the algorithm in [39], named Tseng, the line-search is computed as in [39, equation (2.4)], with $\theta = 0.995$, $\sigma = 1$, and $\beta = 0.5$. We consider the following stopping criteria for AFBF and Tseng's algorithms:

$$\|u_k\| \leq 10^{-2}, \text{ with } u_k \text{ defined in (45).}$$

n	p	m	AFBF		Tseng [39]		
			ITER	CPU	ITER	LSE	CPU
10^3	10^3	250	3914	36.09	15298	91513	387.4
10^3	10^3	500	7563	131.8	23400	140070	1179.3
10^3	10^3	10^3	19044	597.6	37932	227029	3570.4
10^3	10^3	$2 \cdot 10^3$	44039	2900.1	63143	377963	12990
10^4	10^4	125	4705	195.5	3351	19963	418.6
10^4	10^4	250	6131	475.2	4888	29209	1178
10^4	10^4	500	8862	1329	7240	43319	3398
10^4	10^4	750	11380	1821	8670	51893	4251
10^3	500	250	4992	66.9	14750	88223	590.9
10^3	500	500	11069	288.7	25741	154114	2068.7
10^3	500	10^3	24460	1192.7	45654	273360	7010.4
10^3	500	$2 \cdot 10^3$	59762	5939	*	*	*
10^4	$5 \cdot 10^3$	125	5318	336	3428	20412	689.8
10^4	$5 \cdot 10^3$	250	7445	895.3	4762	28452	1864
10^4	$5 \cdot 10^3$	500	11515	2711	11271	67514	8647
10^4	$5 \cdot 10^3$	750	15719	3655.4	14073	84324	10462

TABLE 1. CPU time (sec) and number of iterations (ITER) for solving synthetic QCQPs of the form (78) with AFBF and Tseng's [39] algorithms: strongly convex case (top) and convex case (bottom).

The CPU time (in seconds) and the number of iterations (ITER) required by each algorithm for solving problem (78) are given in Table 5.1, where "*" means that the corresponding algorithm needs more than 5 hours to solve the problem. Moreover, for Tseng's algorithm, we also report the number of line-search evaluations (LSE). The first half of the table corresponds to strongly convex functions ($Q_i > 0$, for every $i \in \{0, \dots, m\}$) and the other half is for convex functions ($Q_i \geq 0$, for every $i \in \{0, \dots, m\}$). As we can notice in Table 5.1, AFBF outperforms Tseng's algorithm (sometimes even 10× faster). Comparisons with Gurobi software are not included in Table 5.1, since we observed that its performance is quite poor on these large test cases.

5.2. Solving multiple kernel learning in support vector machine In this section, we test AFBF on Support Vector Machine (SVM) with multiple kernel learning using real data, which can also be formulated as a convex QCQP. Let us briefly describe the problem (our presentation follows [15]). Given a set of n_{dat} data points $\mathcal{S} = \{(d_j, l_j)\}_{1 \leq j \leq n_{\text{dat}}}$ where, for every $j \in \{1, \dots, n_{\text{dat}}\}$

$d_j \in \mathbb{R}^{n_d}$ is the input vector and $l_j \in \{-1, 1\}$ is its class label, SVM searches for a hyperplane that can best separate the points from the two classes. When the data points cannot be separated in the original space \mathbb{R}^{n_d} , we can search in a feature space \mathbb{R}^{n_f} , by mapping the input data space \mathbb{R}^{n_d} to the feature space through a function $\varphi : \mathbb{R}^{n_d} \rightarrow \mathbb{R}^{n_f}$. Using function φ , we can define a kernel function $\kappa : \mathbb{R}^{n_d} \times \mathbb{R}^{n_d} \rightarrow \mathbb{R}$ as $\kappa(d_j, d_{j'}) := \langle \varphi(d_j), \varphi(d_{j'}) \rangle$ for every $(d_j, d_{j'}) \in (\mathbb{R}^{n_d})^2$, where $\langle \cdot, \cdot \rangle$ denotes the inner product of \mathbb{R}^{n_f} . One popular choice of the kernel function is the Gaussian kernel:

$$\kappa_{\text{GAU}}(d_j, d_{j'}) = \exp\left(-\frac{\|d_j - d_{j'}\|^2}{2\bar{\sigma}^2}\right), \quad \forall (j, j') \in \{1, \dots, n_{\text{dat}}\}^2$$

with $\bar{\sigma} > 0$. We separate the given set \mathcal{S} into a training set, $\mathcal{S}_{\text{tr}} = \{(d_j, l_j)\}_{1 \leq j \leq n_{\text{tr}}}$ and a testing set, $\mathcal{S}_{\text{te}} = \{(d_j, l_j)\}_{1 \leq j \leq n_{\text{te}}}$, such that $n_{\text{tr}} + n_{\text{te}} = n_{\text{dat}}$. Choosing a set of kernel functions $(\kappa_i)_{1 \leq i \leq m}$, the SVM classifier is learned by solving the following convex QCQP problem on the training set \mathcal{S}_{tr} :

$$\begin{aligned} \min_{x \in \mathbb{R}^{n_{\text{tr}}}, x_0 \in \mathbb{R}, x \geq 0} \quad & \frac{1}{2}x^\top Q_0 x - e^\top x + R x_0 \\ \text{s.t.} \quad & \frac{1}{2}x^\top \left(\frac{1}{R_i} G_i(K_{i,\text{tr}}) \right) x - x_0 \leq 0 \quad \forall i \in \{1, \dots, m\}, \quad \sum_{j=1}^{n_{\text{tr}}} l_j x_j = 0, \end{aligned} \quad (79)$$

where $Q_0 = C^{-1}I_{n_{\text{tr}}}$, C being a parameter related the soft margin criteria, and the vector e denotes a vector of all ones. In addition, for every $i \in \{1, \dots, m\}$, $K_{i,\text{tr}} \in \mathbb{R}^{n_{\text{tr}} \times n_{\text{tr}}}$ is a symmetric positive semidefinite matrix, whose (j, j') element is defined by the kernel function: $[K_{i,\text{tr}}]_{j,j'} := \kappa_i(d_j, d_{j'})$. The matrix $G_i(K_{i,\text{tr}}) \in \mathbb{R}^{n_{\text{tr}} \times n_{\text{tr}}}$ in the i -th quadratic constraint of (79) is a symmetric positive semidefinite matrix, its (j, j') element being $[G_i(K_{i,\text{tr}})]_{j,j'} = l_j l_{j'} [K_{i,\text{tr}}]_{j,j'}$. Moreover, R and $(R_i)_{1 \leq i \leq m}$ are given positive constants. Clearly, (79) is an instance of problem (24). In our experiments, we employed a predefined set of Gaussian kernel functions $(\kappa_i)_{1 \leq i \leq m}$, with the corresponding $(\bar{\sigma}_i^2)_{1 \leq i \leq m}$ values. Following the pre-processing strategy outlined in [15], we normalized each matrix $K_{i,\text{tr}}$ such that $R_i = \text{trace}(K_{i,\text{tr}})$ was set to 1, thus restricting $R = \sum_{i=1}^m R_i = m$. For each dataset, the $\bar{\sigma}_i^2$'s were set to m different grid points (generating m kernels) within the interval $[10^{-1}, 10]$ for the first five datasets and $[10^{-2}, 10^2]$ for the last one, with two different values for the number of grid points, namely $m = 3$ and $m = 5$. Additionally, we set $C = 1$. In order to give a better overview of the advantages offered by the multiple kernel SVM approach, we also learn a single Gaussian kernel SVM classifier with $\bar{\sigma}^2$ set a priori to 7, by solving the following QP problem:

$$\min_{x \in [0, C]^{n_{\text{tr}}}} \frac{1}{2}x^\top G(K_{\text{tr}})x - e^\top x, \quad \text{s.t.} \quad \sum_{j=1}^{n_{\text{tr}}} l_j x_j = 0. \quad (80)$$

We consider the following stopping criterion for AFBF and Tseng’s algorithms:

$$\max \left(|f(x) - f^*|, \left| \sum_{j=1}^{n_{tr}} l_j x_j \right|, \max_{1 \leq i \leq m} (0, g_i(x)) \right) \leq 10^{-4},$$

with f^* computed by Gurobi solver and the starting point chosen as the null vector. Moreover, for Tseng’s algorithm the line-search was computed as in [39, equation (2.4)], with $\theta = 0.99$, $\sigma = 1$, and $\beta = 0.1$. Table 5.2 presents a comparison between AFBF algorithm, Tseng’s algorithm [39], and Gurobi solver [22] in terms of CPU time for solving the QCQP of the form (79) using 6 real datasets Ozone-level-8hr, mfeat-fourier, USPS, isolet, semeion, and Ovarian from <https://www.openml.org>. Each dataset was divided into a training set comprising 80% of the data and a testing set of the remaining 20%. For each dataset, we also provided the nonzero optimal dual multiplier value y^* corresponding to the unique active quadratic inequality constraint and the corresponding value of $\bar{\sigma}^2$ corresponding to that active constraint. Finally, the table presents a comparison between the Testing Set Accuracies on the remaining testing datasets obtained by the multiple Gaussian kernel SVM classifier with $\bar{\sigma}^2$ derived from (79), named TSA, and the single Gaussian kernel SVM classifier with $\bar{\sigma}^2 = 7$, named TSA0.

Dataset (n, n_d)	m	TSA0	TSA	$\bar{\sigma}^2$	AFBF		TSENG		Gurobi	
					CPU	y^*	CPU	y^*	CPU	y^*
Ozone-level-8hr (2534, 72)	3	52.7	91.7	5.05	31.18	3.1	58.09	2.99	95.61	3
	5		91.7	2.575	49.9	5.04	61.38	5	339.88	5
mfeat-fourier (2000, 76)	3	87.7	89	5.05	11.82	3.04	21.5	2.99	40.56	3
	5		89	2.575	20.54	5.02	35.06	4.99	170.06	5
USPS (1424, 256)	3	60.2	91.5	10	4	3	5.23	3	232.98	3
	5		92.2	10	3.95	5	8.33	5	1106.7	5
isolet (600, 617)	3	57.5	95	10	0.59	3	1.35	3	10.8	3
	5		95.8	10	0.68	4.97	2.23	5	25.09	5
semeion (319, 256)	3	47.6	77.8	10	0.75	2.98	1.43	2.97	1.37	3
	5		84.1	10	0.89	5.02	3.19	4.99	4.12	5
Ovarian (253, 15154)	3	66	78	100	0.38	3.04	1.72	2.99	0.82	3
	5		88	100	0.47	4.96	2.48	4.99	2.31	5

TABLE 2. Comparison between our algorithm AFBF, Tseng’s algorithm [39] and Gurobi solver [22] in terms of CPU time (in seconds) to solve QPQCs of the form (79) for various real datasets and two different choices of $m = 3, 5$. Additionally, TSA’s are provided for (79) and (80).

5.3. Fractional programming In this final set of experiments, we consider the linear fractional program (30), where the objective function is pseudo-convex. We compare our algorithm with [37, Algorithm 1] developed for solving non-Lipschitzian and pseudo-monotone variational inequalities. We implemented [37, Algorithm 1] with the parameters $\mu = 0.995$, $\gamma = 1$, and $l = 0.001$. Note that μ in Algorithm 1 in [37] has a different meaning from our μ in Assumption 1.vi. Actually, according to Example 3.4, we have in our case $\mu = 2$ for this fractional programming problem.

Hence, we use the first choice of the stepsize. From Theorem 1 in [30], when the vector $r = \eta d$ with $\eta \geq 0$, the objective function f in (30) is pseudo-convex on D .




FIGURE 1. Evolution of Algorithm 1 in [37] (called here FBF) and our AFBF algorithm in function values along time for two linear fractional programs of the form (30) with data generated randomly, $\eta = 1$ and $\eta = 10$, and dimension $n = 10^6$.

In our simulations, the components of the vector d and the constant h_0 were drawn independently from a standard normal distribution $\mathcal{N}(0, 1)$, vector r was chosen as $r = \eta d$, with $\eta > 0$, vector h was taken as a perturbation of vector d , i.e., $h = d + 0.01v$. Vector v and constant d_0 were generated from a uniform distribution. Moreover, we chose the starting point x_0 as $x_0 = \text{proj}_D(t)$, vector t being generated from a standard normal distribution $\mathcal{N}(0, 1)$. The results are displayed in Figure 1, where we plot the evolution of function values along time (in sec). Note that, AFBF is faster than Algorithm 1 from [37] (named here FBF) for chosen values of η .

6. Conclusion In this paper, we have addressed the problem of finding a zero of a pseudo-monotone operator. We have made the assumption that this operator can be split as a sum of three operators: the first continuous operator A satisfies a generalized Lipschitz inequality, the second operator B is Lipschitzian, and the third one C is maximally monotone. For solving this challenging problem, our solution relied upon the forward-backward-forward algorithm, which requires however the use of an iteration-dependent stepsize. In this context, we designed two novel adaptize stepsize strategies. We also derived asymptotic sublinear convergence properties under the considered assumptions. Additionally, when $A + B + C$ satisfies a uniform pseudo-monotonicity condition, the convergence rate becomes even linear. Preliminary numerical results confirm the good performance of our algorithm.

For future research, it would be intriguing to investigate the possibility of achieving more precise convergence rates. For instance, in Example 1, when $g = 0$ and $L = I_n$, [34] introduces a universal

gradient method with a convergence rate of order $O(\epsilon^{-2/(1+\nu)})$ for the convex (i.e., maximally monotone) case, where ν is the constant from Definition 2 (note that $\mu = 2\nu$ in this scenario). Conversely, in the nonconvex (i.e., nonmonotone) case under the same settings, [40] examines a gradient-type method with an adaptive stepsize and achieves a convergence rate of order $O(\epsilon^{-\frac{1+\nu}{\nu}})$ in the norm of the gradient. On the other hand, the convergence rate obtained in Theorem 4 within the general nonmonotone framework we considered is of order $O(\epsilon^{-2/\nu})$ in the norm of the gradient, which is not as favorable as the rate in [40].

Appendix

[32, Lemma 8.(i)] Let $\zeta > 0$ and $(\Delta_k)_{k \geq 0}$ be a decreasing sequence of positive numbers satisfying the following recurrence:

$$(\forall k \geq 0) \quad \Delta_k - \Delta_{k+1} \geq \Delta_k^{\zeta+1}.$$

Then, $\Delta_k \rightarrow 0$ with sublinear rate:

$$(\forall k \geq 0) \quad \Delta_k \leq \frac{\Delta_0}{\left(\zeta \Delta_0^\zeta k + 1\right)^{\frac{1}{\zeta}}} \leq \left(\frac{1}{\zeta k}\right)^{\frac{1}{\zeta}}.$$

References

- [1] S. Adly, L. Bourdin, and F. Caubet. On a decomposition formula for the proximal operator of the sum of two convex functions. *Journal of Convex Analysis*, 26(3):699–718, 2019.
- [2] Chris Aholt, Sameer Agarwal, and Rekha Thomas. A qcqp approach to triangulation. In Andrew Fitzgibbon, Svetlana Lazebnik, Pietro Perona, Yoichi Sato, and Cordelia Schmid, editors, *Computer Vision – ECCV 2012*, pages 654–667, Berlin, Heidelberg, 2012. Springer Berlin Heidelberg.
- [3] A. Alacaoglu, D. Kim, and S.J. Wright. Extending the reach of first-order algorithms for nonconvex min-max problems with cohypomonotonicity, 2024. Preprint.
- [4] D. Aussel. Subdifferential properties of quasiconvex and pseudoconvex functions: unified approach. *Journal of Optimization Theory and Applications*, 97(1):29–45, 1998.
- [5] H. Bauschke and P. Combettes. *Convex analysis and monotone operator theory in Hilbert spaces*. Springer, second edition, 2017.
- [6] J. M. Borwein and A. S. Lewis. *Convex analysis and nonlinear optimization: theory and examples*. Springer Science and Business Media, 2006. New York.
- [7] R. I. Bot, E. R. Csetnek, and P. T. Vuong. The forward-backward-forward method from continuous and discrete perspective for pseudo-monotone variational inequalities in hilbert spaces. *European Journal of Operational Research*, 287(1):49–60, 2020.

- [8] L.M. Briceno-Arias and D. Davis. Forward-backward-half forward algorithm for solving monotone inclusions. *SIAM Journal on Optimization*, 28:2839–2871, 2018.
- [9] F. E. Browder. Convergence theorems for sequences of nonlinear operators in banach spaces. *Mathematische Zeitschrift*, 100:201–225, 1967.
- [10] M. N. Bui and P. L. Combettes. Multivariate monotone inclusions in saddle form. *Mathematics of Operations Research*, 47:1082–1109, 2022.
- [11] Y. Cai and W. Zheng. Accelerated single-call methods for constrained min-max optimization. In *International Conference on Learning Representations*, 2022.
- [12] A. Cambini, J.-P. Crouzeix, and Laura Martein. On the pseudoconvexity of a quadratic fractional function. *Optimization*, 51(4):677–687, 2002.
- [13] L. Carosi and L. Martein. On the pseudoconvexity and pseudolinearity of some classes of fractional functions. *Optimization*, 56:385–398, 2007.
- [14] A. Chambolle and T. Pock. A first-order primal-dual algorithm for convex problems with applications to imaging. *J. Mathematical Imaging and Vision*, 40:120–145, 2011.
- [15] R. Chen and A. L. Liu. A distributed algorithm for high-dimension convex quadratically constrained quadratic programs. *Comput. Optim. and Appl.*, 80:781–830, 2021.
- [16] P. L. Combettes. The geometry of monotone operator splitting methods. *Acta Numerica*, 33:487–632, 2024.
- [17] P. L. Combettes and J.-C. Pesquet. Primal-dual splitting algorithm for solving inclusions with mixtures of composite, lipschitzian, and parallel-sum type monotone operators. *Set-Valued*, 20:307–330, 2012.
- [18] L. Condat. A primal-dual splitting method for convex optimization involving Lipschitzian, proximable and linear composite terms. *Journal of Optimization Theory and Applications*, 158:460–479, 2013.
- [19] D. Davis and W. Yin. A three-operator splitting scheme and its optimization applications. *Set-Valued and Variational Analysis*, 25(4):829–858, 2017.
- [20] Antonio De Maio, Yongwei Huang, Daniel P. Palomar, Shuzhong Zhang, and Alfonso Farina. Fractional qcqp with applications in ml steering direction estimation for radar detection. *IEEE Transactions on Signal Processing*, 59(1):172–185, 2011.
- [21] J. Diakonikolas, C. Daskalakis, and M. Jordan. Efficient methods for structured nonconvex-nonconcave min-max optimization. In *International Conference on Artificial Intelligence and Statistics*, volume 130, pages 2746–2754, 2021.
- [22] Gurobi. Gurobi optimizer reference manual. <https://www.gurobi.com>.
- [23] A. Hassouni and A. Jaddar. On pseudoconvex functions and applications to global optimization. In *ESAIM*, volume 20, pages 138–148, 2007.
- [24] O. Hinder, A. Sidford, and N. Sohoni. Near-optimal methods for minimizing star-convex functions and beyond. In *Conference on Learning Theory*, volume 125, pages 1894–1938, 2020.

- [25] Y. Huang and D. P. Palomar. Randomized algorithms for optimal solutions of double-sided qcqp with applications in signal processing. *IEEE Transactions on Signal Processing*, 62(5):1093–1108, 2014.
- [26] X. Jiang and L. Vandenberghe. Bregman three-operator splitting methods. *Journal of Optimization Theory and Applications*, 196:936–972, 2023.
- [27] S. Karamardian and S. Schaible. Seven kinds of monotone maps. *Journal of Optimization Theory and Applications*, 66:37–46, 1990.
- [28] G. R. Lanckriet, N. Cristianini, P. L. Bartlett, L. E. Ghaoui, and M. Jordan. Learning the kernel matrix with semi-definite programming. *Journal of Machine Learning Research*, 5:27–72, 2004.
- [29] O. L. Mangasarian. Pseudo-convex functions. *SIAM Journal on Control and Optimization*, 3(2):281–290, 1965.
- [30] L. Martein and L. Carosi. The sum of a linear and a linear fractional function: pseudoconvexity on the nonnegative orthant and solution methods. *Bulletin of Malaysian Mathematical Sciences Society*, 35(2A):591–599, 2012.
- [31] Panayotis Mertikopoulos, Christos Papadimitriou, and Georgios Piliouras. Cycles in adversarial regularized learning. In *ACM-SIAM Symposium on Discrete Algorithms*, pages 2703–2717, 2018.
- [32] I. Necoara and F. Chorobura. Efficiency of stochastic coordinate proximal gradient methods on nonseparable composite optimization. *Mathematics of Operations Research*, doi: 10.1287/moor.2023.0044, 2024.
- [33] A. Nemirovski. Prox-method with rate of convergence $O(1/t)$ for variational inequalities with Lipschitz continuous monotone operators and smooth convex-concave saddle point problems. *SIAM Journal on Optimization*, 15:229–251, 2004.
- [34] Y. Nesterov. Universal gradient methods for convex optimization problems. *Mathematical Programming*, 152(1-2):381–404, 2015.
- [35] T. Pethick, P. Latafat, P. Patrinos, O. Fercoq, and V. Cevher. Escaping limit cycles: global convergence for constrained nonconvex-nonconcave minimax problems. In *International Conference on Learning Representations*, 2022.
- [36] S. Schaible. Quasiconvex, pseudoconvex, and strictly pseudoconvex quadratic functions. *Journal of Optimization Theory and Application*, 35(3):303–338, 1981.
- [37] D. V. Thong and P. T. Vuong. Modified tseng’s extragradient methods for solving pseudo-monotone variational inequalities. *Optimization*, 68:2207–2226, 2019.
- [38] B. Tongnoi. A modified tseng’s algorithm with extrapolation from the past for pseudo-monotone variational inequalities. *Taiwanese J. of Mathematics*, 28(1):187–210, 2024.
- [39] P. Tseng. A modified forward-backward splitting method for maximal monotone mappings. *SIAM Journal on Control and Optimization*, 38(2):431–446, 2000.
- [40] M. Yashtini. On the global convergence rate of the gradient descent method for functions with Holder continuous gradients. *Optimization Letters*, 10(6):1361–1370, 2016.