

Dirac operators twisted by ramified Euclidean line bundles

Gorapada Bera

Thomas Walpuski

2025-03-03

Abstract

This article is concerned with the analysis of Dirac operators D twisted by ramified Euclidean line bundles (Z, \mathbb{I}) —motivated by their relation with harmonic $\mathbb{Z}/2\mathbb{Z}$ spinors, which have appeared in various context in gauge theory and calibrated geometry. The closed extensions of D are described in terms of the Gelfand–Robbin quotient $\dot{\mathbb{H}}$. Assuming that the branching locus Z is a closed cooriented codimension two submanifold, a geometric realisation of $\dot{\mathbb{H}}$ is constructed. This, in turn, leads to an L^2 regularity theory.

Contents

1	Introduction	2
2	The Gelfand–Robbin quotient, I: abstract theory	4
2.1	The minimal and maximal extension	4
2.2	Closed extensions and residue conditions	6
2.3	The Green’s form and adjoint extensions	6
2.4	Fredholm extensions	8
2.5	Chirality operators	10
3	The Gelfand–Robbin quotient, II: geometric realisation	11
3.1	The blow-up of X along Z	12
3.2	The model operator	13
3.3	The model Gelfand–Robbin quotient	16
3.4	Spectral decomposition	18
3.5	Leading order terms	21
3.6	Assembly of the residue map	24
3.7	Spectral and local residue conditions	26
4	Regularity theory	27
4.1	Adapted Sobolev spaces, I: definition	28
4.2	Elliptic regularity and estimates	30
4.3	Fredholm extensions in higher regularity	32
4.4	Adapted Sobolev spaces, II: Morrey embedding and polyhomogeneity	33
4.5	Adapted Sobolev spaces, III: spectral description and tameness	36

1 Introduction

Let (X, g) be an oriented closed Riemannian manifold of dimension n and (S, γ, ∇) a Dirac bundle over (X, g) ; see [LM89, Chapter II Definition 5.2].

Definition 1.1. A **ramified Euclidean line bundle** over X consists of:

- (1) a closed subset $Z \subset X$, the **branching locus**, and
- (2) a Euclidean line bundle \mathbb{I} over $X \setminus Z$

such that

- (3) if $W \subset Z$ is closed and \mathbb{I} extends over $X \setminus W$, then $W = Z$. •

This article is concerned with the analysis of the Dirac operator associated with (S, γ, ∇) twisted by a ramified Euclidean line bundle \mathbb{I}

$$D: H^1\Gamma(X \setminus Z, S \otimes \mathbb{I}) \rightarrow L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})$$

and its extensions.

The authors' motivation for this stems from the following. Taubes has observed that the failure of compactness for a wide variety of generalised Seiberg–Witten equations—e.g.: stable flat $\mathrm{PSL}_2(\mathbb{C})$ -connections in dimension three [Tau13a], anti-self-dual $\mathrm{SL}_2(\mathbb{C})$ -connections in dimension four [Tau13b], the Seiberg–Witten equation with multiple spinors [Tau16], the Vafa–Witten equation [Tau17], and the Kapustin–Witten equation [Tau22]—leaves behind evidence in the form of a **harmonic $\mathbb{Z}/2\mathbb{Z}$ spinor**. The latter is a pair $(Z, \mathbb{I}; \Phi)$ consisting of a ramified Euclidean line bundle (Z, \mathbb{I}) and a harmonic spinor $\Phi \in \ker D$. Harmonic $\mathbb{Z}/2\mathbb{Z}$ spinors also appear in Donaldson's work on adiabatic limits of coassociative Kovalev–Lefschetz fibrations of G_2 -manifolds [Don17] and He's work on branched double covers of special Lagrangian submanifolds [He22].

In light of this, it is important to understand the universal moduli space of harmonic $\mathbb{Z}/2\mathbb{Z}$ spinors (allowing for g , γ , and ∇ to vary). The fundamental issue is that D is only left semi-Fredholm (under mild assumptions; see Hypothesis 2.1), but not Fredholm—except in edge cases, e.g., if $Z = \emptyset$ or $n = 2$. The naive expectation is that the ∞ -dimensional cokernel of D can be compensated by wiggling the branching locus Z . In his PhD thesis, Takahashi [Tak15; Tak17] has made some initial progress in this direction. Donaldson [Don21] and Parker [Par23] have developed a (partial) deformation theory for harmonic $\mathbb{Z}/2\mathbb{Z}$ 1-forms and harmonic $\mathbb{Z}/2\mathbb{Z}$ spinors on spin 3-manifolds respectively. There is work in progress by He, Parker and Walpuski to address this problem a bit more systematically. The present article should be considered infrastructure for this project (and, hopefully, other applications as well).

Here is a summary of the results contained in this article. Section 2 considers D as an unbounded operator D_{\min} on $L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})$, the **minimal extension**, and systematically studies its closed extensions. The adjoint $D_{\max} := D_{\min}^*$ is the **maximal extension** of D_{\min} . The closed

extensions D_B of D_{\min} are classified by **residue conditions**, that is: closed subspaces $B \subset \check{H}$ of the **Gelfand–Robbin quotient**

$$\check{H} := \frac{\text{dom}(D_{\max})}{\text{dom}(D_{\min})}.$$

Moreover, \check{H} is equipped with a symplectic form G , the **Green’s form**, which controls the formation of adjoints. Within this framework it is also possible to describe which extensions D_B are Fredholm. The entire discussion only relies on D_{\min} being closed, densely defined, and symmetric as well as left semi-Fredholm. It is confined to the realm of abstract functional analysis and its purpose is to separate what is true for formal reasons from what is true for geometric reasons. Most of the observations in [Section 2](#) can be found in [[MS98](#), Exercise 2.17; [BF98](#), §3; [SWo8](#), Appendix B; [BS18](#), Exercises 6.3.3 and 6.5.11] in some shape or form.

Assuming that $Z \subset X$ is a closed cooriented submanifold of codimension two, [Section 3](#) constructs an isomorphism of symplectic Hilbert spaces

$$\text{res}: (\check{H}, G) \cong (\check{H}\Gamma(Z, \check{S}), \check{Q}),$$

the **residue map**, between the Gelfand–Robbin quotient and a Hilbert space of sections of a symplectic vector bundle over Z . The residue map extracts the leading order behavior of $\phi \in \text{dom}(D_{\max})$ which is shown to be (at worst) comparable to $\bar{z}^{-1/2}$ transversely to Z . With the help of the above it is possible to define spectral residue conditions, analogous to the APS boundary condition [[APS75](#)], as well as local residue conditions. As by product this yields a variant of the bordism theorem, whose significance remains somewhat mysterious to the authors. Evidently the construction in [Section 3](#) is inspired by Bär and Ballmann’s magnificent article [[BB12](#)] on boundary value problems for Dirac operators.

Finally, [Section 4](#) develops an L^2 regularity theory on the scale of **adapted Sobolev spaces** $(H_a^k\Gamma(X \setminus Z, S \otimes I))_{k \in \mathbb{N}_0}$. This scale is defined via the ring of differential operators generated by conormal differential operators and the Dirac operator D . It gives rise to a graded Fréchet space $H_a^\infty\Gamma(X \setminus Z, S \otimes I)$ which is tame in the sense of Hamilton [[Ham82](#), Part II Definition 1.3.2]—a prerequisite for using Nash–Moser theory. Moreover, spinors $\phi \in H_a^\infty\Gamma(X \setminus Z, S \otimes I)$ extend smoothly across Z after untwisting by $\bar{z}^{1/2}$ and, therefore, have well-behaved polyhomogeneous expansions near Z . Crucially, it is proved that if a residue condition $B \subset \check{H}\Gamma(Z, \check{S})$ is ∞ -**regular**, then the extension D_B satisfies a variant of elliptic regularity together with elliptic estimates. For local residue conditions B , ∞ -regularity can be verified using straight-forward symbolic criterion. In particular, this criterion applies to the Lagrangian local residue condition which is secretly at the heart of [[Tak15](#); [Par23](#)]. It is quite plausible that these results can be cobbled together using the powerful machines developed by Mazzeo [[Maz91](#)], Mazzeo and Vertman [[MV14](#)], and Albin and Gell-Redman [[AG16](#); [AG23](#)]. However, the arguments in [Section 4](#) are almost elementary and there should be some value in that.

It should be possible, with suitable modifications, to extend the work in the present article to higher rank ramified Euclidean local systems; in particular: to flat Hermitian line bundles. In fact, Ammann–Große have on-going work in progress in this direction.

Acknowledgements. The authors thank Siqi He and Greg Parker for helpful discussions and Jacek Rzemieniecki for his careful proofreading an earlier version of this article. TW is indebted

to Dietmar Salamon for teaching him about the Gelfand–Robbin quotient almost two decades ago.

This material is based in part upon work carried while the authors were in residence at the Simons Laufer Mathematical Sciences Institute (previously known as MSRI) Berkeley, California, during the Fall 2022 semester.

Conventions. Choose a cut-off function $\chi \in C^\infty([0, \infty), [0, 1])$ with $\chi|_{[0, 1/4]} = 1$ and $\text{supp}(\chi) \subset [0, 1/2)$. The bracket $\langle - \rangle : \mathbf{R} \rightarrow [1, \infty)$ is defined by $\langle x \rangle := (1 + x^2)^{1/2}$.

2 The Gelfand–Robbin quotient, I: abstract theory

This section studies the closed extensions of the Dirac operator $D : H^1\Gamma(X \setminus Z, S \otimes \mathbf{I}) \rightarrow L^2\Gamma(X \setminus Z, S \otimes \mathbf{I})$, considered as an unbounded operator, following [MS98, Exercise 2.17; BF98, §3; SWo8, Appendix B; BS18, Exercises 6.3.3 and 6.5.11]; Throughout, assume the following analytic condition on the branching locus Z .

Hypothesis 2.1. *There is an $r \in C^\infty(X \setminus Z, (0, \infty))$, uniformly comparable to the Riemannian distance to Z , such that following **borderline Hardy inequality** holds: for every $\phi \in H^1\Gamma(X \setminus Z, S \otimes \mathbf{I})$, $r^{-1}\phi \in L^2\Gamma(X \setminus Z, S \otimes \mathbf{I})$ and*

$$\|r^{-1}\phi\|_{L^2} \lesssim \|\phi\|_{H^1}.$$

Remark 2.2. Hypothesis 2.1 holds if Z is a codimension two submanifold; see Takahashi [Tak15, Lemma 2.6] or Lemma 3.2. Moreover, it holds in the situation considered by Haydys, Mazzeo, and Takahashi [HMT23] where $Z \subset X$ is a graph embedded in a 3–manifold. ♣

2.1 The minimal and maximal extension

Proposition 2.3. *The bounded operator $D : H^1\Gamma(X \setminus Z, S \otimes \mathbf{I}) \rightarrow L^2\Gamma(X \setminus Z, S \otimes \mathbf{I})$ is left semi-Fredholm; that is: $\ker D$ is finite-dimensional and $\text{im } D$ is closed.*

The proof relies on the following consequences of the borderline Hardy inequality.

Lemma 2.4. *The following hold:*

- (1) $H^1\Gamma(X \setminus Z, S \otimes \mathbf{I}) = H_0^1\Gamma(X \setminus Z, S \otimes \mathbf{I})$.
- (2) *The inclusion $H^1\Gamma(X \setminus Z, S \otimes \mathbf{I}) \hookrightarrow L^2\Gamma(X \setminus Z, S \otimes \mathbf{I})$ is a compact operator.*

Proof. For $\varepsilon > 0$ set $\chi_\varepsilon := \chi(r/\varepsilon)$. Let $\phi \in H^1\Gamma(X \setminus Z, S \otimes \mathbf{I})$. Since $|\text{rd}\chi_\varepsilon| \lesssim 1$,

$$\|\nabla(\chi_\varepsilon\phi)\|_{L^2} \leq \|(\text{rd}\chi_\varepsilon)r^{-1}\phi\|_{L^2} + \|\chi_\varepsilon\nabla\phi\|_{L^2} \lesssim \left(\int_{\text{supp}(\text{d}\chi_\varepsilon)} |r^{-1}\phi|^2 + |\nabla\phi|^2 \right)^{1/2}.$$

Therefore, by Hypothesis 2.1 and monotone convergence,

$$\lim_{\varepsilon \downarrow 0} \|\nabla(\chi_\varepsilon\phi)\|_{L^2} = 0.$$

This implies (1).

Let $(\phi_n) \in H^1\Gamma(X \setminus Z, S \otimes \mathbb{I})^{\mathbb{N}}$ with $\|\phi_n\|_{H^1} = 1$ for every $n \in \mathbb{N}$. For every $\varepsilon > 0$, a subsequence of $((1 - \chi_\varepsilon)\phi_n)$ converges in $L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})$. By the borderline Hardy inequality,

$$\|\chi_\varepsilon\phi_n\|_{L^2} \lesssim \varepsilon.$$

Therefore, the (2) follows from a diagonal sequence argument. \blacksquare

Proof of Proposition 2.3. By Lemma 2.4 (1), the Schrödinger–Lichnerowicz formula

$$D^2 = \nabla^* \nabla + \gamma(F_\nabla)$$

implies the elliptic estimate

$$(2.5) \quad \|\phi\|_{H^1} \lesssim \|D\phi\|_{L^2} + \|\phi\|_{L^2}$$

for every $\phi \in H^1\Gamma(X \setminus Z, S \otimes \mathbb{I})$. Therefore and by Lemma 2.4 (2), D is left semi-Fredholm. \blacksquare

With the exception of a few edge cases—e.g.: if $Z = \emptyset$ or $Z \subset X$ is a finite subset of a surface [DW24, §3.4.2; HMT23, §4.1–4.5]—the operator $D: H^1\Gamma(X \setminus Z, S \otimes \mathbb{I}) \rightarrow L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})$ is not Fredholm: its cokernel is ∞ -dimensional. Therefore, it is useful to consider D as an unbounded operator and systematically study its closed extensions; cf. [BS18, Chapter 6].

Definition 2.6. The **minimal extension**

$$D_{\min}: \text{dom}(D_{\min}) \rightarrow L^2\Gamma(X \setminus Z, S \otimes \mathbb{I});$$

is the operator $D: H^1\Gamma(X \setminus Z, S \otimes \mathbb{I}) \rightarrow L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})$ considered as unbounded operator on $L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})$. \bullet

Proposition 2.7. D_{\min} is closed, densely defined, and symmetric.

Proof. Evidently, D_{\min} is densely defined. Since D is formally self-adjoint,

$$\langle D\phi, \psi \rangle = \langle \phi, D\psi \rangle$$

for every $\phi, \psi \in \Gamma_c(X \setminus Z, S \otimes \mathbb{I})$. Therefore, by Lemma 2.4 (1) and since $\Gamma_c(X \setminus Z, S \otimes \mathbb{I}) \subset H_0^1\Gamma(X \setminus Z, S \otimes \mathbb{I})$ is dense, D_{\min} is symmetric. By (2.5) the Sobolev norm $\|\cdot\|_{H^1}$ and the graph norm $\|\cdot\|_D := (\|\cdot\|_{L^2}^2 + \|D\cdot\|_{L^2}^2)^{1/2}$ are equivalent. Therefore and since $H^1\Gamma(X \setminus Z, S \otimes \mathbb{I})$ is complete, D_{\min} is closed. \blacksquare

Definition 2.8. The **maximal extension**

$$D_{\max}: \text{dom}(D_{\max}) \rightarrow L^2(X \setminus Z, S \otimes \mathbb{I})$$

is the adjoint of D_{\min} in the sense of unbounded operators; that is:

$$\text{dom}(D_{\max}) := \{\phi \in L^2\Gamma(X \setminus Z, S \otimes \mathbb{I}) : \langle \phi, D_{\min} \cdot \rangle_{L^2} : \text{dom}(D_{\min}) \rightarrow \mathbf{R} \text{ is } \|\cdot\|_D\text{-bounded}\}$$

and for every $\phi \in \text{dom}(D_{\max})$ and $\psi \in \text{dom}(D_{\min})$

$$\langle D_{\max}\phi, \psi \rangle = \langle \phi, D_{\min}\psi \rangle.$$

$D_{\max}\phi$ exists by the Hahn–Banach Theorem and the Riesz Representation Theorem, and is unique because $\text{dom}(D_{\min})$ is dense. \bullet

Remark 2.9. It is convenient to consider $D : H_{\text{loc}}^1 \Gamma(X \setminus Z, S \otimes \mathbf{I}) \rightarrow L_{\text{loc}}^2 \Gamma(X \setminus Z, S \otimes \mathbf{I})$. From this perspective,

$$\text{dom}(D_{\text{max}}) = \{\phi \in H_{\text{loc}}^1 \Gamma(X \setminus Z, S \otimes \mathbf{I}) : \phi, D\phi \in L^2 \Gamma(X \setminus Z, S \otimes \mathbf{I})\};$$

and, it is excusable to drop the subscripts from $D_{\text{min}}\phi$, $D_{\text{max}}\phi$, etc. ♣

2.2 Closed extensions and residue conditions

The closed extensions of D_{min} can be systematically understood as follows.

Definition 2.10. The Gelfand–Robbin quotient is the Hilbert space

$$\check{\mathbf{H}} := \frac{\text{dom}(D_{\text{max}})}{\text{dom}(D_{\text{min}})}.$$

Since D_{min} is closed, $\text{dom}(D_{\text{min}}) \subset \text{dom}(D_{\text{max}})$ is a $\|\cdot\|_D$ -closed subspace. Denote the canonical projection map by

$$[\cdot] : \text{dom}(D_{\text{max}}) \rightarrow \check{\mathbf{H}}. \quad \bullet$$

Remark 2.11. $\check{\mathbf{H}}$ is localised on Z in the following sense: $[\phi] = [\chi(r/\varepsilon)\phi]$ for every $\varepsilon > 0$ and $\phi \in \text{dom}(D_{\text{max}})$. ♣

Definition 2.12. A residue condition is a closed subspace $B \subset \check{\mathbf{H}}$. •

Proposition 2.13 (closed extension=residue condition; cf. [BF98, Lemma 3.3(a)]). *If $B \subset \check{\mathbf{H}}$ is a residue condition, then*

$$D_B := D_{\text{max}}|_{\text{dom}(D_B)} \quad \text{with} \quad \text{dom}(D_B) := [\cdot]^{-1}(B)$$

is a closed extension of D_{min} . Moreover, every closed extension of D_{min} is of this form.

Proof. Let $B \subset \check{\mathbf{H}}$ be a residue condition. The canonical projection $[\cdot] : \text{dom}(D_{\text{max}}) \rightarrow \check{\mathbf{H}}$ is bounded. Therefore, $\text{dom}(D_B) := [\cdot]^{-1}(B) \subset \text{dom}(D_{\text{max}})$ is a $\|\cdot\|_D$ -closed subspace; hence: D_B is closed.

Let \bar{D} be a closed extension of D_{min} . Since $\text{dom}(\bar{D}) \subset \text{dom}(D_{\text{max}})$ is a $\|\cdot\|_D$ -closed subspace, $B := [\text{dom}(\bar{D})] = \frac{\text{dom}(\bar{D})}{\text{dom}(D_{\text{min}})} \subset \check{\mathbf{H}}$ is a closed subspace. Since $\text{dom}(D_{\text{min}}) \subset \text{dom}(\bar{D})$, $\text{dom}(\bar{D}) = [\cdot]^{-1}(B)$; hence: $\bar{D} = D_B$. ■

2.3 The Green's form and adjoint extensions

The Gelfand–Robbin quotient carries a symplectic structure related to the construction of adjoint extensions.

Definition 2.14. The Green's form $G \in \text{Hom}(\Lambda^2 \check{\mathbf{H}}, \mathbf{R})$ is defined by

$$G([\phi] \wedge [\psi]) := \langle D\phi, \psi \rangle_{L^2} - \langle \phi, D\psi \rangle_{L^2}. \quad \bullet$$

Proposition 2.15 (cf. [BF98, Lemma 3.1, Proposition 3.2; SWo8, Remark B.1(ii)]). G is symplectic; that is: it induces a Hilbert space isomorphism

$$J: \check{\mathbf{H}} \rightarrow \hat{\mathbf{H}} := \mathcal{L}(\check{\mathbf{H}}, \mathbf{R})$$

$$[\phi] \mapsto G([\phi] \wedge -).$$

Moreover: if $\sharp: \hat{\mathbf{H}} \rightarrow \check{\mathbf{H}}$ denotes the isomorphism induced by the inner product, then $\sharp \circ J$ is an isometric complex structure.

Proof. The canonical projection induces an isometry $[-]: \text{dom}(D_{\min})^{\perp D} \cong \check{\mathbf{H}}$. Here \perp_D indicates the orthogonal complement with respect to the graph inner product $\langle -, - \rangle_D := \langle -, - \rangle_{L^2} + \langle D-, D- \rangle_{L^2}$. By direct inspection,

$$\begin{aligned} \text{dom}(D_{\min})^{\perp D} &= \{\phi \in \text{dom}(D_{\max}) : \langle \phi, \psi \rangle_{L^2} + \langle D\phi, D\psi \rangle_{L^2} = 0 \text{ for every } \psi \in \text{dom}(D_{\min})\} \\ &= \{\phi \in \text{dom}(D_{\max}) : D\phi \in \text{dom}(D_{\max}) \text{ and } D^2\phi = -\phi\}. \end{aligned}$$

Therefore, D induces an isometric complex structure $D: \text{dom}(D_{\min})^{\perp D} \rightarrow \text{dom}(D_{\min})^{\perp D}$.

The diagram

$$\begin{array}{ccc} \text{dom}(D_{\min})^{\perp D} & \xrightarrow{D} & \text{dom}(D_{\min})^{\perp D} \\ [-] \downarrow & & \downarrow [-] \\ \check{\mathbf{H}} & \xrightarrow{\sharp \circ J} & \check{\mathbf{H}} \end{array}$$

commutes because for every $\phi, \psi \in \text{dom}(D_{\min})^{\perp D}$

$$G([\phi] \wedge [\psi]) = \langle D\phi, \psi \rangle_{L^2} - \langle \phi, D\psi \rangle_{L^2} = \langle D\phi, \psi \rangle_{L^2} + \langle D^2\phi, D\psi \rangle_{L^2} = \langle D\phi, \psi \rangle_D.$$

This proves the assertion. ■

Proposition 2.16 (cf. [BF98, Lemma 3.3(b)]). Let $B \subset \check{\mathbf{H}}$ be a residue condition. The adjoint D_B^* of D_B is the closed extension D_{B^G} associated with the symplectic complement

$$B^G := \{[\phi] \in \check{\mathbf{H}} : G([\phi], [\psi]) = 0 \text{ for every } [\psi] \in B\}.$$

In particular, D_B is self-adjoint if and only if B is Lagrangian.

Proof. A moment's thought shows that

$$\begin{aligned} \text{dom}(D_B^*) &= \{\phi \in \text{dom}(D_{\max}) : \langle D\phi, \psi \rangle_{L^2} = \langle \phi, D\psi \rangle_{L^2} \text{ for every } \psi \in \text{dom}(D_B)\} \\ &= \{\phi \in \text{dom}(D_{\max}) : G([\phi] \wedge [\psi]) = 0 \text{ for every } [\psi] \in B\}. \end{aligned}$$

This proves the assertion. ■

Example 2.17. The Calderón subspace

$$\Lambda := [\ker D_{\max}] \subset \check{\mathbf{H}}$$

is a Lagrangian residue condition. Indeed, $\Lambda \subset \Lambda^G$ because $G([\phi] \wedge [\psi]) = \langle D\phi, \psi \rangle_{L^2} - \langle \phi, D\psi \rangle_{L^2} = 0$ for every $\phi, \psi \in \ker D_{\max}$. Moreover, if $[\phi] \in \Lambda^G$, then $D\phi \perp_{L^2} \ker D_{\max} = (\text{im } D_{\min})^{\perp L^2}$; therefore, there is a $\psi \in \text{dom}(D_{\min})$ with $D\psi = D\phi$; hence: $[\phi] = [\phi - \psi] \in \Lambda$. ♠

Example 2.18. As a consequence of [Proposition 2.15](#), the orthogonal complement of the Calderón subspace

$$\Lambda^\perp = \sharp \circ J(\Lambda) \subset \check{\mathbf{H}}$$

is a Lagrangian residue condition. ♣

Example 2.19. Suppose that S carries a parallel orthogonal complex structure i which commutes with γ ; that is: (S, γ, ∇) is a **complex Dirac bundle**. Evidently, D is complex linear and $\check{\mathbf{H}}$ inherits i as an isometric complex structure i . This induces an orthogonal decomposition

$$\check{\mathbf{H}} = B_+ \oplus B_- \quad \text{with} \quad B_\pm := \{[\phi] \in \check{\mathbf{H}} : \sharp \circ J[\phi] = \pm i[\phi]\}.$$

Since i and $\sharp \circ J$ commute, $B_\pm \subset \check{\mathbf{H}}$ are complex subspaces and, therefore, B_\pm are mutually adjoint:

$$B_\pm^G = B_\mp. \quad \spadesuit$$

Proposition 2.20 (Spectral theory). *Let $B \subset \check{\mathbf{H}}$ be a residue condition. If $B \subset \check{\mathbf{H}}$ is a Lagrangian and $\text{dom}(D_B) \hookrightarrow L^2\Gamma(X \setminus Z, S \otimes \mathbf{I})$ is compact, then $\text{spec}(D_B)$ consists only of point spectrum, is contained in \mathbf{R} and discrete, and for every $\lambda \in \text{spec}(D_B)$ the eigenspace $\ker(D_B - \lambda \cdot \mathbf{1})$ is finite-dimensional; moreover: $L^2\Gamma(X \setminus Z, S \otimes \mathbf{I})$ decomposes as a (Hilbert space) direct sum*

$$L^2\Gamma(X \setminus Z, S \otimes \mathbf{I}) = \bigoplus_{\lambda \in \text{spec}(D_B)} \ker(D_B - \lambda \cdot \mathbf{1}).$$

Proof. By assumption, D_B is self-adjoint and has compact resolvent. The assertion, therefore, follows from the spectral theory of such operators; see, e.g., [\[BS18, Theorem 6.3.13\]](#). ■

2.4 Fredholm extensions

The following characterises residue conditions $B \subset \check{\mathbf{H}}$ which correspond to Fredholm extensions D_B in terms of the relation between B and the Calderón subspace Λ .

Definition 2.21. Let $B \subset \check{\mathbf{H}}$ be a residue condition. Denote by

$$\delta_B: \Lambda \rightarrow \check{\mathbf{H}}/B \quad \text{and} \quad \delta^B: B \rightarrow \check{\mathbf{H}}/\Lambda$$

the compositions of the canonical inclusions and projections. •

Proposition 2.22 (cf. [\[SW08, Lemma B.3\]](#)). *Let $B \subset \check{\mathbf{H}}$ be a residue condition. The closed extension D_B is Fredholm if and only if δ_B is Fredholm if and only if δ^B is Fredholm; moreover:*

$$\text{index } D_B = \text{index } \delta_B = \text{index } \delta^B.$$

The proof relies on the following observation.

Lemma 2.23. *For every residue condition $B \subset \check{\mathbf{H}}$, there are short exact sequences*

$$\ker D_{\min} \hookrightarrow \ker D_B \rightarrow \ker \delta_B \quad \text{and} \quad \text{coker } \delta_B \hookrightarrow \text{coker } D_B \twoheadrightarrow \text{coker } D_{\max}.$$

Proof. The Snake Lemma applied to

$$\begin{array}{ccccc} \ker D_{\min} & \xlongequal{\quad} & \ker D_{\min} & & \\ \downarrow & & \downarrow & & \\ \ker D_B & \hookrightarrow & \ker D_{\max} & \xrightarrow{\delta_B \circ [-]} & \check{\mathbf{H}}/B \end{array}$$

yields an exact sequence

$$\frac{\ker D_B}{\ker D_{\min}} \hookrightarrow \frac{\ker D_{\max}}{\ker D_{\min}} \cong \Lambda \xrightarrow{\delta_B} \check{\mathbf{H}}/B.$$

This induces the first short exact sequence.

The Snake Lemma applied to

$$\begin{array}{ccccc} \text{dom}(D_B) & \hookrightarrow & \text{dom}(D_{\max}) & \longrightarrow & \check{\mathbf{H}}/B \\ \downarrow D_B & & \downarrow D_{\max} & & \\ L^2\Gamma(X \setminus Z, S \otimes \mathbb{I}) & \xlongequal{\quad} & L^2\Gamma(X \setminus Z, S \otimes \mathbb{I}) & & \end{array}$$

yields an exact sequence

$$\ker D_B \hookrightarrow \ker D_{\max} \xrightarrow{\delta_B \circ [-]} \check{\mathbf{H}}/B \rightarrow \text{coker } D_B \rightarrow \text{coker } D_{\max}.$$

Since $\text{coker } \delta_B \circ [-] = \text{coker } \delta_B$, this induces the second short exact sequence. \blacksquare

Proof of Proposition 2.22. A moment's thought shows that

$$\ker \delta_B = \Lambda \cap B = \ker \delta^B \quad \text{and} \quad \text{coker } \delta_B \cong \frac{\check{\mathbf{H}}}{\Lambda + B} \cong \text{coker } \delta^B.$$

By Proposition 2.3, $\ker D_{\min} \cong \text{coker } D_{\max}$ is finite-dimensional. Therefore, the assertion is an immediate consequence of Lemma 2.23. \blacksquare

Example 2.24. Every complement B of the Calderón subspace Λ , in particular: $B = \Lambda^\perp$, produces a Fredholm extension of index 0 because $\delta^B: B \rightarrow \check{\mathbf{H}}/\Lambda \cong B$ is an isomorphism. \spadesuit

Example 2.25. The residue conditions B_\pm defined in Example 2.19 satisfy

$$\ker D_{B_\pm} = \ker D_{\min}$$

and, therefore, produce Fredholm extension of index 0; indeed: if $\phi \in \ker D_{B_\pm}$, then

$$0 = 2\langle D\phi, i\phi \rangle_{L^2} = \langle D\phi, i\phi \rangle_{L^2} - \langle \phi, Di\phi \rangle_{L^2} = G([\phi], i[\phi]) = \langle \# \circ J[\phi], i[\phi] \rangle_{L^2} = \pm \|\phi\|_{L^2}^2. \quad \spadesuit$$

The following are occasionally useful to compute or relate indices.

Proposition 2.26 (Nested Fredholm residue conditions). *Let $B_1 \subset B_2 \subset \check{\mathbf{H}}$ be residue conditions. If $\delta_{B_1}, \delta_{B_2}$ are Fredholm, then*

$$\text{index } D_{B_2} = \text{index } D_{B_1} + \dim B_2/B_1.$$

Corollary 2.27. *Let $B \subset \check{H}$ be a residue condition. If $B \subset B^G$ and δ_B is Fredholm then $\text{index } D_B = -\frac{1}{2} \dim B^G/B$; in particular: if B is Lagrangian, then $\text{index } D_B = 0$. ■*

Proposition 2.26 is an immediate consequence of the following.

Lemma 2.28 (Nested residue conditions). *Let $B_1 \subset B_2 \subset \check{H}$ be residue conditions. There is an exact sequence*

$$\ker \delta_{B_1} \hookrightarrow \ker \delta_{B_2} \rightarrow B_2/B_1 \rightarrow \text{coker } \delta_{B_1} \rightarrow \text{coker } \delta_{B_2}.$$

Proof. The exact sequence follows from the Snake Lemma applied to

$$\begin{array}{ccc} \Lambda & \xlongequal{\quad} & \Lambda \\ \downarrow \delta_{B_1} & & \downarrow \delta_{B_2} \\ B_2/B_1 & \hookrightarrow & \check{H}/B_1 \twoheadrightarrow \check{H}/B_2. \end{array}$$

This implies the assertion by the above proposition. ■

Proposition 2.29 (Deformation of residue conditions). *Let B be a Hilbert space. Let $\iota_- : [0, 1] \rightarrow \mathcal{L}(B, \check{H})$ be a continuous path of embeddings. If D_{B_t} with $B_t := \iota_t(B)$ is Fredholm for every $t \in [0, 1]$, then*

$$\text{index } D_{B_0} = \text{index } D_{B_1}.$$

Proof. By assumption, $t \mapsto \delta^{B_t} \circ \iota_t$ is a continuous path of Fredholm operators and $\iota_t : B \rightarrow B_t$ is an isomorphism for every $t \in [0, 1]$; therefore: $t \mapsto \text{index } D_{B_t} = \text{index } \delta^{B_t} = \text{index } \delta^{B_t} \circ \iota_t$ is constant. ■

2.5 Chirality operators

In the presence of a chirality operator the theory discussed above refines as follows.

Definition 2.30. A chirality operator for (S, γ, ∇) is a self-adjoint parallel isometry $\varepsilon \in \Gamma(X, \text{O}(S))$ such that

$$\gamma \varepsilon + \varepsilon \gamma = 0. \quad \bullet$$

Example 2.31. If $\dim X = 0 \pmod{4}$, then $\varepsilon := \gamma(\text{vol}_g)$ is a chirality operator. If $\dim X = 0 \pmod{2}$ and (S, γ, ∇) is a complex Dirac bundle as in [Example 2.19](#), then $\varepsilon := i\gamma(\text{vol}_g)$ is a chirality operator. ♠

Proposition 2.32 (Chirality operators induce a $\{\pm 1\}$ -grading). *If ε is a chirality operator for (S, γ, ∇) , then the following hold:*

- (1) *The subbundles $S^\pm := \ker(1 \pm \varepsilon) \subset S$ are parallel, S orthogonally decomposes as*

$$S = S^+ \oplus S^-,$$

and $\gamma \in \Gamma(X, \text{Hom}(TX, \text{Hom}(S^+, S^-) \oplus \text{Hom}(S^-, S^+)))$.

(2) The minimal and maximal extensions decompose as

$$D_{\min} = \begin{pmatrix} 0 & D_{\min}^- \\ D_{\min}^+ & 0 \end{pmatrix} \quad \text{and} \quad D_{\max} = \begin{pmatrix} 0 & D_{\max}^- \\ D_{\max}^+ & 0 \end{pmatrix}$$

with

$$D_{\min}^\pm : \text{dom}(D_{\min}^\pm) := H^1\Gamma(X \setminus Z, S^\pm \otimes \mathbb{I}) \rightarrow L^2\Gamma(X \setminus Z, S^\mp \otimes \mathbb{I}) \quad \text{and} \\ D_{\max}^\pm : \text{dom}(D_{\max}^\pm) := \text{dom}(D_{\max}) \cap L^2\Gamma(X \setminus Z, S^\pm \otimes \mathbb{I}) \rightarrow L^2\Gamma(X \setminus Z, S^\mp \otimes \mathbb{I}).$$

(3) \check{H} orthogonally decomposes as

$$\check{H} = \check{H}^+ \oplus \check{H}^- \quad \text{with} \quad \check{H}^\pm := \frac{\text{dom}(D_{\max}^\pm)}{\text{dom}(D_{\min}^\pm)};$$

moreover, $\check{H}^\pm \subset \check{H}$ are Lagrangian. In particular, every residue condition $B \subset \check{H}$ decomposes as $B = B^+ \oplus B^-$.

(4) If $B^+ \subset \check{H}^+$ is a closed subspace, a **positive chirality residue condition**, then there is a unique closed subspace $B^- \subset \check{H}^-$ such that $B := B^+ \oplus B^- \subset \check{H}$ is a Lagrangian residue condition.

(5) Let $B^+ \subset \check{H}^+$ be a **positive chirality residue condition** and B^- as above. The operator δ^B is Fredholm if and only if $\delta^{B^+} : B^+ \rightarrow \check{H}^+/\Lambda^+$ is Fredholm.

Proof. (1), (2), and (3) are an immediate consequence of [Definition 2.30](#).

A moment's thought shows that (4) holds with $B^- := (B^+)^G \cap \check{H}^-$.

Evidently, $\delta^B = \delta^{B^+} \oplus \delta^{B^-}$ is Fredholm if and only if δ^{B^+} and δ^{B^-} are Fredholm. The Green's form G induces isomorphisms

$$\Lambda^- \cong (\check{H}^+/\Lambda^+)^* \quad \text{and} \quad B^+ \cong (\check{H}^-/B^-)^*.$$

This identifies the dual of δ^{B^+} with $\delta_{B^-} : \Lambda^- \rightarrow \check{H}^-/B^-$. By the closed image theorem, if δ^{B^+} is Fredholm, then δ_{B^-} is Fredholm. As in the proof of [Proposition 2.22](#), δ_{B^-} is Fredholm if and only if δ^{B^-} is Fredholm. This proves (5). \blacksquare

3 The Gelfand–Robbin quotient, II: geometric realisation

The usefulness of the theory laid out in [Section 2](#) hinges upon being able to understand \check{H} , e.g., to specify interesting residue conditions. Since \check{H} localises on Z as explained in [Remark 2.11](#), it is plausible that it admits a more geometric description. The purpose of this section is to develop such a description, assuming the following geometric condition on the branching locus Z throughout.

Hypothesis 3.1. $Z \subset X$ is a closed cooriented submanifold of codimension two.

Lemma 3.2 (borderline Hardy inequality; Takahashi [Tak15, Lemma 2.6]). *Hypothesis 3.1 implies Hypothesis 2.1.*

Proof. Let $r > 0$. Denote by \mathbb{I} the non-trivial Euclidean line bundle over $rS^1 := \{z \in \mathbb{C} : |z| = r\}$. A moment's thought and a scaling consideration show that

$$\int_{rS^1} |r^{-1}s|^2 \lesssim \int_{rS^1} |\nabla s|^2$$

for every $s \in \Gamma(rS^1, \mathbb{I})$. This immediately implies the assertion. \blacksquare

3.1 The blow-up of X along Z

It is convenient to blow-up X along Z ; that is: to replace $Z \subset X$ with the following $U(1)$ -principal bundle.

Definition 3.3. Since Z is cooriented, $NZ \rightarrow Z$ is a Hermitian line bundle. Its **frame bundle** is

$$\pi: F := \{v \in NZ : |v| = 1\} \rightarrow Z$$

together with $F \cup U(1)$ defined by $v \cdot e^{i\alpha} := e^{i\alpha}v$. Denote the Levi-Civita connection on F by $i\theta \in \Omega^1(F, i\mathbb{R})$. \bullet

Remark 3.4. The **tautological section** $\partial_r \in \Gamma(F, \pi^*NZ)$ and $\partial_\alpha := i\partial_r$ trivialise π^*NZ . \clubsuit

In order to replace $Z \subset X$ with F a choice is required.

Definition 3.5. Set $U := [0, 1) \cdot F \subset NZ$. A **tubular neighbourhood** $j: U \hookrightarrow X$ of $Z \subset X$ is an embedding such that $j \circ 0 = \text{id}_Z$ and the composition

$$NZ \hookrightarrow 0^*TNZ \xrightarrow{Tj} TX|_Z \rightarrow NZ$$

is the identity. Here $0: Z \rightarrow NZ$ denotes the zero section. \bullet

Choose a tubular neighbourhood $j: U \hookrightarrow X$.

Definition 3.6. Set $\hat{U} := [0, 1) \times F$. The **blow-up** of X along Z is the manifold with boundary

$$\hat{X} := \hat{U} \cup_j (X \setminus Z)$$

obtained by gluing \hat{U} and $X \setminus Z$ along j . The **blow-down map** $\beta: \hat{X} \rightarrow X$ is defined by $\beta(r, v) := j(rv)$ for $(r, v) \in \hat{U}$ and $\beta(x) := x$ for $x \in X \setminus Z$. \bullet

Henceforth, identify $U \subset NZ$ and $j(U) \subset X$; moreover, identify $\partial\hat{X} = F$.

Definition 3.7. Set

$$\hat{S} := \beta^*S \quad \text{and} \quad \underline{S} := \hat{S}|_F = \pi^*(S|_Z).$$

Endow \underline{S} with the quaternionic structure

$$I := \gamma(\text{vol}_{NZ}), \quad J := \gamma(\partial_r), \quad \text{and} \quad K := \gamma(\partial_\alpha) \in \Gamma(F, \text{End}(\underline{S})).$$

Since $X \setminus Z \hookrightarrow \hat{X}$ is a homotopy-equivalence, \mathbb{I} extends uniquely to a Euclidean line bundle

$$\hat{\mathbb{I}} \rightarrow \hat{X}.$$

Set

$$\underline{\mathbb{I}} := \hat{\mathbb{I}}|_F. \quad \bullet$$

3.2 The model operator

The purpose of this subsection is to construct a model \mathring{D} for D near Z . This construction relies on the following.

Definition 3.8 (Restriction of Dirac bundles). Denote the second fundamental form of Z by $\Pi \in \Gamma(Z, \text{Hom}(TZ, \text{Hom}(TZ, NZ)))$. The **restriction** of (S, γ, ∇) to Z is the triple $(S|_Z, \gamma|_{TZ}, \nabla|_Z + \frac{1}{2}\gamma(\Pi))$ with

$$\gamma(\Pi)(v) := \frac{1}{2} \sum_{i=1}^{n-2} \gamma(\Pi(v)e_i)\gamma(e_i).$$

Here (e_1, \dots, e_{n-2}) denotes a local orthonormal frame of TZ . •

Proposition 3.9. $(S|_Z, \gamma|_{TZ}, \nabla|_Z + \frac{1}{2}\gamma(\Pi))$ is a Dirac bundle over $(Z, g|_Z)$.

Proof. Evidently, $(S|_Z, \gamma|_{TZ})$ forms a Clifford module bundle over $(Z, g|_Z)$. Denote by ∇^\parallel the connection on TZ induced by the Levi-Civita connection. Since (S, γ, ∇) is a Dirac bundle over (X, g) , for every $v, w \in \text{Vect}(Z)$

$$[\nabla_v, \gamma(w)] = \gamma(\nabla_v^\parallel w) + \gamma(\Pi(v, w));$$

moreover, by direct computation,

$$\begin{aligned} [\gamma(\Pi)(v), \gamma(w)] &= \sum_{i=1}^{n-2} [\gamma(\Pi(v, e_i))\gamma(e_i), \gamma(w)] \\ &= \sum_{i=1}^{n-2} \gamma(\Pi(v, e_i))(\gamma(e_i)\gamma(w) + \gamma(w)\gamma(e_i)) = -2\gamma(\Pi(v, w)). \end{aligned} \quad \blacksquare$$

Proposition 3.10. Denote by $\Pi: U \rightarrow Z$ the projection map and identify $TU = \Pi^*(TZ \oplus NZ)$ using the Levi-Civita connection. Denote by g^\parallel and g^\perp the Euclidean metrics on TZ and NZ induced by g . Consider $U \subset X$ equipped with the Riemannian metric

$$\mathring{g} := \Pi^*(g^\parallel \oplus g^\perp).$$

The following hold:

- (1) Denote by ∇^\parallel and ∇^\perp the covariant derivatives on TZ and NZ induced by the Levi-Civita connection. The affine connection

$$\mathring{\nabla}^T := \Pi^*(\nabla^\parallel \oplus \nabla^\perp).$$

is \mathring{g} -orthogonal, and its torsion Tor vanishes along Z and satisfies $\text{Tor} = \text{pr}_F^*(d\theta) \otimes \partial_\alpha \in \Omega^2(U, TU)$.

- (2) The triple $(\mathring{S}, \mathring{\gamma}, \mathring{\nabla})$ consisting of

$$\mathring{S} := \Pi^*(S|_Z), \quad \mathring{\gamma} := \Pi^*(\gamma|_Z), \quad \text{and} \quad \mathring{\nabla} := \Pi^*(\nabla|_Z + \frac{1}{2}\gamma(\Pi)).$$

is a Dirac bundle (with torsion) over (U, \mathring{g}) ; that is:

- (a) $(\mathring{S}, \mathring{\gamma})$ is a Clifford module bundle over (U, \mathring{g}) and
- (b) $\mathring{\gamma}$ is parallel with respect to $\mathring{\nabla}^T$ and $\mathring{\nabla}$.

Proof. Since ∇^{\parallel} is g^{\parallel} -orthogonal and ∇^{\perp} is g^{\perp} -orthogonal, $\mathring{\nabla}^T$ is \mathring{g} -orthogonal.

Denote by $\tilde{\cdot}: \text{Vect}(Z) \rightarrow \text{Vect}(U \setminus Z)$ and $\tilde{\cdot}: \Gamma(Z, NZ) \rightarrow \text{Vect}(U \setminus Z)$ the lifting maps. For $u, v \in \text{Vect}(Z)$ and $n, m \in \Gamma(Z, NZ)$, by direct computation,

$$\mathring{\nabla}_{\tilde{u}}^T \tilde{v} - \mathring{\nabla}_{\tilde{v}}^T \tilde{u} - [\tilde{u}, \tilde{v}] = \widetilde{[u, v]} - [\tilde{u}, \tilde{v}] = (\text{pr}_F^* d\theta)(\tilde{u}, \tilde{v}) \otimes \partial_{\alpha};$$

moreover,

$$\mathring{\nabla}_{\tilde{n}}^T \tilde{m} - \mathring{\nabla}_{\tilde{m}}^T \tilde{n} - [\tilde{n}, \tilde{m}] = 0 \quad \text{and} \quad \mathring{\nabla}_{\tilde{v}}^T \tilde{n} - \mathring{\nabla}_{\tilde{n}}^T \tilde{v} - [\tilde{v}, \tilde{n}] = 0.$$

This proves (1).

(2.a) is obvious. (2.b) follows from the computation in the proof of Proposition 3.9. \blacksquare

Definition 3.11. Denote by $\mathring{\mathbb{I}}$ the pullback of \mathbb{I} along the projection $U \setminus Z \cong F \times (0, 1) \rightarrow F$. The model Dirac operator

$$\mathring{D}: H_{\text{loc}}^1 \Gamma(U \setminus Z, \mathring{S} \otimes \mathring{\mathbb{I}}) \rightarrow L_{\text{loc}}^2 \Gamma(U \setminus Z, \mathring{S} \otimes \mathring{\mathbb{I}})$$

is the Dirac operator (with torsion) associated with $(\mathring{S}, \mathring{\nabla}, \mathring{\gamma})$ twisted by $\mathring{\mathbb{I}}$. \bullet

Remark 3.12. More explicitly, the model Dirac operator \mathring{D} is of the form

$$\mathring{D} = J(\partial_r - r^{-1} I \mathring{\nabla}_{\partial_{\alpha}}) + D_Z \quad \text{with} \quad D_Z := \sum_{i=1}^{n-2} \mathring{\gamma}(\tilde{e}_i) \mathring{\nabla}_{\tilde{e}_i}.$$

with $(\tilde{e}_1, \dots, \tilde{e}_{n-2})$ denoting the horizontal lift of a local $g|_Z$ -orthonormal frame. \clubsuit

Proposition 3.13 (cf. Bismut [Bis89, Theorem 1.10]). *The model Dirac operator \mathring{D} has the following properties*

- (1) \mathring{D} is formally self-adjoint; in fact: for every $\phi, \psi \in H_{\text{loc}}^1 \Gamma(U \setminus Z, \mathring{S} \otimes \mathring{\mathbb{I}})$

$$\langle \mathring{D}\phi, \psi \rangle - \langle \phi, \mathring{D}\psi \rangle = \text{div}_{\mathring{g}}(v) \quad \text{with} \quad v := \sum_{i=1}^n \langle \mathring{\gamma}(\mathring{e}_i) \phi, \psi \rangle \cdot \mathring{e}_i.$$

Here $(\mathring{e}_1, \dots, \mathring{e}_n)$ denotes a local \mathring{g} -orthonormal frame.

- (2) \mathring{D} satisfies the Schrödinger–Lichnerowicz formula

$$\mathring{D}^2 = \mathring{\nabla}^* \mathring{\nabla} + \tau \mathring{\nabla} + \gamma(F_{\mathring{\nabla}})$$

with $\tau \in \Gamma(U, \text{Hom}(T^*U \otimes \mathring{S}, \mathring{S}))$ depending linearly on Tor .

Proof. The following argument can be found in [Bis89, Proof of Theorem 1.10] and is repeated here only for the readers' convenience.

The Levi-Civita connection $\overset{\circ}{\nabla}^{\text{LC}}$ of $(U, \overset{\circ}{g})$ and $\overset{\circ}{\nabla}^T$ are related by Koszul's formula

$$\langle \overset{\circ}{\nabla}_u^T v, w \rangle = \langle \overset{\circ}{\nabla}_u^{\text{LC}} v, w \rangle + \frac{1}{2} (\langle \text{Tor}(u, v), w \rangle - \langle \text{Tor}(v, w), u \rangle + \langle \text{Tor}(w, u), v \rangle).$$

By direct computation,

$$\text{div}_{\overset{\circ}{g}} v = \sum_{i=1}^n \mathcal{L}_{\overset{\circ}{e}_i} \overset{\circ}{g}(v, \overset{\circ}{e}_i) = \langle \overset{\circ}{D} \phi, \psi \rangle - \langle \phi, \overset{\circ}{D} \psi \rangle + \sum_{i=1}^n \langle \overset{\circ}{\gamma}(\overset{\circ}{\nabla}_{\overset{\circ}{e}_i}^T \overset{\circ}{e}_i) \phi, \psi \rangle.$$

By Koszul's formula,

$$\sum_{i=1}^n \langle \overset{\circ}{\nabla}_{\overset{\circ}{e}_i}^T \overset{\circ}{e}_i, \overset{\circ}{e}_j \rangle = \sum_{i=1}^n \langle \text{Tor}(\overset{\circ}{e}_j, \overset{\circ}{e}_i), \overset{\circ}{e}_i \rangle = 0.$$

This proves (1).

By direct computation,

$$\begin{aligned} \overset{\circ}{D}^2 &= \sum_{i,j=1}^n \overset{\circ}{\gamma}(\overset{\circ}{e}_i) \overset{\circ}{\nabla}_{\overset{\circ}{e}_i} \overset{\circ}{\gamma}(\overset{\circ}{e}_j) \overset{\circ}{\nabla}_{\overset{\circ}{e}_j} = \sum_{i,j=1}^n \overset{\circ}{\gamma}(\overset{\circ}{e}_j) \overset{\circ}{\gamma}(\overset{\circ}{e}_i) \overset{\circ}{\nabla}_{\overset{\circ}{e}_i} \overset{\circ}{\nabla}_{\overset{\circ}{e}_j} + \overset{\circ}{\gamma}(\overset{\circ}{e}_i) \overset{\circ}{\gamma}(\overset{\circ}{\nabla}_{\overset{\circ}{e}_i}^T \overset{\circ}{e}_j) \overset{\circ}{\nabla}_{\overset{\circ}{e}_j} \\ &= \overset{\circ}{\nabla}^* \overset{\circ}{\nabla} + \overset{\circ}{\gamma} F_{\overset{\circ}{\nabla}} + \overset{\circ}{\gamma}(\overset{\circ}{e}_i) \overset{\circ}{\gamma}(\overset{\circ}{\nabla}_{\overset{\circ}{e}_i}^T \overset{\circ}{e}_j) \overset{\circ}{\nabla}_{\overset{\circ}{e}_j}. \end{aligned}$$

Since

$$\langle \overset{\circ}{\nabla}_{\overset{\circ}{e}_i}^T \overset{\circ}{e}_j, \overset{\circ}{e}_k \rangle = \frac{1}{2} (\langle \text{Tor}(\overset{\circ}{e}_i, \overset{\circ}{e}_j), \overset{\circ}{e}_k \rangle - \langle \text{Tor}(\overset{\circ}{e}_j, \overset{\circ}{e}_k), \overset{\circ}{e}_i \rangle + \langle \text{Tor}(\overset{\circ}{e}_k, \overset{\circ}{e}_i), \overset{\circ}{e}_j \rangle),$$

(2) follows. ■

Choose an isometry $\overset{\circ}{S} \cong S|_U$ which agrees with $\text{id}_{S|_Z}$ over Z , and an isometry $\overset{\circ}{\mathbb{I}} \cong \mathbb{I}|_{U \setminus Z}$; moreover, *henceforth*, regard these as identifications.

Proposition 3.14. *The error term*

$$\text{Err} := D - \overset{\circ}{D}: H_{\text{loc}}^1 \Gamma(U \setminus Z, \overset{\circ}{S} \otimes \overset{\circ}{\mathbb{I}}) \rightarrow L_{\text{loc}}^2 \Gamma(U \setminus Z, \overset{\circ}{S} \otimes \overset{\circ}{\mathbb{I}})$$

is of the form

$$\text{Err} = a \overset{\circ}{\nabla} + b + \overset{\circ}{\gamma}(\tilde{H}_Z)$$

with $a \in \Gamma(U, \text{Hom}(T^*U \otimes \overset{\circ}{S}, \overset{\circ}{S}))$, $b \in \Gamma(U, \text{End}(\overset{\circ}{S}))$, and $\tilde{H}_Z \in \text{Vect}(U)$ denoting the horizontal lift of the mean curvature of Z . Moreover, a and b vanish along Z .

Proof. If $(\overset{\circ}{e}_1, \dots, \overset{\circ}{e}_{n-2})$ is a local orthonormal frame of Π^*TZ and $(\overset{\circ}{e}_{n-1}, \overset{\circ}{e}_n)$ is a local orthonormal frame of Π^*NZ , then

$$\begin{aligned} - \sum_{i=1}^n \overset{\circ}{\gamma}(\overset{\circ}{e}_i) \overset{\circ}{\gamma}(\Pi)(\overset{\circ}{e}_i) &= \frac{1}{2} \sum_{i,j=1}^{n-2} \gamma(\Pi(\overset{\circ}{e}_i) \overset{\circ}{e}_j) \overset{\circ}{\gamma}(\overset{\circ}{e}_i) \overset{\circ}{\gamma}(\overset{\circ}{e}_j) \\ &= \sum_{i,j=1}^{n-2} \gamma(\Pi(\overset{\circ}{e}_i) \overset{\circ}{e}_j) \overset{\circ}{g}(\overset{\circ}{e}_i, \overset{\circ}{e}_j) = \overset{\circ}{\gamma}(\tilde{H}_Z). \end{aligned}$$

Therefore, the assertion follows from the fact that

$$\mathring{g} - g, \quad \mathring{\nabla} - \nabla - \frac{1}{2}\mathring{\gamma}(\Pi), \quad \text{and} \quad \mathring{\gamma} - \gamma$$

vanish along Z . ■

3.3 The model Gelfand–Robbin quotient

By (the proof of) [Proposition 2.7](#), the model minimal extension

$$\mathring{D}_{\min} := \mathring{D}: \text{dom}(\mathring{D}_{\min}) := H_0^1\Gamma(U \setminus Z, \mathring{S} \otimes \mathring{I}) \rightarrow L^2\Gamma(U \setminus Z, \mathring{S} \otimes \mathring{I})$$

is closed, densely defined, and symmetric. A moment's thought show that the domain of the model maximal extension

$$\mathring{D}_{\max} := \mathring{D}_{\min}^*$$

is

$$\text{dom}(\mathring{D}_{\max}) := \{\phi \in H_{\text{loc}}^1\Gamma(U \setminus Z, \mathring{S} \otimes \mathring{I}) : \phi, \mathring{D}\phi \in L^2\Gamma(U \setminus Z, \mathring{S} \otimes \mathring{I})\}.$$

The construction from [Section 2.2](#) and [Section 2.3](#) yields the following.

Definition 3.15. The model Gelfand–Robbin quotient is the Hilbert space

$$\mathring{\mathbf{H}} := \frac{\text{dom}(\mathring{D}_{\max})}{\text{dom}(\mathring{D}_{\min})}$$

equipped with the model Green's form $\mathring{G} \in \text{Hom}(\Lambda^2 \mathring{\mathbf{H}}, \mathbf{R})$ defined by

$$\mathring{G}([\phi] \wedge [\psi]) := \langle \mathring{D}\phi, \psi \rangle_{L^2} - \langle \phi, \mathring{D}\psi \rangle_{L^2}. \quad \bullet$$

By (the proof of) [Proposition 2.15](#), $(\mathring{\mathbf{H}}, \mathring{G})$ is a symplectic Hilbert space. In the sense of [Remark 2.11](#), $\mathring{\mathbf{H}}$ has contributions from $\{0\} \times F \subset \hat{U}$ and $\{1\} \times F$. Only the former is relevant for the purposes of this section.

Proposition 3.16. *The subspace*

$$\mathring{\mathbf{H}}_0 := \frac{\chi(r) \cdot \text{dom}(\mathring{D}_{\max}) + \text{dom}(\mathring{D}_{\min})}{\text{dom}(\mathring{D}_{\min})} \subset \mathring{\mathbf{H}}$$

is closed and symplectic.

Proof. Define the operator $\pi \in \mathcal{L}(\mathring{\mathbf{H}})$ by $\pi([\phi]) := [\chi(r) \cdot \phi]$. Since $\chi(r)(1 - \chi(r)) \cdot \text{dom}(\mathring{D}_{\max}) \subset \text{dom}(\mathring{D}_{\min})$, $\pi^2 = \pi$; that is: π is a projection. Hence, $\mathring{\mathbf{H}}_0 = \text{im } \pi = \ker(1 - \pi)$ is closed.

Since $(1 - \chi(r) - \chi \circ (1 - r)) \cdot \text{dom}(\mathring{D}_{\max}) \subset \text{dom}(\mathring{D}_{\min})$, $(1 - \pi)[\phi] = [\chi \circ (1 - r) \cdot \phi]$. Therefore,

$$\mathring{G}([\phi] \wedge [\psi]) = \mathring{G}(\pi[\phi] \wedge \pi[\psi]) + \mathring{G}((1 - \pi)[\phi] \wedge (1 - \pi)[\psi]).$$

Hence, $\mathring{\mathbf{H}}_0$ is symplectic. ■

Proposition 3.17. *There is a unique isomorphism of symplectic Hilbert spaces*

$$\text{cut-off: } (\check{\mathbf{H}}, G) \cong (\check{\mathbf{H}}_0, \check{G})$$

satisfying $\text{cut-off}([\phi]) = [\chi(r) \cdot \phi]$ for every $\phi \in \text{dom}(D_{\max})$.

The proof requires the following preparation.

Lemma 3.18 ($\text{dom}(D_{\max})$ vs. $\text{dom}(\dot{D}_{\max})$). *The following hold:*

- (1) *If $\phi \in \text{dom}(\dot{D}_{\max})$, then $\chi(r) \cdot \phi \in \text{dom}(D_{\max})$ and $\|\chi(r) \cdot \phi\|_D \lesssim \|\phi\|_{\dot{D}}$.*
- (2) *If $\phi \in \text{dom}(D_{\max})$, then $\chi(r) \cdot \phi \in \text{dom}(\dot{D}_{\max})$ and $\|\chi(r) \cdot \phi\|_{\dot{D}} \lesssim \|\phi\|_D$.*

Proof. Let $\phi \in \text{dom}(\dot{D}_{\max})$. Let $\eta \in C_c^\infty(U \setminus Z, [0, 1])$. By [Proposition 3.13 \(2\)](#),

$$\begin{aligned} \int_{U \setminus Z} \eta^2 |\dot{\nabla}(r\chi(r) \cdot \phi)|^2 &= \int_{U \setminus Z} \eta^2 |\dot{D}(r\chi(r) \cdot \phi)|^2 \\ &\quad - \int_{U \setminus Z} \eta^2 (\langle \tau \dot{\nabla}(r\chi(r) \cdot \phi), r\chi(r) \cdot \phi \rangle + \langle \dot{\gamma}(F_{\dot{\nabla}}) r\chi(r) \cdot \phi, r\chi(r) \cdot \phi \rangle) \\ &\quad + 2 \int_{U \setminus Z} \eta \langle \dot{D}(r\chi(r) \cdot \phi), \dot{\gamma}(d\eta) r\chi(r) \cdot \phi \rangle \\ &\quad - 2 \int_{U \setminus Z} \eta \langle \dot{\nabla}(r\chi(r) \cdot \phi), d\eta \otimes r\chi(r) \cdot \phi \rangle. \end{aligned}$$

Therefore,

$$\int_{U \setminus Z} \eta^2 |\dot{\nabla}(r\chi(r) \cdot \phi)|^2 \lesssim \int_{U \setminus Z} |\dot{D}\phi|^2 + r^2(1 + |d\eta|^2) |\phi|^2.$$

Since $\eta_\varepsilon := 1 - \chi(r/\varepsilon)$ satisfies $r|d\eta_\varepsilon| \lesssim 1$,

$$\int_{U \setminus Z} |\dot{\nabla}(r\chi(r) \cdot \phi)|^2 = \lim_{\varepsilon \downarrow 0} \int_{U \setminus Z} \eta_\varepsilon^2 |\dot{\nabla}(r\phi)|^2 \lesssim \int_{U \setminus Z} |\dot{D}\phi|^2 + |\phi|^2.$$

Therefore, $r\chi(r) \cdot \phi \in \text{dom}(\dot{D}_{\min})$ and

$$\|r\chi(r) \cdot \phi\|_{H^1} \lesssim \|\phi\|_{\dot{D}}.$$

By [Proposition 3.14](#) and the above,

$$\|\text{Err } \chi(r) \cdot \phi\|_{L^2} \lesssim \|\dot{\nabla}(r\phi)\|_{L^2} + \|\phi\|_{L^2} \lesssim \|\dot{D}\phi\|_{L^2} + \|\phi\|_{L^2}.$$

This implies (1). The proof of (2) is similar. ■

Proof of Proposition 3.17. By [Lemma 3.18](#), cut-off is an isomorphism of Hilbert spaces. To prove that cut-off is a symplectomorphism, let $\phi, \psi \in \text{dom}(D_{\max})$ and set

$$v := \sum_{i=1}^n \langle \gamma(e_i)\phi, \psi \rangle e_i \quad \text{and} \quad \check{v} := \sum_{i=1}^n \langle \dot{\gamma}(\check{e}_i)\chi(r) \cdot \phi, \chi(r) \cdot \psi \rangle \check{e}_i$$

with (e_1, \dots, e_n) and (e_1, \dots, e_n) denoting local g - and \mathring{g} -orthonormal frames respectively.

Assume, without loss of generality, that $\text{supp}(\phi) \cup \text{supp}(\psi) \subset (\chi \circ r)^{-1}(1) \subset U \setminus Z$. With η_ε as in the proof of [Lemma 3.18](#),

$$\begin{aligned} (G - \text{cut-off}^* \mathring{G})([\phi] \wedge [\psi]) &= \int_{U \setminus Z} \text{div}_g(v) \cdot \text{vol}_g - \text{div}_{\mathring{g}}(\mathring{v}) \cdot \text{vol}_{\mathring{g}} \\ &= \lim_{\varepsilon \downarrow 0} \int_{U \setminus Z} \eta_\varepsilon \cdot (\text{div}_g(v) \cdot \text{vol}_g - \text{div}_{\mathring{g}}(\mathring{v}) \cdot \text{vol}_{\mathring{g}}) \\ &= - \lim_{\varepsilon \downarrow 0} \int_{U \setminus Z} d\eta_\varepsilon \wedge (i_v \text{vol}_g - i_{\mathring{v}} \text{vol}_{\mathring{g}}). \end{aligned}$$

Since $r|d\eta_\varepsilon| \lesssim 1$,

$$|d\eta_\varepsilon \wedge (i_v \text{vol}_g - i_{\mathring{v}} \text{vol}_{\mathring{g}})| \lesssim |\phi| |\psi|.$$

Therefore,

$$|(G - \text{cut-off}^* \mathring{G})([\phi] \wedge [\psi])| \lesssim \lim_{\varepsilon \downarrow 0} \int_{\text{supp } d\eta_\varepsilon} |\phi| |\psi| \text{vol}_{\mathring{g}} = 0. \quad \blacksquare$$

3.4 Spectral decomposition

This subsection decomposes $(\mathring{H}_0, \mathring{G})$ into concretely understandable summands.

Definition 3.19 (\mathbf{I} determines NZ^λ). The ramified Euclidean line bundle \mathbf{I} determines the following:

- (1) The 2π -periodic vector field ∂_α generating $F \cup U(1)$ uniquely lifts along

$$\rho: \tilde{F} := \{\ell \in \underline{\mathbf{I}} : |\ell| = 1\} \rightarrow F$$

to a 4π -periodic vector field $\frac{1}{2}\partial_\beta$. The 2π -periodic vector field ∂_β generates $\tilde{F} \cup U(1)$ with respect to which $\tilde{\pi}: \tilde{F} \rightarrow Z$ is a $U(1)$ -principal bundle.

- (2) Let $\lambda \in \frac{1}{2}\mathbf{Z}$. The Hermitian line bundle

$$NZ^\lambda := \tilde{F} \times_{U(1)} \mathbf{C}$$

arises from \tilde{F} via the representation $U(1) \cup \mathbf{C}$ of weight 2λ . The Levi-Civita connection on F induces a connection on \tilde{F} and, therefore, a unitary covariant derivative ∇^λ on NZ^λ . •

Remark 3.20. By construction $(NZ^1, \nabla^1) \cong (NZ, \nabla^{\text{LC}})$ and for every $\lambda, \mu \in \frac{1}{2}\mathbf{Z}$

$$(NZ^\lambda, \nabla^\lambda) \otimes_{\mathbf{C}} (NZ^\mu, \nabla^\mu) \cong (NZ^{\lambda+\mu}, \nabla^{\lambda+\mu}). \quad \clubsuit$$

Proposition 3.21. For every $\lambda \in \mathbf{Z} - 1/2$ there is an isomorphism

$$P_\lambda: \pi^*(NZ^\lambda, \nabla^\lambda) \cong (\underline{\mathbf{I}} \otimes \mathbf{C}, \nabla_{\underline{\mathbf{I}} \otimes \mathbf{C}} + i\lambda\theta)$$

of Hermitian line bundles with unitary connections.

Proof. Consider the $U(1)$ -principal bundle $\tilde{F} \times_{\{\pm 1\}} U(1) \rightarrow F$ obtained by extending the $\{\pm 1\}$ -principal bundle $\tilde{F} \rightarrow F$ along the inclusion $\iota: \{\pm 1\} \hookrightarrow U(1)$. The $U(1)$ -principal bundles $\tilde{F} \times_{\{\pm 1\}} U(1) \rightarrow F$ and $\pi^* \tilde{F} \rightarrow F$ are isomorphic via $[f, z] \mapsto [f, \rho(f \cdot z)]$.

Let $\lambda \in \mathbf{Z} - 1/2$. The representation $U(1) \cup \mathbf{C}$ of weight 2λ restricts to the usual representation $\{\pm 1\} \cup \mathbf{C}$ along ι . Therefore, $\pi^* NZ^\lambda$ and $\underline{\mathbb{I}}$ both arise from the representation of weight 2λ . Hence, they are isomorphic as Hermitian line bundles.

The Levi-Civita connection $i\theta$ on $F \rightarrow \mathbf{Z}$ induces the connection $\frac{i}{2}\rho^*\theta$ on $\tilde{F} \rightarrow Z$. Therefore, the induced connection on $\tilde{F} \times_{\{\pm 1\}} U(1) \cong \pi^* F$ is (the descent of) $\frac{i}{2}\rho^*\theta + \mu_{U(1)}$. Here $\mu_{U(1)} \in U(1)$ denotes the Maurer–Cartan form on $U(1)$. The connection on $\tilde{F} \times_{\{\pm 1\}} U(1)$ induced by the flat connection on $\tilde{F} \rightarrow F$ is (the descend) of $\mu_{U(1)}$. This implies the assertion about the covariant derivatives. \blacksquare

Proposition 3.22 (Spectral decomposition of $L^2\Gamma(F, \underline{S} \otimes \underline{\mathbb{I}})$). *For every $\lambda \in \mathbf{Z} - 1/2$ and $\mu \in \mathbf{R}$ set*

$$\begin{aligned} E_{\lambda, \mu} &:= \{\phi \in L^2\Gamma(F, \underline{S} \otimes \underline{\mathbb{I}}) : I\mathring{\nabla}_{\partial_\alpha} \phi = \lambda\phi, D_Z \phi = \mu\phi\} \quad \text{and} \\ \check{E}_{\lambda, \mu} &:= \{\check{\phi} \in L^2\Gamma(Z, S|_Z \otimes_{\mathbf{C}} NZ^\lambda) : D_{S|_Z \otimes_{\mathbf{C}} NZ^\lambda} \check{\phi} = \mu\check{\phi}\}. \end{aligned}$$

Here D_Z is as in [Remark 3.12](#) and $D_{S|_Z \otimes_{\mathbf{C}} NZ^\lambda}$ arises from [Definition 3.8](#) and twisting by $(NZ^\lambda, \nabla^\lambda)$; moreover: the tensor product is with respect to the complex structure I on $S|_Z$. The following hold:

(1) For every $\lambda \in \mathbf{Z} - 1/2$ and $\mu \in \mathbf{R}$, P_λ induces an isomorphism

$$\pi^* \check{E}_{\lambda, \mu} \cong E_{\lambda, \mu}.$$

(2) The subset

$$\sigma := \{(\lambda, \mu) \in (\mathbf{Z} - 1/2) \times \mathbf{R} : E_{\lambda, \mu} \neq 0\}$$

is discrete. Moreover, for every $(\lambda, \mu) \in \sigma$, $E_{\lambda, \mu}$ is finite-dimensional.

(3) For every $(\lambda, \mu) \in \sigma$

$$JE_{\lambda, \mu} = E_{-(\lambda+1), -\mu}.$$

(4) The Hilbert space $L^2\Gamma(F, \underline{S} \otimes \underline{\mathbb{I}})$ decomposes as a (Hilbert space) direct sum

$$L^2\Gamma(F, \underline{S} \otimes \underline{\mathbb{I}}) = \bigoplus_{(\lambda, \mu) \in \sigma} E_{\lambda, \mu}.$$

Proof. By Fourier analysis, the Hilbert space $L^2\Gamma(F, \underline{S} \otimes \underline{\mathbb{I}})$ decomposes as a direct sum

$$L^2\Gamma(F, \underline{S} \otimes \underline{\mathbb{I}}) = \bigoplus_{\lambda \in \mathbf{Z} - 1/2} E_\lambda \quad \text{with} \quad E_\lambda := \{\phi \in L^2\Gamma(F, \underline{S} \otimes \underline{\mathbb{I}}) : I\mathring{\nabla}_{\partial_\alpha} \phi = \lambda\phi\}$$

By [Proposition 3.21](#), P_λ induces an isomorphism $\pi^* L^2\Gamma(Z, S|_Z \otimes_{\mathbf{C}} NZ^\lambda) \cong E_\lambda$. By the spectral theory of Dirac operators, for every $\lambda \in \mathbf{Z} - 1/2$, $\text{spec}(D_{S|_Z \otimes_{\mathbf{C}} NZ^\lambda}) \subset \mathbf{R}$ is discrete and the Hilbert space $L^2\Gamma(Z, S|_Z \otimes_{\mathbf{C}} NZ^\lambda)$ decomposes as a direct sum finite-dimensional eigenspaces $\check{E}_{\lambda, \mu}$ of $D_{S|_Z \otimes_{\mathbf{C}} NZ^\lambda}$. This proves (1), (2), and (4).

(3) holds because J and D_Z anti-commute and $I\mathring{\nabla}_{\partial_\alpha} J = -J(I\mathring{\nabla}_{\partial_\alpha} + 1)$ since $\mathring{\nabla}_{\partial_\alpha} \partial_r = \partial_\alpha$. \blacksquare

Since $\text{vol}_{\check{g}} = dr \wedge r\theta \wedge \text{vol}_{g|_Z}$, by Fubini's theorem and [Proposition 3.22](#),

$$L^2\Gamma(U \setminus Z, \mathring{S} \otimes \mathring{\mathbf{I}}) = L^2((0, 1), r dr; L^2\Gamma(F, \underline{S} \otimes \underline{\mathbf{I}})) = \bigoplus_{\lambda \in \mathbb{Z} - 1/2} \bigoplus_{\mu \in \sigma_\lambda} L^2((0, 1), r dr; E_{\lambda, \mu});$$

moreover, D_{\max} decomposes as follows.

Definition 3.23. Choose a fundamental domain $\check{\sigma} \subset \sigma$ for the involution $(\lambda, \mu) \mapsto -(\lambda+1), -\mu$. Choose a real subspace $E_{-1/2, 0}^{\mathbb{R}} \subset E_{-1/2, 0}$ with respect to J . For every $(\lambda, \mu) \in \check{\sigma}$ set

$$V_{\lambda, \mu} := \begin{cases} E_{-1/2, 0}^{\mathbb{R}} \oplus J E_{-1/2, 0}^{\mathbb{R}} & \text{if } (\lambda, \mu) = (-1/2, 0) \\ E_{\lambda, \mu} \oplus E_{-(\lambda+1), -\mu} & \text{otherwise;} \end{cases}$$

moreover, define $\mathring{D}^{\lambda, \mu}: H_{\text{loc}}^1((0, 1); V_{\lambda, \mu}) \rightarrow L_{\text{loc}}^2((0, 1); V_{\lambda, \mu})$ by

$$\mathring{D}^{\lambda, \mu} := \begin{pmatrix} \mu & J(\partial_r + \frac{\lambda+1}{r}) \\ J(\partial_r - \frac{\lambda}{r}) & -\mu \end{pmatrix}$$

and set

$$\text{dom}(\mathring{D}_{\max}^{\lambda, \mu}) := \left\{ \phi \in H_{\text{loc}}^1((0, 1); V_{\lambda, \mu}) : \phi, \mathring{D}\phi \in L^2((0, 1), r dr; V_{\lambda, \mu}) \right\}. \quad \bullet$$

Remark 3.24. The purpose of the artificial decomposition of $E_{-1/2, 0}$ is to avoid a case distinction in the definition of $\mathring{D}^{\lambda, \mu}$. ♣

Proposition 3.25 (Spectral decomposition of $\text{dom}(\mathring{D}_{\max})$). *The following hold:*

(1) *The Hilbert space $\text{dom}(\mathring{D}_{\max})$ decomposes as a (Hilbert space) direct sum*

$$\text{dom}(\mathring{D}_{\max}) = \bigoplus_{(\lambda, \mu) \in \check{\sigma}} \text{dom}(\mathring{D}_{\max}^{\lambda, \mu}).$$

(2) *The model operator \mathring{D} decomposes as*

$$\mathring{D} = \bigoplus_{(\lambda, \mu) \in \check{\sigma}} \mathring{D}^{\lambda, \mu}.$$

Proof. This is an immediate consequence of [Remark 3.12](#) and [Proposition 3.22](#). ■

Remark 3.26. The ordinary differential equation $\mathring{D}^{\lambda, \mu}\phi = \psi$ can be solved explicitly in terms of modified Bessel functions of the second kind or using the Hankel transform. However, none of this is necessary for the purpose of this article. ♣

Finally, here is the desired decomposition of $(\mathring{\mathbf{H}}_0, \mathring{G})$.

Corollary 3.27 (Spectral decomposition of $(\mathring{\mathbf{H}}_0, \mathring{G})$). *The symplectic Hilbert space $(\mathring{\mathbf{H}}_0, \mathring{G})$ decomposes as a (Hilbert space) direct sum*

$$(\mathring{\mathbf{H}}_0, \mathring{G}) = \bigoplus_{(\lambda, \mu) \in \check{\sigma}} (\mathring{\mathbf{H}}_0^{\lambda, \mu}, \mathring{G}_{\lambda, \mu})$$

with

$$\begin{aligned} \mathring{\mathbf{H}}_0^{\lambda,\mu} &:= \frac{(\chi \circ r) \cdot \text{dom}(\mathring{D}_{\max}^{\lambda,\mu}) + \text{dom}(\mathring{D}_{\min})}{\text{dom}(\mathring{D}_{\min})} \quad \text{and} \\ \mathring{G}_{\lambda,\mu}([\phi] \wedge [\psi]) &:= \langle \mathring{D}^{\lambda,\mu} \phi, \psi \rangle_{L^2} - \langle \phi, \mathring{D}^{\lambda,\mu} \psi \rangle_{L^2}. \end{aligned} \quad \blacksquare$$

3.5 Leading order terms

This subsection determines $(\mathring{\mathbf{H}}_0^{\lambda,\mu}, \mathring{G}_{\lambda,\mu})$ based on the following observation.

Lemma 3.28 (Leading order terms; cf. [BS88, Lemma 2.1; DW24, Lemma 3.50]). *Let $\lambda \in \mathbf{R}$. Let $\phi \in H_{\text{loc}}^1((0, 1))$ with $\phi, (\partial_r - \lambda/r)\phi \in L^2((0, 1), r dr)$. The following hold:*

- (1) *If $\lambda \in (-1, 0)$, then there is a unique $a \in \mathbf{R}$ such that $\lim_{r \downarrow 0} \phi(r) - ar^\lambda = 0$.*
- (2) *If $\lambda = 0$, then $\phi(r) \lesssim_\phi |\log(r)|^{1/2}$.*
- (3) *If $\lambda \notin [-1, 0)$, then $\lim_{r \downarrow 0} \phi(r) = 0$.*
- (4) *If $\lim_{r \downarrow 0} \phi(r) = \lim_{r \uparrow 1} \phi(r) = 0$, then*

$$\int_0^1 \left(|\partial_r \phi|^2 + \frac{\lambda^2}{r^2} |\phi|^2 \right) r dr = \int_0^1 |(\partial_r - \lambda/r)\phi|^2 r dr.$$

Proof. The proof is almost identical to that of [DW24, Lemma 3.50] and is repeated here only for the readers' convenience.

Evidently, $(\partial_r - \lambda/r)r^\lambda = 0$ and $r^\lambda \in L^2((0, 1), r dr)$ if and only if $\lambda > -1$. Set $\psi := (\partial_r - \lambda/r)\phi$. By variation of parameters, there is a unique $a \in \mathbf{R}$ such that

$$\tilde{\phi}(r) := \phi(r) - ar^\lambda = \begin{cases} r^\lambda \int_0^r s^{-(\lambda+1)} \psi(s) s ds & \text{if } \lambda < 0 \\ -r^\lambda \int_r^1 s^{-(\lambda+1)} \psi(s) s ds & \text{if } \lambda \geq 0. \end{cases}$$

Of course, if $\lambda \leq -1$, then $a = 0$.

If $\lambda < 0$, then, by Cauchy–Schwarz and monotone convergence,

$$|\tilde{\phi}(r)|^2 \leq \frac{1}{2|\lambda|} \int_0^r |\psi(s)|^2 s ds = o(1) \quad \text{as } r \downarrow 0.$$

If $\lambda = 0$, then

$$|\tilde{\phi}(r)|^2 \leq |\log(r)| \int_r^1 |\psi(s)|^2 s ds = O(|\log(r)|) \quad \text{as } r \downarrow 0.$$

If $\lambda > 0$, then, by Cauchy–Schwarz, for $r \leq \varepsilon \leq 1$

$$|\tilde{\phi}(r)|^2 \leq \frac{1}{\lambda} \int_0^\varepsilon |\psi(s)|^2 s ds + \frac{(r/\varepsilon)^{2\lambda}}{\lambda} \int_\varepsilon^1 |\psi(s)|^2 s ds =: \text{I}(\varepsilon) + \text{II}(r, \varepsilon).$$

By monotone convergence, $\lim_{\varepsilon \downarrow 0} \text{I}(\varepsilon) = 0$. Evidently, $\lim_{r \downarrow 0} \text{II}(r, \varepsilon) = 0$. Therefore, $\tilde{\phi}(r) = o(1)$ as $r \downarrow 0$. These observations imply (1), (2), and (3).

(4) is a consequence of

$$\int_0^1 |(\partial_r - \lambda/r)\phi|^2 r dr = \int_0^1 \left(|\partial_r \phi|^2 + \frac{\lambda^2}{r^2} |\phi|^2 \right) r dr - \lambda \int_0^1 \partial_r |\phi|^2 dr. \quad \blacksquare$$

Corollary 3.29 (Identification of $\mathring{\mathbf{H}}_0^{\lambda, \mu}$ for $\lambda \neq -1/2$). For every $(\lambda, \mu) \in \check{\sigma}$ with $\lambda \neq -1/2$

$$\mathring{\mathbf{H}}_0^{\lambda, \mu} = 0. \quad \blacksquare$$

Definition 3.30. For $(-1/2, \mu) \in \check{\sigma}$ define the **residue map** $\text{res}_\mu : \text{dom}(\mathring{D}_{\max}^{-1/2, \mu}) \rightarrow V_{-1/2, \mu}$ by

$$\phi - r^{-1/2} \cdot \text{res}_\mu(\phi) \in \text{dom}(\mathring{D}_{\min}),$$

and the symplectic form $\check{\Omega}_\mu \in \text{Hom}(\Lambda^2 V_{-1/2, \mu}, \mathbf{R})$ by

$$\check{\Omega}_\mu(\phi \wedge \psi) := -\langle J\phi, \psi \rangle. \quad \bullet$$

Proposition 3.31 (Identification of $\mathring{\mathbf{H}}_0^{-1/2, \mu}$: symplectic structure). For every $(-1/2, \mu) \in \check{\sigma}$ the residue map induces an isomorphism

$$\text{res}_\mu : (\mathring{\mathbf{H}}_0^{-1/2, \mu}, \mathring{G}^{-1/2, \mu}) \cong (V_{-1/2, \mu}, \check{\Omega}_\mu).$$

Proof. For $\phi, \psi \in \text{dom}(\mathring{D}_{\max}^{-1/2, \mu})$, by direct computation using $\partial_r + \frac{1}{2r} = r^{-1/2} \partial_r r^{1/2}$,

$$\begin{aligned} \mathring{G}^{-1/2, \mu}([\phi] \wedge [\psi]) &= \int_0^1 (\langle J(\partial_r + \frac{1}{2r})\phi, \psi \rangle - \langle \phi, J(\partial_r + \frac{1}{2r})\psi \rangle) r dr \\ &= \int_0^1 \partial_r \langle J r^{1/2} \phi, r^{1/2} \psi \rangle dr \\ &= -\langle J \text{res}_\mu([\phi]), \text{res}_\mu([\psi]) \rangle = \text{res}_\mu^* \check{\Omega}_\mu([\phi] \wedge [\psi]). \end{aligned}$$

This together with [Lemma 3.28](#) immediately implies the assertion. \blacksquare

Although res_μ is an isomorphism, the norms on $\mathring{\mathbf{H}}_0^{-1/2, \mu}$ and $V_{-1/2, \mu}$ are *not uniformly equivalent*. The following discussion rectifies this.

Definition 3.32. Let $(-1/2, \mu) \in \check{\sigma}$.

(1) Define the **branching locus operator** $A_\mu : V_{-1/2, \mu} \rightarrow V_{-1/2, \mu}$ by

$$A_\mu := -J \mathring{D}^{-1/2, \mu} - \partial_r - \frac{1}{2r} = \begin{pmatrix} 0 & J\mu \\ -J\mu & 0 \end{pmatrix}.$$

(2) Define the norm $\|-\|_{\check{H}} : V_{-1/2, \mu} \rightarrow [0, \infty)$ by

$$\|v\|_{\check{H}}^2 := (1 + |\mu|) \cdot \|\mathbf{1}_{(-\infty, 0)}(A_\mu)\phi\|^2 + (1 + |\mu|)^{-1} \cdot \|\mathbf{1}_{[0, \infty)}(A_\mu)\phi\|^2.$$

Here $\mathbf{1}_{(-\infty, 0)}(A_\mu)$ and $\mathbf{1}_{[0, \infty)}(A_\mu)$ denote the orthogonal projection to the negative and non-negative eigenspaces of A_μ respectively.

(3) Define the norm $\|\cdot\|_{H^{-1/2}} : V_{-1/2,\mu} \rightarrow [0, \infty)$ by

$$\|v\|_{H^{-1/2}}^2 := (1 + |\mu|)^{-1} \cdot \|v\|^2. \quad \bullet$$

Proposition 3.33 (Identification of $\mathring{H}_0^{-1/2,\mu}$: uniform norms). *For every $(-1/2, \mu) \in \check{\sigma}$*

$$\|\text{res}_\mu([\phi])\|_{\check{H}} \asymp \|[\phi]\|_{\mathring{H}}.$$

The proof uses the following right inverse of $\text{res}_\mu \circ [\cdot]$.

Definition 3.34. For $(-1/2, \mu) \in \check{\sigma}$ define the **extension map** $\text{ext}_\mu : V_{-1/2,\mu} \rightarrow \text{dom}(\mathring{D}_{\max}^{-1/2,\mu})$ by

$$\text{ext}_\mu(v) := r^{-1/2} e^{-|\mu|r} \cdot v. \quad \bullet$$

Evidently, ext_μ lifts the inverse of $\text{res}_\mu : \mathring{H}_0^{-1/2,\mu} \cong V_{-1/2,\mu}$. Therefore, [Proposition 3.33](#) is an immediate consequence of the following.

Lemma 3.35 (Uniform estimates for res_μ and ext_μ). *Let $(-1/2, \mu) \in \check{\sigma}$. The following hold:*

(1) For every $\phi \in \text{dom}(\mathring{D}_{\max}^{-1/2,\mu})$

$$\|\text{res}_\mu([\phi])\|_{\check{H}} \lesssim \|\phi\|_{\mathring{D}}.$$

(2) For every $v \in V_{-1/2,\mu}$

$$\|\text{ext}_\mu(v)\|_{L^2} \lesssim \|v\|_{H^{-1/2}} \quad \text{and} \quad \|\text{ext}_\mu(v)\|_{\mathring{D}} \lesssim \|v\|_{\check{H}}.$$

Proof. Evidently, $v := \text{res}_\mu([\phi])$ satisfies

$$v = - \int_0^1 \partial_r (r^{1/2} e^{-|\mu|r} \phi) \, dr = - \int_0^1 r^{1/2} (-J\mathring{D}^{-1/2,\mu} - A_\mu - |\mu|) \phi \cdot e^{-|\mu|r} \, dr.$$

Therefore, by Cauchy–Schwarz,

$$\begin{aligned} \|\mathbf{1}_{(-\infty,0)}(A_\mu)v\|^2 &\lesssim \int_0^1 \|\mathring{D}^{-1/2,\mu} \mathbf{1}_{(-\infty,0)}(A_\mu)\phi\|^2 r \, dr \cdot \int_0^1 e^{-2|\mu|r} \, dr \\ &\lesssim \int_0^1 e^{-2|\mu|r} \, dr \cdot \|\mathbf{1}_{(-\infty,0)}(A_\mu)\phi\|_{\mathring{D}}^2 \end{aligned}$$

and

$$\begin{aligned} \|\mathbf{1}_{[0,\infty)}(A_\mu)v\|^2 &\lesssim \int_0^1 \left(\|\mathring{D}^{-1/2,\mu} \mathbf{1}_{[0,\infty)}(A_\mu)\phi\|^2 + |\mu|^2 \|\mathbf{1}_{[0,\infty)}(A_\mu)\phi\|^2 \right) r \, dr \cdot \int_0^1 e^{-2|\mu|r} \, dr \\ &\lesssim (1 + |\mu|)^2 \cdot \int_0^1 e^{-2|\mu|r} \, dr \cdot \|\mathbf{1}_{[0,\infty)}(A_\mu)\phi\|_{\mathring{D}}^2. \end{aligned}$$

The estimate in (1) follows because

$$\int_0^1 e^{-2|\mu|r} \, dr \lesssim (1 + |\mu|)^{-1} \quad \text{and} \quad \|\mathbf{1}_{(-\infty,0)}(A_\mu)\phi\|_{\mathring{D}}^2 + \|\mathbf{1}_{[0,\infty)}(A_\mu)\phi\|_{\mathring{D}}^2 = \|\phi\|_{\mathring{D}}^2.$$

To prove (2), observe that

$$\|\text{ext}_\mu(v)\|_{L^2}^2 \lesssim \int_0^1 r^{-1} e^{-2|\mu|r} \|v\|^2 r dr \lesssim (1 + |\mu|)^{-1} \cdot \|v\|^2$$

and

$$\begin{aligned} \|\mathring{D}^{-1/2, \mu} \text{ext}_\mu(v)\|_{L^2}^2 &= \int_0^1 (r^{-1/2} e^{-|\mu|r})^2 \|(-|\mu| + A_\mu)v\|^2 r dr \\ &\lesssim \int_0^1 (r^{-1/2} |\mu| e^{-|\mu|r})^2 \|\mathbf{1}_{(-\infty, 0]}(A_\mu)v\|^2 r dr \\ &\lesssim \mu^2 (1 + |\mu|)^{-1} \cdot \|v\|^2. \quad \blacksquare \end{aligned}$$

3.6 Assembly of the residue map

This subsection (re)assembles the summands of the decomposition [Corollary 3.27](#) identified in [Section 3.5](#) in a more geometric fashion.

Definition 3.36.

- (1) The residue bundle is

$$\check{S} := S|_Z \otimes_{\mathbb{C}} NZ^{-1/2}.$$

As a consequence of [Proposition 3.22 \(3\)](#) (or by direct inspection), \check{S} inherits the quaternionic structure $I, J, K = IJ$ from \underline{S} . Define the symplectic form $\check{\Omega} \in \Gamma(Z, \text{Hom}(\Lambda^2 \check{S}, \mathbf{R}))$ by

$$\check{\Omega} := -2\pi \langle J-, - \rangle.$$

- (2) The branching locus operator $A: \Gamma(Z, \check{S}) \rightarrow \Gamma(Z, \check{S})$ is defined by

$$A := -JD_{\check{S}}$$

with $D_{\check{S}} := D_{S|_Z \otimes_{\mathbb{C}} NZ^{-1/2}}$ as in [Proposition 3.22](#). Since J and $D_{\check{S}}$ anti-commute, A is (formally) self-adjoint.

- (3) Denote by $\mathbf{1}_{(-\infty, 0)}(A)$ and $\mathbf{1}_{[0, \infty)}(A)$ the orthogonal projection to the negative and non-negative eigenspaces of A respectively. Define the norm $\|-\|_{\check{H}}: \Gamma(Z, \check{S}) \rightarrow [0, \infty)$ by

$$\|\phi\|_{\check{H}} := \|\mathbf{1}_{(-\infty, 0)}(A)\phi\|_{H^{1/2}} + \|\mathbf{1}_{[0, \infty)}(A)\phi\|_{H^{-1/2}}$$

and denote by $\check{H}\Gamma(Z, \check{S})$ the completions of $\Gamma(Z, \check{S})$ with respect to $\|-\|_{\check{H}}$. •

Proposition 3.37. $\check{\Omega}$ extends to a symplectic structure $\check{\Omega} \in \mathcal{L}(\Lambda^2 \check{H}\Gamma(Z, \check{S}), \mathbf{R})$; moreover: the inclusion $V_{-1/2, \mu} \hookrightarrow \Gamma(Z, \check{S})$ assemble into an isomorphism of symplectic Hilbert spaces

$$\bigoplus_{(-1/2, \mu) \in \check{\sigma}} (V_{-1/2, \mu}, \|-\|_{\check{H}}; \check{\Omega}_\mu) \cong (\check{H}\Gamma(Z, \check{S}); \check{\Omega}).$$

Proof. This is an immediate consequence of [Proposition 3.22](#), [Proposition 3.31](#), and [Proposition 3.33](#). The (possibly mysterious) factor 2π arises because $\check{\Omega}_\mu$ is defined using the L^2 inner product on F instead of Z and $\text{vol}(F) = 2\pi\text{vol}(Z)$. \blacksquare

Definition 3.38.

- (1) The **residue map** $\text{res}: \check{\mathbf{H}} \rightarrow \check{H}\Gamma(Z, \check{S})$ obtained as the composition of the following maps

$$\check{\mathbf{H}} \xrightarrow{\text{cut-off}} \check{\mathbf{H}}_0 = \bigoplus_{(-1/2, \mu) \in \check{\sigma}} \check{\mathbf{H}}_0^{-1/2, \mu} \xrightarrow{(\text{res}_\mu)} \bigoplus_{(-1/2, \mu) \in \check{\sigma}} (V_{-1/2, \mu}, \|\cdot\|_{\check{H}}) \cong \check{H}\Gamma(Z, \check{S}).$$

- (2) The **extension map** $\text{ext}: \check{H}\Gamma(Z, \check{S}) \rightarrow \text{dom}(D_{\max})$ is obtained as the composition of the following maps

$$\begin{aligned} \check{H}\Gamma(Z, \check{S}) \cong \bigoplus_{(-1/2, \mu) \in \check{\sigma}} (V_{-1/2, \mu}, \|\cdot\|_{\check{H}}) &\xrightarrow{(\text{ext}_\mu)} \bigoplus_{(-1/2, \mu) \in \check{\sigma}} \text{dom}(D_{\max}^{-1/2, \mu}) \\ &\hookrightarrow \text{dom}(D_{\max}) \xrightarrow{\chi^{\circ r}} \text{dom}(D_{\max}). \quad \bullet \end{aligned}$$

Theorem 3.39. *The following hold:*

- (1) (a) *The residue map is an isomorphism of symplectic Hilbert spaces*

$$\text{res}: (\check{\mathbf{H}}, G) \cong (\check{H}\Gamma(Z, \check{S}), \check{\Omega}).$$

- (b) *The subspace $r^{-1/2}\Gamma(\hat{X}, \hat{S} \otimes \hat{\mathbf{I}}) \cap \text{dom}(D_{\max})$ is dense in $\text{dom}(D_{\max})$; hence: the residue map is uniquely determined by*

$$\pi^* \text{res}[r^{-1/2}\phi] = \phi|_{\partial\hat{X}}$$

for every $r^{-1/2}\phi \in r^{-1/2}\Gamma(\hat{X}, \hat{S} \otimes \hat{\mathbf{I}}) \cap \text{dom}(D_{\max})$.

- (2) (a) *The extension map $\text{ext}: \check{H}\Gamma(Z, \check{S}) \rightarrow \text{dom}(D_{\max})$ is a right-inverse of $\text{res} \circ [-]$.*
(b) *The extension map extends to a bounded linear map*

$$\text{ext}: H^{-1/2}\Gamma(Z, \check{S}) \rightarrow L^2\Gamma(X \setminus Z, S \otimes \mathbf{I}).$$

Proof. (1.a) is an immediate consequence of [Proposition 3.17](#), [Corollary 3.27](#), [Proposition 3.31](#), [Proposition 3.33](#), and [Proposition 3.37](#).

(1.b) is a consequence of [Lemma 3.18](#) and [Proposition 3.25](#).

(2.a) holds by construction and (2.b) follows from [Lemma 3.35 \(2\)](#). \blacksquare

3.7 Spectral and local residue conditions

Theorem 3.39 makes it possible to define a wider variety of residue conditions than those considered in [Section 2](#). Here are some examples.

Example 3.40. The **APS residue condition** is defined by

$$B_{\text{APS}} := \mathbf{1}_{(-\infty, 0)}(A)H^{1/2}\Gamma(Z, \check{S}) \subset \check{H}\Gamma(Z, \check{S});$$

cf. Atiyah, Patodi, and Singer [[APS75](#), (2.3)]. ♠

Proposition 3.41 (Criterion for left semi-Fredholmness). *Let $B \subset \check{H}\Gamma(Z, \check{S})$ be a residue condition. If $B \hookrightarrow H^{-1/2}\Gamma(Z, \check{S})$ is compact, then $\text{dom}(D_B) \hookrightarrow L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})$ is compact and D_B is left semi-Fredholm.*

Proof. By [Lemma 2.4 \(2\)](#), and the assumption, the composition

$$\begin{aligned} \text{dom}(D_B) &\rightarrow H^1\Gamma(X \setminus Z, S \otimes \mathbb{I}) \oplus B \hookrightarrow L^2\Gamma(X \setminus Z, S \otimes \mathbb{I}) \oplus H^{-1/2}\Gamma(Z, \check{S}) \\ \phi &\mapsto (\phi - \text{ext res}[\phi], \text{res}[\phi]) \end{aligned}$$

is compact. Therefore, by [Theorem 3.39 \(2.b\)](#), $\text{dom}(D_B) \hookrightarrow L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})$ is compact. Since for every $\phi \in \text{dom}(D_B)$

$$\|\phi\|_D \lesssim \|D\phi\|_{L^2} + \|\phi\|_{L^2},$$

D_B is left semi-Fredholm. ■

Example 3.42. Since $B_{\text{APS}}^G = B_{\text{APS}} \oplus \ker A$, by [Proposition 2.26](#) and [Proposition 3.41](#), $D_{B_{\text{APS}}}$ is Fredholm of index $-\frac{1}{2} \dim \ker A$. In particular, $\dim \ker A$ is even and inherits a symplectic structure from G . If $L \subset \ker A$ is Lagrangian, then $B_{\text{APS}} \oplus L \hookrightarrow H^{-1/2}\Gamma(Z, \check{S})$ is compact and Lagrangian. In particular, [Proposition 2.20](#) applies. ♠

Definition 3.43. Let $V \subset \check{S}$ be a subbundle. The **local residue condition** associated with V is

$$B_V := \check{H}\Gamma(Z, V) \subset \check{H}\Gamma(Z, \check{S}). \quad \bullet$$

Proposition 3.44. *Let $V \subset \check{S}$ be a subbundle. If V^Ω denotes the symplectic complement of $V \subset \check{S}$, then*

$$B_V^G = B_{V^\Omega}.$$

Proof. This is an immediate consequence of [Proposition 2.16](#) and [Theorem 3.39](#). ■

Example 3.45. Consider the Dirac bundle (S, γ, ∇) corresponding to the Hodge–de Rham operator $d + d^*$; that is: $S := \Lambda T^*X$ with $\gamma(\xi)\phi := \xi \wedge \phi - i_{\xi\#}\phi$. Decompose

$$\check{S} = \check{S}_N \oplus \check{S}_D$$

with $\check{S}_N := S_N \otimes_{\mathbb{C}} NZ^{-1/2}$ and $\check{S}_D := S_D \otimes_{\mathbb{C}} NZ^{-1/2}$, and

$$S_N := (\mathbb{R} \oplus \Lambda^2 N^*Z) \otimes \Lambda T^*Z \quad \text{and} \quad S_D := N^*Z \otimes \Lambda T^*Z.$$

The corresponding residue conditions are Lagrangian. ♠

Example 3.46. Assume that (S, γ, ∇) is a complex Dirac bundle. The MIT bag residue conditions are the local residue conditions arising from the decomposition

$$\check{S} = \check{S}^+ \oplus \check{S}^- \quad \text{with} \quad \check{S}^\pm := \ker(1 \mp iJ);$$

cf. [Joh75]. ♠

Proposition 3.47 (Variation on the bordism theorem). *Assume the situation of Example 3.46. The components A^\pm of A in the decomposition*

$$A =: \begin{pmatrix} 0 & A^- \\ A^+ & 0 \end{pmatrix}$$

satisfy

$$\text{index } A^\pm = 0.$$

Proof. The following proof is essentially identical to the one presented in [BB12, §8.5]. Since $(A^+)^* = A^-$,

$$-\text{index } A^- = \text{index } A^+ = \dim \ker A^+ - \dim \ker A^-.$$

For every $t \in [0, 1]$, set

$$B_t^\pm := \ker A^\pm \oplus (1 \mp tiJ)B_{\text{APS}}.$$

Since $(B_t^+)^G = B_t^-$, by Proposition 3.41, $D_{B_t^\pm}$ is Fredholm. Moreover, by Proposition 2.26,

$$-\text{index } D_{B_t^-} = \text{index } D_{B_t^+} = \text{index } D_{(1 \mp tiJ)B_{\text{APS}}} + \dim \ker A^\pm.$$

In particular,

$$\text{index } A^+ = 2 \text{index } D_{B_0^+}.$$

Therefore, it remains to prove that $\text{index } D_{B_0^+} = 0$. By Proposition 2.29, it suffices to prove that $\text{index } D_{B_1^+} = 0$. Since

$$B_1^\pm = H^{1/2}\Gamma(Z, \check{S}^\pm),$$

analogous to Example 2.25, for every $\phi \in \ker D_{B_1^\pm}$

$$0 = 2\langle D\phi, i\phi \rangle_{L^2} = -2\pi \langle J \text{res } \phi, i \text{res } \phi \rangle = \mp 2\pi \|\text{res } \phi\|_{L^2}^2.$$

Therefore, $\ker D_{B_1^\pm} = \ker D_{\text{min}}$; hence: $\text{index } D_{B_1^\pm} = 0$. ■

4 Regularity theory

This section continues to assume Hypothesis 3.1 throughout. The geometric realisation $\check{H}(Z, \check{S})$ of \check{H} developed in Section 3 and suitable commutator estimates lead to the L^2 regularity theory laid out in the following.

4.1 Adapted Sobolev spaces, I: definition

Here is the scale of Sobolev spaces for which the regularity theory is developed.

Definition 4.1 (Differential operators). Denote by $\text{DiffOp}^\bullet(S \otimes \mathbb{I})$ the \mathbb{N}_0 -filtered ring of differential operators acting on $S \otimes \mathbb{I}$.

- (1) A vector field $v \in \text{Vect}(\hat{X})$ is **conormal** if $v|_{\partial\hat{X}} \in \text{Vect}(\partial\hat{X})$. Denote the subspace of conormal vector fields by $\text{Vect}_b(X \setminus Z)$.
- (2) The filtered subring $\text{DiffOp}_b^\bullet(S \otimes \mathbb{I}) \subset \text{DiffOp}^\bullet(S \otimes \mathbb{I})$ of **conormal** differential operators is generated by $\Gamma(\hat{X}, \text{End}(\hat{S} \otimes \hat{\mathbb{I}}))$ and differential operators of the form ∇_v with $v \in \text{Vect}_b(\hat{X})$.
- (3) The filtered subring $\text{DiffOp}_a^\bullet(S \otimes \mathbb{I}) \subset \text{DiffOp}^\bullet(S \otimes \mathbb{I})$ of **adapted** differential operators is generated by $\text{DiffOp}_b^\bullet(S \otimes \mathbb{I})$ and D . •

Definition 4.2 (Sobolev spaces). Let $k \in \mathbb{N}_0$.

- (b) The **conormal Sobolev space** $H_b^k\Gamma(X \setminus Z, S \otimes \mathbb{I})$ is defined by

$$H_b^k\Gamma(X \setminus Z, S \otimes \mathbb{I}) := \left\{ \phi \in H_{\text{loc}}^k\Gamma(X \setminus Z, S \otimes \mathbb{I}) : \begin{array}{l} P\phi \in L^2\Gamma(X \setminus Z, S \otimes \mathbb{I}) \text{ for} \\ \text{every } P \in \text{DiffOp}_b^k(S \otimes \mathbb{I}) \end{array} \right\}.$$

Choose a finite subset $\mathcal{P}_b^k \subset \text{DiffOp}_b^k(S \otimes \mathbb{I})$ which spans $\text{DiffOp}_b^k(S \otimes \mathbb{I})$ over $\Gamma(\hat{X}, \text{End}(\hat{S} \otimes \hat{\mathbb{I}}))$. Define the norm $\|\cdot\|_{H_b^k} : H_b^k\Gamma(X \setminus Z, S \otimes \mathbb{I}) \rightarrow [0, \infty)$ by

$$\|\phi\|_{H_b^k}^2 := \sum_{P \in \mathcal{P}_b^k} \|P\phi\|_{L^2}^2.$$

- (a) The **adapted Sobolev space** $H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I})$ is defined by

$$H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I}) := \left\{ \phi \in H_{\text{loc}}^k\Gamma(X \setminus Z, S \otimes \mathbb{I}) : \begin{array}{l} P\phi \in L^2\Gamma(X \setminus Z, S \otimes \mathbb{I}) \text{ for} \\ \text{every } P \in \text{DiffOp}_a^k(S \otimes \mathbb{I}) \end{array} \right\}.$$

Choose a finite subset $\mathcal{P}_a^k \subset \text{DiffOp}_a^k(S \otimes \mathbb{I})$ which spans $\text{DiffOp}_a^k(S \otimes \mathbb{I})$ over $\Gamma(\hat{X}, \text{End}(\hat{S} \otimes \hat{\mathbb{I}}))$. Define the norm $\|\cdot\|_{H_a^k} : H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I}) \rightarrow [0, \infty)$ by

$$\|\phi\|_{H_a^k}^2 := \sum_{P \in \mathcal{P}_a^k} \|P\phi\|_{L^2}^2. \quad \bullet$$

$(H_b^k\Gamma(X \setminus Z, S \otimes \mathbb{I}), \|\cdot\|_{H_b^k})$ and $(H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I}), \|\cdot\|_{H_a^k})$ are Hilbert spaces. Evidently, different choices of $\mathcal{P}_b^k, \mathcal{P}_a^k$ lead to equivalent norms. The following discussion leads to particularly convenient choices of $\mathcal{P}_b^k, \mathcal{P}_a^k$.

Definition 4.3 (Convenient vector fields).

- (1) Denote by $\text{Vect}_c(X \setminus Z) \subset \text{Vect}_b(\hat{X})$ the subspace of vector fields supported in $X \setminus Z \subset \hat{X}$.
- (2) Denote by $\text{Vect}_{b;c}(\hat{U}) \subset \text{Vect}_b(\hat{X})$ the subspace of vector fields supported in $\hat{U} \subset \hat{X}$. For $v \in \text{Vect}_{b;c}(\hat{U})$, $\nabla_v \in \text{DiffOp}_b^1(S \otimes \mathbb{I})$.
- (3) Denote by $\text{Vect}_{b;c,0}(\hat{U})$ the subspace of those $v \in \text{Vect}_{b;c}(\hat{U})$ which are $U(1)$ -invariant on ∂X ; that is: $[\partial_\alpha, v|_{\partial \hat{X}}] = 0$. •

Remark 4.4. $\text{Vect}_{b;c}(\hat{U})$ is generated by $\chi(r) \cdot \partial_\alpha$, $\chi(r) \cdot r \partial_r$, vector field of the form $\chi(r) \cdot v$ where v is lifted from Z , and vector field vanishing near Z . ♣

Lemma 4.5 (Commutation relations). *The following commutation relations hold:*

- (1) For every $v \in \text{Vect}_c(X \setminus Z)$, $k \in \mathbb{N}$, and $P \in \text{DiffOp}_a^k(S \otimes \mathbb{I})$

$$[\nabla_v, P] \in \text{DiffOp}_b^k(S \otimes \mathbb{I}).$$

- (2) For every $v \in \text{Vect}_{b;c}(\hat{U})$, $k \in \mathbb{N}$, and $P \in \text{DiffOp}_b^k(S \otimes \mathbb{I})$

$$[\mathring{\nabla}_v, P] \in \text{DiffOp}_b^k(S \otimes \mathbb{I}).$$

- (3) For every $v \in \text{Vect}_{b;c,0}(\hat{U})$

$$[\mathring{\nabla}_v, D] \in \text{DiffOp}_b^1(S \otimes \mathbb{I}) + \text{DiffOp}_b^0(S \otimes \mathbb{I}) \cdot D.$$

Proof. If $v, w \in \text{Vect}(X \setminus Z)$ and $T \in \Gamma(X \setminus Z, S \otimes \mathbb{I})$, then

$$[\nabla_v, T] = \nabla_v T \quad \text{and} \quad [\nabla_v, \nabla_w] = \nabla_{[v,w]} + F_\nabla(v, w).$$

Therefore, for every $k \in \mathbb{N}$ and $P \in \text{DiffOp}_a^k(S \otimes \mathbb{I})$, $[\nabla_v, P] \in \text{DiffOp}_a^k(S \otimes \mathbb{I})$.

If $v \in \text{Vect}_c(X \setminus Z)$, $k \in \mathbb{N}$, and $P \in \text{DiffOp}_a^k(S \otimes \mathbb{I})$, then $\text{supp}([\nabla_v, P]) \subset X \setminus Z$; therefore and by the above observation, $[\nabla_v, P] \in \text{DiffOp}_b^k(S \otimes \mathbb{I})$. This proves (1).

(2) is immediate from the above observation.

Let $v \in \text{Vect}_{b;c,0}(\hat{U})$. By [Proposition 3.14](#), $D - \chi(r) \cdot \mathring{D} \in \text{DiffOp}_b^1(S \otimes \mathbb{I})$. Therefore, it suffices to prove that

$$[\mathring{\nabla}_v, \chi(r) \cdot \mathring{D}] \in \text{DiffOp}_b^1(S \otimes \mathbb{I}) + \text{DiffOp}_b^0(S \otimes \mathbb{I}) \cdot \chi(r) \cdot \mathring{D}.$$

By direct computation,

$$[\mathring{\nabla}_{r \partial_r}, \mathring{\nabla}_{\partial_r} - r^{-1} I \mathring{\nabla}_{\partial_\alpha}] = -(\mathring{\nabla}_{\partial_r} - r^{-1} I \mathring{\nabla}_{\partial_\alpha}) \quad \text{and} \quad [\mathring{\nabla}_{\partial_\alpha}, \mathring{\nabla}_{\partial_r} - r^{-1} I \mathring{\nabla}_{\partial_\alpha}] = 0;$$

moreover, if v is the lift of a vector field along Z , then

$$[\mathring{\nabla}_v, \mathring{\nabla}_{\partial_r} - r^{-1} I \mathring{\nabla}_{\partial_\alpha}] = 0.$$

By [Remark 3.12](#), [Remark 4.4](#) and since $\chi(r) \cdot D_Z \in \text{DiffOp}_b^1(S \otimes \mathbb{I})$ and $\chi(r) \cdot J \in \text{DiffOp}_b^0(S \otimes \mathbb{I})$, this implies (3). ■

Corollary 4.6 (Convenient choices of $\mathcal{P}_b^k, \mathcal{P}_a^k$).

- (1) Set $\mathcal{P}_b^0 := \{\text{id}_{S \otimes \mathbb{I}}\}$. For every $k \in \mathbb{N}$ there is a finite subset $\mathcal{P}_b^k \subset \text{DiffOp}_b^k(S \otimes \mathbb{I})$ which spans $\text{DiffOp}_b^k(S \otimes \mathbb{I})$ over $\Gamma(\hat{X}, \text{End}(\hat{S} \otimes \hat{\mathbb{I}}))$ such that $\mathcal{P}_b^{k-1} \subset \mathcal{P}_b^k$ and every $P \in \mathcal{P}_b^k \setminus \mathcal{P}_b^{k-1}$ is of the form

$$P = \nabla_{u_1} \cdots \nabla_{u_\ell} \overset{\circ}{\nabla}_{v_1} \cdots \overset{\circ}{\nabla}_{v_{k-\ell}}$$

with

$$u_1, \dots, u_\ell \in \text{Vect}_c(X \setminus Z) \quad \text{and} \quad v_1, \dots, v_{k-\ell} \in \text{Vect}_{b;c,0}(\hat{U}).$$

- (2) Let $k \in \mathbb{N}$. If $\mathcal{P}_b^1, \dots, \mathcal{P}_b^k$ are as above then

$$\mathcal{P}_a^k := \bigcup_{\ell=0}^k \{PD^\ell : P \in \mathcal{P}_b^{k-\ell}\}$$

spans $\text{DiffOp}_a^k(S \otimes \mathbb{I})$ over $\Gamma(\hat{X}, \text{End}(\hat{S} \otimes \hat{\mathbb{I}}))$. ■

Henceforth, for every $k \in \mathbb{N}$, $\mathcal{P}_a^k, \mathcal{P}_b^k$ are assumed to be chosen as in [Corollary 4.6](#); in particular,

$$(4.7) \quad \|\phi\|_{H_a^k}^2 = \sum_{\ell=0}^k \|D^\ell \phi\|_{H_b^{k-\ell}}^2.$$

Remark 4.8. Let (W, g) be a Riemannian manifold with boundary equipped with a Dirac bundle (S, γ, ∇) . The conormal Sobolev space $H_b^k \Gamma(W, S)$; see, e.g., [\[Mel93, \(5.42\)\]](#); is an important tool in the study of boundary value problems for Dirac operators. A moment's thought with the above discussion in mind shows that $H_a^k \Gamma(W, S)$, the analogue of the adapted Sobolev space, agrees with the usual Sobolev space $H^k \Gamma(W, S)$. Indeed, these are the appropriate Sobolev spaces for the L^2 regularity theory; see [\[BB12, §6.2\]](#). ♣

4.2 Elliptic regularity and estimates

Here is the fundamental regularity result.

Theorem 4.9 (Elliptic regularity and estimates, I). *Let $k \in \mathbb{N}_0$. If $\phi \in H_{\text{loc}}^{k+1} \Gamma(X \setminus Z, S \otimes \mathbb{I}) \cap \text{dom}(D_{\text{max}})$ satisfies $D\phi \in H_a^k \Gamma(X \setminus Z, S \otimes \mathbb{I})$ and $\text{res}[\phi] \in H^{k+\frac{1}{2}} \Gamma(Z, \check{S})$, then $\phi \in H_a^{k+1} \Gamma(X \setminus Z, S \otimes \mathbb{I})$ and*

$$\|\phi\|_{H_a^{k+1}} \lesssim_k \|D\phi\|_{H_a^k} + \|\phi\|_{L^2} + \|\text{res}[\phi]\|_{H^{k+1/2}}.$$

The proof relies on the following observation.

Lemma 4.10. *The extension map $\text{ext}: \check{H} \Gamma(Z, \check{S}) \rightarrow \text{dom}(D_{\text{max}})$ restricts to a bounded linear map*

$$\text{ext}: H^{1/2} \Gamma(Z, \check{S}) \rightarrow H_a^1 \Gamma(X \setminus Z, S \otimes \mathbb{I}).$$

Proof. By (4.7), it suffices to prove that for every $(-1/2, \mu) \in \check{\sigma}$ and $\phi \in V_{-1/2, \mu} \subset H^{1/2}\Gamma(Z, \check{S})$

$$\|\text{ext}(\phi)\|_{H_b^1} \lesssim (1 + |\mu|)^{1/2} |\phi| \asymp \|\phi\|_{H^{1/2}}.$$

By direct computation,

$$\begin{aligned} \mathring{\nabla}_{r\partial_r}(\chi(r)r^{-1/2}e^{-|\mu|r}\phi) &= (r \cdot \chi'(r) - (\tfrac{1}{2} + |\mu|r) \cdot \chi(r))r^{-1/2}e^{-|\mu|r}\phi \quad \text{and} \\ \mathring{\nabla}_{\partial_\alpha}(\chi(r)r^{-1/2}e^{-|\mu|r}\phi) &= \tfrac{1}{2}\chi(r)r^{-1/2}e^{-|\mu|r}I\phi. \end{aligned}$$

Therefore,

$$\begin{aligned} \|\mathring{\nabla}_{r\partial_r}(\chi(r)r^{-1/2}e^{-|\mu|r}\phi)\|_{L^2}^2 &\lesssim (1 + |\mu|)^2 \int_0^1 e^{-2|\mu|r} dr \cdot \|\phi\|^2 \lesssim (1 + |\mu|) \cdot \|\phi\|^2 \quad \text{and} \\ \|\mathring{\nabla}_{\partial_\alpha}(\chi(r)r^{-1/2}e^{-|\mu|r}\phi)\|_{L^2}^2 &\lesssim (1 + |\mu|)^{-1} \cdot \|\phi\|^2. \end{aligned}$$

Moreover, if v is the lift of a vector field along Z , then

$$\begin{aligned} \|\mathring{\nabla}_v(\chi(r)r^{-1/2}e^{-|\mu|r}\phi)\|_{L^2}^2 &\lesssim_v \|(\chi(r)r^{-1/2}e^{-|\mu|r})\phi\|_{L^2}^2 + \|(\chi(r)r^{-1/2}e^{-|\mu|r})A\phi\|_{L^2}^2 \\ &= (1 + |\mu|)^2 \|(\chi(r)r^{-1/2}e^{-|\mu|r})\phi\|_{L^2}^2 \\ &\lesssim (1 + |\mu|)^{-1} \cdot \|\phi\|^2. \end{aligned}$$

This proves the assertion. ■

Proof of Theorem 4.9. Let $k \in \mathbf{N}_0$, $\phi \in H_{\text{loc}}^{k+1}\Gamma(X \setminus Z, S \otimes \mathbf{I}) \cap \text{dom}(D_{\max})$ with $D\phi \in H_a^k\Gamma(X \setminus Z, S \otimes \mathbf{I})$ and $\text{res}[\phi] \in H^{k+1/2}\Gamma(Z, \check{S})$.

Since $H^1\Gamma(X \setminus Z, S \otimes \mathbf{I}) \hookrightarrow H_a^1\Gamma(X \setminus Z, S \otimes \mathbf{I})$ and by Lemma 4.10,

$$\phi = (\phi - \text{ext res}[\phi]) + \text{ext res}[\phi] \in H_a^1\Gamma(X \setminus Z, S \otimes \mathbf{I}).$$

Moreover,

$$\begin{aligned} \|\phi\|_{H_a^1} &\lesssim \|\phi - \text{ext res}[\phi]\|_D + \|\text{res}[\phi]\|_{H^{1/2}} \\ &\lesssim \|D\phi\|_{L^2} + \|\phi\|_{L^2} + \|\text{res}[\phi]\|_{H^{1/2}}. \end{aligned}$$

This proves the assertion if $k = 0$.

Suppose that $k \in \mathbf{N}$. If $P \in \mathcal{P}_a^k$, then $\|\text{res } P\phi\|_{H^{1/2}} \lesssim_P \|\text{res}[\phi]\|_{H^{k+1/2}}$ and, by Lemma 4.5,

$$\|DP\phi\|_{L^2} \lesssim_P \|D\phi\|_{H_a^k} + \|\phi\|_{H_a^k} + \|\text{res}[\phi]\|_{H^{k+1/2}}.$$

Therefore, by the above, $P\phi \in H_a^1\Gamma(X \setminus Z, S \otimes \mathbf{I})$ and

$$\|P\phi\|_{H_a^1} \lesssim_P \|D\phi\|_{H_a^k} + \|\phi\|_{H_a^k} + \|\text{res}[\phi]\|_{H^{k+1/2}}.$$

This implies the assertion. ■

For suitable residue conditions $B \subset \check{H}\Gamma(Z, \check{S})$, the term $\text{res}[\phi]$ in [Theorem 4.9](#) can be absorbed provided $\text{res}[\phi] \in B$.

Definition 4.11. Let $B \subset \check{H}\Gamma(Z, \check{S})$ be a residue condition.

(1) Let $k \in \mathbb{N}_0$. B is $(k + \frac{1}{2})$ -regular if for every $\phi \in B$

$$\|\phi\|_{H^{k+1/2}} \lesssim_{B,k} \|\mathbf{1}_{(-\infty,0)}(A)\phi\|_{H^{k+1/2}} + \|\phi\|_{\check{H}}.$$

(2) B is ∞ -regular if it is $(k + 1/2)$ -regular for every $k \in \mathbb{N}_0$. •

Example 4.12. The APS residue condition B_{APS} is ∞ -regular. ♠

Theorem 4.13 (Elliptic regularity and estimates, II). *Let $k \in \mathbb{N}_0$. Let B be a $(k + \frac{1}{2})$ -regular residue condition. If $\phi \in H_{\text{loc}}^{k+1}\Gamma(X \setminus Z, S \otimes \mathbb{I}) \cap \text{dom}(D_{\text{max}})$ satisfies $D\phi \in H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I})$ and $\text{res}[\phi] \in B$, then $\phi \in H_a^{k+1}\Gamma(X \setminus Z, S \otimes \mathbb{I})$ and*

$$\|\phi\|_{H_a^{k+1}} \lesssim_{B,k} \|D\phi\|_{H_a^k} + \|\phi\|_{L^2}.$$

The proof requires the following preparation.

Lemma 4.14. *For every $k \in \mathbb{N}_0$ and $\phi \in \text{dom}(D_{\text{max}})$*

$$\|\mathbf{1}_{(-\infty,0)}(A) \text{res}[\phi]\|_{H^{k+1/2}} \lesssim_k \|D\phi\|_{H_b^k} + \|\phi\|_{H_b^k}.$$

Proof. Since $\text{res} \circ [\cdot] : \text{dom}(D_{\text{max}}) \rightarrow \check{H}\Gamma(Z, \check{S})$ is bounded and $\chi(r) \cdot A^k \in \text{DiffOp}_b^k(S \otimes \mathbb{I})$,

$$\begin{aligned} \|\mathbf{1}_{(-\infty,0)}(A) \text{res}[\phi]\|_{H^{k+1/2}} &\lesssim \|A^k \text{res}[\phi]\|_{\check{H}} = \|\text{res}[\chi(r) \cdot A^k \phi]\|_{\check{H}} \lesssim \|\chi(r) \cdot A^k \phi\|_D \\ &\lesssim_k \|D\phi\|_{H_b^k} + \|\phi\|_{H_b^k} \end{aligned}$$

by [Corollary 4.6](#). ■

Proof of Theorem 4.13. By [Lemma 4.14](#) and since B is $(k + \frac{1}{2})$ -regular, for every $\phi \in \text{dom}(D_B)$

$$\|\text{res}[\phi]\|_{H^{k+1/2}} \lesssim_{B,k} \|\mathbf{1}_{(-\infty,0)}(A) \text{res}[\phi]\|_{H^{k+1/2}} + \|\text{res}[\phi]\|_{\check{H}} \lesssim_k \|D\phi\|_{H_b^k} + \|\phi\|_{H_b^k}.$$

This together with [Theorem 4.13](#) implies the assertion. ■

4.3 Fredholm extensions in higher regularity

The following is a consequence of [Proposition 3.41](#).

Corollary 4.15 ($\frac{1}{2}$ -regular implies left semi-Fredholm). *Let $B \subset \check{H}\Gamma(Z, \check{S})$ be a residue condition. If B is $\frac{1}{2}$ -regular, then D_B is left semi-Fredholm.* ■

The discussion in [Section 4.2](#) leads to the following observation.

Definition 4.16. Let $k \in \mathbb{N}_0$. Let $B \subset \check{H}\Gamma(Z, \check{S})$ be a residue condition. Consider the closed subspace

$$H_a^{k+1}\Gamma(X \setminus Z, S \otimes \mathbb{I}; B) := \{\phi \in H_a^{k+1}\Gamma(X \setminus Z, S \otimes \mathbb{I}) : \text{res}[\phi] \in B\}. \quad \bullet$$

Proposition 4.17. *Let $k \in \mathbb{N}_0$. Let B be a $(k + \frac{1}{2})$ -regular residue condition. If D_B is Fredholm, then the restriction $D_B: H_a^{k+1}\Gamma(X \setminus Z, S \otimes \mathbb{I}; B) \rightarrow H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I})$ is Fredholm; in fact:*

$$\ker(D_B: H_a^{k+1}\Gamma(X \setminus Z, S \otimes \mathbb{I}; B) \rightarrow H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I})) = \ker D_B$$

and the canonical map

$$\text{coker}(D_B: H_a^{k+1}\Gamma(X \setminus Z, S \otimes \mathbb{I}; B) \rightarrow H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I})) \rightarrow \text{coker } D_B$$

is an isomorphism.

Proof. The proof is identical to the one of [DW24, Theorem 3.57], but repeated here for the readers' convenience. By Theorem 4.13, $\ker(D_B: H_a^{k+1}\Gamma(X \setminus Z, S \otimes \mathbb{I}; B) \rightarrow H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I})) = \ker D_B$; moreover: the linear map

$$\frac{\text{dom}(D_B)}{H_a^{k+1}\Gamma(X \setminus Z, S \otimes \mathbb{I}; B)} \rightarrow \frac{L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})}{H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I})}$$

induced by D_B is injective. Therefore, by the Snake Lemma, the canonical map

$$\text{coker}(D_B: H_a^{k+1}\Gamma(X \setminus Z, S \otimes \mathbb{I}; B) \rightarrow H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I})) \rightarrow \text{coker } D_B$$

is injective.

Since $H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I})$ is dense in $L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})$ the map

$$H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I}) \rightarrow (\ker D_B)^* \cong \text{coker } D_B$$

is surjective. Since it factors through $\text{coker}(D_B: H_a^{k+1}\Gamma(X \setminus Z, S \otimes \mathbb{I}; B) \rightarrow H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I})) \rightarrow \text{coker } D_B$, the latter must be surjective. \blacksquare

4.4 Adapted Sobolev spaces, II: Morrey embedding and polyhomogeneity

The purpose of the upcoming two subsections is to further understand the scale of adapted Sobolev spaces $(H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I}), \|\cdot\|_{H_a^k})_{k \in \mathbb{N}_0}$. A crucial observation is that the singularities in the elements of $H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I})$ can be removed after untwisting in the following sense.

Definition 4.18. Define the **twist** $\bar{z}^{-1/2} \in \Gamma(U \setminus Z, \text{Hom}_{\mathbb{C}}(\Pi^*\check{S}, \mathring{S} \otimes \mathring{\mathbb{I}}))$ by

$$\bar{z}^{-1/2}\phi = r^{-1/2}P_{-1/2}\phi$$

with $P_{-1/2}$ as in Proposition 3.21. \bullet

Lemma 4.19 (Removable singularities after untwisting). *For every $k_0 \in \mathbb{N}$ the restriction map $H_a^{k_0}\Gamma(X \setminus Z, S \otimes \mathbb{I}) \rightarrow H_{\text{loc}}^{k_0}\Gamma(U \setminus Z, \mathring{S} \otimes \mathring{\mathbb{I}})$ factors through*

$$\bar{z}^{-1/2}H^{k_0}\Gamma(U, \Pi^*\check{S}) \subset H_{\text{loc}}^{k_0}\Gamma(U \setminus Z, \mathring{S} \otimes \mathring{\mathbb{I}}).$$

Proof. Let $k \in \mathbf{N}_0$ and $\phi \in H_a^k \Gamma(X \setminus Z, S \otimes \mathbf{I})$. Set

$$\psi := \bar{z}^{1/2} \phi \in H_{\text{loc}}^k \Gamma(U \setminus Z, \Pi^* \check{S}).$$

Since $|\bar{z}^{1/2}|$ is bounded, $\psi \in r^{1/2} L^2 \Gamma(U, \Pi^* \check{S})$. This proves the assertion for $k = 0$.

Henceforth, suppose that $k \in \mathbf{N}$. Consider the differential operators $\mathfrak{d}_z : H_{\text{loc}}^1 \Gamma(U, \Pi^* \check{S}) \rightarrow L^2 \Gamma(U, \Pi^* \check{S})$ and $\mathring{\mathfrak{d}}_z : H_{\text{loc}}^1 \Gamma(U \setminus Z, \check{S} \otimes \check{\mathbf{I}}) \rightarrow L^2 \Gamma(U \setminus Z, \check{S} \otimes \check{\mathbf{I}})$ defined by

$$\mathfrak{d}_z := J(\partial_r - r^{-1} I \nabla_{\partial_\alpha}) \quad \text{and} \quad \mathring{\mathfrak{d}}_z := J(\partial_r - r^{-1} I \mathring{\nabla}_{\partial_\alpha}).$$

If v is the lift of a vector field on Z , then

$$\nabla_v \psi = \bar{z}^{1/2} \mathring{\nabla}_v \phi \quad \text{and} \quad \mathfrak{d}_z \psi = \bar{z}^{1/2} \mathring{\mathfrak{d}}_z \phi$$

on $U \setminus Z$.

A moment's thought shows that $\nabla_v \psi = \bar{z}^{1/2} \mathring{\nabla}_v \phi$ holds on U in the sense of distributions. In fact, $\mathfrak{d}_z \psi = \bar{z}^{1/2} \mathring{\mathfrak{d}}_z \phi$ also holds on U in the sense of distributions. To see this let η_ε be a suitable cut-off function and τ a test section. By direct computation,

$$\int_U \langle \eta_\varepsilon \tau, \mathfrak{d}_z \psi \rangle = \int_U \langle \eta_\varepsilon \mathring{\mathfrak{d}}_z^* \tau, \psi \rangle + \langle \sigma_{\mathfrak{d}_z}(\mathfrak{d}\eta_\varepsilon) \tau, \psi \rangle$$

and

$$\left| \int_U \langle \sigma_{\mathfrak{d}_z}(\mathfrak{d}\eta_\varepsilon) \tau, \psi \rangle \right| \lesssim_\tau \int_U r^{1/2} |\mathfrak{d}\eta_\varepsilon| r^{-1/2} |\psi| \lesssim_\psi \int_U r |\mathfrak{d}\eta_\varepsilon|^2.$$

Since η_ε can be chosen so that $\mathfrak{d}\eta_\varepsilon$ is supported in $B_{2\varepsilon}(Z)$ and $r |\mathfrak{d}\eta_\varepsilon| \lesssim 1$, it follows that

$$\int_U \langle \tau, \mathfrak{d}_z \psi \rangle - \langle \mathring{\mathfrak{d}}_z^* \tau, \psi \rangle = \lim_{\varepsilon \downarrow 0} \int_U \eta_\varepsilon (\langle \tau, \mathfrak{d}_z \psi \rangle - \langle \mathring{\mathfrak{d}}_z^* \tau, \psi \rangle) = 0.$$

By induction, it follows that if $v_1, \dots, v_{k-\ell}$ are lifts of vector fields on Z , then

$$\nabla_{v_1} \dots \nabla_{v_{k-\ell}} \mathfrak{d}_z^\ell \psi = \bar{z}^{1/2} \mathring{\nabla}_{v_1} \dots \mathring{\nabla}_{v_{k-\ell}} \mathring{\mathfrak{d}}_z^\ell \phi$$

holds on U in the sense of distributions; in particular:

$$\nabla_{v_1} \dots \nabla_{v_{k-\ell}} \mathfrak{d}_z^\ell \psi \in r^{1/2} L^2 \Gamma(U, \Pi^* \check{S}).$$

This implies that $\psi \in H^k \Gamma(U, \Pi^* \check{S})$ because

$$\int_U |\mathfrak{d}_z(\chi \psi)|^2 = \int_U |\partial_r(\chi \psi)|^2 + r^{-2} |\nabla_{\partial_\alpha}(\chi \psi)|^2 + O(|\psi|^2). \quad \blacksquare$$

Set

$$H_a^\infty \Gamma(X \setminus Z, S \otimes \mathbf{I}) := \bigcap_{k \in \mathbf{N}_0} H_a^k \Gamma(X \setminus Z, S \otimes \mathbf{I}).$$

Corollary 4.20 (Smooth after untwisting). *The restriction map $H_a^\infty \Gamma(X \setminus Z, S \otimes \mathbb{I}) \rightarrow \Gamma(U \setminus Z, \mathring{S} \otimes \mathring{\mathbb{I}})$ factors through*

$$\bar{z}^{-1/2} \Gamma(U, \Pi^* \mathring{S}) \subset \Gamma(U \setminus Z, \mathring{S} \otimes \mathring{\mathbb{I}}). \quad \blacksquare$$

Remark 4.21 (Polyhomogeneous expansion). Every $\psi \in \Gamma(U, \Pi^* \mathring{S})$ has a Taylor expansion

$$\psi \sim \sum_{k, \ell=0}^{\infty} \bar{z}^k z^\ell \check{\psi}_{k, \ell} \quad \text{with} \quad \check{\psi}_{k, \ell} \in \Pi^* \Gamma(Z, \mathring{S} \otimes_{\mathbb{C}} NZ^{k-\ell})$$

at Z . Here $\bar{z} \in \Gamma(U, \text{Hom}_{\mathbb{C}}(\Pi^* NZ, \mathbb{C}))$ and $z \in \Gamma(U, \text{Hom}_{\mathbb{C}}(\Pi^* NZ^{-1}, \mathbb{C}))$ denote the tautological sections. Therefore, by [Corollary 4.20](#), every $\phi \in H_a^\infty \Gamma(X \setminus Z, S \otimes \mathbb{I})$ has a polyhomogeneous expansion

$$\phi \sim \sum_{k, \ell=0}^{\infty} \bar{z}^{k-1/2} z^\ell \check{\phi}_{k, \ell} \quad \text{with} \quad \check{\phi}_{k, \ell} \in \Pi^* \Gamma(Z, \mathring{S} \otimes_{\mathbb{C}} NZ^{k-\ell})$$

at Z with $\bar{z}^{k-1/2} := \bar{z}^k \bar{z}^{-1/2}$. Moreover, a moment's thought shows that if $D\phi = 0$, then the leading order term is of the form $\bar{z}^{k-1/2} \check{\phi}_{k, 0}$ for some $k \in \mathbb{N}_0$. \clubsuit

Corollary 4.22. *For every $k \in \mathbb{N}$ with $k \geq n/2$*

$$H_a^k \Gamma(X \setminus Z, S \otimes \mathbb{I}) \subset r^{-1/2} L^\infty \Gamma(X \setminus Z, S \otimes \mathbb{I}). \quad \blacksquare$$

The above observation leads to the following ‘‘poor man’s Weyl law’’.

Proposition 4.23 (Growth of eigenvalues). *Let $k \in \mathbb{N}_0$ with $k > n/2$. If $B \subset \mathring{H} \Gamma(X \setminus Z, S \otimes \mathbb{I})$ is a $(k + \frac{1}{2})$ -regular Lagrangian residue condition, then the **counting function** $N: [0, \infty) \rightarrow \mathbb{N}_0$ defined by*

$$N(\Lambda) := \dim E_{\leq \Lambda} \quad \text{with} \quad E_{\leq \Lambda} := \bigoplus_{\lambda \in [-\Lambda, \Lambda]} \ker(D_B - \lambda \cdot \mathbf{1})$$

satisfies

$$N(\Lambda) \lesssim_{B, k} \langle \Lambda \rangle^{2k}.$$

Proof. The following argument is due to Li [[Li80](#), Lemma 11]. Choose an L^2 orthonormal basis $(\phi_1, \dots, \phi_{N(\Lambda)})$ of $E_{\leq \Lambda}$. The density $d \in C^\infty(X \setminus Z, [0, \infty))$ defined by

$$d := \sum_{i=1}^{N(\Lambda)} |\phi_i|^2$$

does not depend on the choice of L^2 orthonormal basis. By construction

$$N(\Lambda) = \frac{1}{\text{vol}(X)} \int_X d \lesssim \|rd\|_{L^\infty}.$$

Choose an $x \in X$ with $\|rd\|_{L^\infty} \leq 2|rd|(x)$. Since $\text{ev}_x: E_{\leq \Lambda} \rightarrow (S \otimes \mathbb{I})_x$ has rank at most $\text{rk } S$, without loss of generality,

$$|rd|(x) = \sum_{i=1}^{\text{rk } S} r|\phi_i|^2(x).$$

By [Theorem 4.13](#) and [Corollary 4.22](#)

$$\|r^{1/2}\phi_i\|_{L^\infty} \lesssim \|\phi_i\|_{H_a^k} \lesssim_{B,k} \|D^k\phi_i\|_{L^2} + \|\phi_i\|_{L^2} \lesssim \langle \Lambda \rangle^k.$$

This implies the assertion. ■

Corollary 4.24. *Let $k \in \mathbf{N}_0$ with $k > n/2$. If $B \subset \check{H}\Gamma(X \setminus Z, S \otimes \mathbb{I})$ is a $(k + \frac{1}{2})$ -regular Lagrangian residue condition, then for every $t \in (0, \infty)$ the heat operator $h_t := \exp(-tD_B^2)$ is trace class. ■*

Remark 4.25. Assume the situation of [Corollary 4.24](#). If ε is a chirality operator, then for every $t > 0$

$$\text{index } D_B^+ = \dim \ker D_B^+ - \dim \ker D_B^- = \text{str}_\varepsilon h_t.$$

Here str_ε denotes the super trace with respect to ε of the heat operator h_t ; cf. [\[BGV92, §1.3\]](#). For suitable choices of B an analysis of the asymptotic behaviour of the kernel attached to h_t as $t \downarrow 0$ should result in index formulae analogous to the one established by Atiyah, Patodi, and Singer [\[APS75, Theorems 3.10 and 4.2\]](#). It would be interesting to work this out in detail. Also, it should be mentioned that part of the unpublished PhD thesis [\[Yano7, Theorems 1.0.3 and 2.3.4\]](#) discusses such index formulae. ♣

4.5 Adapted Sobolev spaces, III: spectral description and tameness

Proposition 4.26 (Spectral description of $H_a^k\Gamma(X \setminus Z, S \otimes \mathbb{I})$). *Let $B \subset \check{H}\Gamma(Z, \check{S})$ be an ∞ -regular Lagrangian residue condition. Let $(\phi_\alpha)_{\alpha \in \mathbf{N}_0}$ be an L^2 orthonormal basis of $L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})$ consisting of eigenspinors for D_B and denote by $(\lambda_\alpha)_{\alpha \in \mathbf{N}_0}$ the corresponding sequence of eigenvalues. For every $\phi \in L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})$*

$$\|\phi\|_{H_a^k}^2 \asymp_{B,k} \sum_{\alpha=0}^{\infty} \langle \lambda_\alpha \rangle^{2k} \langle \phi, \phi_\alpha \rangle^2.$$

Proof. This is an immediate consequence of [Theorem 4.13](#). ■

Proposition 4.27. *The graded Fréchet space $(H_a^\infty\Gamma(X \setminus Z, S \otimes \mathbb{I}), (\|\cdot\|_{H_a^k})_{k \in \mathbf{N}_0})$ is tame.*

Proof. Consider the graded Fréchet space $(\Sigma(L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})), (\|\cdot\|_k)_{k \in \mathbf{N}_0})$ of exponentially decreasing sequences defined by

$$\Sigma(L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})) := \{(\psi_\beta) \in L^2\Gamma(X \setminus Z, S \otimes \mathbb{I})^{\mathbf{N}_0} : \|\psi_\beta\|_k < \infty \text{ for every } k \in \mathbf{N}_0\}$$

with

$$\|(\psi_\beta)\|_k^2 := \sum_{\beta=0}^{\infty} e^{2k\beta} \|\psi_\beta\|_{L^2}^2;$$

cf. Hamilton [\[Ham82, Part II Example 1.1.4\(b\)\]](#) with $q = 2$.

By [Example 3.42](#), there is an ∞ -regular Lagrangian residue condition $B \subset \check{H}\Gamma(X \setminus Z, S \otimes \mathbb{I})$. Borrowing the notation from [Proposition 4.26](#), define $i: H_a^\infty \Gamma(X \setminus Z, S \otimes \mathbb{I}) \rightarrow \Sigma(L^2 \Gamma(X \setminus Z, S \otimes \mathbb{I}))$ and $p: \Sigma(L^2 \Gamma(X \setminus Z, S \otimes \mathbb{I})) \rightarrow H_a^\infty \Gamma(X \setminus Z, S \otimes \mathbb{I})$ by

$$i(\phi)_\beta := \sum_{\alpha=0}^{\infty} \mathbf{1}_{[e^\beta, e^{\beta+1})}(\langle \lambda_\alpha \rangle) \langle \phi, \phi_\alpha \rangle_{L^2} \phi_\alpha \quad \text{and} \quad p((\psi)_\beta) := \sum_{\alpha, \beta=0}^{\infty} \mathbf{1}_{[e^\beta, e^{\beta+1})}(\langle \lambda_\alpha \rangle) \langle \psi_\beta, \phi_\alpha \rangle_{L^2} \phi_\alpha.$$

A moment's thought shows that $p \circ i = \text{id}$; moreover:

$$\|i(\phi)\|_k^2 = \sum_{\beta=0}^{\infty} e^{2k\beta} \|i(\phi)_\beta\|_{L^2}^2 \leq \sum_{\alpha=0}^{\infty} \langle \lambda_\alpha \rangle^{2k} \langle \phi, \phi_\alpha \rangle_{L^2}^2 \lesssim_k \|\phi\|_{H_a^k}$$

and

$$\begin{aligned} \|p(\psi)\|_{H_a^k}^2 &\lesssim_k \sum_{\alpha=0}^{\infty} \langle \lambda_\alpha \rangle^{2k} \langle p(\psi), \phi_\alpha \rangle^2 \\ &= \sum_{\alpha, \beta=0}^{\infty} \langle \lambda_\alpha \rangle^{2k} \mathbf{1}_{[e^\beta, e^{\beta+1})}(\langle \lambda_\alpha \rangle) \langle \psi_\beta, \phi_\alpha \rangle^2 \leq e^{2k} \sum_{\beta=0}^{\infty} e^{2k\beta} \|\psi_\beta\|_{L^2}^2. \end{aligned}$$

Therefore, $(H_a^\infty \Gamma(X \setminus Z, S \otimes \mathbb{I}), (\|-\|_{H_a^k})_{k \in \mathbb{N}_0})$ is a tame direct summand in the sense of [\[Ham82, Part II Definition 1.3.1\]](#) of $(\Sigma(L^2 \Gamma(X \setminus Z, S \otimes \mathbb{I})), (\|-\|_k)_{k \in \mathbb{N}_0})$. This proves that $(H_a^\infty \Gamma(X \setminus Z, S \otimes \mathbb{I}), (\|-\|_{H_a^k})_{k \in \mathbb{N}_0})$ is tame in the sense of [\[Ham82, Part II Definition 1.3.2\]](#). \blacksquare

4.6 Symbolic criterion for ∞ -regularity

Proposition 4.28 (symbolic criterion for ∞ -regularity). *Let $V \subset \check{S}$ be a subbundle. If*

$$J_Y(\xi)V \subseteq V^\perp$$

for every $\xi \in T^*Z \setminus \{0\}$, then B_V is ∞ -regular.

Proof. The proof relies on the following observation which is already implicit in [\[FS98, Lemma 2\]](#). Denote by $\text{pr}_V: \check{S} \rightarrow \check{S}$ the orthogonal projection onto V . The operator $\text{pr}_V \text{Apr}_V$ is a 0-th order differential operator, because for every $f \in C^\infty(Z)$

$$[\text{pr}_V \text{Apr}_V, f] = -\text{pr}_V J_Y(df) \text{pr}_V = 0.$$

Let $k \in \mathbb{N}_0$. Since $\text{pr}_V \text{Apr}_V$ is a 0-th order differential operator, $\text{pr}_V A^{2k+1} \text{pr}_V$ is a $2k$ -th order differential operator. Therefore, for every $\phi \in \Gamma(Z, V)$ and $\varepsilon > 0$

$$\begin{aligned} \langle A^{2k+1} \phi, \phi \rangle_{L^2} &= \langle \text{pr}_V A^{2k+1} \text{pr}_V \phi, \phi \rangle_{L^2} \lesssim_{B_V, k} \|\phi\|_{H^k}^2 \\ &\lesssim \|\phi\|_{H^{k-1/2}} \|\phi\|_{H^{k+1/2}} \lesssim \varepsilon^{-1} \|\phi\|_{H^{k-1/2}}^2 + \varepsilon \|\phi\|_{H^{k+1/2}}^2; \end{aligned}$$

moreover, by direct inspection,

$$\langle A^{2k+1} \phi, \phi \rangle_{L^2} = -\| |A|^{k+1/2} \mathbf{1}_{(-\infty, 0)}(A) \phi \|_{L^2}^2 + \| |A|^{k+1/2} \mathbf{1}_{[0, \infty)}(A) \phi \|_{L^2}^2.$$

As a consequence, for every $\phi \in B_V$,

$$\|\phi\|_{H^{k+1/2}}^2 \lesssim_{B,k} \|\mathbf{1}_{(-\infty,0)}(A)\phi\|_{H^{k+1/2}}^2 + \|\phi\|_{H^{k-1/2}}^2.$$

By induction, B_V is $(k + 1/2)$ -regular for every $k \in \mathbf{N}_0$; hence: ∞ -regular. \blacksquare

Corollary 4.29. *Let $V \subset \check{S}$ be a subbundle. If*

$$J_Y(\xi)V = V^\perp$$

*for every $\xi \in T^*Z \setminus \{0\}$, then B_V and B_V^G are ∞ -regular and Fredholm.* \blacksquare

Example 4.30. The local residue conditions defined in [Example 3.45](#) satisfy the criterion in [Corollary 4.29](#). \spadesuit

Example 4.31. If

$$L \in \text{Hom}_{\mathbb{C}}(\overline{NZ}, \check{S})$$

is nowhere-vanishing, then

$$V := \text{im } L \subset \check{S}$$

is a complex subbundle. Therefore and since $J_Y(\xi)$ and $IJ_Y(\xi)$ are skew-adjoint, $J_Y(\xi)V \subseteq V^\perp$ for every $\xi \in T^*Z \setminus \{0\}$; that is: V satisfies the criterion in [Proposition 4.28](#). Moreover: if $\text{rk}_{\mathbb{C}} \check{S} = 2$, then $J_Y(\xi)V = V^\perp$ and B_V is self-adjoint. \spadesuit

Remark 4.32. Suppose that $(Z, I; \Phi)$ is a harmonic $Z/2Z$ spinor whose branching locus Z satisfies [Hypothesis 3.1](#). Since $\Phi \in \ker D_{\min}$, by [Remark 4.21](#), the polyhomogeneous expansion of Φ at Z is of the form

$$\Phi \sim \bar{z}^{1/2} \check{\Phi}_{1,0} + \sum_{\substack{k,\ell=0 \\ k+\ell \geq 2}}^{\infty} \bar{z}^{k-1/2} z^\ell \check{\Phi}_{k,\ell}.$$

The leading coefficient $\check{\Phi}_{1,0}$ determines an $L_\Phi \in \Gamma(Z, \text{Hom}_{\mathbb{C}}(\overline{NZ}, \check{S}))$. If $\text{rk } S = 4$ and $\check{\Phi}_{1,0}$ is nowhere vanishing, then this produces an ∞ -regular Lagrangian local residue condition. This is the residue condition behind the scenes in [\[Tak15; Par23\]](#). \clubsuit

References

- [AG16] Pierre Albin and Jesse Gell-Redman. *The index of Dirac operators on incomplete edge spaces*. *SIGMA Symmetry Integrability Geom. Methods Appl.* 12 (2016), Paper No. 089, 45. DOI: [10.3842/SIGMA.2016.089](https://doi.org/10.3842/SIGMA.2016.089). \heartsuit (cit. on p. 3)
- [AG23] Pierre Albin and Jesse Gell-Redman. *The index formula for families of Dirac type operators on pseudomanifolds*. *J. Differential Geom.* 125.2 (2023), pp. 207–343. DOI: [10.4310/jdg/1696432923](https://doi.org/10.4310/jdg/1696432923). \heartsuit (cit. on p. 3)
- [APS75] Michael F. Atiyah, V. K. Patodi, and Isadore M. Singer. *Spectral asymmetry and Riemannian geometry. I*. *Mathematical Proceedings of the Cambridge Philosophical Society* 77 (1975), pp. 43–69. MR: [0397797](https://doi.org/10.1017/S0305004100000000). \heartsuit (cit. on pp. 3, 26, 36)

- [BB12] Christian Bär and Werner Ballmann. *Boundary value problems for elliptic differential operators of first order. Surveys in differential geometry.* 17. International Press, 2012, pp. 1–78. DOI: [10.4310/SDG.2012.v17.n1.a1](https://doi.org/10.4310/SDG.2012.v17.n1.a1). arXiv: [1101.1196](https://arxiv.org/abs/1101.1196). MR: [3076058](https://mathscinet.ams.org/mathscinet-getitem?mr=3076058). Zbl: [1331.58022](https://zbmath.org/?q=1331.58022) (cit. on pp. [3](#), [27](#), [30](#))
- [BGV92] Nicole Berline, Ezra Getzler, and Michèle Vergne. *Heat kernels and Dirac operators.* Grundlehren der Mathematischen Wissenschaften 298. Springer-Verlag, 1992. DOI: [10.1007/978-3-642-58088-8](https://doi.org/10.1007/978-3-642-58088-8). MR: [1215720](https://mathscinet.ams.org/mathscinet-getitem?mr=1215720). Zbl: [0744.58001](https://zbmath.org/?q=0744.58001) (cit. on p. [36](#))
- [Bis89] Jean-Michel Bismut. *A local index theorem for non-Kähler manifolds.* English. *Math. Ann.* 284.4 (1989), pp. 681–699. DOI: [10.1007/BF01443359](https://doi.org/10.1007/BF01443359). Zbl: [0666.58042](https://zbmath.org/?q=0666.58042) (cit. on pp. [14](#), [15](#))
- [BF98] Bernhelm Booß-Bavnbek and Kenro Furutani. *The Maslov index: A functional analytical definition and the spectral flow formula.* English. *Tokyo J. Math.* 21.1 (1998), pp. 1–34. DOI: [10.3836/tjm/1270041982](https://doi.org/10.3836/tjm/1270041982). Zbl: [0932.37063](https://zbmath.org/?q=0932.37063) (cit. on pp. [3](#), [4](#), [6](#), [7](#))
- [BS88] Jochen Brüning and Robert Seeley. *An index theorem for first order regular singular operators.* *Amer. J. Math.* 110.4 (1988), pp. 659–714. DOI: [10.2307/2374646](https://doi.org/10.2307/2374646). MR: [955293](https://mathscinet.ams.org/mathscinet-getitem?mr=955293) (cit. on p. [21](#))
- [BS18] Theo Bühler and Dietmar A. Salamon. *Functional analysis.* Vol. 191. Graduate Studies in Mathematics. American Mathematical Society, 2018. DOI: [10.1090/gsm/191](https://doi.org/10.1090/gsm/191). MR: [3823238](https://mathscinet.ams.org/mathscinet-getitem?mr=3823238) (cit. on pp. [3–5](#), [8](#))
- [DW24] Aleksander Doan and Thomas Walpuski. *Chambered invariants of real Cauchy–Riemann operators.* 2024. arXiv: [2410.21057](https://arxiv.org/abs/2410.21057) (cit. on pp. [5](#), [21](#), [33](#))
- [Don17] Simon K. Donaldson. *Adiabatic limits of co-associative Kovalev–Lefschetz fibrations.* *Algebra, Geometry, and Physics in the 21st Century.* Springer, 2017, pp. 1–29. DOI: [10.1007/978-3-319-59939-7_1](https://doi.org/10.1007/978-3-319-59939-7_1). arXiv: [1603.08391](https://arxiv.org/abs/1603.08391). Zbl: [1409.53044](https://zbmath.org/?q=1409.53044) (cit. on p. [2](#))
- [Don21] Simon K. Donaldson. *Deformations of multivalued harmonic functions.* *The Quarterly Journal of Mathematics* 72.1-2 (2021), pp. 199–235. DOI: [10.1093/qmath/haab018](https://doi.org/10.1093/qmath/haab018) (cit. on p. [2](#))
- [FS98] Simone Farinelli and Günter Schwarz. *On the spectrum of the Dirac operator under boundary conditions.* *J. Geom. Phys.* 28.1-2 (1998), pp. 67–84. DOI: [10.1016/S0393-0440\(98\)00013-8](https://doi.org/10.1016/S0393-0440(98)00013-8).  (cit. on p. [37](#))
- [Ham82] Richard S. Hamilton. *The inverse function theorem of Nash and Moser.* *Bull. Amer. Math. Soc. (N.S.)* 7.1 (1982), pp. 65–222. DOI: [10.1090/S0273-0979-1982-15004-2](https://doi.org/10.1090/S0273-0979-1982-15004-2).  (cit. on pp. [3](#), [36](#), [37](#))
- [HMT23] Andriy Haydys, Rafe Mazzeo, and Ryosuke Takahashi. *An index theorem for $\mathbb{Z}/2$ -harmonic spinors branching along a graph.* 2023. arXiv: [2310.15295](https://arxiv.org/abs/2310.15295) (cit. on pp. [4](#), [5](#))
- [He22] Siqi He. *The branched deformations of the special Lagrangian submanifolds.* 2022. DOI: [10.48550/ARXIV.2202.12282](https://doi.org/10.48550/ARXIV.2202.12282). arXiv: [2202.12282](https://arxiv.org/abs/2202.12282) (cit. on p. [2](#))
- [Joh75] Kenneth A. Johnson. *The M.I.T. Bag Model.* *Acta Phys. Polon. B* 6 (1975), p. 865 (cit. on p. [27](#))

- [LM89] H. Blaine Lawson Jr. and Marie-Louise Michelsohn. *Spin geometry*. Princeton Mathematical Series 38. Princeton University Press, 1989. MR: [1031992](#). Zbl: [0688.57001](#) (cit. on p. 2)
- [Li80] Peter Li. *On the Sobolev constant and the p -spectrum of a compact Riemannian manifold*. *Annales Scientifiques de l'École Normale Supérieure* 13 (1980), pp. 451–468. Zbl: [0466.53023](#) (cit. on p. 35)
- [Maz91] Rafe Mazzeo. *Elliptic theory of differential edge operators. I. Communications in Partial Differential Equations* 16.10 (1991), pp. 1615–1664. DOI: [10.1080/03605309108820815](#). Zbl: [0745.58045](#) (cit. on p. 3)
- [MV14] Rafe Mazzeo and Boris Vertman. *Elliptic theory of differential edge operators, II: Boundary value problems*. *Indiana Univ. Math. J.* 63.6 (2014), pp. 1911–1955. DOI: [10.1512/iumj.2014.63.5435](#).  (cit. on p. 3)
- [MS98] Dusa McDuff and Dietmar A. Salamon. *Introduction to symplectic topology*. 2nd edition. Oxford Mathematical Monographs. Oxford University Press, 1998. MR: [1698616](#). Zbl: [1066.53137](#) (cit. on pp. 3, 4)
- [Mel93] Richard B. Melrose. *The Atiyah–Patodi–Singer index theorem*. Vol. 4. Research Notes in Mathematics. A. K. Peters, 1993. DOI: [10.1016/0377-0257\(93\)80040-i](#). MR: [1348401](#). Zbl: [0796.58050](#) (cit. on p. 30)
- [Par23] Gregory J. Parker. *Deformations of \mathbb{Z}_2 -Harmonic Spinors on 3-Manifolds*. 2023. arXiv: [2301.06245](#) (cit. on pp. 2, 3, 38)
- [SW08] Dietmar A. Salamon and Katrin Wehrheim. *Instanton Floer homology with Lagrangian boundary conditions*. *Geometry and Topology* 12.2 (2008), pp. 747–918. DOI: [10.2140/gt.2008.12.747](#). MR: [2403800](#) (cit. on pp. 3, 4, 7, 8)
- [Tak15] Ryosuke Takahashi. *The moduli space of S^1 -type zero loci for $\mathbb{Z}/2$ -harmonic spinors in dimension 3*. 2015. arXiv: [1503.00767](#) (cit. on pp. 2–4, 12, 38)
- [Tak17] Ryosuke Takahashi. *Index theorem for the zero loci of $\mathbb{Z}/2$ -harmonic spinors*. 2017. arXiv: [1705.01954](#) (cit. on p. 2)
- [Tau13a] Clifford H. Taubes. *PSL(2; \mathbb{C}) connections on 3-manifolds with L^2 bounds on curvature*. *Cambridge Journal of Mathematics* 1.2 (2013), pp. 239–397. DOI: [10.4310/CJM.2013.v1.n2.a2](#). arXiv: [1205.0514](#). MR: [3272050](#). Zbl: [1296.53051](#) (cit. on p. 2)
- [Tau13b] Clifford H. Taubes. *Compactness theorems for SL(2; \mathbb{C}) generalizations of the 4-dimensional anti-self dual equations*. 2013. arXiv: [1307.6447](#) (cit. on p. 2)
- [Tau16] Clifford H. Taubes. *On the behavior of sequences of solutions to U(1) Seiberg–Witten systems in dimension 4*. 2016. arXiv: [1610.07163](#) (cit. on p. 2)
- [Tau17] Clifford H. Taubes. *The behavior of sequences of solutions to the Vafa–Witten equations*. 2017. arXiv: [1702.04610](#) (cit. on p. 2)
- [Tau22] Clifford H. Taubes. *Sequences of Nahm pole solutions to the SU(2) Kapustin–Witten equations*. 2022 (cit. on p. 2)

[Yano07] Fangyun Yang. *Dirac operators and monopoles with singularities*. PhD thesis. Massachusetts Institute of Technology, 2007. MR: [2717453](#).  (cit. on p. 36)