## A Linearly Convergent Frank-Wolfe-type Method for Smooth Convex Minimization over the Spectrahedron

Dan Garber

Technion - Israel Institute of Technology dangar@technion.ac.il

#### Abstract

We consider the problem of minimizing a smooth and convex function over the *n*-dimensional spectrahedron — the set of real symmetric  $n \times n$  positive semidefinite matrices with unit trace, which underlies numerous applications in statistics, machine learning and additional domains. Standard first-order methods often require high-rank matrix computations which are prohibitive when the dimension n is large. The well-known Frank-Wolfe method on the other hand, only requires efficient rank-one matrix computations, however suffers from worst-case slow convergence, even under conditions that enable linear convergence rates for standard methods. In this work we present the first Frank-Wolfe-based algorithm that only applies efficient rank-one matrix computations and, assuming quadratic growth and strict complementarity conditions, is guaranteed, after a finite number of iterations, to converges linearly, in expectation, and independently of the ambient dimension.

## 1 Introduction

This paper is concerned with efficient first-order optimization methods (suitable for high dimensional settings) for minimizing a smooth (Lipschitz continuous gradient) and convex objective function  $f : \mathbb{R}^{n \times n} \to \mathbb{R}$  over the *n*-dimensional spectrahedron — the set of all  $n \times n$  real symmetric positive semidefinite matrices with unit trace  $S^n := \{\mathbf{X} \in \mathbb{S}^n \mid \mathbf{X} \succeq 0, \operatorname{Tr}(\mathbf{X}) = 1\}$ , where  $\mathbb{S}^n$  denotes the space of real symmetric  $n \times n$  matrices, and  $\mathbf{X} \succeq 0$  denotes that  $\mathbf{X}$  is positive semidefinite. This problem underlies numerous applications of interest in diverse fields of science and engineering such as statistics, machine learning, and discrete optimization, including for instance convex relaxations to low-rank matrix recovery problems [5, 26, 3, 4, 20, 28], covariance matrix estimation problems [29, 27, 6], convex relaxations to hard combinatorial problems [30, 15, 25], and many more . More specifically, we are interested in methods inspired by the well known Frank-Wolfe method (FW), aka the conditional gradient method [9, 22, 19]. These type of methods, while applicable more broadly to smooth convex minimization over arbitrary convex and compact subsets of an inner product space, are in particular interesting in case the feasible set is the spectrahedron, see for instance [17, 20, 11, 31, 10]<sup>1</sup>, on a number of accounts. First, while orthogonal projections onto the spectrahderon, that are applied in stanadard projection-based gradient methods, require in worst case a full eigen-decomposition of a symmetric  $n \times n$  matrix, which in practical implementations runs in  $O(n^3)$  time, a linear optimization step over the spectrahedron as required by FW, amounts only to an efficient rank-one matrix computation: a single leading eigenvector computation of a symmetric  $n \times n$  matrix, which often runs in nearly linear time in the size of the matrix using fast iterative eigenvector methods. Second, since FW only applies a rank-one update, in many cases of interest both the objective function value and gradient direction could be updated from one iteration to the next much more efficiently than when the update is of high rank. This is for instance the case when the objective f includes terms such as  $q(\mathcal{A}\mathbf{X})$  (e.g., in matrix sensing problems [31]) or  $g(\mathcal{A}\mathbf{X}^{-1})$  (e.g., in covariance estimation problems [6]), where  $\mathcal{A}: \mathbb{R}^{n \times n} \to \mathbb{R}^n$  is a linear map composed of rank-one matrices:  $\mathcal{A}\mathbf{X} = (\mathbf{a}_1^\top \mathbf{X} \mathbf{b}_1, \dots, \mathbf{a}_m^\top \mathbf{X} \mathbf{b}_m)^\top \in \mathbb{R}^m$ with  $\{(\mathbf{a}_i, \mathbf{b}_i)\}_{i=1}^m \subset \mathbb{R}^n \times \mathbb{R}^n$ , and  $g : \mathbb{R}^m \to \mathbb{R}$  is coordinate-wise separable. Another example, is when the objective function includes a  $\log \det(\mathbf{X})$  component. Third, another consequence of the low-rank updates of FW is that when initialized with a low-rank matrix, and provided that the number of iterations is substantially lower than the dimension n, the iterates produced by the method are low-rank and can be stored in factorized form to further reduce memory and runtime requirements. Finally, FW-based methods are often parameter-free and do not require tuning of any parameters (such as the smoothness or strong convexity parameters), and can often be implemented using simple and efficient line-search.

The major drawback of Frank-Wolfe however is worst-case slow convergence rates. For minimizing a  $\beta$ -smooth convex objective over the spectrahedron, the worst-case number of iterations to guarantee an  $\epsilon$ -approximated solution in function value is  $O(\beta/\epsilon)$  [17, 20, 19]. This does not improve even under standard curvature assumptions such quadratic growth (see for instance lower bound in [18]), which are well known to facilitate linear convergence rates (i.e., convergence rates that scale only with log(1/ $\epsilon$ )) for standard projection-based methods [8, 24].

In recent years, several Frank-Wolfe-inspired methods for optimization over the spectrahedron have been developed that, under conditions such as quadratic growth and/or strict complementarity, enjoy a linear convergence rate. [14] established that in case there exists a rank-one optimal solution which satisfies the strict complementarity condition, the standard FW with line-search, after a finite number of iteration, begins to converge linearly. This result however does not extend to the general case in which there is an optimal solution with rank greater than one. The works [1, 7] proposed methods, which we refer to as block-Frank-Wolfe methods, that converge linearly without restriction on the rank of optimal solutions, however these diverge significantly from the classical FW template: while standard FW only applies rank-one leading eigenvector computations, [1, 7] require on each iteration to compute the r leading components in the singular value decomposition (SVD) of a  $n \times n$  matrix, where the rank parameter r is required to be at least as large as

<sup>&</sup>lt;sup>1</sup>some papers, e.g., [20, 10], consider the closely related problem of optimization over the matrix nuclear norm ball, see [20] for a discussion of the connection between these setups

the highest rank of any optimal solution. As a result, these methods have several inherent and substantial limitations. First, they require knowledge of the rank of optimal solutions which is often unknown and overestimating this quantity results in needlessly expensive SVD computations. Second, these methods are only efficient when optimal solutions are of low-rank, since otherwise the partial SVD computations required are not significantly more efficient than those in projection-based methods. Similarly, in this case also the important FW feature of fast update times of objective function and gradient information is lost. Finally, while FW can often be implemented with simple line-search, the method in [1], aside of the smoothness constant  $\beta$ , requires the quadratic growth constant  $\alpha$  in order to set the step-size, which is often very difficult to estimate. The method in [7] does not require these constants and instead performs a sophisticated multi-variate "step-size" optimization on each iteration, by minimizing the original objective function f over an r-dimensional spectrahedron (where r is the upper-bound on the rank of optimal solutions), for instance by running a nested first-order method. However, this is again may only be efficient for small values of r.

We mention in passing that aside of linear convergence results, the work [11] presented a FW variant for the spectrahedron that also uses only rank-one matrix computations (leading eigenvector) and, under strong convexity of the objective function, is guaranteed to converge with a fast rate of the form  $\min\{\frac{\operatorname{rank}(\mathbf{X}^*)^{2/3}}{t^{4/3}}, \frac{1}{\lambda_{\min}(\mathbf{X}^*)^{2}t^2}\}$ , where t is the iteration counter and  $\lambda_{\min}(\mathbf{X}^*)$  denotes the smallest non-zero eigenvalue of the unique optimal solution  $\mathbf{X}^*$ . While this result does not limit the rank of the optimal solution and clearly improves upon the standard FW rate of O(1/t), it is not a linear convergence rate.

The dichotomy in linear convergence rates between the case of a rank-one optimal solution that can be solved via the standard FW method using only efficient rank-one matrix computations (under strict complementarity condition), and the case of a higher rank optimal solution which currently can only be handled via block-Frank-Wolfe methods that require higher rank matrix computations (with all of the accompanying limitations discussed above) leads us to the following conceptual question:

# Are SVD computations of rank greater than one mandatory (in Frank-Wolfe-based methods) in order to guarantee a linear convergence rate (without restricting the rank of optimal solutions)?

We answer this question in the negative, under the assumption that quadratic growth and strict complementarity conditions hold (see formal definitions in Section 1.2). We propose a novel Frank-Wolfe-based method that is guaranteed, after a finite number of iterations ("burn-in" phase), to converge linearly, in expectation (the expectation is due to the use of randomization in one of the steps of the algorithm). In particular, both the finite number of initial steps and the linear convergence rate are independent of the ambient dimension. Importantly, our method admits an implementation, which aside of three efficient leading eigenvector computations (the same type of rank-one computations applied in the standard FW method and can be executed in parallel), requires only  $O(n^2)$  runtime per iteration. In terms of parameter tuning, our method requires only knowledge of the smoothness constant

Algorithm	SVD	req. parameters	burn-in	convergence rate (after
	rank		phase?	burn-in phase)
Frank-Wolfe [19]	1	-	NO	1/t
Regularized-FW [11]	1	β	NO	$1/t^2$ (in expectation)
Block-FW [1]	r	$r \ge \operatorname{rank}(\mathbf{X}^*), \alpha/\beta$	NO	linear
Spectral-FW [7]	r	$r \ge \operatorname{rank}(\mathbf{X}^*)$	YES	linear
This paper	1	β	YES	linear (in expectation)

Table 1: Comparison of Frank-Wolfe-based methods for the spectrahedron.

of the objective function  $\beta$  (which as opposed to the rank of optimal solutions or the quadratic growth constant, is often relatively easy to bound).

Before moving on, we pause to discuss the strict complementarity assumption in some more detail. While such a condition does not typically appear in linear convergence results for projection-based methods, it was in fact highly instrumental to obtaining linear convergence rates for FW-based methods in a broader context: It was introduced in the classical text [22] for optimization over strongly convex sets, in [16, 12] for obtaining dimension-independent linear rates for polytopes, and in [14, 7] for the spectrahedron setting. Moreover, [7] proved that in our spectrahedron setting with an objective function which admits the highly popular structure  $f(\mathbf{X}) = g(\mathcal{A}\mathbf{X})$ , where  $\mathcal{A}$  is a linear map and g is smooth and strongly convex, strict complementarity further implies that f satisfies quadratic growth over the spectrahedron. [32] further provided numerical evidence for a closely related setting (though not completely identical to ours), that for an objective function with the above structure, linear convergence may be unattainable when strict complementarity does not hold.

#### 1.1 Inspiration from the polytope setting

Aside from our matrix spectrahedron setting, the Frank-Wolfe method has gained significant interest in the past decades also for optimization over convex and compact polytopes. While in this setting also the worst-case convergence of the standard method (with line-search or without) scales with  $1/\epsilon$ , even under favorable conditions such as quadratic growth or strong convexity of the objective function, it was established in a series of works that simple modifications of the method, all of which rely in some way or the other on the concept of incorporating *away steps* into the algorithm, linear convergence can be established, and in particular that a linear rate independent of the ambient dimension could be established when strict complementarity also holds. In the celebrated work [16], which was the first to obtain a linear convergence result for polytopes (under strong convexity and strict complementarity), the authors proved that after a finite number of steps, all iterates of the method must lie inside the optimal face of the polytope, and there it acts as if performing unconstrained minimization and thus, once inside the optimal face, the strong convexity assumption implies linear convergence.

Our method and analysis are highly inspired by [16], but with some considerable differences. The main difficulty is that while polytopes have a finite number of

faces which indeed allows to argue that under strict complementarity, from some iteration, all iterates will lie inside the optimal face, such an argument is not sensible for the spectraehedron which has infinitely many faces<sup>2</sup>. To bypass this difficulty, we apply our arguments w.r.t. to a time-changing face of the spectrahedron, one that is defined according to a principle subspace of the current gradient direction. Once close enough to an optimal solution (which takes a finite number of steps), we can distinguish between two main cases: if our current iterate is sufficiently aligned with this face, we can apply similar arguments to those in [16] (i.e., essentially unconstrained minimization) and argue that either a standard Frank-Wolfe step or a specialized away-step (designed specifically for our spectrahedron setting) will reduce the approximation error by a constant fraction. Otherwise (in case the current iterate is not sufficiently aligned with the current face), we establish that a specialized randomized pairwise step, i.e., a step that replaces a random rankone component supported by the current iterate with a new rank-one component, reduces the error by a constant fraction, in expectation. Thus, an overall linear convergence rate (in expectation) is obtained.

#### **1.2** Notation and Assumptions

We denote matrices by uppercase boldface letters, e.g.,  $\mathbf{X}$ , column vectors by lowercase lightface letters, e.g.,  $\mathbf{v}$ , and scalars are denoted by lightface letters, e.g.,  $r, \alpha, C$ . For matrices we let  $\|\cdot\|$  denote the spectral norm (largest singular value) and for column vectors we let it denote the Euclidean norm. We let  $\|\cdot\|_F$  denote the Frobenius (Euclidean) norm for matrices. For a matrix  $\mathbf{A} \in \mathbb{S}^n$  we let  $\lambda_1(\mathbf{A}), \ldots, \lambda_n(\mathbf{A})$  denote its (real) eigenvalues in non-ascending order. We denote by  $\mathrm{Im}(\mathbf{A})$  and  $\mathrm{Ker}(\mathbf{A})$ its image and kernel, respectively, and by  $\mathbf{A}^{\dagger}$  its Moore-Penrose pseudo inverse. We let  $\mathbf{I}$  denote the identity matrix whose dimension will be clear from context. We let  $\mathbb{S}^n_+$  denote the positive semidefinite cone in  $\mathbb{S}^n$ . We let  $\langle \mathbf{A}, \mathbf{B} \rangle = \mathrm{Tr}(\mathbf{A}^\top \mathbf{B})$  denote the standard matrix inner-product.

We let  $\mathcal{X}^* \subseteq \mathcal{S}^n$  denote the set of minimizers of the objective function f over the spectrahedron  $\mathcal{S}^n$ , and we let  $f^*$  denote the minimal value of f over  $\mathcal{S}^n$ .

Recall we assume that f is convex, and we also assume it is  $\beta$ -smooth over  $S^n$ , i.e., there exists  $\beta \geq 0$  such that for any  $\mathbf{X}, \mathbf{Y} \in S^n$  we have that  $\|\nabla f(\mathbf{X}) - \nabla f(\mathbf{Y})\|_F \leq \beta \|\mathbf{X} - \mathbf{Y}\|_F$ . Recall this further implies the well known inequality (see for instance Lemma 5.7 in [2]):

$$\forall (\mathbf{X}, \mathbf{Y}) \in \mathcal{S}^n \times \mathcal{S}^n : \quad f(\mathbf{Y}) \le f(\mathbf{X}) + \langle \mathbf{Y} - \mathbf{X}, \nabla f(\mathbf{X}) \rangle + \frac{\beta}{2} \|\mathbf{Y} - \mathbf{X}\|_F^2,$$

which we will use extensively.

We now formally present our two main assumptions that will be assumed to hold throughout this work.

Quadratic growth is a standard assumption in the literature on linear convergence rates for first-order methods [8, 24].

<sup>&</sup>lt;sup>2</sup>a face of  $S^n$  is given by the set  $\{\mathbf{VSV}^\top \mid \mathbf{S} \in S^r\}$  for some integer  $r \in \{1, \ldots, n\}$  and matrix  $\mathbf{V} \in \mathbb{R}^{n \times r}$  such that  $\mathbf{V}^\top \mathbf{V} = \mathbf{I}$ 

Assumption 1 (quadratic growth). There exists a scalar  $\alpha > 0$  such that for any  $\mathbf{X} \in S^n$ ,

$$\min_{\mathbf{Y}\in\mathcal{X}^*} \|\mathbf{X} - \mathbf{Y}\|_F^2 \le \frac{2}{\alpha} \left( f(\mathbf{X}) - f^* \right).$$
(1)

Our second assumption is strict complementarity, which as discussed in the Introduction, has played a central role in proving linear convergence rates for Frank-Wolfe-type methods in diverse settings. In case of the spectrahedron it takes the form of a positive eigengap in the gradient direction at optimal points (see also [7, 14, 13]).

Assumption 2 (strict complementarity). There exists an integer  $r^* \in \{1, ..., n\}$ such that for all  $\mathbf{X}^* \in \mathcal{X}^*$  it holds that  $rank(\mathbf{X}^*) = r^*$ , and if  $r^* < n$ , then there exists a scalar  $\delta > 0$  such that

$$\min_{\mathbf{X}^* \in \mathcal{X}^*} \left( \lambda_{n-r^*} (\nabla f(\mathbf{X}^*)) - \lambda_{n-r^*+1} (\nabla f(\mathbf{X}^*)) \right) = \delta.$$
(2)

Following Assumption 2 we denote the minimal value of the smallest non-zero eigenvalue over the optimal set:

$$\lambda_{r^*} := \min_{\mathbf{X}^* \in \mathcal{X}^*} \lambda_{r^*}(\mathbf{X}^*), \tag{3}$$

which is guaranteed to be strictly positive.

Note that Assumption 2 implies that there exists  $\mathbf{V}^* \in \mathbb{R}^{n \times r^*}$  such that  $\mathbf{V}^{*\top}\mathbf{V}^* = \mathbf{I}$  and  $\mathcal{X}^* \subseteq {\mathbf{V}^*\mathbf{S}\mathbf{V}^{*\top} \mid \mathbf{S} \in \mathcal{S}^{r^*}}$ , that is,  $\mathcal{X}^*$  lies on a  $r^*$ -dimensional face of  $\mathcal{S}^n$ .

#### 1.3 Organization of this paper

Section 2 presents our algorithm and its efficient implementation. Section 3 then formally presents and proves the convergence rates of our proposed algorithm. Section 4 presents some numerical experiments. Finally, Section 5 outlines some open questions for future research.

## 2 The Algorithm

Our algorithm is given below as Algorithm 1. As discussed in Section 1.1, our algorithm is inspired by the Away-steps Frank-Wolfe algorithm for optimization over polytopes [16, 21, 12]. Our algorithm applies three different type of steps: I. Standard Frank-Wolfe steps, II. Away/Drop steps, and III. Pairwise steps, all of which are detailed below. In Section 2.1 we discuss in greater detail the efficient implementation of these steps. In particular we show how each step corresponds to a simple leading eigenvector computation. We also describe an implementation that, aside from these eigenvector computations (which can be computed in parallel), only requires  $O(n^2)$  time per iteration.

**Frank-Wolfe steps:** These are based on moving from the current feasible point towards an extreme point of the feasible set that minimizes the inner-product with the current gradient direction. In case of the spectrahedron, these correspond to a rank-one update of the form  $\mathbf{X}_{t+1} \leftarrow (1 - \eta_t)\mathbf{X}_t + \eta_t \mathbf{v}_{t,+} \mathbf{v}_{t,+}^{\mathsf{T}}$ , where  $\mathbf{v}_{t,+}$  is a unit-length eigenvector corresponding to the leading eigenvalue of  $-\nabla f(\mathbf{X}_t)$  (see also [17, 20, 19]). We set the step size  $\eta_t \in [0, 1]$  using line-search.

Away and Drop steps: Away steps perform rank-one updates in which the weight of some rank-one component supported by the current iterate, is lowered in favor of other components. This takes the from:  $\mathbf{X}_{t+1} \leftarrow \frac{1}{1-\eta_t} \left( \mathbf{X}_t - \eta_t \mathbf{v}_{t,-} \mathbf{v}_{t,-}^{\top} \right)$ , where  $\mathbf{v}_{t,-}$  is a unit vector in  $\operatorname{Im}(\mathbf{X}_t)$  that maximizes the quadratic product with the current gradient direction. The step size  $\eta_t$  is set via line search and satisfies  $\eta_t \in [0, (\mathbf{v}_{t,-}^{\top} \mathbf{X}_t^{\dagger} \mathbf{x}_{t,-})^{-1}]$  in order to maintain the feasibility of  $\mathbf{X}_{t+1}$  w.r.t.  $S^n$  (see Lemma 2 in the sequel). In particular, when  $\eta_t = (\mathbf{v}_{t,-}^{\top} \mathbf{X}_t^{\dagger} \mathbf{x}_{t,-})^{-1}$  (i.e., the maximal step size allowed), we have that  $\operatorname{rank}(\mathbf{X}_{t+1}) < \operatorname{rank}(\mathbf{X}_t)$ , and then we refer to this as a *drop step*. It should be mentioned that this type of away steps closely resembles the ones proposed in [10] for optimization over the matrix nuclear norm ball.

**Pairwise steps:** Pairwise steps correspond to rank-two updates, in which a rankone component in the support of the current iterate is replaced with a new rank-one component. The rank-one component to be removed is chosen uniformly at random, and the new rank-one component is chosen according to a *proximal gradient style* rule. Formally, the update takes the form:  $\mathbf{X}_{t+1} \leftarrow \mathbf{X}_t + \gamma_t(\mathbf{u}_{t,+}\mathbf{u}_{t,+}^{\top} - \mathbf{u}_{t,-}\mathbf{u}_{t,-}^{\top})$ , where  $\mathbf{u}_{t,-}$  (the component to be removed) is chosen uniformly at random from the unit sphere in Im( $\mathbf{X}_t$ ), the step-size is set to  $\gamma_t = (\mathbf{u}_{t,-}^{\top}\mathbf{X}_t^{\dagger}\mathbf{u}_{t,-})^{-1}$  (so that feasibility of  $\mathbf{X}_{t+1}$  is guaranteed, see Lemma 2), and  $\mathbf{u}_{t,+}$  is a unit vector chosen according to the rule:

$$\mathbf{u}_{t,+} \leftarrow \operatorname{argmin}_{\|\mathbf{u}\|=1} \mathbf{u}^{\top} \nabla f(\mathbf{X}_t) \mathbf{u} + \frac{\beta \gamma_t}{2} \|\mathbf{u}\mathbf{u}^{\top} - \mathbf{u}_{t,-}\mathbf{u}_{t,-}^{\top}\|_F^2.$$
(4)

Note that since  $\mathbf{u}$  in the RHS is constrained to have unit norm, the above simplifies to

$$\mathbf{u}_{t,+} \leftarrow \operatorname{argmin}_{\|\mathbf{u}\|=1} \mathbf{u}^{\top} \left( \nabla f(\mathbf{X}_t) - \beta \gamma_t \mathbf{u}_{t,-} \mathbf{u}_{t,-}^{\top} \right) \mathbf{u},$$

which in turn implies that  $\mathbf{u}_{t,+}$  is simply a unit-length leading eigenvector of the matrix  $\beta \gamma_t \mathbf{u}_{t,-} \mathbf{u}_{t,-}^{\top} - \nabla f(\mathbf{X}_t)$ . Note that as opposed to previous steps, this step involves both the knowledge of the smoothness parameter  $\beta$  and the use of randomization. Note furthermore that in this step we always take the maximal step-size  $\gamma_t$  that is guaranteed to maintain feasibility (see Lemma 2). In some sense, one can think about the computation of  $\mathbf{u}_{t,+}$  in Eq. (4) as already incorporating the choice of step-size within the choice of  $\mathbf{u}_{t,+}$  (the interested reader can formally see this in the proof of Lemma 4, which analyzes the benefit of applying these steps).

**Choice of step:** On each iteration, our algorithm first attempts to perform a drop step (i.e., an away step with maximal step-size) in order to quickly adapt the

rank of the iterates to that of the optimal face. If this results in a non-increasing objective value, then the algorithm continues to the next iteration. Otherwise, it computes all three possibilities (can be done in parallel): Frank-Wolfe step, Away step, and a Pairwise step. The algorithm then follows the step which reduces the objective value the most.

#### Algorithm 1 Spectrahedron Frank-Wolfe with Away and Pairwise steps

$$\begin{split} \mathbf{X}_{1} \leftarrow \text{arbitrary matrix in } \mathcal{S}^{n} \\ \text{for } t = 1, 2, \dots \text{ do} \\ \mathbf{v}_{t,-} \leftarrow \operatorname{argmax}_{\mathbf{v} \in \operatorname{Im}(\mathbf{X}_{t}): \|\mathbf{v}\| = 1} \mathbf{v}^{\top} \nabla f(\mathbf{X}_{t}) \mathbf{v} \\ \lambda_{t} \leftarrow \left(\mathbf{v}_{t,-}^{\top} \mathbf{X}_{t}^{\dagger} \mathbf{v}_{t,-}\right)^{-1} \\ \text{if } \lambda_{t} < 1 \text{ then} \\ \mathbf{X}_{t+1}^{drop} \leftarrow \frac{1}{1-\lambda_{t}} \left(\mathbf{X}_{t} - \lambda_{t} \mathbf{v}_{t,-} \mathbf{v}_{t,-}^{\top}\right) \\ \text{if } f(\mathbf{X}_{t+1}^{drop}) \leq f(\mathbf{X}_{t}) \text{ then} \\ \mathbf{X}_{t+1} \leftarrow \mathbf{X}_{t+1}^{drop} \left\{ \text{take a drop step} \right\} \\ \text{end if} \\ \text{end if} \\ \text{if drop step was not taken then} \\ \mathbf{v}_{t,+} \leftarrow \text{ leading eigenvector } of -\nabla f(\mathbf{X}_{t}) \\ \eta_{t}^{FW} \leftarrow \operatorname{argmin}_{\eta \in [0,1]} f((1-\eta)\mathbf{X}_{t} + \eta \mathbf{v}_{t,+} \mathbf{v}_{t,+}^{\top}) \\ \mathbf{X}_{t+1}^{FW} \leftarrow (1-\eta_{t}^{FW})\mathbf{X}_{t} + \eta_{t}^{FW}\mathbf{v}_{t,+} \mathbf{v}_{t,+}^{T} \\ \eta_{t}^{AFW} \leftarrow \operatorname{argmin}_{\eta \in [0,\lambda_{t}]} f\left(\frac{1}{1-\eta} \left(\mathbf{X}_{t} - \eta \mathbf{v}_{t,-} \mathbf{v}_{t,-}^{T}\right)\right) \\ \mathbf{X}_{t+1}^{AFW} \leftarrow \left(\frac{1}{1-\eta_{t}^{AFW}} \left(\mathbf{X}_{t} - \eta_{t}^{AFW}\mathbf{v}_{t,-} \mathbf{v}_{t,-}^{T}\right) \\ \mathbf{u}_{t,-} \leftarrow \text{ uniformly dist. random vector in } \left\{\mathbf{z} \in \operatorname{Im}(\mathbf{X}_{t}) \mid \|\mathbf{z}\| = 1\right\} \\ \gamma_{t} \leftarrow \left(\mathbf{u}_{t,-}^{T}\mathbf{X}_{t}^{\dagger}\mathbf{u}_{t,-}\right)^{-1} \\ \mathbf{u}_{t,+} \leftarrow \text{ leading eigenvector } of \beta\gamma_{t}\mathbf{u}_{t,-}\mathbf{u}_{t,-}^{T} - \nabla f(\mathbf{X}_{t}) \\ \mathbf{X}_{t+1}^{PFW} \leftarrow \mathbf{X}_{t} + \gamma_{t} \left(\mathbf{u}_{t,+}\mathbf{u}_{t,+}^{T} - \mathbf{u}_{t,-}\mathbf{u}_{t,-}^{T}\right) \\ \mathbf{X}_{t+1} \leftarrow \operatorname{argmin}_{\mathbf{X} \in \left\{X_{t+1}^{FW}, X_{t+1}^{AFW}, X_{t+1}^{FW}\right\}} f(\mathbf{X}) \\ \text{end if} \end{aligned}$$

**Lemma 1** (feasibility of Algorithm 1). The sequence  $(\mathbf{X}_t)_{t\geq 1}$  produced by Algorithm 1 is always feasible w.r.t.  $S^n$ .

The correctness of this lemma could be easily verified using the following lemma, whose proof is given in the appendix.

**Lemma 2.** Let  $\mathbf{X} \in \mathbb{S}^n_+$  and  $\mathbf{v} \in Im(\mathbf{X})$ . Consider the matrix  $\mathbf{Y} = \mathbf{X} - \lambda \mathbf{v} \mathbf{v}^\top$  for some  $\lambda \ge 0$ . If  $\lambda \le (\mathbf{v}^\top \mathbf{X}^\dagger \mathbf{v})^{-1}$ , then  $\mathbf{Y} \succeq 0$ . Moreover, if  $\lambda = (\mathbf{v}^\top \mathbf{X}^\dagger \mathbf{v})^{-1}$ , then rank $(\mathbf{Y}) < rank(\mathbf{X})$  and in particular  $\mathbf{X}^\dagger \mathbf{v} \in Ker(\mathbf{Y})$ .

#### 2.1 Implementation details

We now discuss in more detail the efficient implementation of Algorithm 1. Let us fix some iteration t of the algorithm and denote by  $\Pi_t \in \mathbb{S}^n_+$  the projection matrix onto the subspace  $\text{Im}(\mathbf{X}_t)$ . Note that  $\Pi_t = \mathbf{X}_t \mathbf{X}_t^{\dagger}$ . Let us assume for now that both matrices  $\mathbf{X}_t^{\dagger}, \Pi_t$  are explicitly given (we shall discuss their efficient maintenance throughout the run of the algorithm in the sequel).

Computing the vector  $\mathbf{v}_{t,-}$ :  $\mathbf{v}_{t,-}$  could be computed by a standard leading eigenvector computation w.r.t. the matrix  $\mathbf{M} = \Pi_t \nabla f(\mathbf{X}_t) \Pi_t + (1+\zeta) \|\nabla f(\mathbf{X}_t)\| \Pi_t$ , for some arbitrary positive constant  $\zeta > 0$ . Observe that by construction,  $\mathbf{M}$  is positive semidefinite and its leading eigenvectors, which correspond to a strictly positive leading eigenvalue, indeed must lie in  $\mathrm{Im}(\mathbf{X}_t)$ . In particular,  $\mathbf{M}$  need not be computed explicitly: standard fast iterative leading eigenvector methods (such as Lanczos-type algorithms) rely only on computing matrix-vector products w.r.t. to the matrix  $\mathbf{M}$ , and thus, given  $\Pi_t$  (or  $\mathbf{X}_t^{\dagger}$ ), these could be implemented efficiently in  $O(n^2)$  time per such matrix-vector product without explicitly computing  $\mathbf{M}$ .

Computing the vector  $\mathbf{u}_{t,-}$ :  $\mathbf{u}_{t,-}$  is a random vector distributed uniformly over the unit sphere in the space  $\operatorname{Im}(\mathbf{X}_t)$ . Using the rotation invariance of the multivariate standard Gaussain distribution  $\mathcal{N}(\mathbf{0}, \mathbf{I}_n)$ , we have that given a random Gaussian vector  $\mathbf{z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$ , the vector  $\mathbf{u}_{t,-} = \frac{\prod_t \mathbf{z}}{\|\prod_t \mathbf{z}\|}$  is distributed uniformly over the unit sphere in  $\operatorname{Im}(\mathbf{X}_t)$ .

Computing the vector  $\mathbf{u}_{t,+}$ : As already explained above (see discussion following Eq. (4)),  $\mathbf{u}_{t,+}$  can be computed using a standard leading eigenvector computation w.r.t. the matrix  $\beta \gamma_t \mathbf{u}_{t,-} \mathbf{u}_{t,-}^\top - \nabla f(\mathbf{X}_t)$ .

All computations discussed above require maintaining the projection matrix  $\Pi_t$ . Additionally, the computations of the scalars  $\lambda_t$ ,  $\gamma_t$  in Algorithm 1 require also access to the pseudo inverse matrix  $\mathbf{X}_t^{\dagger}$ . We now discuss how it is possible to efficiently update either  $\Pi_t$  of  $\mathbf{X}_t^{\dagger}$  (or both) throughout the run of the algorithm.

Maintaining only the pseudo inverse matrix  $\mathbf{X}_{t}^{\dagger}$ : Similarly to the well known Sherman-Morrison-Woodbury formula for the fast update of the matrix inverse after a rank-one update, [23] established a formula for updating the pseudo inverse matrix after a rank-one update. While the update in case of the pseudo inverse is substantially more involved, the computational complexity is similar. In particular, on each iteration t of Algorithm 1, given  $\mathbf{X}_{t}^{\dagger}$  and the quantities  $\mathbf{v}_{t,-}$ ,  $\lambda_{t}$ ,  $\mathbf{v}_{t,+}$ ,  $\mathbf{u}_{t,-}$ ,  $\mathbf{u}_{t,+}$ ,  $\eta_{t}^{\text{FW}}$ ,  $\eta_{t}^{\text{AFW}}$ ,  $\eta_{t}^{\text{PFW}}$ ,  $\gamma_{t}$ , the pseudo inverse matrix of the next iterate,  $\mathbf{X}_{t+1}^{\dagger}$ , can be computed in  $O(n^{2})$  time <sup>3</sup>. For more concrete details see Lemma 9 in the appendix. Since we have that  $\Pi_{t+1} = \mathbf{X}_{t+1}\mathbf{X}_{t+1}^{\dagger}$ , it is not required to also maintain  $\Pi_{t+1}$  explicitly. This leads to the following theorem.

**Theorem 1.** Algorithm 1 admits an implementation such that on each iteration t, aside of the computation of the eigenvectors  $\mathbf{v}_{t,-}, \mathbf{v}_{t,+}, \mathbf{u}_{t,+}$ , all other computations can be carried out in  $O(n^2)$  time.

 $<sup>^{3}</sup>$ while [23] considers the general case of complex non-square matrices which leads to 6 different types of updates for the pseudo inverse, here since we consider symmetric matrices, only 3 of the 6 possibilities need to be considered

Maintaining only the projection matrix  $\Pi_t$ : A different possibility is not to maintain the pseudo inverse  $\mathbf{X}_t^{\mathsf{T}}$  explicitly, but only the projection matrix  $\Pi_t$ . Let us consider the four possible cases. I. In case a drop step is performed, then according to Lemma 2 we have that the unit vector  $\mathbf{z} = \mathbf{X}_t^{\dagger} \mathbf{v}_{t,-}$  satisfies  $\mathbf{z} \in \text{Im}(\mathbf{X}_t)$  and  $\mathbf{z} \in \operatorname{Ker}(\mathbf{X}_{t+1})$ . This implies that  $\Pi_{t+1} = \Pi_t - \frac{1}{\|\mathbf{z}\|^2} \mathbf{z} \mathbf{z}^{\top}$ . The vector  $\mathbf{z}$  could be approximated to arbitrary accuracy by solving the simple least squares problem:  $\min_{\mathbf{v}\in\mathbb{R}^n} \|\mathbf{X}_t\mathbf{y}-\mathbf{v}_{t,-}\|^2$ . II. In case a standard Frank-Wolfe step is preformed and  $\eta_t^{\text{FW}} < 1$ , defining  $\mathbf{z} := \mathbf{v}_{t,+} - \Pi_t \mathbf{v}_{t,+}$ , we have that  $\Pi_{t+1} = \Pi_t + \frac{1}{\|\mathbf{z}\|^2} \mathbf{z} \mathbf{z}^{\top}$ . In case  $\eta_t^{\text{FW}} = 1$ , we set  $\Pi_{t+1} = \mathbf{v}_{t,+} \mathbf{v}_{t,+}^{\top}$ . III. In case an away step is taken (which is not a drop step), we have that  $Im(\mathbf{X}_{t+1}) = Im(\mathbf{X}_t)$ , and so  $\Pi_{t+1} = \Pi_t$ . IV. If a pairwise step is taken, then we update  $\Pi_t$  using two steps: a drop step (as in step I above) w.r.t. the direction  $\mathbf{u}_{t,-}$ , and then adding the direction  $\mathbf{u}_{t,+}$  (as in step II above). That is, denoting  $\mathbf{w} := \mathbf{X}_t^{\dagger} \mathbf{u}_{t,-}, \ \Pi_{t+1/2} := \Pi_t - \frac{1}{\|\mathbf{w}\|^2} \mathbf{w} \mathbf{w}^{\top}$ , and then denoting  $\mathbf{z} := \mathbf{u}_{t,+} - \Pi_{t+1/2} \mathbf{u}_{t,+}$ , we have that  $\Pi_{t+1} = \Pi_{t+1/2} + \frac{1}{\|\mathbf{z}\|^2} \mathbf{z} \mathbf{z}^{\top}$ . Finally, as with the computation of the product  $\mathbf{X}_t^{\dagger} \mathbf{v}_{t,-}$  in step I, computing the scalars  $\lambda_t, \gamma_t$ , which requires to compute a matrix-vector product with the pseudo inverse matrix  $\mathbf{X}_{t}^{\dagger}$ , can be done by solving a simple least squares problem.

## **3** Convergence Rate Analysis

We turn to formally state and prove our main result — the linear convergence rate of Algorithm 1. Throughout this section we let  $h_t := f(\mathbf{X}_t) - f^*$  denote the approximation error of Algorithm 1 on any iteration  $t \ge 1$ .

**Theorem 2** (convergence of Algorithm 1). The sequence of iterates  $(\mathbf{X}_t)_{t\geq 1}$  produced by Algorithm 1 satisfies:

$$\forall t \ge 1: \qquad \qquad h_{t+1} \le h_t. \tag{5}$$

$$\forall t \ge rank(\mathbf{X}_1) + 2: \qquad h_t \le \frac{8\beta}{t - rank(\mathbf{X}_1) + 4}.$$
(6)

Moreover, there exists scalars  $\mathcal{E} = O\left(\min\left\{\frac{\alpha\delta^2}{\beta^2}, \frac{\delta^2}{\beta}, \delta, \alpha\lambda_{r^*}^2, \frac{\delta^3}{\beta^2}\right\}\right), \mathcal{E}_1 = O\left(\frac{\alpha\delta^2}{\beta^2}\right), \mathcal{E}_2 = O(\alpha\lambda_{r^*}^2)$  such that on each iteration t in which a <u>drop step is not taken</u>: If  $r^* = 1$  and  $h_t < \mathcal{E}_1$  then

$$h_{t+1} \le h_t \left( 1 - \min\{\frac{\delta}{12\beta}, \frac{1}{2}\} \right). \tag{7}$$

If  $r^* = n$  and  $h_t < \mathcal{E}_2$  then

$$h_{t+1} \le h_t \left( 1 - \frac{\alpha}{512\beta n} \right). \tag{8}$$

If  $n > r^* \ge 2$  and  $h_t < \mathcal{E}$  then,

$$\mathbb{E}[h_{t+1}|\mathbf{X}_t] \le h_t \left(1 - \frac{1}{r^*} \min\left\{\frac{\delta}{48G} \min\{\frac{\delta}{3\beta}, \lambda_{r^*}\}, \frac{\alpha}{512\beta}\right\}\right),\tag{9}$$

where the expectation is w.r.t. the random choice of  $\mathbf{u}_{t,-}$ , and  $G := \sup_{\mathbf{X} \in S^n} \|\nabla f(\mathbf{X})\|$ .

Finally, up to any iteration t, the overall number of drop steps cannot exceed  $(t + rank(\mathbf{X}_1) - 2)/2$ .

Before proving the theorem let us make a few comments. Result (5) follows in a straightforward manner from the design of the algorithm. Results (6), (7) follow essentially from already known analyses of the Frank-Wolfe method [19] and [14]. The main novel result is the (expected) linear convergence rate in the case  $n > r^* \ge 2$  given in Eq. (9). Note that the rate in (9) depends explicitly on the rank of optimal solutions  $r^*$ , which is well known to by unavoidable in worst case, see for instance [18]. Also note that the  $\frac{\alpha}{\beta}$  term inside the min in the RHS of (9) corresponds (up to a universal constant) to the standard linear convergence rate of (unaccelerated) gradient methods, such as the block Frank-Wolfe method [1]. Finally, Result (8) for the case  $r^* = n$ , i.e., the optimal solutions lie in the relative interior of  $S^n$ , follows as a simplified case of Result (9).

#### 3.1 Proof of Theorem 2

Additional notation for the proof of Theorem 2		
$\nabla_t$	$ abla f(\mathbf{X}_t)$	
$\delta_t$	$\lambda_{n-r^*}(\nabla f(\mathbf{X}_t)) - \lambda_{n-r^*+1}(\nabla f(\mathbf{X}_t)) \text{ (when } r^* < n)$	
$\delta_{t,r^*}$	$\lambda_{n-r^*+1}( abla f(\mathbf{X}_t)) - \lambda_n( abla f(\mathbf{X}_t))$	
$r_t$	$\operatorname{rank}(\mathbf{X}_t)$	
$\mathbf{P}_t \in \mathbb{S}^n_+$	projection matrix onto span of $r^*$ eigenvectors of $\nabla_t$ corresponding	
	to smallest eigenvalues of $\nabla_t$	
$\mathbf{P}_t^\perp \in \mathbb{S}^n_+$	projection matrix onto span of $n-r^*$ eigenvectors of $\nabla_t$ correspond-	
	ing to largest eigenvalues of $\nabla_t$ . If $r^* = n$ set $\mathbf{P}_t^{\perp} = 0_{n \times n}$	
$\mathbb{E}_t[\cdot]$	$\mathbb{E}_{\mathbf{u}_{t-1}}[\cdot \mathbf{X}_t]$ — the expectation w.r.t. $\mathbf{u}_{t-1}$ conditioned on $\mathbf{X}_t$	

The proof of Theorem 2 is constructed from several lemmas that analyze the benefit of each one of the steps employed by Algorithm 1. It is then established how their combination yields the theorem.

The following lemma establishes conditions under-which a drop step will be taken.

**Lemma 3** (drop step). Fix an iteration t of Algorithm 1 for which  $h_t < \min\{\frac{\delta_t^2}{18\beta}, \frac{\delta_t}{6}\}$ . If  $rank(\mathbf{X}_t) > r^*$ , then  $f(\mathbf{X}_{t+1}^{drop}) \leq f(\mathbf{X}_t)$ .

*Proof.* Recall that  $\mathbf{X}_{t+1}^{\text{drop}} = \frac{1}{1-c} (\mathbf{X}_t - c \mathbf{v} \mathbf{v}^{\top})$  for  $c = (\mathbf{v}^{\top} \mathbf{X}_t^{\dagger} \mathbf{v})^{-1}$  and  $\mathbf{v} \in \operatorname{argmax} \{ \mathbf{v}^{\top} \nabla_t \mathbf{v} \mid \mathbf{v} \in \operatorname{Im}(\mathbf{X}_t), \|\mathbf{v}\| = 1 \}$ . Using the smoothness of f we have that,

$$f(\mathbf{X}_{t+1}^{\text{drop}}) \leq f(\mathbf{X}_t) + \langle \frac{1}{1-c} (\mathbf{X}_t - c \mathbf{v} \mathbf{v}^{\top}) - \mathbf{X}_t, \nabla_t \rangle + \frac{\beta}{2} \| \frac{1}{1-c} (\mathbf{X}_t - c \mathbf{v} \mathbf{v}^{\top}) - \mathbf{X}_t \|_F^2$$
$$= f(\mathbf{X}_t) + \frac{c}{1-c} \langle \mathbf{X}_t - \mathbf{v} \mathbf{v}^{\top}, \nabla_t \rangle + \frac{\beta c^2}{(1-c)^2}.$$
(10)

Since dim Im( $\mathbf{X}_t$ )  $\geq r^* + 1$ , it follows from the Poincaré separation theorem that  $\mathbf{v}^\top \nabla_t \mathbf{v} \geq \lambda_{n-r^*}(\nabla_t) = \lambda_{n-r^*+1} + \delta_t$ . On the other-hand, we argue that  $\langle \mathbf{X}_t, \nabla_t \rangle \leq \delta_{n-r^*}(\nabla_t) = \delta_{n-r^*+1} + \delta_t$ .

 $\lambda_{n-r^*+1} + \delta_t/2$ . To see why the latter is true, let us assume by way of contradiction that it is not, and let us denote the matrix  $\mathbf{Y}_{\eta} = (1 - \eta)\mathbf{X}_t + \eta \mathbf{v}_{t,+} \mathbf{v}_{t,+}^{\top}$  for some  $\eta \in [0,1]$  to be specified later, and  $\mathbf{v}_{t,+}$  as defined in Algorithm 1. Using the smoothness of f again we have that,

$$f(\mathbf{Y}_{\eta}) - f^* \leq f(\mathbf{X}_t) + \eta \langle \mathbf{v}_{t,+} \mathbf{v}_{t,+}^{\top} - \mathbf{X}_t, \nabla_t \rangle + \frac{\beta \eta^2}{2} \| \mathbf{v}_{t,+} \mathbf{v}_{t,+}^{\top} - \mathbf{X}_t \|_F^2$$
$$\leq f(\mathbf{X}_t) - f^* - \eta \left( \frac{\delta_t}{2} - \eta \beta \right).$$

Setting  $\eta = \min\{\frac{\delta_t}{3\beta}, 1\}$  yields then

$$f(\mathbf{Y}_{\eta}) - f^* \le f(\mathbf{X}_t) - f^* - \min\{\frac{\delta_t}{3\beta}, 1\}\frac{\delta_t}{6}$$

Now we see that if  $h_t < \min\{\frac{\delta_t^2}{18\beta}, \frac{\delta_t}{6}\}$ , as the lemma assumes, we indeed get a contradiction.

Going back to (10), we can now write

$$f(\mathbf{X}_{t+1}^{\text{drop}}) - f^* \le f(\mathbf{X}_t) - f^* - \frac{\delta_t c}{2(1-c)} + \frac{\beta c^2}{(1-c)^2}.$$
 (11)

Observe that if  $\frac{c}{1-c} \leq \frac{\delta_t}{2\beta}$ , then indeed  $f(\mathbf{X}_{t+1}^{\text{drop}}) \leq f(\mathbf{X}_t)$ . Note furthermore that it must be the case that  $\frac{c}{1-c} \leq \frac{\delta_t}{3\beta}$ : the function  $g(c) = \frac{c}{1-c}$  is monotone increasing on the interval [0, 1), and thus we can in principle replace the scalar c with some  $c' \leq c$  such that  $\frac{c'}{1-c'} < \frac{\delta_t}{3\beta}$ . Denoting the resulting matrix by  $\mathbf{X}'_{t+1} = \frac{1}{1-c'}(\mathbf{X}_t - c'\mathbf{v}\mathbf{v}^{\top})$  and replacing c with c' in (11) will then yield,

$$f(\mathbf{X}'_{t+1}) - f^* \le f(\mathbf{X}_t) - f^* - \frac{\delta_t^2}{18\beta}$$

However, using the assumption of the lemma that  $h_t < \frac{\delta_t^2}{18\beta}$ , this would lead to a contradiction. Thus, we can conclude that indeed  $f(\mathbf{X}_{t+1}^{\text{drop}}) \leq f(\mathbf{X}_t)$ .

Since drop steps, as analyzed in the previous lemma, do not necessarily lower the objective value by a bounded amount, we have the following simple observation that upper-bounds the number of such steps possible until any iteration  $t \ge 1$ .

**Observation 1.** The overall number of drop steps up to the beginning of any iteration t of Algorithm 1 cannot be larger than  $(t + \operatorname{rank}(\mathbf{X}_1) - 2)/2$ .

*Proof.* According to Lemma 2, each drop step reduces the rank of the resulting matrix. On the other-hand, all other steps increase the rank by at most one. Thus, denoting the number of drop steps up to the beginning of some iteration t by  $t_{\rm drop}$ , it must hold that

$$1 \leq \operatorname{rank}(\mathbf{X}_t) \leq \operatorname{rank}(\mathbf{X}_1) - t_{\operatorname{drop}} + (t - 1 - t_{\operatorname{drop}}),$$

which yields the desired bound.

The following lemma (deterministically) bounds the reduction in function value due to a pairwise step. It is complemented by Lemma 5 which further lower-bounds the expectation of the random variable  $\|\mathbf{P}_t^{\perp}\mathbf{u}_{t,-}\|^2$ .

**Lemma 4** (pairwise step). Fix some iteration t of Algorithm 1 for which it holds that  $h_t \leq \min\{\frac{\delta_t^3}{24\beta^2}, \frac{\delta_t^2}{12\beta}\}$ . Then,

$$f(\mathbf{X}_{t+1}^{PFW}) \le f(\mathbf{X}_t) - \min\{\frac{\delta_t^2}{9\beta}, \frac{\gamma_t \delta_t}{3}\} \|\mathbf{P}_t^{\perp} \mathbf{u}_{t,-}\|^2.$$

*Proof.* Write  $\mathbf{u}_{t,-} = \tau \mathbf{z} + \sqrt{1 - \tau^2} \mathbf{w}$ , where  $\mathbf{z} = \frac{\mathbf{P}_t^{\perp} \mathbf{u}_{t,-}}{\|\mathbf{P}_t^{\perp} \mathbf{u}_{t,-}\|}$ ,  $\tau = \|\mathbf{P}_t^{\perp} \mathbf{u}_{t,-}\|$ , and  $\mathbf{w} = \frac{\mathbf{P}_t \mathbf{u}_{t,-}}{\|\mathbf{P}_t \mathbf{u}_{t,-}\|}$ . Consider now some unit vector  $\mathbf{u} = \tau' \mathbf{z} + \sqrt{1 - \tau'^2} \mathbf{w}$  with  $0 \le \tau' \le \tau$ . Fix some step-size  $\eta \in [0, 1]$  and denote  $\mathbf{Y}_{\eta} = \mathbf{X}_t + \eta(\mathbf{u}\mathbf{u}^{\top} - \mathbf{u}_{t,-}\mathbf{u}_{t,-}^{\top})$ . Using the smoothness of f it holds that,

$$\begin{aligned} f(\mathbf{Y}_{\eta}) &\leq f(\mathbf{X}_{t}) + \eta \langle \mathbf{u}\mathbf{u}^{\top} - \mathbf{u}_{t,-}\mathbf{u}_{t,-}^{\top}, \nabla_{t} \rangle + \frac{\eta^{2}\beta}{2} \|\mathbf{u}\mathbf{u}^{\top} - \mathbf{u}_{t,-}\mathbf{u}_{t,-}^{\top}\|_{F}^{2} \\ &= f(\mathbf{X}_{t}) + \eta \left( (\tau'^{2} - \tau^{2})\mathbf{z}^{\top}\nabla_{t}\mathbf{z} - (\tau'^{2} - \tau^{2})\mathbf{w}^{\top}\nabla_{t}\mathbf{w} \right) + \eta^{2}\beta \left( 1 - (\mathbf{u}^{\top}\mathbf{u}_{t,-})^{2} \right) \\ &\leq f(\mathbf{X}_{t}) - \eta (\tau^{2} - \tau'^{2})\delta_{t} + \eta^{2}\beta \left( 1 - (\tau\tau' + \sqrt{1 - \tau^{2}}\sqrt{1 - \tau'^{2}})^{2} \right) \\ &= f(\mathbf{X}_{t}) - \eta (\tau^{2} - \tau'^{2})\delta_{t} \\ &+ \eta^{2}\beta \left( 1 - \left( \tau^{2}\tau'^{2} + (1 - \tau^{2} - \tau'^{2} + \tau^{2}\tau'^{2}) + 2\tau\tau'\sqrt{1 - \tau^{2}}\sqrt{1 - \tau'^{2}} \right) \right) \\ &= f(\mathbf{X}_{t}) - \eta (\tau^{2} - \tau'^{2})\delta_{t} + \eta^{2}\beta \left( \tau^{2} + \tau'^{2} - 2\tau^{2}\tau'^{2} - 2\tau\tau'\sqrt{1 - \tau^{2}}\sqrt{1 - \tau'^{2}} \right) \\ &\leq f(\mathbf{X}_{t}) - \eta (\tau^{2} - \tau'^{2})\delta_{t} + \eta^{2}\beta \left( \tau^{2} + \tau'^{2} - 2\tau^{2}\tau'^{2} - 2\tau\tau'(1 - \tau^{2}) \right) \\ &= f(\mathbf{X}_{t}) - \eta (\tau^{2} - \tau'^{2})\delta_{t} + \eta^{2}\beta (\tau - \tau')^{2} + 2\eta^{2}\beta\tau^{2}\tau'(\tau - \tau') \\ &= f(\mathbf{X}_{t}) - \eta (\tau - \tau') \left( (\tau + \tau')\delta_{t} - \eta\beta \left( (\tau - \tau') + 2\tau^{2}\tau' \right) \right), \end{aligned}$$

where (a) follows from the assumption  $\tau' \leq \tau$ .

First note that setting  $\tau' = 0$  in (12) yields

$$f(\mathbf{Y}_{\eta}) \leq f(\mathbf{X}_{t}) - \eta \tau^{2} \left( \delta_{t} - \eta \beta \right).$$

Using the feasible step-size  $\eta^* = \min\{\frac{\delta_t}{2\beta}, \gamma_t\}$  then gives,

$$f(\mathbf{Y}_{\eta^*}) - f^* \le h_t - \frac{\min\{\frac{\delta_t}{2\beta}, \gamma_t\}\delta_t\tau^2}{2},$$

which implies that

$$\tau^2 \le \frac{2h_t}{\min\{\frac{\delta_t}{2\beta}, \gamma_t\}\delta_t}.$$
(13)

Consider now the case  $\tau' = (1 - c)\tau$  for some scalar  $c \in [0, 1]$  to be specified in the sequel. Using the smoothness of f again, the definition of  $\mathbf{u}_{t,+}$  in Algorithm 1

(see also Eq. (4)), and applying (12) with  $\eta = \gamma_t$  we have that,

$$f(\mathbf{X}_{t+1}^{\text{PFW}}) \leq f(\mathbf{X}_{t}) + \gamma_{t} \langle \mathbf{u}_{t,+} \mathbf{u}_{t,+}^{\top} - \mathbf{u}_{t,-} \mathbf{u}_{t,-}^{\top}, \nabla_{t} \rangle + \frac{\gamma_{t}^{2} \beta}{2} \|\mathbf{u}_{t,+} \mathbf{u}_{t,+}^{\top} - \mathbf{u}_{t,-} \mathbf{u}_{t,-}^{\top}\|_{F}^{2}$$
$$\leq f(\mathbf{X}_{t}) - \gamma_{t} c \tau^{2} \left(\delta_{t} - \gamma_{t} \beta (c + 2\tau^{2})\right).$$

Let us now set  $c = \min\{\frac{\delta_t}{3\gamma_t\beta}, 1\}$  to obtain

$$f(\mathbf{X}_{t+1}^{\text{PFW}}) \leq f(\mathbf{X}_t) - \gamma_t \min\{\frac{\delta_t}{3\gamma_t\beta}, 1\}\tau^2 \left(\frac{2\delta_t}{3} - 2\gamma_t\beta\tau^2\right)$$
$$\leq f(\mathbf{X}_t) - \gamma_t \min\{\frac{\delta_t}{3\gamma_t\beta}, 1\}\tau^2 \left(\frac{2\delta_t}{3} - \frac{4\gamma_t\beta h_t}{\min\{\frac{\delta_t}{2\beta}, \gamma_t\}\delta_t}\right),$$

where the last inequality follows from plugging-in (13).

Thus, recalling that  $\tau^2 = \|\mathbf{P}_t^{\perp} \mathbf{u}_{t,-}\|^2$  and that  $\gamma_t \in (0,1]$ , we have that if  $h_t \leq \frac{\delta_t^2 \min\{\frac{\delta_t}{2\beta},1\}}{12\beta} \leq \frac{\delta_t^2 \min\{\frac{\delta_t}{2\beta},\gamma_t\}}{12\gamma_t\beta}$ , then we indeed have that

$$f(\mathbf{X}_{t+1}^{\text{PFW}}) \le f(\mathbf{X}_t) - \frac{\gamma_t \delta_t}{3} \min\{\frac{\delta_t}{3\gamma_t \beta}, 1\} \|\mathbf{P}_t^{\perp} \mathbf{u}_{t,-}\|^2.$$

**Lemma 5.** Fix some iteration t of Algorithm 1. It holds that  $\mathbb{E}_t[\|\mathbf{P}_t^{\perp}\mathbf{u}_{t,-}\|^2] \geq \frac{1}{r_t} Tr(\mathbf{P}_t^{\perp}\mathbf{X}_t).$ 

*Proof.* Write the eigen-decomposition of  $\mathbf{X}_t$  as  $\mathbf{X}_t = \mathbf{U}_t \Lambda_t \mathbf{U}_t^{\top}$ , where  $\Lambda_t$  is  $r_t \times r_t$  and  $\mathbf{U}_t$  is  $n \times r_t$ . It holds that,

$$\operatorname{Tr}(\mathbf{P}_{t}^{\perp}\mathbf{X}_{t}) = \operatorname{Tr}(\mathbf{P}_{t}^{\perp}\mathbf{U}_{t}\Lambda_{t}\mathbf{U}_{t}^{\top}) = \operatorname{Tr}(\mathbf{U}_{t}^{\top}\mathbf{P}_{t}^{\perp}\mathbf{U}_{t}\Lambda_{t}) \leq \lambda_{1}(\mathbf{X}_{t})\operatorname{Tr}(\mathbf{U}_{t}^{\top}\mathbf{P}_{t}^{\perp}\mathbf{U}_{t})$$
$$\leq \operatorname{Tr}(\mathbf{P}_{t}^{\perp}\mathbf{U}_{t}\mathbf{U}_{t}^{\top}) = \operatorname{Tr}\left(\mathbf{P}_{t}^{\perp}\mathbb{E}_{t}[r_{t}\mathbf{u}_{t,-}\mathbf{u}_{t,-}^{\top}]\right) = \mathbb{E}_{t}[r_{t}\operatorname{Tr}\left(\mathbf{P}_{t}^{\perp}\mathbf{u}_{t,-}\mathbf{u}_{t,-}^{\top}\right)]$$
$$= r_{t}\mathbb{E}_{t}[||\mathbf{P}_{t}^{\perp}\mathbf{u}_{t,-}||^{2}].$$

The following lemma is central to the analysis of the benefit of both standard Frank-Wolfe steps and away steps in Algorithm 1. It lower-bounds the eigengap between the extreme eigenvalues associated with the principal  $r^*$ -dimensional subspace (corresponding to the  $r^*$  lowest eigenvalues) of the current gradient direction  $\nabla_t$ . This is essentially a type of a Polyak-Lojasiewicz (PL) inequality w.r.t. to the face of  $S^n$  corresponding to the this principal subspace of  $\nabla_t$ .

**Lemma 6** (in-face eigengap). Assume  $r^* \geq 2$ . Fix some iteration t of Algorithm 1 and suppose that  $Tr(\mathbf{P}_t^{\perp}\mathbf{X}_t) \leq \frac{h_t}{4\|\nabla_t\|}$ . Then,

$$\lambda_{n-r^*+1}(\nabla_t) - \lambda_n(\nabla_t) \ge \sqrt{\frac{\alpha h_t}{8r^*}}$$

*Proof.* Denote  $\mathbf{X}^* = \arg\min_{\mathbf{Y}\in\mathcal{X}^*} \|\mathbf{Y} - \mathbf{X}_t\|_F$ . Denote  $\mathbf{W} = \mathbf{P}_t(\mathbf{X}_t - \mathbf{X}^*)\mathbf{P}_t$ ,  $\mathbf{W}^{\perp} = \mathbf{P}_t^{\perp}(\mathbf{X}_t - \mathbf{X}^*)\mathbf{P}_t^{\perp}$ , and note that since  $\operatorname{rank}(\mathbf{P}_t) = r^*$  it follows that  $\operatorname{rank}(\mathbf{W}) \leq r^*$ . Using the convexity of f we have that,

$$h_t \le \langle \mathbf{X}_t - \mathbf{X}^*, \nabla_t \rangle = \langle \mathbf{W}, \nabla_t \rangle + \langle \mathbf{W}^{\perp}, \nabla_t \rangle.$$
(14)

It holds that,

$$\langle \mathbf{W}, \nabla_t \rangle \leq \left( \sum_{i \in [n]:\lambda_i(\mathbf{W}) > 0} \lambda_i(\mathbf{W}) \right) \lambda_{n-r^*+1}(\nabla_t) + \left( \operatorname{Tr}(\mathbf{W}) - \sum_{i \in [n]:\lambda_i(\mathbf{W}) > 0} \lambda_i(\mathbf{W}) \right) \lambda_n(\nabla_t) \leq \sqrt{r^*} \|\mathbf{W}\|_F \left( \lambda_{n-r^*+1}(\nabla_t) - \lambda_n(\nabla_t) \right) + \operatorname{Tr}(\mathbf{W}) \lambda_n(\nabla_t) \leq \sqrt{r^*} \|\mathbf{X}_t - \mathbf{X}^*\|_F \left( \lambda_{n-r^*+1}(\nabla_t) - \lambda_n(\nabla_t) \right) + \operatorname{Tr}(\mathbf{W}) \lambda_n(\nabla_t) = \sqrt{r^*} \|\mathbf{X}_t - \mathbf{X}^*\|_F \left( \lambda_{n-r^*+1}(\nabla_t) - \lambda_n(\nabla_t) \right) - \operatorname{Tr}(\mathbf{W}^{\perp}) \lambda_n(\nabla_t) = \sqrt{r^*} \|\mathbf{X}_t - \mathbf{X}^*\|_F \left( \lambda_{n-r^*+1}(\nabla_t) - \lambda_n(\nabla_t) \right) - \operatorname{Tr}(\mathbf{W}^{\perp}) \lambda_n(\nabla_t) + \left( \operatorname{Tr}(\mathbf{P}_t^{\perp} \mathbf{X}^*) - \operatorname{Tr}(\mathbf{P}_t^{\perp} \mathbf{X}_t) \right) \lambda_n(\nabla_t),$$
(15)

where (a) holds since

$$\sum_{i \in [n]:\lambda_i(\mathbf{W}) > 0} \lambda_i(\mathbf{W}) \leq \sum_{i \in [n]} |\lambda_i(\mathbf{W})| \leq \sqrt{\operatorname{rank}(\mathbf{W})} \sqrt{\sum_{i \in [n]} \lambda_i(\mathbf{W})^2} \\ \leq \sqrt{r^*} \|\mathbf{W}\|_F,$$

(b) holds since  $\|\mathbf{W}\|_F \leq \|\mathbf{X}_t - \mathbf{X}^*\|_F$  which is due to the inequality  $\|\mathbf{AB}\|_F \leq \|\mathbf{A}\| \cdot \|\mathbf{B}\|_F$  for any two matrices  $\mathbf{A}, \mathbf{B} \in \mathbb{S}^n$ , and (c) holds since

$$\operatorname{Tr}(\mathbf{W}) = \operatorname{Tr}(\mathbf{P}_t(\mathbf{X}_t - \mathbf{X}^*)) = \operatorname{Tr}((\mathbf{I} - \mathbf{P}_t^{\perp})(\mathbf{X}_t - \mathbf{X}^*))$$
$$= \operatorname{Tr}(\mathbf{X}_t - \mathbf{X}^*) - \operatorname{Tr}(\mathbf{P}_t^{\perp}(\mathbf{X}_t - \mathbf{X}^*)) = -\operatorname{Tr}(\mathbf{P}_t^{\perp}(\mathbf{X}_t - \mathbf{X}^*)) = -\operatorname{Tr}(\mathbf{W}^{\perp}).$$

It also holds that,

$$\langle \mathbf{W}^{\perp}, \nabla_t \rangle = \langle \mathbf{P}_t^{\perp} (\mathbf{X}_t - \mathbf{X}^*) \mathbf{P}_t^{\perp}, \nabla_t \rangle$$
  
 
$$\leq \operatorname{Tr}(\mathbf{P}_t^{\perp} \mathbf{X}_t) \lambda_1(\nabla_t) - \operatorname{Tr}(\mathbf{P}_t^{\perp} \mathbf{X}^*) \lambda_{n-r^*}(\nabla_t).$$
 (16)

Combining (14), (15) and (16) gives,

$$h_{t} \leq \sqrt{r^{*}} \|\mathbf{X}_{t} - \mathbf{X}^{*}\|_{F} (\lambda_{n-r^{*}+1}(\nabla_{t}) - \lambda_{n}(\nabla_{t})) + 2\mathrm{Tr}(\mathbf{P}_{t}^{\perp}\mathbf{X}_{t})\|\nabla_{t}\| + \mathrm{Tr}(\mathbf{P}_{t}^{\perp}\mathbf{X}^{*}) (\lambda_{n}(\nabla_{t}) - \lambda_{n-r^{*}}(\nabla_{t})) \leq \sqrt{r^{*}} \|\mathbf{X}_{t} - \mathbf{X}^{*}\|_{F} (\lambda_{n-r^{*}+1}(\nabla_{t}) - \lambda_{n}(\nabla_{t})) + 2\mathrm{Tr}(\mathbf{P}_{t}^{\perp}\mathbf{X}_{t})\|\nabla_{t}\|.$$

Rearraning the inequality, using the assumption on  $\operatorname{Tr}(\mathbf{P}_t^{\perp}\mathbf{X}_t)$ , and the quadratic growth bound  $\|\mathbf{X}_t - \mathbf{X}^*\|_F \leq \sqrt{\frac{2h_t}{\alpha}}$ , yields the lemma.

The following Lemma, which builds on the result of the previous Lemma 6, establishes the reduction in objective function value for FW / away steps, under a positive in-face eigengap.

**Lemma 7** (FW/away steps under in-face eignegap). Assume  $r^* \geq 2$ . Fix some iteration t of Algorithm 1 and suppose that  $r_t \geq r^*$  and  $\lambda_{r_t}(\mathbf{X}_t) \geq \frac{\delta_{t,r^*}}{16\beta}$ . Then,

$$\min\{f(\mathbf{X}_{t+1}^{FW}), f(\mathbf{X}_{t+1}^{AFW})\} \le f(\mathbf{X}_t) - \frac{\delta_{t,r^*}^2}{64\beta}.$$
(17)

*Proof.* Since rank( $\mathbf{X}_t$ )  $\geq r^*$ , according to the Poincaré separation theorem, we have that there exists a unit vector  $\mathbf{v} \in \text{Im}(\mathbf{X}_t)$  such that  $\mathbf{v}^\top \nabla_t \mathbf{v} \geq \lambda_{n-r^*+1}(\nabla_t)$ , which in turn implies that  $\mathbf{v}_{t,-}^\top \nabla_t \mathbf{v}_{t,-} \geq \lambda_{n-r^*+1}(\nabla_t)$ . Also, we clearly have that  $\mathbf{v}_{t,+}^\top \nabla_t \mathbf{v}_{t,+} = \lambda_n(\nabla_t)$ . This gives,

$$\max\{\langle \mathbf{X}_{t} - \mathbf{v}_{t,+} \mathbf{v}_{t,+}^{\top}, \nabla_{t} \rangle, \langle \mathbf{v}_{t,-} \mathbf{v}_{t,-}^{\top} - \mathbf{X}_{t}, \nabla_{t} \rangle\} \geq \frac{1}{2} \langle \mathbf{v}_{t,-} \mathbf{v}_{t,-}^{\top} - \mathbf{v}_{t,+} \mathbf{v}_{t,+}^{\top}, \nabla_{t} \rangle$$
$$\geq \frac{\delta_{t,r^{*}}}{2}. \tag{18}$$

Using the smoothness of f we have that for any step-size  $\eta \in [0, 1]$ ,

$$f\left(\left(\mathbf{X}_{t}+\eta(\mathbf{v}_{t,+}\mathbf{v}_{t,+}^{\top}-\mathbf{X}_{t})\right) \leq f(\mathbf{X}_{t})+\eta\langle\mathbf{v}_{t,+}\mathbf{v}_{t,+}^{\top}-\mathbf{X}_{t},\nabla_{t}\rangle + \frac{\eta^{2}\beta}{2}\|\mathbf{v}_{t,+}\mathbf{v}_{t,+}^{\top}-\mathbf{X}_{t}\|_{F}^{2} \leq f(\mathbf{X}_{t})+\eta\langle\mathbf{v}_{t,+}\mathbf{v}_{t,+}^{\top}-\mathbf{X}_{t},\nabla_{t}\rangle+\eta^{2}\beta.$$
(19)

Similarly, we have that for any step-size  $\eta \in [0, \lambda_{r_t}(\mathbf{X}_t)] \subseteq [0, (\mathbf{v}_{t,-}^{\top} \mathbf{X}_t^{\dagger} \mathbf{v}_{t,-})^{-1}]$  it holds that,

$$f\left(\frac{1}{1-\eta}\left(\mathbf{X}_{t}-\eta\mathbf{v}_{t,-}\mathbf{v}_{t,-}^{\mathsf{T}}\right)\right) = f\left(\mathbf{X}_{t}+\frac{\eta}{1-\eta}\left(\mathbf{X}_{t}-\mathbf{v}_{t,-}\mathbf{v}_{t,-}^{\mathsf{T}}\right)\right)$$
$$\leq f(\mathbf{X}_{t}) + \frac{\eta}{1-\eta}\langle\mathbf{X}_{t}-\mathbf{v}_{t,-}\mathbf{v}_{t,-}^{\mathsf{T}},\nabla_{t}\rangle + \frac{\eta^{2}\beta}{(1-\eta)^{2}}$$
$$\leq f(\mathbf{X}_{t}) + \eta\langle\mathbf{X}_{t}-\mathbf{v}_{t,-}\mathbf{v}_{t,-}^{\mathsf{T}},\nabla_{t}\rangle + \frac{\eta^{2}\beta}{(1-\eta)^{2}}, \quad (20)$$

where in the last inequality we have used the fact that  $\eta \in [0, 1)$  and  $\langle \mathbf{X}_t - \mathbf{v}_{t,-} \mathbf{v}_{t,-}^\top, \nabla_t \rangle \geq 0$ .

Combining Eq. (19) and Eq. (20) with Eq. (18), and the fact that the step sizes  $\eta_t^{\text{FW}}$ ,  $\eta_t^{\text{AFW}}$  in Algorithm 1 are set via line-search, we have that

$$\forall \eta \in [0, \lambda_{r_t}(\mathbf{X}_t)]: \min\{f(\mathbf{X}_{t+1}^{\text{FW}}), f(\mathbf{X}_{t+1}^{\text{AFW}})\} \le f(\mathbf{X}_t) - \frac{\eta \delta_{t,r^*}}{2} + \frac{\eta^2 \beta}{(1-\eta)^2}.$$
 (21)

In particular, under the assumptions of the lemma that  $\lambda_{r_t}(\mathbf{X}_t) \geq \frac{\delta_{t,r^*}}{16\beta}$  and  $r_t \geq r^* \geq 2$ , which implies that  $\lambda_{r_t}(\mathbf{X}_t) \leq 1/2$ , setting  $\eta = \frac{\delta_{t,r^*}}{16\beta}$  in (21) we have that,

$$\min\{f(\mathbf{X}_{t+1}^{\text{FW}}), f(\mathbf{X}_{t+1}^{\text{AFW}})\} \le f(\mathbf{X}_t) - \frac{\delta_{t,r^*}^2}{32\beta} + \frac{\delta_{t,r^*}^2}{64\beta} = f(\mathbf{X}_t) - \frac{\delta_{t,r^*}^2}{64\beta},$$

which proves the lemma.

We are now finally positioned to complete the proof of Theorem 2.

Proof of Theorem 2. First note that Result (5) (monotonicity of the sequence  $(h_t)_{t\geq 1}$ ) follows in a straightforward manner from the design of Algorithm 1: on each iteration t a drop step is taken only if it does not increase the objective function value, and if a drop step is not taken then, due to the update rule for the next iterate  $\mathbf{X}_{t+1}$  which suggests that  $f(\mathbf{X}_{t+1}) \leq f(\mathbf{X}_{t+1}^{\mathrm{FW}})$  and due to the use of line-search in the computation of  $f(\mathbf{X}_{t+1}^{\mathrm{FW}})$ , we have that  $f(\mathbf{X}_{t+1}) \leq f(\mathbf{X}_{t+1}^{\mathrm{FW}}) \leq f(\mathbf{X}_{t})$ .

We turn to prove Result (6). We use again the observation that on each iteration t which is not a drop step Algorithm 1 reduces the objective function value at least by the amount that a standard Frank-Wolfe step with line-search will reduce  $(f(\mathbf{X}_{t+1}) \leq f(\mathbf{X}_{t+1}^{\mathrm{FW}}))$ . Combining this observation with Result (5), we have that on each iteration t of Algorithm 1, denoting by  $t_{\mathrm{drop}}$  the overall number of drop steps performed up to iteration t, the approximation error  $h_t$  is upper-bounded by  $h_{t-t_{\mathrm{drop}}}^{\mathrm{FW}}$ — the approximation error of the standard FW with line search at the beginning of its  $(t - t_{\mathrm{drop}})$  iteration. Result (6) now follows from using the well-known convergence result for the standard Frank-Wolfe with line-search over the spectrahedron  $S^n$  (i.e.,  $h_t^{\mathrm{FW}} \leq \frac{4\beta}{t+1}$  for all  $t \geq 2$ , see for instance [19]) and the upper-bound on  $t_{\mathrm{drop}}$  in Observation 1.

We now turn to prove Result (7) which deals with the specific case  $r^* = 1$ , i.e., a rank-one optimal solution. In this case, under Assumption 2 there exists a unique optimal solution  $\mathbf{X}^*$ , see [14]. [14] proved (see proof of Theorem 2 in [14]) that on each iteration t for which we have  $\|\nabla_t - \nabla f(\mathbf{X}^*)\|_F \leq \frac{\delta}{3}$ , it holds that

$$f(\mathbf{X}_{t+1}^{\text{FW}}) - f^* \le (f(\mathbf{X}_t) - f^*) \left(1 - \min\{\frac{\delta}{12\beta}, \frac{1}{2}\}\right).$$
 (22)

Using the quadratic growth assumption and the smoothness of f we have that,

$$\|\nabla_t - \nabla f(\mathbf{X}^*)\|_F \le \beta \|\mathbf{X}_t - \mathbf{X}^*\|_F \le \frac{\sqrt{\alpha\beta}}{\sqrt{2}}\sqrt{h_t}.$$

Thus, whenever  $h_t \leq \frac{\alpha \delta^2}{18\beta^2}$ , we indeed have that  $\|\nabla_t - \nabla f(\mathbf{X}^*)\|_F \leq \frac{\delta}{3}$ . Combining this with Eq. (22) and the fact that in case a drop step is not taken we have that  $f(\mathbf{X}_{t+1}) \leq f(\mathbf{X}_{t+1}^{\mathrm{FW}})$ , we obtain Result (7).

We now move on to prove the main result of the theorem — Result (9). Recall in this case we assume  $n > r^* \ge 2$ . Let us fix some iteration t of Algorithm 1 and denote  $\mathbf{X}^* = \arg \min_{\mathbf{Y} \in \mathcal{X}^*} \|\mathbf{Y} - \mathbf{X}_t\|_F$ . It holds that,

$$\forall i \in [n]: \quad |\lambda_i(\nabla_t) - \lambda_i(\nabla f(\mathbf{X}^*))| \le \|\nabla_t - \nabla f(\mathbf{X}^*)\| \le \beta \|\mathbf{X}_t - \mathbf{X}^*\|_F$$
$$\le \frac{\sqrt{2\beta}}{\sqrt{\alpha}}\sqrt{h_t}, \tag{23}$$

where (a) is due the the smoothness of f, and (b) is due to the quadratic growth assumption.

Thus, if  $h_t \leq \epsilon_1 := \frac{\alpha \delta^2}{32\beta^2}$ , we have that

$$\delta_t := \lambda_{n-r^*}(\nabla_t) - \lambda_{n-r^*+1}(\nabla_t) \ge \delta - 2\frac{\sqrt{2}\beta}{\sqrt{\alpha}}\sqrt{\epsilon_1} = \frac{\delta}{2}.$$
 (24)

Denote  $\epsilon_2 = \min\{\frac{(\delta/2)^2}{18\beta}, \frac{\delta/2}{6}\}$ . Using Lemma 3 we have that if  $h_t < \min\{\epsilon_1, \epsilon_2\}$ and  $\operatorname{rank}(\mathbf{X}_t) > r^*$ , then a drop step will be preformed. Note also that if  $h_t < \min\{\epsilon_1, \epsilon_2\}$  and a drop step is not taken, then it must hold that  $\operatorname{rank}(\mathbf{X}_t) \leq r^*$ . Note also that if  $h_t < \epsilon_3 := \frac{\alpha \lambda_{r^*}^2}{8}$ , using the quadratic growth assumption we have that,

$$|\lambda_{r^*}(\mathbf{X}_t) - \lambda_{r^*}(\mathbf{X}^*)| \le \|\mathbf{X}_t - \mathbf{X}^*\|_F \le \sqrt{\frac{2h_t}{\alpha}},$$

which implies that

$$\lambda_{r^*}(\mathbf{X}_t) \ge \frac{\lambda_{r^*}}{2},\tag{25}$$

which further implies that  $\operatorname{rank}(\mathbf{X}_t) \geq r^*$ .

Thus, if  $h_t < \min\{\epsilon_1, \epsilon_2, \epsilon_3\}$  and a drop step is not performed, it must hold that  $\operatorname{rank}(\mathbf{X}_t) = r^*$ . Furthermore, it holds that  $\gamma_t \ge \lambda_{r^*}(\mathbf{X}_t) \ge \lambda_{r^*}/2$ .

Define also  $\epsilon_4 := \frac{(\delta/2)^3}{24\beta^2}$  and throughout the sequel suppose that indeed  $h_t < \min\{\epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4\}$  and a drop step is not taken. We consider two cases. First, if  $\operatorname{Tr}(\mathbf{P}_t^{\perp}\mathbf{X}_t) > \frac{h_t}{4G}$ , then it follows from Lemma 4 that

$$\begin{split} \mathbb{E}_t[f(\mathbf{X}_{t+1}^{\text{PFW}})] &\leq f(\mathbf{X}_t) - \mathbb{E}_t\left[\min\{\frac{\delta_t^2}{9\beta}, \frac{\gamma_t \delta_t}{3}\} \|\mathbf{P}_t^{\perp} \mathbf{u}_{t,-}\|^2\right] \\ &\leq f(\mathbf{X}_t) - \min\{\frac{\delta^2}{36\beta}, \frac{\lambda_{r^*} \delta}{12}\} \mathbb{E}_t\left[\|\mathbf{P}_t^{\perp} \mathbf{u}_{t,-}\|^2\right] \\ &\leq f(\mathbf{X}_t) - \frac{\delta}{48Gr^*}\min\{\frac{\delta}{3\beta}, \lambda_{r^*}\}h_t, \end{split}$$

where (a) follows from plugging-in (24), and since  $\gamma_t \ge \lambda_{r^*}/2$ , and (b) follows from Lemma 5.

This implies that

$$\mathbb{E}_{t}[f(\mathbf{X}_{t+1})] - f^{*} = \mathbb{E}_{t}\left[\min\{f(\mathbf{X}_{t+1}^{\mathrm{FW}}), f(\mathbf{X}_{t+1}^{\mathrm{AFW}}), f(\mathbf{X}_{t+1}^{\mathrm{PFW}})\}\right] - f^{*}$$

$$\leq \min\{f(\mathbf{X}_{t+1}^{\mathrm{FW}}), f(\mathbf{X}_{t+1}^{\mathrm{AFW}}), \mathbb{E}_{t}\left[f(\mathbf{X}_{t+1}^{\mathrm{PFW}})\right]\} - f^{*}$$

$$\leq h_{t}\left(1 - \frac{\delta}{48Gr^{*}}\min\{\frac{\delta}{3\beta}, \lambda_{r^{*}}\}\right).$$
(26)

In the complementing case that  $\operatorname{Tr}(\mathbf{P}_t^{\perp} \perp \mathbf{X}_t) \leq \frac{h_t}{4G}$ , it follows from Lemma 6 that

$$\lambda_{n-r^*+1}(\nabla_t) - \lambda_n(\nabla_t) \ge \sqrt{\frac{\alpha h_t}{8r^*}}.$$
(27)

Additionally, using Eq. (23) again we have that,

$$\lambda_{n-r^*+1}(\nabla_t) - \lambda_n(\nabla_t) \leq \lambda_{n-r^*+1}(\nabla f(\mathbf{X}^*)) - \lambda_n(\nabla f(\mathbf{X}^*)) + \frac{2\sqrt{2\beta}}{\sqrt{\alpha}}\sqrt{h_t}$$
$$= \frac{2\sqrt{2\beta}}{\sqrt{\alpha}}\sqrt{h_t} \leq \frac{2\sqrt{2\beta}}{\sqrt{\alpha}}\sqrt{\epsilon_3} = \beta\lambda_{r^*} \leq 2\beta\lambda_{r^*}(\mathbf{X}_t), \quad (28)$$

where (a) follows since due to the optimality of  $\mathbf{X}^*$  it holds that  $\lambda_n(\nabla f(\mathbf{X}^*)) = \lambda_{n-r^*+1}(\nabla f(\mathbf{X}^*))$  (see also Lemma 5.2 in [13]), and (b) follows from Eq. (25).

Thus, combining Lemma 7 with Eq. (27), we have that

$$f(\mathbf{X}_{t+1}) - f^* \le \min\{f(\mathbf{X}_{t+1}^{\text{FW}}), f(\mathbf{X}_{t+1}^{\text{AFW}})\} - f^* \le h_t \left(1 - \frac{\alpha}{512\beta r^*}\right).$$
(29)

Combining (26) and (29) yields Result (7).

Finally, it remains to prove Result (8) for the case  $r^* = n$ , which follows as a simplified case of Result (7). If  $h_t < \epsilon_3$ , then we have that Eq. (25) and Eq. (28) hold, and as a concequence the assumptions of Lemma 6 and Lemma 7 hold. Thus, if follows that Eq. (29) also holds.

### 4 Numerical Experiments

We conduct numerical experiments with synthetic data for the problem of recovering a ground truth matrix in  $S^n$  from random linear measurements. We use a smiller setup and methodology to the one considered in previous works [14, 7]. Concretely, we consider the following optimization problem:

$$\min_{\mathbf{X}\in\mathcal{S}^n} \{f(\mathbf{X}) := \frac{1}{2} \sum_{i=1}^m \left( \mathbf{a}_i^\top (\tau \mathbf{X}) \mathbf{a}_i - \mathbf{b}_i \right)^2 \}.$$
(30)

We let  $\mathbf{X}_{\sharp} = \mathbf{U}_{\sharp} \mathbf{U}_{\sharp}^{\top} \in S^{n}$  denote the ground truth matrix, where  $\mathbf{U}_{\sharp} \in \mathbb{R}^{n \times r^{*}}$  is first set to a random matrix with standard Gaussian entries and then normalized to have unit Frobenius norm. Each  $\mathbf{a}_{i} \in \mathbb{R}^{n}$  is a random vector with standard Gaussian entries. We set  $\mathbf{b} = \mathbf{b}_{\sharp} + \mathbf{z}$ , where  $\mathbf{b}_{\sharp}(i) = \mathbf{a}_{i}^{\top} \mathbf{X}_{\sharp} \mathbf{a}_{i}$ , and  $\mathbf{z} = \frac{\|\mathbf{b}_{\sharp}\|}{2} \mathbf{v}$  for a random unit vector  $\mathbf{v}$ .

In order to avoid over-fitting the noise we set the scaling parameter  $\tau$  in (30) to  $\tau = 0.5$  (as also been done in [14, 7]).

We solve Problem (30) to accuracy of the order  $10^{-11}$  or lower and denote our estimate for the optimal solution by  $\widehat{\mathbf{X}^*}^4$ . We set the smoothness parameter  $\beta$  to  $\beta = n^2/2^5$ .

<sup>&</sup>lt;sup>4</sup>this is verified by computing the dual gap for the candidate solution  $\widehat{\mathbf{X}^*}$ , which is given by  $\langle \widehat{\mathbf{X}^*}, \nabla f(\widehat{\mathbf{X}^*}) \rangle - \lambda_n (\nabla f(\widehat{\mathbf{X}^*}))$  and is always an upper-bound on the approximation error  $f(\widehat{\mathbf{X}^*}) - f^*$  due the convexity of f

<sup>&</sup>lt;sup>5</sup>while this might seem conservative, note that a simple calculation shows that  $\beta \leq \sum_{i=1}^{m} \|\mathbf{a}_i\|^4$ and thus we should roughly take  $\beta \approx mn^2$ , however this turns out to be an overly pessimistic estimate (see also experiments in [7])

Each one of the figures presented below is the average of 10 i.i.d. runs. In particular, we verify that in each of our randomized experiments the strict complementarity condition indeed holds with substantial positive values (see also Section 2.2.1 in [14] for a detailed discussion about verifying the strict complementarity assumption). An exception is Figure 4 which examines a setting in which the strict complementarity condition intentionally does not hold.

#### 4.1 Comparison to standard Frank-Wolfe

We begin by comparing our Algorithm 1 (ALG 1 in the figures below) to the standard Frank-Wolfe with line-search method (FW in the figures below). Recall that FW has a worst case convergence rate of O(1/t) [20]. In [14] it was established that under strict complementarity and in the special case  $r^* = 1$ , it enjoys a linear convergence rate, while it remained unknown if this is also the case for  $r^* \geq 2$ . Figure 1, which plots for each method the value  $\log(f(\mathbf{X}_t) - f(\widehat{\mathbf{X}^*}))$  vs the number of iteration t, shows that indeed in case  $r^* = 1$  both FW and our Algorithm 1 clearly exhibit a linear convergence rate. However, once we increase  $r^*$ , we see that while our Algorithm 1 maintains a linear convergence rate, FW converges with a sub-linear rate. This suggests that, similarly to the state-of-affairs in the case of optimization over polytopes [12], the standard FW does not exhibit linear convergence once the optimal solution is not an extreme point of the feasible set.



Figure 1: Comparison of Frank-Wolfe with line-search and our Algorithm 1. We set n = 100 and  $m = 15nr^*$ .

#### 4.2 Comparison of variants of Algorithm 1

We turn to provide empirical motivation for the design choices in our Algorithm 1. In particular, Algorithm 1 prioritises drop steps over other steps (even if these will reduce the objective value slower) and in addition to away steps (which are a classical concept in Frank-Wolfe-based methods [16, 21]), uses also randomized proximal pairwise steps (which also require an estimate for the smoothness parameter  $\beta$ ).

Figure 2 demonstrates how each one of the above mentioned design choices contributes significantly to the fast convergence our Algorithm 1. In particular, we see that the  $ALG \ 1$  - away variant, which is the natural extension of the *Frank-Wolfe with away steps* method and is known to converge linearly for polytopes [16, 21, 12], converges only with a sub-linear rate in our experiments.

algorithm	description
ALG 1	Algorithm 1 without modifications
ALG 1 - away	Algorithm 1 without drop steps and pairwise steps
ALG 1 - nodrop	Algorithm 1 without drop steps
ALG 1 - det	Deterministic version of Algorithm 1: instead of using the randomized vector $\mathbf{u}_{t,-}$ in the pairwise step on iteration $t$ , it is replaced with the deterministic vector $\mathbf{v}_{t,-}$ (used for the drop and away steps)

Table 2: Description of variants of Algorithm 1 used in the numerical comparison.



Figure 2: Comparison of different variants of our Algorithm 1. We set n = 100,  $r^* = 5$ , and  $m = 15nr^*$ .

#### 4.3 Comparison to the Block Frank-Wolfe method

We also compare our method to Block Frank-Wolfe methods, that is, methods that on each iteration perform an update using a rank-r matrix, for some  $r \ge r^*$ , rather than a rank-one update as in the standard Frank-Wolfe [1, 7]. We compare with the method from [1] (Block-FW in Figure 3) which performs updates of the form  $\mathbf{X}_{t+1} \leftarrow (1 - \eta)\mathbf{X}_t + \eta \mathbf{V}_t$ , where  $\mathbf{V}_t \in S^n$  is a rank-r matrix that is obtained by projecting onto  $S_n$  the best rank-r approximation (in Frobenius norm) to the matrix  $\mathbf{M}_t := \mathbf{X}_t - \frac{1}{\eta\beta}\nabla f(\mathbf{X}_t)$ , and can be computed using only the top r components in the eigen-decomposition of  $\mathbf{M}_t$ . While this method converges linearly (under quadratic growth), it requires a tight estimation of  $r^*$  (otherwise it suffers from computationally-expensive high-rank eigen-decompositions), and it requires a good estimate of both  $\alpha, \beta$  to set the step size to  $\eta = O(\alpha/\beta)$ , which is difficult to properly tune in practice. Here we consider a somewhat ideal implementation of this method: I. we set  $r = r^*$  (i.e., assume exact knowledge of  $r^*$ ), and II. we manually tune the step-size parameter  $\eta$  for best performance and set  $\eta = 0.3$ .

We can see in Figure 3 (left panel) that when comparing the (log) approximation error vs. number of iterations, as expected, the Block-FW method converges faster than our Algorithm 1 which only uses rank-one matrix computations. However, when we examine the approximation error vs. the number of rank-one updates (recall our implementation of the Block-FW method does a rank- $r^*$  update on each iteration, while our Algorithm 1 does at most a rank-two update), see right panel in Figure 3, we see that our Algorithm 1 is in fact faster.



Figure 3: Comparison of the Block-FW method and our Algorithm 1 for  $r^* = 10$ . We set n = 150 and  $m = 20nr^*$ .

#### 4.4 What if strict complementarity does not hold?

Finally, we examine what happens when we solve Problem (30) when setting  $\tau = \text{Tr}(\mathbf{X}_{\sharp}) = 1$  and without measurement noise, i.e.,  $\mathbf{b} = \mathbf{b}_{\sharp}$ . In this case, the ground truth matrix  $\mathbf{X}_{\sharp}$  is an optimal solution of Problem (30) and in particular  $\nabla f(\mathbf{X}_{\sharp}) = \mathbf{0}$ , meaning the strict complementarity condition does not hold. We see in Figure 4 that in this case, our Algorithm 1 maintains its linear convergence, while the standard Frank-Wolfe with line-search method converges with a sub-linear rate. It remains an open question if the quadratic growth condition alone suffices to prove linear convergence for our method, see also Section 5.



Figure 4: Comparison of Frank-Wolfe with line-search and our Algorithm 1 in case strict complementarity does not hold. We set n = 100,  $r^* = 3$ , and  $m = 10nr^*$ .

## 5 Future Directions

It is interesting to study whether there exists a first-order method similar to our Algorithm 1, which only relies on a constant number of efficient rank-one matrix computations per iteration, that enjoys any of the following features: I. Converges linearly (in particular without explicit dependence on the ambient dimension) without the strict complementarity assumption (though still assuming quadratic growth), II. Completely deterministic, and III. Independent of the choice of norm and in particular does not require the knowledge of the smoothness parameter  $\beta$ .

## References

- [1] Zeyuan Allen-Zhu, Elad Hazan, Wei Hu, and Yuanzhi Li. Linear convergence of a frank-wolfe type algorithm over trace-norm balls. *Advances in neural information processing systems*, 30, 2017.
- [2] Amir Beck. *First-Order Methods in Optimization*. Society for Industrial and Applied Mathematics, Philadelphia, PA, 2017.
- [3] Emmanuel J Candes, Yonina C Eldar, Thomas Strohmer, and Vladislav Voroninski. Phase retrieval via matrix completion. SIAM review, 57(2):225– 251, 2015.
- [4] Emmanuel J Candès, Xiaodong Li, Yi Ma, and John Wright. Robust principal component analysis? *Journal of the ACM (JACM)*, 58(3):1–37, 2011.
- [5] Emmanuel J Candes and Yaniv Plan. Matrix completion with noise. Proceedings of the IEEE, 98(6):925–936, 2010.
- [6] Lior Danon and Dan Garber. Frank-wolfe-based algorithms for approximating tyler's m-estimator. Advances in Neural Information Processing Systems, 35:3637–3648, 2022.
- [7] Lijun Ding, Yingjie Fei, Qiantong Xu, and Chengrun Yang. Spectral frankwolfe algorithm: Strict complementarity and linear convergence. In *International conference on machine learning*, pages 2535–2544. PMLR, 2020.
- [8] Dmitriy Drusvyatskiy and Adrian S Lewis. Error bounds, quadratic growth, and linear convergence of proximal methods. *Mathematics of Operations Re*search, 43(3):919–948, 2018.
- [9] Marguerite Frank, Philip Wolfe, et al. An algorithm for quadratic programming. Naval research logistics quarterly, 3(1-2):95–110, 1956.
- [10] Robert M Freund, Paul Grigas, and Rahul Mazumder. An extended frankwolfe method with "in-face" directions, and its application to low-rank matrix completion. SIAM Journal on optimization, 27(1):319–346, 2017.
- [11] Dan Garber. Faster projection-free convex optimization over the spectrahedron. In D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 29. Curran Associates, Inc., 2016.
- [12] Dan Garber. Revisiting frank-wolfe for polytopes: Strict complementarity and sparsity. Advances in Neural Information Processing Systems, 33:18883–18893, 2020.
- [13] Dan Garber. On the convergence of projected-gradient methods with low-rank projections for smooth convex minimization over trace-norm balls and related problems. SIAM Journal on Optimization, 31(1):727–753, 2021.

- [14] Dan Garber. Linear convergence of frank-wolfe for rank-one matrix recovery without strong convexity. *Mathematical Programming*, 199(1):87–121, 2023.
- [15] Gauthier Gidel, Fabian Pedregosa, and Simon Lacoste-Julien. Frank-wolfe splitting via augmented lagrangian method. In *International Conference on Artificial Intelligence and Statistics*, pages 1456–1465. PMLR, 2018.
- [16] Jacques Guélat and Patrice Marcotte. Some comments on wolfe's 'away step'. Mathematical Programming, 35(1):110–119, 1986.
- [17] Elad Hazan. Sparse approximate solutions to semidefinite programs. In *Latin* American symposium on theoretical informatics, pages 306–316. Springer, 2008.
- [18] Martin Jaggi. Convex optimization without projection steps. arXiv preprint arXiv:1108.1170, 2011.
- [19] Martin Jaggi. Revisiting frank-wolfe: Projection-free sparse convex optimization. In International conference on machine learning, pages 427–435. PMLR, 2013.
- [20] Martin Jaggi, Marek Sulovsk, et al. A simple algorithm for nuclear norm regularized problems. In Proceedings of the 27th international conference on machine learning (ICML-10), pages 471–478, 2010.
- [21] Simon Lacoste-Julien and Martin Jaggi. On the global linear convergence of frank-wolfe optimization variants. Advances in neural information processing systems, 28, 2015.
- [22] Evgeny S Levitin and Boris T Polyak. Constrained minimization methods. USSR Computational mathematics and mathematical physics, 6(5):1–50, 1966.
- [23] Carl D Meyer, Jr. Generalized inversion of modified matrices. Siam journal on applied mathematics, 24(3):315–323, 1973.
- [24] Ion Necoara, Yu Nesterov, and Francois Glineur. Linear convergence of first order methods for non-strongly convex optimization. *Mathematical Program*ming, 175:69–107, 2019.
- [25] Chi Bach Pham, Wynita Griggs, and James Saunderson. A scalable frankwolfe-based algorithm for the max-cut sdp. In *International Conference on Machine Learning*, pages 27822–27839. PMLR, 2023.
- [26] Nathan Srebro, Jason Rennie, and Tommi Jaakkola. Maximum-margin matrix factorization. Advances in neural information processing systems, 17, 2004.
- [27] Ying Sun, Prabhu Babu, and Daniel P Palomar. Robust estimation of structured covariance matrix for heavy-tailed elliptical distributions. *IEEE Trans*actions on Signal Processing, 64(14):3576–3590, 2016.

- [28] Joel A Tropp. Convex recovery of a structured signal from independent random linear measurements. Sampling theory, a renaissance: compressive sensing and other developments, pages 67–101, 2015.
- [29] Ami Wiesel. Geodesic convexity and covariance estimation. *IEEE transactions* on signal processing, 60(12):6182–6189, 2012.
- [30] Alp Yurtsever, Olivier Fercoq, and Volkan Cevher. A conditional-gradientbased augmented lagrangian framework. In *International Conference on Machine Learning*, pages 7272–7281. PMLR, 2019.
- [31] Alp Yurtsever, Madeleine Udell, Joel Tropp, and Volkan Cevher. Sketchy decisions: Convex low-rank matrix optimization with optimal storage. In Artificial intelligence and statistics, pages 1188–1196. PMLR, 2017.
- [32] Zirui Zhou and Anthony Man-Cho So. A unified approach to error bounds for structured convex optimization problems. *Mathematical Programming*, 165:689–728, 2017.

## A Proof of Lemma 2

We first restate the lemma and then prove it.

**Lemma 8.** Let  $\mathbf{X} \in \mathbb{S}^n_+$  and  $\mathbf{v} \in Im(\mathbf{X})$ . Consider the matrix  $\mathbf{Y} = \mathbf{X} - \lambda \mathbf{v} \mathbf{v}^\top$  for some  $\lambda \geq 0$ . If  $\lambda \leq (\mathbf{v}^\top \mathbf{X}^\dagger \mathbf{v})^{-1}$ , then  $\mathbf{Y} \succeq 0$ . Moreover, if  $\lambda = (\mathbf{v}^\top \mathbf{X}^\dagger \mathbf{v})^{-1}$ , then  $rank(\mathbf{Y}) < rank(\mathbf{X})$  and  $\mathbf{X}^\dagger \mathbf{v} \in Ker(\mathbf{Y})$ .

Proof. It holds that,

$$1 - \lambda \mathbf{v}^{\top} \mathbf{X}^{\dagger} \mathbf{v} \ge 0 \Longrightarrow 1 - \lambda_{1} \left( \lambda \mathbf{X}^{\dagger 1/2} \mathbf{v} \mathbf{v}^{\top} \mathbf{X}^{\dagger 1/2} \right) \ge 0$$
$$\Longrightarrow \lambda_{n} \left( \mathbf{I} - \lambda \mathbf{X}^{\dagger 1/2} \mathbf{v} \mathbf{v}^{\top} \mathbf{X}^{\dagger 1/2} \right) \ge 0$$
$$\Longrightarrow \mathbf{I} - \lambda \mathbf{X}^{\dagger 1/2} \mathbf{v} \mathbf{v}^{\top} \mathbf{X}^{\dagger 1/2} \succeq 0$$
$$\Longrightarrow \mathbf{X}^{1/2} \left( \mathbf{I} - \lambda \mathbf{X}^{\dagger 1/2} \mathbf{v} \mathbf{v}^{\top} \mathbf{X}^{\dagger 1/2} \right) \mathbf{X}^{1/2} \succeq 0$$
$$\Longrightarrow \mathbf{X} - \lambda \mathbf{v} \mathbf{v}^{\top} \succeq 0,$$

where (a) relies on the assumption that  $\mathbf{v} \in \text{Im}(\mathbf{X})$  and so  $\mathbf{X}^{1/2}\mathbf{X}^{\dagger^{1/2}}\mathbf{v} = \mathbf{v}$ .

We continue to prove the second part of the lemma. Suppose now that  $\lambda = (\mathbf{v}^{\top} \mathbf{X}^{\dagger} \mathbf{v})^{-1}$ . Note that since  $\mathbf{v} \in \text{Im}(\mathbf{X})$ , we have that  $\text{Im}(\mathbf{Y}) \subseteq \text{Im}(\mathbf{X})$ . Consider now the vector  $\mathbf{u} = \mathbf{X}^{\dagger} \mathbf{v} \in \text{Im}(\mathbf{X})$ . It holds that,

$$\mathbf{Y}\mathbf{u} = \left(\mathbf{X} - \frac{1}{\mathbf{v}^{\top}\mathbf{X}^{\dagger}\mathbf{v}}\mathbf{v}\mathbf{v}^{\top}\right)\mathbf{X}^{\dagger}\mathbf{v} = \mathbf{X}\mathbf{X}^{\dagger}\mathbf{v} - \mathbf{v} = 0,$$

implying that  $\mathbf{u} \notin \text{Im}(\mathbf{Y})$  and so,  $\text{rank}(\mathbf{Y}) < \text{rank}(\mathbf{X})$ .

## **B** Fast Update of the Pseudo Inverse Matrix

The following lemma is based on [23]. While [23] considers a more general case of non-square complex matrices, the result for symmetric real matrices can be significantly simplified.

**Lemma 9.** Let  $\mathbf{A} \in \mathbb{S}^n$  and let  $\mathbf{c}, \mathbf{d}$  be linearly dependent vectors in  $\mathbb{R}^n$ . Define the following:  $\mathbf{k} = \mathbf{A}^{\dagger}\mathbf{c}$  (column vector),  $\mathbf{h} = \mathbf{d}^{\top}\mathbf{A}^{\dagger}$  (row vector),  $\mathbf{u} = (\mathbf{I} - \mathbf{A}\mathbf{A}^{\dagger})\mathbf{c}$ (column vector),  $\mathbf{v} = \mathbf{d}^{\top}(\mathbf{I} - \mathbf{A}^{\dagger}\mathbf{A})$  (row vector), and  $\zeta = 1 + \mathbf{d}^{\top}\mathbf{A}^{\dagger}\mathbf{c}$  (scalar). Additionally, if  $\zeta \neq 0$ , define:  $\mathbf{p} = -\frac{\|\mathbf{k}\|_2^2}{\zeta}\mathbf{v}^{\top} - \mathbf{k}$  (column vector),  $\mathbf{q} = \frac{\|\mathbf{v}\|_2^2}{\zeta}\mathbf{A}^{\dagger}\mathbf{k} - \mathbf{h}^{\top}$ (column vector), and  $\sigma = \|\mathbf{k}\|_2^2\|\mathbf{v}\|_2^2 + \zeta^2$ . Then, it holds that

$$\mathbf{B}^{\dagger} = \begin{cases} \mathbf{A}^{\dagger} - \mathbf{k}\mathbf{u}^{\dagger} - \mathbf{v}^{\dagger}\mathbf{h} + \zeta\mathbf{v}^{\dagger}\mathbf{u}^{\dagger} & \text{if } \mathbf{c} \notin Im(\mathbf{A}); \\ \mathbf{A}^{\dagger} + \frac{1}{\zeta}\mathbf{v}^{\top}\mathbf{k}^{\top}\mathbf{A}^{\dagger} - \frac{\zeta}{\sigma}\mathbf{p}\mathbf{q}^{\top} & \text{if } \mathbf{c} \in Im(\mathbf{A}) \text{ and } \zeta \neq 0; \\ \mathbf{A}^{\dagger} - \mathbf{k}\mathbf{k}^{\dagger}\mathbf{A}^{\dagger} - \mathbf{A}^{\dagger}\mathbf{h}^{\dagger}\mathbf{h} + (\mathbf{k}^{\dagger}\mathbf{A}^{\dagger}\mathbf{h}^{\dagger})\mathbf{k}\mathbf{h} & \text{if } \mathbf{c} \in Im(\mathbf{A}) \text{ and } \zeta = 0. \end{cases}$$

In particular, given  $\mathbf{A}^{\dagger}$ , one can compute  $\mathbf{B}^{\dagger}$  in  $O(n^2)$  time.