

PLANAR LATTICES AND EQUILATERAL ODD-GONS

AKIRA IINO AND MASASHI SAKIYAMA

ABSTRACT. For a planar integral lattice L , let $\nu(L)$ denote the square-free part of the integer $D(L)^2$, where $D(L)$ stands for the area of a fundamental parallelogram of L . For each odd integer n with $3 \leq n < 29$, a planar lattice L contains an equilateral n -gon if and only if L is similar to an integral lattice L' such that $\nu(L') \equiv 3 \pmod{4}$ and the largest prime factor p of $\nu(L')$ satisfies $p \leq n$. Moreover, such L contains a convex equilateral n -gon, which answers a problem posed by Maehara.

1. INTRODUCTION

In 1973, Ball [1] considered the existence of equilateral polygons with vertices on the square lattice and proved the following.

Theorem 1.1 ([1, Theorems 1 and 6]). *The square lattice \mathbb{Z}^2 does not contain an equilateral n -gon if n is odd. The lattice \mathbb{Z}^2 contains a convex equilateral n -gon if n is even.*

Maehara [2, 3] considered the existence of equilateral polygons with vertices on a general planar lattice. He showed many interesting results, some of which are listed below. (The definitions of some terms will be explained in the next section.)

Theorem 1.2 ([3, Theorem 5.1]). *A planar lattice L contains a convex equilateral n -gon for some $n \neq 4$ if and only if L is similar to an integral lattice.*

Theorem 1.3 ([3, Theorem 4.1]). *Every planar integral lattice L contains a convex equilateral n -gon for every even $n \geq 4$. A planar integral lattice L contains an equilateral n -gon for some odd $n \geq 3$ if and only if $\nu(L) \equiv 3 \pmod{4}$.*

Theorem 1.4 ([3, Theorem 6.1]). *For a planar lattice L , the following three are equivalent. (i) L contains an equilateral triangle. (ii) L contains a convex equilateral n -gon for every $n \geq 3$. (iii) L is similar to an integral lattice L' with $\nu(L') = 3$.*

From Theorems 1.2 and 1.3, we see that for each even $n \neq 4$, a planar lattice L contains an equilateral n -gon if and only if L is similar to an integral lattice. In this paper, we consider the following problem.

Problem 1.5. *Let n be an odd integer with $n \geq 3$. Find the condition on a planar lattice L so that L contains an equilateral n -gon.*

Theorem 1.4 answers Problem 1.5 in the case where $n = 3$. In this paper, we extend Theorem 1.4 and answer Problem 1.5 when n is small. We also give an affirmative answer to the following problem, which is posed by Maehara [2, 3].

Date: March 5, 2025.

2020 Mathematics Subject Classification. 52C05, 11H06, 51M04.

Key words and phrases. planar lattice, integral lattice, equilateral polygon.

Problem 1.6 ([3, Problem 1.1]). *Is there a planar integral lattice L with $\nu(L) \neq 3$ that contains a convex equilateral n -gon for some odd $n > 3$?*

2. PRELIMINARIES

In this paper, an element of a Euclidean plane \mathbb{R}^2 is called a *vector* as well as a *point*. For linearly independent vectors \mathbf{a}, \mathbf{b} in the plane \mathbb{R}^2 , $L[\mathbf{a}, \mathbf{b}]$ denotes the *planar lattice* generated by \mathbf{a}, \mathbf{b} , that is, $L[\mathbf{a}, \mathbf{b}] = \{m\mathbf{a} + n\mathbf{b} \mid m, n \in \mathbb{Z}\} \subset \mathbb{R}^2$. If a lattice L is contained in another lattice L' , then L is said to be a *sublattice* of L' . A lattice L is said to be *similar* to L' if there is a $\lambda > 0$ such that $\lambda L = \{\lambda\mathbf{x} \mid \mathbf{x} \in L\}$ is isometric to L' . A planar lattice L is called an *integral lattice*, if for every $\mathbf{x}, \mathbf{y} \in L$, the inner product $\mathbf{x} \cdot \mathbf{y}$ is an integer.

For a planar lattice L , let $D(L)$ denote the area of a fundamental parallelogram of the lattice L . Thus, if $L = L[\mathbf{a}, \mathbf{b}]$, then $D(L) = |\det(\mathbf{a}, \mathbf{b})|$. Although there are many choices of vectors \mathbf{a}, \mathbf{b} that generate the planar lattice L , the area $D(L) = |\det(\mathbf{a}, \mathbf{b})|$ is independent of the choice of generating vectors \mathbf{a}, \mathbf{b} of L . If L is a planar integral lattice, then $D(L)^2$ is an integer. In fact, if $L = L[\mathbf{a}, \mathbf{b}]$, then

$$D(L)^2 = \det(\mathbf{a}, \mathbf{b})^2 = \begin{vmatrix} \mathbf{a} \cdot \mathbf{a} & \mathbf{a} \cdot \mathbf{b} \\ \mathbf{b} \cdot \mathbf{a} & \mathbf{b} \cdot \mathbf{b} \end{vmatrix} \in \mathbb{Z}.$$

Let us denote the square-free part of $D(L)^2$ by $\nu(L)$, that is, $\nu(L)$ is the square-free integer that satisfies $D(L)^2 = k^2\nu(L)$ for some integer k .

For a square-free integer $m > 0$, we denote the rectangular lattice $L[(1, 0), (0, \sqrt{m})]$ by $\Lambda(m)$.

An *equilateral polygon* is a polygon whose sides are all equal in length. If every vertex of a polygon P is in a set S , then we say S *contains* the polygon P .

For an equilateral n -gon $A_1A_2 \dots A_n$, the vectors $\mathbf{e}_1 = \overrightarrow{A_nA_1}$, $\mathbf{e}_2 = \overrightarrow{A_1A_2}$, \dots , $\mathbf{e}_n = \overrightarrow{A_{n-1}A_n}$ are called the *edge-vectors* of the polygon. Clearly $\mathbf{e}_1 + \mathbf{e}_2 + \dots + \mathbf{e}_n = \mathbf{0}$ and $|\mathbf{e}_1| = |\mathbf{e}_2| = \dots = |\mathbf{e}_n|$.

Maehara [2, 3] proved the following results. We will use them in the next section.

Lemma 2.1 ([2, Lemma 3]). *A planar lattice L contains a convex equilateral n -gon if and only if L contains n distinct vectors $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ such that $|\mathbf{e}_1| = |\mathbf{e}_2| = \dots = |\mathbf{e}_n|$ and $\mathbf{e}_1 + \mathbf{e}_2 + \dots + \mathbf{e}_n = \mathbf{0}$.*

Lemma 2.2 ([3, Corollary 4.1]). *A planar integral lattice L contains an equilateral n -gon (resp. a convex equilateral n -gon) if and only if $\Lambda(\nu(L))$ contains an equilateral n -gon (resp. a convex equilateral n -gon).*

Lemma 2.3 ([3, Lemma 4.1]). *If $\Lambda(m)$ contains an equilateral n -gon (resp. a convex equilateral n -gon), then it contains an equilateral $(n+2)$ -gon (resp. a convex equilateral $(n+2)$ -gon).*

3. RESULTS

Let us consider the condition on a planar lattice L so that L contains an equilateral n -gon (or a convex equilateral n -gon) for each odd n . First, we prove the following theorem.

Theorem 3.1. *Let $n \geq 3$ be an odd number. If a planar integral lattice L contains an equilateral n -gon, not necessarily convex, then $n \geq p$ for every prime factor p of $\nu(L)$.*

Proof. Assume that L contains an equilateral n -gon and p is a prime factor of $\nu(L)$. By Lemma 2.2, $\Lambda(\nu)$ with $\nu = \nu(L)$ contains an equilateral n -gon. Let P be one of the n -gons contained in $\Lambda(\nu)$ and let $\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n$ be the edge-vectors of P . Let $\mathbf{e}_1 = (s, t\sqrt{\nu})$, where $s, t \in \mathbb{Z}$. By the linear transformation f induced by the matrix $\begin{pmatrix} s & t\sqrt{\nu} \\ -t\sqrt{\nu} & s \end{pmatrix}$, the vector \mathbf{e}_1 is mapped to $f(\mathbf{e}_1) = (s^2 + t^2\nu, 0)$. Since $\begin{pmatrix} s & t\sqrt{\nu} \\ -t\sqrt{\nu} & s \end{pmatrix} \begin{pmatrix} a \\ b\sqrt{\nu} \end{pmatrix} = \begin{pmatrix} sa+tb\nu \\ (-ta+sb)\sqrt{\nu} \end{pmatrix}$, f maps $\Lambda(\nu)$ into $\Lambda(\nu)$. Hence the vectors $f(\mathbf{e}_1), f(\mathbf{e}_2), \dots, f(\mathbf{e}_n)$ are in $\Lambda(\nu)$. Since f is a similarity, the vectors $f(\mathbf{e}_1), f(\mathbf{e}_2), \dots, f(\mathbf{e}_n)$ determines an equilateral n -gon P' , which is similar to P . Since the side length of P' is the integer $s^2 + t^2\nu$, there exists an equilateral n -gon contained in $\Lambda(\nu)$ whose side length is an integer.

Let Q be one of the equilateral n -gons contained in $\Lambda(\nu)$ whose side length is the minimal integer, and let $\mathbf{v}_1, \dots, \mathbf{v}_n$ be the edge-vectors of Q . Let $\mathbf{v}_i = (a_i, b_i\sqrt{\nu})$ ($a_i, b_i \in \mathbb{Z}$) and let k be the side length of Q . Since each side has length k , we have

$$a_i^2 + b_i^2\nu = k^2 \quad (1)$$

for each $i \in \{1, 2, \dots, n\}$. Assume that k is a multiple of p . Since k and ν are multiples of p , it follows from equation (1) that a_i^2 is a multiple of p for each i . Since p is prime, a_i is a multiple of p . Now, since a_i^2 and k^2 are multiples of p^2 , it follows again from equation (1) that $b_i^2\nu$ is a multiple of p^2 . Since ν is square-free, ν is not a multiple of p^2 . Hence b_i^2 must be a multiple of p , and so is b_i . Now since $a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_n, k$ are multiples of p , $(1/p)\mathbf{v}_i$ is a vector in $\Lambda(\nu)$ for each i , and hence an n -gon determined by the edge-vectors $(1/p)\mathbf{v}_1, \dots, (1/p)\mathbf{v}_n$ can be contained in $\Lambda(\nu)$ and has side length k/p , which is an integer smaller than k . This contradicts the minimality of k . Therefore k is not a multiple of p .

Since ν is a multiple of p , it follows from (1) that $a_i^2 \equiv k^2 \pmod{p}$, and hence $(a_i - k)(a_i + k) \equiv 0 \pmod{p}$. Since p is prime, either $a_i \equiv k \pmod{p}$ or $a_i \equiv -k \pmod{p}$ holds. Now let $n' = \#\{i \mid a_i \equiv k \pmod{p}\}$. We have $\#\{i \mid a_i \not\equiv k \pmod{p}\} = n - n'$. Since $i \in \{i \mid a_i \not\equiv k \pmod{p}\}$ implies $a_i \equiv -k \pmod{p}$, we have $\sum_{i=1}^n a_i \equiv n'k + (n - n')(-k) \equiv (2n' - n)k \pmod{p}$. Since $\sum_{i=1}^n \mathbf{v}_i = \mathbf{0}$, we have $\sum_{i=1}^n a_i = 0$, and hence $\sum_{i=1}^n a_i \equiv 0 \pmod{p}$. Now we have $(2n' - n)k \equiv 0 \pmod{p}$. Since k is not a multiple of p , $2n' - n$ must be a multiple of p . Since $0 \leq n' \leq n$, we have $-n \leq 2n' - n \leq n$. Since n is odd, we have $2n' - n \neq 0$. Therefore, the interval $[-n, n]$ contains a nonzero multiple of p , which shows $n \geq p$. \square

From Theorems 1.3 and 3.1, we know that for each odd integer $n \geq 3$, for a planar integral lattice L to contain an equilateral n -gon it is necessary that $\nu(L) \equiv 3 \pmod{4}$ and the largest prime factor p of $\nu(L)$ satisfies $p \leq n$. Is this condition also sufficient? We will prove that the answer is yes if $n < 17$.

Proposition 3.2. *Let m be a square-free positive integer with $m \equiv 3 \pmod{4}$ and let p be the largest prime factor of m . If $p < 17$, then the rectangular lattice $\Lambda(m)$ contains a convex equilateral p -gon.*

Proof. Each square-free positive integer m satisfying $p < 17$ is expressed as $m = 2^a \cdot 3^b \cdot 5^c \cdot 7^d \cdot 11^e \cdot 13^f$, where $a, b, c, d, e, f \in \{0, 1\}$. Since $2^a \cdot 3^b \cdot 5^c \cdot 7^d \cdot 11^e \cdot 13^f \equiv 2^a \cdot (-1)^b \cdot 1^c \cdot (-1)^d \cdot (-1)^e \cdot 1^f \equiv 2^a \cdot (-1)^{b+d+e} \pmod{4}$, m of this form satisfies $m \equiv 3 \pmod{4}$ if and only if $a = 0$ and $b + d + e \equiv 1 \pmod{2}$. Hence it is possible to list all

of the square-free positive integers m satisfying $p < 17$ as in Table 1. For each of such m , $\Lambda(m)$ contains p distinct vectors of equal length with sum $\mathbf{0}$, which are illustrated in Table 1. From Lemma 2.1, we conclude that $\Lambda(m)$ contains a convex equilateral p -gon. \square

TABLE 1. square-free m with $m \equiv 3 \pmod{4}$ and $p < 17$, and p distinct vectors of equal length in $\Lambda(m)$ with sum $\mathbf{0}$

m	prime factors	p	p distinct vectors of equal length in $\Lambda(m)$ with sum $\mathbf{0}$
3	3	3	$(2, 0), (-1, \pm\sqrt{m})$
15	3, 5	5	$(483724, 0), (445129, 48887\sqrt{m}), (-379901, 77315\sqrt{m}),$ $(-483631, -2449\sqrt{m}), (-65321, -123753\sqrt{m})$
7	7	7	$(88, 0), (81, \pm 13\sqrt{m}), (-38, \pm 30\sqrt{m}), (-87, \pm 5\sqrt{m})$
35	5, 7	7	$(17226, 0), (13194, \pm 1872\sqrt{m}), (-4726, \pm 2800\sqrt{m}),$ $(-17081, \pm 377\sqrt{m})$
11	11	11	$(90, 0), (57, \pm 21\sqrt{m}), (35, \pm 25\sqrt{m}), (-9, \pm 27\sqrt{m}),$ $(-42, \pm 24\sqrt{m}), (-86, \pm 8\sqrt{m})$
55	5, 11	11	$(728, 0), (717, \pm 17\sqrt{m}), (552, \pm 64\sqrt{m}), (-273, \pm 91\sqrt{m}),$ $(-658, \pm 42\sqrt{m}), (-702, \pm 26\sqrt{m})$
231	3, 7, 11	11	$(800, 0), (569, \pm 37\sqrt{m}), (415, \pm 45\sqrt{m}), (-124, \pm 52\sqrt{m}),$ $(-520, \pm 40\sqrt{m}), (-740, \pm 20\sqrt{m})$
1155	3, 5, 7, 11	11	$(22678, 0), (22447, \pm 95\sqrt{m}), (3967, \pm 657\sqrt{m}),$ $(-5273, \pm 649\sqrt{m}), (-15283, \pm 493\sqrt{m}),$ $(-17197, \pm 435\sqrt{m})$
39	3, 13	13	$(440, 0), (401, \pm 29\sqrt{m}), (310, \pm 50\sqrt{m}), (154, \pm 66\sqrt{m}),$ $(-275, \pm 55\sqrt{m}), (-392, \pm 32\sqrt{m}), (-418, \pm 22\sqrt{m})$
195	3, 5, 13	13	$(1666, 0), (1601, \pm 33\sqrt{m}), (1406, \pm 64\sqrt{m}), (119, \pm 119\sqrt{m}),$ $(-791, \pm 105\sqrt{m}), (-1519, \pm 49\sqrt{m}), (-1649, \pm 17\sqrt{m})$
91	7, 13	13	$(5890, 0), (5877, \pm 41\sqrt{m}), (3875, \pm 465\sqrt{m}),$ $(-402, \pm 616\sqrt{m}), (-1767, \pm 589\sqrt{m}), (-5043, \pm 319\sqrt{m}),$ $(-5485, \pm 225\sqrt{m})$
455	5, 7, 13	13	$(4104, 0), (3519, \pm 99\sqrt{m}), (3064, \pm 128\sqrt{m}), (646, \pm 190\sqrt{m}),$ $(-2214, \pm 162\sqrt{m}), (-3306, \pm 114\sqrt{m}), (-3761, \pm 77\sqrt{m})$
143	11, 13	13	$(6384, 0), (4902, \pm 342\sqrt{m}), (3472, \pm 448\sqrt{m}),$ $(-532, \pm 532\sqrt{m}), (-2391, \pm 495\sqrt{m}), (-2534, \pm 490\sqrt{m}),$ $(-6109, \pm 155\sqrt{m})$
715	5, 11, 13	13	$(571778, 0), (492153, \pm 10885\sqrt{m}), (424943, \pm 14307\sqrt{m}),$ $(353157, \pm 16817\sqrt{m}), (-472382, \pm 12048\sqrt{m}),$ $(-526722, \pm 8320\sqrt{m}), (-557038, \pm 4824\sqrt{m})$
3003	3, 7, 11, 13	13	$(4496798, 0), (4468081, \pm 9259\sqrt{m}), (4017631, \pm 36859\sqrt{m}),$ $(-1091473, \pm 79605\sqrt{m}), (-1638877, \pm 76415\sqrt{m}),$ $(-3823202, \pm 43200\sqrt{m}), (-4180559, \pm 30229\sqrt{m})$
15015	3, 5, 7, 11, 13	13	$(456688, 0), (446522, \pm 782\sqrt{m}), (323828, \pm 2628\sqrt{m}),$ $(21097, \pm 3723\sqrt{m}), (-198122, \pm 3358\sqrt{m}),$ $(-373297, \pm 2147\sqrt{m}), (-448372, \pm 708\sqrt{m})$

Remark 3.3. The vectors illustrated in Table 1 can be found in a quite simple search, using a computer, which we explain here. To obtain p distinct vectors of equal length contained in $\Lambda(m)$ with sum $\mathbf{0}$, first we search for p distinct unit vectors contained in $\mathbb{Q} \times \sqrt{m}\mathbb{Q} = \{(s, t\sqrt{m}) \mid s, t \in \mathbb{Q}\}$ with sum $\mathbf{0}$. For each positive integer N , let $U(m)_N$ be the set $\{(a/c, b\sqrt{m}/c) \mid a, b, c \in \mathbb{Z}, 1 \leq c \leq N, a^2 + b^2m = c^2\}$, which is consisting of unit vectors in $\mathbb{Q} \times \sqrt{m}\mathbb{Q}$. For a fixed N , we use a computer to search for p distinct

vectors in $U(m)_N$ whose sum is $\mathbf{0}$. (An exhaustive search is possible since $U(m)_N$ is a finite set.) By setting N by $N = 10^7$, we can find desired unit vectors. By multiplying them by an appropriate integer, we obtain p distinct vectors of equal length in $\Lambda(m)$ with sum $\mathbf{0}$.

Since $\nu(\Lambda(m)) = m$ for a square-free positive integer m , Proposition 3.2 answers Problem 1.6 in the affirmative. It also gives us the following corollary.

Corollary 3.4. *If a planar integral lattice L satisfies $\nu(L) \equiv 3 \pmod{4}$ and the largest prime factor p of $\nu(L)$ satisfies $p < 17$, then L contains a convex equilateral n -gon for every integer n with $n \geq p$.*

Proof. Suppose that a planar integral lattice L satisfies $\nu(L) \equiv 3 \pmod{4}$ and the largest prime factor p of $\nu(L)$ satisfies $p < 17$. Then, from Proposition 3.2, $\Lambda(\nu(L))$ contains a convex equilateral p -gon. Hence by Lemma 2.2, L contains a convex equilateral p -gon. By Lemma 2.3, L contains a convex equilateral n -gon for every odd $n \geq p$. Since L contains a convex equilateral n -gon for every even n by Theorem 1.3, the assertion follows. \square

Now we can prove the following theorem, which generalizes Theorem 1.4.

Theorem 3.5. *Let n be an odd integer with $3 \leq n < 17$. For a planar lattice L , the following three are equivalent.*

- (i) L contains an equilateral n -gon.
- (ii) L contains a convex equilateral k -gon for every $k \geq n$.
- (iii) L is similar to an integral lattice L' such that $\nu(L') \equiv 3 \pmod{4}$ and the largest prime factor p of $\nu(L')$ satisfies $p \leq n$.

Proof. (ii) \Rightarrow (i): Obvious.

(i) \Rightarrow (iii): Suppose that (i) holds. By Theorem 1.2, L is similar to an integral lattice L' . Since L' contains an equilateral n -gon, $\nu(L') \equiv 3 \pmod{4}$ from Theorem 1.3, and the largest prime factor p of $\nu(L')$ satisfies $p \leq n$ from Theorem 3.1.

(iii) \Rightarrow (ii): Suppose that (iii) holds. Since L' satisfies $\nu(L') \equiv 3 \pmod{4}$ and the largest prime factor p of $\nu(L')$ satisfies $p \leq n (< 17)$, L' contains a convex equilateral k -gon for every $k \geq p$ from Corollary 3.4. Since $p \leq n$, L' contains a convex equilateral k -gon for every $k \geq n$. Since L is similar to L' , L contains a convex equilateral k -gon for every $k \geq n$. \square

Theorem 3.5 answers Problem 1.5 in the case where $n < 17$.

Remark 3.6. Proposition 3.2 remains true if we replace “If $p < 17$ ” by “If $p < 29$ ”. To confirm this fact, it is sufficient to search for p distinct unit vectors in $\mathbb{Q} \times \sqrt{m}\mathbb{Q}$ with sum $\mathbf{0}$ as in Remark 3.3 with setting N by $N = 10^{14}$. (Table 2 illustrates p distinct vectors of equal length in $\Lambda(m)$ with sum $\mathbf{0}$ for some m .) From this, we can see that Theorem 3.5 holds for odd n with $3 \leq n < 29$, which answers Problem 1.5 in the case where $n < 29$.

For general n , Problem 1.5 is unsolved.

Acknowledgement. The authors would like to thank Hiroshi Maehara and the referee for their helpful advice.

TABLE 2. p distinct vectors of equal length in $\Lambda(m)$ with sum $\mathbf{0}$ for some m with $m \equiv 3 \pmod{4}$ and $p < 29$

m	prime factors	p	p distinct vectors of equal length in $\Lambda(m)$ with sum $\mathbf{0}$
51	3, 17	17	$(1430, 0), (1243, \pm 99\sqrt{m}), (971, \pm 147\sqrt{m}),$ $(325, \pm 195\sqrt{m}), (70, \pm 200\sqrt{m}),$ $(-406, \pm 192\sqrt{m}), (-695, \pm 175\sqrt{m}),$ $(-1001, \pm 143\sqrt{m}), (-1222, \pm 104\sqrt{m})$
255255	3, 5, 7, 11, 13, 17	17	$(41516416, 0), (37725824, \pm 34304\sqrt{m}),$ $(20300416, \pm 71680\sqrt{m}),$ $(16021856, \pm 75808\sqrt{m}),$ $(13864624, \pm 77456\sqrt{m}),$ $(7027156, \pm 80988\sqrt{m}),$ $(-34068899, \pm 46961\sqrt{m}),$ $(-40195019, \pm 20567\sqrt{m}),$ $(-41434166, \pm 5170\sqrt{m})$
19	19	19	$(770, 0), (751, \pm 39\sqrt{m}), (675, \pm 85\sqrt{m}),$ $(561, \pm 121\sqrt{m}), (238, \pm 168\sqrt{m}),$ $(-275, \pm 165\sqrt{m}), (-427, \pm 147\sqrt{m}),$ $(-446, \pm 144\sqrt{m}), (-693, \pm 77\sqrt{m}),$ $(-769, \pm 9\sqrt{m})$
1616615	5, 7, 11, 13, 17, 19	19	$(1306809216, 0), (1306710264, \pm 12648\sqrt{m}),$ $(1099481216, \pm 555520\sqrt{m}),$ $(971630064, \pm 687312\sqrt{m}),$ $(43693054, \pm 1027226\sqrt{m}),$ $(-138916611, \pm 1021977\sqrt{m}),$ $(-694638111, \pm 870573\sqrt{m}),$ $(-863551284, \pm 771420\sqrt{m}),$ $(-1071771296, \pm 588064\sqrt{m}),$ $(-1306041904, \pm 35216\sqrt{m})$
23	23	23	$(1872, 0), (1826, \pm 86\sqrt{m}), (1803, \pm 105\sqrt{m}),$ $(1044, \pm 324\sqrt{m}), (676, \pm 364\sqrt{m}),$ $(78, \pm 390\sqrt{m}), (-336, \pm 384\sqrt{m}),$ $(-819, \pm 351\sqrt{m}), (-911, \pm 341\sqrt{m}),$ $(-1072, \pm 320\sqrt{m}), (-1509, \pm 231\sqrt{m}),$ $(-1716, \pm 156\sqrt{m})$
111546435	3, 5, 7, 11, 13, 17, 19, 23	23	$(46923183273602, 0),$ $(46918331222398, \pm 63889920\sqrt{m}),$ $(46179809577838, \pm 787694056\sqrt{m}),$ $(45535252265413, \pm 1072579179\sqrt{m}),$ $(32540987084087, \pm 3200887671\sqrt{m}),$ $(1360169729983, \pm 4440962273\sqrt{m}),$ $(-18098964883063, \pm 4099034551\sqrt{m}),$ $(-21535244754542, \pm 3947292712\sqrt{m}),$ $(-33327326200258, \pm 3127514608\sqrt{m}),$ $(-36400586649517, \pm 2803613027\sqrt{m}),$ $(-41444796116242, \pm 2083271992\sqrt{m}),$ $(-45189222912898, \pm 1196604960\sqrt{m})$

REFERENCES

- [1] Ball, D. G.: The constructibility of regular and equilateral polygons on a square pinboard. *Math. Gaz.* **57**, 119–122 (1973).
- [2] Maehara, H.: Equilateral polygons and planar integral lattices. *Educ. Dev.* **6**, 121–132 (2012).
- [3] Maehara, H.: Planar lattices and equilateral polygons. *Eur. J. Comb.* **80**, 277–286 (2019).

NIPPON HYORON SHA, CO., LTD., 3-12-4 MINAMI-OTSUKA, TOSHIMA-KU, TOKYO, 170-8474,
JAPAN

Email address: iino@nippyo.co.jp

THE KAISEI ACADEMY, 4-2-4 NISHI-NIPPORI, ARAKAWA-KU, TOKYO, 116-0013, JAPAN

Email address: sakiyama-ms@kaiseigakuen.jp