

Nonreciprocity of hydrodynamic electron transport in noncentrosymmetric conductors

E. Kirkinis,¹ L. Bonds,² A. Levchenko,³ and A. V. Andreev²

¹*Center for Computation and Theory of Soft Materials,
Robert R. McCormick School of Engineering and Applied Science,
Northwestern University, Evanston IL 60208 USA*

²*Department of Physics, University of Washington, Seattle, Washington 98195, USA*

³*Department of Physics, University of Wisconsin-Madison, Madison, Wisconsin 53706, USA*

(Dated: March 3, 2025)

We show that the nonreciprocity of hydrodynamic electron transport in noncentrosymmetric conductors with broken time-reversal symmetry (TRS) is significantly enhanced compared to the disorder-dominated regime. This enhancement is caused by the linear dependence of the viscosity of the electron liquid on the flow velocity, which is allowed in the absence of TRS and Galilean invariance. The resulting nonlinear flows do not possess dynamical similarity and are characterized by two dimensionless parameters: the Reynolds number and the nonreciprocity number. The latter is linear in velocity but independent of system size. We determine the nonlinear conductance of a Hall bar and show that the nonreciprocal correction to the current can be of comparable magnitude to its reciprocal counterpart.

According to the Onsager reciprocity principle [1], the linear two-terminal conductance G_0 must be invariant under the time-reversal symmetry (TRS), which changes the sign of the magnetic field \mathbf{B} and the magnetization of the system \mathbf{M} . In contrast, the nonlinear two-terminal transport need not be reciprocal. In noncentrosymmetric conductors, i.e. nonpolar systems lacking inversion symmetry, due to the existence of an invariant that is linear in both the electric field \mathbf{E} and the TRS breaking pseudovectors \mathbf{B} or \mathbf{M} , the nonreciprocal contribution to the current density appears already in second order in the electric field \mathbf{E} [2–4]. Thus, in a two-terminal setup, the electric current I through the system may be expressed as

$$I = G_0 V + G_2 V^2, \quad (1)$$

where V is the voltage bias, and the nonlinear part of the conductance, G_2 , has a nonreciprocal (odd in the TRS-breaking perturbation) part. Nonreciprocity of nonlinear electron transport was the subject of extensive research, particularly in the context of mesoscopic systems such as quantum dots, carbon nanotubes, and quantum wires [5–9]. In superconducting systems with broken TRS, nonreciprocity arises in both equilibrium current, i.e. the so called superconducting diode effect [10], and the dissipative regime [11]. For a recent review of nonreciprocal transport and optical phenomena in quantum materials see Ref. 12.

In this article, we extend the theory [13–17] of nonreciprocal electron transport in noncentrosymmetric conductors with broken TRS from the disorder-dominated Drude regime to the hydrodynamic regime dominated by momentum-conserving electron-electron scattering. The nondissipative effects of TRS breaking in the hydrodynamics of liquids, such as ferrofluids [18, 19], can be described by introducing nondissipative kinetic coefficients of the liquid [20], which are odd under TRS as required

by the Onsager symmetry. For instance, significant attention has been devoted to the study of odd, or Hall, viscosity of quantum liquids [21–25] and active matter systems [26–29] with broken TRS.

We consider the effects of TRS breaking on the dissipative nonlinear hydrodynamic transport. In most high-mobility semiconductor heterostructures and graphene devices where the hydrodynamic electron transport has been demonstrated (see, e.g., reviews [30, 31] and references therein), the electron liquid lacks Galilean invariance. Some of the recent experiments suggest the electron liquid may spontaneously break TRS and inversion symmetry [32–35]. We show that in the absence of Galilean invariance of the electron liquid, the nonreciprocity of dissipative transport is strongly enhanced in comparison to that in the Drude regime.

Nonlinear hydrodynamic flows of conventional Newtonian liquids obey dynamical similarity [36, 37], which enables modeling of large-scale hydrodynamic phenomena in a lab; flows in homothetic systems can be mapped to one another by rescaling of the hydrodynamic velocity. All equivalent flows have the same Reynolds number $\mathcal{R} = U_0 L / \nu$, where U_0 and L are characteristic velocity and length scales, respectively, and ν is the kinematic viscosity. We show that nonlinear flows of noncentrosymmetric electron liquids with broken time-reversal invariance do not possess dynamical similarity. Besides the Reynolds number, they need to be characterized by an additional dimensionless parameter, which is proportional to U_0 and is odd under time-reversal.

The existence of the nonreciprocal correction G_2 in Eq. (1) requires breaking not only TRS but also a spatial symmetry, which distinguishes the flows in the forward and reverse directions. Similar to the situation in mesoscopic systems, this symmetry breaking may be associated with the device geometry [38], e.g. a flow in a funnel. Here we focus on the two-dimensional (2D)

Poiseuille flow in the Hall, which does not break the symmetry between the forward and reverse bias. In this case, the required spatial symmetry breaking arises in the non-centrosymmetric electron liquid itself. Below we study nonreciprocal transport for two different types of spatial symmetry breaking relevant to experiments: *i*) symmetry breaking of a vector type realized in 2D electron liquids with spin-orbit interaction in the presence of an in-plane Zeeman field, *ii*) symmetry breaking of a tensor type realized in valley-polarized graphene [32–35]. The breaking of forward/reverse symmetry is described in *i*) by a vector, which is linear in the Zeeman field, and in *ii*) by an invariant rank-3 tensor that is allowed by the C_3 rotation symmetry in graphene. In the latter situation, the magnitude of the nonreciprocal conductance depends on the orientation of the Hall bar relative to the graphene lattice.

The enhancement of nonreciprocity of electron transport in the hydrodynamic regime is caused by the linear dependence of the viscosity of the electron liquid on the hydrodynamic velocity \mathbf{v} . Such a dependence is allowed by symmetry in noncentrosymmetric conductors with broken TRS. Its existence can be established in the framework of the Boltzmann equation for model systems of vector and tensor type symmetry breaking mentioned above. In a general situation, the liquid viscosity is described by a rank-4 tensor $\hat{\eta}$, which relates the viscous stress tensor

$$\sigma_{ij} = \eta_{ijkl} V_{kl}, \quad V_{ij} = \frac{1}{2} (\partial_j v_i + \partial_i v_j), \quad (2)$$

to the strain rate tensor V_{ij} . For Galilean-invariant liquids, the viscosity tensor cannot depend on the flow velocity, and for isotropic incompressible liquids in two dimensions it can be reduced to the shear, η , and odd, η_H , viscosities [21]. When both TRS and Galilean invariance are broken, a linear in the flow velocity correction to the viscous stress tensor is permitted,

$$\delta\sigma_{ij} = T_{ijklm} v_k \partial_l v_m. \quad (3)$$

In vector-type symmetry breaking, where the TRS is broken by an external magnetic field \mathbf{B} , the rank-5 tensor T_{ijklm} may be constructed using \mathbf{B} , and the invariant tensors of Levi-Civita, ϵ_{ij} , and Kronecker, δ_{ij} . This yields

$$\delta\sigma_{ij} = \eta [V_{ij} (\alpha \delta_{kl} B_l v_k + \tilde{\alpha} \epsilon_{lk} B_k v_l) + \beta v_i B_k \partial_k v_j + \tilde{\beta} B_i v_k \partial_k v_j + \dots]. \quad (4)$$

Here we retained only several representative terms in the constitutive law and introduced phenomenological parameters α, β, \dots whose values must be determined from a microscopic theory.

In tensor-type symmetry breaking, which is realized in the recently discovered quarter-metal state in rhombohedral trilayer graphene [32], the TRS breaking is caused by

valley and spin polarization. Consequently, the system does not possess an in-plane vector breaking the TRS. However, due to the trigonal warping of the electron spectrum, the rotational symmetry of the electron liquid is lowered to C_3 , which allows for an invariant rank-3 tensor R_{ijk} . The linear in \mathbf{v} tensor T_{ijklm} in Eq. (3) can be constructed using R_{ijk} , δ_{ij} , and ϵ_{ij} . It is more convenient to express this tensor relation in the chiral complex coordinates $z = x + iy$ and $\bar{z} = x - iy$. In these coordinates, a traceless stress tensor of an incompressible liquid is described by a single element $\delta\sigma_{zz} = \delta\sigma_{xx} - \delta\sigma_{yy} + 2i\delta\sigma_{xy}$, and the invariant rank-3 tensor R has only two non-vanishing components: $r \equiv R_{zzz}$ and $\bar{r} \equiv R_{\bar{z}\bar{z}\bar{z}}$ [39] (since in Cartesian coordinates the tensor R is real, \bar{r} is the complex-conjugate of r). Upon a rotation by $2\pi/3$ the chiral coordinates are transformed as $z \rightarrow \epsilon z$, $\bar{z} \rightarrow \epsilon^{-1} \bar{z}$, where $\epsilon = e^{2\pi i/3}$. Therefore, in the complex coordinates the nonreciprocal part of the viscous stress tensor in Eq. (4) has the form

$$\delta\sigma_{zz} = \eta r V_{\bar{z}\bar{z}} v_z, \quad \delta\sigma_{\bar{z}\bar{z}} = \eta \bar{r} V_{zz} v_{\bar{z}}. \quad (5)$$

The characteristic ratio of the nonreciprocal stress in Eqs. (4), and Eq. (5) to the standard viscous stress $(\sigma_0)_{ij} = 2\eta V_{ij}$ in a given flow defines a dimensionless parameter, which is proportional to the characteristic flow velocity U_0 but is distinct from the Reynolds number,

$$\mathcal{N} \sim \frac{\delta\sigma}{\sigma_0} \propto U_0. \quad (6)$$

We refer to \mathcal{N} as the *nonreciprocity number*. It can be interpreted as the ratio of the typical flow velocity U_0 to the characteristic velocity scale associated with TRS breaking. In vector-type symmetry breaking the latter is proportional to the strength of spin-orbit coupling and the Zeeman field. In tensor-type, assuming that the degree of valley polarization is of order unity, the characteristic velocity is given by the Fermi velocity modulation caused by the trigonal warping. Since \mathcal{N} is proportional to the flow velocity, but independent of system size, it breaks dynamical similarity of nonlinear flows.

To obtain quantitative results for nonreciprocal electron transport in the hydrodynamic regime, we consider the experimentally relevant two-dimensional Poiseuille flow in the Hall bar geometry shown in Fig. 1. Transport of momentum in a steady-state hydrodynamic flow is described by the force balance equation [36]

$$ne\mathbf{E}_i = \partial_j \Pi_{ij}, \quad (7)$$

where n is the density of electrons, e is the electron charge, \mathbf{E} is the electric field, and Π_{ij} is the momentum flux tensor density. The latter is conventionally expressed in terms of the pressure $P\delta_{ij}$ and the viscous stress tensor σ_{ij} , as

$$\Pi_{ij} = P\delta_{ij} + \rho v_i v_j - 2\eta V_{ij} - \delta\sigma_{ij}. \quad (8)$$




FIG. 1. 2D nonreciprocal flow in a Hall bar of width $2d$. Nonreciprocal velocity profile $\mathbf{v} = u(y)\hat{\mathbf{x}}$ is controlled by the nonreciprocity number \mathcal{N} in (6), which depends on the orientation of the in-plane magnetic field B in vector-type symmetry breaking, and on the valley polarization and the orientation of the graphene lattice in tensor-type. Panel (a) displays the velocity profile (10) for $\mathcal{N} = 0.3$ and 0.9 , while panel (b) displays the velocity profiles for the time-reversed values $\mathcal{N} = -0.3$ and -0.9 . In both cases the Poiseuille profile is restored for $\mathcal{N} \rightarrow 0$.

In the absence of Galilean invariance, the expression $P\delta_{ij} + \rho v_i v_j$ for the equilibrium part of the momentum flux tensor in Eq. (8) should be viewed as a formal expansion of momentum flux density in an equilibrium state of uniform flow to second order in the powers of the flow velocity \mathbf{v} with higher-order terms omitted.

For a current flowing through a channel of width $2d$, as displayed in Fig. 1, the velocity field has only one nonzero component $\mathbf{v} = u(y)\hat{\mathbf{x}}$, which depends only on the transverse coordinate y . The viscous force density is fully described by the single component of the viscous stress, $\sigma_{xy}(y)$. As a result, Eq. (7) reduces to

$$\frac{en\mathcal{E}}{U_0} + \partial_y \left[\eta \left(1 + \mathcal{N} \frac{u}{U_0} \right) \partial_y \frac{u}{U_0} \right] = 0, \quad (9)$$

where we absorbed the pressure gradient $\partial_x P$ into the electromotive force (EMF) density $ne\mathcal{E} = neE - \partial_x P$. The latter is related to the voltage drop, $\mathcal{E} = V/L$, with L being the length of the Hall bar [40]. For vector- and tensor-type of symmetry breaking the value of the nonreciprocity parameter \mathcal{N} in Eq. (9) is easily determined from Eqs. (4) and (5). For vector-type the precise form of \mathcal{N} depends on the mutual orientation of the field with respect to the flow velocity. For tensor-type $\mathcal{N} = (r + \bar{r})U_0/2$. Notice that a rotation of crystalline

axes by an angle θ relative to the flow changes $r \rightarrow re^{3i\theta}$. Therefore, the combination $r + \bar{r}$ in the nonreciprocity parameter exhibits a periodic modulation $\propto \cos(3[\theta - \theta_0])$. This is consistent with the three-fold rotation symmetry observed in nonreciprocity measurements in valley-polarized multilayer graphene [34, 35].

Equation (9) can be readily integrated with the integration constants fixed by the boundary condition imposed on the flow. For simplicity, we apply no-slip conditions $u(\pm d) = 0$. The more general case of a flow with a finite slip length does not alter the essential physics underlying nonreciprocity. Thus, the profile of the flow can be found in the form

$$u(y) = \frac{U_0}{\mathcal{N}} \left[\sqrt{1 + \mathcal{N} \left(1 - \frac{y^2}{d^2} \right)} - 1 \right], \quad (10)$$

where we introduced $U_0 = en\mathcal{E}d^2/\eta$. Equation (10) is valid for both positive and negative values of \mathcal{N} . At $\mathcal{N} \rightarrow 0$ it reproduces the Poiseuille flow profile, see Fig. 1 for illustration.

From the flow profile, we can compute the total current [41]. The resulting expression for the total current ($B > 0$) takes a relatively simple form

$$I = \int_{-d}^{+d} enu(y)dy = I_0 f(\mathcal{N}), \quad I_0 = \frac{2}{3} \frac{(en)^2 \mathcal{E} d^3}{\eta}, \quad (11)$$

where the dimensionless function is given by

$$f(\mathcal{N}) = \frac{3}{2\mathcal{N}^{3/2}} \left[(1 + \mathcal{N}) \arcsin \sqrt{\frac{\mathcal{N}}{1 + \mathcal{N}}} - \sqrt{\mathcal{N}} \right]. \quad (12)$$

The field-independent linear conductance, $G_0 = (I/V)_{\mathcal{N} \rightarrow 0}$, takes the value

$$G_0 = \frac{2e^2}{3} (nd^2) \frac{n}{\eta} \frac{d}{L}. \quad (13)$$

It is inversely proportional to the viscosity of the electron fluid, which is the manifestation of the Gurzhi effect [42]. This behavior was confirmed experimentally [43–45]. Working to leading order in the strength of the TRS breaking perturbation, $\mathcal{N} \ll 1$, we extract the nonlinear nonreciprocal correction to conductance of the system. In the notations of Eq. (1) it takes the form

$$G_2 = -\frac{G_0 nd^2}{5 \eta L} \begin{cases} (e\alpha B), & \text{vector-type,} \\ e|r| \cos(3[\theta - \theta_0]), & \text{tensor-type.} \end{cases} \quad (14)$$

This is obtained by expanding the expression $I/I_0 = f(\mathcal{N}) \approx 1 - \mathcal{N}/5$ in (11), where f is given by Eq. (12), and keeping only leading order terms in the nonreciprocity number \mathcal{N} . For simplicity, in vector-type symmetry breaking, we also took a field orientation $\mathbf{B} = B\hat{\mathbf{x}} \parallel \mathbf{v}$ so that nonreciprocity number simplifies to $\mathcal{N} = \alpha BU_0$. In Fig. 2 we display the function $f(\mathcal{N})$ vs. \mathcal{N} , along with the limiting cases of weak and strong nonreciprocity.




FIG. 2. Plot of the dimensionless current $I/I_0 = f(N)$ in Eq. (11) (continuous curve) where f is given by Eq. (12), vs. the nonreciprocity number for $N > 0$, cf. Fig. 1(a). The dashed curve on the left denotes its linear approximation, $I/I_0 \approx 1 - N/5$, whose nonreciprocal correction to the conductance was calculated in Eq. (14). The dashed curve on the right denotes its high N asymptotic expansion $I/I_0 \approx 3\pi/(4\sqrt{N})$. Inset: Plot of the dimensionless current I/I_0 calculated for the negative branch of the flow profile displayed in Fig. 1(b).

We now compare the magnitude of nonreciprocity in the hydrodynamic regime to that in the disorder-dominated regime. In the latter case, the ratio of the nonreciprocal to reciprocal current can be expressed as

$$\frac{\delta I_{\text{dis}}}{I_0} \sim \varsigma \frac{e\mathcal{E}\ell_{\text{ei}}}{\epsilon_{\text{F}}}, \quad (15)$$

where $\ell_{\text{ei}} = v_{\text{F}}\tau_{\text{ei}}$ is the disorder-limited electron mean-free path, and ς is a dimensionless strength of TRS breaking. For example, for carbon nanotubes considered in Ref. 13, $\varsigma \sim \Phi/\Phi_0$, where Φ is the magnetic flux through the nanotube and $\Phi_0 = hc/e$ is the flux quantum. In the hydrodynamic regime, the corresponding estimate, for the relative magnitude of the nonreciprocal current is significantly enhanced in comparison to its disorder-dominated counterpart. In valley-polarized graphene, the degree of nonreciprocity of the electron spectrum depends on the energy scale Δ associated with the trigonal warping. The corresponding dimensionless parameter $\varsigma \sim \Delta/\epsilon_{\text{F}}$, may in principle reach values of order unity. For the case of 2D electron liquid with Rashba spin-orbit coupling, this parameter can be estimated as a ratio of the Zeeman energy to the spin-orbit splitting. Using the estimate for the typical flow velocity from Eq. (10), $U_0 \sim \frac{d^2}{\eta} en\mathcal{E}$, and estimating the shear viscosity in the Fermi liquid regime as $\eta \sim np_{\text{F}}\ell_{\text{ee}}$, where $\ell_{\text{ee}} = v_{\text{F}}\tau_{\text{ee}}$ is the inelastic mean free path, we obtain

$$\frac{\delta I_{\text{hydro}}}{I_0} \sim \varsigma \frac{d}{\ell_{\text{ee}}} \frac{e\mathcal{E}d}{\epsilon_{\text{F}}}. \quad (16)$$

Comparing Eqs. (16) and (15) we see that in the hydrodynamic regime the elastic electron mean free path

is effectively replaced by the channel width, $\ell_{\text{ei}} \rightarrow d$, and an additional large factor $d/\ell_{\text{ee}} \gg 1$ arises from the dependence of the flow velocity on the channel width. Thus, the relative nonreciprocal correction in the hydrodynamic regime is enhanced in comparison to the disorder-dominated regime by a large factor

$$\frac{\delta I_{\text{hydro}}}{\delta I_{\text{dis}}} \sim \frac{d^2}{\ell_{\text{ee}}\ell_{\text{ei}}}. \quad (17)$$

In the presence of impurity scattering in the bulk, the hydrodynamic approach used above is valid provided d does not exceed the Gurzhi length, $\ell_{\text{G}} = \sqrt{\ell_{\text{ei}}\ell_{\text{ee}}}$. For wide channels ($d > \ell_{\text{G}}$), electron flow is dominated by the scattering in the bulk of the device, leading to a transition from hydrodynamic Poiseuille flow to Ohmic behavior. As expected, at the crossover boundary between the two regimes, $d \sim \ell_{\text{G}}$, the results for nonreciprocity in Eqs. (16) and (15) match.

We thank B. Spivak for useful discussions. The work of L. B. and A. V. A. was supported by the National Science Foundation Grant No. DMR-2424364, the Thouless Institute for Quantum Matter (TIQM), and the College of Arts and Sciences at the University of Washington. The work of A. L. was supported by NSF Grant No. DMR-2203411 and H. I. Romnes Faculty Fellowship provided by the University of Wisconsin-Madison Office of the Vice Chancellor for Research and Graduate Education with funding from the Wisconsin Alumni Research Foundation.

-
- [1] Lars Onsager, “Reciprocal relations in irreversible processes. I.” *Phys. Rev.* **37**, 405–426 (1931); “Reciprocal relations in irreversible processes. II.” *Phys. Rev.* **38**, 2265–2279 (1931).
 - [2] B. Sturman and V. Fridkin, *Photovoltaic and Photorefractive effects in noncentrosymmetric materials*, Ferroelectricity and related phenomena (Taylor and Francis, 1992).
 - [3] G. L. J. A. Rikken, J. Fölling, and P. Wyder, “Electrical magnetochiral anisotropy,” *Phys. Rev. Lett.* **87**, 236602 (2001).
 - [4] E. L. Ivchenko and B. Spivak, “Chirality effects in carbon nanotubes,” *Phys. Rev. B* **66**, 155404 (2002).
 - [5] A. Löfgren, C. A. Marlow, I. Shorubalko, R. P. Taylor, P. Omling, L. Samuelson, and H. Linke, “Symmetry of two-terminal nonlinear electric conduction,” *Phys. Rev. Lett.* **92**, 046803 (2004).
 - [6] Jiang Wei, Michael Shimogawa, Zenghui Wang, Iuliana Radu, Robert Dormaier, and David Henry Cobden, “Magnetic-field asymmetry of nonlinear transport in carbon nanotubes,” *Phys. Rev. Lett.* **95**, 256601 (2005).
 - [7] R. Leturcq, D. Sánchez, G. Götz, T. Ihn, K. Ensslin, D. C. Driscoll, and A. C. Gossard, “Magnetic field symmetry and phase rigidity of the nonlinear conductance in a ring,” *Phys. Rev. Lett.* **96**, 126801 (2006).
 - [8] D. M. Zumbühl, C. M. Marcus, M. P. Hanson, and A. C.

- Gossard, “Asymmetry of nonlinear transport and electron interactions in quantum dots,” *Phys. Rev. Lett.* **96**, 206802 (2006).
- [9] L. Angers, E. Zakka-Bajjani, R. Deblock, S. Guéron, H. Bouchiat, A. Cavanna, U. Gennser, and M. Polianski, “Magnetic-field asymmetry of mesoscopic *dc* rectification in Aharonov-Bohm rings,” *Phys. Rev. B* **75**, 115309 (2007).
- [10] Muhammad Nadeem, Michael S. Fuhrer, and Xiaolin Wang, “The superconducting diode effect,” *Nature Reviews Physics* **5**, 558–577 (2023).
- [11] T. Liu, M. Smith, A. V. Andreev, and B. Z. Spivak, “Giant nonreciprocity of current-voltage characteristics of noncentrosymmetric superconductor–normal metal–superconductor junctions,” *Phys. Rev. B* **109**, L020501 (2024).
- [12] Naoto Nagaosa and Youichi Yanase, “Nonreciprocal transport and optical phenomena in quantum materials,” *Annual Review of Condensed Matter Physics* **15**, 63–83 (2024).
- [13] B. Spivak and A. Zyuzin, “Signature of the electron-electron interaction in the magnetic-field dependence of nonlinear I-V characteristics in mesoscopic systems,” *Phys. Rev. Lett.* **93**, 226801 (2004).
- [14] David Sánchez and Markus Büttiker, “Magnetic-field asymmetry of nonlinear mesoscopic transport,” *Phys. Rev. Lett.* **93**, 106802 (2004).
- [15] M. L. Polianski and M. Büttiker, “Mesoscopic fluctuations of nonlinear conductance of chaotic quantum dots,” *Phys. Rev. Lett.* **96**, 156804 (2006).
- [16] A. De Martino, R. Egger, and A. M. Tsvelik, “Nonlinear magnetotransport in interacting chiral nanotubes,” *Phys. Rev. Lett.* **97**, 076402 (2006).
- [17] E. Deyo, B. Spivak, and A. Zyuzin, “Signature of the electron-electron interaction in the magnetic-field dependence of nonlinear I-V characteristics in mesoscopic conductors,” *Phys. Rev. B* **74**, 104205 (2006).
- [18] MI Shliomis, “Hydrodynamics of a liquid with intrinsic rotation,” *Soviet Journal of Experimental and Theoretical Physics* **24**, 173 (1967).
- [19] Stefan Odenbach, *Magnetoviscous effects in ferrofluids* (Berlin ; New York : Springer, 2002).
- [20] E. M. Lifshitz and L. P. Pitaevskii, *Course of theoretical physics. Vol. 10: Physical Kinetics* (Pergamon Press, 1981).
- [21] J. E. Avron, “Odd viscosity,” *Journal of Statistical Physics* **92**, 543–557 (1998).
- [22] F. D. M. Haldane, “Hall viscosity” and intrinsic metric of incompressible fractional Hall fluids,” (2009), arXiv:0906.1854 [cond-mat.str-el].
- [23] Barry Bradlyn, Moshe Goldstein, and N. Read, “Kubo formulas for viscosity: Hall viscosity, Ward identities, and the relation with conductivity,” *Phys. Rev. B* **86**, 245309 (2012).
- [24] Carlos Hoyos and Dam Thanh Son, “Hall viscosity and electromagnetic response,” *Phys. Rev. Lett.* **108**, 066805 (2012).
- [25] Yonatan Messica, Alex Levchenko, and Dmitri B. Gutman, “Stokes flow in the electronic fluid with odd viscosity,” (2025), arXiv:2501.10890 [cond-mat.mes-hall].
- [26] Debarghya Banerjee, Anton Souslov, Alexander G. Abanov, and Vincenzo Vitelli, “Odd viscosity in chiral active fluids,” *Nature Communications* **8**, 1573 (2017).
- [27] A. Aggarwal, E. Kirkinis, and M. Olvera de la Cruz, “Thermocapillary migrating odd viscous droplets,” *Physical Review Letters* **131**, 198201 (2023).
- [28] E. Kirkinis and M. Olvera de la Cruz, “Taylor columns and inertial-like waves in a three-dimensional odd viscous liquid,” *Journal of Fluid Mechanics* **973**, A30 (2023).
- [29] E. Kirkinis and M. Olvera de la Cruz, “Evanescent and inertial-like waves in rigidly rotating odd viscous liquids,” *Journal of Fluid Mechanics* **996**, A13 (2024).
- [30] Boris N. Narozhny, “Hydrodynamic approach to two-dimensional electron systems,” *La Rivista del Nuovo Cimento* **45**, 661–736 (2022).
- [31] L. Fritz and T. Scaffidi, “Hydrodynamic electronic transport,” *Annual Review of Condensed Matter Physics* **15**, 17–44 (2024).
- [32] Haoxin Zhou, Tian Xie, Areg Ghazaryan, Tobias Holder, James R. Ehrets, Eric M. Spanton, Takashi Taniguchi, Kenji Watanabe, Erez Berg, Maksym Serbyn, and Andrea F. Young, “Half- and quarter-metals in rhombohedral trilayer graphene,” *Nature* **598**, 429–433 (2021).
- [33] Naiyuan James Zhang, Jiang-Xiazi Lin, Dmitry V. Chichinadze, Yibang Wang, Kenji Watanabe, Takashi Taniguchi, Liang Fu, and J. I. A. Li, “Angle-resolved transport non-reciprocity and spontaneous symmetry breaking in twisted trilayer graphene,” *Nature Materials* **23**, 356–362 (2024).
- [34] Jiang-Xiazi Lin, Yibang Wang, Naiyuan J. Zhang, Kenji Watanabe, Takashi Taniguchi, Liang Fu, and J. I. A. Li, “Spontaneous momentum polarization and diodicity in Bernal bilayer graphene,” (2023), arXiv:2302.04261 [cond-mat.mes-hall].
- [35] Dmitry V. Chichinadze, Naiyuan James Zhang, Jiang-Xiazi Lin, Xiaoyu Wang, Kenji Watanabe, Takashi Taniguchi, Oskar Vafeek, and J. I. A. Li, “Observation of giant nonlinear Hall conductivity in Bernal bilayer graphene,” (2024), arXiv:2411.11156 [cond-mat.mes-hall].
- [36] L. D. Landau and E. M. Lifshitz, *Fluid Mechanics*, 2nd ed., Course of Theoretical Physics Series, Vol. 6 (Butterworth-Heinemann, Oxford, 1987).
- [37] G.K. Batchelor, *An introduction to fluid dynamics* (Cambridge University Press, 1967).
- [38] A. V. Andreev and L. I. Glazman, “Nonlinear magnetoconductance of a classical ballistic system,” *Phys. Rev. Lett.* **97**, 266806 (2006).
- [39] L. D. Landau and E. M. Lifshitz, *Theory of Elasticity*, 3rd ed., Course of Theoretical Physics Series, Vol. 7 (Butterworth-Heinemann, Oxford, 1986).
- [40] It is worth noting that for Galilean-invariant liquids, the Navier-Stokes equation predicts a linear current-voltage dependence for the Poiseuille flow at arbitrary Reynolds numbers. Therefore, in this case all nonlinearity is non-reciprocal.
- [41] In systems lacking Galilean invariance, the total current density consists of the usual convective term, *enu*, and an additional contribution arising from the finite intrinsic conductivity [46]. Assuming the particle density is sufficiently high, we neglect the latter.
- [42] R. N. Gurzhi, “Hydrodynamic effects in solids at low temperature,” *Soviet Physics Uspekhi* **11**, 255 (1968).
- [43] R. Krishna Kumar, D. A. Bandurin, F. M. D. Pellegrino, Y. Cao, A. Principi, H. Guo, G. H. Auton, M. Ben Shalom, L. A. Ponomarenko, G. Falkovich, K. Watanabe, T. Taniguchi, I. V. Grigorieva, L. S. Levitov, M. Polini, and A. K. Geim, “Superballistic flow of

- viscous electron fluid through graphene constrictions,” *Nature Physics* **13**, 1182–1185 (2017).
- [44] Zachary J. Krebs, Wyatt A. Behn, Songci Li, Keenan J. Smith, Kenji Watanabe, Takashi Taniguchi, Alex Levchenko, and Victor W. Brar, “Imaging the breaking of electrostatic dams in graphene for ballistic and viscous fluids,” *Science* **379**, 671–676 (2023).
- [45] Yihang Zeng, Haoyu Guo, Olivia M. Ghosh, Kenji Watanabe, Takashi Taniguchi, Leonid S. Levitov, and Cory R. Dean, “Quantitative measurement of viscosity in two-dimensional electron fluids,” (2024), arXiv:2407.05026 [cond-mat.mes-hall].
- [46] Songci Li, A. V. Andreev, and Alex Levchenko, “Hydrodynamic electron transport in graphene Hall-bar devices,” *Phys. Rev. B* **105**, 155307 (2022).