Cosmic Acceleration from Nothing

Michael R.R. $Good^{1,2,*}$ and Eric V. Linder^{3,†}

¹Physics Department & Energetic Cosmos Laboratory, Nazarbayev University,

Astana 010000, Qazaqstan

²Leung Center for Cosmology and Particle Astrophysics, National Taiwan University,

Taipei 10617, Taiwan

³Berkeley Center for Cosmological Physics & Berkeley Lab, University of California,

Berkeley CA 94720, USA.

(Dated: March 5, 2025)

We demonstrate that if the universe started as a vacuum fluctuation rather than from a singular Big Bang state, the universe must have a late-time cosmic acceleration. This is required by a "cosmological sum rule" derived using the Schwarzian form of the Friedmann equations. We discuss possible connections to conformal and Möbius transformations, and also compute that the best fit present cosmic data is consistent with the necessary crossing of the Schwarzian through zero having occurred (while it would not yet have happened in a Λ CDM cosmology).

I. INTRODUCTION

Cosmic acceleration may arise from a vacuum energy, i.e., a cosmological constant or a new scalar field component. We do not know any reason why these would *have* to exist and cause the observed late-time acceleration of the cosmic expansion.

Here, we explore changing the vacuum structure in the early universe so that the universe arises as a quantum fluctuation in empty space rather than directly from a hot, dense Big Bang and how this might enforce a latetime acceleration.

In Section II, we introduce the Schwarzian derivative in cosmology and explore its implications for the universe's evolution. Section III discusses the Schwarzian's connection to fundamental symmetries, such as Lorentz and Möbius transformations, its use in conformal field theory and gravitational dynamics, and motivation for exploring its role in cosmology. Section IV uses the Schwarzian to derive a cosmological sum rule, which imposes a necessary condition on the universe's expansion history. Leveraging this sum rule, we compute constraints on cosmic acceleration in Section V, showing that late-time acceleration follows from a vacuum fluctuation origin, and compare to current data. Finally, Section VI summarizes the main findings.

II. COSMIC EXPANSION

The Schwarzian derivative plays a key role in many fields of physics, including connections to Lorentz transformations and conformal field theory (see Section III). Here we will apply it to standard cosmology, revealing intriguing implications connecting the early and late universe. Reviewing the standard quantities of cosmic expansion, we start with the Robertson-Walker metric of a homogeneous, isotropic spacetime,

$$ds^{2} = -dt^{2} + a^{2}(t) \left[\frac{dr^{2}}{1 - kr^{2}} + r^{2}(d\theta^{2} + \sin^{2}\theta d\phi^{2}) \right],$$
(1)

where a(t) is the expansion factor, k the spatial curvature constant, and $\{t, r, \theta, \phi\}$ are the coordinates. Using the Einstein field equations of general relativity yields the Friedmann equations determining the expansion behavior a(t) in terms of the energy-momentum contents (and k).

Those equations of motion can be written in terms of combinations of derivatives of the expansion factor, e.g.,

$$H(a) \equiv \frac{\dot{a}}{a} , \qquad (2)$$

$$q(a) \equiv -\frac{a\ddot{a}}{\dot{a}^2} , \qquad (3)$$

called the Hubble parameter H, or logarithmic expansion rate, and the deceleration parameter q. While third derivatives do not enter the field equations directly, it will be useful to define the jerk parameter,

$$j(a) \equiv \frac{a^2 \ddot{a}}{\dot{a}^3} . \tag{4}$$

Now we introduce the Schwarzian, denoted here by brackets,

$$\{a,t\} \equiv \frac{\ddot{a}}{\dot{a}} - \frac{3}{2} \left(\frac{\ddot{a}}{\dot{a}}\right)^2 \,. \tag{5}$$

This can be written in terms of the previously defined expansion quantities as (e.g. [1]),

$$\{a,t\} = H^2 \left(j - \frac{3}{2}q^2\right) . \tag{6}$$

We note that this has several important properties, both cosmological – it only involves a(t) and has no explicit dependence on spatial curvature k – and involving physical symmetries (discussed in the next section).

^{*} michael.good@nu.edu.kz

[†] evlinder@lbl.gov

Before proceeding, let us connect these quantities to simple Friedmann cosmologies. For energy-momentum contributions from noninteracting components, each with dimensionless energy density (as a fraction of the critical energy density) Ω_w and equation of state (pressure to energy density) ratio w, we have

$$q = \frac{1}{2} \sum (1+3w)\Omega_w \ . \tag{7}$$

We can see explicitly that spatial curvature does not enter into q(a), since it effectively has $w_k = -1/3$. That is, $\Omega_k \equiv -ka^{-2}/(H_0^2)$, where $H_0 = H(a = 1)$ is the expansion rate today, follows the standard energy density dilution with expansion as $a^{-3(1+w)} = a^{-2}$.

Furthermore, the jerk,

$$j = q + 2q^2 - q' , (8)$$

where prime denotes $d/d \ln a$, is also immune to spatial curvature. For illustration we write here the expressions for the case of matter (which has w = 0) plus one additional component, for a critical density universe (so $\Omega_m + \Omega_w = 1$). Then

$$q = \frac{1}{2} + \frac{3}{2}w\Omega_w$$
(9)

$$j = 1 + \frac{9}{2}w(1+w)\Omega_w - \frac{3}{2}w'\Omega_w , \qquad (10)$$

$$\{a,t\} = \frac{H^2}{8} \left[5 + 18w(1+2w)\Omega_w - 12w'\Omega_w - 27w^2\Omega_w^2 \right].$$
(11)

Note that when the additional component is a cosmological constant (w = -1), called Λ CDM cosmology, then $q = (1 - 3\Omega_w)/2$, ranging from 1/2 in matter domination ($\Omega_w = 0$) to -1 in cosmological constant domination ($\Omega_w = 1$, or de Sitter asymptote). However, j = 1 for all times for Λ CDM.

III. SCHWARZIANS AND SYMMETRIES

The Schwarzian derivative as defined in Equation (5) has close connections with the symmetry group SL(2, R) and hence Lorentz transformations, conformal transformations, and Möbius transformations (see e.g. [2]). The symmetry gives invariance under physical actions like boosts, dilations, and rotations.

Among other uses in physics, the Schwarzian appears in conformal field theory (CFT) and AdS_2 gravity, governing boundary dynamics in Jackiw-Teitelboim (JT) gravity (see e.g. [3, 4]). It controls the low-energy sector of the SYK model, a maximally chaotic system (see e.g. [5, 6]), by describing the soft reparametrization modes that emerge from the breaking of conformal symmetry. It measures deviations breaking Möbius symmetry and plays a role in conformal transformations in cosmology [1]. In black hole evaporation analogs, such as the moving mirror model [7, 8], it determines the energy flux (the stress tensor) of quantum particles radiated by an accelerating mirror (see e.g. [9, 10]). Its relation to Möbius transformations plays a central role in analyzing invariant energy flux, particle spectrum, and entanglement entropy of de Sitter and black hole horizons [11].

Given the usefulness of the Schwarzian, its appearance relating basic quantities of cosmic expansion in Equation (5) merits investigation of possible implications.

IV. COSMOLOGICAL SUM RULE

We find that employment of the Schwarzian reveals a constraint known in physics as a sum rule, or integral constraint, here applied to the cosmic expansion. Let us parallel the use of the Schwarzian in the moving mirror case (also known as an accelerating boundary correspondence to a spacetime or black hole horizon). There, it proved that a unitary solution (and hence an evaporating black hole without information loss) gave a horizon sum rule (in that case, requiring some epoch where energy flux was negative) [12]. In our cosmological case, we will find a sum rule that implies that the universe must have some epoch of accelerating expansion.

In [11, 12] the Schwarzian and the resulting sum rule appeared most clearly with use of the rapidity, as in special relativity $\eta = \tanh^{-1}(dx/dt)$, but in spacetime null coordinates $\{u, v\} = \{t - x, t + x\}$ so

$$\eta = \frac{1}{2} \ln \frac{dv}{du} \,. \tag{12}$$

Therefore we define

$$\chi \equiv \frac{1}{2} \ln \dot{a} , \qquad (13)$$

and stay restricted to an expanding universe, $\dot{a} > 0$. We can now write the Schwarzian as

$$\{a,t\} = \frac{1}{2} \left[\ddot{\chi} - \dot{\chi}^2 \right] = \frac{1}{2} e^{\chi} \frac{d}{dt} \left(e^{-\chi} \dot{\chi} \right) .$$
(14)

Removing the (always positive) prefactor and integrating over time gives a total derivative,

$$\int_{0}^{+\infty} dt \, \dot{a}^{-1/2} \left\{ a, t \right\} = \left(\dot{a}^{-1/2} \frac{\ddot{a}}{\dot{a}} \right) \Big|_{t=0}^{t=+\infty} \,. \tag{15}$$

Suppose that in the future $a \sim t^n$ as $t \to +\infty$. Then the quantity in parentheses goes as $n^{-1/2}(n-1)t^{-(1+n)/2}$, and so for expansion (n > 0) the end point term will vanish. (It will also vanish for a de Sitter state, $a \sim e^{Ht}$.) At early times, the boundary term will vanish if $\ddot{a} = 0$, i.e. constant \dot{a} , hence constant χ , or $a \sim t$.

Thus, we have a cosmological sum rule,

$$\int_0^{+\infty} dt \, \dot{a}^{-1/2} \{a, t\} = 0 \,, \qquad (16)$$

if the early universe began as a Milne universe, a state empty of all contents (and with negative curvature k),

having $a \sim t$. Note that the Milne universe is, with a coordinate transform, a patch of Minkowski spacetime (which has k = 0), also a vacuum solution¹. In either case, if the universe begins not with a hot, dense Big Bang state but as a quantum fluctuation about empty spacetime, then the cosmological sum rule holds.

Since $\dot{a}^{-1/2} = e^{-\chi} > 0$ then the Schwarzian $\{a, t\}$ must cross zero to satisfy the sum rule, and hence be negative for some time during its evolution. Note also that since for a Milne universe the Schwarzian is zero (indeed q = 0 = j), then we can extend the lower limit of the integral indefinitely, if the universe arose from a Milne state.

V. COSMIC ACCELERATION

The results of the previous section are completely general, requiring only expansion and vanishing boundary terms, and do not rely on any assumptions about splitting energy density into matter plus dark energy. They imply that $j - (3/2)q^2$ must cross zero. This does not mean that $j(a) = (3/2)q^2(a)$ for all times (for which the only expanding solution is $a \sim t$, e.g. an eternal Milne universe), but rather that the Schwarzian must cross zero.

To give a flavor of this, and be quantitative, in this section we will explore the case of matter plus some other component, as in Equation (11), initially with constant w.

First, let's consider a matter plus cosmological constant (Λ CDM) universe. Then $(8/H^2)\{a,t\} = 5+18\Omega_w - 27\Omega_w^2$. As the universe evolves from early to late time, Ω_w increases from 0 to 1, and the right-hand side varies from 5 to -4, indeed crossing zero. Next, we can allow the component adding to matter to have an arbitrary constant w. In this case, at early times ($\Omega_w \to 0$) the Schwarzian will always be positive. At late times ($\Omega_w \to 1$) we have an equation for the crossing given by (5 + 3w)(1 + 3w) = 0. When w > -1/3 no crossing can occur, i.e. we cannot satisfy the cosmological sum rule.

Thus, if the universe starts as a quantum fluctuation from an empty, Milne state, it must have late-time acceleration $(w < -1/3, \Omega_w \to 1)!$

However, for w < -5/3 the Schwarzian stays positive at late times and never crosses zero. Thus, we are forced by the sum rule to have

$$-5/3 < w < -1/3$$
. (17)

We can evaluate when the crossing occurs by solving for

the roots of Equation (11) in terms of Ω_w ,

$$\Omega_w^{\text{cross}} = \frac{1}{3w} \left[1 + 2w - \sqrt{(1+2w)^2 + 5/3} \right] , \quad (18)$$

where the other root would have $\Omega_w < 0$ for w < 0.

Figure 1 shows at what Ω_w the crossing occurs as a function of w. For w = -1, it occurs at $\Omega_w = 0.878$, i.e. in the future in our universe. Outside the range -5/3 < w < -1/3, a crossing would require $\Omega_w > 1$ (i.e. a negative matter density or positive curvature).



FIG. 1. The sum rule requires the Schwarzian to cross zero, hence the presence of a component with -5/3 < w < -1/3. We plot the value of Ω_w^{cross} , i.e. when in the expansion history this occurs, for each w, from Equation (18). For Λ CDM (w = -1), for example, the crossing will occur when $\Omega_w = 0.878$.

Figure 2 plots the Schwarzian (actually $8\{a,t\}/H^2$) for Λ CDM cosmology. Recall that we can consider Ω_w as a measure of time, going from $\Omega_w = 0$ in the early universe to $\Omega_w = 1$ in the late universe.



FIG. 2. The Schwarzian, actually $8\{a,t\}/H^2$, is plotted for Λ CDM cosmology, showing the required zero crossing.

Relaxing the constancy of the equation of state w, we see that including the w' term from Equation (11) imposes a constraint on the dark energy dynamics such that we require

$$w' > \frac{5 + 18w(1 + 2w)\Omega_w - 27w^2\Omega_w^2}{12\Omega_w} , \qquad (19)$$

¹ One could also regard Minkowski spacetime as the limit $h \to 0$ in $a \sim e^{ht}$, i.e. a static limit. In this case Minkowski contributes $h^{1/2}e^{-ht/2}$ to the integral Eq. (15), which vanishes as $h \to 0$, as required.

at some time. While we cannot test this for the future, i.e. all the way to $\Omega_w = 1$, we might evaluate the expression to see if our universe is at least close to satisfying the crossing required by the cosmological sum rule. As one example, in Λ CDM we see from Figure 2 that even at $\Omega_w = 0.7$ we cannot tell the crossing is close. However, for observational data results similar to DESI [13] with a large $w' \sim 1$, $w \sim -0.7$ the inequality is satisfied today, seeming to indicate the crossing has already been accomplished, as the sum rule requires.

VI. CONCLUSIONS

If the universe started from nothing – specifically a Milne state – with a quantum fluctuation generating matter (including radiation), then we find that at some later time the cosmic expansion must accelerate. This is in sharp contrast to the usual hot, dense Big Bang picture where the universe "starts" with matter and radiation, but there is no guarantee that expansion will ever accelerate – that has to be put in by hand.

The structure of the Friedmann equations in Einstein's general relativity, under the symmetry properties of the Schwarzian and the resulting cosmological sum rule we derived, do guarantee that any quantum fluctuation from a Milne state generates not only matter and radiation (or at least a component with an equation of state w > -1/3) but also an additional component with -5/3 < w < -1/3 that will dominate at late times and accelerate cosmic expansion. The Schwarzian alone does not say how the quantum production process works.

- [1] G. W. Gibbons, Dark Energy and the Schwarzian Derivative, arXiv version (2014), arXiv:1403.5431 [hep-th].
- [2] A. Kitaev, Notes on SL(2, ℝ) representations, arXiv version (2017), arXiv:1711.08169 [hep-th].
- [3] T. G. Mertens and G. J. Turiaci, Solvable models of quantum black holes: a review on Jackiw–Teitelboim gravity, Living Rev. Rel. 26, 4 (2023), arXiv:2210.10846 [hep-th].
- [4] T. G. Mertens, The Schwarzian theory origins, JHEP 05, 036, arXiv:1801.09605 [hep-th].
- [5] J. Maldacena and D. Stanford, Remarks on the Sachdev-Ye-Kitaev model, Phys. Rev. D 94, 106002 (2016), arXiv:1604.07818 [hep-th].
- [6] Y.-H. Qi, Y. Seo, S.-J. Sin, and G. Song, Correlation functions in Schwarzian liquid, Phys. Rev. D 99, 066001 (2019), arXiv:1804.06164 [hep-th].
- [7] B. S. DeWitt, Quantum Field Theory in Curved Space-Time, Phys. Rept. 19, 295 (1975).
- [8] S. A. Fulling and P. C. W. Davies, Radiation from a

The Milne state can persist indefinitely, until the quantum transition, and since during it $a \sim t$ then there is still the possibility of a singular origin in the past, a = 0 – a Big Bang but an empty one with no matter to witness creation. Once the quantum fluctuation generates the matter, cosmic evolution can follow the standard hot, dense early cosmology, satisfying all the usual constraints of primordial nucleosynthesis, the cosmic microwave background radiation, and growth of large scale structure, but now with late-time acceleration guaranteed. We also note that nothing prevents multiple epochs of acceleration, with the Schwarzian $j - (3/2)q^2$ crossing zero several times².

This is an attractive scenario, where the origin of the universe out of nothing, utilizing the symmetry properties inherent to Einstein's equations of motion, automatically provides for a late time accelerated expansion such as we observe. The prevalence of the Schwarzian and its ties to so many fields of physics offer the hope that the quantum origin itself may be part of a deeper theory, whether in conformal field theory, quantum gravity, or symmetry breaking. Finally, Equation (19) offers a predictive constraint on the dark energy equation of state behavior that can be tested with precision observations.

ACKNOWLEDGMENTS

M.G. is supported in part by the FY2024-SGP-1-STMM Faculty Development Competitive Research Grant (FDCRGP) no.201223FD8824 and SSH20224004 at Nazarbayev University in Qazaqstan.

moving mirror in two dimensional space-time: Conformal anomaly, Proc. R. Soc. Lond. A **348**, 393 (1976).

- [9] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, UK, 1982).
- [10] A. Fabbri and J. Navarro-Salas, *Modeling black hole evap*oration (World Scientific, Singapore, 2005).
- [11] M. R. R. Good and E. V. Linder, Möbius mirrors, Class. Quant. Grav. **39**, 105003 (2022), arXiv:2108.07451 [gr-qc].
- [12] M. R. R. Good, E. V. Linder, and F. Wilczek, Moving mirror model for quasithermal radiation fields, Phys. Rev. D 101, 025012 (2020), arXiv:1909.01129 [gr-qc].
- [13] A. G. Adame *et al.* (DESI), DESI 2024 VI: cosmological constraints from the measurements of baryon acoustic oscillations, JCAP **02**, 021, arXiv:2404.03002 [astroph.CO].

tribution from radiation after inflation ends, in much the same way that a late-time de Sitter state still leads to a vanishing boundary term.

² Note that inflation is unlikely to satisfy the sum rule on its own, since the exponentially small $\dot{a}^{-1/2}$ in the integral means its negative Schwarzian would be quickly overcome by the positive con-