arXiv:2503.02418v1 [cond-mat.stat-mech] 4 Mar 2025

Measurement-Induced Crossover of Quantum Jump Statistics in Postselection-Free Many-Body Dynamics

Kazuki Yamamoto^{1, *} and Ryusuke Hamazaki²

 1 Department of Physics, Institute of Science Tokyo, Meguro, Tokyo 152-8551, Japan †

²Nonequilibrium Quantum Statistical Mechanics RIKEN Hakubi Research Team,

RIKEN Cluster for Pioneering Research (CPR), RIKEN iTHEMS, Wako, Saitama 351-0198, Japan

(Dated: March 5, 2025)

We reveal a nontrivial crossover of subsystem fluctuations of quantum jumps in continuously monitored many-body systems, which have a trivial maximally mixed state as a steady-state density matrix. While the fluctuations exhibit the standard volume law $\propto L$ following Poissonian statistics for sufficiently weak measurement strength, anomalous yet universal scaling law $\propto L^{\alpha}$ ($\alpha \sim 2.7$) indicating super-Poissonian statistics appears for strong measurement strength. This drastically affects the precision of estimating the rate of quantum jumps: for strong (weak) measurement, the estimation uncertainty is enhanced (suppressed) as the system size increases. We demonstrate that the anomalous scaling of the subsystem fluctuation originates from an integrated many-body autocorrelation function and that the transient dynamics contributes to the scaling law rather than the Liouvillian gap. The measurement-induced crossover is accessed only from the postselection-free information obtained from the time and the position of quantum jumps and can be tested in ultracold atom experiments.

Introduction-Dissipative quantum many-body phenomena induced by the coupling to environments offer a rich possibility to investigate physics unique to nonequilibrium situations [1-3]. One of the key ingredients is the measurement, which has been actively investigated in the context of open quantum systems [4, 5]. Readout of measurement brings about nonequilibrium phenomena that cannot be seen in unconditional open quantum systems, such as measurement-induced phase transitions of the entanglement entropy [6-33], unconventional non-Hermitian criticality of correlation functions in the no-jump limit [34-37], and measurement-altered universality and phase transitions based on boundary conformal field theory [38-42]. While these works demonstrate the significance to study an interplay between measurements and manybody effects, the so-called postselection problem makes it difficult to experimentally observe measurement-induced guantum many-body phenomena despite efforts to evade the problem [30, 43-51]. In contrast, in quantum simulations of manybody trajectory dynamics under continuous monitoring, we can access the time and the position of quantum jumps without postselection [3].

The statistics of stochastic trajectories has been widely investigated both in classical systems [52, 53] and in open quantum systems [54, 55], where various properties have been demonstrated such as the dynamical phase transitions using large deviations [52, 56] and thermodynamic uncertainty relations [53, 57–59]. For instance, full counting statistics of quantum jumps in open systems are useful to capture manybody physics [60–64] as well as few-body physics [65]. In particular, the variance of the number of jumps contains information that is not obtained from an average of physical quantities [65]; it is related with autocorrelations and relaxation rates and has been investigated even in open quantum many-body systems [66-69]. For example, in driven dissipative many-body systems that break the detailed-balance conditions, diverging current fluctuations at the dissipative phase transition are studied [70, 71]. Recently, it has been demonstrated that space-time correlations of measurement outcomes can reveal dynamical heterogeneity before thermalization in kinetically constrained many-body models [72].

Despite previous efforts to understand quantum jump statistics in open systems, little is known about whether they can exhibit nontrivial universal many-body phenomena in the steady state even when the (unconditional) density matrix heats up to a thermal state. Such a situation naturally occurs in quantum many-body systems under particle-number measurement [21, 30], where the trivial infinite temperature state appears. In particular, it is intriguing if the measurement strength affects the fluctuations of quantum jumps even though the steady-state density matrix is independent of the measurement strength.

In this Letter, we introduce the subsystem fluctuation of quantum jumps (SFQJ) and discover its measurement-induced crossover in the steady state of continuously monitored manybody systems. In particular, we demonstrate that SFQJ obeys the Poissonian-type scaling $\propto L$ (*L* is the system size) for



FIG. 1. (a) Schematic figure of our setup. We count the number of jumps (red cross marks) in a half chain $N_{\rm jump}^{\rm half}$ and take its variance along trajectory realizations. (b) Table of the main results for the variance V and estimation uncertainty $T_{\rm th}$ concerning the number of jumps. Subsystem fluctuation of quantum jumps (SFQJ) for strong measurement exhibits anomalous yet universal super-Poissonian statistics, and the estimation uncertainty is enhanced as the system size is increased, in stark contrast to the case for the weak measurement or the total fluctuation of quantum jumps.



FIG. 2. Numerical results for the variance of quantum jumps for the Heisenberg model for 832 trajectories and T = 990. (a) Variance of net quantum jumps in the whole system [Inset: $V[N_{jump}^{total}(T)]/(\gamma L)$], (b) SFQJ, (c) $V[N_{jump}^{half}(T)]/L^2$ [Inset: $V[N_{jump}^{half}(T)]/(\gamma L^2)$], and (d) $V[N_{jump}^{half}(T)]/L$. The SFQJ exhibits a measurement-induced crossover of the system-size scaling. Data are plotted against γ for system sizes $L = 8, 10, \cdots, 18$ in (a)-(c) and against L for measurement strengths $\gamma = 0.05, 0.1, \cdots, 0.95, 1$ from bottom to top in (d). We take the average over 20 time intervals for $t \in [200, 1190], [1190, 2180], \cdots, [19010, 2 \times 10^4]$.

sufficiently weak measurement, whereas anomalous yet universal super-Poissonian-type scaling $\propto L^{\alpha}$ ($\alpha \sim 2.7$) appears for strong measurement, although the unconditional steady state is always the trivial infinite-temperature state. This crossover results in two distinct regimes in estimating the rate of quantum jumps: for strong (weak) measurement, the uncertainty is enhanced (suppressed) when the system size increases. Strikingly, the crossover is unique to SFQJ and absent for fluctuations in the whole system, where the Poissoniantype scaling always appears. We elucidate that the anomalous scaling stems from the integrated many-body autocorrelation functions for the unconditional dynamics and that the universal exponents are not determined by the Liouvillian gap. Measurement-induced many-body phenomena presented in our work can be accessed in ultracold atom experiments without postselections. See Fig. 1 for the summary of our work.

Continuous monitoring and SFQJ–We consider interacting hard-core boson chains of length L with the periodic boundary condition:

$$H = \sum_{j=1}^{L} \frac{J_{xy}}{2} (b_{j+1}^{\dagger} b_j + b_{j+1} b_j^{\dagger}) + \sum_{j=1}^{L} J_z n_{j+1} n_j, \quad (1)$$

where b_j is the bosonic annihilation operator satisfying the hard-core constraint $b_j^2 = 0$, and $n_j = b_j^{\dagger}b_j$ is the particle number operator. As the Hamiltonian (1) is exactly mapped to the XXZ spin chain with the anisotropy J_z/J_{xy} , we call this model the XXZ spin chain and set $J_{xy} = 1$ unless otherwise specified. Then, we study the dynamics under continuous monitoring of a local particle number by employing the quantum trajectory method [21, 30]. The stochastic Schrödinger equation that governs the dynamics reads [4]

$$d|\psi(t)\rangle = -iH|\psi(t)\rangle dt + \sum_{j=1}^{L} \left(\frac{n_j|\psi(t)\rangle}{\sqrt{\langle n_j\rangle}} - |\psi(t)\rangle\right) dN_j,$$
(2)

where $\langle \cdot \rangle$ denotes a quantum expectation value for the state $|\psi(t)\rangle$. Here, a discrete random variable $dN_j = 0, 1$ that counts the increment of a jump at site j is chosen according to

 $dN_j dN_k = \delta_{jk} dN_j$ and $E[dN_j] = \gamma \langle n_j \rangle dt$, where $E[\cdot]$ represents an ensemble average over the stochastic process. In the following, we assume that the initial state is prepared in the Néel state $|1010\cdots\rangle$. Importantly, the state $\rho(t)$ averaged over the measurement outcomes becomes a maximally mixed state $\rho_{ss} = I/D_0$ after long times, irrespective of measurement strengths γ . Here, I is the identity matrix and D_0 is the dimension of the Hilbert space in the half-filling sector.

As a primary quantity of interest, we introduce the net number of quantum jumps for the half-chain subsystem [see Fig. 1(a)],

$$N_{\rm jump}^{\rm half}(T) = \sum_{j=1}^{L/2} \int_{t_{\rm ss}}^{t_{\rm ss}+T} dN_j,$$
 (3)

where $t_{\rm ss}$ is sufficiently large such that $\rho(t_{\rm ss}) \simeq \rho_{\rm ss}$. We especially focus on the variance of $N_{\rm jump}^{\rm half}(T)$, or SFQJ, which is defined by

$$V[N_{\text{jump}}^{\text{half}}(T)] = E[N_{\text{jump}}^{\text{half}}(T)^2] - E[N_{\text{jump}}^{\text{half}}(T)]^2.$$
(4)

For comparison, we also define the number of quantum jumps for the whole system, $N_{jump}^{total}(T) = \sum_{j=1}^{L} \int_{t_{ss}}^{t_{ss}+T} dN_j$, and consider its variance. Note that the average jump number for $N_{total}^{jump}(T)$ is given by $E[N_{total}^{jump}(T)] = \gamma \langle \sum_{j=1}^{L} n_j \rangle T = \gamma LT/2$, which also leads to $E[N_{half}^{jump}(T)] = \gamma LT/4$ because of the symmetry. We remark that SFQJ is different from the so-called bipartite fluctuations $\langle n_{half}^2 \rangle - \langle n_{half} \rangle^2$ for $n_{half} = \sum_{i=1}^{L/2} n_i$ [48].

Measurement-induced crossover–We first demonstrate the measurement-induced crossover of SFQJ by numerically studying the Heisenberg model $(J_z = 1)$. As shown in Fig. 2(a), we see that $V[N_{jump}^{total}(T)]$ exhibits a trivial scaling proportional to γL , irrespective of γ . On the other hand, SFQJ shown in Fig. 2(b) does not behave linearly against γ nor L, and we find a crossing in $V[N_{jump}^{half}(T)]/L^2$ around $\gamma_c \sim 0.5$ as seen in Fig. 2(c). This indicates a measurement-induced crossover of SFQJ in the system-size scaling, and in fact in Fig. 2(d), we see that the scaling is estimated to show a crossover from $\propto L^{1.13}$ for $\gamma = 0.05$ to $\propto L^{2.45}$ for $\gamma = 1$. However, we note that universal exponents are different from

these values due to the finite-size effect. This is because SFQJ is given as a sum of two terms whose L dependence is different as detailed later, which leads to the deviations of exponents from the universal ones for finite-system sizes. We can conduct the same numerical simulation for XX ($J_z = 0$) and XXZ ($J_z = 0.5, 1.5$) models as shown in the Supplemental Material [73]. All these models exhibit the measurementinduced crossover, signifying an anomalous enhancement of SFQJ in many-body quantum systems under continuous monitoring. We also mention that such an anomalous scaling does not exist in the Ising model ($J_{xy} = 0, J_z = 1$) [73], because the particle number is a conserved quantity at a single site.

Relation to unconditional dynamics–We examine the measurement-induced crossover with the help of the unconditional dynamics of Eq. (2) and clarify that SFQJ is composed of two terms: dynamical activity with the volume law that governs SFQJ for sufficiently weak measurement strength, and an integrated autocorrelation function that causes the anomalous enhancement of SFQJ. First of all, we should beware that SFQJ (4) is unique to quantum trajectories because measurement outcomes are averaged over all possible sequences in the unconditional dynamics. However, the counting variable is still evaluated by using the Liouvillian, and we can write down SFQJ in terms of the ensemble-averaged dynamics [65]. To start with, we calculate the noise $D_{half}(t) = dV[N_{half}^{half}(t)]/dt$ for a subsystem in the steady state as

$$D_{\text{half}}(T) = K_{\text{half}} + 2 \int_0^T d\tau \left\{ \text{Tr}[\mathcal{J}_{\text{half}} e^{\mathcal{L}\tau} \mathcal{J}_{\text{half}} \rho_{\text{ss}}] - J_{\text{half}}^2 \right\}, \quad (5)$$

where $\mathcal{L}(\rho) = -i[H,\rho] + \gamma \sum_{i} (-\frac{1}{2}\{n_i,\rho\} + n_i\rho n_i)$ is the Liouvillian superoperator for the ensemble-averaged dynamics of Eq. (2), the subsystem superoperator \mathcal{J}_{half} is defined as $\mathcal{J}_{half}\rho(t) \equiv \gamma \sum_{i=1}^{L/2} n_i\rho(t)n_i$, and $K_{half} \equiv \gamma \sum_{i=1}^{L/2} \text{Tr}[n_i\rho_{ss}]$ and $J_{half} \equiv \text{Tr}[\mathcal{J}_{half}\rho_{ss}]$ both reduce to $K_{half} = J_{half} = \gamma L/4$, which is nothing but (half of) the dynamical activity in the steady state, $E[dN_{jump}^{total}(t)]/dt = 2E[dN_{jump}^{half}(t)]/dt = \gamma L/2$. By numerically calculating the long-time dynamics in Eq. (5), we find that the noise becomes constant for large T and does not involve a contribution of $\mathcal{O}(T)$ [65]. Then, we obtain SFQJ as $V[N_{jump}^{half}(T)] = D_{half}T$, which leads to

$$V[N_{\text{jump}}^{\text{half}}(T)] = V_{\text{act}}(T) + V_{\text{anom}}(T),$$

$$= \frac{\gamma LT}{4} + 2\gamma^2 T \int_0^T d\tau \langle n'_{\text{half}}(\tau) n'_{\text{half}} \rangle_{\infty},$$

(6)

where $n'_{\text{half}}(\tau) = n_{\text{half}}(\tau) - \langle n_{\text{half}} \rangle_{\infty}$, the Heisenberg picture for the adjoint Liouvillian $\tilde{\mathcal{L}}$ is represented as $n_{\text{half}}(\tau) = e^{\tilde{\mathcal{L}}\tau} n_{\text{half}}$, and $\langle \cdot \rangle_{\infty}$ stands for the expectation value with respect to ρ_{ss} . As shown below, we demonstrate that the super-Poissonian-type anomalous enhancement of SFQJ originates

from the integrated autocorrelation function

$$\mathcal{C}_{\text{auto}}^{\text{half}} \equiv \int_0^T d\tau C_{\text{auto}}^{\text{half}}(\tau) \equiv \int_0^T d\tau \langle n'_{\text{half}}(\tau) n'_{\text{half}} \rangle_{\infty}.$$
 (7)

Here, the positive (zero) correlation $V[N_{jump}^{half}(T)] > E[N_{jump}^{half}(T)]$ ($V[N_{jump}^{half}(T)] = E[N_{jump}^{half}(T)]$) is referred to as the super-Poissonian (Poissonian) statistics. We remark that, as the particle number in the whole chain is a conserved quantity, the anomalous term is absent for the whole system, and the variance $V[N_{jump}^{total}(T)]$ reduces to the dynamical activity, which shows a Poissonian-type volume law irrespective of the measurement strength.

Origin of the measurement-induced crossover–On the basis of the quantum trajectory method, we numerically simulate the Heisenberg chain, obtaining the contribution of V_{anom} and V_{act} to SFQJ [74]. As shown in Fig. 3(a), we find that V_{anom} and V_{act} cross at the critical measurement strength γ_c , and the leading contribution to SFQJ is changed from V_{act} (Poissonian) to V_{anom} (super-Poissonian) as γ is increased. This causes a measurement-induced crossover of SFQJ illustrated in Fig. 2. In Fig. 3(b) and also in other parameters of J_z [73], we find by performing the finite-size scaling analysis for $L = 8, 10, \dots, 18$ that

$$V_{\rm anom} \propto L^{\alpha},$$
 (8)

where $\alpha = 2.69, 2.71, 2.73, 2.68$ for the Heisenberg, XX, and XXZ ($J_z = 0.5, 1.5$) model for $\gamma = 1$, respectively. Remarkably, this indicates that the anomalous scaling of SFQJ is characterized by the universal exponent $\alpha \sim 2.7$. Since V_{anom} is also proportional to $\gamma^2 c(\gamma)$, where $c(\gamma)$ is an increasing polynomial function of γ with $c(0) \geq 0$ (see Supplemental Material [73] for the γ -dependence of SFQJ), SFQJ is dominated by $V_{\text{anom}} \propto L^{\alpha}$ for strong $\gamma = \mathcal{O}(L^0)$ instead of $V_{\text{act}} \propto L$ for weak γ [which should be $o(L^0)$], the fact of which is the direct origin of the crossover of SFQJ [see Fig. 1(b)]. In agreement with this, we see that the crossing value γ_c shifts towards small γ as seen in Fig. 3(a). Physically, such an enhancement of SFQJ is regarded as a kind of Zeno effects [75], which indicate that, if a quantum jump occurs in a subsystem, the next jump is also likely to occur in the same subsystem.



FIG. 3. Numerical results for $V_{\rm anom}$ and $V_{\rm act}$ based on the quantum trajectory method for the Heisenberg model. (a) The leading contribution of $V_{\rm anom}$ and $V_{\rm act}$ changes at a critical measurement strength. (b) System-size scaling of $V_{\rm anom}/L$. Data are shifted by 2000 as L is increased as $L = 8, 10, \cdots, 18$ in (a) to improve visualization, and measurement strengths are $\gamma = 0.05, 0.1, \cdots, 0.95, 1$ from bottom to top in (b). The parameters and methods are the same as in Fig. 2.



FIG. 4. (a) V_{anom} and V_{act} for the Heisenberg model for system sizes L = 4, 6, 8 obtained with the use of Eq. (14) [76]. The leading contribution of V_{anom} and V_{act} changes at a critical measurement strength consistent with Fig. 3. (b) Exponents a, b, and c of the Liouvillian gap $\Delta \propto 1/L^a$ (blue), $C_{\text{auto}}^{\text{single}} \propto L^b$ (green), and $C_{\text{half}}^{\text{half}} \propto L^c$ (red) obtained from the finite-size scaling analysis with L = 8, 10, 12, 14 for the Heisenberg model. The anomalous scaling of SFQJ cannot be captured by the Liouvillian gap. Data are shifted by 1000 as L is increased in (a).

Impact on the precision–One of the interesting consequences of the above anomalous scaling is the drastically enhanced uncertainty in estimating the rate of quantum jumps occurring in the subsystem. Let us estimate the rate of quantum jumps in the subsystem by $\bar{I}_{half}(T) = N_{jump}^{half}(T)/T$ [65]. We then obtain

$$E[\bar{I}_{\text{half}}(T)] = \frac{E[N_{\text{jump}}^{\text{half}}(T)]}{T} = J_{\text{half}} \propto \gamma L, \qquad (9)$$

$$V[\bar{I}_{\text{half}}(T)] = \frac{V[N_{\text{jump}}^{\text{half}}(T)]}{T^2} = \frac{D_{\text{half}}}{T} \propto \frac{\gamma L + \gamma^2 c(\gamma) L^{\alpha}}{T}.$$
(10)

To estimate the true rate $E[\bar{I}_{half}(T)]$ accurately, we shall evaluate the estimation uncertainty defined by the (squared) coefficient of variation

$$\frac{V[\bar{I}_{\text{half}}(T)]}{(E[\bar{I}_{\text{half}}(T)])^2} = \frac{D_{\text{half}}}{J_{\text{half}}^2 T} \equiv \frac{T_{\text{th}}}{T},$$
(11)

where $T_{\rm th} \equiv D_{\rm half}/J_{\rm half}^2$ is frequently discussed in thermodynamic and kinetic uncertainty relations [65]; here, we rather focus on the *L*-dependence in the large-*T* dynamics. For sufficiently weak γ , we find

$$T_{\rm th} \propto L^{-1}.$$
 (12)

This means that the error for evaluating $E[\bar{I}_{half}(T)]$ diminishes fast enough, and $E[\bar{I}_{half}(T)]$ is accurately determined for larger system sizes. On the other hand, for strong measurement satisfying $\gamma \sim \mathcal{O}(L^0)$, we obtain

$$T_{\rm th} \propto L^{\alpha - 2},$$
 (13)

with $\alpha \sim 2.7$, which means that the uncertainty in estimating $E[\bar{I}_{\rm half}(T)]$ is enhanced as the system size is increased [see Fig. 1(b)]. We remark that, as the estimation uncertainty in Eq. (13) is proportional to $L^{\alpha-2}$, particular super-Poissonian-type variance satisfying $\alpha > 2$ enhances $T_{\rm th}$ with increasing L. Moreover, the estimation uncertainty (11) is related to the

ergodicity of a quantum trajectory for the jump statistics [77], i.e., the concept that the long-time average of a single realization yields the ensemble average in almost all trajectories. Equation (13) means that the ergodicity of a quantum trajectory tends to be disturbed for larger system sizes in XXZ spin chains under continuous monitoring. We note that the fluctuation for a whole system always follows the trivial Poissonian statistics, whose estimation uncertainty reduces to Eq. (12) for arbitrary γ .

Spectral decomposition of the Liouvillian–To further analyze the anomalous behavior of SFQJ, we calculate the Liouvillian gap as well as V_{anom} and V_{act} by using the spectral decomposition of the Liouvillian for the unconditional dynamics of Eq. (2), assuming the diagonalizability of \mathcal{L} . By employing the basis transformation to the doubled Hilbert space for the density matrix $\rho = \sum_{i,j} \rho_{ij} |i\rangle \langle j|$ as $|i\rangle \langle j| \mapsto |ij\rangle :=$ $|i\rangle \otimes |j\rangle \in \mathcal{H} \otimes \mathcal{H}$ [65, 78–80], Eq. (6) is rewritten as

$$V[N_{\text{jump}}^{\text{half}}(T)] = \frac{\gamma LT}{4} + \frac{2\gamma^2 T}{D_0} \sum_{j \neq 0} \frac{e^{\lambda_j T} - 1}{\lambda_j} (n_{\text{half}}'|\rho_j^R) (\rho_j^L|n_{\text{half}}'), \quad (14)$$

where the spectral decomposition of the Liouvillian is given by $\mathcal{L} = \sum_{j \neq 0} \lambda_j |\rho_j^R| (\rho_j^L)$, the inner product is defined as $(A|B) = \text{Tr}[A^{\dagger}B]$, and we have restricted both the bra and the ket space to a half-filling sector. As shown in Fig. 4(a), we obtain V_{anom} and V_{act} with the use of Eq. (14). Clearly, we find a crossing of V_{anom} and V_{act} and the critical γ_c quantitatively agrees well with that in Fig. 3(a) for L = 8.

Finally, since the asymptotic behavior of the autocorrelation function is dominated by the Liouvillian gap as $|C_{auto}^{single}(\tau)| \equiv |\langle n'_1(\tau)n'_1\rangle_{\infty}| \sim e^{-\Delta\tau} \ (\tau \to \infty)$ [81, 82], it is illustrative to what extent the scaling (8) is affected by the Liouvillian gap by comparing the exponent of $C_{auto}^{single} \equiv \int_0^T d\tau C_{auto}^{single}(\tau)$ with that solely extracted from the Liouvillian gap; $C_{auto}^{single} \stackrel{?}{\sim} \int_0^T e^{-\Delta\tau} \sim 1/\Delta$ [83]. By sorting the eigenvalues to satisfy $0 = \lambda_0 > \operatorname{Re}[\lambda_1] \ge \operatorname{Re}[\lambda_2] \ge \cdots \ge \operatorname{Re}[\lambda_{D_0^2-1}]$, where we confirm that the unconditional steady state is unique, the Liouvillian gap Δ that dominates the asymptotic decay rate [84, 85] is extracted from the largest real part of the eigenvalue except for λ_0 as $\Delta = -\operatorname{Re}\lambda_1$.

In Fig. 4(b), we depict the exponent of the inverse Liouvillian gap obtained from the exact diagonalization and that of C_{auto}^{single} calculated from the quantum trajectory method for the Heisenberg model. We find that, the Liouvillian gap estimated from the finite-size scaling analysis reads $\Delta \propto 1/L^{1.95}$ for $\gamma = 1$, which is close to the analytical result $\Delta \propto 1/L^2$ for $\gamma \gg 1/L$ [86–88]. On the other hand, we obtain $C_{auto}^{single} \propto L^{0.74}$ for $\gamma = 1$, which indicates that the scaling for the integrated autocorrelation function is largely governed by the transient dynamics [82] rather than the Liouvillian gap. We emphasize that, as the dynamical activity is proportional to L^0 for a single site, the exponent for C_{auto}^{single} is also anomalous. Moreover, the uncertainty in estimating the local quantity $E[\bar{I}_{single}(T)] \equiv E[\int_{t_{ss}}^{t_{ss}+T} dN_1]/T$ is enhanced when L

is increased. As for C_{auto}^{half} shown in Fig. 4(b), which reads $C_{auto}^{half} \propto L^{2.65}$ for $\gamma = 1$ [89], we find that not only the diagonal elements corresponding to C_{auto}^{single} but also off-diagonal components $C_{auto}^{ij} \equiv \int_0^T d\tau \langle n'_i(\tau)n'_j \rangle_{\infty}$ $(i \neq j)$ therein contribute to the scaling of C_{auto}^{half} because $C_{auto}^{half} \gg L/2 \cdot C_{auto}^{single}$ holds. Thus, we cannot estimate the anomalous scaling exponent for C_{auto}^{half} given in Eq. (8) by the Liouvillian gap. Such a discrepancy between the scalings of C_{auto}^{single} and the Liouvillian gap stems from the contribution from eigenvalues governing the transient dynamics in Eq. (14).

Experimental proposal and conclusions–Our main discoveries, i.e., the measurement-induced crossover of SFQJ and its anomalous scaling, are accessible in experiments without postselection because the only information we need is the time and the position of quantum jumps. For example, in ultracold atoms, SFQJ is obtained by recording the photon scattering with a probe light [90, 91], which can be routinely done with quantum-gas microscopes [92, 93]. We then realize the measurement-induced crossover by tuning the light intensity. We also suggest to change the system size instead by fixing the probe intensity; SFQJ shows the volume law for sufficiently small system sizes, whereas it exhibits anomalous universal scaling $\propto L^{\alpha}$ ($\alpha \sim 2.7$) for large system sizes.

In this Letter, we have discovered hitherto unnoticed crossover of measurement-induced many-body properties free from postselection, even though the unconditional steady state trivially becomes a maximally mixed state insensitive to measurement strength. To highlight this, we have shown that SFQJ serves as a new indicator for quantum jumps in manybody systems under measurement, in stark contrast to the fluctuation in the whole system. The next step to proceed with would be the evaluation of the universality class and investigation of the impact of dissipative phase transitions [94] on the anomalous scaling in prototypical many-body models. The generalization of our setup to quantum circuits under measurement also deserves future study. In addition, it is worth studying the relation between our results and well-known universal scalings in many-body problems, e.g., long-time tails in hydrodynamics [95].

Acknowledgments-We are grateful to Hironobu Yoshida and Kazuya Fujimoto for fruitful discussions. This work was supported by JST ERATO Grant No. JPMJER2302, and KAKENHI Grants No. JP23K19031 and No. JP24K16982. K.Y. was also supported by Toyota RIKEN Scholar Program, Murata Science and Education Foundation, Public Promoting Association Kura Foundation, Hirose Foundation, and the Precise Measurement Technology Promotion Foundation. The numerical calculations were partly carried out with the help of QuSpin [96].

- * yamamoto@phys.sci.isct.ac.jp
- [†] Former Tokyo Institute of Technology. From October 2024, Tokyo Institute of Technology and Tokyo Medical and Den-

tal University have merged and become Institute of Science Tokyo.

- M. Müller, S. Diehl, G. Pupillo, and P. Zoller, Engineered open systems and quantum simulations with atoms and ions, Adv. Atom. Mol. Opt. Phys. 61, 1 (2012).
- P. M. Harrington, E. J. Mueller, and K. W. Murch, Engineered dissipation for quantum information science, Nat. Rev. Phys. 4, 660 (2022).
- [3] R. Fazio, J. Keeling, L. Mazza, and M. Schiro, Many-body open quantum systems, arXiv:2409.10300.
- [4] A. J. Daley, Quantum trajectories and open many-body quantum systems, Adv. Phys. 63, 77 (2014).
- [5] Y. Ashida, Z. Gong, and M. Ueda, Non-Hermitian physics, Adv. Phys. 69, 249 (2020).
- [6] M. P. Fisher, V. Khemani, A. Nahum, and S. Vijay, Random quantum circuits, Annu. Rev. Condens. Matter Phys. 14, 335 (2023).
- [7] Y. Li, X. Chen, and M. P. A. Fisher, Quantum Zeno effect and the many-body entanglement transition, Phys. Rev. B 98, 205136 (2018).
- [8] A. Chan, R. M. Nandkishore, M. Pretko, and G. Smith, Unitary-projective entanglement dynamics, Phys. Rev. B 99, 224307 (2019).
- [9] B. Skinner, J. Ruhman, and A. Nahum, Measurement-Induced Phase Transitions in the Dynamics of Entanglement, Phys. Rev. X 9, 031009 (2019).
- [10] Y. Li, X. Chen, and M. P. A. Fisher, Measurement-driven entanglement transition in hybrid quantum circuits, Phys. Rev. B 100, 134306 (2019).
- [11] C. Noel, P. Niroula, D. Ahu, A. Risinger, L. Egan, D. Biswas, M. Cetina, A. V. Gorshkov, M. Gullans, D. A. Huse, and C. Monroe, Measurement-induced quantum phases realized in a trapped-ion quantum computer, Nat. Phys. 18, 760 (2022).
- [12] J. M. Koh, S.-N. Sun, M. Motta, and A. J. Minnich, Measurement-induced entanglement phase transition on a superconducting quantum processor with mid-circuit readout, Nat. Phys. 19, 1314 (2023).
- [13] Google Quantum AI, Measurement-induced entanglement and teleportation on a noisy quantum processor, Nature 622, 481 (2023).
- [14] U. Agrawal, J. Lopez-Piqueres, R. Vasseur, S. Gopalakrishnan, and A. C. Potter, Observing Quantum Measurement Collapse as a Learnability Phase Transition, Phys. Rev. X 14, 041012 (2024).
- [15] X. Cao, A. Tilloy, and A. De Luca, Entanglement in a fermion chain under continuous monitoring, SciPost Phys. 7, 024 (2019).
- [16] O. Alberton, M. Buchhold, and S. Diehl, Entanglement Transition in a Monitored Free-Fermion Chain: From Extended Criticality to Area Law, Phys. Rev. Lett. **126**, 170602 (2021).
- [17] X. Turkeshi, A. Biella, R. Fazio, M. Dalmonte, and M. Schiró, Measurement-induced entanglement transitions in the quantum Ising chain: From infinite to zero clicks, Phys. Rev. B 103, 224210 (2021).
- [18] X. Turkeshi, M. Dalmonte, R. Fazio, and M. Schirò, Entanglement transitions from stochastic resetting of non-Hermitian quasiparticles, Phys. Rev. B 105, L241114 (2022).
- [19] G. Piccitto, A. Russomanno, and D. Rossini, Entanglement transitions in the quantum Ising chain: A comparison between different unravelings of the same Lindbladian, Phys. Rev. B 105, 064305 (2022).
- [20] Q. Tang and W. Zhu, Measurement-induced phase transition: A case study in the nonintegrable model by density-matrix renormalization group calculations, Phys. Rev. Research 2,

013022 (2020).

- [21] Y. Fuji and Y. Ashida, Measurement-induced quantum criticality under continuous monitoring, Phys. Rev. B 102, 054302 (2020).
- [22] M. Szyniszewski, A. Romito, and H. Schomerus, Universality of Entanglement Transitions from Stroboscopic to Continuous Measurements, Phys. Rev. Lett. **125**, 210602 (2020).
- [23] O. Lunt and A. Pal, Measurement-induced entanglement transitions in many-body localized systems, Phys. Rev. Res. 2, 043072 (2020).
- [24] S.-K. Jian, C. Liu, X. Chen, B. Swingle, and P. Zhang, Measurement-Induced Phase Transition in the Monitored Sachdev-Ye-Kitaev Model, Phys. Rev. Lett. **127**, 140601 (2021).
- [25] M. Van Regemortel, Z.-P. Cian, A. Seif, H. Dehghani, and M. Hafezi, Entanglement Entropy Scaling Transition under Competing Monitoring Protocols, Phys. Rev. Lett. **126**, 123604 (2021).
- [26] E. V. H. Doggen, Y. Gefen, I. V. Gornyi, A. D. Mirlin, and D. G. Polyakov, Generalized quantum measurements with matrix product states: Entanglement phase transition and clusterization, Phys. Rev. Research 4, 023146 (2022).
- [27] T. Minato, K. Sugimoto, T. Kuwahara, and K. Saito, Fate of Measurement-Induced Phase Transition in Long-Range Interactions, Phys. Rev. Lett. **128**, 010603 (2022).
- [28] T. Müller, S. Diehl, and M. Buchhold, Measurement-Induced Dark State Phase Transitions in Long-Ranged Fermion Systems, Phys. Rev. Lett. **128**, 010605 (2022).
- [29] M. Buchhold, Y. Minoguchi, A. Altland, and S. Diehl, Effective Theory for the Measurement-Induced Phase Transition of Dirac Fermions, Phys. Rev. X 11, 041004 (2021).
- [30] K. Yamamoto and R. Hamazaki, Localization properties in disordered quantum many-body dynamics under continuous measurement, Phys. Rev. B 107, L220201 (2023).
- [31] M. Szyniszewski, O. Lunt, and A. Pal, Disordered monitored free fermions, Phys. Rev. B 108, 165126 (2023).
- [32] K. Mochizuki and R. Hamazaki, Measurement-Induced Spectral Transition, Phys. Rev. Lett. 134, 010410 (2025).
- [33] T. Matsubara, K. Yamamoto, and A. Koga, to be submitted.
- [34] Y. Ashida, S. Furukawa, and M. Ueda, Quantum critical behavior influenced by measurement backaction in ultracold gases, Phys. Rev. A 94, 053615 (2016).
- [35] Y. Ashida, S. Furukawa, and M. Ueda, Parity-time-symmetric quantum critical phenomena, Nat. Commun. 8, 15791 (2017).
- [36] K. Yamamoto, M. Nakagawa, M. Tezuka, M. Ueda, and N. Kawakami, Universal properties of dissipative Tomonaga-Luttinger liquids: Case study of a non-Hermitian XXZ spin chain, Phys. Rev. B 105, 205125 (2022).
- [37] K. Yamamoto and N. Kawakami, Universal description of dissipative Tomonaga-Luttinger liquids with SU(N) spin symmetry: Exact spectrum and critical exponents, Phys. Rev. B **107**, 045110 (2023).
- [38] S. J. Garratt, Z. Weinstein, and E. Altman, Measurements conspire nonlocally to restructure critical quantum states, Phys. Rev. X 13, 021026 (2023).
- [39] X. Sun, H. Yao, and S.-K. Jian, New critical states induced by measurement, arXiv:2301.11337.
- [40] Q. Tang and X. Wen, A critical state under weak measurement is not critical, arXiv:2411.13705.
- [41] Y. Ashida, S. Furukawa, and M. Oshikawa, Systemenvironment entanglement phase transitions, Phys. Rev. B 110, 094404 (2024).
- [42] Y. Liu, S. Murciano, D. F. Mross, and J. Alicea, Boundary transitions from a single round of measurements on gapless

quantum states, arXiv:2412.07830.

- [43] M. J. Gullans and D. A. Huse, Scalable Probes of Measurement-Induced Criticality, Phys. Rev. Lett. 125, 070606 (2020).
- [44] Y. Li, Y. Zou, P. Glorioso, E. Altman, and M. P. A. Fisher, Cross Entropy Benchmark for Measurement-Induced Phase Transitions, Phys. Rev. Lett. 130, 220404 (2023).
- [45] S. J. Garratt and E. Altman, Probing postmeasurement entanglement without postselection, PRX Quantum 5, 030311 (2024).
- [46] M. Ippoliti and V. Khemani, Postselection-free entanglement dynamics via spacetime duality, Phys. Rev. Lett. 126, 060501 (2021).
- [47] T.-C. Lu and T. Grover, Spacetime duality between localization transitions and measurement-induced transitions, PRX Quantum 2, 040319 (2021).
- [48] A. G. Moghaddam, K. Pöyhönen, and T. Ojanen, Exponential shortcut to measurement-induced entanglement phase transitions, Phys. Rev. Lett. **131**, 020401 (2023).
- [49] G. Passarelli, X. Turkeshi, A. Russomanno, P. Lucignano, M. Schirò, and R. Fazio, Many-body dynamics in monitored atomic gases without postselection barrier, Phys. Rev. Lett. 132, 163401 (2024).
- [50] M. McGinley, Postselection-free learning of measurementinduced quantum dynamics, PRX Quantum 5, 020347 (2024).
- [51] X. Feng, J. C., S. Kourtis, and B. Skinner, Postselection-free experimental observation of the measurement-induced phase transition in circuits with universal gates, arXiv:2502.01735.
- [52] J. P. Garrahan, R. L. Jack, V. Lecomte, E. Pitard, K. van Duijvendijk, and F. van Wijland, Dynamical First-Order Phase Transition in Kinetically Constrained Models of Glasses, Phys. Rev. Lett. 98, 195702 (2007).
- [53] J. P. Garrahan, Simple bounds on fluctuations and uncertainty relations for first-passage times of counting observables, Phys. Rev. E 95, 032134 (2017).
- [54] R. Chetrite and H. Touchette, Nonequilibrium Markov processes conditioned on large deviations, in *Annales Henri Poincaré*, Vol. 16 (Springer, 2015) pp. 2005–2057.
- [55] J. P. Garrahan, Aspects of non-equilibrium in classical and quantum systems: Slow relaxation and glasses, dynamical large deviations, quantum non-ergodicity, and open quantum dynamics, Physica A: Stat. Mech. Appl. **504**, 130 (2018).
- [56] J. P. Garrahan and I. Lesanovsky, Thermodynamics of Quantum Jump Trajectories, Phys. Rev. Lett. 104, 160601 (2010).
- [57] A. C. Barato and U. Seifert, Thermodynamic Uncertainty Relation for Biomolecular Processes, Phys. Rev. Lett. 114, 158101 (2015).
- [58] F. Carollo, R. L. Jack, and J. P. Garrahan, Unraveling the Large Deviation Statistics of Markovian Open Quantum Systems, Phys. Rev. Lett. **122**, 130605 (2019).
- [59] J. M. Horowitz and T. R. Gingrich, Thermodynamic uncertainty relations constrain non-equilibrium fluctuations, Nat. Phys. 16, 15 (2020).
- [60] M. Žnidarič, Exact Large-Deviation Statistics for a Nonequilibrium Quantum Spin Chain, Phys. Rev. Lett. 112, 040602 (2014).
- [61] J. M. Hickey, C. Flindt, and J. P. Garrahan, Trajectory phase transitions and dynamical Lee-Yang zeros of the Glauber-Ising chain, Phys. Rev. E 88, 012119 (2013).
- [62] I. Lesanovsky, M. van Horssen, M. Guță, and J. P. Garrahan, Characterization of Dynamical Phase Transitions in Quantum Jump Trajectories Beyond the Properties of the Stationary State, Phys. Rev. Lett. **110**, 150401 (2013).
- [63] F. Carollo, J. P. Garrahan, I. Lesanovsky, and C. Pérez-

Espigares, Making rare events typical in markovian open quantum systems, Phys. Rev. A 98, 010103 (2018).

- [64] Z.-K. Liu, K.-H. Sun, A. Cabot, F. Carollo, J. Zhang, Z.-Y. Zhang, L.-H. Zhang, B. Liu, T.-Y. Han, Q. Li, Y. Ma, H.-C. Chen, I. Lesanovsky, D.-S. Ding, and B.-S. Shi, Emergence of subharmonics in a microwave driven dissipative rydberg gas, Phys. Rev. Res. 6, L032069 (2024).
- [65] G. T. Landi, M. J. Kewming, M. T. Mitchison, and P. P. Potts, Current Fluctuations in Open Quantum Systems: Bridging the Gap Between Quantum Continuous Measurements and Full Counting Statistics, PRX Quantum 5, 020201 (2024).
- [66] C. Ates, B. Olmos, J. P. Garrahan, and I. Lesanovsky, Dynamical phases and intermittency of the dissipative quantum ising model, Phys. Rev. A 85, 043620 (2012).
- [67] B. Buča and T. Prosen, Exactly Solvable Counting Statistics in Open Weakly Coupled Interacting Spin Systems, Phys. Rev. Lett. 112, 067201 (2014).
- [68] M. Žnidarič, Anomalous nonequilibrium current fluctuations in the Heisenberg model, Phys. Rev. B 90, 115156 (2014).
- [69] M. Žnidarič, Large-deviation statistics of a diffusive quantum spin chain and the additivity principle, Phys. Rev. E 89, 042140 (2014).
- [70] M. J. Kewming, M. T. Mitchison, and G. T. Landi, Diverging current fluctuations in critical Kerr resonators, Phys. Rev. A 106, 033707 (2022).
- [71] M. Matsumoto, M. Baggioli, and Z. Cai, Dissipative quantum phase transitions monitored by current fluctuations, arXiv:2502.01136.
- [72] M. Cech, M. Cea, M. C. Bañuls, I. Lesanovsky, and F. Carollo, Space-time correlations in monitored kinetically constrained discrete-time quantum dynamics, arXiv:2408.09872.
- [73] See Supplemental Material, which includes Refs. [97-99], for the numerical results on SFQJ, $V_{\rm anom}$ and $V_{\rm act}$ in other manybody models, the γ -dependence of V_{anom} , and the integrated autocorrelation functions and the Liouvillian gap in Heisenberg and XX models.
- [74] We calculate $V_{act}(T)$ and $V_{anom}(T)$ with the help of $V[N_{\text{jump}}^{\text{total}}(T)]$ as $V_{\text{act}}(T) = V[N_{\text{jump}}^{\text{total}}(T)]/2$ and $V_{\text{anom}}(T) = V[N_{\text{jump}}^{\text{half}}(T)] - V[N_{\text{jump}}^{\text{total}}(T)]/2.$
- [75] W. M. Itano, D. J. Heinzen, J. J. Bollinger, and D. J. Wineland, Quantum Zeno effect, Phys. Rev. A 41, 2295 (1990).
- [76] We set T = 990 and have numerically checked that C_{auto}^{half} becomes almost constant for large T; we arrive at the same result in the $T \to \infty$ limit.
- [77] J. Cresser, Ergodicity of quantum trajectory detection records, in Directions in Quantum Optics: A Collection of Papers Dedicated to the Memory of Dan Walls Including Papers Presented at the TAMU-ONR Workshop Held at Jackson, Wvoming, USA, 26-30 July 1999 (Springer, 2001) pp. 358-369.
- [78] N. Yoshioka and R. Hamazaki, Constructing neural stationary states for open quantum many-body systems, Phys. Rev. B 99, 214306 (2019).
- [79] N. Shibata and H. Katsura, Dissipative spin chain as a non-Hermitian Kitaev ladder, Phys. Rev. B 99, 174303 (2019).
- [80] K. Yamamoto, M. Nakagawa, N. Tsuji, M. Ueda, and N. Kawakami, Collective Excitations and Nonequilibrium Phase Transition in Dissipative Fermionic Superfluids, Phys. Rev. Lett. 127, 055301 (2021).
- [81] T. Mori and T. Shirai, Symmetrized Liouvillian Gap in Markovian Open Quantum Systems, Phys. Rev. Lett. 130, 230404 (2023).
- [82] T. Shirai and T. Mori, Accelerated Decay due to Operator Spreading in Bulk-Dissipated Quantum Systems, Phys. Rev.

- Lett. **133**, 040201 (2024). [83] We obtain the same C_{auto}^{single} for site 2 instead of taking site 1. Thus, we find a translational invariance for C_{auto}^{single} .
- [84] T. Mori and T. Shirai, Resolving a Discrepancy between Liouvillian Gap and Relaxation Time in Boundary-Dissipated Quantum Many-Body Systems, Phys. Rev. Lett. 125, 230604 (2020).
- [85] T. Haga, M. Nakagawa, R. Hamazaki, and M. Ueda, Liouvillian Skin Effect: Slowing Down of Relaxation Processes without Gap Closing, Phys. Rev. Lett. 127, 070402 (2021).
- [86] Z. Cai and T. Barthel, Algebraic versus Exponential Decoherence in Dissipative Many-Particle Systems, Phys. Rev. Lett. 111, 150403 (2013).
- [87] M. Žnidarič, Relaxation times of dissipative many-body quantum systems, Phys. Rev. E 92, 042143 (2015).
- [88] As the Lindblad equation is invariant under the transformation by a real constant in the particle-number jump operator [100], we refer to the Liouvillian gap in the Heisenberg model with dephasing [86, 87].
- [89] The exponent 2.65 deviates a little from the universal exponent $\alpha = 2.69$ for the Heisenberg model. This stems from the system-size difference used in the calculation, where the latter exponent is obtained from the finite-size scaling analysis for $L = 8, \dots, 18$ in Fig. 3(b), while the former one for $L = 8, \dots, 14$ in Fig. 4(b). In the former calculation, we need to compare it with the exponent of the Liouvillian gap, where the exact diagonalization of the Liouvillian numerically limits the system size.
- [90] H. P. Lüschen, P. Bordia, S. S. Hodgman, M. Schreiber, S. Sarkar, A. J. Daley, M. H. Fischer, E. Altman, I. Bloch, and U. Schneider, Signatures of many-body localization in a controlled open quantum system, Phys. Rev. X 7, 011034 (2017).
- [91] Y. S. Patil, S. Chakram, and M. Vengalattore, Measurement-Induced Localization of an Ultracold Lattice Gas, Phys. Rev. Lett. 115, 140402 (2015).
- [92] W. S. Bakr, J. I. Gillen, A. Peng, S. Fölling, and M. Greiner, A quantum gas microscope for detecting single atoms in a hubbard-regime optical lattice, Nature 462, 74 (2009).
- [93] J. F. Sherson, C. Weitenberg, M. Endres, M. Cheneau, I. Bloch, and S. Kuhr, Single-atom-resolved fluorescence imaging of an atomic mott insulator, Nature 467, 68 (2010).
- [94] F. Minganti, A. Biella, N. Bartolo, and C. Ciuti, Spectral theory of Liouvillians for dissipative phase transitions, Phys. Rev. A 98, 042118 (2018).
- [95] A. Bohrdt, C. B. Mendl, M. Endres, and M. Knap, Scrambling and thermalization in a diffusive quantum many-body system, New J. Phys. 19, 063001 (2017).
- [96] P. Weinberg and M. Bukov, QuSpin: a Python package for dynamics and exact diagonalisation of quantum many body systems part I: spin chains, SciPost Phys. 2, 003 (2017).
- [97] K. Yamamoto, Y. Ashida, and N. Kawakami, Rectification in nonequilibrium steady states of open many-body systems, Phys. Rev. Res. 2, 043343 (2020).
- [98] T. Haga, M. Nakagawa, R. Hamazaki, and M. Ueda, Quasiparticles of decoherence processes in open quantum many-body systems: Incoherentons, Phys. Rev. Res. 5, 043225 (2023).
- [99] T. Ishiyama, F. Kazuya, and T. Sasamoto, Exact density profile in a tight-binding chain with dephasing noise, arXiv:2501.07095 (2025).
- [100] H.-P. Breuer and F. Petruccione, The theory of open quantum systems (Oxford University Press, USA, 2002).

Supplemental Material for "Measurement-Induced Crossover of Quantum Jump Statistics in Postselection-Free Many-Body Dynamics"

Numerical results for subsystem fluctuations of quantum jumps in other many-body models



FIG. S1. Numerical results for the variance of quantum jumps to demonstrate the measurement-induced crossover of SFQJ in the XX model for 832 trajectories and T = 990. Data are plotted against γ for system sizes $L = 8, 10, \dots, 18$ in (a)-(c) and against L for measurement strengths $\gamma = 0.05, 0.1, \dots, 0.95, 1$ from bottom to top in (d). (a) Variance of net quantum jumps in the whole system [Inset: $V[N_{jump}^{total}(T)]/(\gamma L)$], (b) SFQJ, (c) $V[N_{jump}^{half}(T)]/L^2$ [Inset: $V[N_{jump}^{half}(T)]/(\gamma L^2)$], and (d) $V[N_{jump}^{half}(T)]/L$. We take the average over 20 time intervals for $t \in [200, 1190], [1190, 2180], \dots, [19010, 2 \times 10^4]$.

We give additional numerical results for the variance of quantum jumps in several many-body models including XX ($J_{xy} = 1$, $J_z = 0$), XXZ ($J_{xy} = 1$, $J_z = 0.5, 1.5$), and Ising ($J_{xy} = 0$, $J_z = 1$) models. We note that, as the dissipator in the Liouvillian cannot be expressed in terms of quadratic operators, we shall include the XX model with dephasing as part of manybody models [97–99]. In Fig. S1, we show the variance for the XX chain obtained from the quantum trajectory method. In Fig. S1(a), as expected, the variance for the whole chain does not exhibit the nontrivial behavior and follows the scaling given by $V[N_{jump}^{total}(T)] = \gamma LT/2$. On the other hand, we find that the subsystem fluctuation of quantum jumps (SFQJ) indicates an anomalous scaling different from the volume law as shown in Fig. S1(b). Then, we see in Fig. S1(c) that $V[N_{jump}^{half}(T)]/L^2$ shows a crossing around $\gamma_c \sim 0.7$ as the measurement rate γ is increased. However, the crossing point is rather disturbed compared to the one in Fig. 2(c) in the main text. This comes from the fact that SFQJ in the XX model for weak measurement strength exhibits the stepwise behavior [see the inset in Fig. S1(c)], which may come from the specific integrability of the noninteracting XX spin chain in the unitary limit. Accordingly, we see an oscillation in $V[N_{jump}^{half}(T)]/L$ for weak measurement strength as shown in Fig. S1(d). It is also worth noting that this oscillation comes from V_{anom} that reflects nontrivial subsystem fluctuations. Moreover, we find that measurement-induced crossover of SFQJ emerges in Fig. S1(d), where the scaling is estimated as $V[N_{\text{jump}}^{\text{half}}(T)] \propto L^{2.40 (0.96)}$ for $\gamma = 1 (0.05)$. We note that, for large L, these exponents reduce to the pure ones for $V_{\rm anom}$ coming from integrated autocorrelation functions. However, for the finite size, they include the correction coming from V_{act} .



FIG. S2. Numerical results for the variance of quantum jumps to demonstrate the measurement-induced crossover of SFQJ in the XXZ $(J_z = 0.5)$ model for 832 trajectories and T = 990. Data are plotted against γ for system sizes $L = 8, 10, \dots, 18$ in (a)-(c) and against L for measurement strengths $\gamma = 0.05, 0.1, \dots, 0.95, 1$ from bottom to top in (d). (a) Variance of net quantum jumps in the whole system [Inset: $V[N_{jump}^{total}(T)]/(\gamma L)]$, (b) SFQJ, (c) $V[N_{jump}^{half}(T)]/L^2$ [Inset: $V[N_{jump}^{half}(T)]/(\gamma L^2)$], and (d) $V[N_{jump}^{half}(T)]/L$. We take the average over 20 time intervals for $t \in [200, 1190], [1190, 2180], \dots, [19010, 2 \times 10^4]$.

In Figs. S2 and S3, we depict for the XXZ model ($J_z = 0.5, 1.5$) the same quantities as those in Fig. 2 in the main text and in Fig. S1. We find that the behavior is similar to that of the Heisenberg model shown in Fig. 2 in the main text, and the measurement-induced crossover of SFQJ is detected. We find a crossing of $V[N_{\text{jump}}^{\text{half}}(T)]/L^2$ around $\gamma_c \sim 0.6$ for $J_z = 0.5$ and



FIG. S3. Numerical results for the variance of quantum jumps to demonstrate the measurement-induced crossover of SFQJ in the XXZ $(J_z = 1.5)$ model for 832 trajectories and T = 990. Data are plotted against γ for system sizes $L = 8, 10, \dots, 18$ in (a)-(c) and against L for measurement strengths $\gamma = 0.05, 0.1, \dots, 0.95, 1$ from bottom to top in (d). (a) Variance of net quantum jumps in the whole system [Inset: $V[N_{jump}^{total}(T)]/(\gamma L)$], (b) SFQJ, (c) $V[N_{jump}^{half}(T)]/L^2$ [Inset: $V[N_{jump}^{half}(T)]/(\gamma L^2)$], and (d) $V[N_{jump}^{half}(T)]/L$. We take the average over 20 time intervals for $t \in [200, 1190], [1190, 2180], \dots, [19010, 2 \times 10^4]$.

 $\gamma_c \sim 0.35$ for $J_z = 1.5$ in Figs. S2 and S3, respectively. Also, we estimate the scaling of SFQJ as $V[N_{\text{jump}}^{\text{half}}(T)] \propto L^{2.43 \,(1.08)}$ for $\gamma = 1 \,(0.05)$ for $J_z = 0.5$ and $V[N_{\text{jump}}^{\text{half}}(T)] \propto L^{2.49 \,(1.20)}$ for $\gamma = 1 \,(0.05)$ for $J_z = 1.5$, but the exponents include the finite-size contribution from V_{act} as in the case of the XX model.



FIG. S4. Numerical results for the variance of quantum jumps in the Ising model for 832 trajectories and T = 990, where no measurementinduced crossover is found even for SFQJ. Data are plotted against γ for system sizes $L = 8, 10, \dots, 16$ in (a) and (b) and against L for measurement strengths $\gamma = 0.05, 0.1, \dots, 0.95, 1$ from bottom to top in (c). (a) Variance of net quantum jumps in the whole system [Inset: $V[N_{\text{jump}}^{\text{total}}(T)]/(\gamma L)$] and (b), (c) SFQJ. We take the average over 20 time intervals for $t \in [200, 1190], [1190, 2180], \dots, [19010, 2 \times 10^4]$.

We also give numerical results for the Ising model, which is a trivial yet definite example that does not exhibit the measurement-induced crossover of SFQJ. As shown in Figs. S4(a) and (b), we find nontrivial behavior in the variance neither for the total system nor the subsystem. Though the scaling of SFQJ shown in Fig. S4(b) seems to be different from the one of $V[N_{jump}^{total}(T)]/2 = \gamma LT/4$, they agree with each other in the thermodynamic limit. In fact, the stepwise behavior of SFQJ in Fig. S4(c) reflects the particle-number conservation at a single site in the Ising chain. As the particle number for the half chain is given by L/4 (L/2: even) and L/4 + 1/2 (L/2: odd) reflecting the initial Néel state, they agree in the thermodynamic limit, and we cannot distinguish the scaling of SFQJ from the one of the total system. We remark that V_{anom} becomes zero because the particle number conservation at a single site immediately gives zero autocorrelation in Eq. (6) in the main text.

Numerical results for $V_{\rm anom}$ and $V_{\rm act}$ in other many-body models

In Figs. S5(a), (c), and (e), we show V_{anom} and V_{act} for XX ($J_z = 0$) and XXZ ($J_z = 0.5, 1.5$) models obtained from the quantum trajectory method. We find that the dominant contribution of SFQJ changes from $V_{act} \propto L$ (Poissonian statistics) to $V_{anom} \propto L^{\alpha}$ (super-Poissonian statistics) as indicated by the crossing point. In Figs. S5(b), (d), and (f), from the finite-size scaling analysis with $L = 8, 10, \dots, 18$, the anomalous scaling $V_{anom} \propto L^{\alpha}$ is estimated for $\gamma = 1$ as $\alpha = 2.71, 2.73, 2.68$ for XX and XXZ ($J_z = 0.5, 1.5$) models. Remarkably, these values are close to each other including the exponent $\alpha = 2.69$ for the Heisenberg model and indicate the universality of SFQJ in many-body open quantum systems. We note that, in Fig. S5(b) for weak measurement strength, we find an oscillation of V_{anom}/L with respect to the system size, the behavior of which may stem from the specific integrability of the noninteracting XX chain in the unitary limit. We also remark that, as the average particle number thermalizes and is distributed homogeneously in long times, such an oscillation is not coming from the contribution of the initial Néel state.



FIG. S5. Numerical results for V_{anom} and V_{act} based on the quantum trajectory method for (a), (b) XX, (c), (d) XXZ ($J_z = 0.5$), and (e), (f) XXZ ($J_z = 1.5$) models. Data are shifted by 2000 as L is increased as $L = 8, 10, \dots, 18$ to improve visualization in (a),(c), and (e), and measurement strengths are $\gamma = 0.05, 0.1, \dots, 0.95, 1$ from bottom to top in (b), (d), and (f). (a), (c), (e) The leading contribution of V_{anom} and V_{act} changes at a critical measurement strength. (b), (d), (f) System-size scaling of V_{anom}/L . The parameters and methods are the same as in Figs. S1-S3.

Numerical results for the analysis on the $\gamma\text{-dependence of }V_{\mathrm{anom}}$



FIG. S6. Numerical results for V_{anom} in the half chain to analyze the γ - and L-dependence in (a), (b) Heisenberg, (c), (d) XX, (e), (f) XXZ ($J_z = 0.5$), and (g), (h) XXZ ($J_z = 1.5$) models for 832 trajectories and T = 990. (a), (c), (e), (g) V_{anom}/γ^2 and (b), (d), (f), (h) V_{anom}/L^2 are plotted against γ for system sizes $L = 8, 10, \cdots, 18$. We take the average over 20 time intervals for $t \in [200, 1190], [1190, 2180], \cdots, [19010, 2 \times 10^4]$.

We give additional numerical results to analyze the γ -dependence of V_{anom} in SFQJ for the half chain and also compare the behavior of V_{anom} with SFQJ. In Figs. S6(a), (e), and (g), we find that V_{anom}/γ^2 in the Heisenberg ($J_z = 1$) and XXZ ($J_z = 0.5, 1.5$) model almost monotonically increases as the measurement strength is increased, but V_{anom} is not simply proportional to γ^{β} with an exponent β (> 2) because V_{anom}/γ^2 seems to become a positive constant as we approach the $\gamma \rightarrow 0$ limit. On the other hand, in Fig. S6(c) for the XX model ($J_z = 0$), V_{anom}/γ^2 seems to approach 0 in the $\gamma \rightarrow 0$ limit for odd L/2, while it shows the nonmonotonic behavior for even L/2 for small γ . We note that though the data for L = 18 seem to deviate from 0 in the $\gamma \rightarrow 0$ limit due to numerical limitation, we still see that those for strong γ are proportional to γ . Then, as the nonmonotonic behavior for even L/2 seems to vanish as the system size is increased, we predict that V_{anom} would become an increasing function of γ for large system sizes possibly passing through the origin (V_{anom}, γ) = (0,0). One possible scaling to

explain the behavior in Figs. S6(a), (c), (e), and (g) is $V_{anom} \propto \gamma^2 c(\gamma) \propto \gamma^2 \sqrt{J_z^2 + \gamma^2}$, but the data are rather rough to discuss the precise γ -dependence of V_{anom} , and we need more sophisticated study. We also see in Figs. S6(b), (d), (f), and (h) that the crossing found in $V[N_{jump}^{half}]/L^2$ [see Fig. 2(c) in the main text, Figs. S1(c), S2(c), and S3(c)] does not appear in V_{anom}/L^2 . This means that the system-size scaling of V_{anom} does not change too much with respect to the measurement strength, in contrast with the measurement-induced crossover of SFQJ found in Fig. 2(d) in the main text and in Figs. S1(d), S2(d), and S3(d). It is worth exploring the phenomena that exhibit a measurement-induced crossover or a measurement-induced phase transition in V_{anom} itself, but we leave it for future study.

Further numerical results for the integrated autocorrelation functions and the Liouvillian gap



FIG. S7. Numerical results for the integrated autocorrelation functions and the Liouvillian gap for Heisenberg and XX models. (a) [(b)] Integrated autocorrelation functions for the half chain [Inset: $L/2 \cdot C_{auto}^{single}$] for the Heisenberg (XX) model and (c) V_{anom} and V_{act} for system sizes up to L = 8 obtained with the use of the spectral decomposition (14) in the main text. (d) Exponents *a*, *b*, and *c* of the Liouvillian gap $\Delta \propto 1/L^a$ (blue, exact diagonalization), $C_{auto}^{single} \propto L^b$ (green, quantum trajectory method), and $C_{auto}^{half} \propto L^c$ (red, quantum trajectory method) obtained from the finite-size scaling analysis with L = 8, 10, 12, 14. The leading contribution of V_{anom} and V_{act} changes at a critical measurement strength in (c) consistent with Fig. S5(a). The anomalous scaling of SFQJ in (d) cannot be captured by the Liouvillian gap. Data are shifted by 1000 as L is increased in (c) to improve visualization.

We numerically investigate the integrated autocorrelation functions and the Liouvillian gap in the XX model and also give a supplemental figure for the Heisenberg model. First, we depict in Fig. S7(a) the integrated half-chain autocorrelation function for the Heisenberg model [Inset: $L/2 \cdot C_{auto}^{single}$] calculated from Eq. (14) in the main text. Though it seems that $C_{auto}^{half} \sim L/2 \cdot C_{auto}^{single}$ is satisfied for the Heisenberg model up to L = 8, this is a finite-size effect because $C_{auto}^{half} \gg L/2 \cdot C_{auto}^{single}$ holds for larger system sizes. Moreover, we have numerically checked that off-diagonal terms $C_{auto}^{ij} \equiv \int_0^T d\tau \langle n'_i(\tau)n'_j \rangle_{\infty}$ $(i \neq j)$ take positive- and negative-mixed values. Numerically, up to the system size L = 8, off-diagonal terms C_{auto}^{ij} cancel out with each other so as not to contribute to C_{auto}^{half} . However, such terms give a non-negligible contribution for larger system sizes as shown in Fig. 4(b) in the main text. We also see that integrated autocorrelation functions become increasing functions of γ for system sizes with $L \geq 8$.

In Fig. S7(b), we show the integrated half-chain autocorrelation function for the XX model up to the system size L = 8 obtained by using the spectral decomposition (14) in the main text. In the inset, we depict $L/2 \cdot C_{auto}^{single}$ for comparison. We find that C_{auto}^{half} oscillates as the system size is increased for small γ and $C_{auto}^{half} \neq L/2 \cdot C_{auto}^{single}$ for L = 6. This behavior may reflect the specific integrability of the free XX spin chain in the unitary limit. Moreover, we have numerically checked that off-diagonal terms C_{auto}^{ij} ($i \neq j$) take positive- and negative-mixed values and can contribute to C_{auto}^{half} . It is worth noting that, in Fig. S7(b) for odd L/2, C_{auto}^{half} goes to zero for $\gamma \to 0$, which comes from the negative contribution of off-diagonal elements C_{auto}^{ij} . We also obtain V_{anom} and V_{act} with the use of Eq. (14) in the main text as shown in Fig. S7(c). We see that there exists a crossing of V_{anom} and V_{act} and the critical point γ_c quantitatively agrees well with that in Fig. S5(a) for L = 8.

Next, we compare the exponent of the Liouvillian gap with that of C_{auto}^{single} . We remind that the asymptotic behavior of the local autocorrelation function is dominated by the Liouvillian gap as $|C_{auto}^{single}(\tau)| \sim e^{-\Delta\tau} (\tau \to \infty)$ [81, 82], and if the scaling of the integrated autocorrelation function were solely determined by the Liouvillian gap, we could estimate it as $C_{auto}^{single} \sim \int_0^T e^{-\Delta\tau} \sim 1/\Delta$. In Fig. S7(d), we depict the exponent of the inverse Liouvillian gap (blue curve) obtained from the exact diagonalization of the Liouvillian \mathcal{L} . We also show C_{auto}^{single} (green curve) calculated from the quantum trajectory method. By employing the finite-size scaling analysis with L = 8, 10, 12, 14, we find that the Liouvillian gap is scaled as $\Delta \propto 1/L^{2.18}$ for $\gamma = 1$, which is close to the value $\Delta \propto 1/L^2$ for $\gamma \gg 1/L$ [86, 87]. On the other hand, for C_{auto}^{single} , we obtain $C_{auto}^{single} \propto L^{0.75}$ for $\gamma = 1$. This means that the scaling for the integrated autocorrelation functions (red curve) obtained from the finite-size scaling analysis with L = 8, 10, 12, 14, we find that the scaling [82]. As for the exponent of the integrated half-chain autocorrelation functions (red curve) obtained from the finite-size scaling analysis with L = 8, 10, 12, 14, we find $C_{auto}^{half} \propto L^{2.59}$ for $\gamma = 1$. We have numerically checked that the off-diagonal elements C_{auto}^{ij} with $i \neq j$ as well as diagonal components in C_{auto}^{half} give a finite contribution to the scaling as $C_{auto}^{half} \gg L/2 \cdot C_{auto}^{single}$ for large γ . Thus,

we cannot estimate the anomalous scaling exponent for C_{auto}^{half} given in Eq. (8) in the main text by the Liouvillian gap in the XX model, either.