Minimum spanning blob-trees

Katharina Klost¹, Marc van Kreveld², Daniel Perz³, Günter Rote¹, and Josef Tkadlec⁴

- 1 Freie Universität Berlin, Institut für Informatik {kathklost,rote}@inf.fu-berlin.de
- 2 Department of Information and Computing Sciences, Utrecht University m.j.vankreveld@uu.nl
- 3 University of Perugia daniel.perz@unipg.it
- 4 Charles University josef.tkadlec@iuuk.mff.cuni.cz

— Abstract

We investigate blob-trees, a new way of connecting a set of points, by a mixture of enclosing them by cycles (as in the convex hull) and connecting them by edges (as in a spanning tree). We show that a minimum-cost blob-tree for n points can be computed in $O(n^3)$ time.

1 Introduction

Any introductory course on computational geometry will treat convex hulls and minimum spanning trees (MSTs) for a set P of n points in the plane. These are the least-cost structures that enclose or connect the points. In this paper we investigate a new structure, the *blobtree*, that combines the ideas of enclosing and connecting. A *blob* is a simple polygon, and all points of P enclosed in a blob are considered to be connected to each other. A *tree-edge* is a segment between two points of P, and it connects its two endpoints. A blob-tree is a collection of blobs and tree-edges that collectively connects all of P, see Figure 1. The convex hull and the MST of P are special cases of a blob-tree.

We are looking for the blob-tree that minimizes the total length of all edges drawn: the perimeter of the blobs plus the tree-edges. As we show in Lemma 1 below, the blobs are convex and disjoint. Morever, when contracting the blobs to vertices, the tree-edges form a tree, so the name "blob-tree" is warranted.

While blob-trees appear interesting in their own right, similar structures have been used in information visualization, in particular in KelpFusion [7]. This visualization style is based



Figure 1 The optimal blob-trees on two point sets (computed by our implementation [4]). Blob edges are blue, tree-edges are green. The cost of the solution is the total length of blue and green. MST edges are black (and green, if they are part of the blob-tree).

41st European Workshop on Computational Geometry, Liblice, Czech republic, April 9–11, 2025. This is the full version of a paper that will be presented at EuroCG'25.



Figure 2 Three variants of blob-trees for the same point set, and their costs.

on shortest-path graphs [5], a one-parameter family of connected structures whose extremes are the convex hull and the MST, without regard to a specific optimization criterion More precisely, the shortest path graph for a parameter $t \in [1, \infty)$ uses as the distance between two points p and q the Euclidean distance to the power t. The shortest path graph is the union of all shortest paths between pairs of points; for t = 1 this is the complete graph and for $t \to \infty$ this is the Euclidean MST. For KelpFusion, this graph is then converted into a connected set of edges (not necessarily convex blobs) by filling in faces. Other related work considers placing shortest fences for subsets of points [1, 2, 3].

As seen in Figures 1 and 2, an optimal solution might involve a tree-edge that crosses a boundary of a blob. If we disallow such crossings (version (ii)), the optimal solution could get longer and might include non-convex blobs. If we further required the blobs to be convex (version (iii)), the optimal solution might become even longer, see Figure 2. A different version ignores the parts of tree-edges that lie inside a blob when computing the length of a solution. This is less natural. The solution might choose edges that run deep into the blob, just to get a shorter distance to the blob boundary.

In this paper, we concentrate on the original problem. It turns out that the optimal solution can be obtained from the MST by replacing several of its subtrees by blobs (Lemma 2); the other two variants (ii) and (iii) do not have this property. As our main result we show that the optimal blob-tree can be computed in $O(n^3)$ time, and we implement the algorithm in PYTHON [4]. Efficient computation of optimal blob-trees in variants (ii) or (iii) is still open.

Our algorithm is based on dynamic programming and builds upon ideas from [6]. That paper shows how to compute a single optimal convex polygon (or blob) in a point set that has certain properties, using dynamic programming. For example, it can compute the smallest area or perimeter convex polygon with k points on the boundary, or the smallest area convex polygon containing at least k points inside or on the boundary in $O(kn^3)$ time. We show how to deal with multiple blobs and with the minimum spanning connecting structure to compute an overall minimum solution.

All proofs that are not given in the main part are in the appendix.

2 Preliminaries

We assume that the point set P is in general position. In particular, we assume that the pairwise distances between points are unique and no three points are on a line. This implies that the MST of the point set is unique. We also assume that no two points have the same x- or y-coordinate.



Figure 3 A minimum spanning tree, a potential subset of non-tree edges (fat blue), and the resulting blob-tree.

▶ Lemma 1. In any optimal solution: all blobs are convex polygons with vertices at P; any two blobs are disjoint; and when contracting the blobs to vertices, the tree-edges form a tree.

Proof. The claims follows from the triangle inequality. First, we can replace any blob that does not have the claimed property with the convex hull of the points of P contained in it, creating a blob-tree with a strictly smaller total cost. Similarly, two intersecting blobs B_1 and B_2 can be replaced by the convex hull of the union $B_1 \cup B_2$.

Contraction of blobs creates an abstract graph G in which every blob and every singleton point that is not in a blob is represented by a vertex. For every tree-edge, G contains an edge between the corresponding vertices. Since the blob-tree is connected and spans all points, G is connected. If G contained a cycle, then any tree-edge corresponding to an edge from such a cycle could be removed from the blob-tree. So the graph G is a tree.

Throughout the paper, T will denote the minimum spanning tree (MST) of P. We select the lowest point r as the root and direct all edges of T towards r. For a node u, we denote its parent in T by u' and the directed edge from u to its parent by uu'.

3 Structural insights

The following key lemma with its immediate corollary shows that the optimal solution can be obtained from the MST by replacing some of its subtrees by the respective blobs.

▶ Lemma 2. Let T be the MST. Then, in any optimal blob-tree S, every blob is a convex hull of some subtree of T and every tree-edge is an edge of T.

Lemmas 1 and 2 immediately yield the following corollary.

 \blacktriangleright Corollary 3. Let B be a blob in any optimal blob-tree S. Let T be the MST. Then:

- (a) A path in T that leaves B never comes back to B.
- (b) The subgraph of T induced by the points in B is connected.
- (c) No MST edge with both endpoints outside B intersects B.

Lemma 2 and Corollary 3 are the foundation of a straightforward exponential-time algorithm for computing the optimum blob-tree, see Figure 3. After computing the MST T, we guess which of the n - 1 MST edges are tree-edges (the black edges in the middle of Figure 3) and which edges are inside blobs (the blue edges). Removing the black edges cuts the tree into blue components. We take the convex hull of each blue component as a blob



Figure 4 A rooted, directed blob tree, two types of exits, and a bottom-vertex triangulation of a blob with a dynamic programming order through it towards the root.

and add the black edges, as shown in the right of Figure 3. In total, there are 2^{n-1} candidate solutions, and we take the one that minimizes the total cost.

The structure of optimal blob-trees can be used to develop a polynomial-time algorithm by dynamic programming. We extend the structure in two simple ways.

Firstly, we use the direction of all edges of T towards the root r. If we were to contract each blob to a single point, then we would have a normal rooted tree. Consequently, every blob that does not contain the root has one unique exit edge and every blob has zero or more entry edges. The exit from a blob towards the root can be via an edge from one of its hull vertices or via an edge from an interior blob point that crosses a hull edge. Similarly, blob entrances come in two types, see Figure 4.

Secondly, we consider a bottom-vertex triangulation of each blob, where the lowest point is connected to all other vertices on the boundary of the blob. We call the diagonals of the triangulation *chords* and the exterior edges *walls*. The chords and triangles serve to have simple (constant size) elements to be used to define substructures in the dynamic program. Optimal solutions to smaller subproblems are "transported" through a blob from any entrance to the unique exit, using the chords in between, to build solutions to larger subproblems.

There will be three types of subproblems: (i) edge subproblems, for each MST edge that is used as a blob-tree edge, (ii) chord subproblems, and (iii) wall subproblems. There are n-1 of the first type and $O(n^2)$ of the latter two types. Walls and chords do not have an implied orientation (unlike the edges of T). Each segment must be considered as a potential wall or chord separately in both orientations.

For uniformity, we draw chords from the bottom vertex of a blob also to its neighbors on the blob boundary, cutting off two degenerate *digons* from the blob, see Figure 5. Thus, a blob with k boundary points has k - 1 chords and k walls, and it is cut into k - 2 triangles, each bounded by two chords and one wall, and two digons, bounded by one chord and one wall.

Every chord has a *forward side*, towards the tree root, and a *backward side*; both are defined later in detail. Our algorithm proceeds from the leaves towards the root, as indicated by the red arrows in Figure 4. Thus, the subproblem associated to a chord consists of all points that are on the backward side, including the points that are in other blobs hanging off the current blob in the blob tree. It turns out that the points that are involved in this subproblem can be uniquely identified with the help of the MST, without knowing the



Figure 5 A blob is divided into triangles and digons.

optimum blob tree.

Let a, b be a pair of points. Let X_{ab} be the set of edges of T that cross the line segment ab, and let X_{ab}^+ denote the edges in X_{ab} plus the edges of T that are incident to a or b.

Sidedness. Assume that a is below b and consider the boundary curve that is obtained from the union of the vertical ray below a and the ray from a through b, see Figure 6a. This curve subdivides the plane into a left and a right side. We call a point v that is a vertex of an edge in X_{ab} a left or right endpoint, depending on if it lies left or right of the boundary curve. The endpoints of edges in $X_{ab}^+ \setminus X_{ab}$ that are not a or b are classified analogous.

Lemma 4. Every edge in X_{ab} has exactly one left and one right endpoint.

Valid chords. Removing X_{ab}^+ from T gives a forest. We call the segment ab a valid chord if no component in $T \setminus X_{ab}^+$ contains both a left and a right endpoint. For a valid chord, the components of $T \setminus X_{ab}^+$ can be partitioned into *left* and *right components*, depending on the characterization of the vertices from X_{ab}^+ contained in the component. Note that this partitions $P \setminus \{a, b\}$.

The side assigned to the root r is the *forward* side of the chord and the other side the *backward* side. (If the lower point a of the chord is r, we arbitrarily declare the right side as the forward side. This convention simplifies the treatment of the "root blob".) If the forward side is the right side, then ab is a *right-facing* chord, otherwise it is a *left-facing* chord.

▶ Lemma 5. All chords in an optimal solution are valid.

Walls. Walls are the edges on the boundary of a blob in a counterclockwise traversal. Let b, c be a pair of points. Then $uu' \in X_{bc}$ is an *entry edge* for \overrightarrow{bc} if it crosses \overrightarrow{bc} from right to left (from outside the blob to inside) and an *exit edge* otherwise.



Figure 6 (a) Labeling endpoints of MST-edges X_{ab}^+ crossing ab or incident to a or b as left (L) and right (R). The neighbors of a and b are classified as right (R) with respect to ab if the edges emanate in the pink region, otherwise they are left (L). (The shown edges cannot all be MST edges simultaneously.) (b) Proof of Lemma 4.



Figure 7 Six entry edges and one exit edge for the triangle *abc*. The four edges that cross *bc* are associated to the wall *bc*. sd



Figure 8 An example with one exit triangle, several LR- and RL-triangles, and two entry digons.

Triangles and Digons. Among the non-root triangles in to which a blob is decomposed from the lowest vertex, there is a unique wall, where the edge towards the root exits. If this wall is part of a triangle, it is an *exit triangle* the other triangles are LR-triangles (left-to-right), RL-triangles (right-to-left), depending on the direction in which the root lies, see Figure 8. Points adjacent to vertices of the blob are assigned to the triangles or digons according to the extension of the triangles by the rays through *a* and the other two vertices.

For every triangle abc with lowest point a, we can, by analyzing the tree edges that cross the sides or are incident to the vertices a, b, v, classify it as a potential LR-triangle, RL-triangle or exit triangle. A similar classification is obtained for digons. The details are given in Appendix D.

The following lemma shows that this concept of valid triangles and digons agrees with the intended meaning:

▶ Lemma 6. Let B be a blob in an optimal solution. Then,

- (a) There is exactly one exit triangle or digon.
- (b) Every other triangle and digon in the decomposition of the blob from the lowest point is a valid LR-triangle, RL-triangle or a valid entry digon.

4 The algorithm

The algorithm does not actually check that the blobs that we form are convex. The only property that we implicitly enforce is that each blob is star-shaped around the lowest vertex a. Thus, the algorithm considers also nonconvex blobs for potential solutions. However, we know from Lemma 1 that such solutions cannot be optimal. As mentioned, we consider three type of subproblems.

Edge problems. For each MST edge uu', we define a problem edge[u], for the optimum solution in which uu' is used as a tree-edge. In addition, for the root r, we have edge[r] for

25:8 Minimum spanning blob-trees

the overall problem. We define V_u as the set of points in the subtree of T rooted at u. We define the *size* of the problem as the cardinality of V_u .

Chord problems. For each valid chord ab, we define a problem chord[a, b] for the optimum solution on the backward side of ab, supposing that ab occurs as a chord of a blob in the solution. We define V_{ab} as the set of all points that are in a backward component of $T \setminus X_{ab}^+$. We define the *size* of the problem chord[a, b] as the cardinality of V_{ab} . For clarity, we will sometimes write V_{ab} as \overleftarrow{V}_{ab} for a right-facing chord and as \overrightarrow{V}_{ab} for a left-facing chord.

Wall problems. We denote by W_{abc} the set of endpoint of entry edges for abc that lie outside of the blob. We will frequently need the values $\sum_{u \in W_{abc}} edge[u]$. We split this into

$$\sum_{u \in W_{abc}} \mathsf{edge}[u] = \sum_{u \in W_{bc}} \mathsf{edge}[u] + \sum_{u \in W_{abc}, \ uu' \text{ entry edge into } b \text{ or } c} \mathsf{edge}[u]$$

To speed up the evaluation of this term, the first sum is stored as the solution of an auxiliary subproblem, wall[bc]; the second sum can be computed in constant time, because there are only a constant number of entry edges incident to b and c, because the degree of a Euclidean MST is bounded by 6.

For each pair of points b, c, we denote by W_{bc} the set of points w where $ww' \in X_{bc}$ is an entry edge for bc. The size of this problem is defined as the sum of the sizes of the problems edge[u]. If bc is a wall in an optimal solution, the subtrees rooted at the nodes $w \in W_{bc}$ are disjoint. However, we don't check this condition. Hence the size can be bigger than n.

Relation between subproblems. In Appendix G, we describe the relations between the sets of points that define the subproblems (Lemmas 9 and 10). These lemmas ensure that the solution of every subproblem depends only on problems of smaller size.

Preprocessing. In a preprocessing phase, we determine the size of each subproblem, in order to know the order in which we have to solve the subproblems. In Appendix H, we give details about the extra information that is gathered in the preprocessing step (Lemma 11).

Chord problems. For each valid chord *ab*, we determine its sidedness. If it is right-facing, we consider all possibilities for an LR-triangle *abc*, as well as the possibility that *ab* is a left entering digon. Left-facing chords are analogous. More details are in Appendix I.

Edge problems. For each MST edge uu', we have two possibilities. If u is not in a blob, all incoming MST edges of u must be tree-edges, and we can accumulate the values of the corresponding subproblems. If u is in a blob, we consider all potential exit triangles abc for which the uu' is the exiting edge crossing the wall bc, as well as the analogous possibility of an exit digon. Details are given in Appendix I.

▶ **Theorem 7.** The dynamic program above correctly solves the minimum blob-tree problem in $O(n^3)$ time.

Proof. For each size, we first solve the edge problems, then the wall problems, and then the chord problems. Lemmas 9 and 10 in Appendix G show that every subproblem solution that is needed is already computed, and correctness follows from these lemmas.

The straightforward running time analysis is given in Appendix J.

Acknowledgments. We thank Bettina Speckmann for co-proposing this problem, and Philipp Kindermann for hosting GG Week 2024 in Trier, where this research started.

	References	
--	------------	--

- 1 Mikkel Abrahamsen, Panos Giannopoulos, Maarten Löffler, and Günter Rote. Geometric multicut: shortest fences for separating groups of objects in the plane. *Discrete Comput. Geom.*, 64:575–607, 2020. doi:10.1007/s00454-020-00232-w.
- 2 Esther M. Arkin, Samir Khuller, and Joseph S. B. Mitchell. Geometric knapsack problems. *Algorithmica*, 10(5):399–427, 1993. doi:10.1007/BF01769706.
- 3 Therese Biedl, Éric Colin de Verdière, Fabrizio Frati, Anna Lubiw, and Günter Rote. Finding a shortest curve that separates few objects from many. In Oswin Aichholzer and Haitao Wang, editors, 41st International Symposium on Computational Geometry (SoCG 2025), volume 332 of Leibniz International Proceedings in Informatics (LIPIcs). Schloss Dagstuhl-Leibniz-Zentrum für Informatik, 2025. To appear.
- 4 Blobtrees. Github repository. URL: https://github.com/guenterrote/blobtrees.
- 5 Mark de Berg, Wouter Meulemans, and Bettina Speckmann. Delineating imprecise regions via shortest-path graphs. In *Proceedings of the 19th ACM SIGSPATIAL international* conference on advances in geographic information systems, pages 271–280, 2011.
- **6** David Eppstein, Mark Overmars, Günter Rote, and Gerhard Woeginger. Finding minimum area *k*-gons. *Discrete Comp. Geom.*, 7:45–58, 1992. doi:10.1007/BF02187823.
- 7 Wouter Meulemans, Nathalie Henry Riche, Bettina Speckmann, Basak Alper, and Tim Dwyer. Kelpfusion: A hybrid set visualization technique. *IEEE Transactions on Visualization and Computer Graphics*, 19(11):1846–1858, 2013.
- 8 Michael Ian Shamos and Dan Hoey. Closest-point problems. In 16th Annual Symposium on Foundations of Computer Science (FoCS 1975), pages 151–162, 1975. doi:10.1109/ SFCS.1975.8.

A Proof of Lemma 2

Proof. First, consider an MST-edge e = u'u'' that is not a tree-edge or inside a blob in a solution S. We show how to construct a better solution \hat{S} , and therefore S cannot be optimal. The removal of e splits T into two subtrees T' and T'' with $u' \in T'$ and $u'' \in T''$. There has to be a connection between u' and u'' in S, consisting of an alternating sequence of paths and blobs, see Figure 9 for an illustration. More precisely, there are $k \geq 1$ paths $u' = v_0 \rightarrow w_0, v_1 \rightarrow w_1, \ldots, v_k \rightarrow w_k = u''$ consisting of tree-edges, such that w_i and v_{i+1} are in a common blob. It can happen that the first or last path is trivial, i.e., $u' = w_0$ or $v_k = u''$, but there must be at least one non-trivial path, because otherwise u' and u'' are in the same blob, contrary to our assumption.

At some point in this sequence, there must be a switch from T' to T''. If this switch occurs at an edge \bar{e} of one of the paths, we can exchange \bar{e} by e and reduce the cost. Suppose that the switch occurs in one of the blobs, say between $w_i \in T'$ and $v_{i+1} \in T''$. Then we know that the distance between w_i and v_{i+1} is larger than the length of e. In this case, we obtain the shorter solution \hat{S} by forming a single blob from the convex hull of all paths and blobs on the way between u' and u''. To see that this is shorter than S, we first form the outer contour K of all paths and blobs together with the edge e. We have to pay the length of e, but we save the parts of the blob perimeter that lie inside the contour. In particular, consider the last edge xw_i of the path $v_i \to w_i$ and the first edge $v_{i+1}y$ of the path $v_{i+1} \to w_{i+1}$. Let \bar{x} and \bar{y} be the intersections of the edges xw_i and $v_{i+1}y'$ with the boundary of B. (It is possible that $\bar{x} = w_i$ or $\bar{y} = v_{i+1}$.) When comparing the blob with



Figure 9 The situation for Lemma 2. Adding e and then taking the orange region (that is the convex hull of the union of the blue blobs and green edges) as a single blob gives a shorter solution, as the red dotted path between x and y is now inside the blob and does not count towards the length of the solution.

the outer contour K, we save at least the parts $w_i \bar{x}$ and $\bar{y} w_{i+1}$ of the edges plus a part of the boundary of B from \bar{x} to \bar{y} , forming a curve connecting w_i with v_{i+1} (highlighted in red). This curve is at least as long as $||w_i v_{i+1}||$, which is longer than e. Taking the convex hull can only make the curve even shorter. Thus, S cannot be optimal. See Figure 9 for an illustration.

So far we showed that in an optimal solution, any MST edge is either a tree-edge or it is contained in a blob. It remains to show that any blob in an optimal solution is a convex hull of a subtree of the MST (as opposed to being a convex hull of a set of points that is disconnected in the MST).

For contradiction, suppose there exist two points u, v that belong to the same blob B and the path from u to v in the MST does not lie inside B. Let $x\bar{x}$ be the MST edge on this path that leaves B for the first time. If we remove this edge from S, \bar{x} is still connected to x: First, there is the MST path from \bar{x} to v, and every edge on this path is either a tree-edge, or its endpoints are in the same blob. Secondly, v is in the same blob as x Therefore the solution S was not optimal.

B Proof of Lemma 4

Proof. A line segment xy could intersect the boundary separating the left and right area twice. However, as the angle between ab and the vertical downward ray from a is obtuse, the lens defined by the disks with radius ||xy|| centered at x and y always contains a, implying that xy is not an edge of the MST and thus not contained in X_{ab} . See Figure 6b.

C Proof of Lemma 5

Proof. Suppose ab is a chord of some blob B in an optimal solution. First observe that, by Corollary 3 (c) for any edge $uu' \in X_{ab}$, at least one of u and u' must lie in B.



Figure 10 A few endpoints of edges X_{ab} for a chord *ab* of a blob *B*.

1) If an endpoint u of an edge $uu' \in X_{ab}^+$ lies outside B (such as u_1, u_3 , or u_7 in Figure 10), it cannot be connected to any other endpoint v of an edge $v\bar{v} \in X_{ab}^+$, no matter how the two points are labeled. The reason is that the partner u' of u is in B, and either v or its partner \bar{v} is in B. So we would have an MST path that starts in B (at u'), goes out of B (at u) and comes back into B (ending in v or \bar{v}), contradicting Corollary 3 (a).

2) If a left endpoint u and a right endpoint v of two edges in X_{ab}^+ lie both in B, (such as u_2 and u_5 in Figure 10), they cannot be connected in $T \setminus X_{ab}^+$. The reason is that the MST must connect u and v within B, by Corollary 3 (b), and this is impossible without cutting the chord ab or going through a or b.

D Classification of triangles and digons

Let abc be a counterclockwise triangle with lowest point a and valid chords ab, ac. Note that a lies on the left side of \overline{bc} . Let O_{abc} be the region defined by the intersection of the right part of \overrightarrow{bc} with the wedge defined by the rays ab and ac, see the orange area in Figure 7. The **entry edges** for abc are the entry edges for bc plus the MST edges uu' with $u' \in \{b, c\}$ and $u \in O_{abc}$. The **exit edges** for abc are the exit edges for bc plus the MST edges uu' with $u \in \{b, c\}$ and $u \in O_{abc}$.

We call abc an LR-triangle (for "left-to-right" triangle), if ab and ac are right-facing and there are no exit edges for abc. Analogously, abc is an RL-triangle if ab and ac are left-facing chords and there are no exits edges for abc. The triangle abc is an **exit-triangle**, if ab is left facing, ac is right-facing, and in addition, there is exactly one exit edge for abc. LR-triangles, RL-triangles, and exit-triangles are **valid triangles**. For a valid triangle, let $X_{abc}^+ = X_{ab}^+ \cup X_{ac}^+ \cup X_{bc}^+$.

Digons are more tricky to define, because they must simultaneously fulfill the conditions of valid chords and entry or exit walls. A valid chord ab with $a \neq r$ is a **valid entry digon** if the directed edges $uu' \in X_{ab}^+$ satisfy the following condition: For every edge uu' in X_{ab}^+ , u = a or u is on the backward side, and u' = b or u' is on the forward side. It is a **left entry digon** if ab is right-facing (see Figure 11) and a **right entry digon** if it is left-facing.

If there is exactly one edge $uu' \in X_{ab}^+$ for which the above condition is fulfilled, then ab is a *valid exit digon*. It is a *right exit digon* if it is right-facing and a *left exit digon*



Figure 11 A left entry digon (a) and a left exit digon (b), together with the outline of a blob to which these digons might belong. (In (b), the edge from *a* to the point marked R shows that this digon can never be part of a valid blob. However, this is not tested when considering the segment *ab*: This edge will preclude the existence of a matching right entry digon.)

if it is left-facing (see Figure 11). (If $|X_{ab}^+| = 1$ then it is simultaneously an entry digon and an exit digon, but of opposite sides.)

The case a = r is special: By definition, all chords rb are right-facing. Any chord rb for which all edges in X_{rb} cross from the right to the left side is a *valid root digon*. In order to handle root digons consistent with other digons, we assume that a root digon is a right exit digon with a virtual exit edge rr' with ||rr'|| = 0.

E Proof of Lemma 6

Proof. (a) We look at the exit edges from B, i.e., the MST edges uu' with u' not in B that cross a wall of B or that emanate from a vertex u of B into the direction outside B. We claim that B has at most one exit edge, and it has one exit edge if and only if B does not contain the root.

Suppose B has two exit edges uu' and vv'. We know from Corollary 3 (b) that u and v are connected by a tree path within B, i.e., without going through u' and v'. But this is inconsistent with the orientation of uu' and vv'. The same contradiction is obtained if B contains the root and has an exit edge. Of course, if B does not contain the root, there must be at least one exit edge. Thus, the claim has been established.

If B contains the root, it is the lowest point a of the blob, by our assumption the right digon is a right exit digon and none of the other digon and triangles are exit digons or triangles.

Otherwise, there is a unique exit edge. If this edge cuts a wall, this defines a unique exit triangle or digon. If the exit edge emanates from a vertex of B, the partition of the outgoing directions into disjoint sectors (Figure 5) ensures that our convention of assigning the exit tree edge to a triangle (Figure 7) or digon identifies exactly one triangle or digon as the exit triangle or digon.

(b) We have already shown that all chords are valid (Lemma 5). We still need to show that



Figure 12 The path π_1 connectes a right endpoint in X_{ab}^+ with a left endpoint in X_{ac}^+ .

all chords to the right of the exit triangle or digon are left-facing, and all chords to the left of the exit triangle or digon are right-facing. If the root is in B, then, by convention, all chords are right-facing, and this is the desired result.

Consider now the case that the root is not in B, and let uu' be the exit edge. The point u lies inside B. If u = a, one can check directly that all chords are oriented correctly, using the edge $uu' \in X_{ab}$.

Consider the case $u \neq a$. For each chord ab, it is sufficient for find *one* edge of X_{ab} whose endpoint is labeled correctly as the forward side. All chords ab that are crossed by uu' (or touched by uu', in case u = b) are oriented correctly, towards u', because the path from u' to the root does not enter B and therefore does not use edges of X_{ab} .

All remaining chords ab are directed towards u, which is seen as follows. Take the tree path π from b to the root. By Corollary 3 (b), the points in B are connected by tree edges, and hence the path must leave B via the edge uu'. Thus, the last vertex of X_{ab} that is visited by π must be on the side facing u. For the chords ab that are not intersected or touched by uu', the direction towards u agrees with the the direction towards the exit triangle, and hence the proof is completed.

F Interaction between the chords of a valid triangle

The following lemma gives additional structural insight on how the sides of valid triangles interact.

Lemma 8. Let abc be an LR- or RL-triangle and let uu' be an entry edge of abc. Then the following holds:

- (a) Every point p in a left component of ac lies in a left component of ab. Every point p in a right component of ab lies in a right component of ac.
- (b) No point in the left component of ac or in the right component of ab lies in the same connected component of $T \setminus X_{abc}^+$ as u.
- **Proof.** (a) We show the statement for the case of LR-triangles and left components. The remaining three cases are symmetric. Let abc be a LR-triangle. Assume that there is a point p that is in a left component with regard to ac but in a right component with regard to ab. Let π be the path from p to the root.

25:14 Minimum spanning blob-trees

Consider the subpaths $\pi \setminus (X_{ab}^+ \cup X_{ac}^+)$ of π and a subset $\Pi = \pi_1, \ldots \pi_k$ of these paths. Π contains the first path of $\pi \setminus (X_{ab}^+ \cup X_{ac}^+)$ if it ends in an endpoint of X_{ac}^+ and all paths that connect an end point in X_{ab}^+ to an end point in X_{ac}^+ . Note that $\Pi \neq \emptyset$ as otherwise π would not use an edge in X_{ac}^+ . As *ac* is a right-facing chord, this would imply that *p* is in a right component of *ac*.

First note that if $xx' \in X_{ab} \cap X_{ac}$, then x is a left end point of an edge in X_{ab} if and only if it is a left end point of an edge in X_{ac} . Additionally, let e be an edge adjacent to a and \overline{a} the other endpoint of this edge. Then, if \overline{a} is a right endpoint with regard to ab it is also a right endpoint with regard to ac. Symmetrically, if \overline{a} is a left endpoint with regard to ac it is also a left endpoint with regard to ab.

We claim, that all paths in Π contain vertices that are in a left component of ac and in a right component of ab. Furthermore, each path in Π that does not start with p starts and ends with a vertex in $X_{ab}^+ \oplus X_{ac}^+$. We show this in an inductive fashion, starting with π_1 . If π_1 starts with p the claim follows by the definition of p. In the other case let $\pi_1 = (v_1, \ldots, v_\ell)$. Then v_1 is an end point of an edge in X_{ab}^+ and v_ℓ is an endpoint of an edge in X_{ac}^+ . As the prefix of π that ends with v_ℓ did not use any edges in X_{ac}^+ , v_ℓ cannot be a right end point of an edge in X_{ac}^+ since p is in the left component of ac. As there are no exit edges in abc, the point v_1 cannot be a left end point of X_{ab}^+ . This shows that all points on π_1 are in a left component of ac and in a right component of ab. By the observation above, they cannot be in $X_{ab}^+ \cap X_{ac}^+$.s This argument can be extended inductively to show the claim for all $\pi_i \in \Pi$.

The claim implies that π_k ends with a left endpoint of an edge in $X_{ac}^+ \setminus X_{ab}^+$ or a right end point of $X_{ab}^+ \setminus X_{ac}^+$. If π_k ends with a right end point x of an edge in X_{ab}^+ , then all points on π that are successors of x, in particular r are in a left component of ac. This is a contradiction to the assumption that ac is a right facing chord. In the other case, π_k ends with a left end point of X_{ac}^+ and the path does not use any edges of X_{ab}^+ anymore. Consider the suffix of π starting at the last point of π_k . All points on this suffix are in a right component of ab. Let yy' be the last edge of this suffix that uses an edge in X_{abc}^+ . This edge has to be a left-to-right edge of X_{ac}^+ , as ac is a right-facing chord. Thus, y'lies on the right side of ac. There are three possible regions on the right side of ac for y', all leading to a contradiction. If y' lies on the right side of ab this would make y a point in a left component of ab, a contradiction. If it lies in O_{abc} this would make yy' an exit edge. Finally, if it lies in the triangle, the suffix of π connecting y' to r lies completely in abc. This is only possible if a is the root. Consider the last edge $\bar{\tau}r$ on π . This edge is in X_{ab}^+ , a contradiction to the assumption, that the suffix does not use any such edges.

(b) W.l.o.g assume that p is in the right component of ab. We show that u is in the left component of ab.

Let π be the path connecting u to r in T. As abc is not an exit triangle, there exists an edge vv' such that v' is the first point of π after u that is outside of the triangle abc. Note that vv' = uu' is possible. This also means that vv' is either in X_{ab}^+ or in X_{ac}^+ .

We consider two cases based on the position of v'. If $vv' \in X_{ab}^+$ then v is in the left component of ab by definition.

If $vv' \in X_{ac}^+$ then v' is a left endpoint and thus in a left component of ac. By part (a) of this lemma, v' is then also in a left component of ab. However the subpath of π between u and v' does not go through X_{ab}^+ by assumption. Hence, v is also in the left component of ab.

In both cases we get that v is in the left component of ab. As the path between u and v does not use any edges of X_{abc}^+ , this implies that u also in in a left component of ab.



Figure 13 (left) if π_k ends with a right endpoint in X_{ab}^+ then *ac* is not right-facing. (right) the last edge in X_{ac}^+ ends in one of the three regions, all lead to a contradiction.



Figure 14 Illustration of the proof of Lemma 8 (ii).

As p is in a right component of ab, they cannot be in the same connected component of $T \setminus X_{abc}^+$. The other cases follow similarly.

G Relation between subproblems

Here we describe the relations between the sets of points that define the subproblems. This is important to show that the subproblems can be computed in order of increasing sizes of V_{ab} and V_u . Δ_{abc} denotes the set of points in the interior of the triangle *abc*, and W_{abc} is the set of points *w* such that ww' is an entry edges for *abc*.

▶ Lemma 9. (a) If abc is an LR-triangle, then

$$V_{ab} = \{c\} \cup V_{ac} \cup \Delta_{abc} \cup \bigcup_{w \in W_{abc}} V_w$$

Moreover, V_{ab} is a strict superset of V_{ac} .

(b) If abc is an RL-triangle, then

$$V_{ac} = \{b\} \cup \vec{V}_{ab} \cup \Delta_{abc} \cup \bigcup_{w \in W_{abc}} V_w$$

Moreover, V_{ac} is a strict superset of V_{ab} . (c) If abc is an exit triangle with exit edge uu', then

$$V_u = \{a, b, c\} \cup V_{ab} \cup V_{ac} \cup \Delta_{abc} \cup \bigcup_{w \in W_{abc}} V_w$$

Moreover, V_u is a strict superset of \overleftarrow{V}_{ab} , \overrightarrow{V}_{ac} and $\bigcup_{w \in W_{abc}} V_w$.

25:16 Minimum spanning blob-trees

Proof. Note that no path in T joins a point inside *abc* to a point outside *abc* without passing through X_{abc}^+ .

- (a) \subseteq : First we show that every point $p \in V_{ab}$ is at least one of the sets on the right hand side. If p = c we are done. Otherwise, let r_p be the first vertex of the path π from p to r that is an endpoint in X^+_{abc} . Note that r_p exists, as p is on the backward side of ab and thus π has to use at least one edge of X^+_{ab} . If $r_p \in \Delta_{abc}$, then p also lies in Δ_{abc} as the path can only leave the triangle by using an edge in X^+_{abc} . In the other case, if r_p is the left endpoint of an edge in X^+_{ac} , then $p \in V_{ac}$ as it lies in a backward component of this edge. Similarly, if $r_p \in W_{abc}$ the statement follows directly.
 - \supseteq : Now we show that all sets on the right are subsets of V_{ab} . Let $p \in V_{ac}$. Then by definition, p is in a left component of the chord ac and in a right component of ab. By Lemma 8 we get $p \in V_{ab}$.

Similarly, if $p \in V_w$ for some $w \in W_{abc}$ but not in V_{ab} consider the path π from p to r. Let ww' be the entry edge defined by w. Let C be the component of $T \setminus X_{abc}^+$ that contains w and $\pi' = (x', \ldots, w)$ the subpath of π in C. Then x' lies outside of the triangle abc and either x' = p or x' is an endpoint for an edge in $X_{ab}^+ \cup X_{ac}^+$. By Lemma 8, x' cannot be in a left component of ac or a right component of ab. Thus x' is in a right component of ac and a left component of ab. If x' = p, then we have $p \in V_{ab}$. If $x' \neq p$, note that the only endpoints that define such components are in the wedge defined by the rays through ab and ac. As x' does not lie inside the triangle, xx' crosses bc from the inside to the outside, making it an exit edge. This would mean that abc is an exit triangle but abc is an LR-triangle. Hence, we always have x' = p and therefor $p \in V_{ab}$.

Finally, let $p \in \Delta_{abc}$. Assume to the contrary that $p \notin V_{ab}$. Then the first edge yy' in X_{ab}^+ on the path π from p to the root crosses ab from right to left. Thus π has to leave the triangle before yy'. As there are no exit edges, the edge xx' leaving the triangle is in X_{ac}^+ , making x' a point right of ab but left of ac, a contradiction to Lemma 8. Hence, we get $p \in V_{ab}$.

To see that $c \in V_{ab}$, consider the edge cc' on the path from c to r. If $cc' \in X_{ab}$, then $c \in V_{ab}$ by definition. In the other case c is in the same connected component of $T \setminus X_{ab}^+$ as c'. As abc is not an exit triangle, c' either is left of ac, implying $c' \in V_{abc}$ or in Δ_{abc} implying $c' \in \Delta_{abc}$. In both cases we showed above, that $c' \in V_{ab}$ and thus also $c \in V_{ab}$.

The strict inclusion follows as $c \notin V_{ac}$ by definition.

- (b) Analogous.
- (c) \subseteq : First, we show that every point in V_u is contained in at least one of the sets on the right side of the equation. Let $p \in V_u$. If $p \in \{a, b, c\}$ we are done. In the other case, again let r_p be the first endpoint of an edge in X_{abc}^+ on the path from p to the root. As uu' is the exit edge of the triangle and $p \in V_u$, r_p exists. If $r_p \in \Delta_{abc}$, then $p \in \Delta_{abc}$ as the part of π connecting p to r_p does not leave the triangle. In the other case, r_p is a right endpoint of an edge in X_{ab}^+ , a left endpoint of an edge in X_{ac}^+ or an outward point of an edge in W_{abc}^+ . By the same arguments as in (a) it follows, that $p \in V_{ab}, p \in V_{ac}$ or $p \in V_{r_p}$ respectively.
 - ⊇: Now we show that every point in one of the sets on the right is also contained in V_u . Let $p \in V_{ab} \cup V_{ac}$ and let xx' be the last edge on the path π from p to r that is in $X_{ab}^+ \cup X_{ac}^+$. Then x' is a right endpoint in X_{ac}^+ or a left endpoint in X_{ab}^+ . Thus $x \notin X_{ab}^+ \cap X_{ac}^+$. If xx' is also in X_{bc}^+ , then xx' = uu' as uu' is the only exit edge of abc.

K. Klost, M. van Kreveld, D. Perz, G. Rote, and J. Tkadlec

In the other case x' is in the interior of abc and π has to exit the triangle through bc. Again, uu' is the only exit edge and thus π goes through u.

If $p \in \Delta_{abc}$, the same argument holds, as either the path exits the triangle directly through the exit edge, or it goes through an edge in $X_{ab}^+ \cup X_{ac}^+$ to point x in $V_{ab} \cup V_{ac}$. Then p and x are in the same subtree of T and thus $p \in V_u$.

Now assume $p \in \{a, b, c\}$ and let pp' be the first edge on the path from p to r. Then $pp' \in X_{abc}^+$. If $pp' \in X_{ab}^+ \cup X_{ac}^+$ or $p' \in \Delta_{abc}$, then $p' \in V_{ab} \cup V_{ac} \cup \Delta_{abc}$. From the argument above it follows that $p' \in V_u$ implying that also $p \in V_u$. In the other case, $pp' \in W_{abc}$ and thus pp' is the exit edge, implying p = u. As $u \in V_u$ by definition the statement follows.

 V_{ab} and V_{ac} are strict subsets of V_u as none of them contain a, b or c. Furthermore, we have that $V_w \subseteq V_u$ for each $w \in W_{abc}$. As the unique outgoing edge of w is an entry edge and the unique outgoing edge of u is an exit edge for abc, we have $u \neq w$ and thus $V_w \subset V_u$ for all $w \in W_{abc}$.

There is an analogous statement for digons:

▶ Lemma 10. (a) If ab is a valid entry digon, then

$$V_{ab} = \bigcup_{ww' \in X_{ab}^+} V_w$$

If $|X_{ab}^+| = 1$ then $V_{ab} = V_w$ for the unique backward endpoint w. (b) If ab is a valid exit digon with exit edge uu', then

$$V_u = \{a, b\} \cup V_{ab} \cup \bigcup_{ww' \in X_{ab}^+ \setminus \{uu'\}} V_w$$

 V_u is a proper superset of each V_w on the right side.

- Proof. (a) Let p ∈ V_{ab} and r_p be the first endpoint of an edge in X⁺_{ab} on the path from p to the root. As ab is an entry digon, r_p lies on the backward side of ab, and thus p ∈ V_{r_p}. If there is more than one edge in X⁺_{ab} then the backwards endpoints do not lie in the same connected components of T \ X⁺_{ab} and thus V_{ab} is a proper super set of all sets V_w.
 (b) This proof is similar to Lemma 0.
- (b) This proof is similar to Lemma 9.
 - \subseteq : First, we show that every point in V_u is contained in at least one of the sets on the right side of the equation. Let $p \in V_u$. If p = a or p = b we are done. In the other case, let x be the first endpoint of an edge in X_{ab}^+ on the path π from p to the root. As uu' is the exit edge of the digon and $p \in V_u$, x exists. Then x is either in the backward or the forward component of $T \setminus X_{ab}^+$. If it is in the backward component, then $x \in V_{ab}$ and therefore als $p \in V_{ab}$. If x is in the forward component, let x' be next point on π . Then $xx' \in X_{ab}^+ \setminus \{uu'\}$. Hence, the path from p to x is a subpath of V_x . Therefore $p \in V_x$.
 - \supseteq : Now we show that every point in one of the sets on the right is also contained in V_u . Let $p \in V_{ab}$ and let xx' be the last edge on the path π from p to r that is in X_{ab}^+ . Then x' is a point on the forward side in X_{ab}^+ since p is on the backward side. This means that xx' is an exit edge and hence xx' = uu'. So u is on π and therefore $p \in V_u$.

If $p \in \{a, b\}$, then consider the path π from p to r. Since uu' is the unique exit edge of the chord ab, π contains u. Hence, $p \in V_u$.

Now let $p \in V_w$ for some $ww' \in X_{ab}^+ \setminus \{uu'\}$. If w' is a endpoint on the backward side of X_{ab} , then $w' \in V_{ab} \subset V_u$ and $p \in V_u$. If $w' \in \{a, b\}$, then $w' \in V_u$ by the case above. Hence, $p \in V_u$.

Note that $u \in V_u$ but $u \notin V_w$ for any w with $ww' \in X_{ab}^+ \setminus \{uu'\}$

H Preprocessing

▶ Lemma 11. The following preprocessing information can be easily computed in $O(n^3)$ total time and stored in $O(n^3)$ space:

- The MST T rooted at the lowest point r.
- For every point u, the size of the subtree V_u of the MST rooted at u.
- = For every pair of points a, b, the sets X_{ab}^+ and X_{ab} .
- For every potential chord ab, the validity of ab as a chord, the partition of $V \setminus \{a, b\}$ into backward and forward components, the set V_{ab} of vertices in the backward component, and its size.
- For each ordered pair (b, c) of points, the classification of X_{bc} into entry and exit edges as well as W_{bc} and $\bigcup_{w \in W_{ab}} V_w$. wall[b, c]
- For every edge ww' of the MST, the set ET_w of directed segments bc such that ww' is the unique exiting edge in X_{bc} .

Proof. The MST T can be computed in $O(n \log n)$ time [8]. A post order-tree traversal directly gives the sets V_u . As the total size of these sets is $O(n^2)$, reporting all sets takes $O(n^2)$ time. To compute X_{ab}^+ and X_{ab} , the MST T is traversed once for every pair a, b of points, explicitly testing all edges in T for containment in the sets. For each edge in the set, we store which endpoint is the left and which is the right endpoint. During the traversal we keep track and store the connected components of T in X_{ab}^+ . Whenever an edge $uu' \in X_{ab}^+$ is encountered, the children of u are in a new connected component. Each connected component is then traversed. Consider the component that contains r. If all leaves in this component are on the same side, then this side will be the forward side. Otherwise, ab is not a valid chord. For all other component, check if its root and all leaves are on the same side. If this is not the case, then ab is not a valid chord. In the other case, assign the component to the side defined by the vertices. If they are in the backward side, add all vertices of the component to V_{ab} . This takes O(n) time for each pair and thus $O(n^3)$ time overall.

Now consider a directed pair (b, c) of points and iterate over X_{bc} . Every edge ww' with w left of the directed line through bc is an entry edge. Add w to the set W_{bc} . The remaining edges are exit edges. If a directed pair (b, c) has only one exit edge, then add bc to ET_w . For each entry edge ww' traverse the subtree of T rooted at w to collect $\bigcup_{w \in W_{ab}} V_w$. This can be done by one traversal of w, if the vertices w are stored in W_{bc} in a traversal order on the tree. Thus again, considering one directed pairs (b, c) takes O(n) time, and the overall time of $O(n^3)$ follows.

I Solution of chord and edge problems

The chord and edge subproblems are solved as follows:

Chord problems. Let ab be a valid chord. Assume without loss of generality that it is right-facing. For the case that ab is an interior chord, we have to consider all c such that abc is a counterclockwise triangle. We can check whether abc is a valid LR-triangle by checking that (a) ac is a valid right-facing chord, (b) the wall bc has no exit edges, and in addition, (c) there are no outgoing MST edges from b or c into the region O_{abc} . Conditions (a) and (b) have been tested during preprocessing, and the last condition can be checked in constant time because the degree of a Euclidean MST is bounded by 6.

Then: The optimum length of the solution that has abc as a valid LR-triangle in the blob is the sum of the length of all problems wall[bc] plus the length of all problems edge[w] for $w \in W_{abc} \setminus W_{bc}$ plus the length of chord[a, c] plus length of the side bc.

The other possibility is that ab forms a left entering digon: This possibility is valid if all edges in X_{ab} cross from the backwards to the forward side. Then the solution of the subproblem chord[a, b] is the sum of edge[u] for $uu' \in X_{ab}$ plus the length of the line segment ab.

The value chord[a, b] is then the minimum value over all *LR*-triangles and the digon case.

Edge problems. Let $u \in P$ and uu' be the edge in the MST that is defined by edge[u]. Then in any solution, u is either incident to only tree-edges or it connects a blob. In the first case, the length of the solution is simply ||uu'|| plus the value of edge[v] for all children of v in the MST. In the second case, there is an exit triangle or exit digon with uu' being the unique exiting edge. Then the length of the solution is the minimum over all valid exit triangles abc with exiting edge uu'. Let abc be a fixed exit triangle such that uu' is the exit edge. Then the size of the solution is the $||bc|| + chord[a, b] + chord[a, c] + \sum_{w \in W_{abc}} edge[w]$. For an exit digon ab with exit edge uu' the size of the optimal solution is ||ab|| added to the sum of edge[w] for all $w \in X_{ab}$ where w is on the forward side of ab.

The problem for the root can be considered as a degenerate edge problem with a virtual predecessor r' with ||rr'|| = 0.

J Runtime analysis, proof of Theorem 7

For the running time, note that by Lemma 11 the MST and all relevant information can be preprocessed in $O(n^3)$ time. There is a total of $O(n^2)$ subproblems of the type chord[a, b]. For each the at most O(n) elements c in V_{ab} are traversed in O(n) total time. As the degree of any vertex in the MST is at most 6, only O(1) edge problems are considered in condition to one chord and one wall problem. Thus the overall time for the chord problems is $O(n^3)$. For the $O(n^2)$ wall problems, each takes O(n) edge-subproblems, for an overall of $O(n^3)$ time.

On the other hand, there are O(n) subproblems edge[u]. For each of them, the first case of the recurrence uses O(n) smaller subproblems. Here we take the minimum over all candidate walls where uu' is an exit edge. This set is precomputed by Lemma 11. Aditionally, we consider the O(n) candidate walls where one endpoint is adjacent to u. As each wall can only have one edge exiting, over all subproblems edge[u], each wall is considered for exactly one edge problem. For each candidate wall, the O(n) possible vertices a are considered. The recurrence again only refers to O(1) subproblems. Thus, $O(n^3)$ total time is needed for all edge problems.

Contents

1	Introduction	1
2	Preliminaries	2
3	Structural insights	3
4	The algorithm	7
A	Proof of Lemma 2	9
в	Proof of Lemma 4	10
С	Proof of Lemma 5	10
D	Classification of triangles and digons	11
\mathbf{E}	Proof of Lemma 6	12
F	Interaction between the chords of a valid triangle	13
G	Relation between subproblems	15
н	Preprocessing	18
Ι	Solution of chord and edge problems	18
J	Runtime analysis, proof of Theorem 7	19