## Eells-Sampson type result of Symphonic maps

Xiang-Zhi Cao\*†

March 6, 2025

#### Abstract

In this paper, we obtained Eells-Sampson type result of Symphonic maps.

### 1 Introduction

Let  $f: (M,g) \to (N,h)$  be a smooth map between Riemannian manifold. Consider the following symphonic energy of a smooth map

$$E_{sym}(f) = \int_M \frac{\|f^*h\|^2}{2} dv_g.$$

Let  $\{e_i\}_{i=1}^m$  be the local orthonormal frame. Its Euler-Lagrange equation of  $E_{sym}(f)$  is as follows (cf. [2, 10])

$$\operatorname{div}_{q} \sigma_{f} = 0. \tag{1.1}$$

where  $\sigma_f(X) = \sum_j h(df(X), df(e_j)) df(e_j)$ .

In 2011, Nakauchi and Takenaka [10] defined symphonic map, which is defined as the critical point of the above functional. One can refer to [3, 5, 4, 9, 8, 7, 6] for the related results for symphonic map.

**Theorem 1** (cf.[1]). Let  $f: (M^{m\geq 2}, g) \to (N, h)$  be a harmonic map between Riemannian manifolds. Assume that M is closed and that there exists K > 0 such that

 $\operatorname{Ric}_g \ge (m-1)Kf^*h$  and  $\operatorname{sec}_h \le K$ 

<sup>\*</sup>School of Information Engineering, Nanjing Xiaozhuang University, Nanjing 211171, China

<sup>&</sup>lt;sup>†</sup>Supported by General Project of Natural Science (Basic Science) Research of Colleges and Universities in Jiangsu Province (No. 22KJD110004)

Then f is a totally geodesic map. In particular, either i) f is constant, or ii) f is a homothetic immersion, g has positive constant curvature and on M and f(M), respectively

$$\operatorname{Ric}_q = (m-1)Kf^*h$$
 and  $\operatorname{sec}_h(\Pi) = K$ 

for any 2-plane  $\Pi$  contained in  $df(TM) \subseteq TN$ .

About symphonic map, Nakauchi has obtained the similar theorems:

**Theorem 2** (Compact case,[3]). Let (M, g) be a compact connected Riemannian manifold without boundary, and let (N, h) be a Riemannian manifold. Suppose that the Ricci curvature of M is non-negative, and that the sectional curvature of N is non-positive. Let f be any symphonic map of class C<sup>3</sup> from M into N.

#### Then

(a) The pullback  $f^*h$  is parallel, i.e.,  $\nabla(f^*h) = 0$ .

(b) If the Ricci curvature of M is positive at a point, then f is a constant map. (c) If N has negative sectional curvature everywhere, then either f is a constant map or df(x) has at most rank one for all  $x \in M$ .

The main purpose of this paper are to generalize this theorem to weak curvature condition.

**Theorem 3.** Let  $(M^m, g)$  be a compact connected Riemannian manifold without boundary, and let (N, h) be a Riemannian manifold with curvature tensor

$$R^{N}(X, Y, Z, W) = a(h(X, Z)h(Y, W)) - h(X, W)h(Y, Z)$$

Let f be any symphonic map of class  $C^3$  from M into N. Suppose that

$$\operatorname{Ric}_q \ge (m-1)Kf^*h$$
 and  $a \le K$ .

Then the pullback  $f^*h$  is parallel, i.e.,  $\nabla(f^*h) = 0$ . Moreover, f is constant map.

**Remark 1.** In [9], The author got the similar theorem, but the conditions is different here, although we both weaken the upper bound of sectional curvature of the target manifold.

Using the argument in the proof of Theorem 3, we immediately get

**Theorem 4** (Noncompact case ). Let  $M^m$  be a complete connected non-compact Riemannian manifold and let N be a Riemannian manifold with curvature tensor

$$R^{N}(X, Y, Z, W) = a(h(X, Z)h(Y, W)) - h(X, W)h(Y, Z).$$

Suppose that

$$\operatorname{Ric}_q \ge (m-1)Kf^*h$$
 and  $a \le K$ 

If  $E_{sym}(f) < \infty$ , then f is a constant map.

*Proof.* From the proof of Theorem 3, we know

$$\frac{1}{4}\Delta \|f^*h\|^2 \ge \operatorname{div}_g \alpha_f + \frac{1}{2} \|\nabla (f^*h)\|^2.$$

the rest proof proceeds as in [3].

Acknowledgement: This work is Supported by General Project of Natural Science (Basic Science) Research of Colleges and Universities in Jiangsu Province (No. 22KJD110004)

In this paper, we use notation:  $R^N(X, Y, Z, W) = \langle R(X, Y)W, Z \rangle$ 

### 2 the proof of main theorem

**Lemma 1.** (1) If symmetric matrices  $A = (a_{ij})$  and  $B = (b_{ij})$  are positive semi-definite, then we have

$$\operatorname{tr}\left(A^{t}B\right) = \sum_{i,j=1}^{m} a_{ij}b_{ij} \ge 0$$

(2) If A is positive definite and B is positive semi-definite, then tr  $(A^tB) = 0$  implies B = 0.

**Lemma 2** (Formula of Bochner type, cf.[3]). For any map f of class C<sup>3</sup> from M into N, the following equality holds:

$$\begin{split} &\frac{1}{4} \Delta \|f^*h\|^2 \\ &= \operatorname{div}_g \alpha_f - h\left(\tau_f, \operatorname{div}_g \sigma_f\right). \\ &+ \sum_{i,j,k=1}^m h\left(\left(\nabla_{e_k} df\right)(e_i), \left(\nabla_{e_k} df\right)(e_j)\right) h\left(df\left(e_i\right), df\left(e_j\right)\right) \\ &+ \frac{1}{2} \|\nabla\left(f^*h\right)\|^2 \\ &+ \sum_{i,j,k=1}^m h\left(df\left(\operatorname{Ric}^M\left(e_i, e_k\right)(e_k)\right), df\left(e_j\right)\right) h\left(df\left(e_i\right), df\left(e_j\right)\right) \\ &- \sum_{i,j,k=1}^m h\left(R^N\left(df\left(e_i\right), df\left(e_k\right)\right) df\left(e_k\right), df\left(e_j\right)\right) h\left(df\left(e_i\right), df\left(e_j\right)\right), \end{split}$$

where  $\{e_i\}$  denotes a local orthonormal frame around the point of interest in M. Proof of Theorem 3. Let

$$a_{ik} = R_{ik} - (m - 1)Kb_{ik}$$

$$c_{ik} = \sum_{j=1}^{m} h(df(e_k), df(e_j)) h(df(e_i), df(e_j)).$$

$$d_{ij} = -\sum_{k=1}^{m} h(R^N(df(e_i), df(e_k)) df(e_k), df(e_j))$$

$$b_{ij} = h(df(e_i), df(e_j))$$

It is easy to see that the symmetric matrix  $(a_{ij})$  is positive semi-definite. Notice that

$$\sum_{i,k=1}^{m} R_{ik} \sum_{j=1}^{m} h\left(df\left(e_{k}\right), df\left(e_{j}\right)\right) h\left(df\left(e_{i}\right), df\left(e_{j}\right)\right) \\ - (m-1)K \sum_{k=1}^{m} h\left((df\left(e_{i}\right), df\left(e_{k}\right)\right) df\left(e_{k}\right), df\left(e_{j}\right)\right) b_{ij} \\ = R_{ik} b_{kj} b_{ij} - (m-1)K b_{ik} b_{kj} b_{ij} \\ = (R_{ik} - (m-1)K b_{ik}) b_{kj} b_{ij} \\ \ge 0.$$

and

$$(m-1)K\sum_{k=1}^{m} h\left((df(e_i), df(e_k)) df(e_k), df(e_j)\right) b_{ij}$$
  
$$-\sum_{i,j,k=1}^{m} h\left(R^N\left(df(e_i), df(e_k)\right) df(e_k), df(e_j)\right) b_{ij}$$
  
$$=(m-1)Kb_{ik}b_{kj}b_{ij} - \sum_{i,j,k=1}^{m} h\left(R^N\left(df(e_i), df(e_k)\right) df(e_k), df(e_j)\right) b_{ij}$$
  
$$=(m-1)Kb_{ik}b_{kj}b_{ij} - a(b_{ik}b_{kj} - b_{ij}b_{kk})b_{ij}$$
  
$$=((m-1)K - a)b_{ik}b_{kj}b_{ij} + b_{ij}^2b_{kk}$$

By divergence theorem on compact manifold, we have

$$\begin{split} 0 &\geq \int_{M} \sum_{i,j,k=1}^{m} h\left(\left(\nabla_{e_{k}} df\right)(e_{i}\right), \left(\nabla_{e_{k}} df\right)(e_{j})\right) h\left(df\left(e_{i}\right), df\left(e_{j}\right)\right) \\ &+ \int_{M} \frac{1}{2} \left\|\nabla\left(f^{*}h\right)\right\|^{2} \\ &+ \int_{M} \sum_{i,j,k=1}^{m} h\left(df\left(Ric^{M}\left(e_{i}, e_{k}\right)(e_{k})\right), df\left(e_{j}\right)\right) h\left(df\left(e_{i}\right), df\left(e_{j}\right)\right) \\ &- (m-1)K \sum_{k=1}^{m} h\left(\left(df\left(e_{i}\right), df\left(e_{k}\right)\right) df\left(e_{k}\right), df\left(e_{j}\right)\right) b_{ij} dv \\ &+ \int_{M} (m-1)K \sum_{k=1}^{m} h\left(\left(df\left(e_{i}\right), df\left(e_{k}\right)\right) df\left(e_{k}\right), df\left(e_{j}\right)\right) b_{ij} \\ &- \sum_{i,j,k=1}^{m} h\left(R^{N}\left(df\left(e_{i}\right), df\left(e_{k}\right)\right) df\left(e_{k}\right), df\left(e_{j}\right)\right) h\left(df\left(e_{i}\right), df\left(e_{j}\right)\right) dv \\ &\geq \int_{M} (R_{ik} - (m-1)Kb_{ik})b_{kj}b_{ij} + \left((m-1)K - a\right)b_{ik}b_{kj}b_{ij} + b_{ij}^{2}b_{kk}dv_{g} \end{split}$$

We can conclude from the above inequality that  $\nabla(f^*h) = 0$  and df = 0.

# References

 COLOMBO, G., MARIANI, M., AND RIGOLI, M. A sharp Eells-Sampson type theorem under positive sectional curvature upper bounds. J. Math. Anal. Appl. 540, 1 (2024), Paper No. 128584, 10. 1

- [2] KAWAI, S., AND NAKAUCHI, N. Some results for stationary maps of a functional related to pullback metrics. *Nonlinear Anal.* 74, 6 (2011), 2284–2295. 1
- [3] KAWAI, S., AND NAKAUCHI, N. Results of Liouville type for symphonic maps. Differential Geom. Appl. 65 (2019), 147–159. 1, 2, 3
- [4] MISAWA, M., AND NAKAUCHI, N. A Hölder continuity of symphonic maps into the spheres. *Calc. Var. Partial Differential Equations* 55, 1 (2016), Art. 16, 20. 1
- [5] MISAWA, M., AND NAKAUCHI, N. Global existence for the heat flow of symphonic maps into spheres. Adv. Differential Equations 23, 9-10 (2018), 693–724. 1
- [6] MISAWA, M., AND NAKAUCHI, N. On the finite-time blow-up of symphonic map flows. *Differential Integral Equations* 36, 1-2 (2023), 93–131.
- [7] NAKAUCHI, N. Stress energy tensor for symphonic maps. Boll. Unione Mat. Ital. 12, 3 (2019), 431–440.
- [8] NAKAUCHI, N. Rotationally symmetric symphonic maps. Ann. Global Anal. Geom. 62, 1 (2022), 83–92. 1
- [9] NAKAUCHI, N. Two results for symphonic maps under assumptions on msymphonic energy. Results Math. 77, 6 (2022), Paper No. 216, 23. 1, 2
- [10] NAKAUCHI, N., AND TAKENAKA, Y. A variational problem for pullback metrics. *Ric. Mat. 60*, 2 (2011), 219–235. 1