

A NOTE ON SPECTRAL VOLUME COMPARISON FOR MANIFOLDS WITH WEAKLY CONVEX BOUNDARY

JIA LI

ABSTRACT. We establish a new boundary version of spectral volume comparison theorem in this paper. Let (M^n, g) ($n \geq 3$) be a connected, smooth compact n -dimensional manifold with weakly convex boundary. If there exist a positive function $w \in C^\infty(M)$ that satisfies:

$$\begin{cases} -\frac{n-1}{n-2}\Delta w + \text{Ric } w \geq (n-1)w, \\ \frac{\partial w}{\partial \eta} = 0. \end{cases}$$

then $\text{Vol}(M) \leq \text{Vol}(\mathbb{S}_+^n)$, and equality holds if and only if M^n is isometric to the unit round hemisphere $\mathbb{S}_+^n = \{x \in \mathbb{R}^{n+1} : |x| = 1, x_{n+1} \geq 0\} \subset \mathbb{R}^{n+1}$.

1. INTRODUCTION

A classical Bonnet-Myers theorem states that if a complete n -dimensional Riemannian manifold M has Ricci curvature Ric_M at least $(n-1)K$, then the diameter of M is at most $\frac{\pi}{\sqrt{K}}$, moreover, it follows from Bishop-Gromov volume comparison theorem that for a fixed $p \in M$, the ratio $\frac{\text{Vol}(B_r(p))}{\text{Vol}(B^K(r))}$ is nonincreasing for any $r \in (0, \infty)$. When $K > 0$, the equality holds if and only if M is isometric to $\mathbb{S}^n(\frac{1}{\sqrt{K}})$. S. Y. Cheng in [2] proved a rigidity theorem which states that if the diameter is equal to $\frac{\pi}{\sqrt{K}}$, then M is isometric to the n -sphere with constant sectional curvature K . Recently, G. Antonelli and K. Xu [1] prove a sharp and rigid spectral generalization of both Bonnet-Myers theorem and classical Bishop-Gromov volume comparison theorem, which is precisely stated as follows.

Theorem 1 ([1]). *Let M^n , $n \geq 3$, be an n -dimensional compact smooth manifold with $n \geq 3$, and let $0 \leq \theta \leq \frac{n-1}{n-2}$, $\lambda > 0$. Let $\text{Ric}(x) := \inf_{v \in T_p M, |v|=1} \text{Ric}_x(v, v)$ be the smallest eigenvalue of the Ricci tensor. Assume there is a positive function $w \in C^\infty(M)$ such that:*

$$\theta \Delta_M w \leq \text{Ric } w - (n-1)\lambda w,$$

Let \tilde{M} be the universal cover of M , endowed with the pull-back metric. Then:

- A diameter bound

$$\text{diam}(M) \leq \frac{\pi}{\sqrt{\lambda}} \left(\frac{\max w}{\min w} \right)^{\frac{n-3}{n-1}\theta},$$

in particular, $\pi_1(M)$ is finite.

- A sharp volume bound

$$\text{Vol}(\tilde{M}) \leq \lambda^{-\frac{n}{2}} \text{Vol}(\mathbb{S}^n).$$

Moreover, if equality holds, then every function w is constant, and \tilde{M} is isometric to the round sphere of radius $\lambda^{-\frac{1}{2}}$.

This theorem is intimately connected with stable Bernstein problem. In order to address the stable Bernstein problem in \mathbb{R}^5 , C. Chodosh, C. Li, P. Minter and D. Stryker [5] established a volume comparison theorem via weighted isoperimetric profile, but their method works only for 3-dimensional μ -bubbles. It is remarkable that the weakness has been overcome by the unequally weighted isoperimetric profile in [1], which paved the way to study the stable Bernstein problem. Soon, combining the strategy of [5] with the Theorem 1, L. Mazet successfully solved the stable Bernstein problem in \mathbb{R}^6 . Additionally, the spectral volume comparison theorem is also a useful tool in other aspects such as the structure of manifolds.

For compact manifold with boundary and positive Ricci curvature, F. Hang and X. Wang [8] proved a boundary rigidity result as follows:

Theorem 2. *Let (M^n, g) ($n \geq 2$) be a compact manifold with nonempty boundary $\Sigma = \partial M$. Suppose*

- $\text{Ric} \geq (n-1)g$,
- $(\Sigma, g|_\Sigma)$ is isometric to the standard sphere $\mathbb{S}^{n-1} \subset \mathbb{R}^n$,
- Σ is weakly convex in the sense that its second fundamental form A is nonnegative.

Then (M^n, g) is isometric to the hemisphere.

It is natural to investigate whether the theorem 1 can be extended to compact manifolds with boundary. Based on the method in [1], the author in [9] has obtained the spectral diameter estimate of 3-dimensional compact manifold with convex boundary, in order to prove the non-existence of weakly stable free boundary CMC hypersurfaces in 5-manifolds. For general n -dimensional manifold, we establish a spectral generalization of Bonnet-Myers and Bishop-Gromov volume comparison theorem for compact manifold with weakly convex boundary.

Theorem 3. *Let M^n , $n \geq 3$, be a compact connected manifold with weakly convex boundary ∂M , and let $0 \leq \theta \leq \frac{n-1}{n-2}$, $\lambda > 0$. We denote by $\text{Ric}(x) := \inf_{v \in T_x M, |v|=1} \text{Ric}_x(v, v)$ the smallest eigenvalue of the Ricci tensor. Assume there is a positive function $w \in C^\infty(M)$ satisfying Neumann boundary condition $\frac{\partial w}{\partial \eta} = \langle \nabla^M w, \eta \rangle = 0$, and the following inequality holds:*

$$\theta \Delta_M w \leq \text{Ric} w - (n-1)\lambda w,$$

then we have the diameter estimate:

$$\text{diam}(M) \leq \frac{\pi}{\sqrt{\lambda}} \left(\frac{\max w}{\min w} \right)^{\frac{n-3}{n-1}\theta},$$

and the volume estimate:

$$\text{Vol} M \leq \lambda^{-\frac{n}{2}} \text{Vol}(\mathbb{S}_+^n).$$

Moreover, if equality holds, then every function w is constant, and \tilde{M} is isometric to the round hemisphere of radius $\lambda^{-\frac{1}{2}}$.

1.1. organization. In section 2, we give some basic definitions and properties about isoperimetric profiles. In section 3, we will use free boundary μ -bubbles to obtain the diameter estimate of n -dimensional compact manifold with weakly convex boundary, which is the generalization of Theorem 32 in [9]. In section 4, we establish the unequally weighted free boundary isoperimetric profile, get a differential inequality in the barrier sense, finally we obtain the volume bound.

Acknowledgements. *The author would like to thank his advisor, Professor Chao Xia, for introducing this problem and for his continuous guidance, support, and invaluable insights throughout this research. His encouragement and expertise have been instrumental in the completion of this work.*

2. PRELIMINARIES

First, we assume all manifolds are complete in this paper. We now introduce the notions of μ -bubbles and isoperimetric profile of a Riemannian manifold. It's remarkable that it can be defined in any space with suitable notions of volume and perimeter as in metric measure spaces.

2.1. free boundary unequally warped μ -bubbles.

Definition 4. We say that a manifold $(M^n, \partial M)$ has weakly convex boundary if and only if $A \geq 0$, where A is the second fundamental form of ∂M with respect to η , and η is the co-normal of ∂M in M .

Given a compact n -dimensional Riemannian manifold (N^n, g) with weakly convex boundary $\partial N = \partial_0 N \cup \partial_- N \cup \partial_+ N$ ($\partial_i N$ is nonempty for $i \in \{0, -, +\}$), where $\partial_- N$ and $\partial_+ N$ are disjoint and each of them intersect with $\partial_0 N$ at angles no more than $\frac{\pi}{8}$ inside M . We fix a smooth function $w > 0$ on N and a smooth function h on $N \setminus (\partial_- N \cup \partial_+ N)$, with $h \rightarrow \pm\infty$ on $\partial_\pm N$. We pick a regular value c_0 of h on $N \setminus (\partial_- N \cup \partial_+ N)$ and take $\Omega_0 = h^{-1}((c_0, \infty))$ as the reference set, and consider the following area functional:

$$\mathcal{A}(\Omega) := \int_{\partial^* \Omega} w^\theta d\mathcal{H}^{n-1} - \int_N (\chi_\Omega - \chi_{\Omega_0}) h w^\alpha d\mathcal{H}^n,$$

for all Caccioppoli sets Ω with $\Omega \Delta \Omega_0 \Subset \overset{\circ}{N}$ and $k \in \mathbb{N}$, where $\theta, \alpha \geq 0$, and the reference set Ω_0 with smooth boundary satisfies

$$\partial \Omega_0 \subset \overset{\circ}{N}, \quad \partial_+ N \subset \Omega_0.$$

If there exist Ω can minimizes \mathcal{A} in this class, we call it a free boundary unequally warped μ -bubble.

About the existence and regularity of a minimizer of \mathcal{A} among all Caccioppoli sets, we refer to [3, 7, 13, 14] for more details. We only elaborate the conclusions and omit the detailed proof.

Proposition 5. *There exists a smooth minimizer Ω for \mathcal{A} such that $\Omega \Delta \Omega_0$ is compactly contained in $\overset{\circ}{N} \cup \partial_0 N$. The minimizer has smooth boundary which intersect with $\partial_0 N$ orthogonally.*

Let ν_Σ denote by the unit normal of Σ , and let $\phi \in C^\infty(N)$. For an arbitrary variation $\{\Omega_t\}_{t \in (-\epsilon, \epsilon)}$ with $\Omega_0 = \Omega$ and the variational vector field $\dot{F}_t = \phi \nu_\Sigma$ at $t = 0$, we assume that $\Sigma = \partial\Omega$ is a critical point of the area functional and calculate its first variation and the second variation.

Lemma 6. *If Ω_t is a smooth 1-parameter family of regions with $\Omega_0 = \Omega$ and the normal variational vector field \dot{F}_t at $t = 0$ is $\phi \nu_\Sigma$, then*

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} \mathcal{A}(\Omega_t) &= \int_{\partial\Omega_t} \theta w^{\theta-1} \langle \nabla^M w, \nu_{\Sigma_t} \rangle \phi + w^\theta H \phi - h w^\alpha \phi + \int_{\partial\Sigma_t} w^\theta \langle \dot{F}_t, \nu_{\partial\Sigma_t} \rangle \\ &= \int_{\Sigma} (H_\Sigma + \theta w^{-1} \langle \nabla^M w, \nu_\Sigma \rangle - h w^{\alpha-\theta}) w^\theta \phi \end{aligned}$$

where ν_Σ is the outward pointing unit normal vector field and H is the mean curvature of $\partial\Omega_t$. In particular, a μ -bubble Ω with $\partial\Omega = \Sigma$ satisfies

$$H_\Sigma = h w^{\alpha-\theta} - \theta w^{-1} \langle \nabla^M w, \nu_\Sigma \rangle, \quad \dot{F}_0 \perp \nu_{\partial\Sigma} \quad \text{for } x \in \partial\Sigma \subset \partial_0 N.$$

Lemma 7. *Assume Ω is a minimizer of \mathcal{A} in the settings above, then we can have the following the second variational formula:*

$$\begin{aligned} \frac{d^2}{dt^2} \Big|_{t=0} (\mathcal{A}(\Omega_t)) &= \int_{\Sigma} [-\Delta_\Sigma \phi - |\text{II}_\Sigma|^2 \phi - \text{Ric}_M(\nu_\Sigma, \nu_\Sigma) \phi - \theta w^{-2} \langle \nabla^M w, \nu_\Sigma \rangle^2 \phi \\ &\quad + \theta w^{-1} \phi (\Delta_M w - \Delta_\Sigma w - H_\Sigma \langle \nabla^M w, \nu_\Sigma \rangle - \theta w^{-1} \langle \nabla^\Sigma w, \nabla^\Sigma \phi \rangle \\ &\quad - \phi \langle \nabla^M h, \nu_\Sigma \rangle w^{\alpha-\theta} + (\theta - \alpha) w^{\alpha-\theta-1} h \phi \langle \nabla^M w, \nu_\Sigma \rangle] w^\theta \phi + \int_{\partial\Sigma} w^\theta \phi \frac{\partial \phi}{\partial \nu_{\partial\Sigma}} - A_{\partial\Sigma}(\nu_\Sigma, \nu_\Sigma) \phi^2 w^\theta \end{aligned}$$

2.2. free boundary unequally warped isoperimetric profile.

Definition 8. Let (M^n, g) be a Riemannian manifold. The isoperimetric profile of M is the function I_M that assigns, to each $v \in (0, |M|)$, the value

$$I_M(v) = \inf\{P(E) : E \text{ is measurable, } |E| = v\}.$$

The isoperimetric profile of M will be often denoted by I .

Definition 9. Let (M, g) be a Riemannian manifold. We say that a set $E \subset M$ is isoperimetric or that is an isoperimetric region if

$$P(E) = I_M(|E|).$$

If $|E| = v$, then we say that E is an isoperimetric region of volume v .

We notice that the isoperimetric profile can be understood as an optimal isoperimetric inequality in M , since for any subset $F \subset M$ of volume $0 < |F| < |M|$, $P(F) \geq I_M(|F|)$, with equality precisely for isoperimetric sets.

Regularity results for sets minimizing perimeter under a volume constraint were obtained by F. Morgan, who proved in Corollary 3.7 and 3.8 of [6] in the following:

Theorem 10 ([6]). *Let E be a measurable set of finite volume minimizing perimeter under a volume constraint in a smooth n -dimensional Riemannian manifold M . There are two kinds of conclusions depended on dimension:*

- *If $n \leq 7$, then the boundary S of E is a smooth hypersurface.*
- *If $n > 7$, then the boundary S of E is the union of a smooth hypersurface S and a closed singular set S_0 of Hausdorff dimension at most $n - 8$ (i.e., $\mathcal{H}^{n-8+\gamma}(S_0) = 0$ for all $\gamma > 0$).*

When trying to get geometric information on the boundary of a set minimizing perimeter under a volume constraint, the following technical result proved by Sternberg and Zumbrun in Lemma 2.4 of [12] allows us to focus just on the regular part S of the boundary.

Lemma 11 ([12]). *Let $E \subset M$ be a bounded minimizer of perimeter under a volume constraint in a smooth Riemannian manifold M . Let S be the regular part of the boundary of E and S_0 its singular part.*

Then for every $\epsilon > 0$, there exist open sets $U' \Subset S \subset M$ with $S_0 \subset U'$, $U \subset M$ contained in an open tubular neighborhood in M of S_0 of radius ϵ , and a smooth function $\varphi_\epsilon : M \rightarrow \mathbb{R}$ such that $0 \leq \varphi_\epsilon \leq 1$,

$$\varphi_\epsilon(x) = 0 \text{ in } U', \quad \varphi_\epsilon(x) = 1 \text{ in } M \setminus U,$$

and

$$\int_S |\nabla_S \varphi_\epsilon|^2 dS \leq C\epsilon,$$

for some constant $C > 0$ depending on E but independent of ϵ .

Given an open set $\Omega \subset M$, we may consider the minimization problem

$$\inf\{P(E) : E \subset \bar{\Omega}, |E| = v\}.$$

We emphasize that the perimeter here is the one in M . There is a regularity result about this restricted profile, which can be found in [11].

Theorem 12. *Let (M, g) be a complete Riemannian manifold and $\Omega \subset M$ be an open subset with C^2 boundary. Then for any $0 < v < |\Omega|$, there exist an isoperimetric region E of volume v in Ω . Moreover, the boundary of E satisfies the following properties:*

- *∂E is of class $C^{1,1}$ near $\partial\Omega$.*
- *$\partial E \cap \Omega$ is the union of a smooth hypersurface S with constant mean curvature H and a closed singular set S_0 of Hausdorff dimension at most $n - 8$.*

3. DIAMETER ESTIMATES FOR MANIFOLDS WITH CONVEX BOUNDARY

For now on, we always assume that $n \geq 3$ throughout the paper. We first consider n -dimensional compact manifold with weakly convex boundary, then we can give the diameter estimates in the sense of spectrum condition and weakly convex boundary condition.

3.1. for case $3 \leq n \leq 7$.

Theorem 13. *Let M^n , $n \geq 3$, be a compact connected manifold with weakly convex boundary ∂M , and let $0 \leq \theta \leq \frac{n-1}{n-2}$, $\lambda > 0$. We denote by $\text{Ric}(x) := \inf_{v \in T_x M, |v|=1} \text{Ric}_x(v, v)$ the smallest eigenvalue of the Ricci tensor. If there exist a positive function $w \in C^\infty(M)$ that satisfies:*

$$\begin{cases} \theta \Delta_M w \leq \text{Ric} w - (n-1)\lambda w, \\ \frac{\partial w}{\partial \eta} = 0. \end{cases}$$

then we have the diameter estimate: $\text{diam}(M) \leq \frac{\pi}{\sqrt{\lambda}} \left(\frac{\max w}{\min w} \right)^{\frac{n-3}{n-1} \theta}$.

Proof. Let $\Omega_+ \subset \Omega_- \subset M$ be two domains with nonempty boundaries, such that $\overline{\Omega_-} \setminus \Omega_+$ is compact. Suppose $h(x) \in C^\infty(\overline{\Omega_-} \setminus \Omega_+)$ satisfies

$$(1) \quad \lim_{x \rightarrow \partial \Omega_+} h(x) = +\infty, \quad \lim_{x \rightarrow \partial \Omega_-} h(x) = -\infty.$$

For an arbitrary fixed domain Ω_0 with $\Omega_+ \subset \subset \Omega_0 \subset \subset \Omega_-$, consider the functional

$$\mathcal{A}(\Omega_t) = \int_{\partial^* \Omega_t} w^\theta - \int_{\Omega_t} h w^\alpha,$$

defined on sets of finite perimeter, where $\partial^* \Omega$ is the reduced boundary of Ω . According to the Lemma 6 and Lemma 7, we have the second variation formula in the following:

$$\begin{aligned} & \frac{d^2}{dt^2} \Big|_{t=0} \mathcal{A}(\Omega_t) \\ &= \int_{\Sigma} [-\Delta_{\Sigma} \phi - |\text{II}_{\Sigma}|^2 \phi - \text{Ric}_M(\nu_{\Sigma}, \nu_{\Sigma}) \phi - \theta w^{-2} \langle \nabla^M w, \nu_{\Sigma} \rangle^2 \phi \\ & \quad + \theta w^{-1} \phi (\Delta_M w - \Delta_{\Sigma} w - H_{\Sigma} \langle \nabla^M w, \nu_{\Sigma} \rangle - \theta w^{-1} \langle \nabla^{\Sigma} w, \nabla^{\Sigma} \phi \rangle \\ & \quad - \phi \langle \nabla^M h, \nu_{\Sigma} \rangle w^{\alpha-\theta} + (\theta - \alpha) w^{\alpha-\theta-1} h \phi \langle \nabla^M w, \nu_{\Sigma} \rangle] w^\theta \phi + \int_{\partial \Sigma} w^\theta \phi \frac{\partial \phi}{\partial \nu_{\partial \Sigma}} - A_{\partial \Sigma}(\nu_{\Sigma}, \nu_{\Sigma}) \phi^2 w^\theta \end{aligned}$$

We choose $\phi = w^{-\theta}$, then we have

$$\int_{\Sigma} -\Delta_{\Sigma} \phi = \theta \int_{\partial \Sigma} w^{-\theta-1} \langle \nabla^{\Sigma} w, \nu_{\partial \Sigma} \rangle$$

Integrating by parts, we have

$$\begin{aligned}
 & -\theta \int_{\Sigma} w^{-1} w^{-\theta} \Delta_{\Sigma} w + w^{-1} \langle \nabla^{\Sigma} w, \nabla^{\Sigma} (w^{-\theta}) \rangle \\
 &= -\theta \int_{\Sigma} w^{-\theta-1} \Delta_{\Sigma} w + w^{-1} \langle \nabla^{\Sigma} w, \nabla^{\Sigma} (w^{-\theta}) \rangle + \langle \nabla^{\Sigma} w, \nabla^{\Sigma} (w^{-1}) \rangle w^{-\theta} \\
 &\quad - \langle \nabla^{\Sigma} w, \nabla^{\Sigma} (w^{-1}) \rangle w^{-\theta} \\
 &= -\theta \int_{\Sigma} w^{-\theta-1} + \langle \nabla^{\Sigma} w, \nabla^{\Sigma} (w^{-\theta-1}) \rangle + \theta \int_{\Sigma} \langle \nabla^{\Sigma} w, \nabla^{\Sigma} (w^{-1}) \rangle w^{-\theta} \\
 &= -\theta \int_{\partial \Sigma} \langle \nabla^{\Sigma} w, \nu_{\partial \Sigma} \rangle w^{-\theta-1} - \theta \int_{\Sigma} |\nabla^{\Sigma} w|^2 w^{-2-\theta} \\
 &\leq -\theta \int_{\partial \Sigma} \langle \nabla^{\Sigma} w, \nu_{\partial \Sigma} \rangle w^{-\theta-1}.
 \end{aligned}$$

On the other hand, we have

$$|\Pi_{\Sigma}|^2 \geq \frac{1}{n-1} H_{\Sigma}^2 = \frac{1}{n-1} \left(h - \frac{4}{4-k} w^{-1} \langle \nabla^M w, \nu_{\Sigma} \rangle \right)^2,$$

and

$$\langle \nabla^{\Sigma} w, \nu_{\partial \Sigma} \rangle = \langle \nabla^M w, \nu_{\partial \Sigma} \rangle = 0, \quad A \geq 0 \text{ on } \partial \Sigma,$$

which is guaranteed by the weakly convexity of ∂M . Combining these consequences with nonnegativity of the second variation, we obtain that

$$\begin{aligned}
 0 &\leq \int_{\Sigma} -|\Pi_{\Sigma}|^2 w^{-\theta} - \text{Ric}_M(\nu_{\Sigma}, \nu_{\Sigma}) w^{-\theta} - \theta w^{-2} \langle \nabla^M w, \nu_{\Sigma} \rangle^2 \\
 &\quad + \theta w^{-1-\theta} (\Delta_M w - H_{\Sigma} \langle \nabla^M w, \nu_{\Sigma} \rangle) - \langle \nabla^M h, \nu_{\Sigma} \rangle w^{\alpha-2\theta} + (\theta - \alpha) h w^{\alpha-2\theta-1} \langle \nabla^M w, \nu_{\Sigma} \rangle \\
 &\leq w^{-\theta} \left[-\frac{H_{\Sigma}^2}{n-1} - (n-1)\lambda - \theta (w^{-1} \langle \nabla^M w, \nu_{\Sigma} \rangle)^2 - \theta H_{\Sigma} (w^{-1} \langle \nabla^M w, \nu_{\Sigma} \rangle) + |\nabla h| w^{\alpha-\theta} \right. \\
 &\quad \left. + (\theta - \alpha) h w^{\alpha-\theta} (w^{-1} \langle \nabla^M w, \nu_{\Sigma} \rangle) \right]
 \end{aligned}$$

Setting $X = h w^{\alpha-\theta}$, $Y = w^{-1} \langle \nabla^M w, \nu_{\Sigma} \rangle$, then $H_{\Sigma} = X - \theta Y$, we have

$$\begin{aligned}
 0 &\leq \int_{\Sigma} w^{-\theta} \left[-\frac{X^2}{n-1} + \frac{2\theta}{n-1} XY - \frac{\theta^2}{n-1} Y^2 - (n-1)\lambda - \theta Y^2 - \theta(X - \theta Y)Y + |\nabla h| w^{\alpha-\theta} + (\theta - \alpha)XY \right] \\
 &\leq \int_{\Sigma} w^{-\theta} \left[-\frac{X^2}{n-1} - (n-1)\lambda + |\nabla h| w^{\alpha-\theta} + \left(\frac{2\theta}{n-1} - \alpha \right) XY + \left(\frac{n-2}{n-1} \theta^2 - \theta \right) Y^2 \right]
 \end{aligned}$$

If we set $\alpha = \frac{2\theta}{n-1}$, for $0 \leq \theta \leq \frac{n-1}{n-2}$. Additionally, if h satisfies

$$|\nabla h| w^{\alpha-\theta} < \frac{h^2 w^{2\alpha-2\theta}}{n-1} + (n-1)\lambda,$$

then we get a contradiction. Therefore, we are going to construct such h satisfying above condition. Then we get the estimate of the diameter:

$$\text{diam}(M) \leq \frac{\pi}{\sqrt{\lambda}} \cdot \left(\frac{\sup(w)}{\inf(w)} \right)^{\frac{n-3}{n-1}\theta}.$$

First, notice that h is a smooth function on M such that

$$(2) \quad |\nabla h| < \frac{\sup(w)^{2\alpha-2\theta}}{(n-1)\inf(w)^{\alpha-\theta}} h^2 + \frac{(n-1)\lambda}{\inf(w)^{\alpha-\theta}} =: Ch^2 + D,$$

then

$$|\nabla h|w^{\alpha-\theta} \leq |\nabla h|\inf(w)^{\alpha-\theta} < \frac{h^2w^{2\alpha-2\theta}}{n-1} + (n-1)\lambda.$$

Moreover, we notice that

$$CD = \lambda \cdot \left(\frac{\sup(w)}{\inf(w)} \right)^{\frac{6-2n}{n-1}\theta}.$$

We can use the same method derived from [1]. Suppose by contradiction the above estimate does not hold, then there is a $\epsilon > 0$ such that

$$(3) \quad \text{diam}(M) > \frac{\pi}{\sqrt{\lambda}} \cdot \left(\frac{\sup(w)}{\inf(w)} \right)^{\frac{n-3}{n-1}\theta} \cdot (1+\epsilon)^2 + 2\epsilon.$$

Let us fix a point $p \in \partial M$ and take $\Omega_+ := B_\epsilon(p)$, and let $d : M \setminus \Omega_+ \rightarrow \mathbb{R}$ be a smoothing of $d(\cdot, \partial\Omega_+)$ such that

$$d|_{\partial\Omega_+} = 0, \quad |\nabla d| \leq 1 + \epsilon, \quad d \geq \frac{d(\cdot, \partial\Omega_+)}{1 + \epsilon}.$$

We can define

$$h(x) := \sqrt{\frac{D}{C}} \cot\left(\frac{\sqrt{CD}}{1+\epsilon}d(x)\right),$$

which satisfies inequality (2). Note we have that

$$\mathcal{O} := \left\{d > \frac{2(1+\epsilon)\pi}{\sqrt{2\epsilon_0}}\right\} \supset \left\{d(\cdot, p) > \epsilon + \frac{2(1+\epsilon)^2\pi}{\sqrt{2\epsilon_0}}\right\} \neq \emptyset,$$

due to (3.1). Set $\Omega_- := M \setminus \bar{\mathcal{O}}$, then we have found two domains $\Omega_+ \subset\subset \Omega_- \subset\subset M$ and $h(x) \in C^\infty(\Omega_- \setminus \bar{\Omega}_+)$ which satisfies (1). Let Ω be an free boundary μ -bubble minimizing the functional, which has smooth boundary. Since the inequality (2) is in force, we get a contradiction with the minimizing property of $\mathcal{A}(\Omega_t)$. \square

3.2. for case $n \geq 8$.

Remark 14. We notice that above proof works for $3 \leq n \leq 7$ for free boundary μ -bubble, there will exist singularity formation in higher dimensions, the diameter estimate for $n \geq 8$ can also be obtained by modifying above argument, we refer readers to [1, Appendix A].

4. VOLUME COMPARISON FOR MANIFOLD WITH WEAKLY CONVEX BOUNDARY

4.1. for case $3 \leq n \leq 7$.

4.1.1. *Unequally weighted isoperimetric profile.* Let $n \geq 3$, and (M^n, g) be a compact manifold Riemannian manifold with weakly convex boundary. Let $0 \leq \theta \leq \frac{n-1}{n-2}$, and set

$$\alpha := \frac{2\theta}{n-1}.$$

For an open set $E \subset M$ with smooth boundary, we can define unequally weighted area and volume functional by

$$A(E) = \int_{\partial^* E} w^\theta \text{ and } V(E) = \int_E w^\alpha,$$

where w is the smooth positive first eigenfunction of $-\Delta - \mathbf{V}$ in M . Namely, w satisfies

$$-\Delta w \geq \theta^{-1}((n-1)\lambda - \text{Ric})w, \text{ and } \langle \nabla^M w, \eta \rangle = 0.$$

Let $V_0 := \int_M w^\alpha \in (0, \infty]$, and define the unequally weighted isoperimetric profile

$$(4) \quad I(v) := \inf \left\{ \int_{\partial^* E} w^\theta : E \subset\subset M \text{ has finite perimeter, and } \int_E w^\alpha = v \right\},$$

for all $v \in [0, V_0)$.

Remark 15. *We note that I is continuous. On the one hand, by the compactness theory for Caccioppoli sets and the lower semi-continuity, we have $\liminf_{v \rightarrow v_0} I(v) \geq I(v_0)$. On the other hand, we will show that there exists a continuous upper barrier function for I at v_0 for any $v_0 \in (0, V)$, we also have $\limsup_{v \rightarrow v_0} I(v) \leq I(v_0)$.*

4.1.2. *First variation.* We compute the first variation of the functionals A and V . Let E_t be a smooth family of open sets with sooth boundary whose variational vector field along $\Gamma = \partial E = \partial E_0$ is $\varphi\nu$, where ν is the outward pointing unit normal vector field along ∂E . We can directly compute its first variational formulas.

Proposition 16.

$$\frac{d}{dt} \Big|_{t=0} V(E_t) = \int_\Gamma w^\theta \varphi \text{ and } \frac{d}{dt} \Big|_{t=0} A(E_t) = \int_\Gamma w^\theta \varphi (H + \theta w^{-1} \langle \nabla^M w, \nu \rangle).$$

4.1.3. *Second variation.* We proceed to compute the second variation of the functionals A and V . The process is similar to the free boundary minimal hypersurfaces.

Proposition 17.

$$\frac{d^2}{dt^2} \Big|_{t=0} V(E_t) = \int_\Gamma (H + \alpha w^{-1} \langle \nabla^M w, \nu \rangle) w^\alpha \varphi^2 + w^\alpha \varphi \langle \nabla^M \varphi, \nu \rangle.$$

$$\begin{aligned} \frac{d^2}{dt^2} \Big|_{t=0} A(E_t) &= \int_\Gamma (-\Delta_\Gamma \varphi - \text{Ric}(\nu, \nu)\varphi - |\text{II}_\Gamma|^2 \varphi - \theta w^{-2} \langle \nabla^M w, \nu \rangle^2 \varphi) w^\theta \varphi \\ &\quad + [\theta w^{-1} (\Delta w - \Delta_\Gamma w - H \langle \nabla^M w, \nu \rangle) \varphi - \theta w^{-1} \langle \nabla_\Gamma w, \nabla_\Gamma \varphi \rangle] w^\theta \varphi \\ &\quad + (\theta w^{\alpha-1} \langle \nabla^M w, \nu \rangle \varphi + w^\alpha \varphi \langle \nabla^M \varphi, \nu \rangle + H w^\alpha \varphi^2) w^{\theta-\alpha} (H + \theta w^{-1} \langle \nabla^M w, \nu \rangle) \\ &\quad + \int_{\partial\Gamma} w^\theta \varphi \langle \nabla^M \varphi, \nu_{\partial\Gamma} \rangle - A_{\partial\Gamma}(\nu, \nu) \varphi^2 w^\theta \end{aligned}$$

4.1.4. *Differential inequality in the barrier sense.* Fix a $v_0 \in (0, V_0)$. Let E be a weighted isoperimetric surface with free boundary in M for the problem $I(v_0)$. From now on, we fix $\varphi = w^{-\theta}$. We notice that $V(t) := V(E_t)$ is a smooth function. By the first variation information, we have

$$V'(0) = \frac{d}{dt}\Big|_{t=0} V(E_t) = \int_{\partial E} w^{\alpha-\theta} > 0,$$

hence $V(t)$ is a strictly monotone in t in a neighborhood of v_0 . By the inverse function theorem, there is some small $\sigma > 0$ and a smooth function

$$t : (v_0 - \sigma, v_0 + \sigma) \longrightarrow \mathbb{R},$$

that is the inverse of $V(t)$.

Let $u : (v_0 - \sigma, v_0 + \sigma) \rightarrow \mathbb{R}$ be defined by $u(v) = A(t(v))$. Note that $u(v_0) = A(E_0) = I(v_0)$. Moreover, since $V(E_{t(v)}) = v$, we have $A(v) \geq I(v)$ for all $v \in (v_0 - \sigma, v_0 + \sigma)$. Let primes denote derivatives with respect to v and dots denote derivatives with respect to t .

Lemma 18. *Let $(M^n, \partial M)$ be a complete, connected compact manifold with weakly boundary. Assume $w \in C^\infty(M)$ satisfies $\inf(w) = 1$, $\langle \nabla^M w, \eta \rangle = 0$, and*

$$(5) \quad \theta \Delta w \leq w \operatorname{Ric} - (n-1)\lambda w.$$

Suppose for fixed $v_0 \in (0, V_0)$, there exists a bounded set E with finite perimeter, such that $\int_E w^\alpha = v_0$ and $\int_{\partial^ E} = I(v_0)$. Then I satisfies*

$$I'' I \leq -\frac{(I')^2}{n-1} - (n-1)\lambda$$

in the viscosity sense at v_0 .

Proof. We first recall the formula for the first and second derivatives of an inverse function:

$$t'(v) = \frac{1}{\dot{v}}(t(v)) \quad \text{and} \quad t''(v) = -\frac{\ddot{v}(t(v))}{\dot{v}(t(v))^3},$$

then we have

$$t'(v_0) = \left(\int_{\partial E} w^{\alpha-\theta} \right)^{-1}$$

and

$$t''(v_0) = -\left(\int_{\partial E} w^{\alpha-\theta} \right)^{-3} \int_{\partial E} (H + \alpha w^{-1} \langle \nabla^M w, \nu \rangle w^\alpha \varphi^2 + w^\alpha \varphi \langle \nabla^M \varphi, \nu \rangle)$$

We note that

$$\frac{d}{dv} A(t(v)) = \dot{A}(t(v)) t'(v)$$

and

$$\frac{d^2}{dv^2} A(t(v)) = \ddot{A}(t(v)) (t'(v))^2 + \dot{A}(t(v)) t''(v).$$

Since E is a volume-constrained minimizer, we can conclude that $\frac{d}{dt}|_{t=0}A(E_t) = 0$ whenever $\frac{d}{dt}|_{t=0}V(E_t) = 0$. Therefore $A'(v_0) = w^{\theta-\alpha}(H + \theta w^{-1} \langle \nabla^M w, \nu \rangle)$ is constant. By Proposition 16 and 17 with $\varphi = w^{-\theta}$, we have

$$\begin{aligned} \frac{d^2}{dt^2}|_{t=0}A(E_t) &= \int_{\Gamma} (-\Delta_{\Gamma}(w^{-\theta}) - \text{Ric}(\nu, \nu)w^{-\theta} - |\Pi_{\Gamma}|^2w^{-\theta} - \theta w^{-2-\theta} \langle \nabla^M w, \nu \rangle^2 \\ &\quad + \theta w^{-\theta-1}(\Delta w - \Delta_{\Gamma}w - H \langle \nabla^M w, \nu \rangle) - \theta w^{-1} \langle \nabla_{\Gamma}w, \nabla_{\Gamma}(w^{-\theta}) \rangle \\ &\quad + (Hw^{\alpha-2\theta})w^{\theta-\alpha}(H + \theta w^{-1} \langle \nabla^M w, \nu \rangle) \\ &\quad + \int_{\partial\Gamma} \langle \nabla^M(w^{-\theta}), \nu_{\partial\Gamma} \rangle - A_{\partial\Gamma}(\nu, \nu)w^{-\theta} \end{aligned}$$

Using the weakly convexity of ∂M , integrating by parts and rearranging, we obtain that

$$\begin{aligned} \frac{d^2}{dt^2}|_{t=0}A(E_t) &\leq \int_{\Gamma} -\text{Ric}(\nu, \nu)w^{-\theta} - |\Pi_{\Gamma}|^2w^{-\theta} + \theta w^{-\theta-1}\Delta w - \theta w^{-2-\theta} \langle \nabla^M w, \nu \rangle^2 \\ &\quad - \theta Hw^{-\theta-1} \langle \nabla^M w, \nu \rangle + Hw^{-\theta}(H + \theta w^{-1} \langle \nabla^M w, \nu \rangle) \end{aligned}$$

According to inequality (5) and set $X = w^{\alpha-\theta}A'(v_0)$, $Y = w^{-1} \langle \nabla^M w, \nu \rangle$, thus $H = X - \theta Y$, and use the trace inequality $|\Pi_{\Gamma}|^2 \geq H^2/(n-1)$, we have

$$\begin{aligned} \frac{d^2}{dt^2}|_{t=0}A(E_t) &\leq \int_{\Gamma} -(n-1)\lambda w^{-\theta} + w^{-\theta} \left(-\frac{X^2}{n-1} + \frac{2\theta XY}{n-1} - \frac{\theta^2 Y^2}{n-1} - \theta Y^2 \right. \\ &\quad \left. - \theta XY + \theta^2 Y^2 + X^2 - \theta XY \right). \end{aligned}$$

Therefore, we obtain

$$\begin{aligned} \left(\int_{\Gamma} w^{\alpha-\theta} \right)^2 \cdot A''(v_0) &= (\ddot{A}(0)(t'(v_0))^2 + \dot{A}(0)t''(v_0)) \left(\int_{\Gamma} w^{\alpha-\theta} \right)^2 \\ &\leq \int_{\Gamma} -(n-1)\lambda w^{-\theta} + w^{-\theta} \left(-\frac{X^2}{n-1} + \frac{2\theta XY}{n-1} - \frac{\theta^2 Y^2}{n-1} - \theta Y^2 \right. \\ &\quad \left. - \theta XY + \theta^2 Y^2 + X^2 - \theta XY \right) \\ &\quad - \left(\int_{\Gamma} w^{\alpha-\theta} \right)^{-1} \left(\int_{\Gamma} w^{\alpha-\theta} A'(v_0) \right) \cdot \left(\int_{\Gamma} (X + \alpha Y - 2\theta Y) w^{\alpha-2\theta} \right) \\ &= \int_{\Gamma} -(n-1)\lambda w^{-\theta} + w^{-\theta} \left(-\frac{X^2}{n-1} + \frac{2\theta XY}{n-1} - \frac{\theta^2 Y^2}{n-1} - \theta Y^2 \right. \\ &\quad \left. - \theta XY + \theta^2 Y^2 + X^2 - \theta XY - X^2 - \alpha XY + 2\theta XY \right) \\ &= \int_{\Gamma} w^{-\theta} \left[-\frac{X^2}{n-1} + \left(\frac{2\theta}{n-1} - \alpha \right) XY + \left(\frac{n-2}{n-1} \theta^2 - \theta \right) Y^2 \right] - (n-1)\lambda w^{\theta} \\ &\leq \int_{\Gamma} -\frac{X^2}{n-1} w^{-\theta} - (n-1)\lambda w^{-\theta} \\ &= \int_{\Gamma} -\frac{A'(v_0)}{n-1} w^{2\alpha-3\theta} - (n-1)\lambda w^{-\theta}, \end{aligned}$$

where we used the fact that $\alpha = \frac{2\theta}{n-1}$, and $0 \leq \theta \leq \frac{n-1}{n-2}$. On the one hand, since $\alpha \leq \theta$, $w \geq 1$. Hence we have

$$\left(\int_{\Gamma} w^{\alpha-\theta}\right)^2 \cdot A''(v_0) \leq -\left(\frac{A'^2(v_0)}{n-1} + (n-1)\lambda\right) \int_{\Gamma} w^{2\alpha-3\theta}.$$

On the other hand, because $A(v_0) \int_{\Gamma} w^{\theta}$, we can conclude that

$$\left(\int_{\Gamma} w^{\alpha-\theta}\right)^2 \leq A(v_0) \int_{\Gamma} w^{2\alpha-3\theta}.$$

Putting them together, we have

$$(6) \quad A'(v_0)A''(v_0) \leq -\frac{A'^2(v_0)}{n-1} - (n-1)\lambda.$$

Therefore $A(v)$ is an upper barrier of $I(v)$ which satisfies the inequality (6). \square

We consider a power of I and A to simplify the corresponding differential inequality. Let $\mathcal{F}(v) = I(v)^{\frac{n}{n-1}}$, combined with above lemma 18, we have the following result.

Proposition 19. *For any $v_0 \in (0, V)$, there is a smooth function $U : (v_0 - \sigma, v_0 + \sigma) \rightarrow \mathbb{R}$ satisfying*

- $U(v_0) = \mathcal{F}(v_0)$,
- $U(v) \geq \mathcal{F}(v)$ for all $v \in (v_0 - \sigma, v_0 + \sigma)$,
- $U''(v_0) \leq -\lambda n U''(v_0)^{\frac{n}{n-1}}$.

Proof. We take $U(v) = A(v)^{\frac{n}{n-1}}$ as in 18, and compute that

$$U'(v) = \frac{n}{n-1} A(v)^{\frac{1}{n-1}} A'(v)$$

and

$$\begin{aligned} U''(v) &= \frac{n}{(n-1)(n-1)} A^{\frac{2-n}{n-1}}(v) A'^2(v) + \frac{n}{n-1} A^{\frac{1}{n-1}}(v) A''(v) \\ &= \frac{n}{n-1} A^{\frac{2-n}{n-1}}(v) (A'(v)A''(v) - (n-1)\lambda) + \frac{n}{n-1} A^{\frac{1}{n-1}}(v) A''(v) \\ &= -\lambda n U^{\frac{2-n}{n}}. \end{aligned}$$

\square

4.1.5. *Volume bound.* In this section, we will estimate the volume of manifold in the spectral sense. First, We begin by establishing an asymptotic volume expansion estimate for a small geodesic ball centered at a boundary point.

Lemma 20. *Suppose that M^n is a complete, connected compact manifold with weakly convex boundary, $w \in C^\infty(M)$ is positive. Assume $x \in \partial M$ satisfies $w(x) = \inf(w) = 1$. Then, if I is defined as in (4), we have*

$$(7) \quad \limsup_{v \rightarrow 0} v^{-\frac{n-1}{n}} I(v) \leq n \text{Vol}(\mathbb{B}_+^n)^{\frac{1}{n}},$$

where \mathbb{B}_+^n is the unit half ball in \mathbb{R}^n .

Proof. For a small r_0 , the functions $V(r) = \int_{B(x,r)} w^\alpha$ and $A(r) = \int_{\partial B(x,r)} w^\theta$ are smooth and increasing in $(0, r_0)$, where the geodesic ball of radius r is centered at $x \in \partial M$. We have the asymptotics

$$V(r) = \text{Vol}(\mathbb{B}_+^n) r^n + O(r^{n+1}),$$

and

$$A(r) = n \text{Vol}(\mathbb{B}_+^n) r^{n-1} + O(r^n),$$

hence the function $A \circ V^{-1}(v) = n \text{Vol}(\mathbb{B}_+^n)^{\frac{1}{n}} v^{\frac{n-1}{n}} + o(v^{\frac{n-1}{n}})$, and $I(v) \leq A \circ V^{-1}(v)$. \square

Theorem 21. *Let $V \in (0, \infty)$, and let $I : [0, V) \rightarrow \mathbb{R}$ be a continuous function such that $I(0) = 0$, and $I(v) > 0$ for every $v \in (0, V)$. Assume that for some $\lambda > 0$ we have*

$$I'' I \leq -\frac{(I')^2}{n-1} - (n-1)\lambda,$$

and

$$\limsup_{v \rightarrow 0^+} v^{-\frac{n-1}{n}} I(v) \leq n \text{Vol}(\mathbb{B}_+^n)^{\frac{1}{n}}.$$

Then we have $V \leq \lambda^{-\frac{n}{2}} \text{Vol}(\mathbb{S}_+^n)$, where \mathbb{S}_+^n denotes the unit half sphere in \mathbb{R}^{n+1} .

Proof. According to Lemma 18 and Proposition 19, we know that if we set $\mathcal{F}(v) = I(v)^{\frac{n}{n-1}}$, then $\mathcal{F}(v)$ satisfies a differential inequality in the barrier sense:

$$\mathcal{F}''(v) \leq -\lambda n \mathcal{F}^{\frac{n-2}{n}}(v).$$

We first study solutions to the ODE

$$(8) \quad f''(v) = -\lambda n f^{\frac{2-n}{n}}.$$

Since $-\lambda n f^{\frac{2-n}{n}}$ is increasing in f , it follows from a standard ODE comparison theorem that there no solution to (8) can touch $\mathcal{F}(v)$ from below unless they are equal. If f satisfies (8), then we obtain

$$(f'^2 + \lambda n^2 f^{\frac{2}{n}})' = 0.$$

For $z > 0$, let $f_z(v)$ be the solution to 8 satisfying $f'_z(0) = 0$ and $f_z(0) = z^{\frac{n}{n-1}}$. Then we have

$$(9) \quad f'_z(v)^2 = \lambda n^2 z^{\frac{2}{n-1}} - \lambda n^2 f_z(v)^{\frac{2}{n}}.$$

Let $\beta(z) > 0$ be the maximal real number so that $f_z(v) > 0$ on $(-\beta(z), \beta(z))$. We see by (9) that

$$f'_z(v) = -\sqrt{\lambda n^2 z^{\frac{2}{n-1}} - \lambda n^2 f_z(v)^{\frac{2}{n}}} \text{ for } v > 0.$$

Therefore, we can integrate the derivative of the inverse of f_z to find

$$\begin{aligned}
\beta(z) &= - \int_0^{z^{\frac{n}{n-1}}} (f_z^{-1})'(x) dx \\
&= \int_0^{z^{\frac{n}{n-1}}} \frac{1}{\sqrt{\lambda}} \frac{1}{n} (z^{\frac{2}{n-1}} - x^{\frac{2}{n}})^{-\frac{1}{2}} dx \\
&= \int_0^{\frac{\pi}{2}} \frac{1}{\sqrt{\lambda}} z \sin^{n-1} r dr \\
&= \frac{z}{2\sqrt{\lambda}} \frac{\sqrt{\pi} \Gamma(\frac{n}{2})}{\Gamma(\frac{n+1}{2})} \\
&= \frac{z}{2\sqrt{\lambda}} \frac{\text{Vol}(\mathbb{S}^n)}{\text{Vol}(\mathbb{S}^{n-1})},
\end{aligned}$$

where we used the substitution $x = z^{\frac{n}{n-1}} \sin^{n-1}(r)$.

Assume for the sake of contradiction that $V_0 = \int_M w^\alpha > \lambda^{-\frac{n}{2}} \text{Vol}(\mathbb{S}_+^n)$.

Claim. *There is a $\delta > 0$, so that for $z = \xi + \delta$, where $\xi = \lambda^{-\frac{n-1}{2}} \text{Vol}(\mathbb{S}_+^{n-1})$ we have*

$$(10) \quad \mathcal{F}(v) \geq f_z(v - \beta(z))$$

for all $v \in (0, \lambda^{-\frac{n}{2}} \text{Vol}(\mathbb{S}_+^n))$.

Proof. Let $\delta > 0$ and $\epsilon > 0$ be sufficiently small such that $2\beta(z) + \epsilon z < V$ for $z \in (0, \xi + \delta)$, which is possible since $V > \lambda^{-\frac{n}{2}} \text{Vol}(\mathbb{B}_+^n) = 2\beta(\xi)$. Consider the graph of

$$g_z(v) = f_z(v - \beta(z) - \epsilon z)$$

for $v \in [\epsilon z, 2\beta(z) + \epsilon z]$. Note that

$$g_z(\epsilon z) = g_z(2\beta(z) + \epsilon z) = 0 < \min\{\mathcal{F}(\epsilon z), \mathcal{F}(2\beta(z) + \epsilon z)\}.$$

Moreover, g_z converges uniformly to zero as $z \rightarrow 0$. Hence, if $g_{z^*}(v^*) > \mathcal{F}(v^*)$ for some z^* and v^* , then there must be some $0 < z \leq z^*$ so that g_z touches \mathcal{F} from below, which contradicts Proposition 19. Therefore, we have $\mathcal{F} \geq g_z$ for every $z \in (0, \xi + \delta)$. We take z to $\xi + \delta$, and conclude the claim since ϵ can be arbitrary small. \square

We study the asymptotic behavior of \mathcal{F} and $f_{\xi+\delta}(v - (\xi + \delta))$ as $v \rightarrow 0$. According to (9), since $f_{\xi+\delta}(-\beta(\xi + \delta)) = 0$, we have

$$f'_{\xi+\delta}(-\beta(\xi + \delta)) = \sqrt{\lambda} n (\lambda^{-\frac{n-1}{2}} \text{Vol}(\mathbb{S}_+^{n-1}))^{\frac{1}{n-1}},$$

Therefore, we have

$$(11) \quad f_{\xi+\delta}(v - \beta(\xi + \delta)) = \sqrt{\lambda} n (\lambda^{-\frac{n-1}{2}} \text{Vol}(\mathbb{S}_+^{n-1}) + \delta)^{\frac{1}{n-1}} v + o(v), \quad \text{as } v \rightarrow 0.$$

On the other hand, by Lemma 20, if we take x_0 such that $w(x_0) = \min w = 1$, we have the asymptotics

$$V(r) = \text{Vol}(\mathbb{B}_+^n) r^n + O(r^{n+1}),$$

and

$$A(r) = n \text{Vol}(\mathbb{B}_+^n) r^{n-1} + O(r^n),$$

hence $A \circ V^{-1}(v) = n \operatorname{Vol}(\mathbb{B}_+^n)^{\frac{1}{n}} v^{\frac{n-1}{n}} + o(v^{\frac{n-1}{n}})$, and we can obtain

$$(12) \quad \mathcal{F}(v) \leq n^{\frac{n}{n-1}} \operatorname{Vol}(\mathbb{B}_+^n)^{\frac{1}{n-1}} v + o(v)$$

$$(13) \quad = n \operatorname{Vol}(\mathbb{S}_+^{n-1}) v + o(v)$$

However, combing (10), (11) with (12), we conclude that $\sqrt{\lambda} n (\lambda^{-\frac{n-1}{2}} \operatorname{Vol}(\mathbb{S}_+^{n-1}) + \delta)^{\frac{1}{n-1}} \leq n \operatorname{Vol}(\mathbb{S}_+^{n-1})$, which is a contradiction. Therefore, since we normalized so that $\min w = 1$, we have

$$\operatorname{Vol}(M) \leq \int_M w^\alpha = V_0 \leq \lambda^{-\frac{n}{2}} \operatorname{Vol}(\mathbb{S}_+^n).$$

□

4.2. **for case $n \geq 8$.**

Remark 22. *Because the minimizer E may have singularity for $n \geq 8$ as mentioned in Theorem 12, the volume estimate for $n \geq 8$ can also be obtained by modifying above argument, we refer readers to [1, Appendix A] for details.*

4.3. **Rigidity result.** Finally, because M has weakly convex boundary, we can also establish a rigidity result following the method in [1]. We can assume that $\lambda = 1$ after rescaling.

Theorem 23. *Let M^n , $n \geq 3$, be a compact connected manifold with weakly convex boundary ∂M , and let $0 \leq \theta \leq \frac{n-1}{n-2}$. Assume that there exist a positive function $w \in C^\infty(M)$ satisfies:*

$$\begin{cases} -\theta \Delta w + \operatorname{Ric} w \geq (n-1)w, \\ \frac{\partial w}{\partial \eta} = 0. \end{cases}$$

If $\operatorname{Vol}(M) = \operatorname{Vol}(\mathbb{S}_+^n)$, then w must be constant, and M is isometric to the unit round hemisphere \mathbb{S}_+^n .

REFERENCES

- [1] G. Antonelli and K. Xu, New spectral Bishop-Gromov and Bonnet-Myers theorems and applications to isoperimetry, arXiv:2405.08918v2 [math.DG].
- [2] S. Y. Cheng, Eigenvalue comparison theorems and its geometric applications, Math. Z. 143(3), 289-297(1975). MR 0378001.
- [3] O. Chodosh and C. Li, Generalized soap bubbles and the topology of manifolds with positive scalar curvature, Ann. of Math. (2)199(2024), no. 2, 707-740. MR 4713021.
- [4] O. Chodosh, C. Li, P. Minter and D. Stryker, Stable minimal hypersurfaces in \mathbb{R}^5 , arXiv:2401.01492v1[math.DG].
- [5] O. Chodosh, C. Li, P. Minter and D. Stryker, Stable minimal hypersurfaces in \mathbb{R}^5 , arXiv:2401.01492v1[math.DG].
- [6] F. Morgan, Regularity of isoperimetric hypersurfaces in Riemannian manifolds, Trans. Amer. Math. Soc. 355(12), 5041-5052(2003).
- [7] M. Gromov, Four lectures on scalar curvature, arXiv: 1908.10612[math.DG].
- [8] F. Hang and X. Wang, Rigidity theorems for compact manifolds with boundary and positive Ricci curvature, J. Geom. Anal. 19(2009), 628-642.
- [9] J. Li, The structure of weakly stable CMC hypersurfaces with free boundary, arXiv:2502.11739[math.DG].
- [10] L. Mazet, Stable minimal hypersurfaces in \mathbb{R}^6 , arXiv:2405.14676v1[math.DG].

- [11] M. Ritoré, Isoperimetric inequalities in Riemannian manifolds, Vol(348), Progress in Mathematics, ISBN: 978-3-031-37900-0.
- [12] P. Sternberg and K. Zumbrun, On the connectivity of boundaries of sets minimizing perimeter subject to a volume constraint. *Comm. Anal. Geom.* 7(1), 199-220(1999).
- [13] Y. Wu, Free boundary stable minimal hypersurfaces in positively curved 4-manifolds, arXiv:2308.08103v1[math.DG].
- [14] J. Zhu, Width estimate and doubly warped product, *Trans. Amer. Math. Soc.* 374(2021), no. 2, 1497-1511. MR 4196400.

SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, 361005, XIAMEN, P.R. CHINA
Email address: lijiamath@stu.xmu.edu.cn