

Uncovering hidden resonances in non-Hermitian systems with scattering thresholds

Fridtjof Betz,¹ Felix Binkowski,¹ Jan David Fischbach,² Nick Feldman,^{3,4}

Carsten Rockstuhl,^{2,5} A. Femius Koenderink,⁴ and Sven Burger^{1,6}

¹*Zuse Institute Berlin, Takustraße 7, 14195 Berlin, Germany*

²*Institute of Nanotechnology, Karlsruhe Institute of Technology, Kaiserstrasse 12, 76131 Karlsruhe, Germany*

³*Advanced Research Center for Nanolithography, Science Park 106, 1098 XG Amsterdam, The Netherlands*

⁴*Department of Information in Matter and Center for Nanophotonics,*

AMOLF, Science Park 104, NL1098XG Amsterdam, The Netherlands

⁵*Institute of Theoretical Solid State Physics, Karlsruhe Institute of Technology, Kaiserstrasse 12, 76131 Karlsruhe, Germany*

⁶*JCMwave GmbH, Bolivarallee 22, 14050 Berlin, Germany*

The points where diffraction orders emerge or vanish in the propagating spectrum of periodic non-Hermitian systems are referred to as scattering thresholds. Close to these branch points, resonances from different Riemann sheets can tremendously impact the optical response. However, these resonances were so far elusive for two reasons. First, their contribution to the signal is partially obscured, and second, they are inaccessible for standard computational methods. Here, we explore the interplay of scattering thresholds with resonances and introduce a multi-valued rational approximation to access the hidden resonances. Our theoretical and numerical approach is used to analyze the resonances of a plasmonic line grating. This work elegantly explains the occurrence of pronounced spectral features at scattering thresholds applicable to many nanophotonic systems of contemporary and future interest.

Introduction.—In the early years of the twentieth century, Wood was intrigued by sudden changes in the intensity of light diffracted from a grating. He noticed that, under certain conditions, a small change in the wavelength could drop the intensity from maximum to minimum [1]. Wood described these features of the spectrum as the most interesting problem he had encountered so far and termed them anomalies, since they could not be explained by ordinary grating theory. A few years later, in a note on this phenomenon, Lord Rayleigh came to the conclusion that anomalies were to be expected at the scattering thresholds, that is, whenever a higher diffraction order appears in the propagating spectrum [2]. In 1941, Fano attributed another type of Wood’s anomalies to surface plasmons, and thus established a connection to resonances [3]. Eventually, in 1965, Hessel and Oliner clearly distinguished two types of Wood’s anomalies: square-root-type singularities at the scattering thresholds and those that can be attributed to resonances [4]. They commented that the derivative of the square root results in an infinite slope, which they anticipated would boost resonance effects. Although each of these aspects is well-established in nanophotonics, it is the combination of resonances and diffraction that currently attracts interest and has led to the development of metasurfaces with Mie resonances, plasmon resonances, and optical bound states in the continuum (BICs) [5–11]. In fact, it has been proposed to exploit BICs close to a scattering threshold to maximize the sensitivity of refractive index sensing [12].

Recently, a mathematical model for the entries of the scattering matrix in the vicinity of scattering thresholds has been presented [13]. Whereas the model accurately reproduces the square root behavior, it does not include resonances. In contrast, two established formalisms, known as quasinormal mode expansion [14–16]

and resonant state expansion [17], respectively, have been put forward to describe resonant behavior. However, their application to grating systems is known to be subtle. Indeed, for such formalisms, it has been pointed out that algorithms identify a plethora of additional resonances without a direct physical interpretation, but that account for the discontinuity of the derivative at the scattering threshold [18, 19]. These numerical resonances, as explicitly stated in [18], discretize an integral. With an arbitrary black-box solver for Maxwell’s equation at hand, distinguishing physical resonances from those representing the branch cut is a non-trivial task. An alternative approach that explicitly evaluates resonances in the vicinity of scattering thresholds describes the optical response as a function of a wave vector component instead of frequency [20, 21]. The corresponding transformed space does not contain branch points.

In this work, we use such a transformed space to access the relevant resonances on different Riemann sheets of periodic non-Hermitian systems and reconstruct the optical response across several scattering thresholds with a multi-valued rational approximation. The results enable an instructive interpretation of the interaction between resonances and square-root-type singularities. Using the example of a plasmonic line grating, upon changing a design parameter, we observe resonances vanish behind a branch point, and others appear in the spectrum that had been hidden previously. In this way, a distinctive feature seemingly unrelated to resonances is accurately described with inherent properties of the system.

Multi-valued rational approximation.—The AAA algorithm [22] has proven to be an efficient method to reconstruct optical response functions with rational functions $r(k)$, $k \in \mathbb{C}$ [23]. The schematic example provided in Fig. 1(a) shows the specular reflection R_0 of an oblique incident plane wave off the metallic line grating sketched




FIG. 1. Branch points. (a) The specular reflection spectrum $R_0(k)$ shows a distinct feature where its first derivative is discontinuous, indicating a branch point. (b) In the example at hand, the branch point originates from the appearance of the first-order reflection $R_{-1}(k)$. (c) The square root function, which has multiple values, can have its branch point adjusted to align with the anticipated branch point of the reflection spectrum. (d) To encompass all possible values of the complex square root, it is necessary to consider the entire Riemann surface, which spans 4π in polar coordinates.

in Fig. 1(b). The spectrum is shown as a function of the incident light wave number k . We expect sudden drops in intensity due to Wood's anomalies, that is, resonance effects and square-root-type singularities, as well as their interactions. The spectrum in Fig. 1(a) shows two pronounced dips. The broad minimum at lower wave numbers corresponds to a plasmon resonance of the system. At higher wave numbers, there is a sharp feature with a discontinuous derivative. This feature at the point labeled $k_{b,-1}$ corresponds to the appearance of the first-order reflection R_{-1} . A rational function $r(k)$ can approximate the plasmon resonance with a single pole, but not the second feature. As an alternative to the wave number k , the z -component of the wave vector can be used to parameterize the response [20]. In this transformed representation, the specular reflection can be fully approximated with a low-order rational function. Subsequently, $r(k)$ is obtained by a coordinate transformation.

Given a structure periodic in x and y , the z -component k_z of the diffracted light wave vector is constrained by the condition

$$k_z = \sqrt{k^2 - |\mathbf{k}_{\parallel} + \mathbf{G}|^2}, \quad (1)$$

where \mathbf{G} is a reciprocal lattice vector and \mathbf{k}_{\parallel} the components of the wave vector parallel to the periodic structure. Each diffraction order corresponds to a specific \mathbf{G} and will contribute to the propagating spectrum if the term inside the square root is positive. Imaginary k_z contribute to the evanescent spectrum. We are interested in the branch points $k_{b,n}$ with $n \in \mathcal{N} \subset \mathbb{Z}$ that mark the emergence of new diffraction orders in the propagating spectrum. They are solutions of $k_{b,n} - |\mathbf{k}_{\parallel} + \mathbf{G}| = 0$. Using $N = |\mathcal{N}|$ of these solutions, we define the coordinate transform

$$\tilde{k} = \frac{1}{N} \sum_{n \in \mathcal{N}} \sqrt{k^2 - k_{b,n}^2}. \quad (2)$$

This transformation asymptotically approximates k and shows locally, around its branch points $k_{b,n}$, the square root behavior of k_z .

The complex variable \tilde{k} is a multi-valued function of k . Figure 1(c) shows the graph of one of its summands for real k , which has a positive and a negative branch due to the square root. The complete two-sheeted Riemann surface is shown in Fig. 1(d). It is easy to see that to return to the starting point, the branch point $k_{b,n}$ must be encircled twice. It is common to define a single-valued function through the definition of a branch cut. Approaching this cut from different sides results in different function values. A rational function of \tilde{k} can have poles on any Riemann sheet, and therefore, introducing a branch cut would hide information. Unlike [24], where resonances are evaluated directly on a four-sheeted Riemann surface, we solve the resonance problem here in a transformed space using the AAA algorithm [22]. The algorithm takes sampling points and greedily adds terms to a rational approximation of the optical response function. The resulting poles can be identified with the resonances of the physical system [23] and are mapped back to the $2N$ Riemann sheets introduced with \tilde{k} . This back transformation is crucial to interpret their impact on the physical response near branch points, and therefore we use the term *multi-valued approach*.

Application.—We investigate the interaction between resonances and scattering thresholds using the example of a line grating sketched in the inset of Fig. 2(a) that supports pronounced grating anomalies [19]. A Drude-Lorentz model specifies the relative permittivity of gold $\varepsilon(\omega) = \varepsilon_0 - \varepsilon_0 \omega_p^2 (\omega^2 + i\omega\gamma)^{-1}$, with the vacuum permittivity ε_0 and the plasma frequency $\omega_p = 1.26 \times 10^{16} \text{ rad s}^{-1}$, the damping factor $\gamma = 0.0112 \omega_p$, and the angular frequency $\omega = ck$ that is related to the wave number k through the speed of light c . Working in the optical




FIG. 2. The specular reflection spectrum $R_0(k)$ across 3 branch points located at $0.5 k_{b,1}$, $k_{b,1}$ and $1.5 k_{b,1}$, respectively, with $k_{b,1} = 1.074 \times 10^7 \text{ m}^{-1}$. The gold grating, sketched in the circular inset, is illuminated by a TM-polarized plane wave at an incidence angle of $\Theta = 30^\circ$. (a) The approximated spectrum together with reference points and a decomposition into six modal contributions and a residual contribution. (b) Resonances of the system: six resonances (blue crosses) are found at identical positions with and without the coordinate transformation \tilde{k} . In the single-valued approach (top), additional resonances cluster at wave numbers close to the branch points. Conversely, the multi-valued approach (bottom) reveals three additional resonances. The labels of the resonances within the circle refer to Fig. 3.

regime, we can equate the permeability $\mu(\omega) = \mu_0$ with that in the vacuum. The grating period p , which defines the one-dimensional lattice vector $G = 2\pi n/p$ with $n \in \mathbb{Z}$, is 600 nm, and the depth of the rectangular grooves is 350 nm. The width w of 51 nm maximizes the interaction between the square-root-type singularity and resonances at the angle of incidence $\Theta = 30^\circ$. For the definition of the angle, we refer to Fig. 1(b).

Samples of the optical response at real-valued frequencies are the only prerequisites for the proposed method. Here, we rely on the finite element method (FEM) and its implementation in the JCMsuite software package [25] to solve the second-order Maxwell equation

$$\nabla \times \mu^{-1} \nabla \times \mathbf{E} - \omega^2 \epsilon \mathbf{E} = i\omega \mathbf{J}, \quad (3)$$

with the electric current density \mathbf{J} and the electric field strength \mathbf{E} . With plane wave illumination, the upward directed Fourier spectrum of \mathbf{E} yields the specular reflection R_0 , which is the absolute value squared of the zero-order coefficient.




FIG. 3. Resonances passing a branch point. (a) The three resonances closest to the 50 support points (black dots) are displayed as a function of the groove width w , which is varied in steps of 1 nm from $w_1 = 42 \text{ nm}$ to $w_5 = 60 \text{ nm}$. The resonances inside the green boxes are not directly connected to the real axis above them, i.e., they are hidden resonances. (b) Specular reflection spectra corresponding to the labeled subset of resonances. (c) The magnetic field patterns corresponding to the three relevant resonances at selected widths.

The condition $k_{b,n} - |k_{b,n} \sin \Theta + 2\pi n/p| = 0$ for the scattering thresholds $k_{b,n}$ follows from Eq. (1). Figure 2(a) shows the specular reflection $R_0(k) - 1$ in a range that spans the three scattering thresholds $k_{b,-1}$, $k_{b,1}$, and $k_{b,-2}$, as obtained by a usual driven diffraction calculation. In addition, the modal contributions of six dominant resonances in the spectrum are displayed. This modal expansion of R_0 , which is quadratic in the electric field, is based on the residues of the Fourier coefficient and evaluations of its rational approximation at the complex conjugated resonance frequencies following [26]. Subtracting the sum of the six modal contributions from the full spectrum gives the residual contribution that exhibits the square-root-type Wood's anomalies.

The lower part of Fig. 2(b) shows the resonances k_n of a rational approximation without coordinate transformation. In this case, the algorithm clusters many additional resonances along characteristic lines close to the scattering thresholds. The contributions of these resonances approximate integrals along branch cuts, as discussed in [18] and give rise to the discontinuities in the derivative. We will show that the feature at $k_{b,1}$ that particularly catches the eye is the result of two hidden resonances on different Riemann sheets.




FIG. 4. Multi-valued specular reflection. (a) The contributions of the resonances b_3 and c_3 explain the shape of the specular reflection (black line) at the respective sides of the branch point. (b) The Riemann surface of the double-valued specular reflection. The positions of the resonances are projected to the complex plane in the bottom part of the graph. The colors refer to the argument of $k - k_{b,1}$ that ranges from -2π to 2π to uniquely define every point of the double-valued function.

The lower section of Fig. 2(b) shows resonances k_m derived from the poles of the rational approximation $\tilde{r}(\tilde{k})$ that results from an application of the AAA algorithm in the transformed space. Given the poles \tilde{k}_m of $\tilde{r}(\tilde{k})$ and the single-valued definition of the square root with a branch cut parallel to the negative imaginary axis, the resonances marked with green crosses do not solve the radical equation

$$\tilde{k}_m = \frac{1}{N} \sum_{n=1}^N \sqrt{k_m^2 - k_{b,n}^2} \quad (4)$$

that follows from Eq. (2). These hidden resonances do not appear on the sheet that contains the physical signal, but they do impact the response. It should be noted that the pole above the real axis enters the physical sheet at $k_{b,-1}$ with a negative imaginary part if a parameter is changed accordingly. For a better understanding of hidden resonances, we will study in the following how the two resonances near the branch point $k_{b,1}$ labeled b and c behave if the width of the grooves w is varied.

The trajectories of three resonances labeled a , b , and c when varying w are presented in Fig. 3(a). The hidden resonances that require both Riemann sheets to solve

Eq. (4) are highlighted with green boxes. A comparison with the spectra at selected widths shown in Fig. 3(b) reveals that when w is increased, the spectral feature linked to the resonance c approaches the branch point and is eventually replaced by the broader spectral feature of b , which had previously been hidden. Hence, instead of using a large number of numerical resonances to represent the branch cut, few resonances fully explain the physical response. In the current scenario, only two resonances are needed. A special case arises for the width w_3 , as both b_3 and c_3 are hidden resonances, and the signal is highly asymmetric, with one flank steeper than the other. Although the first derivative of the physical response is highly discontinuous, Fig. 3(c) shows that the field patterns of the resonances continuously vary as w changes. This observation emphasizes that Fig. 3(b) does not show the signal of a single pole crossing the branch point but rather two distinct resonances.

We conclude the investigation of resonances in the vicinity of branch points with a picture visualizing their effect on the optical response measured along the real axis. We have observed in Fig. 3 that the resonances b and c are responsible for the asymmetric shape of the spectral feature. This finding is confirmed in Fig. 4(a). At smaller wave numbers, the flank of the feature follows the resonance contribution of c even though the resonance itself is shifted to higher wave numbers, where the shape of the feature can be explained with the contribution of b . The complete Riemann surface of the specular reflection in Fig. 4(b), with colors visualizing the phase $\varphi = \arg(k - k_{b,1})$, explains why the peak at the point b mainly influences the spectrum at wave numbers larger than $k_{b,1}$, even though $\text{Re}(b) < k_{b,1}$. Since the angles of the physical signal are $-\pi$ and zero, the position b with $\varphi > \pi/2$ hardly affects the signal in $\varphi = -\pi$, where the resonance at c contributes dominantly.

Conclusion.—We demonstrated that multi-valued rational approximations accurately describe resonances near scattering thresholds. The resulting low-order model explains the characteristic sharp features with resonances on different Riemann sheets. We show how these hidden resonances can strongly enhance sharp features in the scattering response. The model accurately describes the discontinuity of the first derivative of the optical response function. Therefore, it is possible to reconstruct complex spectra with different types of singularity, as presented in Fig. 2(a), with a single rational approximation.

The proposed method is particularly easy to implement, as it only requires samples of the optical response at real-valued frequencies, from numerical or experimental data.

We envision a wide use of our method in a variety of fields, including optics and electromagnetism, as well as acoustics. Engineering of response functions by hybridization of square-root-type singularities and resonances is a very common motif in the design of plasmonic and dielectric structures for wavefront control, fluores-

cence control, sensing, and nonlinear optics. In the plasmonic domain this includes, for instance, the field of surface lattice resonances, i.e., high quality collective plasmon resonances in periodic metal nanoparticle lattices that have been developed for fluorescence control, surface enhanced Raman scattering and lasing [10, 27, 28]. In the field of dielectric metasurfaces and photonic crystals there is a strong interest in Fano resonances and quasi-bound states in the continuum [29, 30], in which Mie resonances [31], square-root-type singularities and guided mode resonances interplay. These structures are pursued for their high quality factors and strong local field confinement, with demonstrated applications in IR spectroscopy [32], nonlinear optics and high harmonic generation [33, 34], and nonlocal metasurfaces for wavefront control [35]. Our method provides an efficient and intuitive approach to attribute the response of these systems to just a few resonant contributions, including hidden resonances. An interesting prospect is to track these resonances to enable optimization schemes based on their positions [36], even when they are partially obscured by

the branch point.

Data availability.—Supplementary data tables, details on the convergence and source code for the numerical experiments of this work can be found in the open-access data publication [37].

Acknowledgments.—FB, FB, and SB acknowledge funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy - The Berlin Mathematics Research Center MATH+ (EXC-2046/1, project 390685689) and by the German Federal Ministry of Education and Research (BMBF Forschungscampus MODAL, project 05M20ZBM). JDF and CR acknowledge financial support by the Helmholtz Association in the framework of the innovation platform “Solar TAP”. The work of NF and AFK is part of the Dutch Research Council (NWO) and was performed at the research institutes AMOLF and ARCNL. ARCNL (Advanced Research Center for Nanolithography) is a public-private partnership between the University of Amsterdam, Vrije Universiteit Amsterdam, University of Groningen, NWO, and the semiconductor equipment manufacturer ASML.

-
- [1] R. W. Wood, *Lond. Edinb. Dubl. Phil. Mag.* **4**, 396 (1902).
- [2] L. Rayleigh, *Lond. Edinb. Dubl. Phil. Mag.* **14**, 60 (1907).
- [3] U. Fano, *J. Opt. Soc. Am.* **31**, 213 (1941).
- [4] A. Hessel and A. A. Oliner, *Appl. Opt.* **4**, 1275 (1965).
- [5] V. E. Babicheva and A. B. Evlyukhin, *Adv. Opt. Photon.* **16**, 539 (2024).
- [6] F. Ding, Y. Yang, R. A. Deshpande, and S. I. Bozhevolnyi, *Nanophotonics* **7**, 1129 (2018).
- [7] K. Koshelev, S. Lepeshov, M. Liu, A. Bogdanov, and Y. Kivshar, *Phys. Rev. Lett.* **121**, 193903 (2018).
- [8] F. J. García de Abajo, *Rev. Mod. Phys.* **79**, 1267 (2007).
- [9] F. J. García-Vidal, L. Martín-Moreno, T. W. Ebbesen, and L. Kuipers, *Rev. Mod. Phys.* **82**, 729 (2010).
- [10] V. G. Kravets, A. V. Kabashin, W. L. Barnes, and A. N. Grigorenko, *Chem. Rev.* **118**, 5912 (2018).
- [11] W. Wang, M. Ramezani, A. I. Väkeväinen, P. Törmä, J. G. Rivas, and T. W. Odom, *Materials Today* **21**, 303 (2018).
- [12] D. N. Maksimov, V. S. Gerasimov, A. A. Bogdanov, and S. P. Polyutov, *Phys. Rev. A* **105**, 033518 (2022).
- [13] C. C. Wojcik, H. Wang, M. Orenstein, and S. Fan, *Phys. Rev. Lett.* **127**, 277401 (2021).
- [14] P. Lalanne, W. Yan, K. Vynck, C. Sauvan, and J.-P. Hugonin, *Laser Photonics Rev.* **12**, 1700113 (2018).
- [15] S. Franke, S. Hughes, M. Kamandar Dezfouli, P. T. Kristensen, K. Busch, A. Knorr, and M. Richter, *Phys. Rev. Lett.* **122**, 213901 (2019).
- [16] A. Nicolet, G. Demézy, F. Zolla, C. Campos, J. E. Roman, and C. Geuzaine, *Eur. J. Mech. A Solids* **100**, 104809 (2023).
- [17] E. A. Muljarov, W. Langbein, and R. Zimmermann, *EPL* **92**, 50010 (2010).
- [18] M. B. Doost, W. Langbein, and E. A. Muljarov, *Phys. Rev. A* **87**, 043827 (2013).
- [19] A. Gras, W. Yan, and P. Lalanne, *Opt. Lett.* **44**, 3494 (2019).
- [20] A. Akimov, N. A. Gippius, and S. G. Tikhodeev, *JETP Lett.* **93**, 427 (2011).
- [21] L. J. Armitage, M. B. Doost, W. Langbein, and E. A. Muljarov, *Phys. Rev. A* **89**, 053832 (2014).
- [22] Y. Nakatsukasa, O. Sète, and L. N. Trefethen, *SIAM J. Sci. Comput.* **40**, A1494 (2018).
- [23] F. Betz, M. Hammerschmidt, L. Zschiedrich, S. Burger, and F. Binkowski, *Laser Photonics Rev.* **18**, 2400584 (2024).
- [24] R. Smith, S. Houde-Walter, and G. Forbes, *IEEE J. Quantum Electron.* **28**, 1520 (1992).
- [25] J. Pomplun, S. Burger, L. Zschiedrich, and F. Schmidt, *Phys. Status Solidi B* **244**, 3419 (2007).
- [26] F. Betz, F. Binkowski, M. Hammerschmidt, L. Zschiedrich, and S. Burger, *Phys. Status Solidi A* **220**, 2200892 (2023).
- [27] J. Guan, J.-E. Park, S. Deng, M. J. H. Tan, J. Hu, and T. W. Odom, *Chemical Reviews* **122**, 15177 (2022).
- [28] G. Lozano, D. J. Louwers, S. R. Rodríguez, S. Murai, O. T. Jansen, M. A. Verschuuren, and J. Gómez Rivas, *Light Sci. Appl.* **2**, e66 (2013).
- [29] M. F. Limonov, M. V. Rybin, A. N. Poddubny, and Y. S. Kivshar, *Nat. Photonics* **11**, 543 (2017).
- [30] K. Koshelev, S. Lepeshov, M. Liu, A. Bogdanov, and Y. Kivshar, *Phys. Rev. Lett.* **121**, 193903 (2018).
- [31] K. Koshelev and Y. Kivshar, *ACS Photonics* **8**, 102 (2021).
- [32] A. Tittl, A. Leitis, M. Liu, F. Yesilkoy, D. Choi, D. N. Neshev, Y. S. Kivshar, and H. Altug, *Science* **360**, 1105 (2018).
- [33] H. Liu, C. Guo, G. Vampa, J. L. Zhang, T. Sarmiento, M. Xiao, P. H. Bucksbaum, J. Vučković, S. Fan, and D. A. Reis, *Nat. Phys.* **14**, 1006 (2018).

- [34] K. Koshelev, Y. Tang, K. Li, D.-Y. Choi, G. Li, and Y. Kivshar, [ACS Photonics](#) **6**, 1639 (2019).
- [35] A. Overvig and A. Alù, [Adv. Photonics](#) **3**, 026002 (2021).
- [36] F. Binkowski, F. Betz, M. Hammerschmidt, P.-I. Schneider, L. Zschiedrich, and S. Burger, [Commun. Phys.](#) **5**, 202 (2022).
- [37] F. Betz, F. Binkowski, J. D. Fischbach, N. Feldmann, C. Rockstuhl, F. Koenderink, and S. Burger, [Source code and simulation results: Uncovering hidden resonances in non-Hermitian systems with scattering thresholds](#), Zenodo (2025), doi: 10.5281/zenodo.14651613.