

ON A CHARACTER CORRESPONDENCE ASSOCIATED TO \mathfrak{F} -PROJECTORS

MARÍA JOSÉ FELIPE, IRIS GILABERT, AND LUCIA SANUS

Dedicated to the memory of I. M. Isaacs.

ABSTRACT. We study the conditions under which the head characters of a solvable group, as defined by I. M. Isaacs, behave well with respect to restriction. We also determine the intersection of the kernels of all head characters of the group. Using G. Navarro's definition of \mathfrak{F}' -characters, we generalize these results for any saturated formation \mathfrak{F} containing the nilpotent groups.

1. INTRODUCTION

If G is a solvable group, I. M. Isaacs constructed in 2022 [10] a canonical subset of the complex irreducible characters of G associated to the linear characters of a Carter subgroup C of G . Recall that a Carter subgroup of G is a self-normalizing nilpotent subgroup of G . He called these the *head characters* of G , and proved that the number of those was $|C/C'|$, the number of linear characters of C . Long time before, E. C. Dade and D. Gajendragadkar had already constructed in [2, 4] another subset of $\text{Irr}(G)$, this time associated to the system normalizers of the solvable group G . This subset was described by Gajendragadkar from the better known *fully factorable characters*.

The head characters, on the other hand, remain mysterious, and no general properties are known of them. In our first main result, we prove the following key property.

Theorem A. Let G be a solvable group, C a Carter subgroup of G and N normal in G . If χ is a head character of G , then the restriction χ_{NC} contains a head character of NC . In particular, χ_N contains a C -invariant character of N .

Theorem A was conjectured by G. Navarro in private communications with I. M. Isaacs, and its proof depends on new deep properties of the character theory of fully ramified characters. (See Lemma 3.5.)

In Theorem B, we turn our attention to the kernels of the head characters and prove the following.

2010 Mathematics Subject Classification: primary 20C15, secondary 20D10.

Key words: head characters, \mathfrak{F} -projectors, saturated formations.

This research is partially supported by the Generalitat Valenciana (CIAICO/2021/163). The second author is supported by a grant (PAID-01-23 funded by the Universitat Politècnica de València and subsequently CIACIF/2023/389 funded the Generalitat Valenciana). The second and third authors are partially supported by the Spanish Ministerio de Ciencia e Innovación (PID2022-137612NB-I00 funded by MCIN/AEI/10.13039/501100011033 and 'ERDF A way of making Europe'). The first and second authors are partially supported by the Universitat Politècnica de València (PAID-06-23).

We would like to thank Gabriel Navarro for bringing this problem to our attention and for useful conversations on the subject.

Theorem B. Let G be a solvable group and C be a Carter subgroup of G . The intersection of the kernels of the head characters of G is the largest normal subgroup N of G such that $N \cap C$ is contained in C' .

Theorem B is the exact analog of an unpublished result of Navarro which we present here with his permission.

Theorem C (Navarro). Let p be a prime, let G be p -solvable and let P be a Sylow p -subgroup of G . The intersection of the kernels of the p' -characters of G is the largest normal subgroup N of G such that $\mathbf{N}_N(P)$ is contained in P' .

What is the relationship between the head characters and the p' -degree characters of a solvable group? If \mathfrak{F} is a saturated formation and G is a solvable group, Navarro defined in [14] a canonical subset $\text{Irr}_{\mathfrak{F}'}(G)$ of the irreducible characters of G of size $|\text{Irr}(N_G(H)/H')|$, where H is an \mathfrak{F} -projector of G . If \mathfrak{F} is the formation of nilpotent groups, then H is a Carter subgroup, and he proved that $\text{Irr}_{\mathfrak{F}'}(G)$ are the head characters of G ; in particular, reproving Isaacs result. If \mathfrak{F} is the formation of p -groups, then H is a Sylow p -subgroup and this reproved the McKay conjecture for solvable groups.

For solvable groups, Carter subgroups are the \mathfrak{F} -projectors associated to the formation of nilpotent groups, in the same way that system normalizers are the nilpotent case of \mathfrak{F} -normalizers. The latter are a conjugacy class of subgroups that was used by Dade and Gajendragadkar in a more general way to define their subset of $\text{Irr}(G)$ when \mathfrak{F} is a saturated formation containing the nilpotent groups.

In fact, our Theorems A and B are also far more general. In the same framework as Dade and Gajendragadkar (that is, if \mathfrak{F} is a saturated formation containing the nilpotent groups), we shall prove convenient versions of Theorems A and B for $\text{Irr}_{\mathfrak{F}'}(G)$ (see respectively Theorems 6.2 and 7.5) at the same time that we drastically simplify Isaacs' construction (see Theorem 5.4).

There are still several questions on the head characters and Carter subgroups that remain open: among them if the character table of G determines the set of its head characters. We are confident that our results might be helpful in solving these and other related questions.

2. SOME PROPERTIES ON PROJECTORS

A class of groups \mathfrak{F} is called a *formation* if it is closed under quotients and satisfies the following property: whenever $G/N, G/M \in \mathfrak{F}$ for some finite group G , it follows that $G/(M \cap N) \in \mathfrak{F}$. A formation \mathfrak{F} is said to be *saturated* if $G \in \mathfrak{F}$ if and only if $G/\Phi(G) \in \mathfrak{F}$, where $\Phi(G)$ denotes the Frattini subgroup of G . Examples of saturated formations are the classes of finite p -groups, finite nilpotent groups, and finite supersolvable groups. As basic references on this topic, see [3] and [15].

The notion of formations was first introduced by Gaschütz in [5] as a context where both definitions of Carter and Sylow subgroups could be unified under the more general concept of \mathfrak{F} -covering subgroups. Gaschütz later proved that, when \mathfrak{F} is saturated, these subgroups coincide with \mathfrak{F} -projectors, a preferable term in formation theory.

Lemma 2.1. *Let \mathfrak{F} be a saturated formation and G a solvable group. There exists exactly one conjugacy class of subgroups H of G such that, for every normal subgroup N of G , the subgroup HN/N satisfies the following properties:*

- (a) $HN/N \in \mathfrak{F}$;

- (b) if HN/N is a subgroup of the subgroup V/N of G/N with $V/N \in \mathfrak{F}$, then $HN/N = V/N$.

In other words, HN/N is \mathfrak{F} -maximal in G/N for all normal subgroups N of G . We say that these subgroups H are the \mathfrak{F} -projectors of G .

As previously mentioned, Carter subgroups can be found through the above definition by choosing the formation to be that of nilpotent groups, which we will note \mathfrak{N} from now on. Similarly, p -Sylow and π -Hall subgroups are the \mathfrak{F} -projectors for the saturated formations of p - and π -groups respectively. Listed below are some fundamental properties of \mathfrak{F} -projectors which we will need in our proofs.

Proposition 2.2. *Let \mathfrak{F} be a saturated formation and G be a solvable group. Suppose H is an \mathfrak{F} -projector of G . Then*

- (a) if U is a subgroup of G containing H , then H is also an \mathfrak{F} -projector of U .

Moreover, if N is a normal subgroup of G , then

- (b) HN/N is an \mathfrak{F} -projector of G/N ;
(c) if U/N is an \mathfrak{F} -projector of G/N , then $U = H^g N$ for some $g \in G$;
(d) $\mathbf{N}_G(NH) = N\mathbf{N}_G(H)$.

Finally, if $\mathfrak{N} \subseteq \mathfrak{F}$, then

- (e) if U is a subgroup of G containing H , then $\mathbf{N}_G(U) = U$. In particular, H is self-normalizing.

Proof. Assertions (a) to (c) are proved in Section 9.5 of [15]. If $g \in \mathbf{N}_G(HN)$, then both H and H^g are \mathfrak{F} -projectors of $HN = H^g N$. Using (c), there exists $n \in HN$ which can be assumed to be in N and such that $H^n = H^g$. Then $gn^{-1} \in \mathbf{N}_G(H)$. Thus $\mathbf{N}_G(HN) \leq N\mathbf{N}_G(H)$. Since the converse is trivial, this proves (d).

Now suppose $\mathfrak{N} \subseteq \mathfrak{F}$ and let H be an \mathfrak{F} -projector of a solvable group G . Let U be a subgroup of G containing H . By (a), H is also an \mathfrak{F} -projector of $\mathbf{N}_G(U)$, and thus by (b) we have that U/U is an \mathfrak{F} -projector of $\mathbf{N}_G(U)/U$. By definition of projectors, this means that $\{1\}$ is \mathfrak{F} -maximal in $\mathbf{N}_G(U)/U$. If $U \neq \mathbf{N}_G(U)$, then there exists some non trivial element $x \in \mathbf{N}_G(U)/U$ and the subgroup $\langle x \rangle$ contradicts the \mathfrak{F} -maximality of $\{1\}$. \square

The \mathfrak{F} -residual $G^{\mathfrak{F}}$ of G is its smallest normal subgroup such that the quotient $G/G^{\mathfrak{F}}$ lies in \mathfrak{F} . The following result shows a connection between an \mathfrak{F} -projector of a group and its \mathfrak{F} -residual, when the latter is abelian.

Theorem 2.3 (Theorem IV.5.18 of [3]). *Let G be a finite solvable group and $G^{\mathfrak{F}}$ its \mathfrak{F} -residual. If $G^{\mathfrak{F}}$ is abelian, then $G^{\mathfrak{F}}$ is complemented in G , and its complements are the \mathfrak{F} projectors of G .*

Note that, if \mathfrak{F} is a saturated formation, $K = G^{\mathfrak{F}}$ and $L \trianglelefteq G$ is such that K/L is abelian, then by the previous result it is not difficult to prove that $G = KH$ and $K \cap LH = L$, where H is an \mathfrak{F} -projector. It is thus natural to consider this group structure as a hypothesis for some of the theorems in further sections.

We finish this section with a result that can be found as Satz 2.1 of [6] or Theorem IV.5.4 of [3].

Lemma 2.4. *Let G be a finite solvable group and H an \mathfrak{F} -projector of G for a saturated formation \mathfrak{F} . If N, M are normal subgroups of G , then*

$$MN \cap H = (M \cap H)(N \cap H).$$

3. GOOD EXTENSIONS

In this section, we introduce results on extensions that, for our purposes, exhibit desirable behavior with respect to certain subgroups and formations. These extensions can be regarded, in some sense, as “*good extensions*”, and they will be of particular interest in Section 6 to address the question of when the restriction of an \mathfrak{F}' -character to a subgroup containing an \mathfrak{F} -projector of a solvable group G contains an \mathfrak{F}' -character of that subgroup.

Let us first recall two results on character correspondences.

Lemma 3.1 (Corollary 4.3 of [8]). *Let $N \trianglelefteq G$ and $K \subseteq G$ with $NK = G$ and $N \cap K = M$. Let $\varphi \in \text{Irr}(M)$ be invariant in K , and assume $\theta = \varphi^N$ is irreducible. Then induction defines a bijection $\text{Irr}(K \mid \varphi) \rightarrow \text{Irr}(G \mid \theta)$.*

Lemma 3.2 (Corollary 4.2 of [8]). *Let $N \trianglelefteq G$ and $K \subseteq G$ with $NK = G$ and $N \cap K = M$. Let $\theta \in \text{Irr}(N)$ be invariant in G , and assume $\varphi = \theta_M$ is irreducible. Then restriction defines a bijection $\text{Irr}(G \mid \theta) \rightarrow \text{Irr}(K \mid \varphi)$.*

Given an \mathfrak{F} -projector H of a solvable group G , we use the usual notation $\text{Irr}_H(N)$ for the H -invariant irreducible characters of an H -invariant subgroup N of G . The following theorem, which appears in [14] and is due to Navarro, is key to our objectives.

Theorem 3.3. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Suppose G is a finite solvable group. Assume that K/L is abelian, where K and L are normal in G . Let H be an \mathfrak{F} -projector of G , and assume that $KH = G$ and $K \cap HL = L$. We have the following.*

- (a) *If $\theta \in \text{Irr}_H(K)$, then there exists a unique φ in $\text{Irr}_H(L)$ under θ .*
- (b) *If $\varphi \in \text{Irr}_H(L)$, then there exists a unique θ in $\text{Irr}_H(K)$ over φ .*

Now fix φ and θ as before.

- (c) *We have that θ extends to G if and only if φ extends to LH .*

Proof. This result is Theorem 3.5 of [14], taking into account that, since $\mathfrak{N} \subseteq \mathfrak{F}$, the \mathfrak{F} -projectors of G are self-normalizing by Proposition 2.2 (e). \square

Let \mathfrak{F} be a saturated formation. Suppose that K is a normal subgroup of a solvable group G such that $G = KH$, where H is an \mathfrak{F} -projector of G . If L is a normal subgroup of H such that K/L is abelian and $K \cap HL = L$, then we say that (KH, K, L) satisfies the *Navarro condition*. Whenever \mathfrak{F} contains \mathfrak{N} and the Navarro condition is satisfied, we can apply Theorem 3.3.

We are interested in studying whether there is any relationship between the two sets of H -invariant character extensions that intervene in the previous theorem. The result below was generously communicated to us by G. Malle, G. Navarro and P. H. Tiep [13]. We are thankful to them for allowing us to present it here.

Theorem 3.4 (Malle, Navarro, Tiep). *Suppose G is a finite solvable group, $K, L \trianglelefteq G$ such that K/L is abelian. Suppose that H/L is a self-normalizing nilpotent subgroup of G/L , and suppose that $G = KH$ and $K \cap H = L$. Let $\theta \in \text{Irr}_H(L)$ and let $\varphi \in \text{Irr}_H(K)$ lying over φ . Then the following hold.*

- (a) *If $\chi \in \text{Irr}(G)$ extends θ , then there is an irreducible constituent of χ_H extending φ .*
- (b) *If $\nu \in \text{Irr}(H)$ extends φ , then there is an irreducible constituent of ν^G extending θ .*

In the proof of the following lemma, we will use Isaacs' notation from [7].

Lemma 3.5. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Suppose that K and L are normal subgroups of a solvable group G such that K/L is a chief factor of G , $K \cap LH = L$, and $G = KH$, where H is an \mathfrak{F} -projector of G . Moreover, assume that G has odd order whenever $\mathfrak{N} \neq \mathfrak{F}$. Let $\varphi \in \text{Irr}(L)$ be fully ramified with respect to K/L , and let $\theta \in \text{Irr}(K)$ be the unique H -invariant irreducible character lying over φ .*

- (a) *If η is an irreducible extension of φ to LH , then there exists an extension $\chi \in \text{Irr}(G)$ of θ lying over η .*
- (b) *If χ is an irreducible extension of θ to G , then there exists an extension $\eta \in \text{Irr}(LH)$ of φ lying under χ .*

Proof. We have that (G, L, φ) is a character triple. By a character triple isomorphism, we can assume that L is a central subgroup of G and that φ is faithful. Additionally, each coset of L contains an element x such that $\langle x \rangle \cap L = 1$ by Theorem 8.2 in [7].

We also have that K/L is a p -group. If $|G|$ is odd, then $\mathcal{C} = (G, K, H, L, \theta, \varphi)$ is a constellation with prime p (see Definition 4.5 of [7]). Let ψ be the unique canonical character arising from \mathcal{C} that exists by Theorem 5.3 of [7]. By Corollary 5.9 of [7], ψ contains as an irreducible constituent the trivial character 1_H . Since for every extension of θ to G there is a unique extension ν of φ to H such that $\chi_H = \psi\nu$, (a) follows easily.

But the number of extensions of θ to G is the same as the number of extensions of φ to H and thus (b) follows, by Corollary 3.6 of [7].

Now, we consider the case in which $\mathfrak{F} = \mathfrak{N}$. Then H is nilpotent and the result follows by Theorem 3.4 with $H = CL$. \square

We prove that H -invariant extensions to KH , as presented in Theorem 3.3, lie over H -invariant extensions to LH if G is of odd order or \mathfrak{F} is the formation of nilpotent groups.

Theorem 3.6. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Suppose that K and L are normal subgroups of a group G , such that K/L is an abelian factor of G , $K \cap LH = L$, and $G = KH$, where H is an \mathfrak{F} -projector of G . Further assume that G has odd order whenever $\mathfrak{N} \neq \mathfrak{F}$. Let $\varphi \in \text{Irr}_H(L)$, and let $\theta \in \text{Irr}_H(K)$ be the unique H -invariant irreducible character over φ .*

- (a) *If η is an irreducible extension of φ to LH , then there exists an extension $\chi \in \text{Irr}(G)$ of θ lying over η .*
- (b) *If χ is an irreducible extension of θ , then there exists an extension $\eta \in \text{Irr}(LH)$ of φ lying under χ .*

Proof. Working by induction on $|K : L|$ and $|G : K|$, we can assume that K/L is a G -chief factor. Otherwise, let K/M be a chief factor of G , and $\mu \in \text{Irr}_H(M)$ be the unique H -invariant irreducible character under θ and over φ (this follows from Theorem 3.3).

By induction, we have that if η is an irreducible extension of φ to LH , then there exists an irreducible extension γ of the character μ to HM lying over η . Then, again by induction, we have that there is an irreducible extension χ of the character θ to G lying over γ . Hence, χ lies over η and we have proved (a).

Conversely, if χ is an irreducible extension of θ to G , then, by induction, there exists an irreducible extension γ of the character μ to HM lying under χ . Then, again by induction, we have that there is an extension η of the character φ to LH lying under η . Therefore, η lies under χ and we have (b).

We may also assume that φ is G -invariant. Otherwise, since the inertia subgroup $I_G(\varphi) = LH$ is normal in G and K/L is a G -chief factor, we have that $I_K(\varphi) = L$. Then, by Lemma 3.1, induction defines a bijection from $\text{Irr}(LH \mid \varphi)$ onto $\text{Irr}(G \mid \theta)$, and (a) and (b) follow.

Now, by the “going down” theorem, we have that either θ is fully ramified with respect to K/L or θ is an irreducible extension of φ to K . In the latter case, we have that the restriction defines a bijection $\text{Irr}(G \mid \theta) \rightarrow \text{Irr}(K \mid \varphi)$ by Lemma 3.2, and the result follows.

So, we may assume that θ is fully ramified with respect to K/L . Therefore the result holds by Lemma 3.5. \square

4. \mathfrak{F}' -CHARACTERS

In this section, we present the \mathfrak{F}' -characters defined in [14] for the particular case where $\mathfrak{N} \subseteq \mathfrak{F}$. Recall that saturated formations containing the nilpotent groups share the property of their respective projectors being self-normalizing. Thus, we may begin by noting that in this situation, if G is a solvable group and H is an \mathfrak{F} -projector of G , any normal subgroup N of G such that $NH \trianglelefteq G$ satisfies in fact that $NH = G$ (see Proposition 2.2 (d)).

The construction in [14] relies on Navarro’s definition of $G^{\mathfrak{F}^n}$, the unique minimal normal subgroup of G such that $G^{\mathfrak{F}^n}H \trianglelefteq G$. It follows from the above paragraph that in our context $G^{\mathfrak{F}^n}$ coincides with $G^{\mathfrak{F}}$, the \mathfrak{F} -residual. This particularly means that $G^{\mathfrak{F}}$ is a proper subgroup of G .

Using the notation from [14], we observe that the tuple $(U_i, K_i, L_i, L_iH, U_i)$ satisfies $U_i = L_iH$ for all i . We now introduce the notation that will be used throughout this paper.

Let $K_0 = G^{\mathfrak{F}}$, $L_0 = K'_0$, $K_i = (L_{i-1}H)^{\mathfrak{F}}$, and $L_i = K'_i$. Note that, since the \mathfrak{F} -residual is a proper subgroup, there exists a non-negative integer m such that the following chain of subgroups holds:

$$G > L_0H > L_1H > \cdots > L_{m-2}H > L_{m-1}H = H.$$

At this point, $K_m = (L_{m-1}H)^{\mathfrak{F}} = H^{\mathfrak{F}} = 1$. Thus, we have the following H -invariant subnormal series:

$$G \triangleright K_0 \triangleright L_0 \triangleright \cdots \triangleright K_{m-1} \triangleright L_{m-1} \triangleright 1.$$

Note that the subgroups in this chain satisfy $G = K_0H$, $L_iH = K_{i+1}H$ for all $i \geq 0$, and that the Navarro condition is satisfied for (K_iH, K_i, L_i) .

We now proceed to describe the process for computing the characters. To this end, we introduce the following notation: we write $(L, \varphi) \triangleleft (K, \theta)$ to indicate that L is a normal subgroup of K , and the irreducible character θ of K lies over the irreducible character φ of L .

We say that χ is an \mathfrak{F}' -character, and we write $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$, if there exist characters $\theta_i \in \text{Irr}(K_i)$ and $\varphi_i \in \text{Irr}(L_i)$, extending to K_iH and L_iH , respectively, such that

$$(1, 1) \triangleleft (L_{m-1}, \varphi_{m-1}) \triangleleft (K_{m-1}, \theta_{m-1}) \triangleleft \cdots \triangleleft (L_0, \varphi_0) \triangleleft (K_0, \theta_0) \triangleleft (G, \chi).$$

By the construction of the \mathfrak{F}' -characters, if $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$, then the extension ψ_i of φ_i to L_iH , which coincides with the extension of θ_{i+1} to $K_{i+1}H = L_iH$, is also an \mathfrak{F}' -character of L_iH . Moreover, it follows from the construction that χ_{K_0} is an irreducible character.

Next, we provide information about the number of \mathfrak{F}' -characters of a solvable group in terms of its \mathfrak{F} -projector.

Proposition 4.1. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Let G be a solvable group, and let H be an \mathfrak{F} -projector of G . Then*

$$|\text{Irr}_{\mathfrak{F}'}(G)| = |\text{Irr}(H/H')|.$$

That is, the number of \mathfrak{F}' -characters of G coincides with the number of linear characters of H .

Proof. This follows from Theorem A of [14]. □

5. STRONG PAIR SERIES

In this section, we aim to present an alternative characterization of the \mathfrak{F}' -characters. This characterization is analogous to the one given by Isaacs in [10] for the formation \mathfrak{N} of nilpotent groups, but it extends to all saturated formations containing \mathfrak{N} .

Let G be a solvable group and H an \mathfrak{F} -projector of G . Suppose that $\{S_i\}_{0 \leq i \leq r}$ is an H -composition series for G . This is an H -invariant, subnormal series such that S_{i+1}/S_i is H -simple (and thus abelian) for $0 \leq i \leq r-1$. The H -composition length r is an invariant of G .

We say that $\{(S_i, \theta_i)\}_{0 \leq i \leq r}$ is an H -pair series for G that is associated with the H -composition series $\{S_i\}_{0 \leq i \leq r}$, if the characters $\theta_i \in \text{Irr}(S_i)$ are H -invariant and θ_i lies under θ_{i+1} for $0 \leq i < r$. Sometimes, we will write the H -pair series as

$$(1, 1) = (S_0, \theta_0) \triangleleft (S_1, \theta_1) \triangleleft \cdots \triangleleft (S_r, \theta_r)$$

to simplify the understanding of the chain of pairs being used.

We say that an H -pair series $\{(S_i, \theta_i)\}_{0 \leq i \leq r}$ for G is *strong* if each of the characters θ_i extends irreducibly to $S_i H$. An irreducible character χ of G is said to be a *head character* of G if (G, χ) appears as (S_r, θ_r) in a strong H -pair series.

It is worth noting that these (irreducible) extensions define, in turn, head characters of $S_i H$. Indeed, if θ_i extends to $\eta_i \in \text{Irr}(S_i H)$ and $\{T_j\}_{0 \leq j \leq m}$ is such that $T_0 = S_i$, $T_m = S_i H$ and T_{j+1}/T_j is an H -composition factor for all j , then one can easily verify that

$$(S_0, \theta_0) \triangleleft \cdots \triangleleft (S_i, \theta_i) = (T_0, \eta_{i|_{T_0}}) \triangleleft \cdots \triangleleft (T_j, \eta_{i|_{T_j}}) \triangleleft \cdots \triangleleft (T_m, \eta_i) = (S_i H, \eta_i)$$

is a strong H -pair series for $S_i H$.

Lemma 5.1. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Let G be a solvable group, and let H be an \mathfrak{F} -projector of G . Suppose that M/U is an H -composition factor of G . Let α be an irreducible character of M lying over the irreducible character θ of U . If α and θ extend irreducibly to MH and UH respectively, then θ is the unique H -invariant irreducible constituent of U lying under α . Moreover, if $MH = UH$, then $\alpha_U = \theta$.*

Proof. We have that $M \cap UH$ is a normal subgroup of MU that contains U and is contained in M . Since M/U is an MH -chief factor, it follows that $M \cap UH$ is either U or M .

If $M \cap UH = U$, then (MH, M, U) satisfies the Navarro condition, and by Theorem 3.3, we have that θ is the unique H -invariant character of U lying under α .

In the latter case, where $M \cap UH = M$, we have $MH = UH$. Let $\tilde{\theta}$ be an extension of θ to MH . Then $\tilde{\theta}_M$ is irreducible. Since M/U is abelian, by Gallagher's theorem, we have $\alpha = \lambda \tilde{\theta}_M$ for some $\lambda \in \text{Lin}(M/U)$. Therefore, $\alpha_U = \theta$, which completes the proof. □

Lemma 5.2. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Let G be a solvable group, and let H be an \mathfrak{F} -projector of G . Suppose that M/V and M/U are two distinct H -composition factors, and let $D = U \cap V$. If*

$$(D, \gamma) \triangleleft (U, \theta) \triangleleft (M, \alpha),$$

with γ , θ , and α extending irreducibly to DH , UH , and MH , respectively, then there exists an H -invariant irreducible character β of V , which extends irreducibly to VH and satisfies

$$(D, \gamma) \triangleleft (V, \beta) \triangleleft (M, \alpha).$$

Proof. Observe that both M/U and M/V are MH -chief factors; therefore, they are abelian, and M/D is abelian as well. Moreover, $M \cap UH$ is either M or U . The same holds with V in place of U . Since $U/D \cong M/V$, we deduce that $U \cap DH$ can be either U or D . Analogously, the same applies to V/D .

Assume first that $MH = VH = DH = UH$. By Lemma 5.1, it follows that $\alpha_U = \theta$ and $\theta_D = \gamma$. Therefore, α is an extension of γ . It suffices to consider $\beta = \alpha_V$.

Now, suppose that $MH = UH$ but $VH < MH$. In this situation, we have that $VH \cap M = V$ and that $DH < UH$, which implies $DH \cap U = D$. Since $D = DH \cap U \subseteq VH \cap U \subseteq MH \cap U = U$, it follows that $VH = DH$. Therefore, $M \cap DH = M \cap VH = V$. Since θ extends irreducibly to $MH = UH$ and α lies over θ , and because M/U is abelian, as a consequence of Gallagher's theorem, we have that $\alpha_U = \theta$. Note that (MH, M, V) satisfies the Navarro condition. Since U/D is a chief factor of $UH = MH$, it follows that either γ is U -invariant and hence M -invariant, or $\theta = \gamma^U$.

Therefore, there exists a unique irreducible character β that is H -invariant below α and extends to VH , because α does so. In the first case, we have that γ is the only irreducible constituent of α_D , so it is clear that β is a character lying above γ . Suppose γ induces irreducibly to U . By applying Lemma 3.1, it follows that $\beta \in \text{Irr}(V | \gamma)$, as desired.

Next, suppose that $MH = VH$ but $UH < MH$. Similarly to the previous paragraph, by interchanging U and V , we have $DH = UH$ and $U = M \cap UH$. By Lemma 5.1, it follows that $\theta_D = \gamma$. Observe that (VH, V, D) satisfies the Navarro condition. Consequently, there exists a unique character $\beta \in \text{Irr}_H(V | \gamma)$. We want to show that β is a constituent of α_V .

If θ is M -invariant, we can apply Lemma 3.2 to deduce that $\alpha_V \in \text{Irr}(V | \gamma)$. Thus, $\beta = \alpha_V$. Assume now that $\alpha = \theta^M$. In this case, β is an irreducible constituent of $\gamma^V = \theta_D^V = \theta_V^M = \alpha_V$.

Now we consider the case where $DH < MH = VH = UH$. We have $DH \cap U = D$ and $DH \cap V = D$, and we also have $DH = XH$, where $X = M \cap DH$. This is the second case, considering X instead of V . Hence, there exists $\beta_0 \in \text{Irr}(X | \gamma)$ that extends to XH and is a constituent of α_X .

Now, by taking X instead of U , we have $MH = VH$ and $DH = XH$. Then we are in the situation described in the previous case, where we know that there exists $\beta \in \text{Irr}(V | \gamma)$ that extends to VH . Moreover, we have $\alpha_V = \beta$.

The final case is when $UH < MH$ and $VH < MH$. It follows that $UH \cap M = U$ and $VH \cap M = V$. Observe that $DH < UH$, as otherwise we would have $U = M \cap UH = M \cap DH \subseteq M \cap VH = V$, which leads to a contradiction. (Similarly, $DH < VH$.) Since both (MH, M, V) and (VH, V, D) satisfy the Navarro condition, the theorem follows as a direct consequence of applying Theorem 3.3 twice. \square

The proof of the theorem below follows that of Theorem 5.5 of [10], differing only in the final step.

Theorem 5.3. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Let G be a solvable group, and let H be an \mathfrak{F} -projector of G . Suppose $\{T_i \mid 0 \leq i \leq r\}$ is an arbitrary H -composition series for G . Let $\chi \in \text{Irr}(G)$ be a head character of G . Then the series $\{T_i\}$ is associated with some unique strong H -pair series $\{(T_i, \varphi_i) \mid 0 \leq i \leq r\}$ such that $\varphi_r = \chi$.*

Proof. Since χ is a head character of G , by definition, there exists a strong H -pair series

$$(1, 1) = (S_0, \theta_0) \triangleleft (S_1, \theta_1) \triangleleft \cdots \triangleleft (S_r, \theta_r) = (G, \chi).$$

We want to show the existence of characters $\varphi_i \in \text{Irr}(T_i)$ satisfying that

$$(1, 1) = (T_0, \varphi_0) \triangleleft (T_1, \varphi_1) \triangleleft \cdots \triangleleft (T_r, \varphi_r) = (G, \chi)$$

is a strong H -pair series.

Now, note that $S_r = G = T_r$. Therefore, there exists a unique smallest non-negative integer m such that $S_i = T_i$ for all i satisfying $m \leq i \leq r$. If $m = 0$, then $T_i = S_i$ for all i , and we can set $\varphi_i = \theta_i$, concluding the proof. Thus, we assume $m > 0$ and proceed by downward induction on m .

We have $S_m = T_m$, but $S_{m-1} \neq T_{m-1}$. Since $S_0 = 1 = T_0$, it follows that $m - 1 > 0$, and hence $m \geq 2$. Define $M = S_m = T_m$, $U = S_{m-1}$, and $V = T_{m-1}$. Write $D = U \cap V$. Clearly, U and V are nontrivial and distinct. Furthermore, M/U and M/V are H -composition factors of G , so U and V are maximal among H -invariant normal subgroups of M . As $U \neq V$, it follows that $UV = M$. And, U/D is H -isomorphic to M/V , thus U/D is an H -composition factor of G , and similarly, V/D is an H -composition factor of G .

The H -composition length of $U = S_{m-1}$ is $m - 1$, implying that the H -composition length of D is $m - 2$. Thus, we can choose an H -composition series

$$1 = D_0 \triangleleft D_1 \triangleleft \cdots \triangleleft D_{m-2} = D$$

for D . Appending the subgroups S_i for $m - 1 \leq i \leq r$ to this series yields a new H -composition series for G . Denote this series by $\{N_i\}$, where $N_i = D_i$ for $0 \leq i \leq m - 2$, and $N_i = S_i$ for $m - 1 \leq i \leq r$. In particular, $N_{m-1} = S_{m-1} = U$ and $N_{m-2} = D_{m-2} = D$.

Using the inductive hypothesis with the series $\{N_i\}$ in place of $\{T_i\}$ and $m - 1$ in place of m , we conclude that $\{N_i\}$ is associated with a strong H -pair series with head character χ . Moreover, for $i \geq m$, we have $T_i = S_i = N_i$. Hence, we may replace $\{S_i\}$ with $\{N_i\}$ without loss of generality, and we can assume $N_i = S_i$ for all i . In particular, $D = D_{m-2} = N_{m-2} = S_{m-2}$.

Next, construct a new H -composition series for G by modifying $\{S_i\}$, replacing $S_{m-1} = U$ with V . Note that $S_{m-2} = D \triangleleft V \triangleleft M = S_m$. And since V/D is H -isomorphic to $M/U = S_m/S_{m-1}$ and M/V is H -isomorphic to T_m/T_{m-1} , the series is an H -composition series. We denote this new series by $\{X_i\}$, where $X_i = S_i$ for $i \neq m - 1$ and $X_{m-1} = V$.

Finally, by Theorem 5.3 we have that $\{X_i\}$ is associated with a strong H -pair series with head character χ . Assuming this is established, note that $T_i = S_i = X_i$ for $i \geq m$ and $T_{m-1} = V = X_{m-1}$. Thus, the inductive hypothesis applies to $\{X_i\}$ in place of $\{S_i\}$ with $m - 1$ in place of m . Therefore the original series $\{T_i\}$ is associated with a strong H -pair series with head character χ .

The uniqueness of this strong H -pair series follows from Lemma 5.1. \square

Theorem 5.4. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Let G be a solvable group, and let $\chi \in \text{Irr}(G)$. Then the following are equivalent:*

- (a) χ is an \mathfrak{F}' -character of G ;

(b) χ is a head character of G .

Proof. Suppose χ is an \mathfrak{F}' -character of G . By construction, there exist characters $\theta_i \in \text{Irr}(K_i)$ and $\varphi_i \in \text{Irr}(L_i)$, extending to K_iH and L_iH , respectively, such that

$$(1, 1) \triangleleft (L_{m-1}, \varphi_{m-1}) \triangleleft (K_{m-1}, \theta_{m-1}) \triangleleft \cdots \triangleleft (L_0, \varphi_0) \triangleleft (K_0, \theta_0) \triangleleft (G, \chi).$$

where $K_0 = G^{\mathfrak{F}}$, $L_0 = K'_0$, $K_i = (L_{i-1}H)^{\mathfrak{F}}$, and $L_i = K'_i$.

From this series we can obtain an H -subnormal series. Consider $(U, \alpha) \triangleleft (V, \beta)$, where U/V is H -simple. If $L_i \leq U < V \leq K_i$, then (VH, V, U) satisfies the Navarro condition. By applying Theorem 3.3 to each H -composition factor between L_i and K_i , it follows that α and β are H -invariant and extend irreducibly to UH and VH , respectively.

If $K_{i+1} \leq U < V \leq L_i$, it follows that $\alpha = (\varphi_i)_U$ and $\beta = (\varphi_i)_V$, which clearly extend irreducibly to UH and VH , respectively.

The converse follows by applying Theorem 5.3, considering an H -composition series passing through

$$G \triangleright K_0 \triangleright L_0 \triangleright \cdots \triangleright K_{m-1} \triangleright L_{m-1} \triangleright 1.$$

□

6. THEOREM A

The purpose of this section is to study how \mathfrak{F}' -characters of G behave with respect to restriction, when G is a solvable group and \mathfrak{F} is a saturated formation. If $\mathfrak{N} \subseteq \mathfrak{F}$, there is a relationship between the \mathfrak{F}' -characters of G and those of NH , where N is a normal subgroup of G and H is an \mathfrak{F} -projector of G .

Proposition 6.1. *Let \mathfrak{F} be a saturated formation such that $\mathfrak{N} \subseteq \mathfrak{F}$, and let G be a solvable group. Suppose H is an \mathfrak{F} -projector of G , and assume that G has odd order whenever $\mathfrak{N} \neq \mathfrak{F}$. Let $K = G^{\mathfrak{F}}$ be the \mathfrak{F} -residual of a solvable group G . Suppose that $M \triangleleft G$ is such that K/M is a G -chief factor.*

- (a) *If $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$, then there exists $\mu \in \text{Irr}_{\mathfrak{F}'}(MC)$ lying under χ .*
- (b) *If $\mu \in \text{Irr}_{\mathfrak{F}'}(MC)$, then there exists $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$ lying over μ .*

Proof. The \mathfrak{F} -residual residual of G/M is K/M , which is abelian, and is therefore complemented by HM/M (by Theorem 2.3). From this, we see that Theorem 3.6 can be applied to $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$. Thus, χ_{MH} has an irreducible constituent μ that restricts irreducibly to M . We have that $\varphi = \mu_M$ is the unique H -invariant constituent of $(\chi_K)_M$ (by Theorem 3.3). Now, we can consider the following strong H -pair series, which passes through K and M :

$$(1, 1) = (S_0, \varphi_0) \triangleleft \cdots \triangleleft (M, \varphi) = (S_m, \varphi_m) \triangleleft (K, \chi_K) = (S_{m+1}, \varphi_{m+1}) \triangleleft \cdots \triangleleft (S_r, \varphi_r) = (G, \chi).$$

From the remarks we made when defining a strong H -pair series and the fact that μ extends φ , it follows that $\mu \in \text{Irr}_{\mathfrak{F}'}(MH)$.

On the other hand, if $\mu \in \text{Irr}_{\mathfrak{F}'}(MH)$, again by Theorem 3.6, there exists $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$ lying over μ that restricts irreducibly to K , with $\theta = \chi_K$ being the unique irreducible H -invariant constituent of φ^K where $\varphi = \mu_M$ (by Theorem 3.3).

Now, let $(1, 1) = (S_0, \varphi_0) \triangleleft \cdots \triangleleft (M, \mu) = (S_m, \varphi_m) \triangleleft \cdots \triangleleft (S_l, \varphi_l) = (MC, \mu)$ be a strong H -pair series for μ . We can find an H -composition series for G that passes through M and K . Let us write it $1 = T_0 \triangleleft \cdots \triangleleft T_t = M \triangleleft T_{t+1} = K \triangleleft \cdots \triangleleft T_r = G$.

We may then construct a strong H -pair series having χ as the head character and passing through K and M as follows:

$$(1, 1) = (S_0, \varphi_0) \triangleleft \cdots \triangleleft (M, \mu) = (S_m, \varphi_m) \triangleleft (K, \chi_K) = (T_{t+1}, \chi_{T_{t+1}}) \triangleleft \cdots \triangleleft (T_r, \chi_{T_r}) = (G, \chi).$$

Therefore $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$, as desired. \square

This allows us to prove a generalization of Theorem A to saturated formations containing \mathfrak{N} , as presented below.

Theorem 6.2. *Let \mathfrak{F} be a saturated formation such that $\mathfrak{N} \subseteq \mathfrak{F}$, and let G be a solvable group. Suppose H is an \mathfrak{F} -projector of G , and assume that G has odd order whenever $\mathfrak{N} \neq \mathfrak{F}$. If $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$ and N is normal subgroup of G , then the restriction of χ to NH has an irreducible constituent that is an \mathfrak{F}' -character of NH .*

Proof. Let $K = G^{\mathfrak{F}}$. We have that $K \cap N$ is normal in G and is contained in K . Suppose that there is $K \cap N \trianglelefteq M \trianglelefteq K$ where K/M is an H -composition factor. Note that if $K = MH \cap K$, then G/M would be in \mathfrak{F} , contradicting the fact that K is the \mathfrak{F} -radical. Therefore, $K \cap (MH) = M$.

We prove this result by induction on $|G : NH|$. We can assume that $K/(K \cap N)$ is an H -composition factor. Otherwise, let K/M be an H -composition factor. By the Theorem 3.6, there exists an irreducible constituent α of χ_{MH} that is an \mathfrak{F}' -character of MH . Now, we have two possibilities: $MH = NH$, and we are done, or $MH \neq NH$, and hence $|MH : NH| < |G : NH|$. Applying induction, we conclude that α_{NH} has an irreducible constituent that is an \mathfrak{F}' -character of NH . Hence, $\alpha \in \text{Irr}_{\mathfrak{F}'}(NH)$ is an irreducible constituent of χ_{NH} , as desired. \square

Theorem 6.2 considers restriction of \mathfrak{F}' -characters χ of G to subgroups of the form NH . Note that, since the \mathfrak{F}' -characters are constructed in relation to an \mathfrak{F} -projector, we are only interested in \mathfrak{F}' -characters of subgroups having as an \mathfrak{F} -projector the same one we fixed for G , that is, subgroups containing H .

When considering restriction of \mathfrak{F}' -characters χ of G to normal subgroups N that might not contain H , we prove instead that there always exists an H -invariant constituent of χ_N .

If G is of odd order whenever $\mathfrak{N} \neq \mathfrak{F}$, this is a direct consequence of Theorem 6.2, using the fact that \mathfrak{F}' -characters of NH restrict irreducibly to $(NH)^{\mathfrak{F}}$ and that, by definition, $(NH)^{\mathfrak{F}}$ is contained in N . Nonetheless, the result remains true without this hypothesis, as we state in Proposition 6.3 below. We point out that Theorem 6.2 does, however, require G having odd order if $\mathfrak{N} \neq \mathfrak{F}$. The group `SmallGroup(48,28)` of order 48 and the saturated formation of supersolvable groups provide a counterexample to removing this hypothesis.

Proposition 6.3. *Let \mathfrak{F} be a saturated formation such that $\mathfrak{N} \subseteq \mathfrak{F}$, and let G be a solvable group. Suppose H is an \mathfrak{F} -projector of G . If $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$ and N is a normal subgroup of G , then χ_N has an H -invariant constituent.*

Proof. We can find an H -composition series that passes through G via N . Let us consider

$$1 = S_0 \triangleleft \cdots \triangleleft S_m = N \triangleleft \cdots \triangleleft S_r = G.$$

By Theorem 5.3 and Theorem 5.4, this series is associated with a strong H -pair series $\{(S_i, \theta_i)\}_{0 \leq i \leq r}$ such that $\theta_r = \chi$. Then θ_m is a constituent of χ_N which extends to NH . In particular, it is H -invariant. \square

7. THEOREM B

In this final section, we prove Theorem B in the more general context of saturated formations containing all nilpotent groups. This is formulated in Theorem 7.5. We first present some preliminary results.

Lemma 7.1. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Let G be a solvable group, and let H be an \mathfrak{F} -projector of G . Let M, T be H -invariant subgroups of G such that $T \triangleleft M$ and M/T is H -simple. Suppose $M \cap H \subseteq H'$. If φ is an irreducible H -invariant character of M with T in its kernel that extends to MH , then φ is the trivial character.*

Proof. Since M/T is H -simple, it is abelian, and thus $M/T(H \cap M)$ is also abelian. Notice too that, since H is an \mathfrak{F} -projector of G and in particular of MH , we have that $HT(H \cap M)/T(H \cap M) = HT/T(H \cap M)$ is an \mathfrak{F} -projector of $MH/T(H \cap M)$. Let φ be an irreducible H -invariant character of M that extends to MH and has T in its kernel. By applying Theorem 3.3 (a) to MH , we know that there exists a unique $\xi \in \text{Irr}_H(T(H \cap M))$ lying under φ . Moreover, ξ extends to some $\eta \in \text{Irr}(TH)$ by Theorem 3.3(c).

Since φ has T in its kernel, ξ also has T in its kernel. Notice that, since M/T is abelian, φ is linear, and thus so are ξ and η . Then η can be seen as a character in $\text{Lin}(HT/T)$ with $H'T/T$ in its kernel. Thus, $H'T \subseteq \ker(\eta)$. But $(H \cap M)T \subseteq H'T$ since $H \cap M \subseteq H'$. Hence, $\xi = \eta_{(H \cap M)T} = 1_{(H \cap M)T}$. Since φ is by Theorem 3.3 (a) the unique H -invariant character lying over ξ , it must be 1_M . \square

Lemma 7.2. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Let G be a solvable group, and let H be an \mathfrak{F} -projector of G . Suppose that N is a normal subgroup of G such that $N \cap H \subseteq H'$. Then $N \subseteq \ker(\chi)$ for all $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$.*

Proof. Let us consider a fixed $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$. By Theorem 5.4, there exists a strong H -pair series of the form

$$(1, 1) = (S_0, \varphi_0) \trianglelefteq \cdots \trianglelefteq (S_m, \varphi_m) = (N, \varphi_m) \trianglelefteq \cdots \trianglelefteq (S_r, \varphi_r) = (G, \chi).$$

Every $1 \leq i \leq m$ satisfies that S_i and S_{i-1} are H -invariant, S_i/S_{i-1} is H -simple and $S_i \cap H \subseteq H'$. Moreover, $\varphi_0 = 1_{S_0}$. Then, by repeatedly applying Lemma 7.1, we deduce that $\varphi_i = 1_{S_i}$ for every $0 \leq i \leq m$. In particular, $\varphi_m = 1_N$. But $N \triangleleft G$ and φ_m is a constituent of χ_N . Therefore $N \subseteq \ker(\chi)$. \square

If N is a normal subgroup of G , every irreducible character $\tilde{\chi}$ of the quotient group G/N can be regarded, by inflation, as a character of G whose kernel contains N . In this case, we denote the corresponding character in $\text{Irr}(G)$ by χ , where $N \subseteq \ker(\chi)$, and vice versa (Lemma (2.2) of [9]).

Lemma 7.3. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Suppose that N is a normal subgroup of a solvable group G . Then*

- (a) *if $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$ such that $N \subseteq \ker(\chi)$, then $\tilde{\chi} \in \text{Irr}_{\mathfrak{F}'}(G/N)$;*
- (b) *if $\tilde{\chi} \in \text{Irr}_{\mathfrak{F}'}(G/N)$, then $\chi \in \text{Irr}(G)$ is an \mathfrak{F}' -character.*

Proof. Let H be an \mathfrak{F} -projector of G . First suppose that $\chi \in \text{Irr}_{\mathfrak{F}'}(G)$ is such that $N \subseteq \ker(\chi)$. By Theorem 5.4, there exists a strong H -pair series

$$(1, 1) = (S_0, \varphi_0) \trianglelefteq (S_1, \varphi_1) \trianglelefteq \cdots \trianglelefteq (S_m, \varphi_m) = (N, \varphi_m) \trianglelefteq \cdots \trianglelefteq (S_r, \varphi_r) = (G, \chi).$$

Since $S_m \subseteq \ker(\chi)$, and φ_m lies under χ , we have that $\varphi_m = 1_N$ is the trivial character. Thus, $\varphi_i = 1_{S_i}$ for every $0 \leq i \leq m$. Now, let $\bar{G} = G/N$. We have that $\bar{H} = HN/N$ is an \mathfrak{F} -projector of \bar{G} . Adopting the bar convention for the quotients and the tilde convention for their irreducible characters, we can consider the following \bar{H} -pair series,

$$(\bar{1}, \tilde{1}) = (\bar{S}_m, \tilde{\varphi}_m) \trianglelefteq \cdots \trianglelefteq (\bar{S}_i, \tilde{\varphi}_i) \trianglelefteq \cdots \trianglelefteq (\bar{S}_r, \tilde{\varphi}_r) = (\bar{G}, \tilde{\chi})$$

which is a strong \bar{H} -pair series as it is not difficult to verify. Hence, $\tilde{\chi}$ is an \mathfrak{F}' -character of \bar{G} .

Now, suppose that $\tilde{\chi} \in \text{Irr}_{\mathfrak{F}'}(G/N)$, then there is a strong \bar{H} -pair series

$$(\bar{1}, \tilde{1}) = (\bar{S}_0, \tilde{\varphi}_0) \triangleleft \cdots \triangleleft (\bar{S}_i, \tilde{\varphi}_i) \triangleleft \cdots \triangleleft (\bar{S}_r, \tilde{\varphi}_r) = (\bar{G}, \tilde{\chi}).$$

Now, let us consider an H -composition series of N

$$1 = R_0 \triangleleft R_1 \triangleleft \cdots \triangleleft R_s = N.$$

It suffices to consider the following series, viewing the characters of the quotient as characters of the group with N in the kernel, as we have mentioned previously:

$$(1, 1) = (R_0, 1) \trianglelefteq \cdots \trianglelefteq (R_s, 1_{R_s}) = (N, 1_N) = (S_0, \varphi_0) \trianglelefteq \cdots \trianglelefteq (S_r, \varphi_r) = (G, \chi),$$

which again may be verified to be a strong H -pair series without difficulty. \square

Lemma 7.4. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Let G be a solvable group G . Suppose that N is a normal subgroup in G contained in $\bigcap_{\chi \in \text{Irr}_{\mathfrak{F}'}(G)} \ker(\chi)$. Then*

$$|\text{Irr}_{\mathfrak{F}'}(G)| = |\text{Irr}_{\mathfrak{F}'}(G/N)|.$$

Moreover, $N \cap H \subseteq H'$ for any \mathfrak{F} -projector H of G .

Proof. The previous lemma give us an injection $\text{Irr}_{\mathfrak{F}'}(G) \rightarrow \text{Irr}_{\mathfrak{F}'}(G/N)$. Hence

$$|H/H'| = |\text{Irr}_{\mathfrak{F}'}(G)| \leq |\text{Irr}_{\mathfrak{F}'}(G/N)| = |H/D|,$$

where H is any \mathfrak{F} -projector of G and $D/(H \cap N) = (H/(H \cap N))'$. Then, it follows that $|H'| \geq |D|$, but also $H' \subseteq D$ since $D/(H \cap N) = H'(H \cap N)/(H \cap N)$. Then $|H'| = |D|$ and we have the first part of the lemma. We conclude for the second part by noting that $H \cap N \subseteq D$ and $D = H'$. \square

Theorem 7.5. *Let \mathfrak{F} be a saturated formation with $\mathfrak{N} \subseteq \mathfrak{F}$. Let G be a solvable group, and let H be an \mathfrak{F} -projector of G . Then*

$$\bigcap_{\chi \in \text{Irr}_{\mathfrak{F}'}(G)} \ker(\chi)$$

is the largest normal subgroup M of G satisfying $M \cap H \subseteq H'$.

Proof. Let $M = \bigcap_{\chi \in \text{Irr}_{\mathfrak{F}'}(G)} \ker(\chi)$. It is clear that $M \cap H \subseteq H'$ from the previous lemma.

Suppose N and K are two normal subgroups of G such that $N \cap H \subseteq H'$ and $K \cap H \subseteq H'$. It follows that $NK \cap H = (N \cap H)(K \cap H) \subseteq H'$, by Lemma 2.4. Thus, there exists a maximal subgroup N satisfying $N \cap H \subseteq H'$. The only remaining step is now to prove that if N is a normal subgroup of G such that $N \cap H \subseteq H'$, then $N \subseteq M$. This follows from Lemma 7.2, completing the proof of the theorem. \square

The above theorem is the exact analog of an unpublished result by Navarro, which we present here with his permission. Note that, for a prime p and a solvable group G , the Sylow p -subgroups of G and the characters of G of p' -degree correspond respectively to the \mathfrak{F} -projectors and the \mathfrak{F}' -characters of G when \mathfrak{F} is the formation of p -groups. This is a saturated formation not containing \mathfrak{N} .

Theorem 7.6 (Navarro). *Let G be a finite group, let p be a prime, and let $P \in \text{Syl}_p(G)$. Then*

$$K = \bigcap_{\chi \in \text{Irr}_{p'}(G)} \ker(\chi)$$

is the largest normal subgroup K of G satisfying $\mathbf{N}_K(P) \subseteq P'$.

Proof. If N, M are normal subgroups of G , by Lemma 2.1 of [11], we have that

$$\mathbf{N}_{NM}(P) = \mathbf{N}_M(P)\mathbf{N}_N(P).$$

Therefore there exists a largest normal subgroup N of G such that $\mathbf{N}_N(P)$ is contained in P' . Since $N \cap P = \mathbf{N}_N(P) \subset P'$, by Tate's theorem (Theorem 6.31 of [9]), it follows that $\mathbf{O}^p(PN) \cap P = \mathbf{O}^p(P) = 1$. Therefore, PN has a normal p -complement, and N also has one. Let W be the normal p -complement of N . Then, we have $\mathbf{C}_W(P) = 1$ because $\mathbf{C}_W(P) \subseteq \mathbf{N}_N(P) \cap W \subseteq P' \cap W = 1$. Let $\chi \in \text{Irr}_{p'}(G)$, and let $\nu \in \text{Irr}(W)$ be a P -invariant irreducible constituent of the restriction χ_W , which exists because χ has p' -degree. By Glauberman's correspondence (Theorem 2.9 of [12]), we have that ν is the trivial character of W . Hence, $W \subseteq \ker(\chi)$. So, working in G/W we may assume that N is a p -group contained in P' . Now, let $\epsilon \in \text{Irr}(P)$ be a linear constituent of χ_P . Since 1_N is a constituent of ϵ_N , we have that $N \subseteq \ker(\chi)$.

Let

$$K = \bigcap_{\chi \in \text{Irr}_{p'}(G)} \ker(\chi).$$

It only remains to prove that $\mathbf{N}_K(P) \subseteq P'$. By Theorem 7.7 of [12], K has a normal p -complement Y . If $\delta \in \text{Irr}(Y)$ is P -invariant, then there is an extension η of δ to YP (by Corollary 6.28 of [9]), and therefore δ lies under some irreducible character of G of p' -degree (using that η^G has p' -degree). Thus δ is the trivial character of Y , and hence $\mathbf{C}_Y(P) = 1$ by Glauberman's correspondence. Thus $\mathbf{N}_K(P) = P \cap K$. We may assume that K is a p -group. Suppose that K is not contained in P' . Then some linear $\lambda \in \text{Irr}(P)$ does not contain K in its kernel. By inducing up to G , we obtain some $\chi \in \text{Irr}_{p'}(G)$ over λ and therefore a contradiction. \square

REFERENCES

- [1] R. W. Carter, Nilpotent self-normalizing subgroups of soluble groups, *Math. Z.* 75, 136-139 (1960/61).
- [2] E. C. Dade, Caractères venant des \mathfrak{F} -Normalisateurs d'un group fini résoluble, *J. reine angew. Math.* 307/308, 53-112 (1979).
- [3] K. Doerk, T. O. Hawkes, *Finite Soluble Groups*. De Gruyter, 1992.
- [4] D. Gajendragadkar, A characterization of characters which arise from \mathfrak{F} -normalizers, *J. reine und angew. Math.*, 319, 17-195 (1980).
- [5] W. Gaschütz, Zur Theorie der endlichen auflösbaren Gruppen, *Math. Zeitschr.* 80, 300-305 (1963).
- [6] B. Huppert, Zur Theorie der Formationen. *Arch. Math* 19, 561-574 (1969).
- [7] I. M. Isaacs, Characters of solvable and symplectic groups, *Am. J. Math.*, 95(3), 594-635 (1973).
- [8] I. M. Isaacs, Characters of π -separable groups, *J. Algebra* 86, 98-128 (1984).

- [9] I. M. Isaacs, *Character theory of finite groups*. Corrected reprint of the 1976 original. AMS Chelsea Publishing, Providence, RI, 2006.
- [10] I. M. Isaacs, Carter subgroups, characters and composition series. *Trans. Amer. Math. Soc. Series B*, Volume 9, Pages 470-498 (2022).
- [11] I. M. Isaacs, G. Navarro, A characteristic subgroup and kernels of Brauer characters, *Bulletin of the Australian Mathematical Society*, 72(3), 381-384 (2005).
- [12] G. Navarro, *Character Theory and the McKay Conjecture*. Cambridge University Press, 2018.
- [13] G. Malle, G. Navarro, P. H. Tiep, Degrees and Constituents of Characters, Preprint (2025).
- [14] G. Navarro, A McKay bijection for projectors, *Rev. Mat. Iberoam.* 38 no. 3, 1013-1028 (2022).
- [15] D. J. S. Robinson, *A Course in the Theory of Groups*, Graduate Texts in Mathematics 80, Springer-Verlag, New York-Berlin, 1982.

MARÍA JOSÉ FELIPE, INSTITUT UNIVERSITARI DE MATEMÀTICA PURA I APLICADA,
UNIVERSITAT POLITÈCNICA DE VALÈNCIA, VALÈNCIA, SPAIN.

Email address: mfelipe@mat.upv.es

IRIS GILABERT, INSTITUT UNIVERSITARI DE MATEMÀTICA PURA I APLICADA,
UNIVERSITAT POLITÈCNICA DE VALÈNCIA, VALÈNCIA, SPAIN.

Email address: igilman@posgrado.upv.es

LUCIA SANUS, DEPARTAMENT DE MATEMÀTIQUES, FACULTAT DE MATEMÀTIQUES,
UNIVERSITAT DE VALÈNCIA, 46100 BURJASSOT, VALÈNCIA, SPAIN.

Email address: lucia.sanus@uv.es