

Repeated Eigenvalues Imply Nodes? A Problem of Planar Differential Equations

Kenzi Odani
Department of mathematics, Aichi University of Education

March 7, 2025

Abstract


Poincaré gave a criterion which determines the shape of equilibrium for planar differential equations. In his statement, he excluded the case of repeated eigenvalues. In fact, in such a case, we can give a C^1 counter-example to his assertion. In this note, we show that if we strengthen the condition to $C^{1,\alpha}$ ($0 < \alpha < 1$), his assertion becomes true even in case of repeated eigenvalues.

1 Introduction.

Consider a planar differential equation below:

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^2, \tag{1}$$

where \mathbf{f} is a 2-dimensional function. We call a point \mathbf{p} an *equilibrium* if $\mathbf{f}(\mathbf{p}) = \mathbf{0}$. We call an equilibrium *repelling* (*attracting*) if all nearby trajectories are outgoing (incoming). We call a repelling or an attracting equilibrium \mathbf{p} a *node* (*focus*) if the direction $\mathbf{d}(t) = \mathbf{f}(\mathbf{x}(t))/|\mathbf{f}(\mathbf{x}(t))|$ converges to a constant vector (rotates infinite times) as $\mathbf{x}(t) \rightarrow \mathbf{p}$. We call an equilibrium a *saddle* if it has two outgoing and two incoming trajectories, which appear alternately. We give their phase portraits below.



The notions such as node, saddle and focus were introduced by H. Poincaré [2]. He gave a criterion which determines the shape of equilibrium. Let \mathbf{p} be an equilibrium of Eq.(1), and λ_1, λ_2 the eigenvalues of Jacobian matrix. He stated that when \mathbf{f} is a polynomial vector and $\lambda_1 \neq \lambda_2$, the following assertions hold:

- (i) If λ_1, λ_2 are real with the same sign, then \mathbf{p} is a node.
- (ii) If λ_1, λ_2 are real with opposite signs, then \mathbf{p} is a saddle.
- (iii) If λ_1, λ_2 are imaginary with nonzero real part, then \mathbf{p} is a focus.

In case of linear systems, we can prove his statement even when $\lambda_1 = \lambda_2$. The proof can be found in many textbooks. See [1, §20.4], for example.

In case of nonlinear systems, he did not provide any complete proofs. However, we can prove his statement for nonlinear C^1 systems.

Theorem 1. *Suppose that Eq.(1) is of class C^1 . If $\lambda_1 \neq \lambda_2$, then the assertions (i), (ii) and (iii) hold.*

The condition $\lambda_1 \neq \lambda_2$ is critical. In fact, if it is dropped, the following counter-example appears.

Example. *Consider a C^1 nonlinear system below:*

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ \varepsilon & -1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} - \frac{2}{\ln(x^2 + y^2)} \begin{bmatrix} -y \\ x \end{bmatrix}, \quad (2)$$

where $0 \leq \varepsilon < 2$. Then the equilibrium $\mathbf{0}$ is a focus.

The above example shows that if we drop the condition $\lambda_1 \neq \lambda_2$, Theorem 1 becomes false. However, such an example is very fragile. If we slightly strengthen the C^1 condition, Theorem 1 becomes true even when $\lambda_1 = \lambda_2$.

Theorem 2. *Suppose that Eq.(1) is of class $C^{1,\alpha}$ ($\alpha > 0$). Then the assertions (i), (ii) and (iii) hold.*

Here, we say that the system (1) is of class $C^{1,\alpha}$ if \mathbf{f} are of class C^1 and their first partial derivatives are α -Hölder continuous. The $C^{1,\alpha}$ condition ($0 < \alpha < 1$) is stronger than C^1 and weaker than C^2 .

2 Proofs.

Regarding the proof of Theorem 1, the assertion (ii) is guaranteed by Grobman-Hartman Theorem, which is found in [3, §5], for example. Unfortunately, the theorem can not distinguish (i) and (iii) because nodes and foci are topologically equivalent. See [1, §22.6]. Instead, we can prove (i) and (iii) by diagonalizing Jacobian matrix and by transforming it to a form of polar coordinates. We can carry out the proof without any difficulties. So we leave the proof to the reader.

In this section, we shall prove Example and Theorem 2.

Proof of Example. We transform Eq.(2) to a form of polar coordinates:

$$r'(t) = -r + \varepsilon r \cos \theta \sin \theta, \quad \theta'(t) = \varepsilon \cos^2 \theta - \frac{1}{\ln r}, \quad (3)$$

By evaluating the first of Eq.(3), we have that

$$-k_1 r \leq r'(t) \leq -k_2 r, \quad k_1 := 1 + \frac{1}{2}\varepsilon, \quad k_2 := 1 - \frac{1}{2}\varepsilon.$$

Take $r_0 = r(0)$ so small that $r_0 \in [0, 1)$. By integrating the above, we have that

$$r_0 e^{-k_1 t} \leq r(t) \leq r_0 e^{-k_2 t} < 1. \quad (4)$$

Therefore, $r(t) \rightarrow 0$ as $t \rightarrow \infty$. Hence $\mathbf{0}$ is attracting.

By taking logarithm of Eq.(4), we have that

$$-(k_1 t - \ln r_0) \leq \ln r(t) < 0. \quad (5)$$

By applying Eq.(5) to the second of Eq.(3), we have that

$$\theta'(t) \geq -\frac{1}{\ln r(t)} \geq \frac{1}{k_1 t - \ln r_0} > 0,$$

By integrating it, we have that

$$\theta(t) - \theta(0) \geq \frac{1}{k_1} \{ \ln(k_1 t - \ln r_0) - \ln(-\ln r_0) \}.$$

Therefore, $\theta(t) \rightarrow \infty$ as $t \rightarrow \infty$. Hence $\mathbf{0}$ is a focus. \square

Proof of Theorem 2. By translating \mathbf{p} to $\mathbf{0}$, we make Eq.(1) the following form:

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} u(x, y) \\ v(x, y) \end{bmatrix}, \quad (6)$$

where u, v are nonlinear terms. Remark that the square matrix of Eq.(6) is Jacobian matrix and denote it by J .

Consider only the case $\lambda_1 = \lambda_2$. This is because other cases are already established in Theorem 1. Put $\lambda = \lambda_1$, and assume that $\lambda < 0$. (It does not hurt the generality.) We divide the proof into two cases.

1. The case when J is diagonalizable. In such a case, there is a matrix P such that $P^{-1}JP = \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix}$. So we can transform Eq.(6) into the following form:

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} p(x, y) \\ q(x, y) \end{bmatrix}.$$

Also, we transform it to a form of polar coordinates:

$$r'(t) = \lambda r + r\varphi(r, \theta), \quad \theta'(t) = \psi(r, \theta), \quad (7)$$

Since both φ, ψ are α -Hölder continuous, there are reals $c > 0, m > 0$ such that

$$|\varphi(r, \theta)| < mr^\alpha, \quad |\psi(r, \theta)| < mr^\alpha \quad \text{on } [0, c] \times \mathbb{R}. \quad (8)$$

We take a real d so small that $md^\alpha < \frac{1}{2}|\lambda|$ and $d \leq c$. If we take $r(t) \in [0, d]$, then we evaluate the first of Eq.(7) as follows:

$$r'(t) < (-|\lambda| + \frac{1}{2}|\lambda|)r = -kr, \quad k := \frac{1}{2}|\lambda|.$$

If we take $r_0 = r(0)$ so small that $r_0 \in [0, d]$, then we have that

$$r(t) < r_0 e^{-kt}. \quad (9)$$

Therefore, $r(t) \rightarrow 0$ as $t \rightarrow \infty$. Hence \mathbf{p} is attracting.

By applying Eqs.(8), (9) to the second of Eq.(7), we have that

$$|\theta'(t)| < m(r_0 e^{-kt})^\alpha = mr_0^\alpha e^{-lt}, \quad l := \alpha k. \quad (10)$$

By integrating Eq.(10) from 0 to t ($0 \leq t$), we have that

$$|\theta(t) - \theta(0)| \leq -\frac{m}{l}r_0^\alpha(e^{-lt} - 1) < \frac{m}{l}r_0^\alpha.$$

Since $\theta(t)$ is bounded, there is a sequence $\{t_n\}$ diverging to ∞ such that $\theta(t_n)$ converges to a limit θ^* . By integrating Eq.(10) from t to t_n ($t \leq t_n$), we have that

$$|\theta(t_n) - \theta(t)| \leq -\frac{m}{l}r_0^\alpha(e^{-lt_n} - e^{-lt}) < \frac{m}{l}r_0^\alpha e^{-lt}.$$

By taking $n \rightarrow \infty$, we have that

$$|\theta^* - \theta(t)| \leq \frac{m}{l}r_0^\alpha e^{-lt}.$$

Therefore, $\theta(t)$ converges to θ^* . Hence \mathbf{p} is a node.

2. The case when J is not diagonalizable. In such a caes, there is a matrix P such that $P^{-1}JP = \begin{bmatrix} \lambda & 0 \\ \varepsilon & \lambda \end{bmatrix}$ (Jordan normal form). Put $\varepsilon = -\lambda$. So we can transform Eq.(6) into the following form:

$$\frac{d}{dt} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} \lambda & 0 \\ -\lambda & \lambda \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} p(x, y) \\ q(x, y) \end{bmatrix}.$$

Also, we transform it to a form of polar coordinates:

$$r'(t) = \lambda r - \lambda r \sin \theta \cos \theta + r\varphi(r, \theta), \quad \theta'(t) = -\lambda \cos^2 \theta + \psi(r, \theta). \quad (11)$$

Since both φ, ψ are α -Hölder continuous, there are reals $c > 0, m > 0$ such that

$$|\varphi(r, \theta)| < mr^\alpha, \quad |\psi(r, \theta)| < mr^\alpha \quad \text{on } [0, c] \times \mathbb{R}. \quad (12)$$

We take a real d so that $md^\alpha < \frac{1}{4}|\lambda|$ and $d \leq c$. If we take $r(t) \in [0, d]$, then we evaluate the first of Eq.(11) as follows:

$$r'(t) \leq (-|\lambda| + \frac{1}{2}|\lambda| + \frac{1}{4}|\lambda|)r = -kr, \quad k := \frac{1}{4}|\lambda|.$$

If we take $r_0 = r(0)$ so small that $r_0 \in [0, d]$, then we have that

$$r(t) \leq r_0 e^{-kt}. \quad (13)$$

Therefore, $r(t) \rightarrow 0$ as $t \rightarrow \infty$. Hence \mathbf{p} is attracting.

By applying Eqs.(12), (13) to the second of Eq.(11), we have that

$$\begin{aligned} a \cos^2 \theta - be^{-lt} &\leq \theta'(t) \leq a \cos^2 \theta + be^{-lt}, \\ a &:= |\lambda|, \quad b := mr_0^\alpha, \quad l := \alpha k. \end{aligned} \quad (14)$$

Take a real $t_0 \in [0, \infty)$ so large that $e^{lt_0} > 16ab/l^2$.

Firstly, consider the case when $h(t) < 0$ for every $t \in [t_0, \infty)$, where

$$h(t) = a \cos^2 \theta(t) - be^{-lt}.$$

In this case, we have that $\cos \theta(t) \rightarrow 0$ as $t \rightarrow \infty$. Therefore, there is an integer n such that $\theta(t) \rightarrow (n + \frac{1}{2})\pi$ as $t \rightarrow \infty$. Hence \mathbf{p} is a node.

Secondly, consider the case when $h(t_1) \geq 0$ for some $t_1 \in [t_0, \infty)$. In this case, we shall prove that $h(t) \geq 0$ for every $t \in [t_1, \infty)$. To prove it, assume that there is a real $t_2 \in (t_1, \infty)$ satisfying $h(t_2) < 0$.

Take the largest $s \in [t_1, t_2)$ satisfying $h(s) = 0$. This implies that $h(t) < 0$ on $(s, t_2]$. By the definition of derivative, we have that

$$h'(s) = \lim_{t \rightarrow s+0} \frac{h(t) - h(s)}{t - s} \leq 0. \quad (15)$$

By differentiating $h(t)$, and by putting $t = s$, we have that

$$h'(s) = -2a \cos \theta(s) \sin \theta(s) \theta'(s) + lbe^{-ls}. \quad (16)$$

By putting $h(s) = 0$ to Eq.(14), we have that

$$0 \leq \theta'(s) \leq a \cos^2 \theta(s) + be^{-ls}. \quad (17)$$

By using Eqs.(15), (16), we have that

$$\cos \theta(s) \sin \theta(s) \theta'(s) = \frac{1}{2a} \{lbe^{-ls} - h'(s)\} > 0. \quad (18)$$

By applying Eq.(17) to Eq.(18), we have that

$$\cos \theta(s) \sin \theta(s) > 0, \quad \theta'(s) > 0. \quad (19)$$

By applying Eqs.(17), (19) to Eq.(16), we have that

$$h'(s) \geq -2a \cos \theta(s) \sin \theta(s) (a \cos^2 \theta(s) + be^{-ls}) + lbe^{-ls}. \quad (20)$$

Since $a \cos^2 \theta(s) = be^{-ls}$, we have that

$$\begin{aligned} & (lbe^{-ls})^2 - \{2a \cos \theta(s) \sin \theta(s) (a \cos^2 \theta(s) + be^{-ls})\}^2 \\ &= (lbe^{-ls})^2 - 4a^2 \frac{be^{-ls}}{a} \left(1 - \frac{be^{-ls}}{a}\right) (be^{-ls} + be^{-ls})^2 \\ &= l^2 b^2 e^{-3ls} (e^{ls} - 16ab/l^2) + 16b^4 e^{-4ls}. \end{aligned} \quad (21)$$

Since $e^{ls} \geq e^{lt_0} > 16ab/l^2$, we have that Eq.(21) > 0 . By applying it to Eq.(20), we have that $h'(s) > 0$. It contradicts to Eq.(14). Therefore, we have that $h(t) \geq 0$ for every $t \in [t_1, \infty)$.

By using Eq.(14), we have that $\theta(t)$ is monotone increasing on $[t_1, \infty)$. Moreover, $\theta(t)$ can not cross $(n + \frac{1}{2})\pi$ for any integer n . This is because if $\theta(t_2) = (n + \frac{1}{2})\pi$ for some real $t_2 \in [t_1, \infty)$, then $h(t_2) = -be^{-kt_2} < 0$, a contradiction. Therefore, $\theta(t)$ is bounded. Since $\theta(t)$ is monotone increasing and bounded, it converges to a limit. Hence \mathbf{p} is a node. \square

References

- [1] Arnold VI. Ordinary Differential Equations, 3rd Edition. Springer; 1992. ISBN: 3540548130.
- [2] Poincaré H. Mémoire sur les courbes définies par une équation différentielle. J Math Pures Appl Ser 3. 1881; 7, 375–422.
- [3] Robinson C. Dynamical Systems: Stability, Symbolic Dynamics, and Chaos, 2nd Edition. CRC Press; 1998. ISBN: 0849384958.