BI-LIPSCHITZ ANSATZ FOR ANTI-SYMMETRIC FUNCTIONS

NADAV DYM, JIANFENG LU, AND MATAN MIZRACHI

ABSTRACT. Motivated by applications for simulating quantum many body functions, we propose a new universal ansatz for approximating anti-symmetric functions. The main advantage of this ansatz over previous alternatives is that it is bi-Lipschitz with respect to a naturally defined metric. As a result, we are able to obtain quantitative approximation results for approximation of Lipschitz continuous antisymmetric functions. Moreover, we provide preliminary experimental evidence to the improved performance of this ansatz for learning antisymmetric functions.

1. INTRODUCTION AND MAIN RESULTS

The search for an ansatz for quantum many-body wave functions dates back to the early days of quantum mechanics [Sla29], and has been a central task in quantum chemistry [SO96]. In recent years, it has received renewed excitement primarily due to the advances of neural network-based ansatz. While it is natural to utilize the powerful and universal representation power of neural networks to parameterize many-body wave functions, the symmetry requirements from quantum mechanics make the problem different from usual applications of neural networks in deep learning. In particular, for ferminionic systems, such as electrons, the Pauli's exclusion principle requires that the wave function is totally anti-symmetric, i.e., for any $x = (x_1, x_2, ..., x_n) \in \mathbb{R}^{d \times n}$ and $\sigma \in S_n$ the symmetric group on *n* letters, we have

(1)
$$\Psi(x) = (-1)^{\sigma} \Psi(\sigma x)$$

where $(-1)^{\sigma}$ indicates the sign of permutation σ and we denote $\sigma x = (x_{\sigma(1)}, x_{\sigma(2)}, \dots, x_{\sigma(n)})$. As a result, special neural newtork-based ansatz that respects the symmetry have to be developed.

Most ansatz currently used for anti-symmetric functions are built upon the backflow ansatz, dates back to [FC56], where the antisymmetry requirement is fulfilled using a determinant, and each component of the determinant is further parameterized as neural networks. Variational approaches based on such ansatz have achieved impressive empirical results, see e.g., [HZE19, PSMF20, HSN20, CMC20].

On the theoretical side, while it has been established that such ansatz is universal [HLL⁺22], the early approximation result requires linear combination of exponential number of determinants, thus suffering from the curse of dimensionality. Although it was argued in [Hut20] that only one determinant is enough, the ansatz constructed is discontinuous, and hence impractical to use for actual algorithms. Indeed, it was later proved in [HLL23] that within the polynomial

Date: March 7, 2025.

category, the representation based on backflow ansatz would require exponential number of determinants. Of course, this does not exclude the possibility of efficient representation by less regular ansatz. A more efficient ansatz was recently constructed by [CL23] and later improved by [YLG⁺24]. This ansatz is continuous and involves a combination of symmetric functions, with a polynomial number of anti-symmetric Vandermonde determinants of the form

(2)
$$f_{y_k}(x) = \prod_{1 \le i < j \le n} y_k^T (x_i - x_j), \quad k = 1, \dots, K,$$

where $K \ge nd + 1$ (the optimal cardinality is actually a bit lower, see [DLS25]).

A disadvantage of methods based on (2) is that they are very unstable: e.g., scaling *x* by t > 0 will lead to scaling f_{y_k} by $t^{\binom{n}{2}}$. This can be handled by straightforward normalization, as suggested in [YLG⁺24], but this solution leads to an ansatz which is not continuous by construction. To amend this, our goal is to construct alternative antisymmetric models which are stable, by addressing the following theoretical question:

Is it possible to efficiently represent antisymmetric functions using bi-Lipschitz continuous ansatz?

To formalize this goal, we will need to consider a notion of distance on $\mathbb{R}^{d \times n}$ which is appropriate for antisymmetric models. We note that any antisymmetric model is by construction invariant to the action of permutations whose sign is positive. The group of such permutations is known as the alternating group A_n . Accordingly, we will choose a semi-metric on $\mathbb{R}^{d \times n}$ which takes this symmetry into account, namely:

$$d_+(x, y) = \min_{\sigma \in A_n} ||x - \sigma y||_1$$

where recall that $\sigma x = (x_{\sigma(1)}, x_{\sigma(2)}, \dots, x_{\sigma(n)})$, and the 1-norm of a matrix *M* is $||M||_1 = \sum_{ij} |M_{ij}|$.

Our goal is to define a function $F : \mathbb{R}^{d \times n} \to \mathbb{R}^m$, for appropriate *m*, such that

- (1) **Invariance:** $F(x) = F(\sigma x)$ for all $\sigma \in A_n$ and $x \in \mathbb{R}^{d \times n}$.
- (2) **Bi-Lipschtiz** There exist positive c_1, c_2 such that for all $x, y \in \mathbb{R}^{d \times n}$,

$$c_1 d_+(x, y) \le ||F(x) - F(y)||_1 \le c_2 d_+(x, y)$$

Given such an A_n invariant function F, we can approximate a general anti-symmetric function by

(3)
$$h(x) = \frac{1}{2} (N \circ F(x) - N \circ F(\tau_0 x))$$

where τ_0 is a fixed permutation with negative sign, and *N* is a multilayer perceptron (MLP) or some other neural network architectures. By construction, such ansatz is anti-symmetric. Compared with [CL23], while the ansatz is similar, the advantage of the current construction is that *F* will be bi-Lipschitz. This is expected to lead to improved generalization capabilities and more stable learning. Indeed, for symmetric functions, the practical advantages of bi-Lipschitz symmetric models was demonstrated in [AD24, SDDA24, DD24].

An additional advantage of antisymmetric ansatz based on bi-Lipschitz models, is that they lead to quanitative approximation results:

Proposition 1. Let $K \subseteq \mathbb{R}^{d \times n}$ be a compact set which is A_n invariant. Assume that $F : K \to \mathbb{R}^m$ is bi-Lipschitz with constants (c_1, c_2) . Then there exist $C = C(c_1, c_2, n, d, K)$ such that for all $\varepsilon > 0$, and every antisymmetric function $f : \mathbb{R}^{d \times n} \to \mathbb{R}$ which is 1-Lipschitz, there exists a function h as in (3), defined by an MLP N with $\leq C\varepsilon^{-m/2}$ parameters, such that

$$\max_{x \in K} |f(x) - h(x)| < \epsilon$$

Proof. Fix an antisymmetric function $f : \mathbb{R}^{d \times n} \to \mathbb{R}$ that is 1-Lipschitz and anti-symmetric. Let us identify f and F with functions \hat{f} and \hat{F} defined on the quotient space K/A_n . By definition \hat{F} is bi-Lipschitz. The fact that f is both 1-Lipschitz and antisymmetric, implies that \hat{f} is also 1-Lipschitz. To see this, choose $x, y \in K$ and let $\tau \in A_n$ be the minimizer of $||x - \tau y||_1$. Then

$$\|\hat{f}([x]) - \hat{f}([y])\|_1 = \|f(x) - f(\tau y)\|_1 \le \|x - \tau y\|_1 = d_+(x, y),$$

where we used the notation [x] for the A_n orbit of $x \in \mathbb{R}^{d \times n}$. We now obtain

$$f(x) = \hat{f}([x]) = \hat{f} \circ \hat{F}^{-1} \circ \hat{F}([x]) = \hat{f} \circ \hat{F}^{-1} \circ F(x), \quad \forall x \in K$$

The function $\hat{f} \circ \hat{F}^{-1} : F(K) \subseteq \mathbb{R}^m \to \mathbb{R}$ is Lipschitz, since \hat{f} is Lipschitz and \hat{F} is bi-Lipschitz, with a Lipschitz constant of c_1^{-1} . Since F is continuous and K is compact, F(K) is compact. By rescaling f(x) using positive α, β to a function of the form $\alpha f(\beta x)$, we can assume without loss of generality that $f \circ F^{-1}$ is 1-Lipschitz, and F(K) is contained in the unit cube $[0,1]^m$. Finally, by the Kirszbraun theorem, we can extend $f \circ F^{-1}$ to a 1-Lipschitz function on all of $[0,1]^m$. It is then known [Yar18] that the function can be approximated by an MLP N with $\sim \epsilon^{-m/2}$ parameters to obtain an ϵ approximation. We then have for all $x \in K$,

$$|f(x) - N \circ F(x)| = |\hat{f} \circ \hat{F}^{-1} \circ F(x) - N \circ F(x)| < \epsilon,$$

and therefore

$$|f(x) - h(x)| = \frac{1}{2} |(f(x) - f(\tau_0 x) - N \circ F(x) + N \circ F(\tau_0 x))|$$

$$\leq \frac{1}{2} |f(x) - N \circ F(x)| + \frac{1}{2} |f(\tau_0 x) - N \circ F(\tau_0 x)| < \epsilon.$$

Remark: We note that the size of the network in this bound depends exponentially on *m*. It is known [AGA⁺24, JBM⁺23] that *m* is lower bounded via $m \ge n \cdot d$, which is the dimension of the space. In our construction, we can guarantee bi-Lipschitz-ness with m = 2nd + 1, as stated in Theorem 3, which gives an approximation rate of $\sim e^{-(nd+1/2)}$. moreover, if the domain *K* has low intrinsic dimension: that is, it is a semialgebraic subset of dimension $\ll d \cdot n$, then we can take $m = 2 \dim(K) + 1$ and the approximation rate will improve accordingly. Finally, we note that the intrinsic dimension of F(K) is $\leq \dim(K)$. Based on this observation, we believe it should ultimately be possible to acheive an approximation rate with exponent proportional to $\dim(K)/2$ rather than $(2\dim(K) + 1)/2$. However, while approximation results for ReLU networks, which depend on the intinsic dimension of a 'data manifold' are available [SCC18, CJLZ19, NI20], F(K) will generally not be a manifold, leading to technical difficulties which we will not address in this paper, as this paper focuses mostly on finding a bi-Lipschitz construction.

We now present the construction of our bi-Lipschitz *F*. We will first in Section 2 give the construction for d = 1 to present the key idea in a simple setting. In this setting, the upper and lower Lipschitz constants are 1,2 respectively. We then generalize our construction to d > 1 in Section 3. In this setting we provide a bi-Lipschitz *F* but do not specify estimates for the bi-Lipschitz constants c_1, c_2 .

2. CONSTRUCTION FOR d = 1

For d = 1, our construction is based on two components. The first is a *symmetric* bi-Lipschitz function. That is, a symmetric function which is bi-Lipschitz with respect to the metric obtained by quotienting over the group S_n of all permutations, namely

(4)
$$\mathbf{d}_{\pm}(x, y) = \min_{\sigma \in S_n} \|x - \sigma y\|_1$$

As shown in [AGA⁺24], commonly used symmetric models based on summation, as suggested in Deepsets [ZKR⁺17], are not bi-Lipschitz. In contrast, following [BHS22, DD24] we choose the symmetric function sort : $\mathbb{R}^n \to \mathbb{R}^n$ that maps each vector x to the unique vector $\hat{x} = \text{sort}(x)$ satisfying that $\hat{x}_i \leq \hat{x}_{i+1}$ for all i = 1, ..., n - 1.¹ We note that while \hat{x} is uniquely defined, the permutation taking x to \hat{x} is not uniquely defined if x is in the set

$$\Omega := \{ x \in \mathbb{R}^n \mid x_i = x_j \text{ for some } 1 \le i < j \le n \}.$$

In this case, there will be permutations of both positive and negative signs which will take *x* to \hat{x} . However, if *x* is not in Ω , the permutation sorting *x* is unique, and we denote it by τ_x .

The sort function is continuous and piecewise linear. It is not only bi-Lipschitz but it is also an *isometry*, namely

$$\| \operatorname{sort}(x) - \operatorname{sort}(y) \|_1 = d_{\pm}(x, y).$$

The second component of our construction is the function $Q: \mathbb{R}^n \to \mathbb{R}$ defined as

$$Q(x) = \left(\prod_{1 \le i < j \le n} \operatorname{sign}(x_j - x_i)\right) \cdot \min_{1 \le i < j \le n} |x_j - x_i|$$

This function is a continuous piecewise linear function, and it is antisymmetric. Morever we note that Q(x) = 0 if and only if x is in the set Ω . Next, we note that if $\hat{x} = \operatorname{sort}(x)$, then the sign of Q(x) will be positive. These observations give us an efficient method for computing Q: for every $x \in \mathbb{R}^n$, if x is in Ω then Q(x) = 0. Otherwise, there is a unique permutation, which we denote as before by τ_x , such that $\hat{x} = \operatorname{sort}(x) = \tau_x x$. We then deduce that

$$Q(x) = (-1)^{\tau_x} Q(\tau_x x) = (-1)^{\tau_x} \min_{1 \le i < j \le n} |\hat{x}_j - \hat{x}_i| = (-1)^{\tau_x} \min_{1 \le j \le n} (\hat{x}_{j+1} - \hat{x}_j).$$

It follows that Q(x) can be computed by first sorting *x*, and then applying the formula above which requires only O(n) operations. The total complexity of this procedure is then $O(n \log n)$.

¹The idea of sorting has also been used in the anti-symmetric construction in [Hut20] in one dimension.

Finally, we combine the two components of our construction to obtain the function $F : \mathbb{R}^n \to \mathbb{R}^{n+1}$ defined by

(5)
$$F(x_1,...,x_n) = [\operatorname{sort}(x_1,...,x_n);Q(x)],$$

and we prove that this function is bi-Lipschitz with explicit, very moderate constants:

Theorem 2. F is bi-Lipschitz, in the sense that

$$d_{+}(x, y) \le ||F(x) - F(y)|| \le 2d_{+}(x, y), \quad \forall x, y \in \mathbb{R}^{n}$$

Proof. We first note that for all $x, y \in \mathbb{R}^n$, if τ_x, τ_y are permutations satisfying

$$\hat{x} = \operatorname{sort}(x) = \tau_x x, \quad \hat{y} = \operatorname{sort}(y) = \tau_y y,$$

then

$$\|F(x) - F(y)\|_{1} = \|\operatorname{sort}(x) - \operatorname{sort}(y)\|_{1} + |Q(x) - Q(y)|$$

= $d_{\pm}(x, y) + |(-1)^{\tau_{x}}Q(\hat{x}) - (-1)^{\tau_{y}}Q(\hat{y})|$
= $d_{\pm}(x, y) + |(-1)^{\tau_{x}}(-1)^{\tau_{y}}Q(\hat{x}) - Q(\hat{y})|$

We now divide into two cases:

case 1: If τ_x and τ_y can be chosen so that they have the same sign. Note this case includes the case where either *x* or *y* are in Ω . In this case we have that

$$d_{\pm}(x, y) = d_{\pm}(x, y) \le |F(x) - F(y)|$$

In the other direction, if we assume without loss of generality that $Q(\hat{x}) > Q(\hat{y})$ then we obtain

$$\|F(x) - F(y)\|_1 = \mathbf{d}_{\pm}(x, y) + |(-1)^{\tau_x} (-1)^{\tau_y} Q(\hat{x}) - Q(\hat{y})| = \mathbf{d}_{+}(x, y) + Q(\hat{x}) - Q(\hat{y}).$$

It is thus sufficient to show that $Q(\hat{x}) - Q(\hat{y}) < d_+(x, y)$. Indeed, if we choose *s* for which

$$Q(\hat{y}) = \min_{1 \le j \le n} (\hat{y}_{j+1} - \hat{y}_j) = y_{s+1} - y_s.$$

Then

$$Q(\hat{x}) - Q(\hat{y}) = \min_{1 \le j \le n} (\hat{x}_{j+1} - \hat{x}_j) - (\hat{y}_{s+1} - \hat{y}_s)$$

$$\le \hat{x}_{s+1} - \hat{x}_s - (\hat{y}_{s+1} - \hat{y}_s)$$

$$= \hat{x}_{s+1} - \hat{y}_{s+1} + \hat{y}_s - \hat{x}_s$$

$$\le \|\hat{x} - \hat{y}\|_1 = \mathbf{d}_{\pm}(x, y) = \mathbf{d}_{+}(x, y).$$

case 2: We now consider the case where τ_x and τ_y cannot be chosen to have the same sign. In this case

$$\|F(x) - F(y)\|_1 = d_{\pm}(x, y) + Q(\hat{x}) + Q(\hat{y}) \le d_{+}(x, y) + Q(\hat{x}) + Q(\hat{y}),$$

thus to obtain the required upper Lipschitz constant of 2 it is sufficient to show that $Q(\hat{x}) + Q(\hat{y}) \le d_+(x, y)$.

There exists a permutation τ with a negative sign such that

$$\mathbf{d}_{+}(x, y) = \|\hat{x} - \tau \,\hat{y}\|_{1}.$$

Since \hat{y} is sorted and has no repeated entries, $\tau \hat{y}$ is not sorted, and so necessarily there is some index *i* such that $(\hat{y})_{\tau(i+1)} - (\hat{y})_{\tau(i)} < 0$. Thus

$$Q(\hat{x}) + Q(\hat{y}) \le \hat{x}_{i+1} - \hat{x}_i + (\hat{y})_{\tau(i)} - (\hat{y})_{\tau(i+1)}$$
$$\le ||x - \tau y||_1 = \mathbf{d}_+(x, y)$$

For the lower Lipschitz bound, let *i*, *j* be the indices such that

$$Q(\hat{x}) = \hat{x}_{i+1} - \hat{x}_i, \quad Q(\hat{y}) = \hat{y}_{j+1} - \hat{y}_j$$

and let τ_i denote the permutation (i, i+1) which swaps i with i+1. Similarly denote $\tau_j = (j, j+1)$. Then

$$\begin{aligned} 2\mathbf{d}_{+}(x,y) &\leq \|\tau_{i}\hat{x} - \hat{y}\|_{1} + \|\hat{x} - \tau_{j}\hat{y}\|_{1} \\ &\leq \|\tau_{i}\hat{x} - \hat{x}\|_{1} + 2\|\hat{x} - \hat{y}\|_{1} + \|\hat{y} - \tau_{j}\hat{y}\|_{1} \\ &= 2\mathbf{d}_{\pm}(x,y) + 2(\hat{x}_{i+1} - \hat{x}_{i}) + 2(\hat{y}_{j+1} - \hat{y}_{j}) \\ &= 2\left[\mathbf{d}_{\pm}(x,y) + Q(\hat{x}) + Q(\hat{y})\right] = 2\|F(x) - F(y)\|_{1}. \end{aligned}$$

This concludes the proof.

3. The construction for general d

Next we consider the construction of bi-Lipschitz, A_n invariant function on $\mathbb{R}^{d \times n}$, when d > 1. We consider parametric functions of the form

$$\psi(x_1,\ldots,x_n;a,b) = b \cdot F(a \cdot x_1,\ldots,a \cdot x_n)$$

where $a \in \mathbb{R}^d$, $b \in \mathbb{R}^{n+1}$, the notation $x \cdot y$ denotes the standard inner product between x and y, and F is the function defined in (5) for the d = 1 construction. We claim that if we clone F enough times, we will get an injective function (up to A_n symmetries), which will also be bi-Lipschtiz with respect to d_+ .

Theorem 3. Set m = 2nd + 1. For Lebesgue almost every $a_1, ..., a_m \in \mathbb{R}^d$ and $b_1, ..., b_m \in \mathbb{R}^{n+1}$ the function

$$\Psi(x_1,...,x_n|a_1,...,a_m,b_1,...,b_m) = \begin{pmatrix} \psi(x_1,...,x_n;a_1,b_1) \\ \vdots \\ \psi(x_1,...,x_n;a_m,b_m) \end{pmatrix}$$

is one-to-one (up to A_n symmetries) and bi-Lipschitz.

Proof. We first show injectivity, up to A_n symmetries. We will use Theorem 2.7 in [DG24] which guarantees injectivity (separation in the language of that paper) using m = 2nd + 1 generic vectors, under the condition that ψ is *strongly separating*. This means that, for every x_1, \ldots, x_n and y_1, \ldots, y_n which are not related by an A_n symmetry, we have that

$$\psi(x_1,\ldots,x_n;a,b)\neq\psi(y_1,\ldots,y_n;a,b)$$

for Lebesgue almost every *a*, *b*. Indeed, fix some $x_1, ..., x_n$ and $y_1, ..., y_n$ which are not related by an A_n symmetry. For almost all *a*, we have that the inner product of *a* with any two distinct elements in the set $\{x_1, ..., x_n, y_1, ..., y_n\}$ will give two distinct outputs. For all such *a*, if $x_1, ..., x_n$ and $y_1, ..., y_n$ are not related by a permutation then sort $(a \cdot x_1, ..., a \cdot x_n) \neq$ sort $(a \cdot y_1, ..., a \cdot y_n)$, and if $x_1, ..., x_n$ and $y_1, ..., y_n$ are related by a permutation, but not by a permutation with positive sign, then we necessarily know that all elements in both sets are distinct, and moreover that

$$\frac{Q(a \cdot x_1, \dots, a \cdot x_n)}{Q(a \cdot y_1, \dots, a \cdot y_n)} < 0$$

so that in particular they are not equal. We have shown that for almost every *a* we have that

$$F(a \cdot x_1, \ldots, a \cdot x_n) \neq F(a \cdot y_1, \ldots, a \cdot y_n).$$

Next, we note that once we fix any *a* fulfilling this inequality, we will have for almost every $b \in \mathbb{R}^{n+1}$ that

$$b \cdot F(a \cdot x_1, \dots, a \cdot x_n) \neq b \cdot F(a \cdot y_1, \dots, a \cdot y_n).$$

This concludes the proof of strong separation, and hence of injectivity of *F*.

Next, we prove bi-Lipschitzness. For this, we will use Lemma 3.4 in [SDDA24]. This lemma considers two piecewise linear non-negative functions which have the same zero sets, in our case these functions are

$$d_+(X, Y)$$
 and $\|\Psi(X; a_1, \dots, a_m, b_1, \dots, b_m) - \Psi(Y; a_1, \dots, a_m, b_1, \dots, b_m)\|_1$

where the parameter of *F* are such that *F* is injective, so that the two functions have the same zero set. The lemma says that these two functions will be bi-Lipschitz equivalent on a fixed compact polyhedral domain, which in our case we will take to be the unit ball $\{(X, Y)| \|X\|_1 + \|Y\|_1 \le 1\}$. Homogeneity of d₊ and Ψ then implies bi-Lipschitzness on the whole domain.

Remark: The cardinality of *m* chosen here can probably be improved a bit, e.g., using the scale and translation invariance of the question at hand. Moreover, if we assume that *X*, *Y* come from some semi-algebraic subset of $\mathbb{R}^{d \times n}$ of dimension *v*, then we can get a bound of m = 2v + 1 essentially automatically.

4. EXPERIMENTS

To experimentally evaluate the efficiency of our bi-Lipschitz continuous ansatz, we compare it against the Vandermonde determinant (2) based approach from [YLG⁺24], and a naive Multi-Layer-Perceptron (MLP) estimator which does not enforce antisymmetry. We use these different models to address a simple antisymmetric regression task; approximating the antisymmetric function $f(A) = \det A : \mathbb{R}^{n \times n} \to \mathbb{R}$. 4.1. **Setup.** We generated 3 training sets of 110,00 matrices of orders $n \in \{10, 15, 20\}$, with a corresponding validation set of 15,000 matrices and a test set of 20,000 matrices, all uniformly sampled from $[0, 1.1]^{n \times n}$. We optimize the parameters with respect to the $\|\cdot\|_{l_1}$ metric with ADAM, using the same learning rate and the same learning rate scheduler settings for all methods. We implemented the symmetric functions required for the ansatz [YLG⁺24] (which we refer to as $\tilde{O}(n^2)$ model) using the Deep Sets model [ZKR⁺17], and sampled $n^2 + 1$ constant vectors from S^{n-1} for the Vandermonde determinant computation.

	<i>n</i> = 10	<i>n</i> = 15	<i>n</i> = 20	
Ansatz				
Bi-Lipschitz (ours)	79,741	121,501	275,121	
MLP	223,489	78,593	993,793	
$\widetilde{O}(n^2)$ [YLG ⁺ 24]	704,677	1,649,122	3,054,417	

TABLE 1. Ansatz's number of parameters

For the MLP method we used the same depth as the MLP used within in our construction 3, with a width similar to, or wider than, our hidden layers. Note that the MLP ansatz does not respect the anti-symmetry. The number of parameters in each model is summarized in Table 1.

4.2. **Results.** The accuracy of the different models is shown in Table 2 with respect to both the Mean Absolute Error (which was used for training) and the Mean Absolute Relative Error. In Figure 1 we show the convergence rates of the different methods for this experiment. For all values of *n* tried, our model was more accurate than the competing models, sometimes by a large margin. Additionally, convergence of our method was much faster. The results also show that for n = 15,20, the performance of the MLP is substantially lower than the other, antisymmetric models, indicating the importance of imposing the antisymmetric structure.



FIGURE 1. Training-Validation convergence comparison of our Ansatz (black), against [YLG⁺24]'s (red) and an MLP (green), for approximating the determinant of matrices of different orders (denoted by n). Our ansatz exhibits faster convergence and better or comparable accuracy.

BI-LIPSCHITZ ANSATZ FOR ANTI-SYMMETRIC FUNCTIONS

	<i>n</i> = 10		<i>n</i> = 15		<i>n</i> = 20	
	MAE	MARE	MAE	MARE	MAE	MARE
Ansatz						
Bi-Lipschitz (ours)	0.063375	0.000500	0.140079	0.000515	0.662697	0.000501
MLP	0.067679	0.000534	0.576995	0.002123	538.833417	0.407969
$\widetilde{O}(n^2)$ [YLG ⁺ 24]	0.066667	0.000526	0.142137	0.000523	2.004496	0.001517

TABLE 2. Mean Absolute Error (MAE) and Mean Absolute Relative Error (MARE) evaluation of the ansatzs, on the test sets, with n being the order of the matrices evaluated. Best of each measurement is in bold.

Acknowledgements N.D. and M.M. acknowledge the support of ISF grant 272/23. The work of J.L. is supported in part by the National Science Foundation via awards DMS-2309378 and IIS-2403276.

REFERENCES

- [AD24] Tal Amir and Nadav Dym, Fourier sliced-wasserstein embedding for weighted sets and point clouds, arXiv preprint arXiv:2405.16519 (2024).
- [AGA⁺24] Tal Amir, Steven Gortler, Ilai Avni, Ravina Ravina, and Nadav Dym, Neural injective functions for multisets, measures and graphs via a finite witness theorem, Advances in Neural Information Processing Systems 36 (2024).
- [BHS22] Radu Balan, Naveed Haghani, and Maneesh Singh, *Permutation invariant representations with applications to graph deep learning*, 2022.
- [CJLZ19] Minshuo Chen, Haoming Jiang, Wenjing Liao, and Tuo Zhao, *Efficient approximation of deep relu networks for functions on low dimensional manifolds*, Advances in neural information processing systems **32** (2019).
- [CL23] Ziang Chen and Jianfeng Lu, *Exact and efficient representation of totally anti-symmetric functions*, arXiv preprint arXiv:2311.05064 (2023).
- [CMC20] Kenny Choo, Antonio Mezzacapo, and Giuseppe Carleo, *Fermionic neural-network states for ab-initio electronic structure*, Nature communications **11** (2020), no. 1, 2368.
- [DD24] Yair Davidson and Nadav Dym, On the Hölder stability of multiset and graph neural networks, arXiv preprint arXiv:2406.06984 (2024).
- [DG24] Nadav Dym and Steven J Gortler, *Low-dimensional invariant embeddings for universal geometric learning*, Foundations of Computational Mathematics (2024), 1–41.
- [DLS25] Nadav Dym, Hannah Lawrence, and Jonathan W Siegel, *Equivariant frames and the impossibility of continuous canonicalization*, Proceedings of the 41st international conference on machine learning, 2025.
- [FC56] RP Feynman and Michael Cohen, Energy spectrum of the excitations in liquid helium, Physical Review 102 (1956), no. 5, 1189.
- [HLL23] Hang Huang, Joseph M Landsberg, and Jianfeng Lu, *Geometry of backflow transformation ansatze for quantum many-body fermionic wavefunctions*, Communications in Mathematical Sciences **21** (2023), no. 5, 1447–1453.
- [HLL⁺22] Jiequn Han, Yingzhou Li, Lin Lin, Jianfeng Lu, Jiefu Zhang, and Linfeng Zhang, Universal approximation of symmetric and anti-symmetric functions, Communications in Mathematical Sciences 20 (2022), no. 5, 1397– 1408.
- [HSN20] Jan Hermann, Zeno Schätzle, and Frank Noé, Deep-neural-network solution of the electronic schrödinger equation, Nature Chemistry 12 (2020), no. 10, 891–897.

NADAV DYM, JIANFENG LU, AND MATAN MIZRACHI

[Hut20] Marcus Hutter, On representing (anti) symmetric functions, arXiv preprint arXiv:2007.15298 (2020).

- [HZE19] Jiequn Han, Linfeng Zhang, and Weinan E, Solving many-electron schrödinger equation using deep neural networks, Journal of Computational Physics 399 (2019), 108929.
- [JBM⁺23] Chaitanya K. Joshi, Cristian Bodnar, Simon V. Mathis, Taco Cohen, and Pietro Liò, On the expressive power of geometric graph neural networks, Proceedings of the 40th international conference on machine learning, 2023.
 - [NI20] Ryumei Nakada and Masaaki Imaizumi, Adaptive approximation and generalization of deep neural network with intrinsic dimensionality, Journal of Machine Learning Research 21 (2020), no. 174, 1–38.
- [PSMF20] David Pfau, James S Spencer, Alexander GDG Matthews, and W Matthew C Foulkes, Ab initio solution of the many-electron schrödinger equation with deep neural networks, Physical Review Research 2 (2020), no. 3, 033429.
- [SCC18] Uri Shaham, Alexander Cloninger, and Ronald R Coifman, *Provable approximation properties for deep neural networks*, Applied and Computational Harmonic Analysis **44** (2018), no. 3, 537–557.
- [SDDA24] Yonatan Sverdlov, Yair Davidson, Nadav Dym, and Tal Amir, *Fsw-gnn: A bi-lipschitz wl-equivalent graph neural network*, 2024.
 - [Sla29] John C Slater, The theory of complex spectra, Physical review 34 (1929), no. 10, 1293.
 - [SO96] Attila Szabo and Neil S Ostlund, *Modern quantum chemistry: introduction to advanced electronic structure theory*, Courier Corporation, 1996.
 - [Yar18] Dmitry Yarotsky, Optimal approximation of continuous functions by very deep relu networks, Conference on learning theory, 2018, pp. 639–649.
- [YLG⁺24] Haotian Ye, Ruichen Li, Yuntian Gu, Yiping Lu, Di He, and Liwei Wang, $O(N^2)$ representation of general continuous anti-symmetric function, arXiv preprint arXiv:2402.15167 (2024).
- [ZKR⁺17] Manzil Zaheer, Satwik Kottur, Siamak Ravanbakhsh, Barnabas Poczos, Russ R Salakhutdinov, and Alexander J Smola, *Deep sets*, Advances in neural information processing systems **30** (2017).

FACULY OF MATHEMATICS, FACULTY OF COMPUTER SCIENCE, TECHNION- ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA, ISRAEL

Email address: nadavdym@technion.ac.il

DEPARTMENT OF MATHEMATICS, DEPARTMENT OF PHYSICS, AND DEPARTMENT OF CHEMISTRY, DUKE UNIVERSITY, DURHAM, NC 27708 USA

Email address: jianfeng@math.duke.edu

FACULTY OF MATHEMATICS, TECHNION- ISRAEL INSTITUTE OF TECHNOLOGY, HAIFA, ISRAEL *Email address*: mmizrachi@campus.technion.ac.il

10