

# The excenters of bicentric polygons are concyclic

Norbert Hungerbühler<sup>1</sup>, Clemens Pohle<sup>2</sup>, and Yun Zhang<sup>3</sup>

<sup>1</sup>Department of Mathematics, ETH, Zürich, Switzerland

<sup>2</sup>Kantonsschule Hohe Promenade, Zürich, Switzerland

<sup>3</sup>Xi'an, Shaanxi Province, China

March 10, 2025

## Abstract

We show that the centers of the excircles of a bicentric polygon  $B$  are concyclic on a circle  $E$ . The center of the circumscribed circle  $K$  of  $B$  is the midpoint of the center of  $E$  and the center of the inscribed circle  $C$  of  $B$ . The radius of  $E$  is given by a simple formula in terms of the radii of  $C$  and  $K$  and the distance between their centers.

**Key words.** bicentric polygons, excircles, Feuerbach's nine-point circle, Euler line.

**Mathematics Subject Classification.** 51M04, 51M15

## 1 Introduction

A polygon that is inscribed in a circle  $K$  and circumscribed around a circle  $C$  is called a bicentric polygon. Clearly all triangles are bicentric. If  $R_K$  and  $R_C$  are the radii of the circumcircle  $K$  and incircle  $C$  of a triangle, and  $d$  the distance between their centers, then the Chapple-Euler relation

$$\frac{1}{R_K - d} + \frac{1}{R_K + d} = \frac{1}{R_C} \quad (1)$$

holds (see [8] and [1]). A convex quadrilateral is bicentric if it satisfies Fuss's condition

$$\frac{1}{(R_K - d)^2} + \frac{1}{(R_K + d)^2} = \frac{1}{R_C^2} \quad (2)$$

(see, e.g., [6, Problem 39]). Fuss [10], Euler [8], and Steiner [21] found formulas relating the radii of  $C$  and  $K$  and the distance between their centers such that the circles carry a bicentric polygon for up to 10 vertices. Later, Jacobi [14], Richelot [19] and Kerawala [16] used elliptic functions to solve the problem for a general number of vertices. Also note that the circles  $K$  and  $C$  of a bicentric polygon form a Poncelet pair, i.e., every point of  $K$  is the vertex of a bicentric polygon inscribed in  $K$  and circumscribed around  $C$ . Poncelet's porism holds more generally for pairs of conics, not necessarily disjoint and possibly nested (see, e.g., [12] for a simple proof, and [7, 9] for a comprehensive treatment of Poncelet's porism). Bicentric polygons have been studied from various viewpoints, we refer the interested reader

---

<sup>1</sup>norbert.hungerbuehler@math.ethz.ch

<sup>2</sup>clemens.pohle@kshp.ch

<sup>3</sup>yunzhangmath@126.com

to the respective literature: see [18] and the references therein, or [2, 20, 11]. In this article we would like to add a geometric property of bicentric polygons that seems to be new.

It is well known that the centers of the excircles of a convex quadrilateral are concyclic. See Figure 1 for a short proof. By an excircle of a convex polygon we mean a circle that is tangent to one side of the polygon and the extensions of the two adjacent sides<sup>1</sup>. Below, we will extend the notion of excircles to general, non-convex and even self-intersecting polygons. This is necessary so that we can state our main Theorem 1 in its full generality. Notice that the centers of the excircles of a convex polygon with more than four vertices are in general not concyclic. However, it turns out that bicentric polygons have exactly this property (see Theorem 1 in Section 2, and Figure 3 for the case of a bicyclic pentagon). In Section 3, we examine the case of bicentric quadrilaterals more closely. And in Section 4, we show that the ratio of the areas of the bicentric polygon  $B$  and the polygon  $P$  given by the centers of the excircles equals the ratio of the radii of the incircle of  $B$  and the circumcircle of  $P$ .




Figure 1: The centers of the orange excircles of a convex quadrilateral  $A_1A_2A_3A_4$  are concyclic. The thin black lines are the external angle bisectors of the quadrilateral. Their intersections, the points  $M_1, M_2, M_3, M_4$ , are the centers of the excircles. The excircle with center  $M_i$  touches the side  $a_i$  and the extended sides  $a_{i-1}$  and  $a_{i+1}$ . The sum of opposite angles, e.g. in  $M_1$  and  $M_3$ , equals  $\pi$ . Hence,  $M_1, M_2, M_3, M_4$  lie on a circle.

## 2 Main result and proof

Before we state the main result it is necessary to fix the general framework. For the definition of a polygon, we follow [3, §3.1]: For  $n \geq 3$ , a polygon or  $n$ -gon consists of  $n$  points (the vertices)  $A_1, A_2, \dots, A_n$  and  $n$  straight lines (or sides)  $a_1, a_2, \dots, a_n$  such that for each  $i \in \{1, \dots, n\}$  we have

<sup>1</sup>Sometimes such circles are called escribed circles.

- $A_i$  and  $A_{i+1}$  are distinct and lie on  $a_i$ ,
- $a_i$  and  $a_{i+1}$  are distinct and meet in  $A_{i+1}$ .

Here and in the sequel indices are always read cyclically. Note that we allow our polygons to be non-convex and even self-intersecting, and that the sides are extended straight lines, not only the segments between two vertices. The polygon is called cyclic if all its vertices lie on a circle (the circumcircle)  $K$ . The polygon has an inscribed circle  $C$  if all its sides are tangent to  $C$ . The polygon is bicyclic, if it has both a circumcircle and an inscribed circle. In every vertex of a bicyclic polygon, we have two orthogonal angle bisectors of the sides meeting in that vertex. One of them is incident with the center of the inscribed circle and is called internal angle bisector, the other one is called external angle bisector. The external angle bisectors in adjacent points  $A_i, A_{i+1}$  meet in the center  $M_i$  of the excircle touching the sides  $a_{i-1}, a_i$ , and  $a_{i+1}$  (see Figure 2).




Figure 2: Internal angle bisectors (blue) meeting in the center  $M_C$  of the inscribed circle  $C$ . External angle bisectors (black) meeting in the center  $M_i$  of the excircle (orange) that touches the sides  $a_{i-1}, a_i, a_{i+1}$ .

Now we can state the main result.

**Theorem 1.** *Let  $B$  be a bicentric polygon, inscribed in a circle  $K$  and circumscribed around a circle  $C$ . Then the centers of the excircles of  $B$  are concyclic on a circle  $E$ . The center of  $K$  is the midpoint of the centers of  $C$  and  $E$ . The radius  $R_E$  of  $E$  is given by*

$$R_E = \frac{|R_K^2 - d^2|}{R_C}, \quad (3)$$

where  $R_K, R_C$  are the radii of  $K$  and  $C$ , respectively, and  $d$  is the distance between their centers.

In the simplest case, consider a triangle  $A_1A_2A_3$  with incircle  $C$ , circumcircle  $K$ , and centers  $M_1, M_2, M_3$  of its excircles. Then, the center of  $C$  is the orthocenter of the triangle  $M_1M_2M_3$ , and the center of  $K$  is the center of Feuerbach's nine-point circle of the triangle  $M_1M_2M_3$ . Hence the theorem reduces in this case to the well known properties of the Euler line. In particular, combining (3) and (1) we get

$$\begin{aligned} R_E^2 &\stackrel{(3)}{=} \frac{(R_K^2 - d^2)^2}{R_C^2} \stackrel{(1)}{=} \\ &= \frac{(R_K^2 - d^2)^2 (2R_K)^2}{(R_K - d)^2 (R_K + d)^2} = 4R_K^2. \end{aligned}$$




Figure 3: Illustration for the theorem. The centers of the orange **excircles** of the brown **bicentric pentagon** are concyclic on the red circle  $E$ . The center  $M_K$  of the circumcircle  $K$  is the midpoint of the center  $M_E$  of  $E$  and the center  $M_C$  of the inscribed circle  $C$ .

This is the fact that the radius of the Feuerbach circle is half the radius of the circumcircle of a triangle.

Similarly, for convex quadrilaterals, we get by (3) and (2) a simple formula which directly links  $R_E$  to  $R_K$  and  $d$ :

$$\begin{aligned}
 R_E^2 &\stackrel{(3)}{=} \frac{(R_K^2 - d^2)^2}{R_C^2} \stackrel{(2)}{=} \\
 &= \frac{(R_K^2 - d^2)^2 (2R_K^2 + 2d^2)}{(R_K - d)^2 (R_K + d)^2} = 2(R_K^2 + d^2). \tag{4}
 \end{aligned}$$

The case of a bicentric convex quadrilateral will be addressed again in Section 3.

In the proof of Theorem 1 we need the following lemma which is of interest in itself.

**Lemma 1.** *Let  $C$  be a circle with radius  $R_C$  and center  $M_C$ ,  $K$  a circle with radius  $R_K$  and center  $M_K$ , and  $d$  the distance between  $M_C$  and  $M_K$ . Let  $t$  be a tangent to  $C$  which intersects  $K$  in the points  $P_1, P_2$ . Then the center  $M_D$  of the circumcircle  $D$  of the triangle  $M_C P_1 P_2$  lies on a circle  $F$  with center  $M_K$  and radius*

$$R_F = \frac{|R_K^2 - d^2|}{2R_C}, \tag{5}$$

*independent of  $t$  (see Figure 4).*




Figure 4: Illustration for the lemma.

*Remark.* If  $d < R_K = R_F$  in (5) then  $C$  and  $F$  form a Poncelet pair for triangles. This follows by comparing (1) and (5).

*Proof of Lemma 1.* We assume that  $d < R_K$  in the following arguments. The case  $d > R_K$  is similar, and the case  $d = R_K$  is trivial.

Let  $Q$  be the intersection of the line  $M_D M_K$  with  $t$ , and  $R_D$  the radius of  $D$ . In the right triangles  $P_1 Q M_D$  (used in the first line) and  $P_1 Q M_K$  (used in the second line) we have

$$\begin{aligned}
 R_D^2 &= |P_1 M_D|^2 = |P_1 Q|^2 + |M_D Q|^2 \\
 &= (R_K^2 - |M_K Q|^2) + (|M_D M_K| \pm |M_K Q|)^2 \\
 &= R_K^2 + |M_D M_K|^2 \pm 2|M_D M_K||M_K Q|.
 \end{aligned} \tag{6}$$

Here, depending on which side of  $t$  the point  $M_K$  lies, one chooses the sign in the term  $\pm|M_K Q|$  in the second line.

Similarly, if we denote the point at which the tangent  $t$  touches the circle  $C$  by  $P$ , we find

$$\begin{aligned}
 R_D^2 &= |M_C M_D|^2 = |PQ|^2 + (R_C + |M_D M_K| \pm |M_K Q|)^2 \\
 &= (d^2 - (R_C \pm |M_K Q|)^2) + (R_C + |M_D M_K| \pm |M_K Q|)^2 \\
 &= d^2 + |M_D M_K|^2 + 2|M_D M_K|(R_C \pm |M_K Q|).
 \end{aligned} \tag{7}$$

Equating the terms in (6) and (7) and simplifying gives

$$R_K^2 = d^2 + 2|M_D M_K|R_C. \tag{8}$$

Hence  $|M_D M_K| =: R_F$  does not depend on  $t$  and (5) follows from (8).  $\square$

Now we are ready for the proof of the Theorem.

*Proof.* Consider one side of the bicentric polygon, say  $P_1P_2$ , as indicated in Figure 5. The black lines in the vertices of the brown bicentric polygon are the external angle bisectors, and their intersections the centers of the excircles of the bicentric polygon, e.g., the point  $N$ . The dotted internal angle bisectors meet in  $M_C$  and are orthogonal to the external angle bisectors. Hence the circumcircle  $D$  of the triangle  $M_C P_1 P_2$  is the circle of Thales over the segment  $M_C N$ . According to Lemma 1 its center  $M_D$  lies on the circle  $F$  around  $M_K$  with a fixed radius  $R_F$  given by (5), independent of the side of the bicentric polygon. It follows by a homothety with center  $M_C$  and factor 2 that  $N$  lies on a circle  $E$  around the point  $M_E$  with radius  $2R_F$ . This finishes the proof.  $\square$




Figure 5: Proof of the theorem.

Notice that it is not necessary that  $C$  lies inside  $K$ . Figure 6 illustrates the theorem for a bicentric pentagon if  $C$  and  $K$  intersect. Figure 7 shows the situation for a bicentric octagon if  $C$  lies outside of  $K$ .

As a last remark we mention the following dynamic aspect of the main theorem. Since the inscribed circle  $C$  and the circumcircle  $K$  of a bicentric polygon form a Poncelet pair, we can choose any point on  $K$  as a vertex of a bicentric polygon with inscribed circle  $C$  and circumcircle  $K$ . It follows from the main theorem that the centers of the excircles of this new bicyclic polygon still lie on the same circle  $E$ . Stated differently, if we move a vertex along  $K$  the centers of the excircles roll along  $E$ .




Figure 6: The theorem with  $C$  and  $K$  intersecting. The thin black lines are the external angle bisectors in the vertices of the brown bicentric pentagon. Their intersections, the centers of the excircles  $M_1, \dots, M_5$  of the pentagon, are concyclic on the circle  $E$ . To avoid overloading the figure, only the orange excircle with center  $M_1$  has been drawn.

### 3 Bicentric convex quadrilaterals

Bicentric convex quadrilaterals have been widely studied, see, e.g., [13, 4, 15, 5, 17]. Here we want to interpret the relation (4) geometrically. We start with the following lemma which is of some independent interest.

**Lemma 2.** *Let  $A_1A_2A_3A_4$  be a bicentric convex quadrilateral,  $M_C$  the center of its incircle, and  $M_1, M_2, M_3, M_4$  the centers of the excircles. Then the diagonals of the quadrilateral  $M_1M_2M_3M_4$  intersect perpendicularly in  $M_C$ .*

*Proof.* As we have seen in the proof of Lemma 1,  $A_i$  and  $A_{i+1}$  lie on the circle of Thales with diameter  $M_iM_C$  (see Figure 8). Therefore we have

$$\begin{aligned}
 \sphericalangle M_4M_CM_1 &= \sphericalangle M_4M_CA_1 + \sphericalangle A_1M_CM_1 \\
 &= \sphericalangle M_4A_4A_1 + \sphericalangle A_1A_2M_1 \\
 &= \left(\frac{\pi}{2} - \frac{\sphericalangle A_1A_4A_3}{2}\right) + \left(\frac{\pi}{2} - \frac{\sphericalangle A_3A_2A_1}{2}\right) \\
 &= \pi - \frac{1}{2}(\sphericalangle A_1A_4A_3 + \sphericalangle A_3A_2A_1) \\
 &= \frac{\pi}{2}.
 \end{aligned}$$

By shifting the indices cyclically, we get the desired result.  $\square$

Now we can interpret the relation (4) which connects the radii of the circles  $E$  and  $K$  with the distance  $d$  of their centers geometrically:

$$R_E^2 = 2(R_K^2 + d^2). \quad (4)$$




Figure 7: The theorem with circle  $C$  outside of  $K$ . The thin black lines are the external angle bisectors in the vertices of the brown bicentric octagon. Their intersections, the centers of the excircles of the octagon, are concyclic on the circle  $E$ .

*Proof of (4).* According to Poncelet's theorem, we can move  $A_1$  along  $K$  until it coincides with the line  $\ell$  through the points  $M_C, M_K, M_E$ . In this position, the points  $A_2, A_4$  lie symmetrically with respect to  $\ell$  and hence  $A_3$  also lies on  $\ell$  (see Figure 9). By Lemma 2 the triangle  $M_1M_CM_4$  is now an isosceles right-angled triangle.  $A_1$  is the midpoint of the segment  $M_1M_4$ , so the two triangles  $M_1M_CA_1$  and  $A_1M_CM_4$  are also isosceles right-angled triangles. Pythagoras in the triangle  $A_1M_1M_E$  yields

$$\begin{aligned} R_E^2 &= |M_1M_E|^2 = |M_EA_1|^2 + |A_1M_1|^2 = (R_K + d)^2 + |A_1M_C|^2 = \\ &= (R_K + d)^2 + (R_K - d)^2 = 2(R_K^2 + d^2), \end{aligned}$$

as desired. □






Figure 8: Proof of Lemma 2.

## 4 The area ratio of a bicentric convex polygon and its excenters polygon

It turns out that bicentric polygons have another nice property. This observation was originally formulated as a conjecture by a colleague.

**Theorem 2.** *Let  $A_1A_2 \dots A_n$  be a bicentric convex polygon,  $r_C$  the center of its incircle,  $M_1M_2 \dots M_n$  the polygon formed by the centers of the excircles, and  $r_E$  the radius of its circumcircle. Then, we have*

$$\frac{\text{area } A_1A_2 \dots A_n}{\text{area } M_1M_2 \dots M_n} = \frac{r_C}{r_E}.$$

*Proof.* As before, we denote the center of the incircle of  $A_1A_2 \dots A_n$  by  $M_C$  and the center of the circumcircle of  $M_1M_2 \dots M_n$  by  $M_E$ . Observe that the lines  $M_EM_i$  are perpendicular to the sides  $a_i = A_iA_{i+1}$  of the bicentric polygon. To see this, look again at the homothety with center  $M_C$  and factor 2 that we used in the proof of Theorem 1, and go back to Figure 5: The line  $M_KM_D$  is the perpendicular bisector of the line segment  $P_1P_2$ . By the homothety, the line  $M_KM_D$  is mapped to the line  $M_EN$ , which is therefore also perpendicular to  $P_1P_2$ .

Consider now the orthodiagonal quadrilateral  $M_EA_iM_iA_{i+1}$  (see Figure 10 where the case  $i = 1$  is shown). Its area is given by

$$\text{area } M_EA_iM_iA_{i+1} = \frac{1}{2} \cdot a_i \cdot r_E.$$




Figure 9: Proof of relation (4).

Here  $a_i$  denotes the length of the side  $A_iA_{i+1}$ . Similarly we have for the area of the triangle  $M_C A_i A_{i+1}$

$$\text{area } M_C A_i A_{i+1} = \frac{1}{2} \cdot a_i \cdot r_C.$$

This leads us to

$$\frac{\text{area } M_C A_i A_{i+1}}{\text{area } M_E A_i M_i A_{i+1}} = \frac{r_C}{r_E}$$

and then, by taking the sum over  $i$ , to the desired result

$$\frac{\text{area } A_1 A_2 \dots A_n}{\text{area } M_1 M_2 \dots M_n} = \frac{r_C}{r_E}.$$

□

## References

- [1] William Chapple. An essay on the properties of triangles inscribed in and circumscribed about two given circles. *Miscellanea Curiosa Mathematica*, 4:117—124, 1746.
- [2] Junhao Cheng, Long Ma, and Yuanfeng Zhou. A new method for researching and constructing spherical bicentric polygons based on geometric mapping. *Comput. Aided Geom. Design*, 105:Paper No. 102232, 13, 2023.
- [3] H. S. M. Coxeter and S. L. Greitzer. *Geometry revisited*, volume 19 of *New Mathematical Library*. Random House, Inc., New York, 1967.
- [4] Ana Diakvnishvili. On an explicit construction of bicentric quadrilaterals. *Proc. I. Vekua Inst. Appl. Math.*, 72:16–22, 2022.




Figure 10: The ratio of the red and the blue area is independent of the position of the starting point  $A_1$  of the Poncelet polygon  $A_1A_2 \dots A_n$  on  $K$ .

- [5] Michael Diao and Andrew Wu. The radical axis of the circumcircle and incircle of a bicentric quadrilateral. *Forum Geom.*, 19:39–43, 2019.
- [6] Heinrich Dörrie. *100 great problems of elementary mathematics*. Dover Publications, Inc., New York, fifth edition, 1982. Reprint of the 1965 edition.
- [7] Vladimir Dragović and Milena Radnović. *Poncelet porisms and beyond*. Frontiers in Mathematics. Birkhäuser/Springer Basel AG, Basel, 2011.
- [8] Leonhard Euler. Solutio facilis problematum quorundam geometricorum difficillimorum. *Novi Commentarii academiae scientiarum imperialis Petropolitanae*, 11:103–123, 1767.
- [9] Leopold Flatto. *Poncelet's theorem*. American Mathematical Society, Providence, RI, 2009.
- [10] Nicolas Fuss. De polygonis symmetrice irregularibus circulo simul inscriptis et circumscriptis. *Nova Acta Petropol.*, 13:166–189, 1798.
- [11] Grigori Giorgadze and Giorgi Khimshiashvili. Remarks on bicentric polygons. *Bull. Georgian Natl. Acad. Sci. (N.S.)*, 7(3):5–10, 2013.
- [12] Lorenz Halbeisen and Norbert Hungerbühler. A simple proof of Poncelet's theorem (on the occasion of its bicentennial). *Amer. Math. Monthly*, 122(6):537–551, 2015.
- [13] Hans Humenberger. On six collinear points in bicentric quadrilaterals. *Math. Mag.*, 96(3):285–295, 2023.
- [14] Carl Gustav Jacob Jacobi. Ueber die Anwendung der elliptischen Transcendenten auf ein bekanntes Problem der Elementargeometrie: "Die Relation zwischen der Distanz der Mittelpunkte und den Radien zweier Kreise zu finden, von denen der eine einem unregelmäßigen Polygon eingeschrieben, der andere demselben umgeschrieben ist". *J. Reine Angew. Math.*, 3:376–389, 1828.

- [15] Martin Josefsson. New characterisations of bicentric quadrilaterals. *Math. Gaz.*, 106(567):414–426, 2022.
- [16] Sulaiman M. Kerawala. Poncelet porism in two circles. *Bull. Calcutta Math. Soc.*, 39:85–105, 1947.
- [17] G. Khimshiashvili. Remarks on bicentric quadrilaterals. *Proc. A. Razmadze Math. Inst.*, 168:41–52, 2015.
- [18] Mirko Radić. About two characteristic points concerning two nested circles and their use in research of bicentric polygons. *Forum Geom.*, 15:129–157, 2015.
- [19] Friedrich Julius Richelot. Anwendung der elliptischen Transcendenten auf die sphärischen Polygone, welche zugleich einem kleinen Kreise der Kugel eingeschrieben und einem andern umgeschrieben sind. *J. Reine Angew. Math.*, 5:250–267, 1830.
- [20] Pedro Roitman, Ronaldo Garcia, and Dan Reznik. New invariants of Poncelet-Jacobi bicentric polygons. *Arnold Math. J.*, 7(4):619–637, 2021.
- [21] Jakob Steiner. Aufgaben und Lehrsätze, erstere aufzulösen, letztere zu beweisen. Geometrische Lehrsätze. *Journal für die reine und angewandte Mathematik*, 2:96–98, 1827.