GENUS g VIRASORO CORRELATION FUNCTIONS FOR VERTEX OPERATOR ALGEBRAS

MICHAEL P. TUITE AND MICHAEL WELBY[†]

ABSTRACT. For a simple, self-dual, strong CFT-type vertex operator algebra (VOA) of central charge c, we describe the Virasoro n-point correlation function on a genus g marked Riemann surface in the Schottky uniformisation. We show that this n-point function determines the correlation functions for all Virasoro vacuum descendants. Using our recent work on genus g Zhu recursion, we show that the Virasoro n-point function is determined by a differential operator \mathcal{D}_n acting on the genus g VOA partition function normalised by the Heisenberg partition function to the power of c. We express \mathcal{D}_n as the sum of weights over certain Virasoro graphs where the weights explicitly depend on c, the classical bidifferential of the second kind, the projective connection, holomorphic 1-forms and derivatives with respect to any 3g - 3 locally independent period matrix elements. We also describe the modular properties of \mathcal{D}_n under a homology base change.

1. INTRODUCTION

A vertex operator algebra (VOA) (e.g. [K, LL, MT1]) is an algebraic theory closely related to conformal field theory (CFT) in physics e.g. [DFMS]. An essential ingredient in both theories is the Virasoro vector whose vertex operator modes generate a Virasoro algebra of some central charge c. Virasoro n-point correlation functions on genus zero and one Riemann surfaces have long been studied in the CFT and VOA literature. For instance, they occur in describing the Kac determinant and genus zero CFT Ward identities (e.g. [DFMS]) and VOA modular differential equations arising from genus one Zhu recursion [Z1]. In this paper we apply recent results from [TW1] concerning Zhu recursion theory for correlation functions on Riemann surfaces of genus $q \ge 2$ constructed by a Schottky uniformatisation. In particular, we consider the genus g Virasoro n-point function from which all correlation functions for Virasoro vacuum descendants can be determined. We show that this is expressible as an order n differential operator acting on a suitably normalised genus q partition function where the differential operator is explicitly described as a sums of certain weights of Virasoro graphs previously introduced in [HT] for genus zero and one Virasoro *n*-point functions. Here the graph weights are defined in terms of some classical differential forms and differential operators with respect to 3q-3locally independent period matrix elements on the Riemann surface.

In Section 2 we briefly review the classical Schottky uniformisation of a genus g marked Riemann surface $\mathcal{S}^{(g)}$ [Fo, FK, Bo], where we sew g handles to a Riemann sphere, expressed in terms of 3g Schottky sewing parameters. We review the classical bidifferential of the second kind, the projective connection, holomorphic 1-forms and the period matrix. These feature in some differential equations later in Section 2 and also enter into the description of Virasoro *n*-point functions in Section 4. We review properties of the Bers quasidifferential (N, 1 - N)-form $\Psi_N(x, y)$ [Be1, Be2, TW2]. Ψ_N and an associated spanning set of

[†]Supported by an Irish Research Council Government of Ireland Postgraduate Scholarship.

holomorphic N forms appear in the genus g Zhu recursion formula of [TW1]. $\Psi_2(x, y)$ plays a crucial role in this paper since we exploit Zhu recursion for Virasoro vector ω insertions. Using $\Psi_2(x, y)$ and its associated 2-form spanning set, we define the fundamental differential operators $\nabla(x)$ and $\nabla_{\boldsymbol{y}}^{(\boldsymbol{m})}(x)$ for variations in the Schottky parameters and local coordinates $\boldsymbol{y} = y_1, \ldots, y_n$ on $\mathcal{S}^{(g)}$. $\nabla_{\boldsymbol{y}}^{(\boldsymbol{m})}(x)$ maps any meromorphic form of weight $(\boldsymbol{m}) = (m_1, \ldots, m_n)$ in $\boldsymbol{y} = y_1, \ldots, y_n$ to a meromorphic form of weight $(2, \boldsymbol{m})$ in x, \boldsymbol{y} [TW2]. Section 2 also contains some refinements of [TW2] where, using global SL(2, \mathbb{C}) Möbius invariance, we introduce differential operators $\nabla_{\mathfrak{M}}(x)$ and $\nabla_{\mathfrak{M},\boldsymbol{y}}^{(\boldsymbol{m})}(x)$ in terms of local variations in Schottky space \mathfrak{S}_g (the SL(2, \mathbb{C}) quotient of the Schottky parameter space) and local coordinates \boldsymbol{y} on $\mathcal{S}^{(g)}$. We further show that $\nabla_{\mathfrak{M}}(x)$ can be expressed in terms of variations of any local coordinates on moduli space \mathfrak{M} . Lastly, using these operators, we describe some differential equations for classical differentials [TW1, TW2].

Section 3 begins with a brief review of Vertex Operator Algebra (VOA) theory. For a VOA V which is simple, self-dual (V isomorphic to its dual module V') of strong CFT-type, we define the partition function Z_V and n-point correlation function $\mathcal{F}_V(\boldsymbol{v}, \boldsymbol{z})$ on a genus g Riemann surface where $\boldsymbol{v} = v_1, \ldots, v_n \in V$ are inserted at $\boldsymbol{z} = z_1, \ldots, z_n$, respectively. We review genus g Zhu recursion [TW1] which is an expansion of a (n + 1)-point correlation function $\mathcal{F}_V(u, x; \boldsymbol{v}; \boldsymbol{z})$, for $u \in V$ quasiprimary of weight N, in terms of n-point functions dependent on \boldsymbol{z} with universal coefficients given by z_k derivatives of the Bers quasiform $\Psi_N(x, z_k)$ and holomorphic N-forms in x.

In Section 4 we consider the Virasoro *n*-point correlation function $\mathcal{F}_V(\boldsymbol{\omega}, \boldsymbol{z})$ with *n* insertions of the Virasoro vector $\boldsymbol{\omega}$. We show that $\mathcal{F}_V(\boldsymbol{\omega}, \boldsymbol{z})$ is a generating function for all Virasoro vacuum descendant correlation functions. Genus *g* Zhu recursion implies a recursive conformal Ward identity for $\mathcal{F}_V(\boldsymbol{\omega}, \boldsymbol{z})$ involving the differential operators $\nabla(x)$ and $\nabla_{\boldsymbol{z}}^{(\boldsymbol{m})}(x)$ of Section 2 e.g. $\mathcal{F}_V(\boldsymbol{\omega}, \boldsymbol{x}) = \nabla(x)Z_V$. The main result of this paper is an explicit expression for the normalised *n*-point function $G_n(\boldsymbol{z}) := \mathcal{F}_V(\boldsymbol{\omega}, \boldsymbol{x})Z_M^{-c}$ where *M* is the rank 1 Heisenberg VOA and *c* the central charge of *V*. We show that $G_n(\boldsymbol{z}) = \mathcal{D}_n(\boldsymbol{z})\Theta_V$ for normalized partition function $\Theta_V := Z_V Z_M^{-c}$ and where $\mathcal{D}_n(\boldsymbol{z})$ is an order *n* differential operator with respect to 3g - 3 locally independent period matrix elements. $\mathcal{D}_n(\boldsymbol{z})$ is symmetric under permutations of \boldsymbol{z} and is defined as the sum of weights of so-called Virasoro graphs [HT] where the weights are defined in terms of *c*, the classical differentials of Section 2 and variations of the 3g - 3 independent period matrix elements.

In Section 5 we consider a change of Riemann surface marking in the Schottky scheme described by the action of the symplectic modular group $\operatorname{Sp}(2g,\mathbb{Z})$ on the given homology basis. We show that $\nabla_{\mathfrak{M}}(x)$ and $\nabla_{\mathfrak{M},\boldsymbol{y}}^{(\boldsymbol{m})}(x)$ are modular invariant whereas $\mathcal{D}_n(\boldsymbol{z})$ is invariant up to a *c*-dependent Teichmüller form-like automorphic factor. We conclude with some remarks about the significance of our results.

2. DIFFERENTIAL STRUCTURES ON RIEMANN SURFACES

2.1. Notational conventions. Define the following indexing sets for integers $g, N \ge 1$

(1)
$$\mathcal{I} := \{-1, \dots, -g, 1, \dots, g\}, \quad \mathcal{I}_+ := \{1, \dots, g\}, \quad \mathcal{L}_N := \{0, 1, \dots, 2N - 2\}.$$

For functions f(x) and g(x, y) and integers $i, j \ge 0$ we define

$$f^{(i)}(x) := \partial^{(i)} f(x) := \partial^{(i)}_x f(x) := \frac{1}{i!} \frac{\partial^i}{\partial x^i} f(x)$$

$$g^{(i,j)}(x,y) := \partial_x^{(i)} \partial_y^{(j)} g(x,y).$$

2.2. The Schottky uniformisation of a Riemann surface. Consider a compact marked Riemann surface $\mathcal{S}^{(g)}$ of genus g, e.g. [FK, Mu1, Fa, Bo], with canonical homology basis α_a , β_a for $a \in \mathcal{I}_+$. We review the construction of a genus g Riemann surface $\mathcal{S}^{(g)}$ using the Schottky uniformisation where we sew g handles to the Riemann sphere $\mathcal{S}^{(0)} \cong \widehat{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$ e.g. [Fo, Bo]. Every Riemann surface can be (non-uniquely) Schottky uniformised [Be2].

For $a \in \mathcal{I}$ let $\mathcal{C}_a \subset \mathcal{S}^{(0)}$ be 2g non-intersecting Jordan curves where $z \in \mathcal{C}_a, z' \in \mathcal{C}_{-a}$ for $a \in \mathcal{I}_+$ are identified by a sewing relation

(2)
$$\frac{z' - W_{-a}}{z' - W_a} \cdot \frac{z - W_a}{z - W_{-a}} = q_a, \quad a \in \mathcal{I}_+,$$

for q_a with $0 < |q_a| < 1$ and $W_{\pm a} \in \widehat{\mathbb{C}}$. Thus $z' = \gamma_a z$ for $a \in \mathcal{I}_+$ with

$$\gamma_a := \sigma_a^{-1} \begin{pmatrix} q_a^{1/2} & 0\\ 0 & q_a^{-1/2} \end{pmatrix} \sigma_a, \quad \sigma_a := (W_{-a} - W_a)^{-1/2} \begin{pmatrix} 1 & -W_{-a}\\ 1 & -W_a \end{pmatrix}.$$

The points $\sigma_a(W_{-a}) = 0$ and $\sigma_a(W_a) = \infty$ are, respectively, attractive and repelling fixed points of $Z \to Z' = q_a Z$ for $Z = \sigma_a z$ and $Z' = \sigma_a z'$. W_{-a} and W_a are the corresponding fixed points for γ_a . We identify the standard homology cycles α_a with \mathcal{C}_{-a} and β_a with a path connecting $z \in \mathcal{C}_a$ to $z' = \gamma_a z \in \mathcal{C}_{-a}$.

The genus g Schottky group Γ is the free group with generators γ_a for $a \in \mathcal{I}_+$. Define $\gamma_{-a} := \gamma_a^{-1}$. The independent elements of Γ are reduced words of length k of the form $\gamma = \gamma_{a_1} \dots \gamma_{a_k}$ where $a_i \neq -a_{i+1}$ for each $i = 1, \dots, k-1$. We let $\Lambda(\Gamma)$ denote the limit set¹ of Γ i.e. the set of limit points of the action of Γ on $\widehat{\mathbb{C}}$. Then $\mathcal{S}^{(g)} \simeq \Omega_0 / \Gamma$ where $\Omega_0 := \widehat{\mathbb{C}} - \Lambda(\Gamma)$.

Define $w_a := \gamma_{-a}(\infty)$. Using (2) we find

(3)
$$w_a = \frac{W_a - q_a W_{-a}}{1 - q_a}, \quad a \in \mathcal{I},$$

where we define $q_{-a} := q_a$. Then (2) is equivalent to

(4)
$$(z'-w_{-a})(z-w_{a}) = \rho_{a}, \quad a \in \mathcal{I}_{+},$$

with

(5)
$$\rho_a := -\frac{q_a(W_{-a} - W_a)^2}{(1 - q_a)^2} = -\frac{q_a(w_{-a} - w_a)^2}{(1 + q_a)^2}.$$

Equation (4) implies

$$\gamma_a z = w_{-a} + \frac{\rho_a}{z - w_a}.$$

It is convenient (but not necessary) to choose the Jordan curve C_a to be the boundary of the disc Δ_a with centre w_a and radius $|\rho_a|^{\frac{1}{2}}$. Then γ_a maps the exterior (interior) of Δ_a to the interior (exterior) of Δ_{-a} since

$$|\gamma_a z - w_{-a}| |z - w_a| = |\rho_a|.$$

¹Note that for g = 1, $\Lambda(\Gamma)$ is the empty set.

Furthermore, the discs Δ_a , Δ_b are non-intersecting if and only if

(6)
$$|w_a - w_b| > |\rho_a|^{\frac{1}{2}} + |\rho_b|^{\frac{1}{2}}, \quad \forall \ a \neq b.$$

We define \mathfrak{C}_g to be the set $\{(w_a, w_{-a}, \rho_a) | a \in \mathcal{I}_+\} \subset \mathbb{C}^{3g}$ satisfying (6). We refer to \mathfrak{C}_g as the Schottky parameter space.

The cross ratio (2) is Möbius invariant for $\sigma = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{SL}_2(\mathbb{C})$ with $(z, z', W_a, q_a) \rightarrow (\sigma z, \sigma z', \sigma W_a, q_a)$ giving a global Möbius $\mathrm{SL}_2(\mathbb{C})$ action on \mathfrak{C}_g as follows

$$\sigma: (w_a, \rho_a) \mapsto \left(\frac{(Aw_a + B)(Cw_{-a} + D) - \rho_a AC}{(Cw_a + D)(Cw_{-a} + D) - \rho_a C^2}, \frac{\rho_a}{((Cw_a + D)(Cw_{-a} + D) - \rho_a C^2)^2}\right).$$

Furthermore, $\sigma z' = \sigma \gamma_a \sigma^{-1}(\sigma z)$ so that Γ is mapped to the conjugate Schottky group $\sigma \Gamma \sigma^{-1}$. We define Schottky space as $\mathfrak{S}_g := \mathfrak{C}_g/\mathrm{SL}_2(\mathbb{C})$ which provides a natural covering space for the moduli space \mathfrak{M} of genus g Riemann surfaces (of complex dimension 1 for g = 1 and 3g - 3 for $g \ge 2$). We exploit the \mathfrak{C}_g parametrisation throughout because the sewing relation (4) is more readily implemented in the theory of vertex operators.

2.3. Some classical differentials on a Riemann surface. Let $\mathcal{S}^{(g)}$ be a marked compact genus g Riemann surface with canonical homology basis α_a, β_a for $a \in \mathcal{I}_+$. The meromorphic bidifferential form of the second kind is a unique symmetric form [Mu1, Fa]

(8)
$$\omega(x,y) = \left(\frac{1}{(x-y)^2} + \text{regular terms}\right) dxdy$$

for local coordinates x, y where $\oint_{\alpha_a} \omega(x, \cdot) = 0$ for all $a \in \mathcal{I}_+$. It follows that

(9)
$$\nu_a(x) := \oint_{\beta_a} \omega(x, \cdot), \quad a \in \mathcal{I}_+,$$

is a holomorphic 1-form normalised by $\oint_{\alpha_a} \nu_b = 2\pi i \delta_{ab}$. $\{\nu_a(x)\}$ is a basis for the space of holomorphic 1-forms \mathcal{H}_1 . The $g \times g$ symmetric period matrix is given by $\Omega = \frac{1}{2\pi i} \tau$ where

(10)
$$\tau_{ab} := \oint_{\beta_a} \nu_b, \qquad a, b \in \mathcal{I}_+.$$

The projective connection s(x) is defined by

(11)
$$s(x) := 6 \lim_{y \to x} \left(\omega(x, y) - \frac{dxdy}{(x - y)^2} \right).$$

The Bers quasiform of weight (N, 1 - N) for $g \ge 2$ and $N \ge 2$ is defined by the following Poincaré series [Be1, Be2, McIT, TW2]

(12)
$$\Psi_N(x,y) := \sum_{\gamma \in \Gamma} \Pi_N(\gamma x, y), \quad x, y \in \Omega_0,$$

(13)
$$\Pi_N(x,y) := \Pi_N(x,y; \mathbf{A}) := \frac{1}{x-y} \prod_{\ell \in \mathcal{L}_N} \frac{y-A_\ell}{x-A_\ell} dx^N dy^{1-N},$$

for index set \mathcal{L}_N of (1) and where $\mathbf{A} := A_0, \ldots, A_{2N-2} \in \Lambda(\Gamma)$ are distinct limit points of Γ . Π_N is globally Möbius invariant with

(14)
$$\Pi_N(\sigma x, \sigma y; \boldsymbol{\sigma} \boldsymbol{A}) = \Pi_N(x, y; \boldsymbol{A}),$$

for all $\sigma \in SL_2(\mathbb{C})$. $\Psi_N(x, y)$ is meromorphic for $x, y \in \Omega_0$ with simple poles of residue one at $y = \gamma x$ for all $\gamma \in \Gamma$. It is an N-differential in x since

$$\Psi_N(\gamma x, y) = \Psi_N(x, y), \quad \gamma \in \Gamma,$$

by construction, and is a quasiperiodic (1 - N)-form in y with [TW2]

(15)
$$\Psi_N(x, \gamma_a y) - \Psi_N(x, y) = -\sum_{\ell \in \mathcal{L}_N} \Theta_{N,a}^{\ell}(x) (y - w_a)^{\ell} dy^{1-N},$$

for each Schottky group generator γ_a and where $\{\Theta_{N,a}^{\ell}(x)\}_{a\in\mathcal{I}_+}^{\ell\in\mathcal{L}_N}$ spans the vector space \mathcal{H}_N of holomorphic N-forms. We note that $\dim \mathcal{H}_N = (g-1)(2N-1)$ for $g, N \ge 2$ by the Riemann-Roch Theorem.

For N = 1 and $g \ge 2$ we define $\Psi_1(x, y) := \sum_{\gamma \in \Gamma} \prod_1(\gamma x, y)$ with

$$\Pi_1(x,y) := \left(\frac{1}{x-y} - \frac{1}{x-A_0}\right) dx,$$

where here $A_0 \in \Omega_0$. Then we find [Bu]

(16)
$$\omega(x,y) = \Psi_1^{(0,1)}(x,y)dy.$$

Hence $\Psi_1(x, y) = \omega_{y-A_0}(x) = \int_{A_0}^y \omega(x, \cdot)$, the classical differential of the third kind which is not a Bers quasiform since it has simple poles both at x = y and $x = A_0$. Lastly, for $N \ge 1$, we define a canonical symmetric meromorphic (N, N)-form $\omega_N(x, y)$ with a pole of order 2N at x = y, generalising (16) as follows:

(17)
$$\omega_N(x,y) := \Psi_N^{(0,2N-1)}(x,y) dy^{2N-1} = \sum_{\gamma \in \Gamma} \frac{d(\gamma x)^N dy^N}{(\gamma x - y)^{2N}}.$$

2.4. Differential operators on meromorphic forms. We review some first order differential operators \mathcal{D}^p , $\mathcal{D}^{p,(m)}_{\boldsymbol{y}}$, $\nabla(x)$ and $\nabla^{(m)}_{\boldsymbol{y}}(x)$ introduced in [TW2] and developed in a VOA context in [TW1]. We first define the following basis for the tangent space $T(\mathfrak{C}_g)$:

(18)
$$\hat{c}_a^0 := \partial_{w_a}, \quad \hat{c}_a^1 := \rho_a \partial_{\rho_a}, \quad \hat{c}_a^2 := \rho_a \partial_{w_{-a}}, \text{ for } a \in \mathcal{I}_+.$$

The generator \mathcal{D}^p of any global Möbius transformation (7) is determined by a quadratic polynomial p(z) where [TW2, TW1]

(19)
$$\mathcal{D}^{p} := \sum_{a \in \mathcal{I}} p(W_{a}) \partial_{W_{a}} = \sum_{a \in \mathcal{I}} \left(p(w_{a}) \partial_{w_{a}} + p^{(1)}(w_{a}) \rho_{a} \partial_{\rho_{a}} + p^{(2)}(w_{a}) \rho_{a} \partial_{w_{-a}} \right)$$
$$= \sum_{a \in \mathcal{I}_{+}} \sum_{\ell \in \mathcal{L}_{2}} p_{a}^{\ell} \partial_{a}^{\ell},$$

with² $p_a^{\ell} := p^{(\ell)}(w_a) + \rho_a^{1-\ell} p^{(2-\ell)}(w_{-a})$. We next define a differential operator mapping differentiable functions on \mathfrak{C}_g to \mathcal{H}_2 :

(20)
$$\nabla(x) := \sum_{a \in \mathcal{I}_+} \sum_{\ell \in \mathcal{L}_2} \Theta_{2,a}^{\ell}(x) \partial_a^{\ell};$$

²Note that p_a^{ℓ} is defined with opposite sign in [TW1, TW2].

for \mathcal{H}_2 spanning set $\{\Theta_{2,a}^{\ell}(x)\}$ associated with quasi-periodicity $\Psi_2(x, y)$ in (15). $\Psi_2(x, y)$ depends on the choice of the 3 limit points $\mathbf{A} = A_0, A_1, A_2$ in (13). Any pair of limit points \mathbf{A} and $\hat{\mathbf{A}}$ are related by a Möbius map so that $\Psi_2(x, y)$ is unique up to global Möbius maps using (14). This has important consequences for the operator $\nabla(x)$ and its generalisation below. Let $\hat{\Psi}_2(x, y)$ be the Bers quasiform for alternative limit points $\hat{\mathbf{A}}$. Then $\hat{\Psi}_2(x, y) - \Psi_2(x, y)$ is holomorphic with

(21)
$$\widehat{\Psi}_2(x,y) = \Psi_2(x,y) + \sum_{r=1}^{3g-3} \Phi_r(x) p_r(y) dy^{-1},$$

for some \mathcal{H}_2 basis $\{\Phi_r(x)\}\$ and quadratic polynomials p_r . For the corresponding \mathcal{H}_2 spanning set $\{\widehat{\Theta}_{2,a}^{\ell}(x)\}\$ from (15) and with $(p_r)_a^{\ell}$ of (19) we find

(22)
$$\widehat{\Theta}_{2,a}^{\ell}(x) = \Theta_{2,a}^{\ell}(x) + \sum_{r=1}^{3g-3} (p_r)_a^{\ell} \Phi_r(x),$$

and differential operator

(23)
$$\widehat{\nabla}(x) = \nabla(x) + \sum_{r=1}^{3g-3} \Phi_r(x) \mathcal{D}^{p_r}$$

for Möbius generators \mathcal{D}^{p_r} . Thus $\nabla(x)$ determines a unique tangent vector field $\nabla_{\mathfrak{M}}(x)$, independent of A, on the Schottky tangent space $T(\mathfrak{S}_g) \cong T(\mathfrak{M})$. We therefore find:

Lemma 2.1. Let $F(\boldsymbol{\eta})$ be a differentiable function on \mathfrak{M} for any local coordinates $\boldsymbol{\eta} := \eta_1, \ldots, \eta_{3g-3}$. Then

(24)
$$\nabla(x)F(\boldsymbol{\eta}) = \nabla_{\mathfrak{M}}(x)F(\boldsymbol{\eta}).$$

Let $\mathcal{M}^{(m)}$ denote the space of meromorphic forms $H^{(m)}(\boldsymbol{y})$ in *n* variables $\boldsymbol{y} := y_1, \ldots, y_n$ of weight (\boldsymbol{m}) for $\boldsymbol{m} := m_1, \cdots, m_n$ i.e. $H^{(\boldsymbol{m})}(\boldsymbol{y}) = h(\boldsymbol{y})dy_1^{m_1} \ldots dy_n^{m_n}$ for some meromorphic function $h(\boldsymbol{y})$. We define the following differential operator on $\mathcal{M}^{(\boldsymbol{m})}$:

(25)
$$\nabla_{\boldsymbol{y}}^{(\boldsymbol{m})}(x) := \nabla(x) + \sum_{k=1}^{n} \left(\Psi_2(x, y_k) \, d_{y_k} + m_k d_{y_k} \left(\Psi_2(x, y_k) \right) \right),$$

where $d_y(f(y)) := \partial_y f(y) dy$.

Remark 2.1. We note that $\nabla_{\boldsymbol{y}}^{(\boldsymbol{m})}(x)$ satisfies a Leibniz product rule (with adjusted weights) e.g. for $G(y) \in \mathcal{M}^{(m)}$ and $H(z) \in \mathcal{M}^{(n)}$ we have

$$\nabla_{y,z}^{(m,n)}(x)(G(y)H(z)) = \left(\nabla_y^{(m)}(x)G(y)\right)H(z) + G(y)\nabla_z^{(n)}(x)H(z).$$

The definition (25) is unambiguous if any \boldsymbol{y} variables are equal e.g. for $G(y, z) \in \mathcal{M}^{(m,n)}$ regular at y = z, then $G(y, y) \in \mathcal{M}^{(m+n)}$ with

$$\nabla_{y,y}^{(m,n)}(x)G(y,y) = \nabla_y^{(m+n)}(x)G(y,y).$$

Choosing alternative limit points \hat{A} so that (21) holds, then (26) generalises to

(26)
$$\widehat{\nabla}_{\boldsymbol{y}}^{(\boldsymbol{m})}(x) = \nabla_{\boldsymbol{y}}^{(\boldsymbol{m})}(x) + \sum_{r=1}^{3g-3} \Phi_r(x) \mathcal{D}_{\boldsymbol{y}}^{(\boldsymbol{m}),p_r}$$

where

(27)
$$\mathcal{D}_{\boldsymbol{y}}^{p,(\boldsymbol{m})} := \mathcal{D}^p + \sum_{k=1}^n \left(p(y_k) \partial_{y_k} + m_k p^{(1)}(y_k) \right),$$

for any quadratic polynomial p(y). We find [TW2]

Proposition 2.1. Let $H^{(m)}(y) \in \mathcal{M}^{(m)}$. Then

- (i) $\mathcal{D}_{\boldsymbol{y}}^{p,(\boldsymbol{m})}$ generates a global Möbius transformation with $\mathcal{D}_{\boldsymbol{y}}^{p,(\boldsymbol{m})}H^{(\boldsymbol{m})}(\boldsymbol{y}) = 0;$
- (ii) $\nabla_{\boldsymbol{y}}^{(\boldsymbol{m})}(x)H^{(\boldsymbol{m})}(\boldsymbol{y}) \in \mathcal{M}^{(2,\boldsymbol{m})}$, the space of meromorphic forms of weight $(2,\boldsymbol{m})$; (iii) Commutativity: $\nabla_{x,\boldsymbol{y}}^{(2,\boldsymbol{m})}(z)\nabla_{\boldsymbol{y}}^{(\boldsymbol{m})}(x)H^{(\boldsymbol{m})}(\boldsymbol{y}) = \nabla_{z,\boldsymbol{y}}^{(2,\boldsymbol{m})}(x)\nabla_{\boldsymbol{y}}^{(\boldsymbol{m})}(z)H^{(\boldsymbol{m})}(\boldsymbol{y}).$

Eqns (26), (27) and Proposition 2.1 (i), (ii) imply the following generalisation of Lemma 2.1

Lemma 2.2. Let $H^{(m)}(\boldsymbol{y}) \in \mathcal{M}^{(m)}$. Then

$$\nabla_{\boldsymbol{y}}^{(\boldsymbol{m})}(x)H^{(\boldsymbol{m})}(\boldsymbol{y}) = \nabla_{\mathfrak{M},\boldsymbol{y}}^{(\boldsymbol{m})}(x)H^{(\boldsymbol{m})}(\boldsymbol{y}),$$

for

(28)
$$\nabla_{\mathfrak{M},\boldsymbol{y}}^{(\boldsymbol{m})}(x) := \nabla_{\mathfrak{M}}(x) + \sum_{k=1}^{n} \left(\Psi_{\mathfrak{M}}(x,y_k) d_{y_k} + m_k d_{y_k} \left(\Psi_{\mathfrak{M}}(x,y_k) \right) \right),$$

where $\Psi_{\mathfrak{M}}(x,y)$ is an **A** independent weight (2,-1) quasiform.

We apply Lemma 2.2 to some differential equations³ discussed in [TW1, TW2, O] where $\nabla_{\mathbf{y}}^{(m)}(x)$ acts on the period matrix, 1-forms, bidifferential and projective connection of (8)-(11). Applying Lemma 2.2 these may be written as follows:

(29)
$$\nabla_{\mathfrak{M}}(x)\tau_{ab} = \nu_a(x)\nu_b(x)$$

(30)
$$\nabla_{\mathfrak{M},y}^{(1)}(x)\,\nu_a(y) = \omega(x,y)\nu_a(x),$$

(31)
$$\nabla_{\mathfrak{M},y_1,y_2}^{(1,1)}(x)\,\omega(y_1,y_2) = \omega(x,y_1)\omega(x,y_2),$$

(32)
$$\frac{1}{6}\nabla_{\mathfrak{M},y}^{(2)}(x)s(y) = \omega(x,y)^2 - \omega_2(x,y),$$

where $\omega_2(x, y)$ is the weight (2, 2) meromorphic form of (17). Eqn. (29) is equivalent to Rauch's formula [R]. Eqn. (30) implies an explicit formula for $\Psi_{\mathfrak{M}}(x, y)$:

Lemma 2.3. For all $a, b \in \mathcal{I}_+$ with $a \neq b$ we have

(33)
$$\Psi_{\mathfrak{M}}(x,y) = \frac{\omega(x,y) \begin{vmatrix} \nu_{a}(y) & \nu_{a}(x) \\ \nu_{b}(y) & \nu_{b}(x) \end{vmatrix} - \begin{vmatrix} \nu_{a}(y) & \nabla_{\mathfrak{M}}(x)\nu_{a}(y) \\ \nu_{b}(y) & \nabla_{\mathfrak{M}}(x)\nu_{b}(y) \end{vmatrix}}{\begin{vmatrix} \nu_{a}(y) & d_{y}\nu_{a}(y) \\ \nu_{b}(y) & d_{y}\nu_{b}(y) \end{vmatrix}}.$$

Proof. From (30) we find $\begin{vmatrix} \nu_a(y) & \nabla^{(1)}_{\mathfrak{M},y}(x)\nu_a(y) \\ \nu_b(y) & \nabla^{(1)}_{\mathfrak{M},y}(x)\nu_b(y) \end{vmatrix} = \omega(x,y) \begin{vmatrix} \nu_a(y) & \nu_a(x) \\ \nu_b(y) & \nu_b(x) \end{vmatrix}$ from which we solve for $\Psi_{\mathfrak{M}}(x, y)$.

³Eqns (30) and (31) appear in [O] for a postulated weight-(2, -1) quasiform $\Psi_{\mathfrak{M}}$.

2.5. Bers Potentials and Variations of Moduli. Let $\lambda(z, \overline{z})dzd\overline{z}$ be the Poincaré metric and \langle , \rangle be the associated non-singular positive definite Petersson product on \mathcal{H}_2 . A Bers potential function $f(z, \overline{z})$ for $\Phi(z) = \phi(z)dz^2 \in \mathcal{H}_2$ is a differentiable function on Ω_0 such that [Be1, TW2]

(34)
$$\frac{1}{\pi}\partial_{\overline{z}}f = \overline{\phi(z)}\,\lambda(z,\overline{z})^{-1},$$

where $\lim_{z\to 0} |z^2 f(z^{-1})| < \infty$. A potential f(z) can be constructed from a Bers quasiform $\Psi_2(x, z)$ as follows

(35)
$$f(z) = -\langle \Psi_2(\cdot, z), \Phi(\cdot) \rangle dz.$$

The potential f for Φ is unique up to an additive quadratic polynomial in z. This follows from global Möbius invariance of (2) and (14).

The Ahlfors map [A] describes a bijective antilinear map between \mathcal{H}_2 and the moduli tangent space $T(\mathfrak{M})$. This map can be realized in the Schottky parameterization as follows [TW2]. Choose local coordinates $\boldsymbol{\eta} := \eta_1, \ldots, \eta_{3g-3}$ on \mathfrak{M} giving a local $T(\mathfrak{M})$ basis $\{\partial_{\eta_r}\}$. For a given choice for r, consider a small moduli deformation $\eta_r \to \eta_r + \varepsilon$ of the Riemann surface with corresponding quasiconformal map e.g. [GL]

$$z \to w_r = z + \frac{\varepsilon}{\pi} f_r + O(\varepsilon^2),$$

for some $f_r(z, \overline{z})$. The Beltrami equation [GL] implies that

$$\mu_r := \frac{1}{\pi} \partial_{\overline{z}} f_r,$$

gives a harmonic Beltrami differential i.e. f_r is a Bers potential for $\Phi_r := \overline{\mu}_r \lambda dz^2 \in \mathcal{H}_2$ for each r. Since the moduli are independent, $\{\Phi_r\}_{r=1}^{3g-3}$ is a \mathcal{H}_2 -basis.

The deformed Riemann surface is uniformized by some Schottky group Γ^{ε} where for each Schottky generator $\gamma_a \in \Gamma$ for $a \in \mathcal{I}_+$ we define a generator $\gamma_a^{\varepsilon} \in \Gamma^{\varepsilon}$ via the compatibility condition $\gamma_a^{\varepsilon} w_r(z) = w_r(\gamma_a z)$. This implies for each $\gamma \in \Gamma$ that⁴ [TW2]

(36)
$$d(\gamma z)^{-1}\partial_{\eta_r}(\gamma_a z) = \frac{1}{\pi} \Xi_r[\gamma_a](z), \quad a \in \mathcal{I}_+,$$

where $\Xi_r[\gamma_a](z) = f_r(\gamma_a z)d(\gamma_a z)^{-1} - f_r(z)dz^{-1}$ is an Eichler cocycle that uniquely determines a potential for $\Phi_r \in \mathcal{H}_2$ via the bijective antilinear Bers map [Be1, Be2, TW2]. Eqn. (36) allows us to identify a canonical bijective antilinear Ahlfors map between ∂_{η_r} and Φ_r for each $r = 1, \ldots, 3g - 3$ in the Schottky scheme. Let $\{\Phi_r^{\vee}\}$ be the dual basis to $\{\Phi_r\}$ with respect to the Petersson product. Then $\sum_{r=1}^{3g-3} \Phi_r^{\vee}(x)\partial_{\eta_r} \in T(\mathfrak{M})$ is independent of the choice of local coordinates. Recalling $\nabla_{\mathfrak{M}}(x)$ of (24) we find

Proposition 2.2. $\nabla_{\mathfrak{M}}(x) = -\pi \sum_{r=1}^{3g-3} \Phi_r^{\vee}(x) \partial_{\eta_r}$

Proof. From (15), (18) and (20) we find $\partial_b^\ell(\gamma_a z) = -\delta_{ab}(z - w_a)^\ell \partial_z(\gamma_a z)$ so that [TW2] (37) $d(\gamma_a z)^{-1} \nabla(x)(\gamma_a z) = \Psi_2(x, \gamma_a z) - \Psi_2(x, z), \quad a \in \mathcal{I}_+.$

⁴Each $\gamma_a \in \Gamma$ depends on 3g Schottky parameters which we can locally describe as functions of the moduli $\{\eta_s\}$ and 3 global Möbius parameters.

For Φ_r paired with ∂_{η_r} under the canonical Ahlfors map we define

(38)
$$\partial_r := \langle \nabla(\cdot), \Phi_r(\cdot) \rangle = \sum_{b \in \mathcal{I}_+} \sum_{\ell \in \mathcal{L}_2} \langle \Theta_{2,b}^{\ell}, \Phi_r \rangle \partial_b^{\ell} \in T(\mathfrak{C}_g).$$

Then (37) implies

(39)
$$d(\gamma_a z)^{-1}\partial_r(\gamma_a z) = \langle \Psi_2(\cdot, \gamma_a z), \Phi_r(\cdot) \rangle - \langle \Psi_2(\cdot, z), \Phi_r(\cdot) \rangle = -\widehat{\Xi}_r[\gamma_a](z),$$

where $\hat{\Xi}_r[\gamma_a](z) = \hat{f}_r(\gamma_a z) d(\gamma_a z)^{-1} - \hat{f}_r(z) dz^{-1}$ is the Eichler cocycle for Φ_r with a potential \hat{f}_r obtained from (35). Hence $f_r = \hat{f}_r + p(z)$ for some quadratic p(z) where f_r is the Φ_r potential determined in (36). Comparing (36) and (39) we find

$$\left(\partial_r + \pi \partial_{\eta_r}\right)\left(\gamma_a z\right) = p(\gamma_a z) - p(z)\partial_z(\gamma_a z) = \mathcal{D}^p(\gamma_a z), \quad a \in \mathcal{I}_+,$$

for a global Möbius generator \mathcal{D}^p of (19). Thus $\partial_r + \pi \partial_{\eta_r} = \mathcal{D}^p$ and hence

(40)
$$\partial_r F(\boldsymbol{\eta}) = -\pi \partial_{\eta_r} F(\boldsymbol{\eta}),$$

for any differentiable function F on \mathfrak{M} . Using Lemma 2.1 and the chain rule we find

$$\nabla_{\mathfrak{M}}(x)F(\boldsymbol{\eta}) = \nabla(x)F(\boldsymbol{\eta}) = \sum_{s=1}^{3g-3} \Theta_s(x)\partial_{\eta_s}F(\boldsymbol{\eta}),$$

for $\Theta_s(x) := \nabla(x)\eta_s \in \mathcal{H}_2$. But (38) and (40) imply $\langle \Theta_s, \Phi_r \rangle = -\pi \delta_{rs}$ so that $\Theta_s(x) = -\pi \Phi_s^{\vee}(x)$ and the result follows.

We may choose 3g - 3 locally independent components of the period matrix τ of (10) as local coordinates on \mathfrak{M} (the Schottky problem). Let $\tau_{\mathcal{K}} := \ldots, \tau_{ab}, \ldots$ for $(a, b) \in \mathcal{K}$, a τ label set of cardinality 3g - 3 for such a choice. Recalling the 1-form basis $\{\nu_a\}_{a \in \mathcal{I}_+}$ of (9) and defining $\partial_{ab} := \partial_{\tau_{ab}} = \frac{1}{2\pi i} \partial_{\Omega_{ab}}$ we find

Lemma 2.4. Let $\tau_{\mathcal{K}}$ be local coordinates on \mathfrak{M} for some independent period matrix elements with label set \mathcal{K} . Then

(41)
$$\nabla_{\mathfrak{M}}(x) = \sum_{(a,b)\in\mathcal{K}} \nu_a(x)\nu_b(x)\partial_{ab}$$

where $\{\nu_a(x)\nu_b(x)\}_{(a,b)\in\mathcal{K}}$ is a \mathcal{H}_2 -basis which is the Petersson dual of the \mathcal{H}_2 -basis paired with the local $T(\mathfrak{M})$ -basis $\{-\frac{1}{\pi}\partial_{ab}\}_{(a,b)\in\mathcal{K}}$ by the canonical Ahlfors map.

Proof. Let $F(\tau_{\mathcal{K}})$ be a locally differentiable function on \mathfrak{M} . Eqn. (41) follows from the chain rule and (29) where we find

$$\nabla_{\mathfrak{M}}(x)F(\tau_{\mathcal{K}}) = \sum_{(a,b)\in\mathcal{K}} \left(\nabla_{\mathfrak{M}}(x)\tau_{ab}\right)\partial_{ab}F(\tau_{\mathcal{K}}) = \sum_{(a,b)\in\mathcal{K}} \nu_{a}(x)\nu_{b}(x)\partial_{ab}F(\tau_{\mathcal{K}}).$$

From Proposition 2.2 we find that $\{\nu_a(x)\nu_b(x)\}_{(a,b)\in\mathcal{K}}$ is a local \mathcal{H}_2 -basis dual to the \mathcal{H}_2 -basis paired with $\{-\frac{1}{\pi}\partial_{ab}\}_{(a,b)\in\mathcal{K}}$.

Finally, we remark that for any differentiable function $F(\tau)$ of the full period matrix τ then Lemma 2.1, the chain rule and (29) directly imply that

$$\nabla_{\mathfrak{M}}(x)F(\tau) = \sum_{1 \leq a \leq b \leq g} \nu_a(x)\nu_b(x)\partial_{ab}F(\tau).$$

3. Vertex Operator Algebras and Genus g Zhu Recursion

3.1. Vertex operator algebras. For indeterminates x, y we adopt the binomial expansion convention that for $m \in \mathbb{Z}$

$$(x+y)^m = \sum_{k \ge 0} \binom{m}{k} x^{m-k} y^k$$

We review some aspects of vertex operator algebras e.g. [K, FHL, LL, MT1]. A vertex operator algebra (VOA) is a quadruple $(V, Y(\cdot, \cdot), \mathbb{1}, \omega)$ consisting of a graded vector space $V = \bigoplus_{n \ge 0} V_n$, with dim $V_n < \infty$, with two distinguished elements: the vacuum vector $\mathbb{1} \in V_0$ and the Virasoro conformal vector $\omega \in V_2$. For each $v \in V$ there exists a vertex operator, a formal Laurent series in z, given by

$$Y(u,z) = \sum_{n \in \mathbb{Z}} u(n) z^{-n-1},$$

for modes $u(n) \in \text{End}(V)$. For each $u, v \in V$ we have u(n)v = 0 for all $n \gg 0$, known as *lower truncation*, and $u = u(-1)\mathbb{1}$ and $u(n)\mathbb{1} = 0$ for all $n \ge 0$, known as *creativity*. The vertex operators also obey *locality*:

$$(x-y)^{N}[Y(u,x),Y(v,y)] = 0, \quad N \gg 0.$$

For the Virasoro conformal vector

$$Y(\omega, z) = \sum_{n \in \mathbb{Z}} L(n) z^{-n-2}$$

where the operators $L(n) = \omega(n+1)$ satisfy the Virasoro algebra

$$[L(m), L(n)] = (m-n)L(m+n) + \frac{c}{2}\binom{m+1}{3}\delta_{m,-n}\mathrm{Id}_V,$$

for a constant *central charge* $c \in \mathbb{C}$. Vertex operators satisfy the *translation property*:

$$Y(L(-1)u, z) = \partial Y(u, z).$$

Finally, $V_n = \{v \in V : L(0)v = nv\}$ where $v \in V_n$ is the *(conformal) weight* wt(v) = n. We quote a number of basic VOA properties e.g. [K, FHL, LL, MT1]. For $u \in V$ of weight N we have

$$u(j): V_k \to V_{k+N-j-1}.$$

The commutator identity: for all $u, v \in V$ we have

$$[u(k), Y(v, z)] = \left(\sum_{j \ge 0} Y(u(j)v, z)\partial_z^{(j)}\right) z^k.$$

The associativity identity: for each $u, v \in V$ there exists $M \ge 0$ such that

$$(x+y)^{M}Y(Y(u,x)v,y) = (x+y)^{M}Y(u,x+y)Y(v,y).$$

Associated with the formal Möbius map $z \to \rho/z$, for a given scalar $\rho \neq 0$, we define an adjoint vertex operator [FHL, L]

$$Y_{\rho}^{\dagger}(u,z) := \sum_{n \in \mathbb{Z}} u_{\rho}^{\dagger}(n) z^{-n-1} = Y\left(e^{\frac{z}{\rho}L(1)} \left(-\frac{\rho}{z^2}\right)^{L(0)} u, \frac{\rho}{z}\right).$$

We write $Y^{\dagger}(u, z)$ for the adjoint when $\rho = 1$. For quasiprimary u (i.e. L(1)u = 0) of weight N we have

(42)
$$u_{\rho}^{\dagger}(n) = (-1)^{N} \rho^{n+1-N} u(2N-2-n),$$

e.g. $L^{\dagger}_{\rho}(n) = \rho^n L(-n)$. A bilinear form $\langle \cdot, \cdot \rangle_{\rho}$ on V is said to be invariant if

$$\langle Y(u,z)v,w\rangle_{\rho} = \langle v,Y_{\rho}^{\dagger}(u,z)w\rangle_{\rho}, \quad \forall u,v,w \in V.$$

If $\rho = 1$ then we omit the ρ subscripts. $\langle \cdot, \cdot \rangle_{\rho}$ is symmetric and $\langle u, v \rangle_{\rho} = 0$ for wt $(u) \neq$ wt(v) [FHL] with

$$\langle u, v \rangle_{\rho} = \rho^N \langle u, v \rangle, \quad N = \operatorname{wt}(u) = \operatorname{wt}(v).$$

We assume throughout this paper that V is of strong CFT-type i.e. $V_0 = \mathbb{C}\mathbb{1}$ and L(1)V = 0. Then the bilinear form with normalisation $\langle \mathbb{1}, \mathbb{1} \rangle_{\rho} = 1$ is unique [L]. We also assume that V is simple and isomorphic to the contragredient V-module V' [FHL]. Then the bilinear form is non-degenerate [L]. We refer to this unique invariant non-degenerate bilinear form as the Li-Zamolodchikov (Li-Z) metric.

3.2. Genus g correlation functions. Define the genus zero *n*-point (correlation) function for $\boldsymbol{v} := v, \ldots, v_n$ inserted at $\boldsymbol{z} := z, \ldots, z_n$, respectively, by⁵

$$Z^{(0)}(\boldsymbol{v},\boldsymbol{z}) := Z^{(0)}(\ldots;v_k,z_k;\ldots) = \langle \mathbbm{1}, \boldsymbol{Y}(\boldsymbol{v},\boldsymbol{z}) \mathbbm{1} \rangle,$$

for $(\rho = 1)$ Li-Z metric $\langle \cdot, \cdot \rangle$ and

$$\boldsymbol{Y}(\boldsymbol{v},\boldsymbol{z}) := Y(v_1,z_1)\dots Y(v_n,z_n)$$

 $Z^{(0)}(\boldsymbol{v}, \boldsymbol{z})$ can be extended to a rational function in \boldsymbol{z} in the domain $|z_1| > \ldots > |z_n|$.

We next define genus g correlation functions in terms of certain infinite sums of genus zero correlation functions based on a formal version of the Schottky sewing scheme. For each $a \in \mathcal{I}_+$, let $\{b_a\}$ denote a homogeneous V-basis and let $\{\overline{b}_a\}$ be the dual basis with respect to the Li-Z metric $\langle \cdot, \cdot \rangle$ i.e. with $\rho = 1$. Define

(43)
$$b_{-a} := \rho_a^{\operatorname{wt}(b_a)} \overline{b}_a, \quad a \in \mathcal{I}_+$$

for formal ρ_a (later identified with a Schottky sewing parameter). Then $\{b_{-a}\}$ is a dual basis for the Li-Z metric $\langle \cdot, \cdot \rangle_{\rho_a}$ with adjoint (cf. (42))

$$u_{\rho_a}^{\dagger}(m) = (-1)^N \rho_a^{m+1-N} u(2N-2-m),$$

for u quasiprimary of weight N. Let $\mathbf{b}_+ = b_1 \otimes \ldots \otimes b_g$ denote an element of a $V^{\otimes g}$ -basis. Let w_a for $a \in \mathcal{I}$ be 2g formal variables (later identified with the canonical Schottky parameters). Consider the genus zero rational 2g-point function

$$Z^{(0)}(\boldsymbol{b}, \boldsymbol{w}) = Z^{(0)}(b_1, w_1; b_{-1}, w_{-1}; \dots; b_g, w_g; b_{-g}, w_{-g})$$

for $\boldsymbol{b}, \boldsymbol{w} = b_1, w_1, b_{-1}, w_{-1}, \dots, b_g, w_g, b_{-g}, w_{-g}$. Define the genus g partition function by

(44)
$$Z_V := Z_V(\boldsymbol{w}, \boldsymbol{\rho}) = \sum_{\boldsymbol{b}_+} Z^{(0)}(\boldsymbol{b}, \boldsymbol{w})$$

for $\boldsymbol{w}, \boldsymbol{\rho} = w_1, w_{-1}, \rho_1, \dots, w_g, w_{-g}, \rho_g$ and where the sum is over any basis $\{\boldsymbol{b}_+\}$ of $V^{\otimes g}$. This definition is motivated by the sewing relation (4) and ideas in [MT2, MT3, T1, TW2]. This is similar to the sewing analysis employed in [Z2, C, DGT, G]. We suppress the genus

⁵The superscript (0) on $Z^{(0)}(\boldsymbol{v}, \boldsymbol{z})$ refers to the genus.

superscript label (g) except for genus zero. The genus g partition function is formally Möbius invariant where for \mathcal{D}^p of (19) we find [TW1]

Proposition 3.1. $\mathcal{D}^p Z_V = 0$ for any quadratic polynomial p.

Define the genus g formal n-point function for $v_1, \ldots, v_n \in V$ inserted at z_1, \ldots, z_n by

$$Z_V(v, z) := Z_V(v, z; w, \rho) = \sum_{b_+} Z^{(0)}(v, z; b, w)$$

for rational genus zero (n + 2g)-point functions

$$Z^{(0)}(\boldsymbol{v}, \boldsymbol{z}; \boldsymbol{b}, \boldsymbol{w}) = Z^{(0)}(v_1, z_1; \dots; v_n, z_n; b_{-1}, w_{-1}; \dots; b_g, w_g).$$

We also define the corresponding genus g formal n-point correlation differential form

$$\mathcal{F}_V(\boldsymbol{v}, \boldsymbol{z}) := Z_V(\boldsymbol{v}, \boldsymbol{z}) \boldsymbol{d} \boldsymbol{z}^{ ext{wt}(\boldsymbol{v})}.$$

From (43) we note that

$$\mathcal{F}_{V}(\boldsymbol{v},\boldsymbol{z}) = \sum_{n_{1},\dots,n_{g} \ge 0} \rho_{1}^{n_{1}} \dots \rho_{g}^{n_{g}} \sum_{b_{1} \in V_{n_{1}}} \dots \sum_{b_{g} \in V_{n_{g}}} Z^{(0)}(\boldsymbol{v},\boldsymbol{z};b_{1},w_{1},\dots,\overline{b}_{g},w_{-g}) \boldsymbol{d}\boldsymbol{z}^{\mathrm{wt}(\boldsymbol{v})}.$$

Thus the expansion of $\mathcal{F}_V(\boldsymbol{v}, \boldsymbol{z})$ in $\boldsymbol{\rho}$ to order $\rho_1^{n_1} \dots \rho_g^{n_g}$ has rational coefficient functions of w_a and \boldsymbol{z} and is convergent for all $(w_{\pm a}, \rho_a) \in \mathfrak{C}_g$ and for all $z_i \neq z_j \in \Omega_0$ for $i \neq j$.

Remark 3.1. If V is C_2 -cofinite then Gui's Theorem 13.1 [G] implies that $\mathcal{F}_V(\boldsymbol{v}, \boldsymbol{z})$ is absolutely and locally uniformly convergent in the Schottky sewing domain. Here we treat $\mathcal{F}_V(\boldsymbol{v}, \boldsymbol{z})$ formally since Zhu recursion does not require C_2 -cofiniteness.

The ρ expansion of Z_V has rational coefficient functions of w_a convergent for all $(w_{\pm a}, \rho_a) \in \mathfrak{C}_g$. Recalling, (3) and (5) relating $\rho_a, w_{\pm a}$ to the original Schottky parameters, we may also consider the expansion of Z_V in $\boldsymbol{q} := q_1, \ldots, q_q$. We then find

Proposition 3.2. The expansion of Z_V to any finite order $q_1^{n_1} \dots q_g^{n_g}$ is convergent on Schottky space \mathfrak{S}_g .

Proof. Since $w_a = W_a + O(q_a)$ and $\rho_a = -(W_{-a} - W_a)^2 q_a + O(q_a^2)$ we may expand Z_V in \boldsymbol{q} to order $q_1^{n_1} \dots q_g^{n_g}$ from its $\boldsymbol{\rho}$ expansion to order $\rho_1^{n_1} \dots \rho_g^{n_g}$. Furthermore, the coefficients in the \boldsymbol{q} expansion are convergent rational functions of $W_{\pm a}$ on \mathfrak{C}_g . Proposition 3.1 implies that the coefficient functions are Möbius invariant since $(W_a, q_a) \to (\sigma W_a, q_a)$ for $\sigma \in \mathrm{SL}_2(\mathbb{C})$. Thus the result follows.

3.3. Genus g Zhu recursion. We review the genus g correlation function Zhu recursion formula⁶ for a VOA V [TW1]. This generalises the original Zhu [Z1] recursion at genus zero and one. Zhu recursion is essentially a formal version of a residue expansion for meromorphic forms on Riemann surfaces [TW1, TW2].

Theorem 3.1. Let V be a simple VOA of strong CFT-type with V isomorphic to V'. The genus g correlation form for quasiprimary u of weight $N \ge 1$ inserted at $x \in S^{(g)}$ and $v_1, \ldots, v_n \in V$ inserted at $z_1, \ldots, z_n \in S^{(g)}$, respectively, satisfies

(45)
$$\mathcal{F}_{V}(u, x; \boldsymbol{v}, \boldsymbol{z}) = \sum_{a \in \mathcal{I}_{+}} \sum_{\ell \in \mathcal{L}_{N}} \Theta_{N, a}^{\ell}(x) \operatorname{Res}_{a}^{\ell} \mathcal{F}_{V}(u; \boldsymbol{v}, \boldsymbol{z})$$

 $^{^{6}}$ Zhu recursion for V-modules is discussed in ref. [TW1].

+
$$\sum_{k=1}^{n} \sum_{j \ge 0} \Psi_{N}^{(0,j)}(x, z_{k}) \mathcal{F}_{V}(\dots; u(j)v_{k}, z_{k}; \dots) dz_{k}^{j}$$

where $\operatorname{Res}_{a}^{\ell} \mathcal{F}_{V}(u; \boldsymbol{v}, \boldsymbol{z}) := \operatorname{Res}_{x-w_{a}} (x-w_{a})^{\ell} \mathcal{F}_{V}(u, x; \boldsymbol{v}, \boldsymbol{z}), \Psi_{N}(x, z)$ is the Bers quasiform (12) and $\{\Theta_{N,a}^{\ell}(x)\}$ of (15) spans \mathcal{H}_{N} .

4. Genus g Virasoro Correlation Functions

4.1. The Virasoro Ward Identity. We now apply Theorem 3.1 to compute all Virasoro correlation functions $\mathcal{F}_V(\boldsymbol{\omega}, \boldsymbol{z})$ where the Virasoro vector $\boldsymbol{\omega}$ is inserted at $\boldsymbol{z} = z_1, \ldots, z_n$. We find $\operatorname{Res}_a^{\ell} \mathcal{F}_V(\boldsymbol{\omega}) = \partial_a^{\ell} Z_V$ for ∂_a^{ℓ} of (18) so that [TW1]

(46)
$$\mathcal{F}_V(\omega, x) = \nabla(x) Z_V,$$

with $\nabla(x)$ of (20). For the rank 1 Heisenberg VOA M, we may also compute $\mathcal{F}_M(\omega, x)$ in an alternative way to obtain the following differential equation for Z_M [TW1]

(47)
$$\nabla(x)Z_M = \frac{1}{12}s(x)Z_M,$$

for projective connection s(x) of (11). In general we find that Theorem 3.1 implies the following Ward identity [TW1]:

Proposition 4.1.

(48)
$$\mathcal{F}_{V}(\omega, x; \boldsymbol{\omega}, \boldsymbol{z}) = \nabla_{\boldsymbol{z}}^{(2)}(x) \mathcal{F}_{V}(\boldsymbol{\omega}, \boldsymbol{z}) + \frac{c}{2} \sum_{k=1}^{n} \omega_{2}(x, z_{k}) \mathcal{F}_{V}(\dots; \widehat{\boldsymbol{\omega}, z_{k}}; \dots),$$

for n-tuple $\mathbf{2} = 2, ..., 2$ and where the caret denotes omission of the ω insertion at z_k and $\omega_2(xzy)$ is the symmetric meromorphic (2, 2)-form of (17).

4.2. Virasoro *n*-point generating functions.

Proposition 4.2. $\mathcal{F}_V(\boldsymbol{\omega}, \boldsymbol{z})$ is symmetric in z_i and is a generating function for all genus g n-point correlation functions for Virasoro vacuum descendants.

Proof. The proof follows that for corresponding results in [HT, GT]. $\mathcal{F}_V(\boldsymbol{z})$ is symmetric in z_1, \ldots, z_n by locality. Consider the *n*-point function for *n* Virasoro vacuum descendants $v_i = L(-k_{i1}) \ldots L(-k_{im_i}) \mathbb{1}$ for $k_{ij} \ge 2$ inserted at \boldsymbol{z}

$$Z_V(v_1, z_1; \ldots; v_n, z_n) = \sum_{\boldsymbol{b}_+} \langle \mathbb{1}, Y(v_1, z_1) \ldots Y(v_n, z_n) \boldsymbol{Y}(\boldsymbol{b}, \boldsymbol{w}) \mathbb{1} \rangle.$$

Then $\langle \mathbb{1}, Y(v_1, z_1) \dots Y(v_n, z_n) \boldsymbol{Y}(\boldsymbol{b}, \boldsymbol{w}) \mathbb{1} \rangle$ is the coefficient of $\prod_{i=1}^n \prod_{j=1}^{m_i} (x_{ij})^{k_{ij}-2}$ in

$$\langle \mathbb{1}, Y(Y(\omega, x_{11}) \dots Y(\omega, x_{1m_1})\mathbb{1}, z_1) \dots Y(Y(\omega, x_{n1}) \dots Y(\omega, x_{nm_n})\mathbb{1}, z_n) Y(\boldsymbol{b}, \boldsymbol{w})\mathbb{1} \rangle.$$

Using associativity and lower truncation (e.g. [K, LL, MT1]), we find for $N \gg 0$ that $\prod_{i=1}^{n} \prod_{j=1}^{m_{i}} (M_{ij} = M_{ij}) = M_{ij} =$

$$\prod_{i=1}^{n} \prod_{j=1}^{m} (x_{ij} + z_i)^N Y(Y(\omega, x_{11}) \dots Y(\omega, x_{1m_1}) \mathbb{1}, z_1) \dots Y(Y(\omega, x_{n1}) \dots Y(\omega, x_{nm_n}) \mathbb{1}, z_n)$$

=
$$\prod_{i=1}^{n} \prod_{j=1}^{m_i} (x_{ij} + z_i)^N Y(\omega, z_1 + x_{11}) \dots Y(\omega, z_1 + x_{1m_1}) \dots Y(\omega, z_n + x_{n1}) \dots Y(\omega, z_n + x_{nm_n})$$

Thus $Z_V(v_1, z_1; \ldots; v_n, z_n)$ is the coefficient of $\prod_{i=1}^n \prod_{j=1}^{m_i} (x_{ij})^{k_{ij}-2}$ of the formal expansion of $Z_V(\omega, z_1 + x_{11}; \ldots; \omega, z_n + x_{nm_n})$.

4.3. Genus g Virasoro graphs for Virasoro n-point functions. The recursive Ward identity (48) is not manifestly symmetric in z_1, \ldots, z_n . We now describe a symmetric expression for $\mathcal{F}_V(\boldsymbol{\omega}, \boldsymbol{z})$. We exploit the differential equations (29)–(32) and (47) to develop a symmetric graph-theoretic description. A similar approach is given in [HT] for g = 0, 1 and in [GT] for the g = 2 surface formed by sewing two tori.

We follow [HT] to define an order n Virasoro graph g_n to be a directed graph with n vertices labelled z_1, \ldots, z_n . Each z_i -vertex has degree $\deg(z_i) = 0, 1$ or 2. The degree-1 vertices can have either unit indegree or outdegree whereas the degree-2 vertices have both unit indegree and outdegree. The connected subgraphs of g_n consist of r-cycles, with $r \ge 1$ degree-2 vertices, and chains with two degree-1 end vertices with all other chain vertices of degree 2. We regard a single disconnected degree-0 vertex as a degenerate chain. Virasoro graphs are in 1-1 correspondence with the set of partial permutations on n objects, i.e. injective partial mappings from $\{z_1, \ldots, z_n\}$ to itself [HT]. Thus the number of Virasoro graphs of a given order n is given by $\sum_{i=0}^n i! {n \choose i}^2$ e.g. for n = 1 we obtain 2 graphs and for n = 2 we obtain 7 graphs.

We define weights on the components of g_n as follows. For each directed edge we define an edge weight

(49)
$$\mathcal{E}(z_i, z_j) = \begin{cases} \frac{1}{6}s(z_i) & \text{for } z_i = z_j, \\ \omega(z_i, z_j) & \text{for } z_i \neq z_j. \end{cases}$$

We note $\mathcal{E}(z_i, z_j) = \mathcal{E}(z_j, z_i)$. Suppose g_n contains M disconnected chains \mathcal{C}_m with initial vertex x_m and final vertex y_m where $m = 1, \ldots, M$

$$x_m \bigcirc \longrightarrow \cdots \longrightarrow \bigcirc y_m \cdots$$

Choose $\tau_{\mathcal{K}} := \ldots, \tau_{ab}, \ldots$ for $(a, b) \in \mathcal{K}$ as local coordinates on \mathfrak{M} as in Lemma 2.4. Define a weight associated with the M chains given by a differential operator on \mathfrak{M} expressed in terms of $\partial_{ab} := \partial_{\tau_{ab}}$ for $(a, b) \in \mathcal{K}$ as follows

(50)
$$\Delta_M(\boldsymbol{x}|\boldsymbol{y}) := \sum_{(\boldsymbol{a},\boldsymbol{b})} \boldsymbol{\nu}_{\boldsymbol{a}\boldsymbol{b}}(\boldsymbol{x},\boldsymbol{y}) \partial_{\boldsymbol{a}\boldsymbol{b}}$$

where the $(\boldsymbol{a}, \boldsymbol{b})$ sum is over all $(a_1, b_1), \ldots, (a_M, b_M) \in \mathcal{K}$ and with $\nu_{ab}(x, y) := \nu_a(x)\nu_b(y)$

$$\boldsymbol{\nu_{ab}}(\boldsymbol{x},\boldsymbol{y}) := \nu_{a_1b_1}(x_1,y_1)\dots\nu_{a_Mb_M}(x_M,y_M), \quad \partial_{\boldsymbol{ab}} := \partial_{a_1b_1}\dots\partial_{a_Mb_M}.$$

The degenerate chain $\bigcirc z$ has weight $\Delta_1(z|z) = \nabla_{\mathfrak{M}}(z)$ from Lemma 2.4.

We now define the total weight a Virasoro graph g_n containing L cycles and M chains, with endpoints x_m , y_m for $m = 1, \ldots, M$, by the following differential operator:

(51)
$$\mathcal{D}_{g_n}(\boldsymbol{z}) := \left(\frac{c}{2}\right)^L \prod_{\text{edges } (i,j)} \mathcal{E}(z_i, z_j) \Delta_M(\boldsymbol{x}|\boldsymbol{y}),$$

where the product ranges over all the edges of g_n . $\mathcal{D}_{g_n}(\boldsymbol{z})$ depends on the central charge c, the classical differentials $\omega(z_i, z_j)$, $s(z_i)$ and $\nu_a(z_i)$ and ∂_{ab} for $(a, b) \in \mathcal{K}$. Summing over the weights of inequivalent order n Virasoro graphs g_n , we define the differential operator

(52)
$$\mathcal{D}_n(\boldsymbol{z}) := \sum_{g_n} \mathcal{D}_{g_n}(\boldsymbol{z}), \quad n \ge 1.$$

It is useful to also define $\mathcal{D}_0 = 1$ and $\mathcal{D}_{-1} = 0$. $\mathcal{D}_n(\boldsymbol{z})$ is symmetric under permutations of \boldsymbol{z} variables since the set of partial permutations is so symmetric. It is instructive to

consider the first two examples. For n = 1 there are two Virasoro graphs with weights $W(z_1 \circ) = \Delta_1(z_1|z_1) = \nabla_{\mathfrak{M}}(z_1)$ and $W(z_1 \circ) = \frac{c}{12}s(z_1)$ (where W(g) denotes the weight of the displayed graph g). Hence

(53)
$$\mathcal{D}_1(z_1) = \nabla_{\mathfrak{M}}(z_1) + \frac{c}{12}s(z_1).$$

 $\mathcal{D}_2(z_1, z_2)$ is the sum of 7 Virasoro graph weights. These satisfy the following identities:

$$W(z_{1} \odot \circ z_{2}) + W(z_{1} \odot \circ z_{2}) + W(z_{2} \odot \circ z_{1}) = \nabla_{\mathfrak{M},z_{1}}^{(2)}(z_{2})\nabla_{\mathfrak{M}}(z_{1}),$$

$$W(z_{1} \circ \circ z_{2}) = \frac{c}{2}\omega(z_{1}, z_{2})^{2} = \frac{c}{12}\nabla_{\mathfrak{M},z_{1}}^{(2)}(z_{2})s(z_{1}) + \frac{c}{2}\omega_{2}(z_{1}, z_{2}),$$

$$W(z_{1} \circ \circ z_{2}) + W(z_{1} \circ \circ z_{2}) + W(z_{1} \circ \circ z_{2}) + W(z_{1} \circ z_{2}),$$

$$= \frac{c}{12}s(z_{1})\nabla_{\mathfrak{M}}(z_{2}) + \frac{c}{12}s(z_{2})\nabla_{\mathfrak{M}}(z_{1}) + \frac{c^{2}}{144}s(z_{1})s(z_{2}),$$

using Lemma 4.1 and (32). Summing we find

(54)
$$\mathcal{D}_2(z_1, z_2) = \left(\nabla^{(2)}_{\mathfrak{M}, z_1}(z_2) + \frac{c}{12}s(z_2)\right)\mathcal{D}_1(z_1) + \frac{c}{2}\omega_2(z_1, z_2)\mathcal{D}_0.$$

Notice that the order 2 Virasoro graphs result from adjoining the z_2 vertex to the order one graphs $\circ z_1$ and $z_1 \bigcirc$ in all possible ways. In general, all order n+1 graphs are of four Types determined by how vertex z_{n+1} is adjoined to an order n graph g_n as follows:

- (I) Vertex z_{n+1} is of degree zero disconnected from $g_n: \bigcirc z_{n+1}$.
- (II) Vertex z_{n+1} is of degree two disconnected from g_n : $z_{n+1} \bigcirc$.
- (III) Vertex z_{n+1} is inserted into a g_n edge $\ldots z_i \bigcirc \longrightarrow \bigcirc z_j \cdots$ giving

$$\ldots z_i \bigcirc \xrightarrow{z_{n+1}} \bigcirc z_j \cdots$$

(IV) Vertex z_{n+1} is joined to the end vertices z_i, z_j of a g_n chain giving two chains:

$$z_{n+1} \bigcirc \xrightarrow{z_i} \cdots \longrightarrow \bigcirc z_j$$
 and $z_i \bigcirc \xrightarrow{z_j} \cdots \longrightarrow \bigcirc z_{n+1}$.

Eqns. (53) and (53) are examples of an important iterative formula.

Theorem 4.1. $\mathcal{D}_n(\boldsymbol{z})$ obeys the following recursive identity for all $n \ge 0$:

(55)
$$\mathcal{D}_{n+1}(\boldsymbol{z}, z_{n+1}) = \left(\nabla_{\mathfrak{M}, \boldsymbol{z}}^{(2)}(z_{n+1}) + \frac{c}{12}s(z_{n+1})\right)\mathcal{D}_{n}(\boldsymbol{z}) + \frac{c}{2}\sum_{k=1}^{n}\omega_{2}(z_{k}, z_{n+1})\mathcal{D}_{n-1}(\dots, \hat{z}_{k}, \dots),$$

for n-tuple $\mathbf{2} = 2, ..., 2$ and where the caret denotes omission of the z_k entry and $\omega_2(x, y)$ is the symmetric meromorphic (2, 2)-form of (17).

To prove Theorem 4.1 we need the following lemma concerning derivatives of Δ_M :

Lemma 4.1.

$$\nabla_{\mathfrak{M},\boldsymbol{x},\boldsymbol{y}}^{(1)}(z)\Delta_{M}(\boldsymbol{x}|\boldsymbol{y}) = \Delta_{M+1}(\boldsymbol{x}, z|\boldsymbol{y}, z) + \sum_{m=1}^{M} \mathcal{E}(z, x_{m})\Delta_{M}(\dots, z, \dots|\boldsymbol{y})$$
$$+ \sum_{m=1}^{M} \mathcal{E}(y_{m}, z)\Delta_{M}(\boldsymbol{x}|\dots, z, \dots),$$

for 2M-tuple⁷ $\mathbf{1} = 1, ..., 1$ and where in the summands, the vertex label x_m (respectively y_m) is replaced by z in $\Delta_M(\boldsymbol{x}|\boldsymbol{y})$. *Proof.* From Lemma 2.4 and (20) and an applying the Leibniz rule of Bernerly 2.1 we find

Proof. From Lemma 2.4 and (30) and on applying the Leibniz rule of Remark 2.1 we find

$$\nabla_{\mathfrak{M},\boldsymbol{x},\boldsymbol{y}}^{(1)}(z)\Delta_{M}(\boldsymbol{x}|\boldsymbol{y}) = \sum_{(\boldsymbol{a},\boldsymbol{b})} \sum_{(a_{M+1},b_{M+1})} \nu_{a_{1}b_{1}}(x_{1},y_{1}) \dots \nu_{a_{M+1}b_{M+1}}(x_{M+1},y_{M+1})\partial_{\boldsymbol{a}\boldsymbol{b}}\partial_{a_{M+1}b_{M+1}} + \sum_{(\boldsymbol{a},\boldsymbol{b})} \nabla_{\mathfrak{M},\boldsymbol{x},\boldsymbol{y}}^{(1)}(z) \left(\nu_{a_{1}b_{1}}(x_{1},y_{1})\dots\nu_{a_{M}b_{M}}(x_{M},y_{M})\right)\partial_{\boldsymbol{a}\boldsymbol{b}}$$

$$= \Delta_{M+1}(\boldsymbol{x},z|\boldsymbol{y},z) + \sum_{m=1}^{M} \sum_{(\boldsymbol{a},\boldsymbol{b})} \left(\dots \omega(z,x_{m})\nu_{a_{m}b_{m}}(z,y_{m})\dots\right)\partial_{\boldsymbol{a}\boldsymbol{b}}$$

$$+ \sum_{m=1}^{M} \sum_{(\boldsymbol{a},\boldsymbol{b})} \left(\dots \omega(y_{m},z)\nu_{a_{m}b_{m}}(x_{m},z)\dots\right)\partial_{\boldsymbol{a}\boldsymbol{b}},$$

which is equivalent to the stated result on using (49).

Proof of Theorem 4.1. We first observe that the $\nabla_{\mathfrak{M},\mathbf{z}}^{(2)}(z_{n+1})\mathcal{D}_n(\mathbf{z})$ term in (55) contains a $\sum_{(a,b)\in\mathcal{K}}\nu_a(z_{n+1})\nu_b(z_{n+1})\mathcal{D}_n(\mathbf{z})\partial_{ab}$ contribution which is the sum all Type (I) graph weights. Furthermore, the $\frac{c}{12}s(z_{n+1})\mathcal{D}_n(\mathbf{z})$ term is the sum all Type (II) graph weights. The remaining Type (III) and (IV) graph weights in (55) arise from the action of $\nabla_{\mathfrak{M},\mathbf{z}}^{(2)}(z_{n+1})$ on the \mathbf{z} dependent parts of $\mathcal{D}_n(\mathbf{z})$. Consider g_{n+1} of Type (III) where g_n contains a 1-cycle $z_k \bigcirc$ together with some disconnected graph g_{n-1} so that $W(g_n) = \frac{c}{12}s(z_k)W(g_{n-1})$.

Using (32) we find the RHS of (55) gives rise to a contribution

$$W(g_{n-1})\frac{c}{12}\nabla_{\mathfrak{M},z_k}^{(2)}(z_{n+1})s(z_k) = \frac{c}{2}\omega(z_k,z_{n+1})^2W(g_{n-1}) - \frac{c}{2}\omega_2(z_k,z_{n+1})W(g_{n-1}).$$

The $-\frac{c}{2}\omega_2(z_{n+1}, z_k)W(g_{n-1})$ term is canceled by a summand term in (55). Thus we obtain the weights of all Type (III) graphs of the form $z_{n+1} \odot c_k$ with disconnected g_{n-1} . The remaining Type (III) terms come from any g_n edge with vertices $z_i \neq z_j$. We find such a contribution arises from terms in the RHS of (55) of the form

$$\nabla_{\mathfrak{M},z_i,z_j}^{(1,1)}(z_{n+1})\omega(z_i,z_j) = \omega(z_i,z_{n+1})\omega(z_{n+1},z_j) = W_{\mathcal{E}}(z_i \bigcirc z_{n+1}) \longrightarrow z_j),$$

using (31) and where $W_{\mathcal{E}}$ denotes the edge weights of the displayed graph.

Finally, we consider differential operators arising from the weights of disconnected chains in g_n . Thus if g_n contains a single degenerate chain $\circ z_i$, Lemma 4.1 implies

$$\nabla_{z_i}^{(2)}(z_{n+1})W(\circ z_i) = \nabla_{z_i}^{(2)}(z_{n+1})\Delta_1(z_i|z_i)$$

⁷By Remark 2.1, there is no inconsistency for any degenerate chain with $x_m = y_m$.

GENUS g VIRASORO CORRELATION FUNCTIONS FOR VERTEX OPERATOR ALGEBRAS 17

$$= \Delta_2(z_i, z_{n+1}|z_i, z_{n+1}) + \mathcal{E}(z_{n+1}, z_i)\Delta(z_{n+1}|z_i) + \mathcal{E}(z_i, z_{n+1})\Delta(z_i|z_{n+1})$$

= W($z_i \circ \cdots \circ z_{n+1}$) + W($z_{n+1} \circ \cdots \circ \circ z_i$) + W($z_i \circ \cdots \circ \circ z_{n+1}$),

corresponding to adjoining z_{n+1} with $\bigcirc z_i$. In general, if g_n contains M chains with end vertices x_m, y_m for $m = 1, \ldots, M$, then Lemma 4.1 implies

$$\nabla_{\mathfrak{M},\boldsymbol{x},\boldsymbol{y}}^{(\boldsymbol{k})}(z_{n+1})W_{\mathcal{C}}\begin{pmatrix}x_{1}\bigcirc\cdots\cdots\bigtriangledown\bigtriangledown y_{1}\\\vdots\\x_{M}\bigcirc\cdots\cdots\ldots\boxdot\lor y_{M}\end{pmatrix} = W_{\mathcal{C}}\begin{pmatrix}x_{1}\bigcirc\cdots\cdots\frown\circlearrowright y_{1}\\\vdots\\x_{M}\circlearrowright\cdots\boxdot\circlearrowright y_{M}\end{pmatrix}$$
$$+\sum_{m=1}^{M}W_{\mathcal{E}}(y_{m}\bigcirc\cdots\odot\subz_{n+1})W_{\mathcal{C}}(x_{m}\circlearrowright\cdots\cdots\circlearrowright y_{m}\circlearrowright\cdots\ldots\lor y_{m})$$
$$+\sum_{m=1}^{M}W_{\mathcal{E}}(z_{n+1}\circlearrowright\cdots\circlearrowright x_{m})W_{\mathcal{C}}(z_{n+1}\circlearrowright\cdots\circlearrowright y_{m})$$

where $W_{\mathcal{C}}$ denotes the chain weight contribution. Note that the first term is of Type (I) as already discussed whereas the additional $W_{\mathcal{E}}$ terms are all of Type (IV) found by adjoining z_{n+1} to chain endpoint vertices x_m or y_m . Thus, altogether, we confirm that (55) holds since all order n + 1 graph weights occur in the RHS of (55).

We define normalised partition and Virasoro generating n-point functions by

$$\Theta_V := Z_V Z_M^{-c}, \quad G_n(\boldsymbol{z}) := \mathcal{F}_V(\boldsymbol{\omega}, \boldsymbol{z}) Z_M^{-c},$$

where Z_M is the rank one genus g Heisenberg VOA partition function and c is the central charge for the given VOA V. The normalising factor will allow us to exploit the differential equations (29)–(32) and the Ward identity Proposition 4.1 in Theorem 4.1 below. From Proposition 3.2 we know that the expansion of Θ_V to any finite order in q_1, \ldots, q_g is convergent on \mathfrak{S}_g . Thus we may use Lemmas 2.1 and 2.2 to replace the action on Θ_V of $\nabla_{\boldsymbol{y}}^{(\boldsymbol{m})}(x)$ by $\nabla_{\mathfrak{M},\boldsymbol{y}}^{(\boldsymbol{m})}(x)$ throughout. We again choose $\tau_{\mathcal{K}} := \ldots, \tau_{ab}, \ldots$ for $(a, b) \in \mathcal{K}$ as local coordinates on \mathfrak{M} .

Theorem 4.2. The order n genus g Virasoro generating function is given by

(56)
$$G_n(\boldsymbol{z}) = \mathcal{D}_n(\boldsymbol{z})\Theta_V.$$

Proof. We prove the result by induction in n. Eqn. (56) is trivially true for n = 0 since $\mathcal{D}_0 = 1$. For n = 1 we know from (46) and (47) that

$$G_1(z_1) = Z_M^{-c} \nabla_{\mathfrak{M}}(z_1) \left(Z_M^c \Theta_V \right) = \mathcal{D}_1(z_1) \Theta_V,$$

recalling (53). The Virasoro Ward identity (48) implies

$$G_{n+1}(\boldsymbol{z}, z_{n+1}) = Z_M^{-c} \nabla_{\mathfrak{M}, \boldsymbol{z}}^{(2)}(z_{n+1}) \left(G_n(\boldsymbol{z}) Z_M^c \right) + \frac{c}{2} \sum_{k=1}^n \omega_2(z_{n+1}, z_k) G_{n-1}(\dots; \hat{z}_k; \dots)$$

= $\left(\nabla_{\mathfrak{M}, \boldsymbol{z}}^{(2)}(z_{n+1}) + \frac{c}{12} s(z_{n+1}) \right) G_n(\boldsymbol{z}) + \frac{c}{2} \sum_{k=1}^n \omega_2(z_k, z_{n+1}) G_{n-1}(\dots; \hat{z}_k; \dots),$

using (47) again. By induction, we assume that $G_n(\boldsymbol{z}) = \mathcal{D}_n(\boldsymbol{z})\Theta_V$ and $G_{n-1}(\boldsymbol{z}) = \mathcal{D}_{n-1}(\boldsymbol{z})\Theta_V$. Then we find $G_{n+1}(\boldsymbol{z}, z_{n+1}) = \mathcal{D}_{n+1}(\boldsymbol{z}, z_{n+1})\Theta_V$ using Theorem 4.1.

Remark 4.1. In [TW1] we discuss genus g partition and correlation functions for given V-modules (subject to some mild fusion rule assumptions) where the g V-basis sums of (44) are generalised to V-module basis sums together with suitable intertwiner vertex operator insertions. In particular, there is a natural generalisation of the Ward identity Proposition 4.1 and the Virasoro generating function formula of Theorem 4.1 where Θ_V is replaced by the normalised partition function for the given V-modules.

5. Modular transformations

5.1. Modular transformations of classical differentials. Consider the change of homology basis from the original marking (α, β) to $(\tilde{\alpha}, \tilde{\beta})$ where e.g. [Mu1, Fa]

(57)
$$\begin{bmatrix} \widetilde{\boldsymbol{\beta}} \\ \widetilde{\boldsymbol{\alpha}} \end{bmatrix} = \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} \boldsymbol{\beta} \\ \boldsymbol{\alpha} \end{bmatrix},$$

for a symplectic modular matrix $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \text{Sp}(2g, \mathbb{Z})$ i.e.

(58)
$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} D^T & -B^T \\ -C^T & A^T \end{bmatrix},$$

and A, B, C, D obey the relations:

(59)
$$AD^T - BC^T = A^T D - C^T B = \mathbf{I}_g,$$

(60) AB^T, A^TC, DC^T, D^TB are symmetric,

for identity matrix I_q .

Let $\nu(x) := [\nu_1(x), \dots, \nu_q(x)]$ be the row vector of 1-forms (9). It is convenient to define

(61)
$$\mathcal{M} := C\Omega + D, \quad \mathcal{N} := \mathcal{M}^{-1}.$$

Using $\oint_{\alpha_a} \nu_b = 2\pi i \, \delta_{ab}$ then (57) implies $\nu(x)$ transforms as

(62)
$$\widetilde{\nu}(x) = \nu(x)\mathcal{N}$$

This implies the period matrix (10) transforms as

(63)
$$\widetilde{\Omega} = (A\Omega + B)\mathcal{N}.$$

Lemma 5.1. $\omega(x, y)$ and s(x) transform under $\operatorname{Sp}(2g, \mathbb{Z})$ as follows:

(64)
$$\widetilde{\omega}(x,y) = \omega(x,y) - \frac{1}{2\pi i} \nu(x) \mathcal{N} C \nu^T(y),$$

(65)
$$\widetilde{s}(x) = s(x) - \frac{3}{\pi i} \nu(x) \mathcal{N} C \nu^T(x),$$

where $\mathcal{N}C$ is symmetric.

Proof. $f(x,y) := \widetilde{\omega}(x,y) - \omega(x,y)$ is a holomorphic symmetric (1,1) form in (x,y) from (8). Thus $f(x,y) = \nu(x)\mathcal{X}\nu^T(y)$ for a symmetric matrix \mathcal{X} determined by

$$\frac{1}{2\pi \mathrm{i}} \oint_{\alpha_a} \widetilde{\omega}(x, \cdot) = \frac{1}{2\pi \mathrm{i}} \oint_{\alpha_a} f(x, \cdot) = (\nu(x)\mathcal{X})_a.$$

But (58) implies
$$\alpha_a = \sum_{c \in \mathcal{I}_+} \left(-\widetilde{\beta}_c C_{ca} + \widetilde{\alpha}_c A_{ca} \right)$$
 so that

$$\oint_{\alpha_a} \widetilde{\omega}(x, \cdot) = -\sum_{c \in \mathcal{I}_+} \widetilde{\nu}_c(x) C_{ca} + 0 = -\left(\nu(x) \mathcal{N} C\right)_a,$$

using (9). Thus $\mathcal{X} = -\frac{1}{2\pi i}\mathcal{N}C$ giving (64). Eqn. (65) follows using (11).

 \mathcal{N} enjoys the following properties:

Lemma 5.2.

(i) $\mathcal{N} = A^T - C^T \widetilde{\Omega}$. (ii) $\mathcal{N}C$ is given by

$$(\mathcal{N}C)_{ab} = \begin{cases} \partial_{\Omega_{aa}} \log \det \mathcal{M} \text{ for } a = b\\ \frac{1}{2} \partial_{\Omega_{ab}} \log \det \mathcal{M} \text{ for } a \neq b. \end{cases}$$

Proof. $A^T - C^T \widetilde{\Omega} = (A^T (C\Omega + D) - C^T (A\Omega + B))\mathcal{N} = \mathcal{N}$ using (59) and (60). Thus, as already found, $\mathcal{N}C = A^T C - C^T \widetilde{\Omega} C$ is symmetric from (60). The dependence of det \mathcal{M} on Ω_{aa} arises in the *a*-th column of \mathcal{M} so that

$$\partial_{\Omega_{aa}} \det \mathcal{M} = \sum_{b \in \mathcal{I}_+} (C_{ba} + 0) \operatorname{cofact}(\mathcal{M})_{ba} = (\operatorname{adj}(\mathcal{M})C)_{aa},$$

for adjugate $\operatorname{adj}(\mathcal{M})$. Thus $\partial_{\Omega_{aa}} \log \det \mathcal{M} = (\mathcal{N}C)_{aa}$ since $\mathcal{N} := \mathcal{M}^{-1} = \operatorname{adj}(\mathcal{M})/\det \mathcal{M}$. A similar analysis applied for the $a \neq b$ cases where the dependence of $\det \mathcal{M}$ in $\Omega_{ab} = \Omega_{ba}$ arises both in the *a*-th and *b*-th columns of \mathcal{M} .

Combining Lemmas 5.1 and 5.2 and recalling Lemma 2.4 we obtain [Fa]:

Corollary 5.1. $\omega(x, y)$ and s(x) transform under $\operatorname{Sp}(2g, \mathbb{Z})$ as follows

(66)
$$\widetilde{\omega}(x,y) = \omega(x,y) - \frac{1}{2} \sum_{1 \leq a \leq b \leq g} (\nu_a(x)\nu_b(y) + \nu_b(x)\nu_a(y)) \,\partial_{ab} \log \det \mathcal{M}$$
$$\widetilde{s}(x) = s(x) - 6 \sum_{1 \leq a \leq b \leq g} \nu_a(x)\nu_b(x)\partial_{ab} \log \det \mathcal{M}$$
(67)
$$= s(x) - 6 \,\nabla_{\mathfrak{M}}(x) \log \det \mathcal{M},$$

where $\hat{\partial}_{ab} := \hat{\partial}_{\tau_{ab}} = \frac{1}{2\pi i} \hat{\partial}_{\Omega_{ab}}.$

We now consider the modular properties of the differential operators $\nabla_{\mathfrak{M},\mathbf{y}}^{(m)}(x)$ and \mathcal{D}_n that appear in (29)–(32) and (56) respectively. The change of homology basis (57) determines a new marking $(\widetilde{\boldsymbol{\alpha}}, \widetilde{\boldsymbol{\beta}})$ on an isomorphic Riemann surface for which there exists a Schottky uniformisation for some choice of parameters $\widetilde{w}_a, \widetilde{w}_{-a}$ and $\widetilde{\rho}_a$, for $a \in \mathcal{I}_+$. We let $\widetilde{\Psi}_{\mathfrak{M}}(x, y)$ denote the corresponding weight (2, -1) quasiform of Lemma 2.2 and $\widetilde{\nabla}_{\mathfrak{M}}(x)$ and $\widetilde{\nabla}_{\mathfrak{M},\mathbf{y}}^{(m)}(x)$ the differential operators corresponding to Lemma 2.1 and (28), respectively.

Proposition 5.1. Under $Sp(2,\mathbb{Z})$ modular transformations of the homology basis we find

$$(i) \ \nabla_{\mathfrak{M}}(x) = \nabla_{\mathfrak{M}}(x),$$

$$(ii) \ \widetilde{\Psi}_{\mathfrak{M}}(x, y) = \Psi_{\mathfrak{M}}(x, y)$$

$$(iii) \ \widetilde{\nabla}_{\mathfrak{M}, \mathbf{y}}^{(\mathbf{m})}(x) = \nabla_{\mathfrak{M}, \mathbf{y}}^{(\mathbf{m})}(x).$$

Proof. We first prove (i). Rauch's formula (29) in the $(\tilde{\alpha}, \hat{\beta})$ marking gives

(68)
$$\nabla_{\mathfrak{M}}(x)\widetilde{\tau}_{ab} = \widetilde{\nu}_a(x)\widetilde{\nu}_b(x)$$

Consider

$$\sum_{b\in\mathcal{I}_{+}}\nabla_{\mathfrak{M}}(x)\left(\widetilde{\Omega}_{ab}\right)\mathcal{M}_{bc} = \sum_{b\in\mathcal{I}_{+}}\nabla_{\mathfrak{M}}(x)\left((A\Omega+B)\mathcal{N}\right)_{ab}\mathcal{M}_{bc}$$
$$= \nabla_{\mathfrak{M}}(x)(A\Omega+B)_{ac} - \sum_{b\in\mathcal{I}_{+}}\widetilde{\Omega}_{ab}\nabla_{\mathfrak{M}}(x)\mathcal{M}_{bc}$$

using $\mathcal{NM} = I_g$. Rauch's formula in the original (α, β) marking implies

$$2\pi \mathrm{i} \sum_{b \in \mathcal{I}_+} \nabla_{\mathfrak{M}}(x) \left(\widetilde{\Omega}_{ab}\right) \mathcal{M}_{bc} = (\nu(x)(A^T - C^T\widetilde{\Omega}))_a \nu_c(x) = \widetilde{\nu}_a(x)\nu_c(x),$$

using Lemma 5.2 and (62). Multiplying by \mathcal{N} we therefore find $\nabla_{\mathfrak{M}}(x)\widetilde{\tau}_{ab} = \widetilde{\nu}_a(x)\widetilde{\nu}_b(x)$. Thus comparing to (68) we find $\nabla_{\mathfrak{M}}(x)\widetilde{\tau}_{ab} = \widetilde{\nabla}_{\mathfrak{M}}(x)\widetilde{\tau}_{ab}$. Choosing any 3g - 3 locally independent components of $\widetilde{\tau}$ it follows that $\nabla_{\mathfrak{M}}(x) = \widetilde{\nabla}_{\mathfrak{M}}(x)$.

We next prove (ii). In the $(\tilde{\alpha}, \tilde{\beta})$ marking, (30) gives

(69)
$$\nabla_{\mathfrak{M}}(x)\,\widetilde{\nu}_{a}(y) + d_{y}\left(\widetilde{\Psi}_{\mathfrak{M}}(x,y)\,\widetilde{\nu}_{a}(y)\right) = \widetilde{\omega}(x,y)\widetilde{\nu}_{a}(x),$$

using part (i). Consider

$$\sum_{a \in \mathcal{I}_{+}} \nabla_{\mathfrak{M}, y}^{(1)}(x) \left(\widetilde{\nu}_{a}(y)\right) \mathcal{M}_{ab} = \sum_{a \in \mathcal{I}_{+}} \nabla_{\mathfrak{M}}(x) \left(\widetilde{\nu}_{a}(y)\right) \mathcal{M}_{ab} + d_{y} \left(\Psi_{\mathfrak{M}}(x, y) \nu_{b}(y)\right)$$
$$= \nabla_{\mathfrak{M}}(x) \nu_{b}(y) - \sum_{a \in \mathcal{I}_{+}} (\nu(y) \mathcal{N})_{a} \nabla_{\mathfrak{M}}(x) \mathcal{M}_{ab} + d_{y} \left(\Psi_{\mathfrak{M}}(x, y) \nu_{b}(y)\right)$$
$$= \nabla_{\mathfrak{M}, y}^{(1)}(x) \nu_{b}(y) - \frac{1}{2\pi \mathrm{i}} \nu(y) \mathcal{N}C \nu^{T}(x) \nu_{b}(x),$$

much as before. Multiplying by \mathcal{N} and using (30) we obtain

$$\nabla_{\mathfrak{M},y}^{(1)}(x)\widetilde{\nu}_{a}(y) = \left(\omega(x,y) - \frac{1}{2\pi \mathrm{i}}\nu(y)\mathcal{N}C\nu^{T}(x)\right)\widetilde{\nu}_{a}(x) = \widetilde{\omega}(x,y)\widetilde{\nu}_{a}(x),$$

from (62). Comparing with (69) we conclude that for all $a \in \mathcal{I}_+$

$$d_y\left(\left(\widetilde{\Psi}_{\mathfrak{M}}(x,y) - \Psi_{\mathfrak{M}}(x,y)\right) \widetilde{\nu}_a(y)\right) = 0$$

 $\widetilde{\Psi}_{\mathfrak{M}}(x,y) - \Psi_{\mathfrak{M}}(x,y)$ is holomorphic in x and y so that $\widetilde{\Psi}_{\mathfrak{M}}(x,y) - \Psi_{\mathfrak{M}}(x,y) = \Theta(x)\widetilde{\nu}_{a}(y)^{-1}$ for some $\Theta \in \mathcal{H}_{2}$. But $\widetilde{\nu}_{a}(y)^{-1}$ cannot be holomorphic in y by the Riemann-Roch theorem⁸. Therefore (ii) holds. (i) and (ii) imply (iii) from definition (28).

Remark 5.1. One can also prove $\widetilde{\Psi}_{\mathfrak{M}}(x, y) = \Psi_{\mathfrak{M}}(x, y)$ directly from the expression (33) together with (62), (64) and that $\nabla_{\mathfrak{M}}(x)\mathcal{N} = -\mathcal{N}(\nabla_{\mathfrak{M}}(x)\mathcal{M})\mathcal{N} = -\mathcal{N}C\nu(x)^T\nu(x)\mathcal{N}$.

Lastly, we consider the modular properties of $\mathcal{D}_n(\boldsymbol{z})$. $\mathcal{D}_n(\boldsymbol{z})$ depends on $c, \omega(x, y), \nu_a(x), s(x)$ and ∂_{ab} for 3g - 3 locally independent τ components $\tau_{\mathcal{K}} := \ldots, \tau_{ab}, \ldots$ for $(a, b) \in \mathcal{K}$. Let $\widetilde{\mathcal{D}}_n(\boldsymbol{z})$ denote the Sp $(2g, \mathbb{Z})$ transformation (57) of $\mathcal{D}_n(\boldsymbol{z})$ using (62)-(65).

⁸The space of holomorphic weight -1 forms is trivial for $g \ge 2$.

Theorem 5.1. Let $F(\tau_{\mathcal{K}})$ be a locally differentiable function on \mathfrak{M} .

(70)
$$\widetilde{\mathcal{D}}_n(\boldsymbol{z}) \left(\det(\mathcal{M})^{c/2} F(\tau_{\mathcal{K}}) \right) = \det(\mathcal{M})^{c/2} \mathcal{D}_n(\boldsymbol{z}) F(\tau_{\mathcal{K}}).$$

Proof. We prove the result by induction in n. The result is trivially true for n = 0. For n = 1 we have

$$\widetilde{\mathcal{D}}_{1}(z_{1})\left(\det(\mathcal{M})^{c/2}F\right) = \left(\nabla_{\mathfrak{M}}(z_{1}) + \frac{c}{12}\widetilde{s}(z_{1})\right)\left(\det(\mathcal{M})^{c/2}F\right)$$
$$= \det(\mathcal{M})^{c/2}\mathcal{D}_{1}(z_{1})F,$$

using Proposition 5.1 (i) and (67). By induction let us assume that (70) holds for n and n-1. Then Theorem 4.1 implies that

$$\widetilde{\mathcal{D}}_{n+1}(\boldsymbol{z}, z_{n+1}) \det(\mathcal{M})^{c/2} F = \left(\nabla_{\mathfrak{M}, \boldsymbol{z}}^{(2)}(z_{n+1}) + \frac{c}{12} \widetilde{s}(z_{n+1})\right) \left(\det(\mathcal{M})^{c/2} \mathcal{D}_n(\boldsymbol{z}) F\right) \\ + \frac{c}{2} \sum_{k=1}^n \omega_2(z_k, z_{n+1}) \det(\mathcal{M})^{c/2} \mathcal{D}_{n-1}(\dots, \widehat{z}_k, \dots) F,$$

where from Proposition 5.1 we know $\widetilde{\nabla}^{(2)}_{\mathfrak{M},\mathbf{z}}(z_{n+1}) = \nabla^{(2)}_{\mathfrak{M},\mathbf{z}}(z_{n+1})$ and $\widetilde{\omega}_2(x,y) = \omega_2(x,y)$ (since $\widetilde{\Psi}_{\mathfrak{M}}(x,y) = \Psi_{\mathfrak{M}}(x,y)$). Then (67) implies

$$\begin{aligned} \widetilde{\mathcal{D}}_{n+1}(\boldsymbol{z}, z_{n+1}) \det(\mathcal{M})^{c/2} F \\ &= \det(\mathcal{M})^{c/2} \left(\left(\nabla_{\mathfrak{M}, \boldsymbol{z}}^{(2)}(z_{n+1}) + \frac{c}{12} s(z_{n+1}) \right) \mathcal{D}_n + \frac{c}{2} \sum_{k=2}^n \omega_2(z_1, z_k) \mathcal{D}_{n-1} \right) F \\ &= \det(\mathcal{M})^{c/2} \mathcal{D}_{n+1}(\boldsymbol{z}, z_{n+1}) F, \end{aligned}$$

by Theorem 4.1 again.

Corollary 5.2. Suppose that $F_k(\tau_{\mathcal{K}})$ is a weight k Teichmüller form on \mathfrak{M} for $\operatorname{Sp}(2g, \mathbb{Z})$ with a multiplier system. Then, with c = 2k, $\mathcal{D}_n(\boldsymbol{z})F_k(\tau_{\mathcal{K}})$ also transforms like a weight k Teichmüller form with the same multiplier system.

Proof. Let $\tilde{\tau}_{\mathcal{K}}$ denote an Sp(2g, Z) transformation on $\tau_{\mathcal{K}}$. Then $F_k(\tilde{\tau}_{\mathcal{K}}) = \varepsilon \det(\mathcal{M})^k F_k(\tau_{\mathcal{K}})$ for some multiplier ε . Therefore (70) implies that

$$\widetilde{\mathcal{D}}_{n}(\boldsymbol{z})F_{k}(\widetilde{\tau}_{\mathcal{K}}) = \varepsilon \widetilde{\mathcal{D}}_{n}(\boldsymbol{z})\left(\det(\mathcal{M})^{k}F_{k}(\tau_{\mathcal{K}})\right) = \varepsilon \det(\mathcal{M})^{k}\mathcal{D}_{n}(\boldsymbol{z})F_{k}(\tau_{\mathcal{K}}).$$

We conclude with some brief remarks on the significance of Theorems 4.1 and 5.1 in forthcoming work [T2]. Following the seminal work of Friedan and Shenker in [FS], it is believed that the conformal anomaly in CFT obstructs the existence of a global partition function on moduli space for $g \ge 2$. The genus g partition Z_M^2 function [T1] for the rank 2 Heisenberg VOA M^2 is given by the Montonen-Zograf product formula [Mo, Zo] which is the holomorphic part of the Hodge line bundle on \mathfrak{S}_g . Z_M^2 is a convergent function on \mathfrak{S}_g but by Mumford's theorem [Mu2] cannot be projected down to moduli space \mathfrak{M} (due to a non-zero Chern class which corresponds to the conformal anomaly in physics language). However, according to Friedan and Shenker, all (suitable) CFTs of a given central charge c are believed to share the same conformal anomaly. In the language of this paper, it is conjectured that $\Theta_V = Z_V Z_M^{-c}$ can be globally defined on \mathfrak{M} for some suitable class of VOAs. For a lattice VOA V_L , for a rank c even Euclidean lattice L, we know [T1] that Θ_{V_L} is the Siegel lattice theta series $\sum_{\lambda \in L^g} e^{i\pi \lambda \cdot \Omega \cdot \lambda}$ (where the sum is over $(\lambda_1, \ldots, \lambda_g)$ for all $\lambda_a \in L$). Θ_{V_L} is a Teichmüller modular form, with a multiplier system, of weight c/2 and globally defined on \mathfrak{M} . It is also been shown in [C] that for a holomorphic V then Θ_V is a Teichmüller form of weight c/2.

For many rational VOAs, such as Virasoro minimal models (e.g. [DFMS]), there exists a singular Virasoro descendant vector v of weight $2k \ge 4$ with zero 1-point genus one correlation function giving rise to a modular differential equation of order k for the genus one partition function for V and its modules. Modular differential equations based on Virasoro vectors can also be constructed by other methods e.g. [T3]. These approaches can be extended to genus g partition functions. In particular, we may extract $Z_V(v, x)Z_M^{-c}$ from $\mathcal{D}_k(z)\Theta_V$ as described in the proof of Proposition 4.2. Then, for a Virasoro singular vector, $Z_V(v, x) = 0$ determines an order k partial differential equation for Θ_V whose coefficients are holomorphic 2k-forms. Following Remark 4.1, we also find that the normalized partition function for any V-modules, satisfy the same partial differential equation. Lastly, provided the various partition functions are globally defined on \mathfrak{M} , they form a vector-valued Teichmüller form, with a multiplier system, of weight c/2using Corollary 5.2 to Theorem 5.1. Examples of VOAs with order 4 partial differential equations for Θ_V (and V-module partition functions) will be discussed in [T2].

References

- [A] Ahlfors, L., Some remarks on Teichmüller's space of Riemann surfaces, Ann. Math. 74 (1961) 171–191.
- [Be1] Bers, L., Inequalities for finitely generated Kleinian groups, J. Anal. Math. 18 23–41 (1967).
- [Be2] Bers, L., Automorphic forms for Schottky groups, Adv. Math. 16 332–361 (1975).
- [Bo] Bobenko, A., Introduction to compact Riemann surfaces, in *Computational Approach to Rie*mann Surfaces, edited Bobenko, A. and Klein, C., Springer-Verlag (Berlin, Heidelberg, 2011).
- [Bu] Burnside, W., On a class of automorphic functions, Proc. L. Math. Soc. 23 49–88 (1891).
- [C] Codogni, G., Vertex algebras and Teichmüller modular forms, arXiv:1901.03079.
- [DFMS] Di Francesco, P., Mathieu, P., Senechal, D.: Conformal Field Theory (Springer-Verlag, New York1997)
- [DGT] Damiolini, C., Gibney, A. and Tarasca, N., On factorization and vector bundles of conformal blocks from vertex algebras, Ann. Sci. Ec. Norm. Sup.57 241--292 (2024).
- [Fa] Fay, J.D., Theta Functions on Riemann Surfaces, Lecture Notes in Mathematics, Vol. 352. Springer-Verlag, (Berlin-New York, 1973).
- [FHL] Frenkel, I., Lepowsky, J. and Huang, Y.-Z., On Axiomatic Approaches to Vertex Operator Algebras and Modules, Mem. AMS 104 No. 494, (1993).
- [FK] Farkas, H.K. and I. Kra, I., *Riemann Surfaces*, Springer-Verlag (New York, 1980).
- [FS] Friedan, D. and Shenker, S.: The analytic geometry of two-dimensional conformal field theory. Nucl.Phys. B281 509–545 (1987).
- [Fo] Ford, L.R., Automorphic Functions, AMS-Chelsea, (Providence, 2004).
- [G] Gui, B., Convergence of sewing conformal blocks, Comm. Contemp. Math. **26** 2350007 (2024).
- [GL] Gardiner, F.P. and Lakic, N.: *Quasiconformal Teichmüller Theory*, Mathematical Surveys and Monographs **76**, AMS, Providence (1991).
- [GT] Gilroy, T. and Tuite, M.P., Genus two Zhu theory for vertex operator algebras, arXiv:1511.07664.
- [HT] Hurley, D. and Tuite, M.P., Virasoro correlation functions for vertex operator algebras, Int. J. Mathematics. 23 1250106 (2012)
- [K] Kac, V., Vertex Algebras for Beginners, Univ. Lect. Ser. 10, AMS, (1998).
- [L] Li, H., Symmetric invariant bilinear forms on vertex operator algebras, J. Pure. Appl. Alg. 96 279–297 (1994).

- [LL] Lepowsky, J. and Li, H., Introduction to Vertex Operator Algebras and Their Representations, Progress in Mathematics Vol. 227, Birkhäuser, (Boston, 2004).
- [McIT] McIntyre, A. and Takhtajan, L.A., Holomorphic factorization of determinants of Laplacians on Riemann surfaces and a higher genus generalization of Kronecker's first limit formula, GAFA, Geom. Funct. Anal. 16 1291–1323 (2006).
- [Mo] Montonen, C.: Multiloop amplitudes in additive dual-resonance models, Nuovo Cim. **19** 69–89 (1974).
- [MT1] Mason, G. and Tuite, M.P., Vertex operators and modular forms, A Window into Zeta and Modular Physics, eds. K. Kirsten and F. Williams, Cambridge University Press, (Cambridge, 2010), MSRI Publications 57 183–278 (2010).
- [MT2] Mason, G. and Tuite, M.P., Free bosonic vertex operator algebras on genus two Riemann surfaces I, Comm. Math. Phys. 300 673–713 (2010).
- [MT3] Mason, G. and Tuite, M.P. Free bosonic vertex operator algebras on genus two Riemann surfaces II, in *Conformal Field Theory, Automorphic Forms and Related Topics*, Contributions in Mathematical and Computational Sciences 8 183–225, Springer Verlag, (Berlin, 2014).
- [Mu1] Mumford, D., Tata Lectures on Theta I and II, Birkhäuser, (Boston, 1983).
- [Mu2] Mumford, D., Stability of projective varieties, L. Ens. Math. 23 39–110 (1977).
- [O] Odesskii, A., Deformations of complex structures on Riemann surfaces and integrable structures of Whitham type hierarchies, arXiv:1505.07779.
- [R] Rauch, H.E. On the transcendental moduli of algebraic Riemann surfaces, Proc. Nat. Acad. Sc. 11 42–48 (1955).
- [T1] Tuite, M.P., The Heisenberg generalized vertex operator algebra on a Riemann surface, Contemp. Math. 768 321–342 (2021).
- [T2] Tuite, M.P., To appear.
- [T3] Tuite, M.P, Exceptional vertex operator algebras and the Virasoro algebra, Contemp. Math. 497 213—225 (2009).
- [TW1] Tuite, M.P. and Welby, M., Genus g Zhu recursion for vertex operator algebras and their modules, arXiv:2312.13717 (2023), to appear J.Alg.Applic.
- [TW2] Tuite, M.P. and Welby, M., Some properties and applications of the Bers quasiform on a Riemann surface, arXiv:2306.08404 (2023).
- [Z1] Zhu, Y., Modular-invariance of characters of vertex operator algebras, J. Amer. Math. Soc. 9 237–302 (1996).
- [Z2] Zhu, Y., Global vertex operators on Riemann surfaces. Comm. Math. Phys. 165 485–531 (1994).
- [Zo] Zograf, P.G.: Liouville action on moduli spaces and uniformization of degenerate Riemann surfaces, (Russian) Algebra i Analiz 1 (1989) 136–160; translation in Leningrad Math. J. 1 (1990) 941–965.

School of Mathematical and Statistical Sciences, University of Galway, Galway H91 TK33, Ireland

Email address: michael.tuite@universityofgalway.ie

Email address: michael.welby5@gmail.com