NOTES ON CERTAIN BINOMIAL HARMONIC SUMS OF SUN'S TYPE

YAJUN ZHOU

ABSTRACT. We prove and generalize some recent conjectures of Z.-W. Sun on infinite series whose summands involve products of harmonic numbers and several binomial coefficients. We evaluate various classes of infinite sums in closed form by interpreting them as automorphic objects on the moduli spaces for Legendre curves $Y^{g+1} = (1 - X)^g X (1 - tX)$ of positive genera $g \in \{1, 2, 3, 5\}$.

Contents

1. Introduction	1
2. Legendre functions and Sun's series	4
2.1. Frobenius-Zagier process and Legendre-Sun series	5
2.2. Modular parametrizations of Legendre–Sun series	6
2.3. Derivatives and integrals related to Legendre–Sun series	8
3. Clausen couplings and Ramanujan–Sun series	13
3.1. Clausen couplings of Legendre functions	14
3.2. Ramanujan-Sun series	16
3.3. Some series descending from Clausen couplings	18
4. Green-Epstein variations on Sun's series	20
4.1. Derivatives of Legendre functions with respect to their degrees	23
4.2. Automorphic Green's functions and Epstein zeta functions of weight 4	24
4.3. Integral representations of certain ${}_{4}F_{3}$ series	28
5. Clebsch–Gordan couplings and binomial harmonic sums	32
5.1. Clebsch–Gordan couplings of Legendre functions	33
5.2. Bailey-Meijer formulations of Clebsch-Gordan couplings	35
5.3. Generalizations to associated Legendre functions	40
Acknowledgments	41
References	41

1. INTRODUCTION

Set binomial coefficients $\binom{m}{k} := \frac{m!}{k!(m-k)!}$ for $m \in \mathbb{Z}_{\geq 0}, k \in \mathbb{Z} \cap [0, m]$ and harmonic numbers $H_m^{(r)} := \sum_{k=1}^m \frac{1}{k^r}$ of order $r \in \mathbb{Z}_{>0}$ for $m \in \mathbb{Z}_{\geq 0}$. Customarily, one abbreviates $H_m^{(1)}$ into H_m , and simply refers to it as the *m*-th harmonic number. In this note, *binomial harmonic sums* are convergent series in the form of

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k}^{\mathscr{N}}}{k^{s}} \left(\frac{T}{2^{2\mathscr{N}}}\right)^{k} \left[\prod_{j=1}^{M} \mathsf{H}_{k-1}^{(r_{j})}\right] \left[\prod_{j'=1}^{M'} \mathsf{H}_{2k-1}^{(r'_{j'})}\right],\tag{1.1}$$

Date: March 10, 2025.

Keywords: Complex multiplication, automorphic Green's functions, Epstein zeta functions, binomial coefficients, harmonic numbers, Legendre functions

MSC 2020: 11F67, 11M06, 33C05

^{*} This research was supported in part by the Applied Mathematics Program within the Department of Energy (DOE) Office of Advanced Scientific Computing Research (ASCR) as part of the Collaboratory on Mathematics for Mesoscopic Modeling of Materials (CM4).

YAJUN ZHOU

where $\mathcal{N}, s \in \mathbb{Z}, M, M' \in \mathbb{Z}_{\geq 0}$, and |T| < 1. We will be equally interested in some variations of (1.1), as considered by Z.-W. Sun in his large collections of recent conjectures [70, 71]. For example, we may do one or more of the following things:

- allow |T| = 1 when $s \in \mathbb{Z}_{>1}$;
- trade $\binom{2k}{k}^{\mathscr{N}} k^{-s}, k \in \mathbb{Z}_{>0}$ for $\binom{2k}{k}^{\mathscr{N}} (2k+1)^{-s}, k \in \mathbb{Z}_{\geq 0}$;
- accommodate to other binomial coefficients like (^{3k}_k), (^{4k}_{2k}), and (^{6k}_{3k});
 invite harmonic numbers like H^(r)_{3k-1}, H^(r)_{4k-1}, and H^(r)_{6k-1} to the summands.

In our recent work [101, §3], we have produced closed-form evaluations for (1.1) under the additional constraint that $\mathcal{N} \in \{-1,0,1\}$, by studying integrals over Goncharov's multiple polylogarithms [45, 46], which are effectively motivic periods on the moduli space of genus-zero curves [18, 19, 20]. Here, the condition $\mathcal{N} \in \{-1, 0, 1\}$ is relevant to ε -expansions in quantum electrodynamics and quantum chromodynamics [1, 2, 3, 4, 5, 13, 16, 17, 39, 40, 58, 59, 83]. Explicit evaluations of (1.1) for $\mathcal{N} \in \{-1, 1\}$ are also available when the central binomial coefficients $\binom{2k}{k}$ in the summands are replaced by $\binom{3k}{k}$ or $\binom{4k}{2k}$ [72, 100].

In the current work devoted (mainly) to cases where $\mathcal{N} \in \{2,3,4\}$, our analytic tools will be the key players in Ramanujan's elliptic function theory [12, Chapter 33], namely

$$\sum_{k=0}^{\infty} {\binom{2k}{k}}^2 \left(\frac{t}{2^4}\right)^k = P_{-1/2}(1-2t) = \frac{1}{\pi} \int_0^1 \frac{\mathrm{d}X}{\sqrt{(1-X)X(1-tX)}},\tag{1.2}$$

$$\sum_{k=0}^{\infty} \binom{2k}{k} \binom{3k}{k} \left(\frac{t}{3^3}\right)^k = P_{-1/3}(1-2t) = \frac{\sqrt{3}}{2\pi} \int_0^1 \frac{\mathrm{d}\,X}{\sqrt[3]{(1-X)^2 X(1-tX)}},\tag{1.3}$$

$$\sum_{k=1}^{\infty} \binom{2k}{k} \binom{4k}{2k} \left(\frac{t}{2^6}\right)^k = P_{-1/4}(1-2t) = \frac{1}{\sqrt{2\pi}} \int_0^1 \frac{\mathrm{d}\,X}{\sqrt[4]{(1-X)^3}X(1-tX)},\tag{1.4}$$

$$\sum_{k=1}^{\infty} \binom{3k}{k} \binom{6k}{3k} \left(\frac{t}{2^4 3^3}\right)^k = P_{-1/6}(1-2t) = \frac{1}{2\pi} \int_0^1 \frac{\mathrm{d}\,X}{\sqrt[6]{(1-X)^5 X(1-tX)}},\tag{1.5}$$

for |t| < 1. These abelian integrals on Legendre curves $Y^{g+1} = (1-X)^g X(1-tX)$ of positive genera $g \in \{1, 2, 3, 5\}$ correspond to Sun's binomial harmonic sums for the $\mathcal{N} = 2, s = 0$ scenario [71, §2]. Hypergeometric deformations of the Legendre functions P_{ν} will generate harmonic numbers in the summands. By considering products of several Legendre functions, one can produce further analogs and extensions with different values of \mathcal{N} and s in Sun's binomial harmonic sums.

In §2, we exploit the Taylor expansion of the hypergeometric series

$${}_{2}F_{1}\left(\left.\begin{array}{c}a_{1},a_{2}\\b_{1}\end{array}\right|t\right) = 1 + \sum_{k=1}^{\infty} \frac{(a_{1})_{k}(a_{2})_{k}}{(b_{1})_{k}} \frac{t^{k}}{k!}, \quad |t| \le 1, t \ne 1,$$
(1.6)

with $(A)_k = \prod_{m=0}^{k-1} (A + m)$ being the rising factorial, and investigate the following perturbations (Lemma 2.2)

$$_{2}F_{1}\begin{pmatrix} -\nu+\varepsilon,\nu+1+\varepsilon \\ 1 \end{pmatrix} \text{ and } _{2}F_{1}\begin{pmatrix} -\nu,\nu+1 \\ 1+\varepsilon \end{pmatrix} t$$
 (1.7)

to the Legendre functions

$$P_{\nu}(1-2t) \coloneqq {}_{2}F_{1}\left(\left. \begin{array}{c} -\nu,\nu+1\\ 1 \end{array} \right| t \right) = -\frac{\sin(\nu\pi)}{\pi} \int_{0}^{1} \left[\frac{X(1-tX)}{1-X} \right]^{\nu} \frac{\mathrm{d}X}{1-X}$$
(1.8)

in the $\varepsilon \to 0$ regime. This investigation brings us, among other things, series identities like¹

$$\sum_{k=1}^{\infty} {\binom{2k}{k}}^2 \left(\frac{t}{2^4}\right)^k \left(2\mathsf{H}_{2k} - \mathsf{H}_k\right) = -\frac{P_{-1/2}(1-2t)}{2}\log(1-t),\tag{1.9}$$

$$\sum_{k=1}^{\infty} {\binom{2k}{k}}^2 \left(\frac{t}{2^4}\right)^k \mathsf{H}_k = \frac{\pi P_{-1/2}(2t-1)}{2} - \frac{P_{-1/2}(1-2t)}{2} \log \frac{2^4(1-t)}{t},\tag{1.10}$$

where $|t| \leq 1$ and $t \notin [-1,0] \cup \{1\}$. All these *t*-parametrized formulae cover and extend Sun's recent experimental discoveries [71, Conjectures 10, 12–16] as well as their analytic verifications by Wei [80, 81]. In the same section, we will also introduce modular parametrizations of series identities, to pave the way for the rest of this note.

In §3, we treat infinite sums whose addends contain $\binom{2k}{k}^3$ in their numerators. Our discussions there will also apply to other cubic forms of binomial coefficients like $\binom{2k}{k}^2\binom{3k}{k}$, $\binom{2k}{k}^2\binom{4k}{2k}$, and $\binom{2k}{k}\binom{3k}{3k}\binom{6k}{3k}$. The starting point in §3 is Clausen's coupling formula (Lemma 3.2) that represents the products of two Legendre functions as generalized hypergeometric ${}_{3}F_{2}$ series, where the Taylor expansion

$${}_{p}F_{q}\begin{pmatrix}a_{1},\ldots,a_{p}\\b_{1},\ldots,b_{q}\end{vmatrix}t) := 1 + \sum_{k=1}^{\infty} \frac{\prod_{\ell=1}^{p}(a_{\ell})_{k}}{\prod_{m=1}^{q}(b_{m})_{k}} \frac{t^{k}}{k!}$$
(1.11)

extends (1.6). A midst various applications of Clausen's coupling formula and its $\varepsilon\text{-expansions},$ we will demonstrate

$$\sum_{k=0}^{\infty} \binom{2k}{k} \binom{3k}{k} \binom{6k}{3k} \left(-\frac{1}{2^{18} \cdot 3^3 \cdot 5^3 \cdot 23^3 \cdot 29^3} \right)^k \left[2 \cdot 3^2 \cdot 7 \cdot 11 \cdot 19 \cdot 127 \cdot 163 + 3(2 \cdot 3^2 \cdot 7 \cdot 11 \cdot 19 \cdot 127 \cdot 163k + 13 \cdot 1045493)(2H_{6k} - H_{3k} - H_k) \right]$$

$$- \frac{2^7 \sqrt{3 \cdot 5^3 \cdot 23^3 \cdot 29^3}}{\pi} \log \frac{1}{2^{18} \cdot 3^3 \cdot 5^3 \cdot 23^3 \cdot 29^3}$$
(1.12)

as Sun's companion [71, Conjecture 29(iv)] to the Chudnovsky–Chudnovsky series [36, (1.5)] for $\frac{1}{\pi}$.

=

In §4 and §5, we study some binomial harmonic series that are related to Sun's studies but are not part of Sun's conjectures.

In §4, we consider integral representations for derivatives of the Legendre functions $P_{\nu}(1-2t)$ with respect to their degrees $\nu \in (-1,0)$, or equivalently, leading-order ε -expansions of

$${}_{2}F_{1}\begin{pmatrix} -\nu-\varepsilon,\nu+1+\varepsilon \\ 1 \\ \end{pmatrix}.$$
(1.13)

These efforts lead us to infinitely many analogs of Sun's series that represent automorphic Green's functions (Theorem 4.1) and Epstein zeta functions (Theorem 4.2) on the moduli spaces for the Legendre curves $Y^{g+1} = (1 - X)^g X(1 - tX)$ of genera $g \in \{1, 2, 3, 5\}$. One particular example is the following series evaluation:

$$\sum_{k=1}^{\infty} \frac{\binom{2k}{k}^2}{2^{5k}} \left[\mathsf{H}_{2k}^{(2)} - \frac{1}{4} \mathsf{H}_k^{(2)} \right] = \frac{\sqrt{\pi} \left[\Gamma(\frac{1}{4}) \right]^2}{4} \left(\frac{1}{8} - \frac{G}{\pi^2} \right), \tag{1.14}$$

¹In this note, we prescribe the complex logarithms of non-zero numbers as Re log $\xi = \log |\xi|$ and Im log $\xi = \arg \xi \in (-\pi, \pi]$, while setting fractional powers as $\xi^{\beta} := e^{\beta \log \xi}$ for $\xi \in \mathbb{C} \setminus \{0\}$.

YAJUN ZHOU

where $\Gamma(s) := \int_0^\infty e^{-t} t^{s-1} dt$ defines Euler's gamma function for s > 0 and $G := \sum_{k=0}^\infty \frac{(-1)^k}{(2k+1)^2}$ is Catalan's constant [31, 32]. In the same section, we will also revisit the Guillera–Rogers theory [54] for the $\mathcal{N} = -s = -3$ case of (1.1), using automorphic representations of Epstein zeta functions (Theorem 4.3).

In §5, we give a new perspective on our previous works [95, 96] computing certain integrals over the products of three Legendre functions. The closed forms of these integral formulae, when properly reinterpreted, grant us access to series evaluations like

$$\sum_{k=0}^{\infty} \frac{\binom{2k}{k}^4}{2^{8k}} \frac{4k+1}{(k+1)(1-2k)} = \frac{8}{\pi^2},$$
(1.15)

$$\sum_{k=0}^{\infty} \frac{\binom{2k}{k}^4}{2^{8k}} \frac{1}{4k+1} = \frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^8}{96\pi^5},\tag{1.16}$$

$$\sum_{k=0}^{\infty} \frac{\binom{2k}{k}^4}{2^{8k}} \frac{1}{4k+1} \left[\frac{2}{(4k+1)^2} - \frac{\pi^2}{3} - 2\mathsf{H}_{2k}^{(2)} + \mathsf{H}_k^{(2)} \right] = \frac{\left[\Gamma\left(\frac{1}{4}\right) \right]^8}{4\pi^5} \left(G - \frac{5\pi^2}{48} \right). \tag{1.17}$$

2. Legendre functions and Sun's series

In §2.1, we prove the theorem below by differentiating (1.7) with respect to ε .

Theorem 2.1 (Legendre–Sun series). (a) For $|t| \le 1$ and $t \ne 1$, we have the following identities:

$$\mathsf{S}_{-1/2}^{L}(t) \coloneqq 2\sum_{k=1}^{\infty} \binom{2k}{k}^{2} \left(\frac{t}{2^{4}}\right)^{k} (2\mathsf{H}_{2k} - \mathsf{H}_{k}) = -P_{-1/2}(1 - 2t)\log(1 - t), \tag{2.1}$$

$$\mathsf{S}_{-1/3}^{L}(t) \coloneqq \sum_{k=1}^{\infty} \binom{2k}{k} \binom{3k}{k} \left(\frac{t}{3^3}\right)^k (3\mathsf{H}_{3k} - \mathsf{H}_k) = -P_{-1/3}(1 - 2t)\log(1 - t), \tag{2.2}$$

$$\mathsf{S}_{-1/4}^{L}(t) \coloneqq 2\sum_{k=1}^{\infty} \binom{2k}{k} \binom{4k}{2k} \left(\frac{t}{2^{6}}\right)^{k} (2\mathsf{H}_{4k} - \mathsf{H}_{2k}) = -P_{-1/4}(1 - 2t)\log(1 - t), \tag{2.3}$$

$$\mathsf{S}_{-1/6}^{L}(t) \coloneqq \sum_{k=1}^{\infty} \binom{3k}{k} \binom{6k}{3k} \left(\frac{t}{2^4 3^3}\right)^k (6\mathsf{H}_{6k} - 3\mathsf{H}_{3k} - 2\mathsf{H}_{2k} + \mathsf{H}_k) = -P_{-1/6}(1 - 2t)\log(1 - t). \quad (2.4)$$

(b) For $|t| \le 1$ and $t \notin [-1,0] \cup \{1\}$, we have the following identities:

$$\widetilde{\mathsf{S}}_{-1/2}^{L}(t) \coloneqq -\sum_{k=1}^{\infty} \binom{2k}{k}^{2} \left(\frac{t}{2^{4}}\right)^{k} \mathsf{H}_{k} = -\frac{\pi P_{-1/2}(2t-1)}{2} + \frac{P_{-1/2}(1-2t)}{2} \log \frac{2^{4}(1-t)}{t}, \tag{2.5}$$

$$\widetilde{\mathsf{S}}_{-1/3}^{L}(t) \coloneqq -\sum_{k=1}^{\infty} \binom{2k}{k} \binom{3k}{k} \left(\frac{t}{3^3}\right)^k \mathsf{H}_k = -\frac{\pi P_{-1/3}(2t-1)}{\sqrt{3}} + \frac{P_{-1/3}(1-2t)}{2} \log \frac{3^3(1-t)}{t}, \qquad (2.6)$$

$$\widetilde{\mathsf{S}}_{-1/4}^{L}(t) \coloneqq -\sum_{k=1}^{\infty} \binom{2k}{k} \binom{4k}{2k} \left(\frac{t}{2^6}\right)^k \mathsf{H}_k = -\frac{\pi P_{-1/4}(2t-1)}{\sqrt{2}} + \frac{P_{-1/4}(1-2t)}{2} \log \frac{2^6(1-t)}{t}, \qquad (2.7)$$

$$\widetilde{\mathsf{S}}_{-1/6}^{L}(t) \coloneqq -\sum_{k=1}^{\infty} \binom{3k}{k} \binom{6k}{3k} \left(\frac{t}{2^4 3^3}\right)^k \mathsf{H}_k = -\pi P_{-1/6}(2t-1) + \frac{P_{-1/6}(1-2t)}{2} \log \frac{2^4 3^3(1-t)}{t}.$$
 (2.8)

For
$$t \in [-1,0)$$
, one has $\widetilde{\mathsf{S}}_{\nu}^{L}(t) = \lim_{\varepsilon \to 0^{+}} \widetilde{\mathsf{S}}_{\nu}^{L}(t \pm i\varepsilon^{+})$ for $\nu \in \{-\frac{1}{2}, -\frac{1}{3}, -\frac{1}{4}, -\frac{1}{6}\}$.

Remark We note that $P_{\nu}(0)$ simplifies to $\frac{\sqrt{\pi}}{\Gamma(\frac{1-\nu}{2})\Gamma(\frac{\nu+2}{2})}$ [9, §3.4(20)]. Setting $t = \frac{1}{2}$ in (2.1)–(2.4) while employing some standard transformations for Euler's gamma function, we recover Sun's identities [71, (33) and (38) for $S_{-1/4}^{L}(0)$ and $S_{-1/6}^{L}(0)$, respectively], which previously followed from the combinatorial analyses by Wei [81]. Plugging the left halves of (1.3) and (1.4) into (2.2) and (2.3), one can check that Sun's conjectures [71, (29) for $S_{-1/3}^{L}(-1/8)$, (34) for $S_{-1/4}^{L}(8/9)$, (35) for $S_{-1/4}^{L}(1/9)$, (36) for $S_{-1/4}^{L}(-1/3)$, and (37) for $S_{-1/4}^{L}(-1/63)$] hold true, as noted by Wei [80, Theorem 2.1]. The right-hand sides of [71, (29) and (34)–(37)] will be further reduced after Corollary 2.3 below.

Remark Let $\mathbf{K}(\sqrt{t}) = \frac{\pi}{2}P_{-1/2}(1-2t)$ be the complete elliptic integral of the first kind. Its relations to Legendre functions of degrees $-\frac{1}{3}$, $-\frac{1}{4}$, and $-\frac{1}{6}$ are elucidated by Ramanujan's theories for elliptic functions to alternative bases [12, Chapter 33].

Furthermore, the Chowla–Selberg formula [34, (7)] allows one to evaluate $\mathbf{K}(\sqrt{t})$ in closed form (in terms of special values of Euler's gamma function at rational arguments) whenever $z = \frac{i}{2}\mathbf{K}(\sqrt{1-t})/\mathbf{K}(\sqrt{t})$ represents a CM point [such that both Re z and (Im z)² are rational numbers, while Im z > 0]. This brings us an infinite family of series identities, such as (cf. [14, Appendix A.3])

$$\mathsf{S}_{-1/2}^{L}\left(\left(\sqrt{2}-1\right)^{2}\right) = -\frac{\Gamma(\frac{1}{8})\Gamma(\frac{3}{8})}{4\pi\sqrt{\pi}}\sqrt{\frac{2+\sqrt{2}}{2}}\left[\log 2 + \log\left(\sqrt{2}-1\right)\right],\tag{2.9}$$

$$\mathsf{S}_{-1/2}^{L}\left(\frac{2-\sqrt{3}}{4}\right) = -\frac{\left[\Gamma\left(\frac{1}{3}\right)\right]^{3}}{2\pi^{2}}\frac{\sqrt[4]{3}}{\sqrt[3]{2}}\left[\log\left(2+\sqrt{3}\right)-2\log 2\right],\tag{2.10}$$

$$\mathsf{S}_{-1/2}^{L}\left(\left(\sqrt{2}-1\right)^{4}\right) = -\frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^{2}}{8\pi\sqrt{\pi}}\left(2+\sqrt{2}\right)\left[2\log\left(\sqrt{2}-1\right)+\frac{5}{2}\log 2\right],\tag{2.11}$$

$$\mathsf{S}_{-1/2}^{L}\left(\frac{8-3\sqrt{7}}{16}\right) = -\frac{\Gamma(\frac{1}{7})\Gamma(\frac{2}{7})\Gamma(\frac{4}{7})}{2\pi^{2}}\frac{\log\left(8+3\sqrt{7}\right)-4\log 2}{\sqrt[4]{7}},\tag{2.12}$$

thereby generalizing Sun's conjectures.

Remark Sun's empirical evaluations of infinite series [71, Conjectures 10, 12–16] are motivated and accompanied by their supercongruence counterparts modulo p^2 . It is not the purpose of this note to touch upon the *p*-adic structure of the complete elliptic integrals [64] or hypergeometric ${}_2F_1$ series in general [43, 48, 73], yet the localization of Theorem 2.1 over finite fields is perhaps a worthwhile direction for future research.

In §2.2, we give modular parametrizations of the infinite series in Theorem 2.1. The modular functions and modular forms appearing therein will also be used later in §3 and §4.

In §2.3, we evaluate more binomial harmonic sums, by differentiating and integrating the formulae in Theorem 2.1.

2.1. Frobenius–Zagier process and Legendre–Sun series. Consider Euler's integral representation for hypergeometric functions [6, Theorem 2.2.1]

$${}_{2}F_{1}\left(\begin{array}{c}a,b\\c\end{array}\right|t)$$

$$=\frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)}\int_{0}^{1}\frac{X^{b-1}(1-X)^{c-b-1}}{(1-tX)^{a}}\,\mathrm{d}\,X,\quad \operatorname{Re}\,c>\operatorname{Re}\,b>0, -\pi<\arg(1-t)<\pi.$$
(2.13)

The next lemma shows that certain derivatives of $_2F_1$ with respect to its parameters a, b, and c can be expressed through Legendre functions and logarithms.

Lemma 2.2 (Frobenius–Zagier). For $v \in (-1,0)$ and $t \in (\mathbb{C} \setminus \mathbb{R}) \cup (0,1)$, the following identities hold:

$$\frac{\partial}{\partial \varepsilon} {}_{2}F_{1} \begin{pmatrix} -\nu + \varepsilon, \nu + 1 + \varepsilon \\ 1 \\ \end{pmatrix} \Big|_{\varepsilon=0}$$

$$= -P_{\nu}(1-2t)\log(1-t),$$

$$(2.14)$$

$$\frac{\partial}{\partial \varepsilon} {}_{2}F_{1} \left(\left. \begin{array}{c} -\nu, \nu + 1 \\ 1 + \varepsilon \end{array} \right| t \right) \right|_{\varepsilon=0} = \frac{\pi P_{\nu}(2t-1)}{2\sin(\nu\pi)} + \frac{P_{\nu}(1-2t)}{2} \left[-2\gamma_{0} - \psi^{(0)}(-\nu) - \psi^{(0)}(\nu+1) + \log \frac{1-t}{t} \right],$$
(2.15)

where $\psi^{(0)}(s) := \frac{d}{ds} \log \Gamma(s)$ is the digamma function, and $\gamma_0 := -\psi^{(0)}(1)$ is the Euler-Mascheroni constant.

Proof. Both (2.14) and (2.15) have been proved in [99, Lemma 2.2.1]. The main idea is as follows:

• If a function $f_{\gamma}(t) \in C^2(0,1)$ satisfies

$$\frac{\mathrm{d}}{\mathrm{d}\,t} \left[t(1-t)\frac{\mathrm{d}\,f_{\nu}(t)}{\mathrm{d}\,t} \right] + \nu(\nu+1)f_{\nu}(t) = 0 \tag{2.16}$$

for a certain $v \in (-1,0)$ and every $t \in (0,1)$, then it is a linear combination of $P_v(1-2t)$ and $P_v(2t-1)$.

• If it is also true that $f_{\nu}(0^+)$ and $f_{\nu}(1^-)$ are both finite, then $f_{\nu}(t) \equiv 0, t \in (0,1)$.

The difference between the two sides of (2.14) [or (2.15)] fits the description in the two items above, and the corresponding identity analytically continues to $t \in (\mathbb{C} \setminus \mathbb{R}) \cup (0,1)$.

Remark As shown in [99, Lemma 2.2.1 and Remark 2.2.1.1], the lemma above is inspired by the classical Frobenius process [69, (1.3.8)] and Zagier's solution for the $v = -\frac{1}{3}$ case [87], hence the first half in the title of this subsection. In [80, §2], Wei proved (2.14) via Euler's hypergeometric transformation [cf. (3.18) below], without explicitly invoking the Frobenius process. In [81, §2], Wei's approach also resulted in two special values of (2.15) for $v \in \{-\frac{1}{4}, -\frac{1}{3}\}$ and $t = \frac{1}{2}$. **Remark** By analytic continuation, the validity of (2.14) extends to $t \in (-\infty, 0]$. Moreover, the right-hand side of (2.15) remains effective for $t \in (-\infty, 0]$, so long as one reinterprets its right-hand side through limit procedures.

It is an elementary exercise on Euler's (di)gamma functions to transcribe Lemma 2.2 into the Theorem 2.1, when $\nu \in \left\{-\frac{1}{6}, -\frac{1}{4}, -\frac{1}{3}, -\frac{1}{2}\right\}$.

2.2. Modular parametrizations of Legendre–Sun series. In the following corollary to Theorem 2.1, we consider $N = 4 \sin^2(\nu \pi) \in \{1, 2, 3, 4\}$ corresponding to $\nu \in \{-\frac{1}{6}, -\frac{1}{4}, -\frac{1}{3}, -\frac{1}{2}\}$. We say that a Legendre–Sun series $S_{\nu}^{L}(t)$ or $\widetilde{S}_{\nu}^{L}(t)$ has complex multiplication if $0 < |t| \le 1$, $t \ne 1$ and

$$z_{t,N} \coloneqq \lim_{\varepsilon \to 0^+} \frac{iP_{\nu}(2(t+i\varepsilon)-1)}{\sqrt{N}P_{\nu}(1-2(t+i\varepsilon))} \in \mathfrak{H} \coloneqq \{w \in \mathbb{C} | \operatorname{Im} w > 0\}$$
(2.17)

is a CM point (namely, it is a point in the upper half-plane \mathfrak{H} that generates a quadratic extension of \mathbb{Q} , satisfying Im $z_{t,N} > 0$ and $[\mathbb{Q}(z_{t,N}) : \mathbb{Q}] = 2$). We denote the totality of algebraic numbers by $\overline{\mathbb{Q}}$.

Corollary 2.3 (Some arithmetic properties of Legendre–Sun series). *Every Legendre–Sun series with complex multiplication can be expressed in closed form:*

$$S_{\nu}^{L}(t) = P_{\nu}(1 - 2t)\mathfrak{L}_{\nu}(t), \qquad (2.18)$$

$$\widetilde{\mathsf{S}}_{\nu}^{L}(t) = P_{\nu}(1 - 2t)\widetilde{\mathfrak{L}}_{\nu}(t), \qquad (2.19)$$

along with effective algorithms to compute the left-hand sides of the following relations:

$$\mathfrak{L}_{\mathcal{V}}(t) \in \log \overline{\mathbb{Q}},\tag{2.20}$$

$$\widetilde{\mathfrak{L}}_{\nu}(t) - \pi i z_{t,N} \in \log \overline{\mathbb{Q}},$$
(2.21)

$$\log P_{\nu}(1-2t) \in \log \overline{\mathbb{Q}} + \mathbb{Q}\log \pi + \operatorname{span}_{\mathbb{Q}}\{\log \Gamma(r) | r \in \mathbb{Q} \cap (0,1)\}.$$
(2.22)

Proof. Define modular invariants

$$\alpha_N(z) := \left\{ 1 + \frac{1}{N^{6/(N-1)}} \left[\frac{\eta(z)}{\eta(Nz)} \right]^{24/(N-1)} \right\}^{-1} = 1 - \alpha_N \left(-\frac{1}{Nz} \right), \tag{2.23}$$

$$j(z) := \frac{2^8 \{1 - \alpha_4(z/2) + [\alpha_4(z/2)]^2\}^3}{[\alpha_4(z/2)]^2 [1 - \alpha_4(z/2)]^2} = j\left(-\frac{1}{z}\right),\tag{2.24}$$

for $N \in \{2,3,4\}$ by the Dedekind eta function $\eta(z) \coloneqq e^{\pi i z/12} \prod_{n=1}^{\infty} (1 - e^{2\pi i n z}), z \in \mathfrak{H}$. According to [68, §12, Lemmata 2 and 3], we have $\alpha_N(z) \in \overline{\mathbb{Q}} \cup \{\infty\}, N \in \{2,3,4\}$ and $j(z) \in \overline{\mathbb{Q}}$ when $[\mathbb{Q}(z) : \mathbb{Q}] = 2$. This verifies (2.20) and (2.21) for $t \in (\mathbb{C} \setminus \mathbb{R}) \cup (0,1)$.

By Ramanujan's base changes [12, Chapter 33, Corollary 3.4, Theorem 9.9 and Theorem 11.6], we have

$$[\eta(z)]^{24} = \begin{cases} \frac{1}{j(z)} \lim_{\varepsilon \to 0^{+}} \left[P_{-1/6} \left(\sqrt{[j(z+i\varepsilon) - 1728]/j(z+i\varepsilon)} \right) \right]^{12} \\ \frac{\alpha_{2}(z)[1-\alpha_{2}(z)]^{2}}{2^{6}} \left[P_{-1/4} (1-2\alpha_{2}(z)) \right]^{12} \\ \frac{\alpha_{3}(z)[1-\alpha_{3}(z)]^{3}}{3^{3}} \left[P_{-1/3} (1-2\alpha_{3}(z)) \right]^{12} \\ \frac{\alpha_{4}(z)[1-\alpha_{4}(z)]^{4}}{2^{4}} \left[P_{-1/2} (1-2\alpha_{4}(z)) \right]^{12} \end{cases}$$
(2.25)

if the ranges of the context-dependent z are specified by

Im
$$z > 0$$
, $|\operatorname{Re} z| < \frac{1}{2}$, $\left| z + \frac{1}{N} \right| > \frac{1}{N}$, $\left| z - \frac{1}{N} \right| > \frac{1}{N}$ (2.26)

for $N = 4 \sin^2(\nu \pi) \in \{2, 3, 4\}$, and

Im
$$z > 0$$
, $|\operatorname{Re} z| < \frac{1}{2}$, $|z| \ge 1$ (2.27)

for $N = 4\sin^2(-\frac{\pi}{6}) = 1$. The Chowla–Selberg theory [34, 68] essentially provides an explicit formula for every CM value of the Dedekind eta function $\eta(z)$, such that $\log \eta(z) \in \log \overline{\mathbb{Q}} + \mathbb{Q} \log \pi + \operatorname{span}_{\mathbb{Q}} \{\log \Gamma(r) | r \in \mathbb{Q} \cap (0,1)\}$. Therefore, we have confirmed (2.22) for $v \in \{-\frac{1}{6}, -\frac{1}{4}, -\frac{1}{3}, -\frac{1}{2}\}$ and $t \in (\mathbb{C} \setminus \mathbb{R}) \cup (0,1)$.

Thus far, our discussions are confined to the scenarios where $t \notin [-1,0)$ in the Legendre–Sun series with complex multiplications. The cases of $t \in [-1,0)$ can be handled by the limit procedures in (2.17). We leave the details to diligent readers.

Remark The left-hand sides of [71, (29) and (34)—(37)] represent Legendre–Sun series with complex multiplications, corresponding to $z_{t,3} = \frac{1+i\sqrt{3}}{2}$ and $z_{t,4} \in \left\{\frac{i}{2}, i, \frac{1+i\sqrt{3}}{2}, \frac{1+i\sqrt{7}}{2}\right\}$. After invoking Ramanujan's base changes and the Chowla–Selberg theory, we may reduce the right-hand sides of [71, (29) and (34)–(37)] to

$$\mathsf{S}_{-1/3}^{L}\left(-\frac{1}{8}\right) = -\frac{\left[\Gamma(\frac{1}{3})\right]^{3}}{2\pi^{2}}\log\frac{8}{9},\tag{S-29'}$$

$$\mathsf{S}_{-1/4}^{L}\left(\frac{8}{9}\right) = -\frac{\sqrt{6}\left[\Gamma\left(\frac{1}{4}\right)\right]^{2}}{2\pi\sqrt{\pi}}\log 3,\tag{S-34'}$$

YAJUN ZHOU

$$\mathsf{S}_{-1/4}^{L}\left(\frac{1}{9}\right) = -\frac{\sqrt{3}\left[\Gamma\left(\frac{1}{4}\right)\right]^{2}}{4\pi\sqrt{\pi}}\log\frac{9}{8},\tag{S-35'}$$

$$S_{-1/4}^{L}\left(-\frac{1}{3}\right) = -\frac{\sqrt{3}\left[\Gamma\left(\frac{1}{3}\right)\right]^{3}}{2^{11/6}\pi^{2}}\log\frac{3}{4},$$
(S-36')

$$\mathsf{S}_{-1/4}^{L}\left(-\frac{1}{63}\right) = -\frac{\sqrt{3}\Gamma(\frac{1}{7})\Gamma(\frac{2}{7})\Gamma(\frac{4}{7})}{4\sqrt{2}\pi^{2}}\log\frac{63}{64}.$$
 (S-37')

By similar procedures, one can construct

$$S_{-1/6}^{L}\left(\frac{1}{2} - \frac{7}{10}\sqrt{\frac{2}{5}}\right) = -\frac{\Gamma(\frac{1}{8})\Gamma(\frac{3}{8})}{4\pi\sqrt{\pi}}\sqrt[4]{\frac{5}{2}}\log\left(\frac{1}{2} + \frac{7}{10}\sqrt{\frac{2}{5}}\right),$$
(2.9')

$$\mathsf{S}_{-1/6}^{L}\left(\frac{1}{2} - \frac{11}{10\sqrt{5}}\right) = -\frac{\left[\Gamma\left(\frac{1}{3}\right)\right]^{3}}{4\pi^{2}} \frac{\sqrt{3}\sqrt[4]{5}}{\sqrt[3]{2}} \log\left(\frac{1}{2} + \frac{11}{10\sqrt{5}}\right),\tag{2.10'}$$

$$\mathsf{S}_{-1/6}^{L}\left(\frac{1}{2} - \frac{21}{22}\sqrt{\frac{3}{11}}\right) = -\frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^{2}}{4\pi\sqrt{2\pi}}\sqrt[4]{33}\log\left(\frac{1}{2} + \frac{21}{22}\sqrt{\frac{3}{11}}\right),\tag{2.11'}$$

$$\mathsf{S}_{-1/6}^{L}\left(\frac{1}{2} - \frac{171}{170}\sqrt{\frac{21}{85}}\right) = -\frac{\Gamma(\frac{1}{7})\Gamma(\frac{2}{7})\Gamma(\frac{4}{7})}{8\pi^2}\sqrt[4]{\frac{255}{7}}\log\left(\frac{1}{2} + \frac{171}{170}\sqrt{\frac{21}{85}}\right),\tag{2.12'}$$

as companions to (2.9)-(2.12).

2.3. Derivatives and integrals related to Legendre–Sun series.

2.3.1. Derivatives with respect to degrees of Legendre functions. Evaluating the higher-order derivatives [9, §3.4(20)]

$$\frac{\partial^{2m}}{\partial \nu^{2m}}\Big|_{\nu=-1/2} P_{\nu}(0) = \left. \frac{\partial^{2m}}{\partial \nu^{2m}} \right|_{\nu=-1/2} \frac{\sqrt{\pi}}{\Gamma\left(\frac{1-\nu}{2}\right)\Gamma\left(\frac{\nu+2}{2}\right)}, \quad m \in \mathbb{Z}_{>0}$$
(2.28)

via the $t = \frac{1}{2}$ case of

$$P_{\nu}(1-2t) = \sum_{k=0}^{\infty} \frac{(-\nu)_k (\nu+1)_k}{(k!)^2} t^k,$$
(2.29)

one obtains an infinite family of series identities. One can use (2.14)-(2.15) to compute the same set of higher-order derivatives, some of which are displayed in Table 1.

Here, the first entry in Table 1 [namely the m = 1 scenario of (2.28) and (2.29)] was singled out by Chu [35, §5, p. 588] and rediscovered in Sun's numerical experiments [71, Conjecture 9], before being reworked by Wei [80, §3].

Taking the second-order derivative of [9, \$3.4(6)]

$$\Gamma(1-\varepsilon)P_{\nu}^{\varepsilon}(0) = {}_{2}F_{1}\left(\begin{vmatrix} -\nu,\nu+1\\ 1-\varepsilon \end{vmatrix} \middle| \frac{1}{2} \right) = \frac{2^{\varepsilon}\Gamma(1-\varepsilon)\Gamma\left(\frac{1+\varepsilon+\nu}{2}\right)\cos\frac{(\nu+\varepsilon)\pi}{2}}{\sqrt{\pi}\Gamma\left(\frac{2-\varepsilon+\nu}{2}\right)}$$
(2.30)

with respect to ε , before specializing to $\varepsilon = 0$ and $v = -\frac{1}{2}$, we get

$$\sum_{k=0}^{\infty} \frac{\binom{2k}{k}^2}{2^{5k}} \left[\mathsf{H}_k^{(2)} + \mathsf{H}_k^2 \right] = \frac{2 \left[\Gamma(\frac{1}{4}) \right]^2}{\pi \sqrt{\pi}} \left[G + \left(\frac{\pi}{4} - \log 2 \right)^2 - \frac{\pi^2}{12} \right],\tag{2.31}$$

as observed by Chu [35, §5, p. 587].

TABLE 1. Selected series involving $\binom{2k}{k}^2$, where $h_k^{(r)} := \mathsf{H}_{2k}^{(r)} - \frac{1}{2^r} \mathsf{H}_k^{(r)}$, $\lambda := \log 2$, $\zeta(s) := \sum_{k=1}^{\infty} \frac{1}{k^s}$, $\beta(s) := \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)^s}$, and $G := \beta(2)$

$$\begin{split} \sum_{k=0}^{\infty} h_k^{(2)} \frac{\binom{2k}{2}^2}{2^{5k}} &= -\frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^2}{4\pi\sqrt{\pi}} \left(G - \frac{\pi^2}{8}\right) \\ &\sum_{k=0}^{\infty} \left\{h_k^{(4)} - \left[h_k^{(2)}\right]^2\right\} \frac{\binom{2k}{2}^2}{2^{5k}} &= -\frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^2}{4\pi\sqrt{\pi}} \left[\beta\left(4\right) + \frac{1}{2}\left(G - \frac{\pi^2}{8}\right)^2 - \frac{\pi^4}{96}\right] \\ &\sum_{k=0}^{\infty} \left\{h_k^{(6)} - \frac{3h_k^{(4)} - \left[h_k^{(2)}\right]^2}{2}h_k^{(2)}\right\} \frac{\binom{2k}{2}^2}{2^{5k}} &= -\frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^2}{4\pi\sqrt{\pi}} \left\{\beta\left(6\right) + \frac{3}{4}\left[\beta\left(4\right) - \frac{\pi^4}{96}\right] \left(G - \frac{\pi^2}{8}\right) \\ &+ \frac{1}{8}\left(G - \frac{\pi^2}{8}\right)^3 - \frac{\pi^6}{960}\right\} \\ &\sum_{k=0}^{\infty} \left[h_k^{(3)} - h_k^{(1)}h_k^{(2)}\right] \frac{\binom{2k}{2}^2}{2^{5k}} &= \frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^2}{16\pi\sqrt{\pi}} \left(G - \frac{\pi^2}{8}\right)\lambda \\ &\sum_{k=0}^{\infty} \left\{h_k^{(5)} - \frac{h_k^{(1)}h_k^{(4)}}{2} - h_k^{(2)}h_k^{(3)} + \frac{h_k^{(1)}\left[h_k^{(2)}\right]^2}{2}\right\} \frac{\binom{2k}{2^{5k}}^2}{2^{5k}} &= \frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^2}{32\pi\sqrt{\pi}} \left[\beta\left(4\right) + \frac{1}{2}\left(G - \frac{\pi^2}{8}\right)^2 - \frac{\pi^4}{96}\right]\lambda \\ &\sum_{k=0}^{\infty} h_k^{(2)}H_k \frac{\binom{2k}{2}^2}{2^{5k}} &= \frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^2}{8\pi\sqrt{\pi}} \left[7\zeta\left(3\right) + G\left(4\lambda - \pi\right) - \frac{\left(4\lambda + \pi\right)\pi^2}{8}\right] \\ &\sum_{k=0}^{\infty} \left\{h_k^{(4)} - \left[h_k^{(2)}\right]^2\right\} H_k \frac{\binom{2k}{2}^2}{2^{5k}} &= \frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^2}{16\pi\sqrt{\pi}} \left\{31\zeta\left(5\right) + 2\left[\beta\left(4\right) - \frac{G^2}{2}\right] \left(4\lambda - \pi\right) + 14\zeta\left(3\right)\left(G - \frac{\pi^2}{8}\right) - \frac{\pi^2 G\left(4\lambda + \pi\right)}{4} - \frac{\pi^4 \lambda}{48} - \frac{7\pi^5}{192}\right\} \end{split}$$

Applying Clausen's formula [37, last equation]

$$\frac{(\arcsin\sqrt{x})^2}{x} = {}_3F_2\left(\begin{array}{c} 1,1,1\\\frac{3}{2},2 \end{array}\right| x\right)$$
(2.32)

to Campbell's identity [24, (5.2)]

$$\sum_{k=0}^{\infty} \frac{\binom{2k}{k}^2}{2^{5k}} \mathsf{H}_k^{(2)} = \frac{\sqrt{\pi} \left[\Gamma(\frac{1}{4}) \right]^2}{12} - \frac{2\sqrt{2}}{\pi} \int_0^1 \frac{(\arcsin\sqrt{x})^2}{\sqrt{x}\sqrt{1-x^2}} \, \mathrm{d} \, x, \tag{2.33}$$

we obtain

$$\sum_{k=0}^{\infty} \frac{\binom{2k}{k}^2}{2^{5k}} \mathsf{H}_k^{(2)} = \frac{\sqrt{\pi} \left[\Gamma(\frac{1}{4}) \right]^2}{12} - \frac{\left[\Gamma(\frac{1}{4}) \right]^2}{9\pi\sqrt{\pi}} {}_4F_3 \left(\frac{1,1,1,\frac{3}{2}}{\frac{7}{4},\frac{7}{4},2} \right| 1 \right) - \frac{8\sqrt{\pi}}{\left[\Gamma(\frac{1}{4}) \right]^2} {}_4F_3 \left(\frac{\frac{1}{2},\frac{1}{2},1,1}{\frac{5}{4},\frac{5}{4},\frac{3}{2}} \right| 1 \right)$$
(2.34)

through termwise integration of the Taylor expansion for $_{3}F_{2}$ [see (1.11)]. Currently, we do not know how to reduce the two $_{4}F_{3}$ expressions in (2.34) to well-known mathematical constants. Numerical experiments based on the PSLQ algorithm seem to suggest that both

$$\left\{ {}_{4}F_{3} \left(\left. \begin{array}{c} 1,1,1,\frac{3}{2} \\ \frac{7}{4},\frac{7}{4},2 \end{array} \right| 1 \right), G, \pi^{2}, \pi \log 2, \log^{2} 2 \right\}$$
(2.35)

and

$$\left\{ {}_{4}F_{3}\left(\left. \frac{1}{2}, \frac{1}{2}, 1, 1 \right| \\ \frac{5}{4}, \frac{5}{4}, \frac{3}{2} \right| 1 \right), G, \pi^{2}, \pi \log 2, \log^{2} 2 \right\}$$
(2.36)

are linearly independent over \mathbb{Q} .

Unlike our previous analysis [101, §3] of (1.1) for arbitrary polynomials of harmonic numbers in the summands, we have fewer degrees of freedom when $\binom{2k}{k}^2$ is brought into play. Discounting the ${}_4F_3$ expressions in (2.34), we are still unable to render the individual sums [26, 35]

$$\sum_{k=0}^{\infty} \frac{\binom{2k}{k}^2}{2^{5k}} \mathsf{H}_k^2, \quad \sum_{k=0}^{\infty} \frac{\binom{2k}{k}^2}{2^{5k}} \mathsf{H}_k^{(2)}, \quad \sum_{k=0}^{\infty} \frac{\binom{2k}{k}^2}{2^{5k}} \mathsf{H}_{2k}^2, \quad \sum_{k=0}^{\infty} \frac{\binom{2k}{k}^2}{2^{5k}} \mathsf{H}_{2k}^{(2)}$$
(2.37)

into closed forms.

2.3.2. Integrals involving Legendre–Sun series. By termwise integrations over the series representations of $S_{\nu}^{L}(t)$ and $\tilde{S}_{\nu}^{L}(t)$, one can evaluate several classes of non-trivial infinite sums, as described in the next two corollaries.

Corollary 2.4 (Binomial modulations of Legendre-Sun series). (a) We have

$$\int_{0}^{1} \frac{\mathsf{S}_{-1/2}^{L}(t)}{\sqrt{1-t}} \,\mathrm{d}\,t = 4\sum_{k=0}^{\infty} \frac{\binom{2k}{k}}{2^{2k}} \frac{2\mathsf{H}_{2k} - \mathsf{H}_{k}}{2k+1} = 4\pi\log 2,\tag{2.38}$$

$$\int_{0}^{1} \frac{\mathsf{S}_{-1/3}^{L}(t)}{\sqrt{1-t}} \,\mathrm{d}\,t = 2\sum_{k=0}^{\infty} \frac{2^{2k} \binom{3k}{k}}{3^{3k}} \frac{3\mathsf{H}_{3k} - \mathsf{H}_{k}}{2k+1} = 9(2 - \log 3),\tag{2.39}$$

$$\int_{0}^{1} \frac{\mathsf{S}_{-1/4}^{L}(t)}{\sqrt{1-t}} \,\mathrm{d}\,t = 4\sum_{k=0}^{\infty} \frac{\binom{4k}{2k}}{2^{4k}} \frac{2\mathsf{H}_{4k} - \mathsf{H}_{2k}}{2k+1} = 4\sqrt{2}(2-\log 2),\tag{2.40}$$

$$\int_{0}^{1} \frac{\mathsf{S}_{-1/6}^{L}(t)}{\sqrt{1-t}} \,\mathrm{d}\,t = 2\sum_{k=0}^{\infty} \frac{\binom{3k}{k}\binom{6k}{3k}}{(2^{2}3^{3})^{k}\binom{2k}{k}} \frac{\mathsf{6H}_{6k} - 3\mathsf{H}_{3k} - 2\mathsf{H}_{2k} + \mathsf{H}_{k}}{2k+1} = \frac{3\sqrt{3}}{2} \left(3 + \log\frac{2^{4}}{3^{3}}\right). \tag{2.41}$$

More generally, for $m \in \mathbb{Z}_{\geq 0}$, we have

$$\int_{0}^{1} \frac{\mathsf{S}_{-1/2}^{L}(t)}{(1-t)^{\frac{1}{2}-m}} \,\mathrm{d}\,t = 4 \sum_{k=0}^{\infty} \frac{\binom{2k}{k}}{2^{2k}} \frac{(2\mathsf{H}_{2k} - \mathsf{H}_{k})(2k-1)!!(2m-1)!!}{(2k+2m+1)!!} \\ \in \mathbb{Q}\pi + \mathbb{Q}\pi\log 2,$$
(2.42)

$$\int_{0}^{1} \frac{\mathsf{S}_{-1/3}^{L}(t)}{(1-t)^{\frac{1}{2}-m}} \,\mathrm{d}\,t = 2\sum_{k=0}^{\infty} \frac{2^{2k} \binom{3k}{k}}{3^{3k}} \frac{(3\mathsf{H}_{3k}-\mathsf{H}_{k})(2k-1)!!(2m-1)!!}{(2k+2m+1)!!}$$

$$\in \mathbb{Q} + \mathbb{Q}\log 3,$$
(2.43)

$$\int_{0}^{1} \frac{\mathsf{S}_{-1/4}^{L}(t)}{(1-t)^{\frac{1}{2}-m}} \,\mathrm{d}\,t = 4\sum_{k=0}^{\infty} \frac{\binom{4k}{2k}}{2^{4k}} \frac{(2\mathsf{H}_{4k}-\mathsf{H}_{2k})(2k-1)!!(2m-1)!!}{(2k+2m+1)!!} \\ \in \mathbb{Q}\sqrt{2} + \mathbb{Q}\sqrt{2}\log 2,$$

$$(2.44)$$

$$\int_{0}^{1} \frac{\mathsf{S}_{-1/6}^{L}(t)}{(1-t)^{\frac{1}{2}-m}} \,\mathrm{d}\,t = 2\sum_{k=0}^{\infty} \frac{\binom{6k}{3k}\binom{3k}{k}}{(2^{2}3^{3})^{k}\binom{2k}{k}} \frac{(6\mathsf{H}_{6k} - 3\mathsf{H}_{3k} - 2\mathsf{H}_{2k} + \mathsf{H}_{k})(2k-1)!!(2m-1)!!}{(2k+2m+1)!!} \\ \in \mathbb{Q}\sqrt{3} + \mathbb{Q}\sqrt{3}\log\frac{2^{4}}{3^{3}}, \tag{2.45}$$

where the coefficients in the Q-linear combinations are explicitly computable. [Here, double factorials of odd integers are defined as (-1)!! := 1 and $(2n+1)!! := \frac{(2n+1)!}{2^n n!}$ for $n \in \mathbb{Z}_{\geq 0}$.]

(b) We have

$$\int_{0}^{1} \frac{\mathsf{S}_{-1/2}^{L}(t) + \widetilde{\mathsf{S}}_{-1/2}^{L}(t)}{\pi\sqrt{t(1-t)}} \,\mathrm{d}\,t = \sum_{k=0}^{\infty} \frac{\binom{2k}{k}^{3}}{2^{6k}} (4\mathsf{H}_{2k} - 3\mathsf{H}_{k}) = \frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^{4}}{2\pi^{3}}\log 2,\tag{2.46}$$

$$\int_{0}^{1} \frac{\mathsf{S}_{-1/3}^{L}(t) + \widetilde{\mathsf{S}}_{-1/3}^{L}(t)}{\pi\sqrt{t(1-t)}} \,\mathrm{d}\,t = \sum_{k=0}^{\infty} \frac{\binom{2k}{k}^{2}\binom{3k}{k}}{2^{2k}3^{3k}} (3\mathsf{H}_{3k} - 2\mathsf{H}_{k}) = \frac{9\sqrt[3]{2} \left[\Gamma\left(\frac{1}{3}\right)\right]^{6}}{16\pi^{4}} \log 2,\tag{2.47}$$

$$\int_{0}^{1} \frac{\mathsf{S}_{-1/4}^{L}(t) + \widetilde{\mathsf{S}}_{-1/4}^{L}(t)}{\pi\sqrt{t(1-t)}} \,\mathrm{d}\,t = \sum \frac{\binom{2k}{k}^{2}\binom{4k}{2k}}{2^{8k}} (4\mathsf{H}_{4k} - 2\mathsf{H}_{2k} - \mathsf{H}_{k}) = \frac{2\pi\log 2}{\left[\Gamma(\frac{5}{8})\Gamma(\frac{7}{8})\right]^{2}},\tag{2.48}$$

$$\int_{0}^{1} \frac{\mathsf{S}_{-1/6}^{L}(t) + \widetilde{\mathsf{S}}_{-1/6}^{L}(t)}{\pi\sqrt{t(1-t)}} \,\mathrm{d}\,t = \sum_{k=0}^{\infty} \frac{\binom{2k}{k}\binom{3k}{k}\binom{6k}{3k}}{2^{6k}3^{3k}} (6\mathsf{H}_{6k} - 3\mathsf{H}_{3k} - 2\mathsf{H}_{2k}) = \frac{2\pi\log 2}{\left[\Gamma(\frac{7}{12})\Gamma(\frac{11}{12})\right]^2}.$$
 (2.49)

More generally, for $m \in \mathbb{Z}_{\geq 0}$, we have

$$\begin{split} \int_{0}^{1} \frac{S_{-1/2}^{L}(t) + \widetilde{S}_{-1/2}^{L}(t)}{\pi[t(1-t)]^{\frac{1}{2}-m}} dt &= \sum_{k=0}^{\infty} \frac{\binom{2k}{k}^{3}}{2^{6k+3m}} \frac{(4H_{2k} - 3H_{k})(2k + 2m - 1)!!}{\binom{k+2m}{k}(2k - 1)!!m!} \tag{2.50} \\ &\in \mathbb{Q} \frac{\left[\Gamma(\frac{1}{4})\right]^{4}}{\pi^{3}} + \mathbb{Q} \frac{\left[\Gamma(\frac{1}{4})\right]^{4}}{\pi^{3}} \log 2, \\ \int_{0}^{1} \frac{S_{-1/3}^{L}(t) + \widetilde{S}_{-1/3}^{L}(t)}{\pi[t(1-t)]^{\frac{1}{2}-m}} dt &= \sum_{k=0}^{\infty} \frac{\binom{2k}{k}^{2}\binom{3k}{k}}{2^{2k+3m}3^{3k}} \frac{(3H_{3k} - 2H_{k})(2k + 2m - 1)!!}{(k+2m)(2k - 1)!!m!} \\ &\in \mathbb{Q} \frac{\sqrt[3]{2}[\Gamma(\frac{1}{3})]^{6}}{\pi^{4}} + \mathbb{Q} \frac{\sqrt[3]{2}[\Gamma(\frac{1}{3})]^{6}}{\pi^{4}} \log 2, \\ \int_{0}^{1} \frac{S_{-1/4}^{L}(t) + \widetilde{S}_{-1/4}^{L}(t)}{\pi[t(1-t)]^{\frac{1}{2}-m}} dt &= \sum_{k=0}^{\infty} \frac{\binom{2k}{k}\binom{2k}{2k}}{2^{2k+3m}} \frac{(4H_{4k} - 2H_{2k} - H_{k})(2k + 2m - 1)!!}{(k+2m)(2k - 1)!!m!} \\ &\in \mathbb{Q} \frac{\pi}{[\Gamma(\frac{5}{8})\Gamma(\frac{7}{8})]^{2}} + \mathbb{Q} \frac{\pi \log 2}{[\Gamma(\frac{5}{8})\Gamma(\frac{7}{8})]^{2}}, \\ \int_{0}^{1} \frac{S_{-1/6}^{L}(t) + \widetilde{S}_{-1/6}^{L}(t)}{\pi[t(1-t)]^{\frac{1}{2}-m}} dt &= \sum_{k=0}^{\infty} \frac{\binom{2k}{k}\binom{3k}{k}\binom{3k}{3k}}{2^{6k+3m}3^{3k}} \frac{(6H_{6k} - 3H_{3k} - 2H_{2k})(2k + 2m - 1)!!}{(k+2m)(2k - 1)!!m!} \\ &\in \mathbb{Q} \frac{\pi}{[\Gamma(\frac{5}{8})\Gamma(\frac{7}{8})]^{2}} + \mathbb{Q} \frac{\pi \log 2}{[\Gamma(\frac{5}{8})\Gamma(\frac{7}{8})]^{2}}, \\ &\int_{0}^{1} \frac{S_{-1/6}^{L}(t) + \widetilde{S}_{-1/6}^{L}(t)}{\pi[t(1-t)]^{\frac{1}{2}-m}} dt &= \sum_{k=0}^{\infty} \frac{\binom{2k}{k}\binom{3k}{k}\binom{3k}{3k}}}{2^{6k+3m}3^{3k}} \frac{(6H_{6k} - 3H_{3k} - 2H_{2k})(2k + 2m - 1)!!}{(k+2m)(2k - 1)!!m!} \\ &\in \mathbb{Q} \frac{\pi}{[\Gamma(\frac{7}{12})\Gamma(\frac{11}{12})]^{2}} + \mathbb{Q} \frac{\pi \log 2}{[\Gamma(\frac{5}{12})\Gamma(\frac{11}{12})]^{2}}, \end{aligned}$$

where the coefficients in the $\mathbb Q$ -linear combinations are explicitly computable.

Proof. (a) We paraphrase [47, item 7.127] (which reproduces [10, §18.1(15)]) as

$$\int_{0}^{1} P_{\nu}(1-2t)(1-t)^{\sigma} dt = \frac{[\Gamma(1+\sigma)]^{2}}{\Gamma(2+\nu+\sigma)\Gamma(1-\nu+\sigma)}, \quad \text{Re}\,\sigma > -1.$$
(2.54)

YAJUN ZHOU

In particular, for each $m \in \mathbb{Z}_{\geq 0}$, the derivative of the formula above at $\sigma = m - \frac{1}{2}$ brings us

$$\int_{0}^{1} \frac{P_{\nu}(1-2t)}{(1-t)^{\frac{1}{2}-m}} \log(1-t) dt$$

$$= \frac{\left[\Gamma(m+\frac{1}{2})\right]^{2}}{\Gamma(m-\nu+\frac{1}{2})\Gamma(m+\nu+\frac{3}{2})} \left[-\psi^{(0)}\left(m-\nu+\frac{1}{2}\right) - \psi^{(0)}\left(m+\nu+\frac{3}{2}\right) + 2\psi^{(0)}\left(m+\frac{1}{2}\right)\right]$$
(2.55)

for $\nu \neq -1/2$. Exploring the $\nu \to -1/2$ limit and the special cases of $\nu \in \{-\frac{1}{3}, -\frac{1}{4}, -\frac{1}{6}\}$, while bearing in mind that

$$\int_{0}^{1} \frac{t^{k} dt}{(1-t)^{\frac{1}{2}-m}} = \frac{2^{1+2k}}{\binom{2k}{k}} \frac{(2k-1)!!(2m-1)!!}{(2k+2m+1)!!},$$
(2.56)

we may turn Theorem 2.1(a) into (2.38)-(2.45).

(b) We specialize [47, item 7.132.1] (which corrects [10, §18.1(16)]) into

$$\int_{0}^{1} P_{\nu}(1-2t) [t(1-t)]^{\sigma} dt$$

$$= \frac{\pi}{2^{1+2\sigma}} \frac{[\Gamma(1+\sigma)]^{2}}{\Gamma(\frac{\nu+3}{2}+\sigma)\Gamma(\sigma-\frac{\nu-2}{2})\Gamma(\frac{2+\nu}{2})\Gamma(\frac{1-\nu}{2})}, \quad \text{Re}\,\sigma > -1.$$
(2.57)

For each $m \in \mathbb{Z}_{\geq 0}$, the zeroth- and first-order derivatives of the equation above at $\sigma = m - \frac{1}{2}$ leave us

$$\int_{0}^{1} \frac{P_{\nu}(1-2t)}{[t(1-t)]^{\frac{1}{2}-m}} dt = \int_{0}^{1} \frac{P_{\nu}(2t-1)}{[t(1-t)]^{\frac{1}{2}-m}} dt$$

$$= \frac{\pi \left[\Gamma(m+\frac{1}{2})\right]^{2}}{2^{2m}\Gamma(\frac{1-\nu}{2})\Gamma(\frac{\nu+2}{2})\Gamma(m-\frac{\nu-1}{2})\Gamma(m+\frac{\nu+1}{2})} =: M_{m,\nu}$$
(2.58)

and

$$\int_{0}^{1} \frac{P_{\nu}(1-2t)}{[t(1-t)]^{\frac{1}{2}-m}} [\log t + \log(1-t)] dt$$

$$= \left[2\psi^{(0)} \left(m + \frac{1}{2} \right) - \psi^{(0)} \left(m + \frac{\nu+1}{2} \right) - \psi^{(0)} \left(m - \frac{\nu-1}{2} \right) - 2\log 2 \right] M_{m,\nu}.$$
(2.59)

As in our proof of (2.38)–(2.45), we may set $v \in \left\{-\frac{1}{2}, -\frac{1}{3}, -\frac{1}{4}, -\frac{1}{6}\right\}$ in the last two equations and integrate over the identities in Theorem 2.1, to verify (2.46)–(2.53).

Remark The series evaluation in (2.38) is a special case of [101, Corollary 3.8]. One may also compute (2.39) and (2.40) by extending some arguments in [100, §3].

Corollary 2.5 (Harmonic modulations of Legendre–Sun series). We have

$$-\int_{0}^{1} S_{-1/2}^{L}(t) \log(1-t) dt = 2 \sum_{k=0}^{\infty} \frac{\binom{2k}{k}^{2}}{2^{4k}} \frac{(2H_{2k} - H_{k})H_{k+1}}{k+1}$$

$$= \frac{96}{\pi} - \frac{8\pi}{3} - \frac{128\log 2}{\pi} + \frac{64\log^{2} 2}{\pi},$$
(2.60)

$$-\int_{0}^{1} S_{-1/3}^{L}(t) \log(1-t) dt = \sum_{k=0}^{\infty} \frac{\binom{2k}{k}\binom{3k}{k}}{3^{3k}} \frac{(3H_{3k}-H_k)H_{k+1}}{k+1}$$

$$= \frac{567\sqrt{3}}{8\pi} - \frac{9\sqrt{3}\pi}{4} - \frac{243\sqrt{3}\log 3}{4\pi} + \frac{81\sqrt{3}\log^2 3}{4\pi},$$

$$-\int_{0}^{1} S_{-1/4}^{L}(t) \log(1-t) dt = 2\sum_{k=0}^{\infty} \frac{\binom{2k}{k}\binom{4k}{2k}}{2^{6k}} \frac{(2H_{4k}-H_{2k})H_{k+1}}{k+1}$$

$$= \frac{3328\sqrt{2}}{27\pi} - \frac{40\sqrt{2}\pi}{9} - \frac{512\sqrt{2}\log 2}{3\pi} + \frac{96\sqrt{2}\log^2 2}{\pi},$$

$$-\int_{0}^{1} S_{-1/6}^{L}(t) \log(1-t) dt = \sum_{k=0}^{\infty} \frac{\binom{3k}{k}\binom{6k}{3k}}{2^{4k}3^{3k}} \frac{(6H_{6k}-3H_{3k}-2H_{2k}+H_k)H_{k+1}}{k+1}$$

$$= \frac{40176}{125\pi} - \frac{66\pi}{5} - \frac{5184\log 2}{25\pi} - \frac{3888\log 3}{25\pi} + \frac{288\log^2 2}{5\pi} + \frac{432(\log 2)(\log 3)}{5\pi} + \frac{162\log^2 3}{5\pi}.$$
(2.61)
(2.61)
(2.61)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.62)
(2.

Proof. All these are consequences of

$$\frac{\partial^2}{\partial\sigma^2}\Big|_{\sigma=0} \int_0^1 P_{\nu}(1-2t)(1-t)^{\sigma} \,\mathrm{d}\,t = \frac{\partial^2}{\partial\sigma^2}\Big|_{\sigma=0} \frac{[\Gamma(1+\sigma)]^2}{\Gamma(2+\nu+\sigma)\Gamma(1-\nu+\sigma)} \tag{2.64}$$

and integrations over the formulae in Theorem 2.1(a).

Remark One can of course extend the service above to Theorem 2.1(b) and also consider higherorder derivatives in σ . However, the resulting patterns are less systematic than Corollary 2.4, so we omit the details.

3. Clausen couplings and Ramanujan-Sun series

In response to a question raised by Sun on MathOverflow, Au posted on https://mathoverflow.net/question a sketched proof of [71, Conjecture 29]. In this section, we will give a detailed exposition of Au's answer. To facilitate the statement of Au's results in the theorem below, we recall the Legendre-Ramanujan function R_{ν} from [97, (2.2.11)]:

$$R_{\nu}(\xi) := \frac{1-\xi^{2}}{P_{\nu}(\xi)} \frac{\mathrm{d} P_{\nu}(\xi)}{\mathrm{d} \xi} + \frac{1-\xi^{2}}{P_{\nu}(-\xi)} \frac{\mathrm{d} P_{\nu}(-\xi)}{\mathrm{d} \xi} - \frac{\sin(\nu\pi)}{\pi} \left\{ \frac{1}{[P_{\nu}(\xi)]^{2} \operatorname{Im} \frac{iP_{\nu}(-\xi)}{P_{\nu}(\xi)}} - \frac{1}{[P_{\nu}(-\xi)]^{2} \operatorname{Im} \frac{iP_{\nu}(\xi)}{P_{\nu}(-\xi)}} \right\},$$
(3.1)

which is well-defined for $\xi \in (\mathbb{C} \setminus \mathbb{R}) \cup (-1,1)$, $P_{\nu}(\xi) \neq 0$, and $P_{\nu}(-\xi) \neq 0$. If $\nu \in \{-\frac{1}{2}, -\frac{1}{3}, -\frac{1}{4}, -\frac{1}{6}\}$, then the analytic expression in (3.1) extends continuously to all $\xi \in \mathbb{C}$, being smooth across the branch cut of $P_{\nu}(\xi), \xi \in \mathbb{C} \setminus (-\infty, -1]$:

$$\begin{cases} R_{\nu}(1) = -R_{\nu}(-1) := \lim_{\xi \to 1} R_{\nu}(\xi) = 0; \\ R_{\nu}(\xi) = -R_{\nu}(-\xi) := R_{\nu}(\xi + i0^{+}) \equiv R_{\nu}(\xi - i0^{+}), \quad \xi > 1. \end{cases}$$
(3.2)

Theorem 3.1 (Ramanujan–Sun series). For complex numbers a, b, t satisfying $0 < |4t(1-t)| \le 1$, and a sequence $(c_k)_{k \in \mathbb{Z}_{>0}}$ with at most $O(\log k)$ growth rate, define convergent series

$$\Sigma_{-1/2}\left(a,b;c_k;X_{-1/2}(t)=\frac{t(1-t)}{2^4}\right) \coloneqq \sum_{k=0}^{\infty} \binom{2k}{k}^3 (ak+b)c_k [X_{-1/2}(t)]^k,\tag{3.3}$$

YAJUN ZHOU

$$\Sigma_{-1/3}\left(a,b;c_k;X_{-1/3}(t)=\frac{t(1-t)}{3^3}\right) \coloneqq \sum_{k=0}^{\infty} \binom{2k}{k}^2 \binom{3k}{k} (ak+b)c_k [X_{-1/3}(t)]^k, \tag{3.4}$$

$$\Sigma_{-1/4}\left(a,b;c_k;X_{-1/4}(t) = \frac{t(1-t)}{2^6}\right) \coloneqq \sum_{k=0}^{\infty} \binom{2k}{k}^2 \binom{4k}{2k} (ak+b)c_k [X_{-1/4}(t)]^k, \tag{3.5}$$

$$\Sigma_{-1/6}\left(a,b;c_k;X_{-1/6}(t)=\frac{t(1-t)}{2^43^3}\right) := \sum_{k=0}^{\infty} \binom{2k}{k} \binom{3k}{k} \binom{6k}{3k} (ak+b)c_k [X_{-1/6}(t)]^k.$$
(3.6)

For each series of Ramanujan's type

$$\Sigma_{\nu}(2(2t-1), -R_{\nu}(1-2t); 1; X_{\nu}(t)) = \frac{2\sin(\nu\pi)}{\pi \operatorname{Im} \frac{iP_{\nu}(2t-1)}{P_{\nu}(1-2t)}},$$
(3.7)

there is a corresponding series of Sun's type

$$\Sigma_{\nu}(2(2t-1), -R_{\nu}(1-2t); c_{\nu,k}; X_{\nu}(t)) + \Sigma_{\nu}(0, 2(2t-1); 1; X_{\nu}(t)) = -\frac{2\sin(\nu\pi)\log X_{\nu}(t)}{\pi \operatorname{Im} \frac{iP_{\nu}(2t-1)}{P_{\nu}(1-2t)}} - 2i \frac{\operatorname{Re} \frac{iP_{\nu}(2t-1)}{P_{\nu}(1-2t)}}{\operatorname{Im} \frac{iP_{\nu}(2t-1)}{P_{\nu}(1-2t)}},$$
(3.8)

where

$$c_{\nu,k} := \begin{cases} 6(\mathsf{H}_{2k} - \mathsf{H}_k), & \nu = -\frac{1}{2}, \\ 3\mathsf{H}_{3k} + 2\mathsf{H}_{2k} - 5\mathsf{H}_k, & \nu = -\frac{1}{3}, \\ 4(\mathsf{H}_{4k} - \mathsf{H}_k), & \nu = -\frac{1}{4}, \\ 3(2\mathsf{H}_{6k} - \mathsf{H}_{3k} - \mathsf{H}_k), & \nu = -\frac{1}{6}. \end{cases}$$
(3.9)

Remark If

$$z_{t,4\sin^2(\nu\pi)} \coloneqq \lim_{\varepsilon \to 0^+} \frac{P_{\nu}(2(t+i\varepsilon)-1)}{2i\sin(\nu\pi)P_{\nu}(1-2(t+i\varepsilon))}$$
(2.17')

defines a CM point for a certain $v \in \left\{-\frac{1}{6}, -\frac{1}{4}, -\frac{1}{3}, -\frac{1}{2}\right\}$, such that $\operatorname{Im} z_{t,N} > 0$ and $[\mathbb{Q}(z_{t,N}) : \mathbb{Q}] = 2$ for $N = 4 \sin^2(v\pi) \in \{1, 2, 3, 4\}$, then both t and $R_v(1 - 2t)$ are algebraic numbers, being explicitly solvable in radical forms [97, Proposition 2.2.2]. Such statements about algebraicity follow from Ramanujan's elliptic function theories to alternative bases [12, Chapter 33] and the classical knowledge for complex multiplication [89, §6].

After dealing with the analytic and modular parts of Theorem 3.1 in §§3.1–3.2, we will construct more series identities in §3.3, by differentiating and integrating (precursors to) Theorem 3.1.

3.1. Clausen couplings of Legendre functions. We begin with the study of some infinite series concerning

$$s_{\nu,k} := \binom{2k}{k} \frac{(-\nu)_k (\nu+1)_k}{(k!)^2}$$
(3.10)

as well as

$$\varsigma_{\nu,k} \coloneqq \psi^{(0)}(k-\nu) + \psi^{(0)}(k+\nu+1) - 3\psi^{(0)}(k+1) + \psi^{(0)}\left(k+\frac{1}{2}\right)$$
(3.11)

and

$$\tau_{\nu,k} := (\varsigma_{\nu,k})^2 + \psi^{(1)}(k-\nu) + \psi^{(1)}(k+\nu+1) - 3\psi^{(1)}(k+1) + \psi^{(1)}\left(k+\frac{1}{2}\right), \quad (3.12)$$

where $\psi^{(m)}(z) \coloneqq \frac{d^{m+1}}{dz^{m+1}} \log \Gamma(z)$. In particular, we have expressions involving binomial coefficients

$$s_{\nu,k} = \begin{cases} \frac{\binom{2k}{k}^{2}}{2^{6k}}, & \nu = -\frac{1}{2}, \\ \frac{\binom{2k}{k}^{2}\binom{2k}{k}}{2^{2k}3^{3k}}, & \nu = -\frac{1}{3}, \\ \frac{\binom{2k}{k}^{2}\binom{2k}{2k}}{2^{8k}}, & \nu = -\frac{1}{4}, \\ \frac{\binom{2k}{k}\binom{3k}{k}\binom{6k}{3^{3k}}}{2^{6k}3^{3k}}, & \nu = -\frac{1}{6}, \end{cases}$$
(3.10')

and harmonic numbers

$$\varsigma_{\nu,k} = \begin{cases}
6 (H_{2k} - H_k - \log 2), & \nu = -\frac{1}{2}, \\
3H_{3k} + 2H_{2k} - 5H_k - 2\log 2 - 3\log 3, & \nu = -\frac{1}{3}, \\
4 (H_{4k} - H_k - 2\log 2), & \nu = -\frac{1}{4}, \\
3 (2H_{6k} - H_{3k} - H_k - 2\log 2 - \log 3), & \nu = -\frac{1}{6}, \\
\end{cases}$$

$$\tau_{\nu,k} - (\varsigma_{\nu,k})^2 = \begin{cases}
6 \left[-2H_{2k}^{(2)} + H_k^{(2)} + \frac{\pi^2}{6}\right], & \nu = -\frac{1}{2}, \\
-9H_{3k}^{(2)} - 4H_{2k}^{(2)} + 5H_k^{(2)} + \frac{4\pi^2}{3}, & \nu = -\frac{1}{3}, \\
4 \left[-4H_{4k}^{(2)} + H_k^{(2)} + \frac{\pi^2}{2}\right], & \nu = -\frac{1}{4}, \\
3 \left[-12H_{6k}^{(2)} + 3H_{3k}^{(2)} + H_k^{(2)} + \frac{4\pi^2}{3}\right], & \nu = -\frac{1}{6}.
\end{cases}$$
(3.11')

Lemma 3.2 (Series representation for products of Legendre functions). For $v \in (-1,0)$ and $t \in (0,\frac{1}{2}]$, we have

$$[P_{\nu}(1-2t)]^{2} = \sum_{k=0}^{\infty} [t(1-t)]^{k} s_{\nu,k} =: \sigma_{\nu}(4t(1-t)), \qquad (3.13)$$

$$P_{\nu}(1-2t) \left[\frac{\pi P_{\nu}(2t-1)}{\sin(\nu\pi)} + P_{\nu}(1-2t) \log \frac{1}{4t(1-t)} \right]$$

= $\sum_{k=0}^{\infty} [t(1-t)]^{k} s_{\nu,k} \varsigma_{\nu,k},$ (3.14)

and

$$\left[\frac{\pi P_{\nu}(1-2t)}{\sin(\nu\pi)}\right]^{2} + \left[\frac{\pi P_{\nu}(2t-1)}{\sin(\nu\pi)} + P_{\nu}(1-2t)\log\frac{1}{4t(1-t)}\right]^{2}$$

$$= \sum_{k=0}^{\infty} [t(1-t)]^{k} s_{\nu,k} \tau_{\nu,k}.$$
(3.15)

Proof. Consider the following quadratic transformation [6, (3.1.3)] of hypergeometric functions for $t \in (0, \frac{1}{2}]$:

$${}_{2}F_{1}\begin{pmatrix} -\nu+\varepsilon,\nu+1+\varepsilon\\1+\varepsilon \end{pmatrix} t = {}_{2}F_{1}\begin{pmatrix} \frac{-\nu+\varepsilon}{2},\frac{\nu+1+\varepsilon}{2}\\1+\varepsilon \end{pmatrix} 4t(1-t) \end{pmatrix},$$
(3.16)

where ε is a free parameter for the moment. Combining this with Clausen's identity for squares of ${}_{2}F_{1}$ [6, Exercise 2.13 or 3.17(d)], we have

$$\begin{bmatrix} {}_{2}F_{1} \begin{pmatrix} -\nu + \varepsilon, \nu + 1 + \varepsilon \\ 1 + \varepsilon \end{pmatrix} \end{bmatrix}^{2} = {}_{3}F_{2} \begin{pmatrix} -\nu + \varepsilon, \nu + 1 + \varepsilon, \frac{1}{2} + \varepsilon \\ 1 + \varepsilon, 1 + 2\varepsilon \end{pmatrix} 4t(1-t)$$
(3.17)

for $2\varepsilon \notin \mathbb{Z}_{<0}$. Setting $\varepsilon = 0$ in (3.17), we arrive at (3.13).

Thanks to the Frobenius–Zagier process (Lemma 2.2), we can differentiate (3.17) with respect to ε at $\varepsilon = 0$, which verifies (3.14).

In fact, using Euler's hypergeometric transformation [6, (2.2.7)]

$${}_{2}F_{1}\begin{pmatrix} -\nu+\varepsilon,\nu+1+\varepsilon\\1+\varepsilon \end{pmatrix} t \left[t(1-t)\right]^{\varepsilon/2}$$

$$= {}_{2}F_{1}\begin{pmatrix} -\nu,\nu+1\\1+\varepsilon \end{pmatrix} t \left(\frac{t}{1-t}\right)^{\varepsilon/2} =: \Gamma(1+\varepsilon)P_{\nu}^{-\varepsilon}(1-2t),$$
(3.18)

one may rework the derivations of (3.13)-(3.14) by differentiating the associated Legendre function $P^{\mu}_{\nu}(x), x \in (-1,1)$ with respect to its order μ [23].

Furthermore, for $\varepsilon \notin \mathbb{Z}$, one can deduce

$$\frac{\varepsilon\pi}{\sin(\varepsilon\pi)}P_{\nu}^{-\varepsilon}(1-2t)P_{\nu}^{\varepsilon}(1-2t) = {}_{3}F_{2}\left(\begin{array}{c}-\nu,\nu+1,\frac{1}{2}\\1+\varepsilon,1-\varepsilon\end{array}\right|4t(1-t)\right)$$
(3.19)

from a companion to Clausen's identity [6, Exercise 2.13]. As we reexpress

$$\lim_{\varepsilon \to 0} \frac{\partial^2}{\partial \varepsilon^2} \left\{ \frac{\left[\Gamma(1+\varepsilon) P_{\nu}^{-\varepsilon} (1-2t) \right]^2}{[t(1-t)]^{\varepsilon}} - \frac{\varepsilon \pi}{\sin(\varepsilon \pi)} P_{\nu}^{-\varepsilon} (1-2t) P_{\nu}^{\varepsilon} (1-2t) \right\}$$
(3.20)

through termwise derivatives of the generalized hypergeometric series [cf. (3.17) and (3.19)]

$$\frac{\partial^2}{\partial \varepsilon^2} \left[{}_3F_2 \left(\begin{array}{c} -\nu + \varepsilon, \nu + 1 + \varepsilon, \frac{1}{2} + \varepsilon \\ 1 + \varepsilon, 1 + 2\varepsilon \end{array} \right| 4t(1-t) \right) - {}_3F_2 \left(\begin{array}{c} -\nu, \nu + 1, \frac{1}{2} \\ 1 + \varepsilon, 1 - \varepsilon \end{array} \right| 4t(1-t) \right) \right] \right|_{\varepsilon=0}, \quad (3.21)$$

we get (3.15).

Remark Equivalent forms of (3.13)–(3.15) have appeared in Kam Cheong Au's answer (posted under the username "pisco") to Zhi-Wei Sun's question on https://mathoverflow.net/questions/436205/about [71, Conjecture 29]. Our proof here, being slightly different from Au's argument, follows the presentations of Chudnovsky–Chudnovsky [36, (1.3)] and Chan–Wan–Zudilin [33, §2] more closely.

Remark For $\nu = -\frac{1}{2}$, both (3.13) and (3.14) were derived by Watson [79], who also computed derivatives with respect to a free parameter ε .

Remark Clausen's identity has a natural analog over finite fields [42], which may be helpful in future investigations of Sun's supercongruences in [71, Conjecture 29]. \Box

3.2. Ramanujan–Sun series. Exploiting the Wrońskian determinant

$$W[P_{\nu}(x), P_{\nu}(-x)] := P_{\nu}(x) \frac{\mathrm{d} P_{\nu}(-x)}{\mathrm{d} x} - P_{\nu}(-x) \frac{\mathrm{d} P_{\nu}(x)}{\mathrm{d} x} = -\frac{2\sin(\nu\pi)}{\pi(1-x^2)},$$
(3.22)

one can compute

$$\frac{\pi \operatorname{Im} \frac{iP_{\nu}(-\xi)}{P_{\nu}(\xi)}}{2\sin(\nu\pi)} \left[(1-\xi^2) \frac{\mathrm{d}}{\mathrm{d}\,\xi} - R_{\nu}(\xi) \right] \begin{pmatrix} [P_{\nu}(\xi)]^2 \\ P_{\nu}(\xi)P_{\nu}(-\xi) \\ [P_{\nu}(-\xi)]^2 \end{pmatrix} = \begin{pmatrix} 1 \\ -i\operatorname{Re} \frac{iP_{\nu}(-\xi)}{P_{\nu}(\xi)} \\ -\left| \frac{iP_{\nu}(-\xi)}{P_{\nu}(\xi)} \right|^2 \end{pmatrix}$$
(3.23)

for $v \in (-1,0)$, where the function $R_v(\xi)$ is defined in (3.1) for $\xi \in (\mathbb{C} \setminus \mathbb{R}) \cup (-1,1)$, $P_v(\xi) \neq 0$, and $P_v(-\xi) \neq 0$, with no restrictions on v. As pointed out in Au's aforementioned post, the first (resp. second) entry on the right-hand side of the equation above is responsible for Ramanujan's celebrated series representations of $\frac{1}{\pi}$ [38, Chapter 14] (resp. Sun's infinite series in [71, Conjecture 29]). In particular, with the notations in Theorem 3.1, we have

$$\Sigma_{\nu}(a,b;c_k;X_{\nu}(t)) = \sum_{k=0}^{\infty} [t(1-t)]^k (ak+b)c_k s_{\nu,k}, \qquad (3.24)$$

where $s_{\nu,k}$ is given in (3.10') for $\nu \in \{-\frac{1}{2}, -\frac{1}{3}, -\frac{1}{4}, -\frac{1}{6}\}$, so

$$\Sigma_{\nu}\left(-2\xi, -R_{\nu}(\xi); 1; X_{\nu}\left(\frac{1-\xi}{2}\right)\right) = \left[\left(1-\xi^{2}\right)\frac{\mathrm{d}}{\mathrm{d}\,\xi} - R_{\nu}(\xi)\right] \left[P_{\nu}(\xi)\right]^{2} = \frac{2\sin(\nu\pi)}{\pi\,\mathrm{Im}\,\frac{iP_{\nu}(-\xi)}{P_{\nu}(\xi)}} \tag{3.25}$$

explains (3.7), while

$$\begin{split} & \Sigma_{\nu} \left(-2\xi, -R_{\nu}(\xi); \varsigma_{\nu,k}; X_{\nu} \left(\frac{1-\xi}{2} \right) \right) + \Sigma_{\nu} \left(0, -2\xi; 1; X_{\nu} \left(\frac{1-\xi}{2} \right) \right) \\ &= \left[\left(1-\xi^{2} \right) \frac{\mathrm{d}}{\mathrm{d}\xi} - R_{\nu}(\xi) \right] \left\{ P_{\nu}(\xi) \left[\frac{\pi P_{\nu}(-\xi)}{\sin(\nu\pi)} + P_{\nu}(\xi) \log \frac{1}{1-\xi^{2}} \right] \right\} \\ &- \left[P_{\nu}(\xi) \right]^{2} \left(1-\xi^{2} \right) \frac{\mathrm{d}}{\mathrm{d}\xi} \log \frac{1}{1-\xi^{2}} \\ &= - \frac{2\sin(\nu\pi) \log \left(1-\xi^{2} \right)}{\pi \operatorname{Im} \frac{iP_{\nu}(-\xi)}{P_{\nu}(\xi)}} - 2i \frac{\operatorname{Re} \frac{iP_{\nu}(-\xi)}{P_{\nu}(\xi)}}{\operatorname{Im} \frac{iP_{\nu}(-\xi)}{P_{\nu}(\xi)}} \end{split}$$
(3.26)

accounts for (3.8).

Furthermore, after careful analytic continuation, one may extend the Ramanujan–Sun series in Theorem 3.1 to cases where $t \in (\frac{1-\sqrt{2}}{2}, 0)$. For example, a record-breaking [66] formula of Chudnovsky–Chudnovsky [36, (1.5)]

$$\Sigma_{-1/6} (a_{163}, b_{163}; 1; X_{163}) \coloneqq \sum_{k=0}^{\infty} \binom{2k}{k} \binom{3k}{k} \binom{6k}{3k} (a_{163}k + b_{163}) (X_{163})^k = \frac{2^7 \sqrt{3 \cdot 5^3 \cdot 23^3 \cdot 29^3}}{\pi}$$
(3.27)

for

$$\begin{cases} a_{163} \coloneqq 2 \cdot 3^2 \cdot 7 \cdot 11 \cdot 19 \cdot 127 \cdot 163, \\ b_{163} \coloneqq 13 \cdot 1045493, \\ X_{163} \coloneqq -\frac{1}{2^{18} \cdot 3^3 \cdot 5^3 \cdot 23^3 \cdot 29^3} \end{cases}$$
(3.28)

(resp. its counterpart

$$\Sigma_{-1/6} (a_{163}, b_{163}; 3(2H_{6k} - H_{3k} - H_k); X_{163}) + \Sigma_{-1/6} (0, a_{163}; 1; X_{163})$$

$$\coloneqq \sum_{k=0}^{\infty} \binom{2k}{k} \binom{3k}{k} \binom{6k}{3k} [3 (a_{163}k + b_{163}) (2H_{6k} - H_{3k} - H_k) + a_{163}] (X_{163})^k$$

$$= -\frac{2^7 \sqrt{3 \cdot 5^3 \cdot 23^3 \cdot 29^3}}{\pi} \log |X_{163}|$$
(3.29)

in Sun's context [71, Conjecture 29(iv)]) follows from an application of the first (resp. second) entry on the right-hand side of (3.23) to an analytic continuation of (3.13) [resp. (3.14)] for a particular choice of parameters

$$\nu = -\frac{1}{6}, \quad t = \frac{1 - \xi_{163}}{2} = \frac{1}{2} - \frac{1}{2}\sqrt{\frac{3^3 \cdot 7^2 \cdot 11^2 \cdot 19^2 \cdot 127^2 \cdot 163}{2^{12} \cdot 5^3 \cdot 23^3 \cdot 29^3}}$$
(3.30)

that is associated with a modular invariant

$$X_{163} = \frac{1 - \xi_{163}^2}{1728} = \frac{1}{j\left(\frac{1 + i\sqrt{163}}{2}\right)}.$$
(3.31)

Here, to compute

$$R_{-1/6}(\xi_{163}) = \frac{13 \cdot 1045493}{2^6 \sqrt{3 \cdot 5^3 \cdot 23^3 \cdot 29^3 \cdot 163}}$$
(3.32)

for the specific $\xi_{163} \in (1, \infty)$ given in (3.30), one may set $z = \frac{1+i\sqrt{163}}{2}$ in [97, (2.2.20)]

$$R_{-1/6}\left(\sqrt{\frac{j(z)-1728}{j(z)}}\right)\sqrt{\frac{j(z)-1728}{j(z)}} = -\frac{E_6(z)}{3[E_4(z)]^2}\left[E_2(z)-\frac{E_6(z)}{E_4(z)}\right]$$
(3.33)

for

Im
$$z > 0$$
, $|\operatorname{Re} z| \le \frac{1}{2}$, $|z| \ge 1$, (3.34)

where the Eisenstein series are given by

$$\begin{cases} E_2(z) \coloneqq 1 - \frac{3}{\pi \operatorname{Im} z} - 24 \sum_{n=1}^{\infty} \frac{n e^{2\pi i n z}}{1 - e^{2\pi i n z}}, \\ E_4(z) \coloneqq 1 + 240 \sum_{n=1}^{\infty} \frac{n^3 e^{2\pi i n z}}{1 - e^{2\pi i n z}}, \\ E_6(z) \coloneqq 1 - 504 \sum_{n=1}^{\infty} \frac{n^5 e^{2\pi i n z}}{1 - e^{2\pi i n z}}. \end{cases}$$
(3.35)

Likewise, for $\nu \in \left\{-\frac{1}{4}, -\frac{1}{3}, -\frac{1}{2}\right\}$ and $N = 4\sin^2(\nu\pi) \in \{2, 3, 4\}$, the following relations [97, (2.2.21)]

$$R_{\nu}(1 - 2\alpha_N(z)) = -\frac{N-1}{6} \frac{N^2 [E_2(Nz)]^2 - N^2 E_4(Nz) - [E_2(z)]^2 + E_4(z)}{[N E_2(Nz) - E_2(z)]^2}$$
(3.36)

will be useful in the analytic continuation of the corresponding Ramanujan-Sun series.

3.3. **Some series descending from Clausen couplings**. Before closing this section, we mention some simple consequences of Lemma 3.2 with regard to binomial harmonic sums that lie outside the scope of Theorem 3.1.

3.3.1. Derivatives and differential equations for products of two Legendre functions. One may construct Table 2 by evaluating (2.28) with (3.13)-(3.14). The bulkier identities associated with (3.15) are omitted from this table.

There is a class of infinite series similar to (3.13) that can be evaluated in closed form, despite its absence from Sun's recent conjectures [71]. Consider the following third-order differential equation [95, (16)]

$$\frac{\mathrm{d}}{\mathrm{d}\xi} \left\{ \left(1 - \xi^2\right) \frac{\mathrm{d}}{\mathrm{d}\xi} \left[\left(1 - \xi^2\right) \frac{\mathrm{d}f(\xi)}{\mathrm{d}\xi} \right] + 4\nu(\nu+1) \left(1 - \xi^2\right) f(\xi) \right\} + 4\nu(\nu+1)\xi f(\xi) = 0$$
(3.37)

TABLE 2. Selected series involving $\binom{2k}{k}^3$, where $h_k^{(r)} := \mathsf{H}_{2k}^{(r)} - \frac{1}{2^r} \mathsf{H}_k^{(r)}$, $\lambda := \log 2$, $\zeta(s) := \sum_{k=1}^{\infty} \frac{1}{k^s}$, $\beta(s) := \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)^s}$, and $G := \beta(2)$

$$\begin{split} \sum_{k=0}^{\infty} h_k^{(2)} \frac{\binom{2k}{k}^3}{2^{6k}} &= -\frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^4}{4\pi^3} \left(G - \frac{\pi^2}{8}\right) \\ & \sum_{k=0}^{\infty} \left\{h_k^{(4)} - \left[h_k^{(2)}\right]^2\right\} \frac{\binom{2k}{k}^3}{2^{6k}} &= -\frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^4}{4\pi^3} \left[\beta\left(4\right) + \left(G - \frac{\pi^2}{8}\right)^2 - \frac{\pi^4}{96}\right] \\ & \sum_{k=0}^{\infty} \left\{h_k^{(6)} - \frac{3h_k^{(4)} - \left[h_k^{(2)}\right]^2}{2}h_k^{(2)}\right\} \frac{\binom{2k}{k}^3}{2^{6k}} &= -\frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^4}{4\pi^3} \left\{\beta\left(6\right) + \frac{3}{2}\left[\beta\left(4\right) - \frac{\pi^4}{96}\right] \left(G - \frac{\pi^2}{8}\right) \\ & + \frac{1}{2}\left(G - \frac{\pi^2}{8}\right)^3 - \frac{\pi^6}{960}\right\} \\ & \sum_{k=0}^{\infty} \left[h_k^{(3)} - \frac{3}{2}\left(H_{2k} - H_k - \lambda\right)h_k^{(2)}\right] \frac{\binom{2k}{k}^3}{2^{6k}} &= \frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^4}{32\pi^3}\left[7\zeta\left(3\right) - 2\pi G\right] \\ & \sum_{k=0}^{\infty} \left\{h_k^{(5)} - h_k^{(2)}h_k^{(3)} + 3\left(H_{2k} - H_k - \lambda\right)\frac{\left[h_k^{(2)}\right]^2 - h_k^{(4)}}{4}\right\} \frac{\binom{2k}{k}^3}{2^{6k}} &= \frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^4}{32\pi^3}\left\{\frac{31\zeta\left(5\right)}{4} - \pi\left[\beta\left(4\right) + G^2\right] + 7\zeta\left(3\right)\left(G - \frac{\pi^2}{8}\right)\right\} \end{split}$$

satisfied by $f(\xi) = [P_{\nu}(\xi)]^2$, which allows us to find the anti-derivative of $-\frac{\xi}{2}[P_{\nu}(\xi)]^2$ in the context of (3.13), namely

$$\frac{1-\xi^{2}}{4\nu(\nu+1)}\frac{\mathrm{d}}{\mathrm{d}\xi}\left[\left(1-\xi^{2}\right)P_{\nu}(\xi)\frac{\mathrm{d}P_{\nu}(\xi)}{\mathrm{d}\xi}\right] + \frac{1-\xi^{2}}{2}[P_{\nu}(\xi)]^{2}$$

$$=\frac{\left(1-\xi^{2}\right)^{2}}{4\nu(\nu+1)}\left[\frac{\mathrm{d}P_{\nu}(\xi)}{\mathrm{d}\xi}\right]^{2} + \frac{1-\xi^{2}}{4}[P_{\nu}(\xi)]^{2}$$

$$=\sum_{k=0}^{\infty}\binom{2k}{k}\frac{(-\nu)_{k}(\nu+1)_{k}}{k!(k+1)!}\left(\frac{1-\xi^{2}}{4}\right)^{k+1} = \frac{1-\xi^{2}}{4}{}_{3}F_{2}\left(\frac{1}{2},-\nu,\nu+1\right)\left|1-\xi^{2}\right|$$
(3.38)

for $v \in (-1,0)$, Re $\xi > 0$, and $|1 - \xi^2| < 1$. In the $\xi \to 0$ limit, this brings us (cf. [28, (2.10)] for the case where v = -1/2)

$${}_{3}F_{2}\left(\begin{array}{c}\frac{1}{2},-\nu,\nu+1\\1,2\end{array}\right|1\right) = \sum_{k=0}^{\infty}\frac{\binom{2k}{k}}{2^{2k}}\frac{(-\nu)_{k}(\nu+1)_{k}}{k!(k+1)!} = \frac{\nu(\nu+1)\pi}{4\left[\Gamma\left(\frac{2-\nu}{2}\right)\Gamma\left(\frac{\nu+3}{2}\right)\right]^{2}} + \frac{\pi}{\left[\Gamma\left(\frac{1-\nu}{2}\right)\Gamma\left(\frac{\nu+2}{2}\right)\right]^{2}}$$
(3.39)

for $\nu \in (-1, 0)$.

3.3.2. *Integrals over products of two Legendre functions*. Thanks to the closed-form evaluations (cf. [47, items 7.113.1, 7.113.3, and 7.112.3])

$$\int_{0}^{1} [P_{\nu}(\xi)]^{2} d\xi = \frac{1}{2\nu+1} \left\{ 1 + \frac{\sin(\nu\pi)}{\pi} \left[\psi^{(0)} \left(\frac{\nu+2}{2} \right) - \psi^{(0)} \left(\frac{\nu+1}{2} \right) \right] \right\}, \quad (3.40)$$

$$\int_0^1 P_{\nu}(\xi) P_{\nu}(-\xi) \,\mathrm{d}\,\xi = \frac{\cos(\nu\pi)}{2\nu+1},\tag{3.41}$$

$$\int_{-1}^{1} [P_{\nu}(\xi)]^2 \,\mathrm{d}\,\xi = \frac{2}{2\nu+1} \left[1 - \frac{2\sin^2(\nu\pi)}{\pi^2} \psi^{(1)}(\nu+1) \right],\tag{3.42}$$

it is possible to produce some series identities by manipulating (3.13)-(3.15) in a similar fashion as Corollaries 2.4–2.5, as shown below.

Corollary 3.3 (Integral couplings of two Legendre functions). We have

$$2\int_{0}^{1/2} \sigma_{\nu}(4t(1-t)) dt = \sum_{k=0}^{\infty} \frac{(-\nu)_{k}(\nu+1)_{k}}{(2k+1)(k!)^{2}}$$

$$= \begin{cases} \frac{1}{2\nu+1} \left\{ 1 + \frac{\sin(\nu\pi)}{\pi} \left[\psi^{(0)} \left(\frac{\nu+2}{2} \right) - \psi^{(0)} \left(\frac{\nu+1}{2} \right) \right] \right\}, \quad \nu \neq -\frac{1}{2}, \\ \nu = -\frac{1}{2}, \end{cases}$$

$$2\int_{0}^{1/2} \left[\varsigma_{\nu}(4t(1-t)) - \sigma_{\nu}(4t(1-t)) \log \frac{1}{4t(1-t)} \right] dt$$

$$= \sum_{k=0}^{\infty} \frac{(-\nu)_{k}(\nu+1)_{k}}{(2k+1)(k!)^{2}} \left[2\gamma_{0} - 2H_{k} - \frac{2}{2k+1} + \psi^{(0)}(k-\nu) + \psi^{(0)}(k+\nu+1) \right] \qquad (3.44)$$

$$= \begin{cases} \frac{\pi \cot(\nu\pi)}{2\nu+1}, \quad \nu \neq -\frac{1}{2}, \\ -\frac{\pi^{2}}{2}, \quad \nu = -\frac{1}{2}, \end{cases}$$

$$2\int_{0}^{1/2} \left[\tau_{\nu}(4t(1-t)) - 2\varsigma_{\nu}(4t(1-t)) \log \frac{1}{4t(1-t)} + \sigma_{\nu}(4t(1-t)) \log^{2} \frac{1}{4t(1-t)} \right] dt$$

$$= \sum_{k=0}^{\infty} \frac{(-\nu)_{k}(\nu+1)_{k}}{(2k+1)(k!)^{2}} \left\{ \tau_{\nu,k} - 4 \left(\mathsf{H}_{2k} - \mathsf{H}_{k} + \frac{1}{2k+1} - \log 2 \right) \varsigma_{\nu,k} + 4 \left(\mathsf{H}_{2k} - \mathsf{H}_{k} + \frac{1}{2k+1} - \log 2 \right)^{2} + 4\mathsf{H}_{2k}^{(2)} - 2\mathsf{H}_{k}^{(2)} + \frac{4}{(2k+1)^{2}} - \frac{\pi^{2}}{3} \right\}$$

$$= \begin{cases} \frac{2}{2\nu+1} \left[\frac{\pi^{2}}{\sin^{2}(\nu\pi)} - 2\psi^{(1)}(\nu+1) \right], \quad \nu \neq -\frac{1}{2}, \\ 28\zeta(3), \qquad \nu = -\frac{1}{2}. \end{cases}$$
(3.45)

Here, for v = -1/2, the corresponding infinite series can be evaluated by independent methods [101, Corollary 3.16].

In §5, we will encounter more sophisticated corollaries to Clausen couplings.

4. GREEN-EPSTEIN VARIATIONS ON SUN'S SERIES

The automorphic Green's function (cf. [50, p. 207], [51, pp. 238–239] and [49, p. 544]) of weight $k \in 2\mathbb{Z}_{>1}$ and level $N \in \mathbb{Z}_{>0}$ is defined by

$$G_{k/2}^{\mathfrak{H}/\overline{\Gamma}_{0}(N)}(z_{1}, z_{2}) \coloneqq -\sum_{\substack{a,b,c,d \in \mathbb{Z} \\ ad-Nbc=1}} Q_{\frac{k}{2}-1} \left(1 + \frac{\left| z_{1} - \frac{az_{2}+b}{Ncz_{2}+d} \right|^{2}}{2 \operatorname{Im} z_{1} \operatorname{Im} \frac{az_{2}+b}{Ncz_{2}+d}} \right),$$
(4.1)

with $Q_{\nu}(X) := \int_0^\infty (X + \sqrt{X^2 - 1} \cosh u)^{-\nu - 1} du$ being the Legendre function of the second kind for $X > 1, \nu > -1$.

In this section, we will construct analogs of Sun's series that can be evaluated via automorphic Green's functions of weight 4 and levels $N \in \{1,2,3\}$, corresponding to $Q_1(X) := -1 + \frac{X}{2} \log \frac{X+1}{X-1}, X > 1$ in (4.1). Here, the infinite series in (4.1) evaluates to a finite number if the two points $z_1, z_2 \in \mathfrak{H} := \{w \in \mathbb{C} | \operatorname{Im} w > 0\}$ satisfy $\alpha_N(z_1) \neq \alpha_N(z_2) \in \mathbb{C} \cup \{\infty\}$ when $N \in \{2,3\}$ and $j(z_1) \neq j(z_2)$ when N = 1, with modular invariants defined in (2.23)–(2.24).

Theorem 4.1 (Green–Sun series). For $|t| \le 1$ and $t \notin [-1,0] \cup \{1\}$, we have the following integral representations of infinite series:

$$S_{-1/3}^{G}(t) \coloneqq -\frac{3\sqrt{3}}{\pi} \sum_{k=1}^{\infty} {\binom{2k}{k}} {\binom{3k}{k}} \left(\frac{t}{3^3}\right)^k \left[\psi^{(0)}\left(k+\frac{2}{3}\right) - \psi^{(0)}\left(k+\frac{1}{3}\right) - \frac{\pi}{\sqrt{3}}\right]$$

$$= \int_{-1/3}^{1} \left[P_{-1/3}\left(x\right) - \frac{\pi}{\sqrt{3}}\right] \left[P_{-1/3}\left(x\right) - \frac{\pi}{\sqrt{3}}\right]$$

$$(4.2)$$

$$= \int_{1-2t} [r_{-1/3}(-\zeta)r_{-1/3}(1-2t) - r_{-1/3}(\zeta)r_{-1/3}(2t-1)]r_{-1/3}(\zeta) d\zeta,$$

$$S_{-1/4}^{G}(t) \coloneqq -\frac{2\sqrt{2}}{\pi} \sum_{k=1}^{\infty} {\binom{2k}{k} \binom{4k}{2k} \left(\frac{t}{2^6}\right)^k \left[\psi^{(0)}\left(k+\frac{3}{4}\right) - \psi^{(0)}\left(k+\frac{1}{4}\right) - \pi\right]}$$
(4.3)

$$= \int_{1-2t}^{1} [P_{-1/4}(-\xi)P_{-1/4}(1-2t) - P_{-1/4}(\xi)P_{-1/4}(2t-1)]P_{-1/4}(\xi) d\xi,$$

$$S_{-1/6}^{G}(t) \coloneqq -\frac{3}{2\pi} \sum_{k=1}^{\infty} {\binom{3k}{k}} {\binom{6k}{3k}} \left(\frac{t}{2^4 3^3}\right)^k \left[\psi^{(0)}\left(k+\frac{5}{6}\right) - \psi^{(0)}\left(k+\frac{1}{6}\right) - \sqrt{3\pi}\right]$$

$$= \int_{1-2t}^{1} [P_{-1/6}(-\xi)P_{-1/6}(1-2t) - P_{-1/6}(\xi)P_{-1/6}(2t-1)]P_{-1/6}(\xi) d\xi.$$
(4.4)

These series are related to automorphic Green's functions by the following sum rules for |t| < 1 and |1-t| < 1:

$$\operatorname{Re}\left[\frac{\sin(\nu\pi)}{\operatorname{Im}\frac{iP_{\nu}(1-2t)}{P_{\nu}(2t-1)}}\frac{\mathsf{S}_{\nu}^{G}(t)}{P_{\nu}(2t-1)} + \frac{\sin(\nu\pi)}{\operatorname{Im}\frac{iP_{\nu}(2t-1)}{P_{\nu}(1-2t)}}\frac{\mathsf{S}_{\nu}^{G}(1-t)}{P_{\nu}(1-2t)}\right] - \frac{\sin(2\nu\pi)}{2\nu+1}$$

$$= \frac{1-2\sin^{2}(2\nu\pi)}{4(2\nu+1)^{2}\pi\cos^{2}(\nu\pi)}G_{2}^{\mathfrak{H}/\overline{\Gamma}_{0}(4\sin^{2}(\nu\pi))}\left(\frac{P_{\nu}(2t-1)}{2i\sin(\nu\pi)P_{\nu}(1-2t)}, \frac{1-i\cot(\nu\pi)}{2}\right)$$

$$= \frac{1-2\sin^{2}(2\nu\pi)}{4(2\nu+1)^{2}\pi\cos^{2}(\nu\pi)}G_{2}^{\mathfrak{H}/\overline{\Gamma}_{0}(4\sin^{2}(\nu\pi))}\left(\frac{P_{\nu}(1-2t)}{2i\sin(\nu\pi)P_{\nu}(2t-1)}, \frac{1-i\cot(\nu\pi)}{2}\right),$$

$$(4.5)$$

$$= \frac{1-2\sin^{2}(2\nu\pi)}{4(2\nu+1)^{2}\pi\cos^{2}(\nu\pi)}G_{2}^{\mathfrak{H}/\overline{\Gamma}_{0}(4\sin^{2}(\nu\pi))}\left(\frac{P_{\nu}(1-2t)}{2i\sin(\nu\pi)P_{\nu}(2t-1)}, \frac{1-i\cot(\nu\pi)}{2}\right),$$

$$(4.5)$$

where $N = 4\sin^2(\nu\pi) \in \{1, 2, 3\}$ for $\nu \in \{-\frac{1}{6}, -\frac{1}{4}, -\frac{1}{3}\}.$

Remark As we may recall from the proof of Corollary 2.3, for $N = 4 \sin^2(\nu \pi) \in \{2,3\}$, the statements in the theorem above can be paraphrased by Ramanujan's modular parametrization

$$z = \frac{iP_{\nu}(2\alpha_{N}(z) - 1)}{\sqrt{N}P_{\nu}(1 - 2\alpha_{N}(z))},$$
(4.6)

which is valid for $\alpha_N(z) \in (\mathbb{C} \setminus \mathbb{R}) \cup (0,1)$ and z in the range specified by (2.26); for N = 1, one has

$$z = \lim_{\varepsilon \to 0^+} \frac{iP_{-1/6} \left(-\sqrt{[j(z+i\varepsilon) - 1728]/j(z+i\varepsilon)} \right)}{P_{-1/6} \left(\sqrt{[j(z+i\varepsilon) - 1728]/j(z+i\varepsilon)} \right)}$$
(4.7)

when *z* satisfies (2.27). Furthermore, extending the definition of harmonic numbers $H_n := \sum_{k=1}^n \frac{1}{k}, n \in \mathbb{Z}_{\geq 0}$ to $H_w := \sum_{k=1}^{\infty} \left(\frac{1}{k} - \frac{1}{k+w}\right) = \psi^{(0)}(w+1) + \gamma_0, w \in \mathbb{C} \setminus \mathbb{Z}_{<0}$, one may view (4.2)–(4.4) as generalizations of the binomial harmonic sums in (1.1).

Remark The original formulation of automorphic Green's function by Gross–Zagier [51] involves localization of the Néron–Tate heights over non-archimedean places. For future work, it is perhaps worthwhile to explore some p-adic components of Sun's conjectures [71] using the non-archimedean local heights of Gross–Zagier [51, I.4 and III] and/or Disegni's construction of the p-adic Gross–Zagier formula [41, Theorem B and §5].

YAJUN ZHOU

Define the automorphic Green's function $G_2^{\mathfrak{H}/\overline{\Gamma}_0(4)}(z_1, z_2)$ of weight 4 and level 4 for $z_1, z_2 \in \mathfrak{H}$ satisfying $\alpha_4(z_1) \neq \alpha_4(z_2) \in \mathbb{C}$, by specialization of (4.1) to s = 2, N = 4. Its asymptotic expansion [51, p. 240, (2.19)]

$$G_2^{\mathfrak{H}/\overline{\Gamma}_0(4)}(z_1, z_2) = -\frac{4\pi}{3} \frac{E^{\Gamma_0(4)}(z_2, 2)}{\operatorname{Im} z_1} + O(e^{-(2\pi - 0^+)\operatorname{Im} z_1}), \quad z_1 \to i\infty$$
(4.8)

involves the Epstein zeta function $E^{\Gamma_0(4)}(z,2)$. More generally, one has

$$E^{\Gamma_0(N)}(z,s) = -\lim_{z' \to i\infty} \frac{(2s-1)(\operatorname{Im} z')^{s-1}}{4\pi} G_s^{\mathfrak{H}/\overline{\Gamma}_0(N)}(z,z')$$
(4.9)

for $s \in \mathbb{Z}_{>1}$ and $N \in \{1, 2, 3, 4\}$.

Another goal of the current section is to establish series representations for certain $E^{\Gamma_0(4)}(z,2)$, as stated in the next theorem.

Theorem 4.2 (Epstein–Sun series). For $|t| \le 1$ and $t \notin [-1,0] \cup \{1\}$, we have

$$S_{-1/2}^{E}(t) := \frac{8}{\pi} \sum_{k=1}^{\infty} {\binom{2k}{k}}^{2} \left(\frac{t}{2^{4}}\right)^{k} \left[\mathsf{H}_{2k}^{(2)} - \frac{1}{4}\mathsf{H}_{k}^{(2)}\right] = \frac{16}{\pi^{3}} \int_{0}^{t} \mathbf{K}(\sqrt{s}) \left[\mathbf{K}(\sqrt{1-s})\mathbf{K}(\sqrt{t}) - \mathbf{K}(\sqrt{s})\mathbf{K}(\sqrt{1-t})\right] \mathrm{d}\,s,$$
(4.10)

where $\mathbf{K}(\sqrt{t}) = \frac{\pi}{2}P_{-1/2}(1-2t)$ is the complete elliptic integral of the first kind. There is a sum rule for |t| < 1 and |1-t| < 1:

$$2 - \operatorname{Re}\left[\frac{1}{\operatorname{Im}\frac{i\mathbf{K}(\sqrt{t})}{\mathbf{K}(\sqrt{1-t})}}\frac{\mathsf{S}_{-1/2}^{E}(t)}{\mathbf{K}(\sqrt{1-t})} + \frac{1}{\operatorname{Im}\frac{i\mathbf{K}(\sqrt{1-t})}{\mathbf{K}(\sqrt{t})}}\frac{\mathsf{S}_{-1/2}^{E}(1-t)}{\mathbf{K}(\sqrt{t})}\right]$$

$$= \frac{32}{3}E^{\Gamma_{0}(4)}\left(-\frac{\mathbf{K}(\sqrt{t})}{2\left[i\mathbf{K}(\sqrt{1-t}) - \mathbf{K}(\sqrt{t})\right]}, 2\right),$$
(4.11)

where the Epstein zeta function $E^{\Gamma_0(4)}(z,2)$ is given by (4.8).

The proofs and applications of the last two theorems will occupy §§4.1–4.2.

In §4.3, we will rework the Guillera–Rogers theory [54] via automorphic representations of the Epstein zeta functions $E^{\Gamma_0(N)}(z,2)$ for $N \in \{2,3,4\}$, based on the modular invariants $\alpha_N(z), N \in \{2,3,4\}$ in (2.23), the Eisenstein series in (3.35), as well as the Legendre–Ramanujan functions $R_V(1-2\alpha_N(z))$ in (3.36).

Theorem 4.3 (Guillera–Rogers summation formulae). If $|4\alpha_N(z)[1-\alpha_N(z)]| \ge 1$, $\alpha_N(z) \neq \frac{1}{2}$, and z satisfies the inequality constraints in (2.26), then we have (cf. [54, Proposition 2])

$$\sum_{k=1}^{\infty} \frac{(k!)^{2} [1 - 2\alpha_{N}(z)] \left[2k - \frac{R_{\nu}(1 - 2\alpha_{N}(z))}{1 - 2\alpha_{N}(z)} \right]}{k^{3} {\binom{2k}{k}} (-\nu)_{k} (\nu + 1)_{k} \{\alpha_{N}(z) [1 - \alpha_{N}(z)]\}^{k}}$$

$$= -\frac{8\pi^{2}}{3} \left[E^{\Gamma_{0}(N)}(z, 2) - E^{\Gamma_{0}(N)} \left(-\frac{1}{Nz}, 2 \right) \right]$$

$$+ \frac{4\pi^{2}i}{\operatorname{Im} z} \operatorname{Re} \int_{z}^{\frac{\operatorname{Re} z}{2|\operatorname{Re} z|} + \frac{i}{2} \sqrt{\frac{4 - N}{N}}} \frac{E_{4}(w) - N^{2} E_{4}(Nw)}{N^{2} - 1} (z - w) (\overline{z} - w) \, \mathrm{d} w$$
(4.12)

for $v \in \{-\frac{1}{4}, -\frac{1}{3}, -\frac{1}{2}\}$ and $N = 4\sin^2(v\pi) \in \{2, 3, 4\}$. The same is true (by continuity) when the conditions $|\operatorname{Re} z| < \frac{1}{2}, |z + \frac{1}{N}| > \frac{1}{N}$, and $|z - \frac{1}{N}| > \frac{1}{N}$ in (2.26) are relaxed to $|\operatorname{Re} z| \le \frac{1}{2}, |z + \frac{1}{N}| \ge \frac{1}{N}$, and $|z - \frac{1}{N}| \ge \frac{1}{N}$.

N = NHere, the real part of (4.12) can be reformulated with (see [51, p. 240, (2.16)])

$$E^{\Gamma_0(N)}(z,2) - E^{\Gamma_0(N)}\left(-\frac{1}{Nz},2\right) = \frac{E^{\Gamma_0(1)}(Nz,2) - E^{\Gamma_0(1)}(z,2)}{N^2 - 1}$$
(4.13)

for $N \in \{2,3,4\}$. Meanwhile, the imaginary part of (4.12) is equal to

$$\frac{4\pi^2}{3\operatorname{Im} z} \left(\operatorname{Re} z - \frac{\operatorname{Re} z}{2|\operatorname{Re} z|}\right) \left[\left(\operatorname{Re} z - \frac{\operatorname{Re} z}{2|\operatorname{Re} z|}\right)^2 + 3(\operatorname{Im} z)^2 + \frac{12 - N}{4N} \right] + \frac{4\pi^2 \operatorname{Re} \left[\widetilde{\mathscr{E}}_4(Nz) - N\widetilde{\mathscr{E}}_4(z)\right]}{N(N^2 - 1)\operatorname{Im} z},$$
(4.14)

where the real part of

$$\widetilde{\mathscr{E}}_4(z) := \int_z^{i\infty} [1 - E_4(w)](z - w)(\overline{z} - w) \,\mathrm{d}\,w \tag{4.15}$$

satisfies (cf. [54, Proposition 3])

$$\operatorname{Re}\widetilde{\mathscr{E}}_{4}(z) = \begin{cases} 0, & 2\operatorname{Re} z \in \mathbb{Z}, \\ -\frac{\operatorname{Re} z}{3} \left[\frac{|z|^{2} + 2(\operatorname{Im} z)^{2}}{|z|^{4}} + |z|^{2} + 2(\operatorname{Im} z)^{2} - 5 \right], & 2\operatorname{Re} \frac{1}{z} \in \mathbb{Z}. \end{cases}$$
(4.16)

4.1. Derivatives of Legendre functions with respect to their degrees. For any complex-valued degree $v \in \mathbb{C}$, the Legendre function $P_v(x), -1 < x \leq 1$ is the unique $C^2(-1,1]$ solution to the Legendre differential equation

$$\widehat{L}_{\nu}f(x) := \frac{d}{dx} \left[(1 - x^2) \frac{df(x)}{dx} \right] + \nu(\nu + 1)f(x) = 0$$
(2.16')

equipped with a natural boundary condition f(1) = 1.

The next lemma exploits the differential equation (2.16') and its boundary condition in the construction of an integral representation for $\partial P_{\nu}(x)/\partial \nu$.

Lemma 4.4 (Variation of parameters for $\partial P_{\nu}(x)/\partial \nu$). For $\nu \in (-1,0)$ and $x \in \mathbb{C} \setminus (-\infty, -1]$, we have

$$\frac{\partial P_{\nu}(x)}{\partial \nu} = \frac{(2\nu+1)\pi}{2\sin(\nu\pi)} \int_{x}^{1} [P_{\nu}(-\xi)P_{\nu}(x) - P_{\nu}(\xi)P_{\nu}(-x)]P_{\nu}(\xi) \,\mathrm{d}\,\xi, \tag{4.17}$$

where the path of integration evades the branch cuts of the integrand.

Proof. Differentiating the Legendre differential equation in (2.16') with respect to the degree ν , we get

$$0 = \frac{\partial}{\partial \nu} \left[\widehat{L}_{\nu} P_{\nu}(x) \right] = \widehat{L}_{\nu} \frac{\partial P_{\nu}(x)}{\partial \nu} + (2\nu + 1) P_{\nu}(x).$$
(4.18)

In view of the Wrońskian determinant (3.22), we may use the standard variation of parameters [56, §5.23] for inhomogeneous differential equations to check that

$$\widehat{L}_{\nu}\left\{c_{+}P_{\nu}(x) + c_{-}P_{\nu}(-x) - \frac{\pi}{2\sin(\nu\pi)}\int_{x}^{1} [P_{\nu}(-\xi)P_{\nu}(x) - P_{\nu}(\xi)P_{\nu}(-x)]h(\xi)\,\mathrm{d}\,\xi\right\} = h(x) \quad (4.19)$$

for two constants c_{\pm} , along with a suitably regular function $h(x), x \in (-1, 1)$.

Now that the natural boundary condition $P_{\nu}(x) = 1 + O(x - 1)$ stipulates that

$$\lim_{x \to 1-0^+} \frac{\partial P_{\nu}(x)}{\partial \nu} = 0, \qquad (4.20)$$

while

$$P_{\nu}(-x) = \frac{\sin(\nu\pi)}{\pi} \left[\psi^{(0)}(\nu+1) + \psi^{(0)}(-\nu) + 2\gamma_0 + \log\frac{1-x}{2} \right] + O((1-x)\log(1-x))$$
(4.21)

as $x \to 1 - 0^+$, we must pick $c_+ = c_- = 0$ and $h(\xi) = -(2\nu + 1)P_{\nu}(\xi)$ in the operand of (4.19) for a correct integral representation of $\partial P_{\nu}(x)/\partial \nu$, as declared in (4.17).

For $v \in \left\{-\frac{1}{3}, -\frac{1}{4}, -\frac{1}{6}\right\}, |t| \le 1 \text{ and } t \notin [-1,0] \cup \{1\}, \text{ spelling out}$ $S_{v}^{G}(t) \coloneqq \frac{2\sin(v\pi)}{(2v+1)\pi} \frac{\partial P_{v}(1-2t)}{\partial v}$ $= \frac{2\sin(v\pi)}{(2v+1)\pi} \sum_{k=0}^{\infty} \frac{(-v)_{k}(v+1)_{k}}{(1)_{k}} \left[\pi \cot(v\pi) - \psi^{(0)}(k-v) + \psi^{(0)}(k+v+1)\right] \frac{t^{k}}{k!} \qquad (4.22)$ $= \int_{1-2t}^{1} \left[P_{v}(-\xi)P_{v}(1-2t) - P_{v}(\xi)P_{v}(2t-1)\right]P_{v}(\xi) \,\mathrm{d}\xi,$

we get (4.2)-(4.4) in Theorem 4.1.

From either the symmetry $P_{\nu}(x) = P_{-\nu-1}(x)$ or the $\nu = -1/2$ case of Lemma 4.4, we know that

$$\left. \frac{\partial P_{\nu}(x)}{\partial \nu} \right|_{\nu = -1/2} = 0. \tag{4.23}$$

Differentiating (4.17) again in ν , we get

$$\frac{\partial^2 P_{\nu}(x)}{\partial \nu^2} \bigg|_{\nu=-1/2} = -\pi \int_x^1 [P_{-1/2}(-\xi)P_{-1/2}(x) - P_{-1/2}(\xi)P_{-1/2}(-x)]P_{-1/2}(\xi) \,\mathrm{d}\xi, \qquad (4.24)$$

which entails (4.10) in Theorem 4.2.

4.2. Automorphic Green's functions and Epstein zeta functions of weight 4. The integral forms of Green–Sun series $S_{\nu}^{G}(t)$ are pertinent to the representation of $G_{2}^{\mathfrak{H}/\overline{\Gamma}_{0}(N)}(z,z')$ [defined in (4.1)], where $z' = \frac{1}{2} + \frac{i}{2}\sqrt{\frac{4-N}{N}}$ is a special CM point satisfying j(z') = 0 (for N = 1) or $\alpha_{N}(z') = \infty$ (for $N \in \{2,3\}$).

Proposition 4.5 (Some automorphic Green's functions of weight 4). The following identities are true:

$$G_{2}^{5/\overline{\Gamma}_{0}(1)}\left(z,\frac{1+i\sqrt{3}}{2}\right) = G_{2}^{5/\overline{\Gamma}_{0}(1)}\left(-\frac{1}{z},\frac{1+i\sqrt{3}}{2}\right)$$

$$= -\frac{4\pi}{3\operatorname{Im} z}\operatorname{Re}\int_{-\sqrt{\frac{j(z)-1728}{j(z)}}}^{1} [P_{-1/6}(x)]^{2} dx + \frac{8\pi\operatorname{Re} z}{3\operatorname{Im} z}\operatorname{Re}\int_{\sqrt{\frac{j(z)-1728}{j(z)}}}^{1} iP_{-1/6}(x)P_{-1/6}(-x) dx \qquad (4.25)$$

$$-\frac{4\pi|z|^{2}}{3\operatorname{Im} z}\operatorname{Re}\int_{\sqrt{\frac{j(z)-1728}{j(z)}}}^{1} [P_{-1/6}(x)]^{2} dx, \text{ where } \operatorname{Im} z > 0, |z| \ge 1, |\operatorname{Re} z| < \frac{1}{2},$$

$$\begin{aligned} G_{2}^{55/\overline{\Gamma}_{0}(2)}\left(z,\frac{1+i}{2}\right) &= G_{2}^{55/\overline{\Gamma}_{0}(2)}\left(-\frac{1}{2z},\frac{1+i}{2}\right) \\ &= -\frac{\pi}{4\operatorname{Im} z}\operatorname{Re}\int_{2\alpha_{2}(z)-1}^{1}\left[P_{-1/4}(x)\right]^{2}\mathrm{d}\,x + \frac{\pi\operatorname{Re}\,z}{\sqrt{2}\operatorname{Im}\,z}\operatorname{Re}\int_{1-\alpha_{2}(z)}^{1}iP_{-1/4}(x)P_{-1/4}(-x)\,\mathrm{d}\,x \end{aligned} \tag{4.26} \\ &- \frac{\pi|z|^{2}}{2\operatorname{Im}\,z}\operatorname{Re}\int_{1-\alpha_{2}(z)}^{1}\left[P_{-1/4}(x)\right]^{2}\mathrm{d}\,x, \text{ where }\operatorname{Im}\,z > 0, |\operatorname{Re}\,z| < \frac{1}{2}, \left|z + \frac{1}{2}\right| > \frac{1}{2}, \left|z - \frac{1}{2}\right| > \frac{1}{2}, \\ &G_{2}^{55/\overline{\Gamma}_{0}(3)}\left(z,\frac{3+i\sqrt{3}}{6}\right) = G_{2}^{55/\overline{\Gamma}_{0}(3)}\left(-\frac{1}{3z},\frac{3+i\sqrt{3}}{6}\right) \\ &= -\frac{\pi}{9\operatorname{Im}\,z}\operatorname{Re}\int_{2\alpha_{3}(z)-1}^{1}\left[P_{-1/3}(x)\right]^{2}\mathrm{d}\,x + \frac{\pi\operatorname{Re}\,z}{3\sqrt{3}\operatorname{Im}\,z}\operatorname{Re}\int_{1-\alpha_{3}(z)}^{1}iP_{-1/3}(x)P_{-1/3}(-x)\,\mathrm{d}\,x \end{aligned} \tag{4.27} \\ &- \frac{\pi|z|^{2}}{3\operatorname{Im}\,z}\operatorname{Re}\int_{1-\alpha_{3}(z)}^{1}\left[P_{-1/3}(x)\right]^{2}\mathrm{d}\,x, \text{ where }\operatorname{Im}\,z > 0, |\operatorname{Re}\,z| < \frac{1}{3}, \left|z + \frac{1}{3}\right| > \frac{1}{3}, \left|z - \frac{1}{3}\right| > \frac{1}{3}. \end{aligned}$$

Proof. These are specializations of [97, (2.2.14) and (2.2.15)].

One can also verify (4.25)–(4.27) from first principles, as follows. Check that the difference between the series representation [cf. (4.1)] of $G_2^{\mathfrak{H}/\overline{\Gamma}_0(N)}\left(z, \frac{1}{2} + \frac{i}{2}\sqrt{\frac{4-N}{N}}\right)$ and its claimed integral representation extends to a bounded and smooth function on $Y_0(N) := \Gamma_0(N) \setminus \mathfrak{H}$ that is annihilated by the differential operator

$$\Delta_z^{\mathfrak{H}} - 2 \equiv (\operatorname{Im} z)^2 \left[\frac{\partial^2}{(\partial \operatorname{Re} z)^2} + \frac{\partial^2}{(\partial \operatorname{Im} z)^2} \right] - 2, \tag{4.28}$$

and tends to zero as z approaches any cusp in $\Gamma_0(N)\setminus(\mathbb{Q}\cup\{i\infty\})$. Since the Laplacian $\Delta_z^{\mathfrak{H}}$ in this context does not possess a positive eigenvalue [57, §4.1], the aforementioned difference must be identically vanishing on $\Gamma_0(N)\setminus\mathfrak{H}$.

At this point, it is clear that Theorem 4.1 issues directly from Lemma 4.4 and Proposition 4.5.

According to some particular cases of the conjectures by Gross–Zagier [51, p. 317] and Gross–Kohnen–Zagier [49, p. 556], we have

$$\exp\left[\operatorname{Im} z \operatorname{Im} z' G_2^{\mathfrak{H}/\overline{\Gamma}_0(N)}(z,z')\right] \in \overline{\mathbb{Q}} \quad \text{if } [\mathbb{Q}(z):\mathbb{Q}] = [\mathbb{Q}(z'):\mathbb{Q}] = 2 \text{ and } N \in \{1,2,3,4\}, \quad (4.29)$$

with the understanding that $\exp(-\infty) = 0 \in \overline{\mathbb{Q}}$ whenever the automorphic Green's function $G_2^{\mathfrak{H}/\overline{\Gamma}_0(N)}(z,z')$ diverges to $-\infty$ for either j(z) = j(z'), N = 1 or $\alpha_N(z) = \alpha_N(z'), N \in \{2,3,4\}$. This long-standing algebraicity conjecture (in its full generality) has been recently settled by Bruinier-Li-Yang [22]—a remarkable feat that builds upon the milestones of Zhang [94], Mellit [65], Viazovska [74, 75, 76, 77], Bruinier-Ehlen-Yang [21], and Li [62, 63].

Therefore, we have an advanced analog of Corollary 2.3, as given below.

Corollary 4.6 (Some arithmetic properties of Green–Sun series). For each CM point $\frac{P_{\nu}(2t-1)}{2i\sin(\nu\pi)P_{\nu}(1-2t)}$ satisfying |t| < 1, |1 - t| < 1, and $\nu \in \{-\frac{1}{6}, -\frac{1}{4}, -\frac{1}{3}\}$, we have an explicitly computable algebraic number

$$\exp\left\{\frac{\pi}{\tan(\nu\pi)}\operatorname{Re}\left[\left|\frac{P_{\nu}(2t-1)}{P_{\nu}(1-2t)}\right|^{2}\frac{\mathsf{S}_{\nu}^{G}(t)}{P_{\nu}(2t-1)}+\frac{\mathsf{S}_{\nu}^{G}(1-t)}{P_{\nu}(1-2t)}\right]-\frac{2\pi\cos^{2}(\nu\pi)}{2\nu+1}\frac{\operatorname{Im}\frac{iP_{\nu}(2t-1)}{P_{\nu}(1-2t)}}{\sin(\nu\pi)}\right\}\in\overline{\mathbb{Q}},\ (4.30)$$

whose factorization follows from [22, Theorem 1.2].

Remark Table 3 supplies some concrete examples for Theorem 4.1 and Corollary 4.6. Here, to keep the table within manageable size, we have picked a small fraction of the modular invariants listed in [38, Tables 14.3-14.5].

z	$\frac{1}{j(z)}$	$\frac{\sqrt{3}\operatorname{Im} z}{2} G_2^{\mathfrak{H}/\overline{\Gamma}_0(1)}\left(z, \frac{1+i\sqrt{3}}{2}\right)$
$i\sqrt{2}$	$\frac{1}{2^6 \cdot 5^3}$	$3\log\frac{\sqrt{6}-1}{\sqrt{6}+1} - 6\log(\sqrt{2}+\sqrt{3})$
		$= \frac{8\pi^2\sqrt{\pi}}{\sqrt{3}\Gamma(\frac{1}{8})\Gamma(\frac{3}{8})} \sqrt[4]{\frac{2}{5}} \left[\sqrt{2}S^G_{-1/6}\left(\frac{1}{2} - \frac{7}{10}\sqrt{\frac{2}{5}}\right) + S^G_{-1/6}\left(\frac{1}{2} + \frac{7}{10}\sqrt{\frac{2}{5}}\right)\right] - 3\sqrt{2}\pi$
$i\sqrt{3}$	$\frac{1}{2^4\cdot 3^3\cdot 5^3}$	$-6\log 5$
		$= \frac{8\pi^3}{3\left[\Gamma\left(\frac{1}{3}\right)\right]^3} \frac{\sqrt{2}}{\sqrt[4]{5}} \left[\sqrt{3}S^G_{-1/6}\left(\frac{1}{2} - \frac{11}{10\sqrt{5}}\right) + S^G_{-1/6}\left(\frac{1}{2} + \frac{11}{10\sqrt{5}}\right)\right] - 3\sqrt{3}\pi$
	()[]	
z	$\frac{\alpha_2(z)[1-\alpha_2(z)]}{2^6}$	$\frac{\operatorname{Im} z}{2} G_2^{\mathfrak{H}/\overline{\Gamma}_0(2)} \left(z, \frac{1+i}{2} \right)$
i	$\frac{1}{2^3 \cdot 3^4}$	$-\log 3$
_		$= \frac{\pi^{-}\sqrt{\pi}}{2\sqrt{3}\left[\Gamma\left(\frac{1}{4}\right)\right]^{2}} \left[\sqrt{2}S_{-1/4}^{G}\left(\frac{1}{9}\right) + S_{-1/4}^{G}\left(\frac{\mathfrak{o}}{9}\right)\right] - \frac{\pi}{2}$
$i\sqrt{\frac{3}{2}}$	$\frac{1}{2^8 \cdot 3^2}$	$-\log\left(\sqrt{2}+\sqrt{3}\right)$
		$=\frac{\pi^2\sqrt{\pi}}{2\Gamma(\frac{1}{24})\Gamma(\frac{11}{24})}\sqrt{\frac{3}{\sqrt{3}-1}}\left[\sqrt{3}S^G_{-1/4}\left(\frac{1}{2}-\frac{\sqrt{2}}{3}\right)+S^G_{-1/4}\left(\frac{1}{2}+\frac{\sqrt{2}}{3}\right)\right]-\frac{3\pi}{2\sqrt{6}}$
z	$\frac{\alpha_3(z)[1-\alpha_3(z)]}{3^3}$	$\frac{\sqrt{3}\operatorname{Im} z}{6}G_2^{\mathfrak{H}/\overline{\Gamma}_0(3)}\left(z,\frac{3+i\sqrt{3}}{6}\right)$
$i\sqrt{\frac{2}{3}}$	$\frac{1}{2^3 \cdot 3^3}$	$-\frac{\log\left(\sqrt{2}+1\right)}{3}$
		$ = \frac{2\sqrt{2}\pi^2\sqrt{\pi}}{9\sqrt{3}\Gamma(\frac{1}{24})\Gamma(\frac{11}{24})}\sqrt{\frac{\sqrt{2}}{\sqrt{3}-1}} \left[\sqrt{2}S^G_{-1/3}\left(\frac{1}{2}-\frac{1}{2\sqrt{2}}\right) + S^G_{-1/3}\left(\frac{1}{2}+\frac{1}{2\sqrt{2}}\right)\right] - \frac{\pi}{3\sqrt{6}} $
$\frac{2i}{\sqrt{3}}$	$\frac{1}{2\cdot 3^6}$	$\frac{\log 2}{3} - \frac{\log 3}{2}$
,		$=\frac{4\sqrt[3]{2}\pi^{3}}{81\left[\Gamma\left(\frac{1}{3}\right)\right]^{3}}\left[2S_{-1/3}^{G}\left(\frac{1}{2}-\frac{5}{6\sqrt{3}}\right)+S_{-1/3}^{G}\left(\frac{1}{2}+\frac{5}{6\sqrt{3}}\right)\right]-\frac{\pi}{3\sqrt{3}}$

TABLE 3. Selected representations of $G_2^{\mathfrak{H}/\overline{\Gamma}_0(N)}\left(z, \frac{1}{2} + \frac{i}{2}\sqrt{\frac{4-N}{N}}\right)$ for $N \in \{1, 2, 3\}$

For all $z \in \mathfrak{H}$, the modular invariant $\alpha_4(z)$ evaluates to a complex number with finite magnitude. Therefore, a natural N = 4 analog of Proposition 4.5 will invoke asymptotic behavior of automorphic Green's functions [cf. (4.8)], which in turn, ushers in the Epstein zeta function.

Proposition 4.7 (Integral and series representations of Epstein zeta functions). If $z \in \mathfrak{H}$ satisfies the following inequalities:

$$\left| \operatorname{Re}\left(z + \frac{1}{2}\right) \right| < \frac{1}{2}, \quad \left| z + \frac{1}{4} \right| > \frac{1}{4}, \quad \left| z + \frac{3}{4} \right| > \frac{1}{4}, \tag{4.31}$$

TABLE 4. Selected representations of $E^{\Gamma_0(4)}(-\frac{1}{4z},2)$, where $\text{Li}_2(w) := \sum_{k=1}^{\infty} \frac{w^k}{k^2}$ and $G = \text{Im Li}_2(i)$

$z + \frac{1}{2}$	$\frac{\alpha_4 \left(z+\frac{1}{2}\right) \left[1-\alpha_4 \left(z+\frac{1}{2}\right)\right]}{2^4}$	$E^{\Gamma_0(4)}\left(-\frac{1}{4z},2\right)$
$\frac{i}{2}$	$\frac{1}{2^6}$	$= \frac{\frac{3G}{2\pi^2}}{\frac{3}{16}} - \frac{3\sqrt{\pi}}{4[\Gamma(\frac{1}{7})]^2} S_{-1/2}^E\left(\frac{1}{2}\right)$
$\frac{i\sqrt{2}}{2}$	$\frac{\left(\sqrt{2}-1\right)^3}{2^3}$	$= \frac{3}{2\sqrt{2}\pi^2} \left[4 \operatorname{Im} \operatorname{Li}_2(e^{\pi i/4}) - 3G \right]$ = $\frac{3}{16} - \frac{3\sqrt{\pi}}{\sqrt{2\pi^2}} \left[\sqrt{2} \operatorname{S}_{-1/2}^E((\sqrt{2}-1)^2) + \operatorname{S}_{-1/2}^E(2(\sqrt{2}-1)) \right]$
$\frac{i\sqrt{3}}{2}$	$\frac{1}{2^8}$	$= \frac{3}{16} - \frac{\sqrt[3]{2}\sqrt[3]{3}\pi}{\sqrt[3]{3}\pi^2} \left[\sqrt[3]{3}S_{-1/2}^E \left(\frac{2 - \sqrt{3}}{2} \right) + S_{-1/2}^E \left(\frac{2 + \sqrt{3}}{2} \right) \right]$
i	$\frac{\left(\sqrt{2}-1\right)^6}{2\sqrt{2}}$	$ = \frac{3}{16} - \frac{3\sqrt{\pi}}{56} \left[2S_{-1/2}^{E} \left(\sqrt{2} - 1 \right)^{4} \right] + S_{-1/2}^{E} \left(2\sqrt{2} \sqrt{2} - 1 \right)^{2} \right] $
$\frac{i\sqrt{5}}{2}$	$\frac{1}{2^6} \left(\frac{\sqrt{5}-1}{2}\right)^6$	$\frac{16}{4(2+\sqrt{2})} \left[\Gamma(\frac{1}{4}) \right]^2 \left[\Gamma(\frac{1}{4}) \left[\Gamma(\frac{1}{4}) \right]^2 \left[\Gamma(\frac{1}{4}) \right]^2 \left[\Gamma(\frac{1}{4}) \right]^2 \left[\Gamma(\frac{1}{4}) \right]^2 \left[\Gamma(\frac{1}{4}) \left[\Gamma(\frac{1}{4}) \right]^2 \left[\Gamma(\frac{1}{4}) \right]^2 \left[\Gamma(\frac{1}{4}) \left[\Gamma(\frac{1}{4}) \left[\Gamma(\frac{1}{4}) \right]^2 \left$
		$ = \frac{3}{16} - \frac{3\sqrt[8]{5}\Gamma(\frac{1}{10})}{4\sqrt[5]{2^3}[\Gamma(\frac{1}{20})]^2} \frac{\sqrt[4]{\sqrt{5}-2}}{\sqrt{1+\sqrt{5}-\sqrt{5}+2\sqrt{5}}} \left[\sqrt{5}S^E_{-1/2}\left(\frac{\left(1-\sqrt{\sqrt{5}-2}\right)}{2^3\left(\frac{\sqrt{5}-1}{2}\right)}\right) + S^E_{-1/2}\left(\frac{2\left(\frac{\sqrt{5}-1}{2}\right)^7}{\left(1-\sqrt{\sqrt{5}-2}\right)^4}\right)\right] $
$\frac{i\sqrt{6}}{2}$	$\frac{(2-\sqrt{3})^3(\sqrt{3}-\sqrt{2})^3}{2^3(\sqrt{2}-1)^2}$	$\frac{3}{4\pi^{2}\sqrt{6}} \left[7 \operatorname{Im}\operatorname{Li}_{2} \left(e^{\pi i/3} \right) + 36 \operatorname{Im}\operatorname{Li}_{2} \left(e^{\pi i/12} \right) - 12 \operatorname{Li}_{2} \left(e^{5\pi i/12} \right) - 16G \right]$ = $\frac{3}{16} - \frac{3\sqrt{\pi}\sqrt[4]{3}\sqrt[4]{2} - \sqrt{3}}{4\Gamma(\frac{1}{24})\Gamma(\frac{11}{24})} \frac{\sqrt{\sqrt{3} - \sqrt{2}}}{\sqrt{\sqrt{2} - 1}} \left[\sqrt{6} S^{E}_{-1/2} \left((\sqrt{3} - \sqrt{2})^{2} (2 - \sqrt{3})^{2} \right) + S^{E}_{-1/2} \left(\frac{2(\sqrt{3} - \sqrt{2})(2 - \sqrt{3})}{\sqrt{\sqrt{2} - 1}} \right) \right]$
$\frac{i\sqrt{7}}{2}$	$\frac{1}{2^{12}}$	$= \frac{3}{16} - \frac{3\pi}{8\sqrt[4]{7}\Gamma(\frac{1}{7})\Gamma(\frac{4}{7})} \left[\sqrt{7}S_{-1/2}^{E} \left(\frac{8-3\sqrt{7}}{2^{4}} \right) + S_{-1/2}^{E} \left(\frac{8+3\sqrt{7}}{2^{4}} \right) \right]$

then we have

$$\begin{split} & E^{\Gamma_{0}(4)}\left(-\frac{1}{4z},2\right) \\ &= \frac{3}{4\pi^{3}\operatorname{Im} z}\operatorname{Re}\int_{0}^{1-\alpha_{4}\left(z+\frac{1}{2}\right)}\left[\mathbf{K}(\sqrt{t})\right]^{2}\mathrm{d}\,t - \frac{3\operatorname{Re}\left(z+\frac{1}{2}\right)}{\pi^{3}\operatorname{Im} z}\operatorname{Re}\int_{0}^{\alpha_{4}\left(z+\frac{1}{2}\right)}i\mathbf{K}(\sqrt{t})\mathbf{K}(\sqrt{1-t})\,\mathrm{d}\,t \quad (4.32) \\ &\quad + \frac{3\left|z+\frac{1}{2}\right|^{2}}{\pi^{3}\operatorname{Im} z}\operatorname{Re}\int_{0}^{\alpha_{4}\left(z+\frac{1}{2}\right)}\left[\mathbf{K}(\sqrt{t})\right]^{2}\mathrm{d}\,t, \end{split}$$

hence (4.11) in Theorem 4.2.

Proof. The integral representation in (4.32) is a reformulation of [98, (1.14)]. Its proof draws on the spectral analysis of the Laplacian $\Delta_z^{\mathfrak{H}}$ on $Y_0(4) := \Gamma_0(4) \setminus \mathfrak{H}$, akin to the situation in Proposition 4.5.

Remark We note that (see [51, p. 240, (2.16)] or [98, (3.4)])

$$E^{\Gamma_0(4)}(z,2) = \frac{1}{15} \left[E^{\Gamma_0(1)}(4z,2) - \frac{E^{\Gamma_0(1)}(2z,2)}{4} \right],$$
(4.33)

where solvable CM values of the Epstein zeta function [50, p. 207]

$$E^{\Gamma_0(1)}(z,2) = \frac{45}{\pi^4} \sum_{\substack{m,n \in \mathbb{Z} \\ m^2 + n^2 \neq 0}} \frac{(\operatorname{Im} z)^2}{|mz + n|^4}$$
(4.34)

follow from standard procedures [14, 44, 82, 85, 86, 88, 102]. This explains the entries in Table 4 that involve special values of the dilogarithm function $\text{Li}_2(w)$.

4.3. Integral representations of certain $_4F_3$ series. We give a new proof for the main identities of Guillera–Rogers [54, Propositions 2 and 3].

Proposition 4.8 (Ramanujan series upside-down). For $v \in (-1,0)$, $|4t(t-1)| \ge 1$, and $t \ne \frac{1}{2}$, the following formula holds true:

$$\sum_{k=1}^{\infty} \frac{(k!)^2}{k^3 \binom{2k}{k} (-\nu)_k (\nu+1)_k [t(1-t)]^k}$$

$$= \frac{\pi^2}{2\sin^2(\nu\pi)} \int_0^1 [P_\nu (1-2s)]^2 \left(\frac{1}{s-t} + \frac{1}{s-1+t}\right) \mathrm{d} s,$$
(4.35)

where the path of integration runs along the unit interval on the real axis. Particular cases of (4.35) allow us to recover the real part of (4.12).

For $v \in \left\{-\frac{1}{4}, -\frac{1}{3}, -\frac{1}{2}\right\}$ and $N = 4\sin^2(\nu\pi) \in \{2, 3, 4\}$, we have a modular parametrization

$$\frac{\pi^2}{2\sin^2(\nu\pi)} \int_0^1 \frac{[P_{\nu}(1-2s)]^2}{[P_{\nu}(1-2\alpha_N(z))]^2} \left[\frac{1}{s-\alpha_N(z)} + \frac{1}{s-1+\alpha_N(z)}\right] \mathrm{d}s$$

$$= 4\pi^3 i \int_z^{\frac{\mathrm{Re}\,z}{2|\mathrm{Re}\,z|} + \frac{i}{2}\sqrt{\frac{4-N}{N}}} \frac{N^2 E_4(Nw) - E_4(w)}{N^2 - 1} (z-w)^2 \,\mathrm{d}w$$
(4.35*)

for z satisfying (2.26) and $\text{Re } z \neq 0$, hence the imaginary part of (4.12).

Proof. Consider the homogeneous Appell differential equation [7]

$$\widehat{\mathsf{A}}_{\nu,t}f(t) \coloneqq \frac{\partial}{\partial t} \left\{ \left[t(1-t)\frac{\partial}{\partial t} \right]^2 + 4\nu(\nu+1)t(1-t) \right\} f(t) - 2\nu(\nu+1)(1-2t)f(t) = 0$$
(4.36)

whose solution space is spanned by $[P_{\nu}(2t-1)]^2$, $P_{\nu}(2t-1)P_{\nu}(1-2t)$, and $[P_{\nu}(1-2t)]^2$. The left-hand side of (4.35), being equal to [cf. (1.11)]

$$\mathscr{F}_{\nu}(t) \coloneqq \frac{{}_{4}F_{3}\left(\begin{array}{c} 1,1,1,1\\ \frac{3}{2},1-\nu,\nu+2 \\ \end{array} \middle| \frac{1}{4t(1-t)} \right)}{-2t(1-t)\nu(\nu+1)}, \tag{4.37}$$

is annihilated by

$$\begin{split} \widehat{\mathsf{B}}_{\nu,t} &\coloneqq t^{3}(1-t)^{3}(1-2t)\frac{\partial^{4}}{\partial t^{4}} + 2t^{2}(1-t)^{2}[3-11t(1-t)]\frac{\partial^{3}}{\partial t^{3}} \\ &+ t(1-t)(1-2t)\{7+2[2\nu(\nu+1)-15]t(1-t)\}\frac{\partial^{2}}{\partial t^{2}} \\ &+ (1+2t(1-t)\{5\nu(\nu+1)-7-2[8\nu(\nu+1)-9]t(1-t)\})\frac{\partial}{\partial t} + 2\nu(\nu+1)(1-2t)^{3} \\ &= \left\{t(1-t)(1-2t)\frac{\partial}{\partial t} + [1-2t(1-t)]\right\}\widehat{\mathsf{A}}_{\nu,t}. \end{split}$$
(4.38)

In other words, there is a constant C such that

$$\widehat{\mathsf{A}}_{\nu,t}\mathscr{F}_{\nu}(t) = C \frac{1-2t}{t(1-t)}.$$
(4.39)

One can find C = 1 by comparing the asymptotic expansions of both sides in the $t \to \infty$ regime.

From (4.36), one can easily check that the operator $\widehat{A}_{\nu,t} = -\widehat{A}_{\nu,t}^*$ is anti-symmetric. Direct computations also show that $\widehat{A}_{\nu,t}\frac{1}{s-t} = -\widehat{A}_{\nu,s}\frac{1}{s-t}$ and $\widehat{A}_{\nu,t}\frac{1}{s-1+t} = \widehat{A}_{\nu,s}\frac{1}{s-1+t}$. Integrating by parts with respect to *s*, and exploiting the asymptotic behavior of $P_{\nu}(1-2s), s \to 1^-$ [cf. (4.21)], we can verify that

$$\widehat{\mathsf{A}}_{\nu,t} \int_0^1 [P_{\nu}(1-2s)]^2 \left(\frac{1}{s-t} + \frac{1}{s-1+t}\right) \mathrm{d}\,s = \frac{2\sin^2(\nu\pi)}{\pi^2} \frac{1-2t}{t(1-t)}.\tag{4.40}$$

Thus, for |4t(1-t)| > 1 and Im $t \neq 0$, the two sides of (4.35) differ by the restriction of a function²

$$\varphi_{\nu}(t) = \begin{cases} c_1^+ [P_{\nu}(2t-1)]^2 + c_2^+ P_{\nu}(1-2t)P_{\nu}(2t-1) + c_3^+ [P_{\nu}(1-2t)]^2, & \operatorname{Im} t > 0, \\ c_1^- [P_{\nu}(2t-1)]^2 + c_2^- P_{\nu}(1-2t)P_{\nu}(2t-1) + c_3^- [P_{\nu}(1-2t)]^2, & \operatorname{Im} t < 0, \end{cases}$$
(4.41)

where $c_j^{\pm}, j \in \{1, 2, 3\}$ are constants. Since both sides of (4.35) are even functions of 1 - 2t that remain holomorphic for |4t(1-t)| > 1, the function $\varphi_{\nu}(t)$ must be identically vanishing, which establishes (4.35) in full.

To verify the real part of (4.12) for z subject to the constraints in (2.26), simply note that the identities

$$\begin{cases} E^{\Gamma_0(N)}(z,2) = +\frac{3}{16\sin^2(\nu\pi)} \operatorname{Re}\left\{ \left[2t(1-t)\frac{\mathrm{d}}{\mathrm{d}t} + R_{\nu}(1-2t) \right] \int_0^1 \frac{\left[P_{\nu}(1-2s)\right]^2}{s-1+t} \,\mathrm{d}s \Big|_{t=\alpha_N(z)} \right\} \\ E^{\Gamma_0(N)}\left(-\frac{1}{Nz},2\right) = -\frac{3}{16\sin^2(\nu\pi)} \operatorname{Re}\left\{ \left[2t(1-t)\frac{\mathrm{d}}{\mathrm{d}t} + R_{\nu}(1-2t) \right] \int_0^1 \frac{\left[P_{\nu}(1-2s)\right]^2}{s-t} \,\mathrm{d}s \Big|_{t=\alpha_N(z)} \right\} \end{cases}$$
(4.42)

follow from the integral representations of automorphic Green's functions $G_2^{\mathfrak{H}/I_0(N)}(z,z')$ (cf. [97, (2.2.13) and (2.2.15)]) along with their asymptotic behavior in the $z' \to i \infty$ limit [cf. (4.9)].

For $v \in (-1,0)$ and $t \notin \mathbb{R}$, one can prove

$$0 = \int_{0}^{1} [P_{\nu}(1-2s)]^{2} \left(\frac{1}{s-t} + \frac{1}{s-1+t}\right) ds$$

$$-\int_{t}^{\infty} [P_{\nu}(1-2s)P_{\nu}(2t-1) - P_{\nu}(2s-1)P_{\nu}(1-2t)]^{2} \frac{2s-1}{s(1-s)} ds$$
(4.43)

²Recall that $P_{\nu}(1-2t), t \in \mathbb{C} \setminus [1,\infty)$ is holomorphic and $P_{\nu}(1-2t+i0^{+}) - P_{\nu}(1-2t-i0^{+}) = 2i \sin(\nu\pi)P_{\nu}(2t-1)$ for t > 1. Therefore, if the function $\varphi_{\nu}(t)$ in (4.41) satisfies $\varphi_{\nu}(t+i0^{+}) = \varphi_{\nu}(t-i0^{+})$ for $t \in \mathbb{R} \setminus [0,1]$, then $\varphi_{\nu}(t)$ has to be a constant multiple of $[P_{\nu}(2t-1)]^{2} - [P_{\nu}(1-2t)]^{2} - \frac{2i \operatorname{Im} t}{|\operatorname{Im} t|} \sin(\nu\pi)P_{\nu}(1-2t)P_{\nu}(2t-1)$ for $\operatorname{Im} t \neq 0$. If one further knows that $\varphi_{\nu}(t) = \varphi_{\nu}(1-t)$ is an even function of 1-2t, then $\varphi_{\nu}(t)$ must vanish identically.

(where the path of integration in the second integral does not intersect the real axis) by checking that the right-hand side enjoys three properties: (1) It is an even function of 1-2t; (2) It is annihilated by \widehat{A}_{ν} ; (3) It lacks jump discontinuities³ for $t \in \mathbb{R} \setminus [0,1]$. To relate (4.43) to (4.35^{*}), simply differentiate Ramanujan's modular parametrization in (4.6) with the help from the Wrońskian determinant in (3.22):

$$\frac{\mathrm{d}\,z}{\mathrm{d}\,\alpha_N(z)} = \frac{1}{2\pi i \alpha_N(z) [1 - \alpha_N(z)] [P_\nu(1 - 2\alpha_N(z))]^2},\tag{4.44}$$

and observe that (cf. [38, Theorem 4.31])

$$[1 - 2\alpha_N(z)][P_\nu(1 - 2\alpha_N(z))]^4 = \frac{N^2 E_4(Nz) - E_4(z)}{N^2 - 1}.$$
(4.45)

On the right-hand side of (4.35^{*}), one can pick a path connecting z to $\frac{\text{Re}z}{2|\text{Re}z|} + \frac{i}{2}\sqrt{\frac{4-N}{N}}$ that does not cross the positive Im *z*-axis, on which $\alpha_N(z) \in (0,1)$ corresponds to the branch cut for the left-hand side of (4.35^{*}).

To derive the imaginary part of (4.12) from (4.35^*) , note that

$$\left[2t(1-t)\frac{\mathrm{d}}{\mathrm{d}\,t} + R_{\nu}(1-2t)\right]f(t)\bigg|_{t=\alpha_{N}(z)} = -\frac{\mathrm{Im}\,z}{\pi}\frac{\partial}{\partial\,\mathrm{Im}\,z}\left\{\frac{1}{\mathrm{Im}\,z}\frac{f(\alpha_{N}(z))}{[P_{\nu}(1-2\alpha_{N}(z))]^{2}}\right\}$$
(4.46)

for z satisfying (2.26).

The imaginary part of (4.12) splits into three summands:

• an elementary contribution

$$-\frac{4\pi^{2}}{\mathrm{Im}\,z}\,\mathrm{Re}\int_{z}^{\frac{\mathrm{Re}\,z}{2|\mathrm{Re}\,z|}+\frac{i}{2}\sqrt{\frac{4-N}{N}}}(z-w)(\overline{z}-w)\,\mathrm{d}\,w$$

$$=\frac{4\pi^{2}}{3\,\mathrm{Im}\,z}\left(\mathrm{Re}\,z-\frac{\mathrm{Re}\,z}{2|\,\mathrm{Re}\,z|}\right)\left[\left(\mathrm{Re}\,z-\frac{\mathrm{Re}\,z}{2|\,\mathrm{Re}\,z|}\right)^{2}+3(\mathrm{Im}\,z)^{2}-\frac{3(4-N)}{4N}\right];$$
(4.47)

• an expression

$$\frac{4\pi^{2}}{\operatorname{Im} z} \operatorname{Re} \int_{z}^{i\infty} \frac{E_{4}(w) - N^{2} E_{4}(Nw) - 1 + N^{2}}{N^{2} - 1} (z - w) (\overline{z} - w) \, \mathrm{d} \, w$$

$$= \frac{4\pi^{2} \operatorname{Re} \left[\widetilde{\mathscr{E}}_{4}(Nz) - N \widetilde{\mathscr{E}}_{4}(z) \right]}{N(N^{2} - 1) \operatorname{Im} z} \tag{4.48}$$

involving Eichler-type integrals defined in (4.15);

³For two fixed real numbers Re *s*, Re *t* > 1, we have $\lim_{\text{Im } s \to 0^+, \text{Im } t \to 0^+} [P_{\nu}(1-2s)P_{\nu}(2t-1) - P_{\nu}(2s-1)P_{\nu}(1-2t)]^2 = \lim_{\text{Im } s \to 0^-, \text{Im } t \to 0^-} [P_{\nu}(1-2s)P_{\nu}(2t-1) - P_{\nu}(2s-1)P_{\nu}(1-2t)]^2 \in \mathbb{R}$. One has a similar argument for two fixed real numbers Re *s*, Re *t* < 0.

• an integral

$$\frac{4\pi^{2}}{\mathrm{Im}\,z}\operatorname{Re}\int_{i\infty}^{\frac{\mathrm{Re}z}{2|\mathrm{Re}z|}+\frac{i}{2}\sqrt{\frac{4-N}{N}}} \frac{E_{4}(w) - N^{2}E_{4}(Nw) - 1 + N^{2}}{N^{2} - 1} (z - w)(\overline{z} - w) \,\mathrm{d}\,w$$

$$= \frac{8\pi^{2}}{\mathrm{Im}\,z} \left(\operatorname{Re}\,z - \frac{\mathrm{Re}\,z}{2|\mathrm{Re}\,z|}\right) \operatorname{Re}\int_{i\infty}^{\frac{\mathrm{Re}z}{2|\mathrm{Re}\,z|}+\frac{i}{2}\sqrt{\frac{4-N}{N}}} \frac{E_{4}(w) - N^{2}E_{4}(Nw) - 1 + N^{2}}{N^{2} - 1} \left(w - \frac{\mathrm{Re}\,z}{2|\mathrm{Re}\,z|}\right) \,\mathrm{d}\,w$$

$$= \frac{8}{\mathrm{Im}\,z} \left(\operatorname{Re}\,z - \frac{\mathrm{Re}\,z}{2|\mathrm{Re}\,z|}\right) \int_{1}^{\infty} \frac{x[P_{\nu}(x)]^{4} - 1}{(1 - x^{2})[P_{\nu}(x)]^{2}} \left[\frac{\pi\cot(\nu\pi)}{2} - \frac{Q_{\nu}(x)}{P_{\nu}(x)}\right] \,\mathrm{d}\,x$$

$$= \frac{8\pi^{2}}{\mathrm{Im}\,z} \left(\operatorname{Re}\,z - \frac{\mathrm{Re}\,z}{2|\mathrm{Re}\,z|}\right) \frac{6 - N}{6N}$$
(4.49)

where the identities

$$\int_{1}^{\infty} \frac{x[P_{\nu}(x)]^{4} - 1}{(1 - x^{2})[P_{\nu}(x)]^{2}} dx = \lim_{\delta \to 0^{+}} \left\{ \int_{1+\delta}^{\infty} \frac{x[P_{\nu}(x)]^{2} dx}{1 - x^{2}} + \frac{Q_{\nu}(1 + \delta)}{P_{\nu}(1 + \delta)} \right\}$$

$$= \lim_{\delta \to 0^{+}} \left\{ \int_{1+\delta}^{\infty} \frac{x[P_{\nu}(x)]^{2} dx}{1 - x^{2}} - \frac{2\gamma_{0} + 2\psi^{(0)}(\nu) + \log\frac{\delta}{2}}{2} \right\} = \frac{\pi \cot(\nu\pi)}{2},$$

$$\int_{1}^{\infty} \frac{x[P_{\nu}(x)]^{4} - 1}{(1 - x^{2})[P_{\nu}(x)]^{2}} \frac{Q_{\nu}(x)}{P_{\nu}(x)} dx = \lim_{\delta \to 0^{+}} \left\{ \int_{1+\delta}^{\infty} \frac{xP_{\nu}(x)Q_{\nu}(x) dx}{1 - x^{2}} + \frac{[Q_{\nu}(1 + \delta)]^{2}}{2[P_{\nu}(1 + \delta)]^{2}} \right\}$$

$$= \lim_{\delta \to 0^{+}} \left\{ \int_{1+\delta}^{\infty} \frac{xP_{\nu}(x)Q_{\nu}(x) dx}{1 - x^{2}} + \frac{[2\gamma_{0} + 2\psi^{(0)}(\nu) + \log\frac{\delta}{2}]^{2}}{8} \right\} = -\frac{\pi^{2}}{12},$$

$$(4.51)$$

are deducible from the methods in [97, §3.1].

These can be rearranged into (4.14). When $2 \operatorname{Re} z \in \mathbb{Z}$, the integrand of $\widetilde{\mathscr{E}}_4(z)$ is real-valued along an integration path parallel to the Im *z*-axis, thus making $\widetilde{\mathscr{E}}_4(z)$ purely imaginary; when $2 \operatorname{Re} \frac{1}{z} \in \mathbb{Z}$, we may divide Ramanujan's reflection formula [11, p. 276]

$$\mathscr{E}_4(z) - z^2 \mathscr{E}_4\left(-\frac{1}{z}\right) = -\frac{z^4 - 5z^2 + 1}{3z} + \frac{30i\zeta(3)\left(z^2 - 1\right)}{\pi^3}$$
(4.52)

for $\mathscr{E}_4(z) := \int_z^{i\infty} [1 - E_4(w)](z - w)^2 dw = \frac{60i}{\pi^3} \sum_{k=1}^{\infty} \frac{1}{k^3(e^{-2\pi i k z} - 1)}$ by Im *z* before differentiating with respect to Im *z*, so that the resulting reflection formula

$$\frac{|z|^2}{(\operatorname{Im} z)^2} \operatorname{Re} \widetilde{\mathscr{E}}_4 \left(-\frac{1}{z} \right) - \frac{\operatorname{Re} \widetilde{\mathscr{E}}_4 (z)}{(\operatorname{Im} z)^2} = \frac{\operatorname{Re} z}{3(\operatorname{Im} z)^2} \left[\frac{|z|^2 + 2(\operatorname{Im} z)^2}{|z|^4} + |z|^2 + 2(\operatorname{Im} z)^2 - 5 \right]$$
(4.53)

brings us a closed-form evaluation of Re $\widetilde{\mathscr{E}}_4(z)$ in (4.16).

Remark As pointed out by Guillera–Rogers [54, §4.1] and Sun [71, §3], particular cases of (4.12) have been investigated by several authors, including Zeilberger [93], Guillera [52, 53], and Hessami Pilehrood–Hessami Pilehrood [55].

Remark As noted by Guillera–Rogers [54, around (43)], there are no automorphic interpretations for the $\nu = -\frac{1}{6}$ case. This comes as no surprise, since the case violates $\Gamma_0(1)$ symmetry in two

aspects: (1) The following integral representation (cf. [97, (2.2.13)-(2.2.14)])

$$E^{\Gamma_0(1)}(z,2) = -\frac{3}{4} \operatorname{Re} \left[2t(1-t)\frac{\mathrm{d}}{\mathrm{d}t} + R_{-1/6}(1-2t) \right] \int_0^1 [P_{-1/6}(1-2s)]^2 \\ \times \left(\frac{1}{s-t} - \frac{1}{s-1+t} \right) \mathrm{d}s \bigg|_{t=\frac{1}{2} - \frac{1}{2}\sqrt{\frac{j(z)-1728}{j(z)}}}$$
(4.54)

is incompatible with the real part for the right-hand side of (4.35); (2) When $t = \frac{1}{2} - \frac{1}{2}\sqrt{\frac{j(z)-1728}{j(z)}}$ for a certain z satisfying Im z > 0, $|\operatorname{Re} z| < \frac{1}{2}$, |z| > 1, the expression [cf. (4.45)]

$$(1-2t)[P_{-1/6}(1-2t)]^4 = \sqrt{\frac{j(z) - 1728}{j(z)}}E_4(z)$$
(4.55)

disqualifies the right-hand side of (4.35) from being an Eichler integral for a holomorphic modular form of weight 4.

5. Clebsch–Gordan couplings and binomial harmonic sums

Using the Mehler–Dirichlet integral representation for the Legendre functions

$$P_{\nu}(\cos\theta) \coloneqq \frac{2}{\pi} \int_{0}^{\theta} \frac{\cos\frac{(2\nu+1)\beta}{2} \,\mathrm{d}\beta}{\sqrt{2(\cos\beta - \cos\theta)}}, \qquad \qquad \theta \in (0,\pi)$$
(5.1)

of arbitrary complex degrees $v \in \mathbb{C}$, we define the generalized Clebsch–Gordan integrals

$$T_{\mu,\nu} := \int_{-1}^{1} P_{\mu}(x) P_{\nu}(x) P_{\nu}(-x) \,\mathrm{d}\,x, \qquad \mu, \nu \in \mathbb{C}, \qquad (5.2)$$

About a decade ago, closed forms were found [95, 96] for $T_{\nu,\nu}$, $T_{2\nu-1,\nu}$, and \tilde{T}_{ν} , along with some generalized hypergeometric representations for $T_{\mu,\nu}$. Recently, interest in these Clebsch–Gordan integrals has been revived by Cantarini [29] and Campbell [24, 25, 27].

In §§5.1–5.2, we review and re-derive the main results from [95, 96], using automated manipulations of generalized hypergeometric series. Some series identities therein can be reinterpreted as closed-form evaluations of binomial harmonic sums, as in (1.15)-(1.17).

In §5.3, we extend the Clebsch–Gordan couplings to associated Legendre functions [see (3.18)], and study yet another type of integral over the products of three Legendre functions, namely

$$U_{\mu,\nu} := \int_{-1}^{1} \frac{P_{\mu}(x) + P_{\mu}(-x)}{2} [P_{\nu}(x)]^2 \,\mathrm{d}\,x, \qquad \mu, \nu \in \mathbb{C}.$$
(5.4)

These efforts will culminate in a new theorem that is not found in [95, 96].

Theorem 5.1 (Correspondence between $T_{\mu,\nu}$ and $U_{\mu,\nu}$). If we define

$$c_{\mu,\nu}(\varepsilon) \coloneqq \frac{\Gamma(\frac{1}{2} + \varepsilon)\Gamma(-\nu + \varepsilon)\Gamma(\nu + 1 + \varepsilon)}{\Gamma(1 + \varepsilon)\Gamma(\frac{2-\mu}{2} + \varepsilon)\Gamma(\frac{\mu+3}{2} + \varepsilon)},$$
(5.5)

then we have

$$T_{\mu,\nu} = \frac{2\sin(\mu\pi)\sin(\nu\pi)}{\mu(\mu+1)\pi^2} \left. \frac{\partial}{\partial\varepsilon} \right|_{\varepsilon=0} \left[\frac{c_{\mu,\nu}(\varepsilon)}{c_{\mu,\nu}(0)} {}_4F_3 \begin{pmatrix} 1, \frac{1}{2} + \varepsilon, -\nu + \varepsilon, \nu + 1 + \varepsilon \\ 1 + \varepsilon, \frac{2-\mu}{2} + \varepsilon, \frac{\mu+3}{2} + \varepsilon \\ 1 \end{pmatrix} \right], \tag{5.6}$$

$$U_{\mu,\nu} = \frac{\sin(\mu\pi)\sin^2(\nu\pi)}{\mu(\mu+1)\pi^3} \left. \frac{\partial^2}{\partial\varepsilon^2} \right|_{\varepsilon=0} \left[\frac{c_{\mu,\nu}(\varepsilon)}{c_{\mu,\nu}(0)} {}_4F_3 \left(\begin{array}{c} 1, \frac{1}{2} + \varepsilon, -\nu + \varepsilon, \nu + 1 + \varepsilon \\ 1 + \varepsilon, \frac{2-\mu}{2} + \varepsilon, \frac{\mu+3}{2} + \varepsilon \end{array} \right] 1 \right], \tag{5.7}$$

when $\mu \notin \mathbb{Z}$.

5.1. Clebsch–Gordan couplings of Legendre functions. In this subsection, we recall some results from our previous works [95, 96], and compare them to certain consequences of Lemma 3.2.

Proposition 5.2 (Evaluations of $T_{\mu,\nu}$). If $2\mu \notin \mathbb{Z}$ and $2\nu \notin \mathbb{Z}$, then we have the following representations

$$T_{\mu,\nu} = \frac{2}{\pi} \frac{\sin(\mu\pi)\cos(\nu\pi)}{2\nu+1} \left[\frac{1}{\mu} {}_{5}F_{4} \left(\frac{\frac{1}{2}, \frac{1}{2}, -\frac{\mu}{2}, -\nu, \nu+1}{1, \frac{2-\mu}{2}, \frac{1-2\nu}{2}, \frac{2\nu+3}{2}} \right| 1 \right) - \frac{1}{\mu+1} {}_{5}F_{4} \left(\frac{\frac{1}{2}, \frac{1}{2}, \frac{1+\mu}{2}, -\nu, \nu+1}{1, \frac{3+\mu}{2}, \frac{1-2\nu}{2}, \frac{2\nu+3}{2}} \right| 1 \right) \right]$$
(5.8a)

$$= \frac{2}{\pi^2} \frac{\sin(\mu\pi)\sin(\nu\pi)}{\mu(\mu+1)} \left[\frac{1}{\nu} {}_{4}F_3\left(\frac{1,\frac{1-\mu}{2},\frac{\mu+2}{2},-\nu}{\frac{2-\mu}{2},\frac{\mu+3}{2},1-\nu} \right) 1 \right] - \frac{1}{\nu+1} {}_{4}F_3\left(\frac{1,\frac{1-\mu}{2},\frac{\mu+2}{2},\nu+1}{\frac{2-\mu}{2},\frac{\mu+3}{2},\nu+2} \right) \right]$$
(5.8b)

$$= \frac{2}{\pi^{2}} \frac{\sin(\mu\pi)\sin(\nu\pi)}{\mu(\mu+1)} \left\{ \frac{\partial}{\partial\varepsilon} \Big|_{\varepsilon=0} {}_{4}F_{3} \begin{pmatrix} 1, \frac{1}{2} + \varepsilon, -\nu + \varepsilon, \nu + 1 + \varepsilon \\ 1 + \varepsilon, \frac{2-\mu}{2} + \varepsilon, \frac{\mu+3}{2} + \varepsilon \end{pmatrix} \Big| 1 \right\}$$
$$- \left[\psi^{(0)} \left(\frac{\mu+3}{2} \right) + \psi^{(0)} \left(\frac{2-\mu}{2} \right) - \psi^{(0)} (\nu+1) - \psi^{(0)} (-\nu) + 2\log 2 \right]$$
$$\times {}_{3}F_{2} \left(\frac{\frac{1}{2}, -\nu, \nu+1}{\frac{2-\mu}{2}, \frac{\mu+3}{2}} \right) \Big| 1 \right\}$$
(5.8c)

involving generalized hypergeometric series [cf. (1.11)]. In particular, we have

$$T_{\nu,\nu} = \frac{1 + 2\cos(\nu\pi)}{3} \frac{\pi\Gamma(\frac{\nu+1}{2})\Gamma(\frac{3\nu+2}{2})}{\left[\Gamma(\frac{1-\nu}{2})\right]^2 \left[\Gamma(\frac{\nu+2}{2})\right]^3 \Gamma(\frac{3\nu+3}{2})},$$
(5.9)

$$T_{2\nu-1,\nu} = \frac{\sin(\nu\pi)\sin(2\nu\pi)}{\nu^2\pi^2},$$
(5.10)

while interpreting indeterminate forms through l'Hôpital's rule.

Proof. The ${}_{5}F_{4}$ (resp. ${}_{4}F_{3}$) representation in (5.8a) [resp. (5.8b)] comes from [95, (19^{*})] (resp. [95, (19)]).

To prove (5.8c) [which is equivalent to (5.6)], use (3.13)-(3.14) for

$$T_{\mu,\nu} = 2 \int_0^1 P_{\mu} (1-2t) P_{\nu} (1-2t) P_{\nu} (2t-1) dt$$

= $\frac{2\sin(\nu\pi)}{\pi} \int_0^1 P_{\mu} (1-2t) \sum_{k=0}^{\infty} [t(1-t)]^k s_{\nu,k} \left[\varsigma_{\nu,k} - \log \frac{1}{4t(1-t)} \right] dt,$ (5.11)

and then perform termwise integration with the assistance from (2.57).

One can find the closed forms of $T_{\nu,\nu}$ and $T_{2\nu-1,\nu}$ in [95, $(19_{(\nu,\nu)})$] and [95, $(19_{(2\nu-1,\nu)})$].

TABLE 5. Selected series related to $T_{\nu,\nu}$

ν	$T_{\nu,\nu}$
$-\frac{1}{2}$	$\frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^8}{24\pi^5} = 4\sum_{k=0}^{\infty} \frac{\binom{2k}{k}^4}{2^{8k}} \frac{1}{4k+1}$
$-\frac{1}{3}$	$\frac{3\sqrt{3}\left[\Gamma\left(\frac{1}{3}\right)\right]^9}{32\pi^5} = \frac{27\sqrt{3}}{4\pi} \sum_{k=0}^{\infty} \frac{\binom{2k}{k}^3}{\binom{6k}{3k}} \frac{4k+1}{(6k+1)^2(3k+1)}$
$-\frac{1}{4}$	$\frac{\left[\Gamma(\frac{1}{8})\Gamma(\frac{3}{8})\right]^2}{3\pi^3} = \frac{32}{\pi} \sum_{k=0}^{\infty} \frac{\binom{2k}{k}^2}{2^{4k}} \frac{1}{(8k+1)(8k+3)}$
$-\frac{1}{6}$	$\frac{\left[\Gamma\left(\frac{1}{4}\right)\right]^4}{\sqrt{2}\sqrt[4]{3}\pi^3} = \frac{27\sqrt{3}}{\pi} \sum_{k=0}^{\infty} \frac{\binom{2k}{k}\binom{6k}{3k}}{2^{8k}} \frac{4k+1}{(3k+1)(12k+1)(12k+5)}$

TABLE 6. Selected series related to $T_{2\nu-1,\nu}$

Remark The generalized hypergeometric series [cf. (1.11)]

$${}_{p}F_{p-1}\left(\left.\begin{array}{c}a_{1},\ldots,a_{p}\\b_{1},\ldots,b_{p-1}\end{array}\right|1\right) := 1 + \sum_{n=1}^{\infty}\frac{\prod_{j=1}^{p}(a_{j})_{n}}{\prod_{k=1}^{p-1}(b_{k})_{n}}\frac{1}{n!}$$
(5.12)

converges when $(\{a_1,\ldots,a_p\} \cup \{b_1,\ldots,b_{p-1}\}) \cap \mathbb{Z}_{\leq 0} = \emptyset$ and

$$\operatorname{Re}\sum_{j=1}^{p} a_{j} < \operatorname{Re}\sum_{k=1}^{p-1} b_{k}.$$
(5.13)

Therefore, all the generalized hypergeometric series appearing in (5.8a)-(5.8c) are indeed convergent. \Box

Remark Laporta [61, (48)] and Wan [78, (54)–(55)] independently discovered

$$\int_{0}^{1} \left[\mathbf{K}(\sqrt{t}) \right]^{2} \mathbf{K}(\sqrt{1-t}) \, \mathrm{d} \, t = \frac{\left[\Gamma(\frac{1}{4}) \right]^{8}}{384\pi^{2}}$$
(5.14)

through numerical experiments, where $\mathbf{K}(\sqrt{t}) = \frac{\pi}{2}P_{-1/2}(1-2t)$. The same was proved by the present author [95] (via a special case $T_{-1/2,-1/2}$) and by Rogers–Wan–Zucker [67] (as one of various lattice sums representable through modular *L*-values).

Remark In Table 5 (resp. Table 6), we showcase the series representations based on (5.8a) for $T_{\nu,\nu}$ (resp. low-order derivatives of $T_{2\nu-1,\nu}$), with limit procedures $\mu = \nu \rightarrow -\frac{1}{2}$ (resp. $\nu = \frac{\mu+1}{2} \rightarrow -\frac{1}{2}$) wherever necessary. As one may note,

$$\frac{1}{\pi} \frac{\partial T_{2\nu-1,\nu}}{\partial \nu} \bigg|_{\nu=-1/2}, \quad \frac{T_{-1/2,-1/2}}{4}, \quad \frac{1}{8} \left. \frac{\partial^2 T_{\nu,\nu}}{\partial \nu^2} \right|_{\nu=-1/2}$$
(5.15)

can be paraphrased into (1.15)-(1.17) mentioned in the introduction.

To spell out the summands for series (5.8b) and (5.8c), one needs generalized binomial coefficients $\binom{n}{m} \coloneqq \frac{\Gamma(n+1)}{\Gamma(m+1)\Gamma(n-m+1)}$, as well as values of the digamma function at fractional arguments, which do not fit the pattern in (1.1).

Remark Some series identities in Tables 5 and 6 involve $\binom{2k}{k}^4$, which agree in part with a result of Cantarini and D'Aurizio [30, Corollaries 1 and 2]: the relations

$$\pi^{2} \sum_{k=0}^{\infty} \frac{\binom{2k}{k}^{4}}{2^{8k}} \frac{4k+1}{p(k)} \in \mathbb{Q}, \quad \pi^{2} \sum_{k=0}^{\infty} \frac{\binom{2k}{k}^{4}}{2^{8k}} \frac{4k+1}{[p(k)]^{2}} \in \mathbb{Q}$$
(5.16)

hold for infinitely many polynomials $p(k) \in \mathbb{Z}[k]$ with integer coefficients. Formulae for infinite series involving $\binom{2k}{k}^4$ are otherwise rare, being notably absent from Sun's recent conjectures [71, §3].

Remark Cantarini–D'Aurizio [30], Campbell–Cantarini–D'Aurizio [28], Cantarini [29], and Campbell [25] have evaluated a wide variety of binomial harmonic sums using Fourier–Legendre expansions, which are independent of the approach we employ in the present article. Campbell has also published a recent proof [27] of (1.16) by Fourier–Legendre expansions.

Proposition 5.3 (Evaluations of \widetilde{T}_{ν}). If Re $\nu > -1$, then

$$\widetilde{T}_{\nu} = \frac{1}{\pi} \left(\frac{\cos \frac{\nu \pi}{2}}{2^{\nu}} \right)^3 \left[\frac{\Gamma(\frac{1+\nu}{2})}{\Gamma(\frac{2+\nu}{2})} \right]^4$$
(5.17)

$$= \frac{\sin(\nu\pi)\cos\frac{\nu\pi}{2}}{2^{\nu}\pi} \left[\frac{\Gamma(\frac{1+\nu}{2})}{\Gamma(\frac{2+\nu}{2})}\right]^2 \sum_{k=0}^{\infty} \frac{\left(\frac{1+\nu}{2}\right)_k^2 (-\nu)_k}{(k!)^3} \times \left[2\psi^{(0)}\left(k+\frac{\nu+1}{2}\right) + \psi^{(0)}(k-\nu) - 3\psi^{(0)}(k+1)\right].$$
(5.18)

Proof. The last equality in [96, (1.1)] implies (5.17). To show (5.18), apply (3.13)-(3.14) to

$$\widetilde{T}_{\nu} = 2 \int_{0}^{1} \frac{[P_{\nu}(1-2t)]^{2} P_{\nu}(2t-1) \,\mathrm{d} t}{[4t(1-t)]^{\frac{1-\nu}{2}}} = \frac{2\sin(\nu\pi)}{\pi} \int_{0}^{1} \frac{P_{\nu}(1-2t)}{[4t(1-t)]^{\frac{1-\nu}{2}}} \sum_{k=0}^{\infty} [t(1-t)]^{k} s_{\nu,k} \left[\varsigma_{\nu,k} - \log\frac{1}{4t(1-t)}\right] \mathrm{d} t,$$
(5.19)

and then integrate termwise using (2.57).

5.2. **Bailey–Meijer formulations of Clebsch–Gordan couplings**. In this subsection, we give self-contained proofs of (5.9), (5.10) and (5.17), without going through the heavy computations in existent literature [95, 96].

A major tool in this subsection is Bailey's reduction [8, (3.4)] of a convergent and well-poised $_7F_6$ series into (what is currently known as) a special class of the Meijer *G*-function (cf. [103,

Proposition 2] and [15, Figure 3]):

$${}_{7}F_{6}\left(\begin{array}{ccc}a,1+\frac{a}{2}, & b, & c, & d, & e, & f\\ \frac{a}{2},1+a-b,1+a-c,1+a-d,1+a-e,1+a-f\\ 1\end{array}\right)$$

$$=\frac{\Gamma(1+a-b)\Gamma(1+a-c)\Gamma(1+a-d)\Gamma(1+a-e)\Gamma(1+a-f)}{\Gamma(1+a)\Gamma(b)\Gamma(c)\Gamma(d)\Gamma(1+a-b-c)\Gamma(1+a-b-d)\Gamma(1+a-c-d)\Gamma(1+a-e-f)} \quad (5.20)$$

$$\times G_{4,4}^{2,4}\left(1\begin{vmatrix} e+f-a,1-b,1-c,1-d\\ 0,1+a-b-c-d,e-a,f-a\\ \end{vmatrix}\right).$$

Here, on the left-hand side of (5.20), hypergeometric parameters sitting in the same column⁴ all add up to the same number 1 + a (hence the name "well-poised"); on the right-hand side of (5.20), the Meijer *G*-function is defined by an integral of Mellin–Barnes type:

$$G_{p,q}^{m,n}\left(z \begin{vmatrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{vmatrix} := \frac{1}{2\pi i} \int_C \frac{\prod_{j=1}^n \Gamma(1 - a_j - s) \prod_{k=1}^m \Gamma(b_k + s)}{\prod_{j=n+1}^p \Gamma(a_j + s) \prod_{k=m+1}^q \Gamma(1 - b_k - s)} \frac{\mathrm{d}\,s}{z^s},\tag{5.21}$$

where the canonical design of Barnes' contour C ensures that the right-hand side of (5.21) equals the sum over the residues of

$$-\frac{\prod_{j=1}^{n}\Gamma(1-a_{j}-s)\prod_{k=1}^{m}\Gamma(b_{k}+s)}{\prod_{j=n+1}^{p}\Gamma(a_{j}+s)\prod_{k=m+1}^{q}\Gamma(1-b_{k}-s)\frac{1}{z^{s}}}$$
(5.22)

at all the poles in $\prod_{j=1}^{n} \Gamma(1 - a_j - s)$, when the contour closes rightwards. By convention, all the empty products are treated as unity.

Proposition 5.4 (Bailey–Meijer formulations of $T_{\mu,\nu}$). (a) If $2\mu \notin \mathbb{Z}$ and $2\nu \notin \mathbb{Z}$, then we have

$$T_{\mu,\nu} = \frac{2\sin(\mu\pi)\cos(\nu\pi)}{\mu(\mu+1)(2\nu+1)\pi} {}_{7}F_{6} \begin{pmatrix} \frac{1}{2}, \frac{5}{4}, \frac{1}{2}, -\frac{\mu}{2}, \frac{\mu+1}{2}, -\nu, \nu+1\\ \frac{1}{4}, 1, \frac{\mu+3}{2}, \frac{2-\mu}{2}, \frac{2\nu+3}{2}, \frac{1-2\nu}{2} \\ \end{pmatrix}$$
(5.23a)

$$=\frac{2\sin(\mu\pi)\sin(\nu\pi)}{\mu(\mu+1)\nu(\nu+1)\pi^{2}}{}_{6}F_{5}\left(\begin{array}{cc}1,\frac{3}{2},\frac{1-\mu}{2},\frac{\mu+2}{2},&-\nu,\nu+1\\\frac{1}{2},\frac{\mu+3}{2},\frac{2-\mu}{2},\nu+2,1-\nu\end{array}\right|1\right);$$
(5.23b)

if $2\mu \notin \mathbb{Z}$, $2\nu \notin \mathbb{Z}$, and $\operatorname{Re} \nu > -1$, then we have

$$T_{\mu,\nu} = \frac{2(1-2\nu)\sin(\mu\pi)\sin(\nu\pi)}{\mu(\mu+1)\nu\pi^2} {}_{7}F_6 \begin{pmatrix} \frac{1-2\nu}{2}, \frac{5-2\nu}{4}, & 1, & \frac{1}{2}, \frac{\mu-2\nu+1}{2}, -\frac{\mu+2\nu}{2}, -\nu\\ \frac{1-2\nu}{4}, \frac{1-2\nu}{2}, 1-\nu, & \frac{2-\mu}{2}, & \frac{\mu+3}{2}, & \frac{3}{2} \end{pmatrix} | 1 \end{pmatrix}$$

$$= \frac{2(1-2\nu)\sin(\mu\pi)\sin(\nu\pi)}{\mu(\mu+1)\nu\pi^2} {}_{6}F_5 \begin{pmatrix} 1, \frac{5-2\nu}{4}, & \frac{1}{2}, \frac{\mu-2\nu+1}{2}, -\frac{\mu+2\nu}{2}, -\nu\\ \frac{1-2\nu}{4}, 1-\nu, & \frac{2-\mu}{2}, & \frac{\mu+3}{2}, & \frac{3}{2} \end{pmatrix} | 1 \end{pmatrix};$$
(5.23c)

if $2\mu \notin \mathbb{Z}$, $2\nu \notin \mathbb{Z}$, and $\operatorname{Re} \nu < 0$, then we have

$$T_{\mu,\nu} = -\frac{2(2\nu+3)\sin(\mu\pi)\sin(\nu\pi)}{\mu(\mu+1)(\nu+1)\pi^2} {}_{7}F_6 \begin{pmatrix} \frac{2\nu+3}{2}, \frac{2\nu+7}{4}, & 1, & \frac{1}{2}, \frac{\mu+2\nu+3}{2}, \frac{2-\mu+2\nu}{2}, \nu+1 \\ \frac{2\nu+3}{4}, \frac{2\nu+3}{2}, \nu+2, & \frac{2-\mu}{2}, & \frac{\mu+3}{2}, & \frac{3}{2} \end{pmatrix} | 1 \end{pmatrix}$$

$$= -\frac{2(2\nu+3)\sin(\mu\pi)\sin(\nu\pi)}{\mu(\mu+1)(\nu+1)\pi^2} {}_{6}F_5 \begin{pmatrix} 1, \frac{2\nu+7}{4}, & \frac{1}{2}, \frac{\mu+2\nu+3}{2}, \frac{2-\mu+2\nu}{2}, \nu+1 \\ \frac{2\nu+3}{4}, \nu+2, & \frac{2-\mu}{2}, & \frac{\mu+3}{2}, & \frac{3}{2} \end{pmatrix} | 1 \end{pmatrix}.$$
(5.23d)

⁴In this article, we typeset each well-poised series with aligned columns for its hypergeometric parameters, following the practice of Slater [69, §2.4].

Furthermore, for $\mu \notin \mathbb{Z}$ and $2\nu \notin \mathbb{Z}$, we have

$$\begin{split} T_{\mu,\nu} &= \sin(\mu\pi)\sin(\nu\pi) \left(\frac{1}{\nu(\nu+1)\pi^2} {}_{4}F_{3} \left(\begin{array}{c} 1,1,\frac{1-\mu}{2},\frac{\mu+2}{2} \\ \frac{3}{2},1-\nu,\nu+2 \end{array} \right) 1 \right) \\ &+ \frac{\Gamma\left(\frac{\mu+1}{2}\right)\Gamma\left(-\frac{\mu}{2}\right)}{\pi} \left\{ \frac{\tan(\nu\pi)}{2\nu+1} \frac{1}{\Gamma\left(\frac{1-\mu}{2}\right)\Gamma\left(\frac{\mu+2}{2}\right)} {}_{3}F_{2} \left(\begin{array}{c} \frac{1}{2},\frac{1+\mu}{2},-\frac{\mu}{2} \\ \frac{1-2\nu}{2},\frac{2\nu+3}{2} \end{array} \right) 1 \right) \\ &- \frac{\sin(\nu\pi)}{2^{2\nu+1}\pi^2} \frac{[\Gamma(-\nu)\Gamma(2\nu+1)]^2}{\Gamma\left(\frac{2-\mu+2\nu}{2}\right)\Gamma\left(\frac{\mu+2\nu+3}{2}\right)} {}_{3}F_{2} \left(\begin{array}{c} -\frac{\mu+2\nu+1}{2},\frac{\mu-2\nu}{2},-\nu \\ \frac{1-2\nu}{2},-2\nu \end{array} \right) 1 \right) \\ &- \frac{2^{2\nu+1}\sin(\nu\pi)}{\pi^2} \frac{[\Gamma(\nu+1)\Gamma(-2\nu-1)]^2}{\Gamma\left(\frac{\mu-2\nu+1}{2}\right)\Gamma\left(-\frac{\mu+2\nu}{2}\right)} {}_{3}F_{2} \left(\begin{array}{c} \frac{1-\mu+2\nu}{2},\frac{2+\mu+2\nu}{2},\nu+1 \\ \frac{2\nu+3}{2},2\nu+2 \end{array} \right) 1 \right) \bigg\} \bigg). \end{split}$$
(5.23e)

(b) The expression

$$\varphi(\nu) := \frac{T_{\nu,\nu} - \frac{1+2\cos(\nu\pi)}{3} \frac{\pi\Gamma(\frac{\nu+1}{2})\Gamma(\frac{3\nu+2}{2})}{\left[\Gamma(\frac{1-\nu}{2})\right]^2 \left[\Gamma(\frac{\nu+2}{2})\right]^3 \Gamma(\frac{3\nu+3}{2})}{\left[1+2\cos(\nu\pi)\right] \cos\frac{\nu\pi}{2}}$$
(5.24)

extends to a holomorphic function for $\operatorname{Re} v \ge 0$ that vanishes for every $v \in \mathbb{Z}_{\ge 0}$ and has at most $O(e^{\pi |v|/2})$ growth rate for large |v|, hence $\varphi(v) \equiv 0$. (c) For $v \in (-1,0) \setminus \{-\frac{1}{2}\}$, we have

$$T_{2\nu-1,\nu} = \frac{\sin(\nu\pi)\sin(2\nu\pi)}{\nu^2\pi^2} \frac{\nu\pi\cot(\nu\pi)}{(1-2\nu)(1+2\nu)} {}_5F_4\left(\begin{array}{ccc} \frac{1}{2}, \frac{5}{4}, \frac{1}{2}, & -\nu, & \nu\\ \frac{1}{4}, 1, \frac{3}{2}+\nu, \frac{3}{2}-\nu \end{array}\right)$$
(5.25a)

$$=\frac{\sin(\nu\pi)\sin(2\nu\pi)}{\nu^{2}\pi^{2}}\frac{1}{(1+\nu)(1-2\nu)}{}_{5}F_{4}\left(\begin{array}{cc}1,\frac{3}{2},1,&-\nu,\frac{1}{2}+\nu\\\frac{1}{2},1,2+\nu,\frac{3}{2}-\nu\end{array}\right|1\right)$$
(5.25b)

$$=\frac{\sin(\nu\pi)\sin(2\nu\pi)}{\nu^2\pi^2}\tag{5.25c}$$

$$=\frac{\sin(\nu\pi)\sin(2\nu\pi)}{\nu^2\pi^2}\frac{\nu(2\nu+3)}{(\nu+1)(2\nu-1)}{}_{5}F_4\left(\frac{\frac{2\nu+3}{2},\frac{2\nu+7}{4},\ 1,\ \frac{1}{2},1+2\nu}{\frac{2\nu+3}{4},\frac{2\nu+3}{2},2+\nu,\ \frac{3}{2}-\nu}\right|1\right).$$
(5.25d)

Proof. (a) We can rewrite (5.8c) as

$$T_{\mu,\nu} = \frac{2\sin(\mu\pi)\sin(\nu\pi)}{\mu(\mu+1)\pi^2} \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k (-\nu)_k (\nu+1)_k}{\left(\frac{2-\mu}{2}\right)_k \left(\frac{\mu+3}{2}\right)_k k!} \\ \times \left[-\psi^{(0)}\left(k + \frac{\mu+3}{2}\right) - \psi^{(0)}\left(k - \frac{\mu-2}{2}\right) + \psi^{(0)}(k+\nu+1) \right.$$
(5.26)
$$\left. + \psi^{(0)}(k-\nu) - \psi^{(0)}(k+1) + \psi^{(0)}\left(k + \frac{1}{2}\right) \right].$$

For $v \in (-1,0)$, this is equal to

$$-\frac{\Gamma(\frac{\mu+1}{2})\Gamma(-\frac{\mu}{2})\sin(\mu\pi)\sin^{2}(\nu\pi)}{2\pi^{3}\sqrt{\pi}}\lim_{\varepsilon \to 0^{+}}\int_{\varepsilon-i\infty}^{\varepsilon+i\infty}g_{\mu,\nu}(s)\frac{\mathrm{d}\,s}{2\pi i}$$

$$=-\frac{\Gamma(\frac{\mu+1}{2})\Gamma(-\frac{\mu}{2})\sin(\mu\pi)\sin^{2}(\nu\pi)}{2\pi^{3}\sqrt{\pi}}G_{4,4}^{2,4}\left(1\begin{vmatrix}\frac{1}{2},0,\nu+1,-\nu\\0,0,\frac{\mu}{2},-\frac{\mu+1}{2}\end{vmatrix}\right)$$
(5.27)

where

$$g_{\mu,\nu}(s) := \frac{[\Gamma(s)]^2 \Gamma(\frac{1}{2} - s) \Gamma(1 - s) \Gamma(-\nu - s) \Gamma(\nu + 1 - s)}{\Gamma(\frac{2-\mu}{2} - s) \Gamma(\frac{\mu+3}{2} - s)}$$
(5.28)

and the contour closes leftwards.

Due to the invariance of the generalized hypergeometric series and the Meijer *G*-functions under certain permutations of their parameters, there are four (ostensibly) different outcomes after we apply Bailey's reduction formula (5.20) to (5.27), as listed in (5.23a)–(5.23d). One can analytically continue these identities beyond the regime where $\nu \in (-1,0)$, so long as the generalized hypergeometric series in question remain convergent, according to the condition set forth in (5.13).

Alternatively, we may close the contour rightwards in (5.27), after which residue calculus brings us (5.23e) for $\nu \in (-1,0)$. Subsequent analytic continuation confirms (5.23e) in its entirety.

(b) By Dougall's theorem (see [69, (2.3.4.1)] or [6, (2.2.9)]), the well-poised series in (5.23a) reduces to a ratio of gamma values when -^μ/₂, ^{μ+1}/₂, -ν, ν + 1 ∈ Z_{≤0}. These are precisely the cases of [95, (19_(μ,n)) and (19_(2m,ν))], which were previously derived by totally different means.

In particular, the last paragraph allows us to confirm that $\varphi(\nu) = 0$ for all $\nu \in 2\mathbb{Z}_{\geq 0}$.

To show that $\varphi(2n+1) = 0$ for $n \in \mathbb{Z}_{\geq 0}$, we need to verify that the numerator in (5.24) vanishes to the order of $O((\nu - 2n - 1)^2)$ as $\nu \to 2n + 1$. This can be done via a particular case of (5.23b):

$$T_{\nu,\nu} = \frac{2\sin^2(\nu\pi)}{\nu^2(\nu+1)^2\pi^2} {}_{6}F_5 \begin{pmatrix} 1, \frac{3}{2}, \frac{1-\nu}{2}, \frac{\nu+2}{2}, & -\nu, \nu+1 \\ \frac{1}{2}, \frac{\nu+3}{2}, \frac{2-\nu}{2}, \nu+2, 1-\nu \\ \end{pmatrix} \\ = \frac{2\sin^2(\nu\pi)}{\nu^2(\nu+1)^2\pi^2} \sum_{k=0}^{\infty} \frac{\left(\frac{3}{2}\right)_k \left(\frac{1-\nu}{2}\right)_k \left(\frac{\nu+2}{2}\right)_k}{\left(\frac{1}{2}\right)_k \left(\frac{\nu+3}{2}\right)_k \left(\frac{2-\nu}{2}\right)_k} \frac{\nu(\nu+1)}{(\nu-k)(\nu+k+1)},$$
(5.23b')

where the infinite series terminates to a bounded quantity, as v approaches an odd positive integer.

To demonstrate that $\varphi(\nu)$ has a holomorphic extension for $\operatorname{Re} \nu \ge 0$, we still have to check that the numerator of (5.24) vanishes when $\nu = 2n + 1 \pm \frac{1}{3}$ for $n \in \mathbb{Z}_{\ge 0}$. For this purpose, we specialize (5.23c) to two sequences of terminating series

$$T_{2n+\frac{4}{3},2n+\frac{4}{3}} = -\frac{27(12n+5)_{6}F_{5}\begin{pmatrix} 1, \frac{1}{2}, -n+\frac{7}{12}, -n-\frac{1}{6}, -3n-2, -2n-\frac{4}{3} \\ -2n-\frac{1}{3}, -n-\frac{5}{12}, -n+\frac{1}{3}, n+\frac{13}{6}, \frac{3}{2} \end{pmatrix} 1}{8\pi^{2}(3n+2)^{2}(6n+7)}, \qquad (5.23c'_{+})$$

$$T_{2n+\frac{2}{3},2n+\frac{2}{3}} = -\frac{27(12n+1)_{6}F_{5}\begin{pmatrix} 1, \frac{1}{2}, -n+\frac{11}{12}, -n+\frac{1}{6}, -3n-1, -2n-\frac{2}{3} \\ -2n+\frac{1}{3}, -n-\frac{1}{12}, -n+\frac{2}{3}, n+\frac{11}{6}, \frac{3}{2} \end{pmatrix} 1}{8\pi^{2}(3n+1)^{2}(6n+5)},$$
(5.23c'_)

which can be computed through the recursions⁵

$$\begin{cases} T_{2(n+1)+\frac{4}{3},2(n+1)+\frac{4}{3}} = \frac{(n+1)(3n+4)(6n+7)^2}{(2n+3)(3n+5)^2(6n+11)} T_{2n+\frac{4}{3},2n+\frac{4}{3}}, \\ T_{2(n+1)+\frac{2}{3},2(n+1)+\frac{2}{3}} = \frac{(n+1)(3n+2)(6n+5)^2}{(2n+3)(3n+4)^2(6n+7)} T_{2n+\frac{2}{3},2n+\frac{2}{3}}, \end{cases}$$
(5.29)

⁵These recursions can be generated automatically from the Annihilator command in Christoph Koutschan's HolonomicFunctions package [60] for *Mathematica*, which implements the Wilf-Zeilberger algorithm [84, 91, 92]. It turns out that both recursions in (5.29) are valid for non-integer values of n as well [cf. (5.9)].

with initial conditions

$$\begin{cases} T_{\frac{4}{3},\frac{4}{3}} = 0, \\ T_{\frac{2}{3},\frac{2}{3}} = 0. \end{cases}$$
(5.30)

Thus far, we have verified that $\varphi(v)$ remains finite as $v \to 2n + 1 \pm \frac{1}{3}$ for $n \in \mathbb{Z}_{\geq 0}$.

To prove that $\varphi(\nu)$ has at most $O(e^{\pi|\nu|/2})$ growth rate for large $|\nu|$, apply Stirling's formula for Euler's gamma function to the right-hand sides of (5.9) and (5.23b').

By Carlson's theorem [6, Theorem 2.8.1], we conclude that $\varphi(\nu)$ is identically vanishing. (c) Setting $\mu = 2\nu - 1$ in (5.23a)–(5.23d), we arrive at (5.25a)–(5.25d).

For (5.25c), the hypergeometric series terminates after the leading constant term. [Plugging $\mu = 2\nu - 1$ into (5.23e), while employing some standard reduction formulae for $_{3}F_{2}$ and $_{2}F_{1}$, one gets the same result as (5.25c).]

One may also turn (5.25a), (5.25b), and (5.25d) into (5.25c) by a reduction formula (see [69, (2.3.4.5)] or [6, Corollary 3.5.2])

$${}_{5}F_{4}\left(\begin{array}{ccc}a,1+\frac{a}{2}, & b, & c, & d\\ \frac{a}{2},1+a-b,1+a-c,1+a-d\end{array}\right|1\right)$$

$$=\frac{\Gamma(1+a-b)\Gamma(1+a-c)\Gamma(1+a-d)\Gamma(1+a-b-c-d)}{\Gamma(1+a)\Gamma(1+a-b-c)\Gamma(1+a-b-d)\Gamma(1+a-c-d)}$$
(5.31)

for a special class of convergent and well-poised ${}_{5}F_{4}$.

Proposition 5.5 (Bailey–Meijer formulations of \widetilde{T}_{ν}). For $\operatorname{Re} \nu > -1$ and $\nu \notin \mathbb{Z}$, the following identities hold true:

$$\widetilde{T}_{\nu} = \frac{1}{\pi} \left(\frac{\cos \frac{\nu \pi}{2}}{2^{\nu}} \right)^{3} \left[\frac{\Gamma(\frac{1+\nu}{2})}{\Gamma(\frac{2+\nu}{2})} \right]^{2} \frac{2^{2\nu+1}(1-\nu)}{\nu(\nu+1)\cot \frac{\nu \pi}{2}} {}_{5}F_{4} \left(\frac{\frac{1-\nu}{2}}{4}, \frac{\frac{1-\nu}{2}}{4}, -\nu, \frac{\frac{1+\nu}{2}}{2}, 1-\nu \right) 1 \right)$$
(5.32)

$$= \frac{1}{\pi} \left(\frac{\cos\frac{\nu\pi}{2}}{2^{\nu}}\right)^3 \left[\frac{\Gamma(\frac{1+\nu}{2})}{\Gamma(\frac{2+\nu}{2})}\right]^4.$$
(5.33)

Proof. For $v \in (-1,0)$, one may convert (5.18) into

$$\widetilde{T}_{\nu} = -\frac{\sin(\pi\nu)\cos\frac{\nu\pi}{2}}{2^{\nu}\pi \left[\Gamma\left(\frac{2+\nu}{2}\right)\right]^{2}\Gamma(-\nu)}\lim_{\varepsilon \to 0^{+}} \int_{\varepsilon-i\infty}^{\varepsilon+i\infty} \widetilde{g}_{\nu}(s)\frac{\mathrm{d}\,s}{2\pi i}$$

$$= -\frac{\sin(\pi\nu)\cos\frac{\nu\pi}{2}}{2^{\nu}\pi \left[\Gamma\left(\frac{2+\nu}{2}\right)\right]^{2}\Gamma(-\nu)}G_{4,4}^{2,4}\left(1\begin{vmatrix}0,\frac{1-\nu}{2},\frac{1-\nu}{2},\nu+1\\0,0,0,0\end{vmatrix}\right)$$

$$= -\frac{\sin(\pi\nu)\cos\frac{\nu\pi}{2}}{2^{\nu}\pi \left[\Gamma\left(\frac{2+\nu}{2}\right)\right]^{2}\Gamma(-\nu)}G_{3,3}^{2,3}\left(1\begin{vmatrix}\frac{1-\nu}{2},\frac{1-\nu}{2},\nu+1\\0,0,0\end{vmatrix}\right),$$
(5.34)

where

$$\widetilde{g}_{\nu}(s) := \frac{\Gamma(s)^{2} \Gamma(1-s) \Gamma(-s-\nu) \left[\Gamma(\frac{\nu+1}{2}-s) \right]^{2}}{[\Gamma(1-s)]^{2}} = \frac{\Gamma(s)^{2} \Gamma(-s-\nu) \left[\Gamma(\frac{\nu+1}{2}-s) \right]^{2}}{\Gamma(1-s)}.$$
(5.35)

Applying Bailey's reduction (5.20) to the second line in (5.34), we receive two (ostensibly) different outputs. One of them is given in (5.32), where the ${}_{5}F_{4}$ series converges for Re $\nu > -1$; the other is

$$-\frac{4\sin(\nu\pi)\cos\frac{\nu\pi}{2}}{\pi^{3/2}}\frac{\left[\Gamma(\frac{\nu+1}{2})\right]^{3}}{\Gamma(\frac{2+\nu}{2})\Gamma(\nu+2)}{}_{5}F_{4}\left(\begin{array}{c}1+\nu,\frac{3+\nu}{2},\frac{1+\nu}{2},1+\nu,\frac{1+\nu}{2}\\\frac{1+\nu}{2},\frac{3+\nu}{2},1,\frac{3+\nu}{2}\end{array}\right)1\right),\tag{5.32'}$$

where the ${}_{5}F_{4}$ series converges for Re $\nu < 0$. In both (5.32) and (5.32'), the well-poised ${}_{5}F_{4}$ series are reducible by (5.31). This proves (5.33).

5.3. Generalizations to associated Legendre functions.

Lemma 5.6 (Clebsch–Gordan coupling of associated Legendre functions). When $2\varepsilon \notin \mathbb{Z}_{<0}$ and $\operatorname{Re}(\beta + \varepsilon) > \frac{|\operatorname{Re}\delta|}{2} - 1$, we have

$$\int_{0}^{1} (1-x^{2})^{\beta} \left[P_{\mu}^{\delta}(x) + P_{\mu}^{\delta}(-x) \right] \left[P_{\nu}^{-\varepsilon}(x) \right]^{2} dx$$

$$= \frac{2^{\delta-2\varepsilon}\pi\Gamma(1+\beta-\frac{\delta}{2}+\varepsilon)\Gamma(1+\beta+\frac{\delta}{2}+\varepsilon)}{[\Gamma(1+\varepsilon)]^{2}\Gamma(\frac{1-\delta-\mu}{2})\Gamma(\frac{2-\delta+\mu}{2})\Gamma(\beta+\varepsilon-\frac{\mu-2}{2})\Gamma(\beta+\varepsilon+\frac{\mu+3}{2})} \times {}_{5}F_{4} \left(\frac{\frac{1}{2}+\varepsilon,1+\beta-\frac{\delta}{2}+\varepsilon,1+\beta+\frac{\delta}{2}+\varepsilon,\varepsilon-\nu,1+\varepsilon+\nu}{1+\varepsilon,1+2\varepsilon,\beta+\varepsilon-\frac{\mu-2}{2},\beta+\varepsilon+\frac{\mu+3}{2}} \right) \left[1 \right).$$
(5.36)

When $\varepsilon \notin \mathbb{Z}$ and $\operatorname{Re} \beta > \frac{|\operatorname{Re} \delta|}{2} - 1$, we have

$$\int_{0}^{1} (1-x^{2})^{\beta} \left[P_{\mu}^{\delta}(x) + P_{\mu}^{\delta}(-x) \right] P_{\nu}^{\varepsilon}(x) P_{\nu}^{-\varepsilon}(x) dx$$

$$= \frac{\sin(\varepsilon\pi)}{\varepsilon\pi} \frac{2^{\delta}\pi\Gamma(1+\beta-\frac{\delta}{2})\Gamma(1+\beta+\frac{\delta}{2})}{\Gamma(\frac{1-\delta-\mu}{2})\Gamma(\frac{2-\delta+\mu}{2})\Gamma(\beta-\frac{\mu-2}{2})\Gamma(\beta+\frac{\mu+3}{2})}{5} F_{4} \begin{pmatrix} \frac{1}{2},1+\beta-\frac{\delta}{2},1+\beta+\frac{\delta}{2},-\nu,1+\nu\\ 1-\varepsilon,1+\varepsilon,\beta-\frac{\mu-2}{2},\beta+\frac{\mu+3}{2} \end{pmatrix} \left| 1 \right\rangle.$$

$$(5.37)$$

Proof. Instead of (2.57), we need a more general formula (see [47, item 7.132.1], which corrects [10, \$18.1(16)])

$$\int_{0}^{1} \left(1 - x^{2}\right)^{\sigma} \left[P_{\mu}^{\delta}(x) + P_{\mu}^{\delta}(-x)\right] \mathrm{d}x$$

$$= \frac{2^{\delta} \pi \Gamma \left(1 + \sigma + \frac{\delta}{2}\right) \Gamma \left(1 + \sigma - \frac{\delta}{2}\right)}{\Gamma \left(\frac{\mu + 3}{2} + \sigma\right) \Gamma \left(\sigma - \frac{\mu - 2}{2}\right) \Gamma \left(\frac{2 + \mu - \delta}{2}\right) \Gamma \left(\frac{1 - \mu - \delta}{2}\right)}, \quad \operatorname{Re} \sigma > \frac{|\operatorname{Re} \delta|}{2} - 1.$$

$$(2.57')$$

Joining this integral identity with (3.17) and (3.19), while switching the order of summation and integration, we obtain (5.36) and (5.37).

Remark In (5.36), upon trigonometric substitution to the scenario where $\beta = \nu = -\frac{1}{2}$ and $\delta = \varepsilon = \mu = 0$, we have

$$\int_{0}^{\pi/2} \left[\mathbf{K} \left(\sin \frac{\theta}{2} \right) \right]^{2} \mathrm{d}\,\theta = \frac{\pi^{3}}{8} {}_{4}F_{3} \left(\begin{array}{c} \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ 1, 1, 1 \end{array} \right) \mathbf{1} \right).$$
(5.38)

One may compare this to

$$\int_{0}^{\pi/2} \left[\mathbf{K} \left(\sin \theta \right) \right]^{2} \mathrm{d}\,\theta = \frac{\pi^{3}}{4} {}_{4}F_{3} \begin{pmatrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ 1, 1, 1 \end{pmatrix},$$
(5.39)

a formula proved by Wan [78, (35)]. This specific ${}_{4}F_{3}$ series is related to a critical *L*-value (see [67, 32] or [90, §7]) of a weight-4 cusp form attached to the rigid Calabi–Yau threefold $\sum_{j=1}^{4} (x_{j} + \frac{1}{x_{j}}) = 0$ [104, §2].

Now, we are ready to complete the proof of Theorem 5.1 by putting a finishing touch to (5.7).

Proof of (5.7). With

$$a_{\mu}(\varepsilon) \coloneqq \frac{2^{2\varepsilon+1}[\sin(\mu\pi) - \sin(2\varepsilon\pi)]}{\pi^{3/2}(\mu - 2\varepsilon)(\mu + 2\varepsilon + 1)} \frac{\Gamma(\frac{1}{2} + \varepsilon)}{\Gamma(1 + \varepsilon)},$$
(5.40)

$$b_{\mu}(\varepsilon) \coloneqq \frac{2^{2\varepsilon}\pi}{\Gamma(\frac{2-\mu}{2})\Gamma(\frac{\mu+3}{2})\Gamma(\frac{1-\mu-2\varepsilon}{2})\Gamma(\frac{\mu+2-2\varepsilon}{2})},\tag{5.41}$$

we consider the following particular cases of (5.36) and (5.37):

$$\int_{0}^{1} \left[P_{\mu}^{2\varepsilon}(x) + P_{\mu}^{2\varepsilon}(-x) \right] \left[P_{\nu}^{-\varepsilon}(x) \right]^{2} \mathrm{d}x = a_{\mu}(\varepsilon)_{4} F_{3} \begin{pmatrix} 1, \frac{1}{2} + \varepsilon, -\nu + \varepsilon, \nu + 1 + \varepsilon \\ 1 + \varepsilon, \frac{2-\mu}{2} + \varepsilon, \frac{\mu+3}{2} + \varepsilon \\ 1 + \varepsilon, \frac{2-\mu}{2} + \varepsilon, \frac{\mu+3}{2} + \varepsilon \end{pmatrix} \left| 1 \right\rangle, \tag{5.36'}$$

$$\int_{0}^{1} \left[P_{\mu}^{2\varepsilon}(x) + P_{\mu}^{2\varepsilon}(-x) \right] P_{\nu}^{\varepsilon}(x) P_{\nu}^{-\varepsilon}(x) \,\mathrm{d}\, x = b_{\mu}(\varepsilon)_{3} F_{2} \left(\frac{\frac{1}{2}, -\nu, \nu + 1}{\frac{2-\mu}{2}, \frac{\mu+3}{2}} \right| 1 \right), \tag{5.37'}$$

whose right-hand sides involve two generalized hypergeometric series that showed up previously in (5.8c).

Simplifying

$$\lim_{\varepsilon \to 0} \frac{\partial^2}{\partial \varepsilon^2} \int_0^1 \left[P_{\mu}^{2\varepsilon}(x) + P_{\mu}^{2\varepsilon}(-x) \right] \left\{ \frac{\left[P_{\nu}^{-\varepsilon}(x) \right]^2}{a_{\mu}(\varepsilon)} \frac{c_{\mu,\nu}(\varepsilon)}{c_{\mu,\nu}(0)} - \frac{P_{\nu}^{\varepsilon}(x) P_{\nu}^{-\varepsilon}(x)}{b_{\mu}(\varepsilon)} \right\} \mathrm{d} x \tag{5.42}$$

with the assistance from the first-order derivative of any associated Legendre function with respect to its order (cf. [23, (3)])

$$\frac{\partial}{\partial \varepsilon}\Big|_{\varepsilon=0} P_{\nu}^{\varepsilon}(x) = \frac{\psi^{(0)}(\nu+1) + \psi^{(0)}(-\nu)}{2} P_{\nu}(x) - \frac{\pi P_{\nu}(-x)}{2\sin(\nu\pi)},\tag{5.43}$$

and a pair of digamma identities

$$\psi^{(0)}\left(\frac{\mu+3}{2}\right) = \frac{2}{\mu+1} + \frac{\pi}{\tan\frac{\mu\pi}{2}} - \psi^{(0)}\left(\frac{2-\mu}{2}\right) + 2\psi^{(0)}(\mu) - 2\log 2, \tag{5.44}$$

$$\psi^{(0)}\left(\frac{1-\mu}{2}\right) = \frac{2}{\mu} - \pi \tan\frac{\mu\pi}{2} - \psi^{(0)}\left(\frac{\mu+2}{2}\right) + 2\psi^{(0)}(\mu) - 2\log 2, \tag{5.45}$$

we get (5.7).

Acknowledgments. A considerable amount of the materials developed in this work were based on my unpublished notes prepared at Princeton University during 2011–2013, for Prof. Weinan E's working seminar on mathematical physics.

This paper is a write-up of my talk "Sun's series at vanishing and positive genera" presented during the 2nd Workshop on Supercongruences, π Series, and Related Fields, at Nanjing University on Nov. 17–19, 2023. I thank the organizers of this workshop for their hospitality and all the participants for their thoughtful feedback during and after this event.

References

- [1] J. Ablinger. Discovering and proving infinite binomial sums identities. Exp. Math., 26(1):62-71, 2017.
- [2] J. Ablinger. Discovering and proving infinite Pochhammer sum identities. Exp. Math., 31(1):309-323, 2022.
- [3] J. Ablinger, J. Blümlein, C. G. Raab, and C. Schneider. Iterated binomial sums and their associated iterated integrals. *J. Math. Phys.*, 55(11):112301, 57 pages, 2014.
- [4] J. Ablinger, J. Blümlein, and C. Schneider. Harmonic sums and polylogarithms generated by cyclotomic polynomials. *J. Math. Phys.*, 52(10):102301, 52 pages, 2011.
- [5] J. Ablinger, J. Blümlein, and C. Schneider. Analytic and algorithmic aspects of generalized harmonic sums and polylogarithms. *J. Math. Phys.*, 54(8):082301, 74 pages, 2013.

YAJUN ZHOU

- [6] G. E. Andrews, R. Askey, and R. Roy. Special Functions, volume 71 of Encyclopedia of Mathematics and Its Applications. Cambridge University Press, Cambridge, UK, 1999.
- [7] P. Appell. Mémoire sur les équations différentielles linéaires. Ann. sci. Éc. Norm. Supér., 10:391-424, 1881.
- [8] W. N. Bailey. Some transformations of generalized hypergeometric series, and contour integrals of Barnes's type. Quart. J. Math., 3:168–182, 1932.
- [9] H. Bateman. Higher Transcendental Functions, volume I. McGraw-Hill, New York, NY, 1953. (compiled by staff of the Bateman Manuscript Project: Arthur Erdélyi, Wilhelm Magnus, Fritz Oberhettinger, Francesco G. Tricomi, David Bertin, W. B. Fulks, A. R. Harvey, D. L. Thomsen, Jr., Maria A. Weber and E. L. Whitney).
- [10] H. Bateman. Table of Integral Transforms, volume II. McGraw-Hill, New York, NY, 1954. (compiled by staff of the Bateman Manuscript Project: Arthur Erdélyi, Wilhelm Magnus, Fritz Oberhettinger, Francesco G. Tricomi, David Bertin, W. B. Fulks, A. R. Harvey, D. L. Thomsen, Jr., Maria A. Weber and E. L. Whitney).
- [11] B. C. Berndt. Ramanujan's Notebooks (Part II). Springer-Verlag, New York, NY, 1989.
- [12] B. C. Berndt. Ramanujan's Notebooks (Part V). Springer-Verlag, New York, NY, 1998.
- [13] J. M. Borwein, D. J. Broadhurst, and J. Kamnitzer. Central binomial sums, multiple Clausen values, and zeta values. *Experiment. Math.*, 10(1):25–34, 2001.
- [14] J. M. Borwein, M. L. Glasser, R. C. McPhedran, J. G. Wan, and I. J. Zucker. Lattice Sums: Then and Now, volume 150 of Encyclopedia of Mathematics and Its Applications. Cambridge University Press, 2013.
- [15] J. M. Borwein, A. Straub, and J. Wan. Three-step and four-step random walk integrals. *Exp. Math.*, 22(1):1–14, 2013.
- [16] D. J. Broadhurst. On the enumeration of irreducible k-fold Euler sums and their roles in knot theory and field theory. arXiv:hep-th/9604128, 1996.
- [17] D. J. Broadhurst. Massive 3-loop Feynman diagrams reducible to SC* primitives of algebras of the sixth root of unity. *Eur. Phys. J. C Part. Fields*, 8(2):313–333, 1999.
- [18] F. Brown. The massless higher-loop two-point function. Comm. Math. Phys., 287(3):925-958, 2009. arXiv:0804.1660v1 [math.AG].
- [19] F. Brown. On the periods of some Feynman integrals. arXiv:0910.0114v2 [math.AG], 2010.
- [20] F. C. S. Brown. Multiple zeta values and periods of moduli spaces $\overline{\mathfrak{M}}_{0,n}$. Ann. Sci. Éc. Norm. Supér. (4), 42(3):371–489, 2009. arXiv:math/0606419v1 [math.AG].
- [21] J. H. Bruinier, S. Ehlen, and T. Yang. CM values of higher automorphic Green functions for orthogonal groups. *Invent. Math.*, 225(3):693-785, 2021. arXiv:1912.12084 [math.NT].
- [22] J. H. Bruinier, Y. Li, and T. Yang. Deformations of theta integrals and a conjecture of Gross-Zagier. arXiv:2204.10604v2 [math.NT], 2022.
- [23] Y. A. Brychkov. On the derivatives of the Legendre functions $P_{\nu}^{\mu}(z)$ and $Q_{\nu}^{\mu}(z)$ with respect to μ and ν . Integral Transforms Spec. Funct., 21(3–4):175–181, 2010.
- [24] J. M. Campbell. Applications of Caputo operators in the evaluation of Clebsch-Gordan-type multiple elliptic integrals. *Integral Transforms Spec. Funct.*, 34(5):371–383, 2023.
- [25] J. M. Campbell. New Clebsch-Gordan-type integrals involving threefold products of complete elliptic integrals. Math. Methods Appl. Sci., 46(17):17774-17787, 2023. arXiv:2302.05819v1 [math.CA].
- [26] J. M. Campbell. On a conjecture on a series of convergence rate $\frac{1}{2}$. arXiv:2304.00360 [math.NT], 2023.
- [27] J. M. Campbell. On a higher-order version of a formula due to Ramanujan. Integral Transforms Spec. Funct., 35(4):260-269, 2024.
- [28] J. M. Campbell, M. Cantarini, and J. D'Aurizio. Symbolic computations via Fourier-Legendre expansions and fractional operators. *Integral Transforms Spec. Funct.*, 33(2):157–175, 2022.
- [29] M. Cantarini. A note on Clebsch-Gordan integral, Fourier-Legendre expansions and closed form for hypergeometric series. *Ramanujan J.*, 59(2):549–557, 2022.
- [30] M. Cantarini and J. D'Aurizio. On the interplay between hypergeometric series, Fourier-Legendre expansions and Euler sums. *Boll. Unione Mat. Ital.*, 12(4):623-656, 2019. arXiv:1806.08411v1 [math.NT].
- [31] E. Catalan. Mémoire sur la transformation des séries et sur quelques intégrales définies. In Mémoires couronnés et mémoires des savants étrangers, volume 33. l'Académie royale des Sciences, des Lettres et des Beaux-Arts de Belgique, Bruxelles, Belgium, 1865-1867. (50 pages).
- [32] E. Catalan. Recherches sur la constante G, et sur les intégrales Eulériennes. In Mémoires de l'Académie Impériale des sciences de St.-Pétersbourg, volume 31. Commissionaires de l'Académie Impériale des sciences, St. Petersburg, Russia, 1883. (51 pages).
- [33] H. H. Chan, J. Wan, and W. Zudilin. Legendre polynomials and Ramanujan-type series for $1/\pi$. Israel J. Math., 194(1):183–207, 2013.
- [34] S. Chowla and A. Selberg. On Epstein's zeta function (I). Proc. Natl. Acad. Sci. USA, 35:371–374, 1949.
- [35] W. Chu. Infinite series involving harmonic numbers of convergence rate 1/2. Integral Transforms Spec. Funct., 33(7):570–590, 2022.

- [36] D. V. Chudnovsky and G. V. Chudnovsky. Approximations and complex multiplication according to Ramanujan. In *Ramanujan revisited (Urbana-Champaign, Ill., 1987)*, pages 375–472. Academic Press, Boston, MA, 1988.
- [37] T. Clausen. Ueber die Fälle, wenn die Reihe von der Form $y = 1 + \frac{\alpha}{1} \cdot \frac{\beta}{\gamma} x + \frac{\alpha.\alpha+1}{1.2} \cdot \frac{\beta.\beta+1}{\gamma.\gamma+1} x^2 + \text{etc. ein Quadrat von der Form } z = 1 + \frac{\alpha'}{1} \cdot \frac{\beta'}{\gamma'} \cdot \frac{\delta'}{\varepsilon'} x + \frac{\alpha'.\alpha'+1}{1.2} \cdot \frac{\beta'.\beta'+1}{\gamma'.\gamma'+1} \cdot \frac{\delta'.\delta'+1}{\varepsilon'.\varepsilon'+1} x^2 + \text{etc. hat. } J. Reine Angew. Math., 3:89–91, 1828.$
- [38] S. Cooper. Ramanujan's theta functions. Springer, Cham, Switzerland, 2017.
- [39] A. I. Davydychev and M. Y. Kalmykov. New results for the ε-expansion of certain one-, two- and three-loop Feynman diagrams. *Nuclear Phys. B*, 605(1-3):266-318, 2001.
- [40] A. I. Davydychev and M. Y. Kalmykov. Massive Feynman diagrams and inverse binomial sums. Nuclear Phys. B, 699(1-2):3-64, 2004.
- [41] D. Disegni. The universal p-adic Gross-Zagier formula. Invent. Math., 230(2):509-649, 2022.
- [42] R. Evans and J. Greene. Clausen's theorem and hypergeometric functions over finite fields. *Finite Fields Appl.*, 15(1):97–109, 2009.
- [43] J. Fuselier, L. Long, R. Ramakrishna, H. Swisher, and F.-T. Tu. Hypergeometric functions over finite fields. Mem. Amer. Math. Soc., 280(1382):vii+124 pages, 2022.
- [44] M. L. Glasser and I. J. Zucker. Lattice sums. In *Theoretical Chemistry: Advances and Perspectives*, pages 67–139. Academic Press, New York, NY, 1980.
- [45] A. B. Goncharov. The double logarithm and Manin's complex for modular curves. Math. Res. Lett., 4(5):617–636, 1997.
- [46] A. B. Goncharov. Multiple polylogarithms, cyclotomy and modular complexes. Math. Res. Lett., 5(4):497–516, 1998.
- [47] I. S. Gradshteyn and I. M. Ryzhik. *Table of Integrals, Series, and Products*. Academic Press, Burlington, MA, 7th edition, 2007. (Translated from Russian by Scripta Technica, Inc., edited by Alan Jeffrey and Daniel Zwillinger).
- [48] J. Greene and D. Stanton. A character sum evaluation and Gaussian hypergeometric series. J. Number Theory, 23(1):136–148, 1986.
- [49] B. Gross, W. Kohnen, and D. Zagier. Heegner points and derivatives of L-series. II. Math. Ann., 278:497–562, 1987.
- [50] B. H. Gross and D. B. Zagier. On singular moduli. J. Reine Angew. Math., 355:191-220, 1985.
- [51] B. H. Gross and D. B. Zagier. Heegner points and derivatives of L-series. Invent. Math., 84(2):225-320, 1986.
- [52] J. Guillera. Hypergeometric identities for 10 extended Ramanujan-type series. Ramanujan J., 15(2):219–234, 2008.
- [53] J. Guillera. WZ-proofs of "divergent" Ramanujan-type series. In Advances in combinatorics, pages 187–195. Springer, Heidelberg, 2013.
- [54] J. Guillera and M. Rogers. Ramanujan series upside-down. J. Aust. Math. Soc., 97(1):78-106, 2014.
- [55] K. Hessami Pilehrood and T. Hessami Pilehrood. Bivariate identities for values of the Hurwitz zeta function and supercongruences. *Electron. J. Combin.*, 18(2):Paper 35, 30 pages, 2011.
- [56] E. L. Ince. Ordinary differential equations. Dover Publications, New York, NY, 1956.
- [57] H. Iwaniec. Topics in Classical Automorphic Forms, volume 17 of Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 1997.
- [58] M. Y. Kalmykov and O. Veretin. Single-scale diagrams and multiple binomial sums. Phys. Lett. B, 483(1-3):315– 323, 2000.
- [59] M. Y. Kalmykov, B. F. L. Ward, and S. A. Yost. Multiple (inverse) binomial sums of arbitrary weight and depth and the all-order ε-expansion of generalized hypergeometric functions with one half-integer value of parameter. *J. High Energy Phys.*, 2007(10):048, 2007.
- [60] C. Koutschan. Creative telescoping for holonomic functions. In Computer algebra in quantum field theory, Texts Monogr. Symbol. Comput., pages 171–194. Springer, Vienna, Austria, 2013.
- [61] S. Laporta. Analytical expressions of three- and four-loop sunrise Feynman integrals and four-dimensional lattice integrals. *Internat. J. Modern Phys. A*, 23(31):5007–5020, 2008. arXiv:0803.1007v4 [hep-ph].
- [62] Y. Li. Average CM-values of higher Green's function and factorization. Amer. J. Math., 144(5):1241–1298, 2022. arXiv:1812.08523v2 [math.NT].
- [63] Y. Li. Algebraicity of higher Green functions at a CM point. Invent. Math., 234(1):375-418, 2023. arXiv:2106.13653v2 [math.NT].
- [64] E. McSpirit and K. Ono. Hypergeometry and the AGM over finite fields. Contemp. Math., 2025+. (to appear) arXiv:2302.10387v2 [math.NT].
- [65] A. Mellit. Higher Green's Functions for Modular Forms. PhD thesis, Rheinische Friedrich-Wilhelms-Universität Bonn, June 2008. arXiv:0804.3184v1 [math.NT].
- [66] R. Preston. Profiles: The mountains of pi. The New Yorker, pages 36-67, March 2, 1992.

YAJUN ZHOU

- [67] M. Rogers, J. G. Wan, and I. J. Zucker. Moments of elliptic integrals and critical L-values. Ramanujan J., 37(1):113-130, 2015. arXiv:1303.2259v2 [math.NT].
- [68] A. Selberg and S. Chowla. On Epstein's zeta-function. J. Reine Angew. Math., 227:86–110, 1967.
- [69] L. J. Slater. Generalized Hypergeometric Functions. Cambridge University Press, Cambridge, UK, 1966.
- [70] Z.-W. Sun. New series involving binomial coefficients. arXiv:2307.03086v6 [math.NT], 2023.
- [71] Z.-W. Sun. Series with summands involving harmonic numbers. In M. B. Nathanson, editor, Combinatorial and Additive Number Theory VI, volume 464 of Springer Proc. Math. Stat. Springer, Cham, Switzerland, 2025. arXiv:2210.07238v9 [math.NT].
- [72] Z.-W. Sun and Y. Zhou. Evaluations of $\sum_{k=1}^{\infty} \frac{x^k}{k^2 \binom{3k}{k}}$ and related series. arXiv:2401.12083v1 [math.CO], 2024.
- [73] F.-T. Tu and Y. Yang. Evaluation of certain hypergeometric functions over finite fields. SIGMA Symmetry Integrability Geom. Methods Appl., 14:Paper No. 050, 18 pp., 2018.
- [74] M. Viazovska. Modular Functions and Special Cycles. PhD thesis, Rheinische Friedrich-Wilhelms-Universität Bonn, November 2013.
- [75] M. Viazovska. CM values of higher Green's functions and regularized Petersson products. In Arithmetic and geometry, volume 420 of London Math. Soc. Lecture Note Ser., pages 493–503. Cambridge Univ. Press, Cambridge, UK, 2015.
- [76] M. Viazovska. Petersson inner products of weight one modular forms. J. Reine Angew. Math., 749:133–159, 2019. arXiv:1211.4715v2 [math.NT].
- [77] M. Viazovska. CM values of higher Green's functions. In J. Fresán, D. Jetchev, P. Jossen, and R. Pink, editors, Motives and Complex Multiplication, Progress in Mathematics, Boston, MA, 2023+. Birkhäuser. arXiv:1110.4654v1 [math.NT].
- [78] J. G. Wan. Moments of products of elliptic integrals. Adv. Appl. Math., 48:121-141, 2012. arXiv:1101.1132v1 [math.CA].
- [79] G. N. Watson. Three triple integrals. Quart. J. Math., 10:266-276, 1939.
- [80] C. Wei. On some conjectural series containing binomial coefficients and harmonic numbers. arXiv:2306.02641v1 [math.CO], 2023.
- [81] C. Wei. On some conjectures of Z.-W. Sun involving harmonic numbers. arXiv: 2304.09753v1 [math.CO], 2023.
- [82] T. Wei and X. Guo. Solvable lattice sums and quadratic Dirichlet L-values. Acta Arith., 203(3):227–238, 2022.
- [83] S. Weinzierl. Expansion around half integer values, binomial sums and inverse binomial sums. J. Math. Phys., 45:2656-2673, 2004.
- [84] H. S. Wilf and D. Zeilberger. Rational functions certify combinatorial identities. J. Amer. Math. Soc., 3(1):147–158, 1990.
- [85] K. S. Williams. Some Lambert series expansions of products of theta functions. Ramanujan J., 3:367-384, 1999.
- [86] D. Zagier. Hyperbolic manifolds and special values of Dedekind zeta-functions. Invent. Math., 83(2):285–301, 1986.
- [87] D. Zagier. A modular identity arising from mirror symmetry. In M.-H. Saito, Y. Shimizu, and K. Ueno, editors, *Integrable Systems and Algebraic Geometry (Proceedings of the Taniguchi Symposium 1997)*, pages 477–480. World Scientific, Singapore, 1998.
- [88] D. Zagier. The dilogarithm function. In P. Cartier, B. Julia, P. Moussa, and P. Vanhove, editors, Frontiers in Number Theory, Physics, and Geometry II: On Conformal Field Theories, Discrete Groups and Renormalization, pages 3–65. Springer, Berlin, Germany, 2007.
- [89] D. Zagier. Elliptic modular forms and their applications. In K. Ranestad, editor, The 1-2-3 of Modular Forms, Lectures at a Summer School in Nordfjordeid, Norway, pages 1–103. Springer-Verlag, Berlin, Germany, 2008.
- [90] D. Zagier. The arithmetic and topology of differential equations. In *European Congress of Mathematics*, pages 717–776. Eur. Math. Soc., Zürich, 2018.
- [91] D. Zeilberger. A fast algorithm for proving terminating hypergeometric identities. *Discrete Math.*, 80(2):207–211, 1990.
- [92] D. Zeilberger. The method of creative telescoping. J. Symbolic Comput., 11(3):195–204, 1991.
- [93] D. Zeilberger. Closed form (pun intended!). In A tribute to Emil Grosswald: number theory and related analysis, volume 143 of Contemp. Math., pages 579-607. Amer. Math. Soc., Providence, RI, 1993.
- [94] S. Zhang. Heights of Heegner cycles and derivatives of L-series. Invent. Math., 130:99–152, 1997.
- [95] Y. Zhou. Legendre functions, spherical rotations, and multiple elliptic integrals. *Ramanujan J.*, 34(3):373–428, 2014. arXiv:1301.1735v4 [math.CA].
- [96] Y. Zhou. On some integrals over the product of three Legendre functions. *Ramanujan J.*, 35:311–326, 2014. arXiv:1304.1606v2 [math.CA].
- [97] Y. Zhou. Kontsevich–Zagier integrals for automorphic Green's functions. I. Ramanujan J., 38(2):227–329, 2015. arXiv:1312.6352v4 [math.CA].

- [98] Y. Zhou. Ramanujan series for Epstein zeta functions. Ramanujan J., 40(2):367-388, 2016. arXiv:1410.8312v2 [math.CA].
- [99] Y. Zhou. Kontsevich–Zagier integrals for automorphic Green's functions. II. Ramanujan J., 42(3):623–688, 2017.
 [see Ramanujan J. 49(1):231–235, 2019 for erratum/addendum] arXiv:1506.00318v3 [math.NT].
- [100] Y. Zhou. Sun's series via cyclotomic multiple zeta values. SIGMA Symmetry Integrability Geom. Methods Appl., 19:Paper No. 074, 20 pp., 2023. arXiv: 2306.04638v3 [math.NT].
- [101] Y. Zhou. Hyper-Mahler measures via Goncharov-Deligne cyclotomy. *Contemp. Math.*, 2025+. (to appear) arXiv:2210.17243v4 [math.NT].
- [102] I. J. Zucker and M. M. Robertson. Further aspects of the evaluation of $\sum_{(m.n\neq 0,0)} (am^2 + bnm + cn^2)^{-s}$. Math. Proc. Cambridge Philos. Soc., 95(1):5–13, 1984.
- [103] W. Zudilin. Arithmetic of linear forms involving odd zeta values. J. Théor. Nombres Bordeaux, 16(1):251-291, 2004.
- [104] W. Zudilin. A hypergeometric version of the modularity of rigid Calabi-Yau manifolds. SIGMA Symmetry Integrability Geom. Methods Appl., 14:Paper No. 086, 16 pp., 2018.

PROGRAM IN APPLIED AND COMPUTATIONAL MATHEMATICS (PACM), PRINCETON UNIVERSITY, PRINCETON, NJ 08544

Email address: yajunz@math.princeton.edu

Current address: ACADEMY OF ADVANCED INTERDISCIPLINARY STUDIES (AAIS), PEKING UNIVERSITY, BEIJING 100871, P. R. CHINA

Email address: yajun.zhou.1982@pku.edu.cn