ISOMETRIC CLASSIFICATION OF THE L^p-SPACES OF INFINITE DIMENSIONAL LEBESGUE MEASURE

DANIEL L. RODRÍGUEZ-VIDANES AND JUAN CARLOS SAMPEDRO

ABSTRACT. In this paper, we investigate the isometric structure of the L^p -spaces associated with an infinite-dimensional analogue of the Lebesgue measure $(\mathbb{R}^{\mathbb{N}}, \mu)$. Our primary result establishes that under the Continuum Hypothesis (CH), $L^p(\mu)$ is isometrically isomorphic to $\ell^p(\mathfrak{c}, L^p[0, 1])$, where \mathfrak{c} denotes the cardinality of the continuum. In the general case, without assuming CH, we prove that $L^p(\mu)$ contains an isometric and complemented copy of $\ell^p(\mathfrak{c}, L^p[0, 1])$. Furthermore, we characterize isometries between L^p -spaces and establish necessary and sufficient conditions for an isometric isomorphism of the form $L^p(\nu) \cong \ell^p(\kappa, L^p[0, 1])$ for some cardinal κ . In particular, we classify all possible isometric isomorphisms between such spaces.

CONTENTS

1. Introduction	1
2. Isometry under the Continuum Hypothesis	3
3. Study of the general case	7
3.1. Study of G^p	8
3.2. Study of B^p	10
4. Characterization of isometries	12
References	17
Appendix A. Some properties of the measure μ	18

1. INTRODUCTION

The isometric classification of L^p -spaces has been a central topic of research throughout the 20th and 21st centuries (see, for instance, [5, 6, 8, 12–18, 20–22, 24–26, 29], among many others). A fundamental consequence of Maharam's theorem (see, for example, [8, Th. 14, p. 130], [14, Th. 332B], or the original work [21]) states that, for any measure space (X, Σ, ν) , there exist two sets Γ, I and two families of cardinals $\tau_i, \kappa_i, i \in I$, such that the following isometric isomorphism holds:

(1.1)
$$L^{p}(\nu) \cong \ell^{p}(\Gamma) \oplus_{p} \left(\bigoplus_{i \in I} \ell^{p}(\tau_{i}, L^{p}([0, 1]^{\kappa_{i}})) \right)_{p}.$$

²⁰¹⁰ Mathematics Subject Classification. 46B04 (primary), 28C20, 46G12 (secondary).

Key words and phrases. L^p-spaces, Continuum Hypothesis, Lebesgue measure, Isometric Classification. The second author has been supported by the Research Grant PID2021–123343NB-I00 of the Spanish Ministry of Science and Innovation and by the Institute of Interdisciplinar Mathematics of Complutense University.

Throughout this paper, given a family $(X_j)_{j \in J}$ of Banach spaces, we define the space

$$\left(\bigoplus_{j\in J} X_j\right)_p := \left\{ (x_j)_{j\in J} \in \bigotimes_{j\in J} X_j : \sum_{j\in J} \|x_j\|_{X_j}^p < \infty \right\},\$$

with norm

$$||(x_j)_{j\in J}||_p := \left(\sum_{j\in J} ||x_j||_{X_j}^p\right)^{\frac{1}{p}}.$$

In the case where $X_j = X$ for every $j \in J$, we denote $\ell^p(J, X) := \left(\bigoplus_{j \in J} X\right)_p$.

In this paper, we study the isometric structure of the L^p -spaces associated with an infinitedimensional analogue of the Lebesgue measure. To define this measure, we consider the constant sequence of measure spaces $\{(\mathbb{R}, \mathcal{B}, \lambda)\}_{i \in \mathbb{N}}$, where \mathcal{B} is the Borel σ -algebra of \mathbb{R} and λ is the Lebesgue measure on \mathbb{R} , and we construct the product measure space

$$\left(\bigotimes_{i \in \mathbb{N}} \mathbb{R}, \bigotimes_{i \in \mathbb{N}} \mathcal{B}, \bigotimes_{i \in \mathbb{N}} \lambda \right),\$$

as described in [28]. For notational simplicity, we denote $\mathbb{R}^{\mathbb{N}} := \bigotimes_{i \in \mathbb{N}} \mathbb{R}$, $\mathcal{B}_{\infty} := \bigotimes_{i \in \mathbb{N}} \mathcal{B}$, and $\mu := \bigotimes_{i \in \mathbb{N}} \lambda$.

For completeness, we recall the definition of the measure space $(\mathbb{R}^{\mathbb{N}}, \mathcal{B}_{\infty}, \mu)$. The σ -algebra \mathcal{B}_{∞} is generated by the *cylinder sets*

$$\left\{ \sum_{i=1}^{m} C_i \times \sum_{i=m+1}^{\infty} \mathbb{R} : C_i \in \mathcal{B}, \ \forall i \in \{1, 2, ..., m\}, \ m \in \mathbb{N} \right\}.$$

It is worth noting that \mathcal{B}_{∞} coincides with the Borel σ -algebra of $\mathbb{R}^{\mathbb{N}}$ when endowed with the product topology. By $\mathcal{F}(\mathcal{B}, \lambda)$ we denote the set of *finite rectangles* on $\mathbb{R}^{\mathbb{N}}$ defined by

$$\mathcal{F}(\mathcal{B},\lambda) := \left\{ \bigotimes_{i \in \mathbb{N}} C_i : C_i \in \mathcal{B}, \ \forall i \in \mathbb{N}, \ \prod_{i \in \mathbb{N}} \lambda(C_i) \in [0,\infty) \right\},\$$

and **vol** for the *volume* map **vol** : $\mathcal{F}(\mathcal{B}, \lambda) \to [0, \infty)$ given by

$$\operatorname{vol}\left(\bigotimes_{i\in\mathbb{N}}C_i\right):=\prod_{i\in\mathbb{N}}\lambda(C_i).$$

We denote by $\mathcal{P}(\mathbb{R}^{\mathbb{N}})$ the set of all subsets of $\mathbb{R}^{\mathbb{N}}$. The measure μ is defined as the restriction to \mathcal{B}_{∞} of the outer measure $\mu^* : \mathcal{P}(\mathbb{R}^{\mathbb{N}}) \to [0, \infty]$ defined, for every $A \in \mathcal{P}(\mathbb{R}^{\mathbb{N}})$, by

$$\mu^*(A) := \inf \left\{ \sum_{n \in \mathbb{N}} \operatorname{vol}(\mathscr{C}_n) : \{\mathscr{C}_n\}_{n \in \mathbb{N}} \subset \mathcal{F}(\mathcal{B}, \lambda) \text{ and } A \subset \bigcup_{n \in \mathbb{N}} \mathscr{C}_n \right\},$$

where we set $\inf \emptyset = \infty$.

The study of translation-invariant measures on $\mathbb{R}^{\mathbb{N}}$ has been a subject of interest in the literature. In [1], Baker constructed for the first time a translation-invariant Borel measure μ_B on $(\mathbb{R}^{\mathbb{N}}, \mathcal{B}_{\infty})$ that behaves as an infinite-dimensional analogue of the Lebesgue measure, using rectangles of the form $X_{i \in \mathbb{N}}(a_i, b_i)$. Later, in [2], Baker introduced the measure μ as described above. (In [2], Baker considered the completion of μ to obtain an infinite-dimensional analogue of the Fubini–Tonelli theorem.) In that work, Baker raised the question of whether the measure spaces corresponding to μ_B and μ coincide. However, in [23], Pantsulaia showed

in Remark 2.2 that while μ_B is absolutely continuous with respect to μ , it is not equivalent to it. In fact, a simple observation shows that $\mu(\{0,2\}^{\mathbb{N}}) = 0$ while $\mu_B(\{0,2\}^{\mathbb{N}}) = \infty$.

It is well known that infinite-dimensional locally convex topological vector spaces do not admit a nontrivial translation-invariant σ -finite Borel measure. Consequently, μ is not σ finite. Moreover, it is known that μ_B is not localizable in the sense of Fremlin [13, Def. 211G], since it is not semi-finite (see [23, Remark 5.2]). In Appendix A, we prove that μ is also not localizable.

For the measure space $(\mathbb{R}^{\mathbb{N}}, \mu)$, determining the family of cardinals involved in the isometric isomorphism (1.1) remains an open problem. The first main result of this paper solves this problem under the validity of the continuum hypothesis (CH).

Theorem 1.1. Let $1 \leq p < \infty$. Then, under CH, the following isometric isomorphism holds:

(1.2)
$$L^p(\mu) \cong \ell^p(\mathfrak{c}, L^p[0, 1]).$$

In the general case in which CH is not guaranteed, we prove that, at least $L^{p}(\mu)$ admits an isometric complemented copy of $\ell^{p}(\mathfrak{c}, L^{p}[0, 1])$ via a cube decomposition of $\mathbb{R}^{\mathbb{N}}$.

Throughout this article we will use standard notation and notions from Set Theory (see, for instance, [3, 19]), unless stated otherwise.

This paper is organized as follows. In Section 2, we prove Theorem 1.1 by carefully choosing an appropriate partition of $\mathbb{R}^{\mathbb{N}}$ whose existence strongly depends on CH. Section 3 is devoted to study the general case, that is, when CH is not guaranteed. We prove that, in the general case, $L^{p}(\mu)$ contains an isometric and complemented copy of $\ell^{p}(\mathfrak{c}, L^{p}[0, 1])$. This analysis is performed via a cube decomposition of $\mathbb{R}^{\mathbb{N}}$ that splits $L^{p}(\mu)$ into two complemented subspaces G^{p} and B^{p} . We prove that the space G^{p} is isometrically isomorphic to $\ell^{p}(\mathfrak{c}, L^{p}[0, 1])$ while B^{p} contains an isometric and complemented copy of $\ell^{p}(\mathfrak{c}, L^{p}[0, 1])$. The final Section 4 characterizes when $L^{p}(\nu)$ for some measure ν , is isometrically isomorphic to $\ell^{p}(\kappa, L^{p}[0, 1])$ for some cardinal number κ . Moreover, in the case in which these spaces are isometrically isomorphic, we describe all the possible isometric isomorphisms between these spaces. Finally, we include Appendix A in which we prove some properties and curiosities of the measure μ not present in the literature.

2. Isometry under the Continuum Hypothesis

In this section we prove that, under the Continuum Hypothesis, CH, $L^{p}(\mu)$ is isometrically isomorphic to $\ell^{p}(\mathfrak{c}, L^{p}[0, 1])$. Therefore, throughout this section we work under the validity of the CH.

Under this assumption, we have that $|\mathcal{F}(\mathcal{B},\mu)| = \mathfrak{c}$, and therefore, we can consider an enumeration $\{R_{\xi} : \xi < \omega_1\}$ of $\mathcal{F}(\mathcal{B},\lambda)$. For each $\xi < \omega_1$, we denote

(2.1)
$$B_{\xi} := R_{\xi} \setminus \bigcup_{\zeta < \xi} R_{\zeta}.$$

Note that each B_{ξ} is Borel as $\bigcup_{\zeta < \xi} R_{\zeta}$ is a countable union and $\mu(B_{\xi}) < \infty$ as $B_{\xi} \subset R_{\xi}$. Moreover, for distinct $\xi_1, \xi_2 < \omega_1, B_{\xi_1} \cap B_{\xi_2} = \emptyset$. Therefore,

$$\mathbb{R}^{\mathbb{N}} = \biguplus_{\xi < \omega_1} B_{\xi}.$$

Throughout this article, the notation \uplus stands for the disjoint union.

The first result of this section provides a partition of $\mathbb{R}^{\mathbb{N}}$ into finite rectangles that satisfies a property that will be crucial for proving the main result of this section.

Proposition 2.1. There exists a partition $\mathcal{F} = \{C_{\xi} : \xi < \omega_1\} \subset \mathcal{F}(\mathcal{B}, \mu)$ of $\mathbb{R}^{\mathbb{N}}$ into continuum many finite rectangles such that for every Borel $B \in \mathcal{B}_{\infty}$ of finite measure the set $\mathcal{F}(B) := \{\xi < \omega_1 : B \cap C_{\xi} \neq \emptyset\}$ is countable.

Proof. Take $B \in \mathcal{B}_{\infty}$ with finite measure. Then, by the definition of the measure μ , there exists a countable subset $\{R_{\xi_n}\}_{n < \omega} \subset \mathcal{F}(\mathcal{B}, \lambda)$ such that

$$B \subset \bigcup_{n < \omega} R_{\xi_n} = \bigcup_{n < \omega} \bigcup_{\eta \le \xi_n} B_{\eta}$$

Consequently, $\{\xi < \omega_1 : B \cap B_{\xi} \neq \emptyset\}$ is countable. Now, by (2.1), we deduce that

$$B_{\xi} = R_{\xi} \cap \bigcap_{\zeta < \xi} R_{\zeta}^{c} = R_{\xi} \cap \bigcap_{\zeta < \xi} \biguplus_{n \in \mathbb{N}} F_{\zeta, n};$$

for some

$$F_{\zeta,n} \in \mathcal{R}(\mathcal{B}) := \left\{ \bigotimes_{i \in \mathbb{N}} C_i : C_i \in \mathcal{B}, \ \forall i \in \mathbb{N} \right\}.$$

For the last equality, it is convenient to recall the following set theoretical identity: if $\mathscr{C} = \underset{i \in \mathbb{N}}{\underset{i \in \mathbb{N}}{C_i \in \mathcal{R}(\mathcal{B})}}$, then

$$\mathscr{C}^{c} = \biguplus_{n \in \mathbb{N}} \left(\bigotimes_{i=1}^{n-1} C_{i} \times C_{n}^{c} \times \bigotimes_{i=n+1}^{\infty} \mathbb{R} \right).$$

Therefore, we deduce that

$$B_{\xi} = \biguplus_{n \in \mathbb{N}} C_{\xi, n},$$

where

$$C_{\xi,n} := R_{\xi} \cap \bigcap_{\zeta < \xi} F_{\zeta,n} \in \mathcal{F}(\mathcal{B}, \lambda).$$

Let $\mathcal{F} = \{C_{\xi} : \xi < \omega_1\} \subset \mathcal{F}(\mathcal{B}, \mu)$ be a enumeration of $\{C_{\xi,n} : \xi < \omega_1, n \in \mathbb{N}\}$. Then, clearly \mathcal{F} is a partition of $\mathbb{R}^{\mathbb{N}}$ and since $\{\xi < \omega_1 : B \cap B_{\xi} \neq \emptyset\}$ is countable for every Borel $B \in \mathcal{B}_{\infty}$ with finite measure, necessarily $\mathcal{F}(B) = \{\xi < \omega_1 : B \cap C_{\xi} \neq \emptyset\}$ is also countable. The proof is concluded.

The next result is the key to prove the required isometric isomorphism.

Proposition 2.2. Let $1 \le p < \infty$. Then, for every $f \in L^p(\mu)$, it holds that

$$\int_{\mathbb{R}^N} |f|^p \ d\mu = \sum_{\xi < \omega_1} \int_{C_{\xi}} |f|^p \ d\mu.$$

Proof. Let $f \in L^p(\mu)$. Then $\operatorname{supp}(f)$ is σ -finite. Indeed,

$$\operatorname{supp}(f) = \bigcup_{n \in \mathbb{N}} [|f| > 1/n]$$

where

$$\left[|f| > 1/n\right] := \left\{ x \in \mathbb{R}^{\mathbb{N}} : |f(x)| > 1/n \right\},$$

and, by Chebyshev's inequality,

$$\mu\left([|f|^{p} > 1/n]\right) \le n^{p} \int_{\mathbb{R}^{\mathbb{N}}} |f|^{p} d\mu = n^{p} ||f||_{L^{p}}^{p} < \infty.$$

This implies that there exists an $\alpha_n < \omega_1$ such that

$$[|f|^p > 1/n] \subset \bigcup_{\xi < \alpha_n} R_{\xi} = \biguplus_{\xi < \alpha_n} B_{\xi}.$$

Consequently, as the cofinality of ω_1 is ω_1 , there exists $\alpha < \omega_1$ and $\beta < \omega_1$ such that

$$\operatorname{supp}(f) \subset \bigcup_{\xi < \alpha} R_{\xi} = \biguplus_{\xi < \alpha} B_{\xi} = \biguplus_{\xi < \beta} C_{\xi}$$

Therefore, we deduce that

$$\int_{\mathbb{R}^{\mathbb{N}}} |f|^{p} d\mu = \int_{\biguplus_{\xi < \beta} C_{\xi}} |f|^{p} d\mu = \sum_{\xi < \beta} \int_{C_{\xi}} |f|^{p} d\mu = \sum_{\xi < \omega_{1}} \int_{C_{\xi}} |f|^{p} d\mu.$$

The proof is concluded.

Before presenting the main result of this section, we need the following lemma.

Lemma 2.3. Let $M \in \mathcal{B}_{\infty}$ be such that $\mu(M) < \infty$. Then, $L^{p}(M, \mu)$ is separable.

Proof. Let $M \in \mathcal{B}_{\infty}$ with $\mu(M) < \infty$. By the definition of the Borel σ -algebra \mathcal{B}_{∞} , it is apparent that

$$\mathcal{B}_{\infty} := \sigma\left(\mathcal{Q}\right),$$

where

$$\mathcal{Q} := \left\{ \bigotimes_{i=1}^{m} B_i \times \bigotimes_{i=m+1}^{\infty} \mathbb{R} : B_i \in \mathcal{B}_r, \ \forall i \in \{1, 2, \dots, m\} \text{ and } m \in \mathbb{N} \right\}$$

with $\mathcal{B}_r := \{(a, b) : a, b \in \mathbb{Q}, a < b\}$. In particular, $\mathcal{B}_{\infty} \cap M$ is countably generated and $\mu|_M$ is a finite measure. Then, Proposition 3.4.5 of Cohn [4] concludes the proof.

We proceed to state and prove the main result of this section.

Theorem 2.4. Let $1 \leq p < \infty$. Then, the space $L^p(\mu)$ is isometrically isomorphic to $\ell^p(\mathfrak{c}, L^p[0,1])$. Symbolically,

$$L^p(\mu) \cong \ell^p(\mathfrak{c}, L^p[0,1]).$$

Proof. Consider the linear operator

$$T: L^{p}(\mu) \to \left(\bigoplus_{\xi < \omega_{1}} L^{p}(C_{\xi})\right)_{p}$$
$$f \mapsto Tf := (f_{\xi})_{\xi < \omega_{1}},$$

where

$$f_{\xi} := f \uparrow C_{\xi}$$

for every $\xi < \omega_1$. The operator T is well defined as a consequence of Proposition 2.2. Indeed, note that

$$||Tf||_{\ell^p}^p = \sum_{\xi < \omega_1} ||f_\xi||_{L^p(C_\xi)}^p = \sum_{\xi < \omega_1} \int_{C_\xi} |f|^p \, d\mu = \int_{\mathbb{R}^N} |f|^p \, d\mu < \infty.$$

Moreover, this proves that T is an isometry. Now, let $F \equiv (f_{\xi})_{\xi < \omega_1} \in \left(\bigoplus_{\xi < \omega_1} L^p(C_{\xi})\right)_p$. Note that $\operatorname{supp}(F)$ is countable, where $\operatorname{supp}(F) := \{\xi < \omega_1 : \|f_{\xi}\|_{L^p} \neq 0\}$. Define $f : \mathbb{R}^{\mathbb{N}} \to \mathbb{R}$ as

$$f := \sum_{\xi \in \operatorname{supp}(F)} f_{\xi} \chi_{B_{\xi}}.$$

Then, it follows that $f \in L^p(\mu)$ as

$$\|f\|_{L^p}^p = \sum_{\xi \in \text{supp}(F)} \int_{C_{\xi}} |f_{\xi}|^p \, d\mu = \sum_{\xi \in \text{supp}(F)} \|f_{\xi}\|_{L^p}^p = \sum_{\xi < \omega_1} \|f_{\xi}\|_{L^p}^p = \|F\|_{\ell^p}^p < \infty.$$

Moreover, T(f) = F. Therefore, T is an isometric isomorphism.

Now, as $\mu(C_{\xi}) < \infty$ for each $\xi < \omega_1$, by Lemma 2.3, it follows that $L^p(C_{\xi})$ is separable. Hence, by [8, Cr. p. 128], $L^p(C_{\xi})$ is isometrically isomorphic to $L^p[0, 1]$ for each $1 \le p < \infty$. Therefore, we conclude that $L^p(\mu)$ is isometrically isomorphic to $\ell^p(\mathfrak{c}, L^p[0, 1])$.

A direct consequence of Theorem 2.4 is the following.

Corollary 2.5. The space $L^2(\mu)$ is isometrically isomorphic to $\ell^2(\mathfrak{c})$.

Proof. This follows from the following chain of isometric isomorphisms:

$$L^2(\mu) \cong \ell^2(\mathfrak{c}, L^2[0, 1]) \cong \ell^2(\mathfrak{c}, \ell^2) \cong \ell^2(\mathfrak{c}).$$

The proof is concluded.

Theorem 2.4 provides a finite-dimensional reduction formula for the computation of the integral of a μ -integrable function. For the statement of this result, we need some notation. Given the finite rectangle $C_{\xi} = \bigotimes_{i \in \mathbb{N}} C_{\xi,i} \in \mathcal{F}(\mathcal{B}, \lambda)$, we denote

$$C^n_{\xi} := \bigotimes_{i=1}^n C_{\xi,i},$$

for every $n \in \mathbb{N}$. Note that μ restricted to C_{ξ} is the product measure $\bigotimes_{i \in \mathbb{N}} (\lambda \upharpoonright C_{\xi,i})$. We denote $\mu_{\xi}^n := \bigotimes_{i=1}^n (\lambda \upharpoonright C_{\xi,i})$ for every $n \in \mathbb{N}$.

Theorem 2.6. Let $f \in L^1(\mu)$. Then, there exists a family of integrable functions

$$\{f_{\xi,n} \in L^1(\mu_{\xi}^n) : \xi < \omega_1, n \in \mathbb{N}\},\$$

such that

(2.2)
$$\int_{\mathbb{R}^N} f \, d\mu = \lim_{n \to \infty} \sum_{\xi < \omega_1} \int_{C^n_{\xi}} f_{\xi,n} \, d\mu^n_{\xi}.$$

Proof. Let $f \in L^1(\mu)$. Then, by Proposition 2.2,

$$\int_{\mathbb{R}^N} |f| \, d\mu = \sum_{\xi < \omega_1} \int_{C_{\xi}} |f| \, d\mu.$$

As $f \upharpoonright C_{\xi} \in L^1(C_{\xi})$ and $\mu(C_{\xi}) < \infty$, by [27, Th. 3.4], there exists a sequence $(f_{\xi,n})_{n \in \mathbb{N}}$ such that

$$\int_{C_{\xi}} |f| \ d\mu = \lim_{n \to \infty} \int_{C_{\xi}^n} |f_{\xi,n}| \ d\mu_{\xi}^n.$$

Finally, by Beppo–Levi theorem,

$$\int_{\mathbb{R}^{\mathbb{N}}} |f| \, d\mu = \sum_{\xi < \omega_1} \int_{C_{\xi}} |f| \, d\mu = \sum_{\xi < \omega_1} \lim_{n \to \infty} \int_{C_{\xi}^n} |f_{\xi,n}| \, d\mu_{\xi}^n = \lim_{n \to \infty} \sum_{\xi < \omega_1} \int_{C_{\xi}^n} |f_{\xi,n}| \, d\mu_{\xi}^n.$$

This concludes the proof of (2.2) for $f \ge 0$. In the general case, rewrite f as $f = f_+ - f_$ and apply the same argument to $f_+, f_- \ge 0$.

3. Study of the general case

In this section, we study the isometric structure of $L^p(\mu)$ in the general case, that is, if we do not impose the validity of CH. We prove that $L^p(\mu)$ contains an isometric and complemented copy of $\ell^p(\mathfrak{c}, L^p[0, 1])$. This analysis is performed via a cube decomposition of $\mathbb{R}^{\mathbb{N}}$ that splits $L^p(\mu)$ into two complemented subspaces G^p and B^p .

Let us denote $I := \mathbb{Z}^{\mathbb{N}}$. For each $\mathfrak{a} = (a_n)_{n \in \mathbb{N}} \in I$, we introduce the unit cube

$$\mathcal{C}_{\mathfrak{a}} := \bigotimes_{n \in \mathbb{N}} [a_n, a_n + 1).$$

Note that $\mathcal{C}_{\mathfrak{a}} \in \mathcal{F}(\mathcal{B}, \lambda)$ and $\mu(\mathcal{C}_{\mathfrak{a}}) = \operatorname{vol}(\mathcal{C}_{\mathfrak{a}}) = 1$ for every $\mathfrak{a} \in I$. Moreover, it is apparent that $\mathcal{C}_{\mathfrak{a}} \cap \mathcal{C}_{\mathfrak{b}} = \emptyset$ for each $\mathfrak{a} \neq \mathfrak{b}$ and

$$\mathbb{R}^{\mathbb{N}} = \biguplus_{\mathfrak{a} \in I} \mathcal{C}_{\mathfrak{a}}.$$

The following result studies the behavior of a function $f \in L^p(\mu)$ with respect this decomposition.

Theorem 3.1. Let $1 \le p < \infty$ and $f \in L^p(\mu)$. Then the set

$$\mathcal{O}_f := \left\{ \mathfrak{a} \in I : \int_{\mathcal{C}_\mathfrak{a}} |f| \ d\mu \neq 0 \right\}$$

is countable and

3.1)
$$\sum_{\mathfrak{a}\in\mathcal{O}_f}\int_{\mathcal{C}_{\mathfrak{a}}}|f|^p \ d\mu = \int_{\mathcal{F}_f}|f|^p \ d\mu \le \int_{\mathbb{R}^N}|f|^p \ d\mu,$$

where

(

$$\mathcal{F}_f := \biguplus_{\mathfrak{a} \in \mathcal{O}_f} \mathcal{C}_{\mathfrak{a}}.$$

Proof. If \mathcal{O}_f is uncountable, there are $\varepsilon > 0$ and infinite $F \subset \mathcal{O}_f$ such that $\int_{\mathcal{C}_a} |f| d\mu \ge \varepsilon$ for all $\mathfrak{a} \in F$. Then,

$$\int_{\mathbb{R}^{\mathbb{N}}} |f|^p \ d\mu \ge \sum_{\mathfrak{a} \in F} \int_{\mathcal{C}_{\mathfrak{a}}} |f|^p \ d\mu \ge \sum_{\mathfrak{a} \in F} \left(\int_{\mathcal{C}_{\mathfrak{a}}} |f| \ d\mu \right)^p \ge \sum_{\mathfrak{a} \in F} \varepsilon^p = \infty,$$

contradicting $f \in L^p(\mu)$. This concludes the proof, as countability of \mathcal{O}_f clearly implies (3.1).

It would be desirable that inequality (3.1) were in fact an equality. However, as we shall see in subsection 3.2, this is not always true (as we have uncountably many sets $C_{\mathfrak{a}}$ and the integrals are only countably additive with respect to integrated sets). This fact motivates the definition of the following subsets of $L^{p}(\mu)$, $1 \leq p < \infty$:

$$G^p := \left\{ f \in L^p(\mu) : \int_{\mathbb{R}^N} |f|^p \ d\mu = \sum_{\mathfrak{a} \in \mathcal{O}_f} \int_{\mathcal{C}_\mathfrak{a}} |f|^p \ d\mu = \int_{\mathcal{F}_f} |f|^p \ d\mu \right\},$$

and

$$B^p := \left\{ f \in L^p(\mu) : \int_{\mathcal{C}_{\mathfrak{a}}} |f| \ d\mu = 0 \text{ for each } \mathfrak{a} \in I \right\}$$

The next two subsections are devoted to the study of these sets.

3.1. Study of G^p . In this subsection, we proceed to study the subsets G^p , $1 \le p < \infty$. We are able to characterize completely its structure. We start by proving that G^p is a subspace of $L^p(\mu)$. We need a preliminary lemma.

Lemma 3.2. For every $f_1, f_2 \in L^p(\mu)$, we have that $\mathcal{O}_{f_1+f_2} \subset \mathcal{O}_{f_1} \cup \mathcal{O}_{f_2}$, $\mathcal{F}_{f_1+f_2} \subset \mathcal{F}_{f_1} \cup \mathcal{F}_{f_2}$ and

$$\int_{\mathcal{F}_{f_1+f_2}} |f_1 + f_2|^p \ d\mu = \int_{\mathcal{F}_{f_1} \cup \mathcal{F}_{f_2}} |f_1 + f_2|^p \ d\mu.$$

Proof. Let $\mathfrak{a} \in \mathcal{O}_{f_1+f_2}$. Then,

$$0 < \int_{\mathcal{C}_{\mathfrak{a}}} |f_1 + f_2| \ d\mu \leq \int_{\mathcal{C}_{\mathfrak{a}}} |f_1| \ d\mu + \int_{\mathcal{C}_{\mathfrak{a}}} |f_2| \ d\mu.$$

Therefore,

$$\int_{\mathcal{C}_{\mathfrak{a}}} |f_1| \ d\mu \neq 0 \quad \text{or} \quad \int_{\mathcal{C}_{\mathfrak{a}}} |f_2| \ d\mu \neq 0.$$

and this implies that $\mathfrak{a} \in \mathcal{O}_{f_1} \cup \mathcal{O}_{f_2}$, completing the proof of inclusions. The equation follows from the countability of $\mathcal{F}_{f_1} \cup \mathcal{F}_{f_2}$ and the fact that $\int_{\mathcal{C}_\mathfrak{a}} |f_1 + f_2| d\mu = 0$ for every $\mathfrak{a} \in \mathcal{O}_{f_1} \cup \mathcal{O}_{f_2}$ not in $\mathcal{O}_{f_1+f_2}$.

The following result proves that G^p is a subspace of $L^p(\mu)$.

Lemma 3.3. For each $1 \leq p < \infty$, G^p is a normed space.

Proof. It is enough to prove that G^p is a vector subspace of $L^p(\mu)$. To see this, fix $f_1, f_2 \in G^p$ and notice that

$$\left(\int_{(\mathcal{F}_{f_1} \cup \mathcal{F}_{f_2})^c} |f_1 + f_2|^p \, d\mu \right)^{1/p} \leq \left(\int_{(\mathcal{F}_{f_1} \cup \mathcal{F}_{f_2})^c} |f_1|^p \, d\mu \right)^{1/p} + \left(\int_{(\mathcal{F}_{f_1} \cup \mathcal{F}_{f_2})^c} |f_2|^p \, d\mu \right)^{1/p} \\ \leq \left(\int_{\mathcal{F}_{f_1}^c} |f_1|^p \, d\mu \right)^{1/p} + \left(\int_{\mathcal{F}_{f_2}^c} |f_2|^p \, d\mu \right)^{1/p} = 0,$$

where the equality follows from the fact that $f_1, f_2 \in G^p$. So,

$$\int_{(\mathcal{F}_{f_1} \cup \mathcal{F}_{f_2})^c} |f_1 + f_2|^p \, d\mu = 0.$$

By this and Lemma 3.2

$$\begin{split} \int_{\mathbb{R}^{\mathbb{N}}} |f_1 + f_2|^p \ d\mu &\leq \int_{\mathcal{F}_{f_1} \cup \mathcal{F}_{f_2}} |f_1 + f_2|^p \ d\mu + \int_{(\mathcal{F}_{f_1} \cup \mathcal{F}_{f_2})^c} |f_1 + f_2|^p \ d\mu \\ &= \int_{\mathcal{F}_{f_1 + f_2}} |f_1 + f_2|^p \ d\mu. \end{split}$$

This proves that $f_1 + f_2 \in G^p$. If $f \in G^p$ and $k \in \mathbb{R}$, it is obvious that $kf \in G^p$. Therefore, G^p is a vector subspace of $L^p(\mu)$.

The next result characterizes the spaces G^p , $1 \leq p < \infty$ up to isometric isomorphism. Firstly, we need some preliminary definitions. For each $\mathfrak{a} = (a_n)_{n \in \mathbb{N}} \in I$, we define the translation $T_{\mathfrak{a}} : \mathcal{C}_{\mathfrak{a}} \to \mathcal{H}$ as

$$T(x_n)_{n\in\mathbb{N}}\mapsto (x_n-a_n)_{n\in\mathbb{N}},$$

where $\mathcal{H} := [0,1)^{\mathbb{N}}$ is the Hilbert cube. Then, it is clear that $T_{\mathfrak{a}}(\mathcal{C}_{\mathfrak{a}}) = \mathcal{H}$. Moreover, since μ is translation invariant, we have that

(3.2)
$$\mu(A) = \mu(T_{\mathfrak{a}}(A)) = \mu(T_{\mathfrak{a}}^{-1}(A)) \text{ for each } A \in \mathcal{B}_{\infty}.$$

Theorem 3.4. The following spaces are isometrically isomorphic for $1 \le p < \infty$,

$$G^p \cong \ell^p(\mathfrak{c}, L^p[0, 1]).$$

Proof. Fix $1 \leq p < \infty$ and define the operator

$$\Psi: G^p \to \ell^p(I, L^p(\mathcal{H})),$$
$$f \mapsto \left(f \circ T_{\mathfrak{a}}^{-1}\right)_{\mathfrak{a} \in I}.$$

Given $f \in G^p$, we have

$$||f||_{L^p}^p = \int_{\mathbb{R}^N} |f|^p \ d\mu = \sum_{\mathfrak{a} \in \mathcal{O}_f} \int_{\mathcal{C}_\mathfrak{a}} |f|^p \ d\mu.$$

For each $\mathfrak{a} \in I$, by the change of variable formula and identity (3.2), we deduce

$$\int_{\mathcal{C}_{\mathfrak{a}}} |f|^p \ d\mu = \int_{T_{\mathfrak{a}}(\mathcal{C}_{\mathfrak{a}})} |(f \circ T_{\mathfrak{a}}^{-1})(x)|^p \ d\mu(T_{\mathfrak{a}}^{-1}(x)) = \int_{\mathcal{H}} |f \circ T_{\mathfrak{a}}^{-1}|^p \ d\mu$$

Therefore,

$$\|f\|_{L^p}^p = \sum_{\mathfrak{a}\in\mathcal{O}_f} \int_{\mathcal{C}_\mathfrak{a}} |f|^p \ d\mu = \sum_{\mathfrak{a}\in\mathcal{O}_f} \int_{\mathcal{H}} |f \circ T_\mathfrak{a}^{-1}|^p \ d\mu = \sum_{\mathfrak{a}\in I} \|f \circ T_\mathfrak{a}^{-1}\|_{L^p(\mathcal{H})}^p = \|\Psi(f)\|_{\ell^p}^p.$$

Consequently Ψ is well defined and it is an isometry. Finally, we see that Ψ is onto. Let $\overline{f} = (f_{\mathfrak{a}})_{\mathfrak{a}\in I} \in \ell^p(I, L^p(\mathcal{H}))$ and denote by $\operatorname{supp}(\overline{f})$ the support of \overline{f} , that is,

$$\operatorname{supp}(\bar{f}) := \{ \mathfrak{a} \in I : \| f_\mathfrak{a} \|_{L^p(\mathcal{H})} \neq 0 \}.$$

Clearly, by the definition of $\ell^p(I, L^p(\mathcal{H}))$, the set $\operatorname{supp}(\bar{f})$ is countable. Define

$$f := \sum_{\mathfrak{a} \in \operatorname{supp}(\bar{f})} (f_{\mathfrak{a}} \circ T_{\mathfrak{a}}) \cdot \chi_{\mathcal{C}_{\mathfrak{a}}}.$$

By the monotone convergence theorem and the change of variable formula, it follows that

$$\int_{\mathbb{R}^{\mathbb{N}}} |f|^{p} d\mu = \int_{\mathbb{R}^{\mathbb{N}}} \sum_{\mathfrak{a} \in \operatorname{supp}(\bar{f})} |f_{\mathfrak{a}} \circ T_{\mathfrak{a}}|^{p} \cdot \chi_{\mathcal{C}_{\mathfrak{a}}} d\mu = \sum_{\mathfrak{a} \in \operatorname{supp}(\bar{f})} \int_{\mathcal{C}_{\mathfrak{a}}} |f_{\mathfrak{a}} \circ T_{\mathfrak{a}}|^{p} d\mu$$
$$= \sum_{\mathfrak{a} \in \operatorname{supp}(\bar{f})} \int_{\mathcal{H}} |f_{\mathfrak{a}}|^{p} d\mu = \|\bar{f}\|_{\ell^{p}}^{p} < \infty.$$

Therefore, $f \in L^p(\mu)$, and it is straightforward to verify that $\Psi(f) = \overline{f}$. Hence, Ψ is an isometric isomorphism.

Finally, as $\mu(\mathcal{H}) = 1$, by Lemma 2.3, it follows that $L^p(\mathcal{H})$ is separable. Hence, by [8, Cr. p. 128], $L^p(\mathcal{H})$ is isometrically isomorphic to $L^p[0, 1]$ for each $1 \leq p < \infty$. Therefore, we conclude that G^p is isometrically isomorphic to $\ell^p(\mathfrak{c}, L^p[0, 1])$.

3.2. Study of B^p . This subsection is devoted to the study of the sets B^p for $1 \le p < \infty$. These sets contain the *pathological* behavior of functions in $L^p(\mu)$. We describe some structure result of these sets. We start by proving that B^p are Banach spaces.

Lemma 3.5. For every $1 \le p < \infty$, B^p is a Banach space.

Proof. It is easy to verify that B^p is a vector subspace of $L^p(\mu)$. It remains to prove that it is closed. Let $(f_n)_{n \in \mathbb{N}} \subset B^p$ such that $f_n \to f$ as $n \to \infty$ in $L^p(\mu)$ for some $f \in L^p(\mu)$. Then, we obtain

$$\left| \int_{\mathcal{C}_{\mathfrak{a}}} |f_n|^p \, d\mu - \int_{\mathcal{C}_{\mathfrak{a}}} |f|^p \, d\mu \right| \leq \int_{\mathcal{C}_{\mathfrak{a}}} |f_n - f|^p \, d\mu \leq \int_{\mathbb{R}^N} |f_n - f|^p \, d\mu \xrightarrow[n \to \infty]{} 0.$$

Therefore, we deduce

$$0 = \lim_{n \to \infty} \int_{\mathcal{C}_{\mathfrak{a}}} |f_n|^p \ d\mu = \int_{\mathcal{C}_{\mathfrak{a}}} |f|^p \ d\mu,$$

for each $\mathfrak{a} \in I$. This proves that $f \in B^p$ and the proof is concluded.

The main result of this section is the following.

Theorem 3.6. For each $1 \leq p < \infty$, B^p contains an isometric and complemented copy of $\ell^p(\mathfrak{c}, L^p[0, 1])$.

Proof. Recall that two infinite subsets $A, B \subset \mathbb{N}$ are called *almost disjoint* if $|A \cap B| < \infty$. It is well known that there exists a family of almost disjoint subsets \mathcal{A} of \mathbb{N} with $|\mathcal{A}| = \mathfrak{c}$.¹ For every $A \in \mathcal{A}$, let us denote

$$A := \{ n_1^A < n_2^A < \cdots \}.$$

Then, for each $A \in \mathcal{A}$ and each $k \in \mathbb{N}$, let us consider two subintervals $I_k^A \subset (n_k^A - 1, n_k^A)$, $J_k^A \subset (-n_k^A, 1 - n_k^A)$ of length 2^{-k-1} . Consider the Borel set

$$\mathcal{I}_A := \biguplus_{k \in \mathbb{N}} (I_k^A \uplus J_k^A) \in \mathcal{B}$$

Then, by the σ -additivity of the Lebesgue measure λ , we obtain that

$$\lambda(\mathcal{I}_A) = \sum_{k \in \mathbb{N}} (\lambda(I_k^A) + \lambda(J_k^A)) = \sum_{k \in \mathbb{N}} 2^{-k} = 1.$$

For each $A \in \mathcal{A}$, consider the associated rectangle

$$C_A := \bigotimes_{n \in \mathbb{N}} \mathcal{I}_A.$$

Then, $C_A \in \mathcal{F}(\mathcal{B}, \lambda)$ and

$$\mu(C_A) = \prod_{n \in \mathbb{N}} \lambda(\mathcal{I}_A) = 1.$$

¹For every $x \in \mathbb{R}$, fix a sequence of rational numbers $(q_{x,n})_{n \in \mathbb{N}}$ that converges to x. Then, by considering f a bijection from \mathbb{Q} to \mathbb{N} , we have that the sets $A_x := \{f(q_{x,n}) : n \in \mathbb{N}\}$ form an almost disjoint family of subsets of \mathbb{N} having cardinality \mathfrak{c} .

Note that given two distinct sets $A, B \in \mathcal{A}$, necessarily $|A \cap B| < \infty$ and, therefore, $\mathcal{I}_A \cap \mathcal{I}_B$ is a finite union of intervals. Consequently $\lambda(\mathcal{I}_A \cap \mathcal{I}_B) < 1$. This implies that

$$\mu(C_A \cap C_B) = \mu\left(\left(\bigotimes_{n \in \mathbb{N}} \mathcal{I}_A\right) \cap \left(\bigotimes_{n \in \mathbb{N}} \mathcal{I}_B\right)\right)$$
$$= \mu\left(\bigotimes_{n \in \mathbb{N}} \mathcal{I}_A \cap \mathcal{I}_B\right) = \prod_{n \in \mathbb{N}} \lambda(\mathcal{I}_A \cap \mathcal{I}_B) = 0.$$

On the one hand, note that, by definition, every

$$\bar{f} = (f_A)_{A \in \mathcal{A}} \in \left(\bigoplus_{A \in \mathcal{A}} L^p(C_A)\right)_p$$

has countable support $\operatorname{supp}(\bar{f})$, where

$$\operatorname{supp}(\bar{f}) := \left\{ A \in \mathcal{A} : \|f_A\|_{L^p(C_A)} \neq 0 \right\}.$$

Define the linear operator

$$T: \left(\bigoplus_{A \in \mathcal{A}} L^p(C_A)\right)_p \to B^p$$

as follows:

$$T(f_A)_{A \in \mathcal{A}} = \sum_{A \in \operatorname{supp}(\bar{f})} f_A \chi_{C_A}.$$

Since for each $\mathfrak{a} \in I$ and each $A \in \mathcal{A}$, the identity $\mu(\mathcal{C}_{\mathfrak{a}} \cap C_A) = 0$ holds, we deduce that

$$\int_{\mathcal{C}_{\mathfrak{a}}} |T(f_A)_{A \in \mathcal{A}}|^p \ d\mu = \sum_{A \in \operatorname{supp}(\bar{f})} \int_{\mathcal{C}_{\mathfrak{a}} \cap C_A} |f_A|^p \ d\mu = 0,$$

which implies that $T(f_A)_{A \in \mathcal{A}} \in B^p$. Moreover,

$$\|T(f_A)_{A\in\mathcal{A}}\|_{L^p}^p = \int_{\mathbb{R}^N} \left| \sum_{A\in\mathrm{supp}(\bar{f})} f_A \chi_{C_A} \right|^p d\mu = \sum_{A\in\mathrm{supp}(\bar{f})} \|f_A\|_{L^p(C_A)}^p = \|(f_A)_A\|_{\ell^p}^p$$

This proves that T is well defined and that it is an isometry. On the other hand, as $\mu(C_A) = 1$ for each $A \in \mathcal{A}$, by Lemma 2.3, $L^p(C_A)$ is isometrically isomorphic to $L^p[0, 1]$. Therefore,

$$\left(\bigoplus_{A\in\mathcal{A}}L^p(C_A)\right)_p\cong\ell^p(\mathfrak{c},L^p[0,1]).$$

Consequently, $\ell^p(\mathfrak{c}, L^p[0, 1])$ embeds isometrically into B^p .

Finally, let us prove that $V := \operatorname{Im}(T) = \ell^p(\mathfrak{c}, L^p[0, 1])$ is a complemented subspace of B^p . Let us start by showing that for each $f \in B^p$, the support $\operatorname{supp}(f)$ is countable. If $\operatorname{supp}(f)$ is uncountable, then there are $\varepsilon > 0$ and infinite countable $F \subset \operatorname{supp}(f)$ such that $\int_{C_A} |f| d\mu \ge \varepsilon$ for all $A \in F$. Then

$$\int_{\mathbb{R}^{\mathbb{N}}} |f|^p \ d\mu \ge \sum_{A \in F} \int_{C_A} |f|^p \ d\mu \ge \sum_{A \in F} \varepsilon^p = \infty,$$

contradicting $f \in L^p(\mu)$.

Given $f \in B^p$, we denote

$$\mathfrak{A}_f := \bigcup_{A \in \mathrm{supp}(f)} C_A.$$

Consider the linear projection

$$P: B^p \to V$$

defined by

$$Pf := f \upharpoonright \mathfrak{A}_f = \sum_{A \in \mathrm{supp}(f)} (f \upharpoonright C_A) \chi_{C_A}.$$

Then, for each $f \in B^p$,

$$\|Pf\|_{L^p}^p = \int_{\mathbb{R}^N} |f\chi_{\mathfrak{A}_f}|^p \, d\mu \le \int_{\mathbb{R}^N} |f|^p \, d\mu = \|f\|_{L^p}^p.$$

This proves that V is complemented and concludes the proof.

The last result of this section gives the desired decomposition of $L^p(\mu)$ in the complemented subspaces G^p and B^p .

Theorem 3.7. For each $1 \leq p < \infty$, the following decomposition into complemented subspaces holds,

$$L^p(\mu) = G^p \oplus B^p \cong \ell^p(\mathfrak{c}, L^p[0, 1]) \oplus B^p.$$

Proof. Since G^p and B^p are closed subspaces, it is enough to prove the algebraic decomposition. Each $f \in L^p(\mu)$ can be decomposed as

$$f = f\chi_{\mathcal{F}_f} + f(1 - \chi_{\mathcal{F}_f})$$

It is easily verified that $f\chi_{\mathcal{F}_f} \in G^p$ and $f(1-\chi_{\mathcal{F}_f}) \in B^p$.

Therefore, we obtain the following.

Corollary 3.8. For $1 \le p < \infty$, the space $L^p(\mu)$ contains an isometric and complemented copy of $\ell^p(\mathfrak{c}, L^p[0, 1])$.

4. CHARACTERIZATION OF ISOMETRIES

In this final section, we characterize when $L^{p}(\nu)$ for some measure ν , is isometrically isomorphic to $\ell^{p}(\kappa, L^{p}[0, 1])$ for some cardinal number κ , symbolically,

$$L^p(\nu) \cong \ell^p(\kappa, L^p[0, 1])$$

Moreover, in the case in which these spaces are isometrically isomorphic, we describe all the isometric isomorphisms between these spaces.

We begin by introducing two definitions.

Definition 4.1 (κ -virtual partition). Let (X, Σ, ν) be a measure space and κ a cardinal number. A family $\{X_i : i \in J\} \subset \Sigma$ of σ -finite measurable subsets is a κ -virtual partition if $|J| = \kappa$,

(4.1)
$$\nu(X_i \cap X_j) = 0 \text{ for distinct } i, j \in J,$$

and for every $E \in \Sigma$ of finite measure,

(4.2)
$$\nu(E) = \sum_{i \in J} \nu(E \cap X_i).$$

Definition 4.2 (κ -virtual envelope). Let (X, Σ, ν) be a measure space and κ a cardinal number. A family $\{X_i : i \in J\} \subset \Sigma$ of σ -finite measurable subsets with $\nu(X_i) > 0$ for each $i \in J$ is a κ -virtual envelope if $|J| = \kappa$,

(4.3)
$$\nu(X_i \cap X_j) = 0 \text{ for distinct } i, j \in J,$$

and for every $E \in \Sigma$ of finite measure with $E \subset \bigcup_{i \in J} X_i$,

(4.4)
$$\nu(E) = \sum_{i \in J} \nu(E \cap X_i)$$

The term *virtual* in Definitions 4.1 and 4.2 references to the fact that equations (4.1) and (4.3) are satisfied, that is, the pairwise intersection of the sets of the family is not necessarily empty but it is small in the sense of the measure ν .

The first result of this section shows that if a measure space admits a κ -virtual partition, then $L^p(\nu) \cong \ell^p(\kappa, L^p[0, 1])$.

Theorem 4.3. Let $1 \le p < \infty$. If a measure space (X, Σ, ν) admits a κ -virtual partition, then $L^p(X, \nu)$ is isometrically isomorphic to $\ell^p(\kappa, L^p[0, 1])$.

Proof. Let $\{X_i : i \in J\}$ be a κ -virtual partition and consider the operator

(4.5)
$$T: L^p(\nu) \to \left(\bigoplus_{i \in J} L^p(X_i)\right)_p$$

defined as

$$Tf = (f \upharpoonright X_i)_{i \in J}.$$

The operator (4.5) is well-defined. Indeed, for every $F \subset J$ finite and $i \in F$, we have that

$$\nu\left(X_i \cap \bigcup_{j \in F \setminus J} X_j\right) = \nu\left(\bigcup_{j \in F \setminus J} (X_i \cap X_j)\right) \le 0.$$

Thus, for any $F \subset J$ finite and $f \in L^p(\nu)$,

$$\sum_{i \in F} \int_{X_i} |f|^p \ d\nu \le \int_X |f|^p \ d\nu$$

Hence, for every $f \in L^p(\nu)$, it follows that

$$||Tf||_{\ell^{p}}^{p} = \sum_{i \in J} ||f| \upharpoonright X_{i}||_{L^{p}(X_{i})}^{p} = \sum_{i \in J} \int_{X_{i}} |f|^{p} d\nu$$
$$= \sup \left\{ \sum_{i \in F} \int_{X_{i}} |f|^{p} d\nu : F \subset J \text{ finite} \right\} \leq \int_{X} |f|^{p} d\nu < \infty.$$

For every $E \in \Sigma$ with finite measure, consider the characteristic function χ_E . Then, by (4.2), we deduce that

$$\|\chi_E\|_{L^p}^p = \int_X |\chi_E|^p \, d\nu = \nu(E) = \sum_{i \in J} \nu(E \cap X_i) = \sum_{i \in J} \int_{X_i} |\chi_E|^p \, d\nu$$
$$= \sum_{i \in J} \|\chi_E \upharpoonright X_i\|_{L^p(X_i)}^p = \|T\chi_E\|_{\ell^p}^p.$$

This property extends easily to $\mathcal{X} := \operatorname{span}\{\chi_E : E \in \Sigma^f\}$. That is, for each $f \in \mathcal{X}$,

(4.6)
$$||f||_{L^p} = ||Tf||_{\ell^p}^p.$$

Now, take a sequence $(f_n)_{n \in \mathbb{N}} \subset \mathcal{X}$ of functions that satisfy (4.6) and such that $f_n \to f$ in L^p as $n \to \infty$. Then,

$$||f||_{L^p}^p = \lim_{n \to \infty} \int_X |f_n|^p \, d\nu = \lim_{n \to \infty} \sum_{i \in J} \int_{X_i} |f_n|^p \, d\nu.$$

Since for each $n \in \mathbb{N}$, the sum is finite, there exist countable subsets $J_n \subset J$ such that $\|f_n\|_{L^p(X_i)} = 0$ for each $i \in J \setminus J_n$. Now consider $\hat{J} := \bigcup_{n \in \mathbb{N}} J_n$ (which is countable) and

$$\mathcal{J} := \bigcup \{ X_i : i \in \hat{J} \} \in \Sigma.$$

Hence, by convergence, $||f||_{L^p(X_i)} = 0$ for each $i \in J \setminus \mathcal{J}$. So,

$$\int_{\mathcal{J}} |f|^p \, d\nu = \sum_{i \in \mathcal{J}} \int_{X_i} |f|^p \, d\nu = \sum_{i \in J} \int_{X_i} |f|^p \, d\nu.$$

Thus,

$$\|f\|_{L^p}^p = \lim_{n \to \infty} \sum_{i \in J_n} \int_{X_i} |f_n|^p \, d\nu = \lim_{n \to \infty} \int_{\mathcal{J}} |f_n|^p \, d\nu = \int_{\mathcal{J}} |f|^p \, d\nu = \sum_{i \in J} \int_{X_i} |f|^p \, d\nu = \|Tf\|_{\ell^p}^p.$$

Therefore, by the density of \mathcal{X} in $L^p(\nu)$, we deduce that property (4.6) holds for every $f \in L^p(\nu)$. Consequently, T is an isometry. It remains to prove that it is onto. Let

$$(f_i)_{i\in J} \in \left(\bigoplus_{i\in J} L^p(X_i)\right)_p.$$

Then, there exists a countable subset $C \subset J$ such that $||f_i||_{L^p(X_i)} = 0$ for every $i \in J \setminus C$. (In particular, $f_i \upharpoonright X_i = 0$ ν -a.e. for every $i \in J \setminus C$.) Observe that for each $i \in C$ we have that

$$\nu\left(X_i \cap \bigcup_{j \in C \setminus \{i\}} X_j\right) = \nu\left(\bigcup_{j \in C \setminus \{i\}} (X_i \cap X_j)\right) \le \sum_{j \in C \setminus \{i\}} \nu(X_i \cap X_j) = 0.$$

Now consider the function $f: X \to \mathbb{R}$ defined as:

$$f := \sum_{i \in C} f_i \ \chi_{X_i}.$$

Note that $f \upharpoonright X_i = f_i \upharpoonright X_i \nu$ -a.e. for any $i \in J$, and $f \in L^p(\nu)$ since

$$||f||_{L^p}^p = \int_X |f|^p \, d\nu = \sum_{i \in C} \int_{X_i} |f_i|^p \, d\nu = ||(f_i)_{i \in J}||_{\ell^p}^p < \infty.$$

Clearly, it yields that $Tf = (f_i)_{i \in J}$. This proves that T is an isometric isomorphism. \Box

In particular, since every strictly localizable measure space (X, ν) admits a κ -virtual partition for some cardinal $\kappa \leq |X|$, we deduce the following result.

Corollary 4.4. Let (X, Σ, ν) be a strictly localizable measure space. Then, the space $L^p(\nu)$ is isometrically isomorphic to $\ell^p(\kappa, L^p[0, 1])$ for some cardinal $\kappa \leq |X|$.

The following result gives somehow the reverse implication of Theorem 4.3. We prove that if $L^p(\nu) \cong \ell^p(\kappa, L^p[0, 1])$, for some cardinal κ , then (X, ν) admits a κ -virtual envelope. Additionally, we fully characterize all the isometric isomorphisms between $L^p(\nu)$ and $\ell^p(\kappa, L^p[0, 1])$.

Theorem 4.5. Let (X, Σ, ν) be a measure space and $p \in [1, \infty) \setminus \{2\}$. If $L^p(\nu)$ is isometrically isomorphic to $\ell^p(\kappa, L^p[0, 1])$, then (X, ν) admits a κ -virtual envelope $\mathcal{P} = \{X_i : i \in J\}$ with $|J| = \kappa$. Moreover, every isometric isomorphism $T : L^p(\nu) \to \ell^p(\kappa, L^p[0, 1])$ is of the form

(4.7)
$$Tf = (R_i^{-1}(\theta(f) \upharpoonright X_i))_{i \in J},$$

for some $\theta \in \text{Isom}(L^p(\nu))$ and some family $(R_i)_{i \in J}$ of isometric isomorphisms $R_i : L^p[0,1] \to L^p(X_i)$.

15

Proof. Suppose that $L^{p}(\nu)$ is isometrically isomorphic to $\ell^{p}(\kappa, L^{p}[0, 1])$. Then, there exists an isometric isomorphism

$$T \equiv (T_i)_{i \in J} : L^p(\nu) \to \ell^p(\kappa, L^p[0, 1]),$$

for some set J such that $|J| = \kappa$. Let us denote for each $i \in J$, $f_i := T^{-1}(\mathfrak{e}_{ij})_{j \in J} \in L^p(\nu)$, where $(\mathfrak{e}_{ij})_{j \in J} \in \ell^p(\kappa, L^p[0, 1])$ is given by

$$\mathbf{\mathfrak{e}}_{ij} := \left\{ \begin{array}{ll} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{array} \right.$$

Let $X_i := \operatorname{supp}(f_i) \in \Sigma$ for any $i \in J$. Let us show that $\nu(X_i \cap X_j) = 0$ for each $i \neq j$. Since T is an isometry,

(4.8)
$$\|f_i\|_{L^p}^p = \sum_{j \in J} \|T_j f_i\|_{L^p[0,1]}^p = \sum_{j \in J} \|\mathfrak{e}_{ij}\|_{L^p[0,1]}^p = 1,$$

for any $i \in J$, and

$$\|f_i \pm f_j\|_{L^p}^p = \sum_{k \in J} \|T_k f_i \pm T_k f_j\|_{L^p[0,1]}^p = \sum_{k \in J} \|\mathfrak{e}_{ik} \pm \mathfrak{e}_{jk}\|_{L^p[0,1]}^p = 2$$

provided that $i \neq j$. Identity (4.8) implies that $\nu(X_i) > 0$ for each $i \in J$. Moreover, for $i \neq j$, f_i and f_j attains the equality in the Clarkson inequality, that is,

$$||f_i + f_j||_{L^p}^p + ||f_i - f_j||_{L^p}^p = 2\left(||f_i||_{L^p}^p + ||f_j||_{L^p}^p\right).$$

Therefore, as $p \neq 2$, by [9, Cor. 2.1], $f_i \cdot f_j = 0$ ν -a.e. Hence, $\nu(X_i \cap X_j) = 0$ for $i \neq j$. Moreover, as $f_i \in L^p(\nu)$, X_i is σ -finite. Consequently, for each $i \in J$, there is an isometric isomorphism $R_i : L^p[0,1] \to L^p(X_i,\nu)$. In this way, by composition with T, we obtain an isometric isomorphism

$$\hat{T}: L^p(\nu) \to \left(\bigoplus_{i \in J} L^p(X_i)\right)_p.$$

Now, consider the linear operator

$$R: \left(\bigoplus_{i\in J} L^p(X_i)\right)_p \to L^p(\nu)$$

given by

$$R(f_i)_{i\in J} := \sum_{i\in J_{\mathfrak{f}}} f_i \ \chi_{X_i},$$

where $\mathfrak{f} := (f_i)_{i \in J}$ and $J_{\mathfrak{f}} := \{i \in J : ||f_i||_{L^p(X_i)} \neq 0\}$ is countable. Note that R is well defined and is an isometry since

$$\|R(f_i)_{i\in J}\|_{L^p}^p = \int_X \sum_{i\in J_{\mathfrak{f}}} |f_i|^p \chi_{X_i} \, d\nu = \sum_{i\in J_{\mathfrak{f}}} \int_{X_i} |f_i|^p \, d\nu = \|(f_i)_{i\in J}\|_{\ell^p}^p.$$

We define $\theta := R \circ \hat{T} \in \text{Isom}(L^p(\nu))$. Then, for every $f \in L^p(\nu)$,

(4.9)
$$\theta(f) = (R \circ \hat{T})(f) = \sum_{i \in J_{\mathfrak{f}}} f_i \chi_{X_i},$$

where $\hat{T}f = (f_i)_{i \in J} = \mathfrak{f}$. Equation (4.9) means that there exists $N \in \Sigma$ with $\nu(N) = 0$ such that

$$\theta(f)(x) = \sum_{i \in J_{\mathfrak{f}}} f_i \chi_{X_i}(x), \quad x \in X \setminus N.$$

Since

$$\nu\left(X_i \cap \left(\bigcup_{j \in J_{\mathfrak{f}} \setminus \{i\}} X_j \cup N\right)\right) \le \nu\left(\bigcup_{j \in J_{\mathfrak{f}} \setminus \{i\}} (X_i \cap X_j)\right) + \nu(X_i \cap N) = 0,$$

we have that $f_i = \theta(f) \upharpoonright X_i \nu$ -a.e. and therefore,

(4.10)
$$\hat{T}(f) = (\theta(f) \upharpoonright X_i)_{i \in J}.$$

Let $E \in \Sigma$ with $\nu(E) < \infty$ and $E \subset \bigcup_{i \in J} X_i$. Then, $(\chi_E \upharpoonright X_i)_{i \in J} \in (\bigoplus_{i \in J} L^p(X_i))_p$ since

$$\|(\chi_E \upharpoonright X_i)_{i \in J}\|_{\ell^p}^p = \sum_{i \in J} \nu(E \cap X_i) := \sup\left\{\sum_{j \in F} \nu(E \cap X_j) : F \subset J \text{ finite}\right\} \le \nu(E) < \infty.$$

As \hat{T} is onto, there exists $f_E \in L^p(\nu)$ such that $\hat{T}(f_E) = (\chi_E \upharpoonright X_i)_{i \in J}$. Let us denote

$$J_E := \{ i \in J : \| \chi_E \upharpoonright X_i \|_{L^p(X_i)} \neq 0 \} = \{ i \in J : \nu(E \cap X_i) \neq 0 \}.$$

Hence, since J_E is countable and $E \subset \bigcup_{i \in J} X_i$, by equation (4.9), we deduce that

$$\theta(f_E) = \sum_{i \in J_E} \chi_{E \cap X_i} = \chi_E,$$

and since $\theta \in \text{Isom}(L^p(\nu))$,

$$|f_E||_{L^p}^p = ||\theta(f_E)||_{L^p}^p = ||\chi_E||_{L^p}^p = \nu(E).$$

Finally, as \hat{T} is an isometry,

$$\nu(E) = \|f_E\|_{L^p}^p = \|\hat{T}f_E\|_{\ell^p}^p = \|(\chi_E \upharpoonright X_i)_{i \in J}\|_{\ell^p}^p = \sum_{i \in J} \nu(E \cap X_i).$$

This proves that $\mathcal{P} := \{X_i : i \in J\}$ is a κ -virtual envelope. The factorization (4.7) is a direct consequence of (4.10). The proof is concluded.

The case p = 2, excluded in Theorem 4.5, is simpler. In this case, $L^2(\nu) \cong \ell^2(\kappa)$ for κ the cardinality of the maximal orthonormal set of $L^2(\nu)$. Thus, as κ is infinite,

$$L^{2}(\nu) \cong \ell^{2}(\kappa) \cong \ell^{2}(\kappa, \ell^{2}) \cong \ell^{2}(\kappa, L^{2}[0, 1]).$$

Hence, if κ is the cardinality of the maximal orthonormal set of $L^2(\nu)$, the isometric isomorphism $L^2(\nu) \cong \ell^2(\kappa, L^2[0, 1])$ always holds.

Acknowledgments: The authors would like to thank Prof. Krzysztof C. Ciesielski for his insightful and meaningful comments, which contributed to the main result of this paper, Theorem 2.4. His feedback also provided valuable input toward the proof of Theorem A.1 in the Appendix section and helped improve the formulation of Theorem A.2. Additionally, we appreciate his suggestions, which enhanced the overall presentation of this paper.

We also express our gratitude to Prof. Mingu Jung for his helpful comments regarding the classification of isometric isomorphisms in Theorem 4.5, which contributed to clarifying our discussion on this topic.

References

- R. Baker, "Lebesgue measure" on R[∞], Proc. Amer. Math. Soc. 113 (1991), no. 4, 1023–1029, DOI 10.2307/2048779. MR1062827
- [2] R. L. Baker, "Lebesgue measure" on ℝ[∞]. II, Proc. Amer. Math. Soc. 132 (2004), no. 9, 2577–2591, DOI 10.1090/S0002-9939-04-07372-1.
- [3] K. Ciesielski, Set theory for the working mathematician, London Mathematical Society Student Texts, vol. 39, Cambridge University Press, Cambridge, 1997.
- [4] D. L. Cohn, Measure theory, 2nd ed., Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks], Birkhäuser/Springer, New York, 2013.
- [5] P. Enflo and T. W. Starbird, Subspaces of L¹ containing L¹, Studia Math. 65 (1979), no. 2, 203–225, DOI 10.4064/sm-65-2-203-225.
- [6] P. Enflo and H. P. Rosenthal, Some results concerning $L^p(\mu)$ -spaces, J. Functional Analysis 14 (1973), 325–348, DOI 10.1016/0022-1236(73)90050-5.
- [7] M. Fabian, P. Habala, P. Hájek, V. Montesinos, and V. Zizler, Banach space theory, CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, Springer, New York, 2011. The basis for linear and nonlinear analysis.
- [8] H. E. Lacey, The isometric theory of classical Banach spaces, Die Grundlehren der mathematischen Wissenschaften, vol. Band 208, Springer-Verlag, New York-Heidelberg, 1974.
- [9] J. Lamperti, On the isometries of certain function-spaces, Pacific J. Math. 8 (1958), no. 3, 459–466.
- [10] G. B. Folland, *Real analysis*, 2nd ed., Pure and Applied Mathematics (New York), John Wiley & Sons, Inc., New York, 1999. Modern techniques and their applications; A Wiley-Interscience Publication. MR1681462
- [11] D. Fremlin, Baker's product measures, Miscellaneous research notes.
- [12] D. H. Fremlin, *Measure theory. Vol. 1*, Torres Fremlin, Colchester, 2004. The irreducible minimum; Corrected third printing of the 2000 original.
- [13] _____, Measure theory. Vol. 2, Torres Fremlin, Colchester, 2003. Broad foundations; Corrected second printing of the 2001 original.
- [14] _____, *Measure theory. Vol. 3*, Torres Fremlin, Colchester, 2004. Measure algebras; Corrected second printing of the 2002 original.
- [15] _____, Measure theory. Vol. 4, Torres Fremlin, Colchester, 2006. Topological measure spaces. Part I, II; Corrected second printing of the 2003 original.
- [16] _____, Measure theory. Vol. 5. Set-theoretic measure theory. Part I, Torres Fremlin, Colchester, 2015. Corrected reprint of the 2008 original.
- [17] _____, Measure theory. Vol. 5. Set-theoretic measure theory. Part II, Torres Fremlin, Colchester, 2015. Corrected reprint of the 2008 original.
- [18] U. Haagerup, H. P. Rosenthal, and F. A. Sukochev, On the Banach-isomorphic classification of L_p spaces of hyperfinite von Neumann algebras, C. R. Acad. Sci. Paris Sér. I Math. **331** (2000), no. 9, 691–695, DOI 10.1016/S0764-4442(00)01692-X (English, with English and French summaries).
- [19] T. Jech, Set theory, Springer-Verlag, New York, 2002.
- [20] J. Lindenstrauss and A. Pelczyński, Absolutely summing operators in Lp-spaces and their applications, Studia Math. 29 (1968), 275–326, DOI 10.4064/sm-29-3-275-326.
- [21] D. Maharam, On homogeneous measure algebras, Proc. Nat. Acad. Sci. U.S.A. 28 (1942), 108–111, DOI 10.1073/pnas.28.3.108.
- [22] R. E. A. C. Paley, Some theorems on abstract spaces, Bull. Amer. Math. Soc. 42 (1936), no. 4, 235–240, DOI 10.1090/S0002-9904-1936-06277-4.
- [23] G. R. Pantsulaia, On generators of shy sets on Polish topological vector spaces, New York J. Math. 14 (2008).
- [24] H. P. Rosenthal, *Embeddings of L¹ in L¹*, Conference in modern analysis and probability (New Haven, Conn., 1982), Contemp. Math., vol. 26, Amer. Math. Soc., Providence, RI, 1984, DOI 10.1090/conm/026/737412.
- [25] H. P. Rosenthal, Subspaces of L^p which do not contain L^p isomorphically, Séminaire d'Analyse Fonctionnelle (1978–1979), École Polytech., Palaiseau, 1979, pp. Exp. No. 28, 9.
- [26] H. P. Rosenthal, On subspaces of L^p, Ann. of Math. (2) 97 (1973), 344–373, DOI 10.2307/1970850.
- [27] J. C. Sampedro, On the space of infinite dimensional integrable functions, J. Math. Anal. Appl. 488 (2020), no. 1, DOI 10.1016/j.jmaa.2020.124043.

- [28] _____, Existence of infinite product measures, Real Anal. Exchange 48 (2023), no. 2, 271–284, DOI 10.14321/realanalexch.48.2.1643257005.
- [29] J. C. Sampedro, On the L^p-spaces of projective limits of probability measures, J. Theoret. Probab. 37 (2024), no. 3, 2665–2703, DOI 10.1007/s10959-024-01329-1. MR4784955

Appendix A. Some properties of the measure μ

In this appendix, we prove some properties and curiosities of the measure μ not present in the literature.

We start by proving that the measure μ is not localizable. Recall that a measure space (X, Σ, ν) is localizable if ν is semi-finite and every $\mathcal{E} \subset \Sigma$ admits an essential supremum $H_{\mathcal{E}} \in \Sigma$, which have the following two properties:

- (a) $\nu(E \setminus H_{\mathcal{E}}) = 0$ (i.e., $E \subset^* H_{\mathcal{E}}$) for every $E \in \mathcal{E}$;
- (b) if $G \in \Sigma$ is such that $\nu(E \setminus G) = 0$ (i.e., $E \subset^* G$) for every $E \in \mathcal{E}$, then $\nu(H_{\mathcal{E}} \setminus G) = 0$ (i.e., $H_{\mathcal{E}} \subset^* G$).

Theorem A.1. The measure space $(\mathbb{R}^{\mathbb{N}}, \mathcal{B}_{\infty}, \mu)$ is not localizable.

Proof. To see this, first notice that \mathcal{B}_{∞} contains the following family \mathbb{E} of continuum many pairwise disjoint sets of positive finite μ -measure (in fact, of measure 1):

$$\mathbb{E} = \left\{ \prod_{i \in \mathbb{N}} [s_i, s_i + 1] \colon s \in \{0, 2\}^{\mathbb{N}} \right\}$$

Now, by way of contradiction, assume that μ is localizable and for every $\mathcal{E} \in \mathcal{P}(\mathbb{E})$, let $H_{\mathcal{E}}$ be its essential supremum. Since $|\mathcal{P}(\mathbb{E})| = 2^{2^{\omega}} > 2^{\omega} = |\mathcal{B}_{\infty}|$, there are distinct $\mathcal{E}, \mathcal{E}' \in \mathcal{P}(\mathbb{E})$ with $H_{\mathcal{E}} = H_{\mathcal{E}'}$. Since $\mathcal{E} \neq \mathcal{E}'$, there is an $E \in \mathbb{E}$ that belongs to one of them, but not the other, say $E \in \mathcal{E} \setminus \mathcal{E}'$. Then we must have $E \subset^* H_{\mathcal{E}}$. At the same time, $H_{\mathcal{E}} = H_{\mathcal{E}'} \subset^* H_{\mathcal{E}} \setminus E$, a contradiction, since $H_{\mathcal{E}} \not\subset^* H_{\mathcal{E}} \setminus E$.

The second result of this appendix shows that nontrivial continuous functions $f : \mathbb{R}^{\mathbb{N}} \to \mathbb{R}$ under the product topology of $\mathbb{R}^{\mathbb{N}}$ do not belong to $L^{p}(\mu)$ for every $1 \leq p < \infty$. This is not the case for the usual Lebesgue measure on \mathbb{R}^{n} , $n \in \mathbb{N}$. We start by proving that every nonempty open subset of $\mathbb{R}^{\mathbb{N}}$ has infinite measure.

Theorem A.2. Every nonempty open subset $U \subset \mathbb{R}^{\mathbb{N}}$ satisfies $\mu(U) = \infty$.

Proof. Let U be a nonempty open subset of $\mathbb{R}^{\mathbb{N}}$ and take $x_0 \in U$. As μ is translation invariant, we can suppose without lost of generality that $x_0 = 0$. Then, there exists $\varepsilon > 0$ such that $B_{\varepsilon}(0) \subset U$, where $B_{\varepsilon}(0)$ is the ball in $\mathbb{R}^{\mathbb{N}}$ with center 0 and radius ε . By the definition of the product topology, we deduce the existence of $n_0 \in \mathbb{N}$ and $\delta > 0$ such that

$$\mathcal{Y}_{n_0} := \bigotimes_{n=1}^{n_0} (-\delta, \delta) \times \bigotimes_{n=n_0}^{\infty} \mathbb{R} \subset B_{\varepsilon}(0).$$

This implies that $\mu(U) \ge \mu(\mathcal{Y}_{n_0})$. Note that for each integer $k > n_0$,

$$\mathcal{Y}_{k,n_0} := \bigotimes_{n=1}^{n_0} (-\delta, \delta) \times \bigotimes_{n=n_0}^{\infty} [k, k+1) \subset \mathcal{Y}_{n_0}.$$

Moreover, $\mathcal{Y}_{k,n_0} \in \mathcal{F}(\mathcal{B},\lambda)$ and $\mathcal{Y}_{k_1,n_0} \cap \mathcal{Y}_{k_2,n_0} = \emptyset$ for $k_1 \neq k_2$. Consequently, by the σ -additivity of μ , we deduce that

$$\mu(\mathcal{Y}_{n_0}) \ge \mu\left(\biguplus_{k>n_0} \mathcal{Y}_{k,n_0}\right) = \sum_{k=n_0+1}^{\infty} \mu(\mathcal{Y}_{k,n_0}) = \sum_{k=n_0+1}^{\infty} (2\delta)^{n_0} = \infty.$$

The proof is concluded.

Theorem A.3. Let $f \in \mathcal{C}(\mathbb{R}^{\mathbb{N}})$ and $x_0 \in \mathbb{R}^{\mathbb{N}}$ such that $f(x_0) \neq 0$. Then, $f \notin L^p(\mu)$ for every $1 \leq p < \infty$. In particular, it holds that $\mathcal{C}(\mathbb{R}^{\mathbb{N}}) \cap L^p(\mu) = \{0\}$ for each $1 \leq p < \infty$.

Proof. Let $f \in \mathcal{C}(\mathbb{R}^{\mathbb{N}})$ and $x_0 \in \mathbb{R}^{\mathbb{N}}$ be such that $f(x_0) \neq 0$. As μ is translation invariant, we can suppose without lost of generality that $x_0 = 0$. Then, there exist $\varepsilon > 0$ and $\rho > 0$ such that $B_{\varepsilon}(0) \subset \{|f| > \rho\}$. By the definition of the product topology, we deduce the existence of $n_0 \in \mathbb{N}$ and $\delta > 0$ such that

$$\mathcal{Y}_{n_0} := \bigotimes_{n=1}^{n_0} (-\delta, \delta) \times \bigotimes_{n=n_0}^{\infty} \mathbb{R} \subset B_{\varepsilon}(0).$$

Since f is continuous and we are dealing with the Borel σ -algebra of $\mathbb{R}^{\mathbb{N}}$, it is apparent that f is measurable. Therefore,

$$\int_{\mathbb{R}^{\mathbb{N}}} |f|^p \ d\mu \ge \int_{\mathcal{Y}_{n_0}} |f|^p \ d\mu > \rho^p \mu(\mathcal{Y}_{n_0}) = \infty.$$

This concludes the proof.

DEPARTAMENTO DE MATEMÁTICA APLICADA A LA INGENIERÍA INDUSTRIAL, E.T.S.I.D.I., RONDA DE VALENCIA 3, UNIVERSIDAD POLITÉCNICA DE MADRID, MADRID, 28012, SPAIN.

Email address: dl.rodriguez.vidanes@upm.es

DEPARTAMENTO DE MATEMÁTICA APLICADA A LA INGENIERÍA INDUSTRIAL, E.T.S.I.D.I., RONDA DE VALENCIA 3, UNIVERSIDAD POLITÉCNICA DE MADRID, MADRID, 28012, SPAIN. Email address: juancarlos.sampedro@upm.es