Field theory aspects of η -deformed superstring background

Dibakar Roychowdhury^{a1}

^a Department of Physics, Indian Institute of Technology Roorkee, Roorkee 247667, Uttarakhand, India

Abstract

We explore various field theory aspects of integrable η -deformed geometry in type IIB supergravity by employing several holographic probes. These include the computation of holographic timelike entanglement entropy and estimation of various other field theory observables for example, the flow central charge and the quantum complexity. We also discuss the associated brane set up and compute Page charges. We further use them to calculate the coupling constant in the dual QFTs considering both small and large deformation limits.

¹dibakar.roychowdhury@ph.iitr.ac.in

Contents

1	Ove	erview and motivation	1							
2	Timelike entanglement									
	2.1	Small deformation limit	3							
	2.2	Comments on large deformation limit	5							
3	QFT observables									
	3.1	Flow central charge	6							
	3.2	Complexity	7							
	3.3	Remarks about large κ deformation	8							
4	Page charges and brane set up									
	4.1	Background RR fluxes and Page charges	9							
	4.2	More on probe D branes and couplings	10							
	4.3	Comments on large deformation limit	11							
5	Cor	ncluding remarks	12							

1 Overview and motivation

The integrable η -deformation [1]- [2] of various AdS supergravity backgrounds had witnessed an overwhelming and ever growing interest for the last one decade due to its several novel aspects. The η - deformed backgrounds¹ are primarily of interest because the associated sigma model is integrable [1]- [2], [6]. It has been an open question whether these backgrounds are still solutions of type IIB supergravity equations or not [7]- [14]. It took some time to realise that the answer is positive if the associated R-matrix [15]- [16] satisfies the unimodularity condition [17].

Based on these ideas, the authors of [18] considered a new class of R-matrices that satisfy the unimodularity condition when all the simple roots of the Dynkin diagram are fermionic. These R-matrices correspond to a particular q-deformation of the (super)isometry algebra [5], [19]. While the stringy counterpart of the duality is well established, the dual QFTs living on the holographic screen [20]- [21] is much less explored. It is quite intuitive to argue that constructing an extremal surface in the bulk, that ends on a timelike/spacelike subregion in the boundary, would shed light about the quantum entanglement in these special class of (0 + 1)d QFTs.

The purpose of this paper is to revisit the holographic aspects [22]- [23] of η -deformation in a greater detail. In particular, to calculate various field theory observables in the dual QFTs. This include, the computation of holographic entanglement entropy [24]- [26] following the recent proposal of timelike [27]- [35] hypersurfaces in a novel class of η -deformed backgrounds [18] in type IIB supergravity. The big picture here is to relate the quantum entanglement with

¹This is a generalisation of the η -deformed PCM [3]- [4] and the symmetric space model of [5].

various other field theory observables [36] namely the flow central charge [37]- [40] and the complexity [41]- [44], in a holographic set up pertaining to η -deformed superstring backgrounds.

We begin by reviewing the η -deformed $AdS_2 \times S^2 \times T^6$ background in type IIB [18]

$$ds_{10}^2 = ds_4^2 + ds_{T^6}^2 \tag{1}$$

$$ds_4^2 = \frac{L^2}{(1-\kappa^2\rho^2)} \Big(-(1+\rho^2)dt^2 + \frac{d\rho^2}{1+\rho^2} \Big) + \frac{L^2}{(1+\kappa^2r^2)} \Big((1-r^2)d\phi^2 + \frac{dr^2}{1-r^2} \Big)$$
(2)

$$ds_{T^6}^2 = L^2 d\varphi_i d\varphi_i \; ; \; \kappa = \frac{2\eta}{1 - \eta^2} \; ; \; i = 4, \cdots, 9$$
 (3)

where κ is the (real) deformation parameter that ranges between 0 to ∞ .

The above background (1)-(3) is supported by a dilaton of the form

$$e^{-2\Phi} = e^{-2\Phi_0} \frac{(1-\kappa^2 \rho^2)(1+\kappa^2 r^2)}{1-\kappa^2 (\rho^2 - r^2 - \rho^2 r^2)}.$$
(4)

In what follows, we adopt the following change of coordinates

$$\rho = \sinh \rho \; ; \; r = \cos \theta \tag{5}$$

and rewrite the metric (1)-(3) and the dilaton (4) in global coordinates as

$$ds_4^2 = \frac{L^2}{(1 - \kappa^2 \sinh^2 \varrho)} \Big(-\cosh^2 \varrho dt^2 + d\varrho^2 \Big) + \frac{L^2}{(1 + \kappa^2 \cos^2 \theta)} \Big(d\theta^2 + \sin^2 \theta d\phi^2 \Big)$$
(6)

$$e^{-2\Phi} = e^{-2\Phi_0} \frac{(1 - \kappa^2 \sinh^2 \varrho)(1 + \kappa^2 \cos^2 \theta)}{1 - \kappa^2 (\sinh^2 \varrho \sin^2 \theta - \cos^2 \theta)}.$$
(7)

The first term in (6) is the deformed AdS_2 where ρ is the global radial coordinate. As in the undeformed case, $\rho = \rho_{IR} = 0$ corresponds to the deep IR limit in the bulk. On the other hand, the UV limit corresponds to setting $\Lambda_{UV} = \sinh^{-1}\left(\frac{1}{\kappa}\right)$, which clearly goes to infinity in the limit of the vanishing deformation. We identify Λ_{UV} as the UV cut-off in the bulk, that sets the location of the holographic screen [20]- [21] where the dual (0 + 1)d QFT is living.

Given the above set up, we calculate various field theory observables in the following sections, which we list down below. In section 2, we revisit the RT prescription [24]- [26] of computing entanglement entropy and in particular, the recent proposal of computing timelike entanglement entropy (tEE) [27]- [35]. We provide an RG interpretation of our findings in terms of holographic c function [37]- [40] and complexity [41]- [44], that we compute in the following section 3. Next, in section 4, we explore different Page charges and the associated brane set up [45]. We also calculate the coupling constant in the dual QFTs by probing the geometry with D branes and in particular following the lines of [45]- [48]. Finally, we summarise all our findings in section 5 and conclude the paper along with some future remarks and key observations.

2 Timelike entanglement

Since the boundary theory is (0 + 1)d, therefore quantum entanglement in these QFTs corresponds to integrating out timelike subregions. On the dual gravitational counterpart, this corresponds to considering a class of extremal surfaces in the bulk that ends on a timelike surface in the boundary [27]- [35], which for the present case is the location of the holographic screen. To begin with, we propose $t = t(\varrho)$, which results in a co-dimension one surface

$$ds_9^2 = (g_{\varrho\varrho} - g_{tt}t'^2)d\varrho^2 + g_{\theta\theta}d\theta^2 + g_{\phi\phi}d\phi^2 + g_{\varphi_i\varphi_i}d\varphi_i^2 \tag{8}$$

where we identify the individual metric coefficients as

$$g_{tt} = \frac{L^2 \cosh^2 \varrho}{(1 - \kappa^2 \sinh^2 \varrho)} ; \ g_{\varrho\varrho} = \frac{L^2}{(1 - \kappa^2 \sinh^2 \varrho)}$$
(9)

$$g_{\theta\theta} = \frac{L^2}{(1+\kappa^2\cos^2\theta)} ; \ g_{\phi\phi} = \frac{L^2\sin^2\theta}{(1+\kappa^2\cos^2\theta)} ; \ g_{\varphi_i\varphi_i} = L^2.$$
(10)

The tEE is defined as the area of the co-dimension one hyper-surface, weighted by the dilaton

$$\mathcal{S}^{(tEE)} = \frac{V_T}{4G_N} \int_{S^2_{\kappa}} d\theta d\phi \sqrt{\det g_2} \int_0^{\Lambda_{UV}} d\varrho e^{-2\Phi} \sqrt{g_{\varrho\varrho} - g_{tt} t'^2}$$
(11)

where V_T represents the volume of the six torus (T^6) and det $g_2 = g_{\theta\theta}g_{\phi\phi}$ is the determinant of the metric on the deformed unit two sphere $(S^2_{\kappa}(\theta, \phi))$.

The equation of motion for $t(\varrho)$ yields

$$t'^{2}(\varrho) = \frac{C^{2}g_{\varrho\varrho}}{g_{tt}(C^{2} + e^{-4\Phi}g_{tt})}$$
(12)

where C is the constant of integration.

Substituting (12) into (11), one finds

$$\mathcal{S}^{(tEE)} = \frac{V_T}{4G_N} \int_{S^2_{\kappa}} d\theta d\phi \sqrt{\det g_2} \int_0^{\Lambda_{UV}} d\varrho \frac{e^{-4\Phi} \sqrt{g_{tt}g_{\varrho\varrho}}}{\sqrt{C^2 + e^{-4\Phi}g_{tt}}}.$$
(13)

2.1 Small deformation limit

Expanding in the limit of small deformation ($\kappa \ll 1$), we find

$$\mathcal{S}^{(tEE)} = \frac{\pi V_T L^4}{G_N} \int_0^\infty d\varrho \frac{e^{-4\Phi_0} \cosh \varrho}{\sqrt{C^2 + L^2 e^{-4\Phi_0} \cosh^2 \varrho}} - \frac{\pi \kappa^2 V_T L^4}{4G_N} \int_0^\pi d\theta \sin \theta \int_0^\infty d\varrho \frac{e^{-4\Phi_0} \cosh \varrho \Gamma(\theta, \varrho)}{\sqrt{C^2 + L^2 e^{-4\Phi_0} \cosh^2 \varrho}}$$
(14)

where we denote the above function as

$$\Gamma(\theta, \varrho) = \frac{1}{(C^2 e^{4\Phi_0} + L^2 \cosh^2 \varrho)} \Big[2C^2 e^{4\Phi_0} \left(\cos 2\theta \sinh^2 \varrho + \cos^2 \theta \right) \\ + L^2 \cosh^2 \varrho \left(\cos 2\theta \cosh^2 \varrho + 1 \right) \Big].$$
(15)

Performing the θ integral independently, one finds

$$\mathcal{S}^{(tEE)} = \frac{\pi V_T L^4}{G_N} \int_0^\infty d\varrho \frac{e^{-4\Phi_0} \cosh \varrho}{\sqrt{C^2 + L^2 e^{-4\Phi_0} \cosh^2 \varrho}} \left(1 + \frac{\kappa^2}{12} \bar{\Gamma}(\varrho)\right) \tag{16}$$

$$\bar{\Gamma}(\varrho) = \frac{2C^2 e^{4\Phi_0} (\cosh 2\varrho - 3) + L^2 \cosh^2 \varrho (\cosh 2\varrho - 5)}{(C^2 e^{4\Phi_0} + L^2 \cosh^2 \varrho)}.$$
(17)

In order to find out the turning point in the bulk one sets $C = i\tilde{C}$, such that $C^2 = -\tilde{C}^2 < 0$ [34]. On top of that, we introduce the following change of variables

$$\cosh \varrho = \frac{1}{z} \tag{18}$$

such that, z = 1 stands for the deep IR limit in the bulk. On the other hand, $z_{UV} = \epsilon \ll 1$ corresponds to the UV cut-off in the bulk. Using this change of variable (18), one obtains

$$\mathcal{S}^{(tEE)} = \frac{\pi V_T L^4}{G_N} \int_{\epsilon}^1 \frac{dz}{z^2} \frac{e^{-4\Phi_0}}{\sqrt{1-z^2}} \frac{1}{\sqrt{\frac{L^2}{z^2}e^{-4\Phi_0} - \tilde{C}^2}} \Big(1 + \frac{\kappa^2}{12}\bar{\Gamma}(z)\Big).$$
(19)

Clearly, the turning point $(z = z_0)$ is obtained by setting $\tilde{C} = \frac{L}{z_0}e^{-2\Phi_0}$, where $z_0 < 1$. The next step is to substitute it into (19), which finally reveals $\mathcal{S}^{(tEE)} = -iz_0\mathcal{S}^{(\text{Im})}$ where

$$\mathcal{S}^{(\mathrm{Im})} = \frac{2\pi V_T L^3}{G_N} \int_{z_0}^1 \frac{dz}{z} \frac{e^{-2\Phi_0}}{\sqrt{1-z^2}\sqrt{z^2-z_0^2}} \left(1 + \frac{\kappa^2}{12}\bar{\Gamma}(z)\right)$$
(20)

$$\bar{\Gamma}(z) = \frac{-8z^4 + 6z^2 z_0^2 + 4z^2 - 2z_0^2}{3z^2(z^2 - z_0^2)}.$$
(21)

Let us check asymptotic properties of (20). To begin with, we push the tip (z_0) of the extremal surface near UV, $z_0 \sim \epsilon \sim 0$. This yields the following expression for the tEE (20)

$$|\mathcal{S}^{(\mathrm{Im})}|_{z_0 \sim \epsilon} = \frac{2\pi V_T L^3}{G_N} e^{-2\Phi_0} \int_{\epsilon}^1 dz \frac{\kappa^2 + (9 - 2\kappa^2) z^2}{9z^4 \sqrt{1 - z^2}} = \frac{2\pi \kappa^2 V_T L^3}{27G_N \epsilon^3} e^{-2\Phi_0} \left(1 + \frac{9\epsilon^2}{\kappa^2} (6 - \kappa^2) + \cdots\right)$$
(22)

which clearly exhibits a pole of order 3, in limit $\epsilon \to 0$. Comparing with the undeformed ($\kappa = 0$) example, which reveals a pole of order one, one might therefore conclude that the tEE diverges at a faster rate (near the asymptotic infinity) in the presence of non zero η -deformation.

Using (22), we finally note down

$$\mathcal{S}^{(tEE)}|_{z_0 \sim \epsilon} = -\frac{2\pi i \kappa^2 V_T L^3}{27G_N \epsilon^2} e^{-2\Phi_0} \left(1 + \mathcal{O}(\epsilon^2/\kappa^2)\right)$$
(23)

which clearly exhibits a pole of order 2 near asymptotic infinity ($\epsilon \rightarrow 0$). As we show in the subsequent section, this is precisely the pole structure appearing in the holographic flow central charge (c_{flow}) near UV. Combing them, it is quite suggestive to argue that tEE is measure of degrees of freedom for (0 + 1)d QFTs that are dual to η -deformed superstring backgrounds.

On the other hand, considering a deep IR $(z_0 \sim 1)$ limit in the bulk, one finds

$$|\mathcal{S}^{(\mathrm{Im})}|_{z_0 \sim 1} = \frac{\pi V_T L^3}{18G_N} e^{-2\Phi_0} \Big[\frac{\kappa^2}{z_0^2} (1 - z_0^2) - 3(6 - \kappa^2) \log \left(\frac{\Lambda_{IR} z_0^2}{1 - z_0^2}\right) \Big]_{z_0 \sim 1}$$
(24)

where, $\Lambda_{IR} \sim 0$ is a small number such that the ratio $\frac{\Lambda_{IR}z_0^2}{1-z_0^2}$ is finite and $\mathcal{O}(1)$ in the deep IR limit. The first term, on the other hand, vanishes identically as we approach $z_0 \sim 1$.

Combining the above facts together, one might therefore argue that tEE (20) decreases in a RG flow from UV ($z_0 \sim 0$) to deep IR ($z_0 \sim 1$). This precisely reflects the properties of a cfunction [37]- [40] in a RG flow, which we further justify in the subsequent section. The UV ($z_0 \sim \epsilon \sim 0$) divergence is an artefact of the large area contribution due to the extremal surface. The corresponding subsystem length at the boundary turns out to be

$$t_{sub} = 2 \int_0^1 \frac{dz}{\sqrt{1-z^2}} = \pi.$$
(25)

2.2 Comments on large deformation limit

Let us briefly comment about the large $\kappa \gg 1$ regime, which corresponds to a holographic screen located at a finite radial cut-off, $\Lambda_{UV} = \frac{1}{\kappa} + \mathcal{O}(\kappa^{-3})$ in the bulk. The associated metric functions (9)-(10) and the background dilaton (7) read as

$$g_{tt} = -\frac{L^2 \coth^2 \varrho}{\kappa^2} \; ; \; g_{\varrho\varrho} = -\frac{L^2 \mathrm{csch}^2 \varrho}{\kappa^2} \tag{26}$$

$$g_{\theta\theta} = \frac{L^2 \sec^2 \theta}{\kappa^2} ; \ g_{\phi\phi} = \frac{L^2 \tan^2 \theta}{\kappa^2} ; \ g_{\varphi_i\varphi_i} = L^2$$
(27)

$$e^{-2\Phi} = -e^{-2\Phi_0} \left[\frac{\kappa^2}{\operatorname{csch}^2 \varrho - \tan^2 \theta} - \frac{\left(\sin^2 \theta \sinh^2 \varrho \left(\sinh^2 \varrho - \cos^2 \theta \right) + \cos^4 \theta \right)}{\left(\cos^2 \theta - \sin^2 \theta \sinh^2 \varrho \right)^2} \right].$$
(28)

Considering a large κ expansion of the entities in (13), we find at LO in $1/\kappa$

$$\mathcal{S}^{(tEE)} = \frac{-i\pi V_T L^3}{2G_N \kappa} e^{-2\Phi_0} \int_0^\pi \frac{\sin\theta}{\cos^2\theta} d\theta \int_0^{1/\kappa} \frac{d\varrho}{\sinh\varrho} \frac{1}{\left(\operatorname{csch}^2 \varrho - \tan^2\theta\right)}$$
(29)

which after performing the θ integral independently yields

$$\mathcal{S}^{(tEE)}|_{\kappa\gg1} = \frac{i\pi V_T L^3}{4G_N \kappa} e^{-2\Phi_0} \int_0^{1/\kappa} d\varrho \left(\log\left(e^{-3\varrho}\right) - \log\left(e^\varrho\right)\right) \operatorname{sech}\varrho$$
$$= -\frac{i\pi V_T L^3}{2G_N \kappa^3} e^{-2\Phi_0} + \cdots .$$
(30)

Clearly, tEE vanishes when the deformation is large enough ($\kappa \gg 1$). The vanishing entanglement (30) is an artefact of the vanishing area contribution due to the extremal surface in the bulk counterpart. One should think of this as a limiting situation in which the holographic screen is pushed deep inside the bulk and as a result the bulk entangling surface sizes to exist.

3 QFT observables

The purpose of this section is to calculate various observables in the dual QFT, that count the number of degrees of freedom in a RG flow from UV to deep IR. These are generally denoted in terms of the flow central charge [36]- [40] and the complexity [41]- [44].

3.1 Flow central charge

For generic QFTs, the c function or flow central charge (c_{flow}) counts the number of degrees of freedom that is monotonically decreasing in a RG flow, namely $\frac{dc_{flow}}{d\varrho} > 0$, where $0 < \varrho < \infty$. In case of a RG flow between two fixed points (UV and IR), the flow function (c_{flow}) precisely represents the corresponding central charges such that, $c_{UV} > c_{IR}$.

The calculation that we carry out, precisely follows the algorithm developed by authors in a series of papers [36]- [40]. To begin with, we re-express the type IIB background as

$$ds_{10}^2 = \frac{L^2}{(1 - \kappa^2 \sinh^2 \varrho)} \Big(-\cosh^2 \varrho dt^2 + d\varrho^2 \Big) + \mathcal{G}_{ab} d\omega^a d\omega^b \tag{31}$$

$$\mathcal{G}_{ab}d\omega^a d\omega^b = \frac{L^2}{(1+\kappa^2\cos^2\theta)} \left(d\theta^2 + \sin^2\theta d\phi^2\right) + L^2 d\varphi_i \varphi_i \tag{32}$$

where \mathcal{G}_{ab} is the metric of the eight dimensional internal manifold.

For a (0 + 1)d QFT, the flow central charge is given by the volume of the internal eight manifold weighted by the dilaton

$$c_{flow} = \frac{1}{G_N} \int d^8 \omega e^{-2\Phi} \sqrt{\det \mathcal{G}_{ab}}$$
$$= \frac{2\pi V_T}{G_N} \int_0^\pi d\theta e^{-2\Phi} \sqrt{\det g_2}.$$
(33)

Considering a small $\kappa \ll 1$ expansion, we find

$$c_{flow} = \frac{2\pi V_T L^2}{G_N} e^{-2\Phi_0} \int_0^\pi d\theta \sin\theta (1 - \kappa^2 \cos^2\theta \cosh^2\varrho).$$
(34)

Performing the θ integral independently and using (18), we find

$$|c_{flow}| = \frac{4\pi V_T L^2}{G_N} e^{-2\Phi_0} \Big| 1 - \frac{\kappa^2}{3z^2} \Big|.$$
(35)

Clearly, the flow central charge (35) diverges in the UV ($z \sim \epsilon \sim 0$). It decreases as we move in a RG flow from the UV asymptotic into the deep IR ($z \sim 1$). Taking a UV limit we find

$$|c_{flow}|_{z\sim\epsilon} = \frac{4\pi\kappa^2 V_T L^2}{3G_N \epsilon^2} e^{-2\Phi_0} \tag{36}$$

which reveals an identical pole structure as that of (23).

Comparing (23) and (36), we map these entities near the UV asymptotic

$$\mathcal{S}^{(tEE)}|_{UV} = -\frac{iL}{18}|c_{flow}|_{UV}.$$
(37)

3.2 Complexity

Complexity is yet another measure of number of degrees of freedom in a QFT. These degrees of freedom are represented by number of elementary gates that are required to build up a quantum circuit. In what follows, in our analysis, we follow the CV proposal that was outlined and successfully implemented in a series of papers [40]- [44]. The basic idea is to compute the volume of the nine manifold weighted by the dilaton, that is defined at a fixed boundary time.

We express the 10d metric (1) in the following form

$$ds_{10}^2 = -g_{tt}dt^2 + ds_9^2. aga{38}$$

Following the holographic prescription [40]- [44], complexity in the dual QFT is defined as the maximal volume associated with the spacelike hyper-surface at constant time, $t = t_0$

$$C_V = \frac{1}{G_N} \int d^9 x e^{-2\Phi} \sqrt{\det g_9}$$

= $c_{flow} \int_0^{\Lambda_{UV}} d\varrho \sqrt{g_{\varrho\varrho}}$
= $\frac{L}{8} c_{flow} \mathcal{F}(\Lambda_{UV})$ (39)

where we denote the above function as

$$\mathcal{F}(\Lambda_{UV}) = \frac{\kappa^2}{2} e^{2\Lambda_{UV}} - 2\left(\kappa^2 - 4\right) \Lambda_{UV}.$$
(40)

Using (37) and (39), we finally relate tEE and complexity in the UV limit of dual QFT

$$\mathcal{C}_V = \frac{9i}{4} \mathcal{S}^{(tEE)}|_{UV} \mathcal{F}(\Lambda_{UV}).$$
(41)

3.3 Remarks about large κ deformation

Let us now investigate above entities in the limit of large deformations ($\kappa \gg 1$). Expanding (33) in the large $\kappa \gg 1$) regime, we obtain the flow central charge as

$$c_{flow} = -\frac{2\pi V_T L^2}{G_N} e^{-2\Phi_0} \int_0^\pi d\theta \frac{\sin\theta \sinh^2\varrho}{(\cos^2\theta - \sin^2\theta \sinh^2\varrho)}$$
(42)

which by virtue of the θ integral yields

$$c_{flow}|_{\rho \sim \Lambda_{UV}} = \frac{4\pi V_T L^2}{G_N \kappa^2} e^{-2\Phi_0}.$$
(43)

Clearly, (43) is vanishingly small at LO in $1/\kappa$ expansion. In fact, there is no notion of RG flow in the large deformation limit. This stems from the fact that the holographic screen is pushed well inside the bulk interior and the corresponding RG flow sizes to exist.

A close comparison between (30) and (43) reveals

$$\mathcal{S}^{(tEE)}|_{\kappa\gg1} = -\frac{iL}{8\kappa} c_{flow}|_{\varrho\sim\Lambda_{UV}}.$$
(44)

On a similar note, we estimate complexity (39) in the limit of large deformation

$$C_V|_{\kappa\gg1} = c_{flow} \int_{\Lambda_{IR}}^{1/\kappa} d\varrho \sqrt{g_{\varrho\varrho}}$$
$$= \frac{L}{\kappa} c_{flow} \left(\log\left(\frac{1}{2\kappa\Lambda_{IR}}\right) - \frac{1}{12\kappa^3} \right)$$
(45)

where $\Lambda_{IR} \sim 0$ is some IR cut-off in the bulk such that $\kappa \Lambda_{IR} \sim \mathcal{O}(1)$.

Finally, comparing with (44), we relate complexity with tEE near the UV limit

$$\mathcal{C}_V|_{\kappa\gg1} = 8i\mathcal{S}^{(tEE)}|_{\kappa\gg1} \left(\log\left(\frac{1}{2\kappa\Lambda_{IR}}\right) - \frac{1}{12\kappa^3}\right).$$
(46)

4 Page charges and brane set up

The purpose of this section is to go one step further and construct the quantised Page charges associated with background RR fluxes. To do that, it is customary to note down background RR fields that couple to Dp branes. These background RR fluxes are constructed using R-matrix that satisfies unimodularity condition [18] and hence they solve a type II supergravity equation.

Let us briefly summarise what we find below. The η -deformed supergravity background possesses non zero Page charges for the D5 and D3 brane and yields a zero Page charge for NS5 branes. One could think of these D3 brane world-volume directions wrapping the three cycles of the six torus (T^6), that are localised along the bulk radial direction ($\rho = \rho_0$). On a similar token, D5 brane world-volume directions wrap the five cycles of the six torus (T^6) and are expanded along $R_t \times T^5$, as shown in the Table below. The dual QFT is what we identify as 1d defect [47] living inside a 4d QFT describing D3 branes. However, such an interpretation breaks down in the large deformation ($\kappa \gg 1$) limit, in which D3 branes sizes to exist.

Dp	t	ρ	r	ϕ	φ_4	φ_5	φ_6	φ_7	φ_8	$arphi_9$
D3	х				х	х	х			
D5	х				х	х	х	х	х	

4.1 Background RR fluxes and Page charges

The RR three form field strength is given by [18]

$$\mathcal{F}_{3} = -\mathcal{N}\kappa r (1 + \kappa^{2}r^{2})d\rho \wedge (d\varphi_{4} \wedge d\varphi_{5} + d\varphi_{6} \wedge d\varphi_{7} + d\varphi_{8} \wedge d\varphi_{9}) - \mathcal{N}\kappa \rho (1 - \kappa^{2}\rho^{2})dr \wedge (d\varphi_{4} \wedge d\varphi_{5} + d\varphi_{6} \wedge d\varphi_{7} + d\varphi_{8} \wedge d\varphi_{9}) + \mathcal{N}\kappa \rho (1 - r^{2})d\rho \wedge dt \wedge d\phi - \mathcal{N}\kappa r (1 + \rho^{2})dr \wedge dt \wedge d\phi$$
(47)

where the pre-factor can be expressed as

$$\mathcal{N} = \frac{\sqrt{1+\kappa^2}}{\sqrt{1-\kappa^2\rho^2}\sqrt{1+\kappa^2r^2(1-\kappa^2(\rho^2-r^2-\rho^2r^2))}}.$$
(48)

The Page charge counts the number of Dp branes that the background RR fluxes are coupled to. Given a (8 - p) background RR flux \mathcal{F}_{8-p} , this can be expressed as [45]- [48]

$$\mathcal{Q}_{Dp} = \frac{1}{(2\pi)^{7-p}} \int_{\Sigma^{(8-p)}} \mathcal{F}_{8-p}.$$
(49)

Setting t = constant, we have number of D5 branes

$$Q_{D5} = \frac{1}{(2\pi)^2} \int_{\Sigma^{(3)}} \mathcal{F}_3.$$
 (50)

Considering a small $\kappa \ll 1$ expansion, this further yields at LO

$$\mathcal{Q}_{D5} = -\frac{3\kappa}{(2\pi)^2} \Big[\int r d\rho \times (2\pi)^2 + \int \rho dr \times (2\pi)^2 \Big] + \mathcal{O}(\kappa^3)$$

$$= -3\kappa \int d(\rho r) + \mathcal{O}(\kappa^3)$$

$$= 3\kappa \rho_{max} + \mathcal{O}(\kappa^3)$$

$$= 3 + \mathcal{O}(\kappa^3)$$
(51)

where we set, $\rho_{max} = \frac{1}{\kappa} \gg 1$ as the location of the holographic screen.

RR five form field strength, on the other hand, is given by [18]

$$\mathcal{F}_5 = \mathcal{N}((1+\kappa^2 r^2)d\rho - \kappa^2 \rho r (1+\rho^2)dr) \wedge dt \wedge \operatorname{Re}\Omega_3 - \mathcal{N}(\kappa^2 \rho r (1-r^2)d\rho + (1-\kappa^2 \rho^2)dr) \wedge d\phi \wedge \operatorname{Im}\Omega_3$$
(52)

where $\Omega_3 = (d\varphi_4 - id\varphi_5) \wedge (d\varphi_6 - id\varphi_7) \wedge (d\varphi_8 - id\varphi_9)$. This clearly reveals that²

$$\operatorname{Im}\Omega_{3} = -d\varphi_{4} \wedge (d\varphi_{6} \wedge d\varphi_{9} + d\varphi_{7} \wedge d\varphi_{8}) - d\varphi_{5} \wedge (d\varphi_{6} \wedge d\varphi_{8} - d\varphi_{7} \wedge d\varphi_{9}).$$
(53)

Setting t = constant, we find the Page charge associated with D3 branes as

$$\begin{aligned} \mathcal{Q}_{D3} &= \frac{1}{(2\pi)^4} \int_{\Sigma^{(5)}} \mathcal{F}_5 \\ &= -\frac{1}{(2\pi)^4} \Big[\kappa^2 \int \rho r (1-r^2) d\rho \\ &+ \int \Big(1 + \frac{1}{2} \kappa^2 \left(\rho^2 - (2\rho^2 + 3) r^2 + 1 \right) \Big) dr \Big] \int_0^{2\pi} d\phi \int \mathrm{Im}\Omega_3 \\ &= -\frac{4}{(2\pi)^4} \Big[-2 + \frac{\kappa^2}{2} \int d \Big(r (\rho^2 + 1)(1-r^2) \Big) + \frac{\kappa^2}{2} \int \rho^2 r^2 dr \Big] \times (2\pi)^4 \\ &= 8 - 2\kappa^2 \rho_0^2 \int_1^{-1} r^2 dr \\ &= 8 + \frac{4}{3} \kappa^2 \rho_0^2 \end{aligned}$$
(54)

where D3 branes are considered to be localised along the bulk radial direction, $\rho = \rho_0$. The radial locations ρ_0 are discrete numbers, such that the entire Page charge is an integer.

Finally, we note down possibilities for any NS5 branes in the background. The corresponding Page charge can be expressed as the flux of background NS-NS three form

$$\mathcal{Q}_{NS5} = \frac{1}{(2\pi)^2} \int_{\Sigma^3} \mathcal{H}_3 = 0 \tag{55}$$

which vanishes identically by virtue of background NS-NS two form as found in [18].

4.2 More on probe D branes and couplings

To summarise, η -deformed AdS_2 superstring background contains D3 and D5 branes as a part of full type IIB solution. The background NS-NS fluxes, on the other hand, couple only to the F1 string and there are no NS5 branes present in the background. Below, we consider these D branes and their world-volume theories in the probe limit.

We begin by considering probe D3 brane in the background that is located at a fixed radial distance $\rho = \rho_0$ and wrapping the three cycle of the six torus (T^6). The corresponding DBI action could be formally expressed as

$$S_{D3} = T_{D3} \int d^4x \Big[e^{-\Phi} \sqrt{-\det(g + 2\pi\alpha' F_2)} - \mathcal{C}_4 \Big]$$
(56)

²While integrating over cycles $0 \le \varphi_i \le 2\pi$ of a six torus, each term contributes a factor of $(2\pi)^3$.

where $T_{D3} = \frac{1}{(2\pi)^3 g_s \alpha'^2}$. The background RR potential relevant to our analysis is given by [18]

$$C_4 = \frac{L^4 e^{-\Phi_0} \rho_0 \sqrt{1 + \kappa^2}}{\sqrt{1 - \kappa^2 (\rho_0^2 - r^2 - \rho_0^2 r^2)}} dt \wedge \text{Re}\Omega_3.$$
(57)

Turning on a world-volume field strength tensor $F_2 = F_{t\varphi} dt \wedge d\varphi$, that is considered to be small enough and thereby allowing a Taylor exampliant of the world-volume action [48], yields

$$e^{-\Phi}\sqrt{-\det(g+2\pi\alpha'F_2)} - \mathcal{C}_4 \simeq \frac{3\pi\alpha'^2}{2\rho_0}\mathcal{Q}_{D3}e^{-\Phi_0}F_{t\varphi}^2$$
 (58)

where, we set $r = \pm 1$ and take the boundary limit $\rho_0 \gg 1$, such that the number of D3 branes $N_{D3} = Q_{D3} \simeq \frac{4}{3} \kappa^2 \rho_0^2$ is kept finite in the limit, $\kappa \ll 1$.

Using (58) and integrating over the three cycles of the six torus, we finally obtain

$$S_{D3} = \frac{3\pi Q_{D3}}{2g_s \rho_0} e^{-\Phi_0} \int dt (\partial_t \Phi)^2 = \frac{1}{g_{eff}^2} \int dt (\partial_t \Phi)^2$$
(59)

where g_s is the string coupling and g_{eff}^2 is the effective coupling of (0 + 1)d QFT that is living inside the D3 brane world-volume. Here, we identify $A_{\varphi} = \Phi$ as the scalar of the reduced (effective) one dimensional theory living inside the D3 brane world-volume.

Clearly, in the near boundary limit $\rho_0 \to \infty$, the effective one dimensional theory becomes strongly coupled $(g_{eff}^2 \sim \rho_0 \to \infty)$, which we identify as the (0 + 1)d QFT that is dual to η deformed $AdS_2 \times S^2 \times T^6$ supergravity background. In what follows, we identify this reduced QFT as one dimensional defect within a four dimensional QFT inside D3 brane world-volume.

4.3 Comments on large deformation limit

As a final remark, we note down background RR fluxes in the large deformation ($\kappa \gg 1$) limit. Notice that, in the large deformation limit, the radial coordinate is bounded between $0 < \rho < \rho_{max}$ where $\rho_{max} \sim \frac{1}{\kappa} \sim 0$. In other words, $\kappa \rho < 1$ for the interval considered. On the other hand, one must consider $\kappa r \gg 1$, given the range $-1 \le r \le 1$.

Considering above facts, we note down components of background RR fluxes that are relevant for the computation of Page charges. To begin with, we note down the three form flux

$$\mathcal{F}_3 = -\kappa d\rho \wedge (d\varphi_4 \wedge d\varphi_5 + d\varphi_6 \wedge d\varphi_7 + d\varphi_8 \wedge d\varphi_9) + \mathcal{O}(1/\kappa) \tag{60}$$

which yields the Page charge associated with D5 branes as

$$\mathcal{Q}_{D5}\Big|_{\kappa\gg1} = -\frac{1}{(2\pi)^2} \int_{\Sigma^{(3)}} \mathcal{F}_3 = 3\kappa \int_0^{1/\kappa} d\rho = 3.$$
(61)

In other words, the number of D5 branes is preserved. This stems from the fact that the RR three form flux increases linearly with the deformation parameter ($\sim \kappa$), while on the other other hand, the radius decreases inversely as $\sim \frac{1}{\kappa}$, keeping the product finite.

On a similar note, for the five form flux we find (for constant t)

$$\mathcal{F}_5 = \left(1 - \frac{1}{r^2}\right) \rho d\rho \wedge d\phi \wedge \operatorname{Im}\Omega_3 + \mathcal{O}(1/\kappa^2)$$
(62)

which yields a vanishing flux in the limit of large deformations

$$\mathcal{Q}_{D3}\Big|_{\kappa\gg1} = -\frac{1}{(2\pi)^4} \int_{\Sigma^{(5)}} \mathcal{F}_5 = \frac{2}{\kappa^2} \left(1 - \frac{1}{r^2}\right) \simeq 0.$$
(63)

In other words, the number of D3 branes vanishes, which is consistent with the fact that the dual QFT looses all its degrees of freedom in the large deformation limit. D5 brane, on the other hand, wraps the six torus (T^6) and does not feel the radial shrinking. To summarise, one might therefore conclude that η - deformed backgrounds possess well defined dual description as long as the deformation parameter is small enough. In the limit of large deformations, the superstring theory looses its dual interpretation in the form QFTs living in (0 + 1)d.

5 Concluding remarks

We now summarise the key findings of the paper. In the first part of the paper, we compute timelike entanglement entropy (tEE) in the dual QFT₁ both in the limit of small ($\kappa \ll 1$) as well as large deformations ($\kappa \gg 1$). Since the boundary theory is (0 + 1)d, it is therefore expected that tEE is the only possible way of defining the entanglement between different (timelike) subsystems in the dual QFT₁. As our analysis reveals, for small deformations, the imaginary component of tEE precisely captures the notion of the flow central charge (c_{flow}), in the sense that it decreases in a RG flow from UV to deep IR. In other words, the imaginary tEE is a measure of the number of degrees of freedom in the dual QFT₁ living on the holographic screen. This result should follow from dimension reduction of a parent 2d QFT, where the flow central charge in (0+1)d would correspond to either of the left or right c functions of the parent QFT₂.

To be precise, we map these entities near the UV asymptotic (37). This follows quite naturally from the unique pole (of order 2) structure appearing in both the entities near the UV asymptotic. On a similar token, one could also relate holographic complexity and tEE near the UV scale of the dual QFT₁ (41). Combining all these three pieces together, we conclude that tEE ($\mathcal{S}^{(tEE)}$), flow central charge (c_{flow}) and complexity (\mathcal{C}_V) are all equally good measure of the degrees of freedom of the boundary QFT₁ near its UV limit.

We also explore the physics of dual QFT₁ in the limit of large deformations ($\kappa \gg 1$). It turns out that all three entities namely tEE ($\mathcal{S}^{(tEE)}$), flow central charge (c_{flow}) and complexity (\mathcal{C}_V) become vanishingly small. This stems from the fact that in the large deformation limit, the holographic screen is pushed well inside the bulk interior. As a result, one does not have much access to the bulk degrees of freedom as they are enormously reduced. This is reflected in the corresponding QFT observables that actually count these degrees of freedom. In this limit, the dual QFT₁ exhausts almost all its degrees of freedom and becomes non-dynamical.

We extend our analysis one step further by computing Page charges using background NS-

NS and RR fluxes. Our analysis reveals the existence of D3 and D5 branes and no NS5 branes as a part of full type IIB solution. This sort of completes the RR sector of η - deformed type IIB AdS_2 superstrings. The question that remains to be be answered is how to engineer a (0 + 1)dquantum mechanical model using these brane set up. We portray these QFT₁ as 1d defect inside a 4d QFT living within D3 brane world-volume. As a preliminary support to our conjecture, we compute the DBI action for N_{D3} probe D3 branes near the boundary, which eventually boils down into a (0 + 1)d QFT living inside D3 brane world-volume. We further show that the effective coupling of this reduced theory becomes very large indicating a strongly coupled theory. These could be further confirmed by computing various other QFT observables/probes for example, the Wilson line and t'Hooft loops [47], which we leave for future investigation.

Acknowledgements : The author is indebted to the authorities of IIT Roorkee for their unconditional support towards researches in basic sciences. The author also acknowledges The Royal Society, UK for financial assistance. The author also acknowledges the Mathematical Research Impact Centric Support (MATRICS) grant (MTR/2023/000005) received from ANRF, India.

References

- [1] F. Delduc, M. Magro and B. Vicedo, "An integrable deformation of the $AdS_5 \times S^5$ superstring action," Phys. Rev. Lett. **112**, no.5, 051601 (2014) doi:10.1103/PhysRevLett.112.051601 [arXiv:1309.5850 [hep-th]].
- [2] F. Delduc, M. Magro and B. Vicedo, "Derivation of the action and symmetries of the q-deformed $AdS_5 \times S^5$ superstring," JHEP **10**, 132 (2014) doi:10.1007/JHEP10(2014)132 [arXiv:1406.6286 [hep-th]].
- [3] C. Klimcik, "Yang-Baxter sigma models and dS/AdS T duality," JHEP 12, 051 (2002) doi:10.1088/1126-6708/2002/12/051 [arXiv:hep-th/0210095 [hep-th]].
- [4] C. Klimcik, "On integrability of the Yang-Baxter sigma-model," J. Math. Phys. 50, 043508 (2009) doi:10.1063/1.3116242 [arXiv:0802.3518 [hep-th]].
- [5] F. Delduc, M. Magro and B. Vicedo, "On classical q-deformations of integrable sigmamodels," JHEP 11, 192 (2013) doi:10.1007/JHEP11(2013)192 [arXiv:1308.3581 [hep-th]].
- [6] D. Roychowdhury, "Analytic integrability for strings on η and λ deformed backgrounds," JHEP **10**, 056 (2017) doi:10.1007/JHEP10(2017)056 [arXiv:1707.07172 [hep-th]].
- [7] G. Arutyunov, R. Borsato and S. Frolov, "S-matrix for strings on η-deformed AdS5 x S5," JHEP 04, 002 (2014) doi:10.1007/JHEP04(2014)002 [arXiv:1312.3542 [hep-th]].
- [8] G. Arutyunov, R. Borsato and S. Frolov, "Puzzles of η -deformed AdS₅× S⁵," JHEP **12**, 049 (2015) doi:10.1007/JHEP12(2015)049 [arXiv:1507.04239 [hep-th]].

- [9] G. Arutyunov, S. Frolov, B. Hoare, R. Roiban and A. A. Tseytlin, "Scale invariance of the η-deformed AdS₅ × S⁵ superstring, T-duality and modified type II equations," Nucl. Phys. B 903, 262-303 (2016) doi:10.1016/j.nuclphysb.2015.12.012 [arXiv:1511.05795 [hep-th]].
- [10] B. Hoare, R. Roiban and A. A. Tseytlin, "On deformations of $AdS_n \ge S^n$ supercosets," JHEP **06**, 002 (2014) doi:10.1007/JHEP06(2014)002 [arXiv:1403.5517 [hep-th]].
- [11] D. Roychowdhury, "On pp wave limit for η deformed superstrings," JHEP **05**, 018 (2018) doi:10.1007/JHEP05(2018)018 [arXiv:1801.07680 [hep-th]].
- [12] O. Lunin, R. Roiban and A. A. Tseytlin, "Supergravity backgrounds for deformations of $AdS_n \times S^n$ supercoset string models," Nucl. Phys. B **891**, 106-127 (2015) doi:10.1016/j.nuclphysb.2014.12.006 [arXiv:1411.1066 [hep-th]].
- [13] T. Araujo, E. Ó Colgáin, J. Sakamoto, M. M. Sheikh-Jabbari and K. Yoshida, "I in generalized supergravity," Eur. Phys. J. C 77, no.11, 739 (2017) doi:10.1140/epjc/s10052-017-5316-5 [arXiv:1708.03163 [hep-th]].
- [14] L. Wulff and A. A. Tseytlin, "Kappa-symmetry of superstring sigma model and generalized 10d supergravity equations," JHEP 06, 174 (2016) doi:10.1007/JHEP06(2016)174 [arXiv:1605.04884 [hep-th]].
- [15] V. G. Drinfeld, "Hopf algebras and the quantum Yang-Baxter equation," Sov. Math. Dokl. 32, 254-258 (1985)
- [16] M. Jimbo, "A q difference analog of U(g) and the Yang-Baxter equation," Lett. Math. Phys. 10, 63-69 (1985) doi:10.1007/BF00704588
- [17] R. Borsato and L. Wulff, "Target space supergeometry of η and λ -deformed strings," JHEP **10**, 045 (2016) doi:10.1007/JHEP10(2016)045 [arXiv:1608.03570 [hep-th]].
- [18] B. Hoare and F. K. Seibold, "Supergravity backgrounds of the η -deformed AdS₂ × S² × T^6 and AdS₅ × S⁵ superstrings," JHEP **01**, 125 (2019) doi:10.1007/JHEP01(2019)125 [arXiv:1811.07841 [hep-th]].
- [19] F. Delduc, S. Lacroix, M. Magro and B. Vicedo, "On q-deformed symmetries as Poisson-Lie symmetries and application to Yang-Baxter type models," J. Phys. A 49, no.41, 415402 (2016) doi:10.1088/1751-8113/49/41/415402 [arXiv:1606.01712 [hep-th]].
- [20] T. Kameyama and K. Yoshida, "A new coordinate system for q-deformed $AdS_5 \times S^5$ and classical string solutions," J. Phys. A **48**, no.7, 075401 (2015) doi:10.1088/1751-8113/48/7/075401 [arXiv:1408.2189 [hep-th]].
- [21] T. Kameyama and K. Yoshida, "Minimal surfaces in q-deformed $AdS_5 \times S^5$ with Poincare coordinates," J. Phys. A **48**, no.24, 245401 (2015) doi:10.1088/1751-8113/48/24/245401 [arXiv:1410.5544 [hep-th]].

- [22] D. Roychowdhury, "Probing η deformed backgrounds with Dp branes," Phys. Lett. B **778**, 167-173 (2018) doi:10.1016/j.physletb.2018.01.020 [arXiv:1706.02625 [hep-th]].
- [23] D. Roychowdhury, "Stringy correlations on deformed $AdS_3 \times S^3$," JHEP **03**, 043 (2017) doi:10.1007/JHEP03(2017)043 [arXiv:1702.01405 [hep-th]].
- [24] S. Ryu and T. Takayanagi, "Holographic derivation of entanglement entropy from AdS/CFT," Phys. Rev. Lett. 96, 181602 (2006) doi:10.1103/PhysRevLett.96.181602 [arXiv:hep-th/0603001 [hep-th]].
- [25] S. Ryu and T. Takayanagi, "Aspects of Holographic Entanglement Entropy," JHEP 08, 045 (2006) doi:10.1088/1126-6708/2006/08/045 [arXiv:hep-th/0605073 [hep-th]].
- [26] V. E. Hubeny, M. Rangamani and T. Takayanagi, "A Covariant holographic entanglement entropy proposal," JHEP 07, 062 (2007) doi:10.1088/1126-6708/2007/07/062 [arXiv:0705.0016 [hep-th]].
- [27] K. Doi, J. Harper, A. Mollabashi, T. Takayanagi and Y. Taki, "Pseudoentropy in dS/CFT and Timelike Entanglement Entropy," Phys. Rev. Lett. 130, no.3, 031601 (2023) doi:10.1103/PhysRevLett.130.031601 [arXiv:2210.09457 [hep-th]].
- [28] K. Doi, J. Harper, A. Mollabashi, T. Takayanagi and Y. Taki, "Timelike entanglement entropy," JHEP 05, 052 (2023) doi:10.1007/JHEP05(2023)052 [arXiv:2302.11695 [hep-th]].
- [29] Z. Li, Z. Q. Xiao and R. Q. Yang, "On holographic time-like entanglement entropy," JHEP 04, 004 (2023) doi:10.1007/JHEP04(2023)004 [arXiv:2211.14883 [hep-th]].
- [30] C. S. Chu and H. Parihar, "Time-like entanglement entropy in AdS/BCFT," JHEP 06, 173 (2023) doi:10.1007/JHEP06(2023)173 [arXiv:2304.10907 [hep-th]].
- [31] K. Narayan and H. K. Saini, "Notes on time entanglement and pseudo-entropy," Eur. Phys. J. C 84, no.5, 499 (2024) doi:10.1140/epjc/s10052-024-12855-x [arXiv:2303.01307 [hep-th]].
- [32] S. Grieninger, K. Ikeda and D. E. Kharzeev, "Temporal entanglement entropy as a probe of renormalization group flow," JHEP 05, 030 (2024) doi:10.1007/JHEP05(2024)030 [arXiv:2312.08534 [hep-th]].
- [33] W. z. Guo, Y. z. Jiang and Y. Jiang, "Pseudo entropy and pseudo-Hermiticity in quantum field theories," JHEP 05, 071 (2024) doi:10.1007/JHEP05(2024)071 [arXiv:2311.01045 [hep-th]].
- [34] M. Afrasiar, J. K. Basak and D. Giataganas, "Timelike entanglement entropy and phase transitions in non-conformal theories," JHEP 07, 243 (2024) doi:10.1007/JHEP07(2024)243 [arXiv:2404.01393 [hep-th]].
- [35] M. Afrasiar, J. K. Basak and D. Giataganas, "Holographic Timelike Entanglement Entropy in Non-relativistic Theories," [arXiv:2411.18514 [hep-th]].

- [36] D. Roychowdhury, "Holographic timelike entanglement and c theorem for supersymmetric QFTs in (0 + 1)d," [arXiv:2502.10797 [hep-th]].
- [37] N. T. Macpherson, C. Núñez, L. A. Pando Zayas, V. G. J. Rodgers and C. A. Whiting, "Type IIB supergravity solutions with AdS₅ from Abelian and non-Abelian T dualities," JHEP 02, 040 (2015) doi:10.1007/JHEP02(2015)040 [arXiv:1410.2650 [hep-th]].
- [38] Y. Bea, J. D. Edelstein, G. Itsios, K. S. Kooner, C. Nunez, D. Schofield and J. A. Sierra-Garcia, "Compactifications of the Klebanov-Witten CFT and new AdS₃ backgrounds," JHEP 05, 062 (2015) doi:10.1007/JHEP05(2015)062 [arXiv:1503.07527 [hep-th]].
- [39] P. Merrikin, C. Nunez and R. Stuardo, "Compactification of 6d N=(1,0) quivers, 4d SCFTs and their holographic dual Massive IIA backgrounds," Nucl. Phys. B 996, 116356 (2023) doi:10.1016/j.nuclphysb.2023.116356 [arXiv:2210.02458 [hep-th]].
- [40] D. Chatzis, A. Fatemiabhari, C. Nunez and P. Weck, "SCFT deformations via uplifted solitons," Nucl. Phys. B 1006, 116659 (2024) doi:10.1016/j.nuclphysb.2024.116659 [arXiv:2406.01685 [hep-th]].
- [41] L. Susskind, "Computational Complexity and Black Hole Horizons," Fortsch. Phys. 64, 24-43 (2016) doi:10.1002/prop.201500092 [arXiv:1403.5695 [hep-th]].
- [42] A. Reynolds and S. F. Ross, "Complexity in de Sitter Space," Class. Quant. Grav. 34, no.17, 175013 (2017) doi:10.1088/1361-6382/aa8122 [arXiv:1706.03788 [hep-th]].
- [43] D. Stanford and L. Susskind, "Complexity and Shock Wave Geometries," Phys. Rev. D 90, no.12, 126007 (2014) doi:10.1103/PhysRevD.90.126007 [arXiv:1406.2678 [hep-th]].
- [44] A. Fatemiabhari and C. Nunez, "From conformal to confining field theories using holography," JHEP 03, 160 (2024) doi:10.1007/JHEP03(2024)160 [arXiv:2401.04158 [hep-th]].
- [45] Y. Lozano, C. Nunez, A. Ramirez and S. Speziali, "New AdS_2 backgrounds and $\mathcal{N} = 4$ conformal quantum mechanics," JHEP **03**, 277 (2021) doi:10.1007/JHEP03(2021)277 [arXiv:2011.00005 [hep-th]].
- [46] C. Núñez, D. Roychowdhury, S. Speziali and S. Zacarías, "Holographic aspects of four dimensional N = 2 SCFTs and their marginal deformations," Nucl. Phys. B 943, 114617 (2019) doi:10.1016/j.nuclphysb.2019.114617 [arXiv:1901.02888 [hep-th]].
- [47] Y. Lozano, C. Nunez and A. Ramirez, "AdS₂ × S² × CY₂ solutions in Type IIB with 8 supersymmetries," JHEP **04**, 110 (2021) doi:10.1007/JHEP04(2021)110 [arXiv:2101.04682 [hep-th]].
- [48] Y. Lozano and C. Núñez, "Field theory aspects of non-Abelian T-duality and $\mathcal{N} = 2$ linear quivers," JHEP **05**, 107 (2016) doi:10.1007/JHEP05(2016)107 [arXiv:1603.04440 [hep-th]].