CONTRACTIVE PROJECTIONS ON H^p SPACES

XIANGDI FU, KUNYU GUO, AND DILONG LI

ABSTRACT. This paper investigates contractive projections on closed subspaces X of L^p with 0 . One of the main results states that, subject to certain mild conditions, every contractive projection <math>P on X preserving constants coincides with a conditional expectation on $L^{\infty} \cap P^{-1}(L^{\infty})$. It results in some interesting applications concerning contractive idempotent coefficient multipliers for analytic function spaces and translation-invariant subspaces of $L^p(G)$, where G is a compact Abelian group. Focusing specifically on descriptions of boundedness and contractivity of conditional expectations on the Hardy space $H^p(\mathbb{T})$ with $0 , we give a complete characterization of contractive idempotent coefficient multipliers for <math>H^p(\mathbb{T}^d)$ with $0 , which complements a remarkable result due to Brevig, Ortega-Cerdà, and Seip characterizing such multipliers on <math>H^p(\mathbb{T}^d)$ for $1 \le p \le \infty$.

1. INTRODUCTION

Let (Ω, Σ, μ) be a complete probability space. Suppose \mathcal{F} is a complete sub σ -algebra of Σ . The conditional expectation associated with \mathcal{F} is an operator

$$\mathbb{E}(\cdot|\mathcal{F}): L^1(\Omega, \Sigma, \mu) \to L^1(\Omega, \mathcal{F}, \mu); f \mapsto \mathbb{E}(f|\mathcal{F})$$

uniquely determined by the following identities:

$$\int_{F} \mathbb{E}(f|\mathcal{F}) d\mu = \int_{F} f d\mu, \quad \forall F \in \mathcal{F}.$$

Indeed, the existence and the uniqueness of $\mathbb{E}(f|\mathcal{F})$ follow from the Radon–Nikodym theorem. The conditional expectation $\mathbb{E}(\cdot|\mathcal{F})$ is a contractive linear projection from $L^1(\Omega, \Sigma, \mu)$ onto its closed subspace $L^1(\Omega, \mathcal{F}, \mu)$. Actually, it is also contractive on $L^p(\Omega, \Sigma, \mu)$ for all $1 \leq p \leq \infty$. In particular, when p = 2, $\mathbb{E}(\cdot|\mathcal{F})$ is the orthogonal projection from $L^2(\Omega, \Sigma, \mu)$ onto $L^2(\Omega, \mathcal{F}, \mu)$. The conditional expectation, introduced by A. Kolmogorov in 1933, is a fundamental concept in probability theory. We refer to [Du10] for further properties and applications of conditional expectation.

²⁰²⁰ Mathematics Subject Classification. Primary: 30H10, 46E30; Secondary: 30J05.

Key words and phrases. Hardy spaces, contractive projections, conditional expectations, coefficient multipliers, finite Blaschke product.

As well known, conditional expectations are contractive projections on $L^p(\Omega, \Sigma, \mu)$, $1 \leq p \leq \infty$, which leave constants invariant. In his seminal paper [Do65], Douglas showed that when p = 1, this operator-theoretic property characterizes conditional expectations. Later, Douglas's result was extended by Andô [An66] to the general L^p case with $p \neq 2$. The case p = 2 is eliminated since every orthogonal projection on $L^2(\Omega, \Sigma, \mu)$ is contractive. Below we reformulate their result as the Douglas-Andô theorem.

Theorem 1.1 (The Douglas-Andô Theorem). Let (Ω, Σ, μ) be a complete probability space. Suppose P is a contractive projection on $L^p(\Omega, \Sigma, \mu)$ satisfying $P\mathbf{1} = \mathbf{1}$. If $1 \leq p < \infty$ and $p \neq 2$, then P is a conditional expectation. If 0 , then <math>P is the identity.

As a modification of Theorem 1.1, they also characterized the contractive projection on $L^p(\Omega, \Sigma, \mu)$ which fails to satisfy $P\mathbf{1} = \mathbf{1}$ (the case p = 1 is due to Douglas and the case $p \neq 1$ is due to Andô), see [An66, Theorem 2].

The study of contractive projections is an important topic in Banach space theory. An interesting problem in this field is to characterize contractive projections on some classical function spaces. There is a lot of work concerning contractive projections on various Banach spaces X, including the spaces of continuous functions [FR82], vector-valued function spaces [Ray04], and some non-commutative Banach spaces [AF92,AR24]. We also refer to the excellent survey [Ran01] for historical remarks and a detailed bibliography on this topic. However, as mentioned in [LRR05], little is known about contractive projections on spaces of analytic functions, even when $X = H^p(\mathbb{T})$, which serves as the most classic example of analytic function spaces.

To study contractive projections on the Hardy spaces, we begin with two natural ways to produce contractive projections on $H^p(\mathbb{T})$ for $1 \leq p < \infty, p \neq 2$. The first involves restricting a conditional expectation $\mathbb{E}(\cdot|\mathcal{F})$ to $H^p(\mathbb{T})$. For this purpose, $\mathbb{E}(\cdot|\mathcal{F})$ should leave $H^p(\mathbb{T})$ invariant. Such a conditional expectation was fully characterized by Aleksandrov [Al86], who showed that a conditional expectation $\mathbb{E}(\cdot|\mathcal{F})$ leaves $H^p(\mathbb{T})$ invariant if and only if the σ -algebra \mathcal{F} is generated by an inner function vanishing at the origin. The second involves idempotent coefficient multipliers. Let Λ be a subset of $\mathbb{N}_0 = \{0, 1, 2, \ldots\}$ and let P_{Λ} be the associated idempotent coefficient multiplier densely defined on the space of analytic polynomials by

$$P_{\Lambda} : \sum_{n \ge 0} a_n z^n \mapsto \sum_{n \in \Lambda} a_n z^n.$$
(1.1)

It is known that P_{Λ} extends to a contraction on $H^p(\mathbb{T})$ if Λ consists of a single point or $\Lambda = \{km + r : m = 0, 1, ...\}$ for some integers $0 \leq r < k$. Additionally, based on the previous two methods, Lancien, Randrianantoanina, and Ricard [LRR05] constructed some other contractive projections on $H^p(\mathbb{T})$.

This paper, primarily motivated by [BOS21], investigates the connection between contractive projections, conditional expectations, and contractive idempotent coefficient multipliers in the framework of H^p spaces.

We first establish a general theorem concerning contractive projections on closed subspaces of L^p .

Theorem 1.2. Suppose that $0 , <math>p \notin 2\mathbb{N}$, where \mathbb{N} is the set of all positive integers. Let X be a closed subspace of $L^p(\Omega, \Sigma, \mu)$ containing the constants. If P is a contractive projection on X satisfying $P\mathbf{1} = \mathbf{1}$, then there exists a complete sub σ -algebra Σ' of Σ such that for every $f \in L^{\infty} \cap P^{-1}(L^{\infty})$, it holds that

$$Pf = \mathbb{E}(f|\mathbf{\Sigma}').$$

In particular, if $L^{\infty} \cap P^{-1}(L^{\infty})$ is dense in X, then P is a restriction of some conditional expectation.

Theorem 1.2 is stated in a general form. The following special case is of interest as it characterizes the restrictions of conditional expectations on certain subspaces of $L^p(\Omega, \Sigma, \mu)$.

Theorem 1.3. Suppose that $0 , <math>p \notin 2\mathbb{N}$, and let X be a closed subspace of $L^p(\Omega, \Sigma, \mu)$ such that $\mathbf{1} \in X$ and $X \cap L^{\infty}$ is dense in X. Let P be a contractive projection on X. Then P is a restriction of some conditional expectation to X if and only if $P\mathbf{1} = \mathbf{1}$ and P maps bounded functions to bounded functions.

The proof of Theorem 1.2 will be presented in Section 2, and an example will be provided to show that, in both Theorem 1.2 and Theorem 1.3, the condition $p \notin 2\mathbb{N}$ can not be omitted in general. We also illustrate some applications of Theorem 1.2 related to the idempotent coefficient multipliers for some typical function spaces, such as L^p -subspaces on compact Abelian groups, and weighted Hardy spaces. As regards idempotent coefficient multipliers, there exist some new fascinating behaviors in higher-dimensional case, as shown in [BOS21]. However in dimension d = 1, there seems to be of considerable consistency for generalized Bergman spaces on the closed unit disk (including the Hardy space), as shown in Proposition 2.7 of Section 2. It reads as follows. Suppose that $1 \leq p < \infty, p \neq 2$, the idempotent coefficient multiplier P_{Λ} defined by (1.1) extends to a contraction on the generalized Bergman space $L^p_a(d\mu)$ if and only if Λ is a single point or $\Lambda = (l + k\mathbb{Z}) \cap \mathbb{N}_0$ with $k \ge 1$, where μ is a rotation-invariant probability measure on $\overline{\mathbb{D}}$.

Let $0 , <math>p \notin 2\mathbb{N}$, and let P be a contractive projection on $H^p(\mathbb{T})$ such that $P\mathbf{1} = \mathbf{1}$ and $Pz^n \in H^\infty(\mathbb{T})$ for each $n \ge 1$. Applying Theorem 1.2 to the case $X = H^p(\mathbb{T})$, we conclude that every such contractive projection P is of the form

$$Pf = \mathbb{E}(f|\eta), \quad \forall f \in H^{\infty}(\mathbb{T}),$$

where η is an inner function vanishing at the origin. When $p \geq 1$, the conditional expectation $\mathbb{E}(\cdot|\eta)$ is always contractive on $H^p(\mathbb{T})$. In contrast, when 0 , $<math>\mathbb{E}(f|\eta)$ is only well-defined for integrable functions f and the operator $\mathbb{E}(\cdot|\eta)$ may be unbounded in $H^p(\mathbb{T})$. The following theorem, which will be proved in Section 3, examines when the conditional expectation $\mathbb{E}(\cdot|\eta)$ can be boundedly extended to $H^p(\mathbb{T})$ for 0 .

Theorem 1.4. For $0 , the conditional expectation <math>\mathbb{E}(\cdot|\eta)$ is bounded on $H^p(\mathbb{T})$ if and only if η is a finite Blaschke product. Furthermore, for a finite Blaschke product η with $\eta(0) = 0$, we have

$$\|\mathbb{E}(\cdot|\eta)\|_{H^p} = \|\mathbb{E}(\cdot|\eta)\|_{L^p} = \|\eta'\|_{\infty}^{\frac{1}{p}-1}.$$

The proof of Theorem 1.4 is based on a pointwise representation of conditional expectations deduced by the Arc Length Formula. Note that if η is a finite Blaschke product with $\eta(0) = 0$, then $\|\eta'\|_{\infty} \leq 1$ if and only if $\eta = cz$ for some unimodular constant c. As a consequence, when $0 , the identity is the only contractive projection on <math>H^p(\mathbb{T})$ satisfying $P\mathbf{1} = \mathbf{1}$ and $Pz^n \in H^{\infty}(\mathbb{T})$ for all $n \geq 1$.

For $1 \leq d \leq \infty$, we denote by \mathbb{T}^d the *d*-dimensional (countably infinite if $d = \infty$) torus equipped with the normalized Haar measure m_d . For $0 , the Hardy space <math>H^p(\mathbb{T}^d)$ is defined as the closure of analytic polynomials in $L^p(\mathbb{T}^d)$. In Section 4 we apply results from Sections 2 and 3 to characterize the contractive idempotent coefficient multipliers on $H^p(\mathbb{T}^d)$ for $0 , <math>1 \leq d \leq \infty$. Let $\mathbb{Z}^d = \bigoplus_{j=1}^d \mathbb{Z}$ be the dual group of \mathbb{T}^d and let $\mathbb{N}_0^d = \bigoplus_{j=1}^d \mathbb{N}_0$ be the positive cone of \mathbb{Z}^d . For a nonempty subset Λ of \mathbb{N}_0^d , the associated idempotent coefficient multiplier P_Λ is defined on the space of analytic polynomials by

$$P_{\Lambda}: \sum_{\alpha \in \mathbb{N}_0^d} c_{\alpha} z^{\alpha} \mapsto \sum_{\alpha \in \Lambda} c_{\alpha} z^{\alpha}.$$
(1.2)

Theorem 1.5. Suppose that $0 and <math>1 \leq d \leq \infty$. Then P_{Λ} defined by (1.2) extends to a contraction on $H^p(\mathbb{T}^d)$ if and only if there exists a subset

$$J \subseteq \{1, 2, \dots, d\} \ (J \subseteq \{1, 2, \dots\} \ if \ d = \infty) \ , \ such \ that$$
$$\Lambda = \{\alpha \in \mathbb{N}_0^d : \alpha_j = 0 \ \text{for} \ j \in J\}.$$

This complements a remarkable result of Brevig, Ortega-Cerdà and Seip [BOS21] which characterizes such multipliers on $H^p(\mathbb{T}^d)$ for $1 \le p \le \infty$ and $1 \le d \le \infty$.

Notations. In this paper, (Ω, Σ, μ) always denotes a complete probability space. The completeness means Σ contains all μ -null sets and their subsets. For a family \mathcal{G} of measurable functions on $(\Omega, \Sigma, \mu), \Sigma(\mathcal{G})$ denotes the smallest complete sub σ -algebra that makes every function in \mathcal{G} measurable. For simplicity, if $\mathcal{G} = \{g_1, g_2, \ldots\}$ is countable, we write $\Sigma(g_1, g_2, \ldots)$ instead of $\Sigma(\{g_1, g_2, \ldots\})$. For a Σ -measurable function η , we write

$$\mathbb{E}(\cdot|\eta) := \mathbb{E}(\cdot|\boldsymbol{\Sigma}(\eta)).$$

Multi-indices are used in the usual way. For example, for $z = (z_1, z_2, \ldots, z_d) \in \mathbb{T}^d$ and $\gamma = (\gamma_1, \gamma_2, \ldots, \gamma_d) \in \mathbb{Z}^d$, we define

$$z^{\gamma} := \prod_{k=1}^d z_k^{\gamma_k}.$$

2. Contractive projections on closed subspaces of L^p

In this section, the space $L^p(\Omega, \Sigma, \mu)$ will be simply denoted by L^p . To prove Theorem 1.2, we start with the following standard variation lemma.

Lemma 2.1. If $f, g \in L^{\infty}$, and $1/g \in L^{\infty}$, then for 0 the function $<math>\varphi: z \mapsto \int_{\Omega} |g + zf|^p d\mu, \quad z \in \mathbb{C},$

is differentiable in an open neighborhood of the origin and

$$\frac{\partial \varphi}{\partial x}(0) = p \int_{\Omega} |g|^{p-2} \operatorname{Re}(\bar{f}g) d\mu, \quad \frac{\partial \varphi}{\partial y}(0) = p \int_{\Omega} |g|^{p-2} \operatorname{Im}(\bar{f}g) d\mu.$$

Recall that a linear operator P on a Banach space (or more generally, a quasi-Banach space) X is a contractive projection if $||P|| \leq 1$ and $P^2 = P$. Now suppose that P is a contractive projection and $f \in \text{Ker } P, g \in \text{Ran } P$. Then $||g||_p = ||P(g + zf)||_p \leq ||g + zf||_p$ holds for all $z \in \mathbb{C}$. Thus the function

$$\varphi: z \mapsto \int_{\Omega} |g+zf|^p d\mu, \quad z \in \mathbb{C},$$

achieves its minimum value at z = 0. If we further assume that both f and g satisfy the conditions of Lemma 2.1, then

$$\int_{\Omega} |g|^{p-2} \bar{f}g d\mu = 0.$$
(2.1)

The identity (2.1) is important for the contractive projection problem. We remark that when p > 1, the identity (2.1) holds for all $f \in \text{Ker } P$ and $g \in \text{Ran } P$ [Sha71, Section 4.2]. However, to get (2.1) when p = 1, some additional conditions on fand g must be assumed, see [KR65]. Nevertheless, the conditions $f, g \in L^{\infty}$ and $1/g \in L^{\infty}$ are enough to guarantee that the identity (2.1) holds for all p > 0. The proof of Lemma 2.1 involves a straightforward calculation, with details similar to those in [KR65].

We also need the following simple fact concerning the σ -algebra generated by a family of functions.

Lemma 2.2. Let \mathcal{G} be a family of measurable functions on (Ω, Σ, μ) . Denote by $\Sigma(\mathcal{G})$ the σ -algebra generated by \mathcal{G} . For any $A \in \Sigma(\mathcal{G})$, there exist countably many functions $\{g_i\}_{i=1}^{\infty}$ in \mathcal{G} , such that $A \in \Sigma(g_1, g_2, \ldots)$.

Proof. We present a proof for the sake of completeness. Consider the union

$$\mathcal{F}:=igcup_{\substack{\mathcal{C}\subseteq\mathcal{G},\ \mathcal{C} ext{ is countable}}}\Sigma(\mathcal{C})$$

The result follows once if we show $\mathcal{F} = \Sigma(\mathcal{G})$. Since $\mathcal{F} \subseteq \Sigma(\mathcal{G})$, it is enough to show \mathcal{F} is a complete σ -algebra. Clearly, \mathcal{F} is closed under complement and contains all null sets. If $A \in \Sigma(\mathcal{C})$ and $B \in \Sigma(\mathcal{C}')$, then $A \cap B \in \Sigma(\mathcal{C} \cup \mathcal{C}') \subseteq \mathcal{F}$. Similarly, if $A_n \in \Sigma(\mathcal{C}_n)$ for $n \ge 1$, then $\bigcup_{n\ge 1} A_n \in \Sigma(\bigcup_{n\ge 1} \mathcal{C}_n)$. Thus \mathcal{F} is also closed under intersection and countable unions. This completes the proof. \Box

Now we are ready to prove Theorem 1.2.

Proof of Theorem 1.2. Suppose that $\{f_k : k \ge 1\}$ is a countable subset of $L^{\infty} \cap P^{-1}(L^{\infty})$, and $g_k = Pf_k \in L^{\infty}$. Since P is a contractive projection and $P\mathbf{1} = \mathbf{1}$, for any given positive integer n, complex numbers w_1, w_2, \ldots, w_n , and any fixed k with $1 \le k \le n$ we have

$$\int_{\Omega} \left| 1 + \sum_{j=1}^{n} w_j g_j + z g_k \right|^p d\mu \le \int_{\Omega} \left| 1 + \sum_{j=1}^{n} w_j g_j + z f_k \right|^p d\mu.$$

Since $g_j \in L^{\infty}$, there exists a $\delta > 0$ such that $1 + \sum_{j=1}^{n} w_j g_j$ is invertible in L^{∞} when $|w_j| \leq \delta, 1 \leq j \leq n$. It follows from Lemma 2.1 and (2.1) that

$$\int_{\Omega} |1 + \sum_{j=1}^{n} w_j g_j|^{p-2} (1 + \sum_{j=1}^{n} w_j g_j) (\bar{f}_k - \bar{g}_k) d\mu = 0.$$
 (2.2)

Since $p \notin 2\mathbb{N}$, we have the following infinite series expansion

$$\left|1 + \sum_{j=1}^{n} w_j g_j\right|^{p-2} \left(1 + \sum_{j=1}^{n} w_j g_j\right) = \left(1 + \sum_{j=1}^{n} w_j g_j\right)^{p/2} \left(1 + \sum_{j=1}^{n} \bar{w}_j \bar{g}_j\right)^{p/2-1}$$

$$= \sum_{\gamma} \sum_{\ell} {p/2 \choose \gamma} {p/2-1 \choose \ell} g^{\gamma} \bar{g}^{\ell} w^{\gamma} \bar{w}^{\ell}.$$
(2.3)

Here, the multi-indices $\gamma = (\gamma_1, \ldots, \gamma_n)$ and $\ell = (\ell_1, \ldots, \ell_n)$ run over \mathbb{N}_0^n and

$$\binom{x}{\gamma} = \frac{\Gamma(x+1)}{\Gamma(\gamma_1+1)\cdots\Gamma(\gamma_n+1)\Gamma(x-|\gamma|+1)}, \quad x \notin \mathbb{Z}$$

where $|\gamma| := \gamma_1 + \cdots + \gamma_n$. Because each g_j is bounded for $1 \leq j \leq n$, we can choose δ to be small enough to ensure the series (2.3) converges uniformly on $w \in \delta \overline{\mathbb{D}}^n$ for almost all $x \in \Omega$. Multiplying $\overline{f}_k - \overline{g}_k$ in two sides of (2.3) and then integrating with respect to $x \in \Omega$, it follows from Equation (2.2) that

$$\int_{\Omega} g^{\gamma} \bar{g}^{\ell} (\bar{f}_k - \bar{g}_k) d\mu = 0 \quad \forall \gamma, \ell \in \mathbb{N}_0^n.$$
(2.4)

Define $\Delta = \{ w \in \mathbb{C}^n : |w_j| \le ||g_j||_{\infty}, 1 \le j \le n \}$ and

$$G: \Omega \to \Delta; x \mapsto (g_1(x), g_2(x), \dots, g_n(x))$$

Then the map

$$\Lambda: q \mapsto \int_{\Omega} q \circ G \cdot (\bar{f}_k - \bar{g}_k) d\mu$$

is a bounded linear functional on $C(\Delta)$. We deduce from (2.4) and the Stone-Weierstrass theorem that Λ is zero. As a consequence, the push-forward measure $G_*((\bar{f}_k - \bar{g}_k)d\mu)$ is zero. Thus for any Borel subset $B \subseteq \Delta$ it holds that

$$\int_{G^{-1}(B)} (\bar{f}_k - \bar{g}_k) d\mu = \int_{\Omega} \mathbf{1}_B \circ G \cdot (\bar{f}_k - \bar{g}_k) d\mu = 0.$$
(2.5)

Clearly, each $A \in \Sigma(g_1, g_2, \ldots, g_n)$ is of the form $A = G^{-1}(B) \triangle C$, where B is a Borel subset of Δ and C is a μ -null set. So the identity (2.5) implies $\mathbb{E}((\bar{f}_k - \bar{g}_k) | \Sigma(g_1, g_2, \ldots, g_n)) = 0$, and hence

$$\mathbb{E}(f_k|\boldsymbol{\Sigma}(g_1,g_2,\ldots,g_n)) = g_k, \quad 1 \le k \le n.$$

Letting n tend to the infinity, an application of Doob's martingale convergence theorem yields

$$\mathbb{E}(f_k|\boldsymbol{\Sigma}(g_1, g_2, \ldots)) = g_k, \quad k \ge 1.$$

Now let Σ' be the complete sub σ -algebra of Σ generated by all functions in $P(L^{\infty} \cap P^{-1}(L^{\infty}))$. We claim that for each $f \in L^{\infty} \cap P^{-1}(L^{\infty})$ it holds that

$$Pf = \mathbb{E}(f|\mathbf{\Sigma}').$$

In the virtual of Lemma 2.2, for any $E \in \Sigma'$, there exist countably many functions $\{g_k : k \geq 1\}$ in $P(L^{\infty} \cap P^{-1}(L^{\infty}))$, such that $E \in \Sigma(g_1, g_2, \ldots)$. In particular,

 $E \in \Sigma(g_0, g_1, g_2, \ldots)$, where $g_0 := Pf$. By our previous reasoning,

$$\mathbb{E}(f|\boldsymbol{\Sigma}(g_0,g_1,g_2,\ldots)) = Pf.$$

The definition of conditional expectation gives

$$\int_E f d\mu = \int_E P f d\mu$$

Since $E \in \Sigma'$ is arbitrary, we obtain $Pf = \mathbb{E}(f|\Sigma')$. The proof is completed. \Box

We remark that Theorem 1.2 is not true in general when $p = 2k, k \in \mathbb{N}$. An example is a closed subspace of $L^{2k}(\mathbb{T})$.

Example 2.3. Let p = 2k be an even positive integer and let X be the closed subspace of $L^{2k}(\mathbb{T})$ spanned by $\{1, z, z^{k+1}, z^{k+2}, \ldots\}$. Let P be the projection on X defined by

$$P: a + bz + z^{k+1}p(z) \mapsto a + bz.$$

For any $a, b \in \mathbb{C}$ and any analytic polynomial p, one has

$$(a + bz + z^{k+1}p(z))^k = (a + bz)^k + z^{k+1}q(z)$$

so that

$$\begin{aligned} \|a + bz\|_{2k} &= \|(a + bz)^k\|_2^{1/k} \\ &\leq \|(a + bz)^k + z^{k+1}q(z)\|_2^{1/k} \\ &= \|(a + bz + z^{k+1}p(z))^k\|_2^{1/k} \\ &= \|a + bz + z^{k+1}p(z)\|_{2k} \end{aligned}$$

Thus P extends to a contractive projection on X. But P is not a restriction of any conditional expectation. Indeed, if $P = \mathbb{E}(\cdot|\mathcal{F})$, then the coordinate function z is \mathcal{F} -measurable, and hence \mathcal{F} consists of all Lebesgue measurable sets. This forces P to be the identity.

Theorem 1.2 exhibits considerable generality as it can be applied to various closed subspaces of L^p . In what follows we provide several applications of Theorem 1.2 concerning contractive idempotent coefficient multipliers on some typical function spaces.

The first example involves function spaces on compact Abelian groups.

Definition 2.4. Let G be a compact Abelian group, endowed with the normalized Haar measure μ , and let Γ denote its dual group. Suppose Λ is a non-empty subset of Γ . The idempotent coefficient multiplier associated with Λ , denoted by P_{Λ} , is defined for trigonometric polynomials as follows:

$$P_{\Lambda}: \sum_{\gamma \in \Gamma} c_{\gamma} \gamma(x) \mapsto \sum_{\gamma \in \Lambda} c_{\gamma} \gamma(x).$$

Such idempotent coefficient multipliers are closely related to idempotent measures on G which constitutes an intriguing topic [Rud90, Chapter 3]. It is well known that every coset Λ of Γ induces a contractive idempotent coefficient multipliers P_{Λ} on $L^{p}(G)$ for $p \geq 1$. Suppose that Γ_{0} is a subgroup of Γ , and let Γ_{0}^{\perp} be its annihilator, a closed subgroup of G, defined by

$$\Gamma_0^{\perp} = \{ x \in G : \gamma(x) = 1, \, \forall \gamma \in \Gamma_0 \}.$$

Set $\Lambda = \gamma' + \Gamma_0$, and write $H = \Gamma_0^{\perp}$. Denote by μ_H the normalized Haar measure of H. Then it follows from the duality relation between H and Γ/Γ_0 that

$$\mathbf{1}_{\Lambda}(\gamma) = \mathbf{1}_{\Gamma_0}(\gamma - \gamma') = \int_G \gamma(x)\overline{\gamma'}(x)d\mu_H(x)$$

An application of the Fourier transform shows that for any trigonometric polynomial f,

$$P_{\Lambda}f = (\gamma' d\mu_H) * f.$$

For $p \geq 1$, P_{Λ} extends to a contraction since

$$||P_{\Lambda}f||_{p} = ||(\gamma'd\mu_{H}) * f||_{p} \le ||\gamma'd\mu_{H}|| \, ||f||_{p} = ||f||_{p}.$$

The above argument also applies to translation-invariant subspaces of $L^p(G)$. It is easy to prove that any translation-invariant subspace is spanned by some characters of G. Suppose that $p \geq 1$ and S is a non-empty subset of Γ . We denote by $L^p_S(G)$ the closed subspace of $L^p(G)$ spanned by $\{\gamma : \gamma \in S\}$. For a function $f \in L^p(G)$, f belongs to $L^p_S(G)$ if and only if the Fourier transform of f vanishes outside S. The above reasoning also shows that if Λ is a coset of Γ , then $P_{\Lambda \cap S}$ is contractive on $L^p_S(G)$.

It is interesting to consider the converse of the above statement. This is to determine when $L_S^p(G)$ $(p \neq 2)$ possesses the so-called *contractive restriction* property. This property means that every subset A of S for which P_A is contractive on $L_S^p(G)$ must be of the form $A = \Lambda \cap S$ where Λ is a coset. When S = Γ , $L^p(G)$ enjoys the contractive restriction property, which can be deduced by suitably combining arguments and results due to Rudin [Rud90, Section 3.2] and Andô [An66]. Recently, Brevig, Ortega-Cerdà and Seip [BOS21] presented a distinct method to establish the contractive restriction property for $L^p(\mathbb{T}^d)$ via developing test function inequalities and a generalized Euclidean algorithm, which also fits to $L^p(G)$. In general, it remains a highly difficult problem to characterize whether $L_S^p(G)$ is of the contractive restriction property. In the case $G = \mathbb{T}^d$, $\Gamma = \mathbb{Z}^d$, and $S = \mathbb{N}_0^d$, the associated subspace of $L^p(\mathbb{T}^d)$ is the Hardy space $H^p(\mathbb{T}^d)$. When $p \in 2\mathbb{N}$, $p \neq 2$, and $d \geq 3$, $H^p(\mathbb{T}^d)$ no longer enjoy the contractive restriction property in general, and in these situations, the contractive projection set Λ exhibits some fascinating geometric behaviors, as shown by [BOS21]. While $p \notin 2\mathbb{N}$, one of the main results in [BOS21] states that $H^p(\mathbb{T}^d)$ enjoys the contractive restriction property.

Next, we generalize this result for $H^p(\mathbb{T}^d)$ $(p \notin 2\mathbb{N})$ to the case $L^p_S(G)$ by applying Theorem 1.2.

Theorem 2.5. Let G be a compact Abelian group with dual group Γ , and let S be a nonempty subset of Γ . If $1 \leq p < \infty$ and $p \notin 2\mathbb{N}$, then for any nonempty $\Lambda \subseteq S$ the following statements are equivalent:

- (i) P_{Λ} extends to a contraction on $L_{S}^{p}(G)$;
- (ii) Λ is a restriction of a coset to S;
- (iii) there exists an idempotent measure ν on G (i.e. $\nu * \nu = \nu$), with $\|\nu\| = 1$ such that $P_{\Lambda}f = \nu * f$, for any $f \in L^p_S(G)$.

Proof. The implication "(ii) \Rightarrow (iii) \Rightarrow (i)" can be deduced in a manner similar to the previous discussion. We next prove that (i) implies (ii). Fix a $\gamma' \in \Lambda$. We define

$$Pf = \overline{\gamma'} \cdot P_{\Lambda} (\gamma' \cdot f).$$

It is easy to verify that P is a contractive projection on $X := L^p_{S-\gamma'}(G)$ and $P\mathbf{1} = \mathbf{1}$. By Theorem 1.2, $Pf = \mathbb{E}(f|\mathcal{F})$ for $f \in X$ where \mathcal{F} is the sub σ -algebra generated by functions in

$$\operatorname{Ran} P = \overline{\operatorname{span}} \{ \gamma : \gamma \in \Lambda - \gamma' \}.$$
(2.6)

Let

$$\Delta := \{ \gamma \in \Gamma : \gamma \text{ is } \mathcal{F} - \text{measurable} \}.$$
(2.7)

Clearly, Δ is a subgroup of Γ . For $\gamma \in S - \gamma'$, it follows from $P\gamma = \mathbb{E}(\gamma | \mathcal{F})$ that $\gamma \in \operatorname{Ran} P$ is and only if γ is \mathcal{F} -measurable. Combining this with (2.6) and (2.7) we obtain

$$(S - \gamma') \cap (\Lambda - \gamma') = (S - \gamma') \cap \Delta.$$

This implies

$$\Lambda = (\gamma' + \Delta) \cap S$$

as desired.

In general, Theorem 2.5 is false if we omit the condition $p \notin 2\mathbb{N}$. Example 2.3 provides an contractive idempotent coefficient multipliers P_{Λ} on $L_S^{2k}(\mathbb{T})$ where $\Lambda = \{0, 1\}$ and $S = \{0, 1, k, k+1, \ldots\}$. It is clear that Λ is not a restriction of any

coset. For this reason, the case $p \in 2\mathbb{N}$ is particularly subtle and requires further separate consideration. The case where $G = \mathbb{T}^d$ and $S = \mathbb{N}_0^d$ was conclusively solved by Brevig, Ortega-Cerdà and Seip [BOS21, Theorem 1.2]. Their results show that when $p = 2k \ge 4$ whether $H^{2k}(\mathbb{T}^d)$ enjoys the contractive restriction property depends not only on k but also on the dimension d. Actually, they provided an effective algorithm [BOS21, Theorem 1.3] to characterize contractive idempotent coefficient multipliers on $H^{2k}(\mathbb{T}^d)$.

The second example focuses on the weighted Hardy space $H_K^p(\mathbb{T}^d)$ for $0 and <math>1 \leq d \leq \infty$. For a non-negative function K satisfying $K \in L^1(\mathbb{T}^d)$ and $\log K \in L^1(\mathbb{T}^d)$, the weighted Hardy space $H_K^p(\mathbb{T}^d)$ is defined as the closure of all analytic polynomials in $L^p(\mathbb{T}^d, Kdm_d)$, where m_d is the normalized Haar measure of \mathbb{T}^d . It is shown in [GNZ24, Proposition 3.2] that $H_K^p(\mathbb{T}^d)$ is contained in the Nevanlinna-Smirnov class $N_*(\mathbb{T}^d)$ and every function f in $H_K^p(\mathbb{T}^d)$ uniquely extends to an analytic function on \mathbb{D}^d (or $\mathbb{D}^\infty \cap \ell^1$ if $d = \infty$) admitting a power series expansion

$$f(z) = \sum_{\alpha \in \mathbb{N}_0^d} c_\alpha z^\alpha,$$

for which the coefficients c_{α} are uniquely determined by $f \in H^p_K(\mathbb{T}^d)$ [GNZ24]. With these notations, the associated result reads as follows.

Theorem 2.6. Suppose that $0 , <math>p \notin 2\mathbb{N}$. Let Λ be a nonempty subset of \mathbb{N}_0^d and denote by

$$P_{\Lambda}: \sum_{\alpha \in \mathbb{N}_0^d} c_{\alpha} z^{\alpha} \mapsto \sum_{\alpha \in \Lambda} c_{\alpha} z^{\alpha}$$

the associated idempotent coefficient multiplier defined on the space of analytic polynomials. If P_{Λ} extends to a contraction on $H^p_K(\mathbb{T}^d)$, then Λ is of the form $(\gamma + \Delta) \cap \mathbb{N}^d_0$, where $\gamma \in \mathbb{Z}^d$ and Δ is a subgroup of \mathbb{Z}^d .

Proof. The proof shares similarity with that of Theorem 2.5. Fix an $\gamma \in \Lambda$. We define an operator

$$T: L^p(\mathbb{T}^d, Kdm_d) \to L^p(\mathbb{T}^d, Kdm_d)$$

by $Tf(z) := z^{-\gamma}f(z)$. Clearly, T is an isometry. Denote by X the range of $H^p_K(\mathbb{T}^d)$ under T. Then the restriction of T on $H^p_K(\mathbb{T}^d)$ induces an isometric isomorphism from $H^p_K(\mathbb{T}^d)$ onto X. Set

$$P = TP_{\Lambda}T^{-1}|_X,$$

then P is a contractive projection on X. Moreover, for any $\alpha \in \mathbb{N}_0^d - \gamma$, we have $z^{\alpha} \in X \cap L^{\infty}(\mathbb{T})$ and

$$Pz^{\alpha} = z^{-\gamma} P_{\Lambda}(z^{\gamma+\alpha}) = \begin{cases} z^{\alpha} & \text{if } \alpha + \gamma \in \Lambda; \\ 0 & \text{otherwise.} \end{cases}$$

By Theorem 1.2, there exists a sub σ -algebra \mathcal{F} , such that

$$Pz^{\alpha} = \mathbb{E}(z^{\alpha}|\mathcal{F}), \ \alpha \in \mathbb{N}_0^d - \gamma.$$

Define

$$\Delta := \{ \alpha \in \mathbb{Z}^d : z^\alpha \text{ is } \mathcal{F} - \text{measurable} \}.$$

Then Δ is a subgroup of \mathbb{Z}^d and

$$(\mathbb{N}_0^d - \gamma) \cap (\Lambda - \gamma) = (\mathbb{N}_0^d - \gamma) \cap \Delta.$$

This implies $\Lambda = (\gamma + \Delta) \cap \mathbb{N}_0^d$.

A simple example is presented below to show the converse of Theorem 2.6 is not true in general. Suppose $0 . Pick a function <math>g \in H^1(\mathbb{T})$ such that |g| is not equal to a constant. Set K = |g| and $\Lambda = \{0\}$. We claim that P_{Λ} cannot extend to a contraction on $H^p_K(\mathbb{T})$. Otherwise, $\mathbf{1} \in \operatorname{Ran} P_{\Lambda}$ and $z^n \in \operatorname{Ker} P_{\Lambda}$ for each $n \geq 1$. By (2.1), it holds that

$$\int_{\mathbb{T}} \bar{z}^n K(z) dm(z) = 0, \ \forall n \ge 1,$$

and hence

$$\int_{\mathbb{T}} z^n K(z) dm(z) = 0, \ \forall n \neq 0.$$

This forces K to be a constant, which leads to a contradiction.

As regards idempotent coefficient multipliers, there exist some new fascinating behaviors in higher-dimensional cases, as shown in [BOS21]. However, in dimension d = 1, there seems to be of considerable consistency for the generalized Bergman spaces on the closed unit disk (including the Hardy space). For a probability measure μ on $\overline{\mathbb{D}}$ not centred on the origin, the generalized Bergman space $L^p_a(\mu)$ is defined to be the closure of analytic polynomials in $L^p(\mu)$. As done below Definition 2.4, when μ is rotation-invariant, it is not difficult to verify that if Λ consists of a single point or $\Lambda = (l + k\mathbb{Z}) \cap \mathbb{N}_0$ with $k \geq 1$, then P_{Λ} extends to a contraction on $L^p_a(\mu)$ for all $1 \leq p < \infty$. Conversely, even if μ is not rotation-invariant, we still have the following conclusion.

Proposition 2.7. Suppose that μ is a probability measure μ on $\overline{\mathbb{D}}$ not centred on the origin, and $1 \leq p < \infty$, $p \neq 2$. Let Λ be a nonempty subset of \mathbb{N}_0 and denote

by

$$P_{\Lambda}: \sum_{n \ge 0} c_n z^n \mapsto \sum_{n \in \Lambda} c_n z^n$$

the associated idempotent coefficient multiplier defined on the space of analytic polynomials. If P_{Λ} extends to a contraction on $L^p_a(\mu)$, then Λ consists of a single point or $\Lambda = (l + k\mathbb{Z}) \cap \mathbb{N}_0$ with $k \geq 1$.

Proof. We assume that Λ is not a single point. Suppose $r \in \Lambda$ and s > 0 such that $r+s \in \Lambda$. Motivated by the identity (2.1) (here we use the version of $p \ge 1$), we consider the integral for $t \in \mathbb{N}_0$,

$$\Phi_t(\zeta) = \int_{\overline{\mathbb{D}}} |z^r + \zeta z^{r+s}|^{p-2} (z^r + \zeta z^{r+s}) \overline{z}^t d\mu(z), \quad \zeta \in \mathbb{D}.$$

When $|\zeta|$ is small enough, this integral can be expressed as

$$\int_{\overline{\mathbb{D}}} |z^r + \zeta z^{r+s}|^{p-2} (z^r + \zeta z^{r+s}) \overline{z}^t d\mu(z)$$

$$= \int_{\overline{\mathbb{D}}} |1 + \zeta z^s|^{p-2} (1 + \zeta z^s)|z|^{rp} \overline{z}^{t-r} d\mu(z)$$

$$= \int_{\overline{\mathbb{D}}} \left(\sum_{m,n\geq 0} c_{m,n} z^{sm} \zeta^m \overline{z}^{sn} \overline{\zeta}^n \right) |z|^{rp} \overline{z}^{t-r} d\mu(z)$$

$$= \sum_{m,n\geq 0} \left(c_{m,n} \int_{\overline{\mathbb{D}}} z^{sm} \overline{z}^{sn+t-r} |z|^{rp} d\mu(z) \right) \zeta^m \overline{\zeta}^n.$$

Here, the coefficients $c_{m,n}$ is determined by binomial expansion as in (2.3). When $t = s(m-n) + r \ge 0$, the coefficient of $\zeta^m \bar{\zeta}^n$ is $c_{m,n} \int_{\overline{\mathbb{D}}} |z|^{rp+2sm} d\mu(z)$. In the case m = 2, n = 0, t = r + 2s, a straightforward calculation shows that the coefficient of ζ^2 equals to $\frac{p(p-2)}{8} \int_{\overline{\mathbb{D}}} |z|^{rp+4s} d\mu(z) \ne 0$, and hence by the identity (2.1), $r + 2s \in \Lambda$. In the case m = 0, n = 1, if $t = r - s \ge 0$, then the coefficient of $\bar{\zeta}$ equals to $\frac{p-2}{2} \int_{\overline{\mathbb{D}}} |z|^{rp} d\mu(z) \ne 0$, and hence $r - s \in \Lambda$. Let k be the smallest positive integer for which there exists $l \in \Lambda$ such that $l + k \in \Lambda$. The above discussions shows $(l + k\mathbb{Z}) \cap \mathbb{N}_0 \subseteq \Lambda$. The minimality of k then implies that $\Lambda = (l + k\mathbb{Z}) \cap \mathbb{N}_0$. The proof is completed.

3. Conditional expectation operators on $H^p(\mathbb{T}), \ 0 .$

Theorem 1.2 provides a sufficient condition by which a contractive projection on a closed subspace X of L^p is restriction of a conditional expectation. Here we emphasize that to obtain contractive projections via restricting conditional expectations to X, two natural problems arise. The first one is

Problem A. Given a closed subspace X of $L^p(\Omega, \Sigma, \mu)$ for $1 \le p < \infty$. For which sub σ -algebra $\mathcal{F} \subseteq \Sigma$, the conditional expectation $\mathbb{E}(\cdot|\mathcal{F})$ leaves X invariant?

For the second, one should be careful to consider the case 0 , sinceconditional expectations are only defined for integrable functions.

Problem B. Let X be a closed subspace of $L^p(\Omega, \Sigma, \mu)$ with $0 . Suppose that <math>X \cap L^1$ is dense in X, and that $\mathbb{E}(\cdot|\mathcal{F})$ leaves $X \cap L^1$ invariant. Then $\mathbb{E}(\cdot|\mathcal{F})$ is a densely defined operator in X. When is $\mathbb{E}(\cdot|\mathcal{F})$ bounded or contractive on X?

It seems to be difficult to completely solve Problem A for general closed subspaces of L^p . To the best of our knowledge, the only progress in this direction is the following interesting result due to Aleksanderov [Al86].

Theorem 3.1 (Aleksanderov's Theorem). The conditional expectation $\mathbb{E}(\cdot|\mathcal{F})$ on $L^1(\mathbb{T})$ leaves $H^1(\mathbb{T})$ invariant if and only if either \mathcal{F} is generated by the constant function **1** or \mathcal{F} is generated by an inner function with $\eta(0) = 0$.

Clearly, $\mathbb{E}(\cdot|\mathcal{F})$ leaves $H^1(\mathbb{T})$ invariant if and only if it leaves $H^p(\mathbb{T})$ invariant for any $1 \leq p \leq \infty$. Moreover, the 'if' part of Aleksanderov's theorem is easy. Actually, if η is an inner function vanishing at the origin, then $\{\eta^k : k \in \mathbb{Z}\}$ forms an orthonormal basis of $L^2(\mathbb{T}, \Sigma(\eta), dm)$. Since $\mathbb{E}(\cdot|\eta)$ is the orthogonal projection from $L^2(\mathbb{T}, dm)$ onto $L^2(\mathbb{T}, \Sigma(\eta), dm)$, we have

$$\mathbb{E}(f|\eta) = \sum_{k \in \mathbb{Z}} \langle f, \eta^k \rangle \, \eta^k$$

for any $f \in L^2(\mathbb{T}, dm)$, and hence $\mathbb{E}(\cdot|\eta)$ leaves $H^2(\mathbb{T})$ invariant. An approximation argument shows that $\mathbb{E}(\cdot|\eta)$ leaves $H^1(\mathbb{T})$ invariant.

Aleksanderov's theorem provides an answer to the Problem A in the case $X = H^p(\mathbb{T})$. The main aim of this section is to prove Theorem 1.4 which solves Problem B in the case $X = H^p(\mathbb{T})$. This theorem will enable us to study contractive idempotent coefficient multipliers on $H^p(\mathbb{T}^d)$ for $0 , <math>1 \le d \le \infty$.

In what follows that we will show that when $0 , the conditional expectation <math>\mathbb{E}(\cdot|\eta)$ is bounded on H^p if and only if η is a finite Blaschke product. Furthermore, if η is a finite Blaschke product, then

$$\|\mathbb{E}(\cdot|\eta)\|_{H^p} = \sup_{w\in\mathbb{T}} |\eta'(w)|^{\frac{1}{p}-1}.$$

We begin with the necessary part. The following lemma is a part of the Duren-Romberg-Shields theorem [DRS69, Theorem 1] characterizing the dual space of $H^p(\mathbb{T})$ for 0 .

Lemma 3.2. Let $0 and let <math>\lambda$ be a continuous linear functional on $H^p(\mathbb{T})$. Then there exists a unique function $g \in A(\mathbb{D})$ such that

$$\lambda(f) = \lim_{r \to 1} \frac{1}{2\pi} \int_0^{2\pi} f(re^{i\theta}) \overline{g(e^{i\theta})} d\theta, \quad f \in H^p(\mathbb{T}).$$

Here, $A(\mathbb{D})$ is the classical disc algebra, consisting of all continuous functions on $\overline{\mathbb{D}}$ that are analytic in \mathbb{D} .

Theorem 3.3. Let $0 , and let <math>\eta$ be an inner function with $\eta(0) = 0$. If $\mathbb{E}(\cdot|\eta)$ is bounded on $H^p(\mathbb{T})$, then η is a finite Blaschke product.

Proof. Write $\eta(z) = z^k \xi(z)$ where ξ is an inner function and $\xi(0) \neq 0$. Denote by $\tau_k : f \mapsto \widehat{f}(k)$, where $\widehat{f}(k)$ is the k-th Fourier coefficient of f. Because both τ_k and $\mathbb{E}(\cdot|\eta)$ are bounded on $H^p(\mathbb{T})$, we see

$$\lambda = \tau_k \circ \mathbb{E}(\cdot|\eta) : f \mapsto \lim_{r \to 1} \frac{1}{2\pi} \int_0^{2\pi} \mathbb{E}(f|\eta)(re^{i\theta}) \cdot \overline{e^{ik\theta}} d\theta$$

is a continuous linear functional on $H^p(\mathbb{T})$. By Lemma 3.2, there exists a unique function $g \in A(\mathbb{D})$, such that

$$\lambda(f) = \lim_{r \to 1} \frac{1}{2\pi} \int_0^{2\pi} f(re^{i\theta}) \overline{g(e^{i\theta})} d\theta, \quad f \in H^p(\mathbb{T}).$$

In particular, for any analytic polynomial q, it holds that

$$\lambda(q) = \int_{\mathbb{T}} q(w) \overline{g(w)} dm(w).$$

Since $\mathbb{E}(q|\eta) \in H^{\infty}(\mathbb{T})$ for any analytic polynomial q, and $\mathbb{E}(\cdot|\eta)$ is self-adjoint as an operator on $L^{2}(\mathbb{T}, dm)$, we have

$$\begin{split} \lambda(q) &= \int_{\mathbb{T}} \mathbb{E}(q|\eta)(w) \cdot \overline{w}^k dm(w) \\ &= \int_{\mathbb{T}} q(w) \cdot \overline{\mathbb{E}(z^k|\eta)(w)} dm(w) \\ &= \int_{\mathbb{T}} q(w) \cdot \overline{\left(\sum_{l \ge 0} \langle z^k, \eta^l \rangle \eta^l(w)\right)} dm(w) \\ &= \int_{\mathbb{T}} q(w) \cdot \xi(0) \overline{\eta(w)} dm(w) \end{split}$$

As a result, the identity

$$\int_{\mathbb{T}} q(w)\overline{g(w)}dm(w) = \int_{\mathbb{T}} q(w) \cdot \xi(0)\overline{\eta(w)}dm(w)$$

holds for any analytic polynomial q. We conclude that $g = \overline{\xi(0)}\eta$. This shows $\eta \in A(\mathbb{D})$ and hence η is a finite Blaschke product.

Next, we consider the converse of Theorem 3.3. We first recall some basic properties for finite Blaschke products. Let $\eta(z) = \prod_{j=1}^{k} \frac{z-a_j}{1-\overline{a_j}z}$ be a finite Blaschke product where $a_j \in \mathbb{D}$, $1 \leq j \leq k$. A direct calculation yields

$$\sum_{j=1}^{k} \frac{1-|a_j|}{1+|a_j|} \le |\eta'(z)| \le \sum_{j=1}^{k} \frac{1+|a_j|}{1-|a_j|}, \ z \in \mathbb{T}.$$

This shows that η is a locally univalent Lipschitz function on \mathbb{T} . Actually, η is a covering map from \mathbb{T} onto itself with degree k. To be more specific, for each $z \in \mathbb{T}$ the preimage $\eta^{-1}(z)$ consists of exactly k distinct points on \mathbb{T} . And if we write $\eta^{-1}(z) = \{w_{\ell} : 1 \leq \ell \leq k\}$ with

$$\operatorname{Arg} w_1 < \operatorname{Arg} w_2 < \dots < \operatorname{Arg} w_k < 2\pi + \operatorname{Arg} w_1,$$

then η maps each arc $[w_{\ell}, w_{\ell+1})$ to \mathbb{T} bijectively $(w_{k+1} := w_1)$.

The following identity, which can be easily established by considering f as the indicator function on a small arc, is crucial to the proof of the sufficiency part of Theorem 1.4.

Lemma 3.4 (The Arc Length Formula). Let η be a finite Blaschke product. Then for any $f \in L^1(\mathbb{T})$,

$$\int_{\mathbb{T}} f(z) |\eta'(z)| dm(z) = \int_{\mathbb{T}} \bigg(\sum_{w \in \eta^{-1}(z)} f(w) \bigg) dm(z).$$

It is also well-known that the preimage $\eta^{-1}(E)$ is a null set if $E \subseteq \mathbb{T}$ is a null set. This can be easily deduced from Lemma 3.4. Indeed, by setting f as the indicator function of $\eta^{-1}(E)$, we obtain

$$C \cdot m(\eta^{-1}(E)) \leq \int_{\mathbb{T}} \mathbf{1}_{\eta^{-1}(E)}(z) |\eta'(z)| dm(z)$$
$$= \int_{\mathbb{T}} \sum_{w \in \eta^{-1}(z)} \mathbf{1}_{\eta^{-1}(E)}(w) dm(z)$$
$$= \int_{\mathbb{T}} k \cdot \mathbf{1}_{E}(z) dm(z)$$
$$= 0,$$

where $C = \inf_{z \in \mathbb{T}} |\eta'(z)| > 0.$

When we fix a finite Blaschke product η , the notation \sim stands for the equivalence induced by η . Namely, for two points $z, w \in \mathbb{T}$, $w \sim z$ if $\eta(w) = \eta(z)$.

Proposition 3.5. Let η be a finite Blaschke product with $\eta(0) = 0$. Then

$$\sum_{w \sim z} \frac{1}{|\eta'(w)|} = 1, \quad \text{for each } z \in \mathbb{T}.$$

In particular, $\|\eta'\|_{\infty} \ge k$, where k is the degree of η .

Proof. We write $F(z) = \sum_{w \sim z} \frac{1}{|\eta'(w)|}$ and $G(z) = \sum_{w \in \eta^{-1}(z)} \frac{1}{|\eta'(w)|}$. Clearly, $F(z) = G(\eta(z))$. For each $n \in \mathbb{Z}$, applying The Arc Length Formula to $f = \eta^n / |\eta'|$ we obtain

$$\int_{\mathbb{T}} \sum_{w \in \eta^{-1}(z)} \frac{\eta^n(w)}{|\eta'(w)|} dm(z) = \int_{\mathbb{T}} \eta^n(z) dm(z) = \begin{cases} 1 & n = 0; \\ 0 & n \neq 0. \end{cases}$$

This implies that

$$\int_{\mathbb{T}} z^n G(z) dm(z) = \begin{cases} 1 & n = 0; \\ 0 & n \neq 0. \end{cases}$$
(3.1)

Observe that G is continuous, it follows from (3.1) that G(z) = 1 for each $z \in \mathbb{T}$. Hence $F(z) = G(\eta(z)) = 1$ for each $z \in \mathbb{T}$.

Proposition 3.6. Let f be a Lebesgue measurable function on \mathbb{T} and let η be a finite Blaschke product. Then the function $F(z) = \sum_{w \sim z} f(w)$ is $\Sigma(\eta)$ -measurable.

Proof. It suffices to prove the case when $f = \mathbf{1}_E$ is an indicator function for any Lebesgue measurable set E. Writing \mathbb{T} as a finite union of arcs on which η is injective, we may assume, without loss of generality that η is injective on E. Then $F = \mathbf{1}_{\eta^{-1}(\eta(E))}$, and we need to show that $\eta^{-1}(\eta(E))$ is $\Sigma(\eta)$ -measurable. Since η is a Lipschitz function on \mathbb{T} , $\eta(E)$ is Lebesgue measurable. Thus $\eta(E) = F_0 \cup F_1$, where F_0 is a null set and F_1 is a Borel set. Note that $\eta^{-1}(F_0)$ is also null. This implies $\eta^{-1}(\eta(E)) = \eta^{-1}(F_0) \cup \eta^{-1}(F_1)$ is $\Sigma(\eta)$ -measurable. \Box

With the above preparations, we are able to give a pointwise expression for the conditional expectation.

Proposition 3.7. Let η be a finite Blaschke product with $\eta(0) = 0$. Then for any $f \in L^1(\mathbb{T})$,

$$\mathbb{E}(f|\eta)(z) = \sum_{w \sim z} \frac{f(w)}{|\eta'(w)|}.$$

Proof. Given a function $f \in L^1(\mathbb{T})$, the function

$$F(z) = \sum_{w \sim z} \frac{f(w)}{|\eta'(w)|}$$

is $\Sigma(\eta)$ -measurable by Proposition 3.6. As a consequence, to show $\mathbb{E}(f|\eta) = F$ it is enough to establish

$$\int_{\mathbb{T}} \eta^n(z) F(z) \, dm(z) = \int_{\mathbb{T}} \eta^n(z) f(z) \, dm(z),$$

for each $n \in \mathbb{Z}$. This can be deduced, by applying the Arc Length Formula, Proposition 3.5, and the following calculations:

$$\begin{split} &\int_{\mathbb{T}} \eta^{n}(z)F(z)\,dm(z) \\ &= \int_{\mathbb{T}} \bigg(\sum_{w \in \eta^{-1}(z)} \frac{\eta^{n}(w) \cdot F(w)}{|\eta'(w)|} \bigg)\,dm(z) \\ &= \int_{\mathbb{T}} z^{n} \cdot \bigg(\sum_{w \in \eta^{-1}(z)} \frac{F(w)}{|\eta'(w)|} \bigg)\,dm(z) \\ &= \int_{\mathbb{T}} z^{n} \cdot \bigg(\sum_{w \in \eta^{-1}(z)} \frac{1}{|\eta'(w)|} \sum_{\zeta \sim w} \frac{f(\zeta)}{|\eta'(\zeta)|} \bigg)\,dm(z) \\ &= \int_{\mathbb{T}} z^{n} \cdot \bigg(\sum_{w \in \eta^{-1}(z)} \frac{1}{|\eta'(w)|} \bigg) \cdot \bigg(\sum_{\zeta \in \eta^{-1}(z)} \frac{f(\zeta)}{|\eta'(\zeta)|} \bigg)\,dm(z) \\ &= \int_{\mathbb{T}} \sum_{\zeta \in \eta^{-1}(z)} \frac{\eta^{n}(\zeta) \cdot f(\zeta)}{|\eta'(\zeta)|}\,dm(z) \\ &= \int_{\mathbb{T}} \eta^{n}(z)f(z)\,dm(z). \end{split}$$

We complete the proof.

Next, we prove the sufficient part of Theorem 1.4.

Theorem 3.8. Suppose $0 and let <math>\eta$ be a finite Blaschke product of degree $k \geq 1$ with $\eta(0) = 0$. Denote by $\|\mathbb{E}(\cdot|\eta)\|_{H^p}$ and $\|\mathbb{E}(\cdot|\eta)\|_{L^p}$ the norms of the conditional expectation operator $\mathbb{E}(\cdot|\eta)$ on $H^p(\mathbb{T})$ and $L^p(\mathbb{T})$, respectively. Then

$$\|\mathbb{E}(\cdot|\eta)\|_{H^p} = \|\mathbb{E}(\cdot|\eta)\|_{L^p} = \|\eta'\|_{\infty}^{\frac{1}{p}-1}.$$

In particular,

$$\|\mathbb{E}(\cdot|z^k)\|_{H^p} = \|\mathbb{E}(\cdot|z^k)\|_{L^p} = k^{\frac{1}{p}-1}.$$

Proof. Clearly, $\|\mathbb{E}(\cdot|\eta)\|_{H^p} \leq \|\mathbb{E}(\cdot|\eta)\|_{L^p}$. We first establish $\|\mathbb{E}(\cdot|\eta)\|_{L^p} \leq \|\eta'\|_{\infty}^{\frac{1}{p}-1}$. For any $f \in L^1(\mathbb{T})$, by proposition 3.7 we have

$$\int_{\mathbb{T}} |\mathbb{E}(f|\eta)(z)|^p dm(z) = \int_{\mathbb{T}} \left| \sum_{w \sim z} \frac{f(w)}{|\eta'(w)|} \right|^p dm(z)$$
$$\leq \int_{\mathbb{T}} \sum_{w \sim z} \frac{|f(w)|^p}{|\eta'(w)|^p} dm(z)$$

$$= \int_{\mathbb{T}} \mathbb{E}(|f|^{p} |\eta'|^{1-p} |\eta)(z) \, dm(z)$$

$$= \int_{\mathbb{T}} |f(z)|^{p} |\eta'(z)|^{1-p} \, dm(z)$$

$$\leq \|f\|_{p}^{p} \|\eta'\|_{\infty}^{1-p}.$$

Thus, $\|\mathbb{E}(\cdot|\eta)\|_{L^p} \leq \|\eta'\|_{\infty}^{\frac{1}{p}-1}$. Next, we show $\|\mathbb{E}(\cdot|\eta)\|_{H^p} \geq \|\eta'\|_{\infty}^{\frac{1}{p}-1}$. Given $z_0 \in \mathbb{T}$ and $\varepsilon > 0$, there exists an open arc I containing z_0 such that for each $z \in I$,

$$\left|\frac{1}{|\eta'(z_0)|} - \frac{1}{|\eta'(z)|}\right| < \varepsilon, \text{ and } \left||\eta'(z_0)| - |\eta'(z)|\right| < \varepsilon.$$

Moreover, we can choose I small enough to ensure that $J := \eta^{-1}(\eta(I))$ consists exactly of k disjoint open arcs. Now for sufficiently large t, there exists a function $f_t \in H^{\infty}(\mathbb{T})$ such that

$$f_t(z)| = \begin{cases} t & z \in I; \\ 1 & z \notin I. \end{cases}$$

It follows from Proposition 3.7 that

$$|\mathbb{E}(f_t|\eta)(z)| = \left|\sum_{w\sim z} \frac{f_t(w)}{|\eta'(w)|}\right|$$
$$\geq t\left(\frac{1}{|\eta'(z_0)|} - \varepsilon\right) - (k-1)\left\|\frac{1}{\eta'}\right\|_{\infty}, \quad z \in J.$$
(3.2)

and

$$m(J) = \int_{\mathbb{T}} \sum_{w \sim z} \mathbf{1}_{I}(w) \, dm(z) = \int_{\mathbb{T}} \mathbb{E} \big(|\eta'| \cdot \mathbf{1}_{I} |\eta \big)(z) dm(z)$$
$$= \int_{\mathbb{T}} |\eta'(z)| \cdot \mathbf{1}_{I}(z) dm(z) \ge \big(|\eta'(z_{0})| - \varepsilon \big) \, m(I).$$
(3.3)

Combining the lower estimations (3.2) and (3.3) we have

$$\|\mathbb{E}(f_t|\eta)\|_p \ge \left(t\left(\frac{1}{|\eta'(z_0)|} - \varepsilon\right) - (k-1)\|\frac{1}{\eta'}\|_{\infty}\right) \cdot \left(|\eta'(z_0)| - \varepsilon\right)^{1/p} m(I)^{1/p}.$$
(3.4)

Moreover, we have a upper bound

$$||f_t||_p \le (t^p m(I) + 1)^{1/p}.$$
 (3.5)

Now by (3.4) and (3.5)

$$\|\mathbb{E}(\cdot|\eta)\|_{H^p} \ge \limsup_{t \to \infty} \frac{\|\mathbb{E}(f_t|\eta)\|_p}{\|f_t\|_p} \ge \left(\frac{1}{|\eta'(z_0)|} - \varepsilon\right) \left(|\eta'(z_0)| - \varepsilon\right)^{\frac{1}{p}}.$$

Letting $\varepsilon \to 0$, we see $\|\mathbb{E}(\cdot|\eta)\|_{H^p} \ge |\eta'(z_0)|^{\frac{1}{p}-1}$. Since $z_0 \in \mathbb{T}$ is arbitrary,

$$\|\mathbb{E}(\cdot|\eta)\|_{H^p} \ge \|\eta'\|_{\infty}^{\frac{1}{p}-1}$$

This finishes the proof.

The following result, which will be used in Section 4, is an immediate consequence of Theorem 1.3 and Theorem 1.4.

Corollary 3.9. Let 0 . Suppose <math>P is a contractive projection on $H^p(\mathbb{T})$ such that $P\mathbf{1} = \mathbf{1}$ and for each $n \in \mathbb{N}$, $Pz^n \in H^{\infty}(\mathbb{T})$, then P is either $\mathbb{E} : f \mapsto f(0)$ or $\mathbb{E}(\cdot|z) : f \mapsto f$.

4. Contractive projection sets for $H^p(\mathbb{T}^d), \ 0$

Let $1 \leq d \leq \infty$, and $0 . Suppose that <math>\Lambda$ is a nonempty subset of \mathbb{N}_0^d and P_{Λ} is the associated idempotent coefficient multiplier defined by (1.2). As we mentioned in previous sections, Brevig, Ortega-Cerdà and Seip [BOS21] completely determined when P_{Λ} extends to a contraction on $H^p(\mathbb{T}^d)$ for $1 \leq p \leq \infty$. In their work, such Λ is termed a contractive projection set for $H^p(\mathbb{T}^d)$. Our main aim in this section is to characterize contractive projection sets for $H^p(\mathbb{T}^d)$ for the remaining case 0 .

Observe that both $\{0\}$ and \mathbb{N}_0 are contractive projection sets for $H^p(\mathbb{T})$ for 0 . In fact, there is no other.

Theorem 4.1. Suppose that $0 and <math>\Lambda$ is a contractive projection set for $H^p(\mathbb{T})$. Then $\Lambda = \{0\}$ or $\Lambda = \mathbb{N}_0$.

Proof. Suppose k is the smallest integer in Λ . If k = 0, then the results follow immediately from Corollary 3.9. Now assume $k \neq 0$ by contradiction. Define

$$Pf = \bar{z}^k P_{\Lambda}(z^k f(z)).$$

Then P is a contractive projection on $H^p(\mathbb{T})$ satisfying the hypotheses of Corollary 3.9. As a consequence, $\Lambda = \{k\}$ or $\Lambda = \{k, k+1, k+2, \ldots\}$. The latter case can be excluded by utilizing Theorem 2.6. To complete the proof, it is sufficient to show c(k,p) > 1 where c(k,p) is the norm of the k-th Fourier coefficient functional on $H^p(\mathbb{T})$. For $f \in H^p(\mathbb{T})$, define $f_k(z) = f(z^k)$ so that $||f||_p = ||f_k||_p$ and $f'(0) = f_k^{(k)}(0)/k!$. Thus $c(k,p) \ge c(1,p)$. Now the proof is completed as the explicit value

$$c(1,p) = \sqrt{\frac{2}{p}} \left(1 - \frac{p}{2}\right)^{\frac{1}{p} - \frac{1}{2}} > 1$$

was established by Brevig and Saksman in [BS20].

For $1 \leq d \leq \infty$ and a subset $J \subseteq \{1, 2, \dots, d\}$ (if $d = \infty, J \subseteq \mathbb{N}$), we define $\mathbb{N}_J = \{\alpha \in \mathbb{N}_0^d : \alpha_j = 0 \text{ for all } j \in J\}$. For an analytic polynomial f and p > 0, the function $|f|^p$ is plurisubharmonic. Using the sub-mean value principle, it is easy to verify that \mathbb{N}_J is a contractive projection set for $H^p(\mathbb{T}^d)$. Actually, when $0 , the collection <math>\{\mathbb{N}_J : J \text{ runs over all subsets}\}$ exhausts all contractive projection sets for $H^p(\mathbb{T}^d)$.

Theorem 4.2. Suppose that $0 and <math>1 \le d \le \infty$, and Λ is a contractive projection set for $H^p(\mathbb{T}^d)$. Then $\Lambda = \mathbb{N}_J$ for some subset J.

Proof. Let $J = \{j : \alpha_j = 0, \forall \alpha \in \Lambda\}$. We will show that $\Lambda = \mathbb{N}_J$. Clearly $\Lambda \subseteq \mathbb{N}_J$. Next, we prove the converse inclusion. Let $\mathbf{e}^{(i)}$ be the canonical basis of \mathbb{N}_0^d determined by $\mathbf{e}_i^{(i)} = 1$ and $\mathbf{e}_j^{(i)} = 0$ for $j \neq i$. For a $\beta \in \mathbb{N}_0^d$, we define $\beta^{(i)}$ by $\beta_i^{(i)} = 0$ and $\beta_j^{(i)} = \beta_j$ for $j \neq i$. In what follows, we may assume $\Lambda \neq \{\mathbf{0}\}$.

We first claim that if $\beta \in \Lambda$ and $\beta_i \neq 0$, then $\beta^{(i)} + k\mathbf{e}^{(i)} \in \Lambda$ for all $k \in \mathbb{N}_0$. Indeed, for any analytic polynomial q in one single variable, one can verify that the function

$$Pq(z) = \bar{z}^{\beta^{(i)}} P_{\Lambda} \left(z^{\beta^{(i)}} q(z_i) \right)$$

only depends on the variable z_i , and hence P extends to a contraction on $H^p(\mathbb{T})$ with the contractive projection set $\{k \in \mathbb{N}_0 : \beta^{(i)} + k\mathbf{e}^{(i)} \in \Lambda\}$. Now our claim follows from Theorem 4.1 because $\beta_i \neq 0$ and β_i belongs to this set.

Take a $\beta = (\beta_j)_{j\geq 1} \in \Lambda$ with $\beta \neq 0$. Since there are only finitely many j such that $\beta_j \neq 0$, iteratively applying the previous claim shows $\sum_{\beta_j\neq 0} k_j \mathbf{e}^{(j)} \in \Lambda$ for all $k_j \in \mathbb{N}_0$. In particular, $\mathbf{0} \in \Lambda$. Together this with Theorem 2.6, Λ is a sub-semigroup of \mathbb{N}_0^d . For $\alpha = (\alpha_j)_{j\geq 1} \in \mathbb{N}_J$, if $\alpha_l \neq 0$, then by definition there exists a $\beta \in \Lambda$ such that $\beta_l \neq 0$, and hence $\alpha_l \mathbf{e}^{(l)} \in \Lambda$. Since Λ is a sub-semigroup of \mathbb{N}_0^d , we see that $\alpha = \sum_l \alpha_l \mathbf{e}^{(l)} \in \Lambda$, completing the proof.

We end this paper by reformulating Theorem 4.2 for the Hardy space of Dirichlet series \mathscr{H}^p which corresponds to the case $d = \infty$ in Theorem 4.2 via the Bohr lift. The Hardy space of Dirichlet series \mathscr{H}^p is defined as the closure of all Dirichlet polynomials $f(s) = \sum_{n=1}^{N} a_n n^{-s}$ under the norm (or quasi-norm when 0),

$$||f||_{\mathscr{H}^p}^p = \lim_{T \to \infty} \frac{1}{2T} \int_{-T}^T |f(it)|^p \, dt.$$

It is known that the elements of \mathscr{H}^p are indeed Dirichlet series absolutely converging on the right half plane $\operatorname{Re} s > 1/2$. Let $p = (p_1, p_2, \ldots)$ be the ordered

prime numbers. According to the fundamental theorem of arithmetic, each integer $n \in \mathbb{N}$ can be uniquely decomposed as

$$n = p^{\kappa(n)} := \prod_{j=1}^{\infty} p_j^{\kappa_j}, \quad \kappa(n) \in \mathbb{N}_0^d.$$

By the Birkhoff-Oxtoby ergodic theorem, the Bohr lift,

$$\mathscr{B}: \mathscr{H}^p \to H^p(\mathbb{T}^\infty), \quad f(s) = \sum_{n=1}^\infty a_n n^{-s} \mapsto \mathscr{B}f(z) = \sum_{n=1}^\infty a_n z^{\kappa(n)}$$

establishes an isometric isomorphism between \mathscr{H}^p and $H^p(\mathbb{T}^\infty)$.

A nonempty subset $\Lambda \subseteq \mathbb{N}$ is termed a *contractive projection set for* \mathscr{H}^p if the operator $P_{\Lambda} : \sum_{n=1}^{\infty} a_n n^{-s} \mapsto \sum_{n \in \Lambda} a_n n^{-s}$, defined on the space of Dirichlet polynomials, extends to a contraction on \mathscr{H}^p .

Theorem 4.3. Suppose that $0 . Then <math>\Lambda$ is a contractive projection set for \mathscr{H}^p if and only if there is a subset J of all prime numbers such that $\Lambda = \{n \in \mathbb{N} : \text{all prime divisors of } n \text{ are in } J\}.$

Acknowledgements

This work was partially supported by NNSF of China (12231005).

References

- [Al86] A. B. Aleksandrov, Measurable partitions of the circle induced by inner functions, J. Soviet Math. 42 (1988), 1610–1613.
- [An66] T. Andô, Contractive projections in L^p-spaces, Pacific J. Math. 17 (1966), 391–405.
- [AF92] J. Arazy, Y. Friedman, Contractive projections in C_p , Mem. Amer. Math. Soc. 95 (1992).
- [AR24] C. Arhancet, Y. Raynaud, 2-positive contractive projections on noncommutative Lpspaces, J. Funct. Anal. 286 (2024).
- [Bo07] V. I. Bogachev, Measure theory, Vol. I. Springer-Verlag, Berlin, 2007.
- [BOS21] O. F. Brevig, J. Ortega-Cerdà, and K. Seip, Idempotent Fourier multipliers acting contractively on H^p spaces, Geom. Funct. Anal. **31** (2021), 1377–1413.
- [BS20] O. F. Brevig, E. Saksman, Coefficient estimates for H^p spaces with 0 , Proc.Amer. Math. Soc.**148**(2020), 3911–3924.
- [Do65] R. G. Douglas, Contractive projections on an \mathfrak{L}_1 space, Pacific J. Math. 15 (1965), 443–462.
- [Du10] R. Durrett, *Probability: Theory and Examples*, Fourth edition, Cambridge University Press, Cambridge, 2010.
- [DRS69] P. L. Duren, B. W. Romberg, and A. L. Shields, *Linear functionals on* H^p spaces with 0 , J. Reine Angew. Math.**238**(1969), 32–60.
- [FR82] Y. Friedman, B. Russo, Contractive projections on $C_0(K)$, Trans. Amer. Math. Soc. **273** (1982), 57–73.

- [GNZ24] K. Guo, J. Ni, and Q. Zhou, Nevanlinna class, Dirichlet series and Szegö's problem, arXiv:2201.01993, (to appear in Annales de l'institut Fourier).
- [KR65] B. R. Kripke, T. J. Rivlin, Approximation in the Metric of $L^1(X, \mu)$, Trans. Amer. Math. Soc. **119**, (1965) 101–122.
- [LRR05] F. Lancien, B. Randrianantoanina, and E. Ricard, On contractive projections in Hardy spaces, Studia Math. 171 (2005), 93–102.
- [Ran01] B. Randrianantoanina, Norm-one projections in Banach spaces, Taiwanese J. Math. 5 (2001), 35–95.
- [Ray04] Y. Raynaud, The range of a contractive projection in $L^p(H)$, Rev. Mat. Complut. 17 (2004), 485–512.
- [Rud90] W. Rudin, Fourier Analysis on Groups, Wiley Classics Library, John Wiley & Sons, Inc., New York, 1990.
- [Sha71] H. S. Shapiro, Topics in Approximation Theory, Springer-Verlag, Berlin-New York, 1971.

XIANGDI FU: SCHOOL OF MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, SHANGHAI, 200433, CHINA

Email address: xdfu20@fudan.edu.cn

Kunyu Guo: School of Mathematical Sciences, Fudan University, Shanghai, 200433, China

Email address: kyguo@fudan.edu.cn

DILONG LI: SCHOOL OF MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, SHANGHAI, 200433, CHINA

Email address: dlli23@m.fudan.edu.cn