Representability for distributive quasi relation algebras via generalised ordinal sums

Andrew Craig^{1,2*}, Wilmari Morton¹ and Claudette Robinson¹

^{1*}Department of Mathematics and Applied Mathematics, University of Johannesburg, PO Box 524, Auckland Park, 2006, South Africa.
²National Institute for Theoretical and Computational Sciences (NITheCS), Johannesburg, South Africa.

*Corresponding author(s). E-mail(s): acraig@uj.ac.za; Contributing authors: wmorton@uj.ac.za; claudetter@uj.ac.za;

Abstract

We extend the work of Galatos (2004) on generalised ordinal sums of residuated lattices. We show that the generalised ordinal sum of an odd quasi relation algebra (qRA) satisfying certain conditions and an arbitrary qRA is again a qRA. In a recent paper by Craig and Robinson (2024), the notion of representability for distributive quasi relation algebras (DqRAs) was developed. For certain pairs of representable DqRAs, we prove that their generalised ordinal sum is again representable. An important consequence of this result is that finite Sugihara chains are finitely representable.

 ${\bf Keywords:}$ quasi relation algebra, representability, Sugihara chain, generalised ordinal sum

MSC Classification: 06F05, 03B47, 03G10

Quasi relation algebras (qRAs) were first described by Galatos and Jipsen [6]. On the one hand they can be viewed as a generalisation of relation algebras, or as involutive FL-algebras with an additional negation-like unary operation. Unlike relation algebras, the variety of qRAs has a decidable equational theory.

In this paper, we first extend the so-called *generalised ordinal sum* of Galatos [5] from residuated lattices to quasi relation algebras. When thinking about the order structure of the lattices, that construction can be considered in the following way: a

residuated lattice \mathbf{L} is inserted into a residuated lattice \mathbf{K} where it replaces the unit of the monoid operation. The requirements placed on \mathbf{L} and \mathbf{K} are discussed in Section 2. We essentially extend the conditions for residuated lattices to the additional unary operations in the signature of qRAs. A consequence of the conditions on \mathbf{K} is that its FL-algebra reduct will be odd (i.e. 0 = 1).

Having shown that the $\mathbf{K}[\mathbf{L}]$ construction gives us a method for constructing new qRAs from existing ones, we then turn to the question of representability.

Abstract relation algebras, first described by Tarski in 1941 [17], were designed to provide an abstract algebraic setting for studying binary relations. Jónsson and Tarski [12] asked whether all algebras were isomorphic to algebras of binary relations (with the monoid operation given by relational composition). This question was answered in the negative by Lyndon [13]. The task of representing relation algebras as algebras of binary relations has been an intriguing area of research ever since.

In [3], two of the current authors gave a definition of *representability* of distributive quasi relation algebras (see Section 3). Partially ordered sets equipped with an equivalence relation (and satisfying certain symmetry requirements) can be used to build distributive quasi relation algebras (DqRAs) as algebras of binary relations. This method was further exploited to give relational representations of Sugihara monoids [4].

In Section 4 we show that if $\mathbf{K} = \mathbf{S}_3$, and if \mathbf{L} is representable, then their generalised ordinal sum $\mathbf{K}[\mathbf{L}]$ will again be representable (Theorem 4.1). This is extended in Corollary 5.4 to the case of \mathbf{K} being any finite odd Sugihara chain. These results are similar in spirit to the result for integral relation algebras where the relation algebra obtained from \mathbf{A} and \mathbf{B} by Comer's construction [1] is representable if and only if \mathbf{A} and \mathbf{B} are representable.

A further application of Theorem 4.1 is given in Section 5, where we show that all finite Sugihara chains can be represented using a set of binary relations on a *finite* poset (Theorem 5.1). This improves the results in [15] and [4] where infinite posets (using copies of \mathbb{Q} , the rational numbers) were required to obtain a relational representation of \mathbf{S}_n for $n \ge 4$.

In Problem 1 we pose an intriguing question about the structures of possible algebras \mathbf{K} for which we can guarantee that $\mathbf{K}[\mathbf{L}]$ will be representable. Specifically we ask whether \mathbf{K} must be a finite Sugihara chain.

1 Preliminaries

In this section we recall the basic definitions of the algebras used in this paper. The reader is referred to [9] for more information.

1.1 Quasi relation algebras

A quasi relation algebra is an expansion of a residuated lattice [6]. Recall that a residuated lattice (RL) is an algebra $\mathbf{A} = \langle A, \wedge, \vee, \cdot, \rangle, /, 1 \rangle$ such that $\langle A, \cdot, 1 \rangle$ is a monoid, $\langle A, \wedge, \vee \rangle$ is a lattice and the monoid operation \cdot is residuated with residuals

 \setminus and /, i.e., for all $a, b, c \in A$,

$$a \cdot b \leqslant c \quad \Longleftrightarrow \quad a \leqslant c/b \quad \Longleftrightarrow \quad b \leqslant a \backslash c$$

A distributive residuated lattice **A** is a residuated lattice whose underlying lattice is distributive. Recall also that a residuated lattice **A** is commutative if $a \cdot b = b \cdot a$ for all $a, b \in A$. An element $a \in A$ is said to be *idempotent* if $a \cdot a = a^2 = a$.

If $a \leq 1$ or $1 \leq a$ for all $a \in A$, then **A** is called *conic*. We note that when the term conic first appeared in [10, Definition 2.3] it was applied to *commutative* residuated pomonoids, but since then the term has been used more generally.

A residuated lattice expanded with a constant 0, that is, $\mathbf{A} = \langle A, \wedge, \vee, \cdot, \rangle, /, 1, 0 \rangle$ is called a *Full Lambek (FL-)algebra* (cf. [9, Chapter 2.2]). No additional properties are assumed about the constant 0. An FL-algebra satisfying 1 = 0 is called *odd*.

On an FL-algebra, two unary operations $\sim : A \to A$ and $- : A \to A$, called *linear* negations, are defined in terms of the residuals and 0 as follows. For $a \in A$,

$$\sim a = a \setminus 0$$
 and $-a = 0/a$.

It follows that $\sim 1 = 1 \setminus 0 = 0 = 0/1 = -1$. By residuation, $\sim (a \lor b) = \sim a \land \sim b$ and $-(a \lor b) = -a \land -b$ for all $a, b \in A$.

We say that an FL-algebra **A** is *cyclic* if $-a = \sim a$ for all $a \in A$. It is easy to see that if an FL-algebra **A** is commutative, then it is cyclic.

If an FL-algbra \mathbf{A} satisfies the condition

$$(\mathsf{In}): \sim -a = a = -\sim a, \text{ for all } a \in A,$$

then it is called an *involutive Full Lambek (InFL-)algebra*. In an InFL-algebra **A** we have that $a \leq b \iff a \cdot (\sim b) \leq -1 \iff (-b) \cdot a \leq -1$, for all $a, b \in A$. Moreover, since both - and \sim are order-reversing, we get that - and \sim are dual lattice isomorphisms in an InFL-algebra.

The dual of \cdot is the binary operation $+ : A \times A \to A$ defined by $a + b = \sim (-a \cdot -b)$ for all $a, b \in A$. In an InFL-algebra we have that $a + b = -(\sim b \cdot \sim a)$ holds.

In [6, Lemma 2.2] Galatos and Jipsen show that an InFL-algebra is term-equivalent to an algebra $\mathbf{A} = \langle A, \wedge, \vee, \cdot, \sim, -, 1 \rangle$ such that $\langle A, \wedge, \vee \rangle$ is a lattice, $\langle A, \cdot, 1 \rangle$ is a monoid, and for all $a, b, c \in A$, we have

$$a \cdot b \leqslant c \quad \iff \quad a \leqslant -(b \cdot \sim c) \quad \iff \quad b \leqslant \sim (-c \cdot a).$$

We will usually write $\mathbf{A} = \langle A, \wedge, \vee, \cdot, \sim, -, 1, 0 \rangle$ for an InFL-algebra, where 0 = -1 = -1. The residuals can be expressed in terms of \cdot and the linear negations as follows:

$$c/b = -(b \cdot \sim c)$$
 and $a \setminus c = \sim (-c \cdot a)$. (1)

An InFL-algebra **A** can be expanded with an additional unary operation $\neg : A \to A$ to form an InFL'-algebra $\mathbf{A} = \langle A, \wedge, \vee, \cdot, \sim, -, \neg, 1, 0 \rangle$ such that $\neg \neg a = a$ for all $a \in A$.

We note that the involution is usually denoted by a prime, i.e., ', but for readability when using subscripts we will use \neg .

If an InFL'-algebra \mathbf{A} additionally satisfies the De Morgan law

$$(\mathsf{Dm}): \neg (a \lor b) = \neg a \land \neg b, \text{ for all } a, b, \in A$$

then **A** is called a *DmInFL'-algebra*. A *quasi relation algebra* (qRA) is a DmInFL'algebra $\mathbf{A} = \langle A, \wedge, \vee, \cdot, \sim, -, \neg, 1, 0 \rangle$ that satisfies, for all $a, b \in A$,

(Di):
$$\neg(\sim a) = -(\neg a)$$
 and (Dp): $\neg(a \cdot b) = \neg a + \neg b$.

The abbreviations (Di) and (Dp) stand for De Morgan involution and De Morgan product, respectively.

We note that if **A** is a qRA, then it can be shown that $\neg 1 = -1 = \sim 1 = 0$. An equivalent definition of a qRA is an FL'-algebra (i.e. an FL-algebra with an involutive unary operation ') that satisfies (Dm), (Di) and (Dp). Notice that if a DmFL'-algebra satisfies (In) and (Dp), then it can be shown that it also satisfies (Di) and therefore is a qRA (cf. [2, Lemma 1]).

Finally, a distributive quasi relation algebra (DqRA) is a quasi relation algebra $\mathbf{A} = \langle A, \wedge, \vee, \cdot, \sim, -, \neg, 1, 0 \rangle$ such that the underlying lattice $\langle A, \wedge, \vee \rangle$ is distributive.

In Figure 1 we give examples of small cyclic qRAs. Black nodes denote idempotent elements. The second diamond, \mathbf{K}_2 , in Figure 1 is an example of an odd qRA (hence -1 = -1 = -1 = -1 = -1). In [3, Figure 1] two examples of non-cyclic qRAs are given. All of these algebras were found with the help of Prover9/Mace4 [16].




Fig. 1 Examples of small qRAs.

An important class of examples of (distributive) quasi relation algebras is covered in more detail in the next subsection.

1.2 Sugihara chains

In Section 2 we will describe a construction, called the genaralised ordinal sum, that 'inserts' one algebra in another. In Section 4 we show that if the algebras satisfy certain conditions and are representable, then their generalised ordinal sum is representable too. Specifically, we will show that this is the case if the algebra into which the other is 'inserted' is a finite odd Sugihara chain. We therefore recall the basic definitions related to finite odd Sugihara chains.

Let $\mathbf{S} = \langle S, \wedge, \vee, \cdot, \rangle, /, 1 \rangle$ be a residuated lattice such that the monoid operation \cdot is commutative. Then \backslash and / collapse into a single operation denoted by \rightarrow , as $c \backslash b = b/c$ for all $b, c \in S$, and we use the signature $\mathbf{S} = \langle S, \wedge, \vee, \cdot, \rightarrow, 1 \rangle$ instead. The algebra \mathbf{S} is called idempotent if $a \cdot a = a$ for all $a \in S$. A commutative residuated lattice can then be expanded with a single linear negation, $\mathbf{S} = \langle S, \wedge, \vee, \cdot, \rightarrow, \cdot, 1 \rangle$, such that $\sim a = a$ and $a \rightarrow (\sim b) = b \rightarrow (\sim a)$ for all $a, b \in S$, to form an involutive commutative residuated lattice \mathbf{S} . A Sugihara monoid $\mathbf{S} = \langle S, \wedge, \vee, \cdot, \rightarrow, \cdot, 1 \rangle$ is an involutive commutative distributive idempotent residuated lattice. A Sugihara chain is a Sugihara monoid such that the underlying lattice $\langle S, \wedge, \vee \rangle$ is linearly ordered.

In the sequel we will only consider finite Sugihara chains. These are algebras $\mathbf{S}_n = \langle S_n, \wedge, \vee, \cdot, \rightarrow, \sim, 1 \rangle$ for $n \in \omega$ and $n \ge 2$. If n = 2k for k > 0 then $S_n = \{a_{-k}, \ldots, a_{-1}, a_1, \ldots, a_k\}$, and if n = 2k + 1 for k > 0 then $S_n = \{a_{-k}, \ldots, a_{-1}, a_0, a_1, \ldots, a_k\}$. The meet and join are defined as expected: $a_i \wedge a_j = a_{\min\{i,j\}}$ and $a_i \vee a_j = a_{\max\{i,j\}}$. The unary operation \sim is defined by $\sim a_j = a_{-j}$. The monoid operation is defined by

$$a_i \cdot a_j = \begin{cases} a_i & \text{if } |j| < |i| \\ a_j & \text{if } |i| < |j| \\ a_{\min\{i,j\}} & \text{if } |j| = |i|. \end{cases}$$

The implication $a_i \to a_j$ is defined by $\sim a_i \lor a_j$ if $i \leq j$ and $\sim a_i \land a_j$ if i > j. If n is odd then, the monoid identity is given by $1 = a_0$, and if n is even, then the monoid identity is $1 = a_1$. See Figure 2 for examples.

Given a finite Sugihara chain \mathbf{S}_n , we can set $0 = \sim 1$ and $-x = \sim x$ and $\neg x = \sim x$, which gives rise to a (distributive) quasi relation algebra. Expanding the signature of the reduct $\langle S_n, \wedge, \vee, \cdot, \sim, 1 \rangle$, we can view \mathbf{S}_n as the DqRA $\langle S_n, \wedge, \vee, \cdot, \sim, \sim, \cdot, 1, 0 \rangle$.

2 Generalised ordinal sums of qRAs

We first recall some definitions which are used in the original generalised ordinal sum result [5, Lemma 6.1] (see also [9, Lemma 9.6.2]). We note that our description makes a small correction to the original requirements for residuated lattices.

An element b in an algebra \mathbf{A} is said to be *irreducible* with respect to an n-ary operation (where n is a positive integer) f of \mathbf{A} if $f(b_1, \ldots, b_n) = b$ implies $b_i = b$ for some $i \in \{1, \ldots, n\}$, for all $b_1, \ldots, b_n \in A$. If $b \in A$ is irreducible with respect to all of the non-nullary operations of \mathbf{A} , then we call b totally irreducible.




Fig. 2 Finite Sugihara chains S_2 to S_5 .

Let $\mathbf{K} = \langle K, \wedge_{\mathbf{K}}, \vee_{\mathbf{K}}, \vee_{\mathbf{K}}, \backslash_{\mathbf{K}}, \mathbf{I}_{\mathbf{K}} \rangle$ and $\mathbf{L} = \langle L, \wedge_{\mathbf{L}}, \vee_{\mathbf{L}}, \cdot_{\mathbf{L}}, \mathbf{I}_{\mathbf{L}}, \mathbf{I}_{\mathbf{L}} \rangle$ be residuated lattices such that $K \cap L = \emptyset$ and $\mathbf{1}_{\mathbf{K}}$ is totally irreducible. Let $K[L] = (K \setminus \{\mathbf{1}_{\mathbf{K}}\}) \cup L$ and $\mathbf{1}_{\mathbf{K}[\mathbf{L}]} = \mathbf{1}_{\mathbf{L}}$. For $m_0, m_1 \in (K \setminus \{\mathbf{1}_{\mathbf{K}}\}) \cup L$ and $\star \in \{\cdot, \backslash, /\}$, we define $\star_{\mathbf{K}[\mathbf{L}]}$ by

$$m_0 \star_{\mathbf{K}[\mathbf{L}]} m_1 = \begin{cases} m_0 \star_{\mathbf{K}} m_1 & \text{if } m_0, m_1 \in K \setminus \{\mathbf{1}_{\mathbf{K}}\} \\ m_0 \star_{\mathbf{L}} m_1 & \text{if } m_0, m_1 \in L \\ m_0 \star_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} & \text{if } m_0 \in K \setminus \{\mathbf{1}_{\mathbf{K}}\} \text{ and } m_1 \in L \\ \mathbf{1}_{\mathbf{K}} \star_{\mathbf{K}} m_1 & \text{if } m_0 \in L \text{ and } m_1 \in K \setminus \{\mathbf{1}_{\mathbf{K}}\}; \end{cases}$$
(2)

and $\wedge_{\mathbf{K}[\mathbf{L}]}$ and $\vee_{\mathbf{K}[\mathbf{L}]}$ are defined by,

$$m_0 \wedge_{\mathbf{K}[\mathbf{L}]} m_1 = \begin{cases} m_0 \wedge_{\mathbf{K}} m_1 & \text{if } m_0, m_1 \in K \setminus \{\mathbf{1}_{\mathbf{K}}\} \\ m_0 \wedge_{\mathbf{L}} m_1 & \text{if } m_0, m_1 \in L \\ m_i & \text{if } m_i \in L, m_{1-i} \in K \setminus \{\mathbf{1}_{\mathbf{K}}\} \text{ and } m_{1-i} \geqslant_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} \\ m_{1-i} \wedge_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} & \text{if } m_i \in L, m_{1-i} \in K \setminus \{\mathbf{1}_{\mathbf{K}}\} \text{ and } m_{1-i} \not\geqslant_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} \end{cases}$$
(3)

and

$$m_0 \vee_{\mathbf{K}[\mathbf{L}]} m_1 = \begin{cases} m_0 \vee_{\mathbf{K}} m_1 & \text{if } m_0, m_1 \in K \setminus \{\mathbf{1}_{\mathbf{K}}\} \\ m_0 \vee_{\mathbf{L}} m_1 & \text{if } m_0, m_1 \in L \\ m_i & \text{if } m_i \in L, m_{1-i} \in K \setminus \{\mathbf{1}_{\mathbf{K}}\} \text{ and } m_{1-i} \leqslant_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} \\ m_{1-i} \vee_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} & \text{if } m_i \in L, m_{1-i} \in K \setminus \{\mathbf{1}_{\mathbf{K}}\} \text{ and } m_{1-i} \notin_{\mathbf{K}} \mathbf{1}_{\mathbf{K}}. \end{cases}$$
(4)

In the original definition of the $\mathbf{K}[\mathbf{L}]$ construction [5], it was stated that in addition to $\mathbf{1}_{\mathbf{K}}$ being totally irreducible, \mathbf{K} also needed to satisfy:

$$k \setminus_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} \neq \mathbf{1}_{\mathbf{K}} \quad \text{and} \quad \mathbf{1}_{\mathbf{K}} / \mathbf{K} \neq \mathbf{1}_{\mathbf{K}} \quad \text{for all } k \in K \setminus \{\mathbf{1}_{\mathbf{K}}\}$$
(5)

or, \mathbf{L} needed to be integral. We prove in Lemma 2.1 below that, in fact, the condition above is not required.

Lemma 2.1. Let $\mathbf{K} = \langle K, \wedge_{\mathbf{K}}, \vee_{\mathbf{K}}, \cdot_{\mathbf{K}}, \rangle_{\mathbf{K}}, 1_{\mathbf{K}} \rangle$ be a residuated lattice with $1_{\mathbf{K}}$ irreducible with respect to $\backslash_{\mathbf{K}}$ and $/_{\mathbf{K}}$. Then \mathbf{K} satisfies (5).

Proof. Assume that $1_{\mathbf{K}}$ is irreducible with respect to $\backslash_{\mathbf{K}}$ and $/_{\mathbf{K}}$, and then suppose that there exists $k \neq 1_{\mathbf{K}}$ such that $1_{\mathbf{K}}/_{\mathbf{K}}k = 1_{\mathbf{K}}$ or $k \backslash_{\mathbf{K}} 1_{\mathbf{K}} = 1_{\mathbf{K}}$.

If $1_{\mathbf{K}/\mathbf{K}}k = 1_{\mathbf{K}}$, we get $1_{\mathbf{K}} \leq 1_{\mathbf{K}/\mathbf{K}}k$, and so by residuation $k = 1_{\mathbf{K}} \cdot_{\mathbf{K}} k \leq 1_{\mathbf{K}}$. Since $/_{\mathbf{K}}$ is order-preserving in the first argument, we have $k/_{\mathbf{K}}k \leq 1_{\mathbf{K}}/_{\mathbf{K}}k = 1_{\mathbf{K}}$. By residuation, we always have $1_{\mathbf{K}} \leq k/_{\mathbf{K}}k$ so $k/_{\mathbf{K}}k = 1_{\mathbf{K}}$. But this is a contradiction since $1_{\mathbf{K}}$ is irreducible with respect to $/_{\mathbf{K}}$.

If $k \setminus_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} = \mathbf{1}_{\mathbf{K}}$, we follow a similar argument to reach a contradiction. Hence, no such k can exist and so **K** satisfies (5).

We further note that the definitions of $\wedge_{\mathbf{K}[\mathbf{L}]}$ and $\vee_{\mathbf{K}[\mathbf{L}]}$ given above differ from the definitions given in [5]. This is to ensure closure of $(K \setminus \{\mathbf{1}_{\mathbf{K}}\}) \cup L$ under meet and join. For example, if $\mathbf{K} = \{a, \mathbf{1}_{\mathbf{K}}, b\}$ with $a < \mathbf{1}_{\mathbf{K}} < b$, then under the original definition for any $c \in L$ we get $b \wedge_{\mathbf{K}[\mathbf{L}]} c = \mathbf{1}_{\mathbf{K}} \notin K[L]$.

Lemma 2.2 ([5, Lemma 6.1]). If **K** and **L** are residuated lattices such that $1_{\mathbf{K}}$ is totally irreducible, then the structure $\mathbf{K}[\mathbf{L}] = \langle K[L], \wedge_{\mathbf{K}[\mathbf{L}]}, \vee_{\mathbf{K}[\mathbf{L}]}, \backslash_{\mathbf{K}[\mathbf{L}]}, |\mathbf{K}_{\mathbf{K}[\mathbf{L}]}, 1_{\mathbf{K}[\mathbf{L}]} \rangle$, as defined above, is a residuated lattice.

The algebra $\mathbf{K}[\mathbf{L}]$ is called the *generalised ordinal sum* of \mathbf{K} and \mathbf{L} . It is easy to see that if both \mathbf{K} and \mathbf{L} are commutative, then so is $\mathbf{K}[\mathbf{L}]$. Any idempotent element of \mathbf{K} or \mathbf{L} is also idempotent in $\mathbf{K}[\mathbf{L}]$.

The following lemma is an immediate consequence of the construction described above.

Lemma 2.3. Let $\mathbf{K}[\mathbf{L}]$ be the generalised ordinal sum of residuated lattices \mathbf{K} and \mathbf{L} such that $\mathbf{1}_{\mathbf{K}}$ is totally irreducible. Let $k_0, k_1 \in K \setminus \{\mathbf{1}_{\mathbf{K}}\}$ and $\ell_0, \ell_1 \in L$, then

$$k_0 \leq_{\mathbf{K}[\mathbf{L}]} \ell_0 \iff k_0 \leq_{\mathbf{K}} 1_{\mathbf{K}}, \qquad \ell_0 \leq_{\mathbf{K}[\mathbf{L}]} k_0 \iff k_0 \geq_{\mathbf{K}} 1_{\mathbf{K}},$$
$$k_0 \leq_{\mathbf{K}[\mathbf{L}]} k_1 \iff k_0 \leq_{\mathbf{K}} k_1, \qquad \ell_0 \leq_{\mathbf{K}[\mathbf{L}]} \ell_1 \iff \ell_0 \leq_{\mathbf{L}} \ell_1.$$

Let $\mathbf{K} = \langle K, \wedge_{\mathbf{K}}, \vee_{\mathbf{K}}, \vee_{\mathbf{K}}, |_{\mathbf{K}}, 1_{\mathbf{K}}, 0_{\mathbf{K}} \rangle$ and $\mathbf{L} = \langle L, \wedge_{\mathbf{L}}, \vee_{\mathbf{L}}, \setminus_{\mathbf{L}}, 1_{\mathbf{L}}, 0_{\mathbf{L}} \rangle$ be FL-algebras. Recall that $K[L] = (K \setminus \{\mathbf{1}_{\mathbf{K}}\}) \cup L$, let $\wedge_{\mathbf{K}[\mathbf{L}]}, \vee_{\mathbf{K}[\mathbf{L}]}, \cdot_{\mathbf{K}[\mathbf{L}]}$ and $\mathbf{1}_{\mathbf{K}[\mathbf{L}]}$ be as defined before and set $\mathbf{0}_{\mathbf{K}[\mathbf{L}]} = \mathbf{0}_{\mathbf{L}}$. If $\mathbf{1}_{\mathbf{K}}$ is totally irreducible, then an immediate consequence of Lemma 2.2 is that the generalised ordinal sum $\mathbf{K}[\mathbf{L}] = \langle K[L], \wedge_{\mathbf{K}[\mathbf{L}]}, \vee_{\mathbf{K}[\mathbf{L}]}, \cdot_{\mathbf{K}[\mathbf{L}]}, \mathbf{1}_{\mathbf{K}[\mathbf{L}]}, \mathbf{0}_{\mathbf{K}[\mathbf{L}]} \rangle$ of \mathbf{K} and \mathbf{L} is an FL-algebra. Recall that for FL-algebras \mathbf{K}, \mathbf{L} and $\mathbf{K}[\mathbf{L}]$, the linear negations, $\sim_{\mathbf{K}}, -_{\mathbf{K}}$ and $\sim_{\mathbf{L}}, -_{\mathbf{L}}$ and $\sim_{\mathbf{K}[\mathbf{L}]}, -_{\mathbf{K}[\mathbf{L}]}$ can be defined on \mathbf{K}, \mathbf{L} and $\mathbf{K}[\mathbf{L}]$, respectively, in terms of the residuals and the 0. In particular, for $m \in K[L]$,

$$\sim_{\mathbf{K}[\mathbf{L}]} m = m \setminus_{\mathbf{K}[\mathbf{L}]} 0_{\mathbf{K}[\mathbf{L}]} \text{ and } -_{\mathbf{K}[\mathbf{L}]} m = 0_{\mathbf{K}[\mathbf{L}]} / _{\mathbf{K}[\mathbf{L}]} m$$

Lemma 2.4. Let $\mathbf{K} = \langle K, \wedge_{\mathbf{K}}, \vee_{\mathbf{K}}, \vee_{\mathbf{K}}, |_{\mathbf{K}}, 1_{\mathbf{K}}, 0_{\mathbf{K}} \rangle$ be an FL-algebra such that $1_{\mathbf{K}}$ is irreducible with respect to the residuals. Then $1_{\mathbf{K}}$ is irreducible with respect to $\sim_{\mathbf{K}}$ and $-_{\mathbf{K}}$.

Proof. We consider the case for $-\mathbf{K}$. Assume $-\mathbf{K}k = 1\mathbf{K}$. Then $0\mathbf{K}\setminus\mathbf{K}k = 1\mathbf{K}$, and so, since $1\mathbf{K}$ is irreducible with respect to $\setminus\mathbf{K}$, we have $0\mathbf{K} = 1\mathbf{K}$ or $k = 1\mathbf{K}$. If $k = 1\mathbf{K}$, we are done, so assume $0\mathbf{K} = 1\mathbf{K}$. Then $k = 1\mathbf{K}\setminus\mathbf{K}k = 0\mathbf{K}\setminus\mathbf{K}k = 1\mathbf{K}$, as required.

The following equivalent statements for an InFL-algebra show that the total irreducibility of the monoid identity is independent of our choice of signature for an InFL-algebra and can be characterised as precisely those InFL-algebras that are odd.

Lemma 2.5. Let $\mathbf{K} = \langle K, \wedge_{\mathbf{K}}, \vee_{\mathbf{K}}, \sim_{\mathbf{K}}, -\mathbf{K}, \mathbf{1}_{\mathbf{K}}, \mathbf{0}_{\mathbf{K}} \rangle$ be an InFL-algebra such that $\mathbf{1}_{\mathbf{K}}$ is irreducible with respect to $\cdot_{\mathbf{K}}$. Then the following statements are equivalent:

- (i) $1_{\mathbf{K}}$ is irreducible with respect to $\sim_{\mathbf{K}}$ and $-_{\mathbf{K}}$.
- (ii) $1_{\mathbf{K}}$ is irreducible with respect to $\setminus_{\mathbf{K}}$ and $/_{\mathbf{K}}$.
- (iii) **K** is odd, i.e., $1_{\mathbf{K}} = 0_{\mathbf{K}}$.

Proof. (i) \Rightarrow (ii): Assume that **K** is an InFL-algebra such that $1_{\mathbf{K}}$ is irreducible with respect to $\cdot_{\mathbf{K}}$, $\sim_{\mathbf{K}}$ and $-_{\mathbf{K}}$ and let $k_1, k_2 \in K$ such that $k_1 \setminus_{\mathbf{K}} k_2 = 1_{\mathbf{K}}$. Then by (1) we have that $k_1 \setminus_{\mathbf{K}} k_2 = \sim_{\mathbf{K}} (-_{\mathbf{K}} k_2 \cdot k_1) = 1_{\mathbf{K}}$ which implies that $-_{\mathbf{K}} k_2 \cdot k_1 = 1_{\mathbf{K}}$, hence $-_{\mathbf{K}} k_2 = 1_{\mathbf{K}}$ or $k_1 = 1_{\mathbf{K}}$ and therefore $k_2 = 1_{\mathbf{K}}$ or $k_1 = 1_{\mathbf{K}}$ since $1_{\mathbf{K}}$ is irreducible with respect to $\cdot_{\mathbf{K}}$, $\sim_{\mathbf{K}}$ and $-_{\mathbf{K}}$, respectively. That is, $1_{\mathbf{K}}$ is irreducible with respect to $\setminus_{\mathbf{K}}$. The proof that $1_{\mathbf{K}}$ is irreducible with respect to $/_{\mathbf{K}}$ follow similarly.

(ii) \Rightarrow (i): This implication holds even in the case of FL-algebras as shown in Lemma 2.4.

(i) \Rightarrow (iii): Assume that **K** is an InFL-algebra such that $1_{\mathbf{K}}$ is irreducible with respect to $\cdot_{\mathbf{K}}$, $\sim_{\mathbf{K}}$ and $-_{\mathbf{K}}$. In any FL-algebra we have $\sim_{\mathbf{K}} 1_{\mathbf{K}} = 0_{\mathbf{K}}$, and so by (ln), $1_{\mathbf{K}} = -_{\mathbf{K}} \sim_{\mathbf{K}} 1_{\mathbf{K}} = -_{\mathbf{K}} 0_{\mathbf{K}}$ and finally since $1_{\mathbf{K}}$ is irreducible with respect to $-_{\mathbf{K}}$ we have that $0_{\mathbf{K}} = 1_{\mathbf{K}}$.

(iii) \Rightarrow (i): Assume that **K** is an odd InFL-algebra such that $\mathbf{1}_{\mathbf{K}}$ is irreducible with respect to $\cdot_{\mathbf{K}}$. Let $k \in K$ such that $\sim_{\mathbf{K}} k = \mathbf{1}_{\mathbf{K}}$. Then $k = -\mathbf{K} \sim_{\mathbf{K}} k = -\mathbf{K} \mathbf{1}_{\mathbf{K}} = \mathbf{0}_{\mathbf{K}} = \mathbf{1}_{\mathbf{K}}$. If $-\mathbf{K}k = \mathbf{1}_{\mathbf{K}}$, then $k = \sim_{\mathbf{K}} -\mathbf{K}k = \sim_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} = \mathbf{0}_{\mathbf{K}} = \mathbf{1}_{\mathbf{K}}$.

Now consider InFL-algebras $\mathbf{K} = \langle K, \wedge_{\mathbf{K}}, \vee_{\mathbf{K}}, -_{\mathbf{K}}, -_{\mathbf{K}}, 1_{\mathbf{K}}, 0_{\mathbf{K}} \rangle$ and $\mathbf{L} = \langle L, \wedge_{\mathbf{L}}, \vee_{\mathbf{L}}, -_{\mathbf{L}}, -_{\mathbf{L}}, 1_{\mathbf{L}}, 0_{\mathbf{L}} \rangle$ such that $1_{\mathbf{K}}$ is totally irreducible (and hence, **K** is odd). If $K[L] = (K \setminus \{1_{\mathbf{K}}\}) \cup L$ with $\wedge_{\mathbf{K}[\mathbf{L}]}, \vee_{\mathbf{K}[\mathbf{L}]}, -_{\mathbf{K}[\mathbf{L}]}, -_{\mathbf{K}[\mathbf{L}]}, 1_{\mathbf{K}[\mathbf{L}]}$ and $0_{\mathbf{K}[\mathbf{L}]}$ defined as before, then $\mathbf{K}[\mathbf{L}] = \langle K[L], \wedge_{\mathbf{K}[\mathbf{L}]}, \vee_{\mathbf{K}[\mathbf{L}]}, \cdots_{\mathbf{K}[\mathbf{L}]}, -_{\mathbf{K}[\mathbf{L}]}, 1_{\mathbf{K}[\mathbf{L}]}, 0_{\mathbf{K}[\mathbf{L}]} \rangle$ is the generalised ordinal sum of **K** and **L**.

Lemma 2.6. Let $\mathbf{K}[\mathbf{L}]$ be the generalised ordinal sum of InFL-algebras \mathbf{K} and \mathbf{L} , as defined above with $\mathbf{1}_{\mathbf{K}}$ totally irreducible (implying that \mathbf{K} is odd). Then, for $m \in K[L]$, we have

$$\sim_{\mathbf{K}[\mathbf{L}]} m = \begin{cases} \sim_{\mathbf{L}} m & \text{if } m \in L \\ \sim_{\mathbf{K}} m & \text{if } m \in K \setminus \{1_{\mathbf{K}}\}, \end{cases} \quad and \quad -_{\mathbf{K}[\mathbf{L}]} m = \begin{cases} -_{\mathbf{L}} m & \text{if } m \in L \\ -_{\mathbf{K}} m & \text{if } m \in K \setminus \{1_{\mathbf{K}}\}. \end{cases}$$

Proof. Let $m \in K[L]$. If $m \in L$, then $\sim_{\mathbf{K}[\mathbf{L}]} m = m \setminus_{\mathbf{K}[\mathbf{L}]} 0_{\mathbf{K}[\mathbf{L}]} = m \setminus_{\mathbf{L}} 0_{\mathbf{L}} = \sim_{\mathbf{L}} m$ and $-_{\mathbf{K}[\mathbf{L}]} m = 0_{\mathbf{K}[\mathbf{L}]} /_{\mathbf{K}[\mathbf{L}]} m = 0_{\mathbf{L}} /_{\mathbf{L}} m = -_{\mathbf{L}} m$.

Recall that since $1_{\mathbf{K}}$ is totally irreducible, **K** is odd. If $m \in K \setminus \{1_{\mathbf{K}}\}$, then $\sim_{\mathbf{K}[\mathbf{L}]} m = m \setminus_{\mathbf{K}[\mathbf{L}]} 0_{\mathbf{K}[\mathbf{L}]} = m \setminus_{\mathbf{K}[\mathbf{L}]} 0_{\mathbf{L}} = m \setminus_{\mathbf{K}} 1_{\mathbf{K}} = m \setminus_{\mathbf{K}} 0_{\mathbf{K}} = \sim_{\mathbf{K}} m$ and $-_{\mathbf{K}[\mathbf{L}]} m = 0_{\mathbf{K}[\mathbf{L}]} / \mathbf{K}_{[\mathbf{L}]} m = 0_{\mathbf{L}} / \mathbf{K}_{[\mathbf{L}]} m = 1_{\mathbf{K}} / \mathbf{K} m = 0_{\mathbf{K}} / \mathbf{K} m = -\mathbf{K} m$.

The following proposition follows immediately from Lemmas 2.2, 2.5 and 2.6 and the fact that \mathbf{K} and \mathbf{L} satisfy (In).

Proposition 2.7. If **K** and **L** are (cyclic) InFL-algebras such that $1_{\mathbf{K}}$ is totally irreducible (implying that **K** is odd), then their generalised ordinal sum $\mathbf{K}[\mathbf{L}]$ is a (cyclic) InFL-algebra.

Proof. Let $m \in \mathbf{K}[\mathbf{L}]$. If $m \in K \setminus \{\mathbf{1}_{\mathbf{K}}\}$ then $\sim_{\mathbf{K}[\mathbf{L}]} -_{\mathbf{K}[\mathbf{L}]} m = \sim_{\mathbf{K}} -_{\mathbf{K}} m = m$ since \mathbf{K} is an InFL-algebra. Similarly for $m \in L$.

Next let $\mathbf{K} = \langle K, \wedge_{\mathbf{K}}, \vee_{\mathbf{K}}, -\mathbf{\kappa}, -\mathbf{\kappa}, -\mathbf{\kappa}, \mathbf{1}_{\mathbf{K}}, \mathbf{0}_{\mathbf{K}} \rangle$ and $\mathbf{L} = \langle L, \wedge_{\mathbf{L}}, \vee_{\mathbf{L}}, -\mathbf{L}, -\mathbf{L}, -\mathbf{L}, \mathbf{1}_{\mathbf{L}}, \mathbf{0}_{\mathbf{L}} \rangle$ be InFL'-algebras with $\mathbf{1}_{\mathbf{K}}$ totally irreducible, then their generalised ordinal sum is the algebra $\mathbf{K}[\mathbf{L}] = \langle K[L], \wedge_{\mathbf{K}[\mathbf{L}]}, \vee_{\mathbf{K}[\mathbf{L}]}, -\mathbf{\kappa}_{\mathbf{K}[\mathbf{L}]}, -\mathbf{\kappa}_{\mathbf{K}[\mathbf{L}]}, \mathbf{1}_{\mathbf{K}[\mathbf{L}]}, \mathbf{0}_{\mathbf{K}[\mathbf{L}]} \rangle$ where the InFL-reduct of $\mathbf{K}[\mathbf{L}]$ is as defined above, and for $m \in K[L]$,

$$\neg_{\mathbf{K}[\mathbf{L}]} m = \begin{cases} \neg_{\mathbf{L}} m & \text{if } m \in L \\ \neg_{\mathbf{K}} m & \text{if } m \in K \setminus \{\mathbf{1}_{\mathbf{K}}\} \end{cases}$$

Given this definition of $\neg_{\mathbf{K}[\mathbf{L}]}$ it is immediate that $\mathbf{K}[\mathbf{L}]$ is an InFL'-algebra whenever \mathbf{K} and \mathbf{L} are.

Finally, assume that **K** are **L** are qRAs. Since (Di) implies (In), it follows as above that if $1_{\mathbf{K}}$ is totally irreducible, then **K** is odd.

Proposition 2.8. If **K** and **L** are qRAs such that $1_{\mathbf{K}}$ is totally irreducible (and hence **K** is odd), then their generalised ordinal sum $\mathbf{K}[\mathbf{L}]$ is a qRA.

Proof. To prove that $\mathbf{K}[\mathbf{L}]$ is a qRA, we need to show that it satisfies (Dm), (Di) and (Dp). Let $m_0, m_1 \in (K \setminus \{1_{\mathbf{K}}\}) \cup L$. We first show that $\mathbf{K}[\mathbf{L}]$ satisfies (Dm).

If $m_0, m_1 \in K \setminus \{1_{\mathbf{K}}\}$ or $m_0, m_1 \in L$, then (Dm) is inherited from **K** or **L**, respectively. For the remaining cases, we can assume without loss of generality that $m_0 \in K \setminus \{1_{\mathbf{K}}\}$ and $m_1 \in L$ since $\wedge_{\mathbf{K}[\mathbf{L}]}$ and $\vee_{\mathbf{K}[\mathbf{L}]}$ are commutative. Now consider two cases. If $m_0 \geq_{\mathbf{K}} 1_{\mathbf{K}}$, then $\neg_{\mathbf{K}} m_0 \leq_{\mathbf{K}} \neg_{\mathbf{K}} 1_{\mathbf{K}} = 0_{\mathbf{K}} = 1_{\mathbf{K}}$. Using this and Lemma 2.3 we have that

 $\neg_{\mathbf{K}[\mathbf{L}]}(m_0 \wedge_{\mathbf{K}[\mathbf{L}]} m_1) = \neg_{\mathbf{K}[\mathbf{L}]} m_1 = \neg_{\mathbf{L}} m_1 = \neg_{\mathbf{K}} m_0 \vee_{\mathbf{K}[\mathbf{L}]} \neg_{\mathbf{L}} m_1 = \neg_{\mathbf{K}[\mathbf{L}]} m_0 \vee_{\mathbf{K}[\mathbf{L}]} \neg_{\mathbf{K}[\mathbf{L}]} m_1.$

On the other hand, assume that $m_0 \not\geq_{\mathbf{K}} 1_{\mathbf{K}}$. Since **K** is odd and **K** satisfies (Dm), we get

$$\neg_{\mathbf{K}[\mathbf{L}]}(m_0 \wedge_{\mathbf{K}[\mathbf{L}]} m_1) = \neg_{\mathbf{K}[\mathbf{L}]}(m_0 \wedge_{\mathbf{K}} 1_{\mathbf{K}}) = \neg_{\mathbf{K}}(m_0 \wedge_{\mathbf{K}} 1_{\mathbf{K}}) = \neg_{\mathbf{K}} m_0 \vee_{\mathbf{K}} \neg_{\mathbf{K}} 1_{\mathbf{K}} = \neg_{\mathbf{K}} m_0 \vee_{\mathbf{K}} 0 \vee_{\mathbf{K}} 1_{\mathbf{K}} = \neg_{\mathbf{K}} m_0 \vee_{\mathbf{K}} 0 \vee_{\mathbf{K}} 1_{\mathbf{K}} = \neg_{\mathbf{K}} m_0 \vee_{\mathbf{K}[\mathbf{L}]} \cdots 1_{\mathbf{K}} m_1 = \neg_{\mathbf{K}[\mathbf{L}]} m_0 \vee_{\mathbf{K}[\mathbf{L}]} \cdots 1_{\mathbf{K}[\mathbf{L}]} m_1.$$

(Di) follows directly from the definition of $\neg_{\mathbf{K}[\mathbf{L}]}$, Lemma 2.6 and the fact that **K** and **L** satisfy (Di).

Finally, we show that $\mathbf{K}[\mathbf{L}]$ satisfies (Dp). As with (Dm), if $m_0, m_1 \in K \setminus \{1_{\mathbf{K}}\}$ or $m_0, m_1 \in L$, then (Dp) is inherited from \mathbf{K} or \mathbf{L} , respectively. Next assume that $m_0 \in K \setminus \{1_{\mathbf{K}}\}$ and $m_1 \in L$. Then,

$$\neg_{\mathbf{K}[\mathbf{L}]}(m_0 \cdot_{\mathbf{K}[\mathbf{L}]} m_1) = \neg_{\mathbf{K}[\mathbf{L}]}(m_0 \cdot_{\mathbf{K}} 1_{\mathbf{K}}) = \neg_{\mathbf{K}[\mathbf{L}]} m_0 = \neg_{\mathbf{K}} m_0.$$

Since (Di) implies (In) we also have

$$\sim_{\mathbf{K}[\mathbf{L}]} (-_{\mathbf{K}[\mathbf{L}]} (\neg_{\mathbf{K}[\mathbf{L}]} m_0) \cdot_{\mathbf{K}[\mathbf{L}]} -_{\mathbf{K}[\mathbf{L}]} (\neg_{\mathbf{K}[\mathbf{L}]} m_1)) = \sim_{\mathbf{K}[\mathbf{L}]} (-_{\mathbf{K}} (\neg_{\mathbf{K}} m_0) \cdot_{\mathbf{K}[\mathbf{L}]} -_{\mathbf{L}} (\neg_{\mathbf{L}} m_1))$$
$$= \sim_{\mathbf{K}[\mathbf{L}]} (-_{\mathbf{K}} (\neg_{\mathbf{K}} m_0) \cdot_{\mathbf{K}} \mathbf{1}_{\mathbf{K}})$$
$$= \sim_{\mathbf{K}[\mathbf{L}]} -_{\mathbf{K}} (\neg_{\mathbf{K}} m_0)$$
$$= \sim_{\mathbf{K}} -_{\mathbf{K}} (\neg_{\mathbf{K}} m_0)$$
$$= \neg_{\mathbf{K}} m_0.$$

The case where $m_0 \in L$ and $m_1 \in K \setminus \{1_{\mathbf{K}}\}$ follows similarly.

Now we give an example of the generalised ordinal sum $\mathbf{K}[\mathbf{L}]$ for qRAs \mathbf{K} and \mathbf{L} .

Example 2.9. Let \mathbf{K}_1 be the three-element Sugihara chain \mathbf{S}_3 (depicted in Figure 3). Then $\mathbf{1}_{\mathbf{K}_1}$ is totally irreducible. Let \mathbf{L}_1 be the four-element algebra also depicted in Figure 3 (also see Figure 1 for a full description). Since \mathbf{K}_1 and \mathbf{L}_1 are qRAs such that $\mathbf{1}_{\mathbf{K}_1}$ is totally irreducible, it follows from Proposition 2.8 that their generalised ordinal sum $\mathbf{K}_1[\mathbf{L}_1]$ (the final algebra depicted in Figure 3) is a qRA. The non-lattice operations are defined via Proposition 2.8.




Fig. 3 qRAs K_1 , L_1 and their generalised ordinal sum, the qRA $K_1[L_1]$.

For our representation results in Section 4 we require that $\mathbf{K}[\mathbf{L}]$ be distributive. Hence we are interested in when distributivity will be preserved.

Proposition 2.10. Let **K** and **L** be qRAs such that |L| > 1 and $1_{\mathbf{K}}$ is totally irreducible. If there exists $k \in K$ such that $k \notin_{\mathbf{K}} 1_{\mathbf{K}}$ and $1_{\mathbf{K}} \notin_{\mathbf{K}} k$, then $\mathbf{K}[\mathbf{L}]$ is non-distributive.

Proof. Let $k \in K$ such that $k \notin_{\mathbf{K}} \mathbf{1}_{\mathbf{K}}$ and $\mathbf{1}_{\mathbf{K}} \notin_{\mathbf{K}} k$. Set $m_0 = k \wedge_{\mathbf{K}} \mathbf{1}_{\mathbf{K}}$ and $m_1 = k \vee_{\mathbf{K}} \mathbf{1}_{\mathbf{K}}$. Then $m_0, m_1 \neq \mathbf{1}_{\mathbf{K}}$ and $m_0, m_1 \neq k$ and $\{k, m_0, m_1, \mathbf{1}_{\mathbf{K}}\}$ is a sublattice of K. Since L is a lattice and |L| > 1, there exist $\ell_0, \ell_1 \in L$ such that $\ell_0 <_{\mathbf{L}} \ell_1$. Clearly $\ell_0 \wedge_{\mathbf{K}[\mathbf{L}]} \ell_1 = \ell_0$ and $\ell_0 \vee_{\mathbf{K}[\mathbf{L}]} \ell_1 = \ell_1$. Also, $m_1 \wedge_{\mathbf{K}[\mathbf{L}]} \ell_i = \ell_i$ and $m_0 \vee_{\mathbf{K}[\mathbf{L}]} \ell_i = \ell_i$ for $i \in \{0, 1\}$. Moreover, for $d_0 \in \{k, m_0\}, d_1 \in \{k, m_1\},$

$$d_0 \wedge_{\mathbf{K}[\mathbf{L}]} \ell_0 = d_0 \wedge_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} = m_0 \qquad \text{and} \qquad d_0 \wedge_{\mathbf{K}[\mathbf{L}]} \ell_1 = d_0 \wedge_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} = m_0$$
$$d_1 \vee_{\mathbf{K}[\mathbf{L}]} \ell_0 = d_1 \vee_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} = m_1 \qquad \text{and} \qquad d_1 \vee_{\mathbf{K}[\mathbf{L}]} \ell_1 = d_1 \vee_{\mathbf{K}} \mathbf{1}_{\mathbf{K}} = m_1.$$

Thus, $\{k, m_0, m_1, \ell_0, \ell_1\}$ is a sublattice of the underlying lattice of $\mathbf{K}[\mathbf{L}]$ isomorphic to N_5 .

Example 2.11. Let \mathbf{K}_2 be the four-element algebra depicted in Figure 4 (also see the second diamond in Figure 1). Then $\mathbf{1}_{\mathbf{K}_2}$ is totally irreducible. Let \mathbf{L}_2 be the two-element Sugihara chain \mathbf{S}_2 (also depicted in Figure 4). Since \mathbf{K}_2 and \mathbf{L}_2 are both qRAs, it follows from Proposition 2.8 that $\mathbf{K}_2[\mathbf{L}_2]$ is again a qRA. However, since $a \leq \mathbf{1}_{\mathbf{K}_2}$ and $\mathbf{1}_{\mathbf{K}_2} \leq a$ it follows from Proposition 2.10 that $\mathbf{K}_2[\mathbf{L}_2]$ is not a DqRA. Indeed, $\mathbf{K}_2[\mathbf{L}_2]$ (also depicted in Figure 4) has its lattice reduct isomorphic to N_5 .




Fig. 4 DqRAs \mathbf{K}_2 , \mathbf{L}_2 and their non-distributive generalised ordinal sum $\mathbf{K}_2[\mathbf{L}_2]$.

Recall that a DqRA **K** is called *conic* if every $a \in K$ satisfies $a \leq_{\mathbf{K}} 1$ or $1 \leq_{\mathbf{K}} a$.

Theorem 2.12. Let **K** and **L** be DqRAs such that **K** is conic and $1_{\mathbf{K}}$ is totally irreducible, then their generalised ordinal sum $\mathbf{K}[\mathbf{L}]$ is a DqRA.

Proof. It follows from Proposition 2.8 that $\mathbf{K}[\mathbf{L}]$ is a qRA so we need only confirm that the underlying lattice is distributive. Since $\mathbf{1}_{\mathbf{K}}$ is both join- and meet-irreducible,

the set $A = \{m \in K \mid m > 1_{\mathbf{K}}\}$ forms a distributive sublattice of $\mathbf{K}[\mathbf{L}]$, as does $B = \{m \in K \mid m < 1_{\mathbf{K}}\}$. For $a \in A$, $b \in B$ and $\ell \in L$ we always have $b < \ell < a$. Now suppose that $P \subseteq K[L]$ forms a sublattice isomorphic to \mathbf{M}_3 or \mathbf{N}_5 . If the lattice formed by P is isomorphic to \mathbf{M}_3 , then the three incomparable join-irreducibles must all be in one of A, B or L. But this would lead to $P \subseteq A$, $P \subseteq B$ or $P \subseteq L$, a contradiction. If the lattice formed by P is isomorphic to \mathbf{n}_2 , which is below n_3) would again all have to be in one of A, B or L. However, since \mathbf{K} and \mathbf{L} have distributive lattice reducts, we cannot have $P \subseteq A$, $P \subseteq B$, or $P \subseteq L$.

We note that if \mathbf{S}_n is a finite odd Sugihara chain, then \mathbf{S}_n can be viewed as an odd DqRA such that its monoid identity $1(=a_0)$ is totally irreducible. Moreover, since \mathbf{S}_n is conic, it follows from Theorem 2.12 that if \mathbf{L} is a DqRA, then the generalised ordinal sum $\mathbf{S}_n[\mathbf{L}]$ of \mathbf{S}_n and \mathbf{L} is a DqRA.

Proposition 2.13. Let $\mathbf{K} = \mathbf{S}_n$ for n odd, and $\mathbf{L} = \mathbf{S}_m$ for $m \ge 2$, then $\mathbf{K}[\mathbf{L}] \cong \mathbf{S}_{n+m-1}$.

Proof. Let $\mathbf{K} = \mathbf{S}_n$ be a Sugihara chain with n = 2k + 1 for $k \ge 1$, and let $\mathbf{L} = \mathbf{S}_m$ with $m = 2\ell$ or $m = 2\ell + 1$ for $\ell \ge 1$. Elements of S_n will be denoted by $a_{-k}^{\mathbf{K}}, \ldots, a_k^{\mathbf{K}}$, elements of S_m by $a_{-\ell}^{\mathbf{L}}, \ldots, a_{\ell}^{\mathbf{L}}$, and elements of S_{n+m-1} by $a_{-k-\ell}, \ldots, a_{k+\ell}$. Define a map $\psi : K[L] \to S_{n+m-1}$ by setting, for each $b \in K[L]$,

$$\psi(b) = \begin{cases} a_{j-\ell} & \text{if } b = a_j^{\mathbf{K}} \text{ and } -k \leq j \leq -1 \\ a_{j+\ell} & \text{if } b = a_j^{\mathbf{K}} \text{ and } 1 \leq j \leq k \\ a_i & \text{if } b = a_i^{\mathbf{L}}. \end{cases}$$

It is straightforward to show that ψ is an isomorphism from $\mathbf{K}[\mathbf{L}]$ to \mathbf{S}_{n+m-1} .

There is evidence to suggest that a conic DqRA with its monoid identity 1 totally irreducible will always be a chain, and further will always be a Sugihara chain. We have not however been able to prove this. When searching for conic DqRAs with 1 totally irreducible, Prover9/Mace4 returns only finite Sugihara chains for models of size less than 20.

Problem 1. Let \mathbf{K} be a DqRA such that $\mathbf{1}_{\mathbf{K}}$ is totally irreducible. If \mathbf{K} is conic, is it always linearly ordered? If \mathbf{K} is finite and conic, is it linearly ordered? When \mathbf{K} is finite, is it always isomorphic to \mathbf{S}_n for some odd n?

3 Representable distributive quasi relation algebras

In this section we recall the construction of distributive quasi relation algebras from partially ordered equivalence relations. This recent work by two of the current authors [3] included a definition of *representability* of a DqRA. The basic idea is to

replicate the way in which a relation algebra can be concretely constructed as an algebra of binary relations, starting from an equivalence relation. In the setting of DqRAs, the algebras are constructed as up-sets of a partially ordered equivalence relation (with the partial order satisfying some additional symmetry conditions).

As observed in [3], the lattice structure in the construction is the same as that used by Galatos and Jipsen [7, 8], and by Jipsen and Šemrl [11]. An essential difference is the additional requirement of the existence of the maps α and β (see below).

Before describing the construction from [3], we recall some basic facts about binary relations. For $R \subseteq X^2$ its converse is $R^{\smile} = \{(x, y) \mid (y, x) \in R\}$. For two binary relations R and S, the composition, R; S, is given by R; $S = \{(x, y) \mid (\exists z \in X) \ ((x, z) \in R \text{ and } (z, y) \in S)\}$. When considering a binary relation, R, it will usually be contained in an equivalence relation E and therefore the complement of R, denoted R^c , will mean $R^c = \{(x, y) \in E \mid (x, y) \notin R\}$. If an equivalence relation is not specified, it should be taken as X^2 .

We define $\operatorname{id}_X = \{(x, x) \mid x \in X\}$. The following familiar equivalences will be used frequently throughout the paper. If E is an equivalence relation on X and $R, S, T \subseteq E$ then we have $(R^{\smile})^{\smile} = R, (R^{\smile})^c = (R^c)^{\smile}$, and $\operatorname{id}_X ; R = R$; $\operatorname{id}_X = R$. Further, (R;S); T = R; (S;T) and $(R;S)^{\smile} = S^{\smile}; R^{\smile}$.

For a function $\gamma: X \to X$ we will use γ to denote either the function or the binary relation that is its graph. Note that the lemma below applies when γ is a bijective function $\gamma: X \to X$. Also recall that R^c is the complement of R in E.

Lemma 3.1 ([3, Lemma 3.4]). Let *E* be an equivalence relation on a set *X*, and let $R, S, \gamma \subseteq E$. If γ satisfies γ^{\frown} ; $\gamma = \operatorname{id}_X$ and γ ; $\gamma^{\frown} = \operatorname{id}_X$ then the following hold: (*i*) $(\gamma; R)^c = \gamma; R^c$ (*ii*) $(R; \gamma)^c = R^c; \gamma$

We now recall from [3] how to construct a DqRA of binary relations. Consider a poset $\mathbf{X} = (X, \leq)$ and E an equivalence relation on X with $\leq \subseteq E$. The equivalence relation E can be partially ordered for all $(u, v), (x, y) \in E$ as follows: $(u, v) \preccurlyeq (x, y)$ iff $x \leq u$ and $v \leq y$. The structure $\mathbf{E} = (E, \preccurlyeq)$ is then a poset, hence the set of up-sets of \mathbf{E} , denoted Up (\mathbf{E}), ordered by inclusion, is a distributive lattice.

If $R, S \in Up(\mathbf{E})$, then $R; S \in Up(\mathbf{E})$. The set of downsets of (E, \preccurlyeq) is denoted Down(\mathbf{E}) and is also closed under composition. Importantly, $R \in Up(\mathbf{E})$ iff $R^c \in$ Down(\mathbf{E}) iff $R^{\sim} \in Down(\mathbf{E})$. We have $\leqslant \in Up(\mathbf{E})$ and it is the identity with respect to ;. Further, the operation of relational composition, ;, is residuated with $R \setminus_{Up(\mathbf{E})} S =$ $(R^{\sim}; S^c)^c$ and $R/_{Up(\mathbf{E})} S = (R^c; S^{\sim})^c$. Hence $\langle Up(\mathbf{E}), \cap, \cup, ;, \setminus_{Up(\mathbf{E})}, /_{Up(\mathbf{E})}, \leqslant \rangle$ is a distributive residuated lattice.

The remaining DqRA operations in the construction (see Theorem 3.2 below) require an order automorphism $\alpha : X \to X$ and a dual order automorphism $\beta : X \to X$. We note briefly some facts regarding set-theoretic operations on Up(**E**). Details can be found in [3, Lemma 3.5]. If $R, S \in Up(\mathbf{E})$, then α ; R, and R; α are elements of Up(**E**). If $R \in Down(\mathbf{E})$, then β ; R; $\beta \in Up(\mathbf{E})$.

For $\alpha: X \to X$ an order automorphism, we let $0 = \alpha; \leq^{c}$. With this we can define \sim and - on Up(E). By [3, Lemma 3.10] we can in fact define the linear negations

without using the residuals. We get $\sim R = R^c$; α and $-R = \alpha$; R^c for all $R \in \mathsf{Up}(\mathbf{E})$. For calculations it is much easier to use these definitions of $\sim R$ and -R than those involving the residuals.

Lastly, $\neg R$ is defined using the dual order automorphism β , as stated below.

Theorem 3.2 ([3, Theorem 3.15]). Let $\mathbf{X} = (X, \leq)$ be a poset and E an equivalence relation on X such that $\leq \subseteq E$. Let $\alpha : X \to X$ be an order automorphism of \mathbf{X} and $\beta : X \to X$ a self-inverse dual order automorphism of \mathbf{X} such that $\alpha, \beta \subseteq E$ and $\beta = \alpha$; β ; α . Set $1 = \leq$ and $0 = \alpha$; \leq^{c} . For $R \in \mathsf{Up}(\mathbf{E})$, define $\sim R = R^{c}$; $\alpha, -R = \alpha$; R^{c} , and $\neg R = \alpha$; β ; R^{c} ; β . Then the algebra $\mathbf{Dq}(\mathbf{E}) = \langle \mathsf{Up}(\mathbf{E}), \cap, \cup, ;, \sim, -, \neg, 1, 0 \rangle$ is a distributive quasi relation algebra. If α is the identity, then $\mathbf{Dq}(\mathbf{E})$ is a cyclic distributive quasi relation algebra.

An algebra of the form described by Theorem 3.2 is called an *equivalence distributive quasi relation algebra* and the class of such algebras is denoted EDqRA. If $E = X^2$, then we refer to the algebra $\mathbf{Dq}(\mathbf{E})$ as a *full distributive quasi relation algebra*, with the class of such algebras denoted by FDqRA. Analogous to the case for relation algebras (cf. [14, Chapter 3]), it was shown [3, Theorem 4.4] that $\mathbb{IP}(\mathsf{FDqRA}) = \mathbb{I}(\mathsf{EDqRA})$. This gives rise to the definition below:

Definition 3.3 ([3, Definition 4.5]). A $DqRA \mathbf{A} = \langle A, \wedge, \vee, \cdot, \sim, -, \neg, 1, 0 \rangle$ is representable if $\mathbf{A} \in \mathbb{ISP}(\mathsf{FDqRA})$ or, equivalently, $\mathbf{A} \in \mathbb{IS}(\mathsf{EDqRA})$.

We say that a DqRA **A** is *finitely* representable if the poset $\langle X, \leq \rangle$ used in the representation of **A** is finite.

Below we give some examples of representable DqRAs. These examples will become useful later on.

Example 3.4. Consider the two-element Sugihara chain \mathbf{S}_2 (viewed as a DqRA). Let $\mathbf{X}_{\mathbf{S}_2}$ be the one element poset with $X_{\mathbf{S}_2} = \{u\}$ and $\leq_{\mathbf{X}_{\mathbf{S}_2}} = \alpha = \beta = \mathrm{id}_{\mathbf{X}_{\mathbf{S}_2}}$. Then the underlying lattice of $\mathsf{Up}(X^2_{\mathbf{S}_2},\preccurlyeq)$ is the two-element chain $(\{\varnothing, X^2_{\mathbf{S}_2}\}, \subseteq)$ that forms a representation of \mathbf{S}_2 . Figure 5 contains a depiction of $\mathbf{X}_{\mathbf{S}_2}, \alpha, \beta$ and the blocks of $X^2_{\mathbf{S}_2}$ as well as the representation of \mathbf{S}_2 .




Fig. 5 The poset X_{S_2} used to represent S_2 (left) and the representation of S_2 (right).

Example 3.5 ([3, Example 5.1]). Recall that \mathbf{S}_3 denotes the three-element Sugihara chain (thought of as a DqRA) with $S_3 = \{a_{-1}, a_0, a_1\}$ such that a_0 is the identity of the monoid operation. Let $\mathbf{X}_{\mathbf{S}_3}$ be the two-element antichain, i.e., $X_{\mathbf{S}_3} = \{x, y\}$ with $\leq_{\mathbf{X}_{\mathbf{S}_3}} = \mathrm{id}_{\mathbf{X}_{\mathbf{S}_3}}$, and consider the order automorphism $\alpha = \{(x, y), (y, x)\}$ and the dual order automorphism $\beta = \mathrm{id}_{\mathbf{X}_{\mathbf{S}_3}}$ of $\mathbf{X}_{\mathbf{S}_3}$. See Figure 6 for a depiction of the poset $\mathbf{X}_{\mathbf{S}_3}$, the order automorphism α , the dual order automorphism β , and the blocks of $X_{\mathbf{S}_3}^2$. Then the underlying lattice of $\mathsf{Up}(X_{\mathbf{S}_3}^2, \preccurlyeq)$ is the 16-element Boolean lattice. It can easily be shown that α ; $(\leq_{\mathbf{X}_{\mathbf{S}_3}}^c)^{\sim} = (\leq_{\mathbf{X}_{\mathbf{S}_3}}^c)^{\sim}$; $\alpha = \sim \leq_{\mathbf{X}_{\mathbf{S}_3}} = \leq_{\mathbf{X}_{\mathbf{S}_3}}$ and that the set of binary relations $\{\emptyset, \leq_{\mathbf{X}_{\mathbf{S}_3}}, X_{\mathbf{S}_3}^2\}$ forms a subuniverse of $\langle \mathsf{Up}(X_{\mathbf{S}_3}^2, \preccurlyeq), \cap, \cup, ;, \sim, -, \neg, \leq_{\mathbf{X}_{\mathbf{S}_3}}, (\alpha; \leq_{\mathbf{X}_{\mathbf{S}_3}}^c) \rangle$, such that the subalgebra $\langle \{\emptyset, \leq_{\mathbf{X}_{\mathbf{S}_3}}, X_{\mathbf{S}_3}^2\}, \cap, \cup, ;, \sim, -, \neg, \leq_{\mathbf{X}_{\mathbf{S}_3}}, \langle \alpha; \leq_{\mathbf{X}_{\mathbf{S}_3}}^c \rangle \rangle$, is a representation of \mathbf{S}_3 (also depicted in Figure 6).




Fig. 6 The poset X_{S_3} used to represent S_3 (left) and the representation of S_3 (right).

Example 3.6. Consider the first diamond \mathbf{L}_1 given in Figure 1. It can be represented over the two-element antichain $\mathbf{X}_{\mathbf{L}_1}$, i.e., $X_{\mathbf{L}_1} = \{u, v\}$ and $\leq_{\mathbf{L}_1} = \operatorname{id}_{X_{\mathbf{L}_1}}$, with $E = \alpha = \beta = \operatorname{id}_{X_{\mathbf{L}_1}}$. Hence, in this example the algebra $\mathbf{Dq}(\mathbf{E})$ is not a full distributive quasi relation algebra. The poset $\mathbf{X}_{\mathbf{L}_1}$, the order automorphism α , the dual order automorphism β and the blocks of E are depicted in Figure 7. Clearly the set of binary relations $\{\varnothing, \{(u, u)\}, \{(v, v)\}, \leq_{\mathbf{L}_1}\}$ forms a subuniverse of the algebra $\langle \mathsf{Up}(E, \preccurlyeq), \cap, \cup, ;, \sim, -, \neg, \leq_{\mathbf{L}_1}, (\alpha; \leq_{\mathbf{L}_1}^{c_{\mathbf{C}}}) \rangle$. We have that the monoid identity $\mathbf{1}_{\mathbf{L}_1}$ is mapped to $\leq_{\mathbf{L}_1} = E$, 0 to \varnothing , a to $\{(u, u)\}$ and b to $\{(v, v)\}$.

4 Representable DqRAs via generalised ordinal sums

In this section we prove that if \mathbf{L} is a representable distributive quasi relation algebra, then the generalised ordinal sum of \mathbf{S}_3 and \mathbf{L} is representable.

Theorem 4.1. Let $\mathbf{L} = \langle L, \wedge_{\mathbf{L}}, \vee_{\mathbf{L}}, \cdots, \vee_{\mathbf{L}}, \neg_{\mathbf{L}}, 1_{\mathbf{L}}, 0_{\mathbf{L}} \rangle$ be a (finitely) representable DqRA. Then $\mathbf{S}_{3}[\mathbf{L}]$ is a (finitely) representable DqRA.




Fig. 7 The poset X_{L_1} used to represent the diamond L_1 (left) and the representation of L_1 (right).

In order to prove Theorem 4.1, we use the representations of \mathbf{S}_3 and \mathbf{L} to construct a suitable candidate with which to represent $\mathbf{S}_3[\mathbf{L}]$. Through a series of lemmas we then show that the construction does indeed produce a representation of the DqRA $\mathbf{S}_3[\mathbf{L}]$ (as summarized in the proof at the end of the section).

Let **K** be the odd three element Sugihara chain described in Section 1, i.e., $\mathbf{K} = \langle \{a_{-1}, a_0, a_1\}, \wedge, \vee, \cdot, \sim, \sim, a_0, a_0 \rangle$ (also see Figure 2). Then a_0 , the monoid identity in **K**, is totally irreducible and **K** is odd, conic and representable. Recall from Example 3.5 that the representation uses the two-element antichain $\mathbf{X}_{\mathbf{K}} = (X_{\mathbf{K}}, \leq_{X_{\mathbf{K}}})$ with $X_{\mathbf{K}} = \{x, y\}$ and the equivalence relation $E_{\mathbf{K}} = X_{\mathbf{K}} \times X_{\mathbf{K}}$. The order automorphism $\alpha_{\mathbf{K}} : X_{\mathbf{K}} \to X_{\mathbf{K}}$ is defined by $\alpha_{\mathbf{K}}(x) = y$ and $\alpha_{\mathbf{K}}(y) = x$, while the dual order automorphism $\beta_{\mathbf{K}} : X_{\mathbf{K}} \to X_{\mathbf{K}}$ is the identity map and $\alpha_{\mathbf{K}}$; $\beta_{\mathbf{K}}$; $\alpha_{\mathbf{K}} = \beta_{\mathbf{K}}$. Elements of $X_{\mathbf{K}}$ will be denoted by w_1, w_2, \ldots or just w. Finally, **K** can be embedded into the algebra of up-sets of $(E_{\mathbf{K}}, \preccurlyeq_{E_{\mathbf{K}}})$ via the embedding $\varphi_{\mathbf{K}} : \mathbf{K} \hookrightarrow \mathbf{Dq}(E_{\mathbf{K}}, \preccurlyeq_{E_{\mathbf{K}}})$ such that $\varphi_{\mathbf{K}}(a_{-1}) = \emptyset$, $\varphi_{\mathbf{K}}(a_0) = \leqslant_{X_{\mathbf{K}}}$ and $\varphi_{\mathbf{K}}(a_1) = E_{\mathbf{K}} = X_{\mathbf{K}} \times X_{\mathbf{K}}$.

Now let $\mathbf{L} = \langle L, \wedge_{\mathbf{L}}, \vee_{\mathbf{L}}, \sim_{\mathbf{L}}, -_{\mathbf{L}}, \neg_{\mathbf{L}}, \mathbf{1}_{\mathbf{L}}, \mathbf{0}_{\mathbf{L}} \rangle$ be any representable DqRA. Then there exists a poset $\mathbf{X}_{\mathbf{L}} = (X_{\mathbf{L}}, \leq_{X_{\mathbf{L}}})$, an equivalence relation $E_{\mathbf{L}} \subseteq X_{\mathbf{L}} \times X_{\mathbf{L}}$ with $\leq_{X_{\mathbf{L}}} \subseteq E_{\mathbf{L}}$, an order automorphism $\alpha_{\mathbf{L}} : X_{\mathbf{L}} \to X_{\mathbf{L}}$ and a self-inverse dual order automorphism $\beta_{\mathbf{L}} : X_{\mathbf{L}} \to X_{\mathbf{L}}$ such that $\alpha_{\mathbf{L}}, \beta_{\mathbf{L}} \subseteq E_{\mathbf{L}}$ and $\alpha_{\mathbf{L}}; \beta_{\mathbf{L}}; \alpha_{\mathbf{L}} = \beta_{\mathbf{L}}$. Elements of the set $X_{\mathbf{L}}$ will be denoted by z_1, z_2, \ldots or just z. Moreover, there exists an embedding $\varphi_{\mathbf{L}} : \mathbf{L} \hookrightarrow \mathbf{Dq} (E_{\mathbf{L}}, \preccurlyeq_{E_{\mathbf{L}}})$ (see Section 3 for the details).

We first describe the poset used to represent $\mathbf{K}[\mathbf{L}]$. For each $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, let $[z]\mathbf{X}_{\mathbf{K}}$ and $[z]\mathbf{X}_{\mathbf{K}}$ each be a copy of $\mathbf{X}_{\mathbf{K}}$ with $[z]X_{\mathbf{K}} = \{x^{[z]}, y^{[z]}\}$ and $[z]X_{\mathbf{K}} = \{x_{[z]}, y_{[z]}\}$. Thus, both $[z]\mathbf{X}_{\mathbf{K}}$ and $[z]\mathbf{X}_{\mathbf{K}}$ are two element antichains with their orderings denoted by $[z] \leq_{X_{\mathbf{K}}}$ and $[z] \leq_{X_{\mathbf{K}}}$, respectively. Now let

$$X_{\mathbf{K}[\mathbf{L}]} := X_{\mathbf{L}} \cup \bigcup \left\{ [z] X_{\mathbf{K}} \mid [z] \in X_{\mathbf{L}} / E_{\mathbf{L}} \right\} \cup \bigcup \left\{ [z] X_{\mathbf{K}} \mid [z] \in X_{\mathbf{L}} / E_{\mathbf{L}} \right\}$$
(6)

and we use u's and v's to denote elements of $X_{\mathbf{K}[\mathbf{L}]}$.

We define $\leq_{X_{\mathbf{K}[\mathbf{L}]}} \subseteq X_{\mathbf{K}[\mathbf{L}]} \times X_{\mathbf{K}[\mathbf{L}]}$ by

$$\leq_{X_{\mathbf{K}[\mathbf{L}]}} := \bigcup \left\{ [z] \leq_{X_{\mathbf{K}}} \middle| [z] \in X_{\mathbf{L}} / E_{\mathbf{L}} \right\} \cup \bigcup \left\{ [z] \leq_{X_{\mathbf{K}}} \middle| [z] \in X_{\mathbf{L}} / E_{\mathbf{L}} \right\} \cup$$

$$\bigcup \left\{ [z] \times {}^{[z]}X_{\mathbf{K}} \middle| [z] \in X_{\mathbf{L}}/E_{\mathbf{L}} \right\} \cup \bigcup \left\{ {}_{[z]}X_{\mathbf{K}} \times [z] \middle| [z] \in X_{\mathbf{L}}/E_{\mathbf{L}} \right\} \cup \\
\bigcup \left\{ {}_{[z]}X_{\mathbf{K}} \times {}^{[z]}X_{\mathbf{K}} \middle| [z] \in X_{\mathbf{L}}/E_{\mathbf{L}} \right\} \cup \leqslant_{X_{\mathbf{L}}}.$$
(7)

Then $\mathbf{X}_{\mathbf{K}[\mathbf{L}]} = (X_{\mathbf{K}[\mathbf{L}]}, \leq_{X_{\mathbf{K}[\mathbf{L}]}})$ is a poset. Intuitively, $\mathbf{X}_{\mathbf{K}[\mathbf{L}]}$ is obtained from $\mathbf{X}_{\mathbf{L}}$ by placing a copy of $\mathbf{X}_{\mathbf{K}}$ above and below ($^{[z]}\mathbf{X}_{\mathbf{K}}$ and $_{[z]}\mathbf{X}_{\mathbf{K}}$, respectively) each block [z] of $\mathbf{X}_{\mathbf{L}}$ modulo $E_{\mathbf{L}}$.

Example 4.2. Let $\mathbf{K} = \mathbf{S}_3$ from Figure 2 and $\mathbf{L} = \mathbf{L}_1$ the first diamond from Figure 1. Then $\mathbf{K}[\mathbf{L}]$ is the DqRA depicted in Figure 3. Recall from Examples 3.5 and 3.6, respectively, that both \mathbf{K} and \mathbf{L} are representable, with the posets used to find their representations depicted in Figures 6 and 7, respectively. Using these posets we can construct the poset $\mathbf{X}_{\mathbf{K}[\mathbf{L}]}$, as described in (6) and (7). The structure $(X_{\mathbf{K}[\mathbf{L}]}, \leqslant_{X_{\mathbf{K}[\mathbf{L}]}})$ is drawn in Figure 8.




Fig. 8 The poset $X_{K[L]}$.

Next, we set

$$E_{\mathbf{K}[\mathbf{L}]} := \bigcup \left\{ \left([z] \cup {}^{[z]}X_{\mathbf{K}} \cup {}_{[z]}X_{\mathbf{K}} \right)^2 \ \Big| \ [z] \in X_{\mathbf{L}}/E_{\mathbf{L}} \right\}.$$
(8)

A block in $X_{\mathbf{K}[\mathbf{L}]}/E_{\mathbf{K}[\mathbf{L}]}$ consists of a block [z] in $X_{\mathbf{L}}/E_{\mathbf{L}}$ along with $[z]\mathbf{X}_{\mathbf{K}}$ and $[z]\mathbf{X}_{\mathbf{K}}$, the copies of $\mathbf{X}_{\mathbf{K}}$ associated with it. Using the fact that $E_{\mathbf{L}}$ is an equivalence relation on $X_{\mathbf{L}}$ we can show that $E_{\mathbf{K}[\mathbf{L}]}$ is an equivalence relation on $X_{\mathbf{K}[\mathbf{L}]}$. Moreover, since $\leq_{X_{\mathbf{L}}} \subseteq E_{\mathbf{L}}$, it follows that $\leq_{\mathbf{K}[\mathbf{L}]} \subseteq E_{\mathbf{K}[\mathbf{L}]}$. Recall that $E_{\mathbf{K}[\mathbf{L}]}$ can be partially ordered as follows, for $(u_1, u_2), (v_1, v_2) \in E_{\mathbf{K}[\mathbf{L}]}$,

$$(u_1, u_2) \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}} (v_1, v_2) \qquad \Longleftrightarrow \qquad v_1 \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} u_1 \quad \text{and} \quad u_2 \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} v_2 \qquad (9)$$

Define $\alpha_{\mathbf{K}[\mathbf{L}]} : X_{\mathbf{K}[\mathbf{L}]} \to X_{\mathbf{K}[\mathbf{L}]}$ by

$$\alpha_{\mathbf{K}[\mathbf{L}]}(u) = \begin{cases} \alpha_{\mathbf{L}}(u) & \text{if } u \in X_{\mathbf{L}} \\ (\alpha_{\mathbf{K}}(w))^{[z]} & \text{if } u = w^{[z]} \text{ for some } w \in X_{\mathbf{K}} \text{ and some } [z] \text{ in } X_{\mathbf{L}}/E_{\mathbf{L}} \\ (\alpha_{\mathbf{K}}(w))_{[z]} & \text{if } u = w_{[z]} \text{ for some } w \in X_{\mathbf{K}} \text{ and some } [z] \text{ in } X_{\mathbf{L}}/E_{\mathbf{L}}, (10) \end{cases}$$

and $\beta_{\mathbf{K}[\mathbf{L}]} : X_{\mathbf{K}[\mathbf{L}]} \to X_{\mathbf{K}[\mathbf{L}]}$ by

$$\beta_{\mathbf{K}[\mathbf{L}]}(u) = \begin{cases} \beta_{\mathbf{L}}(u) & \text{if } u \in X_{\mathbf{L}} \\ (\beta_{\mathbf{K}}(w))_{[z]} & \text{if } u = w^{[z]} \text{ for some } w \in X_{\mathbf{K}} \text{ and some } [z] \text{ in } X_{\mathbf{L}}/E_{\mathbf{L}} \\ (\beta_{\mathbf{K}}(w))^{[z]} & \text{if } u = w_{[z]} \text{ for some } w \in X_{\mathbf{K}} \text{ and some } [z] \text{ in } X_{\mathbf{L}}/E_{\mathbf{L}}, (11) \end{cases}$$

for $u \in X_{\mathbf{K}[\mathbf{L}]}$.

Example 4.3. Let **K** and **L** be as described in Example 4.2 with $\mathbf{X}_{\mathbf{K}[\mathbf{L}]}$ the poset depicted in Figure 8. Then Figure 9 depicts the equivalence classes of $E_{\mathbf{K}[\mathbf{L}]}$ as well as the maps $\alpha_{\mathbf{K}[\mathbf{L}]}$ and $\beta_{\mathbf{K}[\mathbf{L}]}$ described in (8),(10) and (11), respectively.




Fig. 9 $E_{\mathbf{K}[\mathbf{L}]}$, $\alpha_{\mathbf{K}[\mathbf{L}]}$ and $\beta_{\mathbf{K}[\mathbf{L}]}$ on $\mathbf{X}_{\mathbf{K}[\mathbf{L}]}$.

Lemma 4.4. (i) The map $\alpha_{\mathbf{K}[\mathbf{L}]} : X_{\mathbf{K}[\mathbf{L}]} \to X_{\mathbf{K}[\mathbf{L}]}$ is an order automorphism of $\mathbf{X}_{\mathbf{K}[\mathbf{L}]}$ such that $\alpha_{\mathbf{K}[\mathbf{L}]} \subseteq E_{\mathbf{K}[\mathbf{L}]}$.

- (ii) The map $\beta_{\mathbf{K}[\mathbf{L}]} : X_{\mathbf{K}[\mathbf{L}]} \to X_{\mathbf{K}[\mathbf{L}]}$ is a self-inverse dual order automorphism of $\mathbf{X}_{\mathbf{K}[\mathbf{L}]}$ such that $\beta_{\mathbf{K}[\mathbf{L}]} \subseteq E_{\mathbf{K}[\mathbf{L}]}$.
- (*iii*) $\alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}; \alpha_{\mathbf{K}[\mathbf{L}]} = \beta_{\mathbf{K}[\mathbf{L}]}$

Proof. We prove items (ii) and (iii). The proof of item (i) is similar to that of (ii).

To prove (ii) we first show that $\beta_{\mathbf{K}[\mathbf{L}]} \subseteq E_{\mathbf{K}[\mathbf{L}]}$. Let $u, v \in X_{\mathbf{K}[\mathbf{L}]}$ such that $\beta_{\mathbf{K}[\mathbf{L}]}(u) = v$. There are three cases to consider:

<u>Case 1:</u> If $u \in X_{\mathbf{L}}$, then $v \in X_{\mathbf{L}}$ and $v = \beta_{\mathbf{K}[\mathbf{L}]}(u) = \beta_{\mathbf{L}}(u)$, and so $(u, v) \in \beta_{\mathbf{L}} \subseteq E_{\mathbf{L}}$. Hence, $u \in [u]$ and $v \in [u]$, which means $(u, v) \in E_{\mathbf{K}[\mathbf{L}]}$.

<u>Case 2:</u> If $u \in {}^{[z]}X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then $v \in {}_{[z]}X_{\mathbf{K}}$ and there exist elements $w_1, w_2 \in X_{\mathbf{K}}$ such that $u = (w_1)^{[z]}$ and $v = (w_2)_{[z]}$. Since ${}^{[z]}X_{\mathbf{K}} \times {}_{[z]}X_{\mathbf{K}} \subseteq E_{\mathbf{K}[\mathbf{L}]}$ by (8), it follows that $(u, v) \in E_{\mathbf{K}[\mathbf{L}]}$.

<u>Case 3:</u> If $u \in [z]X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then $v \in [z]X_{\mathbf{K}}$ and there exist elements $w_1, w_2 \in X_{\mathbf{K}}$ such that $u = (w_1)_{[z]}$ and $v = (w_2)^{[z]}$. Since $[z]X_{\mathbf{K}} \times [z]X_{\mathbf{K}} \subseteq E_{\mathbf{K}[\mathbf{L}]}$ by (8), it follows that $(u, v) \in E_{\mathbf{K}[\mathbf{L}]}$.

To see that $\beta_{\mathbf{K}[\mathbf{L}]}$ is a dual order automorphism, let $u, v \in X_{\mathbf{K}[\mathbf{L}]}$. There are six cases to consider.

<u>Case 1:</u> If $u, v \in X_{\mathbf{L}}$, then

$$u \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} v \iff u \leqslant_{X_{\mathbf{L}}} v \iff \beta_{\mathbf{L}}(v) \leqslant_{X_{\mathbf{L}}} \beta_{\mathbf{L}}(u) \iff \beta_{\mathbf{K}[\mathbf{L}]}(v) \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} \beta_{\mathbf{K}[\mathbf{L}]}(u).$$

<u>Case 2:</u> If $u, v \in [z]X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then there exist $w_1, w_2 \in X_{\mathbf{K}}$ such that $u = (w_1)^{[z]}$ and $v = (w_2)^{[z]}$. Hence,

$$u \leq_{X_{\mathbf{K}[\mathbf{L}]}} v \iff w_1 \leq_{X_{\mathbf{K}}} w_2 \iff \beta_{\mathbf{K}}(w_2) \leq_{X_{\mathbf{K}}} \beta_{\mathbf{K}}(w_1)$$
$$\iff (\beta_{\mathbf{K}}(w_2))_{[z]} \leq_{X_{\mathbf{K}[\mathbf{L}]}} (\beta_{\mathbf{K}}(w_1))_{[z]}$$
$$\iff \beta_{\mathbf{K}[\mathbf{L}]}(v) \leq_{X_{\mathbf{K}[\mathbf{L}]}} \beta_{\mathbf{K}[\mathbf{L}]}(u).$$

<u>Case 3:</u> If $u, v \in [z]X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, the proof is similar to the proof of the previous case described above.

<u>Case 4:</u> If $u \in X_{\mathbf{L}}$ and $v \in {}^{[u]}X_{\mathbf{K}}$, then $(u, \beta_{\mathbf{L}}(u)) \in \beta_{\mathbf{L}} \subseteq E_{\mathbf{L}}$, i.e., $\beta_{\mathbf{L}}(u) \in [u]$, and there exist $w \in X_{\mathbf{K}}$ such that $v = w^{[u]}$. It follows that

$$u \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} v \iff u \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} w^{[u]} \iff (\beta_{\mathbf{K}}(w))_{[u]} \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} \beta_{\mathbf{L}}(u)$$
$$\iff \beta_{\mathbf{K}[\mathbf{L}]}(v) \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} \beta_{\mathbf{K}[\mathbf{L}]}(u).$$

<u>Case 5:</u> If $u \in [u]X_{\mathbf{K}}$ and $v \in X_{\mathbf{L}}$, the proof is similar to the proof of the previous case described above.

<u>Case 6:</u> If $u \in [z]X_{\mathbf{K}}$ and $v \in [z]X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then there exist $w_1, w_2 \in X_{\mathbf{K}}$ such that $u = (w_1)_{[z]}$ and $v = (w_2)^{[z]}$. Thus,

$$u \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} v \iff (w_1)_{[z]} \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} (w_2)^{[z]} \iff (\beta_{\mathbf{K}}(w_2))_{[z]} \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} (\beta_{\mathbf{K}}(w_1))^{[z]}$$
$$\iff \beta_{\mathbf{K}[\mathbf{L}]}(v) \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} \beta_{\mathbf{K}[\mathbf{L}]}(u).$$

Note that for any other combination of elements $u, v \in X_{\mathbf{K}[\mathbf{L}]}$ it will follow immediately from (7) and (11) that $u \not\leq_{X_{\mathbf{K}[\mathbf{L}]}} v$ and $\beta_{\mathbf{K}[\mathbf{L}]}(v) \not\leq_{X_{\mathbf{K}[\mathbf{L}]}} \beta_{\mathbf{K}[\mathbf{L}]}(u)$. Next we show that $\beta_{\mathbf{K}[\mathbf{L}]}$ is self-inverse. Let $u \in X_{\mathbf{K}[\mathbf{L}]}$. There are three cases.

<u>Case 1:</u> If $u \in X_{\mathbf{L}}$, then $\beta_{\mathbf{K}[\mathbf{L}]}(u) = \beta_{\mathbf{L}}(u) \in X_{\mathbf{L}}$ and so $\beta_{\mathbf{K}[\mathbf{L}]}(\beta_{\mathbf{K}[\mathbf{L}]}(u)) = \beta_{\mathbf{L}}(\beta_{\mathbf{L}}(u))$. Hence, since $\beta_{\mathbf{L}}$ is self-inverse, we get $\beta_{\mathbf{K}[\mathbf{L}]}(\beta_{\mathbf{K}[\mathbf{L}]}(u)) = u$.

<u>Case 2</u>: If $u \in [z]X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then there exists $w \in X_{\mathbf{K}}$ such that $u = (w)_{[z]}$. Hence,

$$\beta_{\mathbf{K}[\mathbf{L}]} \left(\beta_{\mathbf{K}[\mathbf{L}]}(u) \right) = \beta_{\mathbf{K}[\mathbf{L}]} \left(\beta_{\mathbf{K}[\mathbf{L}]}((w)_{[z]}) = \beta_{\mathbf{K}[\mathbf{L}]} \left(\left(\beta_{\mathbf{K}}(w) \right)^{[z]} \right) \\ = \left(\beta_{\mathbf{K}} \left(\beta_{\mathbf{K}}(w) \right) \right)_{[z]} = (w)_{[z]}$$

The last equality follows from the fact that $\beta_{\mathbf{K}}$ is self-inverse.

<u>Case 3:</u> If $u \in [z]X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, the proof is similar to the previous case.

Finally we show that $\beta_{\mathbf{K}[\mathbf{L}]}$ is surjective. Let $u \in X_{\mathbf{K}[\mathbf{L}]}$. We have to find some $v \in X_{\mathbf{K}[\mathbf{L}]}$ such that $\beta_{\mathbf{K}[\mathbf{L}]}(v) = u$. There are three cases to consider:

<u>Case 1:</u> If $u \in [z]X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then there exists $w_1 \in X_{\mathbf{K}}$ such that $u = (w_1)_{[z]}$. Since $\beta_{\mathbf{K}}$ is surjective, there is some $w_2 \in X_{\mathbf{K}}$ such that $\beta_{\mathbf{K}}(w_2) = w_1$. and hence $u = (w_1)_{[z]} = (\beta_{\mathbf{K}}(w_2))_{[z]} = \beta_{\mathbf{K}[\mathbf{L}]} ((w_2)^{[z]})$.

<u>Case 2:</u> If $u \in {}^{[z]}X_{\mathbf{K}}$, the proof is similar to the previous case.

<u>Case 3:</u> If $u \in X_{\mathbf{L}}$, then the surjectivity of $\beta_{\mathbf{L}}$ implies that there is some $v \in X_{\mathbf{L}}$ such that $u = \beta_{\mathbf{L}}(v) = \beta_{\mathbf{K}[\mathbf{L}]}(v)$.

To prove item (iii) let $u \in X_{\mathbf{K}[\mathbf{L}]}$. We consider three cases.

<u>Case 1:</u> If $u \in X_{\mathbf{L}}$, then, since $\alpha_{\mathbf{L}}$; $\beta_{\mathbf{L}}$; $\alpha_{\mathbf{L}} = \beta_{\mathbf{L}}$, we have that

$$\alpha_{\mathbf{K}[\mathbf{L}]}\left(\beta_{\mathbf{K}[\mathbf{L}]}\left(\alpha_{\mathbf{K}[\mathbf{L}]}\left(u\right)\right)\right) = \alpha_{\mathbf{L}}\left(\beta_{\mathbf{L}}\left(\alpha_{\mathbf{L}}\left(u\right)\right)\right) = \beta_{\mathbf{L}}(u) = \beta_{\mathbf{K}[\mathbf{L}]}(u).$$

<u>Case 2:</u> If $u \in {}^{[z]}X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then $u = w^{[z]}$ for some $w \in X_{\mathbf{K}}$. Since $\alpha_{\mathbf{K}}; \beta_{\mathbf{K}}; \alpha_{\mathbf{K}} = \beta_{\mathbf{K}}$, we have that

$$\alpha_{\mathbf{K}[\mathbf{L}]} \left(\beta_{\mathbf{K}[\mathbf{L}]} \left(\alpha_{\mathbf{K}[\mathbf{L}]} \left(u \right) \right) \right) = \alpha_{\mathbf{K}[\mathbf{L}]} \left(\beta_{\mathbf{K}[\mathbf{L}]} \left(\left(\alpha_{\mathbf{K}}(w) \right)^{[z]} \right) \right)$$
$$= \alpha_{\mathbf{K}[\mathbf{L}]} \left(\left(\beta_{\mathbf{K}} \left(\alpha_{\mathbf{K}}(w) \right) \right)_{[z]} \right)$$
$$= \left(\alpha_{\mathbf{K}} \left(\beta_{\mathbf{K}} \left(\alpha_{\mathbf{K}}(w) \right) \right)_{[z]}$$
$$= \left(\beta_{\mathbf{K}}(w) \right)_{[z]} = \beta_{\mathbf{K}[\mathbf{L}]}(u)$$

<u>Case 3:</u> If $u \in [z]X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$ the proof is similar to the previous case.

Lemma 4.4 shows that $\alpha_{\mathbf{K}[\mathbf{L}]}$ and $\beta_{\mathbf{K}[\mathbf{L}]}$ satisfy the conditions of Theorem 3.2, so we obtain the following result.

Theorem 4.5. The algebra $\left\langle \mathsf{Up}\left(E_{\mathbf{K}[\mathbf{L}]}, \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}}\right), \cap, \cup, ;, \sim, -, \neg, \leqslant_{X_{\mathbf{K}[\mathbf{L}]}}, \left(\alpha_{\mathbf{K}[\mathbf{L}]}; \leqslant_{X_{\mathbf{K}[\mathbf{L}]}}^{c_{\smile}}\right) \right\rangle$ is a DqRA.

Next we define a map $\psi : K \setminus \{a_0\} \cup L \to \mathcal{P}\left((X_{\mathbf{K}[\mathbf{L}]})^2\right)$ that will show $\mathbf{K}[\mathbf{L}]$ to be representable. For the sake of readability we let $R \subseteq X_{\mathbf{K}[\mathbf{L}]} \times X_{\mathbf{K}[\mathbf{L}]}$ denote the following binary relation:

$$R := \bigcup \left\{ \begin{bmatrix} z \end{bmatrix} \leqslant_{X_{\mathbf{K}}} \middle| [z] \in X_{\mathbf{L}}/E_{\mathbf{L}} \right\} \cup \bigcup \left\{ \begin{bmatrix} z \end{bmatrix} \leqslant_{X_{\mathbf{K}}} \middle| [z] \in X_{\mathbf{L}}/E_{\mathbf{L}} \right\} \cup \bigcup \left\{ \begin{bmatrix} z \end{bmatrix} \times \begin{bmatrix} z \end{bmatrix} X_{\mathbf{K}} \middle| [z] \in X_{\mathbf{L}}/E_{\mathbf{L}} \right\} \cup \bigcup \left\{ \begin{bmatrix} z \end{bmatrix} X_{\mathbf{K}} \times \begin{bmatrix} z \end{bmatrix} \middle| [z] \in X_{\mathbf{L}}/E_{\mathbf{L}} \right\} \cup \bigcup \left\{ \begin{bmatrix} z \end{bmatrix} X_{\mathbf{K}} \times \begin{bmatrix} z \end{bmatrix} X_{\mathbf{K}} \times \begin{bmatrix} z \end{bmatrix} X_{\mathbf{K}} \right\} \cup \bigcup \left\{ \begin{bmatrix} z \end{bmatrix} X_{\mathbf{K}} \times \begin{bmatrix} z \end{bmatrix} X_{\mathbf{K}} \right\} = \bigcup \left\{ \begin{bmatrix} z \end{bmatrix} X_{\mathbf{K}} \times \begin{bmatrix} z \end{bmatrix} X_{\mathbf{K}} \Big| [z] \in X_{\mathbf{L}}/E_{\mathbf{L}} \right\}.$$
(12)

In fact, $R = \leq_{X_{\mathbf{K}[\mathbf{L}]}} \leq_{X_{\mathbf{L}}}$. Now, for $m \in K \setminus \{a_0\} \cup L$,

$$\psi(m) = \begin{cases} E_{\mathbf{K}[\mathbf{L}]} & \text{if } m = a_1 \\ R \cup \varphi_{\mathbf{L}}(m), & \text{if } m \in L, \\ \varnothing, & \text{if } m = a_{-1} \end{cases}$$
(13)

Before setting out to prove that this construction provides a representation for the generalised ordinal sum of \mathbf{K} and \mathbf{L} , we consider an example that demonstrates how the representation is constructed.

Example 4.6. Let $\mathbf{K} = \mathbf{S}_3$ and $\mathbf{L} = \mathbf{L}_1$ (the first diamond in Figure 1). Then, as per Example 2.9, the six-element algebra from Figure 3 is the generalised ordinal sum of \mathbf{K} and \mathbf{L} . The posets used to represent \mathbf{K} and \mathbf{L} , as well as their representations, are given in Figures 6 and 7, respectively (their representations are also depicted at the top of Figure 10). The poset $\mathbf{X}_{\mathbf{K}[\mathbf{L}]}$ is constructed from these posets and depicted in Figures 8 and 9. Recall that R is defined as in (12). Now, using the labels for elements of $\mathbf{K}[\mathbf{L}]$ from Figure 3, we have that

$$\begin{split} \psi(\bot) &= \varnothing \\ \psi(0) &= R = \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} \backslash \leqslant_{X_{\mathbf{L}}} \\ \psi(a) &= R \cup \{(u, u)\} = \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} \backslash \{(v, v)\} \\ \psi(b) &= R \cup \{(v, v)\} = \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} \backslash \{(u, u)\}, \text{ and} \\ \psi(1) &= R \cup \leqslant_{X_{\mathbf{L}}} = \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} \\ \psi(\top) &= E_{\mathbf{K}[\mathbf{L}]} = \bigcup_{z \in \{u, v\}} \left([z] \cup \{x^{[z]}, y^{[z]}\} \cup \{x_{[z]}, y_{[z]}\} \right)^2 \end{split}$$

The set of binary relations $\{\psi(\perp), \psi(0), \psi(a), \psi(b), \psi(1), \psi(\top)\}$ forms a subuniverse of $\langle \mathsf{Up}(E_{\mathbf{K}[\mathbf{L}]}, \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}}), \cap, \cup, ;, \sim, -, \neg, \leqslant_{X_{\mathbf{K}[\mathbf{L}]}}, R \rangle$ and hence $\mathbf{K}[\mathbf{L}]$ is representable, as depicted in Figure 10.

Having given an example of how the new representation will work, we now prove that ψ does indeed map elements from $K \setminus \{a_0\} \cup L$ to elements in $\mathsf{Up}\left(E_{\mathbf{K}[\mathbf{L}]}, \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}}\right)$.

Lemma 4.7. For all $m \in K \setminus \{a_0\} \cup L$, the set $\psi(m)$ as defined in (13) is an up-set of $(E_{\mathbf{K}[\mathbf{L}]}, \leq_{E_{\mathbf{K}[\mathbf{L}]}})$.

Proof. It is immediate that \varnothing and $E_{\mathbf{K}[\mathbf{L}]}$ are up-sets of $(E_{\mathbf{K}[\mathbf{L}]}, \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}})$. Thus, we only need to prove that $\psi(m)$ is an up-set of $(E_{\mathbf{K}[\mathbf{L}]}, \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}})$ for $m \in L$. Let $(u_1, v_1) \in \psi(m)$ and assume $(u_1, v_1) \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}} (u_2, v_2)$. That is, by (9), we have that $u_2 \preccurlyeq_{X_{\mathbf{K}[\mathbf{L}]}} u_1$ and $v_1 \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} v_2$. There are six cases to consider.

<u>Case 1:</u> If $(u_1, v_1) \in {}^{[z]} \leq_{X_{\mathbf{K}}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then there exists $w_1, w_2 \in X_{\mathbf{K}}$ such that $u_1 = (w_1)^{[z]}, v_1 = (w_2)^{[z]}$ and $w_1 \leq_{X_{\mathbf{K}}} w_2$. Moreover, $v_1 \leq_{X_{\mathbf{K}|\mathbf{L}|}} v_2$ implies




Fig. 10 The representations of $\mathbf{K},\,\mathbf{L}$ and $\mathbf{K}[\mathbf{L}].$

that $v_2 \in [z]X_{\mathbf{K}}$ and there exists $w_3 \in X_{\mathbf{K}}$ such that $v_2 = (w_3)^{[z]}$ and $w_2 \leq_{X_{\mathbf{K}}} w_3$. On the other hand, $u_2 \leq_{X_{\mathbf{K}|\mathbf{L}|}} u_1$ gives rise to three subcases:

- If $u_2 \in [z] \subseteq X_{\mathbf{L}}$, then $(u_2, v_2) \in [z] \times {}^{[z]}X_{\mathbf{K}} \subseteq \psi(m)$.
- If $u_2 \in {}_{[z]}X_{\mathbf{K}}$, then $(u_2, v_2) \in {}_{[z]}X_{\mathbf{K}} \times {}^{[z]}X_{\mathbf{K}} \subseteq \psi(m)$.
- If $u_2 \in [z]X_{\mathbf{K}}$, then there exists $w \in X_{\mathbf{K}}$ such that $u_2 = w^{[z]}$. Since $u_2 \leq_{X_{\mathbf{K}[\mathbf{L}]}} u_1$, it follows that $w \leq_{X_{\mathbf{K}}} w_1$ and we have that $w \leq_{X_{\mathbf{K}}} w_1 \leq_{X_{\mathbf{K}}} w_2 \leq_{X_{\mathbf{K}}} w_3$. This in turn implies that $(u_2, v_2) \in [z] \leq_{X_{\mathbf{K}}} \subseteq \psi(m)$.

<u>Case 2:</u> If $(u_1, v_1) \in [z] \leq X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then there exists $w_1, w_2 \in X_{\mathbf{K}}$ such that $u_1 = (w_1)_{[z]}, v_1 = (w_2)_{[z]}$ and $w_1 \leq_{X_{\mathbf{K}}} w_2$. Moreover, $u_2 \leq_{X_{\mathbf{K}[\mathbf{L}]}} u_1$ implies that $u_2 \in [z]X_{\mathbf{K}}$ and there exists $w \in X_{\mathbf{K}}$ such that $u_2 = w_{[z]}$ and $w \leq_{X_{\mathbf{K}}} w_1$. On the other hand, $v_1 \leq_{X_{\mathbf{K}[\mathbf{L}]}} v_2$ gives rise to three subcases:

- If $v_2 \in [z] \subseteq X_{\mathbf{L}}$, then $(u_2, v_2) \in [z] X_{\mathbf{K}[\mathbf{L}]} \times [z] \subseteq \psi(m)$.
- If $v_2 \in {}^{[z]}X_{\mathbf{K}}$, then $(u_2, v_2) \in {}_{[z]}X_{\mathbf{K}} \times {}^{[z]}X_{\mathbf{K}} \subseteq \psi(m)$.
- If $v_2 \in [z]X_{\mathbf{K}}$, then there exists $w_3 \in X_{\mathbf{K}}$ such that $v_2 = (w_3)_{[z]}$. Since we have $v_1 \leq_{X_{\mathbf{K}[\mathbf{L}]}} v_2$, it follows that $w_2 \leq_{X_{\mathbf{K}}} w_3$ and we get $w \leq_{X_{\mathbf{K}}} w_1 \leq_{X_{\mathbf{K}}} w_2 \leq_{X_{\mathbf{K}}} w_3$. This in turn implies that $(u_2, v_2) \in [z] \leq_{X_{\mathbf{K}}} \subseteq \psi(m)$.

<u>Case 3:</u> If $(u_1, v_1) \in [z] \times {}^{[z]}X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then $v_1 \leq_{X_{\mathbf{K}[\mathbf{L}]}} v_2$ implies that $v_2 \in {}^{[z]}X_{\mathbf{K}}$. On the other hand, $u_2 \leq_{X_{\mathbf{K}[\mathbf{L}]}} u_1$ gives rise to two subcases:

- If $u_2 \in [z]$, then $(u_2, v_2) \in [z] \times {}^{[z]}X_{\mathbf{K}} \subseteq \psi(m)$.
- If $u_2 \in {}_{[z]}X_{\mathbf{K}}$, then $(u_2, v_2) \in {}_{[z]}X_{\mathbf{K}} \times {}^{[z]}X_{\mathbf{K}} \subseteq \psi(m)$.

<u>Case 4</u>: If $(u_1, v_1) \in {}_{[z]}X_{\mathbf{K}} \times [z]$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then $u_2 \leq_{X_{\mathbf{K}[\mathbf{L}]}} u_1$ implies that $u_2 \in {}_{[z]}X_{\mathbf{K}}$. On the other hand, $v_1 \leq_{X_{\mathbf{K}[\mathbf{L}]}} v_2$ gives rise to two subcases:

- If $v_2 \in [z]$, then $(u_2, v_2) \in {}_{[z]}X_{\mathbf{K}} \times [z] \subseteq \psi(m)$.
- If $v_2 \in {}^{[z]}X_{\mathbf{K}}$, then $(u_2, v_2) \in {}_{[z]}X_{\mathbf{K}} \times {}^{[z]}X_{\mathbf{K}} \subseteq \psi(m)$.

<u>Case 5:</u> If $(u_1, v_1) \in {}_{[z]}X_{\mathbf{K}} \times {}^{[z]}X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then $u_2 \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} u_1$ implies that $u_2 \in {}_{[z]}X_{\mathbf{K}}$ and $v_1 \leqslant_{X_{\mathbf{K}[\mathbf{L}]}} v_2$ implies that $v_2 \in {}^{[z]}X_{\mathbf{K}}$. Therefore, it follows that $(u_2, v_2) \in {}_{[z]}X_{\mathbf{K}} \times {}^{[z]}X_{\mathbf{K}} \subseteq \psi(m)$.

<u>Case 6:</u> If $(u_1, v_1) \in \varphi_{\mathbf{L}}(m)$, then $u_1, v_1 \in X_{\mathbf{L}}$ and $(u_1, v_1) \in E_{\mathbf{L}}$. The inequalities $u_2 \leq_{X_{\mathbf{K}[\mathbf{L}]}} u_1$ and $v_1 \leq_{X_{\mathbf{K}[\mathbf{L}]}} v_2$ combined give rise to four subcases:

- If $u_2 \in [u_1]X_{\mathbf{K}}$ and $v_2 \in [u_1]X_{\mathbf{K}}$, then $(u_2, v_2) \in [u_1]X_{\mathbf{K}} \times [u_1]X_{\mathbf{K}} \subseteq \psi(m)$.
- If $u_2 \in [u_1]X_{\mathbf{K}}$ and $v_2 \in [u_1]$, then $(u_2, v_2) \in [u_1]X_{\mathbf{K}} \times [u_1] \subseteq \psi(m)$.
- If $u_2 \in [u_1]$ and $v_2 \in {}^{[u_1]}X_{\mathbf{K}}$, then $(u_2, v_2) \in [u_1] \times {}^{[u_1]}X_{\mathbf{K}} \subseteq \psi(m)$.
- If $u_2 \in [u_1]$ and $v_2 \in [u_1]$, then $u_2 \leq_{X_{\mathbf{L}}} u_1$ and $v_1 \leq_{X_{\mathbf{L}}} v_2$, which means $(u_1, v_1) \preccurlyeq_{E_{\mathbf{L}}} (u_2, v_2)$. But $\varphi_{\mathbf{L}}(m)$ is an upset of $(E_{\mathbf{L}}, \preccurlyeq_{E_{\mathbf{L}}})$. Thus, $(u_1, v_1) \in \varphi_{\mathbf{L}}(m)$ implies $(u_2, v_2) \in \varphi_{\mathbf{L}}(m) \subseteq \psi(m)$.

We now prove a series of lemmas that, when combined, show that the map ψ defined in (13) is a DqRA embedding.

Lemma 4.8. The map $\psi : K \setminus \{a_0\} \cup L \to \mathsf{Up}\left(E_{\mathbf{K}[\mathbf{L}]}, \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}}\right)$ is injective and $\psi(\mathbf{1}_{\mathbf{K}[\mathbf{L}]}) = \leqslant_{X_{\mathbf{K}[\mathbf{L}]}}$.

Proof. Since $\varphi_{\mathbf{L}}(1_{\mathbf{L}}) = \leq_{X_{\mathbf{L}}}$, it follows directly from the definitions of $\psi(m)$ and $\leq_{X_{\mathbf{K}[\mathbf{L}]}}$ in (13) and (7), respectively that $\psi(1_{\mathbf{K}[\mathbf{L}]}) = \leq_{X_{\mathbf{K}[\mathbf{L}]}}$.

To show that ψ is injective, let $m_1, m_2 \in K \setminus \{a_0\} \cup L$ such that $m_1 \neq m_2$. Without loss of generality we can consider four cases.

<u>Case 1</u>: If $m_1, m_2 \in L$, then $\varphi_{\mathbf{L}}(m_1) \neq \varphi_{\mathbf{L}}(m_2)$ since $\varphi_{\mathbf{L}}$ is injective and therefore we get $\psi(m_1) \neq \psi(m_2)$.

<u>Case 2</u>: If $m_1 = a_{-1}$ and $m_2 \in L$, then $\psi(m_1) = \emptyset$. On the other hand, if we let $w \in X_{\mathbf{K}}$ and $z \in X_{\mathbf{L}}$, then $(w_{[z]}, z) \in \psi(m_2)$ which shows $\psi(m_2) \neq \emptyset$ and hence $\psi(m_1) \neq \psi(m_2)$.

<u>Case 3</u>: If $m_1 = a_1$ and $m_2 \in L$, then $\psi(m_1) = E_{\mathbf{K}[\mathbf{L}]}$. Let $w \in X_{\mathbf{K}}$ and $z \in X_{\mathbf{L}}$, then $(z, w_{[z]}) \in E_{\mathbf{K}[\mathbf{L}]} = \psi(m_1)$, but by (13) we have $(z, w_{[z]}) \notin \psi(m_2)$. Hence, it follows that $\psi(m_1) \neq \psi(m_2)$.

<u>Case 4</u>: If $m_1, m_2 \in K \setminus \{a_0\}$, then we may assume, without loss of generality, that $m_1 = a_1$ and $m_2 = a_{-1}$. Then $\psi(m_1) \neq \emptyset$ since $E_{\mathbf{K}[\mathbf{L}]} \neq \emptyset$ and therefore we get $\psi(m_1) \neq \psi(m_2)$.

In order to prove that ψ preserves the linear negations and the involution, we will need the following auxiliary result.

Lemma 4.9. Let $E_{\mathbf{K}[\mathbf{L}]}$ be the relation defined in (8) and $\alpha_{\mathbf{K}[\mathbf{L}]}$ and $\beta_{\mathbf{K}[\mathbf{L}]}$ the maps defined in (10) and (11), respectively. Then, for $\gamma \in \{\alpha_{\mathbf{K}[\mathbf{L}]}, \beta_{\mathbf{K}[\mathbf{L}]}\}$, we have

 $\gamma\,; \varnothing = \varnothing, \quad \varnothing\,; \gamma = \varnothing, \quad \gamma\,; E_{\mathbf{K}[\mathbf{L}]} = E_{\mathbf{K}[\mathbf{L}]}, \quad and \quad E_{\mathbf{K}[\mathbf{L}]}\,; \gamma = E_{\mathbf{K}[\mathbf{L}]}.$

Proof. For $\gamma \in \{\alpha_{\mathbf{K}[\mathbf{L}]}, \beta_{\mathbf{K}[\mathbf{L}]}\}$ it follows directly from $\gamma \subseteq E_{\mathbf{K}[\mathbf{L}]}$ and the transitivity of $E_{\mathbf{K}[\mathbf{L}]}$ that γ ; $E_{\mathbf{K}[\mathbf{L}]} \subseteq E_{\mathbf{K}[\mathbf{L}]}$. For the inclusion in the other direction, let $(u, v) \in E_{\mathbf{K}[\mathbf{L}]}$. Then $(v, u) \in E_{\mathbf{K}[\mathbf{L}]}$ by symmetry and, since $(u, \gamma(u)) \in \gamma \subseteq E_{\mathbf{K}[\mathbf{L}]}$, we have that $(v, \gamma(u)) \in E_{\mathbf{K}[\mathbf{L}]}$ by transitivity. This means $(\gamma(u), v) \in E_{\mathbf{K}[\mathbf{L}]}$, and therefore $(u, v) \in \gamma$; $E_{\mathbf{K}[\mathbf{L}]}$.

The first two equalities are trivially true and the final equality can be proved similarly to the above. $\hfill \Box$

Lemma 4.10. The map $\psi : K \setminus \{a_0\} \cup L \to \mathsf{Up}\left(E_{\mathbf{K}[\mathbf{L}]}, \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}}\right)$ preserves the linear negations $\sim_{\mathbf{K}[\mathbf{L}]}$ and $-_{\mathbf{K}[\mathbf{L}]}$, i.e., for $m \in K \setminus \{a_0\} \cup L$,

$$\psi(-_{\mathbf{K}[\mathbf{L}]}m) = -\psi(m) \quad and \quad \psi(\sim_{\mathbf{K}[\mathbf{L}]}m) = \sim\psi(m).$$

Proof. We prove the claim for $-\mathbf{K}[\mathbf{L}]$. The result for $\sim_{\mathbf{K}[\mathbf{L}]}$ can be proved analogously. If $m \in K \setminus \{a_0\} \cup L$, then $m \in K \setminus \{a_0\}$ or $m \in L$.

<u>Case 1</u>: If $m \in K \setminus \{a_0\}$, then $m = a_{-1}$ and $\psi(m) = \emptyset$, or $m = a_1$ and $\psi(m) = E_{\mathbf{K}[\mathbf{L}]}$. Note that $E^c_{\mathbf{K}[\mathbf{L}]} = \emptyset$ hence $(E^c_{\mathbf{K}[\mathbf{L}]})^{\smile} = \emptyset^{\smile} = \emptyset$. Also $\emptyset^c = E_{\mathbf{K}[\mathbf{L}]}$ hence $(\emptyset^c)^{\smile} = E^{\smile}_{\mathbf{K}[\mathbf{L}]} = E_{\mathbf{K}[\mathbf{L}]}$ since $E_{\mathbf{K}[\mathbf{L}]}$ is symmetric. By Lemmas 2.6 and 4.9 we have

$$\psi(-_{\mathbf{K}[\mathbf{L}]}a_1) = \psi(-_{\mathbf{K}}a_1) = \psi(a_{-1}) = \varnothing = \alpha_{\mathbf{K}[\mathbf{L}]} ; \varnothing = \alpha_{\mathbf{K}[\mathbf{L}]} ; (E^c_{\mathbf{K}[\mathbf{L}]})^{\smile}$$
$$= -E_{\mathbf{K}[\mathbf{L}]} = -\psi(a_1)$$

and

$$\psi(-_{\mathbf{K}[\mathbf{L}]}a_{-1}) = \psi(-_{\mathbf{K}}a_{-1}) = \psi(a_1) = E_{\mathbf{K}[\mathbf{L}]} = \alpha_{\mathbf{K}[\mathbf{L}]} ; E_{\mathbf{K}[\mathbf{L}]} = \alpha_{\mathbf{K}[\mathbf{L}]} ; (\varnothing^c)^{\smile} = -\varnothing = -\psi(a_{-1})$$

<u>Case 2</u>: Let $m \in L$. To prove that $\psi(-_{\mathbf{K}[\mathbf{L}]}m) = -\psi(m)$ we first show that $\psi(-_{\mathbf{K}[\mathbf{L}]}m) \subseteq -\psi(m)$. Let $(u, v) \in \psi(-_{\mathbf{K}[\mathbf{L}]}m)$. Then, by (13) there are six sub-cases to consider.

• Let $(u, v) \in {}^{[z]} \leq_{X_{\mathbf{K}}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$. Since $X_{\mathbf{K}}$ is a two-element antichain this implies that u = v and there exists $w_1 \in X_{\mathbf{K}}$ such that $u = (w_1)^{[z]} = v$. Furthermore, $\alpha_{\mathbf{K}[\mathbf{L}]}(u) = (\alpha_{\mathbf{K}}(w_1))^{[z]} = (w_2)^{[z]}$ where $w_2 \in X_{\mathbf{K}}$ such that $w_2 \neq w_1$. Then,

$$(w_{1}, w_{2}) \in E_{\mathbf{K}} \text{ and } (w_{1}, w_{2}) \notin \leq_{X_{\mathbf{K}}}$$

$$\Rightarrow \left((w_{1})^{[z]}, (w_{2})^{[z]} \right) \in E_{\mathbf{K}[\mathbf{L}]} \text{ and } \left((w_{1})^{[z]}, (w_{2})^{[z]} \right) \notin [z] \leq_{X_{\mathbf{K}}}$$

$$\Rightarrow \left((w_{1})^{[z]}, (\alpha_{\mathbf{K}} (w_{1}))^{[z]} \right) \in E_{\mathbf{K}[\mathbf{L}]} \text{ and } \left((w_{1})^{[z]}, (\alpha_{\mathbf{K}} (w_{1}))^{[z]} \right) \notin \psi(m)$$

$$\Rightarrow (v, \alpha_{\mathbf{K}[\mathbf{L}]}(u)) \in \psi(m)^{c}$$

$$\Rightarrow (\alpha_{\mathbf{K}[\mathbf{L}]}(u), v) \in (\psi(m)^{c})^{\smile}$$

$$\Rightarrow (u, v) \in \alpha_{\mathbf{K}[\mathbf{L}]}; (\psi(m)^{c})^{\smile} = -\psi(m).$$

- Let $(u, v) \in [z] \leq X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$. Then the proof that $(u, v) \in -\psi(m)$ is similar to the proof of the previous subcase.
- Let $(u,v) \in [z] \times {}^{[z]}X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$. Then $\alpha_{\mathbf{K}[\mathbf{L}]}(u) = \alpha_{\mathbf{L}}(u)$ and $(u, \alpha_{\mathbf{L}}(u)) \in \alpha_{\mathbf{L}} \subseteq \alpha_{\mathbf{K}[\mathbf{L}]} \subseteq E_{\mathbf{K}[\mathbf{L}]}$. That is, $\alpha_{\mathbf{L}}(u) \in [z]$. Hence,

$$(v, \alpha_{\mathbf{L}}(u)) \in E_{\mathbf{K}[\mathbf{L}]} \text{ and } (v, \alpha_{\mathbf{L}}(u)) \notin \psi(m) \Rightarrow (v, \alpha_{\mathbf{K}[\mathbf{L}]}(u)) \in \psi(m)^{c}$$
$$\Rightarrow (\alpha_{\mathbf{K}[\mathbf{L}]}(u), v) \in (\psi(m)^{c})^{\smile}$$
$$\Rightarrow (u, v) \in \alpha_{\mathbf{K}[\mathbf{L}]}; (\psi(m)^{c})^{\smile} = -\psi(m).$$

• Let $(u, v) \in [z]X_{\mathbf{K}} \times [z]$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$. Then there exists $w \in X_{\mathbf{K}}$ such that $u = w_{[z]}$ and $\alpha_{\mathbf{K}[\mathbf{L}]}(u) = (\alpha_{\mathbf{K}}(w))_{[z]}$. Hence,

$$(v, (\alpha_{\mathbf{K}}(w))_{[z]}) \in E_{\mathbf{K}[\mathbf{L}]} \text{ and } (v, (\alpha_{\mathbf{K}}(w))_{[z]}) \notin \psi(m)$$

$$\Rightarrow (v, \alpha_{\mathbf{K}[\mathbf{L}]}(u)) \in \psi(m)^{c}$$

$$\Rightarrow (\alpha_{\mathbf{K}[\mathbf{L}]}(u), v) \in (\psi(m)^{c})^{\smile}$$

$$\Rightarrow (u, v) \in \alpha_{\mathbf{K}[\mathbf{L}]}; (\psi(m)^{c})^{\smile} = -\psi(m).$$

- Let $(u, v) \in {}_{[z]}X_{\mathbf{K}} \times {}^{[z]}X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$. Then the proof that $(u, v) \in -\psi(m)$ is similar to the proof of the previous subcase.
- Let $(u, v) \in \varphi_{\mathbf{L}}(-_{\mathbf{K}[\mathbf{L}]}m) = \varphi_{\mathbf{L}}(-_{\mathbf{L}}m)$. Since $\varphi_{\mathbf{L}}$ is an embedding that preserves the linear negations, it follows that

$$\begin{split} &(u,v) \in -\varphi_{\mathbf{L}}(m) = \alpha_{\mathbf{L}} ; (\varphi_{\mathbf{L}}(m)^c)^{\smile} \\ \Rightarrow &(\alpha_{\mathbf{L}}(u),v) \in (\varphi_{\mathbf{L}}(m)^c)^{\smile} \\ \Rightarrow &(v,\alpha_{\mathbf{L}}(u)) \in \varphi_{\mathbf{L}}(m)^c \\ \Rightarrow &(v,\alpha_{\mathbf{L}}(u)) \in E_{\mathbf{L}} \text{ and } (v,\alpha_{\mathbf{L}}(u)) \notin \varphi_{\mathbf{L}}(m) \\ \Rightarrow &(v,\alpha_{\mathbf{L}}(u)) \in E_{\mathbf{K}[\mathbf{L}]} \text{ and } (v,\alpha_{\mathbf{K}[\mathbf{L}]}(u)) \notin \psi(m) \\ \Rightarrow &(v,\alpha_{\mathbf{K}[\mathbf{L}]}(u)) \in \psi(m)^c \\ \Rightarrow &(\alpha_{\mathbf{K}[\mathbf{L}]}(u),v) \in (\psi(m)^c)^{\smile} \\ \Rightarrow &(u,v) \in \alpha_{\mathbf{K}[\mathbf{L}]} ; (\psi(m)^c)^{\smile} = -\psi(m). \end{split}$$

For the inclusion in the other direction, let $(u, v) \in -\psi(m) = \alpha_{\mathbf{K}[\mathbf{L}]}$; $(\psi(m)^c)^{\smile}$. Then,

$$\begin{aligned} (u,v) \in \alpha_{\mathbf{K}[\mathbf{L}]} ; (\psi(m)^c)^{\smile} &\Rightarrow (\alpha_{\mathbf{K}[\mathbf{L}]}(u), v) \in (\psi(m)^c)^{\smile} \\ &\Rightarrow (v, \alpha_{\mathbf{K}[\mathbf{L}]}(u)) \in \psi(m)^c \\ &\Rightarrow \left(v, \alpha_{\mathbf{K}[\mathbf{L}]}(u)\right) \in E_{\mathbf{K}[\mathbf{L}]} \text{ and } (v, \alpha_{\mathbf{K}[\mathbf{L}]}(u)) \notin \psi(m). \end{aligned}$$

Since $\alpha_{\mathbf{K}[\mathbf{L}]} \subseteq E_{\mathbf{K}[\mathbf{L}]}$ and $E_{\mathbf{K}[\mathbf{L}]}$ is symmetric and transitive, we get $(u, v) \in E_{\mathbf{K}[\mathbf{L}]}$. Hence, there exists $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$ such that $(u, v) \in ([z] \cup [z]X_{\mathbf{K}} \cup [z]X_{\mathbf{K}})^2$ by (8).

Suppose $(u, v) \in [{}^{[z]}X_{\mathbf{K}} \times [z]$. Then $v \in [z]$ and there exists $w \in X_{\mathbf{K}}$ such that $u = w^{[z]}$ and $\alpha_{\mathbf{K}[\mathbf{L}]}(u) = (\alpha_{\mathbf{K}}(w))^{[z]} \in [{}^{[z]}X_{\mathbf{K}}$. But then $(v, \alpha_{\mathbf{K}[\mathbf{L}]}(u)) \notin \psi(m)$ means $(v, (\alpha_{\mathbf{K}}(w))^{[z]}) \notin \psi(m)$, which contradicts the fact that $[z] \times [{}^{[z]}X_{\mathbf{K}} \subseteq \psi(m)$ for any $m \in L$ as per the definition in (13). Hence, $-\psi(m)$ does not contain any elements of $[{}^{[z]}X_{\mathbf{K}} \times [z]$. Similarly, we can show that $-\psi(m)$ also does not contain any elements of $[{}^{[z]}X_{\mathbf{K}} \times [z]X_{\mathbf{K}}$ nor of $[z] \times [{}^{[z]}X_{\mathbf{K}}$.

If $(u,v) \in [z] \times {}^{[z]}X_{\mathbf{K}}$ or $(u,v) \in {}_{[z]}X_{\mathbf{K}} \times [z]$ or $(u,v) \in {}_{[z]}X_{\mathbf{K}} \times {}^{[z]}X_{\mathbf{K}}$, it follows immediately from the definition of ψ in (13) that $(u,v) \in \psi(-\mathbf{K}[\mathbf{L}]m)$.

- There are three subcases left to consider.
- Let $(u, v) \in {}^{[z]}X_{\mathbf{K}} \times {}^{[z]}X_{\mathbf{K}}$. Then there exist $w_1, w_2 \in X_{\mathbf{K}}$ such that $u = (w_1)^{[z]}$ and $v = (w_2)^{[z]}$. Since $X_{\mathbf{K}}$ is a two-element antichain, either $w_1 \neq w_2$ and $\alpha_{\mathbf{K}[\mathbf{L}]}(u) = (\alpha_{\mathbf{K}}(w_1))^{[z]} = (w_2)^{[z]} = v$, or $w_1 = w_2$. If $w_1 \neq w_2$, then $(v, \alpha_{\mathbf{K}[\mathbf{L}]}(u)) \notin \psi(m)$ means that $(v, (\alpha_{\mathbf{K}}(w_1))^{[z]}) = ((w_2)^{[z]}, (w_2)^{[z]}) \notin \psi(m)$, which contradicts that ${}^{[z]} \leqslant_{X_{\mathbf{K}}} \subseteq \psi(m)$ for all $m \in L$. Therefore, it must be the case that $w_1 = w_2$ and u = v, and hence $(u, v) \in {}^{[z]} \leqslant_{X_{\mathbf{K}}} \subseteq \psi(-{}_{\mathbf{K}[\mathbf{L}]}m)$.
- Let $(u, v) \in [z]X_{\mathbf{K}} \times [z]X_{\mathbf{K}}$. Then we can use a similar argument to the one used in the previous subcase to conclude that u = v and hence that $(u, v) \in [z] \leq X_{\mathbf{K}} \subseteq \psi(-\mathbf{K}[\mathbf{L}]m)$.
- Let $(u, v) \in [z] \times [z]$. Then, $(v, \alpha_{\mathbf{L}}(u)) \in E_{\mathbf{L}}$, and $(v, \alpha_{\mathbf{K}[\mathbf{L}]}(u)) \notin \psi(m)$ implies $(v, \alpha_{\mathbf{L}}(u)) \notin \varphi_{\mathbf{L}}(m)$. Hence, $(v, \alpha_{\mathbf{L}}(u)) \in \varphi_{\mathbf{L}}(m)^c$, and so $(\alpha_{\mathbf{L}}(u), v) \in (\varphi_{\mathbf{L}}(m)^c)$, which gives $(u, v) \in \alpha_{\mathbf{L}}$; $(\varphi_{\mathbf{L}}(m)^c)$. That is, $(u, v) \in -\mathbf{L}\varphi_{\mathbf{L}}(m) = \varphi_{\mathbf{L}}(-\mathbf{L}m) = \varphi_{\mathbf{L}}(-\mathbf{K}[\mathbf{L}]m) \subseteq \psi(-\mathbf{K}[\mathbf{L}]m)$ by Lemma 2.6 and since $\varphi_{\mathbf{L}}$ is an embedding that preserves the linear negations.

Lemma 4.11. The map $\psi : K \setminus \{a_0\} \cup L \to \mathsf{Up}\left(E_{\mathbf{K}[\mathbf{L}]}, \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}}\right)$ preserves the involutive operation $\neg_{\mathbf{K}[\mathbf{L}]}$, *i.e.*, for $m \in K \setminus \{a_0\} \cup L$, $\psi(\neg_{\mathbf{K}[\mathbf{L}]}m) = \neg\psi(m)$.

Proof. Recall from Theorem 3.2 that $\neg \psi(m) = \alpha_{\mathbf{K}[\mathbf{L}]}$; $\beta_{\mathbf{K}[\mathbf{L}]}$; $\psi(m)^c$; $\beta_{\mathbf{K}[\mathbf{L}]}$. Now let $m \in K \setminus \{a_0\} \cup L$.

<u>Case 1</u>: If $m \in K \setminus \{a_0\}$ there are two subcases to consider:

• If $m = a_{-1}$, then $\psi(m) = \emptyset$ and $\neg_{\mathbf{K}[\mathbf{L}]} m = \neg_{\mathbf{K}} m = a_1$. Thus, $\psi(\neg_{\mathbf{K}[\mathbf{L}]} m) = E_{\mathbf{K}[\mathbf{L}]}$. By Lemma 4.9,

$$\neg \psi(m) = \alpha_{\mathbf{K}[\mathbf{L}]} ; \beta_{\mathbf{K}[\mathbf{L}]} ; \beta^{c} ; \beta_{\mathbf{K}[\mathbf{L}]} = \alpha_{\mathbf{K}[\mathbf{L}]} ; \beta_{\mathbf{K}[\mathbf{L}]} ; E_{\mathbf{K}[\mathbf{L}]} ; \beta_{\mathbf{K}[\mathbf{L}]} = E_{\mathbf{K}[\mathbf{L}]}$$
$$= \psi(\neg_{\mathbf{K}[\mathbf{L}]}m).$$

• If $m = a_1$, then $\psi(m) = E_{\mathbf{K}[\mathbf{L}]}$. Also, $\neg_{\mathbf{K}[\mathbf{L}]}m = \neg_{\mathbf{K}}m = a_{-1}$ and $\psi(\neg_{\mathbf{K}[\mathbf{L}]}m) = \emptyset$. Using Lemma 4.9 again we have

$$\neg \psi(m) = \alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}; E^{c}_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]} = \alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}; \varnothing; \beta_{\mathbf{K}[\mathbf{L}]} = \varnothing = \psi(\neg_{\mathbf{K}[\mathbf{L}]}m)$$

<u>Case 2</u>: Next assume that $m \in L$. Then $\neg_{\mathbf{K}[\mathbf{L}]}m = \neg_{\mathbf{L}}m$ by definition. We first show that $\psi(\neg_{\mathbf{K}[\mathbf{L}]}m) \subseteq \neg\psi(m)$. Let $(u, v) \in \psi(\neg_{\mathbf{K}[\mathbf{L}]}m) = \psi(\neg_{\mathbf{L}}m)$. We consider the six cases that arise from (13).

- If $(u, v) \in [z] \leq_{X_{\mathbf{K}}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then there exists $w_1 \in X_{\mathbf{K}}$ such that $u = (w_1)^{[z]} = v$ since $\mathbf{X}_{\mathbf{K}}$ is a two-element antichain. Hence $\alpha_{\mathbf{K}[\mathbf{L}]}(u) = (\alpha_{\mathbf{K}}(w_1))^{[z]} = (w_2)^{[z]}$ where $w_2 \in X_{\mathbf{K}}$ such that $w_1 \neq w_2$. Furthermore, $\beta_{\mathbf{K}[\mathbf{L}]}(\alpha_{\mathbf{K}[\mathbf{L}]}(u)) = \beta_{\mathbf{K}[\mathbf{L}]}\left((w_2)^{[z]}\right) = (\beta_{\mathbf{K}}(w_2))_{[z]} = (w_2)_{[z]}$, and so $(u, (w_2)_{[z]}) \in \alpha_{\mathbf{K}[\mathbf{L}]}$; $\beta_{\mathbf{K}[\mathbf{L}]}(u)$ Next note that since $(w_2, w_1) \notin \leq_{X_{\mathbf{K}}}$, it follows from (13) that $((w_2)_{[z]}, (w_1)_{[z]}) \notin \psi(m)$ and hence, since $((w_2)_{[z]}, (w_1)_{[z]}) \in E_{\mathbf{K}[\mathbf{L}]}$ we get $((w_2)_{[z]}, (w_1)_{[z]}) \in \psi(m)^c$. Combined with the above, we have that $(u, (w_1)_{[z]}) \in \alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}; \psi(m)^c$. Finally, since we have $((w_1)_{[z]}, (w_1)^{[z]}) \in \beta_{\mathbf{K}[\mathbf{L}]}$ by (11), it follows that $(u, (w_1)^{[z]}) = (u, v) \in \alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}; \psi(m)^c; \beta_{\mathbf{K}[\mathbf{L}]} = \neg \psi(m)$.
- If $(u, v) \in [z] \leq X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then the proof is similar to the previous case.
- If $(u, v) \in [z] \times [z] X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then there exists $w \in X_{\mathbf{K}}$ such that $v = w^{[z]}$. Also, $\alpha_{\mathbf{K}[\mathbf{L}]}(u) = \alpha_{\mathbf{L}}(u)$ and $\beta_{\mathbf{K}[\mathbf{L}]}(\alpha_{\mathbf{L}}(u)) = \beta_{\mathbf{L}}(\alpha_{\mathbf{L}}(u))$ which implies that $(u, \beta_{\mathbf{L}}(\alpha_{\mathbf{L}}(u))) \in \alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}$. Next, by (13), $(\beta_{\mathbf{L}}(\alpha_{\mathbf{L}}(u)), (w)_{[z]}) \notin \psi(m)$, and therefore $(\beta_{\mathbf{L}}(\alpha_{\mathbf{L}}(u)), (w)_{[z]}) \in \psi(m)^c$. Consequently, we have $(u, (w)_{[z]}) \in \alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}; \psi(m)^c$. Finally, since $((w)_{[z]}, (w)^{[z]}) \in \beta_{\mathbf{K}[\mathbf{L}]}$, it follows that $(u, (w)^{[z]}) = (u, v) \in \alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}; \psi(m)^c; \beta_{\mathbf{K}[\mathbf{L}]} = \neg \psi(m)$.
- If $(u, v) \in [z] X_{\mathbf{K}} \times [z]$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then there exists $w_1 \in X_{\mathbf{K}}$ such that $u = (w_1)_{[z]}$. Hence, $\alpha_{\mathbf{K}[\mathbf{L}]}(u) = (\alpha_{\mathbf{K}}(w_1))_{[z]} = (w_2)_{[z]}$ where $w_2 \in X_{\mathbf{K}}$ such that $w_1 \neq w_2$. Furthermore, $\beta_{\mathbf{K}[\mathbf{L}]}((w_2)_{[z]}) = (\beta_{\mathbf{K}}(w_2))^{[z]} = (w_2)^{[z]}$. Thus, $(u, (w_2)^{[z]}) \in \alpha_{\mathbf{K}[\mathbf{L}]}$; $\beta_{\mathbf{K}[\mathbf{L}]}$. Next, as $\beta_{\mathbf{L}}^{-1}(v) \in X_{\mathbf{L}}$ since $v \in [z]$, we have $((w_2)^{[z]}, \beta_{\mathbf{L}}^{-1}(v)) \notin \psi(m)$ by (13). Therefore, $((w_2)^{[z]}, \beta_{\mathbf{L}}^{-1}(v)) \in \psi(m)^c$ and $(u, \beta_{\mathbf{L}}^{-1}(v)) \in \alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}; \psi(m)^c$. Finally, since $(\beta_{\mathbf{L}}^{-1}(v), v) \in \beta_{\mathbf{L}} \subseteq \beta_{\mathbf{K}[\mathbf{L}]}$, it follows that $(u, v) \in \alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}; \psi(m)^c; \beta_{\mathbf{K}[\mathbf{L}]} = \neg \psi(m)$.
- If $(u, v) \in [z] X_{\mathbf{K}} \times [z] X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then there exists $w_1, w_2 \in X_{\mathbf{K}}$ such that $u = (w_1)_{[z]}$ and $v = (w_2)^{[z]}$. It follows that $\alpha_{\mathbf{K}[\mathbf{L}]}(u) = (\alpha_{\mathbf{K}}(w_1))_{[z]} = (w_3)_{[z]}$ where $w_3 \in X_{\mathbf{K}}$ such that $w_1 \neq w_3$. Also, $\beta_{\mathbf{K}[\mathbf{L}]}((w_3)_{[z]}) = (\beta_{\mathbf{K}}(w_3))^{[z]} = (w_3)^{[z]}$. Hence, $(u, (w_3)^{[z]}) \in \alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}$. Next, $((w_3)^{[z]}, (w_2)_{[z]}) \notin \psi(m)$ by (13). Thus, $((w_3)^{[z]}, (w_2)_{[z]}) \in \psi(m)^c$ and $(u, (w_2)_{[z]}) \in \alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}; \psi(m)^c$. Finally, since $\beta_{\mathbf{K}[\mathbf{L}]}((w_2)_{[z]}) = (\beta_{\mathbf{K}}(w_2))^{[z]} = (w_2)^{[z]}$, it follows that $(u, (w_2)^{[z]}) = (u, v) \in \alpha_{\mathbf{K}[\mathbf{L}]}; \beta_{\mathbf{K}[\mathbf{L}]}; \psi(m)^c; \beta_{\mathbf{K}[\mathbf{L}]} = \neg \psi(m)$.
- If $(u, v) \in \varphi_{\mathbf{L}}(\neg_{\mathbf{L}} m)$, then, since $\varphi_{\mathbf{L}}$ is an embedding, we have

$$(u, v) \in \varphi_{\mathbf{L}}(\neg_{\mathbf{L}} m) = \neg \varphi_{\mathbf{L}}(m)$$

= $\alpha_{\mathbf{L}}$; $\beta_{\mathbf{L}}$; $\varphi_{\mathbf{L}}(m)^{c}$; $\beta_{\mathbf{L}}$
 $\subseteq \alpha_{\mathbf{K}[\mathbf{L}]}$; $\beta_{\mathbf{K}[\mathbf{L}]}$; $\psi(m)^{c}$; $\beta_{\mathbf{K}[\mathbf{L}]} = \neg \psi(m)$.

For the containment in the other direction, let $(u, v) \in \neg \psi(m)$. Then we have $(u, v) \in \alpha_{\mathbf{K}[\mathbf{L}]}$; $\beta_{\mathbf{K}[\mathbf{L}]}$; $\psi(m)^c$; $\beta_{\mathbf{K}[\mathbf{L}]}$, i.e., $\left(\beta_{\mathbf{K}[\mathbf{L}]}(\alpha_{\mathbf{K}[\mathbf{L}]}(u)), \beta_{\mathbf{K}[\mathbf{L}]}^{-1}(v)\right) \in \psi(m)^c$, i.e.,

 $\left(\beta_{\mathbf{K}[\mathbf{L}]}(\alpha_{\mathbf{K}[\mathbf{L}]}(u)), \beta_{\mathbf{K}[\mathbf{L}]}^{-1}(v)\right) \in E_{\mathbf{K}[\mathbf{L}]} \text{ and } \left(\beta_{\mathbf{K}[\mathbf{L}]}(\alpha_{\mathbf{K}[\mathbf{L}]}(u)), \beta_{\mathbf{K}[\mathbf{L}]}^{-1}(v)\right) \notin \psi(m).$ Since $\alpha_{\mathbf{K}[\mathbf{L}]}, \beta_{\mathbf{K}[\mathbf{L}]} \subseteq E_{\mathbf{K}[\mathbf{L}]} \text{ and } E_{\mathbf{K}[\mathbf{L}]} \text{ is transitive, we get } (u, v) \in E_{\mathbf{K}[\mathbf{L}]}.$

Now suppose $u \in [z]X_{\mathbf{K}}$ and $v \in [z]$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$. Then there exists $w \in X_{\mathbf{K}}$ such that $u = w^{[z]}$ and $\beta_{\mathbf{K}[\mathbf{L}]}(\alpha_{\mathbf{K}[\mathbf{L}]}(u)) = (\beta_{\mathbf{K}}(\alpha_{\mathbf{K}}(w)))_{[z]}$. Also $\beta_{\mathbf{K}[\mathbf{L}]}^{-1}(v) = \beta_{\mathbf{L}}^{-1}(v) \in [z]$, since $\beta_{\mathbf{L}} \subseteq E_{\mathbf{L}}$. But then $(\beta_{\mathbf{K}[\mathbf{L}]}(\alpha_{\mathbf{K}[\mathbf{L}]}(u)), \beta_{\mathbf{K}[\mathbf{L}]}^{-1}(v)) \in [z]X_{\mathbf{K}} \times [z] \subseteq \psi(m)$, contradicting the previous statement that this pair is not a member of $\psi(m)$. Hence, it cannot be the case that $u \in [z]X_{\mathbf{K}}$ and $v \in [z]$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$. Similarly we can show that it can also not be the case that $u \in [z]$ and $v \in [z]X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$.

On the other hand, if $u \in [z]X_{\mathbf{K}}$ and $v \in [z]$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then it is immediate that $(u, v) \in [z]X_{\mathbf{K}} \times [z] \subseteq \psi(\neg_{\mathbf{K}[\mathbf{L}]}m)$ by (13). Similarly, if $u \in [z]X_{\mathbf{K}}$ and $v \in [z]X_{\mathbf{K}}$, or if $u \in [z]$ and $v \in [z]X_{\mathbf{K}}$, then $(u, v) \in [z]X_{\mathbf{K}} \times [z]X_{\mathbf{K}} \subseteq \psi(\neg_{\mathbf{K}[\mathbf{L}]}m)$ or, respectively, $(u, v) \in [z] \times [z]X_{\mathbf{K}} \subseteq \psi(\neg_{\mathbf{K}[\mathbf{L}]}m)$. Thus we are left with three subcases to consider.

• If $u, v \in X_{\mathbf{L}}$, then [u] = [v] since $(u, v) \in E_{\mathbf{K}[\mathbf{L}]}$. Furthermore, $\beta_{\mathbf{K}[\mathbf{L}]}(\alpha_{\mathbf{K}[\mathbf{L}]}(u)) = \beta_{\mathbf{L}}(\alpha_{\mathbf{L}}(u)) \in [u]$ and $\beta_{\mathbf{K}[\mathbf{L}]}^{-1}(v) = \beta_{\mathbf{L}}^{-1}(v) \in [v]$ since $\beta_{\mathbf{L}} \subseteq \beta_{\mathbf{K}[\mathbf{L}]} \subseteq E_{\mathbf{K}[\mathbf{L}]}$ and $\alpha_{\mathbf{L}} \subseteq \alpha_{\mathbf{K}[\mathbf{L}]} \subseteq E_{\mathbf{K}[\mathbf{L}]}$. Hence,

$$\begin{pmatrix} \beta_{\mathbf{K}[\mathbf{L}]}(\alpha_{\mathbf{K}[\mathbf{L}]}(u)), \beta_{\mathbf{K}[\mathbf{L}]}^{-1}(v) \end{pmatrix} \notin \psi(m) \\ \Rightarrow \left(\beta_{\mathbf{L}}(\alpha_{\mathbf{L}}(u)), \beta_{\mathbf{L}}^{-1}(v) \right) \notin \varphi_{\mathbf{L}}(m) \\ \Rightarrow \left(\beta_{\mathbf{L}}(\alpha_{\mathbf{L}}(u)), \beta_{\mathbf{L}}^{-1}(v) \right) \in \varphi_{\mathbf{L}}(m)^{c} \\ \Rightarrow (u, v) \in \alpha_{\mathbf{L}} ; \beta_{\mathbf{L}} ; \varphi_{\mathbf{L}}(m)^{c} ; \beta_{\mathbf{L}} \\ \Rightarrow (u, v) \in \neg \varphi_{\mathbf{L}}(m) = \varphi_{\mathbf{L}}(\neg_{\mathbf{L}}m) \subseteq \psi(\neg_{\mathbf{K}[\mathbf{L}]}m)$$

since $\varphi_{\mathbf{L}}$ is an embedding that preserves the involution.

- If $u, v \in [z]X_{\mathbf{K}}$, then there exist $w_1, w_2 \in X_{\mathbf{K}}$ such that $u = (w_1)_{[z]}$ and $v = (w_2)_{[z]}$. If $\alpha_{\mathbf{K}}(w_1) = w_3$ where $w_3 \in X_{\mathbf{K}}$ such that $w_1 \neq w_3$, then $\beta_{\mathbf{K}[\mathbf{L}]}(\alpha_{\mathbf{K}[\mathbf{L}]}(u)) = \beta_{\mathbf{K}[\mathbf{L}]}(\alpha_{\mathbf{K}}(w_1)_{[z]}) = (\beta_{\mathbf{K}}(w_3))^{[z]} = (w_3)^{[z]}$. Also, $\beta_{\mathbf{K}[\mathbf{L}]}^{-1}(v) =$ $(\beta_{\mathbf{K}}^{-1}(w_2))^{[z]} = (w_2)^{[z]}$. Hence, $(\beta_{\mathbf{K}[\mathbf{L}]}(\alpha_{\mathbf{K}[\mathbf{L}]}(u)), \beta_{\mathbf{K}[\mathbf{L}]}^{-1}(v)) \notin \psi(m)$ implies $((w_3)^{[z]}, (w_2)^{[z]}) \notin \psi(m)$, which means $w_3 \neq w_2$. But then, since $X_{\mathbf{K}}$ only contains two elements, it must be the case that $w_1 = w_2$ and u = v. Hence, $(u, v) \in [z] \leq X_{\mathbf{K}} \subseteq \psi(\neg_{\mathbf{K}[\mathbf{L}]}m)$ by (13).
- If $u, v \in {}^{[z]}X_{\mathbf{K}}$, then the proof is similar to the previous case.

Lemma 4.12. The map $\psi : (K \setminus \{a_0\}) \cup L \to \mathsf{Up}\left(E_{\mathbf{K}[\mathbf{L}]}, \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}}\right)$ preserves the lattice operations $\wedge_{\mathbf{K}[\mathbf{L}]}$ and $\vee_{\mathbf{K}[\mathbf{L}]}$, *i.e.*, for $m_1, m_2 \in (K \setminus \{a_0\}) \cup L$,

$$\psi(m_1 \wedge_{\mathbf{K}[\mathbf{L}]} m_2) = \psi(m_1) \cap \psi(m_2) \quad and \quad \psi(m_1 \vee_{\mathbf{K}[\mathbf{L}]} m_2) = \psi(m_1) \cup \psi(m_2).$$

Proof. We use the definitions of $\wedge_{\mathbf{K}[\mathbf{L}]}$ and $\vee_{\mathbf{K}[\mathbf{L}]}$ as described in (3) and (4). Their commutativity reduces the number of cases we need to consider.

<u>Case 1:</u> $m_1 = a_{-1}$ or $m_2 = a_{-1}$. Without loss of generality, suppose $m_1 = a_{-1}$. Then

$$\psi(m_1 \wedge_{\mathbf{K}[\mathbf{L}]} m_2) = \psi(a_{-1}) = \varnothing = \varnothing \cap \psi(m_2) = \psi(m_1) \cap \psi(m_2), \text{ and}$$

$$\psi(m_1 \vee_{\mathbf{K}[\mathbf{L}]} m_2) = \psi(m_2) = \varnothing \cup \psi(m_2) = \psi(m_1) \cup \psi(m_2).$$

<u>Case 2</u>: $m_1 = a_1$ or $m_2 = a_1$. Without loss of generality, suppose $m_1 = a_1$. Then

$$\psi(m_1 \wedge_{\mathbf{K}[\mathbf{L}]} m_2) = \psi(m_2) = E_{\mathbf{K}[\mathbf{L}]} \cap \psi(m_2) = \psi(m_1) \cap \psi(m_2), \text{ and}$$

$$\psi(m_1 \vee_{\mathbf{K}[\mathbf{L}]} m_2) = \psi(a_1) = E_{\mathbf{K}[\mathbf{L}]} = E_{\mathbf{K}[\mathbf{L}]} \cup \psi(m_2) = \psi(m_1) \cup \psi(m_2).$$

<u>Case 3:</u> Finally, if $m_1, m_2 \in L$, then $m_1 \wedge_{\mathbf{K}[\mathbf{L}]} m_2 = m_1 \wedge_{\mathbf{L}} m_2$ and $m_1 \vee_{\mathbf{K}[\mathbf{L}]} m_2 = m_1 \wedge_{\mathbf{L}} m_2$.

Recall the relation $R = \leq_{X_{\mathbf{K}[\mathbf{L}]}} \leq_{\mathbf{X}_{\mathbf{L}}} \text{ defined in (12). We have } \psi(m) = R \cup \varphi_{\mathbf{L}}(m)$ for any $m \in L$. Then, since $\varphi_{\mathbf{L}}$ preserves $\wedge_{\mathbf{L}}$ and $\vee_{\mathbf{L}}$, we have

$$\psi(m_1 \wedge_{\mathbf{K}[\mathbf{L}]} m_2) = R \cup \varphi_{\mathbf{L}}(m_1 \wedge_{\mathbf{L}} m_2) = R \cup (\varphi_{\mathbf{L}}(m_1) \cap \varphi_{\mathbf{L}}(m_2))$$
$$= (R \cup \varphi_{\mathbf{L}}(m_1)) \cap (R \cup \varphi_{\mathbf{L}}(m_2)) = \psi(m_1) \cap \psi(m_2),$$

and

$$\psi(m_1 \vee_{\mathbf{K}[\mathbf{L}]} m_2) = R \cup \varphi_{\mathbf{L}}(m_1 \vee_{\mathbf{L}} m_2) = R \cup (\varphi_{\mathbf{L}}(m_1) \cup \varphi_{\mathbf{L}}(m_2))$$
$$= (R \cup \varphi_{\mathbf{L}}(m_1)) \cup (R \cup \varphi_{\mathbf{L}}(m_2)) = \psi(m_1) \cup \psi(m_2).$$

Lastly, we need to show that ψ preserves the monoid operation.

Lemma 4.13. The map $\psi : K \setminus \{a_0\} \cup L \to \bigcup \bigcup (E_{\mathbf{K}[\mathbf{L}]}, \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}})$ preserves the monoid operation, $\cdot_{\mathbf{K}[\mathbf{L}]}$, *i.e.*, for $m_1, m_2 \in K \setminus \{a_0\} \cup L$,

$$\psi(m_1 \cdot_{\mathbf{K}[\mathbf{L}]} m_2) = \psi(m_1) ; \psi(m_2).$$

Proof. Recall the definition of $\cdot_{\mathbf{K}[\mathbf{L}]}$ from (2). Case 1: $m_1 = a_{-1}$ or $m_2 = a_{-1}$. Without loss of generality, assume $m_1 = a_{-1}$. Then

$$\psi(m_1 \cdot_{\mathbf{K}[\mathbf{L}]} m_2) = \psi(a_{-1}) = \emptyset = \emptyset; \ \psi(m_2) = \psi(a_{-1}); \ \psi(m_2) = \psi(m_1); \ \psi(m_2).$$

<u>Case 2</u>: $m_1 = a_1 = m_2$. Then $m_1 \cdot_{\mathbf{K}[\mathbf{L}]} m_2 = m_1 \cdot_{\mathbf{K}} m_2 = a_1$ and $\psi(m_1 \cdot_{\mathbf{K}[\mathbf{L}]} m_2) = \psi(a_1) = E_{\mathbf{K}[\mathbf{L}]} = E_{\mathbf{K}[\mathbf{L}]}$; $E_{\mathbf{K}[\mathbf{L}]} = \psi(m_1)$; $\psi(m_2)$, where the equality $E_{\mathbf{K}[\mathbf{L}]} = E_{\mathbf{K}[\mathbf{L}]}$; $E_{\mathbf{K}[\mathbf{L}]}$ follows from $E_{\mathbf{K}[\mathbf{L}]}$ being reflexive and transitive.

<u>Case 3:</u> If $m_1, m_2 \in L$, then $m_1 \cdot_{\mathbf{K}[\mathbf{L}]} m_2 = m_1 \cdot_{\mathbf{L}} m_2$. We first show that $\psi(m_1 \cdot_{\mathbf{K}[\mathbf{L}]} m_2) \subseteq \psi(m_1); \psi(m_2)$. Let $(u, v) \in \psi(m_1 \cdot_{\mathbf{K}[\mathbf{L}]} m_2)$. By (13), there are six subcases to consider:

• If $(u,v) \in {}^{[z]} \leq_{X_{\mathbf{K}}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then $(u,v) \in \psi(m_1)$ and $(v,v) \in \psi(m_2)$ which implies that $(u,v) \in \psi(m_1)$; $\psi(m_2)$.

- If $(u, v) \in [z] \leq X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$ the proof is similar to the proof of the previous case.
- If $(u,v) \in [z] \times [z] X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then we have $(u,v) \in \psi(m_1)$ and $(v,v) \in [z] \leq_{X_{\mathbf{K}}} \subseteq \psi(m_2)$, which means $(u,v) \in \psi(m_1)$; $\psi(m_2)$.
- If $(u, v) \in [z]X_{\mathbf{K}} \times [z]X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, the proof is similar to the proof of the previous case.
- If $(u,v) \in {}_{[z]}X_{\mathbf{K}} \times [z]$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then $(u,u) \in {}_{[z]} \leq X_{\mathbf{K}} \subseteq \psi(m_1)$, $(u,v) \in \psi(m_2)$ and hence $(u,v) \in \psi(m_1)$; $\psi(m_2)$.
- If $(u, v) \in \varphi_{\mathbf{L}}(m_1 \cdot_{\mathbf{L}} m_2)$, then since $\varphi_{\mathbf{L}}$ preserves $\cdot_{\mathbf{L}}$, $(u, v) \in \varphi_{\mathbf{L}}(m_1)$; $\varphi_{\mathbf{L}}(m_2)$. Therefore, there exist $u_1, v_1 \in X_{\mathbf{L}}$ such that $(u, u_1) \in \varphi_{\mathbf{L}}(m_1) \subseteq \psi(m_1)$ and $(v_1, v) \in \varphi_{\mathbf{L}}(m_2) \subseteq \psi(m_2)$. It follows that $(u, v) \in \psi(m_1)$; $\psi(m_2)$.

For the inclusion in the other direction, let $(u, v) \in \psi(m_1)$; $\psi(m_2)$. Then there exists $u_1 \in X_{\mathbf{K}[\mathbf{L}]}$ such that $(u, u_1) \in \psi(m_1)$ and $(u_1, v) \in \psi(m_2)$. There are three subcases to consider:

- If $u_1 \in {}_{[z]}X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then $(u, u_1) \in \psi(m_1)$ implies that $u \in {}_{[z]}X_{\mathbf{K}}$. On the other hand, $(u_1, v) \in \psi(m_2)$ implies that $v \in {}_{[z]}X_{\mathbf{K}}$, or $v \in [z]$, or $v \in {}^{[z]}X_{\mathbf{K}}$. These three possibilities in turn respectively imply that $(u, v) \in {}_{[z]} \leq X_{\mathbf{K}}$, or $(u, v) \in {}_{[z]}X_{\mathbf{K}} \times [z]$.
- If $u_1 \in {}^{[z]}X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then $(u_1, v) \in \psi(m_2)$ implies that $v \in {}^{[z]}X_{\mathbf{K}}$. On the other hand, $(u, u_1) \in \psi(m_1)$ implies that $u \in {}_{[z]}X_{\mathbf{K}}$, or $u \in [z]$, or $u \in {}^{[z]}X_{\mathbf{K}}$. Respectively, these possibilities imply that $(u, v) \in {}_{[z]}X_{\mathbf{K}} \times {}^{[z]}X_{\mathbf{K}}$, or $(u, v) \in [z] \times {}^{[z]}X_{\mathbf{K}}$, or $(u, v) \in {}^{[z]} \leq_{X_{\mathbf{K}}}$. Hence, $(u, v) \in \psi(m_1 \cdot_{\mathbf{K}[\mathbf{L}]} m_2)$.
- If $u_1 \in [z]$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then there are four possibilities:
 - $-u \in {}_{[z]}X_{\mathbf{K}}, v \in [z] \text{ and hence } (u,v) \in {}_{[z]}X_{\mathbf{K}} \times [z].$
 - $-u \in {}_{[z]}X_{\mathbf{K}}, v \in {}^{[z]}X_{\mathbf{K}} \text{ and hence } (u,v) \in {}_{[z]}X_{\mathbf{K}} \times {}^{[z]}X_{\mathbf{K}}.$
 - $-u \in [z], v \in [z]X_{\mathbf{K}}$ and hence $(u, v) \in [z] \times [z]X_{\mathbf{K}}$.
 - $-u \in [z]$ and $v \in [z]$. Consequently, since $(u, v) \in \psi(m_1)$; $\psi(m_2)$ we get $(u, v) \in \varphi_{\mathbf{L}}(m_1)$; $\varphi_{\mathbf{L}}(m_2) = \varphi_{\mathbf{L}}(m_1 \cdot \mathbf{L} m_2)$.

For all of these possibilities it follows that $(u, v) \in \psi(m_1 \cdot_{\mathbf{K}[\mathbf{L}]} m_2)$.

<u>Case 4:</u> $m_1 = a_1$ and $m_2 \in L$. If $m_1 = a_1$, then $\psi(m_1) = E_{\mathbf{K}[\mathbf{L}]}$. Since $\psi(m_2) \subseteq E_{\mathbf{K}[\mathbf{L}]}$ it follows that $\psi(m_1)$; $\psi(m_2) = E_{\mathbf{K}[\mathbf{L}]}$; $\psi(m_2) \subseteq E_{\mathbf{K}[\mathbf{L}]} = \psi(m_1) = \psi(m_1 \cdot_{\mathbf{K}[\mathbf{L}]} m_2)$. For the inclusion in the other direction, let $(u, v) \in E_{\mathbf{K}[\mathbf{L}]}$.

- If $u \in X_{\mathbf{K}[\mathbf{L}]}$ and $v \in [z]X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then $(v,v) \in \psi(m_2)$ and $(u,v) \in E_{\mathbf{K}[\mathbf{L}]}$; $\psi(m_2)$.
- If $u \in X_{\mathbf{K}[\mathbf{L}]}$ and $v \in [z]X_{\mathbf{K}}$ for some $[z] \in X_{\mathbf{L}}/E_{\mathbf{L}}$, then $(v,v) \in \psi(m_2)$ and $(u,v) \in E_{\mathbf{K}[\mathbf{L}]}$; $\psi(m_2)$.
- If $u \in [v]X_{\mathbf{K}}$ and $v \in X_{\mathbf{L}}$, then $(u, u) \in E_{\mathbf{K}[\mathbf{L}]}$ and $(u, v) \in \psi(m_2)$, and so $(u, v) \in E_{\mathbf{K}[\mathbf{L}]}$; $\psi(m_2)$.
- If $u \in {}^{[v]}X_{\mathbf{K}}$ and $v \in X_{\mathbf{L}}$, then there exists $w \in X_{\mathbf{K}}$ such that $u = w^{[v]}$. Thus, $(u, w_{[v]}) \in E_{\mathbf{K}[\mathbf{L}]}$ and $(w_{[v]}, v) \in \psi(m_2)$, and hence $(u, v) \in E_{\mathbf{K}[\mathbf{L}]}$; $\psi(m_2)$.

• If $u, v \in X_{\mathbf{L}}$, then [u] = [v] in $X_{\mathbf{L}}/E_{\mathbf{L}}$. Let $w \in X_{\mathbf{K}} \neq \emptyset$. Thus we have that $(u, w_{[u]}) \in E_{\mathbf{K}[\mathbf{L}]}$ and $(w_{[u]}, v) \in \psi(m_2)$, and hence $(u, v) \in E_{\mathbf{K}[\mathbf{L}]}$; $\psi(m_2)$.

<u>Case 5:</u> If $m_1 \in L$ and $m_2 = a_1$, then the proof that $\psi(m_1 \cdot \mathbf{K}[\mathbf{L}] m_2) = \psi(m_2) = \psi(m_1)$; $\psi(m_2)$ is similar to the proof of the previous case.

Combining Lemmas 4.8, 4.10, 4.11, 4.12 and 4.13 we have shown that the map ψ is a DqRA embedding from $(K \setminus \{a_0\}) \cup L$ into Up $(E_{\mathbf{K}[\mathbf{L}]}, \preccurlyeq_{E_{\mathbf{K}[\mathbf{L}]}})$. Regarding the finiteness of the representations, it is clear from the construction of $\mathbf{X}_{\mathbf{S}_3[\mathbf{L}]}$ that if \mathbf{L} is finitely representable (i.e. $\mathbf{X}_{\mathbf{L}}$ is finite), then $\mathbf{S}_3[\mathbf{L}]$ is finitely representable. This completes the proof of Theorem 4.1.

5 Finite representations of finite Sugihara chains

In this section, we show how the results of Section 4 lead to finite representations of all finite Sugihara chains, and give some examples. A consequence of this is that the result in Theorem 4.1 can be generalised as in Corollary 5.4 which states that if \mathbf{L} is a representable distributive quasi relation algebra, then the generalised ordinal sum of a finite odd Sugihara chain \mathbf{S}_n and \mathbf{L} is representable.

Proposition 2.13 gives us that $\mathbf{S}_3[\mathbf{S}_{n-2}] \approx \mathbf{S}_n$. Using Examples 3.4 and 3.5 and Theorem 4.1 we can apply a simple inductive argument to obtain the result below.

Theorem 5.1. Every finite Sugihara chain is finitely representable.

The construction used in the proof of Theorem 4.1 provides alternative representations to those obtained in [15] and [4] where the representations for \mathbf{S}_n were infinite for $n \ge 4$.

Example 5.2. Let $\mathbf{K} = \mathbf{S}_3$ and $\mathbf{L} = \mathbf{S}_2$. Then $\mathbf{K}[\mathbf{L}] \approx \mathbf{S}_4$ (by Proposition 2.13) and $\mathbf{K}[\mathbf{L}]$ is representable (by Theorem 4.1). Recall that \mathbf{S}_2 is representable over a oneelement poset and \mathbf{S}_3 over a two-element discrete poset with an order automorphism interchanging the elements of the poset. If $X_{\mathbf{K}} = \{x, y\}$ and $X_{\mathbf{L}} = \{u\}$, then \mathbf{S}_4 is representable over $X_{\mathbf{K}[\mathbf{L}]} = \{u, x^{[u]}, y^{[u]}, x_{[u]}, y_{[u]}\}$ with

$$\leq_{X_{\mathbf{K}[\mathbf{L}]}} = \mathrm{id}_{X_{\mathbf{K}[\mathbf{L}]}} \cup \left(\{x_{[u]}, y_{[u]}\} \times \{u\} \right) \cup \left(\{u\} \times \{x^{[u]}, y^{[u]}\} \right) \cup \left(\{x_{[u]}, y_{[u]}\} \times \{x^{[u]}, y^{[u]}\} \right)$$

and $E_{\mathbf{K}[\mathbf{L}]} = X_{\mathbf{K}[\mathbf{L}]}^2$. Here $\alpha_{\mathbf{K}[\mathbf{L}]} = \{(u, u), (x_{[u]}, y_{[u]}), (y_{[u]}, x_{[u]}), (x^{[u]}, y^{[u]}), (y^{[u]}, x^{[u]})\}$ and $\beta_{\mathbf{K}[\mathbf{L}]} = \{(u, u), (x_{[u]}, x^{[u]}), (x^{[u]}, x_{[u]}), (y_{[u]}, y^{[u]}), (y^{[u]}, y_{[u]})\}$. The posets $\mathbf{X}_{\mathbf{K}}, \mathbf{X}_{\mathbf{L}}$ and $\mathbf{X}_{\mathbf{K}[\mathbf{L}]}$ are depicted in Figure 11. It also shows the representations of \mathbf{S}_3 , \mathbf{S}_2 and \mathbf{S}_4 .

Example 5.3. Let $\mathbf{K} = \mathbf{S}_3$ and $\mathbf{L} = \mathbf{S}_3$. Then $\mathbf{K}[\mathbf{L}] \approx \mathbf{S}_5$ (by Proposition 2.13) and $\mathbf{K}[\mathbf{L}]$ is representable (by Theorem 4.1). If $X_{\mathbf{K}} = \{x, y\}$ and $X_{\mathbf{L}} = \{u, v\}$, then \mathbf{S}_5 is




Fig. 11 The posets used to represent S_3, S_2 and $S_3[S_2] \approx S_4$, respectively, and their respective representations.

representable over $X_{\mathbf{K}[\mathbf{L}]} = \{u, v, x^{[u]}, y^{[u]}, x_{[u]}, y_{[u]}\}$ with $E_{\mathbf{K}[\mathbf{L}]} = X_{\mathbf{K}[\mathbf{L}]}^2$,

$$\leq_{X_{\mathbf{K}[\mathbf{L}]}} = \mathrm{id}_{X_{\mathbf{K}[\mathbf{L}]}} \cup \left(\{x_{[u]}, y_{[u]}\} \times \{u, v\} \right) \cup \left(\{u, v\} \times \{x^{[u]}, y^{[u]}\} \right) \cup \left(\{x_{[u]}, y_{[u]}\} \times \{x^{[u]}, y^{[u]}\} \right)$$

 $\begin{array}{ll} \alpha_{\mathbf{K}[\mathbf{L}]} &= \{(u,v),(v,u),(x_{[u]},y_{[u]}),(y_{[u]},x_{[u]}),(x^{[u]},y^{[u]}),(y^{[u]},x^{[u]})\} & and \ \beta_{\mathbf{K}[\mathbf{L}]} &= \{(u,u),(v,v),(x_{[u]},x^{[u]}),(x^{[u]},x_{[u]}),(y_{[u]},y^{[u]}),(y^{[u]},y_{[u]})\}. \ Figure \ 12 \ shows \ the \ posets \ \mathbf{X}_{\mathbf{K}},\mathbf{X}_{\mathbf{L}} \ and \ \mathbf{X}_{\mathbf{K}[\mathbf{L}]}, \ and \ the \ representation \ of \ \mathbf{S}_{5}. \end{array}$

In Figure 13 we show how to extend the posets from Examples 5.2 and 5.3 to give finite posets which can be used to represent S_6 and S_7 .

Lastly, using Proposition 2.13 we get the isomorphic relationship $\mathbf{S}_n[\mathbf{L}] \approx \mathbf{S}_3[\mathbf{S}_{n-2}[\mathbf{L}]]$. This isomorphism and Theorem 5.1 justify the corollary below.




Fig. 12 The poset used to represent $S_3[S_3] \approx S_5$ and its representation.




Fig. 13 Figure showing the posets used for S_4 up to S_7 .

Corollary 5.4. Let $\mathbf{K} = \mathbf{S}_n$ for $n \ge 3$ and n odd. If \mathbf{L} is a (finitely) representable DqRA, then $\mathbf{S}_n[\mathbf{L}]$ is (finitely) representable.

Acknowledgements. The authors are grateful to Peter Jipsen for drawing our attention to the $\mathbf{K}[\mathbf{L}]$ construction for residuated lattices.

References

- Comer, S.D.: Extension of polygroups by polygroups and their representations using color schemes. In: Universal Algebra and Lattice Theory, LNM 1004, Freese et al. (Eds), 91–103 (1983)
- [2] Craig, A., Jipsen, P., Robinson, C.: Frames and spaces for distributive quasi relation algebras and distributive involutive FL-algebras, In: RAMiCS 2024, LNCS 14787, Fahrenburg et al. (Eds.), 156–174 (2024)

- [3] Craig, A., Robinson, C.: Representable distributive quasi relation algebras, Algebra Universalis, accepted. Available at: https://arxiv.org/abs/2310.11719
- [4] Craig, A., Robinson, C.: Representing Sugihara monoids via weakening relations, submitted. Available at: https://arxiv.org/abs/2310.12935
- [5] Galatos, N.: Minimal varieties of residuated lattices, Algebra Universalis 52, 215– 239 (2004)
- [6] Galatos, N., Jipsen, P.: Relation algebras as expanded FL-algebras, Algebra Universalis 69, 1–21 (2013)
- [7] Galatos, N., Jipsen, P.: The structure of generalized BI-algebras and weakening relation algebras, Algebra Universalis 81:35 (2020)
- [8] Galatos, N., Jipsen, P.: Weakening relation algebras and FL²-algebras, In: RAMiCS 2020, LNCS 12062, U. Fahrenberg et al. (Eds.), 117–133 (2020)
- [9] Galatos, N., Jipsen, P., Kowalski, T., Ono, H.: Residuated Lattices: An Algebraic Glimpse at Substructural Logics, Elsevier (2007)
- [10] Hsieh, A., Raftery, J.G.: Semiconic idempotent residuated structures, Algebra Universalis, 61, 413–430 (2009)
- [11] Jipsen, P., Šemrl, J.: Representable and diagonally representable weakening relation algebras, In: RAMiCS 2023, LNCS 13896, Glück et al. (Eds.), 140–157 (2023)
- [12] Jónsson, B., Tarski, A.: Representation problems for relation algebras, Bull. AMS 54, 89t (1948)
- [13] Lyndon, R.C.: The representation of relation algebras, Ann. Math. 51, 707–729 (1950)
- [14] Maddux, R.D.: Relation algebras, Studies in Logic and the Foundations of Mathematics, Elsevier (2006)
- [15] Maddux, R.D.: Relevance logic and the calculus of relations, Rev. Symb. Log. 3, 41–70 (2010)
- [16] McCune, W.: Prover9 and Mace4, http://www.cs.unm.edu/~mccune/Prover9, 2005-2010.
- [17] Tarski, A.: On the calculus of binary relations, J. Symb. Logic 6, 73–89 (1941)