# Mass spectroscopy of charmonium using a screened potential

Sreelakshmi M $\,\cdot\,$  Akhilesh Ranjan

Received: date / Accepted: date

Abstract In this work, we estimate the mass spectra and decay properties of charmonium  $(c\bar{c})$  using a non-relativistic potential model. We employ a potential model incorporating a Coulomb like term, representing one gluon exchange at short distances, and a screening term representing quark confinement at long distances. Spin-dependent corrections are also added perturbatively. Our results are compared with available experimental data and some other theoretical models. Based on this, we have made some comments on interquark potential.

Keywords Quarks  $\cdot$  QCD  $\cdot$  Hadrons  $\cdot$  Charmonium

## 1 Introduction

There are various experimental facilities across the world, like LHCb [1], Belle [2], CLEO [3], and BaBar [4] etc that investigate the properties of hadrons. Hadrons are composed of quarks and antiquarks and follow the principles of Quantum Chromodynamics (QCD). QCD is the theory of the strong interaction. Studying the properties of hadrons provides deep insight into the strong interaction. Gluons are the mediators of the strong interaction between quarks and also carry color charges.

One unique feature of strong interaction is asymptotic freedom. According to this, the effective coupling constant ( $\alpha_s = g_s/4\pi$ ) decreases logarithmically at short distances, which is also well supported theoretically [5]. Another

Sreelakshmi M

Department of Physics, Manipal Institute of Technology

A. Ranjan

Department of Physics, Manipal Institute of Technology Manipal Academy of Higher Education, Manipal, 576104, Karnataka, India E-mail: ak.ranjan@manipal.edu

Manipal Academy of Higher Education, Manipal, 576104, Karnataka, India E-mail: sreelakshmim.araam@gmail.com

special feature of strong interaction is color confinement. According to color confinement, only color singlet hadrons can freely exist in nature. Color confinement property [6], is supported by lattice simulations but not yet theoretically proven. Any strong interaction theory should incorporate these two features. There are many theoretical methods used in hadron spectroscopy. Some of the highly used methods are the bag model [7][8], QCD sum rules [9], Heavy Quark Effective Theory (HQET) [10], phenomenological potential models [11][12][13], Lattice QCD (LQCD) including lattice gauge theories [14], etc.

Phenomenological potential models have been highly successful in describing the properties of hadrons, including both light and heavy quark systems. These models are inspired by the characteristics of QCD, confinement, and asymptotic freedom. For interactions at shorter distances, one gluon exchange (OGE) dominates, while at large interquark distances, QCD perturbation theory breaks down, and quarks become confined. The Cornell potential is a widely used phenomenological potential that combines a Coulomb-like term to account for one gluon exchange and a linear term to describe quark confinement [11][15]. These models often provide remarkably accurate descriptions of experimental data, making them a significant method for investigating the properties of hadrons. In the case of heavy hadrons, the potential can be treated non-relativistically and solved using the Schrödinger equation. Therefore, in this work, we have used a phenomenological potential model.

The study of quarkonium  $(q\bar{q})$  has been very helpful in understanding the strong interaction after the discovery of  $J/\psi$  [16] at SLAC. Quarkonium is relatively simple to study, as it consists of one quark and its antiquark, which is a two-body system. We are finding many interesting experimental results on exotic multiquark states these days [17][18]. The studies on quarkonium can be the base for understanding such exotic multiquark states like tetraquarks and pentaquarks. This also makes the study of quarkonium systems important.

In this paper, we have estimated the mass and decay properties of charmonium using a non-relativistic screened potential. The paper is organized as follows: a brief introduction in Section 1 is followed by the theoretical formulation of the problem in Section 2. Discussion of the results in Section 3 is followed by conclusions in Section 4.

### **2** Theoretical Formulation

The QCD is a non-abelian and complex theory. Therefore, the real nature of the interquark interaction is still unknown. The QCD vacuum contains virtual quark-antiquark pairs and gluons. A quark inside a hadron attracts an opposite color charge, and a cloud of virtual particles builds up around the quark. This, in turn, leads to a screening effect on other quarks at a distance. Therefore, it is worth using screened potential to study quark interactions. We have used a non-relativistic screened potential for the analysis of quarkonia. A screened potential  $(V_{NR}(r))$  is a combination of the Coulomb term and screening term [19]. Apart from the non-relativistic contribution  $(V_{NR}(r))$ , interaction potential includes spin-dependent correction  $(V_{SD}(r))$ , which is added perturbatively to the Hamiltonian,

$$V(r) = V_{NR}(r) + V_{SD}(r).$$
 (1)

The form of screened potential can be written as [20],

$$V_{NR}(r) = \frac{k_s \alpha_s}{r} + b \left(\frac{1 - \exp^{(-\mu_s r)}}{\mu_s}\right) - c, \qquad (2)$$

where b and c are potential parameters,  $\alpha_s$  is the strong coupling constant and  $\mu_s$  is the parameter for screening effect. The parameter  $k_s$  represents the color factor and denotes a color singlet state. From SU(3) representation,  $3 \otimes \overline{3} = 1 \oplus 8$ , leads to  $-\frac{4}{3}$  for a color singlet state.

A screened potential introduces light quark-antiquark pairs in the vacuum, which softens the interaction at large distances compared to the linear potential. At short distances,  $r \leq 1 fm$ , the screened potential behaves identically to a linear potential. Screening lowers the mass of higher excited states [20].

In the case of heavy hadrons, the kinetic energy of the quarks is less compared to their rest mass energy. Therefore, a non-relativistic approach can be used. The time-independent radial Schrödinger equation is,

$$\left[\frac{1}{2\mu}\left(-\frac{d^2}{dr^2}\right) + \frac{l(l+1)}{r^2} + V(r)\right]y(r) = Ey(r),\tag{3}$$

parameter  $\mu$  is the reduced mass,  $\mu = \frac{m_1 m_2}{m_1 + m_2}$ . The Schrödinger equation is solved by numerical integration with the Runge-Kutta method using Mathematica package [21]. We have fitted the center of weight mass or spin average mass  $(M_{SA,nJ})$  of the ground state with the known experimental values (from Particle Data Group [22]) of pseudoscalar and vector mesons,

$$M_{SA_{,nJ}} = \frac{\sum_{J} (2J+1)M_{nJ}}{\sum_{J} (2J+1)}.$$
(4)

The constant parameters used for the calculations are given in Table 1. Spindependent terms will be considered perturbatively. The correction contains spin-spin, spin-orbit, and tensor interactions. The spin-spin interaction creates hyperfine levels, whereas spin-orbit and tensor interactions are responsible for the fine structure of the quarkonium. The coefficients of these spin-dependent terms can be expressed as [19],

$$V_{SD}(r) = V_{SS}(r) \left[ S(S+1) - \frac{3}{2} \right] + V_{L \cdot S}(r) (\mathbf{L} \cdot \mathbf{S}) + V_T(r) \left[ S(S+1) - 3 \frac{(\mathbf{S} \cdot \mathbf{r})(\mathbf{S} \cdot \mathbf{r})}{r^2} \right],$$
(5)

where S is the total spin and L is total orbital angular momentum. These terms are the result of Lorentz structure  $(\Gamma_{\Sigma} \otimes \Gamma_{\Sigma})$  of inter-quark interaction. Various Lorentz structures of a fermion-anti fermion interaction will give rise to scalar  $(V_s(r))$ , pseudo scalar (zero), vector  $(V_v(r))$ , axial vector  $(V_A(r))$  and tensor  $(V_T(r))$  terms in a non-relativistic potential [23]. Therefore, the effective potential should be considered as the sum of vector and scalar terms,

$$V(r) = V_v(r) + V_s(r).$$
 (6)

The spin-spin term for equal quark masses is given by,

$$V_{SS}(r) = \frac{32\pi\alpha_s}{9m_Q^2} |\psi(0)|^2 \langle \mathbf{S_1} \cdot \mathbf{S_2} \rangle, \tag{7}$$

where  $m_Q$  is heavy quark mass and  $S_1$  and  $S_2$  are the spin of the heavy quark and heavy antiquark, respectively, with,

$$\langle \mathbf{S_1} \cdot \mathbf{S_2} \rangle = \begin{cases} -\frac{3}{4} & S = 0, \text{spin singlets} \\ \frac{1}{4} & S = 1, \text{spin triplets.} \end{cases}$$
(8)

The spin-orbit term is given by,

$$V_{L\cdot S}(r) = \frac{\mathbf{L} \cdot \mathbf{S}}{(2m_Q)^2 r} \left[ \frac{1}{r} \frac{dV_v(r)}{dr} - \frac{d^2 V_v(r)}{dr^2} \right],\tag{9}$$

where L and S are the total orbital angular momentum and total spin of the heavy quark and heavy antiquark, respectively, with,

$$\langle L \cdot S \rangle = \begin{cases} l, & J = l+1 \\ -1, & J = l \\ -(l+1), & J = l-1. \end{cases}$$
(10)

The tensor term is given by,

$$V_T(r) = \frac{\mathbf{S_{12}}}{(12m_Q)^2} \left[ 3\frac{dV_v(r)}{dr} - \frac{dV_s(r)}{dr} \right],$$
(11)

$$\langle S_{12} \rangle_{\frac{1}{2} \otimes \frac{1}{2} \to S=1, l \neq 0} = \begin{cases} -\frac{2l}{2l+3}, & J = l+1\\ 2, & J = l\\ -\frac{2(l+1)}{2l-1}, & J = l-1. \end{cases}$$
(12)

Value of  $\langle S_{12} \rangle$  vanishes when l = 0 and S = 0. Moreover, these values are valid for the spin half particles.

## 2.1 Decay width

In addition to mass spectra, predictions of decay width are also important in hadron spectroscopy. Annihilation decays are very helpful in identifying conventional mesons and multiquark structures [24][25]. Quarkonium annihilation into light particles helps to learn about the strong fine structure constant and its role in understanding the quark forces.

## 2.1.1 Leptonic decays

 ${}^{3}S_{1}$  and  ${}^{3}D_{1}$  states of a quarkonium can decay into a pair of leptons through a virtual photon. The leptonic decay rate can be estimated from the conventional Van-Royen-Weisskopf formula [26]. This formula with first-order radiative corrections is given by [24][27],

$$\Gamma(n^3 S_1 \to e^+ e^-) = \frac{4q^4 \alpha^2 |R_{nS}(0)|^2}{M_{nS}^2} \left(1 - \frac{16\alpha_s}{3\pi}\right).$$
(13)

Similarly,

$$\Gamma(n^3 D_1 \to e^+ e^-) = \frac{25q^2 \alpha^2 |R_{nD}^{''}(0)|^2}{2m_Q^4 M_{nD}^2} \left(1 - \frac{16\alpha_s}{3\pi}\right).$$
 (14)

Here, q is the Coulomb charge of the quark  $(e_c = \frac{2e}{3})$ ,  $\alpha$  is the fine structure constant  $(\alpha = \frac{1}{137})$ ,  $\alpha_s$   $(\alpha_s(c\bar{c}) = 0.318)$  is strong coupling constant,  $m_Q$  denotes the mass of quark  $(m_c = 1.32GeV)$ .  $M_{nS}$  and  $M_{nD}$  are masses of decaying quarkonia states.  $R_{nS}(0)$  and  $R_{nD}(0)$  are radial wave functions at zero for S and D states.  $R_{nD}''(0)$  is the second derivative of the radial wave function at zero.

## 2.1.2 Two photon decay

The decay of different charmonium states into photons is given by [24],

$$\Gamma(n^{1}S_{0} \to \gamma\gamma) = \frac{3q^{4}\alpha^{2}|R_{nS}(0)|^{2}}{m_{Q}^{2}} \left(1 - \frac{3.4\alpha_{s}}{\pi}\right).$$
 (15)

$$\Gamma(n^3 S_1 \to \gamma \gamma \gamma) = \frac{4(\pi^2 - 9)q^6 \alpha^3 |R_{nS}(0)|^2}{3\pi m_Q^2} \left(1 - \frac{12.6\alpha_s}{\pi}\right).$$
(16)

$$\Gamma(n^{3}P_{0} \to \gamma\gamma) = \frac{27q^{4}\alpha^{2}|R_{nP}^{'}(0)|^{2}}{m_{Q}^{4}} \left(1 + \frac{0.2\alpha_{s}}{\pi}\right).$$
 (17)

$$\Gamma(n^{3}P_{2} \to \gamma\gamma) = \frac{36q^{4}\alpha^{2}|R_{nP}^{'}(0)|^{2}}{5m_{Q}^{4}} \left(1 - \frac{16\alpha_{s}}{3\pi}\right).$$
 (18)

Here,  $R'_{nP}(0)$  is the first derivative of the radial wave function at zero. All these formulae include QCD first-order radiative corrections.

## 2.1.3 Two gluon decay

Experimental detection of di-gluon annihilation decay faces challenges because the gluonic state breaks down into multiple hadrons, which makes direct measurement difficult and first principle approximations unreliable. The decay width of a pseudoscalar meson via di-gluon annihilation, including the leading order QCD radiative correction, is given by [24][28],

$$\Gamma(n^{3}P_{0} \to gg) = \frac{6\alpha_{s}^{2}|R_{nP}^{'}(0)|^{2}}{m_{Q}^{4}}\left(1 + \frac{9.5\alpha_{s}}{3\pi}\right).$$
(19)

$$\Gamma(n^{3}P_{2} \to gg) = \frac{8\alpha_{s}^{2}|R_{nP}^{'}(0)|^{2}}{5m_{Q}^{4}} \left(1 - \frac{2.2\alpha_{s}}{3\pi}\right).$$
(20)

#### **3** Results and Discussions

The mass spectra of quarkonia  $(c\bar{c})$  can be calculated using the formula,

$$M_{c\bar{c}} = 2m_c + E_{nl} + E_{spin}.$$
(21)

The parameters used in the calculation to get charmonium masses are given in Table 1.

#### Table 1 Parameters used in the calculation.

Parameters	$\alpha_s$ [25]	b~(GeV)	c~(GeV)	$\mu (GeV)$	$m_q \ (GeV)$
$c\bar{c}$	0.318	0.150	1.00	0.03	1.32

#### 3.1 Mass spectra of $c\bar{c}$

Mass spectra for charmonium states are calculated and shown in Table 2, Table 3 and Table 4. The results are compared with different theoretical results and available experimental results [22]. Here, Chaturvedi and Rai [25][29] Sultan *et al*, [30] and Soni *et al*, [31] used Cornell potential plus spin-dependent corrections. Ebert, Faustov, and Galkin [32] used a relativistic quark model based on the quasipotential approach, and Kalinowski and Wagner [33] used the LQCD method. We have calculated the mass spectra of different states of charmonium, including S, P, and D states. The spectroscopic notation for the states can be represented as  $n^{2S+1}L_J$  (symbols have the usual meaning).

State	Present	PDG [22]	[25]	[29]	[30]	[31]	[32]	[33]
$1^{1}S_{0}$	2.976	$2.9839 \pm 0.0004$	3.004	2.989	2.982	2.989	2.981	2.884
$1^{3}S_{1}$	3.082	$3.096900 \pm 0.000006$	3.086	3.094	3.090	3.094	3.096	3.056
$2^{1}S_{0}$	3.550	$3.6375 \pm 0.0011$	3.645	3.572	3.630	3.602	3.630	3.535
$2^{3}S_{1}$	3.642	$3.68610 \pm 0.00006$	3.708	3.649	3.672	3.681	3.672	3.662
$3^{1}S_{0}$	4.027		4.124	3.998	4.058	-	4.043	
$3^{3}S_{1}$	4.111		4.147	4.062	4.129	-	4.072	
$4^{1}S_{0}$	4.445		4.534	4.372	4.384	4.448	4.384	
$4^{3}S_{1}$	4.523		4.579	4.428	4.406	4.514	4.406	

 ${\bf Table \ 2} \ {\rm Mass \ spectra \ of \ charmonium \ } (S\mbox{-} {\rm states}) \ {\rm in \ } GeV.$ 

Table 3 Mass spectra of charmonium (P- states) in GeV.

State	Present	PDG [22]	[25]	[29]	[30]	[31]	[32]	[33]
$1^{3}P_{0}$	3.435	$3.4147 \pm 0.0030$	3.440	3.473	3.424	3.428	3.413	3.421
$1^{3}P_{1}$	3.479	$3.51067 \pm 0.00005$	3.492	3.506	3.505	3.468	3.511	3.480
$1^{3}P_{2}$	3.498	$3.55617 \pm 0.00007$	3.511	3.551	3.549	3.480	3.555	3.536
$2^{3}P_{0}$	3.852		3.932	3.918	3.852	3.897	3.870	
$2^{3}P_{1}$	3.896		3.932	3.949	3.925	3.938	3.870	
$2^{3}P_{2}$	3.919	$3.9225 \pm 0.0010$	4.007	4.002	3.965	3.943	3.949	4.066
$3^{3}P_{0}$	4.283		4.394	4.306	4.202	4.296	4.301	
$3^{3}P_{1}$	4.326		4.401	4.336	4.271	4.338	4.319	
$3^{3}P_{2}$	4.350		4.427	4.392	4.309	4.358	4.350	

Table 4 Mass spectra of charmonium (D- states) in GeV.

State	Present	PDG [22]	[25]	[29]	[30]	[31]	[32]
$1^{3}D_{3}$	3.706		3.789	3.806	3.805	3.775	3.813
$1^{3}D_{2}$	3.711	$3.8237 \pm 0.0005$	3.814	3.800	3.795	3.772	3.759
$1^{3}D_{1}$	3.705	$3.7737 \pm 0.0004$	3.815	3.785	3.783	3.775	3.783
$2^{3}D_{3}$	4.153		4.273	4.206	4.165	4.176	4.220
$2^{3}D_{2}$	4.160		4.248	4.203	4.158	4.188	4.190
$2^{3}D_{1}$	4.158	$4.1910 \pm 0.0050$	4.245	4.196	4.141	4.182	4.105
$3^{3}D_{3}$	4.551		4.626	4.568	4.481	4.549	4.574
$3^{3}D_{2}$	4.559		4.632	4.566	4.472	4.557	4.554
$3^{3}D_{1}$	4.560		4.627	4.562	4.455	4.553	4.507

Table 2, Table 3, and Table 4 show reasonable agreement with experimental data, while small deviations start to appear at increasing energy levels. Comparisons with other models, such as those from [25], [30], and [32], indicate overall consistency, though differences are more pronounced for the 3S and 4S states. Our results show better matching compared to other studies for  $2^{3}P_{2}$  state in Table 3, whereas other studies overestimated the results. From Table 4, analysis of the  $1^{3}D_{1}$ ,  $1^{3}D_{2}$  and  $1^{3}D_{3}$  states show correspondence between calculated results and theoretical expectations while showing small inconsistencies across different models. Experimental data from the  $1^{3}D_{2}$  and  $1^{3}D_{1}$ 

states show largest deviation with our findings compared to other models. In potential plots shown in Fig. 1, the width of our proposed potential is more than the width of other potentials. Therefore, the energy eigenvalues of our proposed potential will be lowest among all these potentials. This may be the reason for the highest deviation among the compared ones.

**Table 5** Mass splitting for  $c\bar{c}$  (*MeV*).

State	Present	PDG [22]	[25]	[29]	[30]	[31]	[32]	[33]
$m_{J/\psi} - m_{\eta_c}(1S)$	106	$113 \pm 0.7 \pm 0.1$	82	105	108	105	115	172
$\overline{m_{\psi}(2S) - m_{J/\psi}(1S)}$	518	$589.188 {\pm} 0.028$	704	660	690	692	691	778

Fig. 1 Comparison of different potentials (without spin-dependent terms).



From Fig. 1, it is clear that all these potentials have similar well-type structures. Therefore, their mass spectra will also show a similar nature. The energy gap among two consecutive energy levels decreases as we go for higher excited states because, for higher energy states, the width of the potential well gradually increases. Since all these potentials contain spin interaction terms, the energy gap among two consecutive energy levels does not maintain this behavior. This may be because of the significant role of the spin correction term in the mass spectrum of charmonium. From Tables 2 to 4, this nature of the charmonium spectrum can be seen clearly. Especially 1D state data in Table 4 for potential [31]. We have limited experimental data on higher states. This highlights the need for further experimental efforts to establish

the masses of higher states, which would help in the refinement and validation of theoretical models of charmonium spectroscopy.

In the case of S states, there is only spin-spin interaction, while P and D states have spin-orbit and tensor interactions. The mass splitting for different states of the same principal quantum number n and orbital angular momentum l is much smaller, especially for P and D states, indicating that these corrections are perturbative. The experimental study suggests  $c\bar{c}$  [22],  $m_{J/\psi} - m_{\eta_c}(1S) = 113 \pm 0.7 \pm 0.1 \ MeV$ . Similarly,  $m_{\psi}(2S) - m_{J/\psi}(1S) =$ 589.188  $\pm 0.028 \ MeV$ . The difference has been calculated and compared with experimental and other studies and given in Table 5. This way we find that overall our results show good agreement with the experimental results. Hence we can say that the screened potential is very close to the interquark potential.

#### 3.2 Decay width

Different types of annihilation decay of charmonium states were calculated and compared with other studies. Experimental results are available for only a few decay modes. Calculated decay widths are shown in Table 6 to Table 9. The results are compared with other studies. The constants, including quark masses, strong coupling constant, etc, are taken from the study of mass spectra, as mentioned earlier.

3.2.1 Leptonic decay  $(nS_1^3 \rightarrow e^+e^-)$ 

**Table 6** Leptonic decay width of S states of  $c\bar{c}$  (keV).

State	Present	PDG [22]	[25]	[29]	[31]	[34]
$1S_{1}^{3}$	1.085	$5.55{\pm}0.08$	1.957	6.932	2.925	5.47
$2S_{1}^{3}$	0.967	$2.33 {\pm} 0.01$	1.178	3.727	1.533	2.14
$3S_{1}^{3}$	0.529	$0.86 {\pm} 0.01$	0.969	2.994	1.091	0.796
$4S_{1}^{3}$	0.422		0.860	2.638	0.856	0.288

For leptonic decay, experimental results are available for 1S-3S states. When the results for different leptonic decays are compared, it is observed that as the decays proceed to higher excited states, the decay widths become closer to experimental values. Decay rates given by Vinodkumar *et al* [34] have shown good agreement with the experimental results. Vinodkumar used a relativistic model with harmonic oscillator potential. On the other hand, the decay rate given by Chaturvedi and Rai [25] and Soni *et al* [31] and our present calculation show good agreement with each other but a large difference from the experimental results.

Leptonic decay of charmonia can occur due to Quantum Electrodynamics (QED), weak interactions, or strong interactions. In charmonia, the quark

annihilation may be due to all three interactions, but the lepton production will be due to QED and weak interactions only. Therefore, in charmonium decay, the QCD corrections become relevant to account for the interactions of quarks within hadrons. Chaturvedi and Rai [25] employed the NRQCD factorization expressions for various leading orders of decay. In another study, Chaturvedi and Rai [29] applied radiative and quark propagator corrections, which appear to overestimate the values. The present study and study by Soni and others [31] used a similar type of QCD corrections, but the parameters were different and showed different results. Brodsky  $et \ al$  have shown that the running coupling constant can affect the perturbative expansions in QCD [27]. First-order corrections are comparatively small for most processes when the appropriate scale for the coupling constant is used. Our calculation shows a large difference from the experimental data. It indicates that in this formalism, we may have either missed some processes or some correction factors. Again, if we compare it with Vinodkumar's [34] result, we find that relativistic model corrections are more successful than non-relativistic corrections for leptonic decays.

## 3.2.2 Photonic decay $(nS_0^1/nP_0^3/nP_2^3 \rightarrow \gamma\gamma)$

State	Present	PDG [22]	[25]	[29]	[31]	[35]	[36]	[37]
$1S_{0}^{1}$	6.289		6.725	8.246	5.618	7.18		
$2S_{0}^{1}$	5.839		3.178	4.560	2.944	1.71		
$3S_{0}^{1}$	5.596		1.493	3.737	2.095	1.21		
$4S_{0}^{1}$	5.437		0.858	3.340	1.644			
$1P_0^3$	8.752	$2.341 {\pm} 0.189$	4.185	2.692			6.38	7.33
$2P_0^3$	11.261		4.306	4.716				8.70
$3P_0^3$	13.422		4.847	8.078				
$1P_{2}^{3}$	1.095		0.538	1.242			0.57	1.95
$2P_{2}^{3}$	1.409		0.554	1.485				2.32
$3P_{2}^{3}$	1.923		0.626	1.691				

**Table 7** Photonic decay width for  $c\bar{c}$  (keV).

For the di-photonic decay of charmonium, we have only one data point available. Our results show good agreement with Cornell potential [25][29]. However, with experimental data, a big variation is present. Here, we also find that there are huge differences among relativistic and non-relativistic data.

State	Present	[25]	[29]
$1S_{0}^{1}$	1.046	1.022	2.997
$2S_{0}^{1}$	0.972	0.900	1.083
$3S_{0}^{1}$	0.931	0.857	1.046
$4S_0^1$	0.905	0.832	0.487

**Table 8** Tri-photonic decay width of S states of  $c\bar{c} \ (eV) \ (nS_0^1 \rightarrow \gamma\gamma\gamma)$ .

In the tri-photonic decay of charmonium, our results show good agreement with [25]. We do not have experimental data for this process, therefore, it will be too early to comment on it.

3.2.3 Gluonic decay  $(nP_0^3/nP_2^3 \rightarrow gg)$ 

**Table 9** Gluonic decay width of  $c\bar{c}$  states (*MeV*).

State	Present	[29]	[36]	[37]	[38]
$1P_0^3$	25.18	14.19	13.44	32.58	3.337
$2P_0^3$	32.40	24.973		38.70	2.060
$3P_0^3$	38.61	33.876			
$1P_{2}^{3}$	4.708	2.914	1.2	3.38	0.784
$2P_{2}^{3}$	6.059	5.099		4.01	0.504
$3P_{2}^{3}$	7.221	6.867			

In gluonic decay, experimental detection is a big challenge. Unfortunately, we do not have experimental data for this decay mode. Our data shows good agreement with [29], where the authors have used the Cornell potential. Our results show a large difference with [38] and [36] results. They have used relativistic models. In this way, we find a significant difference between the results of relativistic and non-relativistic potentials as observed in the case of photonic decay.

## 4 Conclusions

In the present work, we have found that the mass spectra of charmonium using a non-relativistic screened potential model with spin-dependent corrections show good agreement with the experiment and other non-relativistic models. For relativistic or semi-relativistic models, mass spectra show differences. Comparative study on different potential models, lattice QCD, relativistic, and experimental results indicates that the proposed screened potential with spindependent elements is very close to the interquark potential (the exact form is still unknown). On the other hand, for decay spectra, relativistic models show good agreement with experimental data compared to the non-relativistic models. However, decay spectra show underestimated results with non-relativistic models, which indicates that some processes may have been missed. It needs a separate and detailed analysis which is beyond the scope of this work.

#### Acknowledgements

SM is thankful to Manipal Academy of Higher Education (MAHE) for the financial support under Dr. T. M. A. Pai scholarship program.

#### References

- et al, (LHCb Collaboration), Observation of the Resonant 1. Aaij, R. Character of the  $Z(4430)^{-}$ State. Phys. Rev. Lett. 112222002 (2014).DOI:10.1103/PhysRevLett.112.222002
- 2. Choi, S.K. et al (Belle Collaboration), Observation of a Narrow Charmonium like State in Exclusive  $B^{\pm} \rightarrow K^{\pm}\pi^{+}\pi^{-}J/\psi$  Decays. Phys. Rev. Lett. **91** 262001 (2003). DOI:10.1103/PhysRevLett.91.262001
- 3. Asner, D.M. *et al*, Observation of  $\eta_c$  Production in  $\gamma\gamma$  Fusion at CLEO. *Phys. Rev. Lett.* **92** 142001 (2004). DOI:10.1103/PhysRevLett.92.142001
- 4. Lees, J.P. *et al* (BaBar Collaboration), Study of  $B^{\pm,0} \rightarrow J/\psi K^+ K^- K^{\pm,0}$ and Search for  $B^0 \rightarrow J/\psi \phi$  at BaBar. *Phys. Rev. D.* **91** 012003 (2015). DOI:10.1103/PhysRevD.91.012003
- Gross, D.J., Nobel Lecture: The Discovery of Asymptotic Freedom and the Emergence of QCD. Rev. Mod. Phys. 77 837–849 (2005). DOI:10.1103/RevModPhys.77.837
- Wilson, K.G., Confinement of Quarks. Phys. Rev. D. 10 2445–2459 (1974). DOI:10.1103/PhysRevD.10.2445
- Chodos, A., Jaffe, R.L., Johnson, K., Thorn, C.B., Weisskopf, V.F., New Extended Model of Hadrons. *Phys. Rev. D.* 9 3471–3495 (1974). DOI:10.1103/PhysRevD.9.3471
- Chodos, A., Jaffe, R.L., Johnson, K., Thorn, C.B., Baryon Structure in the Bag Theory. *Phys. Rev. D.* **10** 2599–2604 (1974). DOI:10.1103/PhysRevD.10.2599
- Shifman, M.A., Vainshtein, A.I., Zakharov, V.I., QCD and Resonance Physics. Theoretical Foundations. Nucl. Phys. B 147 385–447 (1979). DOI:10.1016/0550-3213(79)90022-1
- Körner, J.G., Thompson, G., The Heavy Mass Limit in Field Theory and The Heavy Quark Effective Theory. *Phys. Rev. Lett. B* 264 185–192 (1991). DOI:10.1016/0370-2693(91)90725-6
- Eichten, E., Gottfried, K., Kinoshita, T., Kogut, J., Lane, K. D., Yan, T. M., Spectrum of Charmed Quark-Antiquark Bound States. *Phys. Rev. Lett.* **34** 369–372 (1975). DOI:10.1103/PhysRevLett.34.369
- Martin, A., A Simultaneous Fit of b<sub>0</sub>, cc̄,ss̄ (bcs Pairs) and cs̄ Spectra. Phys. Lett. B 100 511–514 (1981). DOI:10.1016/0370-2693(81)90617-1
- Bhanot, G., Rudaz, S., A New Potential for Quarkonium. Phys. Lett. B 78 119–124 (1978). DOI:10.1016/0370-2693(78)90362-3
- Kogut, J.B., The Lattice Gauge Theory Approach to Quantum Chromodynamics. *Rev. Mod. Phys.* 55 775–836 (1983). DOI:10.1103/RevModPhys.55.775
- 15. Kuchin, S.M., Maksimenko, N.V., Theoretical Estimations of the Spin–Averaged Mass Spectra of Heavy Quarkonia and  $B_c$  Mesons. UJPA 7 295–298 (2013). DOI: 10.13189/ujpa.2013.010310
- 16. Augustin, J. E. et al, Discovery of a Narrow Resonance in  $e^+e^-$  Annihilation. Phys. Rev. Lett. **33** 1406–1408 (1974). DOI:10.1103/PhysRevLett.33.1406
- 17. Abazov, V.M. et al (D0 Collaboration), Inclusive Production of the X(4140) State in  $pp^-$  Collisions at D0. Phys. Rev. Lett. **115** 232001 (2015). DOI:10.1103/PhysRevLett.115.232001
- 18. Aaltonen, T. (CDF Collaboration), Observation of the Y(4140) structure in the  $J/\psi\phi$  mass spectrum in  $B^{\pm} \rightarrow J/\psi\phi K^{\pm}$  decays. Mod. Phys. Lett. A. **32** 1750139 (2017). DOI:10.1142/S0217732317501395

- Lucha, W., Bound States of Quarks. Phys. Rep. 200 127–240 (1991). DOI:10.1016/0370-1573(91)90001-3
- Patel, V., Gandhi, K., Rai, A.K., Interpreting Charm-Strange Mesons with a Screened Potential Model. *Few-Body Syst.* 62, 68 (2021). DOI:10.1007/s00601-021-01651-y
- 21. Lucha, W., Schöberl, F.F., Solving the Schrödinger Equation for Bound States with Mathematica 3.0. Int. Jn. Mod. Phys. C 10 607-19 (1998). DOI:10.1142/S0129183199000450
- 22. S. Navas *et al.* (Particle Data Group), *Phys. Rev. D* **110** 030001 (2024). DOI:10.1093/ptep/ptac097
- Lucha, W., Schöberl, F.F., Effective Potential Models for Hadrons, *HEPHY-PUB-621-95, UWTHPH-1995-16* (1996). DOI:arxiv.org/abs/hep-ph/9601263
- Kwong, W., Mackenzie, P.B., Rosenfeld, R., Rosner, J.L., Quarkonium Annihilation Rates. Phys. Rev. D. 37 3210–3215 (1988). DOI:10.1103/PhysRevD.37.3210
- Chaturvedi, R., Rai, A.K., Charmonium Spectroscopy Motivated by General Features of pNRQCD. Int. J. Theor. Phys. 59 3508–32 (2019). DOI:10.1007/s10773-020-04613-y
- Van Royen, R., Weisskopf, V.F., Hardon Decay Processes and the Quark Model. Nuov Cim A. 50 617–645 (1967). DOI:10.1007/BF02823542
- Brodsky, S.J., Lepage, G.P., Mackenzie, P.B., On the Elimination of Scale Ambiguities in Perturbative Quantum Chromodynamics. *Phys. Rev. D.* 28 228–235 (1983). DOI:10.1103/PhysRevD.28.228
- 28. Lansberg, J.P., Pham, T.N., Effective Lagrangian for Two-Photon and Two-Gluon Decays of P- Wave Heavy Quarkonium  $\chi_{c0,2}$  and  $\chi_{b0,2}$  States. Phys. Rev. D. **79** 094016 (2009). DOI:10.1103/PhysRevD.79.094016
- 29. Chaturvedi, R., Kumar Rai, A., Mass Spectra and Decay Properties of the  $c\bar{c}$  Meson. Eur. Phys. J. Plus. **133** 220 (2018). DOI:10.1140/epjp/i2018-12044-8
- 30. Sultan, M.A., Akbar, N., Masud, B., Akram, F., Higher Hybrid Charmonia in an Extended Potential Model. *Phys. Rev. D.* **90** 054001 (2014). DOI:10.1103/PhysRevD.90.054001
- 31. Soni, N.R., Joshi, B.R., Shah, R.P., Chauhan, H.R., Pandya, J.N.,  $Q\bar{Q} \ (Q \in \{b,c\})$  Spectroscopy Using the Cornell Potential. *Eur. Phys. J. C.* **78** 592 (2018). DOI:10.1140/epjc/s10052-018-6068-6
- Ebert, D., Faustov, R.N., Galkin, V.O., Spectroscopy and Regge Trajectories of Heavy Quarkonia and B<sub>c</sub> Mesons. Eur. Phys. J. C. **71** 1825 (2011). DOI:10.1140/epjc/s10052-011-1825-9
- 33. Kalinowski, M., Wagner, M., Masses of D mesons, D<sub>s</sub> Mesons, and Charmonium states from Twisted-Mass Lattice QCD. Phys. Rev. D. 92 094508 (2015). DOI:10.1103/PhysRevD.92.094508
- 34. Vinodkumar, P.C., Pandya, J.N., Bannur, V.M., Khadkikar, S.B., A Unified Scheme for Flavoured Mesons and Baryons. *Eur. J. Phys. A.* 4 83–90 (1999). DOI:10.1007/s100500050206
- Lakhina, O., Swanson, E.S., Dynamic Properties of Charmonium. Phys. Rev. D. 74 014012 (2006). DOI:10.1103/PhysRevD.74.014012
- 36. Gupta, S.N., Johnson, J.M., Repko, W.W., Relativistic Two-Photon and Two-Gluon Decay Rates of Heavy Quarkonia. *Phys. Rev. D.* 54 2075–2080 (1996). DOI:10.1103/PhysRevD.54.2075
- B. K. Patel, Spectroscopy and Decay Properties of Heavy Flavour Hadrons. PhD thesis, Sardar Patel University, 2009. hdl.handle.net/10603/7354
- Bhaghyesh, Vijaya Kumar, K.B., Charmonium Spectra and Decays in a Semirelativistic Model. Int. J. Mod. Phys. A. 27 1250127 (2012). DOI:10.1142/S0217751X12501278