Anomaly Equation of the Large U(1) Chiral Symmetry

Shingo Takeuchi

Faculty of Environmental and Natural Sciences, Duy Tan University, Da Nang, Vietnam

Abstract

In this study, we first heuristically constitute the charges for the chiral transformation associated with the large U(1) gauge symmetry. We name those as the large chiral charges, and the chiral transformation those generate as the large chiral transformations. Then, showing that those can be obtained based on Noether's theorem, we once obtain the anomaly equation associated with those large chiral transformations.

Then, for the one-loop diagrams of the fermionic field coupling to the multiple classical gauge fields (those constitute the effective action of the model in this study with regard to the gauge field), we perform an axailization. Then, defining the BRS transformations for the large U(1)gauge symmetry (we name those as the large BRS transformation), we perform those large BRS transformations to those axialized one-loop diagrams. Then, evaluating those, we show that the anomaly equations mentioned above can be derived. It is also shown that those anomaly equations can be derived in Fujikawa method. The result in this study would be important as a development of the large U(1) gauge symmetry.

1 Introduction

The four-dimensional gravity given by the Einstein-Hilbert action includes the unrenormalizable ultraviolet divergence. In order to overcome this, to extend the gravitational theory was invented, such as the causal dynamical triangulation [1,2], the loop quantum gravity [3,4], and the superstring theory [5–7], etc. The asymptotic safety of gravitational theories has been also studied [8,9].

Meanwhile, the holographic principle is held [10–13]. From that, we can infer there would be some microscopic structure in the gravity, which is considered to be given in one lower dimensional spacetime and appear as the gravity at the low-energy. Holographic properties of the gravity are now being studied, and one of the issues closely related with the holography is the asymptotic symmetry group (ASG).

The ASG in hep-th usually composes of the general coordinate transformations, that maintain the fall-off conditions (usually denoted as $\mathcal{O}(r^{-p})$), in which what are observationally indistinguishable are identified. The ASG now usually investigated in hep-th are some ASG on the asymptotic flat space-time [14, 15] or AdS₃ [16] (some ASG on dS is also considered [18]).

In the ASG on the asymptotic flat spacetime, just an ADM mass [19] or infinite charges [15] can be obtained up to how to take the fall-off conditions. In the latter case, the symmetries of the ASG are broken (their central charges are analyzed [20, 21]), which is one of the keys now being investigated to solve the information paradox [22].

Moreover, the latter case is developed to the study on equivalences between the Ward identities and the gravitational soft theorems [23–27], and the holographic properties of the asymptotic flat spacetime [28, 29]. In addition, some equivalence between soft theorems and the memory effects are discussed [30–34]. The observational studies concerning the memory effects in gravitational waves is reviewed in [35]. There are also studies for the soft-hairs on photon spheres and black hole shadows [36,37].

On the other hand, the Virasoro algebra with a central charge, Brown-Henneaux charge, was obtained in the asymptotic AdS_3 with some ASG corrections, which can be identified with that of the CFT_2 on the boundary of the AdS_3 [16] (derivation by Fujikawa method [17]). Applying this to the near-horizon geometry of the D1-D5-P system, the entropy microscopically obtained in [39, 40] was reproduced [38]. Also, Kerr/CFT correspondence was discovered in the near-horizon geometry of the extremal Kerr black hole [41], and a correspondence analogous to Kerr/CFT was obtained based on an explicit D-brane configuration given by the near-horizon geometry of the rotating extremal NS5-branes [42].

Exploiting the fact that the flat spacetime and AdS are interchanged each other up to the AdS radius, various relations between gravities, thermodynamics, boundary CFT, etc on those spacetimes are studied in connection the ASG [43–46].

The analogous group with the ASG can be also considered in the gauge theory, which is called as the large gauge group. In fact, in the review article [15], the large U(1) charges are introduced based on the U(1) charge by mixing some arbitrary function ε into its expression. Those are defined for each ε . Therefore, the large U(1) charge infinitely exists as ε can be infinitely considered (charge's values determined for each ε are respectively constant for the *t*-direction). The infinite transformations each charge generates form a large gauge symmetry group. As for the studies concerning the large U(1) gauge symmetry, the following studies could be taken [47–56].

Now, let us remind that a chiral transformation can be considered when there is a gauge symmetry. Then, since there are the large gauge symmetries, the chiral transformations associated with that can be considered. However, since those have not been considered yet, we investigate those in this study.

This study would lead to developments of the studies referred above toward studies concerning chiral symmetries and its anomalies. In addition, since anomalies have closed relations with the breaking of the gauge symmetry brought by the regularization of the UV-divergence, this study would relate the large gauge symmetries and those issues. Moreover, from the fact that phenomena caused from gauge anomalies are observables, we can discuss the observability of the anomalies in this study as the extension of those, which is interesting as the observation of the large gauge symmetry. Further, although the anomalies in this study are on the gauge field background, we can take the gravitational field as the background. We discuss those last two topics in Sec. 14, and mention the organization of this paper.

The model addressed in this study is given in Sec. 2, and Noether's second theorem to be used in Sec. 8 are reviewed in Sec. 3. Then, once the U(1) charge is given on the Cartesian coordinates in Sec. 4, it is given again on the Penrose coordinates in Sec. 6. The Penrose coordinates and its relations with the Cartesian coordinates are reviewed in Sec. 5. In Sec. 7, the large U(1) charges are given on the Penrose coordinates based on the U(1) charge given in Sec. 6, then, in Sec. 8, those are given again on the Cartesian coordinates.

In Sec. 9, the chiral charges are once heuristically constituted based on the large U(1) charges given in Sec. 8. We name the chiral transformations those chiral charges generate as the large chiral transformations, then it is shown that the large chiral charges heuristically constituted in Sec. 9 can be defined with Noether's theorem reviewed in Sec. 3. In Sec. 10, the anomaly equation associated with the large chiral symmetry is once heuristically obtained.

In Sec. 11, defining the BRS transformations for the large U(1) gauge symmetry (which we name as the large BRS transformation), the one-loop diagrams obtained by axializing the one-loop diagrams of the fermionic field coupling to the multiple classical gauge fields is considered (those one-loop diagrams constitute the effective action for the model in this study). Then, performing the large BRS transformations to those axialized one-loop diagrams, it is shown that the anomaly equations obtained in Sec. 10 can be derived from the calculations of the e.v. of those.

In Sec. 12, it is shown that the anomaly equation obtained in Sec. 10 and 11 can be derived by the Fujikawa method. In Sec. 13, we comment on the anomaly equation associated with the large U(1) symmetry which we can consider from this study, and in Sec. 14, the summary and future development of this study are discussed. In Appendix. A, the derivation of some equations in Sec. 3 is noted, and, in Appendix. B, the formulas used in Sec. 11 are noted.

2 The model considered in this study

The model considered in this study is given as follows:

$$S_0 = \int d^4x \sqrt{-g} \mathcal{L}_0, \quad \mathcal{L}_0 = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + \bar{\psi} i \not\!\!D \psi, \qquad (1)$$

where $\not{\!\!D} \equiv \gamma^{\mu} D_{\mu} (D_{\mu} \equiv \partial_{\mu} - ieA_{\mu}), \ \bar{\psi} \equiv \psi^{\dagger} \gamma_{0}$. g means the determinant of the metric, A_{μ} is the U(1) gauge field. Thus, $F_{\mu\nu} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu}$. ψ is the Dirac field, which is massless in this study.

3 Noether's second theorem

In this study, we address the charges in the case that transformation parameters depend on coordinates. Therefore, we give the general expression of charges in Noether's second theorem. The result in this section is used in Sec. 6 and 7.

3.1 Noether's first theorem

We begin with Noether's first theorem as it is needed to explain Noether's second theorem. We consider a four-dimensional spacetime Ω patched by the coordinates x^{μ} , and write a given Lagrangian as $L(\phi_A, \phi_{A,\mu})$, where $\phi_A = \phi_A(x)$ means all fields $(A = 1, \dots, N \text{ identify each field})$ and $\phi_{A,\mu} \equiv \partial_{\mu}\phi_A(x)$. Then, considering action: $I = \int_{\Omega} d^4x L$, the Euler-Lagrange (EL) equation for each ϕ_A can be given as

$$[L]^{A} \equiv \frac{\partial L}{\partial \phi_{A}} - \partial_{\mu} \left(\frac{\partial L}{\partial \phi_{A,\mu}}\right).$$
⁽²⁾

We write the constant transformations (global transformations) as follows:

$$\delta x^{\mu} \equiv \sum_{r=1}^{n} \eta^{r} X^{\mu}{}_{r}, \quad \delta \phi_{A} \equiv \sum_{r=1}^{n} \eta^{r} M_{r,A}, \tag{3}$$

where η^r $(r = 1, 2, \dots, n)$ are the constant transformation parameters. $X^{\mu}{}_r$ and $M_{r,A}$ are functions with regard to x^{μ} and ϕ_A $(M_{r,A}$ may generally depend on the coordinates). Then, the variation of I under (3) can be written as $\int d^4x \, \eta^r \, \partial_{\mu}(\cdots)$. From this, the following conservation law can be obtained:

$$\partial_{\mu}J^{\mu}_{r} = 0, \quad J^{\mu}_{r} \equiv \frac{\partial L}{\partial \phi_{A,\mu}} M_{r,A} - T^{\mu}{}_{\nu} X^{\nu}{}_{r}, \quad T^{\mu}{}_{\nu} \equiv \frac{\partial L}{\partial \phi_{A,\mu}} \phi_{A,\nu} - \delta^{\mu}{}_{\nu}L \tag{4}$$

for each r. From (4), one charge for each r can be defined as

$$Q_r^{(G)} \equiv \int_{\Sigma} d^3 x \, J_r^0, \tag{5}$$

where Σ means a x^0 -constant 3D hypersurface on Ω .

3.2 Noether's second theorem

Now, let us write the local transformations as follows:

$$\delta x^{\mu} \equiv \lambda^{r} X^{\mu}{}_{r}, \quad \delta \phi_{A} \equiv \lambda^{r} M_{r,A} + \partial_{\mu} \lambda^{r} N_{r}{}^{\mu}{}_{,A}, \tag{6}$$

where $\lambda^r = \lambda^r(x)$ are the transformation parameters. $X^{\mu}{}_r$, $M_{r,A}$ and $N_r{}^{\mu}{}_{,A}$ are functions of x^{μ} and ϕ_A . We use the same notations as those in (3). If I is invariant under (6), the following equations are held:

$$\partial_{\mu}B^{\mu}{}_{r} = 0, \quad B^{\mu}{}_{r} + \partial_{\nu}C^{\nu\mu}{}_{r} = 0, \quad C^{\mu\nu}{}_{r} + C^{\nu\mu}{}_{r} = 0, \tag{7}$$
$$B^{\mu}{}_{r} \equiv J^{\mu}{}_{r} + [L]^{A}N^{\mu}{}_{r,A}, \quad C^{\mu\nu}{}_{r} \equiv \frac{\partial L}{\partial\phi_{A,\mu}}N^{\nu}{}_{r,A},$$

where J_r^{μ} is that in (4). The derivation of these is given in Appendix. A. From the first one of (7), a conservation law can be obtained for each r as follows:

$$\partial_{\mu} J_r^{\mu} = 0, \tag{8}$$

where the EL equations (2) have been used. Therefore, one charge can be defined for each r as

$$Q_r^{(\mathrm{L})} \equiv \int_{\Sigma} d^3 x \, J_r^0,\tag{9}$$

where Σ means a x^0 -constant hypersurface as well as Σ in (5).

4 The U(1) charge on the Cartesian coordinates

In the previous section, we generally defined the charge with regard to the local transformation. Using that, we give the U(1) charge in our model on the Cartesian coordinates.

The equations of motion with regard to A_{μ} , ψ and $\bar{\psi}$ can be obtained from (1) as follows:

The Lagrangian density $\sqrt{-g} \mathcal{L}_0$ in (1) is invariant under the following U(1) gauge transformation:

$$A_{\mu} \to A_{\mu} + e^{-1} \partial_{\mu} \lambda,$$
 (11a)

$$\psi \to e^{ie\lambda}\psi, \quad \bar{\psi} \to e^{-ie\lambda}\bar{\psi},$$
 (11b)

where $\lambda = \lambda(x)$. Taking λ to the linear-order, we denote (11) by the notations (6). Then, X^{μ}_{r} , $M_{r,A}$ and $N_{r}^{\mu}{}_{,A}$ in (6) are taken as $X^{\mu} = 0$ and

$$(M_{,0}, M_{,1}, M_{,2}, M_{,3}, M_{,4}, M_{,5}) = (0, 0, 0, 0, ie\psi, -ie\bar{\psi}),$$
(12a)

$$(N^{\mu}_{,0}, N^{\mu}_{,1}, N^{\mu}_{,2}, N^{\mu}_{,3}, N^{\mu}_{,4}, N^{\mu}_{,5}) = e^{-1}(\delta^{\mu}_{0}, \delta^{\mu}_{1}, \delta^{\mu}_{2}, \delta^{\mu}_{3}, 0, 0),$$
(12b)

where $\mu = 0, \dots, 3$. We have taken ϕ_A as $(\phi_0, \phi_1, \phi_2, \phi_3, \phi_4, \phi_5) = (A_0, A_1, A_2, A_3, \psi, \bar{\psi})$. We have omitted to denote r as the index n, which is r_{max} , is 1. Applying (12) to (7),

$$B^{\mu} = \frac{\partial(\sqrt{-g}\,\mathcal{L}_0)}{\partial(\partial_{\mu}\psi)}(ie\psi) + \frac{\partial(\sqrt{-g}\,\mathcal{L}_0)}{\partial(\partial_{\mu}\bar{\psi})}(-ie\bar{\psi}) = \sqrt{-g}\,J^{\mu}_{\mathrm{U}(1)}, \ C^{\mu\nu} = \frac{\partial(\sqrt{-g}\,\mathcal{L}_0)}{\partial(\partial_{\mu}A_{\lambda})}e^{-1}\delta^{\nu}{}_{\lambda} = -\sqrt{-g}\,F^{\mu\nu}, (13)$$

where the EL equations (2) have been used. Then, the conservation law for the U(1) gauge transformation (11) can be obtained from the first one of (7) as

$$0 = \nabla_{\mu} (\sqrt{-g} J^{\mu}_{\mathrm{U}(1)}). \tag{14}$$

Now, we consider the following Cartesian coordinates:

$$ds^{2} = (dx^{0})^{2} - (dx^{1})^{2} - (dx^{2})^{2} - (dx^{3})^{2}.$$
(15)

Then, following (9), the U(1) charge on the Cartesian coordinates can be given as

$$Q_{\mathrm{U}(1)}^{(\mathrm{M})} \equiv \int_{\Sigma} d^3 x \, J_{\mathrm{U}(1)}^0 = \sum_{k=1}^3 \int_S d\sigma_k \, F^{0k},\tag{16}$$

where Σ is a x^0 -constant 3D hypersurface, and S means its boundary.

5 The Penrose coordinates

In this study, the chiral anomaly of the large U(1) gauge symmetry is addressed. We call it as *the large chiral anomaly*. We define it on the Cartesian coordinates following the custom that the U(1) chiral anomaly is defined on the Cartesian coordinates. However, the asymptotic symmetry is defined on the Penrose coordinates given with (z, \bar{z}) (the reason to consider the (z, \bar{z}) coordinates is that, by that, it becomes easier to address "the matching condition" [15]). Therefore, we review the link between the Cartesian coordinates, etc. Those are used in the following sections.

We consider the following coordinate transformation:

$$x^{0} = u + r, \quad x^{1} = +r \frac{1 - z\bar{z}}{1 + z\bar{z}}, \quad x^{2} + ix^{3} = +\frac{2rz}{1 + z\bar{z}} \quad \text{for } t \ge 0,$$
 (17a)

$$x^{0} = v - r, \quad x^{1} = -r \frac{1 - z\bar{z}}{1 + z\bar{z}}, \quad x^{2} + ix^{3} = -\frac{2rz}{1 + z\bar{z}} \quad \text{for } t \le 0,$$
 (17b)

where x^{μ} are the Cartesian coordinates defined in (15). $r^2 \equiv (x^1)^2 + (x^2)^2 + (x^3)^2$, and (z, \bar{z}) are the coordinates on a complex plane defined as the destinations of the mappings of the points on the surface of the spatial S^2 in (15). Using (17), ds^2 in (15) can be given as

$$ds^{2} = \begin{cases} du^{2} + 2dudr - 2r^{2} \gamma_{z\bar{z}} dz d\bar{z} & \text{for } t \ge 0, \\ dv^{2} - 2dvdr - 2r^{2} \gamma_{z\bar{z}} dz d\bar{z} & \text{for } t \le 0, \end{cases}$$
(18)

where $\gamma_{z\bar{z}} \equiv \frac{2}{(1+z\bar{z})^2}$. We show the 4D flat spacetime with the coordinates (18) in Fig. 1.



Figure 1: The 4D flat spacetime with the coordinates (18)

We note the points in the Penrose coordinates which are used in the next sections:

- 1) The x^0 -constant hypersurfaces infinitely exist for each x^0 . However, the boundaries of those on the $x^0 \ge 0$ and $x^0 \le 0$ region are a \mathcal{I}^+_+ and a \mathcal{I}^-_+ commonly.
- 2) On the $x^0 \ge 0$ and $x^0 \le 0$ region respectively, the $x^0 \to \infty$ and $x^0 \to -\infty$ hypersurface overlaps the $r \to \infty$ hypersurface, except for $(\mathcal{I}^+_+, \mathcal{I}^+_-)$ and $(\mathcal{I}^-_-, \mathcal{I}^+_-)$.

6 The U(1) charge on the Penrose coordinates

The large U(1) charges are defined by some extention of the U(1) charge on the Penrose coordinates [15]. However, we gave the U(1) charge on the Cartesian coordinates in Sec. 4. Therefore, we transform its coordinates to the Penrose coordinates.

First, we consider the coordinates $y^a \equiv (t, r, z, \overline{z})$, where (z, \overline{z}) and r are those in (17). Meanwhile, we denote the Cartesian coordinates (15) as $x^{\mu} \equiv (t, x^1, x^2, x^3)$. $\tilde{J}^a_{\mathrm{U}(1)}$ and \tilde{F}^{ab} , quantities on the y^a coordinates, are given from those on the x^{μ} coordinates as

$$\tilde{J}^{a}_{U(1)} = h^{a}{}_{\mu}J^{\mu}_{U(1)}, \quad \tilde{F}^{ab} = h^{a}{}_{\mu}h^{b}{}_{\nu}F^{\mu\nu}, \quad h^{a}{}_{\mu} \equiv \partial y^{a}/\partial x^{\mu},$$
(19)

where $J_{U(1)}^m$ is defined in (10). With these, the U(1) charge (16) on the coordinates y^a are given as

$$\tilde{Q}_{\mathrm{U}(1)} \equiv \int_{\Sigma} dr \, d^2 z \, r^2 \, \gamma_{z\bar{z}} \, \tilde{J}^t_{\mathrm{U}(1)} = \int_{S} d^2 z \, r^2 \, \gamma_{z\bar{z}} \, \tilde{F}^{rt}.$$
(20)

We change the coordinates of $\tilde{Q}_{U(1)}$ to the Penrose coordinates (18). We denote the Penrose coordinates (u, r, z, \bar{z}) and (v, r, z, \bar{z}) with w^{μ} , commonly. $\tilde{F}^{\mu\nu}$, quantities on the w^m coordinates, are given from those on the y^a coordinates as

$$\widetilde{J}^{\mu}_{\mathrm{U}(1)} = h^{\mu}{}_{a}\widetilde{J}^{a}_{\mathrm{U}(1)}, \quad \widetilde{F}^{\mu\nu} = h^{\mu}{}_{a}h^{\nu}{}_{b}\widetilde{F}^{ab}, \quad h^{\mu}{}_{a} \equiv \partial w^{\mu}/\partial y^{a}$$
(21)

(we use the same notation "h" as (19), we tells the difference by the indices). With these, the U(1) charge (20) on the w^m coordinates are given as

$$\widetilde{Q}^{+} \equiv \int_{\mathcal{I}_{-}^{+}} d^{2}z \, r^{2} \, \gamma_{z\bar{z}} \, \widetilde{F}_{ru}, \quad \widetilde{Q}^{-} \equiv \int_{\mathcal{I}_{+}^{-}} d^{2}z \, r^{2} \, \gamma_{z\bar{z}} \, \widetilde{F}_{rv}.$$

$$(22)$$

 \widetilde{Q}^+ and \widetilde{Q}^- are those on the $x^0 \ge 0$ and $x^0 \le 0$ regions given by (u, r, z, \overline{z}) and (v, r, z, \overline{z}) , respectively. The boundaries of *t*-constant 3D hypersurfaces are always \mathcal{I}^+_- and \mathcal{I}^+_- on the w^m coordinates, respectively, as can be seen in Fig. 1. In addition, it is obvious that \widetilde{Q}^\pm agree with each other as the integral value for a common x^0 -constant surface at $x^0 = 0$. These guarantee \widetilde{Q}^\pm are conserved quantities.

7 The large U(1) charges on the Penrose coordinates

In the previous section, the U(1) charge was given on the Penrose coordinates. In this section, the large U(1) charges (the large charges) are defined by extending that.

Following [15], the large gauge charges are defined by mixing $\varepsilon = \varepsilon(z, \bar{z})$ into \tilde{Q}^{\pm} in (22) as

$$\widetilde{Q}_{\varepsilon}^{+} \equiv \int_{\mathcal{I}_{-}^{+}} d^{2}z \, r^{2} \, \gamma_{z\bar{z}} \, \varepsilon \, \widetilde{F}_{ru}, \quad \widetilde{Q}_{\varepsilon}^{-} \equiv \int_{\mathcal{I}_{+}^{-}} d^{2}z \, r^{2} \, \gamma_{z\bar{z}} \, \varepsilon \, \widetilde{F}_{rv}, \tag{23}$$

where ε is an arbitrary function, which amplifies/weakens the electro-magnetic field for each angle. Since the function to be considered as ε infinitely exists, the value of $\widetilde{Q}_{\varepsilon}^{\pm}$ infinitely exists as well, and each of which is respectively constant against x^0 (this can be explained in the same way under (22)). In what follows, we give \widetilde{F}_{ru} and \widetilde{F}_{rv} in (23) in terms of the conserved current.

From the Maxwell equations (10), $\nabla^{\mu} \widetilde{F}_{\mu u} = \widetilde{J}_{U(1)u}$ and $\nabla^{\mu} \widetilde{F}_{\mu v} = \widetilde{J}_{U(1)v}$. The metrices on the Penrose coordinates (18) are given as

$$g_{uu} = +g_{ur} = +g_{ru} = 1, \quad g_{z\bar{z}} = g_{\bar{z}z} = -r^2 \gamma_{z\bar{z}} \quad \text{for } t \ge 0,$$
 (24a)

$$g_{vv} = -g_{vr} = -g_{rv} = 1, \quad g_{z\bar{z}} = g_{\bar{z}z} = -r^2 \gamma_{z\bar{z}} \quad \text{for } t \le 0$$
 (24b)

(others are 0, and $\gamma_{z\bar{z}}$ is given in (18)). From these,

$$+ \partial_u \widetilde{F}_{ru} = J_{\mathrm{U}(1)u} - g^{z\bar{z}} \nabla_{\bar{z}} \widetilde{F}_{zu} - g^{\bar{z}z} \nabla_z \widetilde{F}_{\bar{z}u}, \qquad (25a)$$

$$-\partial_v \widetilde{F}_{rv} = J_{\mathrm{U}(1)v} - g^{z\bar{z}} \nabla_{\bar{z}} \widetilde{F}_{zv} - g^{\bar{z}z} \nabla_z \widetilde{F}_{\bar{z}v}.$$
(25b)

From (25), \widetilde{F}_{ru} on \mathcal{I}^+_- and \widetilde{F}_{rv} on \mathcal{I}^-_+ can be given as follows:

$$\widetilde{F}_{ru}|_{\mathcal{I}^+_{-}} = \widetilde{F}_{ru}|_{u=u_0} + \int_{u_0}^{-\infty} du \, (\widetilde{J}_{\mathrm{U}(1)u} - g^{z\bar{z}} \nabla_{\bar{z}} \widetilde{F}_{zu} - g^{\bar{z}z} \nabla_z \widetilde{F}_{\bar{z}u}), \tag{26a}$$

$$\widetilde{F}_{rv}|_{\mathcal{I}^{-}_{+}} = \widetilde{F}_{rv}|_{v=v_0} - \int_{v_0}^{\infty} dv \left(\widetilde{J}_{\mathrm{U}(1)v} - g^{z\bar{z}} \nabla_{\bar{z}} \widetilde{F}_{zv} - g^{\bar{z}z} \nabla_{z} \widetilde{F}_{\bar{z}v} \right),$$
(26b)

where \mathcal{I}^+_- and \mathcal{I}^-_+ can be reached by taking $u \to -\infty$ and $v \to +\infty$ on the infinite r line, therefore, r is being taken to ∞ in (26a) and (26b).

When u_0 and v_0 are some finite, the values of $\widetilde{F}_{ru}|_{u=u_0}$ and $\widetilde{F}_{rv}|_{v=v_0}$ are concretely unclear, and (26a) and (26b) do not concretely make sense. However, the point to be reached by $u \to +\infty$ $(v \to -\infty)$ on the infinite r line is \mathcal{I}^+_+ (\mathcal{I}^-_-) . In lights of this, we take u_0 to $+\infty$ $(v_0$ to $-\infty)$. Then, with the fact that r in (26) is being taken to ∞ (as mentioned above), $\widetilde{F}_{ru}|_{u=u_0}$ and $\widetilde{F}_{rv}|_{v=v_0}$ are reduced to $\widetilde{F}_{ru}|_{\mathcal{I}^+_+}$ and $\widetilde{F}_{rv}|_{\mathcal{I}^-_-}$, respectively. The values of $\widetilde{F}_{ru}|_{\mathcal{I}^+_+}$ and $\widetilde{F}_{rv}|_{\mathcal{I}^-_-}$ have been known as

$$\widetilde{F}_{ru}|_{\mathcal{I}^+_+} = \widetilde{F}_{rv}|_{\mathcal{I}^-_-} = 0 \tag{27}$$

due to the facts and assumptions: 1) electro-magnetic field never reach \mathcal{I}^+_+ and \mathcal{I}^-_- , 2) since the Dirac field in this study is massless, no charge exists at \mathcal{I}^+_+ and \mathcal{I}^-_- , and 3) no charge exists on \mathcal{I}^+_- and \mathcal{I}^-_+ .

As such, taking u_0 and v_0 to $+\infty$ and $-\infty$, respectively, we substitute (26a) and (26b) in Q_{ε}^{\pm} in (23). As a result, the large gauge charges on the Penrose coordinates are finally given as

$$\widetilde{Q}_{\varepsilon}^{+} = \int_{\infty}^{-\infty} du \int_{\mathcal{I}_{-}^{+}} d^{2}z \,\varepsilon \,\Big\{ - \left(\nabla_{z} \widetilde{F}_{\bar{z}u} + \nabla_{\bar{z}} \widetilde{F}_{zu}\right) + \gamma_{z\bar{z}} \,\widetilde{J}_{\mathrm{U}(1)u} \Big\},\tag{28a}$$

$$\widetilde{Q}_{\varepsilon}^{-} = \int_{-\infty}^{\infty} dv \int_{\mathcal{I}_{+}^{-}} d^2 z \, \varepsilon \, \Big\{ + \left(\nabla_z \widetilde{F}_{\bar{z}v} + \nabla_{\bar{z}} \widetilde{F}_{zv} \right) + \, \gamma_{z\bar{z}} \, \widetilde{J}_{\mathrm{U}(1)v} \Big\}.$$
(28b)

We comment on the integral regions of (28a) and (28b). r is taken to infinity in (28a), therefore, by the property 2 in Sec.5, the integral region in (28a) overlaps with the *t*-constant 3D hypersurface at $t \to +\infty$ except for \mathcal{I}^+_+ and \mathcal{I}^-_- (these are the points given by $u \to \infty$ and $-\infty$, respectively). These mismatching at \mathcal{I}^+_+ and \mathcal{I}^-_- would be ignoble by the assumptions and facts commented under (27). Therefore, the integral regions of (28a) can be regarded as a *t*-constant 3D hypersurface. The integral region of (28b) is considered in the same way.

8 The large U(1) charges on the Cartesian coordinates

In the previous section, the large U(1) charges were given on the Penrose coordinates. We transform their coordinates to the Cartesian coordinates x^{μ} .

With (19) and (21),
$$\widetilde{F}_{\overline{z}u}$$
, $\widetilde{F}_{\overline{z}v}$ and $\widetilde{J}_{U(1)u}$ can be given by those on the x^{μ} coordinates as

$$\widetilde{F}_{\overline{z}u} = h_{\overline{z}}{}^{\mu}F_{\mu t}, \quad \widetilde{F}_{zu} = h_{z}{}^{\mu}F_{\mu t}, \quad \widetilde{J}_{\mathrm{U}(1)u} = h_{u}{}^{\mu}J_{\mathrm{U}(1)\mu}.$$
 (29a)

$$\widetilde{F}_{\overline{z}v} = h_{\overline{z}}{}^{\mu}F_{\mu t}, \quad \widetilde{F}_{zv} = h_{z}{}^{\mu}F_{\mu t}, \quad \widetilde{J}_{\mathrm{U}(1)v} = h_{v}{}^{\mu}J_{\mathrm{U}(1)\mu},$$
(29b)

where $J_{\mathrm{U}(1)\mu}$ is defined in (10). Substituting these in (28), the large gauge charges $\widetilde{Q}_{\varepsilon}^{\pm}$ in (28) can be commonly given on the x^{μ} coordinates as

$$Q_{\varepsilon} \equiv \int_{\Sigma} d^3 x \,\varepsilon \left(h_z^{\ \mu} \nabla_{\mu} (h_{\bar{z}}^{\ \nu} F_{\nu t}) + h_{\bar{z}}^{\ \mu} \nabla_{\mu} (h_z^{\ \nu} F_{\nu t}) \right) + \int_{\Sigma} d^3 x \,\varepsilon \, J_{\mathrm{U}(1)t},\tag{30}$$

 Σ means a x^0 -constant 3D hypersurface. The arguments of ε are supposed being given by the Cartesian coordinates. It can be seen that the large U(1) current can be defined as $J^{\mu}_{\varepsilon} \equiv \varepsilon J^{\mu}_{\mathrm{U}(1)}$.

9 The large chiral symmetry

In the previous section, the large U(1) charges were given on the Cartesian coordinates. In this section, we once heuristically constitute the chiral charges based on those. We name those as *the large chiral charges*. Then, we show that the transformations those large chiral charges generate are symmetric in our model and it is possible to define those based on Noether's theorem.

9.1 Heuristic constitution of the large chiral charges

The chiral charge can be obtained from the U(1) charge by changing its integrand $\bar{\psi}\gamma^{\mu}\psi$ (the U(1) current) to $\psi\gamma^{5}\gamma^{\mu}\psi$. This means, giving the U(1) charge in the chiral representation (the form in which the left- and right-hand components are separately written), to flip the sign of the right-hand component. In lights of this way, we change Q_{ε} in (30) to a chiral charge.

For this purpose, taking to the chiral representation: $\gamma^{\mu} = \begin{pmatrix} 0 & \sigma^{\mu} \\ \bar{\sigma}^{\mu} & 0 \end{pmatrix}$ and $\psi = (\xi, \eta)^{T}$, where $\sigma^{\mu} \equiv (I_{2}, \vec{\sigma}), \ \bar{\sigma}^{\mu} \equiv (I_{2}, -\vec{\sigma}) \ (\vec{\sigma} \text{ mean Pauli matrices})$, we give Q_{ε} in (30) as

Contribution of left-hand (L.H.)

$$Q_{\varepsilon} \text{ in } (30) = \frac{1}{2} \int_{\Sigma} d^3 x \, \varepsilon \left(h_z^{\mu} \nabla_{\mu} (h_{\bar{z}}^{\mu} F_{\mu t}) + h_{\bar{z}}^{\mu} \nabla_{\mu} (h_z^{\mu} F_{\mu t}) \right) - e \int_{\Sigma} d^3 x \, \varepsilon \, \eta^{\dagger} \, \eta$$

$$(31)$$

$$+ \frac{1}{2} \int_{\Sigma} d^3 x \, \varepsilon \left(h_z^{\mu} \nabla_{\mu} (h_{\bar{z}}^{\mu} F_{\mu t}) + h_{\bar{z}}^{\mu} \nabla_{\mu} (h_z^{\mu} F_{\mu t}) \right) - e \int_{\Sigma} d^3 x \, \varepsilon \, \xi^{\dagger} \, \xi,$$

where, in the one above, dividing the bosonic part in half, we have included those in the left-hand part (L.H.) and right-hand part (R.H.), respectively. The first and second terms in the L.H. and R.H. are usually referred to as "soft-term" and "hard-term".

We perform the manipulation mentioned in the beginning of this subsection to (31), which is to change the R.H. as $+R.H. \rightarrow -R.H.$, leaving L.H. as it is. Therefore, (31) is changed as follows:

L.H. - R.H. =
$$-e \int_{\Sigma} d^3 x \,\varepsilon \,(\eta^{\dagger} \,\eta - \xi^{\dagger} \,\xi) = \int_{\Sigma} d^3 x \,J_{c\varepsilon}^{\mu} \equiv Q_{c\varepsilon}, \quad J_c^{\mu} \equiv -e \bar{\psi} \gamma_5 \gamma^{\mu} \psi, \quad J_{c\varepsilon}^{\mu} \equiv \varepsilon J_c^{\mu}, \quad (32)$$

where we have defined the obtained quantity as $Q_{c\varepsilon}$, We name it as the large chiral charges. Remarkably, there is no contribution from the soft-term as those are canceled each other.

9.2 The large chiral transformations and their charges by Noether's theorem

In Sec. 9.1, the large chiral charges were heuristically constituted. In this subsection, we firstly define the transformations those generate as *the large chiral symmetry*, then, obtain the large chiral charges once heuristically constituted in Sec. 9.1 as the conserved charges based on Noether's theorem.

First, let us remind the fact mentioned at the last of Sec. 3. Then, the simultaneous Poisson bracket (the Grassmann bracket) for our Dirac field ψ^{α} are given as

$$\{\psi_{\alpha}(t,\vec{x}),\pi^{\beta}(t,\vec{y})\}_{\text{G.B.}} = \delta_{\alpha}{}^{\beta} \delta^{3}(\vec{x}-\vec{y}),\tag{33a}$$

$$\{\psi_{\alpha}(t,\vec{x}),\psi_{\beta}(t,\vec{y})\}_{\rm G.B.} = \{\pi^{\alpha}(t,\vec{x}),\pi^{\beta}(t,\vec{y})\}_{\rm G.B.} = 0,$$
(33b)

where (t, \vec{x}) and (t, \vec{y}) are the Cartesian coordinates. $\{\cdot, \cdot\}_{G.B.}$ and π_{α} are defined as

$$\{X(t,\vec{x}), Y(t,\vec{y})\}_{\text{G.B.}} \equiv \int d^3x \left(\frac{\partial X(t,\vec{x})}{\partial \psi^{\alpha}(t,\vec{z})}\frac{\partial Y(t,\vec{y})}{\partial \pi_{\alpha}(t,\vec{z})} + \frac{\partial X(t,\vec{x})}{\partial \pi_{\alpha}(t,\vec{z})}\frac{\partial Y(t,\vec{x})}{\partial \psi^{\alpha}(t,\vec{z})}\right),\tag{34a}$$

$$\pi^{\alpha}(t,\vec{x}) \equiv \frac{\partial \mathcal{L}_0}{\partial \left(\partial_0 \psi_{\alpha}(t,\vec{x})\right)} = i\psi^{\dagger \alpha}(t,\vec{x}).$$
(34b)

X and Y are some functions and \mathcal{L}_0 is that in (1). With (34b), the following Grassmann bracket can be obtained from (33a):

$$\{\psi_{\alpha}(t,\vec{x}),\psi^{\dagger\beta}(t,\vec{y})\}_{\text{G.B.}} = -i\delta_{\alpha}{}^{\beta}\delta^{3}(\vec{x}-\vec{y}).$$
(35)

Using (33b), (35) and $Q_{c,\varepsilon}$ in (32), the following Grassmann bracket can be obtained:

$$\{Q_{c\varepsilon},\psi\}_{G.B.} = i\gamma_5 \,\varepsilon \,\psi, \quad \{Q_{c\varepsilon},\bar{\psi}\}_{G.B.} = i\bar{\psi}\gamma_5 \,\varepsilon. \tag{36}$$

(36) means that, if $Q_{c\varepsilon}$ are conserved quantities, $Q_{c\varepsilon}$ are the charges associated with the following transformations:

$$\psi \to \psi' = e^{ie\varepsilon\lambda\gamma_5}\psi = \psi + ie\,\varepsilon\lambda\gamma_5\psi + \mathcal{O}(\lambda^2),\tag{37a}$$

$$\bar{\psi} \to \bar{\psi}' = \bar{\psi} e^{ie\varepsilon\lambda\gamma_5} = \bar{\psi} + ie\,\varepsilon\lambda\,\bar{\psi}\gamma_5 + \mathcal{O}(\lambda^2).$$
 (37b)

We name these as the large chiral transformations.

We verify that (37) is symmetric transformation in our model (1). For this purpose, we examine the variation arisen in (1) by (37):

$$\int d^4x \,\bar{\psi} i \,\mathcal{D}\psi \to \int d^4x \,\bar{\psi}' \,i \,\mathcal{D}\psi' = \int d^4x \,\bar{\psi} \,i \,\mathcal{D}\psi - e \int d^4x \,\partial_\mu (\lambda \,\varepsilon) \,\bar{\psi}\gamma_5 \gamma^\mu \psi$$
$$= \int d^4x \,\bar{\psi} \,i \,\mathcal{D}\psi + \underbrace{\int d^4x \,\partial_\mu (\lambda \,\varepsilon J_c^\mu)}_{=0} - \underbrace{\int d^4x \,\lambda \,\varepsilon \,\partial_\mu J_c^\mu}_{=0}, \qquad (38)$$

where J_c^{μ} is defined in (32). Since λ is assumed to be vanished at $r \to \infty$, the second term is vanished. The third term is also vanished as $\bar{\psi}\gamma_5\gamma^{\mu}\psi = \text{const.}$ due to the chiral symmetry. Therefore, (37) is a symmetric transformation in our model (1).

Now, we show that $Q_{c\varepsilon}$ are the charges associated with (37) based on the Noether's theorem. For (37), quantities in (6) can be written as follows:

$$X^{\mu} = N^{\mu}{}_{,A} = 0 \text{ for all } \mu \text{ and } A \tag{39a}$$

$$(M_{,0}, M_{,1}, M_{,2}, M_{,3}, M_{,4}, M_{,5}) = (0, 0, 0, 0, ie\varepsilon\gamma_5\psi, ie\varepsilon\psi\gamma_5),$$
(39b)

where $(\phi_0, \phi_1, \phi_2, \phi_3, \phi_4, \phi_5) = (A_0, A_1, A_2, A_3, \psi, \overline{\psi})$ and n, r_{max} , is 1, therefore we omitted to denote r. Using these in the same way as (13), B^{μ} and $C^{\mu\nu}$ in (7) are obtained as

$$B^{\mu} = J^{\mu}_{c\varepsilon}, \quad C^{\mu\nu} = 0. \tag{40}$$

Therefore, the following conserved law can be obtained by following (7):

$$\partial_{\mu}J^{\mu}_{c\varepsilon} = 0. \tag{41}$$

From this, $Q_{c\varepsilon}$ in (32) can be obtained as the charges for the large chiral transformations (37).

9.3 The large U(1) gauge transformations

In the previous subsection, the large chiral transformations of the fermionic and gauge fields are given. In this subsection, the large U(1) transformations of the fermionic field are given. From (36),

$$\{Q_{\varepsilon},\psi\}_{\mathrm{G.B.}} = i\,\varepsilon\,\psi, \quad \{Q_{\varepsilon},\bar{\psi}\}_{\mathrm{G.B.}} = i\,\varepsilon\,\bar{\psi}.$$
(42)

From these, the large transformed fermionic filed and gauge field can be known as follows:

$$\psi \to \psi' = e^{+ie\varepsilon\lambda}\psi, \quad \bar{\psi} \to \bar{\psi}' = e^{-ie\varepsilon\lambda}\bar{\psi}, \quad A_{\mu} \to A_{\mu} + e^{-1}\partial_{\mu}(\varepsilon\lambda).$$
 (43)

In the one above, the transformation of A_{μ} has not been obtained from the analysis with the Poisson bracket (P.B.) like (42), but obtained by educing it from the transformations of ψ and $\bar{\psi}$. Its reason is that, by the reason noted in the footnote^{*}, it cannot be generally obtained from the analysis of the P.B..

10 The anomaly equation for the large chiral symmetry

In the previous section, we obtained the conservation law (41) associated with the large chiral symmetry (37). In general, conservation laws including γ_5 get quantum corrections, the anomaly. Therefore, we in this section give (41) including the anomaly.

We give (41) as follows:

$$\partial_{\mu}\langle J_{c\varepsilon}^{\mu}\rangle = (\partial_{\mu}\varepsilon)\,\langle J_{c}^{\mu}\rangle + \varepsilon\,\partial_{\mu}\langle J_{c}^{\mu}\rangle = (\partial_{\mu}\varepsilon)\,\langle J_{c}^{\mu}\rangle - e\,\varepsilon\,\partial_{\mu}\langle\bar{\psi}\gamma_{5}\gamma^{\mu}\psi\rangle,\tag{44}$$

where $J^{\mu}_{c\varepsilon}$ and J^{μ}_{c} are defined in (32). $\langle \cdots \rangle$ means the quantum e.v. defined as

^{*}In the global transformation, charges generate transformations as $\{Q_r^{(G)}, \phi_A\}_{P.B.} = M_{r,A}$ $(M_{r,A} \text{ and } Q_r^{(G)} \text{ are those}$ in (3) and (5), respectively). Then, as can be seen from (5) and (9), the forms of the charges in the global and local transformation are the same as each other. From this, it can be seen that the charges in the local transformation can generate the transformations that the charges in the global transformation generate. This means $\{Q_r^{(L)}, \phi_A\}_{P.B.} = M_{r,A}$ (this $M_{r,A}$ and $Q_r^{(L)}$ are those in (6) and (9), respectively).

Concretely, the transformations generated by $\{Q_r^{(L)}, \phi_A\}_{P.B.}$ are the phases of the matter fields (such as (11b)), and the transformations on the gauge fields (such as (11a)) are not be generated by the charges in the gauge transformations.

$$\langle \mathcal{O} \rangle \equiv \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \,\mathcal{D}A_{\mu} \mathcal{D}B \,\mathcal{D}c \mathcal{D}\bar{c} \,\mathcal{O} \,\exp\left[i \int d^4x \sqrt{-g} \,\mathcal{L}_{g.f.}\right]. \tag{45}$$

 $\mathcal{L}_{g.f.}$ means \mathcal{L}_0 with some gauge-fixing terms. While the first term in (44) cannot be generally calculated more than that, the result of the second term is known, using which, (44) can be given as follows:

$$\partial_{\mu}\langle J_{c\varepsilon}^{\mu}\rangle = \left(\partial_{\mu}\varepsilon\right)\langle J_{c}^{\mu}\rangle - \varepsilon \,i\,\frac{\hbar\,e^{3}}{16\pi^{2}}\,\varepsilon^{\mu\nu\rho\sigma}F_{\mu\nu}F_{\rho\sigma}.\tag{46}$$

In this study, \hbar is taken to 1. This is the anomaly equation with regard to the large chiral symmetry.

11 The anomaly equation from the large BRS transformed effective action

In the previous section, we obtained the anomaly equation with regard to the large chiral symmetry as can be seen in (46). In this section, we derive it from the evaluation of the one-loop diagrams.

Specifically, we first define the BRS transformations of the large U(1) symmetry (the large BRS transformations) (Sec. 11.1). Then, by performing those large BRS transformations to the one-loop diagrams constituting the effective action of our model (1), we once obtain anomaly equations associated with the large U(1) current (Sec. 11.2), Then, we change the anomaly equation obtained in Sec. 11.2 by axializing those one-loop diagrams (by changing one of the vector type vertices in each of those one-loop diagrams to the axial-vector type) (Sec. 11.3). Then, evaluating each of those axialized one-loop diagrams, we derive the anomaly equation in (46) (Sec. 11.4-11.6).

11.1 The BRS transformation of the large U(1) gauge transformation

Following the procedure to define a BRS transformation for a given gauge transformation^{\dagger}, we define the BRS transformation of the large gauge transformation from (43) as

$$\boldsymbol{\delta}_{\varepsilon}\psi(x) = ie\,\varepsilon(x)c(x)\psi(x), \quad \boldsymbol{\delta}_{\varepsilon}A_{\mu}(x) = e^{-1}\partial_{\mu}(\varepsilon(x)c(x)), \tag{47}$$

where c(x) is the ghost field. The coordinates are Cartesian. We name these as the large BRS transformations and denote with " δ_{ε} ". These are given in the momentum space as

$$\boldsymbol{\delta}_{\varepsilon}\psi(k) = ie\,\varepsilon(k)c(k)\psi(k), \quad \boldsymbol{\delta}_{\varepsilon}A_{\mu}(k) = -ie^{-1}k_{\mu}\varepsilon(k)c(k), \tag{48}$$

where $\varepsilon(x)c(x)\psi(x) = \int \frac{d^4x}{(2\pi)^4} e^{-ikx} c(k)c(k)\psi(k)$ and $\varepsilon(x)c(x) = \int \frac{d^4x}{(2\pi)^4} e^{-ikx} \varepsilon(k)c(k)$.

11.2 The anomaly equation of the large U(1) current

To derive the anomaly equation of the large chiral current, we first derive the anomaly equation of the large U(1) current from the breaking of the large BRS symmetry (47) in the effective action of (1). We begin with the definition of the effective action:

$$e^{i\Gamma[A]} = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \exp\left[i \int d^4x \sqrt{-g} \mathcal{L}_0\right].$$
(49)

 \mathcal{L}_0 is given in (1). Since the spacetime is flat and the coordinates are Cartesian, $\sqrt{-g} = 1$. This effective action is the one with regard to A_{μ} and can be written as

$$\Gamma[A] = i^{-1} \operatorname{Tr} \ln[\partial \!\!\!/ - ie \mathcal{A}] - \int d^4 x \, \frac{1}{4} \, F_{\mu\nu} F^{\mu\nu}, \qquad (50)$$

[†]It is just a replacement of the gauge transformation parameter with a product of a Grassmann number and a ghost field such as $\lambda \to \theta c$ (θ means a Grassmann number), then take up the part other than θ .

where $\int \mathcal{D}\psi \mathcal{D}\bar{\psi} \exp\left[i\int d^4x \,\bar{\psi}i\mathcal{D}\psi\right] = \text{Det}(-\gamma^0\mathcal{D})$. Tr and Det mean the trace and determinant for the coordinates and the indices of fields.

Operating $\int d^4x \, \boldsymbol{\delta}_{\varepsilon}$ to the effective action ($\boldsymbol{\delta}_{\varepsilon}$ is given in (47)),

$$\int d^4x \, \boldsymbol{\delta}_{\varepsilon} \Gamma[A] = \int d^4x \, \boldsymbol{\delta}_{\varepsilon} A_{\mu} \, \frac{\delta \Gamma[A]}{\delta A_{\mu}} = \int d^4x \, \boldsymbol{\delta}_{\varepsilon} A_{\mu} \, \frac{\delta}{\delta A_{\mu}} \Big(i^{-1} \ln \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \exp\left[i \int d^4x \, \mathcal{L}_0\right] \Big)$$
$$= i^{-1} \int d^4x \, \boldsymbol{\delta}_{\varepsilon} A_{\mu} \left(-i \langle J^{\mu}_{\mathrm{U}(1)} \rangle + i \langle \partial_{\nu} F^{\nu\mu} \rangle \right) = -\int d^4x \, \boldsymbol{\delta}_{\varepsilon} A_{\mu} \, \langle J^{\mu}_{\mathrm{U}(1)} \rangle, \tag{51}$$

where $\partial_{\nu}F^{\nu\mu} = 0$ by the equations of motion. With (47),

r.h.s. of (51) =
$$-e^{-1} \int d^4 x \left(\partial_\mu \varepsilon \, c + \varepsilon \, \partial_\mu c \right) \langle J^\mu_{\mathrm{U}(1)} \rangle = -e^{-1} \int d^4 x \, c \left(\partial_\mu \varepsilon \, \langle J^\mu_{\mathrm{U}(1)} \rangle - \partial_\mu (\varepsilon \, \langle J^\mu_{\mathrm{U}(1)} \rangle) \right),$$
 (52)

where supposing c is vanished at $r \to \infty$, $\int d^4x \, \partial_\mu (\varepsilon c \langle J^{\mu}_{U(1)} \rangle) = 0$. Therefore, (51) is given as follows:

$$\int d^4x \, \boldsymbol{\delta}_{\varepsilon} \Gamma[A] = -e^{-1} \int d^4x \, c \left(\partial_{\mu} \varepsilon \, \langle J^{\mu}_{\mathrm{U}(1)} \rangle - \partial_{\mu} \langle J^{\mu}_{\varepsilon} \rangle \right), \tag{53}$$

where $\varepsilon J_{U(1)}^{\mu} \equiv J_{\varepsilon}^{\mu}$ as defined in (30). From this, we can obtain the equation such as $\partial_{\mu} \langle J_{\varepsilon}^{\mu} \rangle = \cdots$, which is the anomaly equation with regard to the large U(1) gauge symmetry.

11.3 Change the anomaly equation (53) to the one with regard to the large chiral current

As such, we change the equation for $\partial_{\mu} \langle J_{\varepsilon}^{\mu} \rangle$ to the one with regard to the large chiral current in (32). For this purpose, we first give the effective action (50) to the one-loop order by expanding it:

$$i\Gamma[A] = \operatorname{Tr}\left[\ln[\partial] - \sum_{n=1}^{\infty} \frac{1}{n} \left(\partial^{-1} i e \mathcal{A}\right)^{n}\right] = \operatorname{Tr}\ln[\partial] + \sum_{n=1}^{\infty} \Gamma[A]\big|_{n-\mathrm{th}} + \cdots, \qquad (54)$$

where
$$\Gamma[A]|_{n-\text{th}} \equiv \int \left(\prod_{i=1}^{n} \frac{d^4 k_i}{(2\pi)^4}\right) (2\pi)^4 \,\delta^4(K_n) \,\frac{1}{n} \left(\prod_{i=1}^{n} eA_{\mu_n}(-k_n)\right) \Gamma^{(n)},$$
 (55a)

$$\Gamma^{(n)} \equiv \Gamma^{\mu_1 \cdots \mu_n} \equiv -\int \frac{d^4 l}{i(2\pi)^4} \operatorname{tr} \left[\frac{1}{-\ell} \gamma^{\mu_n} \prod_{i=n-1}^{1} \frac{1}{-(\ell+\ell_i)} \gamma^{\mu_i} \right],$$
(55b)

where "..." means the contribution from more than two-loop order, and $K_n \equiv \sum_{i=1}^n k_i$. $\Gamma^{(n)}$ is just an abbreviation of $\Gamma^{\mu_1 \dots \mu_n}$. tr means the trace for spinor's indices. In giving Tr by integrals in (55), we used the Feynman rule[‡]. We ignored $-\int d^4x \frac{1}{4} F_{\mu\nu} F^{\mu\nu}$ in (50) as it is ignorable as can be seen under (51).



Figure 2: The Feynman diagrams of $\Gamma^{(n)}$ for n = 2, 3, 4, 5 in (55b).

We have derived a anomaly equation of the large U(1) current in (53). However, our aim is the anomaly equation associated with the large chiral current. In addition, even if we could eventually

[‡]For the model (1), (i) $(m - k' - i\epsilon)^{-1}$ corresponds to the fermionic inner-lines with momentum k_{μ} $(m = \epsilon = 0)$, (ii) $e\gamma^{\mu}A_{\mu}$ corresponds to the vertexes, and (iii) $-\int \frac{d^4l}{i(2\pi)^4}$ corresponds to the loop-integrate the fermions run around.

obtain $\partial_{\mu}\langle J^{\mu}\rangle$ and $\partial_{\mu}\langle J^{5\mu}\rangle$ from (53), those would not still immediately coincide with the right ones. We explain this point by taking the already known U(1) case.

Suppose the effective action (49) to be separately given by the chirality as $e^{i\Gamma[A]} = \text{Det}(i\mathcal{D}_{R}) \text{Det}(i\mathcal{D}_{L})$. Then, from the equation analogous to (53), $\partial_{\mu}\langle J_{L}^{\mu}\rangle$ and $\partial_{\mu}\langle J_{R}^{\mu}\rangle$ can be obtained. From those, $\partial_{\mu}\langle J^{\mu}\rangle$ and $\partial_{\mu}\langle J^{5\mu}\rangle$ can be obtained as $\partial_{\mu}\langle J_{L}^{\mu}\rangle + \partial_{\mu}\langle J_{R}^{\mu}\rangle$ and $\partial_{\mu}\langle J_{L}^{\mu}\rangle - \partial_{\mu}\langle J_{R}^{\mu}\rangle$, respectively. However, this $\partial_{\mu}\langle J^{\mu}\rangle$ is non-zero and $\partial_{\mu}\langle J^{5\mu}\rangle$ does not coincide with the right chiral anomaly.

However, there is the degree of freedom in the effective action to modify it keeping the correspondence of the effective action to the original theory as it is (as the matter of regularization term), and utilizing that, the effective action can be modified in such a way $\partial_{\mu} \langle J^{\mu} \rangle = 0$ can be realized. At this time, $\partial_{\mu} \langle J^{5\mu} \rangle$ coincides with the right chiral anomaly. However, the analysis of this modification is highly dense[§].

Since this situation would be the same in the case to obtain the anomaly equation of the large chiral current, and as long as we begin with the effective action (49), we would have to perform the modification mentioned above. However, since it is beyond the issue this study addresses, we will obtain $\partial_{\mu} \langle J^{5\mu} \rangle$ by another way.

It is to axialize each of the one-loop diagrams (55b) by changing a γ^{μ} in the each by hands as follows:

$$\gamma^{\mu} \to \gamma^{\mu} \gamma_5. \tag{56}$$

By this, $J^{\mu}_{U(1)}$ and J^{μ}_{ε} in the r.h.s. of (53) are changed to J^{μ}_{c} and $J^{\mu}_{c\varepsilon}$ $(J^{\mu}_{U(1)} \equiv -e\bar{\psi}\gamma^{\mu}\psi$ and $J^{\mu}_{c} \equiv -e\bar{\psi}\gamma_{5}\gamma^{\mu}\psi$, and $J^{\mu}_{\varepsilon} \equiv \varepsilon J^{\mu}_{U(1)}$ and $J^{\mu}_{c\varepsilon} \equiv \varepsilon J^{\mu}_{c}$ as can be seen under (30) and in (32)). Therefore, performing (56) to (53), we can obtain the equation to give $\partial_{\mu}\langle J^{\mu}_{c\varepsilon} \rangle$ by a calculable quantity $\boldsymbol{\delta}_{\varepsilon}\Gamma_{\mathcal{A}}[A]$ as follows:

$$\int d^4x \, \boldsymbol{\delta}_{\varepsilon} \Gamma_{\mathcal{A}}[A] = -e^{-1} \int d^4x \, c \left(\partial_{\mu} \varepsilon \, \langle J^{\mu}_{c} \rangle - \partial_{\mu} \langle J^{\mu}_{c\varepsilon} \rangle \right). \tag{57}$$

where
$$i \Gamma_{\mathcal{A}}[A] \equiv \operatorname{Tr} \ln \left[\partial \right] + \sum_{n=1}^{\infty} \Gamma_{\mathcal{A}}[A] \Big|_{n-\mathrm{th}},$$
 (58)

$$\Gamma_{\mathcal{A}}[A]\Big|_{n-\text{th}} \equiv \int \left(\prod_{i=1}^{n} \frac{d^{4}k_{i}}{(2\pi)^{4}}\right) (2\pi)^{4} \delta^{4}(K_{n}) \frac{1}{n} \left(\prod_{i=1}^{n} eA_{\mu_{i}}(-k_{i})\right) \Gamma_{\mathcal{A}}^{(n)},$$

$$\Gamma_{\mathcal{A}}^{(n)} \equiv \Gamma_{\mathcal{A}}^{\mu_{1}\dots\mu_{n}} \equiv -\int \frac{d^{4}l}{i(2\pi)^{4}} \operatorname{tr}\left[\frac{1}{-\not}\gamma^{\mu_{n}}\gamma_{5}\prod_{i=n-1}^{1} \frac{1}{-(\not+\not K_{i})}\gamma^{\mu_{i}}\right].$$

The equation $\partial_{\mu} \langle J^{\mu}_{c\varepsilon} \rangle = \cdots$ to be obtained from this is what we want to obtain, and the final result is given in (83). With (58), the l.h.s. of (57) can be written as

$$\int d^4x \, \boldsymbol{\delta}_{\varepsilon} \, \Gamma_{\mathcal{A}}[A]|_{n-\text{th}} = \sum_{n=1}^{\infty} \int \left(\prod_{i=1}^{n} \frac{d^4 k_i}{(2\pi)^4} \right) (2\pi)^4 \, \delta^4(K_n) \left(e \, \boldsymbol{\delta}_{\varepsilon}(A_{\mu_n}(-k_n)) \prod_{i=1}^{n-1} e A_{\mu_i}(-k_i) \right) \Gamma_{\mathcal{A}}^{(n)} \\ = \int \frac{d^4 k}{(2\pi)^4} (2\pi)^4 \, \delta^4(k_1) \left(\varepsilon(-k_1) \, c(-k_1) \right) i k_{1\mu} \, \Gamma_{\mathcal{A}}^{(1)}$$
(59a)

$$+ \int \left(\prod_{i=1}^{2} \frac{d^{4}k_{i}}{(2\pi)^{4}}\right) (2\pi)^{4} \,\delta^{4}(K_{2}) \left(\varepsilon(-k_{2}) \,c(-k_{2}) \,eA_{\mu_{1}}(-k_{1})\right) ik_{2\mu} \,\Gamma_{\mathcal{A}}^{(2)} \tag{59b}$$

$$+ \int \left(\prod_{i=1}^{3} \frac{d^4 k_i}{(2\pi)^4}\right) (2\pi)^4 \,\delta^4(K_3) \left(\varepsilon(-k_3) c(-k_3) \prod_{i=1}^{2} eA_{\mu_n}(-k_n)\right) ik_{3\mu} \,\Gamma_{\mathcal{A}}^{(3)} \tag{59c}$$

$$+ \int \left(\prod_{i=1}^{4} \frac{d^4 k_i}{(2\pi)^4}\right) (2\pi)^4 \,\delta^4(K_4) \left(\varepsilon(-k_4) \,c(-k_4) \,\prod_{i=1}^{3} eA_{\mu_n}(-k_n)\right) ik_{4\mu} \,\Gamma_{\mathcal{A}}^{(4)} \tag{59d}$$

[§]As the preparation to perform that modification, we first need to analyze Wess-Zumino consistency-condition. Also, we need to prepare various mathematical techniques (the superfield formalism, Cartan homotopy formula, Chern-Simon forms, etc) to smoothly perform the analysis. We will take care of these in the future work.

$$+ \int \left(\prod_{i=1}^{5} \frac{d^4 k_i}{(2\pi)^4}\right) (2\pi)^4 \delta^4(K_5) \left(\varepsilon(-k_5) c(-k_5) \prod_{i=1}^{4} eA_{\mu_n}(-k_n)\right) ik_{5\mu} \Gamma_{\mathcal{A}}^{(5)}$$
(59e)
+ \cdots . (59f)

Since $\Gamma^{(n)}$ for $n = 1, \dots, 5$ are diverged, we explicitly wrote till n = 5 (as can be seen from (58), the divergence-order of $\Gamma^{(n)}$ is 4-n, but $\Gamma^{(5)}$ potentially includes log-divergence by the regularization scheme we employ as we comment under (66)). "..." means the terms to be surely vanished. In the following subsections, we evaluate each term. The final result is given in (83).

The calculation of $ik_{3\mu}\Gamma_A^{(3)}$ in (59c) 11.4

We calculate (59c), which is given by the *l*-integrate. Since it is UV-diverged, we employ the dimensional regularization. Thus, 4 of the four-dimensional integrate is analytic-continued to some complex number n. We first comment on its treatments:

- n is taken to some complex number when the *l*-integrate is performed. However, in calculations other than that, n is treated as some real number. Here, we will not perform Euclideanization.
- The fermionic field is integrated out before the *l*-integrate by supposing that it is the fermionic field on the real *n*-dimensional Minkowski spacetime.
- γ -matrices are supposed to be on the real *n*-dimensional Minkowski spacetime except for the time that n is taken to some complex number in the *l*-integrate. When n is taken to some complex number, γ -matrices are treated formally, and concrete calculations with γ -matrices, such as $\operatorname{tr}(\gamma_5 \gamma^{\nu} \gamma^{\rho} \gamma^{\alpha} \gamma^{\beta}) = -4i \, \epsilon^{\nu \rho \alpha \beta}$, are performed after performing the *l*-integrate and the $n \to 4$ limit.
- Even when the real *n*-dimensional spacetime is considered $\epsilon^{\mu\nu\sigma\rho}$ and γ_5 work only on the fourdimensional part in that. Therefore, for example, the commutation relations between γ^{μ} and γ_5 on the real *n*-dimension are given as follows:

$$\gamma_5 \gamma^\mu = \begin{cases} -\gamma_5 \gamma^\mu & \text{for } \mu = 0, 1, 2, 3, \\ +\gamma_5 \gamma^\mu & \text{for } \mu = \text{other than } 0, 1, 2, 3. \end{cases}$$
(60)

Then, let us first give the integral variable l_{μ} on the *n*-dimensional as

$$l_{\mu} = \underline{l}_{\mu} + r_{\mu},\tag{61}$$

where $\underline{l}_{\mu} \equiv (l_0, l_1, l_2, l_3, 0, \dots, 0), r_{\mu} \equiv (0, 0, 0, 0, l_4, l_5, \dots, l_{n-1})$ and $l_{\mu} \equiv (l_0, \dots, l_{n-1}).$

- However, even in the *n*-dimension,
 - the elements other than the first four of the external line's momenta k_{μ} remain 0 (namely, for example, $k_{\mu} = (k_0, k_1, k_2, k_3, 0, \dots, 0)$, where k_{μ} ($\mu = 0, \dots, 3$) remain the original values).
 - the first four elements of the external gauge field A_{μ} are also the original values, and others are 0 (namely, $A_{\mu} = (A_0, A_1, A_2, A_3, 0, \dots, 0)$, where A_{μ} remain the original values).

Now, we give $k_{3\mu}$ as the difference between the incoming and outgoing momenta for the vertex, which is $l_{\mu} - (l + K_2)_{\mu}$ (namely, give as $k_{3\mu} = l_{\mu} - (l + K_2)_{\mu}$). Using this in $-k_{3\mu}\gamma^{\mu}\gamma_5$,

$$k_{3\mu}\gamma^{\mu}\gamma_{5} = \not\!\!\!/ \gamma_{5} + \gamma_{5}\left(\not\!\!\!/ + \not\!\!/ K_{2}\right) - 2\not\!\!/ \gamma_{5}, \tag{62}$$

where $\not{l}\gamma_5 = -\gamma_5 \not{l} + 2 \not{r}\gamma_5$ by following (60). Let us rewrte $ik_{\mu}\Gamma_{\mathcal{A}}^{(3)}$ with this; originally, it is given as

$$ik_{3\mu}\Gamma_{\mathcal{A}}^{(3)} = -\int \frac{d^4l}{i(2\pi)^4} \operatorname{tr}\left[\frac{1}{-\ell} (ik_{3\mu}\gamma^{\mu}\gamma_5) \frac{1}{-(\ell+K_2)}\gamma^{\rho} \frac{1}{-(\ell+K_1)}\gamma^{\nu}\right],\tag{63}$$

and we substitute (62) into $ik_{3\mu}\gamma^{\mu}\gamma_5$. Consequently,

$$ik_{3\mu}\widetilde{\Gamma}_{\mathcal{A}}^{(3)} = -\int \frac{d^{n}l}{i(2\pi)^{n}} \operatorname{tr} \left[\underbrace{-i\left(\gamma_{5} \frac{1}{-(\not{l} + \not{K}_{2})} + \frac{1}{-\not{l}}\gamma_{5}\right)\gamma^{\rho} \frac{1}{-(\not{l} + \not{K}_{1})}\gamma^{\nu}}_{+\frac{1}{-\not{l}}(-2i\not{r}\gamma_{5}) \frac{1}{-(\not{l} + \not{K}_{2})}\gamma^{\rho} \frac{1}{-(\not{l} + \not{K}_{1})}\gamma^{\nu} \right],$$
(64)

where $\widetilde{\Gamma}_{\mathcal{A}}^{(3)}$ means the *n*-dimensional version of $\Gamma_{\mathcal{A}}^{(3)} (\equiv \Gamma_{\mathcal{A}}^{\mu\rho\nu})$. Here,

- The first two terms are vanished at arbitrary real n as shown in (65). This is reflecting the BRS symmetry. Therefore, that the term irrelevant of $\not r$ vanishes is universal in $ik_{n\mu}\widetilde{\Gamma}_{\mathcal{A}}^{(n)}$. Along with this, that the term proportional to $\not r$ remains (the third term) is also universal in $ik_{n\mu}\widetilde{\Gamma}_{\mathcal{A}}^{(n)}$.
- The third term is appeared to be vanished at $n \to 4$ as f = 0 at $n \to 4$. However, it includes divergence, and those are multiplied each other. As a result, those coefficients remain at $n \to 4$.
- From these two points, it can be seen that one of the sources of the finite contribution is $-2 \not r \gamma_5$ in (62) (another source is the UV-divergence). It arises when γ_5 is included in the addressed current. Therefore, the finite contribution arises when the current is the chiral type, and vice versa.

$$(\cdots) i\gamma^{\rho} \frac{1}{-(\ell+K_1)} i\gamma^{\nu} \text{ in (64) can be written in the whole (59c) like the following, and is vanished as}$$
$$i\int \left(\prod_{i=1}^{3} \frac{d^n k_i}{(i(2\pi)^4)}\right) (2\pi)^4 \,\delta^4(K_3) \,\varepsilon(-k_3) \,c(-k_3) \,A_{\nu}(-k_1) \,A_{\rho}(-k_2)$$
$$\times \int \frac{d^n l}{i(2\pi)^n} \operatorname{tr} \left[\gamma_5 \left(\frac{1}{\ell+K_2} \gamma^{\rho} \frac{1}{\ell+K_1} \gamma^{\nu} - \frac{1}{\ell+K_1} \gamma^{\nu} \frac{1}{\ell} \gamma^{\rho}\right)\right] = 0. \tag{65}$$

This is because, first, note that $A_{\mu} = (A_0, A_1, A_2, A_3, 0, \dots, 0)$, thus always $\gamma_5 \gamma^{\mu} = -\gamma_5 \gamma^{\mu}$ in (65). Then, in the first term, exchange (ρ, k_2) and (ν, k_1) with each other, then, shift l_{μ} as $l_{\mu} \to l_{\mu} - k_{2\mu}$. As a result, the first term can coincide with the second term with the opposite sign.

With (93a), (64) can be given as follows:

$$(64) = -2i \int \frac{d^{n}l}{i(2\pi)^{n}} \frac{\operatorname{tr}\left[\mathcal{I}(\not r\gamma_{5})(\mathcal{I} + \mathcal{K}_{2})\gamma^{\rho}(\mathcal{I} + \mathcal{K}_{1})\gamma^{\nu}\right]}{l^{2}(l + K_{2})^{2}(l + K_{1})^{2}} = -4i \int_{0}^{1} y \, dy \int_{0}^{1} dx \int \frac{d^{n}l}{i(2\pi)^{n}} \frac{\operatorname{tr}\left[(\gamma_{5}\not r)(\mathcal{I} + \mathcal{K}_{2})\gamma^{\rho}(\mathcal{I} + \mathcal{K}_{1})\gamma^{\nu}\mathcal{I}\right]}{\Omega^{3}}, \qquad (66)$$
$$\Omega \equiv -l^{2} + 2\Delta l + \Sigma^{2}, \quad \Delta^{\mu} \equiv -(xK_{1} - (1 - x)K_{2})^{\mu}y, \quad \Sigma^{2} \equiv -(x(K_{1})^{2} + (1 - x)(K_{2})^{2})y,$$

where, as for how to take Δ^{μ} and Σ^2 , there are other ways than the one above, but any ways do not affect the final result. The divergence-order at (63) is 1, however, it has been changed to 2 as can be seen in (64) or (66) as a result of the implementation of the dimensional regularization (supposing n = 4). The same thing happens in the calculation of (59e).

Using (61) in the numerator of (66), it is given maximally to the third-order of $\not r$ as follows:

$$(66) = -4i \int_{0}^{1} y \, dy \int_{0}^{1} dx \int \frac{d^{n}l}{i(2\pi)^{n}} \frac{\operatorname{tr}\left[(\gamma_{5} \not r)\{\overbrace{(\cdots)}^{\prime} \not r^{1} + \overbrace{(\cdots)}^{\prime} \not r^{0} + \overbrace{(\cdots)}^{\prime} \not r^{2} + \overbrace{(\cdots)}^{\prime} \not r^{3}\}\right]}{\Omega^{3}} \to (76). \quad (67)$$

Each $(\gamma_5 \not r)(\cdots) \not r^p$ (p = 0, 1, 2, 3) includes a product of six γ -matrices. Because of tr $[\gamma_5 \gamma^{\nu} \gamma^{\rho} \gamma^{\alpha} \gamma^{\beta}] = -4i \epsilon^{\nu \rho \alpha \beta}$, two of them should be contracted. Based on this, we evaluate each term in what follows.

• tr $[(\gamma_5 \not r) \{ (\cdots) \not r^1 \}]$ is concretely given as follows:

 $r^{2} \operatorname{tr}[\gamma_{5}\{\gamma^{\rho}(\underline{\ell}+\underline{k}_{1})\gamma^{\nu}\underline{\ell}+(\underline{\ell}+\underline{k}_{2})\gamma^{\rho}\gamma^{\nu}\underline{\ell}+(\underline{\ell}+\underline{k}_{2})\gamma^{\rho}(\underline{\ell}+\underline{k}_{1})\gamma^{\nu}\}]+r_{\alpha}v^{\alpha}r_{\beta}\operatorname{tr}[\gamma_{5}(\cdots)^{\beta\rho\nu}], (68)$

where

- The first term is the contribution by two $\not r$ contracting, where $\not r = r^2$ (the source of those two $\not r$ are that in $(\gamma_5 \not r)$ and that in $\{(\cdots) \not r^1\}$).
- The second term is the not the contribution by two $\not/$ contracting.
 - * v^{α} means some vector other than r^{α} , which is some of K_i (i = 1, 2, 3) or \underline{l}^{α} . Therefore, we can denote v^{α} as \underline{v}^{α} in the same fashion as \underline{l}_{μ} in (61).
 - * Since the second term is finally vanished no matter what $v_{\alpha} \operatorname{tr}[\gamma_5(\cdots)^{\beta\rho\nu}]$ is (as shown in (77)), we do not need to concretely write $v^{\alpha} \operatorname{tr}[\gamma_5(\cdots)^{\beta\rho\nu}]$.
- Each term consists of four γ -matrices by the result that two γ -matrices have been contracted.

The first three terms in (68) can be given as follows:

$$r^{2} \operatorname{tr} \left[\gamma_{5} \gamma^{\rho} (\underline{\ell} + \underline{K}_{1}) \gamma^{\nu} \underline{\ell}\right] = r^{2} \operatorname{tr} \left[\gamma_{5} \gamma^{\rho} \gamma^{\alpha} \gamma^{\nu} \gamma^{\beta}\right] (\underline{\ell} + \underline{K}_{1})_{\alpha} \underline{l}_{\beta}, \tag{69a}$$

$$r^{2} \operatorname{tr} \left[\gamma_{5} \left(\underline{\ell} + \underline{K}_{2}\right) \gamma^{\rho} \gamma^{\nu} \underline{\ell}\right] = r^{2} \operatorname{tr} \left[\gamma_{5} \gamma^{\alpha} \gamma^{\rho} \gamma^{\nu} \gamma^{\beta}\right] \left(\underline{\ell} + \underline{K}_{2}\right)_{\alpha} \underline{l}_{\beta}, \tag{69b}$$

$$r^{2}\operatorname{tr}\left[\gamma_{5}\left(\underline{\ell}+\underline{k}_{2}\right)\gamma^{\rho}(\underline{\ell}+\underline{k}_{1})\gamma^{\nu}\right] = r^{2}\operatorname{tr}\left[\gamma_{5}\gamma^{\alpha}\gamma^{\rho}\gamma^{\beta}\gamma^{\nu}\right](\underline{\ell}+\underline{k}_{2})_{\alpha}(\underline{\ell}+\underline{k}_{1})_{\beta}.$$
(69c)

Summing up (69a)-(69c),

(68) =
$$-r^2 \operatorname{tr} \left[\gamma_5 \gamma^\beta \gamma^\rho \gamma^\alpha \gamma^\nu\right] K_{1\beta} K_{2\alpha} + r_\alpha v^\alpha r_\beta \operatorname{tr} \left[\gamma_5 (\cdots)\right],$$
 (70)

where $K_{1\beta}K_{2\alpha}$ can be given as $k_{1\beta}k_{2\alpha}$ using the fact tr $[\gamma_5\gamma^\beta\gamma^\rho\gamma^\alpha\gamma^\nu] = -4i\,\epsilon^{\beta\rho\alpha\nu}$ at $n \to 4$.

• tr $[(\gamma_5 \not r) \{(\cdots) \not r^0\}]$ is concretely given and vanished as

$$\int \frac{d^n l}{i(2\pi)^n} \frac{\operatorname{tr}\left[(\gamma_5 \not r) \{(\not K_2) \, \gamma^{\rho}(\not K_1) \, \gamma^{\nu} \not \underline{\ell}\,\}\right]}{\Omega^3} = \operatorname{tr}\left[(\gamma_5 \gamma^{\alpha})(\not K_2 \gamma^{\rho} \not K_1 \gamma^{\nu} \gamma^{\beta})\right] \int \frac{d^n l}{i(2\pi)^n} \, \frac{r_{\alpha} \, \underline{l}_{\beta}}{\Omega^3} \xrightarrow[n \to 4]{} 0.$$
(71)

The result of (71) can be obtained by dividing the middle equation into two parts with $r_{\alpha} \underline{l}_{\beta} = (l_{\alpha} - l_{\alpha}) l_{\beta}$, then, using the formulae:

$$\int \frac{d^n k}{i(2\pi)^n} \frac{k_{\alpha} \underline{k}_{\beta}}{\mathcal{B}^3} = \frac{1}{(4\pi)^\eta \Gamma(3)} \left(\overbrace{\Gamma(3-\eta)}^{\text{finite}} \underbrace{\frac{\partial p_{\alpha} \underline{p}_{\beta}}{\mathcal{C}^{3-\eta}}}_{\Gamma(2-\eta)} - \overbrace{\Gamma(2-\eta)}^{\text{divergent}} \right), \quad \eta \equiv \frac{n}{2}, \tag{72a}$$

$$\int \frac{d^n k}{i(2\pi)^n} \frac{\underline{k}_{\alpha} \underline{k}_{\beta}}{\mathcal{B}^3} = \frac{1}{(4\pi)^\eta \,\Gamma(3)} \left(\Gamma(3-\eta) \frac{\underline{p}_{\alpha} \underline{p}_{\beta}}{\mathcal{C}^{3-\eta}} - \Gamma(2-\eta) \frac{\underline{g}_{\alpha\beta}}{2\mathcal{C}^{2-\eta}} \right),\tag{72b}$$

for those parts, respectively. These were obtained from (93b) by multiplying those by $\theta_{\mu\alpha}\underline{\theta}_{\nu\beta}$ and $\underline{\theta}_{\mu\alpha}\underline{\theta}_{\nu\beta}$, respectively ($\underline{\theta}_{\alpha\beta} \equiv \text{diag}_{n-\text{dim.}}(1,1,1,1,0,\cdots,0)$). $\underline{p}_{\alpha} \equiv \underline{\theta}_{\alpha\beta}p^{\beta}$ and $\underline{g}_{\alpha}{}^{\beta} \equiv \underline{\theta}_{\alpha\gamma}g^{\gamma\beta}$. $(\theta_{\gamma\beta}\underline{g}_{\alpha}{}^{\beta} = \underline{\theta}_{\gamma\beta}\underline{g}_{\alpha}{}^{\beta} = \underline{g}_{\alpha\gamma})$.

- The finite parts in the $l_{\alpha}l_{\underline{\beta}}$ and $l_{\underline{\alpha}}l_{\underline{\beta}}$ terms always cancel each other at $n \to 4$ as $p_{\alpha} \xrightarrow[n \to 4]{} \underline{p}_{\alpha}$. This is understandable as these are originally terms in a term (64) proportional to $\not r$.
- In the divergent part, $\Gamma(2-\eta)$ is diverged at $n \to 4$. However, for the case that the numerator is not contracted, the numerator is 0, and there is no contribution for arbitrary n. On the other hand, even in the case that the numerator is contracted, the divergent parts in (72a) and (72b) are the same each other, and its difference is 0 for arbitrary n.

- The existence of the divergent part is caused in the UV-divergence of (59c). Conversely, if the UV-divergence were not in (59c), such a divergent part would not appear. Therefore, such divergent part does not appear in (59f).
- tr $[(\gamma_5 \not r) \{(\cdots) \not r^2\}]$ exists totally $_3C_2 = 3$ patterns, and vanish at $n \to 4$ as follows:

$$\int \frac{d^n l}{i(2\pi)^n} \frac{\operatorname{tr}\left[(\gamma_5 \not r)(\not r) \gamma^{\rho}(\not r) \gamma^{\nu} \not \underline{\ell}\right]}{\Omega^3} = \operatorname{tr}\left[\gamma_5 \gamma^{\rho} \gamma^{\alpha} \gamma^{\nu} \gamma^{\beta}\right] \int \frac{d^n l}{i(2\pi)^n} \frac{r^2 r_{\alpha} \underline{l}_{\beta}}{\Omega^3} \xrightarrow[n \to 4]{} 0, \tag{73a}$$

$$\int \frac{d^n l}{i(2\pi)^n} \frac{\operatorname{tr}\left[(\gamma_5 \not r)(\not r) \gamma^{\rho}(\not \underline{l} + \not K_1) \gamma^{\nu} \not r\right]}{\Omega^3} = \operatorname{tr}\left[\gamma_5 \gamma^{\rho} \gamma^{\alpha} \gamma^{\nu} \gamma^{\beta}\right] \int \frac{d^n l}{i(2\pi)^n} \frac{r^2 (\not \underline{l} + K_1)_{\alpha} r_{\beta}}{\Omega^3} \xrightarrow[n \to 4]{} 0, \quad (73b)$$

$$\int \frac{d^n l}{i(2\pi)^n} \frac{\operatorname{tr}\left[(\gamma_5 \not r)(\not l + \not K_2) \,\gamma^{\rho}(\not r) \,\gamma^{\nu} \not r\right]}{\Omega^3} = \operatorname{tr}\left[\gamma_5 \gamma^{\alpha} \gamma^{\rho} \gamma^{\nu} \gamma^{\beta}\right] \int \frac{d^n l}{i(2\pi)^n} \frac{r^2 \,(\not l + K_2)_{\alpha} \,r_{\beta}}{\Omega^3} \xrightarrow[n \to 4]{} 0, \quad (73c)$$

where $r^2 r_{\alpha} \underline{l}_{\beta} = l^2 (l_{\alpha} \underline{l}_{\beta} - \underline{l}_{\alpha} \underline{l}_{\beta}) - \underline{l}^2 (l_{\alpha} \underline{l}_{\beta} - \underline{l}_{\alpha} \underline{l}_{\beta})$ and $r^2 r_{\beta} = l^2 (l - \underline{l})_{\beta} - \underline{l}^2 (l - \underline{l})_{\beta}$, and the results of (73a)-(73c) can be obtained with the following formulae obtainable from (93c) and (93d):

$$\int \frac{d^n k}{i(2\pi)^n} \frac{k^2 k_{\alpha} \underline{k}_{\beta}}{\mathcal{B}^3} = \frac{1}{\mathcal{D}} \Big(\Gamma(3-\eta) \frac{p^2 \underline{p}_{\alpha} \underline{p}_{\beta}}{\mathcal{C}^{3-\eta}} - \Gamma(2-\eta) \frac{(n+4) \underline{p}_{\alpha} \underline{p}_{\beta} + p^2 \underline{g}_{\alpha\beta}}{2 \mathcal{C}^{2-\eta}} + \Gamma(1-\eta) \frac{(n+2) \underline{g}_{\alpha\beta}}{4 \mathcal{C}^{1-\eta}} \Big),$$
(74a)

$$\int \frac{d^n k}{i(2\pi)^n} \frac{k^2 k_{\underline{\alpha}}}{\mathcal{B}^3} = \frac{1}{\mathcal{D}} \Big(\Gamma(3-\eta) \frac{p^2 \underline{p}_{\underline{\alpha}}}{\mathcal{C}^{3-\eta}} - \Gamma(2-\eta) \frac{(n+2) \underline{p}_{\underline{\alpha}}}{2 \mathcal{C}^{2-\eta}} \Big), \tag{74b}$$

where $p_{\alpha}\underline{p}_{\beta} = \underline{p}_{\alpha}\underline{p}_{\beta}$ (we skip to give other necessary formulae).

• tr $[(\gamma_5 \not r) \{(\cdots) \not r^3\}]$ is concretely given as and can be known to be vanished as

$$\operatorname{tr}\left[(\gamma_5 \not r)(\not r) \gamma^{\rho}(\not r) \gamma^{\nu} \not r\right] = \operatorname{tr}\left[(\gamma_5) \gamma^{\rho} \gamma^{\nu}\right] (r^2)^2 = 0.$$
(75)

Assembling the results of (70), (71), (73) and (75), $ik_{3\mu}\widetilde{\Gamma}_{\mathcal{A}}^{(3)}$ in (67) can be given as follows:

$$ik_{3\mu}\widetilde{\Gamma}_{\mathcal{A}}^{(3)} = -4i\int_{0}^{1} y\,dy\int_{0}^{1} dx\int \frac{d^{n}l}{i(2\pi)^{n}\Omega^{3}} \Big(-\operatorname{tr}\left[\gamma_{5}\gamma^{\beta}\gamma^{\rho}\gamma^{\alpha}\gamma^{\nu}\right]k_{1\beta}k_{2\alpha}\left(l^{2}-\underline{l}^{2}\right) + \operatorname{tr}\left[\gamma_{5}(\cdots)\right]v^{\alpha}r_{\alpha}r_{\beta}\Big).$$
(76)

The integrates in the one above result in as follows:

t

$$\int_{0}^{1} y \, dy \int_{0}^{1} dx \int \frac{d^{n}l}{i(2\pi)^{n}} \frac{l^{2} - \underline{l}^{2}}{\Omega^{3}} = \int_{0}^{1} y \, dy \int_{0}^{1} dx \frac{1}{(4\pi)^{\eta} \Gamma(3)} \Big(\frac{\Gamma(3-\eta)(p^{2}-\underline{p}^{2})}{(\Delta^{2}+\Sigma^{2})^{3-\eta}} - \frac{\Gamma(2-\eta)(n-4)}{2(\Delta^{2}+\Sigma^{2})^{2-\eta}} \Big) \xrightarrow[n \to 4]{} \int_{0}^{1} y \, dy \int_{0}^{1} dx \frac{1}{32\pi^{2}} = \frac{1}{64\pi^{2}},$$
(77a)

$$\int \frac{d^{n}l}{i(2\pi)^{n}} \frac{v^{\alpha} r_{\alpha} r_{\beta}}{\Omega^{3}} = -\frac{\Gamma(2-\eta)}{(4\pi)^{\eta} \Gamma(3)} \frac{v^{\alpha}(g_{\alpha\beta} - \underline{g}_{\alpha\beta})}{2(\Delta^{2} + \Sigma^{2})^{2-\eta}} = -\frac{\Gamma(2-\eta)}{(4\pi)^{\eta} \Gamma(3)} \frac{v_{\beta} - \underline{v}_{\beta}}{2(\Delta^{2} + \Sigma^{2})^{2-\eta}} = 0.$$
(77b)

(77a) can be known from (72). $n = g^{\mu}{}_{\mu}$ and $4 = \underline{g}^{\mu}{}_{\mu}$, and n - 4 converges to 0 at $n \to 4$, which cancels the divergence from $\Gamma(2 - \eta)$, and its coefficient remains. In (77b), since v_{β} can be denoted as \underline{v}_{β} as mentioned under (68), $v_{\beta} - \underline{v}_{\beta}$ is vanished for arbitrary n. Finally, $ik_{\mu}\Gamma^{(3)}_{\mathcal{A}}$ results in as follows:

$$ik_{3\mu}\Gamma^{(3)}_{\mathcal{A}} = \epsilon^{\beta\rho\alpha\nu}k_{1\beta}k_{2\alpha}/4\pi^2.$$
(78)

11.5 The calculations of terms other than (59c)

We turn to the terms other than (59c), these are (59a), (59b), (59d), (59e) and (59f).

- It can be seen immediately that (59a) is vanished because of $\delta^4(k_1) k_{1\mu}$ in that equation.
- As for (59b), its calculation can be proceeded in the same way as that in Sec.11.4, and the following *l*-integrate appears at the stage corresponding to (66):

Four γ -matrices, which becomes less than four after contractions

$$\int \frac{d^{n}l}{i(2\pi)^{n}} \frac{\operatorname{tr}\left[\mathcal{I}(\not r\gamma_{5})(\mathcal{I}+\not K_{1})\gamma^{\rho}\right]}{l^{2}(l+K_{1})^{2}} = \int \frac{d^{n}l}{i(2\pi)^{n}} \frac{\operatorname{tr}\left[\mathcal{I}(\not r\gamma_{5})\mathcal{I}\gamma^{\rho}+\mathcal{I}(\not r\gamma_{5})\not K_{1}\gamma^{\rho}\right]}{l^{2}(l+K_{1})^{2}}.$$
(79)

f is replaced with (61). Then, in order for *l*-integrates to yield finite contributions at $n \to 4$, all f and \underline{f} should be contracted in its numerator (for this, see the points under (72)). As a result, the number of γ -matrices becomes less than four in each term. However, four γ -matrices are needed for tr[$\gamma_5 \cdots$] to remain finitely. From these, it can be seen (79) is vanished at $n \to 4$.

• The calculation of (59d) can be proceeded in the same way as that in Sec.11.4, and it is finally turned to be vanished at $n \to 4$. This can be seen from the equation corresponding to (66):

Four $\gamma\text{-matrices}$ are needed for this to remain

$$\int \left(\prod_{i=1}^{4} \frac{d^4 k_i}{i(2\pi)^4}\right) (2\pi)^4 \,\delta^4(K_4) \left(c(-k_4) \prod_{i=1}^{3} A_{\mu_i}(-k_i)\right) \int \frac{d^4 l}{i(2\pi)^4} \frac{\operatorname{tr}\left[\mathcal{I}\left(\not\!\!\!/ \gamma_5\right) \prod_{i=3}^{4} \left(\mathcal{I} + \not\!\!\!/ K_i\right) \gamma^{\mu_i}\right]}{l^2 (l+K_3)^2 (l+K_2)^2 (l+K_1)^2}.$$
 (80)

As noted above, some four γ -matrices are needed. Three γ -matrices have been already appeared, one γ -matrix should be appeared from the rest part by some contractions:

$$\operatorname{tr}\left[\not l \left(\not \gamma_{5} \right) \prod_{i=3}^{1} (\not l + \not K_{i}) \overbrace{\gamma^{\mu_{i}}}^{\text{Three } \gamma \text{-matrices already}} \right].$$
(81)

The form to be finally realized is $v_{\mu_4} \prod_{i=1}^3 A_{\mu_i}(-k_i) \epsilon^{\mu_4 \mu_3 \mu_2 \mu_1} (v_{\mu_4} \text{ is } k_{i\mu}, (k_i+k_j)_{\mu} \text{ or } (k_i+k_j+k_k)_{\mu}).$ By the flip $(\mu_i, k_i) \leftrightarrow (\mu_j, k_j)$ (i, j = 1, 2 or 3), this can be seen to be vanished.

- From the counting of the divergent-order, the term in (59e) can be regarded as no divergent. However, by the situation mentioned under (66), it includes some possibility to be divergent. However, in (59e), four γ -matrices have already appeared, and the form to finally realize is $\prod_{i=1}^{4} A_{\mu_i}(-k_i) \, \epsilon^{\mu_4 \mu_3 \mu_2 \mu_1}$. By the flip $(\mu_i, k_i) \leftrightarrow (\mu_j, k_j)$, this can be seen to be vanished.
- In the *n*-dimensional calculation for each term in (59f), at the stage corresponding to (64), the term irrelevant to $\not r$ is vanished, and the term proportional to $\not r$ remains, as well as (64). Since terms in (59f) are not divergent, that $\not r$ proportional term yields only finite contributions, and is vanished at $n \to 4$ as $\not r = 0$ at $n \to 4$.

11.6 The result of (59)

With (78) and the results in Sec. 11.5, (59) is finally given as follows:

$$\int d^4x \, \boldsymbol{\delta}_{\varepsilon} \Gamma_{\mathcal{A}}[A] = i \int \left(\prod_{i=1}^3 \frac{d^4 k_i}{(2\pi)^4}\right) (2\pi)^4 \, \delta^4(K_3) \, \varepsilon(-k_3) \, c(-k_3) \, A_{\rho}(-k_1) A_{\nu}(-k_2) \, \frac{e^2 \epsilon^{\beta \rho \alpha \nu} k_{1\beta} k_{2\alpha}}{4\pi^2}$$
$$= -\int d^4x \, \varepsilon(x) c(x) \, \frac{i e^2 \epsilon^{\beta \rho \alpha \nu}}{4\pi^2} \partial_{\beta} A_{\rho}(x) \partial_{\alpha} A_{\nu}(x) = -\int d^4x \, \varepsilon(x) c(x) \, \frac{i e^2 \epsilon^{\beta \rho \alpha \nu}}{16\pi^2} F_{\beta \rho}(x) F_{\alpha \nu}(x), \tag{82}$$

where $A_{\mu}(k) = \int d^4k e^{ikx} A_{\mu}(x)$ and $\varepsilon(k)c(k) = \int d^4k e^{ikx}\varepsilon(x)c(x)$, the inverse of those in Sec. 11.1, and $\int d^4k e^{-ikx} = (2\pi)^4 \delta^4(x)$. Applying this result to (57), the identity agreeing with (46) can be obtained:

$$\varepsilon \, i \, \frac{e^3}{16\pi^2} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma} = \partial_\mu \varepsilon \, \langle J_c^\mu \rangle - \partial_\mu \langle J_{c\epsilon}^\mu \rangle. \tag{83}$$

12 The anomaly equation by the Fujikawa method

The anomaly equation associated with the large chiral symmetry was obtained from the heuristically constituted large chiral charge in Sec. 10 and the evaluation of the BRS transformed one-loop diagrams in and Sec. 11. In this section, we derive it by the Fujikawa method.

Under the large chiral transformation (37) in the Cartesian coordinates, the variations of action (1) and the measure of the Dirac field are given as

$$\delta S_0 = \int d^4x \, i(\bar{\psi}' \not\!\!\!D\psi' - \bar{\psi} \not\!\!D\psi) = e \int d^4x \, \partial_\mu(\lambda \varepsilon) \, \bar{\psi} \gamma^\mu \gamma_5 \psi = \int d^4x \, \lambda \left(\partial_\mu(\varepsilon J_c^\mu) - (\partial_\mu \varepsilon) J_c^\mu \right), \tag{84a}$$

$$\mathcal{D}\bar{\psi}'\mathcal{D}\psi' = \exp\left[-2i\int d^4x \,e\lambda\varepsilon\,\mathcal{A}\right]\mathcal{D}\bar{\psi}\mathcal{D}\psi, \quad \mathcal{A} \equiv \lim_{\Lambda \to \infty} (\sum_{n=-\infty}^{\infty} \varphi_n^{\dagger}\,\gamma_5 \,e^{-(\eta_n/\Lambda)^2}\varphi_n) = -\frac{e^2}{32\pi^2}\,\epsilon^{\mu\nu\rho\sigma}\,F_{\mu\nu}F_{\rho\sigma},$$
(84b)

where λ has been taken to the linear-order and supposed to vanish at $r \to \infty$. J_c^{μ} and $J_{c\varepsilon}^{\mu}$ are defined in (32). φ_n and φ_n^{\dagger} are the eigenvectors as $\not{D}\varphi_n = \eta_n\varphi_n$, and form a complete orthogonal set: $\int d^4x \, \varphi_m^{\dagger}\varphi_n = \delta_{mn}$. With those, ψ and $\bar{\psi}$ have been expanded. $e^{-(\eta_n/\Lambda)^2}$ have the cut-off role for $\eta_n \gg \Lambda$. The calculation of \mathcal{A} in (84b) is well-known, thus we have noted just the results.

Giving $Z_{\psi,A} \equiv \int \mathcal{D}\bar{\psi}\mathcal{D}\psi\mathcal{D}A e^{iS_0}$ with (84a) and (84b),

$$Z_{\psi',A'} = \int \mathcal{D}\bar{\psi}\mathcal{D}\psi\mathcal{D}A \exp\left[\int d^4x \left\{ \mathcal{L}_0 + \lambda \left(\partial_\mu (\varepsilon J_c^\mu) - (\partial_\mu \varepsilon) J_c^\mu + \varepsilon \frac{ie^3}{16\pi^3} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma} \right) \right\} \right].$$
(85)

From the invariance $Z_{\psi',A'} = Z_{\psi,A}$, the following anomaly equation can be obtained:

$$\partial_{\mu}\langle J^{\mu}_{c\varepsilon}\rangle - \partial_{\mu}\varepsilon \langle J^{\mu}_{c}\rangle + \varepsilon \frac{ie^{3}}{16\pi^{2}} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu}F_{\rho\sigma} = 0, \qquad (86)$$

where $\varepsilon J_c^{\mu} \equiv J_{c\varepsilon}^{\mu}$ as seen in (32). We can see this agrees with (46) and (83).

Taking the limit of Λ prior to the summation of n in the middle equation of \mathcal{A} , it is given as $\sum_{n} \varphi_{n}^{\dagger} \gamma_{5} \varphi_{n}$. Then, $\mathcal{D} \gamma_{5} \varphi_{n} = -\eta_{n} \gamma_{5} \varphi_{n}$, thus, $\gamma_{5} \varphi_{n}$ are the eigenvectors associated with the eigenvalues $-\lambda_{n}$, and φ_{n} and $\gamma_{5} \varphi_{n}$ are orthogonal each other. From this, $\int d^{4}x \mathcal{A}$ results in $n_{+} - n_{-}$ (n_{\pm} mean the number of the zero-mode particles with each positive/negative chirality, respectively)[¶].

13 The anomaly equation for the large U(1) symmetry

In this study, we have studied the anomaly equation for the large chiral symmetry. However, we can consider the anomaly equation for the large U(1) symmetry from (53) as

$$\boldsymbol{\delta}_{\varepsilon} \Gamma[A] = -e^{-1} c \left(\partial_{\mu} \varepsilon \left\langle J_{\mathrm{U}(1)}^{\mu} \right\rangle - \partial_{\mu} \left\langle J_{\varepsilon}^{\mu} \right\rangle \right). \tag{87}$$

In this section, we comment on this.

[¶]Since \not{D} and γ_5 can be simultaneously diagonalized, $\int d^4x \, \mathcal{A} = \sum_n \int d^4x \, (\varphi_n^{(+)\dagger} \, \varphi_n^{(-)} - \varphi_n^{(-)\dagger} \, \varphi_n^{(-)}) = \int d^4x \, (\varphi_0^{(+)\dagger} \, \varphi_0^{(+)} - \varphi_n^{(-)\dagger} \, \varphi_n^{(-)})$, where $\gamma_5 \varphi_n = \pm \varphi_n^{(\pm)}$, and by the orthogonality between φ_n and $\gamma_5 \varphi_n$, contributions for $n \neq 0$ vanish.

We can see that $\delta_{\varepsilon}\Gamma[A]$ is vanished. This is because, proceeding the calculation of it in the same way as (57)-(64), at the stage of (64), the term not proportional to \not vanishes for the BRS symmetry and the term proportional to \not does not exist from the beginning as γ^5 does not exist in the large U(1) current. Therefore, Thus, (87) is given as $0 = \partial_{\mu} \varepsilon \langle J^{\mu}_{U(1)} \rangle - \partial_{\mu} \langle J^{\mu}_{\varepsilon} \rangle$.

Certainly, this equation is telling about $\partial_{\mu} \langle J_{\varepsilon}^{\mu} \rangle$ and indicates that J_{ε}^{μ} is not conserved when $\langle J_{U(1)}^{\mu} \rangle \neq 0$. 0. However, this would not be said as anomaly equation. This is because this equation is irrelevant of the breaking of symmetry. Therefore, we have not addressed (87) in this study.

14 Summary

In this study, we proposed the large chiral symmetry based on the large U(1) gauge symmetry, and gave the anomaly equation of that. Then, defining the BRS transformation for the large U(1) gauge symmetry, based on the BRS transformed effective action, we gave the anomaly equation concering the large chiral symmetry given with the BRS transformed one-loop diagrams. Then, evaluating those, we showed that the anomaly equation of the large chiral symmetry can be derived. We also showed that it can be derived by Fujikawa method. The results in this study would be important as a development of the large U(1) gauge symmetry.

The fermionic field in this study was massless. If it were massive, it all starts off from \mathcal{I}^- and reaches \mathcal{I}^+ in Fig.1. In that case, the way to define the soft-charges is changed from the massless case which we performed in Sec. 7. For this reason, this study did not address the massive Dirac field.

As a development of this study, constructing the low-energy effective model (from the condition that it can reproduce the original model's anomaly) and calculating the decay-width of the NG bosons of the symmetry breaking associated with the large anomaly would be taken up.

The way to obtain the effective model for the case of $SU_{color}(3)$ QCD with regard to u, d and s quarks (those are taken as massless) and with the anomaly associated with $SU_L(3) \times SU_R(3) \rightarrow SU_v(3)$ has been already known, which is to solve a kind of differential equation with regard to the effective action obtained from (53) (the effective action to be solved in this differential equation is called Wess-Zumino-Witten term). The decay-widths of the NG bosons associated with the symmetry breaking above, such as $\pi^0 \rightarrow 2\gamma$, etc, are calculable within the effective model. Agreements of them with the experimental data can be considered good^{||}.

This project will be performed applying this way, taking the large $SU_{color}(3)$. However, there is the extra term in the anomalies associated with the large chiral symmetries, which is the first term in (46). Therefore, in this project, the differential equation of the effective action mentioned above should be improved in such a way that it can yield that extra term, which will be a new point in this project. If we could eventually carry out these and calculate decay-widths, it would be interesting in terms of the observation of a phenomenology of the large gauge theory.

Other than the one above, we might consider the fermion coupling to the gravity (the fermion in this study is coupling to the U(1) gauge field). However, the gravity in the asymptotic symmetry is a problem concerning how the spacetime is curved. Therefore, since the anomaly equation is covariant, it could be obtained just with the general coordinate transformation for an anomaly equation calculated on the flat spacetime.

A The derivation of (7)

We first examine the variation of the action for the following general coordinate and field transformations:

^{$\|$}For $\pi^0 \to 2\gamma$, $\Gamma_{\rm th} \simeq 7.64 {\rm eV}$ while $\Gamma_{\rm exp} \simeq 7.69 \pm 0.12 {\rm eV}$, where $\Gamma_{\rm th}$ is the calculated value and $\Gamma_{\rm exp}$ is obtained from [58]. Other decay-widths can also be calculated, which will be presented in the future paper with the description to give the effective model, etc.

$$x^{\mu} \to x^{\prime \mu} = x^{\mu} + \delta x^{\mu}, \quad \phi_A(x) \to \phi_A'(x^{\prime}) = \phi_A(x) + \delta \phi_A(x), \tag{88}$$

where $\phi_A(x)$ is that in Sec. 3.2. δx^{μ} and $\delta \phi_A$ are supposed to be the quantities to the liner-order. The variation of the action under (88) is known to be computed as follows:

$$\delta I = \int_{\Omega'} d^4 x' L(\phi'_A(x'), \phi'_{A,\mu}(x')) - \int_{\Omega} d^4 x L(\phi_A(x), \phi_{A,\mu}(x))$$

=
$$\int_{\Omega} d^4 x \left(\frac{\partial L}{\partial \phi_A} \bar{\delta} \phi_A + \frac{\partial L}{\partial \phi_{A,\mu}} \partial_\mu (\bar{\delta} \phi_A) + \partial_\mu (L \, \delta x^\mu) \right),$$
(89)

where $d^4x' = \left|\frac{\partial x'^{\mu}}{\partial x^{\nu}}\right| d^4x = \left(1 + \frac{\partial \delta x^{\mu}}{\partial x^{\mu}}\right) d^4x$ and $\phi'_{A,\mu}(x') = \frac{\partial x^{\nu}}{\partial x'^{\mu}} \frac{\partial}{\partial x^{\nu}} \phi'_A(x') = \partial_{\mu} \phi_A(x) + \partial_{\mu} (\delta \phi_A(x)) - (\partial_{\mu} \delta x^{\nu}) \partial_{\nu} \phi_A(x)$. $\bar{\delta} \phi_A(x)$ means the Lie derivative for ϕ_A , which is defined as $\bar{\delta} \phi_A(x=a) \equiv \phi'_A(x'=a) - \phi_A(x=a)$, and $\bar{\delta} \phi_A$ can be written as $\bar{\delta} \phi_A = \delta \phi_A - \partial_{\mu} \phi_A \delta x^{\mu}$. Finally, δI can be given as

$$\delta I = \int_{\Omega} d^4 x \left([L]^A \bar{\delta} \phi_A + \partial_\mu \mathcal{J}^\mu \right), \quad \mathcal{J}^\mu \equiv \frac{\partial L}{\partial \phi_{A,\mu}} \delta \phi_A - T^\mu{}_\nu \,\delta x^\nu, \tag{90}$$

where $[L]^A$ and $T^{\mu}{}_{\nu}$ are defined in (2) and (4), respectively.

Applying the local transformation (6) to (90),

$$\delta I|_{(6)} = \int_{\Omega} d^4 x \left\{ \lambda^r \Big([L]^A (M_{r,A} - \phi_{A,\mu} X^{\mu}{}_r) - \partial_{\mu} ([L]^A N_r{}^{\mu}{}_{,A}) \Big) + \partial_{\mu} \Big(\mathcal{J}^{\mu} + [L]^A N_r{}^{\mu}{}_{,A} \lambda^r \Big) \right\},$$
(91)

where J_r^{μ} is defined in (4). Writing out λ^r included in $\delta \phi_A$ and δx^{ν} with (6) in the second part,

$$\partial_{\mu}(B^{\mu}{}_{r}\lambda^{r} + C^{\mu\nu}{}_{r}\lambda^{r}{}_{,\nu}) = \partial_{\mu}B^{\mu}{}_{r}\lambda^{r} + (B^{\mu}{}_{r} + \partial_{\nu}C^{\nu\mu}{}_{r})\lambda^{r}{}_{,\mu} + C^{\mu\nu}{}_{r}\lambda^{r}{}_{,\mu\nu}, \qquad (92)$$

and (7) can be obtained. (On the other hand, it can be known from the first part that there are n equations $(r = 1, \dots, n)$ among N EL equations for ϕ_A $(A = 1, \dots, N)$. This means that the EL equations are N - n missing for determining $N \phi_A$, which leads to the issue of the constrained system.)

B The formulas used in Sec. 11

$$\frac{1}{ABC} = 2 \int_0^1 y \, dy \int_0^1 dx \, \frac{1}{(Ayx + By(1 - x) + C(1 - y))^3},\tag{93a}$$

$$\int \frac{d^n k}{i(2\pi)^n} \frac{k^{\mu} k^{\nu}}{\mathcal{B}^{\alpha}} = \frac{1}{\mathcal{D}} \left(\Gamma(\bar{\alpha}) \frac{p^{\mu} p^{\nu}}{\mathcal{C}^{\bar{\alpha}}} - \Gamma(\bar{\alpha} - 1) \frac{g^{\mu\nu}}{2\mathcal{C}^{\bar{\alpha} - 1}} \right),\tag{93b}$$

$$\int \frac{d^n k}{i(2\pi)^n} \frac{k^{\mu} k^{\nu} k^{\rho}}{\mathcal{B}^{\alpha}} = \frac{1}{\mathcal{D}} \left(\Gamma(\bar{\alpha}) \frac{p^{\mu} p^{\nu} p^{\rho}}{\mathcal{C}^{\bar{\alpha}}} - \Gamma(\bar{\alpha} - 1) \frac{g^{(\mu\nu} p^{\rho)}}{2\mathcal{C}^{\bar{\alpha} - 1}} \right), \tag{93c}$$

$$\int \frac{d^{n}k}{i(2\pi)^{n}} \frac{k^{\mu}k^{\nu}k^{\rho}k^{\sigma}}{\mathcal{B}^{\alpha}} = \frac{1}{\mathcal{D}} \left(\Gamma(\bar{\alpha}) \frac{p^{\mu}p^{\nu}p^{\rho}p^{\sigma}}{\mathcal{C}^{\bar{\alpha}}} - \Gamma(\bar{\alpha}-1) \frac{g^{(\mu\nu}p^{\rho}p^{\sigma)}}{2\mathcal{C}^{\bar{\alpha}-1}} + \Gamma(\bar{\alpha}-2) \frac{g^{(\mu\nu}g^{\rho\sigma)}}{4\mathcal{C}^{\bar{\alpha}-2}} \right), \tag{93d}$$

where $\eta \equiv n/2$, $\mathcal{B} \equiv m^2 + 2k \cdot p - k^2$, $\mathcal{C} \equiv m^2 + p^2$, $\mathcal{D} \equiv (4\pi)^{\eta} \Gamma(\alpha)$, $\bar{\alpha} \equiv \alpha - \eta$ and

$$g^{(\mu\nu}p^{\rho)} = g^{\mu\nu}p^{\rho} + g^{\nu\rho}p^{\mu} + g^{\rho\mu}p^{\nu},$$

$$g^{(\mu\nu}p^{\rho}p^{\sigma)} = g^{\mu\nu}p^{\rho}p^{\sigma} + g^{\mu\rho}p^{\nu}p^{\sigma} + g^{\mu\sigma}p^{\nu}p^{\rho} + g^{\nu\rho}p^{\mu}p^{\sigma} + g^{\nu\sigma}p^{\mu}p^{\rho} + g^{\rho\sigma}p^{\mu}p^{\nu},$$

$$g^{(\mu\nu}g^{\rho\sigma)} = g^{\mu\nu}g^{\rho\sigma} + g^{\mu\rho}g^{\nu\sigma} + g^{\mu\sigma}g^{\nu\rho}.$$

(93a) is a Feynman parameter formula. (93b)-(93d) can be known from Sec. A.4 in [59].

References

- [1] J. Ambjorn, A. Goerlich, J. Jurkiewicz and R. Loll, Phys. Rept. 519, 127-210 (2012) [arXiv:1203.3591 [hep-th]].
- [2] R. Loll, Class. Quant. Grav. 37, no.1, 013002 (2020) [arXiv:1905.08669 [hep-th]].
- [3] R. Gambini and J. Pullin, Oxford University Press, 2011,
- [4] C. Rovelli and F. Vidotto, Cambridge University Press, 2014,
- [5] M. B. Green, J. H. Schwarz and E. Witten, 1988, M. B. Green, J. H. Schwarz and E. Witten, 1988,
- [6] J. Polchinski, Cambridge University Press, 2007, J. Polchinski, Cambridge University Press, 2007,
- [7] K. Hashimoto, Springer, 2012,
- [8] S. Nagy, Annals Phys. **350**, 310-346 (2014) [arXiv:1211.4151 [hep-th]].
- [9] R. Percacci, "An Introduction to Covariant Quantum Gravity and Asymptotic Safety," World Scientific, 2017,
- [10] G. 't Hooft, Conf. Proc. C 930308, 284-296 (1993) [arXiv:gr-qc/9310026 [gr-qc]].
- [11] L. Susskind, J. Math. Phys. 36, 6377-6396 (1995) [arXiv:hep-th/9409089 [hep-th]].
- [12] D. Bigatti and L. Susskind, [arXiv:hep-th/0002044 [hep-th]].
- [13] R. Bousso, Rev. Mod. Phys. **74**, 825-874 (2002) [arXiv:hep-th/0203101 [hep-th]].
- [14] G. Barnich and C. Troessaert, Phys. Rev. Lett. 105, 111103 (2010) [arXiv:0909.2617 [gr-qc]].
- [15] A. Strominger, [arXiv:1703.05448 [hep-th]].
- [16] J. D. Brown and M. Henneaux, Commun. Math. Phys. 104, 207-226 (1986)
- [17] H. Terashima, Phys. Rev. D 64, 064016 (2001) [arXiv:hep-th/0102097 [hep-th]].
- [18] D. Anninos, G. S. Ng and A. Strominger, Class. Quant. Grav. 28, 175019 (2011) [arXiv:1009.4730 [gr-qc]].
- [19] T. Regge and C. Teitelboim, Annals Phys. 88, 286 (1974)
- [20] G. Barnich and C. Troessaert, JHEP 05, 062 (2010) [arXiv:1001.1541 [hep-th]].
- [21] G. Barnich and C. Troessaert, JHEP 12, 105 (2011) [arXiv:1106.0213 [hep-th]].
- [22] A. Strominger, [arXiv:1706.07143 [hep-th]].
- [23] T. He, V. Lysov, P. Mitra and A. Strominger, JHEP 05, 151 (2015) [arXiv:1401.7026 [hep-th]].
- [24] M. Campiglia and A. Laddha, JHEP 12, 094 (2015) [arXiv:1509.01406 [hep-th]].
- [25] A. Laddha and A. Sen, JHEP **10**, 065 (2017) [arXiv:1706.00759 [hep-th]].
- [26] B. Sahoo and A. Sen, JHEP **02**, 086 (2019) [arXiv:1808.03288 [hep-th]].
- [27] S. Atul Bhatkar and B. Sahoo, JHEP **01**, 153 (2019) [arXiv:1809.01675 [hep-th]].

- [28] A. M. Raclariu, [arXiv:2107.02075 [hep-th]].
- [29] S. Pasterski, Eur. Phys. J. C 81, no.12, 1062 (2021) [arXiv:2108.04801 [hep-th]].
- [30] S. Pasterski, A. Strominger and A. Zhiboedov, JHEP **12**, 053 (2016) [arXiv:1502.06120 [hep-th]].
- [31] P. Mao and H. Ouyang, Phys. Lett. B 774, 715-722 (2017) [arXiv:1707.07118 [hep-th]].
- [32] M. Pate, A. M. Raclariu and A. Strominger, Phys. Rev. Lett. 119, no.26, 261602 (2017) [arXiv:1707.08016 [hep-th]].
- [33] Y. Hamada and S. Sugishita, JHEP 11, 203 (2017) [arXiv:1709.05018 [hep-th]].
- [34] C. S. Chu and Y. Koyama, Phys. Rev. D 100, no.10, 104034 (2019) [arXiv:1906.09361 [hep-th]].
- [35] K. Mitman, M. Boyle, L. C. Stein, N. Deppe, L. E. Kidder, J. Moxon, H. P. Pfeiffer, M. A. Scheel, S. A. Teukolsky and W. Throwe, *et al.* Class. Quant. Grav. **41**, no.22, 223001 (2024) [arXiv:2405.08868 [gr-qc]].
- [36] S. Sarkar, S. Kumar and S. Bhattacharjee, Phys. Rev. D 105, no.8, 084001 (2022) [arXiv:2110.03547 [gr-qc]].
- [37] F. L. Lin, A. Patel and H. Y. Pu, JHEP 09, 117 (2022) [arXiv:2202.13559 [gr-qc]].
- [38] A. Strominger, JHEP **02**, 009 (1998) [arXiv:hep-th/9712251 [hep-th]].
- [39] C. G. Callan and J. M. Maldacena, Nucl. Phys. B 472, 591-610 (1996) [arXiv:hep-th/9602043 [hep-th]].
- [40] J. M. Maldacena and L. Susskind, Nucl. Phys. B 475, 679-690 (1996) [arXiv:hep-th/9604042 [hep-th]].
- [41] M. Guica, T. Hartman, W. Song and A. Strominger, Phys. Rev. D 80, 124008 (2009) [arXiv:0809.4266 [hep-th]].
- [42] Y. Nakayama, Phys. Lett. B 673, 272-278 (2009) [arXiv:0812.2234 [hep-th]].
- [43] G. Barnich, A. Gomberoff and H. A. Gonzalez, Phys. Rev. D 86, 024020 (2012) [arXiv:1204.3288 [gr-qc]].
- [44] G. Barnich, JHEP 10, 095 (2012) [arXiv:1208.4371 [hep-th]].
- [45] G. Barnich, A. Gomberoff and H. A. González, Phys. Rev. D 87, no.12, 124032 (2013) [arXiv:1210.0731 [hep-th]].
- [46] G. Barnich and H. A. Gonzalez, JHEP 05, 016 (2013) [arXiv:1303.1075 [hep-th]].
- [47] M. Campiglia and A. Laddha, JHEP 07, 115 (2015) [arXiv:1505.05346 [hep-th]].
- [48] M. Campiglia and A. Laddha, JHEP **11**, 012 (2016) [arXiv:1605.09677 [hep-th]].
- [49] M. Campiglia and A. Laddha, JHEP 10, 287 (2019) [arXiv:1903.09133 [hep-th]].
- [50] S. Atul Bhatkar, JHEP 10, 110 (2020) [arXiv:1912.10229 [hep-th]].
- [51] S. Atul Bhatkar, JHEP **02**, 082 (2021) [arXiv:2007.03627 [hep-th]].
- [52] S. Atul Bhatkar, Phys. Rev. D 105, no.12, 124028 (2022) [arXiv:2108.00835 [hep-th]].

- [53] P. Cheng and P. Mao, Phys. Rev. D 106, no.8, L081702 (2022) [arXiv:2206.11564 [hep-th]].
- [54] N. Banerjee, K. Fernandes and A. Mitra, "Soft photon theorem in the small negative cosmological constant limit," JHEP 08, 105 (2021) [arXiv:2102.06165 [hep-th]].
- [55] K. Fernandes, N. Banerjee and A. Mitra, "Soft factors with AdS radius corrections," JHAP 3, no.4, 5-22 (2023) [arXiv:2310.19299 [hep-th]].
- [56] S. Pasterski, JHEP 09, 154 (2017) [arXiv:1505.00716 [hep-th]].
- [57] R. Uchiyama, "General Theory of Relativity," (Shokabo, 1978),
- [58] S. Navas et al. [Particle Data Group], Phys. Rev. D 110, no.3, 030001 (2024)
- [59] M. E. Peskin and D. V. Schroeder, Addison-Wesley, 1995,