

Towards construction of superintegrable basis in matrix models

Batukhan Azheev^{1*} and Nikita Tselousov^{1,2,3,4†}

¹MIPT, 141701, Dolgoprudny, Russia

²NRC “Kurchatov Institute”, 123182, Moscow, Russia

³IITP RAS, 127051, Moscow, Russia

⁴ITEP, Moscow, Russia

ABSTRACT

We develop methods for systematic construction of superintegrable polynomials in matrix/eigenvalue models. Our consideration is based on a tight connection of superintegrable property of Gaussian Hermitian model and $W_{1+\infty}$ algebra in Fock representation. Motivated by this example, we propose a set of assumptions that may allow one to recover superintegrable polynomials. The main two assumptions are box adding/removing rule (Pierri rule) and existence of Hamiltonian for superintegrable polynomials. We detail our method in case of the Gaussian Hermitian model, and then apply it to the cubic Kontsevich model.

1 Introduction

Matrix models [1,2] are a rapidly developing area of modern mathematical physics. They represent zero-dimensional and therefore often exactly solvable examples of quantum field theories where one can study intriguing phenomena such as *superintegrability* [3]:

$$\langle \chi \rangle \sim \chi(*) \quad (1)$$

Remarkably, quantum correlators of distinguished basis in fields χ turn out to be expressed via simple functions, or even more, via functions $\chi(*)$ themselves in a specific point. The classic example of the superintegrability phenomenon is the simple Gaussian Hermitian one-matrix model [4] (GH model), where integrals are taken over $N \times N$ Hermitian matrices H :

$$\langle f \rangle = \frac{1}{Z} \int dH f(H) \exp\left(-\frac{1}{2} \text{tr} H^2\right) \quad (2)$$

The normalization factor $Z = \int dH \exp\left(-\frac{1}{2} \text{tr} H^2\right)$ is chosen to satisfy $\langle 1 \rangle = 1$. Superintegrable basis χ in this case is given by well-known characters of linear groups – Schur polynomials S_λ that

*e-mail: azheev.bd@phystech.edu

†e-mail: tselousov.ns@phystech.edu

enumerated by Young diagrams $\lambda = [\lambda_1, \lambda_2, \dots, \lambda_{l(\lambda)}]$, where λ_i are positive integers obeying the non-increasing rule $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \dots \geq \lambda_{l(\lambda)} \geq 0$. All basic facts about Schur polynomials are listed in [5]. Example of Young diagram is presented in the following picture:

$$\lambda = \begin{array}{cccccccc} \square & \square & \square & \square & \square & \square & \square & \square \\ \square & \square & \square & \square & \square & \square & & \\ \square & \square & \square & \square & & & & \\ \square & \square & & & & & & \\ \square & & & & & & & \\ \square & & & & & & & \end{array} = [7, 6, 4, 2, 1, 1]$$

Explicit superintegrable formula for this model reads:

$$\langle S_\lambda \rangle = \frac{S_\lambda(N)}{S_\lambda(\delta_{k,1})} \cdot S_\lambda(\delta_{k,2}) \quad (3)$$

where we understand Schur polynomials as functions of p_k variables and our notation $S_\lambda(\delta_{k,1}) := S_\lambda(p_k \rightarrow \delta_{k,1}), S_\lambda(N) := S_\lambda(p_k \rightarrow N)$. Connection to symmetric functions of N variables x_i are given by Miwa transformation $p_k = \sum_{i=1}^N (x_i)^k$. Schur polynomials for single-row Young diagrams $[n]$ can be computed from generating function:

$$\exp\left(\sum_{k=1}^{\infty} \frac{p_k z^k}{k}\right) = \sum_{n=1}^{\infty} z^n \cdot S_{[n]}(p_a) \quad (4)$$

The other polynomials for higher Young diagram can be obtained from Jacobi-Trudi determinant formula:

$$S_\lambda = \det_{1 \leq i, j \leq l(\lambda)} S_{[\lambda_i + j - i]} \quad (5)$$

Ratio of Schur polynomials in special points in formula (3) is a fully factorizable quantity:

$$\frac{S_\lambda(N)}{S_\lambda(\delta_{k,1})} = \prod_{\square \in \lambda} (N + j_\square - i_\square) \quad (6)$$

where product runs over all boxes in the Young diagram λ and (i_\square, j_\square) are the vertical and horizontal coordinates of the box \square . The quantity $\omega_\square = N + j_\square - i_\square$ is called content of the box and it plays an important role in the superintegrable formula (3).

For this simple and well-studied model such as GH model one can offer the following explanation of superintegrability phenomenon. The first key observation is that full partition function of the model:

$$Z_{GH}(p_k) = \frac{1}{Z} \int dH \exp\left(-\frac{1}{2} \text{tr} H^2 + \sum_{k=1}^{\infty} \frac{p_k}{k} \text{tr} H^k\right) \quad (7)$$

admits W -representation [6, 7]:

$$Z_{GH}(p_k) = e^{\hat{W}} \cdot 1 \quad (8)$$

Explicit form of \hat{W} operator reads:

$$\hat{W} = \frac{1}{2} \sum_{a,b=1}^{\infty} (a+b-2) p_a p_b \frac{\partial}{\partial p_{a+b-2}} + ab p_{a+b+2} \frac{\partial}{\partial p_a} \frac{\partial}{\partial p_b} + N \sum_{a=1}^{\infty} a p_{a+2} \frac{\partial}{\partial p_a} + \frac{N^2 p_2}{2} + \frac{N p_1^2}{2} \quad (9)$$


Note, that for simplicity of formulas we add extra factor $\frac{1}{k}$ in front of $p_k \text{tr} H^k$ in the definition of partition function. The operator \hat{W} has grading 2, that is explicitly seen from the following formula:

$$[\hat{D}, \hat{W}] = 2\hat{W} \quad \hat{D} = \sum_{k=1}^{\infty} k p_k \frac{\partial}{\partial p_k} \quad (10)$$

Due to the positive grading 2 the image of operator \hat{W} when acting on polynomials of grading n is given by polynomials of grading $n + 2$. Schur polynomials form *distinguished basis* of graded polynomials where the action of \hat{W} operator has very natural meaning:

$$\hat{W} S_{\lambda} = \frac{1}{2} \cdot \sum_{\square \in \text{Add}_{[2]}(\lambda)} \omega_{\square_1} \omega_{\square_2} S_{\lambda + \square} - \frac{1}{2} \cdot \sum_{\blacksquare \in \text{Add}_{[1,1]}(\lambda)} \omega_{\blacksquare_1} \omega_{\blacksquare_2} S_{\lambda + \blacksquare} \quad (11)$$

Operator \hat{W} acts on Schur polynomials by adding boxes to corresponding Young diagrams. In other words, on the r.h.s. appears only those polynomials which Young diagram is obtained by adding boxes to initial Young diagram λ from the l.h.s. Definitions of sets $\text{Add}_{[2]}(\lambda)$ and $\text{Add}_{[1,1]}(\lambda)$ are clear from the following example:



Quantities $\omega_{\square_{1,2}}$ in formula (11) are contents $N + j_{\square} - i_{\square}$ of added boxes. For example:

$$\hat{W} S_{\square} = \frac{1}{2} (N + 2)(N + 3) S_{\square\square} + \frac{1}{2} (N - 1)(N) S_{\blacksquare} - \frac{1}{2} (N - 1)(N - 2) S_{\blacksquare} \quad (13)$$

Orthogonal properties of Schur polynomials that are expressed in an elegant way via Cauchy identity [5]:

$$\sum_{\lambda} S_{\lambda}(p_a) S_{\lambda}(\bar{p}_a) = \exp \left(\sum_{k=1}^{\infty} \frac{p_k \bar{p}_k}{k} \right) \quad (14)$$

The above relation allows one to express the partition function (7) in the following form:

$$\begin{aligned} Z_{GH}(p_k) &= \left\langle \exp \left(\sum_{k=1}^{\infty} \frac{p_k}{k} \text{tr} H^k \right) \right\rangle = \sum_{\lambda} S_{\lambda}(p_k) \left\langle S_{\lambda} \left(\bar{p}_k = \text{tr} H^k \right) \right\rangle = \\ &= \sum_{\lambda} S_{\lambda}(p_k) S_{\lambda}(\delta_{k,2}) \prod_{\square \in \lambda} (N + j_{\square} - i_{\square}) \end{aligned} \quad (15)$$

At this point we are equipped with the knowledge that exponent of \hat{W} operator generates the whole partition function and the operator acts on Schur polynomials by adding two boxes - domino tiles. It can be seen, if the partition function is split into graded components:

$$Z_{GH}(p_k) = Z_{GH}^{(0)} + Z_{GH}^{(1)} + Z_{GH}^{(2)} + \dots \quad (16)$$

$$Z_{GH}^{(2k)} = \frac{1}{k!} (\hat{W})^k \cdot 1 \quad (17)$$

$$Z_{GH}^{(2k+1)} = 0 \quad (18)$$

We compute small levels explicitly by action of the \hat{W} operator:

$$Z_{GH}^{(0)} = 1 \quad (19)$$

$$Z_{GH}^{(2)} = \hat{W} \cdot 1 = \underbrace{\frac{1}{2} (N)(N+1)}_{\langle S_{\square} \rangle} S_{\square} + \underbrace{\left(-\frac{1}{2}\right) (N)(N-1)}_{\langle S_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \rangle} S_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \quad (20)$$

$$\begin{aligned} Z_{GH}^{(4)} = \frac{1}{2} \hat{W} \cdot Z_{GH}^{(2)} &= \underbrace{\frac{1}{8} (N)(N+1)(N+2)(N+3)}_{\langle S_{\square\square\square\square} \rangle} S_{\square\square\square\square} + \underbrace{\left(-\frac{1}{8}\right) (N)(N-1)(N+1)(N+2)}_{\langle S_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} \rangle} S_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} + \\ &+ \underbrace{\left(\frac{1}{8} + \frac{1}{8}\right) (N)^2 (N-1)(N+1)}_{\langle S_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} \rangle} S_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} + \underbrace{\left(-\frac{1}{8}\right) N(N+1)(N-1)(N-2)}_{\langle S_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} \rangle} S_{\begin{smallmatrix} \square & \square \\ \square & \square \end{smallmatrix}} + \\ &\underbrace{\frac{1}{8} (N)(N-1)(N-2)(N-3)}_{\langle S_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} \rangle} S_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} \end{aligned} \quad (21)$$

Now we better understand the superintegrable formula (3) due to the following consequences of the above formulas:

- The overall product over boxes of content function $\omega_{\square} = N + j_{\square} - i_{\square}$ in formula (3) is generated step by step due to factors in r.h.s. of \hat{W} operator¹ action (11).
- Correlator of Schur polynomial $\langle S_{\lambda} \rangle$ has non-vanishing value if and only if the diagram λ can be constructed only with domino-tiles \square and $\begin{smallmatrix} \square \\ \square \end{smallmatrix}$ and starting from empty diagram \emptyset . More accurately, the number of ways to construct diagram λ from domino-tiles (note, that the vertical tile is weighted with (-1)) is equal to $2^{\frac{|\lambda|}{2}} \cdot \left(\frac{|\lambda|}{2}\right)! \cdot S_{\lambda}(\delta_{k,2})$.

After all, a logical question follows: *how can one guess that Schur polynomials should arise as a superintegrable basis of correlators?* In particular case of Gaussian Hermitian model the answer is known. We formulate it in two steps:

- \hat{W} operator as differential operator in p_k variables belongs to an infinite dimensional algebra $W_{1+\infty}$ [8,9] in Fock representation that acts on space of polynomials $\mathbb{C}[p_1, p_2, \dots]$.
- Commutative family of operators $\hat{\psi}_a$ of $W_{1+\infty}$ in Fock representation have Schur polynomials as a common set of eigenfunctions:

$$\hat{\psi}_a S_{\lambda} = \mathcal{E}_{\lambda,a} S_{\lambda} \quad (22)$$

¹Note, that \hat{W} -operator may act on Schur polynomial of arbitrary Young diagram, however for GH model relevant only diagrams of even size.

Schur polynomials are distinguished by the underlying algebra to which \hat{W} belongs. Due to Schur polynomials one can say that $W_{1+\infty}$ algebra in Fock representation acts on the Hilbert space, where basis vectors enumerated by Young diagrams.

We do not give here details of $W_{1+\infty}$ representation theory and list only some basic formulas concerning Fock representation. We work with generators $\hat{e}_k, \hat{\psi}_k, \hat{f}_k$, where $k = 0, 1, 2, \dots$, that are rising operators, Cartan-like operators and lowering operators respectively. Among them there is a special small set of operators $\hat{e}_0, \hat{\psi}_3, \hat{f}_0$ that have the following form in Fock representation:

$$\hat{e}_0 = p_1 \qquad \hat{f}_0 = \frac{\partial}{\partial p_1} \qquad (23)$$

$$\hat{\psi}_3 = \frac{1}{2} \sum_{a,b=1}^{\infty} (a+b) p_a p_b \frac{\partial}{\partial p_{a+b}} + ab p_{a+b} \frac{\partial}{\partial p_a} \frac{\partial}{\partial p_b} + N \sum_{a=1}^{\infty} a p_a \frac{\partial}{\partial p_a} \qquad (24)$$

The higher operators can be expressed via only these three operators by the following rules:

$$\hat{e}_k = \left(\text{ad}_{\hat{\psi}_3} \right)^k \hat{e}_0 \qquad \hat{f}_k = (-)^k \left(\text{ad}_{\hat{\psi}_3} \right)^k \hat{f}_0 \qquad (25)$$


$$\hat{\psi}_{a+b} = \left[\hat{e}_a, \hat{f}_b \right] \qquad (26)$$

Action of small set operators $\hat{e}_0, \hat{\psi}_3, \hat{f}_0$ on Schur polynomials obeys the following formulas:

$$\begin{aligned} \hat{e}_0 S_\lambda &= \sum_{\square \in \text{Add}(\lambda)} S_{\lambda + \square} \\ \hat{f}_0 S_\lambda &= \sum_{\blacksquare \in \text{Rem}(\lambda)} S_{\lambda - \blacksquare} \end{aligned} \qquad (27)$$

$$\hat{\psi}_3 S_\lambda = \left(\sum_{\square \in \lambda} \omega_\square \right) \cdot S_\lambda \qquad (28)$$

where as usual $\omega_\square = N + j_\square - i_\square$. Operator (24) is a cut-and-join operator [10], however, we call it Hamiltonian of Schur polynomials, due to connection to quantum integrable systems². Formulas (27) for Schur polynomials are called Pierri rules [5]. The sums in the above runs over all possible ways of adding/removing boxes in Young diagram. We provide example of $\text{Add}(\lambda)$ and $\text{Rem}(\lambda)$:



Action of higher \hat{e}_k, \hat{f}_k is a direct consequence of (25):

$$\begin{aligned} \hat{e}_k S_\lambda &= \sum_{\square \in \text{Add}(\lambda)} (\omega_\square)^k S_{\lambda + \square} \\ \hat{f}_k S_\lambda &= \sum_{\blacksquare \in \text{Rem}(\lambda)} (\omega_{\blacksquare})^k S_{\lambda - \blacksquare} \end{aligned} \qquad (30)$$

² β -deformed version of this operator coincides with quantum Hamiltonian of Calogero-Moser many body integrable system with trigonometric interaction [11]

\hat{W} -operator of Gaussian Hermitian model is expressed via \hat{e}_k by the following simple formula:

$$\hat{W} = \frac{1}{2} [\hat{e}_2, \hat{e}_1] \quad (31)$$

Action of \hat{W} operator on Schur polynomials (11) follows from the above formulas by direct computation. Therefore superintegrable formula (3) acquires natural interpretation in terms of $W_{1+\infty}$ algebra.

Remarkable fact is that all components of the above story *simultaneously* admit one parametric deformation widely known as β -deformation. Under this deformation the matrix integral becomes eigenvalue β -ensemble [12–14]; superintegrable property persist with one parametric deformation of Schur polynomials S_λ that bear the name of Jack polynomials J_λ^β ; the underlying algebra becomes affine Yangian $Y(\hat{\mathfrak{gl}}_1)$ [15–18]. This story certainly deserves attention, however, it is under the scope of this paper.

Motivated by the tight relation of superintegrable property of Gaussian Hermitian model and $W_{1+\infty}$ algebra we propose the conjecture: superintegrability is a manifestation of underlying symmetry algebra. However, it seems a difficult question to tell the algebra for a given matrix model. Basing on the above mentioned example we propose a set of assumptions:

- *Diagrams.*

In GH model superintegrable basis was enumerated by Young diagrams, in particular, number of graded polynomials $P(n)$ on the level n of time-variables p_a matches the number of Young diagrams with n boxes.

We suppose that set of time-variables T_a is known for a given matrix model. Then we analyze the number of graded polynomials of corresponding time-variables of this model and choose the proper set of diagrams Λ by counting argument. This step of the algorithm is rather a clever guess. For now we do not have any additional general arguments to systematically extract the shape of diagrams from the given matrix model. However, in each particular case there are may be more arguments specific for a given model.

Superintegrable basis $\text{SIB}_\Lambda(T_a)$ then is enumerated by diagrams Λ .

- *Box adding/removing rule.*

Usually the first (the smallest) time-variable, say T_1 , correspond to first superintegrable polynomial corresponding to just \square diagram. Therefore we suppose that the first time-variable T_1 and its derivative add and remove boxes:

$$\begin{aligned} T_1 \cdot \text{SIB}_\Lambda &= \sum_{\square \in \text{Add}(\Lambda)} C_{\Lambda, \Lambda + \square} \text{SIB}_{\Lambda + \square} \\ \frac{\partial}{\partial T_1} \text{SIB}_\Lambda &= \sum_{\square \in \text{Rem}(\Lambda)} \tilde{C}_{\Lambda, \Lambda - \square} \text{SIB}_{\Lambda - \square} \end{aligned} \quad (32)$$

We do not impose any relations on coefficients $C_{\Lambda, \Lambda + \square}$, $\tilde{C}_{\Lambda, \Lambda - \square}$ and treat them as unknown constants. The main idea here is that the sum on r.h.s. runs only over limited subsets $\text{Add}(\Lambda), \text{Rem}(\Lambda)$ of diagrams of size $|\Lambda| + 1$ and $|\Lambda| - 1$ respectively. These assumptions are rather strong constraints on the form of SIB_Λ polynomials.

Higher time-variables and Virasoro operators of given matrix model may act on superintegrable polynomials by adding and removing rule. However, for a non minimal grading there are in general several different adding/removing rules.

Remarkably, if a set of diagrams Λ contain linear (single-row or single-column) diagrams then this set is closed under operation of removing boxes. We use this fact to define linear system of equation on single-row polynomials.

- *Hamiltonian for SIB_Λ polynomials.*

Box adding/removing rules impose strong constraints on SIB_Λ polynomials, however, they can not fix the form of these polynomials completely. Motivated by the $W_{1+\infty}$ example we postulate that SIB_Λ polynomials should be eigenfunctions of some Hamiltonian operator \hat{H} :

$$\hat{H} SIB_\Lambda = \kappa_\Lambda \cdot SIB_\Lambda \quad (33)$$

with unknown eigenvalues κ_Λ . The only assumption for the Hamiltonian operator is *finite spin constraint*. Where by *spin* we mean the number of time variables T_a and derivatives $\frac{\partial}{\partial T_a}$ that enter each term (in normal ordered form) of the Hamiltonian. For example operator (24) has spin 3. We will consider Hamiltonians of minimal possible spin in our presentation.

We argue that these two main assumptions: *box adding/removing rule* and existence of *Hamiltonian* may allow one to fully determine the form of superintegrable polynomials in some cases.

This paper is organized as follows. In Section 2 we demonstrate in details our method of construction of superintegrable basis. First, we analyze the diagrams. Second, we consider single-row family of diagrams and construct naively overdetermined system of equations for corresponding superintegrable polynomials, that has unique solution. Third, we introduce ansatz for Hamiltonian of finite (minimal) spin with unknown coefficients and partially fix these coefficients from single-row superintegrable polynomials. Finally, using box adding and removing rule we complete the basis and find the form of Hamiltonian. In Section 3 we apply the above mentioned algorithm to cubic Kontsevich matrix model and step by step restore its superintegrable basis – Q-Schur polynomials. In Section 4 we briefly summarize our results.

2 Gaussian Hermitian model

In this section we perform steps of our algorithm without references to Schur polynomials themselves, i.e. pretending that we do not know form of Schur polynomials. Our method starts from partition function of Gaussian Hermitian model that is given by the following integral over $N \times N$ Hermitian matrices H :

$$Z_{GH}(p_k) = \frac{1}{Z} \int dH \exp \left(-\frac{1}{2} \text{tr} H^2 + \sum_{k=1}^{\infty} \frac{p_k}{k} \text{tr} H^k \right) \quad (34)$$

Partition function as a function of infinite family of time-variables p_k obeys infinitely many linear equations (Ward identities) - Virasoro constraints [19–22]:

$$\hat{L}_n Z_{GH} = (n+2) \frac{\partial Z_{GH}}{\partial p_{n+2}}, \quad n = -1, 0, 1, 2, \dots \quad (35)$$

Explicit formulas for Virasoro operators depend on the model and in this particular case they have the following form:

$$\hat{L}_n = \sum_{a=1}^{n-1} a(n-a) \frac{\partial}{\partial p_a} \frac{\partial}{\partial p_{n-a}} + \sum_{k=1}^{\infty} (n+k) p_k \frac{\partial}{\partial p_{n+k}} + 2Nn \frac{\partial}{\partial p_n} + N^2 \delta_{n,0} + N p_1 \delta_{n+1,0} \quad (36)$$

Virasoro operators L_n obey commutation relations of Virasoro algebra:

$$[\hat{L}_n, \hat{L}_m] = (n - m)\hat{L}_{n+m} \quad n, m \geq -1 \quad (37)$$

2.1 Step 1. Diagrams

Time-variables p_k have a parameter - grading $[p_k]$. The gradings are encoded in L_0 Virasoro operator:

$$\hat{L}_0 = \sum_{k=1}^{\infty} k \cdot p_k \frac{\partial}{\partial p_k} + N^2 \quad (38)$$

$$[p_k] = k \quad (39)$$

Usually it is possible to identify polynomial basis in time-variables with the set of diagrams. In case of time-variables p_a with gradings $[p_a] = a$ the set of diagrams is given by Young diagrams.

Level	Young diagrams	Basis polynomials
1		p_1
2		p_2, p_1^2
3		$p_3, p_2 p_1, p_1^3$
4		$p_4, p_3 p_1, p_2^2, p_2 p_1^2, p_1^4$
5		$p_5, p_4 p_1, p_3 p_2, p_3 p_1^2, p_2^2 p_1, p_2 p_1^3, p_1^5$

Table 1: Young diagrams and graded polynomials on small levels 1-5. Young diagrams are in one-to-one correspondence to the graded polynomials in time-variables p_a .

Generating function of number of Young diagrams $P^{YD}(n)$ on level n have the following form:

$$\prod_{k=1}^{\infty} \frac{1}{1 - q^k} = \sum_{n=0}^{\infty} q^n P^{YD}(n) = 1 + q + 2q^2 + 3q^3 + 5q^4 + 7q^5 + \dots \quad (40)$$

2.2 Step 2. System of equations for single-row polynomials

Virasoro operators L_n and time-variables $p_k, \frac{\partial}{\partial p_k}$ have the following gradings:

grading	...	3	2	1	0	-1	-2	-3	...
time-variables	...	p_3	p_2	p_1	1	$\frac{\partial}{\partial p_1}$	$\frac{\partial}{\partial p_2}$	$\frac{\partial}{\partial p_3}$...
Virasoro operators				L_{-1}	L_0	L_1	L_2	L_3	...

According to the list of assumptions (listed in the end of Section 1) we observe the following property of single-row subset of Young diagrams. If we act on a SIB (i.e. SuperIntegrable Basis) polynomial $SIB_{[n]}$ labeled by single-row diagram $[n]$ of length n with operator \mathcal{O}_m of negative grading $[\mathcal{O}_m] = -m$ we get $SIB_{[n-m]}$ without any option, because there is only *one way* to remove

m boxes from single-row diagram. Pictorially this statement looks as follows in a particular example $n = 8, m = 3$:

$$\hat{O}_3 \text{SIB}_{\square\square\square\square\square\square\square\square} \sim \text{SIB}_{\square\square\square\square\square\square\square\square} \quad (41)$$

Here we colored the boxes that are removed after the action of operator. Our approach is based on removing boxes, because adding boxes immediately takes us out from single-row subset. For example, operator of positive grading 1 add a box into second row:

$$\hat{O}_{-1} \text{SIB}_{\square\square\square\square\square\square\square\square} \sim \text{SIB}_{\square\square\square\square\square\square\square\square} + \text{SIB}_{\square\square\square\square\square\square\square\square} \quad (42)$$

We consider the following system of equations:

$$\frac{\partial \text{SIB}_{[n]}}{\partial p_k} = u_{n,k} \text{SIB}_{[n-k]} \quad (43)$$

$$\hat{L}_k \text{SIB}_{[n]} = v_{n,k} \text{SIB}_{[n-k]} \quad (44)$$

This system naively is heavily overdetermined; however it is self-consistent and has a unique solution. We consider the following general form of SIB polynomials:

$$\begin{aligned} \text{SIB}_{[0]} &= 1 \\ \text{SIB}_{[1]} &= p_1 \\ \text{SIB}_{[2]} &= \frac{p_1^2}{2} + a_{2,1} p_2 \\ \text{SIB}_{[3]} &= \frac{p_1^3}{6} + a_{3,1} p_2 p_1 + a_{3,2} p_3 \\ \text{SIB}_{[4]} &= \frac{p_1^4}{24} + a_{4,1} p_2 p_1^2 + a_{4,2} p_2^2 + a_{4,3} p_3 p_1 + a_{4,4} p_4 \\ \text{SIB}_{[5]} &= \frac{p_1^5}{120} + a_{5,1} p_2 p_1^3 + a_{5,2} p_2^2 p_1 + a_{5,3} p_3 p_1^2 + a_{5,4} p_4 p_1 + a_{5,5} p_3 p_2 + a_{5,6} p_5 \\ &\dots \end{aligned} \quad (45)$$

The overall normalization of polynomials is fixed by the first coefficient that we choose to be $1/n!$. This choice makes the coefficients $u_{n,1} = 1$ trivial. The solution of this system is unique and parametrized by one parameter x :

$$\begin{aligned} \text{SIB}_{[0]} &= 1 \\ \text{SIB}_{[1]} &= p_1 \\ \text{SIB}_{[2]} &= \frac{p_1^2}{2} + x \frac{p_2}{2} \\ \text{SIB}_{[3]} &= \frac{p_1^3}{6} + x \frac{p_1 p_2}{2} + x^2 \frac{p_3}{3} \\ \text{SIB}_{[4]} &= \frac{p_1^4}{24} + x \frac{p_1^2 p_2}{4} + x^2 \frac{p_2^2}{8} + x^2 \frac{p_1 p_3}{3} + x^3 \frac{p_4}{4} \\ \text{SIB}_{[5]} &= \frac{p_1^5}{120} + x \frac{p_1^3 p_2}{12} + x^2 \frac{p_1 p_2^2}{8} + x^2 \frac{p_1^2 p_3}{6} + x^3 \frac{p_2 p_3}{6} + x^3 \frac{p_1 p_4}{4} + x^4 \frac{p_5}{5} \\ &\dots \end{aligned} \quad (46)$$

Formulas for coefficients $u_{n,k}, v_{n,k}$ have the following form:

$$u_{n,k} = x^{k-1} \quad (47)$$

$$v_{n,k} = x^{k-2} (k - 1 + 2xN + (n - k)x^2) \quad (48)$$

Note, that free parameter x is nothing but rescaling of time-variables $p_k \rightarrow x^{k-1} p_k$. However, Virasoro operators (36) have fixed normalization of time-variables. We observe, that correct normalization is restored when coefficients $u_{n,k}, v_{n,k}$ do not have power dependence on k . In this case it corresponds to value $\boxed{x = 1}$.

2.3 Step 3. Hamiltonian for SIB polynomials

To complete the polynomials $\text{SIB}_{[n]}$ to complete basis we use additional assumption. The first of them is a finite-spin Hamiltonian \hat{H} that has SIB as the set of eigenfunctions:

$$\hat{H} \text{SIB}_\lambda = \kappa_\lambda \text{SIB}_\lambda \quad (49)$$

Here we use assumption that \hat{H} has the smallest possible spin = number of time-variable and derivatives. In this particular case it is possible to construct non-trivial zero grading operator of spin 3:

$$\hat{H} = \sum_{a,b=1}^{\infty} A_{a,b} \cdot p_a p_b \frac{\partial}{\partial p_{a+b}} + B_{a,b} \cdot p_{a+b} \frac{\partial}{\partial p_a} \frac{\partial}{\partial p_b} \quad (50)$$

where $A_{a,b} = A_{b,a}$ and $B_{a,b} = B_{b,a}$ are some unknown coefficients. Similar formulas were considered in [23]. Solving equation (49) for the already found functions (46):

$$\hat{H} \text{SIB}_{[n]} = \kappa_{[n]} \text{SIB}_{[n]} \quad (51)$$

we can fix the form of coefficients $A_{a,b}, B_{a,b}$. We choose overall normalization of Hamiltonian by setting $B_{1,1} = 1$. Then eigenvalues are restored immediately:

n	0	1	2	3	4	5	6	7	8	...
$\kappa_{[n]}$	0	0	2	6	12	20	30	42	56	...

The resulting formula for eigenvalues reads:

$$\kappa_\lambda = n(n - 1) \quad (52)$$

Coefficients $A_{a,b}$ are *fully* restored from (51):

$A_{a,b}$	1	2	3	4	5	6	...
1	2	3	4	5	6	7	...
2	3	4	5	6	7	8	...
3	4	5	6	7	8	9	...
4	5	6	7	8	9	10	...
5	6	7	8	9	10	11	...
6	7	8	9	10	11	12	...
...

The resulting formula reads:

$$\boxed{A_{a,b} = a + b} \quad (53)$$

The situation with coefficients $B_{a,b}$ is more involved. They can not be fully determined from the above reasoning, however, there are linear equations on them. One equation for each level:

$$B_{1,1} = 1 \quad (54)$$

$$\frac{1}{2} B_{1,2} + \frac{1}{2} B_{2,1} = 2 \quad (55)$$

$$\frac{1}{3} B_{1,3} + \frac{1}{4} B_{2,2} + \frac{1}{3} B_{3,1} = 3 \quad (56)$$

$$\frac{1}{4} B_{1,4} + \frac{1}{6} B_{2,3} + \frac{1}{6} B_{3,2} + \frac{1}{4} B_{4,1} = 4 \quad (57)$$

$$\frac{1}{5} B_{1,5} + \frac{1}{8} B_{2,4} + \frac{1}{9} B_{3,3} + \frac{1}{8} B_{4,2} + \frac{1}{5} B_{5,1} = 5 \quad (58)$$

$$\dots \quad (59)$$

We intentionally do not use property $B_{a,b} = B_{b,a}$ in order to represent this system in a symmetric form. Generally, these equations read:

$$\sum_{a=1}^{n-1} \frac{1}{a(n-a)} \cdot B_{a,n-a} = n \quad (60)$$

2.4 Step 4. Box adding and removing rule

2.4.1 Small levels

For now we have considered two objects:

- Subset of single-row $\text{SIB}_{[n]}$ polynomials (46)
- Spin-3 ansatz for simplest Hamiltonian (50)

that obviously can not determine SIB completely, however, they allow us to *fully* determine complete SIB on 2 and 3 level, because the corresponding coefficients $B_{1,1} = 1$ and $B_{1,2} = B_{2,1} = 2$ are known.

2 level:

$$\text{SIB}_{\square\square} = \frac{p_1^2}{2} + \frac{p_2}{2} \quad \kappa_{\square\square} = 2 \quad (61)$$

$$\text{SIB}_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} = \frac{p_1^2}{2} - \frac{p_2}{2} \quad \kappa_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} = -2 \quad (62)$$

3 level:

$$\text{SIB}_{\square\square\square} = \frac{p_1^3}{6} + \frac{p_1 p_2}{2} + \frac{p_3}{3} \quad \kappa_{\square\square\square} = 6 \quad (63)$$

$$\text{SIB}_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}} = \frac{p_1^3}{3} - \frac{p_3}{3} \quad \kappa_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}} = 0 \quad (64)$$

$$\text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} = \frac{p_1^3}{6} - \frac{p_1 p_2}{2} + \frac{p_3}{3} \quad \kappa_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} = -6 \quad (65)$$

These results are in full agreement with box adding and removing rule. For example:

$$p_1 \cdot \text{SIB}_{\square\square} = \text{SIB}_{\square\square\square} + \text{SIB}_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}} \quad (66)$$

$$\frac{\partial}{\partial p_1} \text{SIB}_{\begin{smallmatrix} \square & \square \\ \square \end{smallmatrix}} = \text{SIB}_{\square\square} + \text{SIB}_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \quad (67)$$

2.4.2 Higher levels

One can proceed further in computing SIB polynomials via box adding rules. Multiplication by p_1 increases the grading by one, therefore we write the following ansatz for adding a box:

$$p_1 \cdot \text{SIB}_{\square\square\square} = \alpha_1 \cdot \text{SIB}_{\square\square\square\square} + \alpha_2 \cdot \text{SIB}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}} \quad (68)$$

with some constants α_1, α_2 and the following general form of new polynomial:

$$\text{SIB}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}} = \frac{p_1^4}{8} + b_1 \cdot p_1^2 p_2 + b_2 \cdot p_2^2 + b_3 \cdot p_1 p_3 + b_4 \cdot p_4 \quad (69)$$

The solution to the above equation is the following:

$$\alpha_1 = 4 - 3\alpha_2 \quad b_1 = \frac{3}{4} - \frac{1}{2\alpha_2} \quad b_2 = \frac{3}{8} - \frac{1}{2\alpha_2} \quad b_3 = 1 - \frac{1}{\alpha_2} \quad b_4 = \frac{3}{4} - \frac{1}{\alpha_2} \quad (70)$$

where only one unknown parameter α_2 remained. We can find it via the Hamiltonian:

$$\hat{H} \text{SIB}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}} = \kappa_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}} \text{SIB}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}} \quad (71)$$

The remaining parameter α_2 is fixed from the above equation, moreover the unknown coefficient $B_{1,3}$ of the Hamiltonian is fixed too:

$$\alpha_2 = 1 \quad B_{1,3} = 3 \quad (72)$$

Now all the coefficients of Hamiltonian of the 4-th level are fixed and the basis of 4-th level is known in full generality:

$$\text{SIB}_{\square\square\square\square} = \frac{p_1^4}{24} + \frac{p_1^2 p_2}{4} + \frac{p_2^2}{8} + \frac{p_1 p_3}{3} + \frac{p_4}{4} \quad \kappa_{\square\square\square\square} = 12 \quad (73)$$

$$\text{SIB}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}} = \frac{p_1^4}{8} + \frac{p_1^2 p_2}{4} - \frac{p_2^2}{8} - \frac{p_4}{4} \quad \kappa_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \\ \hline \end{array}} = 4 \quad (74)$$

$$\text{SIB}_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = \frac{p_1^4}{12} + \frac{p_2^2}{4} - \frac{p_1 p_3}{3} \quad \kappa_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = 0 \quad (75)$$

$$\text{SIB}_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} = \frac{p_1^4}{8} - \frac{p_1^2 p_2}{4} - \frac{p_2^2}{8} + \frac{p_4}{4} \quad \kappa_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} = -4 \quad (76)$$

$$\text{SIB}_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} = \frac{p_1^4}{24} - \frac{p_1^2 p_2}{4} + \frac{p_2^2}{8} + \frac{p_1 p_3}{3} - \frac{p_4}{4} \quad \kappa_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}} = -12 \quad (77)$$

$$(78)$$

One can proceed further using box adding rule with Hamiltonian and level by level compute SIB polynomials. In particular, one finds remaining coefficients of the Hamiltonian:

$$\begin{aligned} B_{1,1} &= 1 \\ \frac{1}{2} B_{1,2} &= \frac{1}{2} B_{2,1} = 1 \\ \frac{1}{3} B_{1,3} &= \frac{1}{4} B_{2,2} = \frac{1}{3} B_{3,1} = 1 \\ \frac{1}{4} B_{1,4} &= \frac{1}{6} B_{2,3} = \frac{1}{6} B_{3,2} = \frac{1}{4} B_{4,1} = 1 \\ &\dots \end{aligned} \quad (79)$$

One can guess the solution for system of equations (60):

$$\boxed{B_{a,b} = a \cdot b} \quad (80)$$

On this step we fully determine the Hamiltonian:

$$\hat{H} = \sum_{a,b=1}^{\infty} (a+b) \cdot p_a p_b \frac{\partial}{\partial p_{a+b}} + ab \cdot p_{a+b} \frac{\partial}{\partial p_a} \frac{\partial}{\partial p_b} \quad (81)$$

which has eigenvalues:

$$\kappa_{\lambda} = 2 \cdot \sum_{\square \in \lambda} j_{\square} - i_{\square} \quad (82)$$

Remarkably, the eigenvalues are integer numbers that is a rather nontrivial property of Hamiltonian. Even if we know the full form of Hamiltonian, it does not determine SIB polynomials completely, because the spectrum is degenerate: eigenvalues start to coincide from 6-th level. However eigenfunction property equipped with box adding/removing rules *determines* SIB_{λ} *completely*.

2.5 Action of Virasoro operators on SIB_{λ} polynomials

Notably, the overall coefficients of SIB_{λ} polynomials can be chosen in a way that the following equations have unit coefficients in front of polynomials in the r.h.s (Pierri rules):

$$p_1 \cdot \text{SIB}_{\lambda} = \sum_{\square \in \text{Add}(\lambda)} \text{SIB}_{\lambda + \square} \quad (83)$$

$$\frac{\partial}{\partial p_1} \text{SIB}_{\lambda} = \sum_{\square \in \text{Rem}(\lambda)} \text{SIB}_{\lambda - \square} \quad (84)$$

Another intriguing property of resulting SIB_{λ} polynomials is that Virasoro operators add and remove boxes. We demonstrate this fact by several examples:


$$\hat{L}_{-1} \text{SIB}_{\lambda} = \sum_{\square \in \text{Add}(\lambda)} (N + j_{\square} - i_{\square}) \cdot \text{SIB}_{\lambda + \square} \quad (85)$$

$$\hat{L}_1 \text{SIB}_{\lambda} = \sum_{\square \in \text{Rem}(\lambda)} (2N + j_{\square} - i_{\square}) \cdot \text{SIB}_{\lambda - \square} \quad (86)$$

Operators \hat{L}_{-1} , \hat{L}_1 have gradings 1 and -1 respectively, therefore they add and remove only one box. The situation is more involved for higher Virasoro operators and higher time-derivatives, they remove skew diagrams that are called *border strips*. It is instructive to consider similar formulas for higher time-variables:

$$k \frac{\partial}{\partial p_k} \text{SIB}_{\lambda} = \sum_{\substack{\text{bs} \in \text{BS}(\lambda) \\ |\text{bs}|=k}} (-)^{h(\text{bs})+1} \text{SIB}_{\lambda - \text{bs}} \quad (87)$$

We denote as $\text{BS}(\lambda)$ a set of border strips of diagram λ . A border strip (also called ribbon or rim-hook) is a skew diagram that is connected, has unit width. An example of border strip $\text{bs} \in \text{BS}([7, 6, 4, 2, 1, 1])$ of size $|\text{bs}| = 7$ and height $h(\text{bs}) = 3$:



(88)

Then formula for L_2 Virasoro operator reads:

$$\begin{aligned} \hat{L}_2 \text{SIB}_\lambda = & \sum_{\substack{\text{bs} \in \text{BS}(\lambda) \\ |\text{bs}|=2}} (-)^{h(\text{bs})+1} (2N + j_{\blacksquare_{\text{bs}}} - i_{\blacksquare_{\text{bs}}}) \text{SIB}_{\lambda-\text{bs}} + \\ & + \sum_{\substack{\text{bs}, \text{bs}' \in \text{BS}(\lambda) \\ |\text{bs}|+|\text{bs}'|=2}} (-)^{h(\text{bs})+h(\text{bs}')} \text{SIB}_{\lambda-\text{bs}-\text{bs}'} \end{aligned} \quad (89)$$

In the above formula there are two sums, the first sum over border strips of size 2 while the second sum runs over pairs of (non-empty) border strips of overall size 2. In this particular case it means the second sum runs over pairs of (non-adjacent) boxes from set $\text{Rem}(\lambda)$.

In the first sum of the above formula content $j_{\blacksquare} - i_{\blacksquare}$ is evaluated in the special point \blacksquare_{bs} that depends on border strip. There are only 2^{k-1} possible border strips of size k and we provide them explicitly for $k = 2$:

$$\begin{array}{cc} \blacksquare & \blacksquare \\ \blacksquare & \blacksquare \end{array} \quad (90)$$

Details of the above formula are clear from explicit example:

$$\hat{L}_2 \text{SIB}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \square \\ \hline \end{array}} = (-)(2N - 2) \text{SIB}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & & \square \\ \hline \end{array}} + (2N + 2) \text{SIB}_{\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}} + \text{SIB}_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} \quad (91)$$

General formula for \hat{L}_k operator action has the following form:

$$\begin{aligned} \hat{L}_k \text{SIB}_\lambda = & \sum_{\substack{\text{bs} \in \text{BS}(\lambda) \\ |\text{bs}|=k}} (-)^{h(\text{bs})+1} (2N + j_{\blacksquare_{\text{bs}}} - i_{\blacksquare_{\text{bs}}}) \text{SIB}_{\lambda-\text{bs}} + \\ & + \sum_{\substack{\text{bs}, \text{bs}' \in \text{BS}(\lambda) \\ |\text{bs}|+|\text{bs}'|=k}} (-)^{h(\text{bs})+h(\text{bs}')} \text{SIB}_{\lambda-\text{bs}-\text{bs}'} \end{aligned} \quad (92)$$

The remaining element of the general formula that does not have yet any description is a position of the special box \blacksquare_{bs} . We show all border strips of size 3 and 4 explicitly:

$$\begin{array}{cccc} \blacksquare & \blacksquare & \blacksquare & \blacksquare \\ \blacksquare & \blacksquare & \blacksquare & \blacksquare \\ \blacksquare & \blacksquare & \blacksquare & \blacksquare \end{array} \quad (93)$$

$$\begin{array}{cccccccc} \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare \\ \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare \\ \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare & \blacksquare \end{array} \quad (94)$$

However, this general formula demonstrates important property: Virasoro operators act on superintegrable polynomials by removing boxes from corresponding diagrams. Similar formulas was considered in [24]

3 Cubic Kontsevich model

In this section we will apply our method and construct superintegrable basis for Kontsevich matrix model [25] in Kontsevich phase. According to the recent progress [26] the superintegrable basis for this model is given by Q-Schur polynomials [5], that are tightly connected to BKP integrability of this model [27, 28]. Our presentation will be parallel to the previous section.

First, we define the partition function for Kontsevich model and write down Virasoro operators that realize the Ward identities. Next we discuss the time-variables of the model and analyze possible families of diagrams, that are combinatorial objects indexing basis graded polynomials.

We start construction of superintegrable basis SIB_λ ³ from the single-row sector, which is closed with respect to the action of the box removing operators. Next we consider minimal spin ansatz for the Hamiltonian (in this case minimal spin is 4) for which SIB polynomials are eigenfunctions. To complete single-row subset to the full basis we impose box adding/removing rule for the minimal time-variable. These assumptions allow us to completely fix the form of SIB polynomials. Remarkably, the resulting Hamiltonian has only integer eigenvalues and higher Virasoro operators act on SIB polynomials by removing boxes.

Partition function of Kontsevich matrix integral in Kontsevich phase is defined by the following integral over Hermitian $N \times N$ matrices H with external matrix Φ :

$$\mathcal{Z}_K(\Phi) = \frac{1}{\mathcal{Z}(\Phi)} \int dH \exp \left(-\frac{1}{3!} \text{tr} H^3 - \frac{1}{2} \text{tr} H^2 \Phi \right) \quad (95)$$

where normalization factor $\mathcal{Z}(\Phi)$ is defined as follows:

$$\mathcal{Z}(\Phi) = \int dH \exp \left(-\frac{1}{2} \text{tr} H^2 \Phi \right) \quad (96)$$

Partition function is a function of an external matrix Φ and in Kontsevich phase it becomes a power series in variables $\text{tr} \Phi^{-k}$ [29–31]. More importantly it depends only on odd negative powers of external matrix that we treat (in the limit $N \rightarrow \infty$) as independent time-variables τ_k :

$$\tau_k = \frac{\text{tr} \Phi^{-2k-1}}{2k+1} \quad k = 0, 1, 2, 3, \dots \quad (97)$$

As a function of time-variables partition function $\mathcal{Z}_K(\tau_k)$ obeys infinitely many linear equations - Virasoro constraints [32, 33]:

$$\hat{\mathcal{L}}_n \mathcal{Z}_K = \frac{\partial \mathcal{Z}_K}{\partial \tau_{n+1}} \quad n = -1, 0, 1, 2, \dots \quad (98)$$

Virasoro operators of this model has the following explicit form [34]:

$$\hat{\mathcal{L}}_n = \frac{1}{8} \sum_{k=0}^{n-1} \frac{\partial}{\partial \tau_k} \frac{\partial}{\partial \tau_{n-k-1}} + \sum_{k=0}^{\infty} \left(k + \frac{1}{2} \right) \tau_k \frac{\partial}{\partial \tau_{n+k}} + \frac{\delta_{n,0}}{16} + \delta_{n+1,0} \frac{\tau_0^2}{2} \quad (99)$$

and obey relations of Virasoro algebra:

$$\left[\hat{\mathcal{L}}_n, \hat{\mathcal{L}}_m \right] = (n-m) \hat{\mathcal{L}}_{n+m} \quad n, m \geq -1 \quad (100)$$

3.1 Step 1. Diagrams

Time-variables τ_k have natural gradings that are encoded in zero Virasoro operator:

$$\hat{\mathcal{L}}_0 = \sum_{k=0}^{\infty} \left(k + \frac{1}{2} \right) \tau_k \frac{\partial}{\partial \tau_k} + \frac{1}{16} \quad (101)$$

$$[\tau_k] = 2 \left(k + \frac{1}{2} \right) = 2k + 1 \quad (102)$$

³We denote superintegrable basis of Kontsevich model by the same name SIB_λ , however they are different from superintegrable basis of Gaussian Hermitian model.

Lvl.	Odd partitions (OP)	Distinct partitions (DP)	Basis polynomials
1			τ_0
2			τ_0^2
3			τ_1, τ_0^3
4			$\tau_1\tau_0, \tau_0^4$
5			$\tau_2, \tau_1\tau_0^2, \tau_0^5$
6			$\tau_2\tau_0, \tau_1^2, \tau_1\tau_0^3, \tau_0^6$

Table 2: Odd partitions, distinct (strict) partition and graded polynomials of time-variables τ_k for small levels 1-6. Odd partitions (OP) have obvious mapping to graded polynomials in time-variables τ_k . Distinct partitions (DP) do not have a clear connection to basis monomials, however their dimensions indeed coincide on each level.

where for simplicity we rescaled all the gradings by 2.

Similar set of time-variables can be obtained by reduction $p_{2k} \rightarrow 0$ of time-variables p_a of GH model from the previous section. The set of τ_k time-variables can be described by diagrams in at least two ways. First option is Young diagrams with odd row lengths – *odd partitions* (OP). Second option is Young diagrams with different row lengths – *distinct partitions* (DP). The last diagrams are also called strict partitions in the literature. Examples of these diagrams are collected in Table 2.

Generating function of number of odd and distinct partitions $P^{OP}(n) = P^{DP}(n)$ ⁴ on level n has the following form:

$$\prod_{k=1}^{\infty} \frac{1}{1 - q^{2k-1}} = \sum_{n=0}^{\infty} q^n P^{DP}(n) = 1 + q + q^2 + 2q^3 + 2q^4 + 3q^5 + 4q^6 + \dots \quad (103)$$

Odd partitions are *not suitable* in our method and we provide the following arguments below. Odd partitions have linear one-parametric subset of one-column diagrams $[1^n]$. According to our assumptions (see Section 1) the first time-variable τ_0 and its derivative $\frac{\partial}{\partial \tau_0}$ acts by adding and removing a box to corresponding diagrams. In particular, subset of one-column diagrams of OP is closed under operation of removing a box, therefore the following equation holds for corresponding polynomials :

$$\frac{\partial}{\partial \tau_0} \text{SIB}_{[1^n]}^{OP} = c_n \text{SIB}_{[1^{n-1}]}^{OP} \quad (104)$$

⁴The idea of proof $P^{OP}(n) = P^{DP}(n)$ is that every natural number can be expressed uniquely as a power of 2 multiplied by an odd number.

with some coefficients c_n . On the other hand, the box adding operation also does not output from the subset of one-column diagrams:

$$\tau_0 \text{SIB}_{[1^n]}^{OP} = c'_n \text{SIB}_{[1^{n+1}]}^{OP} \quad (105)$$

because the only other possible diagram $[2, 1^{n-1}]$ contains even number and therefore is not included in odd partition family. Therefore for OP family there is only trivial solution:

$$\text{SIB}_{[1^n]}^{OP} \sim \tau_0^n \quad (106)$$

We do not consider OP diagrams in this paper and focus on distinct partitions (DP).

3.2 Step 2. System of equations on single-row polynomials

Virasoro operators $\hat{\mathcal{L}}_n$ and variables $\tau_k, \frac{\partial}{\partial \tau_k}$ have the following gradings:

grading	...	3	2	1	0	-1	-2	-3	...
time-variables	...	τ_1	-	τ_0	1	$\frac{\partial}{\partial \tau_0}$	-	$\frac{\partial}{\partial \tau_1}$...
Virasoro operators			$\hat{\mathcal{L}}_{-1}$	-	$\hat{\mathcal{L}}_0$	-	$\hat{\mathcal{L}}_1$	-	...

We assume that single-row subspace $\text{SIB}_{[n]}$ is closed under action of Virasoro operators and derivatives and consider the following system of equations (very similar to (43)):

$$\frac{\partial \text{SIB}_{[n]}}{\partial \tau_k} = u_{n,k} \text{SIB}_{[n-2k-1]} \quad (107)$$

$$\hat{\mathcal{L}}_k \text{SIB}_{[n]} = v_{n,k} \text{SIB}_{[n-2k]} \quad (108)$$

This system is naively overdetermined, however it is self-consistent and has a unique solution. We consider general form of SIB polynomials:

$$\begin{aligned} \text{SIB}_{[0]} &= 1 \\ \text{SIB}_{[1]} &= \tau_0 \\ \text{SIB}_{[2]} &= \tau_0^2 \\ \text{SIB}_{[3]} &= \frac{2\tau_0^3}{3} + a_{3,1} \tau_1 \\ \text{SIB}_{[4]} &= \frac{\tau_0^4}{3} + a_{4,1} \tau_1 \tau_0 \\ \text{SIB}_{[5]} &= \frac{2\tau_0^5}{15} + a_{5,1} \tau_1 \tau_0^2 + a_{5,2} \tau_2 \\ \text{SIB}_{[6]} &= \frac{2\tau_0^6}{45} + a_{6,1} \tau_0^3 \tau_1 + a_{6,2} \tau_1^2 + a_{6,3} \tau_0 \tau_2 \\ &\dots \end{aligned} \quad (109)$$

We put first coefficient $2^{n-1}/n!$ to fix the overall normalization for future convenience. Solution of

the above system is unique and parametrized by one parameter x :

$$\begin{aligned}
\text{SIB}_{[0]} &= 1 \\
\text{SIB}_{[1]} &= \tau_0 \\
\text{SIB}_{[2]} &= \tau_0^2 \\
\text{SIB}_{[3]} &= \frac{2\tau_0^3}{3} + 4x\tau_1 \\
\text{SIB}_{[4]} &= \frac{\tau_0^4}{3} + 8x\tau_1\tau_0 \\
\text{SIB}_{[5]} &= \frac{2\tau_0^5}{15} + 8x\tau_0^2\tau_1 + 16x^2\tau_2 \\
\text{SIB}_{[6]} &= \frac{2\tau_0^6}{45} + \frac{16}{3}x\tau_0^3\tau_1 + 16x^2\tau_1^2 + 32x^2\tau_0\tau_2 \\
&\dots
\end{aligned} \tag{110}$$

From the above formulas one can see that x parameter is a trivial rescaling of time-variables $\tau_k \rightarrow x^k \tau_k$. The explicit formula for coefficients $u_{n,k}, v_{n,k}$:

$$u_{n,k} = 2 \cdot (4x)^k \tag{111}$$

$$v_{n,k} = \frac{1}{2} \cdot (4x)^{k-1} (k + 4x(n - 2k)) \tag{112}$$

According to our observation after formula (47) the true normalization of time-variables is restored when power dependence on k is eliminated from $u_{n,k}, v_{n,k}$ coefficient. In this particular case there

only one solution $\boxed{x = \frac{1}{4}}$.

3.3 Step 3. Hamiltonian for SIB polynomials

In order to complete the basis we consider the simplest Hamiltonian $\hat{\mathcal{H}}$ for SIB_λ polynomials:

$$\hat{\mathcal{H}} \text{SIB}_\lambda = \varepsilon_\lambda \text{SIB}_\lambda \tag{113}$$

For the time-variable τ_k the simplest non-trivial Hamiltonian has spin 4:

$$\begin{aligned}
\hat{\mathcal{H}} &= \sum_{a,b,c=0}^{\infty} A_{a,b,c} \tau_{a+b+c+1} \frac{\partial^3}{\partial \tau_a \partial \tau_b \partial \tau_c} + B_{a,b,c} \tau_a \tau_b \tau_c \frac{\partial}{\partial \tau_{a+b+c+1}} + \\
&\quad + \sum_{a+b=c+d} C_{a,b|c,d} \tau_a \tau_b \frac{\partial^2}{\partial \tau_c \partial \tau_d} + \sum_{a=0}^{\infty} G_a \tau_a \frac{\partial}{\partial \tau_a}
\end{aligned} \tag{114}$$

Coefficients $A_{a,b,c}$ and $B_{a,b,c}$ are absolutely symmetric with respect to permutations of indices a, b, c , while coefficients $C_{a,b|c,d}$ have symmetry $a \leftrightarrow b, c \leftrightarrow d$.

It is instructive to consider grading operator:

$$\hat{D} = \sum_{k=0}^{\infty} (2k + 1) \tau_k \frac{\partial}{\partial \tau_k} \tag{115}$$

then there are two trivial Hamiltonians \hat{D} and \hat{D}^2 among operators of zero grading and spin at most 4. Eigenvalues of these trivial Hamiltonians on single-row polynomials:

$$\hat{D} \text{SIB}_{[n]} = n \cdot \text{SIB}_{[n]} \tag{116}$$

$$\hat{D}^2 \text{SIB}_{[n]} = n^2 \cdot \text{SIB}_{[n]} \tag{117}$$

We impose eigenfunction condition for single-row polynomials:

$$\hat{\mathcal{H}} \text{SIB}_{[n]} = \varepsilon_{[n]} \cdot \text{SIB}_{[n]} \quad (118)$$

and solve corresponding equations. As an output we get eigenvalues $\varepsilon_{[n]}$ and represent them as a functions of $\varepsilon_{[1]}, \varepsilon_{[2]}, \varepsilon_{[3]}$:

$$\begin{aligned} \varepsilon_{[4]} &= 4\varepsilon_{[1]} - 6\varepsilon_{[2]} + 4\varepsilon_{[3]} \\ \varepsilon_{[5]} &= 15\varepsilon_{[1]} - 20\varepsilon_{[2]} + 10\varepsilon_{[3]} \\ \varepsilon_{[6]} &= 36\varepsilon_{[1]} - 45\varepsilon_{[2]} + 20\varepsilon_{[3]} \\ \varepsilon_{[7]} &= 70\varepsilon_{[1]} - 84\varepsilon_{[2]} + 35\varepsilon_{[3]} \\ \varepsilon_{[8]} &= 120\varepsilon_{[1]} - 140\varepsilon_{[2]} + 56\varepsilon_{[3]} \\ \varepsilon_{[9]} &= 189\varepsilon_{[1]} - 216\varepsilon_{[2]} + 84\varepsilon_{[3]} \\ \varepsilon_{[10]} &= 280\varepsilon_{[1]} - 315\varepsilon_{[2]} + 120\varepsilon_{[3]} \\ &\dots \end{aligned} \quad (119)$$

$$\varepsilon_{[n]} = \frac{n(n-2)(n-3)}{2}\varepsilon_{[1]} - \frac{n(n-1)(n-3)}{2}\varepsilon_{[2]} + \frac{n(n-1)(n-2)}{6}\varepsilon_{[3]}, \quad (120)$$

Three parameters $\varepsilon_{[1]}, \varepsilon_{[2]}, \varepsilon_{[3]}$ allow one to vary the form of single-row eigenvalues $\varepsilon_{[n]}$ and we argue that there is one distinguished choice of the parameters. Motivated by eigenvalues (116) we impose:

$$\boxed{\varepsilon_{[n]} = n^3} \quad (121)$$

Notably, this condition is satisfied when $\varepsilon_{[1]} = 1, \varepsilon_{[2]} = 2^3, \varepsilon_{[3]} = 3^3$. From the above equations on single-row polynomials we find $B_{a,b,c}$ coefficients:

$$\begin{aligned} B_{0,0,0} &= 4 \\ B_{0,0,1} &= 4 \cdot 3 \\ B_{0,0,2} &= 4 \cdot 5 \\ B_{0,1,1} &= 4 \cdot 3^2 \\ B_{0,0,3} &= 4 \cdot 7 \\ B_{0,1,2} &= 4 \cdot 3 \cdot 5 \\ B_{1,1,1} &= 4 \cdot 3^3 \\ &\dots \end{aligned} \quad (122)$$

therefore

$$B_{a,b,c} = 4 \cdot (2a+1)(2b+1)(2c+1) \quad (123)$$

We do not provide explicitly all the other constraints on $A_{a,b,c}, C_{a,b|c,d}, G_a$ coefficients but several examples:

$$\begin{aligned} C_{0,0|0,0} &= 3 & G_0 &= 1 \\ C_{0,1|0,1} &= 9 & A_{0,0,0} &= \frac{1}{4}(27 - G_1) \\ 2C_{1,1|0,2} + C_{1,1|1,1} &= 81 & A_{0,0,1} &= \frac{1}{12}(125 - G_2) \\ C_{0,2|1,1} + 2C_{0,2|0,2} &= 45 & A_{0,0,2} + A_{0,1,1} &= \frac{1}{12}(343 - G_3) \\ &\dots & &\dots \end{aligned} \quad (124)$$

3.4 Step 4. Box adding and removing rule

On the next step we impose box adding/removing rules with eigenfunction property (113) for SIB polynomials up to 5 level. Box adding rules read:

$$\begin{aligned}
\tau_0 \cdot \text{SIB}_{\square\square} &= c_1 \text{SIB}_{\square\square\square} + c_2 \text{SIB}_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \\
\tau_0 \cdot \text{SIB}_{\square\square\square} &= c_3 \text{SIB}_{\square\square\square\square} + c_4 \text{SIB}_{\begin{smallmatrix} \square\square \\ \square \end{smallmatrix}} \\
\tau_0 \cdot \text{SIB}_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} &= c_5 \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} \\
\tau_0 \cdot \text{SIB}_{\square\square\square\square} &= c_6 \text{SIB}_{\square\square\square\square\square} + c_7 \text{SIB}_{\begin{smallmatrix} \square\square\square \\ \square \end{smallmatrix}} \\
\tau_0 \cdot \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} &= c_8 \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \end{smallmatrix}} + c_9 \text{SIB}_{\begin{smallmatrix} \square\square \\ \square \\ \square \end{smallmatrix}}
\end{aligned} \tag{125}$$

Box removing rules (we do not mention here single-row polynomials, because they were analyzed earlier):

$$\begin{aligned}
\frac{\partial}{\partial\tau_0} \text{SIB}_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} &= d_1 \text{SIB}_{\square\square} \\
\frac{\partial}{\partial\tau_0} \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} &= d_2 \text{SIB}_{\square\square\square} + d_3 \text{SIB}_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} \\
\frac{\partial}{\partial\tau_0} \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \end{smallmatrix}} &= d_4 \text{SIB}_{\square\square\square\square} + d_5 \text{SIB}_{\begin{smallmatrix} \square\square \\ \square \\ \square \end{smallmatrix}} \\
\frac{\partial}{\partial\tau_0} \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ \square \end{smallmatrix}} &= d_6 \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}}
\end{aligned} \tag{126}$$

Here c_i, d_i are some constants. Solving the above equations and eigenfunction equations (113) simultaneously with respect to polynomial coefficients we obtain the following solution:

$$\begin{aligned}
\text{SIB}_{\begin{smallmatrix} \square \\ \square \end{smallmatrix}} &= \frac{\tau_0^3}{3} + (\gamma - 27) \frac{\tau_1}{12} \\
\text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}} &= \frac{\tau_0^4}{3} + (\gamma - 27) \frac{\tau_0\tau_1}{12} \\
\text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \end{smallmatrix}} &= \frac{\tau_0^5}{5} + (\gamma - 15) \frac{\tau_0^2\tau_1}{20} + (\gamma - 39) \frac{\tau_2}{24} \\
\text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ \square \end{smallmatrix}} &= \frac{2\tau_0^5}{15} + (\gamma - 27) \frac{\tau_0^2\tau_1}{12} + (\gamma - 39)(\gamma - 27) \frac{\tau_2}{288}
\end{aligned} \tag{127}$$

where $\gamma := G_1$ is a free parameter that is not fixed from the above mentioned equations. However, if we consider equations from the next level 6:

$$\begin{aligned}
\tau_0 \cdot \text{SIB}_{\square\square\square\square} &= c_{10} \text{SIB}_{\square\square\square\square\square} + c_{11} \text{SIB}_{\begin{smallmatrix} \square\square\square \\ \square \end{smallmatrix}} \\
\tau_0 \cdot \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \end{smallmatrix}} &= c_{12} \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ \square \end{smallmatrix}} + c_{13} \text{SIB}_{\begin{smallmatrix} \square\square \\ \square \\ \square \end{smallmatrix}} \\
\tau_0 \cdot \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ \square \end{smallmatrix}} &= c_{13} \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ \square \\ \square \end{smallmatrix}} + c_{14} \text{SIB}_{\begin{smallmatrix} \square\square \\ \square \\ \square \\ \square \end{smallmatrix}} \\
\frac{\partial}{\partial\tau_0} \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ \square \end{smallmatrix}} &= d_7 \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ \square \\ \square \end{smallmatrix}} + d_8 \text{SIB}_{\begin{smallmatrix} \square\square \\ \square \\ \square \\ \square \end{smallmatrix}} \\
\frac{\partial}{\partial\tau_0} \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ \square \\ \square \end{smallmatrix}} &= d_9 \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ \square \\ \square \\ \square \end{smallmatrix}} + d_{10} \text{SIB}_{\begin{smallmatrix} \square\square \\ \square \\ \square \\ \square \\ \square \end{smallmatrix}} \\
\frac{\partial}{\partial\tau_0} \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ \square \\ \square \\ \square \end{smallmatrix}} &= d_{11} \text{SIB}_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \end{smallmatrix}}
\end{aligned} \tag{128}$$

this parameter is fixed uniquely $\boxed{\gamma = 15}$. We provide explicit formulas for SIB polynomials un to 6 level:

$$\begin{array}{ll}
\text{SIB}_{\square} = \tau_0 & \varepsilon_{\square} = 1 \\
\text{SIB}_{\square\square} = \tau_0^2 & \varepsilon_{\square\square} = 8 \\
\text{SIB}_{\square\square\square} = \frac{2\tau_0^3}{3} + \tau_1 & \varepsilon_{\square\square\square} = 27 \\
\text{SIB}_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} = \frac{\tau_0^3}{3} - \tau_1 & \varepsilon_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} = 9 \\
\text{SIB}_{\square\square\square\square} = \frac{\tau_0^4}{3} + 2\tau_0\tau_1 & \varepsilon_{\square\square\square\square} = 64 \\
\text{SIB}_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}} = \frac{\tau_0^4}{3} - \tau_0\tau_1 & \varepsilon_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}} = 28 \\
\text{SIB}_{\square\square\square\square\square} = \frac{2\tau_0^5}{15} + 2\tau_0^2\tau_1 + \tau_2 & \varepsilon_{\square\square\square\square\square} = 125 \\
\text{SIB}_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}} = \frac{\tau_0^5}{5} - \tau_2 & \varepsilon_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}} = 64 \\
\text{SIB}_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = \frac{2\tau_0^5}{15} - \tau_0^2\tau_1 + \tau_2 & \varepsilon_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} = 35 \\
\text{SIB}_{\square\square\square\square\square\square} = \frac{2\tau_0^6}{45} + \frac{4}{3}\tau_0^3\tau_1 + \tau_1^2 + 2\tau_0\tau_2 & \varepsilon_{\square\square\square\square\square\square} = 216 \\
\text{SIB}_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} = \frac{4\tau_0^6}{45} + \frac{2}{3}\tau_0^3\tau_1 - \tau_1^2 - \tau_0\tau_2 & \varepsilon_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} = 126 \\
\text{SIB}_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}} = \frac{\tau_0^6}{9} - \frac{2}{3}\tau_0^3\tau_1 + \tau_1^2 & \varepsilon_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}} = 72 \\
\text{SIB}_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square \\ \hline \end{array}} = \frac{\tau_0^6}{45} - \frac{1}{3}\tau_0^3\tau_1 - \tau_1^2 + \tau_0\tau_2 & \varepsilon_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square \\ \hline \end{array}} = 36
\end{array} \tag{129}$$

Now one can see that SIB polynomials are nothing but well-known Q-Schur polynomials [5,26]. On this limited set of examples we observe remarkable "sum over boxes" property of eigenvalues:

$$\varepsilon_{\lambda} = \sum_{\square \in \lambda} j_{\square}^3 - (j_{\square} - 1)^3 = \sum_{i=1}^{l(\lambda)} \lambda_i^3 \tag{130}$$

Here we denote by j_{\square} the horizontal coordinate of a box \square and the first box has coordinate 1. Solving equations like (125), (126), (127) for box adding/removing rules and equations (113) one can level by level compute SIB polynomials. From the other hand, eigenfunction conditions (113) fix the form of Hamiltonian coefficients:

$$\begin{array}{llll}
A_{0,0,0} = 3 & & & G_0 = 1 \\
A_{0,0,1} = 5 & C_{0,0|0,0} = 3 & & G_1 = 15 \\
A_{0,0,2} = 7 & C_{0,1|0,1} = 3 \cdot 3 & C_{0,3|0,3} = 3 \cdot 7 & G_2 = 65 \\
A_{0,1,1} = 7 & C_{0,2|0,2} = 3 \cdot 5 & C_{0,3|1,2} = 3 \cdot 7 & G_3 = 175 \\
A_{0,0,3} = 9 & C_{0,2|1,1} = 3 \cdot 5 & C_{1,2|0,3} = 3 \cdot 3 \cdot 5 & G_4 = 369 \\
A_{0,1,2} = 9 & C_{1,1|0,2} = 3 \cdot 3^2 & C_{1,2|1,2} = 3 \cdot 3 \cdot 5 & G_5 = 671 \\
A_{1,1,1} = 9 & C_{1,1|1,1} = 3 \cdot 3^2 & \dots & \dots \\
\dots & & & \dots
\end{array} \tag{131}$$

These values of coefficients are enough to reconstruct the full dependence:

$$\begin{aligned} A_{a,b,c} &= 2(a+b+c) + 3 \\ C_{a,b|c,d} &= 3(2a+1)(2b+1) \\ G_a &= (2a+1)(2a^2+2a+1) \end{aligned} \tag{132}$$


Now we are able to write the full Hamiltonian for SIB polynomials:

$$\begin{aligned} \hat{\mathcal{H}} &= \sum_{a,b,c=0}^{\infty} (2(a+b+c) + 3) \tau_{a+b+c+1} \frac{\partial^3}{\partial \tau_a \partial \tau_b \partial \tau_c} + \sum_{a+b=c+d} 3(2a+1)(2b+1) \tau_a \tau_b \frac{\partial^2}{\partial \tau_c \partial \tau_d} \\ &\quad + \sum_{a,b,c=0}^{\infty} 4(2a+1)(2b+1)(2c+1) \tau_a \tau_b \tau_c \frac{\partial}{\partial \tau_{a+b+c+1}} + \sum_{a=0}^{\infty} (2a+1)(2a^2+2a+1) \tau_a \frac{\partial}{\partial \tau_a} \end{aligned} \tag{133}$$


One can check that eigenvalues of the above operator are positive integers and obey formula (130). This Hamiltonian itself does not define SIB basis due to degenerate spectrum, for example $\varepsilon_{[8,7]} = 8^3 + 7^3 = 855 = 9^3 + 5^3 + 1^3 = \varepsilon_{[9,5,1]}$. However, with adding/removing rules the basis is fixed completely.

3.5 Action of Virasoro operators on SIB_λ polynomials

Motivated by the action of time-variables and Virasoro operators on SIB polynomials that we discuss in this section it is instructive to slightly change notation of diagrams DP - distinct partitions. For this purpose we represent a Young diagram (turned upside down) as a way of tight packing of square boxes in a corner of a room. In this setting distinct partitions (DP) are in one-to-one correspondence with possible ways of tight packing of square boxes under a ladder. We demonstrate examples of these diagrams, left picture correspond to usual Young diagram, right one correspond to distinct partition:



(a) Usual partition



(a) Distinct partition


(134)

Form now on we represent distinct partitions as Young diagrams under the ladder. With these notations the action of τ_0 and $\frac{\partial}{\partial \tau_0}$ have the following from:

$$\tau_0 \cdot \text{SIB}_\lambda = \sum_{\square \in \text{AddDP}(\lambda)} \text{SIB}_{\lambda + \square} \tag{135}$$

$$\frac{\partial}{\partial \tau_0} \text{SIB}_\lambda = \sum_{\square \in \text{RemDP}(\lambda)} (2 - \delta_{i_\square, j_\square}) \cdot \text{SIB}_{\lambda - \square} \quad (136)$$


Note, that coefficients in the r.h.s. almost always are units except the diagonal $i_\square = j_\square$. The sets $\text{AddDP}(\lambda)$ and $\text{RemDP}(\lambda)$ are defined in the same way as in the previous section (29). We provide explicit example of $\text{AddDP}(\lambda)$ and $\text{RemDP}(\lambda)$:



Derivatives with respect to higher time-variables τ_k act on superintegrable polynomials via the following rules:

$$\frac{\partial}{\partial \tau_k} \text{SIB}_\lambda = \sum_{\substack{\text{bs} \in \text{BS}^*(\lambda) \\ |\text{bs}| = 2k+1}} (-1)^{\frac{|\text{bs}^{(2)}|}{2} + h(\text{bs}^{(1)}) + 1} \cdot (2 - \delta_{i_{\square_{\text{bs}}}, j_{\square_{\text{bs}}}}) \cdot \text{SIB}_{\lambda - \text{bs}} \quad (138)$$

Where $\text{BS}^*(\lambda)$ is a set of *double strips* [5]. A double strip is a skew diagram (i.e. it can be represented as λ/μ , where $\mu \subset \lambda$, λ, μ are distinct partitions), that is connected and has width one or two. Subdiagram of width two always touches on the ladder-border. A double strip can be a usual border strip if it has width one everywhere. We demonstrate two examples of double strips of the same distinct partitions:



On the above diagrams we explicitly mark boxes of subset of unit width $\text{bs}^{(1)}$ with 1 and boxes of subset $\text{bs}^{(2)}$ (width 2) with 2. We denote as $|\text{bs}^{(2)}|$ the number of boxes is subset of width 2. While $h(\text{bs}^{(1)})$ is a height of unit width subset. In our notation $i_{\square_{\text{bs}}}, j_{\square_{\text{bs}}}$ are coordinates of the head of the strip, i.e. the box for which $j_\square - i_\square$ has minimal value (for essentially double strip there are two such boxes, however there is no ambiguity because for both of them $\delta_{i_{\square_{\text{bs}}}, j_{\square_{\text{bs}}}} = 1$). We provide a representative example of $\frac{\partial}{\partial \tau_2}$ action:

$$\frac{\partial}{\partial \tau_2} \text{SIB}_{\begin{array}{c} \square \square \square \\ \square \square \square \\ \square \square \square \end{array}} = \text{SIB}_{\begin{array}{c} \square \square \square \\ \square \square \square \end{array}} + \text{SIB}_{\begin{array}{c} \square \square \square \\ \square \square \square \end{array}} - 2 \cdot \text{SIB}_{\begin{array}{c} \square \square \square \\ \square \square \square \end{array}} \quad (140)$$

Virasoro operators $\hat{\mathcal{L}}_n$ ($n > 0$) act by removing double strips as well as time-derivatives (138), however in this case coefficients are not constants:

$$\hat{\mathcal{L}}_n \text{SIB}_\lambda = \frac{1}{4} \cdot \sum_{\substack{\text{bs} \in \text{BS}^*(\lambda) \\ |\text{bs}| = 2k+1}} (-1)^{\frac{|\text{bs}^{(2)}|}{2} + h(\text{bs}^{(1)}) + 1} \cdot (2 - \delta_{i_{\square_{\text{bs}}}, j_{\square_{\text{bs}}}}) \cdot \left(k + j_{\square_{\text{bs}}} - i_{\square_{\text{bs}}} - \frac{|\text{bs}^{(2)}|}{2} \right) \cdot \text{SIB}_{\lambda - \text{bs}} \quad (141)$$

Note that the box removing rules for the above Virasoro operators and corresponding time-derivatives (138) are the same. While in case of GH model the box removing rules are different for time-derivatives (87) and Virasoro operators (92) (in case of Virasoro operators there is an additional sum over two distinct border strips).

4 Conclusion

In this paper we made a trial of systematic construction of superintegrable basis (SIB) in matrix models for particular examples of Gaussian Hermitian and cubic Kontsevich models. Our construction procedure is based on the main conjecture: *superintegrable formulas are manifestations of underlying "hidden symmetry" algebras.*

Motivated by the example of Gaussian Hermitian model, which symmetry algebra $W_{1+\infty}$ is well-known, we formulate a (conjectural) set of assumptions, that may allow one to reconstruct superintegrable polynomials in other models. These assumptions concern the shape of Young-like diagrams that enumerate SIB polynomials, Pierri rules for adding and removing boxes in these diagrams and existence of Hamiltonian operator that encodes SIB polynomials as its own eigenfunctions.

Our method represents first steps towards systematic construction of SIB polynomials. We applied the method for models where SIB polynomials were already known, while the true success of the method would come after a discovery of SIB polynomials in more complicated models (e.g. generalized Kontsevich models [29, 35, 36] or super-eigenvalue models [37–41]).

We list here disadvantages of our method and possible modifications:

- In our method we used box adding/removing rules only for the simplest time-variables and higher grading operators only for single-row sector. However, as a result we observe that all time-derivatives (87), (138) and Virasoro operators (92), (141) remove boxes via special border strip rules for all diagrams.

In case of more complicated models the above simple reasoning may not allow to completely fix the form of SIB polynomials. It may be useful to impose box removing rules for all time-derivatives and Virasoro operators at the very beginning.

- In our method we impose very special condition of finite spin for Hamiltonian operator (50), (114). It may be wrong for other cases, and the finite spin condition may be replaced with more universal one (e.g. self-adjoint property with respect to proper scalar product, "sum over box" property of eigenvalues, etc.).

Acknowledgments

We are indebted for illuminating discussions to D.Galakhov, A.Mironov, A.Morozov, A.Popolitov, A.Sleptsov.

The work was partially funded within the state assignment of the Institute for Information Transmission Problems of RAS (N.T.). Our work is partly supported by the grants of the Foundation for the Advancement of Theoretical Physics and Mathematics “BASIS” (N.T.).

References

- [1] A. Morozov, “Integrability and matrix models,” *Phys. Usp.* **37** (1994) 1–55, [arXiv:hep-th/9303139](#).
- [2] A. Mironov, “2-d gravity and matrix models. 1. 2-d gravity,” *Int. J. Mod. Phys. A* **9** (1994) 4355–4406, [arXiv:hep-th/9312212](#).
- [3] A. Mironov and A. Morozov, “Superintegrability summary,” *Phys. Lett. B* **835** (2022) 137573, [arXiv:2201.12917 \[hep-th\]](#).
- [4] A. Mironov and A. Morozov, “On the complete perturbative solution of one-matrix models,” *Phys. Lett. B* **771** (2017) 503–507, [arXiv:1705.00976 \[hep-th\]](#).
- [5] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*. Oxford Mathematical Monographs, 1998.
- [6] A. Morozov and S. Shakirov, “Generation of Matrix Models by W-operators,” *JHEP* **04** (2009) 064, [arXiv:0902.2627 \[hep-th\]](#).
- [7] A. Mironov, V. Mishnyakov, A. Morozov, and R. Rashkov, “Matrix model partition function by a single constraint,” *Eur. Phys. J. C* **81** no. 12, (2021) 1140, [arXiv:2105.09920 \[hep-th\]](#).
- [8] C. N. Pope, L. J. Romans, and X. Shen, “ $W(\infty)$ and the Racah-wigner Algebra,” *Nucl. Phys. B* **339** (1990) 191–221.
- [9] H. Awata, M. Fukuma, Y. Matsuo, and S. Odake, “Representation theory of the $W(1+\infty)$ algebra,” *Prog. Theor. Phys. Suppl.* **118** (1995) 343–374, [arXiv:hep-th/9408158](#).
- [10] A. Mironov, A. Morozov, and S. Natanzon, “Complete Set of Cut-and-Join Operators in Hurwitz-Kontsevich Theory,” *Theor. Math. Phys.* **166** (2011) 1–22, [arXiv:0904.4227 \[hep-th\]](#).
- [11] *Calogero–Moser–Sutherland Models*. Springer New York, NY.
- [12] V. Mishnyakov and A. Oreshina, “Superintegrability in β -deformed Gaussian Hermitian matrix model from W-operators,” *Eur. Phys. J. C* **82** no. 6, (2022) 548, [arXiv:2203.15675 \[hep-th\]](#).

- [13] R. Wang, C.-H. Zhang, F.-H. Zhang, and W.-Z. Zhao, “CFT approach to constraint operators for (β -deformed) hermitian one-matrix models,” *Nucl. Phys. B* **985** (2022) 115989, [arXiv:2203.14578 \[hep-th\]](#).
- [14] A. Bawane, P. Karimi, and P. Sułkowski, “Proving superintegrability in β -deformed eigenvalue models,” *SciPost Phys.* **13** no. 3, (2022) 069, [arXiv:2206.14763 \[hep-th\]](#).
- [15] A. Tsybaliuk, “The affine yangian of gl_1 revisited,” *Advances in Mathematics* **304** (2017) 583–645.
- [16] D. Maulik and A. Okounkov, “Quantum Groups and Quantum Cohomology,” [arXiv:1211.1287 \[math.AG\]](#).
- [17] T. Procházka, “ \mathcal{W} -symmetry, topological vertex and affine Yangian,” *JHEP* **10** (2016) 077, [arXiv:1512.07178 \[hep-th\]](#).
- [18] A. Morozov and N. Tselousov, “3-Schurs from explicit representation of Yangian $Y(\hat{\mathfrak{gl}}_1)$. Levels 1–5,” *JHEP* **11** (2023) 165, [arXiv:2305.12282 \[hep-th\]](#).
- [19] F. David, “Loop Equations and Nonperturbative Effects in Two-dimensional Quantum Gravity,” *Mod. Phys. Lett. A* **5** (1990) 1019–1030.
- [20] J. Ambjorn, J. Jurkiewicz, and Y. M. Makeenko, “Multiloop correlators for two-dimensional quantum gravity,” *Phys. Lett. B* **251** (1990) 517–524.
- [21] A. Mironov and A. Morozov, “On the origin of Virasoro constraints in matrix models: Lagrangian approach,” *Phys. Lett. B* **252** (1990) 47–52.
- [22] H. Itoyama and Y. Matsuo, “Noncritical Virasoro algebra of $d < 1$ matrix model and quantized string field,” *Phys. Lett. B* **255** (1991) 202–208.
- [23] A. Morozov and N. Tselousov, “Hunt for 3-Schur polynomials,” *Phys. Lett. B* **840** (2023) 137887, [arXiv:2211.14956 \[hep-th\]](#).
- [24] X. Liu and C. Yang, “Action of W-type operators on Schur functions and Schur Q-functions,” *J. Lond. Math. Soc.* **111** no. 2, (2025) e70080, [arXiv:2206.01066 \[math-ph\]](#).
- [25] M. Kontsevich, “Intersection theory on the moduli space of curves and the matrix Airy function,” *Commun. Math. Phys.* **147** (1992) 1–23.
- [26] A. Mironov and A. Morozov, “Superintegrability of Kontsevich matrix model,” *Eur. Phys. J. C* **81** no. 3, (2021) 270, [arXiv:2011.12917 \[hep-th\]](#).
- [27] A. Alexandrov, “Intersection numbers on $\overline{M}_{g,n}$ and BKP hierarchy,” *JHEP* **09** (2021) 013, [arXiv:2012.07573 \[math-ph\]](#).
- [28] Y. Drachov and A. Zhabin, “Genus expansion of matrix models and \hbar expansion of BKP hierarchy,” *Eur. Phys. J. C* **83** no. 5, (2023) 437, [arXiv:2302.03949 \[hep-th\]](#).
- [29] S. Kharchev, A. Marshakov, A. Mironov, A. Morozov, and A. Zabrodin, “Unification of all string models with $C < 1$,” *Phys. Lett. B* **275** (1992) 311–314, [arXiv:hep-th/9111037](#).

- [30] S. Kharchev, A. Marshakov, A. Mironov, A. Morozov, and A. Zabrodin, “Towards unified theory of 2-d gravity,” *Nucl. Phys. B* **380** (1992) 181–240, [arXiv:hep-th/9201013](#).
- [31] P. Di Francesco, C. Itzykson, and J. B. Zuber, “Polynomial averages in the Kontsevich model,” *Commun. Math. Phys.* **151** (1993) 193–219, [arXiv:hep-th/9206090](#).
- [32] A. Marshakov, A. Mironov, and A. Morozov, “On equivalence of topological and quantum 2-d gravity,” *Phys. Lett. B* **274** (1992) 280–288, [arXiv:hep-th/9201011](#).
- [33] A. S. Alexandrov, A. Mironov, A. Morozov, and P. Putrov, “Partition Functions of Matrix Models as the First Special Functions of String Theory. II. Kontsevich Model,” *Int. J. Mod. Phys. A* **24** (2009) 4939–4998, [arXiv:0811.2825 \[hep-th\]](#).
- [34] A. Alexandrov, “Cut-and-Join operator representation for Kontsevich-Witten tau-function,” *Mod. Phys. Lett. A* **26** (2011) 2193–2199, [arXiv:1009.4887 \[hep-th\]](#).
- [35] A. D. Mironov and A. Morozov, “Generalized Q-functions for GKM,” *Phys. Lett. B* **819** (2021) 136474, [arXiv:2101.08759 \[hep-th\]](#).
- [36] A. Morozov, “A new kind of anomaly: on W-constraints for GKM,” *JHEP* **21** (2020) 213, [arXiv:2108.07198 \[hep-th\]](#).
- [37] L. Alvarez-Gaume, H. Itoyama, J. L. Manes, and A. Zadra, “Superloop equations and two-dimensional supergravity,” *Int. J. Mod. Phys. A* **7** (1992) 5337–5368, [arXiv:hep-th/9112018](#).
- [38] G. Akemann and J. C. Plefka, “The Chiral superEigenvalue model,” *Mod. Phys. Lett. A* **12** (1997) 1745–1758, [arXiv:hep-th/9705114](#).
- [39] I. N. McArthur, “The Partition function for the supersymmetric Eigenvalue model,” *Mod. Phys. Lett. A* **8** (1993) 3355–3362.
- [40] P. Ciosmak, L. Hadasz, M. Manabe, and P. Sułkowski, “Super-quantum curves from super-eigenvalue models,” *JHEP* **10** (2016) 044, [arXiv:1608.02596 \[hep-th\]](#).
- [41] R. Wang, S.-K. Wang, K. Wu, and W.-Z. Zhao, “Correlators in the Gaussian and chiral supereigenvalue models in the Neveu-Schwarz sector,” *JHEP* **11** (2020) 119, [arXiv:2009.02929 \[hep-th\]](#).
- [42] D. Galakhov, A. Morozov, and N. Tselousov, “Super-Hamiltonians for super-Macdonald polynomials,” [arXiv:2501.14714 \[hep-th\]](#).

Appendix A. Another approach.

In both Hermitian and Kontsevich cases the resulting Hamiltonians have a very special property – the eigenvalue κ_λ corresponding to the diagram λ is given by sum over the all boxes:

$$\kappa_\lambda = \sum_{\square \in \lambda} \Omega_\square \tag{142}$$

where Ω_\square depends on the coordinates of the box (see (82) and (130)). Following an algorithm from [42] we can construct infinite set of commuting operators, starting from our Hamiltonians (81) and (133). We briefly remind the algorithm here. On the first step we consider two operators:

$$\hat{E}_0 \text{SIB}_\lambda = \sum_{\square \in \text{Add}(\lambda)} C_{\lambda, \lambda + \square} \text{SIB}_{\lambda + \square} \quad (143)$$

$$\hat{F}_0 \text{SIB}_\lambda = \sum_{\square \in \text{Rem}(\lambda)} \tilde{C}_{\lambda, \lambda - \square} \text{SIB}_{\lambda - \square} \quad (144)$$

In our cases the role of \hat{E}_0, \hat{F}_0 operators play the smallest time variable and corresponding derivative $p_1, \frac{\partial}{\partial p_1}$ and $\tau_0, \frac{\partial}{\partial \tau_0}$. On the second step we consider a tower auxiliary operators:

$$\begin{array}{ccc} & \hat{E}_0 & \hat{F}_0 \\ \hat{E}_1 = [\hat{H}, \hat{E}_0] & & \hat{F}_1 = [\hat{F}_0, \hat{H}] \\ \hat{E}_2 = [\hat{H}, [\hat{H}, \hat{E}_0]] & & \hat{F}_2 = [[\hat{F}_0, \hat{H}], \hat{H}] \\ \dots & & \dots \end{array} \quad (145)$$

that have the following action on polynomials:

$$\hat{E}_k \text{SIB}_\lambda = \sum_{\square \in \text{Add}(\lambda)} (\Omega_\square)^k \cdot C_{\lambda, \lambda + \square} \text{SIB}_{\lambda + \square} \quad (146)$$

$$\hat{F}_k \text{SIB}_\lambda = \sum_{\square \in \text{Rem}(\lambda)} (\Omega_\square)^k \cdot \tilde{C}_{\lambda, \lambda - \square} \text{SIB}_{\lambda - \square} \quad (147)$$

On the final step we consider the following operators:

$$\hat{\mathcal{H}}_{a+b} := [\hat{E}_a, \hat{F}_b] \quad (148)$$

These operators have remarkable properties – they depend only on the sum of indices $a + b$, form commuting family and commute with starting Hamiltonian \hat{H} :

$$[\hat{\mathcal{H}}_a, \hat{\mathcal{H}}_b] = 0 \quad [\hat{\mathcal{H}}_a, \hat{H}] = 0 \quad (149)$$

Therefore, our families of polynomials SIB_λ are their eigenfunctions:

$$\hat{\mathcal{H}}_a \text{SIB}_\lambda = \mathcal{E}_{\lambda, a} \cdot \text{SIB}_\lambda \quad (150)$$

Basing on the above reasoning we propose alternative approach to obtain SIB_λ polynomials in Hermitian and Kontsevich cases:

- We start from fixed-spin ansatz for Hamiltonian (50) and (114), that depend on unknown coefficients $(A_{a,b}, B_{a,b})$ and $(A_{a,b,c}, B_{a,b,c}, C_{a,b,c,d}, G_a)$ respectively.
- Using ansatz for Hamiltonians we compute auxiliary operators \hat{E}_k, \hat{F}_k and then $\hat{\mathcal{H}}_{a,b}$.
- We impose the following conditions:

$$\hat{\mathcal{H}}_{a,b} = \hat{\mathcal{H}}_{a+b} \quad (151)$$

$$[\hat{\mathcal{H}}_a, \hat{H}] = 0 \quad (152)$$

$$[\hat{\mathcal{H}}_a, \hat{\mathcal{H}}_b] = 0 \quad (153)$$

that determine special relations on coefficients $(A_{a,b}, B_{a,b})$ and $(A_{a,b,c}, B_{a,b,c}, C_{a,b,c,d}, G_a)$.

- We observe that these nonlinear system of equations have unique solution up to rescaling of time variables. SIB polynomials are restored up to overall scaling as a common set of eigenfunctions.