Probing the axial symmetry of gamma-ray burst jets using afterglow polarimetry

Thomas Baxter^{[0,1,*}, Shiho Kobayashi,¹

¹Astrophysics Research Institute, Liverpool John Moores University, IC2, Liverpool Science Park, 146 Brownlow Hill, Liverpool L3 5RF, UK

Accepted XXX. Received YYY; in original form ZZZ

ABSTRACT

Polarisation measurements of gamma-ray burst afterglows provide a powerful tool for probing the structure of relativistic jets. In this study, we revisit polarisation signals observed in gamma-ray burst afterglows, focusing on the effects of non-axisymmetric jet structures. To characterize these non-axisymmetric jets, we adopt a simple elliptical jet head model and investigate how deviations from axisymmetry influence the temporal evolution of polarisation properties, particularly around the jet break. Our results show that the polarisation degree curve typically exhibits two peaks for top-hat jets or a single peak for structured jets, even in the presence of an elliptical jet head. In non-axisymmetric jets, a complete drop in polarisation between peaks is generally absent, and the position angle rotation between the peaks can deviate significantly from 90 degrees. In single-peak cases, the polarisation position angle evolves gradually, contrasting with the constant position angle expected in axisymmetric jets. We also explore the implications of these findings for recent GRB events, including GRB 121024A, GRB 091018, GRB 020813, and GRB 210610B.

Key words: gamma-ray bursts - polarization - radiation mechanisms: non-thermal - relativistic processes - magnetic fields.

1 INTRODUCTION

The afterglow of a gamma-ray burst (GRB) emerges as the relativistic jet decelerates due to its interaction with the surrounding circumburst material. This interaction generates a forward shock that propagates as a collimated blast wave. Electrons accelerated within these shocks produce the observed non-thermal emission via synchrotron radiation (Sari et al. 1998).

The simple top-hat jet model has been remarkably successful in explaining many GRB afterglow observations. This success stems from the fact that, for an on-axis observer, the emission from the jet core dominates, resulting in an afterglow light curve that shows no distinct features indicative of the jet's structure. However, the groundbreaking gravitational-wave-triggered detection of a binary neutron star merger, linked to GRB 170817A (Abbott et al. 2017), demonstrated that the afterglows of off-axis events are highly sensitive to the structure of the jet (e.g., Lamb & Kobayashi 2017). Observations have shown that the late-time afterglow light curve of GRB 170817A is consistent with Gaussian jet models (Lazzati et al. 2018; Margutti et al. 2018; Ghirlanda et al. 2019; Troja et al. 2019; Lamb et al. 2019). Meanwhile, a shallow angular profile in relativistic jet models (e.g., power-law jets) has been proposed for GRB 221009A, the brightest burst ever observed (Lesage et al. 2023; Burns et al. 2023). The observed monochromatic steepening of the X-ray and optical light curves strongly suggests a geometrical effect, such as a jet break, although the post-break decay rates are shallower than expected. This discrepancy can be resolved if the afterglow emission originates from a structured jet with a shallow angular energy profile (O'Connor et al. 2023; Birenbaum et al. 2024).

Numerical simulations suggest such jet structures can arise from the interaction between the jet and the confining medium (Gottlieb et al. 2021). Specifically, this occurs as the jet breaks out through the stellar envelope in the collapsar model (e.g. Zhang et al. 2003; Lazzati & Begelman 2005; Gottlieb et al. 2021); or interacts with merger ejecta (e.g. Perego et al. 2014; Nativi et al. 2022). While most afterglow studies to date have focused on axisymmetric jets (either top-hat or structured), some works have explored with uneven energy and velocity distributions in the azimuthal direction (Li et al. 2023; Li et al. 2024), which discuss uneven distribution of energy and velocity within a jet in the azimuth angle direction due to internal non-uniform magnetic dissipation processes or the precession of the central engine (Huang et al. 2019). Recent three-dimensional hydrodynamic simulations of jets in the aftermath of neutron star mergers (Gottlieb et al. 2022; Lamb et al. 2022) further highlight these complexities. These studies show that when a jet breaks out of the ejecta, the resulting outflow exhibits a power-law-like polar energy distribution with rotational inhomogeneity. Notably, the jet head is not perfectly circular, as often assumed in conventional theoretical models, instead, it tends to exhibit a more elliptical shape.

Non-axisymmetric or inhomogeneous structures have been primarily discussed in the context of prompt gamma-ray emission (e.g. Gill & Granot 2024; Lazar et al. 2009). This emission originates from the original ejecta from the central engine, which carries large-scale magnetic fields generated at the central source. While polarisation signals in the prompt gamma-rays are more sensitive to the magnetic field structure within the jet, those in the afterglow phase are predominantly influenced by geometrical effects. Consequently, af-

Email: aritbaxt@ljmu.ac.uk




Figure 1. The shaded area represents the physical extent of the jet head, with the line of sight intersecting it at Point A. The location of Point A is specified by two angles: viewing angle θ_{view} , measured from the jet axis (or equivalently expressed as $\xi = (\theta_{view}/\theta_j)$, and the azimuthal angle χ , measured from the semi-major axis of the jet head. The ring surrounding Point A marks the visible region. Red double-headed arrows indicate the polarisation directions of fluid element emission. The point symmetric to Point A with respect to the jet axis (the origin) is Point B, while the point symmetric to Point A with respect to the x-axis is labelled Point C.


terglow polarimetry offers a more effective means of probing the jet's structure and geometry.

Although the mechanisms behind the acceleration and collimation of GRB jets remain uncertain, understanding their structures can provide valuable insights into these processes. This paper explores how jet asymmetries can be constrained through polarisation observations of afterglows. Our study is partly motivated by recent polarisation measurements of GRB 210610B (Agüí Fernández et al. 2024), which revealed a polarisation position angle (PPA) rotation of $\Delta \phi = 54^{\circ} \pm 9^{\circ}$ around a possible jet break (see section 5.4 for details). This rotation deviates significantly from $\Delta \phi = 90^{\circ}$ predicted for axisymmetric jets (Sari 1999; Ghisellini & Lazzati 1999).

We aim to investigate the signature of jet axial asymmetry imprinted on the temporal evolution of afterglow polarisation. To break the axisymmetry, we model the global jet structure with an elliptical head. While real jets may exhibit more complex geometries, this approach represents a natural starting point for exploring nonaxisymmetric jets in this context. We present our model for evaluating polarisation signals around a jet break in Section 2, where we illustrate the primary polarisation signature of a non-axisymmetric jet. We describe the distributions of polarisation signals when the lineof-sight of an observer intersects the jet head (or shock surface) at random locations in Section 3. In Section 4, we discuss the structured model in the context of non-axisymmetric jets. In Section 5 we present case studies. Finally, in Section 6, we give our conclusions and discussion.

2 POLARISATION BEHAVIOUR AT A JET BREAK

In the afterglow phase, magnetic fields in the blast wave are believed to be produced by shock instabilities, and expected to be highly tangled. However, the precise mechanisms by which magnetic fields are produced in relativistic collisionless shocks remain poorly understood. Since the shock normal (aligned with the radial direction) represents a distinct orientation, the magnetic field components parallel and perpendicular to the shock normal could exhibit



0.3

Figure 2. Top panel: Polarisation as a function of time for e = 0 (blue dashed) and 0.6 (red solid). The polarisation scales almost linearly with the local polarisation degree P_0 , and we have assumed $P_0 = 0.7$. The first peaks, the troughs and second peaks in the polarisation light curves are marked by circles, crosses and squares, respectively. The line of sight is characterised by $\chi = 1$ and $\xi = 0.3$. Bottom panel: Same as the top panel, but showing the the evolution of polarisation signals in the Stokes parameter space (q, u).

significantly different average strengths (Medvedev & Loeb 1999; Sari 1999; Ghisellini & Lazzati 1999).

Due to relativistic beaming, an observer at early times can only see a small portion of the jet head's surface. This visible region is confined to a small area with an angular size of $1/\Gamma$, centred around the point where the line of sight intersects the jet head. Especially, at high frequencies (above the peak synchrotron frequency), most of the emission arises from a ring-like structure, caused by the relativistic limb-brightening effect (Granot et al. 1999). Each small segment of the ring might produce highly polarized radiation due to the local anisotropy in the magnetic field, as discussed. However, the overall symmetry of the ring results in a net polarisation of zero.

As noted independently by Sari 1999 and Ghisellini & Lazzati 1999, net polarisation signals are expected to emerge around the time of a jet break, when the expanding visible region (i.e., the ring) begins to extend beyond the edge of the jet. We here revisit the polarisation signals for a non-axisymmetric homogeneous (top-hat) jet.

Following Sari 1999, we adopt a toy model to approximate the evolution of the polarisation signal. In this model, (1) the jet head is represented by an ellipse with eccentricity e, and the half-opening angle of the jet, particularly along the semi-major axis, is denoted by θ_j . (2) Recent numerical hydrodynamic simulations suggest that lateral jet expansion occurs much more slowly than previously estimated (e.g., Granot & Piran 2012). Thus, we assume the lateral expansion is negligible, meaning both the jet opening angle and eccentricity remain constant. The Lorentz factor of the fluid is related to the observer time t by $t/t_j = (\theta_j \Gamma)^{-8/3}$ where t_j is the characteristic time for a jet break. (3) The (main) visible region is modelled as a thin ring with a radius of Γ^{-1} , centred on the point where the line of sight intersects the jet head. The ring's width is set to 30% of its radius. (4) The emission from each fluid element within the ring is

polarized in the local radial direction (i.e., perpendicular to the ring) with a polarisation degree of P_0 . This polarisation direction assumes that the magnetic field component parallel to the shock normal dominates over the perpendicular component. If the parallel component were weaker, the Stokes parameters Q and U would change sign in the subsequent discussion. However, the discussion is basically identical. (5) The inner region of the ring, corresponding to a circular area with a radius of 0.7 Γ^{-1} , emits much dimmer radiation—10% of the ring's brightness-and is assumed to be unpolarised. (6) Only the portion of the visible region (the ring and the inner circular area) that overlaps with the jet emits photons, uniformly distributed within each area. The region outside the jet does not contribute any emission. Under these assumptions, the evolution of polarisation over time depends on the location of the observer's line of sight within the jet head, characterized by two parameters: the azimuthal angle from the semi-major, χ , and the offset between the observer's line of sight and the centre of the jet, measured in units of the jet opening angle, $\xi = \theta_{view}/\theta_i$.

To estimate the Stokes parameters: Q and U, we divide the visible region (the ring and the inner circle) into many segments. Since we consider synchrotron emission from an optically thin blast wave, the circular polarisation is zero V = 0. Each segment contributes to Q and U as

$$dQ = P_e dL \cos\left(2\phi_e\right) \tag{1}$$

$$dU = P_e dL \sin\left(2\phi_e\right) \tag{2}$$

where the local polarisation degree $P_e = P_0$ when the segment lies within the ring, and $P_e = 0$ when the segment is inside the inner circle. The local luminosity, dL, is proportional to the area of the segment while also incorporating the constraints imposed by conditions (3), (5), and (6) in the previous paragraph. The angle ϕ_e denotes the angle between the positive x-axis (the semi-major axis) and the segment's position vector, measured relative to the light of sight (Point A in Figure 1). Considering condition (6), ΣdQ , ΣdU and ΣdL are evaluated to estimate the net values of the parameters: $q = \Sigma dQ/\Sigma dL$ and $u = \Sigma dU/\Sigma dL$. The net polarisation degree P and position angle ϕ are given by

$$P = \sqrt{q^2 + u^2},\tag{3}$$

$$\phi = \frac{1}{2}\arctan\left(u/q\right). \tag{4}$$

Figure 2 illustrates the differences in polarisation signals between axisymmetric jets and non-axisymmetric jets (in this paper, nonaxisymmetric jets refer to those with elliptical-shaped heads). For axisymmetric jets, the polarisation degree curve (shown as the blue dashed line in the top panel) exhibits two peaks, with the polarisation dropping to zero between the peaks around the jet break time. As seen in the bottom panel, the time evolution of $(q, u) \propto (\cos 2\phi, \sin 2\phi)$ follows a linear trajectory passing through the origin. The polarisation position angle (PPA) ϕ rotates precisely by $\pi/2$ before and after the point where polarisation vanishes, around the jet break time (Sari 1999; Ghisellini & Lazzati 1999). In contrast, jets with elliptical heads generally do not exhibit a complete polarisation drop between the two peaks ($P \neq 0$ at the trough). The PPA rotates more smoothly, though it still undergoes rapid changes near the trough (approximately at the jet break time). The change in PPA between the two peaks can significantly deviate from $\pi/2$; in this example, it is approximately $\Delta \phi = 0.28\pi \sim 50.4$ degrees (it is more convenient to express angles in radians for theoretical discussion, while degrees are preferred in the context of observations. Unless otherwise specified, angles are given in radians).




Figure 3. Distribution functions of deviations from $\pi/2$ in the PPA rotation between polarisation peaks for uniformly sampled jets with eccentricities of e = 0.3 (blue lines), e = 0.45 (orange lines), and e = 0.6 (green lines). Solid lines represent probability density distributions, while dashed lines show their corresponding cumulative distributions.




Figure 4. The PPA deviations from $\pi/2$ are shown in 16 bins of equal angular width χ for jets with eccentricities e = 0.3 (blue), e = 0.45 (orange), e = 0.6 (green). The dot represent the averaged values, and the bars indicate the 1 σ spreads for each bin.

3 THE CHARACTERISATION OF POLARISATION SIGNALS

Consider an observer whose line of sight intersects a jet head, which is characterised by an eccentricity e, at the point (χ, ξ) . This intersection point is labelled as Point A in Figure 1. To characterise the time evolution of the PPA, we examine the PPA rotation $\Delta\phi$ between two polarisation peaks and the ratio $R_{pd} = P_{1st}/P_{tr}$, which quantifies the polarisation degrees at the first peak relative to the trough. In Figure 2, circles and squares denote these polarisation peaks around a jet break. In particular, we analyse $|\Delta\phi - \pi/2|$, which measures deviations from the expected $\pi/2$ rotation in axisymmetric jets. As discussed in section 2, non-axisymmetric jets generally do not exhibit a complete polarisation drop between the two peaks, resulting in lower values of R_{pd} .

We investigate the distribution of polarisation signals under the assumption that the intersection points of an observer's line of sight with the jet head are uniformly distributed. Due to the symmetry of the system, it is unnecessary to sample the entire ellipse ($-\pi < \chi <$




Figure 5. The deviation of the PPA rotation from $\pi/2$ and the peak-to-trough ratio R_{pd} are shown for the same random sample (e = 0.45) that generates the orange lines in Figure 3. The colour map indicates the offset ξ value of each sample.

 π); instead, it is sufficient to consider a single quadrant ($0 < \chi < \pi/2$).

To illustrate this, we first introduce Point B, which is symmetric to Point A with respect to the origin. The distribution of fluid elements around Point B (or the relative position of the jet edge) is identical to that around Point A when rotated by π around Point B. Consequently, given the angular dependences in Eqs. (1) and (2), the polarisation signals at any given time are therefore identical for Points A and B.

Similarly, consider Point C, which is symmetric to Point A with respect to the x-axis. The fluid element distribution around Point C mirrors that of Point A but with a reflection of $\phi \leftrightarrow -\phi$ (flipping vertically with respect to the line passing through Point A and parallel to the x-axis). While the absolute values of the Stokes parameters at Point C match those at Point A, one of them flips sign, specifically (q, -u). Nevertheless, the position angle rotation $|\Delta \phi - \pi/2|$ and the ratio R_{pd} remain the same as at Point A.

The jet head is a part of a spherical blast wave, but the ellipse represented in Figure 2 represents its projection onto the sky. The opening angle of GRB jets or their core size is typically only several degrees. For such small angles, uniform sampling over projected area is approximately equivalent to uniform sampling over the solid angle of the shock surface.

Figure 3 shows the distributions of $|\Delta \phi - \pi/2|$, derived from 5000 uniform sampled line-of-sight realisations across a quadrant of the elliptic jet head for eccentricity e = 0.3, 0.45, and 0.6. For axisymmetric jets (e = 0), the probability distribution is given by a delta function at 0, meaning the PPA rotation $\Delta \phi$ is always $\pi/2$. As eccentricity increases, the probability distribution (solid lines) peaks at larger deviation angles, with the distribution's spread also increasing. The probability density distributions peak at $|\Delta \phi - \pi/2| = 4.3, 9.5$ and 19 degrees, while their cumulative distributions reach 50% at 3.8, 8.9, 18 degrees, and 90% at 7.3, 17, 31 degrees for e = 0.3, 0.45and 0.6, respectively. Since the semi-minor axis is smaller than the semi-major axis by a factor of $\sqrt{1-e^2}$, it is reduced by approximately 5%–20% (or equivalently by a factor of $\sim 0.8 - 0.95$) across this range of eccentricities. Our results show that even a mild deformation of the jet head leads to PPA rotations that deviate significantly from $\pi/2$.

When the line of sight intersects the jet head near its semi-major or

semi-minor axis, the geometry exhibits greater symmetry, leading to a position angle rotation close to $\pi/2$. As expected, Figure 4 shows that the deviation from $\pi/2$ increases with higher eccentricity, particularly for intermediate angles ($\chi \sim \pi/4$). The spread in the probability distributions shown in Figure 3 can be partly attributed to variations in PPA rotation between the more symmetric configurations ($\chi \sim 0$ or $\pi/2$) and the asymmetric configuration ($\chi \sim \pi/4$). In Figure 4, for e = 0.6, the azimuthal bin nearest the semi-minor axis ($\chi \sim 90$ deg) results in negative values for 1σ error. This is due to the significant positive skewness of the distribution.

Additionally, the offset $\xi = \theta_{view}/\theta_i$ also influences the rotation of the position angle. Figure 5 shows $|\Delta \phi - \pi/2|$ and R_{pd} for the same random uniform samples with e = 0.45, where the colour map represents the offset ξ for each sample. For a smaller offset (and intermediate χ), the deviation $|\Delta \phi - \pi/2|$ becomes larger (if the line of sight passes regions near the semi-major or semi-minor axis, the rotation is close to $\pi/2$ regardless of the offset values). The high-deviation tails in the probability density distributions (the solid lines), seen above their peaks in Figure 3, correspond to low- ξ cases $(\xi \lesssim 0.4)$ where the light of sight is relatively close to the central jet axis. At small offsets ξ , the first peak of the polarisation curve becomes less pronounced, leading to a lower R_{pd} . For eccentricity e = 0.3, 0.45 and 0.6, the top 10% of cases with the largest $\Delta \phi$ deviation from $\pi/2$ have R_{pd} values below approximately 86, 38, and 22. Although these extreme cases show lower R_{pd} for higher eccentricities, a jet head with a larger eccentricity e has a higher probability of producing a significant $\Delta \phi$ deviation from $\pi/2$. As discussed in section 5.4, observational measurements of R_{pd} and $\Delta \phi$ can provide valuable constraints on the eccentricity *e*.

For even smaller offsets, the first peak becomes comparable to the trough, resulting in $R_{pd} \sim 1$. To determine whether a polarisation curve exhibits two peaks and to locate them, we have implemented a peak-finding algorithm similar to the one discussed in Li & Fenimore 1996 (see Appendix A for more details). For approximately, 4%, 7% and 11% of uniform random realisations for the e = 0.3, 0.45 and 0.6, respectively, the polarisation curves exhibit only a single peak. These single peak cases have been excluded from the analysis presented in Figs 3, 4 and 5.

4 NON-AXISYMMETRIC STRUCTURED JET

Observations of GRB 170817A demonstrate that some GRB jets deviate from a simple top-hat structure, instead exhibiting a structured profile. To capture this complexity, we extend our model to incorporate a structured luminosity distribution. In this framework, a uniform central core is encircled by a fainter outer region, where the surface luminosity density gradually declines following a power-law profile, extending out to the jet's maximum opening angle.

We assume that the core, the jet edge and brightness contours all have a common elliptical shape with a common eccentricity e, where the jet core θ_c and the jet outer edge angles θ_j are defined along the semi-major axis. Considering the relativistic limb-brightening effect, we adopt the same ring-shaped visible region as in the top-hat jet case. The procedure for estimating polarisation signals remains unchanged, but the local luminosity dL of a segment is now determined by a position-dependent surface luminosity density, which is highest in the core and decreases as a power law with index α in the outer region.

The luminosity reduction factor for a segment in the outer region,




Figure 6. Polarisation signals for a line of sight intersecting the outer region, where the surface luminosity density declines following a power-law with an index of $\alpha = 2$ (red) or 3 (blue). Two offset angles are considered: $\xi = 2\theta_c/\theta_j$ (solid) and $4\theta_c/\theta_j$ (dashed). Top panel: temporal evolution of polarisation degree. Bottom panel: the evolution of the Stokes parameter (q, u). The model assumes e = 0.3, $\chi = \pi/4$, and $\theta_c/\theta_i = 0.13$.




Figure 7. Polarisation signals for a line of sight intersecting the jet core at $\xi = 0.5\theta_c/\theta_j$. All jet properties are identical to those in Figure 6, except where specified. Top panel: polarisation degree curves. Bottom panel: evolution of the Stoke parameters. Curves are shown for $\alpha = 2$ (red) and $\alpha = 3$ (blue). The first peaks, the troughs and second peaks in the polarisation light curves are marked by circles, crosses and squares, respectively.

located at (x, y) on the jet head is given by

$$\left(\frac{x^2}{a_c^2} + \frac{y^2}{b_c^2}\right)^{-\alpha/2} \quad , \tag{5}$$

where $a_c = \theta_c$ and $b_c = \theta_c \sqrt{1 - e^2}$ are the semi-major and semiminor axes of the elliptical jet core.

Figure 6 shows the polarisation signals when the line of sight falls in the outer region. As demonstrated by Rossi et al. 2004 and Birenbaum et al. 2024, the polarisation curves differs significantly from those of a top-hat jet, regardless of axial symmetry. This discrepancy arises from the non-uniform luminosity distribution in the visible region surrounding the line of sight (i.e., the ring), with the intense emission concentrated toward the jet axis. Consequently, the observed polarisation signal is primarily influenced by these segments of the ring, resulting in a single peak in the polarisation curve, which occurs around the time the core becomes visible to the observer (top panel). The polarisation degree curves for the decay indices $\alpha = 2$ and 3 exhibit similar shapes.

Since the the emission from the ring segments closer to the jet axis dominates the polarisation signals, the position angle evolution also differs from those of a top-hat jet. It consistently aligns with the direction toward the central jet axis (see assumption (4) in Section 2). When the line of sight falls on the jet head near its semi-major or semi-minor axis ($\chi \sim 0$ or $\pi/2$), the geometry becomes more symmetric, the position angle remains constant over time, consistent with the results for axisymmetric structured jets (Rossi et al. 2004). However, for a general azimuthal angle χ , the eccentricity of the brightness contour lines leads to a slight evolution of the position angle around the peak of the polarisation degree. This effect is illustrated for $\chi = \pi/4$ in the bottom panel of Figure 6.

The PPA rotation over the full width at half maximum of the polarisation peak is evaluated for $\chi = \pi/4$ (the rotation is maximised for $\chi = \pi/4$). We obtain rotation values of $\Delta \phi = 3.5$, 2.9, 2.6 degrees for e = 0.3, $\Delta \phi = 16$, 13, 12 degrees for e = 0.6, corresponding to $\xi/\langle \theta_c/\theta_j \rangle = 2, 3, 4$, respectively. While these values are evaluated for $\alpha = 3$, the rotation remains insensitive to α , with slightly larger values obtained for $\alpha = 2$.

As discussed in Section 2, the polarisation degree curves exhibit a single peak even for top-hat, non-axisymmetric jets when the line of sight is close to the jet axis. In such cases, the position angle rotations are more pronounced, reaching at least, several tens degrees for our samples with e = 0.3, 0.45 or 0.6.

Figure 7 illustrates polarisation signals when the line of sight passes through the jet core, corresponding to $\xi = 0.5\theta_c/\theta_j$ and $\chi = \pi/4$. The polarisation curves resemble those of top-hat, non-axisymmetric jets (represented by the red solid line in the top panel of Figure 2), exhibiting two peaks without a complete polarisation dip in between. The trajectories in the (q, u) space (the bottom panel) and the PPA rotation between two peaks show little sensitivity to α . Specifically, for $\alpha = 2$ and 3, we obtain $\Delta \phi \sim 79$ degrees and 80 degrees, respectively. In the limit $\alpha \gg 1$ (i.e., the top-hat model), the rotation angle increases slightly to $\Delta \phi \sim 83$ degrees. Overall, the top-hat model serve as a reasonable approximation of the polarisation signals when the line of sight intersects the jet core.

5 CASE STUDIES

Optical linear polarisation measurements have been conducted for many GRB afterglows, typically several hours to a few days after the prompt gamma-ray emission (Covino & Gotz 2016, and references therein). This timeframe corresponds to when a jet break is expected to occur. The observed linear polarisation degree or its upper limits are generally low (\lesssim several percent). Such low polarisation levels suggest that in shock-generated random magnetic fields, neither the component parallel nor perpendicular to the shock normal dominates entirely (e.g., Gill & Granot 2020), though the fields are not fully isotropic. Our model, which assumes a low local polarisation degree parameter P_0 , should be applicable to modelling afterglow polarisation measurements.

5.1 GRB 020813

A bright long-duration burst (> 125 s) detected by HETE-2, located at a redshift of z = 1.255 (Barth et al. 2003), was studied. Polarimetry data obtained with VLT and Keck (Gorosabel et al. 2004) was collected around a jet break, occurring at ~ 13 hours post-burst (Covino et al. 2003). The data is most consistently described by an almost constant degree of linear polarisation at the $\sim 1\%$ level, along with a stable position angle. While a slow evolution of the position angle $\Delta \phi \sim$ a few tens of degrees cannot be ruled out, the observations do not support a sudden 90 degree rotation of the position angle at the jet break, which is a signature of top-hat axisymmetric jets. Given the sparsity of polarisation data points, a single-peaked polarisation curve remains consistent with the observations. The evolution of both the polarisation degree and position angle can explained by structured jets, whether axisymmetric or not. If a gradual rotation of the position angle were confirmed, it would indicate a structured, non-axisymmetric jet.

5.2 GRB 091018

This long burst ($T_{90} = 4.4$ sec and redshift z = 0.971) was detected by Swift (Wiersema et al. 2012). Its optical light curve is well described by a broken power-law, with a break at ~ 9 hours, coinciding with a steepening in the X-ray light curve. While the post-break decay indices (1.54 in the X-ray and 1.33 in the optical) are shallower than predicted from the standard fireball model, similar trends have been observed in other Swift XRT afterglows, where potential jet breaks often exhibit relatively shallow decay indices (Racusin et al. 2009).

Linear polarisation monitoring with VLT (Wiersema et al. 2012) revealed a polarisation curve inconsistent with a single-peaked profile, and instead favouring a two-peak structure with a trough around the suspected jet break. In top-hat jet models, both axisymmetric and non-axisymmetric, the second peak is expected to have a higher amplitude than the first. Although some data points support this trend, significant scatter suggests the presence of a deep, short dip within the second peak. The PPA remains stable and nearly constant during the first peak, but shifts to a higher overall value in the second peak, with significant scatter around the 90-degree rotation. These variations in both the polarisation degree and position angle during the second peak are not fully explained even by non-axisymmetric jets (i.e., elliptical jet heads). As noted by Wiersema et al. 2012, if the scatter in the second peak is caused by an additional component (e.g., a bright patch within the jet or microlensing effects), the presence of a two-peak profile, combined with a nearly constant PPA in the first peak, suggests either a top-hat jet or a view of the core of a structured jet. For a non-axisymmetric jet, the line of sight must be positioned near either the semi-major or semi-minor axis of the jet head. Additionally, the shallow post-break decay indices may imply that the jet core is encased by an outer region with a shallow angular profile.

5.3 GRB 121024A

A long burst with $T_{90} = 69$ sec was detected by Swift, with a redshift of z = 2.298 determined shortly afterwards. The X-ray light curve follows a three-segment power-law evolution, with the final break at $t \sim 10$ hours coinciding with a break in the optical light curve (Wiersema et al. 2014). The monochromatic nature of this break suggests a jet break origin. However, the post-break decay indices remain relatively shallow, at 1.67 in the X-rays and 1.25 in the optical. VLT polarimetry measurements taken 3–6 hours after the burst show linear polarisation degrees ranging from 3–5%, with a stable position angle. However, observations from the following night (after the jet break) reveal a significant shift in the position angle, consistent with a 90 degree rotation (Wiersema et al. 2014). These polarisation properties also support the same jet structure interpretations as for GRB 091018.

5.4 GRB 210610B

A long-duration burst at z = 1.134 was detected by Fermi ($T_{90} =$ 69 sec) and Swift ($T_{90} = 55$ sec). The optical and X-ray light curves are modelled using a broken power-law. The optical light curve remains initially flat until the break at \sim 7.8 hours, after which it steepens to a decay index of 1.85. The X-ray light curve appears to undergo a simultaneous break and shares the same post break decay index, thought its pre-break decay is faster than that of the optical counterpart (Agüí Fernández et al. 2024). Polarimetric observations were conducted with the Calar Alto Telescope and the VLT. The first measurements, taken at around 2.9 hours, revealed a relatively high polarisation degree of $\sim 4.5\%$. During the second observation period (5.8-6.7 hours), which is close to the break time, the polarisation level dropped significantly to 0.18-0.6%. However, after the break, the polarisation degree increased again to $\sim 2\%$ (30.4–30.6 hours). Interestingly, the PPA rotates by 54 degrees between the pre- and post-break measurements, suggesting a significant shift that deviates notably from the 90 degree rotation expected for axisymmetric jets.

This event is a strong candidate for modelling with a nonaxisymmetric jet. Here, we assume top-hat, non-axisymmetric jets for our analysis. While a structured, non-axisymmetric jet could also explain the event if the line of sight intersects the jet core, the constraint on the jet head's eccentricity remain similar. Given the large deviation of $|\Delta \phi - \pi/2| = 36$ degrees, and the distinct trough structure $R_{pd} \ge 10$, we performed 5000 random samplings. For a lower eccentricity (e = 0.3), most cases (87%) yield $R_{pd} > 10$. However, virtually none satisfy $|\Delta \phi - \pi/2| > 36$ degrees. For e = 0.75, the fraction of cases meeting the first, second and both conditions are 64%, 35% and 21%, respectively. The lines of sight that satisfy both conditions are characterised by intermediate azimuthal angles: $0.4 < \chi < 1.5$ and a relatively central region: $\xi < 0.7$. These findings suggest this event could be explained within a non-axisymmetric jet framework. However, there are caveats in the jet-break modelling. 1) The flat optical light curve before the break suggests significant energy injection into the forward shock at early times. Since ejecta from the central engine can carry large-scale magnetic fields (Mundell et al. 2013) unlike forward shocks, which generates highly tangled magnetic fields, the polarisation signals before the break may be significantly influenced by emission from refreshed shocks (energy injection). 2) the post-break decay index of the optical afterglow (1.85 ± 0.04) is close to, but still shallower than, the expected value (~ 2) for the standard fireball model (Agüí Fernández et al. 2024).

6 CONCLUSIONS

We have investigated the afterglow polarisation signals around a jet break for non-axisymmetric jets. For jets with top-hat luminosity density distributions, the polarisation degree curves exhibits two peaks in general, even when the jet head is elliptical. However, a complete polarisation drop between the peaks is generally absent. Unlike axisymmetric jets, where the polarisation position angle (PPA) undergoes a sudden jump at the trough, the PPA in non-axisymmetric jets evolves smoothly. Additionally, the change in PPA between the two peaks can deviate significantly from 90 degrees. In our random line-of-sight sampling, 50% of the cases show deviations of more than 18 degrees, and 10% exceed 31 degrees for e = 0.6.

If the jet's luminosity density gradually declines following a powerlaw profile, the polarisation curves exhibit a single peak. We find the shapes of the curves are similar for $\alpha = 2$ and 3. In this structured jet case, the PPA can show a slight evolution for non-axisymmetric jets, in contrast to a constant PPA expected in axisymmetric jet models. Over the full width at half maximum of the polarisation peak, the PPA rotates by 16 degrees for e = 0.6 with a line of sight at $\chi = \pi/4$ and $\xi = 2(\theta_c/\theta_j)$. The results remain consistent for $\alpha = 2$ and 3.

For structured, non-axisymmetric jets, in which a uniform central core is surrounded by an outer region with a power-law luminosity profile, the polarisation signals depends on the line-of-sight location. This model can account for a wide range of polarisation behaviours, as discussed in the case study section. When the line of sight intersects the uniform jet core, the signals resemble those of top-hat jets. If the line of sight passes through the jet head - whether in the core or outer region - near its semi-major or semi-minor axis, the geometry becomes more symmetrical, making the polarisation signals of non-axisymmetric jets similar to those of axisymmetric jets. Significant deviations in the PPA rotation from 90 degrees (or from a constant PPA in the case of a structured jet) arise when the line of sight has an intermediate azimuthal angle ($\chi \sim \pi/4$) and a relatively small offset angle ($\xi \lesssim 0.4$), i.e., when it is close to the jet axis.

As a simple toy model for a structured jet, we considered a case where the luminosity density in the jet's outer region follows a powerlaw profile. While this simplified model should captures the key features of polarisation signals in structured jets, the angular dependence of luminosity density in a more realistic scenario arises from the lateral energy distribution - where the jet core has a higher energy per solid angle, and the outer regions gradually contain less energy per solid angle. This also induces the angular variations in the Lorentz factor, with the core having a higher Lorentz factor and progressively decreasing toward the outer regions. As a result, the visible region is no longer circular symmetric, toward the jet centre, the relativistic beaming is stronger and luminosity contribution is larger (e.g., Figure 2 in Beniamini et al. 2022). The impact of this deformed visible region will be explored in a future paper.

When constructing the polarisation degree light curves shown in Figs 2, 6 and 7, we assumed the forward shock dynamics $\Gamma \propto t^{-3/8}$ for the ISM. In the wind medium, the curve shapes are slightly modified due to a different forward shock dynamics $\Gamma \propto t^{-1/4}$. However, the overall morphology (e.g. the presence of two peaks or a single peak) remains unchanged and the discussion on $\Delta \phi$ should be similar.

If the jet head has a highly eccentric shape, it can be characterised by two angular scales: θ_j (the opening angle along the semi-major axis) and θ_{j2} (along the semi-minor axis). As long as $\Gamma > 1/\theta_{j2}$, the flux follows the typical blast wave emission evolution, decaying approximately as t^{-1} . When $\Gamma \propto t^{-3/8}$ drops below this first critical threshold, photons begin to be emitted over a larger solid angle, increasing from $\theta_j \theta_{j2}$ to $\Gamma^{-1} \theta_j$, causing the flux to decay as $t^{-11/8}$. As Γ continues to decreases and falls below the 2nd threshold $1/\theta_j$, the emission solid angle expands further to Γ^{-2} , leading to a steeper decay of $t^{-7/4}$. Since these breaks arise from geometrical effects, they are expected to be monochromatic, similar to the usual jet break. The two jet break times t_j and t_{j2} , with $t_{j2} < t_j$, satisfy the relation

$$t_{j2}/t_j \sim \left(1-e^2\right)^{4/3}.$$
 (6)

For axisymmetric jets (e = 0), it simplifies to the standard single

jet break. For a mildly deformed jet head with e = 0.6, the ratio is approximately $t_{j2}/t_j = 0.55$. As a result, the breaks will appear as a smooth transition rather than two distinct ones. While we have assumed a top hat-jet in the ISM to estimate the break time ratio, a wind-like medium would further smooth out each break (Panaitescu & Kumar 2000), making them appear as single gradual break. Detecting these breaks in a light curve (requiring $t_{j2}/t_j \ll 1$) would allow for an estimate of the jet's eccentricity as

$$e \sim \sqrt{1 - (t_{j2}/t_j)^{3/4}}.$$
 (7)

A more detailed discussion of jet breaks in light curves, including estimates for non-axisymmetric structured jets, will be investigated in a future study.

Although our case study section focused on optical afterglow observations, the jet break is a geometrical effect, meaning our results are applicable across different wavelengths (e.g., X-ray, radio). The number of events with measured radio polarisation has recently increased (Corsi et al. 2018; Laskar et al. 2019; Urata et al. 2023). Significant advancement in X-ray polarimetry in recent years (e.g., Astrosat, IXPE, POLAR, XPoSat) suggest that future X-ray satellites may be capable of detecting and analysing polarisation signals days after GRBs. Since both X-ray and optical bands are expected to lie above the typical frequency of the forward shock emission around jet breaks, the relativistic limb-brightening effect should be pronounced in both bands (Granot et al. 1999).

We adopted a conventional scenario in which magnetic fields are amplified by plasma kinetic instabilities at the shock, resulting in a coherence length scale comparable to the plasma skin depth scale (Medvedev & Loeb 1999). Recently, large-scale turbulent magnetic fields driven by magnetohydrodynamic instabilities have also been proposed (e.g., Kuwata et al. 2023). In this scenario, the coherence length scale is comparable to the thickness of the blast wave. In the isotropic field case, both the polarisation degree and the PPA change randomly and continuously over time, resembling the behaviour described in the classical patch shell model (e.g., Gruzinov & Waxman 1999). By incorporating magnetic field anisotropy and the observer's viewing angle, this model can also account for a variety of polarisation behaviour in afterglows. Notably, in this framework, the degree of radio polarisation can exceed that in the optical band (Kuwata et al. 2024). Simultaneous polarimetric observations across multiple wavelengths will be crucial for constraining the magnetic field amplification mechanism and improving our understanding of polarisation signals in afterglows.

ACKNOWLEDGEMENTS

This research was supported by the Sciences and Technology Facilities Research Council (STFC).

DATA AVAILABILITY

The data underlying this article will be shared on reasonable request to the corresponding author.

REFERENCES

Abbott B. P., et al., 2017, ApJL, 848, L12 Agüí Fernández J. F., et al., 2024, A&A, 690, A216 Barth A. J., et al., 2003, ApJ, 584, L47 Beniamini P., Gill R., Granot J., 2022, MNRAS, 515, 555

- Birenbaum G., Gill R., Bromberg O., Beniamini P., Granot J., 2024, ApJ, 974, 308
- Burns E., et al., 2023, ApJL, 946, L31
- Corsi A., et al., 2018, ApJ, 861, L10
- Covino S., Gotz D., 2016, A&AT, 29, 205
- Covino S., et al., 2003, A&A, 404, L5
- Ghirlanda G., et al., 2019, Science, 363, 968
- Ghisellini G., Lazzati D., 1999, MNRAS, 309, L7
- Gill R., Granot J., 2020, MNRAS, 491, 5815
- Gill R., Granot J., 2024, MNRAS, 527, 12178
- Gorosabel J., et al., 2004, A&A, 422, 113
- Gottlieb O., Nakar E., Bromberg O., 2021, MNRAS, 500, 3511
- Gottlieb O., Moseley S., Ramirez-Aguilar T., Murguia-Berthier A., Liska M., Tchekhovskoy A., 2022, ApJL, 933, L2
- Granot J., Piran T., 2012, MNRAS, 421, 570
- Granot J., Piran T., Sari R., 1999, ApJ, 513, 679
- Gruzinov A., Waxman E., 1999, ApJ, 511, 852
- Huang B.-Q., Lin D.-B., Liu T., Ren J., Wang X.-G., Liu H.-B., Liang E.-W., 2019, MNRAS, 487, 3214
- Kuwata A., Toma K., Kimura S. S., Tomita S., Shimoda J., 2023, ApJ, 943, 118
- Kuwata A., Toma K., Tomita S., Shimoda J., 2024, ApJ, 977, 224
- Lamb G. P., Kobayashi S., 2017, MNRAS, 472, 4953
- Lamb G. P., et al., 2019, ApJ, 870, L15
- Lamb G. P., Nativi L., Rosswog S., Kann D. A., Levan A., Lundman C., Tanvir N., 2022, Universe, 8, 612
- Laskar T., et al., 2019, ApJL, 878, L26
- Lazar A., Nakar E., Piran T., 2009, ApJ, 695, L10
- Lazzati D., Begelman M. C., 2005, ApJ, 629, 903
- Lazzati D., Perna R., Morsony B. J., Lopez-Camara D., Cantiello M., Ciolfi R., Giacomazzo B., Workman J. C., 2018, PRL, 120, 241103
- Lesage S., et al., 2023, ApJL, 952, L42
- Li H., Fenimore E. E., 1996, ApJ, 469, L115
- Li J.-D., Gao H., Ai S., Lei W.-H., 2023, MNRAS, 525, 6285
- Li J.-D., Gao H., Ai S., Lei W.-H., 2024, ApJ, 978, 124
- Margutti R., et al., 2018, ApJ, 856, L18
- Medvedev M. V., Loeb A., 1999, ApJ, 526, 697
- Mundell C. G., et al., 2013, Nature, 504, 119
- Nativi L., Lamb G. P., Rosswog S., Lundman C., Kowal G., 2022, MNRAS, 509, 903
- O'Connor B., et al., 2023, SciA, 9, eadi1405
- Panaitescu A., Kumar P., 2000, ApJ, 543, 66
- Perego A., Rosswog S., Cabezón R. M., Korobkin O., Käppeli R., Arcones A., Liebendörfer M., 2014, MNRAS, 443, 3134
- Racusin J. L., et al., 2009, ApJ, 698, 43
- Rossi E. M., Lazzati D., Salmonson J. D., Ghisellini G., 2004, MNRAS, 354, 86
- Sari R., 1999, ApJ, 524, L43
- Sari R., Piran T., Narayan R., 1998, ApJ, 497, L17
- Troja E., et al., 2019, MNRAS, 489, 1919
- Urata Y., et al., 2023, Nat. Astron., 7, 80
- Wiersema K., et al., 2012, MNRAS, 426, 2
- Wiersema K., et al., 2014, Nature, 509, 201
- Zhang W., Woosley S. E., MacFadyen A. I., 2003, ApJ, 586, 356

APPENDIX A: PEAK FINDING ALGORITHM

To identify peaks and their locations in numerical polarisation curves, we use a peak-finding algorithm similar to the one discussion in Li & Fenimore 1996. Originally developed for identifying peaks in prompt gamma-ray light curves, this method is well-suited for our analysis. The detailed procedure is as follows: (1) A numerical polarisation curve consists of discrete data points, representing the polarisation degree [%] as a function of time. A data point is considered a candidate peak if its value is higher than that of its neighbouring points on both sides. The candidate peak has a polarisation degree P_p [%] at

time t_p . (2) For each candidate peak, we search both sides to identify data points with polarisation degree P_1 at $t = t_1 < t_p$ and P_2 at $t = t_2 > t_p$ that satisfy the conditions:

$$P_p - P_i \ge N_v \sqrt{P_p}$$
 (A1)

where i = 1,2 and $N_v = 5$. (3) The search stops when either: both P_1 and P_2 are found, confirming P_p as a true peak, or a polarisation degree higher than P_p is encountered before finding a data point satisfying Eq. (A1) on either side of t_p , in which case P_p is discarded as a false peak.

After this step, all the peaks (one or two peaks in our case) should have been identified. If two peaks are found, the minimum point between them is designated as a trough. This method effectively identifies peaks in polarisation curves while mitigating the effects of small fluctuations due to numerical errors, including the segmentation of the jet head.

This paper has been typeset from a TFX/IATFX file prepared by the author.