# arXiv:2503.07713v1 [hep-ph] 10 Mar 2025

# $\nu$ limits from Super-Kamiokande on dark matter-electron scattering in the Sun

Dhashin Krishna <sup>()</sup>,<sup>1,\*</sup> Rinchen Sherpa <sup>()</sup>,<sup>1,†</sup> Akash Kumar Saha <sup>()</sup>

,<sup>1,‡</sup> Tarak Nath Maity <sup>(b)</sup>,<sup>2,§</sup> Ranjan Laha <sup>(b)</sup>,<sup>1,¶</sup> and Nirmal Raj <sup>(b)</sup>, \*\*

<sup>1</sup>Centre for High Energy Physics, Indian Institute of Science, C. V. Raman Avenue, Bengaluru 560012, India

<sup>2</sup>School of Physics, The University of Sydney, ARC Centre of Excellence for Dark Matter Particle Physics, NSW 2006, Australia

(Dated: March 12, 2025)

Particle dark matter scattering on electrons in the Sun may gravitationally capture and selfannihilate inside it to neutrinos and anti-neutrinos, or other final states that in turn decay to them. Using up-to-date measurements by Super-Kamiokande of the fluxes of atmospheric electron-type and muon-type neutrinos, we set the most stringent limits on the electron scattering cross sections of dark matter down to about  $10^{-40} - 10^{-39}$  cm<sup>2</sup> over a mass range of 4–200 GeV. These outdo direct searches for dark matter-electron scattering and previously set limits at IceCube. We also derive corresponding reaches at Hyper-K, and show that atmospheric neutrino observations restricted to the direction of the Sun can improve sensitivities.

## I. INTRODUCTION

While evidence for dark matter (DM) abounds in astrophysical and cosmological settings, detection of its non-gravitational interactions would mark an epoch in the hunt for its identity [1]. One strategy is to look for the signals of particle DM scattering on Standard Model (SM) particles in terrestrial detector targets [2–5]. In particular, searches for DM scattering on electrons have resulted in stringent null results [6–21].

Another attractive possibility for probing DM-SM scattering interactions, garnering much attention in recent literature, is the use of celestial objects as largevolume detectors. This includes main-sequence stars counting the Sun [22–35], planets [36–50], and Fermidegenerate compact stars [51-75]. One well-studied means by which these bodies probe DM interactions is through their gravitational capture of ambient DM following energy loss by scattering on their constituent particles. The captured DM can either heat the body, form a destructive black hole, or self-annihilate to SM states that escape the celestial object to be detected at terrestrial experiments. In this work, we study the capture of DM in the Sun via scattering on electrons, followed by annihilations to final states that result in a neutrino flux detectable at the Super-Kamiokande experiment as an excess over atmospheric neutrino events. Using this we set the most stringent limits to date on this scenario. Previous limits on DM-solar electron scattering using neutrinos at Super-K and IceCube may be found in Refs. [6, 28].

Our main result is summarized in Fig. 1, which we will describe in detail in the rest of the work. The key takeaway is that Super-K atmospheric  $\nu$  datasets set the

<sup>§</sup> tarak.maity.physics@gmail.com

best bounds on the DM-electron scattering cross section over a wide range of DM masses. In this way our study makes significant headway in probing models of so-called leptophilic DM [6, 36, 56, 57, 76–127]. In the following sections we briefly review the mechanisms of DM solar capture, annihilations to  $\nu$  fluxes, and Super-K measurements, before discussing our results and the scope of our study. Technical details are collected in the Supplemental Material.

### II. SET-UP

# A. Dark matter capture and annihilation in the Sun

For a DM particle of any spin, mass  $m_{\chi}$ , and cross section for elastic scattering on electrons  $\sigma_{\chi e}$ , the rate of capture in the Sun is given by [107]

$$C_{\odot} = 4\pi \int_{0}^{R_{\odot}} dr \ r^{2} \int_{0}^{\infty} du_{\chi} \left(\frac{\rho_{\chi}}{m_{\chi}}\right) \frac{f_{v_{\odot}}(u_{\chi})}{u_{\chi}} w(r)$$
$$\times \int_{0}^{v_{\rm esc}(r)} dv R_{e}^{-}(w \to v), \qquad (1)$$

where  $R_{\odot}$  is the radius of the Sun,  $\rho_{\chi} = 0.3 \text{ GeV/cm}^3$  is the solar neighborhood DM density, and  $f_{v_{\odot}}(u_{\chi})$  is the DM speed distribution in the solar frame. We denote the DM speed at infinity as  $u_{\chi}$ , so that at a distance r from the Sun the speed  $w(r) = \sqrt{u_{\chi}^2 + v_{\rm esc}^2(r)}$  for solar escape speed  $v_{\rm esc}$ . We take from Ref. [107] the differential scattering rate  $R_e^-$  that encodes the probability that a DM particle with speed w scatters to reach a final speed v < w, with the AGSS09 model [131] used for solar electron number density and temperature profiles. Note that DM couplings to electrons would generically give rise to DM-nucleon scattering at the loop level, which may be a sub-dominant effect [6, 117]. We do not consider the model-dependent effect of nucleon scattering in this work and leave to future investigation the exploration of its possible interplay with electron scattering.

<sup>\*</sup> mdhashinkrishna@gmail.com

 $<sup>\</sup>dagger$  rinchens@iisc.ac.in

 $<sup>^\</sup>ddagger$ akashks@iisc.ac.in

<sup>¶</sup> ranjanlaha@iisc.ac.in

<sup>\*\*</sup> nraj@iisc.ac.in



FIG. 1. Limits on DM-electron scattering cross section as a function of DM mass, assuming DM annihilations in the Sun to  $\nu\bar{\nu}$ on the left, and to  $\tau^+\tau^-$  (in turn decaying to  $\nu_{e,\mu,\tau}$ ) on the **right**. Shown here are our bounds from Super-K observations of atmospheric  $\nu_e$  and  $\nu_{\mu}$  fluxes [128] and corresponding future sensitivities at Hyper-K. Also shown in the  $\tau^+\tau^-$  panel are our limits obtained from upper limits on the total integrated flux of  $\nu_{\mu}$  from DM annihilations in the Sun in Ref. [129]. For comparison are shown limits from DeepCore-2021, DeepCore-2016, and IceCube-2016 observations of neutrinos from the solar direction [28], an older Super-K atmospheric neutrino dataset [6], and XENON1T direct searches for DM-electron scattering [9]. The region below the dotted curve denotes the "neutrino mist" where solar atmospheric neutrinos become a major background [130]. Note that the dynamical ranges of the y-axis are different across the panels. See Sec. III for more details.

If the captured DM self-annihilates, the DM population in the solar interior may attain a steady state within the age of the Sun due to equilibrium between capture and annihilation rates. This is insensitive to the annihilation cross sections above certain  $m_{\chi}$ -dependent values [28], and is easily achieved near thermal freeze-out values as allowed by indirect searches. We therefore assume this capture-annihilation equilibrium. Then ignoring the effects of evaporation of the captured DM, as is valid for  $m_{\chi} \gtrsim 4$  GeV [132], the annihilation rate of DM in the Sun is simply [6, 133]

$$\Gamma_{\odot}^{\rm ann} = \frac{C_{\odot}}{2} \ . \tag{2}$$

In this work we consider the leptonic channels  $\chi \bar{\chi} \to \nu \bar{\nu}$ , where by  $\nu \bar{\nu}$  we denote the average over all 3 flavors, and  $\chi \bar{\chi} \to \tau^+ \tau^-$ , which would further decay to produce neutrino final states. The resultant energy flux of neutrinos on the Earth's surface is then

$$E_{\nu}^{2} \frac{d\phi_{\nu}}{dE_{\nu}} = \frac{\Gamma_{\odot}^{\text{ann}}}{4 \pi D_{\odot}^{2}} \times E_{\nu}^{2} \frac{dN_{\nu}}{dE_{\nu}},\tag{3}$$

where  $D_{\odot}$  is the Earth-Sun distance and  $dN_{\nu}/dE_{\nu}$  is the spectrum of the final state neutrinos. Here we assume neutrinos of energies  $\leq 500$  GeV as otherwise their scattering cross sections would be so high as to have them significantly attenuated by the solar medium. As the neutrino energy is bounded from above by the DM mass, we consider only the conservative range of  $m_{\chi} \leq 200$  GeV.

For  $\chi \bar{\chi} \to \nu \bar{\nu}$ , the neutrino spectrum is a line feature at  $m_{\gamma}$ , which will be broadened at Super-K due to its finite energy resolution. To account for this we Gaussian-smear the neutrino flux with a 10% resolution as appropriate for our energies of interest [134–136]. In the  $\chi \bar{\chi} \to \tau^+ \tau^$ channel, the  $\tau^{\pm}$  decay daughters  $\mu^{\pm}$ ,  $K^+$  and  $\pi^+$  may come to rest before decaying, with the mesons giving rise to monochromatic neutrino signals. We have used the package  $\chi aro \nu$  [137] that takes into account these interactions. We have checked that the  $\tau^+\tau^-$  channel produces neutrino spectra with both mono-energetic "spikes" from stopped mesons (resulting in  $\pi$  decay-atrest and K decay-at-rest peaks, seen for only  $\nu_{\mu}$  spectra), and broad "shoulders" from the prompt primary decay of mesons (seen for all  $\nu$  flavors). We compute the effects of neutrino propagation (including oscillations) in the Sun and vacuum using nuSQuIDS [138], and in the atmosphere and rock of the Earth using nuCraft [139], with which we estimate the annual averaged neutrino oscillation probability.

### B. Super-K signals

In Fig. 2 we show an illustration of the flux of  $\nu_e$  and  $\nu_{\mu}$  obtained from DM annihilations in the Sun that may be detected by Super-K for  $\{m_{\chi}, \sigma_{\chi e}\}$  of  $\{20 \text{ GeV}, 2 \times 10^{-39} \text{ cm}^2\}$  and  $\{70 \text{ GeV}, 8 \times 10^{-40} \text{ cm}^2\}$ . For comparison are also shown Super-K data from Runs I–IV [128]



FIG. 2. Atmospheric neutrino flux measured at Super-K, binned in energy and fit by the HKKM11 model, and the neutrino fluxes from  $\chi \bar{\chi} \to \tau^+ \tau^-$  in the Sun for the benchmark parameters indicated. The DM signal combined with the atmospheric flux is seen to well exceed the measured flux near the energies where the signal peaks for  $\nu_e$  data but only slightly for  $\nu_{\mu}$  data. Thus these benchmarks are only ruled out by Super-K  $\nu_e$  flux measurements as seen in Fig. 1. See text for more details.

on the atmospheric  $\nu$  flux and the HKKM11 model that fits it [140]; this flux constitutes our main background. In obtaining this plot, we have imposed an angular cut on the atmospheric  $\nu$  flux, multiplying the per-steradian flux in Ref.[128] by

$$2\pi \int_0^{\delta\theta_{\rm tot}(E_{\nu})} d\theta \,\sin\theta = 4\pi \sin^2\left(\frac{\delta\theta_{\rm tot}(E_{\nu})}{2}\right) \,, \quad (4)$$

where  $\delta\theta_{\rm tot}(E_{\nu})$  is the total angular resolution as a function of incident neutrino energy; see the Supplemental Material. This cut ensures that we only analyze the flux originating from the solar direction and discard the rest of the diffuse atmospheric  $\nu$  background. For our benchmark parameters the combined fluxes of atmospheric and DM-sourced neutrinos are visibly in excess of Super-K  $\nu_e$  measurements in Fig. 2, and may therefore be constrained. We note that these benchmarks are safe from XENON1T constraints as seen in Fig. 1.

### III. RESULTS

In Fig. 1, we show our Super-K limits on the DMelectron scattering cross section vs DM mass. We also show the future reach of Hyper-K [141] via appropriately scaling the detector fiducial mass with respect to Super-K, so that the  $\sigma_{\chi e}$  sensitivities are  $\sqrt{187/22.5} \simeq 2.9$  times stronger. To set these limits we first compute

$$\chi^{2} = \sum_{i=1}^{N_{\rm bin}} \frac{\left(\phi_{\rm d}^{i} - \phi_{\rm atm}^{i} - \phi_{\chi}^{i}\right)^{2}}{\sigma_{\rm d}^{i}} , \qquad (5)$$

where in the  $i^{\text{th}}$  bin  $\phi_{d}^{i}$  is the measured neutrino flux,  $\sigma_{d}^{i}$ its corresponding uncertainty, and  $\phi^i_{\rm atm} \& \phi^i_{\chi}$  are respectively the HKKM11 model & DM signal average flux, and then obtain the 95% C.L. upper limits on  $(\sigma_{\chi e}, m_{\chi})$  by setting  $\chi^2 - \chi^2_{\min} = 2.71$ , where  $\chi^2_{\min}$  is the minimum value of  $\chi^2$  obtained by varying the DM parameters [142]. We use the same energy binning considered in Ref. [128]. We consider energy bins up to  $\pm 20\%$  of  $m_{\gamma}$ for the  $\nu\bar{\nu}$  channel accounting for the detector energy resolution, and up to 1.2  $m_{\chi}$  for the  $\tau^+\tau^-$  channel; in practice the resolution has negligible impact on the broad  $E_{\nu}$ spectrum. In any case, due to the  $\mathcal{O}(100\%)$  width of the energy bins (see Fig. 2) the Super-K energy resolution has a negligible effect on the limits we derive. For the  $\tau^+\tau^-$  channel we also present in our parameter space limits from a dedicated search at Super-K for a  $\nu_{\mu}$  flux from DM annihilations in the Sun [129]. Finally, we also show the "neutrino mist" region where our signal events  $\leq$  events from an irreducible flux of solar atmospheric neutrinos, degrading sensitivities. For this we have used the flux of the "naive" model derived in Ref. [130]. The region shown denotes the  $\nu_{\mu}$  mist, although we expect the  $\nu_e$  mist sensitivities to be comparable. Here we remark that values of  $\sigma_{\chi e}$  at the edge of the neutrino mist region do not satisfy the capture-equilibrium condition that resulted in Eq. (2), so we used the full solution for the annihilation rate as derived in, e.g., Ref. [107].

It may be seen that we constrain new ranges of parameters beyond previous limits. This is mainly because the large density and size of the Sun generically help in capturing DM with small scattering cross sections. For both  $\nu\bar{\nu}$  and  $\tau^+\tau^-$  channels the  $\nu_e$  data is seen to provide stronger limits on  $\sigma_{\chi e}$  than the  $\nu_{\mu}$  data, which is simply due to the smaller  $\nu_e$  flux produced in the atmosphere. Here we note that the net angular resolution in detection, dominated by the scattering angle, does not make a difference in the limits between  $\nu_e$  and  $\nu_{\mu}$  as it is comparable for both channels; see the Supplemental Material.

Our limits in both panels of Fig. 1 exhibit wiggle-like features due to the nature of the binning of Super-K data. In particular, the signal neutrino flux peaks near the DM mass (Fig. 2), and since only the flux up to the corresponding  $E_{\nu}$  drives the limit in Eq. (5), increasing  $m_{\chi}$  while the edge of the signal spectrum stays within a corresponding  $E_{\nu}$  bin weakens the constraints. However, when  $m_{\chi}$  is further increased so that the signal spectrum reaches the next energy bin, the limits strengthen.

### IV. DISCUSSION

In this work we have set stringent, world-leading limits on DM-electron scattering using the most up-to-date measurements of atmospheric neutrinos at Super-K, assuming that DM captured in the Sun annihilates to  $\nu\bar{\nu}$ or  $\tau^+\tau^-$ . Hyper-K will be able to extend these sensitivities by a factor of a few. Despite their smaller volume than Hyper-K, other imminent neutrino telescopes such as DUNE[143, 144] and JUNO[145, 146] would also be interesting targets to study thanks to their complementary detection channels and technologies. Other DM annihilation final states giving rise to neutrino fluxes may also be studied in these contexts, and DM masses higher than considered here may also be constrained by carefully treating the effects of attenuation of neutrinos in the Sun that become more important at high neutrino energies. While we had applied a simple angular cut in Eq. (4)to select neutrinos from the solar direction, more robust limits may be obtained by using the angular dependence of atmospheric neutrinos or, as done in Ref. [129], by applying the angular cut for all Super-K event categories.

Overall, we anticipate improved reaches with larger datasets and better understanding of detector systematics.

### Note added.

While this work was under progress, Ref. [147] appeared on arXiv investigating a scenario similar to ours. The limits obtained in that work on  $\sigma_{\chi e}$  using  $\nu_{\mu}$  detection at Super-K are seen to be stronger by up to an order of magnitude than ours near  $m_{\chi} = 10$  GeV, and are comparable near  $m_{\chi} = 100$  GeV. We believe one source of

this discrepancy is the DM solar capture rate obtained in Ref. [147]. While Ref. [147] uses expressions for the capture rate derived in Ref. [107], the capture rates plotted in Fig. S3 exceed those plotted in Fig. 1 of Ref. [107] by a factor of about 7 for the relevant DM mass range. Our capture rates match those in Ref. [107]. Further, we also differ in our estimates of the angular cut used to reduce the atmospheric neutrino background. Whereas we use Eq. (4), Ref. [147] mentions following Ref. [44], which has used the square of the net angular resolution that underestimates the backgrounds by roughly a factor of 3. The cut in Ref. [147] is stronger than ours also due to a difference in treatment of charged lepton scattering angles at detector events. Reference [147] takes the mean scattering angle as a function of neutrino energy from Refs. [148, 149], which only includes quasielastic neutrino-nucleus scattering. We determine the mean scattering angles by also including resonant and deep inelastic scattering as relevant for high neutrino energies that are of interest here. Moreover, for setting limits Ref. [147] integrates over the flux obtained from the atmospheric model HKKM11 to estimate the total number of background events, and uses Poisson statistics to derive a constraint on the number of DM-induced signal events. On the other hand, ours is a binned analysis that includes the data points and uncertainties in the Super-K measurements, which are dominated by systematics (see Fig. 6 of Ref. [128]). Finally, we note that we set limits using both the  $\nu_e$  and  $\nu_{\mu}$  fluxes measured at Super-K, with  $\nu_e$  giving better limits, whereas Ref. [147] uses only the  $\nu_{\mu}$  flux.

### ACKNOWLEDGMENTS

We thank Bhavesh Chauhan, Marco Cirelli, Raghuveer Garani, Srubabati Goswami, Joachim Kopp, and Thomas Schwetz for useful discussions. We also thank Marta Czurylo, Biprajit Mondal, Danilo Piparo, and Jonas Rembser for help with ROOT. R.S. acknowledges the University Grants Commission (UGC), Government of India, for financial support via the UGC-NET Senior Research Fellowship. A.K.S. acknowledges the Ministry of Human Resource Development, Government of India, for financial support via the Prime Ministers' Research Fellowship (PMRF). The work of T.N.M. is supported by the Australian Research Council through the ARC Centre of Excellence for Dark Matter Particle Physics. R.L. acknowledges financial support from the institute start-up funds and ISRO-IISc STC for the grant no. ISTC/PHY/RL/499.

The first three authors contributed equally to this work.

- Marco Cirelli, Alessandro Strumia, and Jure Zupan, "Dark Matter," (2024), arXiv:2406.01705 [hep-ph].
- [2] Mark W. Goodman and Edward Witten, "Detectability

of Certain Dark Matter Candidates," Phys. Rev. D **31**, 3059 (1985).

- [3] Eugenio Del Nobile, "The Theory of Direct Dark Matter Detection: A Guide to Computations," (2021), 10.1007/978-3-030-95228-0, arXiv:2104.12785 [hep-ph].
- [4] D. S. Akerib *et al.*, "Snowmass2021 Cosmic Frontier Dark Matter Direct Detection to the Neutrino Fog," in *Snowmass 2021* (2022) arXiv:2203.08084 [hep-ex].
- [5] Jodi Cooley, "Dark Matter direct detection of classical WIMPs," SciPost Phys. Lect. Notes 55, 1 (2022), arXiv:2110.02359 [hep-ph].
- [6] Joachim Kopp, Viviana Niro, Thomas Schwetz, and Jure Zupan, "DAMA/LIBRA and leptonically interacting Dark Matter," Phys. Rev. D 80, 083502 (2009), arXiv:0907.3159 [hep-ph].
- [7] Rouven Essig, Jeremy Mardon, and Tomer Volansky, "Direct Detection of Sub-GeV Dark Matter," Phys. Rev. D 85, 076007 (2012), arXiv:1108.5383 [hep-ph].
- [8] Rouven Essig, Marivi Fernandez-Serra, Jeremy Mardon, Adrian Soto, Tomer Volansky, and Tien-Tien Yu, "Direct Detection of sub-GeV Dark Matter with Semiconductor Targets," JHEP 05, 046 (2016), arXiv:1509.01598 [hep-ph].
- [9] E. Aprile *et al.* (XENON), "Light Dark Matter Search with Ionization Signals in XENON1T," Phys. Rev. Lett. 123, 251801 (2019), arXiv:1907.11485 [hep-ex].
- [10] R. Agnese *et al.* (SuperCDMS), "First Dark Matter Constraints from a SuperCDMS Single-Charge Sensitive Detector," Phys. Rev. Lett. **121**, 051301 (2018), [Erratum: Phys.Rev.Lett. **122**, 069901 (2019)], arXiv:1804.10697 [hep-ex].
- [11] P. Agnes *et al.* (DarkSide), "Constraints on Sub-GeV Dark-Matter–Electron Scattering from the DarkSide-50 Experiment," Phys. Rev. Lett. **121**, 111303 (2018), arXiv:1802.06998 [astro-ph.CO].
- [12] P. Agnes *et al.* (DarkSide-50), "Search for dark matter particle interactions with electron final states with DarkSide-50," (2022), arXiv:2207.11968 [hep-ex].
- [13] A. Aguilar-Arevalo *et al.* (DAMIC), "Constraints on Light Dark Matter Particles Interacting with Electrons from DAMIC at SNOLAB," Phys. Rev. Lett. **123**, 181802 (2019), arXiv:1907.12628 [astro-ph.CO].
- [14] Q. Arnaud *et al.* (EDELWEISS), "First germaniumbased constraints on sub-MeV Dark Matter with the EDELWEISS experiment," Phys. Rev. Lett. **125**, 141301 (2020), arXiv:2003.01046 [astro-ph.GA].
- [15] Michael Crisler, Rouven Essig, Juan Estrada, Guillermo Fernandez, Javier Tiffenberg, Miguel Sofo haro, Tomer Volansky, and Tien-Tien Yu (SENSEI), "SENSEI: First Direct-Detection Constraints on sub-GeV Dark Matter from a Surface Run," Phys. Rev. Lett. **121**, 061803 (2018), arXiv:1804.00088 [hep-ex].
- [16] Liron Barak et al. (SENSEI), "SENSEI: Direct-Detection Results on sub-GeV Dark Matter from a New Skipper-CCD," Phys. Rev. Lett. 125, 171802 (2020), arXiv:2004.11378 [astro-ph.CO].
- [17] Shuaijie Li et al. (PandaX), "Search for Light Dark Matter with Ionization Signals in the PandaX-4T Experiment," Phys. Rev. Lett. 130, 261001 (2023), arXiv:2212.10067 [hep-ex].
- [18] I. Arnquist *et al.* (DAMIC-M), "First Constraints from DAMIC-M on Sub-GeV Dark-Matter Particles Interacting with Electrons," Phys. Rev. Lett. **130**, 171003 (2023), arXiv:2302.02372 [hep-ex].

- [19] Prakruth Adari et al. (SENSEI), "First Direct-Detection Results on Sub-GeV Dark Matter Using the SENSEI Detector at SNOLAB," Phys. Rev. Lett. 134, 011804 (2025), arXiv:2312.13342 [astro-ph.CO].
- [20] Jeff A. Dror, Gilly Elor, Robert McGehee, and Tien-Tien Yu, "Absorption of sub-MeV fermionic dark matter by electron targets," Phys. Rev. D 103, 035001 (2021), [Erratum: Phys.Rev.D 105, 119903 (2022)], arXiv:2011.01940 [hep-ph].
- [21] Chen Cheng et al. (PandaX-II), "Search for Light Dark Matter-Electron Scatterings in the PandaX-II Experiment," Phys. Rev. Lett. **126**, 211803 (2021), arXiv:2101.07479 [hep-ex].
- [22] R. Abbasi *et al.* (IceCube), "Multi-year search for dark matter annihilations in the Sun with the AMANDA-II and IceCube detectors," Phys. Rev. D 85, 042002 (2012), arXiv:1112.1840 [astro-ph.HE].
- [23] Nicolás Bernal, Justo Martín-Albo, and Sergio Palomares-Ruiz, "A novel way of constraining WIMPs annihilations in the Sun: MeV neutrinos," JCAP 08, 011 (2013), arXiv:1208.0834 [hep-ph].
- [24] M. G. Aartsen *et al.* (IceCube), "Search for annihilating dark matter in the Sun with 3 years of IceCube data," Eur. Phys. J. C 77, 146 (2017), [Erratum: Eur.Phys.J.C 79, 214 (2019)], arXiv:1612.05949 [astro-ph.HE].
- [25] R. Abbasi *et al.* (IceCube), "Search for GeV-scale dark matter annihilation in the Sun with Ice-Cube DeepCore," Phys. Rev. D 105, 062004 (2022), arXiv:2111.09970 [astro-ph.HE].
- [26] Aman Gupta, Debasish Majumdar, and Ashadul Halder, "KM3NeT upper bounds of detection rates of solar neutrinos from annihilations of dark matter at the solar core," Mod. Phys. Lett. A 37, 2250233 (2022), arXiv:2203.13697 [hep-ph].
- [27] Debajit Bose, Tarak Nath Maity, and Tirtha Sankar Ray, "Solar constraints on captured electrophilic dark matter," Phys. Rev. D 105, 123013 (2022), arXiv:2112.08286 [hep-ph].
- [28] Tarak Nath Maity, Akash Kumar Saha, Sagnik Mondal, and Ranjan Laha, "Neutrinos from the Sun can discover dark matter-electron scattering," (2023), arXiv:2308.12336 [hep-ph].
- [29] Yu-Xuan Chen, Lei Zu, Zi-Qing Xia, Yue-Lin Sming Tsai, and Yi-Zhong Fan, "Multi-frequency test of dark matter annihilation into long-lived particles in Sirius," JCAP 08, 036 (2023), arXiv:2302.09951 [astro-ph.HE].
- [30] Sulagna Bhattacharya, Andrew L. Miller, and Anupam Ray, "Continuous gravitational waves: A new window to look for heavy nonannihilating dark matter," Phys. Rev. D 110, 043006 (2024), arXiv:2403.13886 [hep-ph].
- [31] Isabelle John, Rebecca K. Leane, and Tim Linden, "Dark Branches of Immortal Stars at the Galactic Center," (2024), arXiv:2405.12267 [astro-ph.HE].
- [32] Debajit Bose, Rohan Pramanick, and Tirtha Sankar Ray, "Neutrinos from captured dark matter in galactic stars," (2024), arXiv:2405.07894 [hep-ph].
- [33] Aman Gupta, Pooja Bhattacharjee, and Pratik Majumdar, "Searching for dark matter annihilating into lightlong-lived mediators from stars inside dwarf spheroidal galaxies," (2025), arXiv:2501.04663 [astro-ph.HE].
- [34] Xiaoyong Chu, Raghuveer Garani, Camilo García-Cely, and Thomas Hambye, "Dark matter boundstate formation in the Sun," JHEP 05, 045 (2024), arXiv:2402.18535 [hep-ph].

- [35] Rebecca K. Leane and Joshua Tong, "Optimal celestial bodies for dark matter detection," JCAP 12, 031 (2024), arXiv:2405.05312 [hep-ph].
- [36] Bhavesh Chauhan and Subhendra Mohanty, "Constraints on leptophilic light dark matter from internal heat flux of Earth," Phys. Rev. D 94, 035024 (2016), arXiv:1603.06350 [hep-ph].
- [37] R. Abbasi *et al.* (IceCube), "Search for dark matter from the center of the Earth with ten years of IceCube data," (2024), arXiv:2412.12972 [astro-ph.HE].
- [38] Joseph Bramante, Jason Kumar, Gopolang Mohlabeng, Nirmal Raj, and Ningqiang Song, "Light dark matter accumulating in planets: Nuclear scattering," Phys. Rev. D 108, 063022 (2023), arXiv:2210.01812 [hep-ph].
- [39] Giovanni Renzi and Juan A. Aguilar (IceCube), "Search for dark matter annihilations in the center of the Earth with IceCube," in 38th International Cosmic Ray Conference (2023) arXiv:2308.02920 [astro-ph.HE].
- [40] Raghuveer Garani and Peter Tinyakov, "Constraints on Dark Matter from the Moon," Phys. Lett. B 804, 135403 (2020), arXiv:1912.00443 [hep-ph].
- [41] Rebecca K. Leane and Tim Linden, "First Analysis of Jupiter in Gamma Rays and a New Search for Dark Matter," Phys. Rev. Lett. 131, 071001 (2023), arXiv:2104.02068 [astro-ph.HE].
- [42] George M. French and Marc Sher, "Monoenergetic neutrinos from WIMP annihilation in Jupiter," Phys. Rev. D 106, 115037 (2022), arXiv:2210.04761 [hep-ph].
- [43] Carlos Blanco and Rebecca K. Leane, "Search for Dark Matter Ionization on the Night Side of Jupiter with Cassini," Phys. Rev. Lett. 132, 261002 (2024), arXiv:2312.06758 [hep-ph].
- [44] Sandra Robles and Stephan A. Meighen-Berger, "Extending the Dark Matter Reach of Water Cherenkov Detectors using Jupiter," (2024), arXiv:2411.04435 [hepph].
- [45] Joseph Bramante, Andrew Buchanan, Alan Goodman, and Eesha Lodhi, "Terrestrial and Martian Heat Flow Limits on Dark Matter," Phys. Rev. D 101, 043001 (2020), arXiv:1909.11683 [hep-ph].
- [46] Stephen L. Adler, "Planet-bound dark matter and the internal heat of Uranus, Neptune, and hot-Jupiter exoplanets," Phys. Lett. B 671, 203–206 (2009), arXiv:0808.2823 [astro-ph].
- [47] Carlos Blanco, Rebecca K. Leane, Marianne Moore, and Joshua Tong, "Search for Dark Matter Induced Airglow in Planetary Atmospheres," (2024), arXiv:2408.15318 [hep-ph].
- [48] Rebecca K. Leane and Juri Smirnov, "Exoplanets as Sub-GeV Dark Matter Detectors," Phys. Rev. Lett. 126, 161101 (2021), arXiv:2010.00015 [hep-ph].
- [49] Pooja Bhattacharjee, Francesca Calore, and Pasquale Dario Serpico, "Gamma-ray flux limits from brown dwarfs: Implications for dark matter annihilating into long-lived mediators," Phys. Rev. D 107, 043012 (2023), [Erratum: Phys.Rev.D 109, 129904 (2024)], arXiv:2211.08067 [astro-ph.HE].
- [50] Tim Linden, Thong T. Q. Nguyen, and Tim M. P. Tait, "Indirect Searches for Dark Photon-Photon Tridents in Celestial Objects," (2024), arXiv:2402.01839 [hep-ph].
- [51] Joseph Bramante and Nirmal Raj, "Dark matter in compact stars," Phys. Rept. 1052, 1–48 (2024), arXiv:2307.14435 [hep-ph].
- [52] Masha Baryakhtar et al., "Dark Matter In Extreme As-

trophysical Environments," in *Snowmass 2021* (2022) arXiv:2203.07984 [hep-ph].

- [53] Masha Baryakhtar, Joseph Bramante, Shirley Weishi Li, Tim Linden, and Nirmal Raj, "Dark Kinetic Heating of Neutron Stars and An Infrared Window On WIMPs, SIMPs, and Pure Higgsinos," Phys. Rev. Lett. 119, 131801 (2017), arXiv:1704.01577 [hep-ph].
- [54] Nirmal Raj, Philip Tanedo, and Hai-Bo Yu, "Neutron stars at the dark matter direct detection frontier," Phys. Rev. D 97, 043006 (2018), arXiv:1707.09442 [hep-ph].
- [55] Javier F. Acevedo, Joseph Bramante, Rebecca K. Leane, and Nirmal Raj, "Warming Nuclear Pasta with Dark Matter: Kinetic and Annihilation Heating of Neutron Star Crusts," JCAP 03, 038 (2020), arXiv:1911.06334 [hep-ph].
- [56] Aniket Joglekar, Nirmal Raj, Philip Tanedo, and Hai-Bo Yu, "Relativistic capture of dark matter by electrons in neutron stars," Phys. Lett. B, 135767 (2020), arXiv:1911.13293 [hep-ph].
- [57] Aniket Joglekar, Nirmal Raj, Philip Tanedo, and Hai-Bo Yu, "Dark kinetic heating of neutron stars from contact interactions with relativistic targets," Phys. Rev. D 102, 123002 (2020), arXiv:2004.09539 [hep-ph].
- [58] Raghuveer Garani, Aritra Gupta, and Nirmal Raj, "Observing the thermalization of dark matter in neutron stars," Phys. Rev. D 103, 043019 (2021), arXiv:2009.10728 [hep-ph].
- [59] Giorgio Busoni, "Capture of Dark Matter in Neutron Stars," Moscow Univ. Phys. Bull. 77, 301–305 (2022), arXiv:2201.00048 [hep-ph].
- [60] Tarak Nath Maity and Farinaldo S. Queiroz, "Detecting bosonic dark matter with neutron stars," Phys. Rev. D 104, 083019 (2021), arXiv:2104.02700 [hep-ph].
- [61] Joseph Bramante, Bradley J. Kavanagh, and Nirmal Raj, "Scattering Searches for Dark Matter in Subhalos: Neutron Stars, Cosmic Rays, and Old Rocks," Phys. Rev. Lett. **128**, 231801 (2022), arXiv:2109.04582 [hepph].
- [62] Debajit Bose, Tarak Nath Maity, and Tirtha Sankar Ray, "Neutrinos from captured dark matter annihilation in a galactic population of neutron stars," JCAP 05, 001 (2022), arXiv:2108.12420 [hep-ph].
- [63] Nicole F. Bell, Giorgio Busoni, Sandra Robles, and Michael Virgato, "Thermalization and annihilation of dark matter in neutron stars," JCAP 04, 006 (2024), arXiv:2312.11892 [hep-ph].
- [64] Pooja Bhattacharjee and Francesca Calore, "Probing the Dark Matter Capture Rate in a Local Population of Brown Dwarfs with IceCube Gen 2 †," Particles 7, 489–501 (2024), arXiv:2311.18455 [astro-ph.HE].
- [65] Pooja Bhattacharjee and Amit Dutta Banik, "Confronting the dark matter capture rate with a continuous gravitational wave probe of local neutron stars," Phys. Rev. D 111, 043009 (2025), arXiv:2404.15038 [astroph.HE].
- [66] Peter W. Graham, Ryan Janish, Vijay Narayan, Surjeet Rajendran, and Paul Riggins, "White Dwarfs as Dark Matter Detectors," Phys. Rev. D 98, 115027 (2018), arXiv:1805.07381 [hep-ph].
- [67] Javier F. Acevedo and Joseph Bramante, "Supernovae Sparked By Dark Matter in White Dwarfs," Phys. Rev. D 100, 043020 (2019), arXiv:1904.11993 [hep-ph].
- [68] Nicole F. Bell, Giorgio Busoni, Maura E. Ramirez-Quezada, Sandra Robles, and Michael Virgato, "Im-

proved treatment of dark matter capture in white dwarfs," JCAP **10**, 083 (2021), arXiv:2104.14367 [hep-ph].

- [69] Javier F. Acevedo, Rebecca K. Leane, and Lillian Santos-Olmsted, "Milky Way white dwarfs as sub-GeV to multi-TeV dark matter detectors," JCAP 03, 042 (2024), arXiv:2309.10843 [hep-ph].
- [70] John Coffey, David McKeen, David E. Morrissey, and Nirmal Raj, "Neutron star observations of pseudoscalarmediated dark matter," (2022), arXiv:2207.02221 [hepph].
- [71] Raghuveer Garani, Nirmal Raj, and Javier Reynoso-Cordova, "Could compact stars in globular clusters constrain dark matter?" JCAP 07, 038 (2023), arXiv:2303.18009 [astro-ph.HE].
- [72] Nirmal Raj, "Supernovae and superbursts triggered by dark matter clumps," Phys. Rev. D 109, 123020 (2024), arXiv:2306.14981 [hep-ph].
- [73] Nirmal Raj, Prajwal Shivanna, and Gaurav Niraj Rachh, "Exploring reheated sub-40000 Kelvin neutron stars with JWST, ELT, and TMT," Phys. Rev. D 109, 123040 (2024), arXiv:2403.07496 [astro-ph.HE].
- [74] Ujjwal Basumatary, Nirmal Raj, and Anupam Ray, "Beyond Hawking evaporation of black holes formed by dark matter in compact stars," Phys. Rev. D 111, L041306 (2025), arXiv:2410.22702 [hep-ph].
- [75] Joseph Bramante, Katherine Mack, Nirmal Raj, Lijing Shao, and Narayani Tyagi, "Seeking the nearest neutron stars using a new local electron density map," (2024), arXiv:2411.18647 [astro-ph.HE].
- [76] Lawrence M. Krauss, Salah Nasri, and Mark Trodden,
  "A Model for neutrino masses and dark matter," Phys. Rev. D 67, 085002 (2003), arXiv:hep-ph/0210389.
- [77] E. A. Baltz and L. Bergstrom, "Detection of leptonic dark matter," Phys. Rev. D 67, 043516 (2003), arXiv:hep-ph/0211325.
- [78] Ernest Ma, "Verifiable radiative seesaw mechanism of neutrino mass and dark matter," Phys. Rev. D 73, 077301 (2006), arXiv:hep-ph/0601225.
- [79] Thomas Hambye, Kristjan Kannike, Ernest Ma, and Martti Raidal, "Emanations of Dark Matter: Muon Anomalous Magnetic Moment, Radiative Neutrino Mass, and Novel Leptogenesis at the TeV Scale," Phys. Rev. D 75, 095003 (2007), arXiv:hep-ph/0609228.
- [80] R. Bernabei et al., "Investigating electron interacting dark matter," Phys. Rev. D 77, 023506 (2008), arXiv:0712.0562 [astro-ph].
- [81] Marco Cirelli, Mario Kadastik, Martti Raidal, and Alessandro Strumia, "Model-independent implications of the e+-, anti-proton cosmic ray spectra on properties of Dark Matter," Nucl. Phys. B 813, 1–21 (2009), [Addendum: Nucl.Phys.B 873, 530–533 (2013)], arXiv:0809.2409 [hep-ph].
- [82] Chuan-Ren Chen and Fuminobu Takahashi, "Cosmic rays from Leptonic Dark Matter," JCAP 02, 004 (2009), arXiv:0810.4110 [hep-ph].
- [83] Xiao-Jun Bi, Pei-Hong Gu, Tianjun Li, and Xinmin Zhang, "ATIC and PAMELA Results on Cosmic e+-Excesses and Neutrino Masses," JHEP 04, 103 (2009), arXiv:0901.0176 [hep-ph].
- [84] Qing-Hong Cao, Ernest Ma, and Gabe Shaughnessy, "Dark Matter: The Leptonic Connection," Phys. Lett. B 673, 152–155 (2009), arXiv:0901.1334 [hep-ph].
- [85] Hock-Seng Goh, Lawrence J. Hall, and Piyush Kumar,

"The Leptonic Higgs as a Messenger of Dark Matter," JHEP **05**, 097 (2009), arXiv:0902.0814 [hep-ph].

- [86] Xiao-Jun Bi, Xiao-Gang He, and Qiang Yuan, "Parameters in a class of leptophilic models from PAMELA, ATIC and FERMI," Phys. Lett. B 678, 168–173 (2009), arXiv:0903.0122 [hep-ph].
- [87] Alejandro Ibarra, Andreas Ringwald, David Tran, and Christoph Weniger, "Cosmic Rays from Leptophilic Dark Matter Decay via Kinetic Mixing," JCAP 08, 017 (2009), arXiv:0903.3625 [hep-ph].
- [88] Hooman Davoudiasl, "Dark Matter with Time-Varying Leptophilic Couplings," Phys. Rev. D 80, 043502 (2009), arXiv:0904.3103 [hep-ph].
- [89] A. Dedes, I. Giomataris, K. Suxho, and J. D. Vergados, "Searching for Secluded Dark Matter via Direct Detection of Recoiling Nuclei as well as Low Energy Electrons," Nucl. Phys. B 826, 148–173 (2010), arXiv:0907.0758 [hep-ph].
- [90] Timothy Cohen and Kathryn M. Zurek, "Leptophilic Dark Matter from the Lepton Asymmetry," Phys. Rev. Lett. 104, 101301 (2010), arXiv:0909.2035 [hep-ph].
- [91] Eung Jin Chun, Jong-Chul Park, and Stefano Scopel, "Dirac gaugino as leptophilic dark matter," JCAP 02, 015 (2010), arXiv:0911.5273 [hep-ph].
- [92] Wei Chao, "Pure Leptonic Gauge Symmetry, Neutrino Masses and Dark Matter," Phys. Lett. B 695, 157–161 (2011), arXiv:1005.1024 [hep-ph].
- [93] Naoyuki Haba, Yuji Kajiyama, Shigeki Matsumoto, Hiroshi Okada, and Koichi Yoshioka, "Universally Leptophilic Dark Matter From Non-Abelian Discrete Symmetry," Phys. Lett. B 695, 476–481 (2011), arXiv:1008.4777 [hep-ph].
- [94] Pyungwon Ko and Yuji Omura, "Supersymmetric U(1)B x U(1)L model with leptophilic and leptophobic cold dark matters," Phys. Lett. B 701, 363–366 (2011), arXiv:1012.4679 [hep-ph].
- [95] Christopher D. Carone and Reinard Primulando, "A Froggatt-Nielsen Model for Leptophilic Scalar Dark Matter Decay," Phys. Rev. D 84, 035002 (2011), arXiv:1105.4635 [hep-ph].
- [96] Daniel Schmidt, Thomas Schwetz, and Takashi Toma, "Direct Detection of Leptophilic Dark Matter in a Model with Radiative Neutrino Masses," Phys. Rev. D 85, 073009 (2012), arXiv:1201.0906 [hep-ph].
- [97] Moumita Das and Subhendra Mohanty, "Leptophilic dark matter in gauged  $L_{\mu} L_{\tau}$  extension of MSSM," Phys. Rev. D 89, 025004 (2014), arXiv:1306.4505 [hep-ph].
- [98] P. S. Bhupal Dev, Dilip Kumar Ghosh, Nobuchika Okada, and Ipsita Saha, "Neutrino Mass and Dark Matter in light of recent AMS-02 results," Phys. Rev. D 89, 095001 (2014), arXiv:1307.6204 [hep-ph].
- [99] Joachim Kopp, Lisa Michaels, and Juri Smirnov, "Loopy Constraints on Leptophilic Dark Matter and Internal Bremsstrahlung," JCAP 04, 022 (2014), arXiv:1401.6457 [hep-ph].
- [100] Spencer Chang, Ralph Edezhath, Jeffrey Hutchinson, and Markus Luty, "Leptophilic Effective WIMPs," Phys. Rev. D 90, 015011 (2014), arXiv:1402.7358 [hepph].
- [101] Prateek Agrawal, Zackaria Chacko, and Christopher B. Verhaaren, "Leptophilic Dark Matter and the Anomalous Magnetic Moment of the Muon," JHEP 08, 147 (2014), arXiv:1402.7369 [hep-ph].

- [102] Nicole F. Bell, Yi Cai, Rebecca K. Leane, and Anibal D. Medina, "Leptophilic dark matter with Z' interactions," Phys. Rev. D 90, 035027 (2014), arXiv:1407.3001 [hepph].
- [103] Ayres Freitas and Susanne Westhoff, "Leptophilic Dark Matter in Lepton Interactions at LEP and ILC," JHEP 10, 116 (2014), arXiv:1408.1959 [hep-ph].
- [104] Qing-Hong Cao, Chuan-Ren Chen, and Ti Gong, "Leptophilic dark matter confronts AMS-02 cosmicray positron flux," Chin. J. Phys. 55, 10–15 (2017), arXiv:1409.7317 [hep-ph].
- [105] Sofiane M. Boucenna, Marco Chianese, Gianpiero Mangano, Gennaro Miele, Stefano Morisi, Ofelia Pisanti, and Edoardo Vitagliano, "Decaying Leptophilic Dark Matter at IceCube," JCAP 12, 055 (2015), arXiv:1507.01000 [hep-ph].
- [106] Bo-Qiang Lu and Hong-Shi Zong, "Leptophilic dark matter in Galactic Center excess," Phys. Rev. D 93, 083504 (2016), [Addendum: Phys.Rev.D 93, 089910 (2016)].
- [107] Raghuveer Garani and Sergio Palomares-Ruiz, "Dark matter in the Sun: scattering off electrons vs nucleons," JCAP 05, 007 (2017), arXiv:1702.02768 [hep-ph].
- [108] Guang Hua Duan, Lei Feng, Fei Wang, Lei Wu, Jin Min Yang, and Rui Zheng, "Simplified TeV leptophilic dark matter in light of DAMPE data," JHEP 02, 107 (2018), arXiv:1711.11012 [hep-ph].
- [109] Guang Hua Duan, Xiao-Gang He, Lei Wu, and Jin Min Yang, "Leptophilic dark matter in gauged  $U(1)_{L_e-L_{\mu}}$ model in light of DAMPE cosmic ray  $e^+ + e^-$  excess," Eur. Phys. J. C 78, 323 (2018), arXiv:1711.11563 [hepph].
- [110] Wei Chao, Huai-Ke Guo, Hao-Lin Li, and Jing Shu, "Electron Flavored Dark Matter," Phys. Lett. B 782, 517–522 (2018), arXiv:1712.00037 [hep-ph].
- [111] Tong Li, Nobuchika Okada, and Qaisar Shafi, "Scalar dark matter, Type II Seesaw and the DAMPE cosmic ray  $e^+ + e^-$  excess," Phys. Lett. B **779**, 130–135 (2018), arXiv:1712.00869 [hep-ph].
- [112] Karim Ghorbani and Parsa Hossein Ghorbani, "DAMPE electron-positron excess in leptophilic Z' model," JHEP 05, 125 (2018), arXiv:1712.01239 [hep-ph].
- [113] Yicong Sui and Yongchao Zhang, "Prospects of type-II seesaw models at future colliders in light of the DAMPE  $e^+e^-$  excess," Phys. Rev. D **97**, 095002 (2018), arXiv:1712.03642 [hep-ph].
- [114] Zhi-Long Han, Weijian Wang, and Ran Ding, "Radiative Seesaw Model and DAMPE Excess from Leptophilic Gauge Symmetry," Eur. Phys. J. C 78, 216 (2018), arXiv:1712.05722 [hep-ph].
- [115] Eric Madge and Pedro Schwaller, "Leptophilic dark matter from gauged lepton number: Phenomenology and gravitational wave signatures," JHEP 02, 048 (2019), arXiv:1809.09110 [hep-ph].
- [116] Sam Junius, Laura Lopez-Honorez, and Alberto Mariotti, "A feeble window on leptophilic dark matter," JHEP 07, 136 (2019), arXiv:1904.07513 [hep-ph].
- [117] Nicole F. Bell, Giorgio Busoni, and Sandra Robles, "Capture of Leptophilic Dark Matter in Neutron Stars," JCAP 06, 054 (2019), arXiv:1904.09803 [hep-ph].
- [118] Seyed Yaser Ayazi and Ahmad Mohamadnejad, "DAMPE excess from leptophilic vector dark matter:

a model-independent approach," J. Phys. G **47**, 095003 (2020), arXiv:1909.10729 [hep-ph].

- [119] Sayan Ghosh, Amit Dutta Banik, Eung Jin Chun, and Debasish Majumdar, "Leptophilic-portal dark matter in the light of AMS-02 positron excess," Phys. Rev. D 104, 075016 (2021), arXiv:2003.07675 [hep-ph].
- [120] Sreemanti Chakraborti and Rashidul Islam, "Implications of dark sector mixing on leptophilic scalar dark matter," JHEP 03, 032 (2021), arXiv:2007.13719 [hepph].
- [121] Nicole F. Bell, Giorgio Busoni, Sandra Robles, and Michael Virgato, "Improved Treatment of Dark Matter Capture in Neutron Stars II: Leptonic Targets," JCAP 03, 086 (2021), arXiv:2010.13257 [hep-ph].
- [122] Shun-Ichi Horigome, Taisuke Katayose, Shigeki Matsumoto, and Ipsita Saha, "Leptophilic fermion WIMP: Role of future lepton colliders," Phys. Rev. D 104, 055001 (2021), arXiv:2102.08645 [hep-ph].
- [123] Raghuveer Garani, Federico Gasparotto, Pierpaolo Mastrolia, Henrik J. Munch, Sergio Palomares-Ruiz, and Amedeo Primo, "Two-photon exchange in leptophilic dark matter scenarios," JHEP 12, 212 (2021), arXiv:2105.12116 [hep-ph].
- [124] Basudeb Dasgupta, Ranjan Laha, and Anupam Ray, "Low Mass Black Holes from Dark Core Collapse," Phys. Rev. Lett. **126**, 141105 (2021), arXiv:2009.01825 [astro-ph.HE].
- [125] Debajit Bose, Debtosh Chowdhury, Poulami Mondal, and Tirtha Sankar Ray, "Troubles mounting for multipolar dark matter," JHEP 06, 014 (2024), arXiv:2312.05131 [hep-ph].
- [126] Sulagna Bhattacharya, Basudeb Dasgupta, Ranjan Laha, and Anupam Ray, "Can LIGO Detect Nonannihilating Dark Matter?" Phys. Rev. Lett. 131, 091401 (2023), arXiv:2302.07898 [hep-ph].
- [127] Pouya Asadi, Samuel Homiller, Aria Radick, and Tien-Tien Yu, "Fermion-Portal Dark Matter at a High-Energy Muon Collider," (2024), arXiv:2412.14235 [hepph].
- [128] E. Richard *et al.* (Super-Kamiokande), "Measurements of the atmospheric neutrino flux by Super-Kamiokande: energy spectra, geomagnetic effects, and solar modulation," Phys. Rev. D 94, 052001 (2016), arXiv:1510.08127 [hep-ex].
- [129] K. Choi *et al.* (Super-Kamiokande), "Search for neutrinos from annihilation of captured low-mass dark matter particles in the Sun by Super-Kamiokande," Phys. Rev. Lett. **114**, 141301 (2015), arXiv:1503.04858 [hep-ex].
- [130] Kenny C. Y. Ng, John F. Beacom, Annika H. G. Peter, and Carsten Rott, "Solar Atmospheric Neutrinos: A New Neutrino Floor for Dark Matter Searches," Phys. Rev. D 96, 103006 (2017), arXiv:1703.10280 [astroph.HE].
- [131] Núria Vinyoles, Aldo M. Serenelli, Francesco L. Villante, Sarbani Basu, Johannes Bergström, M. C. Gonzalez-Garcia, Michele Maltoni, Carlos Peña Garay, and Ningqiang Song, "A new Generation of Standard Solar Models," Astrophys. J. 835, 202 (2017), arXiv:1611.09867 [astro-ph.SR].
- [132] Raghuveer Garani and Sergio Palomares-Ruiz, "Evaporation of dark matter from celestial bodies," JCAP 05, 042 (2022), arXiv:2104.12757 [hep-ph].
- [133] Annika H. G. Peter, "Dark matter in the solar system II: WIMP annihilation rates in the Sun," Phys. Rev. D

79, 103532 (2009), arXiv:0902.1347 [astro-ph.HE].

- [134] M. Shiozawa (Super-Kamiokande), "Reconstruction algorithms in the Super-Kamiokande large water Cherenkov detector," Nucl. Instrum. Meth. A 433, 240– 246 (1999).
- [135] E. Drakopoulou, G. A. Cowan, M. D. Needham, S. Playfer, and M. Taani, "Application of machine learning techniques to lepton energy reconstruction in water Cherenkov detectors," JINST 13, P04009 (2018), arXiv:1710.05668 [physics.ins-det].
- [136] M. Jiang *et al.* (Super-Kamiokande), "Atmospheric Neutrino Oscillation Analysis with Improved Event Reconstruction in Super-Kamiokande IV," PTEP **2019**, 053F01 (2019), arXiv:1901.03230 [hep-ex].
- [137] Qinrui Liu, Jeffrey Lazar, Carlos A. Argüelles, and Ali Kheirandish, " $\chi$ aro $\nu$ : a tool for neutrino flux generation from WIMPs," JCAP **10**, 043 (2020), arXiv:2007.15010 [hep-ph].
- [138] Carlos A. Argüelles, Jordi Salvado, and Christopher N. Weaver, "nuSQuIDS: A toolbox for neutrino propagation," Comput. Phys. Commun. 277, 108346 (2022), arXiv:2112.13804 [hep-ph].
- [139] Marius Wallraff and Christopher Wiebusch, "Calculation of oscillation probabilities of atmospheric neutrinos using nuCraft," Comput. Phys. Commun. 197, 185–189 (2015), arXiv:1409.1387 [astro-ph.IM].
- [140] M. Honda, T. Kajita, K. Kasahara, and S. Midorikawa, "Improvement of low energy atmospheric neutrino flux calculation using the JAM nuclear interaction model," Phys. Rev. D 83, 123001 (2011), arXiv:1102.2688 [astroph.HE].
- [141] K. Abe et al. (Hyper-Kamiokande), "Hyper-Kamiokande Design Report," (2018), arXiv:1805.04163 [physics.ins-det].
- [142] Anna Sophia Lamperstorfer, Spectral Features from Dark Matter Annihilations and Decays in Indirect Searches, Ph.D. thesis, Munich, Tech. U. (2015).
- [143] Carsten Rott, DongYoung Jeong, Jason Kumar, and David Yaylali, "Neutrino Topology Reconstruction at DUNE and Applications to Searches for Dark Matter Annihilation in the Sun," JCAP 07, 006 (2019), arXiv:1903.04175 [astro-ph.HE].
- [144] Bhavesh Chauhan, Mary Hall Reno, Carsten Rott, and Ina Sarcevic, "Neutrino constraints on inelastic dark matter captured in the Sun," JCAP 01, 030 (2024), arXiv:2308.16134 [hep-ph].
- [145] Wan-Lei Guo, "Detecting electron neutrinos from solar dark matter annihilation by JUNO," JCAP 01, 039 (2016), arXiv:1511.04888 [hep-ph].
- [146] Angel Abusleme *et al.* (JUNO), "JUNO sensitivity to the annihilation of MeV dark matter in the galactic halo," JCAP **09**, 001 (2023), arXiv:2306.09567 [hep-ex].
- [147] Thong T. Q. Nguyen, Tim Linden, Pierluca Carenza, and Axel Widmark, "Super-Kamiokande Strongly Constrains Leptophilic Dark Matter Capture in the Sun," (2025), arXiv:2501.14864 [astro-ph.HE].
- [148] E. Konishi, Y. Minorikawa, V. I. Galkin, M. Ishiwata, I. Nakamura, N. Takahashi, M. Kato, and A. Misaki, "On the Sensitivity of L/E Analysis of Super-Kamiokande Atmospheric Neutrino Data to Neutrino Oscillation Part~1," (2010), arXiv:1007.3812 [hep-ex].
- [149] E. Konishi, Y. Minorikawa, V. I. Galkin, M. Ishiwata, I. Nakamura, N. Takahashi, M. Kato, and A. Misaki, "On the Relation between the True Directions of Neu-

trinos and the Reconstructed Directions of Neutrinos in L/E Analysis Performed by Super-Kamiokande Collaboration Part1," (2011) arXiv:1108.1064 [hep-ex].

- [150] Tomasz Golan, Cezary Juszczak, and Jan T. Sobczyk, "Final State Interactions Effects in Neutrino-Nucleus Interactions," Phys. Rev. C 86, 015505 (2012), arXiv:1202.4197 [nucl-th].
- [151] C. Andreopoulos *et al.*, "The GENIE Neutrino Monte Carlo Generator," Nucl. Instrum. Meth. A **614**, 87–104 (2010), arXiv:0905.2517 [hep-ph].
- [152] Koun Choi, Study of capture and annihilation of a few GeV WIMPs inside the Sun by using an underground neutrino detector, Ph.D. thesis, Nagoya U. (2015).
- [153] Kaito Hagiwara, Search for Astronomical Neutrinos from Blazar TXS0506+056 in Super-Kamiokande, Ph.D. thesis, Okayama U. (2020).
- [154] S. Desai *et al.* (Super-Kamiokande), "Study of TeV neutrinos with upward showering muons in Super-Kamiokande," Astropart. Phys. **29**, 42–54 (2008), arXiv:0711.0053 [hep-ex].

# Supplemental Material



FIG. 3. Angular uncertainties from the mean scattering angle of the outgoing charged lepton (obtained with NuWro) and detector resolutions of various event types at Super-K as described in Section B of Supplemental Material. The net angular resolution is seen to be dominated by the scattering angle except at high  $\nu_{\mu}$  energies.

### A. Charged current cross sections from NuWro

We use the Monte Carlo neutrino event generator NuWro [150] to obtain neutrino-H<sub>2</sub>O charged current scattering cross sections and the average scattering angle of the final state  $e^{\pm}/\mu^{\pm}$ . In the parameter initialization file, we set SF\_method=1, and neutral current interactions to zero, and generate  $10^5$  events for each neutrino energy. We checked that the total cross sections thus obtained match with the output of GENIE [151], and note that the cross sections for  $\bar{\nu}$  are generally smaller than for  $\nu$ .

### B. Resolutions for the angular cut

To impose the angular cut in Eq. (4) for reducing the atmospheric neutrino background, we must estimate the energy-dependent total angular resolution for neutrino- $H_2O$  scattering events in Super-K, given by

$$\delta\theta_{\rm tot}(E_{\nu}) = \sqrt{\bar{\theta}_{\rm scat}^2(E_{\nu})} + \delta\theta_{\rm det}^2(E_{\nu}) , \qquad (6)$$

where  $\bar{\theta}_{\rm scat}$  is the mean scattering angle of the outgoing charged lepton and  $\delta \theta_{\rm det}$  is the detector angular resolution. We obtain  $\bar{\theta}_{\rm scat}$  by generating 10<sup>5</sup> events as described above. As the mean scattering angle for  $\nu_{\mu}$  ( $\nu_{e}$ ) is larger than that of  $\bar{\nu}_{\mu}$  ( $\bar{\nu}_{e}$ ), we conservatively use the former. We also find that the the mean scattering angle dominates  $\delta \theta_{\rm tot}$  except at the highest  $\nu_{\mu}$  energies considered. Nonetheless, for the sake of completeness we describe here the determination of  $\delta \theta_{\rm det}$ .

The detector angular resolution depends on the type of event in Super-K, broadly classified as:

(i) **Fully contained (FC) events** where the lepton is produced and contained in the inner detector (ID). FC events are further divided into 7 sub-GeV and 6 multi-GeV sub-types on the basis of the species of particle and number of Cherenkov rings produced. (ii) **Par**tially contained (PC) events in which the lepton is produced in the ID but deposits energy in the outer detector (OD). PC event particles can either pass through the OD ("OD through-going") or stop in the OD ("OD stopping"). (iii) **Upward-going-muons** (**Up**- $\mu$ ) are selected to identify neutrinos that interact with the rock surrounding the detector and produce muons. This is to mitigate the background of atmospheric muons. Up- $\mu$ can again either stop in the detector ("upward stopping") or pass through it ("upward-through going"). Further, through-going Up- $\mu$  events can be either showering or non-showering [152]. We take the detector angular resolution for different event types from Refs. [153] and [154]. To obtain conservative estimates we choose the maximum  $\delta\theta_{det}$  possible at a given energy for any event type. We depict the detector angular resolutions of these event types in Fig. 3.