

ASYMPTOTIC SCATTERING RELATION FOR THE TODA LATTICE

AMOL AGGARWAL

ABSTRACT. In this paper we consider the Toda lattice $(\mathbf{p}(t); \mathbf{q}(t))$ at thermal equilibrium, meaning that its variables (p_i) and $(e^{q_i - q_{i+1}})$ are independent Gaussian and Gamma random variables, respectively. We justify the notion from the physics literature that this model can be thought of as a dense collection of solitons (or “soliton gas”) by, (i) precisely defining the locations of these solitons; (ii) showing that local charges and currents for the Toda lattice are well-approximated by simple functions of the soliton data; and (iii) proving an asymptotic scattering relation that governs the dynamics of the soliton locations. Our arguments are based on analyzing properties about eigenvector entries of the Toda lattice’s (random) Lax matrix, particularly, their rates of exponential decay and their evolutions under inverse scattering.

CONTENTS

1. Introduction	1
2. Results	7
3. Miscellaneous Preliminaries	14
4. Comparison Estimates	19
5. Localization Centers	25
6. Eigenvector Analysis for Lax Matrices	31
7. Properties of Localization Centers	38
8. Proof of the Asymptotic Scattering Relation	45
Appendix A. Proofs of Results From Section 3	53
Appendix B. Heuristics for Eigenvalue Velocities	57
References	58

1. INTRODUCTION

The *Toda lattice* is a Hamiltonian dynamical system $(\mathbf{p}(t); \mathbf{q}(t))$, where $\mathbf{p}(t) = (p_i(t))$ and $\mathbf{q}(t) = (q_i(t))$ are indexed by a one-dimensional integer lattice $i \in \mathcal{J}$ that could either be an interval $\mathcal{J} = [N_1, N_2]$, a torus $\mathcal{J} = \mathbb{Z}/N\mathbb{Z}$, or the full line $\mathcal{J} = \mathbb{Z}$. Its Hamiltonian is given by

$$\mathfrak{H}(\mathbf{p}; \mathbf{q}) = \sum_{i \in \mathcal{J}} \left(\frac{p_i^2}{2} + e^{q_i - q_{i+1}} \right),$$

so the dynamics $\partial_t q_i = \partial_{p_i} \mathfrak{H}(\mathbf{p}; \mathbf{q})$ and $\partial_t p_i = -\partial_{q_i} \mathfrak{H}(\mathbf{p}; \mathbf{q})$ are

$$\partial_t q_i(t) = p_i(t), \quad \text{and} \quad \partial_t p_i(t) = e^{q_{i-1}(t) - q_i(t)} - e^{q_i(t) - q_{i+1}(t)}.$$

This model may be thought of as a system of particles moving on the real line, with locations (q_i) and momenta (p_i) . It was originally introduced by Toda [49] as a Hamiltonian dynamic that admits

soliton solutions, which (loosely speaking) are localized, wave-like functions that retain their shape as they propagate in time. Since the works of Flaschka [19] and Manakov [32] exhibiting its full set of conserved quantities, and that of Moser [39] determining its scattering shift, the Toda lattice has become recognized as an archetypal example of a completely integrable system.

A basic question about the Toda lattice is to understand its behavior as the domain J and time t become large. In situations when its initial data either decays or approximates a profile that is smooth almost everywhere, detailed answers have been attained (largely by analyzing associated Riemann–Hilbert problems) in works of Venakides–Deift–Oba [50], Deift–Kamviss–Kriecherbauer–Zhou [12], Deift–McLaughlin [13], Bloch–Golse–Paul–Uribe [5], and Krüger–Teschl [28].

However, much less is known when the initial data is rough and not decaying, or random. These situations were emphasized Zakharov, broadly within the context of classical integrable systems, under the terms “soliton gases” [51] and “integrable turbulence” [52]. They are especially prominent, since invariant measures for integrable dynamics are of this type. For example, perhaps the most natural invariant measure for the Toda lattice, sometimes called *thermal equilibrium*, is when the (p_j) and $(e^{q_j - q_{j+1}})$ are independent Gaussian and Gamma random variables, respectively.

Over the past decade, an extensive framework has emerged in the physics literature for predicting how an integrable system behaves in these situations. It is based on the notion that it can be thought of as a dense collection of many solitons, each possessing an amplitude λ_j and a location $Q_j(t)$, which have two properties that together pinpoint the system’s asymptotics. First, interesting quantities describing the integrable system, such as local charges and currents, are approximable by simple expressions of the solitons (for example, in the Toda lattice, the number of particles in an interval should nearly equal the number of solitons in it). Second, the evolution of these solitons is well-approximated by an explicit equation, which for the Toda lattice reads [16, Section IV]

$$(1.1) \quad Q_k(t) \approx Q_k(0) + \lambda_k t - 2 \sum_{j: Q_j(t) < Q_k(t)} \log |\lambda_k - \lambda_j| + 2 \sum_{j: Q_j(0) < Q_k(0)} \log |\lambda_k - \lambda_j|.$$

The relation (1.1) can be informally interpreted as follows. The k -th soliton, initially at $Q_k(0)$, moves with velocity λ_k until it meets another soliton, say that j -th one. At that moment, the k -th soliton instantaneously moves forward or backward by $2 \log |\lambda_k - \lambda_j|$, depending on whether it met the j -th soliton from the right or left, respectively (this could make it pass another soliton, producing a “cascade” of such interactions, but no two solitons can interact with each other twice in this way). Then, the k -th soliton proceeds at velocity λ_k until meeting another soliton, when the procedure repeats. The reason for the choice $2 \log |\lambda_k - \lambda_j|$ is that it is the Toda lattice’s scattering shift, describing the phase displacement of just two solitons passing through each other. As such, if the collection of solitons is sparse, one may expect them to mainly interact successively in pairs, thereby giving rise to the above dynamics. This reasoning led to the prediction in [51] that a form of (1.1) should hold in this sparse setting, for the Korteweg–de Vries (KdV) equation.

That a relation such as (1.1) should persist if the solitons are dense was first postulated (for various quantum integrable systems) by Bertini–Collura–De Nardis–Fagotti [4] and Castro-Alvaredo–Doyon–Yoshimura [10]. Although less intuitive in this dense regime, some of its consequences for the Toda lattice at thermal equilibrium were investigated through simulations by Cao–Bulchadani–Spohn in [9], and verified to striking numerical accuracy. Below, we refer to (1.1) as the *asymptotic scattering relation*; it is also sometimes called the “collision rate ansatz” or “flea-gas algorithm.”

The question of mathematically making sense of, and justifying, the above framework (for at least some Hamiltonian integrable system) has received considerable interest in recent years. To that

end, the soliton amplitudes λ_j are understood. They are defined to be the conserved quantities for the system, given by the eigenvalues of its *Lax matrix*. For the Toda lattice, this is the tridiagonal, symmetric matrix $\mathbf{L}(t)$ whose diagonal and off-diagonal entries are the (p_i) and $(e^{(q_i - q_{i+1})/2})$, respectively [19, 32]. When \mathcal{J} is large and $(\mathbf{p}(t); \mathbf{q}(t))$ is random, $\mathbf{L}(t)$ becomes a high-dimensional random matrix. Its eigenvalue density then prescribes the distribution of soliton amplitudes in the Toda lattice, whose computation was addressed by Spohn [44] (after initial work of Oppen [40]) under thermal equilibrium and many other invariant measures. He predicted formulas for its limiting density (and derived expectations for local currents), which were later verified in works of Mazucca, Guionnet, and Memin [33, 24, 34].

The above results do not address the evolution of the soliton locations (Q_j) in time. Even a coherent definition of these locations (Q_j) from a given state $(\mathbf{p}; \mathbf{q})$ of the Toda lattice does not seem to exist in the mathematics literature. In fact, we only know of two integrable systems for which such a definition has been given; they are the hard rods model and box-ball system. For the former, the soliton locations are simply the positions of the rods; for the latter, they can be recovered through a combinatorial algorithm of Takahashi–Satsuma [46]. In each case, the analog of the approximate evolution (1.1) becomes exact, and it is an eventual consequence of the inverse scattering that linearizes the system; see the works by Boldrighini–Dubroshin–Suhov [6] and Croydon–Sasada [11].

As most integrable systems lack a definition for their soliton locations from a given state, researchers have turned to studying specific families of solutions for them, called finite-gap solutions (see the survey of Dubrovin–Matveev–Novikov [17]). These are associated with an algebraic curve and allow logical candidates for soliton locations to be incorporated as tunable parameters called phases. The physics work of El [18] proposed a specific scaling limit, involving making the curve’s genus large and the phases random, under which finite-gap solutions to the KdV equation should satisfy the analog of (1.1). While certain infinite genus finite-gap solutions have been studied by mathematicians since the work of McKean–Trubowitz [35], the scaling limit from [18] does not seem to have received a thorough mathematical treatment. Still, in a different direction, the papers by Girotti, Grava, Jenkins, McLaughlin, Minakov, and Nanjudel analyzed finite-gap solutions of low genus with many solitons (allowing for random amplitudes) [20, 22]. By modifying these solutions to include one large soliton passing through the many small ones, they used Riemann–Hilbert methods to prove [21] that the position of this large “tracer” soliton satisfies a version of (1.1).

While finite-gap solutions can sometimes be amenable to analysis, they become very complicated when expressed in terms of the original variables on which the integrable system is defined (which, for the Toda lattice, would be the $(\mathbf{p}; \mathbf{q})$ ones). Partly for this reason, a precise interpretation and proof of (1.1) for Hamiltonian integrable systems under most natural families of random initial data (particularly product measures, such as thermal equilibria) had until now been unavailable.

The task of making sense of, and justifying, the framework behind (1.1) may be viewed as threefold.

- (1) Define the soliton locations $(Q_j(t))$.
- (2) Show that relevant quantities for the integrable system (such as local charges and currents) are approximable by simple functions of the soliton locations $(Q_j(t))$ and amplitudes λ_j .
- (3) Establish that the approximate asymptotic scattering relation (1.1) holds.

In this paper we implement these three tasks for the Toda lattice at thermal equilibrium. See Definition 2.6 for the first; Proposition 2.10 for the second; and Theorem 2.11 for the third.

We should mention that there is also the fourth task of using (1.1) to prove asymptotic results about the Toda lattice. This can be done as well, though we defer its implementation to the sequel paper [1], as the ideas used there are almost entirely orthogonal from the ones introduced here.

Before describing the ideas in this paper, we briefly explain on the setup. Recall that the domain \mathcal{J} of the Toda lattice is either an interval, a torus, or the line. As Proposition 2.5 below, we will show (if the lengths of the interval and torus are sufficiently large) that these three choices lead to approximately equal systems, which differ by an error decaying exponentially in the time parameter t . So, we mainly assume below that $\mathcal{J} = [N_1, N_2]$ is an interval of length $N = N_2 - N_1 + 1 \gg t$. Then, the rows and columns of the Lax matrix $\mathbf{L}(t) = [L_{ij}(t)]$ are indexed by $i, j \in \mathcal{J} = [N_1, N_2]$.

We consider the Toda lattice at thermal equilibrium, with parameters $\beta, \theta > 0$. This means (see Definition 2.4) that we sample the diagonal and superdiagonal entries of the tridiagonal, symmetric Lax matrix $\mathbf{L}(0)$ independently, with probability densities $C_\beta e^{-\beta x^2/2}$ and $C_{\beta, \theta} x^{2\theta-1} e^{-\beta x^2}$, respectively (where $C_\beta, C_{\beta, \theta} > 0$ are normalization constants). The near-invariance of thermal equilibrium implies that this description continues to approximately¹ hold for $\mathbf{L}(t)$, when $t > 0$.

1. *Soliton locations:* First, we must provide a definition for the location $Q_j(t)$ of the j -th soliton at time t . Recall that its amplitude is an eigenvalue λ_j of the Lax matrix $\mathbf{L}(t)$. So, we examine the corresponding unit eigenvector $\mathbf{u}_j(t) = (u_j(N_1; t), \dots, u_j(N_2; t))$. Central to our analysis is the fact that, if $\mathbf{L}(t)$ is under thermal equilibrium, then $\mathbf{u}_j(t)$ is *exponentially localized*. This means that it admits some “center” $\varphi \in [N_1, N_2]$ such that $|u_j(i; t)| \leq C e^{-c|i-\varphi|}$ likely holds for any $i \in [N_1, N_2]$. Such exponential decay of eigenvectors holds generally for random tridiagonal (or bounded band) matrices with independent (or weakly correlated) entries. It was first proven in the context of one-dimensional Anderson localization by Molchanov [37] and Kunz–Souillard [29], though the precise estimates we will use are due to Schenker [42] and Aizenman–Schenker–Friedrich–Hundertmark [2].

Now fix a parameter $\zeta > 0$ that is not too small; we will take $\zeta = e^{-C(\log N)^{3/2}}$ (which decays in N superpolynomially, but not stretched exponentially). Define a ζ -*localization center* of λ_j with respect to $\mathbf{L}(t)$ to be any index $\varphi_t(j) \in [N_1, N_2]$ for which $|u_j(\varphi_t(j); t)| \geq \zeta$; see Definition 2.6. We view this localization center $\varphi_t(j)$ as the index of the particle associated with the j -th soliton, so we then define the location of this soliton on \mathbb{R} to be this particle’s position $Q_j(t) = q_{\varphi_t(j)}(t)$.

Observe that there may be multiple choices for this index $\varphi_t(j)$, and thus for the soliton location $Q_j(t)$. However, exponential localization of $\mathbf{u}_j(t)$ quickly implies that all such choices will be very close to (within $\mathcal{O}(|\log \zeta|) = N^{o(1)}$ of) each other, so will be equivalent for asymptotic purposes.

Before proceeding, let us mention that, guided by quantum mechanics, one might be inclined to instead define the index $\varphi_t(j)$ to be random variable, whose probability of equalling any $k \in [N_1, N_2]$ is $u_j(k; t)^2$. Indeed, this notion was hypothesized in the earlier physics work of Bulchandini–Cao–Moore [7, Section 3.2], who observed this probability distribution concentrates on few indices (also attributing it to Anderson localization). If we had adopted this as the definition of $\varphi_t(j)$ in our context, we would have obtained essentially the same results as below, with similar proofs.

2. *Approximate locality:* Next, we must verify that interesting quantities of the Toda lattice (local charges and currents) are simply expressible through the soliton locations $Q_j(t)$ defined above. This will follow from a more fundamental property, which is the “approximate locality” of eigenvalues. To explain this, a *local* quantity for the Toda lattice is one that is expressible as a function of the Lax matrix entries $L_{ik}(t)$, for i and k in a uniformly bounded subinterval of $[N_1, N_2]$. The

¹Thermal equilibrium is exactly invariant if \mathcal{J} is the torus or line. While this does not quite hold when \mathcal{J} is an interval, the above-mentioned comparison between these domains implies that it is approximately true in this case as well. To ease this introductory exposition, we ignore this subtlety for now and act as if invariance held exactly.

momentum $p_i(t)$ of the i -th particle is an example of one, as it is the diagonal (i, i) -entry $L_{ii}(t)$ of $\mathbf{L}(t)$. On the other hand, eigenvalues λ_j of $\mathbf{L}(t)$ are not local, as they depend on all entries of $\mathbf{L}(t)$.

However, we will see under thermal equilibrium that λ_j is “approximately local” around its localization center $\varphi_t(j)$; informally, this means that the “dependence” of λ_j on the i -th row and column of $\mathbf{L}(t)$ decays exponentially in $|i - \varphi_t(j)|$. A more specific (but slightly simplified) statement is, if $\ell \gg 1$ is a parameter and $\tilde{\mathbf{L}}(t)$ is a tridiagonal matrix whose (i, k) -entry coincides with that of $\mathbf{L}(t)$ whenever $i, k \in [\varphi_t(j) - \ell, \varphi_t(j) + \ell]$, then $\tilde{\mathbf{L}}(t)$ likely has an eigenvalue $\tilde{\lambda}_j$ satisfying $|\lambda_j - \tilde{\lambda}_j| \leq Ce^{-c\ell}$; see Corollary 5.5 and Corollary 5.6 for precise formulations. This approximate locality can be deduced from the exponential eigenvector localization of $\mathbf{L}(t)$. Similar deductions were used by Molchanov [38] and Minami [36] in their studies of spectral statistics for random Schrödinger operators.

Given this approximate locality, let us outline how local charges and currents can be recovered from soliton data. Instead of explaining this in complete generality (which would require us to recall the full family of Toda charges and currents; see Definition 2.3 for that), to simplify the exposition, we consider a specific example, which computes the total momentum of Toda particles in a large interval $\mathcal{J} \subseteq \mathbb{R}$. In this case, the predicted relation reads [16, Section III.B]

$$\sum_{i:q_i(t) \in \mathcal{J}} p_i(t) \approx \sum_{j:Q_j(t) \in \mathcal{J}} \lambda_j.$$

Its left side is the total momentum (which is the first local charge) in \mathcal{J} , and its right side is the total amplitude of all solitons in \mathcal{J} ; see Proposition 2.10 for the general statement. It can be shown that the Toda particles $(q_i(t))$ are nearly ordered, so since $Q_j(t) = q_{\varphi_t(j)}(t)$, the above reduces to

$$(1.2) \quad \sum_{i \in \mathcal{I}} p_i(t) \approx \sum_{j: \varphi_t(j) \in \mathcal{I}} \lambda_j,$$

for any large interval of indices $\mathcal{I} \subseteq [N_1, N_2]$ (whose endpoints are the smallest and largest i for which $q_i(t) \in \mathcal{J}$). The left side of (1.2) is the trace of the matrix $\tilde{\mathbf{L}}(t)$ obtained from $\mathbf{L}(t)$ by setting any entry with indices not in \mathcal{I} to 0. Now, by approximate locality, most eigenvalues λ_j on the right side of (1.2) (namely, those whose localization centers $\varphi_t(j)$ are not too close to an endpoint of \mathcal{I}) are approximately equal to a corresponding eigenvalue $\tilde{\lambda}_j$ of $\tilde{\mathbf{L}}(t)$. This confirms (1.2), as it implies that

$$\sum_{i \in \mathcal{I}} p_i(t) = \text{Tr } \tilde{\mathbf{L}}(t) = \sum_j \tilde{\lambda}_j \approx \sum_{j: \varphi_t(j) \in \mathcal{I}} \lambda_j.$$

3. *Asymptotic scattering relation:* It remains to access the soliton dynamics, by proving (1.1). We first use the inverse scattering relation for the Toda lattice, shown in [39], providing the explicit (linear) evolution for the first entry $u_k(N_1; t)$ of the eigenvector $\mathbf{u}_k(t)$ of $\mathbf{L}(t)$. It states that

$$(1.3) \quad 2 \log |u_k(N_1; t)| = 2 \log |u_k(N_1; 0)| - \lambda_k t + \mathfrak{C}(t),$$

for some constant $\mathfrak{C}(t)$ independent of k . Now, recall since $\mathbf{u}_k(s)$ is exponentially localized for any $s \in [0, t]$ that $\log |u_k(N_1; s)|$ decays linearly in $\varphi_s(k) - N_1$. In fact, there is a *Lyapunov exponent* $\gamma_k \leq 0$ satisfying $\log |u_k(N_1; s)| \approx \gamma_k \cdot (\varphi_s(k) - N_1)$, which admits an exact formula called the *Thouless relation*. Predicted by Thouless [48] and established by Avron–Simon [3], it states

$$(1.4) \quad 2\gamma_k = 2 \cdot \mathbb{E}[\log L_{i, i+1}(s)] - 2 \int_{-\infty}^{\infty} \log |\lambda_k - \lambda| \varrho(\lambda) d\lambda,$$

where ϱ is the limiting eigenvalue density for $\mathbf{L}(s)$ (as its dimension becomes large), and $i \in [N_1, N_2 - 1]$ is any index (since the $L_{i,i+1}(s)$ are identically distributed under thermal equilibrium, $\mathbb{E}[L_{i,i+1}(s)]$ does not depend i). While the relation (1.4) will not be of direct use to us in proving (1.1), a discrete variant of it will be. More precisely, by combining a linear algebraic identity in [48], the transfer matrix arguments of [3], and the approximate locality of eigenvalues, we will show (see Proposition 6.1) that

$$(1.5) \quad 2 \log |u_k(N_1; s)| \approx q_{N_1}(s) - Q_k(s) - 2 \sum_{j: \varphi_s(j) < \varphi_s(k)} \log |\lambda_k - \lambda_j|.$$

Let us briefly indicate the relation between (1.5) and (1.4). Since (by definition) $2 \log L_{i,i+1}(s) = q_i(s) - q_{i+1}(s)$ and $Q_k = q_{\varphi_s(k)}(s)$, one may expect the first two terms on the right side of (1.5) to approximate the first term on the right side of (1.4), multiplied by $\varphi_s(k) - N_1$. Also, since the sum on the right side of (1.5) constitutes $\varphi_s(k) - N_1$ terms (and since ϱ is the limiting density of the (λ_i)), one may expect it to approximate the integral on the right side of (1.4), multiplied by $\varphi_s(k) - N_1$. In this way, (1.5) may be thought of as a discrete form of (1.4).

Inserting (1.5) at $s \in \{0, t\}$ into (1.3) yields

$$(1.6) \quad \begin{aligned} Q_k(t) \approx & Q_k(0) + \lambda_k(t) - 2 \sum_{j: \varphi_t(j) < \varphi_t(k)} \log |\lambda_k - \lambda_j| + 2 \sum_{j: \varphi_0(j) < \varphi_0(k)} \log |\lambda_k - \lambda_j| \\ & + q_{N_1}(t) - q_{N_1}(0) - \mathfrak{C}(t). \end{aligned}$$

It can be shown that $\mathfrak{C}(t) \approx q_{N_1}(t) - q_{N_1}(0)$. Using this in (1.6), with the fact that the (q_j) are ordered (to transform the restrictions on $\varphi_s(j)$ in (1.6) to ones on $q_{\varphi_s(j)}(s) = Q_j(s)$), gives (1.1).

In fact, upon implementing this outline, one finds (see Theorem 2.11) that the error in (1.1) is quite small, of order $(\log N)^C$. This indicates that the asymptotic scattering relation is close to exact, which might explain why the simulations in [9] had such a high degree of numerical accuracy.

Before proceeding, we briefly comment on the potential applicability of the above framework to other initial data or different integrable systems. First, there are many (an infinite-dimensional family of) invariant measures for the Toda lattice other than thermal equilibrium; they are the *generalized Gibbs ensembles* with polynomial potential (see [44, Section 2]). Under these measures, the entries $(p_i, e^{(q_i - q_{i+1})/2})$ of the Lax matrix $\mathbf{L}(t)$ are not independent, but their coupling is of finite range. It was indicated in [42, Section 2] that the arguments of [42] should apply to show that such tridiagonal matrices also have exponentially localized eigenvectors. Thus, it could be possible to establish above results under generalized Gibbs ensembles, as well.

Next, one might inquire about initial data that is not invariant. If this initial data is sampled under a product measure (or one with finite-range couplings), then again the framework of [42] should yield exponential eigenvector localization of the initial Lax matrix $\mathbf{L}(0)$. However, since the initial data is not invariant, this does not immediately extend to later times $t \gg 1$. If one could show that the eigenvectors of $\mathbf{L}(t)$ were still localized for $t \gg 1$, then it may again be possible to show the asymptotic scattering relation (1.1) remains valid away from invariant initial data.

There are numerous other integrable systems admitting Lax matrices that are (skew-)symmetric tridiagonal or of Cantero–Moral–Velázquez (CMV) [8] type. These include the Volterra lattice; generalized Toda lattices studied by Deift–Nanda–Tomei [14]; and Ablowitz–Ladik (and Schur flow) hierarchy considered by Killip–Nenciu [27]. As explained by Grava–Gisonni–Gubbiotti–Mazzuca [23] and Spohn [45], invariant measures analogous to thermal equilibrium exist for these systems. Moreover, relevant eigenvector properties of the associated random matrices are known in many of

these cases (though sometimes in a weaker form than what we use here). For example, exponential eigenvector localization and an analog of the Thouless relation for random CMV matrices were established in [43] by Simon. These properties also hold in certain settings for various continuum integrable equations, such as the KdV equation² [37, 3]. It would be of interest to investigate if the framework developed here can be applied to these contexts, as well.

Acknowledgements. The author heartily thanks Alexei Borodin, Patrick Lopatto, Jeremy Quastel, and Herbert Spohn for valuable conversations. This work was partially supported by a Clay Research Fellowship and a Packard Fellowship for Science and Engineering.

2. RESULTS

We introduce the Toda lattice and its Lax matrix in Section 2.1, and we define its thermal equilibrium initial data in Section 2.2. In Section 2.3, we introduce localization centers, and we explain how they can be used to access locally conserved quantities (also called local charges) and currents. We then state the asymptotic scattering relation for the Toda lattice under thermal equilibrium (Theorem 2.11), in Section 2.4.

Throughout, for any $a, b \in \mathbb{R}$, set $\llbracket a, b \rrbracket = [a, b] \cap \mathbb{Z}$. A vector $\mathbf{v} = (v_1, v_2, \dots, v_n) \in \mathbb{C}^n$ is a unit vector if $\sum_{i=1}^n v_i^2 = 1$; we call it nonnegatively normalized if $v_j > 0$, where $j \in \llbracket 1, n \rrbracket$ is the minimal index such that $v_j \neq 0$. For any real symmetric $n \times n$ matrix \mathbf{M} , let $\text{eig } \mathbf{M} = (\lambda_1, \lambda_2, \dots, \lambda_n)$ denote the eigenvalues of \mathbf{M} , counted with multiplicity and ordered so that $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$.

2.1. Toda Lattice.

2.1.1. *Open Toda Lattice.* In this section we recall the Toda lattice on an interval. Throughout, we fix integers $N_1 \leq N_2$ and set $N = N_2 - N_1 + 1$ (which will prescribe the interval's endpoints and length, respectively).

The state space of the Toda lattice on the interval $\llbracket N_1, N_2 \rrbracket$, also called the *open Toda lattice*, is given by a pair of N -tuples $(\mathbf{p}(t); \mathbf{q}(t)) \in \mathbb{R}^N \times \mathbb{R}^N$, where $\mathbf{p}(t) = (p_{N_1}(t), p_{N_1+1}(t), \dots, p_{N_2}(t))$ and $\mathbf{q}(t) = (q_{N_1}(t), q_{N_1+1}(t), \dots, q_{N_2}(t))$; both are indexed by a real number $t \geq 0$ called the time. Given any *initial data* $(\mathbf{p}(0); \mathbf{q}(0)) \in \mathbb{R}^N \times \mathbb{R}^N$, the joint evolution of $(\mathbf{p}(t); \mathbf{q}(t))$ for $t \geq 0$ is prescribed by the system of ordinary differential equations

$$(2.1) \quad \partial_t q_j(t) = p_j(t), \quad \text{and} \quad \partial_t p_j(t) = e^{q_{j-1}(t) - q_j(t)} - e^{q_j(t) - q_{j+1}(t)},$$

for all $(j, t) \in \llbracket N_1, N_2 \rrbracket \times \mathbb{R}_{\geq 0}$; here, we set $q_{N_1-1}(t) = -\infty$ and $q_{N_2+1}(t) = \infty$ for all $t \geq 0$. One might interpret this as the dynamics for N points (indexed by $\llbracket N_1, N_2 \rrbracket$) moving on the real line, whose locations and momenta at time $t \geq 0$ are given by the $(q_i(t))$ and $(p_i(t))$, respectively.

The system of differential equations (2.1) is equivalent to the Hamiltonian dynamics generated by the Hamiltonian $\mathfrak{H} : \mathbb{R}^N \times \mathbb{R}^N \rightarrow \mathbb{R}$ that is defined, for any $\mathbf{p} = (p_0, p_1, \dots, p_{N-1}) \in \mathbb{R}^N$ and $\mathbf{q} = (q_0, q_1, \dots, q_{N-1}) \in \mathbb{R}^N$, by setting

$$(2.2) \quad \mathfrak{H}(\mathbf{p}; \mathbf{q}) = \sum_{j=0}^{N-1} \left(\frac{p_j^2}{2} + e^{q_j - q_{j+1}} \right),$$

²As observed by Saitoh [41] (and by Matetski–Quastel–Remenik [26]), the KdV equation can also be formally recovered from a limit degeneration of the Toda lattice. It remains to be seen whether this can be justified to allow asymptotic frameworks about the latter to carry over to the former.

where $q_N = \infty$. The existence and uniqueness of solutions to (2.1) for all time $t \geq 0$, under arbitrary initial data $(\mathbf{p}; \mathbf{q}) \in \mathbb{R}^N \times \mathbb{R}^N$, is thus a consequence of the Picard–Lindelöf theorem (see, for example, the proof of [47, Theorem 12.6]).

It will often be useful to reparameterize the variables of the Toda lattice, following [19]. To that end, for any $(i, t) \in \llbracket N_1, N_2 \rrbracket \times \mathbb{R}_{\geq 0}$, define

$$(2.3) \quad r_i(t) = q_{i+1}(t) - q_i(t); \quad a_i(t) = e^{-r_i(t)/2}; \quad b_i(t) = p_i(t).$$

Denoting $\mathbf{a}(t) = (a_{N_1}(t), a_{N_1+1}(t), \dots, a_{N_2}(t)) \in \mathbb{R}_{\geq 0}^N$ and $\mathbf{b}(t) = (b_{N_1}(t), b_{N_1+1}(t), \dots, b_{N_2}(t)) \in \mathbb{R}^N$, the $(\mathbf{a}(t); \mathbf{b}(t))$ are sometimes called *Flaschka variables*; they satisfy $r_{N_2}(t) = q_{N_2+1}(t) - q_{N_2}(t) = \infty$ and $a_{N_2}(t) = e^{-r_{N_2}(t)/2} = 0$. Then, (2.1) is equivalent to the system

$$(2.4) \quad \partial_t a_j(t) = \frac{a_j(t)}{2} \cdot (b_j(t) - b_{j+1}(t)), \quad \text{and} \quad \partial_t b_j(t) = a_{j-1}(t)^2 - a_j(t)^2,$$

for each $(j, t) \in \llbracket N_1, N_2 \rrbracket \times \mathbb{R}_{\geq 0}$.

It will at times be necessary to define the original Toda state space variables $(\mathbf{p}(t); \mathbf{q}(t))$ from the Flaschka variables $(\mathbf{a}(t); \mathbf{b}(t))$; it suffices to do this at $t = 0$, as $(\mathbf{p}(t); \mathbf{q}(t))$ is determined from $(\mathbf{p}(0); \mathbf{q}(0))$, by (2.1). We explain how to do this when $0 \in \llbracket N_1, N_2 \rrbracket$ (as otherwise we may translate (N_1, N_2) to guarantee that this inclusion holds).³ By (2.3), the Flaschka variables $\mathbf{a}(0)$ only specify the differences between consecutive entries in $\mathbf{q}(0)$, so the former only determines the latter up to an overall shift. We will fix this shift by setting $q_0(0) = 0$, that is, we define $(\mathbf{p}(0); \mathbf{q}(0))$ by imposing

$$(2.5) \quad q_0(0) = 0; \quad q_{i+1}(0) - q_i(0) = -2 \log a_i(0); \quad p_i(0) = b_i(0),$$

for each $i \in \llbracket N_1, N_2 \rrbracket$. Then, $(\mathbf{p}(0); \mathbf{q}(0))$ is called the Toda state space initial data associated with $(\mathbf{a}(0); \mathbf{b}(0))$. The evolution $(\mathbf{p}(t), \mathbf{q}(t))$ of this initial data under (2.1) is called the Toda state space dynamics associated with $(\mathbf{a}(t), \mathbf{b}(t))$; observe that we may have $q_0(t) \neq 0$ if $t \neq 0$.

2.1.2. Lax Matrices, Locally Conserved Quantities, and Currents. In this section we recall the Lax matrix, locally charges, and currents associated with the Toda lattice. Throughout, we fix integers $N_1 \leq N_2$ and set $N = N_2 - N_1 + 1$. Let $(\mathbf{a}(t); \mathbf{b}(t)) \in \mathbb{R}_{\geq 0}^N \times \mathbb{R}^N$ be a pair of N -tuples indexed by a real number $t \in \mathbb{R}_{\geq 0}$, where $\mathbf{a}(t) = (a_j(t))$ and $\mathbf{b}(t) = (b_j(t))$ satisfies the system (2.4) for each $(j, t) \in \llbracket N_1, N_2 \rrbracket \times \mathbb{R}$; we assume (as explained above (2.4)) that $a_{N_2}(t) = 0$ for each $t \in \mathbb{R}_{\geq 0}$. Under this notation, the Lax matrix (introduced in [19, 32]) is defined as follows.

Definition 2.1 (Lax matrix). For any real number $t \geq 0$, the *Lax matrix* $\mathbf{L}(t) = [L_{ij}] = [L_{ij}(t)]$ is an $N \times N$ real symmetric matrix, with entries indexed by $i, j \in \llbracket N_1, N_2 \rrbracket$, defined as follows. Set

$$L_{ii} = b_i(t), \quad \text{for each } i \in \llbracket N_1, N_2 \rrbracket; \quad L_{i, i+1} = L_{i+1, i} = a_i(t), \quad \text{for each } i \in \llbracket N_1, N_2 - 1 \rrbracket.$$

Also set $L_{ij} = 0$ for any $i, j \in \llbracket N_1, N_2 \rrbracket$ with $|i - j| \geq 2$.

A fundamental feature of the Lax matrix is that its eigenvalues are preserved under the Toda dynamics (2.4). This was originally due to [19]; see also [39, Section 2].

Lemma 2.2 ([19, 39]). *For any real numbers $t, t' \in \mathbb{R}_{\geq 0}$, we have $\text{eig } \mathbf{L}(t) = \text{eig } \mathbf{L}(t')$.*

Lemma 2.2 provides a large family of conserved quantities for the Toda lattice, given by the eigenvalues of the Lax matrix. However, these are “non-local,” in the sense that they depend on all of the Flaschka variables $(\mathbf{a}(t); \mathbf{b}(t))$, as opposed to only the $(a_i(t), b_i(t))$ for i in some (uniformly)

³In this work, we will usually have N_1 and N_2 be large negative and large positive integers, respectively, and we will be interested in the $(p_i(t); q_i(t))$ for i in some interval containing 0.

bounded interval. To remedy this, observe that Lemma 2.2 implies that $\text{Tr } \mathbf{L}(t)^m$ is conserved for any integer $m \geq 0$. Since $\mathbf{L}(t)$ is tridiagonal, the diagonal entries of $\mathbf{L}(t)^m$ are local quantities (see [45, Section 2.1] for further information), whose total is preserved; they are called local charges and have associated currents. In what follows, for any matrix \mathbf{M} , we let $[M]_{ij}$ denote the (i, j) entry of \mathbf{M} .

Definition 2.3 (Local charges and currents). Fix an integer $m \geq 0$, an index $i \in \llbracket N_1, N_2 \rrbracket$, and a real number $t \geq 0$. Define the m -th local charge $\mathfrak{k}_i^{[m]}(t)$ of $\mathbf{L}(t)$ at i by setting $\mathfrak{k}_i^{[m]}(t) = [\mathbf{L}(t)^m]_{ii}$. Define the associated m -th current $\mathfrak{j}_i^{[m]}(t)$ of $\mathbf{L}(t)$ at i by setting

$$\mathfrak{j}_i^{[m]}(t) = a_{i-1}(t) \cdot [\mathbf{L}(t)^m]_{i,i-1}, \quad \text{if } i \in \llbracket N_1 + 1, N_2 \rrbracket; \quad \mathfrak{j}_{N_1}^{[m]}(t) = 0, \quad \text{if } i \leq N_1 \text{ or } i > N_2.$$

For example, the first local charge $\mathfrak{k}_i^{[1]}(t) = b_i(t) = p_i(t)$ denotes momentum. To explain the currents, using (2.4) it can be verified that (see [45, Equation (2.24)])

$$(2.6) \quad \partial_s \mathfrak{k}_i^{[m]}(s) = \mathfrak{j}_i^{[m]}(s) - \mathfrak{j}_{i+1}^{[m]}(s), \quad \text{for all } i \in \llbracket N_1, N_2 \rrbracket \text{ and } s \geq 0.$$

In this way, the current $\mathfrak{j}_i^{[m]}$ signifies the ‘‘rate of transfer’’ of the local charge $\mathfrak{k}_i^{[m]}$ across the edge connecting $i + 1$ to i .

2.2. Thermal Equilibrium. In this section we describe a class of random initial data that we will study for the Toda lattice; it is sometimes referred to as thermal equilibrium, and is given by independent Gamma random variables for the \mathbf{a} Flaschka variables, and independent Gaussian random variables for the \mathbf{b} ones. In what follows, we fix an integer $N \geq 1$ and real numbers $\beta, \theta > 0$.

Definition 2.4 (Open thermal equilibrium). The (open) thermal equilibrium with parameters $(\beta, \theta; N)$ is the product measure $\mu = \mu_{\beta, \theta} = \mu_{\beta, \theta; N-1, N}$ on $\mathbb{R}^{N-1} \times \mathbb{R}^N$ defined by

$$\mu(d\mathbf{a}; d\mathbf{b}) = \left(\frac{2\beta^\theta}{\Gamma(\theta)} \right)^{N-1} (2\pi\beta^{-1})^{-N/2} \cdot \prod_{j=0}^{N-2} a_j^{2\theta-1} e^{-\beta a_j^2} da_j \prod_{j=0}^{N-1} e^{-\beta b_j^2/2} db_j,$$

where $\mathbf{a} = (a_0, \dots, a_{N-2}) \in \mathbb{R}_{\geq 0}^{N-1}$ and $\mathbf{b} = (b_0, b_1, \dots, b_{N-1}) \in \mathbb{R}^N$. It will be convenient to view $\mu_{\beta, \theta; N-1, N}$ as a measure on $\mathbb{R}^N \times \mathbb{R}^N$ by, if we denote $\hat{\mathbf{a}}(0) = (a_0, a_1, \dots, a_{N-2}, 0) \in \mathbb{R}_{\geq 0}^N$, then also saying $(\hat{\mathbf{a}}; \mathbf{b}) \in \mathbb{R}^N \times \mathbb{R}^N$ is sampled under $\mu_{\beta, \theta; N-1, N}$.

Thermal equilibrium, as described above, is related to invariant measures for the Toda lattice; the latter are measures on the Flaschka variable initial data $(\mathbf{a}(0); \mathbf{b}(0))$ such that, for any $t \geq 0$, the law of $(\mathbf{a}(t); \mathbf{b}(t))$ under the Toda lattice is the same as that of $(\mathbf{a}(0); \mathbf{b}(0))$. The open Toda lattice on a finite interval $\llbracket N_1, N_2 \rrbracket$ admits no nontrivial invariant measures. However, the Toda lattice on the full line \mathbb{Z} does, though this must be made sense of, as it involves infinitely many variables. In this context, the thermal equilibrium product measure of Definition 2.4 (extrapolated to when $N = \infty$) is perhaps the most natural invariant measure.

The following proposition, to be shown in Section 4.3 below (as a consequence of Proposition 4.7, which addresses more general initial data), states this to be the case. Specifically, it shows that at thermal equilibrium the Toda lattice on \mathbb{Z} can be defined by taking a limit of open Toda lattices on growing intervals; confirms that thermal equilibrium is invariant for this infinite Toda lattice; and provides a quantitative approximation for it by open Toda lattices at thermal equilibrium on finite intervals.

Proposition 2.5. *Let $(a_j)_{j \in \mathbb{Z}}$ and $(b_j)_{j \in \mathbb{Z}}$ be mutually independent random variables, whose laws for each $j \in \mathbb{Z}$ are given by*

$$\mu(da_j) = 2\beta^\theta \Gamma(\theta)^{-1} \cdot a_j^{2\theta-1} e^{-\beta a_j^2} da_j, \quad \text{and} \quad \mu(db_j) = (2\pi\beta^{-1})^{-1/2} \cdot e^{-\beta b_j^2/2} db_j.$$

For any integers $N_1 \leq N_2$, let $(\mathbf{a}^{[N_1, N_2]}(t), \mathbf{b}^{[N_1, N_2]}(t))$ denote the open Toda lattice (2.4) on $\llbracket N_1, N_2 \rrbracket$, with initial data $(\mathbf{a}^{[N_1, N_2]}(0); \mathbf{b}^{[N_1, N_2]}(0))$ given by $\mathbf{a}^{[N_1, N_2]}(0) = (a_{N_1}, a_{N_1+1}, \dots, a_{N_2-1})$ and $\mathbf{b}^{[N_1, N_2]}(0) = (b_{N_1}, b_{N_1+1}, \dots, b_{N_2})$.

- (1) For each $(j, t) \in \mathbb{Z} \times \mathbb{R}_{\geq 0}$, the limits $\lim_{N \rightarrow \infty} a_j^{[-N, N]}(t) = a_j(t)$ and $\lim_{N \rightarrow \infty} b_j^{(N)}(t) = b_j(t)$ exist almost surely; are finite; and solve (2.4).
- (2) For each $t \geq 0$, denoting $\mathbf{a}(t) = (a_j(t))_{j \in \mathbb{Z}}$ and $\mathbf{b}(t) = (b_j(t))_{j \in \mathbb{Z}}$, the laws of $(\mathbf{a}(0); \mathbf{b}(0))$ and $(\mathbf{a}(t); \mathbf{b}(t))$ coincide.
- (3) For any $\mathbf{p} \in (0, 1)$, there exist constants $c = c(\beta, \theta, \mathbf{p}) > 0$ and $C = C(\beta, \theta, \mathbf{p}) > 1$ such that the following holds. Let $R \geq 1$ and $T \geq 0$ be real numbers, and $K \geq 1$ and $N_1 \leq -K \leq K \leq N_2$ be integers, such that

$$R \geq \log N, \quad \text{and} \quad RT \leq K \leq N^{\mathbf{p}}, \quad \text{where} \quad N = N_2 - N_1 + 1.$$

Then, with probability at least $1 - Ce^{-cR^2}$, we have

$$\sup_{t \in [0, T]} \max_{j \in \llbracket N_1 + K, N_2 - K \rrbracket} (|a_j(t) - a_j^{[N_1, N_2]}(t)| + |b_j(t) - b_j^{[N_1, N_2]}(t)|) \leq e^{-K/5}.$$

The system $(\mathbf{a}(t); \mathbf{b}(t))$ from Proposition 2.5 is the Toda lattice on \mathbb{Z} , under thermal equilibrium with parameters (β, θ) . Let us briefly interpret the third part of Proposition 2.5, when the interval length N satisfies $T \ll N \ll e^T$. It states that, with high probability (at least $1 - Ce^{-(\log T)^2}$), the Flaschka variables at site j of the Toda lattice on \mathbb{Z} approximately coincide with those of the open Toda lattice on $\llbracket N_1, N_2 \rrbracket$, up to an error that is exponentially small in T , as long as j is in the “bulk” of $\llbracket N_1, N_2 \rrbracket$ (that is, not too close to its endpoints N_1 and N_2 ; otherwise, boundary effects on the interval should become more visible and make this comparison invalid). Thus, asymptotic questions about the Toda lattice run for some large time T , on \mathbb{Z} under thermal equilibrium, can be recovered from those about the open Toda lattice on a finite interval at thermal equilibrium. Hence, we will throughout focus on the open Toda lattice on the finite interval $\llbracket N_1, N_2 \rrbracket$ for times $t \in [0, T]$ with $T \in [N^\delta, N^{1-\delta}]$ for some small constant $\delta > 0$ (though the statements below will permit more flexibility in T), and be interested in its variables at sites j in the bulk of $\llbracket N_1, N_2 \rrbracket$.

Before proceeding, let us mention that one might also be interested in analyzing Toda lattice on the torus $\mathbb{T}_N = \mathbb{Z}/N\mathbb{Z}$, for which thermal equilibrium is also invariant (see Section 3.1 below). It can be deduced from Proposition 4.3 below that a counterpart of Proposition 2.5 holds comparing this Toda lattice on \mathbb{T}_N to the open one on $\llbracket N_1, N_2 \rrbracket$, indicating they are likely very close to each other in the bulk of $\llbracket N_1, N_2 \rrbracket$, for times $T \ll N = N_2 - N_1 + 1$ (for much larger times, we expect the periodicity of the torus to become more visible, again making such a comparison invalid). As such, asymptotic questions about the Toda lattice on \mathbb{T}_N at thermal equilibrium, for times $T \in [N^\delta, N^{1-\delta}]$, also reduce to those about the model on the interval.

2.3. Localization Centers. As explained in Section 2.1.2, while the eigenvalues of the Lax matrix $\mathbf{L}(t)$ are conserved under the Toda dynamics, they are not local, as they depend on all entries of $\mathbf{L}(t)$. Still, we will see under thermal equilibrium that they are “approximately local,” in that they only depend on a few entries of $\mathbf{L}(t)$, up to a small error. These entries will correspond to those on which the associated eigenvectors of $\mathbf{L}(t)$ are mainly supported; we call them localization centers,

given (in a more general context) by the below definition. Here, we again let $N_1 \leq N_2$ be integers and set $N = N_2 - N_1 + 1$.

Definition 2.6 (Localization centers). Let $\mathbf{u} = (u(N_1), u(N_1 + 1), \dots, u(N_2)) \in \mathbb{R}^N$ be a unit vector. For any $\zeta \in \mathbb{R}_{\geq 0}$, we call an index $\varphi \in \llbracket N_1, N_2 \rrbracket$ a ζ -localization center for \mathbf{u} if $|u(\varphi)| \geq \zeta$.

Next, let $\mathbf{M} = [M_{ij}]$ be a symmetric $N \times N$ matrix, with entries indexed by $i, j \in \llbracket N_1, N_2 \rrbracket$. If $\lambda \in \text{eig } \mathbf{M}$, then we call $\varphi \in \llbracket N_1, N_2 \rrbracket$ a ζ -localization center for λ with respect to \mathbf{M} if φ is a ζ -localization center for some unit eigenvector $\mathbf{u} \in \mathbb{R}^N$ of \mathbf{M} with eigenvalue λ . Further let $(\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_N)$ denote an orthonormal eigenbasis of \mathbf{M} . We call a bijection $\varphi : \llbracket 1, N \rrbracket \rightarrow \llbracket N_1, N_2 \rrbracket$ a ζ -localization center bijection for \mathbf{M} if $\varphi(j)$ is a ζ -localization center for \mathbf{u}_j for each $j \in \llbracket 1, N \rrbracket$.

One might first ask whether ζ -localization center bijections exist. The following lemma, whose quick proof appears in Section 5.1 below, verifies this if $\zeta \leq (2N)^{-1}$.

Lemma 2.7. *Any symmetric $N \times N$ matrix \mathbf{M} admits a $(2N)^{-1}$ -localization center bijection.*

One might next ask if such localization centers bijections are unique; this is not always the case. An extreme example is if \mathbf{M} is the adjacency matrix for the discrete Laplacian on a torus of length N , in which case one can verify that any bijection from $\llbracket 1, N \rrbracket$ to $\llbracket N_1, N_2 \rrbracket$ is an $N^{-1/2}$ -localization center bijection.

However, we will show that this uniqueness is true “up to some error,” if \mathbf{M} is the Lax matrix for the Toda lattice run under thermal equilibrium. To make this precise, we require some notation, which will frequently be adopted throughout the remainder of this paper.

Assumption 2.8. Fix real numbers⁴ $\beta, \theta > 0$, and assume that the *stretch parameter*

$$(2.7) \quad \alpha = \log \beta - \frac{\Gamma'(\theta)}{\Gamma(\theta)} \neq 0.$$

For each $t \geq 0$, let $\mathbf{L}(t) = [L_{ij}(t)]$ denote the Lax matrix for the open Toda lattice $(\mathbf{a}(t); \mathbf{b}(t))$ on $\llbracket N_1, N_2 \rrbracket$ (as in Definition 2.1). Set $\text{eig } \mathbf{L}(t) = (\lambda_1, \lambda_2, \dots, \lambda_N)$, which does not depend on t by Lemma 2.2, and for each $k \in \llbracket 1, N \rrbracket$ let $\mathbf{u}_k(t) = (u_k(N_1; t), u_k(N_2; t), \dots, u_k(N_2; t))$ denote the nonnegatively normalized, unit eigenvector of $\mathbf{L}(t)$ with eigenvalue λ_k . At $t = 0$, abbreviate $\mathbf{L} = \mathbf{L}(0)$; $(\mathbf{a}; \mathbf{b}) = (\mathbf{a}(0); \mathbf{b}(0))$; and $\mathbf{u}_k = \mathbf{u}_k(0) = (u_k(N_1), u_k(N_1 + 1), \dots, u_k(N_2))$. Assume that the initial data $(\mathbf{a}; \mathbf{b})$ is sampled under the thermal equilibrium $\mu_{\beta, \theta; N-1, N}$ from Definition 2.4. Let $(\mathbf{p}(s); \mathbf{q}(s))$, over $s \in \mathbb{R}_{\geq 0}$, denote the Toda state space dynamics associated with (Flaschka variable) initial data $(\mathbf{a}(0); \mathbf{b}(0))$, as in Section 2.1.1.

Let $T \geq 1$ and $\zeta \geq 0$ be real numbers satisfying

$$(2.8) \quad N_1 \leq -N(\log N)^{-1} \leq N(\log N)^{-1} \leq N_2; \quad 1 \leq T \leq N(\log N)^{-7}; \quad \zeta \geq e^{-100(\log N)^{3/2}}.$$

For each $s \in \mathbb{R}$ let $\varphi_s : \llbracket 1, N \rrbracket \rightarrow \llbracket N_1, N_2 \rrbracket$ be a ζ -localization center bijection for $\mathbf{L}(s)$, and denote

$$(2.9) \quad Q_j(s) = q_{\varphi_s(j)}(s), \quad \text{for each } (j, s) \in \llbracket 1, N \rrbracket \times \mathbb{R}_{\geq 0}.$$

The reason for the term, “stretch parameter,” is given by Lemma 3.12 below, which indicates that the average spacing $\mathbb{E}[q_i(0) - q_{i+1}(0)] = \alpha$ between points in the Toda lattice at thermal equilibrium is α . The assumption (2.7) states that the stretch parameter α is nonzero, which implies that the Toda particles $(q_j(t))$ do not “accumulate” (with infinite density) around 0.

Let us briefly explain (2.8). Its first bound indicates that 0 is in the bulk of $\llbracket N_1, N_2 \rrbracket$. Its second ensures that the time scale T is sublinear in N , in accordance with the discussion at the end of

⁴Throughout this paper, constants may depend on β and θ , even when not explicitly stated.

Section 2.2 (as this will guarantee that the boundary of $[[N_1, N_2]]$ does not asymptotically affect its bulk under the Toda lattice). Its third ensures that ζ is not too small. In (2.8), the constants 7, 100, and $3/2$ are of little significance (we will need $e^{-(\log N)^2} \ll \zeta \ll N^{-C}$ for C sufficiently large).

Under Assumption 2.8, we view $\varphi_j(s)$ as the “location” of λ_j on the lattice $[[N_1, N_2]]$, and $Q_j(s)$ as the location of the corresponding Toda particle⁵ $q_{\varphi_s(j)}(s)$ on the line \mathbb{R} (the latter may be thought of as the location for the “soliton with amplitude λ_j ,” as described in Section 1).

The following proposition, to be shown in Section 7.1 below, indicates that the location $\varphi_s(j)$ is likely unique, up to a small error (as long as it is in the bulk of $[[N_1, N_2]]$).

Proposition 2.9. *Adopt Assumption 2.8. There exists a constant $c > 0$ such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. For any real number $t \in [0, T]$; eigenvalue $\lambda \in \text{eig } \mathbf{L}(t)$; and two ζ -localization centers $\varphi, \tilde{\varphi} \in [[N_1, N_2]]$ for λ with respect to $\mathbf{L}(t)$, satisfying*

$$(2.10) \quad N_1 + T(\log N)^3 \leq \varphi \leq N_2 - T(\log N)^3,$$

we have $|\varphi - \tilde{\varphi}| \leq (\log N)^3$.

As mentioned above, one may think of the “location” of λ_j on the lattice $[[N_1, N_2]]$ at time s as being $\varphi_j(s)$. One manifestation of this is through Corollary 5.5 (and Corollary 5.6) below, which essentially states the following. Let $\lambda \in \text{eig } \mathbf{L}$ have ζ -localization center φ . If one perturbs the entries of \mathbf{L} , whose indices are sufficiently far from φ , to form a tridiagonal matrix $\tilde{\mathbf{L}}$, then there exists an eigenvalue $\tilde{\lambda} \in \text{eig } \tilde{\mathbf{L}}$ that is close to λ (and whose $N^{-1}\zeta$ -localization center is close to φ).

We will not state that result more precisely in this section, but instead mention the following relevant consequence of it, to be proven in Section 7.3 below. It indicates that the (sums of) local charges and (integrated) currents from Definition 2.3 are well-approximated by local sums of λ_j^m , over j satisfying $Q_j(t) \approx q_i(t)$. Thus, to analyze asymptotics of the former, it suffices to understand the limiting dynamics of the particle locations $Q_j(t)$.

Proposition 2.10. *Adopt Assumption 2.8. For any integer $m \geq 0$, there exists a constant $c = c(m) > 0$ such that the following two statements hold with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. Below, we let $t \in [0, T]$ be any real number and $\mathcal{J} \subset \mathbb{R}$ be any interval satisfying*

$$(2.11) \quad \begin{aligned} \mathcal{J} &\subseteq [\alpha N_1 + (T + |N_1|^{1/2})(\log N)^5, \alpha N_2 - (T + N_2^{1/2})(\log N)^5], & \text{if } \alpha > 0; \\ \mathcal{J} &\subseteq [\alpha N_2 + (T + N_2^{1/2})(\log N)^5, \alpha N_1 - (T + |N_1|^{1/2})(\log N)^5], & \text{if } \alpha < 0. \end{aligned}$$

(1) We have

$$(2.12) \quad \left| \sum_{i:q_i(t) \in \mathcal{J}} \mathfrak{t}_i^{[m]}(t) - \sum_{j:Q_j(t) \in \mathcal{J}} \lambda_j^m \right| \leq (\log N)^{m+6}.$$

(2) For any $k \in \mathcal{J}$, we have

$$(2.13) \quad \left| \int_0^t \mathfrak{j}_k^{[m]}(s) ds + \sum_{j:Q_j(t) < k} \lambda_j^m - \sum_{j:Q_j(0) < k} \lambda_j^m \right| \leq (\log N)^{m+6}.$$

⁵One might ask why we impose that $\varphi_t : [[1, N]] \rightarrow [[N_1, N_2]]$ is a bijection, as opposed to any map such that $\varphi_t(j)$ is a ζ -localization center of $\mathbf{u}(j; t)$ for each $j \in [[1, N]]$. As suggested by Proposition 2.9 below (which addresses arbitrary localization centers, and not only localization center bijections), this would not be necessary for our results below to hold. Still, we adopt it since it has the aesthetic property of exhibiting a one-to-one correspondence between eigenvalues of the Lax matrix and particles in the Toda lattice.

Let us briefly comment on Proposition 2.10. First, the assumption (2.11) on \mathcal{J} implying that \mathcal{J} is bounded away from αN_1 and αN_2 indicates that⁶ this interval only contains particles q_i of index i in the bulk of $\llbracket N_1, N_2 \rrbracket$. Second, it is typical in the context of hydrodynamical limits to consider local charges on an interval \mathcal{J} whose length is of order T . Then, the sums (or integrals) on the left side of (2.12) (or (2.13)) will also be of order T , which can be taken to be much larger than $(\log N)^{m+6}$, so the error on the right side of (2.12) is negligible in comparison.

2.4. Asymptotic Scattering Relation. The following theorem, proven in Section 8.1 below, provides the asymptotic scattering relation that governs the dynamics of the $Q_k(t)$. In what follows, the condition (8.4) ensures that the “initial location” $\varphi_0(k)$ of λ_k is in the bulk of $\llbracket N_1, N_2 \rrbracket$. Observe that the error⁷ $(\log N)^{15}$ on the right side of (2.15) is quite small, negligible in comparison to any power of N . So, Theorem 2.11 indicates that the asymptotic scattering relation is quite close to an equality.

Theorem 2.11 (Asymptotic scattering relation). *Adopt Assumption 2.8. There exists a constant $c > 0$ such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. Let $k \in \llbracket 1, N \rrbracket$ satisfy*

$$(2.14) \quad N_1 + T(\log N)^6 \leq \varphi_0(k) \leq N_2 - T(\log N)^6.$$

Then, for each $t \in [0, T]$, we have

$$(2.15) \quad \left| \lambda_k t - Q_k(t) + Q_k(0) - 2 \operatorname{sgn}(\alpha) \sum_{i: Q_i(t) < Q_k(t)} \log |\lambda_k - \lambda_i| \right. \\ \left. + 2 \operatorname{sgn}(\alpha) \sum_{i: Q_i(0) < Q_k(0)} \log |\lambda_k - \lambda_i| \right| \leq (\log N)^{15}.$$

In Appendix B below, we will provide a heuristic (following the physics literature) for how Theorem 2.11 can be used to evaluate the limiting velocities of the λ_k , as predicted in [16, 45]. While we do not know how to mathematically justify this heuristic directly, its output will be established for sufficiently small θ in a sequel work [1], through a different method.

2.5. Notation. For any integer $n \geq 1$, denote $\mathbb{T}_n = \mathbb{Z}/n\mathbb{Z}$. For any point $z \in \mathbb{C}$ and set $\mathcal{A} \subseteq \mathbb{C}$, denote $\operatorname{dist}(z, \mathcal{A}) = \inf_{s \in \mathcal{A}} |z - s|$. Denote the complement of any event \mathbf{E} by \mathbf{E}^c . Denote the set of $n \times n$ real matrices by $\operatorname{Mat}_{n \times n}$. For any $\mathbf{M} \in \operatorname{Mat}_{n \times n}$, denote its transpose by \mathbf{M}^T . Denote the set of $n \times n$ real symmetric matrices by $\operatorname{SymMat}_{n \times n} = \{\mathbf{M} \in \operatorname{Mat}_{n \times n} : \mathbf{M} = \mathbf{M}^T\}$.

As in Definition 2.1, it will often be convenient to index the rows and columns of $n \times n$ matrices by index sets different from $\llbracket 1, n \rrbracket$. Given a nonempty index set $\mathcal{J} \subset \mathbb{Z}$ of size $n = |\mathcal{J}|$, let $\operatorname{Mat}_{\mathcal{J}}$ denote the set of $n \times n$ real matrices $\mathbf{M} = [M_{ij}]_{i,j \in \mathcal{J}} \in \operatorname{Mat}_{n \times n}$, whose rows and columns are indexed by \mathcal{J} ; also let $\operatorname{SymMat}_{\mathcal{J}} = \operatorname{Mat}_{\mathcal{J}} \cap \operatorname{SymMat}_{n \times n}$ denote the set of real symmetric matrices whose rows and columns are indexed by \mathcal{J} . Given any matrix $\mathbf{M} \in \operatorname{Mat}_{\mathcal{J}}$ and subset $\mathcal{K} \subseteq \mathcal{J}$, let $\mathbf{M}^{(\mathcal{K})} = [M_{ij}^{(\mathcal{K})}]_{i,j \in \mathcal{J}} \in \operatorname{Mat}_{\mathcal{J}}$ denote the matrix obtained from \mathbf{M} by setting all its entries in a row or column indexed by an element of \mathcal{K} to 0, namely, $M_{ij}^{(\mathcal{K})} = \mathbb{1}_{i \notin \mathcal{K}} \cdot \mathbb{1}_{i \notin \mathcal{K}} \cdot M_{ij}$.

Throughout, given some integer parameter $N \geq 1$ and event \mathbf{E}_N depending on N , we will often make statements of the following form. There exists a constant $c > 0$, independent of N , such that

⁶Indeed, since $q_0(0) = 0$ and the stretch parameter α denotes the expected value of $q_{i+1} - q_i$, the extreme particles in the Toda lattice should reside around $q_{N_1}(0) \approx \alpha N_1$ and $q_{N_2}(0) \approx \alpha N_2$.

⁷No effort was made to optimize the exponent 15, which could likely be improved.

$\mathbb{P}[\mathbf{E}_N^{\mathcal{C}}] \leq f(c, N)$ holds for an explicit function $f : \mathbb{R}_{>0} \times \mathbb{Z}_{>0} \rightarrow \mathbb{R}_{\geq 0}$, which is non-decreasing in c and satisfies $\lim_{N \rightarrow \infty} f(c, N) = 0$ and $\lim_{c \rightarrow 0} f(c, N) > 1$ (an example is $f(c, N) = c^{-1} e^{-c(\log N)^2}$). When proving such statements we will often implicitly (and without comment) assume that $N \geq N_0$ is sufficiently large. Indeed, suppose there exist $N_0 \geq 1$ and $c_0 > 0$ such that $\mathbb{P}[\mathbf{E}_N^{\mathcal{C}}] \leq f(c_0, N)$ holds whenever $N \geq N_0$. Since $\lim_{c \rightarrow 0} f(c, N) > 1$, there exists a constant $c_1 > 0$ such that for $N \leq N_0$ we have $f(c_1, N) \geq 1$, in which case $\mathbb{P}[\mathbf{E}_N^{\mathcal{C}}] \leq 1 \leq f(c_1, N)$ holds for $N \leq N_0$. Thus, taking $c = \min\{c_0, c_1\}$ guarantees that $\mathbb{P}[\mathbf{E}_N^{\mathcal{C}}] \leq f(c, N)$ holds for all $N \geq 1$.

2.6. Outline. The remainder of this paper is organized as follows. We begin in Section 3 with some miscellaneous facts about the Toda lattice and random matrices; the results there are mainly known (and those for which we lack a reference are proven in Appendix A). In Section 4 we provide comparison estimates for the Toda lattice on different domains, showing Proposition 2.5 in the process. In Section 5 we show various results (Corollary 5.5 and Corollary 5.6 below) indicating that eigenvalues of the Lax matrix under thermal equilibrium are approximately local, which will be used in Section 7 to prove the results about localization centers stated in Section 2.3. In Section 6 we provide an effective approximation for the rate of exponential decay for entries of random Lax matrix eigenvectors, which is used to establish the asymptotic scattering relation (Theorem 2.11) in Section 8.

3. MISCELLANEOUS PRELIMINARIES

3.1. Periodic Toda Lattice. In this section we recall the periodic Toda lattice on a torus (which will be parallel to Section 2.1.1). Throughout, we fix an integer $N \geq 1$ and a real number $\Upsilon \in \mathbb{R}$ (which will prescribe the torus size and a periodicity constraint, respectively). The state space of the Toda lattice on \mathbb{T}_N , also called the *periodic Toda lattice*, with parameter Υ is given by a pair of N -tuples of real numbers $(\mathbf{p}(t); \mathbf{q}(t)) \in \mathbb{R}^N \times \mathbb{R}^N$, where $\mathbf{p}(t) = (p_0(t), p_1(t), \dots, p_{N-1}(t))$ and $\mathbf{q}(t) = (q_0(t), q_1(t), \dots, q_{N-1}(t))$; both are indexed by a real number $t \geq 0$ called the time. Given any initial data $(\mathbf{p}(0); \mathbf{q}(0)) \in \mathbb{R}^N \times \mathbb{R}^N$, the joint evolution of $(\mathbf{p}(t); \mathbf{q}(t))$ for $t \geq 0$ is prescribed by the system of differential equations (2.1) for all $(j, t) \in \llbracket 0, N-1 \rrbracket \times \mathbb{R}_{\geq 0}$. Here, we have set

$$p_{i+N}(t) = p_i(t), \quad \text{and} \quad q_{i+N}(t) = q_i(t) + \Upsilon, \quad \text{for each } (i, t) \in \mathbb{Z} \times \mathbb{R}_{\geq 0},$$

which corresponds to the periodicity constraints of considering the Toda lattice on the torus \mathbb{T}_N . The system of differential equations (2.1) is equivalent to the Hamiltonian dynamics generated by the Hamiltonian \mathfrak{H} from (2.2) (where again we set $q_N = q_0 + \Upsilon$). The existence and uniqueness of solutions to (2.1) for all time $t \geq 0$, under arbitrary initial data $(\mathbf{p}; \mathbf{q}) \in \mathbb{R}^N \times \mathbb{R}^N$, is thus a consequence of the Picard–Lindelöf theorem; see [47, Theorem 12.6].

We moreover again define $\mathbf{r}(t) = (r_0(t), r_1(t), \dots, r_{N-1}(t))$, as well as the Flaschka variables $\mathbf{a}(t) = (a_0(t), a_1(t), \dots, a_{N-1}(t)) \in \mathbb{R}_{\geq 0}^N$ and $\mathbf{b}(t) = (b_0(t), b_1(t), \dots, b_{N-1}(t)) \in \mathbb{R}^N$, according to (2.3). Then, (2.1) is equivalent to the system (2.4), for each $(j, t) \in \mathbb{Z} \times \mathbb{R}_{\geq 0}$, where the periodicity constraints become

$$a_i(t) = a_{i+N}(t); \quad b_i(t) = b_{i+N}(t); \quad r_i(t) = r_{i+N}(t); \quad \Upsilon = \sum_{j=1}^N r_j(t),$$

for each $(i, t) \in \mathbb{Z} \times \mathbb{R}_{\geq 0}$ (where the fact that Υ is independent of t follows from the fact that $\partial_t r_j(t) = 2a_j(t)^{-1} \cdot \partial_t a_j(t)$, by dividing both sides of the first equation in (2.4) by $a_j(t)$ and then summing over $j \in \llbracket 0, N-1 \rrbracket$). If $0 \in \llbracket N_1, N_2 \rrbracket$, then we can prescribe Toda state space variables $(\mathbf{p}(t); \mathbf{q}(t))$ associated with these Flaschka variables $(\mathbf{a}(t); \mathbf{b}(t))$ as in Section 2.1.1.

Next we recall the Lax matrix associated with the periodic Toda lattice. In what follows, we adopt the notation above, letting $(\mathbf{a}(t); \mathbf{b}(t)) \in \mathbb{R}^N \times \mathbb{R}^N$ denote the periodic Toda lattice.

Definition 3.1 (Periodic Lax matrix). For any $t \in \mathbb{R}_{\geq 0}$, define the *Lax matrix* $\mathbf{L}(t) = [L_{ij}]_{i,j \in \mathcal{J}} = [L_{ij}(t)] \in \text{SymMat}_{\llbracket 0, N-1 \rrbracket}$ as follows, where $L_{ij} = L_{ij}(t)$. Set

$$L_{ii} = b_i(t), \quad \text{for each } i \in \llbracket 0, N-1 \rrbracket.$$

Set

$$L_{i,i+1} = L_{i+1,i} = a_i(t), \quad \text{for each } i \in \llbracket 0, N-2 \rrbracket; \quad L_{0,N-1} = L_{N-1,0} = a_{N-1}(t).$$

Also set $L_{ij} = 0$ if (i, j) with $i \neq j$ is not of the above form.

As in Lemma 2.2 (for the open model), the eigenvalues of the Lax matrix are preserved under the periodic Toda dynamics (2.4). This was originally due to [19]; see also [47, Theorem 12.5].

Lemma 3.2 ([19, 47]). *For any real numbers $t, t' \in \mathbb{R}$, we have $\text{eig } \mathbf{L}(t) = \text{eig } \mathbf{L}(t')$.*

Next we define thermal equilibrium for the periodic Toda lattice (similarly to Definition 2.4).

Definition 3.3 (Periodic thermal equilibrium). Fix $\beta, \theta \in \mathbb{R}_{>0}$. The (*periodic*) *thermal equilibrium* with parameters $(\beta, \theta; N)$ is the product measure $\mu = \mu_{\beta, \theta} = \mu_{\beta, \theta; N}$ on $\mathbb{R}^N \times \mathbb{R}^N$ defined by

$$\mu(d\mathbf{a}; d\mathbf{b}) = \left(\frac{2^{1/2} \beta^{\theta+1/2}}{\pi^{1/2} \Gamma(\theta)} \right)^N \cdot \prod_{j=0}^{N-1} a_j^{2\theta-1} e^{-\beta a_j^2} da_j \prod_{j=0}^{N-1} e^{-\beta b_j^2/2} db_j,$$

where $\mathbf{a} = (a_0, a_1, \dots, a_{N-1}) \in \mathbb{R}_{\geq 0}^N$ and $\mathbf{b} = (b_0, b_1, \dots, b_{N-1}) \in \mathbb{R}^N$.

We now recall the notion of invariance for the periodic Toda lattice. To that end, fix an integer $N \geq 1$; let μ denote a probability measure on $\mathbb{R}_{\geq 0}^N \times \mathbb{R}^N$; and sample $(\mathbf{a}; \mathbf{b}) \in \mathbb{R}_{\geq 0}^N \times \mathbb{R}^N$ under μ . Consider the periodic Toda lattice in the Flaschka variables (2.4) on \mathbb{T}_N , denoted by $(\mathbf{a}(t); \mathbf{b}(t))$, with initial data $(\mathbf{a}(0); \mathbf{b}(0)) = (\mathbf{a}; \mathbf{b})$. If the law of $(\mathbf{a}(t); \mathbf{b}(t))$ coincides with the law μ of $(\mathbf{a}; \mathbf{b})$ for each $t \geq 0$, then we say that the measure μ is *invariant* for the periodic Toda lattice.

The following (known) lemma states the invariance of the above measure for the periodic Toda lattice. Its proof is a quick consequence of the Liouville theorem, with the fact that the quantities

$$(3.1) \quad \sum_{j=0}^{N-1} \log a_j(t), \quad \text{and} \quad \sum_{j=0}^{N-1} (2a_j(t)^2 + b_j(t)^2),$$

are conserved under the Toda dynamics (2.4); see, for example, [23] for further details.

Lemma 3.4 ([23, Sections 2 and 3]). *For any integer $N \geq 1$ and real numbers $\beta, \theta \in \mathbb{R}_{>0}$, the thermal equilibrium $\mu_{\beta, \theta; N}$ is invariant for the periodic Toda lattice on \mathbb{T}_N .*

3.2. Lax Matrix Eigenvector Evolution. Before proceeding, we state the following consequence of Lemma 2.2, to be proven in Appendix A.2 below. It indicates that, over time $t \in \mathbb{R}_{\geq 0}$, the maximum magnitude of an entry of the Lax matrix $\mathbf{L}(t)$ can change by at most a constant factor.

Lemma 3.5. *Adopt the notation of either Definition 2.1 or Definition 3.1, and let $\mathcal{J} = \llbracket N_1, N_2 \rrbracket$ in the first (open) case and $\mathcal{J} = \llbracket 0, N-1 \rrbracket$ in the second (periodic) one. Denote*

$$A(t) = \max_{i \in \mathcal{J}} |a_i(t)|; \quad B(t) = \max_{i \in \mathcal{J}} |b_i(t)|.$$

Then, for any $t, t' \in \mathbb{R}$, we have $A(t) + B(t) \leq 6(A(t') + B(t'))$.

We next discuss how the eigenvectors of Lax matrices evolve under the open Toda lattice. This discussion will only apply to the open case, so for the remainder of this section we restrict to that case, meaning we consider the Toda lattice on $\llbracket N_1, N_2 \rrbracket$. Then $\mathbf{L}(t)$ is for any $t \in \mathbb{R}_{\geq 0}$ a real, symmetric, tridiagonal matrix with nonzero off-diagonal entries; as such, its eigenvalues are mutually distinct (see [15, Proposition 2.40(a)]).

Denote $\text{eig } \mathbf{L}(t) = (\lambda_1, \lambda_2, \dots, \lambda_N)$, which is independent of t by Lemma 2.2. For any $t \in \mathbb{R}_{\geq 0}$ and $j \in \llbracket 1, N \rrbracket$, let $\mathbf{u}_j(t) = (u_j(N_1; t), u_j(N_1 + 1; t), \dots, u_j(N_2; t)) \in \mathbb{R}^N$ denote the nonnegatively normalized, unit eigenvector of $\mathbf{L}(t)$ with eigenvalue λ_j (so that $\mathbf{L}(t) \cdot \mathbf{u}_j(t) = \lambda_j \cdot \mathbf{u}_j(t)$). The following result, due to [39], identifies the evolution of the first entry of these eigenvectors under the Toda dynamics (indicating that their logarithms evolve linearly, upon suitable normalization).

Lemma 3.6 ([39, Section 3]). *For any $t \in \mathbb{R}_{\geq 0}$ and $k \in \llbracket 1, N \rrbracket$, we have*

$$u_k(N_1; t)^2 = e^{-\lambda_k t} u_k(N_1; 0)^2 \cdot \left(\sum_{j=1}^N e^{-\lambda_j t} u_j(N_1; 0)^2 \right)^{-1}.$$

3.3. Resolvents. In this section we recall properties of resolvents. Throughout, we let $\mathcal{J} \subset \mathbb{Z}$ denote a nonempty, finite set of indices, and set $n = |\mathcal{J}|$. Let $\mathbf{M} = [M_{ij}] \in \text{Mat}_{\mathcal{J}}$. Given $z \in \mathbb{C}$, the associated *resolvent* of \mathbf{M} is $\mathbf{G}(z) = (\mathbf{M} - z)^{-1} = [G_{ij}(z)] = [G_{i,j}(z)]$, assuming that this inverse exists. Denote $\text{eig } \mathbf{M} = (\lambda_1, \lambda_2, \dots, \lambda_n)$, and let $(\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n)$ denote an orthonormal family of eigenvectors of \mathbf{M} , so that $\mathbf{u}_k = (u_k(i))_{i \in \mathcal{J}}$ is an eigenvector of \mathbf{M} with eigenvalue λ_k . Then, for any $i, j \in \mathcal{J}$, observe that

$$(3.2) \quad G_{ij}(z) = \sum_{k=1}^n \frac{u_k(i)u_k(j)}{\lambda_k - z},$$

which by the orthonormality of the (\mathbf{u}_k) implies for $\text{Im } z \neq 0$ that

$$(3.3) \quad |G_{ij}(z)| \leq |\text{Im } z|^{-1}.$$

Further observe, for any invertible matrices $\mathbf{A}, \mathbf{B} \in \text{Mat}_{n \times n}$, that we have

$$(3.4) \quad \mathbf{A}^{-1} - \mathbf{B}^{-1} = \mathbf{A}^{-1}(\mathbf{B} - \mathbf{A})\mathbf{B}^{-1} = \mathbf{B}^{-1}(\mathbf{B} - \mathbf{A})\mathbf{A}^{-1}.$$

The following lemma indicates that, if the resolvents of two matrices are close, then their eigenvalues are as well. Its proof will appear in Appendix A.2 below.

Lemma 3.7. *Fix an index $\varphi \in \mathcal{J}$; real numbers $\eta, \zeta, \delta > 0$; and symmetric $n \times n$ matrices $\mathbf{A}, \mathbf{B} \in \text{SymMat}_{\mathcal{J}}$. Fix an eigenvalue $\lambda \in \text{eig } \mathbf{A}$, with an associated unit eigenvector $\mathbf{u} = (u(i))_{i \in \mathcal{J}} \in \mathbb{R}^n$; denote $z = \lambda + i\eta$; and set $\mathbf{G} = [G_{ij}] = (\mathbf{A} - z)^{-1}$ and $\mathbf{H} = [H_{ij}] = (\mathbf{B} - z)^{-1}$. Assume that*

$$(3.5) \quad \delta \leq (2\eta)^{-1}\zeta^2; \quad |u(\varphi)| \geq \zeta; \quad |G_{\varphi\varphi}(z) - H_{\varphi\varphi}(z)| \leq \delta.$$

Then there exists an eigenvalue $\mu \in \text{eig } \mathbf{B}$, with an associated unit eigenvector $\mathbf{v} = (v(i))_{i \in \mathcal{J}} \in \mathbb{R}^n$, such that $|\lambda - \mu| \leq 3n\zeta^{-2}\eta$ and $|v(\varphi)| \geq (6n)^{-1/2}\zeta$.

3.4. Eigenvectors of Tridiagonal Matrices. In this section we provide (mostly known) facts about the eigenvectors of tridiagonal matrices. Throughout, we let $\mathbf{M} = [M_{ij}] \in \text{Mat}_{\llbracket N_1, N_2 \rrbracket}$ denote a tridiagonal, real symmetric $N \times N$ matrix, where $N = N_2 - N_1 + 1$; we assume that $M_{i, i+1} \neq 0$ for each $i \in \llbracket N_1, N_2 - 1 \rrbracket$. The below lemma expresses certain eigenvector entries of \mathbf{M} in terms of the matrix entries of \mathbf{M} ; it was originally due to [48], but we provide its proof in Appendix A.3 below.

Lemma 3.8 ([48, Equation (4)]). *Let $\mu \in \text{eig } \mathbf{M}$, and let $\mathbf{u} = (u_{N_1}, u_{N_1+1}, \dots, u_{N_2}) \in \mathbb{R}^N$ denote a unit eigenvector of \mathbf{M} with eigenvalue μ . Then,*

$$\log |u_{N_1}| + \log |u_{N_2}| = \sum_{i=N_1}^{N_2-1} \log |M_{i,i+1}| - \sum_{\mu' \in \text{eig } \mathbf{M} \setminus \{\mu\}} \log |\mu - \mu'|.$$

Next we discuss more precise behavior for the eigenvectors of \mathbf{M} through the method of transfer matrices, following [3, Section 4]. For any integer $k \in \llbracket N_1, N_2 - 1 \rrbracket$ and real number $E \in \mathbb{R}$, define the *transfer matrix* $\mathbf{S}_k(E) = \mathbf{S}_k(E; \mathbf{M}) \in \text{Mat}_{2 \times 2}$ by setting

$$(3.6) \quad \mathbf{S}_k(E) = \begin{bmatrix} 0 & 1 \\ -M_{k,k+1}^{-1} M_{k-1,k} & M_{k,k+1}^{-1} (E - M_{k,k}) \end{bmatrix}.$$

Moreover, for any subset $\mathcal{K} = (k_1, k_2, \dots, k_m) \subseteq \llbracket N_1, N_2 - 1 \rrbracket$ with $k_1 < k_2 < \dots < k_m$, define $\mathbf{S}_{\mathcal{K}}(E) = \mathbf{S}_{\mathcal{K}}(E; \mathbf{M}) \in \text{Mat}_{2 \times 2}$ by setting

$$(3.7) \quad \mathbf{S}_{\mathcal{K}}(E) = \mathbf{S}_{k_m}(E) \cdot \mathbf{S}_{k_{m-1}}(E) \cdots \mathbf{S}_{k_1}(E).$$

The following (standard) lemma indicates how the above transfer matrices can be used to evaluate eigenvector entries of \mathbf{M} . Its proof is given in Appendix A.3 below.

Lemma 3.9. *Let $\mu \in \text{eig } \mathbf{M}$, and let $\mathbf{u} = (u_{N_1}, u_{N_1+1}, \dots, u_{N_2}) \in \mathbb{R}^N$ denote an eigenvector of \mathbf{M} with eigenvalue μ . For each $k \in \llbracket N_1, N_2 \rrbracket$, set $\mathbf{w}_k = (u_{k-1}, u_k) \in \mathbb{R}^2$, where $u_{N_1-1} = 0$. Then $\mathbf{S}_{\llbracket i, j \rrbracket}(\mu) \cdot \mathbf{w}_i = \mathbf{w}_{j+1}$ for any $i, j \in \llbracket N_1, N_2 - 1 \rrbracket$ with $i \leq j$.*

The next lemma expresses entries of $\mathbf{S}_{\llbracket i, j \rrbracket}(E)$ through eigenvalues of truncations of \mathbf{M} ; its proof appears in [47], but we also provide it in Appendix A.3 below. In what follows, for any integers $k, \ell \in \llbracket N_1, N_2 \rrbracket$ with $k \leq \ell$, we let $\mathbf{M}^{[k, \ell]}$ denote the $(\ell - k + 1) \times (\ell - k + 1)$ matrix obtained by restricting \mathbf{M} to rows and columns indexed by $i, j \in \llbracket k, \ell \rrbracket$; stated equivalently, for each $i, j \in \llbracket k, \ell \rrbracket$, the (i, j) -entry of $\mathbf{M}^{[k, \ell]}$ is equal to the (i, j) -entry of \mathbf{M} .

Lemma 3.10 ([47, Equation (1.65)]). *Set $\text{eig } \mathbf{M}^{[i, j]} = (\mu_1^{[i, j]}, \mu_2^{[i, j]}, \dots, \mu_{j-i+1}^{[i, j]})$ for any $i, j \in \llbracket N_1, N_2 \rrbracket$ with $i \leq j$. Then, for any $i, j \in \llbracket N_1, N_2 - 1 \rrbracket$, denoting $\ell = j - i + 1$ we have*

$$\mathbf{S}_{\llbracket i, j \rrbracket}(E) = \begin{bmatrix} -M_{i-1, i} \prod_{k=i}^{j-1} M_{k, k+1}^{-1} \prod_{h=1}^{\ell-2} (E - \mu_h^{[i+1, j-1]}) & \prod_{k=i}^{j-1} M_{k, k+1}^{-1} \prod_{h=1}^{\ell-1} (E - \mu_h^{[i, j-1]}) \\ -M_{i-1, i} \prod_{k=i}^j M_{k, k+1}^{-1} \prod_{h=1}^{\ell-1} (E - \mu_h^{[i+1, j]}) & \prod_{k=i}^j M_{k, k+1}^{-1} \prod_{h=1}^{\ell} (E - \mu_h^{[i, j]}) \end{bmatrix}.$$

3.5. Spectral Behavior of Random Lax Matrices. In this section we describe spectral properties of Lax matrices whose Flaschka variables are sampled from thermal equilibrium. In what follows, we fix real numbers $\beta, \theta > 0$; the constants below might depend on them, even if not stated explicitly.

The below two lemmas indicate that the stretch parameter α from Assumption 2.8 prescribe the average distance between particles in the Toda lattice under thermal equilibrium. We establish them in Appendix A.1 below.

Lemma 3.11. *Let $\mathbf{a} > 0$ be a random variable with law $\mathbb{P}[\mathbf{a} \in (a, a + da)] = 2\beta\theta \cdot \Gamma(\theta)^{-1} \cdot a^{2\theta-1} e^{-\beta a^2} da$. Denoting $\mathbf{a} = e^{-\tau/2}$, we have that $\mathbb{E}[\tau] = \alpha$.*

Lemma 3.12. *Adopt Assumption 2.8. There exists a constant $c > 0$ such that the following holds. For any distinct indices $i, j \in \llbracket N_1, N_2 \rrbracket$ and real number $R \geq 1$, we have*

$$\mathbb{P}[|q_j(0) - q_i(0) - \alpha(j - i)| \geq R] \leq 2(e^{-cR^2/|i-j|} + e^{-cR}).$$

We next define events on which a symmetric matrix has bounded entries and eigenvalues, and on which its eigenvalues are separated.

Definition 3.13 (Bounded and separated events). Fix real numbers $A, \delta > 0$; let \mathcal{J} denote an index set; and let $\mathbf{M} = [M_{ij}] \in \text{SymMat}_{\mathcal{J}}$. Define the events

$$\text{BND}_{\mathbf{M}}(A) = \left\{ \max_{i,j \in \mathcal{J}} |M_{ij}| \leq A \right\} \cap \left\{ \max_{\lambda \in \text{eig } \mathbf{M}} |\lambda| \leq A \right\}; \quad \text{SEP}_{\mathbf{M}}(\delta) = \left\{ \min_{\substack{\nu, \nu' \in \text{eig } \mathbf{M} \\ \nu \neq \nu'}} |\nu - \nu'| \geq \delta \right\}.$$

Remark 3.14. Adopt the notation of Definition 3.13; let $\mathcal{J} \subseteq \mathcal{J}$ be some index set; and denote $\mathbf{M}' = \mathbf{M}^{(\mathcal{J})}$ (recall Section 2.5). By the Weyl interlacing inequality, we have $\text{BND}_{\mathbf{M}}(A) \subseteq \text{BND}_{\mathbf{M}'}(A)$.

If $\mathbf{L}(t)$ is the Lax matrix for a Toda lattice initialized under thermal equilibrium (recall Definition 2.4 and Definition 3.3), then the following lemma bounds its entries and eigenvalues with high probability. Its proof will be given in Appendix A.4 below.

Lemma 3.15. *There exists a constant $c > 0$ such that the following holds. Fix an integer $N \geq 1$; let $(\mathbf{a}(t); \mathbf{b}(t))$ denote the Flaschka variables for a Toda lattice (2.4), either in the open case on $\llbracket N_1, N_2 \rrbracket$ (as in Section 2.1.1) or in the periodic one on \mathbb{T}_N (as in Section 3.1). Denote the associated Lax matrix by $\mathbf{L}(t) = [L_{ij}(t)]$, as in Definition 2.1 for the open case (on $\llbracket N_1, N_2 \rrbracket$) and Definition 3.1 for the periodic one (on \mathbb{T}_N). Assume that $(\mathbf{a}(0); \mathbf{b}(0))$ is sampled under $\mu_{\beta, \theta; N-1, N}$ in the open case (on $\llbracket N_1, N_2 \rrbracket$), and under $\mu_{\beta, \theta; N}$ in the open case (on \mathbb{T}_N). Then, for any real number $A \geq 1$,*

$$\mathbb{P} \left[\bigcap_{t \in \mathbb{R}_{\geq 0}} \text{BND}_{\mathbf{L}(t)}(A) \right] \geq 1 - c^{-1} N e^{-cA^2}.$$

The following result indicates that the off-diagonal entries in the resolvent of a Lax matrix, of the open Toda lattice under the thermal equilibrium, decay exponentially. The bound (3.8) is a special case of [42, Theorem 4] (and the remark following it) when $z \in \mathbb{R}$ is real; together with [2, Equation (B.8)], this implies that it continues to hold for any complex $z \in \mathbb{C}$.

Lemma 3.16 ([42, 2]). *Adopt Assumption 2.8. For any real number $s \in (0, 1)$, there exists a constant $c = c(s) > 0$ such that the following holds. For any $z \in \mathbb{C}$, denote $\mathbf{G}(z) = [G_{ij}(z)] = (\mathbf{L} - z)^{-1}$. We have*

$$(3.8) \quad \sup_{z \in \mathbb{R}} \mathbb{E}[|G_{ij}(z)|^s] \leq c^{-1} e^{-c|i-j|}.$$

The below corollary, which is due to [2, Theorem A.1] (using Lemma 3.16 to verify its hypotheses), will quickly imply that eigenvectors of \mathbf{L} are localized (see Lemma 5.1 below).

Corollary 3.17 ([2, Theorem A.1]). *Adopt Assumption 2.8. There exists a constant $c > 0$ so that*

$$\max_{k \in \llbracket 1, N \rrbracket} \max_{i, j \in \llbracket N_1, N_2 \rrbracket} \mathbb{E}[|u_k(i)u_k(j)|] \leq c^{-1} e^{-c|i-j|}.$$

The following lemma states that, with high probability, no distinct eigenvalues of \mathbf{L} can be too close (recall Definition 3.13). It is a quick consequence of the Minami estimates [36]; we provide its proof in Appendix A.4 below.

Lemma 3.18. *Adopt Assumption 2.8. There exists a constant $c > 0$ such that $\mathbb{P}[\text{SEP}_{\mathbf{L}}(\delta)] \geq 1 - c^{-1}(\delta N^3 + e^{-cN^2})$ holds for any real number $\delta > 0$.*

4. COMPARISON ESTIMATES

4.1. Comparisons for the Toda Lattice on Different Domains. In this section we compare Toda lattices (through their Flaschka variables (2.3)) on different domains, that initially coincide on a subdomain. Such estimates are variants of the Lieb–Robinson estimates for quantum systems [31]; analogous bounds (in a slightly different form from what we use, but with similar proofs) have also appeared for classical ones, including the Toda lattice [30, 25].

We begin with the below proposition that compares two open Toda lattices on different intervals, assuming that they initially coincide on a subinterval of both.

Proposition 4.1. *Let $\tilde{N}_1 \leq N_1 \leq N_2 \leq \tilde{N}_2$ be integers; set $\tilde{N} = \tilde{N}_2 - \tilde{N}_1 + 1$ and $N = N_2 - N_1 + 1$. For each $t \in \mathbb{R}_{\geq 0}$, fix \tilde{N} -tuples $\tilde{\mathbf{a}}(t), \tilde{\mathbf{b}}(t) \in \mathbb{R}^{\tilde{N}}$ and N -tuples $\mathbf{a}(t), \mathbf{b}(t) \in \mathbb{R}^N$, indexed as*

$$\begin{aligned} \tilde{\mathbf{a}}(t) &= (\tilde{a}_{\tilde{N}_1}(t), \tilde{a}_{\tilde{N}_1+1}(t), \dots, \tilde{a}_{\tilde{N}_2}(t)); & \tilde{\mathbf{b}}(t) &= (\tilde{b}_{\tilde{N}_1}(t), \tilde{b}_{\tilde{N}_1+1}(t), \dots, \tilde{b}_{\tilde{N}_2}(t)); \\ \mathbf{a}(t) &= (a_{N_1}(t), a_{N_1+1}(t), \dots, a_{N_2}(t)); & \mathbf{b}(t) &= (b_{N_1}(t), b_{N_1+1}(t), \dots, b_{N_2}(t)). \end{aligned}$$

For each $s \in \mathbb{R}_{\geq 0}$, also set $\tilde{a}_i(s) = 0 = \tilde{b}_i(s)$ if $i \in \mathbb{Z} \setminus \llbracket \tilde{N}_1, \tilde{N}_2 - 1 \rrbracket$, and set $a_i(s) = 0 = b_i(s)$ if $i \in \mathbb{Z} \setminus \llbracket N_1, N_2 - 1 \rrbracket$. Assume $(\tilde{\mathbf{a}}(t); \tilde{\mathbf{b}}(t))$ satisfies (2.4) for each $(j, t) \in \llbracket \tilde{N}_1, \tilde{N}_2 \rrbracket \times \mathbb{R}_{\geq 0}$, and $(\mathbf{a}(t), \mathbf{b}(t))$ satisfies (2.4) for each $(j, t) \in \llbracket N_1, N_2 \rrbracket \times \mathbb{R}$. For any integers $I \leq J$ and real number $t \geq 0$, let

$$(4.1) \quad \begin{aligned} G_{\llbracket I, J \rrbracket}(t) &= \sup_{s \in [0, t]} \left(\max_{i \in \llbracket I, J \rrbracket} |a_i(s) - \tilde{a}_i(s)| + \max_{i \in \llbracket I, J \rrbracket} |b_i(s) - \tilde{b}_i(s)| \right); \\ H_{\llbracket I, J \rrbracket}(t) &= 6 \cdot \sup_{s \in [0, t]} \left(\max_{i \in \llbracket I, J \rrbracket} |a_i(s)| + \max_{i \in \llbracket I, J \rrbracket} |\tilde{a}_i(s)| + \max_{i \in \llbracket I, J \rrbracket} |b_i(s)| + \max_{i \in \llbracket I, J \rrbracket} |\tilde{b}_i(s)| \right). \end{aligned}$$

Now let $K \geq 1$ and $N'_1 \leq N'_2$ be integers such that $N_1 \leq N'_1 \leq N'_2 \leq N_2$ and $N'_1 + K \leq N'_2 - K$. If $a_j(0) = \tilde{a}_j(0)$ and $b_j(0) = \tilde{b}_j(0)$ for each $j \in \llbracket N'_1, N'_2 \rrbracket$, then for any $T \in \mathbb{R}_{\geq 0}$ we have

$$(4.2) \quad G_{\llbracket N'_1+K, N'_2-K \rrbracket}(T) \leq \frac{T^K}{K!} \cdot G_{\llbracket N'_1, N'_2 \rrbracket}(T) \cdot \prod_{i=0}^{K-1} H_{\llbracket N'_1+i, N'_2-i \rrbracket}(T).$$

To establish this proposition, we first require the following variant of the Grönwall inequality.

Lemma 4.2. *Let $T \geq 0$ be a real number; $K \geq 1$ be an integer; and let $g_k, h_k : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ be nondecreasing functions, for each $k \in \llbracket 0, K \rrbracket$. Assume for each $(k, t) \in \llbracket 0, K - 1 \rrbracket \times [0, T]$ that*

$$(4.3) \quad g_k(t) \leq h_{k+1}(t) \int_0^t g_{k+1}(s) ds.$$

Then, for any $(j, t) \in \llbracket 0, K \rrbracket \times [0, T]$, we have

$$(4.4) \quad g_j(t) \leq \frac{t^{K-j}}{(K-j)!} \cdot g_K(t) \cdot \prod_{i=j+1}^K h_i(t).$$

Proof. We verify (4.4) by induction on $K-j$. Since (4.4) holds at $K-j=0$, let us fix $j_0 \in \llbracket 0, K-1 \rrbracket$ and prove that (4.4) holds for $K-j = K-j_0$, assuming it holds whenever $K-j \leq K-j_0-1$. This follows from the estimates

$$\begin{aligned} g_{j_0}(t) &\leq h_{j_0+1}(t) \int_0^t g_{j_0+1}(s) ds \leq h_{j_0+1}(t) \int_0^t \frac{s^{K-j_0-1}}{(K-j_0-1)!} \cdot g_K(s) \cdot \prod_{i=j_0+2}^K h_i(s) ds \\ &\leq g_K(t) \cdot \prod_{i=j_0+1}^K h_i(t) \cdot \int_0^t \frac{s^{K-j_0-1}}{(K-j_0-1)!} ds \\ &= \frac{t^{K-j_0}}{(K-j_0)!} \cdot g_K(t) \cdot \prod_{i=j_0+1}^K h_i(t), \end{aligned}$$

where the first bound holds by the $k=j_0$ case of (4.3); the second by the inductive hypothesis; the third by the fact that the g_i and h_i are nondecreasing; and the fourth by performing the integration. \square

Proof of Proposition 4.1. Observe from (2.4) that, for any $(i, s) \in \llbracket \tilde{N}_1, \tilde{N}_2 \rrbracket \times \mathbb{R}_{\geq 0}$, we have

$$\begin{aligned} \partial_t(a_i - \tilde{a}_i) &= \frac{1}{2} \cdot (b_i - b_{i+1})(a_i - \tilde{a}_i) + \frac{\tilde{a}_i}{2} \cdot ((b_i - \tilde{b}_i) - (b_{i+1} - \tilde{b}_{i+1})); \\ \partial_t(b_i - \tilde{b}_i) &= (a_{i-1} - \tilde{a}_{i-1})(a_{i-1} + \tilde{a}_{i-1}) - (a_i - \tilde{a}_i)(a_i + \tilde{a}_i), \end{aligned}$$

where we have abbreviated $(\tilde{a}_i, a_i; \tilde{b}_i, b_i) = (\tilde{a}_i(s), a_i(s); \tilde{b}_i(s), b_i(s))$. Using (4.1), it follows that

$$\begin{aligned} |\partial_t a_i(s) - \partial_t \tilde{a}_i(s)| &= \frac{1}{2} \cdot |b_i - b_{i+1}| \cdot G_i(s) + |\tilde{a}_i| \cdot G_{\llbracket i, i+1 \rrbracket}(s) \leq \frac{1}{3} \cdot H_{\llbracket i, i+1 \rrbracket}(s) \cdot G_{\llbracket i, i+1 \rrbracket}(s); \\ |\partial_t b_i(s) - \partial_t \tilde{b}_i(s)| &= |a_{i-1} - \tilde{a}_{i-1}| \cdot G_{i-1}(s) + |a_i + \tilde{a}_i| \cdot G_i(s) \leq \frac{2}{3} \cdot H_{\llbracket i-1, i \rrbracket}(s) \cdot G_{\llbracket i-1, i \rrbracket}(s). \end{aligned}$$

Summing these two bounds, fixing $t_0 \in \mathbb{R}_{\geq 0}$, integrating over $s \in [0, t_0]$, and using (4.1) yields

$$|a_i(t_0) - \tilde{a}_i(t_0)| + |b_i(t_0) - \tilde{b}_i(t_0)| \leq H_{\llbracket i-1, i+1 \rrbracket}(t_0) \int_0^{t_0} G_{\llbracket i-1, i+1 \rrbracket}(s) ds,$$

for any $i \in \llbracket N'_1, N'_2 \rrbracket$, where we have also used that $a_i(0) = \tilde{a}_i(0)$ and $b_i(0) = \tilde{b}_i(0)$ for such i . Taking the supremum over $t_0 \in [0, t]$, we deduce that (4.3) holds for any $(k, t) \in \llbracket 0, K-1 \rrbracket \times \mathbb{R}_{\geq 0}$, if we set

$$g_k(t) = G_{\llbracket N'_1+K-k, N'_2-K+k \rrbracket}(t); \quad h_k(t) = H_{\llbracket N'_1+K-k, N'_2-K+k \rrbracket}(t).$$

The lemma then follows from the $(j, t) = (0, T)$ case of Lemma 4.2. \square

The next proposition compares a Toda lattice $(\mathbf{a}(t); \mathbf{b}(t))$ on a torus to one $(\tilde{\mathbf{a}}(t), \tilde{\mathbf{b}}(t))$ on an interval (of the same size), if the two coincide on a subinterval of their domains. Its proof is entirely analogous to that of Proposition 4.1 and is therefore omitted.

Proposition 4.3. *Let $N_1 \leq N_2$ be integers; set $N = N_2 - N_1 + 1$. For each real number $t \in \mathbb{R}_{\geq 0}$, fix N -tuples $\tilde{\mathbf{a}}(t), \mathbf{a}(t), \tilde{\mathbf{b}}(t), \mathbf{b}(t) \in \mathbb{R}^N$, indexed as*

$$\begin{aligned} \tilde{\mathbf{a}}(t) &= (\tilde{a}_{N_1}(t), \tilde{a}_{N_1+1}(t), \dots, \tilde{a}_{N_2}(t)); & \tilde{\mathbf{b}}(t) &= (\tilde{b}_{N_1}(t), \tilde{b}_{N_1+1}(t), \dots, \tilde{b}_{N_2}(t)); \\ \mathbf{a}(t) &= (a_0(t), a_1(t), \dots, a_{N-1}(t)); & \mathbf{b}(t) &= (b_0(t), b_1(t), \dots, b_{N-1}(t)). \end{aligned}$$

For each $s \in \mathbb{R}_{\geq 0}$, also set $\tilde{a}_i(s) = 0 = \tilde{b}_i(s)$ if $i \notin \llbracket \tilde{N}_1, \tilde{N}_2 - 1 \rrbracket$; additionally set $a_i(s) = a_{i+N}(s)$ and $b_i(s) = b_{i+N}(s)$ for each $(i, s) \in \mathbb{Z} \times \mathbb{R}_{\geq 0}$. Assume that $(\tilde{\mathbf{a}}(t); \tilde{\mathbf{b}}(t))$ satisfies (2.4) for each

$(j, t) \in \llbracket N_1, N_2 \rrbracket \times \mathbb{R}_{\geq 0}$ and that $(\mathbf{a}(t); \mathbf{b}(t))$ satisfies (2.4) for each $(j, t) \in \llbracket 0, N-1 \rrbracket \times \mathbb{R}_{\geq 0}$. Under this notation, for any $I \leq J$ and $t \in \mathbb{R}_{\geq 0}$, define $G_{\llbracket I, J \rrbracket}(t)$ and $H_{\llbracket I, J \rrbracket}(t)$ as in (4.1).

Now let $K \geq 1$ and $N'_1 \leq N'_2$ be integers such that $N_1 \leq N'_1 \leq N'_2 \leq N_2$ and $N'_1 + K \leq N'_2 - K$. If $a_j(0) = \tilde{a}_j(0)$ and $b_j(0) = \tilde{b}_j(0)$ for each $j \in \llbracket N'_1, N'_2 \rrbracket$, then (4.2) holds for any $T \in \mathbb{R}_{\geq 0}$.

4.2. Approximate Thermal Equilibrium for the Open Toda Lattice. Unlike for the periodic Toda lattice, thermal equilibrium $\mu_{\beta, \theta; N-1, N}$ (from Definition 2.4) is not an invariant measure for the open Toda lattice. In this section we establish the below proposition indicating that it, in a certain sense, still “approximately is,” if time scale is much shorter than the domain size. Its proof uses Proposition 4.3 to couple the periodic Toda lattice to the closed one.

Proposition 4.4. *There exists a constant $c > 0$ satisfying the below. Adopt Assumption 2.8, and fix $t \in [0, T]$. There exists a random matrix $\mathbf{M} = [M_{ij}] \in \text{SymMat}_{\llbracket N_1, N_2 \rrbracket}$, whose law coincides with that of $\mathbf{L}(0)$, such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. For any real number $K \geq T \log N$, we have that*

$$(4.5) \quad \max_{i, j \in \llbracket N_1 + K, N_2 - K \rrbracket} |L_{ij}(t) - M_{ij}| \leq e^{-K/5}.$$

To establish Proposition 4.4, we first require the following quick consequence of Proposition 4.1 and Proposition 4.3.

Lemma 4.5. *Adopt the notation and assumptions of either Proposition 4.1 or Proposition 4.3, and suppose that $T \geq 1$. Fix a real number $A \geq 1$, and assume that $K \geq 200AT$ and that $|a_i(0)| + |\tilde{a}_i(0)| + |b_i(0)| + |\tilde{b}_i(0)| \leq A$ for each $i \in \mathbb{Z}$. Then, $G_{\llbracket N'_1 + K, N'_2 - K \rrbracket}(T) \leq e^{-K/4}$.*

Proof. By Lemma 3.5, we have

$$(4.6) \quad G_{\llbracket N'_1, N'_2 \rrbracket}(T) \leq H_{\llbracket N'_1, N'_2 \rrbracket}(T) \leq 36 \cdot \max_{i \in \mathbb{Z}} (|a_i(0)| + |b_i(0)| + |\tilde{a}_i(0)| + |\tilde{b}_i(0)|) \leq 36A.$$

Thus,

$$G_{\llbracket N'_1 + K, N'_2 - K \rrbracket}(T) \leq \frac{T^K}{K!} (36A)^{K+1} \leq 36A \left(\frac{36eAT}{K} \right)^K \leq 36A \cdot 2^{-K} \leq 2^{-K/2} \leq e^{-K/4}.$$

where in the first inequality we used Proposition 4.1 with (4.6); in the second we used the fact that $K! \geq (e^{-1}K)^K$; in the third we used the fact that $36eATK^{-1} \leq 100ATK^{-1} \leq 1/2$; in the fourth we used the fact that $2^{-K/2} \leq 2^{-100AT} \leq (36A)^{-1}$ (as $A, T \geq 1$); and in the fifth we used the fact that $e \leq 4$. This establishes the lemma. \square

Proof of Proposition 4.4. We will compare the open Lax matrix $\mathbf{L}(t)$ to a periodic one $\mathbf{R}(t)$. To prescribe initial data for the latter (through its Flaschka variables), define $(\mathbf{a}(0), \mathbf{b}(0))$, where $\mathbf{a}(0) = (a_j(0))$ and $\mathbf{b}(0) = (b_j(0))$ for $j \in \mathbb{Z}$, as follows. First, for each $i \in \llbracket N_1, N_2 - 1 \rrbracket$, set

$$\mathbf{a}_i(0) = L_{i, i+1}(0) = a_i(0), \quad \text{and} \quad \mathbf{b}_i(0) = L_{i, i}(0) = b_i(0).$$

Letting $\mathbf{r} \in \mathbb{R}_{>0}$ denote a Gamma random variable, independent from $\mathbf{L}(0)$, with density $\mathbb{P}[\mathbf{r} \in (r, r+dr)] = 2\beta^\theta \cdot \Gamma(\theta)^{-1} \cdot r^{2\theta-1} e^{-\beta r^2} dr$, further set $\mathbf{a}_{N_2} = \mathbf{r}$ and $\mathbf{b}_{N_2} = L_{N_2, N_2}(0) = b_{N_2}(0)$. We then extend these Flaschka variables periodically, by imposing $(\mathbf{a}_{j+N}(0); \mathbf{b}_{j+N}(0)) = (\mathbf{a}_j(0); \mathbf{b}_j(0))$ for each $j \in \mathbb{Z}$. In this way, $(\mathbf{a}_j(0); \mathbf{b}_j(0))$ over $j \in \llbracket N_1, N_2 \rrbracket$ is sampled under $\mu_{\beta, \theta; N}$ from Definition 3.3.

Next, let $(\mathbf{a}(t); \mathbf{b}(t))$, where $\mathbf{a}(t) = (a_j(t))$ and $\mathbf{b}(t) = (b_j(t))$, denote the solution to the periodic Toda lattice (2.4) on the torus \mathbb{T}_N , with initial data $(\mathbf{a}(0); \mathbf{b}(0))$. Here, (j, t) ranges over $\mathbb{Z} \times \mathbb{R}$,

imposing the periodicity constraint $(\mathbf{a}_{j+N}(t); \mathbf{b}_{j+N}(t)) = (\mathbf{a}_j(t); \mathbf{b}_j(t))$. It follows from Lemma 3.4 that, for each $t \in \mathbb{R}_{\geq 0}$,

$$(4.7) \quad \text{the law of } (\mathbf{a}(t); \mathbf{b}(t)) \text{ coincides with that of } (\mathbf{a}(0); \mathbf{b}(0)).$$

Define the associated Lax matrix⁸ $\mathbf{R}(t) = [R_{ij}(t)] \in \text{SymMat}_{\llbracket N_1, N_2 \rrbracket}$, by setting $R_{jj}(t) = \mathbf{b}_j(t)$ and $R_{j,j+1}(t) = R_{j+1,j}(t) = \mathbf{a}_j(t)$ for each $j \in \llbracket N_1, N_2 \rrbracket$, where we have denoted $R_{N_2+1, N_2}(t) = R_{N_1, N_2}(t)$ and $R_{N_2, N_2+1}(t) = R_{N_2, N_1}(t)$. If (i, j) is not of the above form, then we set $R_{ij}(t) = 0$.

Now define $\mathbf{M} = [M_{ij}] \in \text{SymMat}_{\llbracket N_1, N_2 \rrbracket}$ by setting the (N_1, N_2) -entry and (N_2, N_1) -entry of $\mathbf{R}(T)$ to 0, namely $M_{ij} = \mathbb{1}_{(i,j) \neq (N_1, N_2)} \cdot \mathbb{1}_{(i,j) \neq (N_2, N_1)} \cdot R_{ij}(T)$. By (4.7), the fact that $(\mathbf{a}(0); \mathbf{b}(0))$ is sampled under $\mu_{\beta, \theta; N}$; and Assumption 2.8, \mathbf{M} has the same law as $\mathbf{L}(0)$.

It therefore remains to confirm that (4.5) holds with high probability. To that end, recalling Definition 3.13, define the event $\mathbf{E}_1 = \bigcap_{s \geq 0} \text{BND}_{\mathbf{R}(s)}(\log N/800)$. By Lemma 3.15, there is a constant $c_1 > 0$ such that $\mathbb{P}[\mathbf{E}_1^c] \leq c_1^{-1} e^{-c_1(\log N)^2}$. So, we will restrict to \mathbf{E}_1 in what follows.

Now we apply Proposition 4.3 and Lemma 4.5. More specifically, since $L_{ij}(0) = R_{ij}(0)$ for $i, j \in \llbracket N_1 + 1, N_2 - 1 \rrbracket$, applying Lemma 4.5 (with the $(A; I, J; K)$ there equal to $(\log N/200; N_1 + 1, N_2 - 1; K - 1)$ here, using the fact that $(K - 1)/4 \leq K/5$ for sufficiently large N , and our restriction to \mathbf{E}_1), yields (4.5), since $M_{ij} = R_{ij}(t)$ for $i, j \in \llbracket N_1 + K, N_2 - K \rrbracket$. \square

4.3. Infinite Volume Limit. Recall that our previous descriptions of the Toda lattice involved a finite number of variables, and were thus defined on a finite domain. In this section we explain conditions under which solutions of the Toda lattice on infinite domains can be realized as limits of those on finite domains. These conditions are described through the below assumption.

Assumption 4.6. Fix real numbers $R > 1 > \mathfrak{p} \geq 0$. For each real number $t \in \mathbb{R}_{\geq 0}$ and integers $n \geq 1 \geq m$, let

$$\begin{aligned} \mathbf{a}^{[m,n]}(t) &= (a_j^{[m,n]}(t))_{j \in \llbracket m, n \rrbracket} \in \mathbb{R}^{m+n+1}; & \mathbf{a}^{(n)}(t) &= (a_j^{(n)}(t))_{j \in \llbracket -n, n \rrbracket} \in \mathbb{R}^{2n+1}; & \mathbf{a} &= (a_j)_{j \in \mathbb{Z}}; \\ \mathbf{b}^{[m,n]}(t) &= (b_j^{[m,n]}(t))_{j \in \llbracket m, n \rrbracket} \in \mathbb{R}^{m+n+1}; & \mathbf{b}^{(n)}(t) &= (b_j^{(n)}(t))_{j \in \llbracket -n, n \rrbracket} \in \mathbb{R}^{2n+1}; & \mathbf{b} &= (b_j)_{j \in \mathbb{Z}}, \end{aligned}$$

be $(m + n + 1)$ -tuples, $(2n + 1)$ -tuples, and infinite sequences of real numbers. Assume that $(\mathbf{a}^{[m,n]}(t); \mathbf{b}^{[m,n]}(t))$ is a solution to the open Toda lattice (2.4) on the interval $\llbracket m, n \rrbracket$, and that $(\mathbf{a}^{(n)}(t); \mathbf{b}^{(n)}(t))$ is a solution to the periodic Toda lattice (2.4) on the torus \mathbb{T}_{2n+1} , which we identify with $\llbracket -n, n \rrbracket$. Further assume for any integers $n \geq 1 \geq m$ and $j \in \mathbb{Z}$ that

$$(4.8) \quad |a_j^{[m,n]}(0)| + |a_j^{[m,n]}(0)| \leq R(|j| + 1)^{\mathfrak{p}}, \quad \text{and} \quad |a_j^{(n)}(0)| + |b_j^{(n)}(0)| \leq R(|j| + 1)^{\mathfrak{p}}.$$

Also assume $(a_j^{(n)}(0), b_j^{(n)}(0)) = (a_j, b_j)$ whenever $j \in \llbracket -n, n \rrbracket$; that $b_j^{[m,n]}(0) = b_j$ whenever $j \in \llbracket m, n \rrbracket$; and that $a_j^{[m,n]}(0) = a_j$ whenever $j \in \llbracket m, n - 1 \rrbracket$ (and $a_n^{[m,n]}(0) = 0$).

Proposition 4.7. *Adopt Assumption 4.6. Then, for each $(j, t) \in \mathbb{Z} \times \mathbb{R}_{\geq 0}$, there exist real numbers $a_j(t)$ and $b_j(t)$ such that*

$$(4.9) \quad \lim_{n \rightarrow \infty} a_j^{[-n,n]}(t) = a_j(t) = \lim_{n \rightarrow \infty} a_j^{(n)}(t); \quad \lim_{n \rightarrow \infty} b_j^{[-n,n]}(t) = b_j(t) = \lim_{n \rightarrow \infty} b_j^{(n)}(t).$$

Moreover, denoting $\mathbf{a}(t) = (a_j(t))_{j \in \mathbb{Z}}$ and $\mathbf{b}(t) = (b_j(t))_{j \in \mathbb{Z}}$, we have that $(\mathbf{a}(t); \mathbf{b}(t))$ solves the Toda lattice (2.4) for each $(j, t) \in \mathbb{Z} \times \mathbb{R}$.

⁸This convention is slightly different from the one in Definition 3.1 (where the rows and columns are indexed by $\llbracket 0, N - 1 \rrbracket$ instead of $\llbracket N_1, N_2 \rrbracket$); we use it for notational convenience.

Proof. Observe by (4.8) and Lemma 3.5 that, for any integers $n \geq 1 \geq m$ and real number $t \in \mathbb{R}_{\geq 0}$,

$$(4.10) \quad \sup_{j \in \llbracket -n, n \rrbracket} (|a_j^{[-n, n]}(t)| + |b_j^{[-n, n]}(t)| + |a_j^{(n)}(t)| + |b_j^{(n)}(t)|) \leq 12R(n+1)^{\mathfrak{p}}.$$

Now let $K \geq 1$ be an integer, and let $N \geq 1$ be an integer. By Proposition 4.3, with the last statement in Assumption 4.6 (and the fact that $72R(n+1)^{\mathfrak{p}} \leq 144Rn^{\mathfrak{p}}$ for $n \geq 1$), we have

$$(4.11) \quad |a_j^{(N)}(t) - a_j^{[-N, N]}(t)| + |b_j^{(N)}(t) - b_j^{[-N, N]}(t)| \leq \frac{t^K}{K!} \cdot (144RN^{\mathfrak{p}})^{K+1},$$

for any $(j, t) \in \mathbb{Z} \times \mathbb{R}$ with $|j| + K \leq N$. Thus, applying (4.11) at $(N, K) = (n, \lfloor n/2 \rfloor)$, we find

$$\lim_{n \rightarrow \infty} (|a_j^{(n)}(t) - a_j^{[-n, n]}(t)| + |b_j^{(n)}(t) - b_j^{[-n, n]}(t)|) \leq \lim_{n \rightarrow \infty} \frac{t^{\lfloor n/2 \rfloor}}{\lfloor n/2 \rfloor!} \cdot (144Rn^{\mathfrak{p}})^{n/2+1} = 0,$$

where in the last equality we used the fact that $\mathfrak{p} < 1$. Hence, it suffices to show that the first limits in both of the pairs of equalities in (4.9) hold.

To that end, fix an integer $J \geq 1$ and a real number $T \geq 0$. For $(j, t) \in \llbracket -J, J \rrbracket \times [0, T]$, we first bound $(a_j^{[-n, n]}(t), b_j^{[-n, n]}(t))$ independently of n . To do this, observe (as used to obtain (4.11)) by Proposition 4.1 with the last statement in Assumption 4.6 that, for any integers $N \geq N' \geq 1$,

$$(4.12) \quad |a_j^{[-N, N]}(t) - a_j^{[-N', N']}(t)| + |b_j^{[N, N]}(t) - b_j^{[-N', N']}(t)| \leq \frac{T^K}{K!} \cdot (144RN^{\mathfrak{p}})^{K+1},$$

for any $(j, t) \in \mathbb{Z} \times \mathbb{R}$ with $|j| + K \leq N'$. Now let $N_0 > N_1 > \dots > N_r$ be integers such that $N_0 = N$, such that $N_i/16 \leq N_{i+1} \leq N_i/4$ for each $i \in \llbracket 0, r-1 \rrbracket$, and such that $N_r \in [4J, 16J]$. Applying (4.12) with the $(N, N'; K)$ there equal to $(N_i, N_{i+1}; N_{i+1}/2)$ here, we find that there exists a constant $C_1 = C_1(\mathfrak{p}, R, T) > 1$ such that

$$|a_j^{[-N_i, N_i]}(t) - a_j^{[-N_{i+1}, N_{i+1}]}(t)| + |b_j^{[-N_i, N_i]}(t) - b_j^{[-N_{i+1}, N_{i+1}]}(t)| \leq \frac{T^K}{K!} \cdot (144RN_i^{\mathfrak{p}})^{K+1} \leq C_1 e^{-N_i},$$

where in the last bound we used the facts that $K = N_{i+1}/2 \geq N_i/32$ and $\mathfrak{p} < 1$. Hence, for any integer $m \in \llbracket 1, r \rrbracket$, we have

$$(4.13) \quad \begin{aligned} & |a_j^{[-N, N]}(t) - a_j^{[-N_m, N_m]}(t)| + |b_j^{[-N, N]}(t) - b_j^{[-N_m, N_m]}(t)| \\ & \leq \sum_{i=0}^{m-1} (|a_j^{[-N_i, N_i]}(t) - a_j^{[-N_{i+1}, N_{i+1}]}(t)| + |b_j^{[-N_i, N_i]}(t) - b_j^{[-N_{i+1}, N_{i+1}]}(t)|) \\ & \leq C_1 \sum_{i=0}^{m-1} e^{-N_i} \leq 2C_1 e^{-N_m}. \end{aligned}$$

Taking $m = r$ and using (4.10) (with the n there equal to $N_r \in [4J, 16J]$ here), it follows that

$$(4.14) \quad \max_{j \in \llbracket -J, J \rrbracket} \sup_{t \in [0, T]} (|a_j^{[-N, N]}(t)| + |b_j^{[-N, N]}(t)|) \leq 2C_1 e^{-N_r} + 12R(N_r + 1)^{\mathfrak{p}} \leq 250RJ + 2C_1.$$

By (4.13) and (4.12), $(a_j^{[-N, N]}(t))$ and $(b_j^{[-N, N]}(t))$ are Cauchy sequences over $N \in \mathbb{Z}_{\geq 0}$. Hence, these sequences admit unique limits $a_j(t)$ and $b_j(t)$, respectively, as N tends to ∞ . This establishes the first part of the proposition.

To establish the second, observe by (4.14) that there exists a constant $C_2 = C_2(\mathfrak{p}, R, J, T) > 1$ such that $|a_j(t)| + |b_j(t)| \leq C_2$ and $|a_j^{[-N, N]}(t)| + |b_j^{[-N, N]}(t)| \leq C_2$, for all $(j, t) \in \llbracket -J, J \rrbracket \times [0, T]$ and $N \in \mathbb{Z}_{\geq 1}$. Since $(\mathbf{a}^{[-N, N]}(t); \mathbf{b}^{[-N, N]}(t))$ satisfies (2.4), these estimates imply that the first two

t -derivatives of $a_j^{[-N,N]}(t)$ and $b_j^{[-N,N]}(t)$ are uniformly bounded, for all $(j, t) \in \llbracket -J, J \rrbracket \times [0, T]$. Thus, the t -derivatives of $a_j(t)$ and $b_j(t)$ are also uniformly bounded for all $(j, t) \in \llbracket -J, J \rrbracket \times [0, T]$. Together with the fact that $(\mathbf{a}^{[-N,N]}(t); \mathbf{b}^{[-N,N]}(t))$ satisfy (2.4), this implies that $(\mathbf{a}; \mathbf{b})$ satisfy (2.4), thereby confirming the second statement of the proposition. \square

The below corollary indicates that the Toda lattice on the full line \mathbb{Z} at thermal equilibrium can be defined by taking a limit of periodic Toda lattices at thermal equilibrium $\mu_{\beta, \theta; N}$ (recall Definition 3.3). Here, for any integer $N \geq 1$, we identify the torus \mathbb{T}_{2N+1} with the interval $\llbracket -N, N \rrbracket$ (instead of $\llbracket 0, 2N \rrbracket$), which will index the Flaschka variables of the associated Toda lattice.

Corollary 4.8. *Fix $\beta, \theta \in \mathbb{R}_{>0}$. Let $a_1, a_2, \dots, b_1, b_2, \dots$ be mutually independent random variables, so that (a_j, b_j) has law $\mu_{\beta, \theta; 1}$ for each $j \in \mathbb{Z}$. For each integer $N \geq 1$, let $(\mathbf{a}^{(N)}(t); \mathbf{b}^{(N)}(t))$ denote the periodic Toda lattice (2.4) on \mathbb{T}_{2N+1} , with initial data $(\mathbf{a}^{(N)}(0); \mathbf{b}^{(N)}(0))$ given by $\mathbf{a}^{(N)}(0) = (a_{-N}, a_{1-N}, \dots, a_N)$ and $\mathbf{b}^{(N)}(0) = (b_{-N}, b_{1-N}, \dots, b_N)$.*

- (1) *For each $(j, t) \in \mathbb{Z} \times \mathbb{R}_{\geq 0}$, the limits $\lim_{N \rightarrow \infty} a_j^{(N)}(t) = a_j(t)$ and $\lim_{N \rightarrow \infty} b_j^{(N)}(t) = b_j(t)$ exist almost surely; are finite; and solve (2.4).*
- (2) *Denoting $\mathbf{a}(t) = (a_j(t))_{j \in \mathbb{Z}}$ and $\mathbf{b}(t) = (b_j(t))_{j \in \mathbb{Z}}$, the law of $(\mathbf{a}(t); \mathbf{b}(t))$ coincides with that of $(\mathbf{a}(0); \mathbf{b}(0))$, for any $t \geq 0$.*

Proof. The second statement of the corollary follows from the first, together with Lemma 3.4. It therefore remains to show the first, to which end it suffices by Proposition 4.7 to verify Assumption 4.6, that is, to show that there almost surely exists a constant $R > 1$ such that $|a_j| + |b_j| \leq R(|j| + 1)^{1/2}$ for all $j \in \mathbb{Z}$. To that end, observe that there exists a constant $C > 1$ such that, for each $j \in \mathbb{Z}$, we have

$$(4.15) \quad \mathbb{P}[|a_j| + |b_j| > 2(|j| + 1)^{1/2}] \leq \mathbb{P}[a_j > (|j| + 1)^{1/2}] + \mathbb{P}[|b_j| > (|j| + 1)^{1/2}] \leq C|j|^{\theta+1} e^{-\beta|j|/2},$$

where in the last inequality we used the fact that (a_j, b_j) has law $\mu_{\beta, \theta; 1}$ given by Definition 3.3. Since the sum of the right side of (4.15) over $j \in \mathbb{Z}$ is finite, the Borel–Cantelli lemma implies the almost sure existence of a constant $R > 1$ such that $|a_j| + |b_j| \leq R(|j| + 1)^{1/2}$ for all $j \in \mathbb{Z}$. \square

We can now establish Corollary 4.8.

Proof of Proposition 2.5. This will follow from using Lemma 4.5 to compare the open Toda lattice $(\mathbf{a}^{[-N, -N]}(t); \mathbf{b}^{[-N, -N]}(t))$ to one on the torus \mathbb{T}_{2N+1} with the same initial data, and then applying Corollary 4.8 (and its proof) to confirm the large N limit of the latter. So let $(\check{\mathbf{a}}(t); \check{\mathbf{b}}(t))$, where $\check{\mathbf{a}}(t) = (\check{a}_{-N}(t), \check{a}_{1-N}(t), \dots, \check{a}_N(t))$ and $\check{\mathbf{b}}(t) = (\check{b}_{-N}(t), \check{b}_{1-N}(t), \dots, \check{b}_N(t))$, denote the periodic Toda lattice (2.4) on \mathbb{T}_{2N+1} , with initial data obtained by setting $\check{a}_j(0) = a_j$ and $\check{b}_j(0) = b_j$ for each $j \in \llbracket -N, N \rrbracket$. Denoting for each integer $k \geq 2$ the event

$$\mathbf{E}_k = \bigcap_{j=-k}^k \left\{ |a_j| + |b_j| \leq \frac{1}{1600} \cdot (R + \log(|k| + 1)) \right\}; \quad \mathbf{E} = \bigcap_{k=N}^{\infty} \mathbf{E}_k,$$

we have from the explicit densities of a_j and b_j that $\mathbb{P}[\mathbf{E}_k^c] \leq c_1^{-1} e^{-c_1(R + \log k)^2}$, for some constant $c_1 > 0$. Hence, a union bound yields $\mathbb{P}[\mathbf{E}^c] \leq c_2^{-1} e^{-c_2 R^2}$ for some constant $c_2 > 0$ (as $R \geq \log N$).

Restricting to \mathbf{E} , we then apply Lemma 4.5, with the (N_1, N_2, A, K) ; $(\mathbf{a}(t); \mathbf{b}(t))$; and $(\tilde{\mathbf{a}}(t); \tilde{\mathbf{b}}(t))$ there equal to $(-N, N, R/200, K)$; $(\check{\mathbf{a}}(t); \check{\mathbf{b}}(t))$; and $(\mathbf{a}(t); \mathbf{b}(t))$ here, respectively. This yields

$$(4.16) \quad \sup_{t \in [0, T]} \max_{j \in [K-N, N-K]} (|a_j^{[-N, N]}(t) - \check{a}_j(t)| + |b_j^{[-N, N]}(t) - \check{b}_j(t)|) \leq e^{-K/4}.$$

Letting N (and then K) tend to ∞ , the first two statements of the proposition therefore follow from Corollary 4.8 (with the $(\mathbf{a}(t); \mathbf{b}(t))$ there equal to $(\check{\mathbf{a}}(t); \check{\mathbf{b}}(t))$ here).

To confirm the third, we assume that $N_2 \geq -N_1$, as the case when $-N_1 > N_2$ is entirely analogous. We will first use (4.13) to estimate the difference between $(\mathbf{a}^{[-N_2, N_2]}(t); \mathbf{b}^{[-N_2, N_2]}(t))$ and $(\mathbf{a}(t); \mathbf{b}(t))$, and then use Lemma 4.5 to estimate the difference between $(\mathbf{a}^{[N_1, N_2]}(t); \mathbf{b}^{[N_1, N_2]}(t))$ and $(\mathbf{a}^{[-N_2, N_2]}(t); \mathbf{b}^{[-N_2, N_2]}(t))$. To implement the former, first observe that the assumption (4.8) holds in Proposition 4.7 (with the R there uniformly bounded for fixed $\mathfrak{p} \in (0, 1)$), since on \mathbf{E} we have $|a_j| \leq (R + \log j)/800 \leq (j^{\mathfrak{p}} + \log j)/800 \leq (|j| + 1)^{\mathfrak{p}}$ for sufficiently large j . Therefore, (4.13) applies, with the (N, N_m) there equal to (\mathfrak{N}, N_2) here for some $\mathfrak{N} \geq 4^m N_2 \geq 4^{m-1} N$. This yields for some constant⁹ $C_1 = C_1(\mathfrak{p}) > 0$ that

$$\sup_{t \in [0, T]} (|a_j^{[-\mathfrak{N}, \mathfrak{N}]}(t) - a_j^{[-N_2, N_2]}(t)| + |b_j^{[-\mathfrak{N}, \mathfrak{N}]}(t) - b_j^{[-N_2, N_2]}(t)|) \leq C_1 e^{-N_2}.$$

Letting m , and thus \mathfrak{N} , tend to ∞ (and using the fact that $N_2 \geq N/3$, as $N_2 \geq -N_1$), we deduce

$$(4.17) \quad \sup_{t \in [0, T]} (|a_j(t) - a_j^{[-N_2, N_2]}(t)| + |b_j(t) - b_j^{[-N_2, N_2]}(t)|) \leq C_1 e^{-N/3}.$$

We next apply Lemma 4.5, with the $(N_1, N_2; \tilde{N}_1, \tilde{N}_2; A, K)$; $(\mathbf{a}(t); \mathbf{b}(t))$; and $(\tilde{\mathbf{a}}(t); \tilde{\mathbf{b}}(t))$ there equal to $(N_1, N_2; -N_2, N_2; R/200, K)$; $(\mathbf{a}^{[-N_2, N_2]}(t); \mathbf{b}^{[-N_2, N_2]}(t))$; and $(\mathbf{a}^{[N_1, N_2]}(t); \mathbf{b}^{[N_1, N_2]}(t))$ here, respectively (using our restriction to \mathbf{E} , and the fact that $R \geq \log N$, to verify its hypotheses). As in the derivation of (4.16), this yields

$$\sup_{t \in [0, T]} \max_{j \in [N_1+K, N_2+K]} (|a^{[N_1, N_2]}(t) - a^{[-N_2, N_2]}(t)| + |a^{[N_1, N_2]}(t) - a^{[-N_2, N_2]}(t)|) \leq e^{-K/4}.$$

Together with (4.17) (and the fact that $K \leq N$), this yields the third part of the corollary. \square

5. LOCALIZATION CENTERS

Throughout this section, we fix $\beta, \theta \in \mathbb{R}_{>0}$; the constants below may depend on them, even if not explicitly stated. We further let $N_1 \leq N_2$ be integers and set $N = N_2 - N_1 + 1$.

5.1. Existence and Speed Bounds for Localization Centers. In this section we discuss localization centers (recall Definition 2.6) and their properties. We begin by confirming their existence, by proving Lemma 2.7.

Proof of Lemma 2.7. Let $(\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_N)$ denote an orthonormal eigenbasis for \mathbf{M} ; assume that the rows and columns of \mathbf{M} are indexed by $\llbracket N_1, N_2 \rrbracket$. For each index $j \in \llbracket 1, N \rrbracket$, denote $\mathbf{u}_j = (u_j(N_1), u_j(N_1 + 1), \dots, u_j(N_2))$ and define the set

$$\mathcal{I}_j = \{i \in \llbracket N_1, N_2 \rrbracket : |u_j(i)| \geq (2N)^{-1}\}.$$

⁹By our assumption $RT \leq K \leq N^{\mathfrak{p}}$, it is quickly verified using (4.12) that this constant is independent of T .

We must show that there exists a bijection $\varphi : \llbracket 1, N \rrbracket \rightarrow \llbracket N_1, N_2 \rrbracket$ such that $\varphi(j) \in \mathcal{I}_j$ for each $j \in \llbracket 1, N \rrbracket$. To do this, it suffices by Hall's theorem to show for any set $\mathcal{J} \subseteq \llbracket 1, N \rrbracket$ that $|\mathcal{I}_{\mathcal{J}}| \geq |\mathcal{J}|$, where $\mathcal{I}_{\mathcal{J}} = \bigcup_{j \in \mathcal{J}} \mathcal{I}_j$. If this statement were false for some subset $\mathcal{J} \subseteq \llbracket 1, N \rrbracket$, then we would have

$$\begin{aligned} |\mathcal{J}| - 1 \geq |\mathcal{I}_{\mathcal{J}}| &= \sum_{i \in \mathcal{I}_{\mathcal{J}}} \sum_{j=1}^N |u_j(i)|^2 \geq \sum_{j \in \mathcal{J}} \sum_{i \in \mathcal{I}_j} |u_j(i)|^2 \geq \sum_{j \in \mathcal{J}} \sum_{i=N_1}^{N_2} |u_j(i)|^2 - |\mathcal{J}|N(2N)^{-2} \\ &\geq |\mathcal{J}| - \frac{1}{4}, \end{aligned}$$

where the first bound follows from the assumption; the second from the orthonormality of the eigenbasis $(\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_N)$; the third from restricting to $j \in \mathcal{J}$; the fourth from using that $|u_j(i)| < (2N)^{-1}$ for $i \notin \mathcal{I}_j$; and the fifth from the fact that each \mathbf{u}_j is a unit vector. This is a contradiction, so a $(2N)^{-1}$ -localization center bijection for \mathbf{M} exists. \square

The next lemma concerns a random Lax matrix corresponding to the open Toda lattice, sampled under thermal equilibrium (recall Section 3.5). It first states that its eigenvectors are exponentially localized around their localization centers; it then states (as a quick consequence) that such localization centers are essentially unique, up to an error of $(\log N)^2/2$.

Lemma 5.1. *There exists a constant $c \in (0, 1)$ such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. Adopt Assumption 2.8, but assume more generally that $\zeta \geq e^{-200(\log N)^{3/2}}$. Fix any index $j \in \llbracket 1, N \rrbracket$, and let $\varphi_j \in \llbracket N_1, N_2 \rrbracket$ denote any ζ -localization center for \mathbf{u}_j .*

- (1) *For any $i \in \llbracket N_1, N_2 \rrbracket$ with $|i - \varphi_j| \geq (\log N)^2/2$, we have $|u_j(i)| \leq e^{-c|i - \varphi_j|}$.*
- (2) *If $i \in \llbracket N_2, N_2 \rrbracket$ satisfies $|i - \varphi_j| \geq (\log N)^2/2$, then i is not a ζ -localization center for \mathbf{u}_j .*

Proof. The first statement of the lemma implies the second (as $c^{-1}e^{-c(\log N)^2/2} < \zeta$ for sufficiently large N), so it suffices to verify the former. To that end, observe for some $c > 0$ that

$$\zeta^{-1} \cdot \mathbb{E}[|u_j(i)|] \leq \mathbb{E}[|u_j(i) \cdot u_j(\varphi_j)|] \leq c^{-1}Ne^{-c|i - \varphi_j|}.$$

Here, the first inequality holds by the definition of φ_j ; the second by summing Corollary 3.17 over $j \in \llbracket N_1, N_2 \rrbracket$ (and examining its (k, j) term equal to (j, φ_j) here). This, with a Markov estimate, a union bound, and the facts that $\zeta \geq e^{-200(\log N)^{3/2}}$ and $|i - \varphi_j| \geq (\log N)^2/2$, yields the first statement of the lemma. \square

The next lemma bounds the ‘‘speed’’ at which localization centers can move, for a Lax matrix initially at thermal equilibrium.

Lemma 5.2. *There exists a constant $c \in (0, 1)$ such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. Adopt Assumption 2.8, and assume more generally that $\zeta \geq e^{-200(\log N)^{3/2}}$. For each $j \in \llbracket 1, N \rrbracket$, let $\varphi_j \in \llbracket N_1, N_2 \rrbracket$ denote any ζ -localization center for $\mathbf{u}_j(0)$. Then, for each $(j, s) \in \llbracket 1, N \rrbracket \times [0, T]$, we have*

$$(5.1) \quad |u_j(m; s)| \leq e^{-c|m - \varphi_j|}, \quad \text{for any } m \in \llbracket N_1, N_2 \rrbracket \text{ with } |m - \varphi_j| \geq T(\log N)^2.$$

Moreover, m is not a localization center for $\mathbf{u}_j(s)$ whenever $|m - \varphi_j| \geq T(\log N)^2$.

Proof. The first statement (5.1) of the lemma implies the second, as $e^{-c|m - \varphi_j|} < \zeta$ for $|m - \varphi_j| \geq T(\log N)^2$ if N is sufficiently large; it therefore suffices to establish (5.1). Recalling Definition 3.13

and letting $c \in (0, 1)$ denote the constant c from Lemma 5.1, and define the event

$$(5.2) \quad \mathbf{E} = \bigcap_{s \geq 0} \text{BND}_{\mathbf{L}(s)}(\log N) \cap \bigcap_{i=N_1}^{N_2} \bigcap_{j=1}^N \left\{ \mathbb{1}_{|i-\varphi_j| \geq (\log N)^2/2} \cdot |u_j(i; 0)| \leq e^{-c|i-\varphi_j|} \right\}.$$

By Lemma 3.15 and Lemma 5.1, there exists a constant $c > 0$ such that $\mathbb{P}[\mathbf{E}^c] \leq e^{-1}e^{-c(\log N)^2}$. Therefore, we may restrict to \mathbf{E} for the remainder of this proof.

We next recall a fact concerning the evolution of the Lax matrix $\mathbf{L}(s)$. For each $s \in \mathbb{R}$, define the tridiagonal skew-symmetric matrix $\mathbf{P}(s) = [P_{ij}(s)] \in \text{Mat}_{[[N_1, N_2]]}$ as follows. For each $i \in [[N_1, N_2 - 1]]$, set $P_{i, i+1}(s) = a_i(s)/2$ and $P_{i+1, i}(s) = -a_i(s)/2$; for all $(i, j) \in [[N_1, N_2]]^2$ not of the above form, set $P_{i, j}(s) = 0$. For each $s \in \mathbb{R}$, further let $\mathbf{V}(s) = [V_{ij}(s)] \in \text{Mat}_{[[N_1, N_2]]}$, satisfying the ordinary differential equation $\partial_s \mathbf{V}(s) = \mathbf{P}(s) \cdot \mathbf{V}(s)$, with initial data $\mathbf{V}(0) = \text{Id}$; the existence of such a matrix $\mathbf{V}(s)$ follows from the Picard–Lindelöf theorem. For any $(i, j) \in [[N_1, N_2]]^2$, its (i, j) -entry is more explicitly given by

$$(5.3) \quad V_{ij}(s) = \text{Id} + \sum_{k=1}^{\infty} \sum_{i_1=N_1}^{N_2} \cdots \sum_{i_{k-1}=N_1}^{N_2} \int_0^s \cdots \int_0^s \mathbb{1}_{s_1 < s_2 < \cdots < s_k} \cdot \prod_{h=1}^k P_{i_{h-1}, i_h}(s_h) ds_h,$$

where we have denoted $(i_0, i_k) = (i, j)$. Then, by [39, Section 2], we have $\mathbf{V}(s)^{-1} \cdot \mathbf{L}(s) \cdot \mathbf{V}(s) = \mathbf{L}(0)$.

This implies that $\mathbf{L}(s) = \mathbf{V}(s) \cdot \mathbf{L}(0) \cdot \mathbf{V}(s)^{\text{T}}$, as $\mathbf{V}(s)$ is orthogonal (since $\mathbf{V}(0) = \text{Id}$, $\partial_s \mathbf{V}(s) = \mathbf{P}(s) \cdot \mathbf{V}(s)$, and $\mathbf{P}(s)$ is skew-symmetric). Hence, letting $\mathbf{U}(s) = [U_{ij}(s)] \in \text{Mat}_{N \times N}$ denote matrix of eigenvectors of $\mathbf{L}(s)$, whose (i, j) -entry is given by $U_{ij}(s) = u_j(i; s)$ for each $(i, j) \in [[N_1, N_2]] \times [[1, N]]$, we have $\mathbf{U}(s) = \mathbf{V}(s) \cdot \mathbf{U}(0)$. In particular,

$$(5.4) \quad u_j(i; s) = \sum_{k=N_1}^{N_2} V_{ik}(s) \cdot u_j(k; 0).$$

Now observe whenever $|i - j| \geq 20T \log N$ that

$$(5.5) \quad |V_{ij}(s)| \leq \sum_{k=|i-j|}^{\infty} \frac{s^k}{k!} \cdot (2 \log N)^k \leq \sum_{k=|i-j|}^{\infty} \left(\frac{2es \log N}{k} \right)^k \leq \sum_{k=|i-j|}^{\infty} e^{-k} \leq 2e^{-|i-j|}.$$

Here, in the first inequality we used (5.3), with the facts that each $|P_{ij}(s_h)| \leq \log N$ (as we have restricted to the event \mathbf{E} from (5.2)) and that $P_{ij} = 0$ whenever $|i - j| \neq 1$ (meaning that there are at most two choices for each i_h that gives rise to a nonzero summand in (5.3)); in the second we used the bound $k! \geq (e^{-1}k)^k$ for each $k \geq 0$; in the third we used the bound $2k^{-1}es \log N \leq e^{-1}$ for $k \geq |i - j| \geq 20T \log N$; and in the fourth we performed the sum. Hence, for $|m - \varphi_j| > T(\log N)^2$,

$$\begin{aligned} |u_j(m; s)| &\leq \sum_{k=N_1}^{N_2} \mathbb{1}_{|k-m| \geq |m-\varphi_j|/2} \cdot |V_{mk}(s)| + \sum_{k=N_1}^{N_2} \mathbb{1}_{|k-m| \leq |m-\varphi_j|/2} \cdot |u_j(k; 0)| \\ &\leq 2 \sum_{k=N_1}^{N_2} \mathbb{1}_{|k-m| \geq |m-\varphi_j|/2} \cdot e^{-|k-m|} + \sum_{k=N_1}^{N_2} \mathbb{1}_{|k-\varphi_j| \geq |m-\varphi_j|/2} \cdot c^{-1} e^{-c|k-\varphi_j|} \\ &\leq 16c^{-2} e^{-c|m-\varphi_j|/2}, \end{aligned}$$

which verifies (5.1). Here, the first inequality follows from (5.4), together with the facts that $|u_j(k; 0)| \leq 1$ (as \mathbf{u}_j is a unit vector) and $|V_{mk}(s)| \leq 1$ (as $\mathbf{V}(s)$ is orthogonal); the second follows from (5.5) (with the fact that $T(\log N)^2 \geq 40T \log N$ for sufficiently large N) and the fact that

we have restricted to the event \mathbf{E} from (5.2) (with the fact that, if $|m - \varphi_j| \geq T(\log N)^2$ and $|k - m| \leq |\varphi_j - m|/2$, then $|k - \varphi_j| \geq |m - \varphi_j|/2 \geq (\log N)^2/2$); and the third follows from performing the sums. \square

5.2. Resolvent Perturbation Estimates. In this section we estimate the effect of perturbing a random Lax matrix, associated with the open Toda lattice sampled from thermal equilibrium, on its eigenvalues. We first require some notation.

Assumption 5.3. Sample $(\mathbf{a}; \mathbf{b})$ under the thermal equilibrium $\mu_{\beta, \theta; N-1, N}$ from Definition 2.4, where $\mathbf{a} = (a_{N_1}, a_{N_1+1}, \dots, a_{N_2-1})$ and $\mathbf{b} = (b_{N_1}, b_{N_1+1}, \dots, b_{N_2})$. Let $\mathbf{L} = [L_{ij}] \in \text{SymMat}_{\llbracket N_1, N_2 \rrbracket}$ denote the associated Lax matrix (as in Definition 2.1), and let $\tilde{\mathbf{L}} = [\tilde{L}_{ij}] \in \text{SymMat}_{\llbracket N_1, N_2 \rrbracket}$ be another tridiagonal matrix. Assume that there is an index set $\mathcal{D} \subseteq \llbracket N_1, N_2 \rrbracket$ and a real number $\delta \in (0, 1)$ satisfying

$$(5.6) \quad \max_{i, j \notin (\llbracket N_1, N_2 \rrbracket \setminus \mathcal{D})^2} |\tilde{L}_{ij}| \leq 2 \log N; \quad \max_{i, j \in \llbracket N_1, N_2 \rrbracket \setminus \mathcal{D}} |L_{ij} - \tilde{L}_{ij}| \leq \delta.$$

For any $z \in \mathbb{C}$, denote $\mathbf{G}(z) = [G_{ij}(z)] = (\mathbf{L} - z)^{-1}$ and $\tilde{\mathbf{G}}(z) = [\tilde{G}_{ij}(z)] = (\tilde{\mathbf{L}} - z)^{-1}$.

The following lemma bounds the difference between entries of the resolvents $\mathbf{G}(z)$ and $\tilde{\mathbf{G}}(z)$, in terms of the distance from their indices to the set \mathcal{D} where \mathbf{L} and $\tilde{\mathbf{L}}$ might substantially disagree.

Lemma 5.4. *There exists a constant $c \in (0, 1)$ such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. Adopt Assumption 5.3; let $\eta \in [\delta, 1]$ be a real number; and define the set $\Omega = \{z \in \mathbb{C} : -N \leq \text{Re } z \leq N, \eta \leq \text{Im } z \leq 1\}$. For any integers $i, j \in \llbracket N_1, N_2 \rrbracket$, we have*

$$(5.7) \quad \sup_{z \in \Omega} |G_{ij}(z) - \tilde{G}_{ij}(z)| \leq e^{(\log N)^2} \eta^{-2} (\delta^{1/4} + e^{-c \text{dist}(i, \mathcal{D}) - c \text{dist}(j, \mathcal{D})}).$$

Proof. Throughout, we suppose $\text{dist}(i, \mathcal{D}) \geq \text{dist}(j, \mathcal{D})$, as the proof in the alternative case is entirely analogous. We may assume that $\text{dist}(i, \mathcal{D}) \geq 1$, as otherwise (5.7) holds deterministically, by (3.3).

We first show a variant of (5.7) for a fixed point $z \in \Omega$. So, fix $z_0 \in \Omega$; abbreviate $G_{km} = G_{km}(z_0)$ and $\tilde{G}_{km} = \tilde{G}_{km}(z_0)$, for any $k, m \in \llbracket N_1, N_2 \rrbracket$. Recalling Definition 3.13, also denote

$$(5.8) \quad \mathbf{E} = \text{BND}_{\mathbf{L}}(\log N), \quad \text{so that} \quad \mathbb{P}[\mathbf{E}^{\mathbf{C}}] \leq c^{-1}e^{-c(\log N)^2},$$

for some constant $c > 0$, by Lemma 3.15. Further let $\mathcal{D}_0 = \{k \in \llbracket N_1, N_2 \rrbracket : \text{dist}(k, \mathcal{D}) \leq 1\}$. Then, for any real number $s \in (0, 1)$, there exist constants $c_1 = c_1(s) > 0$ and $C_1 = C_1(s) > 1$ such that

$$\begin{aligned} \mathbb{E}[\mathbb{1}_{\mathbf{E}} \cdot |G_{ij} - \tilde{G}_{ij}|^s] &\leq \sum_{N_1 \leq k, m \leq N_2} \mathbb{E}[\mathbb{1}_{\mathbf{E}} \cdot |G_{ik}(L_{km} - \tilde{L}_{km})\tilde{G}_{kj}|^s] \\ &\leq \eta^{-s} \sum_{N_1 \leq k, m \leq N_2} \mathbb{E}[\mathbb{1}_{\mathbf{E}} \cdot |G_{ik}|^s \cdot |L_{km} - \tilde{L}_{km}|^s] \\ &\leq \delta^s \eta^{-s} \sum_{k \notin \mathcal{D}_0} \mathbb{E}[|G_{ik}|^s] + \eta^{-s} (3 \log N)^s \cdot \sum_{k \in \mathcal{D}_0} \mathbb{E}[|G_{ik}|^s] \\ &\leq C_1 \delta^s \eta^{-s} \sum_{k \notin \mathcal{D}_0} e^{-c_1|k-i|} + 3C_1 \eta^{-s} (\log N)^s \cdot \sum_{k \in \mathcal{D}_0} e^{-c_1|k-i|} \\ &\leq 8c_1^{-1} C_1 \eta^{-s} (\log N)^s (\delta^s + e^{-c_1 \text{dist}(i, \mathcal{D}_0)}), \end{aligned}$$

where in the first inequality we applied (3.4); in the second we applied (3.3); in the third we used (5.6), with the definition (5.8) of \mathbf{E} (which implies that $\mathbb{1}_{\mathbf{E}} \cdot |L_{km} - \tilde{L}_{km}| \leq 3 \log N$) and the fact

that $\mathbf{L} - \tilde{\mathbf{L}}$ is tridiagonal; in the fourth we used (3.8); and in the fifth we performed the sums. Taking $s = 2/3$ and using the fact that $\text{Im } z_0 \geq \eta$ (as $z_0 \in \Omega$), we therefore deduce by a Markov estimate that there exist constants $c_2 > 0$ and $C_2 > 1$ such that

$$(5.9) \quad \begin{aligned} \mathbb{P}[\mathbb{1}_{\mathbf{E}} \cdot |G_{ij} - \tilde{G}_{ij}| \geq e^{(\log N)^2/2} \eta^{-1} (\delta + e^{-c_1 \text{dist}(i, \mathcal{D})})^{1/4}] \\ \leq e^{-(\log N)^2/3} \eta^{2/3} (\delta + e^{-c_1 \text{dist}(i, \mathcal{D})})^{-1/6} \cdot \mathbb{E}[\mathbb{1}_{\mathbf{E}} \cdot |G_{ij} - \tilde{G}_{ij}|^{2/3}] \\ \leq C_2 e^{-c_2 (\log N)^2} (\delta + e^{-c_1 \text{dist}(i, \mathcal{D})})^{1/2}. \end{aligned}$$

Now, let us extend the estimate (5.9) to simultaneously apply for all $z \in \Omega$. To do this, set $\delta_0 = (\delta + e^{-c_1 \text{dist}(i, \mathcal{D})})^{1/4}$. Observe for any $z, z' \in \Omega$ with $|z - z'| \leq \delta_0$ that

$$(5.10) \quad |G_{ij}(z) - G_{ij}(z')| \leq |z - z'| \sum_{k=1}^N |G_{ik}(z)| \cdot |\tilde{G}_{kj}(z)| \leq \eta^{-2} N \delta_0,$$

where in the first inequality we used (3.4) and in the second we used (3.3) (with the fact that $\min\{\text{Im } z, \text{Im } z'\} \geq \eta$ for $z, z' \in \Omega$). Let $\Omega_0 \subset \Omega$ denote a δ_0 -mesh of Ω , so that $|\Omega_0| \leq 8N\delta_0^{-2}$. Denote the event

$$\mathbf{F} = \left\{ \max_{z \in \Omega_0} |G_{ij}(z) - \tilde{G}_{ij}(z)| \leq e^{(\log N)^2/2} \eta^{-1} \delta_0 \right\}.$$

Applying (5.8) and (5.9) for all $z_0 \in \Omega$, yields constants $c_3 > 0$ and $C_3 > 1$ such that

$$(5.11) \quad \mathbb{P}[\mathbf{F}^c] \leq C_2 e^{-c_2 (\log N)^2} \delta_0^2 \cdot |\Omega_0| + \mathbb{P}[\mathbf{E}^c] \leq 8C_2 N e^{-c_2 (\log N)^2} + \mathfrak{c}^{-1} e^{-c (\log N)^2} \leq C_3 e^{-c_3 (\log N)^2}.$$

Observe for sufficiently large N that $|G_{ij}(z) - \tilde{G}_{ij}(z)| \leq e^{(\log N)^2} \eta^{-2} \delta_0$ holds for all $z \in \Omega$ on the event \mathbf{F} , due to (5.10) and the estimate $e^{(\log N)^2/2} \eta^{-1} \delta_0 + 2\eta^{-2} N \delta_0 \leq e^{(\log N)^2} \eta^{-2} \delta_0$. Since $\text{dist}(i, \mathcal{D}) \geq \text{dist}(j, \mathcal{D})$, this yields (5.11). \square

Using Lemma 5.4 and Lemma 3.7, we can establish the following two corollaries. The first indicates that eigenvalues of \mathbf{L} with localization centers distant from \mathcal{D} are also nearly eigenvalues of $\tilde{\mathbf{L}}$ (with the same localization center), and the second essentially indicates the reverse.

Corollary 5.5. *There exists a constant $c \in (0, 1)$ such that the following holds with probability at least $1 - c^{-1} e^{-c (\log N)^2}$. Adopt Assumption 5.3; assume $\delta \leq e^{-10 (\log N)^2}$, and let $\zeta \geq e^{-200 (\log N)^{3/2}}$ be a real number. Fix $\lambda \in \text{eig } \mathbf{L}$, and let $\varphi \in \llbracket N_1, N_2 \rrbracket$ denote a ζ -localization center of λ with respect to \mathbf{L} , satisfying*

$$(5.12) \quad \text{dist}(\varphi, \mathcal{D}) \geq (\log N)^3.$$

Then there exists an eigenvalue $\tilde{\lambda} \in \text{eig } \tilde{\mathbf{L}}$ such that

$$(5.13) \quad |\lambda - \tilde{\lambda}| \leq e^{(\log N)^2} (\delta^{1/8} + e^{-c \text{dist}(\varphi, \mathcal{D})}),$$

and φ is an $N^{-1}\zeta$ -localization center for $\tilde{\lambda}$ with respect to $\tilde{\mathbf{L}}$.

Proof. Let $\mathfrak{c} \in (0, 1)$ denote the constant c from Lemma 5.4, and set

$$(5.14) \quad \delta_0 = \delta^{1/4} + e^{-c \text{dist}(\varphi, \mathcal{D})}; \quad \eta = (e^{(\log N)^2} \delta_0)^{1/2}.$$

Recalling the event \mathbf{E} from (5.8), also define event

$$(5.15) \quad \mathbf{F} = \mathbf{E} \cap \left\{ \sup_{E \in [-N, N]} |G_{\varphi\varphi}(E + i\eta) - \tilde{G}_{\varphi\varphi}(E + i\eta)| \leq e^{(\log N)^2} \eta^{-2} \delta_0 \right\}.$$

By (5.6) and the min-max principle, on \mathbf{F} , we have $\text{eig } \mathbf{L} \cup \text{eig } \tilde{\mathbf{L}} \subseteq [-N, N]$ for sufficiently large N .

By (5.8) and Lemma 5.4, we may restrict to the event \mathbf{F} . We then apply Lemma 3.7, with the parameters $(\varphi; \eta, \zeta, \delta; \mathbf{A}, \mathbf{B})$ there equal to $(\varphi; \eta, \zeta, e^{(\log N)^2} \eta^{-2} \delta_0; \mathbf{L}, \tilde{\mathbf{L}})$ here. To verify (3.5), observe that the first estimate there follows for sufficiently large N from the fact that

$$(5.16) \quad (2\eta) \cdot e^{(\log N)^2} \eta^{-2} \delta_0 \leq 2e^{(\log N)^2/2} \cdot (\delta^{1/4} + e^{-c \text{dist}(\varphi, \mathcal{D})})^{1/2} \leq e^{-400(\log N)^{3/2}} \leq \zeta^2,$$

where in the first bound we used (5.14); in the second we used the facts that $\delta \leq e^{-10(\log N)^2}$, that $\text{dist}(\varphi, \mathcal{D}) \geq (\log N)^3$, and that N is sufficiently large; and in the third we used the fact that $\zeta \geq e^{-200(\log N)^{3/2}}$. The second estimate in (3.5) follows from the fact that φ is a ζ -localization center for λ with respect to \mathbf{L} , and the third follows from our restriction to \mathbf{F} .

Thus, Lemma 3.7 yields some $\tilde{\lambda} \in \text{eig } \tilde{\mathbf{L}}$ such that $|\lambda - \tilde{\lambda}| \leq 3N\zeta^{-2}\eta$, which by (5.14) (and again the fact that $\zeta \geq e^{-200(\log N)^{3/2}}$) confirms (5.13). Since $(6N)^{-1/2}\zeta \geq N^{-1}\zeta$, Lemma 3.7 also indicates that φ is an $N^{-1}\zeta$ -localization center $\tilde{\lambda}$ with respect to $\tilde{\mathbf{L}}$, establishing the corollary. \square

Corollary 5.6. *There exists a constant $c \in (0, 1)$ such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. Adopt Assumption 5.3; assume that $\delta \leq e^{-10(\log N)^2}$; and let $\zeta \geq Ne^{-200(\log N)^{3/2}}$ be a real number. Fix $\tilde{\lambda} \in \text{eig } \tilde{\mathbf{L}}$, and let $\tilde{\varphi} \in \llbracket N_1, N_2 \rrbracket$ denote a ζ -localization center of $\tilde{\lambda}$ with respect to $\tilde{\mathbf{L}}$; suppose that*

$$(5.17) \quad \text{dist}(\tilde{\varphi}, \mathcal{D}) \geq (\log N)^3.$$

- (1) *There exists a unique eigenvalue $\lambda \in \text{eig } \mathbf{L}$ such that $|\lambda - \tilde{\lambda}| \leq e^{(\log N)^2} (\delta^{1/8} + e^{-c \text{dist}(\tilde{\varphi}, \mathcal{D})})$.*
- (2) *We have that $\tilde{\varphi}$ is an $N^{-1}\zeta$ -localization center of λ with respect to \mathbf{L} , and any $N^{-1}\zeta$ -localization center $\varphi \in \llbracket N_1, N_2 \rrbracket$ satisfies $|\varphi - \tilde{\varphi}| \leq (\log N)^2/2$.*

Proof. Let $c \in (0, 1)$ denote the constant c from Lemma 5.4, and set δ_0 and η as in (5.14). Recalling Definition 3.13 and the event \mathbf{F} from (5.15), define the event $\mathbf{G} = \mathbf{F} \cap \text{SEP}_{\mathbf{L}}(e^{-(\log N)^2/2})$. By (5.8), Lemma 3.18, and Lemma 5.4, there exists $c_1 > 0$ such that $\mathbb{P}[\mathbf{G}^c] \leq c_1^{-1}e^{-c_1(\log N)^2}$. By (5.6) and the min-max principle we have on \mathbf{G} that $\text{eig } \mathbf{L} \cup \text{eig } \tilde{\mathbf{L}} \subseteq [-N, N]$ for sufficiently large N .

Restricting to the event \mathbf{G} , we then apply Lemma 3.7, with the parameters $(\varphi; \eta, \zeta, \delta; \mathbf{A}, \mathbf{B})$ there equal to $(\tilde{\varphi}; \eta, \zeta, e^{(\log N)^2} \eta^{-2} \delta_0; \tilde{\mathbf{L}}, \mathbf{L})$ here. As in the proof of Corollary 5.5, the first bound in (3.5) is verified by (5.16); the second by the fact that $\tilde{\varphi}$ is a ζ -localization center for $\tilde{\lambda}$ with respect to $\tilde{\mathbf{L}}$; and the third by our restriction to \mathbf{G} . Thus, Lemma 3.7 gives some $\lambda \in \text{eig } \mathbf{L}$ such that $|\lambda - \tilde{\lambda}| \leq 3N\zeta^{-2}\eta$, which by (5.14) (and again the bound $\zeta \geq e^{-200(\log N)^{3/2}}$) yields some λ satisfying the first statement of the corollary. As any two distinct eigenvalues of \mathbf{L} differ by at least $e^{-(\log N)^2/2}$ by our restriction to \mathbf{G} , and $6N\zeta^{-2}\eta \leq e^{-(\log N)^2}$ (by (5.14), (5.17), and the facts that $\zeta \geq e^{-200(\log N)^{3/2}}$ and $\delta \leq e^{-10(\log N)^2}$), such an eigenvalue $\lambda \in \text{eig } \mathbf{L}$ satisfying these hypotheses is unique.

Since $(6N)^{-1/2}\zeta \geq N^{-1}\zeta$, Lemma 3.7 also indicates that φ is an $N^{-1}\zeta$ -localization center $\tilde{\lambda}$ with respect to $\tilde{\mathbf{L}}$. Thus, the second part of Lemma 5.1 yields $c_2 > 0$ such that, with probability at least $1 - c_2^{-1}e^{-c_2(\log N)^2}$, any $N^{-1}\zeta$ -localization center $\varphi \in \llbracket N_1, N_2 \rrbracket$ of λ with respect to \mathbf{L} satisfies $|\varphi - \tilde{\varphi}| \leq (\log N)^2/2$. This confirms the second statement of the corollary. \square

6. EIGENVECTOR ANALYSIS FOR LAX MATRICES

In this section we analyze the eigenvectors of random Lax matrices. We begin in Section 6.1 by proving a result, Proposition 6.1, that approximates the exponential rate of decay of the first entries of these eigenvectors, by reducing it to Proposition 6.3 below; the latter indicates that eigenvalues of the Lax matrix are close to those of certain truncations of it. We then show Proposition 6.3 in Section 6.2 and Section 6.3. Throughout this section, we adopt Assumption 2.8.

6.1. Eigenvector Decay for Lax Matrices. The following proposition estimates the behavior of the first entries of eigenvectors of the Lax matrix $\mathbf{L}(t)$. Recall that we adopt Assumption 2.8 throughout.

Proposition 6.1. *Let $t \in [0, T]$ be a real number. There is a constant $c > 0$ so that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. For any integer $k \in \llbracket 1, N \rrbracket$ satisfying*

$$(6.1) \quad N_1 + T(\log N)^5 + N^{1/100} \leq \varphi_t(k) \leq N_2 - T(\log N)^5 - N^{1/100},$$

we have

$$(6.2) \quad \left| \log |u_k(N_1; t)| - \sum_{j=N_1}^{\varphi_t(k)-1} \log L_{j,j+1}(t) + \sum_{i:\varphi_t(i) < \varphi_t(k)} \log |\lambda_i - \lambda_k| \right| \leq (\log N)^6.$$

To establish Proposition 6.1, we use the following lower bounds on $\log |u_k(N_1; t)|$ and $\log |u_k(N_2; t)|$.

Lemma 6.2. *Adopting the notation and assumptions of Proposition 6.1, we have with probability at least $1 - c^{-1}e^{-c(\log N)^2}$ that*

$$(6.3) \quad \begin{aligned} \log |u_k(N_1; t)| &\geq \sum_{j=N_1}^{\varphi_t(k)-1} \log L_{j,j+1}(t) - \sum_{i:\varphi_t(i) < \varphi_t(k)} \log |\lambda_i - \lambda_k| - \frac{1}{2} \cdot (\log N)^6; \\ \log |u_k(N_2; t)| &\geq \sum_{j=\varphi_t(k)}^{N_2-1} \log L_{j,j+1}(t) - \sum_{i:\varphi_t(i) > \varphi_t(k)} \log |\lambda_i - \lambda_k| - \frac{1}{2} \cdot (\log N)^6. \end{aligned}$$

Given Lemma 6.2, we can quickly establish Proposition 6.1 using Lemma 3.8.

Proof of Proposition 6.1. Let \mathbf{F} denote the event on which (6.3) holds. By Lemma 6.2, there exists a constant $c > 0$ such that $\mathbb{P}[\mathbf{F}^c] \leq c^{-1}e^{-c(\log N)^2}$. It suffices to show that (6.2) holds for sufficiently large N on the event \mathbf{F} ; assume to the contrary that this is false. Since the first bound in (6.3) holds but (6.2) does not hold, we must have

$$\log |u_k(N_1; t)| \geq \sum_{j=N_1}^{\varphi_t(k)-1} \log L_{j,j+1}(t) - \sum_{i:\varphi_t(i) < \varphi_t(k)} \log |\lambda_i - \lambda_k| + (\log N)^6.$$

Together with the second bound in (6.3), this yields

$$\log |u_k(N_1; t)| + \log |u_k(N_2; t)| \geq \sum_{j=N_1}^{N_2-1} \log L_{j,j+1}(t) - \sum_{i \neq k} \log |\lambda_i - \lambda_k| + \frac{1}{2} \cdot (\log N)^6,$$

which contradicts Lemma 3.8. This verifies the proposition. \square

To prove Lemma 6.2, we will make use of Lemma 3.9, which expresses the eigenvector entries of $\mathbf{L}(t)$ in terms of transfer matrices. The latter admit an explicit form, given by Lemma 3.10, which involves the eigenvalues of truncations of $\mathbf{L}(t)$. So, we require an estimate on how the eigenvalues of the Lax matrix $\mathbf{L}(t)$ change after setting one of its rows and columns to 0. This is provided by the first part of the following proposition. Its second part states that the localization centers of these eigenvalues cannot differ by too much, and its third improves the bound on this difference (making it independent of T) if one of the localization centers is not too close to an endpoint of the domain $\llbracket N_1, N_2 \rrbracket$. The proof of this proposition will appear in Section 6.3 below.

Proposition 6.3. *Let $t \in [0, T]$ be a real number. There exists a constant $c > 0$ such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. Let $\ell \in \llbracket N_1, N_2 \rrbracket$ be an integer, and assume that*

$$(6.4) \quad N_1 + T(\log N)^4 + N^{1/100} \leq \ell \leq N_2 - T(\log N)^4 - N^{1/100}.$$

Set $\mathbf{P} = \mathbf{L}(t)$ and $\mathbf{M} = \mathbf{P}^{(\ell)}$. Let $\mu \in \text{eig } \mathbf{M}$ be any eigenvalue of \mathbf{M} ; let $\Phi \in \llbracket N_1, N_2 \rrbracket$ be any ζ -localization center for μ with respect to \mathbf{M} . Suppose that $|\Phi - \ell| \geq (\log N)^4$.

- (1) *There exists a unique eigenvalue $\lambda \in \text{eig } \mathbf{P}$ such that $|\lambda - \mu| \leq e^{-c \min\{|\Phi - \ell|, N^{1/100}\}}$.*
- (2) *If $\varphi \in \llbracket N_1, N_2 \rrbracket$ is a ζ -localization center for λ with respect to \mathbf{P} , then we have $|\varphi - \Phi| \leq T(\log N)^3$.*
- (3) *Assume that*

$$(6.5) \quad N_1 + T(\log N)^3 + \frac{1}{2} \cdot N^{1/100} \leq \Phi \leq N_2 - T(\log N)^3 - \frac{1}{2} \cdot N^{1/100}.$$

If $\varphi \in \llbracket N_1, N_2 \rrbracket$ is a ζ -localization center for λ with respect to \mathbf{P} , then $|\varphi - \Phi| \leq (\log N)^2$.

Given Proposition 6.3, we can establish Lemma 6.2.

Proof of Lemma 6.2. We only establish the first bound in (6.3), as the second would then follow from symmetry. Throughout, we set $\mathbf{P} = [P_{ij}] = \mathbf{L}(t)$ and $v_j = u_k(j; t)$ for each $j \in \llbracket N_1, N_2 \rrbracket$.

Recall from Section 3.4 the notation on the transfer matrices $\mathbf{S}_{\mathcal{X}}$, defined by (3.6) and (3.7); abbreviate $\mathbf{S} = [S_{ij}] = \mathbf{S}_{\llbracket N_1, \varphi_t(k) \rrbracket}(\lambda_k; \mathbf{P}) \in \text{Mat}_{2 \times 2}$, whose entries are indexed by $i, j \in \{1, 2\}$. Also denote $\mathbf{Q} = \mathbf{P}^{\llbracket N_1, \varphi_t(k) - 1 \rrbracket}$, where we recall from above Lemma 3.10 that this is the matrix obtained by restricting \mathbf{P} to rows and columns indexed by $\llbracket N_1, \varphi_t(k) - 1 \rrbracket$. Then Lemma 3.9 yields $\mathbf{S} \cdot (0, v_{N_1}) = (v_{\varphi_t(k)}, v_{\varphi_t(k)+1})$, so $S_{12} \cdot v_{N_1} = v_{\varphi_t(k)}$. We thus deduce for sufficiently large N that

$$\begin{aligned} \log |v_{N_1}| &= \log |v_{\varphi_t(k)}| - \log |S_{12}| \\ &\geq \log \zeta - \log |S_{12}| \geq \sum_{j=N_1}^{\varphi_t(k)-1} \log P_{j, j+1} - \sum_{\mu \in \text{eig } \mathbf{Q}} \log |\mu - \lambda_k| - (\log N)^2, \end{aligned}$$

where in the second statement we used the fact that φ_t is a ζ -localization center bijection for \mathbf{P} , and in the third we used Lemma 3.10 and the fact that $\zeta \geq e^{-100(\log N)^{3/2}} \geq e^{-(\log N)^2}$. To verify the first bound in (6.3), it therefore suffices to show that with probability at least $1 - c^{-1}e^{-c(\log N)^2}$ we have

$$(6.6) \quad \sum_{\mu \in \text{eig } \mathbf{Q}} \log |\mu - \lambda_k| \leq \sum_{i: \varphi_t(i) < \varphi_t(k)} \log |\lambda_i - \lambda_k| + \frac{1}{3} \cdot (\log N)^6.$$

To that end, recalling Definition 3.13, define the event $\mathbf{E}_1 = \text{BND}_{\mathbf{P}}(\log N) \cap \text{SEP}_{\mathbf{P}}(e^{-(\log N)^2})$. Then $\mathbb{P}[\mathbf{E}_1^c] \leq c_1^{-1}e^{-c_1(\log N)^2}$ for some constant $c_1 > 0$, due to Lemma 3.15 and Lemma 3.18.

We next apply Proposition 6.3, with the ℓ there equal to $\varphi_t(k)$ here; observe that the estimate (6.4) assumed in that proposition holds by (6.1). Thus, Proposition 6.3 yields a constant $c_2 > 0$ and an event \mathbf{E}_2 with $\mathbb{P}[\mathbf{E}_2^c] \leq c_2^{-1} e^{-c_2(\log N)^2}$ such that the following holds on \mathbf{E}_2 . Let $\mu \in \text{eig } \mathbf{Q}$ be any eigenvalue that admits a ζ -localization center $\Phi \in \llbracket N_1, \varphi_t(k) - 1 \rrbracket$ with respect to \mathbf{Q} satisfying $\Phi \leq \varphi_t(k) - (\log N)^4$.

- (1) There exists a unique eigenvalue $\lambda = \lambda(\mu) \in \text{eig } \mathbf{P}$ such that $|\lambda - \mu| \leq e^{-(\log N)^3}$.
- (2) Letting $\lambda = \lambda_\psi$ for some $\psi \in \llbracket 1, N \rrbracket$, we have $|\varphi_t(\psi) - \Phi| \leq T(\log N)^3$.
- (3) If $\Phi \geq N_1 + T(\log N)^3 + N^{1/100}/2$, then we further have $|\varphi_t(\psi) - \Phi| \leq (\log N)^2$.

In what follows, we restrict to $\mathbf{E}_1 \cap \mathbf{E}_2$ and show (6.6) holds.

To that end, denote $\text{eig } \mathbf{Q} = (\mu_1, \mu_2, \dots, \mu_{\varphi_t(k) - N_1})$, and let $\Phi : \llbracket 1, \varphi_t(k) - N_1 \rrbracket \rightarrow \llbracket N_1, \varphi_t(k) - 1 \rrbracket$ denote a $(2N)^{-1}$ -localization center bijection for \mathbf{Q} , which is guaranteed to exist by Lemma 2.7. Let \mathcal{J} denote the set of indices $j \in \llbracket 1, \varphi_t(k) - N_1 \rrbracket$ such that $\Phi(j) \leq \varphi_t(k) - (\log N)^4$. For each $j \in \mathcal{J}$, let $\psi_j \in \llbracket 1, N \rrbracket$ denote the unique index such that

$$(6.7) \quad |\lambda_{\psi_j} - \mu_{\Phi(j)}| \leq e^{-(\log N)^3}, \quad \text{where } j \text{ satisfies } \varphi_t(\psi_j) < \varphi_t(k).$$

Such an index ψ_j satisfying the first bound in (6.7) exists and is unique by our restriction to \mathbf{E}_2 . To verify that ψ_j satisfies the second observe that, if $\Phi(j) \geq N_1 + T(\log N)^3 + N^{1/100}/2$, then

$$\varphi_t(\psi_j) \leq \Phi(j) + (\log N)^2 \leq \varphi_t(k) - (\log N)^4 + (\log N)^2 < \varphi_t(k),$$

the first bound since we restricted to \mathbf{E}_2 , the second since $j \in \mathcal{J}$, and the third since $N > 1$. If instead $\Phi(j) < N_1 + T(\log N)^3 + N^{1/100}/2$, then

$$\varphi_t(\psi_j) \leq \Phi(j) + T(\log N)^3 \leq N_1 + 2T(\log N)^4 + \frac{1}{2} \cdot N^{1/100} < \varphi_t(k),$$

the first bound by our restriction to \mathbf{E}_2 , the second by our assumption on $\Phi(j)$, and the third by (6.1). This confirms (6.7). We also have that

$$(6.8) \quad \begin{aligned} \log |\mu_h - \lambda_k| &\leq \log N && \text{for all } h \in \llbracket 1, \varphi_t(k) - N_1 - 1z \rrbracket; \\ \log |\mu_j - \lambda_k| &\leq \log |\lambda_{\psi_j} - \lambda_k| + e^{-(\log N)^2}, && \text{for all } j \in \mathcal{J}. \end{aligned}$$

Here, the first bound holds since $|\mu_i - \lambda_i| \leq 2 \log N$ (by our restriction to \mathbf{E}_1 and the fact that $\mathbf{E}_1 \subseteq \text{BND}_{\mathbf{Q}}(\log N)$, by Remark 3.14); the second holds since $|\lambda_{\psi_j} - \mu_j| \leq e^{-(\log N)^3}$, with the bound $|\lambda_{\psi_j} - \lambda_k| \geq e^{-(\log N)^2}$ (by our restriction to \mathbf{E}_1 and the fact from (6.7) that $\psi_j \neq k$). Therefore,

$$\begin{aligned} \sum_{\mu \in \mathbf{Q}} \log |\mu - \lambda_k| &\leq \sum_{j \in \mathcal{J}} \log |\mu_h - \lambda_k| + (\log N)^5 \\ &\leq \sum_{j \in \mathcal{J}} \log |\lambda_{\psi_j} - \lambda_k| + 2(\log N)^5 \leq \sum_{i: \varphi_t(i) < \varphi_t(k)} \log |\lambda_i - \lambda_k| + 3(\log N)^5. \end{aligned}$$

Here, in the first estimate, we used the fact that at most $(\log N)^4$ indices $h \notin \mathcal{J}$ exist, with the first bound in (6.8) for each such h . In the second, we used the second bound in (6.8). In the third we used the fact that $\varphi_t(\psi_j) < \varphi_t(k)$ for each $j \in \mathcal{J}$ (by (6.7)); the bound $\log |\lambda_i - \lambda_k| \leq \log(2 \log N) \leq \log N$ for all indices i with $\varphi_t(i) \in \llbracket N_1, \varphi_t(k) - 1 \rrbracket$ but that are not of the form ψ_j for some $j \in \mathcal{J}$; and the fact that at most $(\log N)^4$ such indices i exist (as is quickly verified from the injectivity of φ_t and ψ , the second statement in (6.7), and the fact that there are at most $(\log N)^4$ indices not in \mathcal{J}). This establishes the second bound in (6.6) and thus the lemma. \square

6.2. Eigenvalues of Truncated Lax Matrices. In this section we begin the proof of Proposition 6.3; we adopt the notation of that proposition throughout. We first address it when (6.5) holds, that is, when Φ is not too close to N_1 or N_2 .

Proposition 6.4. *Proposition 6.3 holds assuming (6.5).*

To prove Proposition 6.4, we first apply Proposition 4.4, which yields a constant $c_0 > 0$; random matrix $\mathbf{Q} = [Q_{ij}] \in \text{SymMat}_{\llbracket N_1, N_2 \rrbracket}$ with the same law as $\mathbf{L}(0)$; and an event \mathbf{E} with $\mathbb{P}[\mathbf{E}^c] \leq c^{-1}e^{-c(\log N)^2}$ such that on \mathbf{E} we have

$$(6.9) \quad \max_{i,j \in \llbracket N_1+K, N_2-K \rrbracket} |P_{ij} - Q_{ij}| \leq e^{-K/5}, \quad \text{for any } K \geq T \log N.$$

Lemma 6.5. *If (6.5) holds, then there exists a constant $c > 0$ such that the following two statements hold with probability at least $1 - c^{-1}e^{-c(\log N)^2}$.*

- (1) *There exists an eigenvalue $\kappa \in \text{eig } \mathbf{Q}$ such that $|\mu - \kappa| \leq e^{-c \min\{|\Phi - \ell|, N^{1/100}\}}$.*
- (2) *The index $\Phi \in \llbracket N_1, N_2 \rrbracket$ is an $N^{-2}\zeta$ -localization center for κ with respect to \mathbf{Q} .*

Proof. Define $\mathbf{R} = [R_{ij}] \in \text{SymMat}_{\llbracket N_1, N_2 \rrbracket}$ by setting $\mathbf{R} = \mathbf{Q}^{(\ell)}$. Throughout this proof, we assume that $\Phi \leq \ell$, as the proof when $\Phi \geq \ell$ is entirely analogous. Then denote by $\mathbf{P}' = [P'_{ij}]$ and $\mathbf{Q}' = [Q'_{ij}]$ the top $(\ell - N_1) \times (\ell - N_1)$ corners of \mathbf{P} and \mathbf{Q} , respectively. In this way, their rows and columns are indexed by $i, j \in \llbracket N_1, \ell - 1 \rrbracket$, and we have $P'_{ij} = P_{ij} = M_{ij}$ and $Q'_{ij} = Q_{ij} = R_{ij}$ for each such (i, j) ; observe that $\text{eig } \mathbf{P}' \subseteq \text{eig } \mathbf{M}$ and $\text{eig } \mathbf{Q}' \subseteq \text{eig } \mathbf{R}$. Since $\Phi \leq \ell$, we also have $\mu \in \text{eig } \mathbf{P}'$.

To establish the lemma, we will first use Corollary 5.6 to show μ is close to some $\nu \in \text{eig } \mathbf{R}$. To that end, recalling Definition 3.13, define the quantities $K, \delta > 0$ and event \mathbf{E}_1 by

$$(6.10) \quad K = \frac{1}{10} \cdot \min\{|\Phi - \ell|, N^{1/100}\} + \frac{T}{2} \cdot (\log N)^3; \quad \delta = e^{-K/4}; \quad \mathbf{E}_1 = \bigcap_{s \geq 0} \text{BND}_{L(s)}\left(\frac{\log N}{600}\right).$$

By Lemma 3.15, there is $c_1 > 0$ so that $\mathbb{P}[\mathbf{E}_1^c] \leq c_1^{-1}e^{-c_1(\log N)^2}$; we restrict to \mathbf{E}_1 in what follows.

Applying Corollary 5.6 requires (through Assumption 5.3) a bound on $|P'_{ij} - Q'_{ij}|$ for i, j sufficiently far from N_1 . This will follow from (6.9); indeed, observe that $N_2 - \ell \geq T(\log N)^4 + N^{1/100} \geq K$, where the first bound holds by (6.4) and the second holds by (6.10). Thus $N_2 - K \geq \ell$, so since $(P'_{ij}, Q'_{ij}) = (P_{ij}, Q_{ij})$ for $i, j \leq \ell$, it follows from (6.9) that

$$(6.11) \quad \max_{i,j \in \llbracket N_1+K, \ell \rrbracket} |P'_{ij} - Q'_{ij}| \leq \delta.$$

Using this, we next verify Assumption 5.3 with the $(\mathbf{L}, \tilde{\mathbf{L}})$ there given by $(\mathbf{Q}', \mathbf{P}')$ here. To that end, observe that $|P_{ij}| \leq (\log N)/50 \leq 2 \log(\ell - N_1)$, where the first bound follows from Lemma 3.5 and our restriction to \mathbf{E}_1 , and the second follows from (6.4). This, together with (6.11) and the fact that \mathbf{Q} has the same law as $\mathbf{L}(0)$, implies that Assumption 5.3 holds, with the $(\mathbf{L}, \tilde{\mathbf{L}}; \mathcal{D}; N)$ there equal to $(\mathbf{Q}', \mathbf{P}'; \llbracket N_1, N_1 + K - 1 \rrbracket; \ell - N_1)$ here. Since (6.5) and (6.10) together yield

$$(6.12) \quad \Phi - N_1 \geq T(\log N)^3 + \frac{1}{2} \cdot N^{1/100} \geq 2K + (\log N)^3,$$

we deduce that $\Phi \geq N_1 + K + (\log N)^3$, verifying (5.17). Since $\mu \in \text{eig } \mathbf{P}'$, the first statement of Corollary 5.6 yields a constant $c_2 > 0$ and an event \mathbf{E}_2 with $\mathbb{P}[\mathbf{E}_2^c] \leq c_2^{-1}e^{-c_2(\log N)^2}$ such that the following holds on \mathbf{E}_2 . There exists an eigenvalue $\nu \in \text{eig } \mathbf{Q}' \subseteq \text{eig } \mathbf{R}$ satisfying

$$(6.13) \quad |\nu - \mu| \leq e^{(\log N)^2} (\delta^{1/16} + e^{-2c_2|\Phi - N_1 - K|}) \leq e^{-c_2K},$$

where in the last inequality we used (6.12) and the bound $K \geq (\log N)^4/10$ (as $|\Phi - \ell| \geq (\log N)^4$). The second statement of Corollary 5.6 further implies that Φ is an $N^{-1}\zeta$ -localization center of ν with respect to \mathbf{Q}' , and thus with respect to \mathbf{R} . We further restrict to \mathbf{E}_2 in what follows.

We now use Corollary 5.6 again to show that ν is close to some $\kappa \in \text{eig } \mathbf{Q}$; this will proceed similarly to above. Since $R_{ij} = Q_{ij}$ unless $\ell \in \{i, j\}$ (and \mathbf{Q} has the same law as $\mathbf{L}(0)$), we have by our restriction to $\mathbf{E} \cap \mathbf{E}_1$ that Assumption 5.3 holds with the $(\mathbf{L}, \tilde{\mathbf{L}}; \delta; \mathcal{D})$ there equal to $(\mathbf{Q}, \mathbf{R}; 0; \{\ell\})$ here. Since $|\Phi - \ell| \geq (\log N)^4$, the first statement of Corollary 5.6 yields a constant $c_3 > 0$ and an event \mathbf{E}_3 with $\mathbb{P}[\mathbf{E}_3^c] \leq c_3^{-1}e^{-c_3(\log N)^2}$ such that following holds on \mathbf{E}_3 . There exists an eigenvalue $\kappa \in \text{eig } \mathbf{Q}$ such that

$$(6.14) \quad |\kappa - \nu| \leq e^{(\log N)^2} \cdot e^{-2c_3|\Phi - \ell|} \leq e^{-c_3|\Phi - \ell|},$$

where in the last inequality we used the fact that $|\Phi - \ell| \geq (\log N)^4$. Thus, (6.13) and (6.14) imply the first statement of the lemma. The second statement of Corollary 5.6 further implies that Φ is an $N^{-2}\zeta$ -localization center for κ with respect to \mathbf{Q} , verifying the second statement of the lemma. \square

Proof of Proposition 6.4. Throughout this proof, we adopt the notation and assumptions from Lemma 6.5 and its proof. In particular, we recall the quantities $K, \delta > 0$ and the event \mathbf{E}_1 from (6.10). Recalling Definition 3.13, we further define the event

$$(6.15) \quad \mathbf{F}_1 = \text{SEP}_{\mathbf{P}}(e^{-(\log N)^2}) \cap \text{SEP}_{\mathbf{Q}}(e^{-(\log N)^2}).$$

By Lemma 2.2, Lemma 3.18, and a union bound, there exists $c_1 > 0$ with $\mathbb{P}[\mathbf{F}_1^c] \leq c_1^{-1}e^{-c_1(\log N)^2}$. We restrict to $\mathbf{E}_1 \cap \mathbf{F}_1$, and further to the event that Lemma 6.5 holds, in what follows.

We will use Corollary 5.5 to show that κ is close to some $\lambda \in \text{eig } \mathbf{P}$. To that end, first observe from (6.9) and our restriction to $\mathbf{E} \cap \mathbf{E}_1$ (with the fact that \mathbf{Q} has the same law as $\mathbf{L}(0)$) that Assumption 5.3 holds, with the $(\mathbf{L}, \tilde{\mathbf{L}}; \mathcal{D})$ there equal to $(\mathbf{Q}, \mathbf{P}; \llbracket N_1, N_2 \rrbracket \setminus \llbracket N_1 + K, N_2 + K \rrbracket)$ here. Also, by (6.5), $\min\{\Phi - N_1 - K, N_2 - \Phi - K\} \geq N^{1/100}/2 \geq (\log N)^3$, which verifies (5.12). Therefore, Corollary 5.5 applies and yields a constant $c_2 > 0$ and an event \mathbf{F}_2 with $\mathbb{P}[\mathbf{F}_2^c] \leq c_2^{-1}e^{-c_2(\log N)^2}$ such that the following holds on \mathbf{F}_2 . There exists $\lambda \in \text{eig } \mathbf{P}$ such that Φ is a $N^{-3}\zeta$ -localization center of λ with respect to \mathbf{P} , and

$$(6.16) \quad |\lambda - \kappa| \leq e^{(\log N)^2} (\delta^{1/8} + e^{-2c_2 \min\{\Phi - N_1 - K, N_2 - \Phi - K\}}) \leq e^{-c_2 K},$$

where in the last inequality we used the facts that $\min\{\Phi - N_1, N_2 - \Phi\} \geq T(\log N)^3 + N^{1/100}/2 \geq 2K + (\log N)^3$ and $|\Phi - \ell| \geq (\log N)^4$. Together with the first statement in Lemma 6.5, this implies the existence of λ satisfying the conditions in the first statement of Proposition 6.3, upon additionally restricting to \mathbf{F}_2 . The uniqueness of such a $\lambda \in \text{eig } \mathbf{P}$ follows from our restriction to the event \mathbf{F}_1 from (6.15) (with the fact that $|\Phi - \ell| \geq (\log N)^4$).

It remains to establish the third part of the proposition (which implies the second, as we have assumed that (6.5) holds), to which we must use Corollary 5.6 (instead of Corollary 5.5). Fix any ζ -localization center $\varphi \in \llbracket N_1, N_2 \rrbracket$ for λ with respect to \mathbf{P} . Let a $\mathbf{u}(t) = (u(N_1; t), u(N_1 + 1; t), \dots, u(N_2; t)) \in \mathbb{R}^N$ be a unit eigenvector of \mathbf{P} with eigenvalue λ . Then, by the definition of φ and Corollary 5.5, we have

$$(6.17) \quad |u(\varphi; t)| \geq \zeta, \quad \text{and} \quad |u(\Phi; t)| \geq N^{-3}\zeta.$$

We must show (5.17), to which end we claim that, with high probability,

$$(6.18) \quad \min\{\varphi - N_1, N_2 - \varphi\} \geq K + (\log N)^3.$$

To do so, observe that Lemma 5.2 yields a constant $c_3 > 0$ and an event F_3 with $\mathbb{P}[F_3^c] \leq c_3^{-1} e^{-c_3(\log N)^2}$ such that the following holds on F_3 . An index $m \in \llbracket N_1, N_2 \rrbracket$ can only be a $N^{-3}\zeta$ -localization center for λ with respect to \mathbf{P} if $|m - \varphi| \leq T(\log N)^2$. Restricting to F_3 , the second bound in (6.17) then implies that we must have $|\Phi - \varphi| < T(\log N)^2$ for sufficiently large N . Together with the estimate (by (6.5) and (6.10))

$$\min\{\Phi - N_1, N_2 - \Phi\} \geq T(\log N)^3 + \frac{1}{2} \cdot N^{1/100} \geq K + T(\log N)^2 + (\log N)^3,$$

this confirms (6.18).

The bound (6.18) (with (6.9)) verifies (5.17), with the parameters $(\mathbf{L}, \tilde{\mathbf{L}}; \tilde{\lambda}, \tilde{\varphi}; \delta; \mathcal{D})$ there equal to $(\mathbf{Q}, \mathbf{P}; \lambda, \varphi; \delta; \llbracket N_1, N_2 \rrbracket \setminus \llbracket N_1 + K, N_2 - K \rrbracket)$ here. Thus, Corollary 5.6 applies and yields a constant $c_4 > 0$ and an event F_4 with $\mathbb{P}[F_4^c] \leq c_4^{-1} e^{-c_4(\log N)^2}$, such that on F_4 there exists $\kappa' \in \text{eig } \mathbf{Q}$ with

$$|\kappa' - \lambda| \leq e^{(\log N)^2} (\delta^{1/8} + e^{-c_4 \min\{\varphi - N_1 - K, N_2 - K - \varphi\}}) \leq e^{-c_4(\log N)^3},$$

where in the last inequality we used the definition (6.10) of δ (with the fact that $|\Phi - \ell| \geq (\log N)^4$) with (6.18). Together with (6.16), the fact that $|\Phi - \ell| \geq (\log N)^4$, and our restriction to the event F_1 from (6.15), this implies that $\kappa = \kappa'$. Hence, the second part of Corollary 5.6 implies that φ is a $N^{-1}\zeta$ -localization center for κ with respect to \mathbf{Q} . Since Φ is an $N^{-2}\zeta$ -localization center for κ with respect to \mathbf{Q} (by Lemma 6.5), it also implies that $|\varphi - \Phi| \leq (\log N)^2$, verifying the third (and thus also the second) statement of the proposition. \square

6.3. Proof of Proposition 6.3. In this section we establish Proposition 6.3, adopting its notation and assumptions throughout. For any $z \in \mathbb{C}$, denote the resolvents of \mathbf{M} and \mathbf{P} by $\check{\mathbf{G}}(z) = [\check{G}_{ij}(z)] = (\mathbf{M} - z)^{-1}$ and $\check{\check{\mathbf{G}}}(z) = [\check{\check{G}}_{ij}(z)] = (\mathbf{P} - z)^{-1}$, respectively. The following lemma estimates the (Φ, Φ) entry of $\check{\mathbf{G}} - \check{\check{\mathbf{G}}}$, assuming that (6.5) does not hold. Its proof is similar to that of Lemma 5.4 (by using Lemma 5.2 in place of Lemma 3.16), so we only outline it.

Lemma 6.6. *There exists a constant $c > 0$ so that the following holds with probability at least $1 - c^{-1} e^{-c(\log N)^2}$. Suppose that (6.5) does not hold, and let $\eta \in \mathbb{R}$ satisfy $e^{-cN^{1/100}} \leq \eta \leq 1$. Denoting $\Omega = \{z \in \mathbb{C} : -N \leq \text{Re } z \leq N, \eta \leq \text{Im } z \leq 1\}$, we have*

$$(6.19) \quad \sup_{z \in \Omega} |\check{\check{G}}_{\Phi\Phi}(z) - \check{G}_{\Phi\Phi}(z)| \leq e^{-cN^{1/100}}.$$

Proof of Lemma 6.6 (Outline). Throughout this proof, recalling Definition 3.13, we set

$$(6.20) \quad \mathbf{F} = \text{BND}_{\mathbf{P}}(\log N), \quad \text{so that} \quad \mathbb{P}[F^c] \leq c_1^{-1} e^{-c_1(\log N)^2},$$

for some constant $c_1 \in (0, 1)$, by Lemma 3.15. We begin by using Lemma 5.2 to estimate the eigenvectors of \mathbf{P} . To that end, first observe, by Lemma 2.2 and the fact that $\mathbf{P} = \mathbf{L}(t)$, that $\text{eig } \mathbf{P} = (\lambda_1, \lambda_2, \dots, \lambda_N)$. For each $j \in \llbracket 1, N \rrbracket$, recall that $\mathbf{u}_j(t) = (u_j(N_1; t), u_j(N_1 + 1; t), \dots, u_j(N_2; t)) \in \mathbb{R}^N$ denotes the nonnegatively normalized, unit eigenvector of \mathbf{P} with eigenvalue λ_j . For each $j \in \llbracket 1, N \rrbracket$, also let $\psi_j \in \llbracket N_1, N_2 \rrbracket$ denote a $N^{-1/2}$ -localization center for the eigenvector \mathbf{u}_j of \mathbf{L} of eigenvalue λ_j (which is guaranteed to exist, as \mathbf{u} is a unit vector). Since (6.5) does not hold, (6.4) implies for sufficiently large N that

$$|\ell - \Phi| \geq K_0, \quad \text{where} \quad K_0 = \frac{1}{2} \cdot (T(\log N)^4 + N^{1/100}).$$

Thus, for any $i \in \{\ell-1, \ell, \ell+1\}$ and $k \in \llbracket N_1, N_2 \rrbracket$, we either have $|\Phi - \psi_j| \geq K_0/2 \geq T(\log N)^2$ or $|i - \psi_j| \geq K_0/2 \geq T(\log N)^2$. Therefore, Lemma 5.2 yields a constant $\mathfrak{c} \in (0, c_1/25)$ and an event \mathbf{E}_0 with $\mathbb{P}[\mathbf{E}_0^c] \leq \mathfrak{c}^{-1}e^{-25\mathfrak{c}(\log N)^2}$, such that

$$(6.21) \quad \mathbb{1}_{\mathbf{E}_0} \cdot \max_{i:|i-\ell|\leq 1} \sum_{k \in \llbracket 1, N \rrbracket} |u_k(\Phi; t)| \cdot |u_k(i; t)| \leq N \cdot e^{-25\mathfrak{c}K_0} \leq e^{-12\mathfrak{c}N^{1/100}}.$$

Set $\mathbf{E} = \mathbf{E}_0 \cap \mathbf{F}$, which satisfies $\mathbb{P}[\mathbf{E}^c] \leq \mathfrak{c}^{-1}e^{-20\mathfrak{c}(\log N)^2}$, by (6.20) and a union bound.

Now, let us show that (6.19) holds with high probability for a fixed point $z \in \Omega$. Fix $z_0 \in \Omega$; abbreviate $\check{G}_{ij} = \check{G}_{ij}(z_0)$ and $\check{G}_{ij} = \check{G}_{ij}(z_0)$ for each $i, j \in \llbracket N_1, N_2 \rrbracket$. For $\eta \geq e^{-\mathfrak{c}N^{1/100}}$, we have

$$(6.22) \quad \begin{aligned} \mathbb{E}[\mathbb{1}_{\mathbf{E}} \cdot |\check{G}_{\Phi\Phi} - \check{G}_{\Phi\Phi}|] &\leq \sum_{\substack{i:|i-\ell|\leq 1 \\ j:|j-\ell|\leq 1}} \mathbb{E}[\mathbb{1}_{\mathbf{E}} \cdot |\check{G}_{\Phi i} \cdot P_{ij} \cdot \check{G}_{j\Phi}|] \\ &\leq 9\eta^{-1} \cdot \log N \cdot \max_{i:|i-\ell|\leq 1} \mathbb{E}[\mathbb{1}_{\mathbf{E}} \cdot |\check{G}_{\Phi i}|] \\ &\leq 9\eta^{-2} \cdot \log N \cdot \max_{i:|i-\ell|\leq 1} \sum_{k=1}^N \mathbb{E}[|u_k(\Phi; t)| \cdot |u_k(i; t)|] \\ &\leq 9\eta^{-2} \cdot \log N \cdot e^{-12\mathfrak{c}N^{1/100}} \leq e^{-9\mathfrak{c}N^{1/100}}, \end{aligned}$$

where the first inequality follows from (3.4), together with the fact that $P_{ij} - M_{ij} \in \{P_{ij}, 0\}$ with $P_{ij} - M_{ij} = 0$ unless $i, j \in \{\ell-1, \ell, \ell+1\}$; the second from (3.3) (with the fact that $\text{Im } z_0 \geq \eta$, as $z_0 \in \Omega$) and the definition (6.20) of \mathbf{F} ; the third from (3.2), together with the bound $|\lambda_k - z_0| \geq \text{Im } z_0 \geq \eta$ for any $k \in \llbracket 1, N \rrbracket$; the fourth from (6.21); and the fifth from the fact that $\eta \geq e^{-\mathfrak{c}N^{1/100}}$. The estimate (6.22), together with a Markov bound, verifies (6.19) at $z = z_0$.

To extend it to all $z \in \Omega$, we first use a union bound to apply (6.22) on an $e^{-3\mathfrak{c}N^{1/100}}$ -mesh $\Omega_0 \subset \Omega$. Then, using (3.4) and (3.3), we approximate the resolvents of \mathbf{M} and \mathbf{P} at an arbitrary points $z \in \Omega$ by those at the nearest point $z_0 \in \Omega_0$. This is very similar to what was done at the end of the proof of Lemma 5.4, so we omit further details. \square

Proof of Proposition 6.3. In what follows, we assume that (6.5) does not hold, as Proposition 6.4 establishes the proposition when it does. Letting $\mathfrak{c} > 0$ denote the constant $c/10$ from Lemma 6.6, set $\eta = e^{-2\mathfrak{c}N^{1/100}}$ and define the event (recalling Definition 3.13)

$$\begin{aligned} \mathbf{G}_1 &= \text{BND}_{\mathbf{P}}(\log N) \cap \text{SEP}_{\mathbf{P}}(e^{-(\log N)^2}) \\ &\cap \left\{ \sup_{E \in [-N, N]} |\check{G}_{\Phi\Phi}(E + i\eta) - \check{G}_{\Phi\Phi}(E + i\eta)| \leq e^{-10\mathfrak{c}N^{1/100}} \right\}. \end{aligned}$$

By Lemma 3.15, Lemma 3.18 (with Lemma 2.2), and Lemma 6.6, there exists a constant $c_1 > 0$ such that $\mathbb{P}[\mathbf{G}_1^c] \leq c_1^{-1}e^{-c_1(\log N)^2}$.

Restricting to \mathbf{G}_1 , we next apply Lemma 3.7 with the parameters $(\lambda, \varphi; \eta, \zeta, \delta; \mathbf{A}, \mathbf{B})$ there equal to $(\mu, \Phi; \eta, \zeta, e^{-10\mathfrak{c}N^{1/100}}; \mathbf{M}, \mathbf{P})$ here. The first bound in (3.5) holds for sufficiently large N by the facts that $\zeta \geq e^{-100(\log N)^{3/2}}$ and $\eta = e^{-2\mathfrak{c}N^{1/100}}$; the second holds since Φ is a ζ -localization center of μ with respect to \mathbf{M} ; and the third holds by our restriction to \mathbf{G}_1 . Hence, Lemma 3.7 yields an eigenvalue $\lambda \in \text{eig } \mathbf{P}$ such that $|\lambda - \mu| \leq 3N\zeta^{-2}\eta \leq e^{-\mathfrak{c}N^{1/100}}$ (again as $\zeta \geq e^{-100(\log N)^{3/2}}$ and $\eta = e^{-2\mathfrak{c}N^{1/100}}$). This shows the first statement of the proposition; it remains to verify the second.

To that end, observe that Lemma 3.7 further indicates that Φ is an $N^{-1}\zeta$ -localization center for λ with respect to \mathbf{P} . Letting $\Psi \in \llbracket N_1, N_2 \rrbracket$ denote an $N^{-1}\zeta$ -localization center of λ with respect to \mathbf{L} , Lemma 5.2 yields a constant $c_2 > 0$ and an event \mathbf{G}_2 with $\mathbb{P}[\mathbf{G}_2^c] \leq c_2 e^{-c(\log N)^2}$ such that the following holds on \mathbf{G}_2 . We have $|m - \Psi| \leq T(\log N)^2$ for any $N^{-1}\zeta$ -localization center $m \in \llbracket N_1, N_2 \rrbracket$ of λ with respect to \mathbf{P} . Applying this for $m \in \{\Phi, \varphi\}$ then yields $|\varphi - \Phi| \leq 2T(\log N)^2 \leq T(\log N)^3$ for sufficiently large N ; this confirms the second statement of the proposition. \square

7. PROPERTIES OF LOCALIZATION CENTERS

We establish Proposition 2.9 in Section 7.1 and Proposition 2.10 in Section 7.3, after showing several estimates on spacings between particles in the Toda lattice in Section 7.2. Throughout this section, we adopt Assumption 2.8.

7.1. Proof of Proposition 2.9. In this section we show Proposition 2.9 by proving the following generalization of it. Recall we adopt Assumption 2.8 throughout.

Proposition 7.1. *Assume more generally that $\zeta \geq N^3 e^{-200(\log N)^{3/2}}$. For any real number $\mathfrak{d} > 0$, there exists a constant $c = c(\mathfrak{d}) > 0$ such that the following holds with probability at least $1 - c^{-1} e^{-c(\log N)^2}$. Fix real numbers $t, \tilde{t} \in [0, T]$; an eigenvalue $\lambda \in \text{eig } \mathbf{L}$; and $N^{-1}\zeta$ -localization centers $\varphi, \tilde{\varphi} \in \llbracket N_1, N_2 \rrbracket$ of λ with respect to $\mathbf{L}(t)$ and $\mathbf{L}(\tilde{t})$, respectively. If φ satisfies (2.10) and $|t - \tilde{t}| \leq e^{-\mathfrak{d}(\log N)^2}$ holds, then $|\varphi - \tilde{\varphi}| \leq (\log N)^3$.*

Proof. We may assume in what follows that $T \geq 3$ and that \mathfrak{d} is sufficiently small, in a way to be determined later. We first restrict to several events.

Let $\mathcal{T} \subseteq [0, T]$ denote an $e^{-\mathfrak{d}(\log N)^2}$ -mesh of $[0, T]$, and fix $s \in \mathcal{T}$. By Proposition 4.4, there exists a constant $c_1 > 0$; an event $\mathbf{E}_1(s)$ with $\mathbb{P}[\mathbf{E}_1(s)^c] \leq c_1^{-1} e^{-c_1(\log N)^2}$; and a random matrix $\mathbf{M}(s) = [M_{ij}(s)] \in \text{SymMat}_{\llbracket N_1, N_2 \rrbracket}$ with the same law as $\mathbf{L}(0)$, such that the following holds on $\mathbf{E}_1(s)$. Denoting $N'_1 = N_1 + T(\log N)^{5/2}$ and $N'_2 = N_2 - T(\log N)^{5/2}$, we have

$$(7.1) \quad \max_{i,j \in \llbracket N'_1, N'_2 \rrbracket} |M_{ij}(s) - L_{ij}(s)| \leq e^{-(\log N)^{5/2}/5}.$$

Corollary 5.6 (with the $(\mathbf{L}, \tilde{\mathbf{L}}; \mathcal{D}; \delta; \tilde{\lambda})$ there equal to $(\mathbf{M}, \mathbf{L}(s); \llbracket N_1, N_2 \rrbracket \setminus \llbracket N'_1, N'_2 \rrbracket; e^{-(\log N)^{5/2}/5}; \lambda)$ here) therefore yields a constant $c_2 > 0$ and an event $\mathbf{E}_2(s)$ for each $s \in \mathcal{T}$, with $\mathbb{P}[\mathbf{E}_2(s)^c] \leq c_2^{-1} e^{-c_2(\log N)^2}$, such that the following holds on $\mathbf{E}_2(s)$. There exists a unique eigenvalue $\mu = \mu(s) \in \text{eig } \mathbf{M}(s)$ so that, for any $N^{-2}\zeta$ -localization center $\varphi_s \in \llbracket N'_1 + (\log N)^3, N'_2 - (\log N)^3 \rrbracket$ for λ with respect to $\mathbf{L}(s)$ and any $N^{-3}\zeta$ -localization center $\psi_s \in \llbracket N_1, N_2 \rrbracket$ for $\mu(s)$ with respect to $\mathbf{M}(s)$, we have $|\varphi_s - \psi_s| \leq (\log N)^2$.

Set $\mathfrak{d}' = \mathfrak{d}/8$. Recalling Definition 3.13, let $\mathbf{E}_3 = \bigcap_{r \geq 0} \text{BND}_{\mathbf{L}(r)}(\log N) \cap \text{SEP}_{\mathbf{L}(0)}(e^{-\mathfrak{d}'(\log N)^2})$, so Lemma 3.15 and Lemma 3.18 give a constant $c_3 = c_3(\mathfrak{d}) > 0$ with $\mathbb{P}[\mathbf{E}_3^c] \leq c_3^{-1} e^{-c_3(\log N)^2}$. Further let \mathbf{E}_4 denote the event on which Lemma 5.2 holds, which satisfies $\mathbb{P}[\mathbf{E}_4^c] \leq c_4^{-1} e^{-c_4(\log N)^2}$ for some $c_4 > 0$. We restrict to the event $\mathbf{E} = \bigcap_{s \in \mathcal{T}} (\mathbf{E}_1(s) \cap \mathbf{E}_2(s)) \cap \mathbf{E}_3 \cap \mathbf{E}_4$ in what follows, which for $\mathfrak{d} < \min\{c_1/2, c_2/2\}$ satisfies $\mathbb{P}[\mathbf{E}^c] \leq c_5^{-1} e^{-c_5(\log N)^2}$ for some $c_5 > 0$, by a union bound.

By Lemma 5.2 (and our restriction to \mathbf{E}_4), we have for any $N^{-1}\zeta$ -localization center φ_0 for λ with respect to $\mathbf{L}(0)$ that $|\varphi - \varphi_0| \leq T(\log N)^2$ and $|\tilde{\varphi} - \varphi_0| \leq T(\log N)^2$. Therefore, $|\varphi - \tilde{\varphi}| \leq 2T(\log N)^2$, so (2.10) implies that

$$(7.2) \quad N_1 + \frac{T}{2} \cdot (\log N)^3 \leq \tilde{\varphi} \leq N_2 - \frac{T}{2} \cdot (\log N)^3.$$

Now, fix $t \in [0, T]$, and let $s \in \mathcal{T}$ satisfy $|s - t| \leq e^{-\mathfrak{d}(\log N)^2}$, so $|s - t'| \leq 2e^{-\mathfrak{d}(\log N)^2}$. We will first apply Lemma 3.7, with the $(\mathbf{A}; \mathbf{B})$ there equal to $(\mathbf{L}(t); \mathbf{L}(s))$ here, to show that φ and $\tilde{\varphi}$ are localization centers for λ with respect to $\mathbf{L}(s)$. Then, we will apply our restriction to $\mathbf{E}_2(s)$ to deduce that φ and $\tilde{\varphi}$ are close to a localization center of $\mu(s)$ with respect to $\mathbf{M}(s)$, and thus are close to each other.

To implement this, we require the third estimate in (3.5). So, for any $r \in [0, T]$ and $z \in \mathbb{C}$, denote the resolvent $G(z; r) = [G_{ij}(z; r)] = (\mathbf{L}(r) - z)^{-1} \in \text{Mat}_{[[N_1, N_2]]}$. Observe that

$$\max_{i, j \in [[N_1, N_2]]} |L_{ij}(s) - L_{ij}(t)| \leq 2e^{-\mathfrak{d}(\log N)^2} \cdot \max_{i, j \in [[N_1, N_2]]} |L'_{ij}(t)| \leq e^{-7\mathfrak{d}'(\log N)^2},$$

where in the second inequality we used (2.1), (2.4), Definition 2.1, and our restriction to \mathbf{E}_3 (which together imply that $|L'_{ij}(t)| \leq 2(\log N)^2$). With (3.3) and (3.4), this implies that

$$\sup_{z \in \Omega} \max_{i, j \in [[N_1, N_2]]} |G_{ij}(z; t) - G_{ij}(z; s)| \leq N^2 \cdot e^{4\mathfrak{d}'(\log N)^2} \cdot e^{-7\mathfrak{d}'(\log N)^2} \leq e^{-2\mathfrak{d}'(\log N)^2},$$

where we have denoted $\Omega = \{z \in \mathbb{C} : e^{-2\mathfrak{d}'(\log N)^2} \leq \text{Im } z \leq 1\}$.

Now we apply Lemma 3.7, with the parameters $(\mathbf{A}, \mathbf{B}; \lambda; \varphi; \eta, \delta)$ there equal to the parameters $(\mathbf{L}(t), \mathbf{L}(s); \lambda; \varphi; e^{-2\mathfrak{d}'(\log N)^2}, e^{-2\mathfrak{d}'(\log N)^2})$ here. This yields an eigenvalue $\lambda' \in \text{eig } \mathbf{L}(s) = \text{eig } \mathbf{L}(t)$ such that $|\lambda - \lambda'| \leq 3N^2\zeta^{-1}e^{-2\mathfrak{d}'(\log N)^2} < e^{-\mathfrak{d}'(\log N)^2}$ and φ is a $N^{-2}\zeta$ -localization center for λ' with respect to $\mathbf{L}(s)$. Due to our restriction to $\mathbf{E}_3 \subseteq \text{SEP}_{\mathbf{L}(s)}(e^{-\mathfrak{d}'(\log N)^2})$, we have that $\lambda = \lambda'$, meaning that φ is an $N^{-2}\zeta$ -localization center for λ with respect to $\mathbf{L}(s)$. By similar reasoning, $\tilde{\varphi}$ is an $N^{-2}\zeta$ -localization center for λ with respect to $\mathbf{L}(s)$.

By our restriction to $\mathbf{E}_2(s)$, and the fact from (2.10) and (7.2) that $\varphi, \tilde{\varphi} \in [[N'_1 + (\log N)^3, N'_2 - (\log N)^3]]$, it follows that for any $N^{-3}\zeta$ -localization center $\psi_s \in [[N_1, N_2]]$ of $\mu(s)$ with respect to $\mathbf{M}(s)$ we have $|\varphi - \psi_s| \leq (\log N)^2$ and $|\tilde{\varphi} - \psi_s| \leq (\log N)^2$. Hence, $|\varphi - \tilde{\varphi}| \leq 2(\log N)^2$, confirming the proposition. \square

Proof of Proposition 2.9. This follows from the $\tilde{t} = t$ case of Proposition 7.1. \square

7.2. Spacing Bounds for the Toda Particles. In this section we prove the following lemma approximating the distances between the Toda particles $q_j(s)$ under thermal equilibrium. Recall we adopt Assumption 2.8 throughout.

Lemma 7.2. *There exists a constant $c > 0$ such that the following two statements hold with probability at least $1 - c^{-1}e^{-c(\log N)^2}$.*

(1) *For any $s \in [0, T]$ and $i, j \in [[N_1 + T(\log N)^3, N_2 - T(\log N)^3]]$, we have*

$$(7.3) \quad |q_i(s) - q_j(s) - \alpha(i - j)| \leq |i - j|^{1/2}(\log N)^2.$$

(2) *For any $s \in [0, T]$ and $i, j \in [[N_1, N_2]]$ with $|i - j| \geq T(\log N)^5$, we have*

$$(7.4) \quad (q_i(s) - q_j(s)) \cdot \text{sgn}(\alpha i - \alpha j) \geq \frac{|\alpha|}{2} \cdot |i - j|.$$

Proof. Observe that it suffices to show for any fixed $s \in [-T, T]$ that the below two statements hold with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. First, for any $i, j \in [[N_1 + T(\log N)^3, N_2 - T(\log N)^3]]$,

$$(7.5) \quad |q_i(s) - q_j(s) - \alpha(i - j)| \leq \frac{1}{2} \cdot |i - j|^{1/2}(\log N)^2.$$

Second, for any $i \in \llbracket N_1, N_2 \rrbracket$ with $|i - j| \geq T(\log N)^5$,

$$(7.6) \quad (q_i(s) - q_j(s)) \cdot \text{sgn}(\alpha i - \alpha j) \geq \frac{3|\alpha|}{4} \cdot |i - j|.$$

Indeed, if this were true then applying a union bound, we may restrict to the event \mathbf{E}_1 on which (7.5) and (7.6) hold for all $s \in \mathcal{T}$, where $\mathcal{T} \subset [-T, T]$ is an N^{-20} -mesh of $[-T, T]$. Recalling Definition 3.13, we may further by Lemma 3.15 restrict to the event $\mathbf{E}_2 = \bigcap_{s \geq 0} \text{BND}_{\mathbf{L}(s)}(\log N)$. In view of (2.1), (2.3), Definition 2.1, and our restriction to \mathbf{E}_2 , we have $|q'_k(s)| \leq \log N$ for all $s \geq 0$ and $k \in \llbracket N_1, N_2 \rrbracket$. Then, the two statements of the lemma hold for any $s \in [-T, T]$. Indeed, letting $s' \in \mathcal{T}$ be such that $|s - s'| \leq N^{-20}$, we have from (7.5) and the above bound on $|q'_k(s)|$ that

$$\begin{aligned} |q_i(s) - q_j(s) - \alpha(i - j)| &\leq |q_i(s') - q_j(s') - \alpha(i - j)| + 2N^{-20} \log N \\ &\leq \frac{1}{2} \cdot |i - j|^{1/2} (\log N)^2 + 2N^{-20} \log N \leq |i - j|^{1/2} (\log N)^2, \end{aligned}$$

which confirms (7.3). The verification of (7.4) from (7.6) is entirely analogous and thus omitted.

Hence, it remains to show (7.5) and (7.6). Throughout the remainder of this proof, we set $K = T(\log N)^3$. By Proposition 4.4, there is a family of random variables $\tilde{\mathbf{a}} = (\tilde{a}_{N_1}, \tilde{a}_{N_1+1}, \dots, \tilde{a}_{N_2-1}) \in \mathbb{R}^{N-1}$ and $\tilde{\mathbf{b}} = (\tilde{b}_{N_1}, \tilde{b}_{N_1+1}, \dots, \tilde{b}_{N_2}) \in \mathbb{R}^N$, such that $(\tilde{\mathbf{a}}; \tilde{\mathbf{b}})$ has the same law as $(\mathbf{a}; \mathbf{b})$ and the following holds. There exists a constant $c_1 > 0$ and an event \mathbf{E}_1 with $\mathbb{P}[\mathbf{E}_1^c] < c_1^{-1} e^{-c_1(\log N)^2}$ so that, on \mathbf{E}_1 , we have

$$(7.7) \quad \max_{i \in \llbracket N_1+K, N_2-K \rrbracket} |a_i(s) - \tilde{a}_i| + \max_{i \in \llbracket N_1+K, N_2-K \rrbracket} |b_i(s) - \tilde{b}_i| \leq 2e^{-(\log N)^3/5}.$$

We restrict to \mathbf{E}_1 in what follows and let $(\tilde{\mathbf{p}}; \tilde{\mathbf{q}}) \in \mathbb{R}^N \times \mathbb{R}^N$ denote the Toda state space initial data associated with $(\tilde{\mathbf{a}}; \tilde{\mathbf{b}})$, as described in Section 2.1.1. Further define event

$$\mathbf{E}_2 = \bigcap_{i, i' \in \llbracket N_1, N_2 \rrbracket} \left\{ |q_i(0) - q_{i'}(0) - \alpha(i - i')| + |\tilde{q}_i - \tilde{q}_{i'} - \alpha(i - i')| \leq \frac{1}{4} \cdot |i - i'|^{1/2} (\log N)^2 \right\},$$

observing that $\mathbb{P}[\mathbf{E}_2^c] \leq c_2^{-1} e^{-c_2(\log N)^2}$ for some constant $c_2 > 0$, by Lemma 3.12. Recalling Definition 3.13, we also define the event

$$\mathbf{E}_3 = \bigcap_{r \geq 0} \text{BND}_{\mathbf{L}(r)}(\log N) \cap \bigcap_{i=N_1}^{N_2} \{a_i(0) \geq e^{-(\log N)^2}\} \cap \{\tilde{a}_i \geq e^{-(\log N)^2}\},$$

which satisfies $\mathbb{P}[\mathbf{E}_3^c] \leq c_3^{-1} e^{-c_3(\log N)^2}$ for some constant $c_3 > 0$, by Lemma 3.15 and the explicit form of the density $\mu_{\beta, \theta; N-1, N}$ for $\mathbf{a}(0)$ (from Definition 2.4). Denoting the event $\mathbf{E} = \mathbf{E}_1 \cap \mathbf{E}_2 \cap \mathbf{E}_3$, we restrict to \mathbf{E} in what follows; it then suffices to show that (7.5) and (7.6) hold.

To that end, first observe that, for any $i, i' \in \llbracket N_1 + K, N_2 - K \rrbracket$, we have

$$\begin{aligned} |q_i(s) - q_{i'}(s) - (\tilde{q}_i - \tilde{q}_{i'})| &\leq 2 \sum_{k=i}^{i'-1} |\log a_i(s) - \log \tilde{a}_i| \\ &\leq 2N \cdot \max_{k \in \llbracket i, i' \rrbracket} |a_i(s) - \tilde{a}_i| \cdot (|a_i(s)|^{-1} + |\tilde{a}_i|^{-1}) \leq N^{-1}, \end{aligned}$$

where in the first bound we used (2.5), and in the second and third we used (7.7) and our restriction to $\mathbf{E}_2 \cap \mathbf{E}_3$. This, together with our restriction to the event \mathbf{E}_2 , establishes (7.5) and thus the first

statement of the lemma. To show the second, we suppose that $\alpha > 0$ and $i \geq j + T(\log N)^5$, as the proof when $\alpha < 0$ or $i \leq j - T(\log N)^5$ is entirely analogous. Then, observe

$$\begin{aligned} q_i(s) - q_j(s) &\geq q_i(0) - q_j(0) - |q_i(s) - q_i(0)| - |q_j(s) - q_j(0)| \\ &\geq \alpha(i - j) - |i - j|^{1/2}(\log N)^2 - 2s \cdot \max_{|s'| \leq s} |b_i(s')| \\ &\geq \frac{4\alpha}{5} \cdot (i - j) - 2T \log N \geq \frac{\alpha}{2} \cdot (i - j), \end{aligned}$$

where in the first and second statements we used our restriction to $\mathbf{E}_2 \cap \mathbf{E}_3$, the first equality in (2.1), and the fact that $p_i(t) = b_i(t)$ by (2.3); in the third we used the fact that $i - j \geq T(\log N)^5$ and our restriction to \mathbf{E}_2 ; and in the fourth we again used the fact that $i - j \geq T(\log N)^5$. This confirms (7.6) and thus the second statement of the lemma when $i \geq j + T(\log N)^5$; since (as mentioned above) the proof when $i \leq j - T(\log N)^5$ is entirely analogous, this establishes the lemma. \square

7.3. Proof of Proposition 2.10. In this section we establish Proposition 2.10. We begin with the following lemma proving a variant of (2.12), when the second sum appearing there is over i such that $\varphi_t(i)$ (as opposed to $q_i(t)$ or $Q_i(t)$) is in a prescribed interval. Recall we adopt Assumption 2.8 throughout.

Proposition 7.3. *For any integer $m \geq 0$, there exists a constant $c = c(m) > 0$ such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. Let $N'_1, N'_2 \in \llbracket N_1, N_2 \rrbracket$ be indices satisfying*

$$(7.8) \quad N_1 + T(\log N)^5 + N^{1/100} \leq N'_1 \leq N'_2 \leq N_2 - T(\log N)^5 - N^{1/100}.$$

Then, the following two statements hold for any $t \in [0, T]$.

(1) *We have*

$$(7.9) \quad \left| \sum_{i=N'_1}^{N'_2} \mathfrak{k}_i^{[m]}(t) - \sum_{i:\varphi_t(i) \in \llbracket N'_1, N'_2 \rrbracket} \lambda_i^m \right| \leq 12(\log N)^{m+3}.$$

(2) *For any $k \in \llbracket N'_1, N'_2 \rrbracket$, we have*

$$(7.10) \quad \left| \sum_{i < k} \mathfrak{k}_i^{[m]}(t) - \sum_{i:\varphi_t(i) \leq k} \lambda_i^m \right| \leq 12(\log N)^{m+3}.$$

Proof. The proofs of (7.9) and (7.10) are very similar, so we only focus on the former. We may assume in what follows that (N'_1, N'_2) is fixed, by a union bound. Let us first establish (7.9) for any deterministic $t \in [0, T]$, and then we will show it holds for all $t \in [0, T]$.

So, fix $t \in [0, T]$. We first apply Proposition 4.4 to deduce the existence of a constant $c_1 > 0$; an event \mathbf{E}_1 with $\mathbb{P}[\mathbf{E}_1^c] \leq c_1^{-1}e^{-c_1(\log N)^2}$; and a random matrix $\mathbf{P} = [P_{ij}] \in \text{SymMat}_{\llbracket N_1, N_2 \rrbracket}$ with the same law as $\mathbf{L}(0)$ such that, on \mathbf{E}_1 , we have

$$(7.11) \quad \max_{i,j \in \llbracket N'_1 - m, N'_2 + m \rrbracket} |P_{ij} - L_{ij}(t)| \leq e^{-(\log N)^3}.$$

where we used the fact that $\llbracket N'_1 - m, N'_2 + m \rrbracket \subseteq \llbracket N_1 + T(\log N)^4, N_2 - T(\log N)^4 \rrbracket$. Further set $N' = N'_2 - N'_1 + 1$ and define the $N' \times N'$ symmetric matrix $\mathbf{Q} = [Q_{ij}] \in \text{SymMat}_{\llbracket N_1, N_2 \rrbracket}$ by setting $Q_{ij} = P_{ij}$ whenever $i, j \in \llbracket N'_1, N'_2 \rrbracket$. Set $\text{eig } \mathbf{Q} = (\mu_1, \mu_2, \dots, \mu_{N'})$, and let $\psi : \llbracket 1, N' \rrbracket \rightarrow \llbracket N'_1, N'_2 \rrbracket$ denote a ζ -localization center bijection for \mathbf{Q} . Recalling Definition 3.13, we define the event $\mathbf{E}_2 =$

$\text{BND}_{\mathbf{L}(t)}(\log N) \cap \text{BND}_{\mathbf{P}}(\log N) \cap \text{BND}_{\mathbf{Q}}(\log N) \cap \text{SEP}_{\mathbf{L}(0)}(e^{-(\log N)^2})$, which by Lemma 3.15 and Lemma 3.18 satisfies $\mathbb{P}[\mathbf{E}_2^c] \leq c_2^{-1} e^{-c_2(\log N)^2}$, for some $c_2 > 0$. We restrict to $\mathbf{E}_1 \cap \mathbf{E}_2$ in the below.

We will proceed by comparing the first sum on the left side of (7.9) to $\text{Tr } \mathbf{Q} = \sum_{\mu \in \text{eig } \mathbf{Q}} \mu^m$; use Corollary 5.6 to approximate the eigenvalues of \mathbf{Q} by those of $\mathbf{L}(t)$; and compare the resulting expression to the second sum on the left side of (7.9). To implement the first task, observe that

$$(7.12) \quad \left| \sum_{i=N'_1}^{N'_2} t_i^{[m]}(t) - \text{Tr } \mathbf{Q}^m \right| = \left| \sum_{i=N'_1}^{N'_2} [\mathbf{L}(t)^m]_{ii} - \text{Tr } \mathbf{Q}^m \right| \\ \leq \left| \sum_{i=N'_1}^{N'_2} [\mathbf{P}^m]_{ii} - \text{Tr } \mathbf{Q}^m \right| + N e^{-(\log N)^3} m (\log N)^m,$$

where we have denoted the (i, j) entry of any matrix \mathbf{M} by $[\mathbf{M}]_{ij}$; here, the first statement holds by Definition 2.3 and the second by (7.11) and our restriction to \mathbf{E}_2 . Next, since \mathbf{P} and \mathbf{Q} are tridiagonal and satisfy $P_{ij} = Q_{ij}$ whenever $i, j \in \llbracket N'_1, N'_2 \rrbracket$, observe that $[\mathbf{P}^m]_{ii} = [\mathbf{Q}^m]_{ii}$ for each $i \in \llbracket N'_1 + m, N'_2 - m \rrbracket$. We further have by our restriction to \mathbf{E}_2 that each entry of \mathbf{P} and \mathbf{Q} is bounded by $\log N$, meaning (again since \mathbf{P} and \mathbf{Q} are tridiagonal) that each entry of \mathbf{P}^m and \mathbf{Q}^m is bounded by $(3 \log N)^m$. It follows that

$$(7.13) \quad \left| \sum_{i=N'_1}^{N'_2} [\mathbf{P}^m]_{ii} - \text{Tr } \mathbf{Q}^m \right| \leq \sum_{\substack{i \in \llbracket N'_1, N'_2 \rrbracket \\ i \notin \llbracket N'_1 + m, N'_2 - m \rrbracket}} (|(\mathbf{P}^m)_{ii}| + |(\mathbf{Q}^m)_{ii}|) \leq 4m(3 \log N)^m.$$

Next, denote $N''_1 = N'_1 + (\log N)^3$ and $N''_2 = N'_2 - (\log N)^3$. Then,

$$(7.14) \quad \left| \text{Tr } \mathbf{Q}^m - \sum_{j: \psi(j) \in \llbracket N''_1, N''_2 \rrbracket} \mu_j^m \right| \leq 2(\log N)^3 \cdot \max_{\mu \in \text{eig } \mathbf{Q}} |\mu|^m \leq 2(\log N)^{m+3},$$

where in the first bound we used that $\text{Tr } \mathbf{Q}^m = \sum_{\mu \in \text{eig } \mathbf{Q}} \mu^m$ and $|\llbracket N'_1, N'_2 \rrbracket \setminus \llbracket N''_1, N''_2 \rrbracket| \leq 2(\log N)^3$, and in the second we used the fact that $|\mu| \leq \log N$ for any $\mu \in \text{eig } \mathbf{Q}$ (by our restriction to \mathbf{E}_2).

Now, we apply Proposition 7.1 and Corollary 5.5, the latter with the $(\mathbf{L}; \tilde{\mathbf{L}}; \delta; \mathcal{D})$ there equal to $(\mathbf{Q}; \mathbf{L}(t); e^{-(\log N)^3}; \llbracket N_1, N_2 \rrbracket \setminus \llbracket N'_1, N'_2 \rrbracket)$ here. This by (7.11) yields a constant $c_2 > 0$ and an event \mathbf{E}_3 with $\mathbb{P}[\mathbf{E}_3^c] \leq c_3^{-1} e^{-c_3(\log N)^2}$, such that the following holds on \mathbf{E}_3 . There exists a function $\kappa: \llbracket 1, N' \rrbracket \rightarrow \llbracket 1, N \rrbracket$ such that, for each $j \in \llbracket 1, N' \rrbracket$ with $\psi(j) \in \llbracket N''_1, N''_2 \rrbracket$,

$$(7.15) \quad |\mu_j - \lambda_{\kappa(j)}| \leq e^{-c_2(\log N)^3}, \quad \text{and} \quad |\psi(j) - \varphi_t(\kappa(j))| \leq (\log N)^3.$$

The second statement in (7.15) holds since Corollary 5.5 implies that $\psi(j)$ is an $N^{-1}\zeta$ -localization center of $\lambda_{\kappa(j)}$ with respect to $\mathbf{L}(t)$; therefore, since $\varphi_t(\kappa(j))$ is as well, Proposition 7.1 implies (as $\psi(j) \in \llbracket N''_1, N''_2 \rrbracket$) that $|\psi(j) - \varphi_t(\kappa(j))| \leq (\log N)^3$. Further observe that, due our restriction to \mathbf{E}_2 , we have $|\lambda_i - \lambda_{i'}| \geq e^{-(\log N)^2} \geq 2e^{-c_2(\log N)^3}$ for any distinct $i, i' \in \llbracket 1, N \rrbracket$; thus, $\kappa(j) \neq \kappa(j')$

for any distinct $j, j' \in \llbracket 1, N' \rrbracket$ with $\psi(j), \psi(j') \in \llbracket N'_1, N'_2 \rrbracket$. Therefore,

$$(7.16) \quad \left| \sum_{j: \psi(j) \in \llbracket N'_1, N'_2 \rrbracket} \mu_j^m - \sum_{j: \varphi_t(j) \in \llbracket N'_1, N'_2 \rrbracket} \lambda_j^m \right| \leq \sum_{j: \psi(j) \in \llbracket N'_1, N'_2 \rrbracket} |\mu_j^m - \lambda_{\kappa(j)}^m| + 2(\log N)^3 \cdot \max_{\lambda \in \text{eig } \mathbf{L}(t)} |\lambda|^m \leq 4(\log N)^{m+3},$$

where the first statement holds by changing variables from j to $\kappa(j)$ in the second sum there whenever $\psi(j) \in \llbracket N'_1, N'_2 \rrbracket$, and using (7.15) with the facts that $\llbracket N'_1 - (\log N)^3, N_2 + (\log N)^3 \rrbracket = \llbracket N'_1, N'_2 \rrbracket$ and that $|\llbracket N'_1, N'_2 \rrbracket \setminus \llbracket N'_1, N'_2 \rrbracket| \leq 2(\log N)^3$; the second follows from (7.15) with the fact that $|\lambda| \leq \log N$ for each $\lambda \in \text{eig } \mathbf{L}(t)$ (as we restricted to \mathbf{E}_2). Summing (7.12), (7.13), (7.14), and (7.16), we deduce that

$$(7.17) \quad \left| \sum_{i=N'_1}^{N'_2} \mathfrak{k}_i^{[m]}(t) - \sum_{i: \varphi_t(i) \in \llbracket N'_1, N'_2 \rrbracket} \lambda_i^m \right| \leq 7(\log N)^{m+3}.$$

This verifies that (7.9) holds with high probability for a fixed $t \in [0, T]$; it remains to show it holds with high probability for all $t \in [0, T]$ simultaneously. To that end, for any $t \in [0, T]$, let $\mathbf{E}(t)$ denote the event on which (7.17) holds. We then have that $\mathbb{P}[\mathbf{E}(t)^c] \leq c_3^{-1} e^{-c_3(\log N)^2}$ for some constant $c_3 > 0$. Denoting $\mathfrak{c} = c_3/2$, it follows from Proposition 7.1 that there exists a constant $c_4 > 0$ and an event \mathbf{E}_4 with $\mathbb{P}[\mathbf{E}_4^c] \leq c_4^{-1} e^{-c_4(\log N)^2}$ such that the following holds on \mathbf{E}_4 . For any $s, s' \in [0, T]$ with $|s - s'| \leq e^{-\mathfrak{c}(\log N)^2}$, we have $|\varphi_s(i) - \varphi_{s'}(i)| \leq (\log N)^3$, whenever $\varphi = \varphi_s(i)$ satisfies (2.10). Letting \mathcal{T} denote an $e^{-\mathfrak{c}(\log N)^2}$ -mesh of $[0, T]$, we restrict to the event $\bigcap_{s \in \mathcal{T}} \mathbf{E}(s) \cap \mathbf{E}_4 \cap \bigcap_{r \geq 0} \text{BND}_{\mathbf{L}(r)}(\log N)$, which we may by a union bound (and Lemma 3.15).

Now let $t \in [0, T]$ be arbitrary, and let $s \in \mathcal{T}$ satisfy $|t - s| \leq e^{-\mathfrak{c}(\log N)^2}$. Then, we claim that

$$(7.18) \quad \left| \sum_{i=N'_1}^{N'_2} |\mathfrak{k}_i^{[m]}(t) - \mathfrak{k}_i^{[m]}(s)| \right| \leq 1; \quad \left| \sum_{i: \varphi_t(i) \in \llbracket N'_1, N'_2 \rrbracket} \lambda_i^m - \sum_{i: \varphi_s(i) \in \llbracket N'_1, N'_2 \rrbracket} \lambda_i^m \right| \leq 4(\log N)^{m+3}.$$

which together with (7.17) (with the t there equal to s here) would imply the proposition.

To verify the first bound in (7.18), observe for any $i, j \in \llbracket N_1, N_2 \rrbracket$ that

$$|L_{ij}(t) - L_{ij}(s)| \leq |s - t| \cdot \sup_{r \in [s, t]} |L'_{ij}(r)| \leq e^{-\mathfrak{c}(\log N)^2} \cdot 2(\log N)^2 \leq e^{-\mathfrak{c}(\log N)^2/2},$$

where in the second inequality we used Definition 2.1, (2.3), (2.4), and our restriction to the event $\bigcap_{r \geq 0} \text{BND}_{\mathbf{L}(r)}(\log N)$. Since the same event implies that each entry of $\mathbf{L}(s)$ and $\mathbf{L}(t)$ is bounded by $\log N$, and both of these matrices are tridiagonal, it follows for any $i \in \llbracket N_1, N_2 \rrbracket$ that

$$\begin{aligned} |\mathfrak{k}_i^{[m]}(t) - \mathfrak{k}_i^{[m]}(s)| &= |[\mathbf{L}(t)^m]_{ii} - [\mathbf{L}(s)^m]_{ii}| \leq \max_{i, j \in \llbracket N_1, N_2 \rrbracket} |L_{ij}(s) - L_{ij}(t)| \cdot m(3 \log N)^m \\ &\leq e^{-\mathfrak{c}(\log N)^2/2} \cdot m(3 \log N)^m \leq N^{-1}. \end{aligned}$$

Summing over $i \in \llbracket N'_1, N'_2 \rrbracket$ confirms the first bound in (7.18). The second bound in (7.18) follows from the fact that $|\lambda| \leq \log N$ for any $\lambda \in \text{eig } \mathbf{L}(t)$ (by our restriction to $\text{BND}_{\mathbf{L}(t)}(\log N)$), with the fact that there are at most $4(\log N)^3$ indices i for which $\varphi_s(i) \in \llbracket N'_1, N'_2 \rrbracket$ and $\varphi_t(i) \notin \llbracket N'_1, N'_2 \rrbracket$ or for which $\varphi_s(i) \notin \llbracket N'_1, N'_2 \rrbracket$ and $\varphi_t(i) \in \llbracket N'_1, N'_2 \rrbracket$ (as $|\varphi_t(i) - \varphi_s(i)| \leq (\log N)^3$, by our restriction to \mathbf{E}_4). This proves (7.18) and thus (7.9).

As mentioned previously, the proof of (7.10) is entirely analogous, and is obtained by replacing the application of Proposition 4.4 and Corollary 5.6 above with Proposition 6.3 (with the ℓ there equal to k here); we omit further details. \square

We can now prove Proposition 2.10.

Proof of Proposition 2.10. We assume that $\alpha > 0$ in what follows, as the proof when $\alpha < 0$ is entirely analogous. The proofs of (2.12) and (2.13) are very similar, so we only focus on the former. Recalling Definition 3.13, define the event $\mathbf{E}_1 = \bigcap_{s \in [0, t]} \mathbf{BND}_{\mathbf{L}(s)}(\log N)$. By Lemma 3.15, there exists a constant $c_1 > 0$ such that $\mathbb{P}[\mathbf{E}_1^c] \leq c_1^{-1} e^{-c_1 (\log N)^2}$, so we restrict to \mathbf{E}_1 in what follows. We further restrict to the event \mathbf{E}_2 on which Lemma 7.2 holds, with the T there both equal to 0 and T here (we may take the T there to be 0 by the $R = (\log N)^2 |i - j|^{1/2}$ case of Lemma 3.12).

Fix $t \in [0, T]$, and let $N'_1 = \min\{i : q_i(t) \in \mathcal{J}\}$ and $N'_2 = \max\{i : q_i(t) \in \mathcal{J}\}$. We claim that (N'_1, N'_2) satisfy (7.8). To verify this, assume to the contrary, so that either $N'_1 < N_1 + T(\log N)^5 + N^{1/100}$ or $N'_2 > N_2 - T(\log N)^5 - N^{1/100}$ holds. The analysis of these two cases is entirely analogous, so we assume the former. Then,

$$\begin{aligned} q_{N'_1}(0) - q_0(0) &\geq q_{N'_1}(t) - T \log N \geq \alpha N_1 + (T + |N_1|^{1/2})(\log N)^5 - T \log N \\ &\geq \alpha N_1 + T(\log N)^5 + \frac{1}{2} \cdot |N_1|^{1/2} (\log N)^5, \end{aligned}$$

where the first statement holds since $q_0(0) = 0$ and $|q'_i(t)| = |p_i(t)| \leq \log N$ (by (2.1), (2.3), Definition 2.1, and our restriction to \mathbf{E}_1); the second holds by (2.11) and the inclusion $q_{N'_1}(t) \in \mathcal{J}$; and the third holds for N sufficiently large, by (2.8). This contradicts (7.3) (at $s = T = 0$), confirming (7.8).

Therefore, Proposition 7.3 applies and yields a constant $c_2 > 0$ and an event \mathbf{E}_3 with $\mathbb{P}[\mathbf{E}_3^c] \leq c_2^{-1} e^{-c_2 (\log N)^2}$, such that (7.9) holds on \mathbf{E}_3 ; we restrict to \mathbf{E}_3 in the below. Then, since $q_i(t) \in \mathcal{J}$ implies that $i \in \llbracket N'_1, N'_2 \rrbracket$, it suffices (by (2.9)) to show that

$$\sum_{i \in \llbracket N'_1, N'_2 \rrbracket : q_i(t) \notin \mathcal{J}} |\mathfrak{k}_i^{[m]}(t)| + \sum_{i : \varphi_t(i) \in \llbracket N'_1, N'_2 \rrbracket, q_{\varphi_t(i)}(t) \notin \mathcal{J}} |\lambda_i|^m \leq (3 \log N)^{m+5}.$$

Changing variables in the second sum from i to $\varphi_t^{-1}(i)$, this is equivalent to

$$\sum_{i \in \llbracket N'_1, N'_2 \rrbracket : q_i(t) \notin \mathcal{J}} |\mathfrak{k}_i^{[m]}(t)| + \sum_{i \in \llbracket N'_1, N'_2 \rrbracket : q_i(t) \notin \mathcal{J}} |\lambda_{\varphi_t^{-1}(i)}|^m \leq (3 \log N)^{m+5}.$$

Using the facts that each entry of $\mathbf{L}(t)^m$ is bounded by $(3 \log N)^m$ (as $\mathbf{L}(t)$ is tridiagonal and each of its entries is bounded by $\log N$) and that $|\lambda| \leq \log N$ for each $\lambda \in \text{eig } \mathbf{L}(t)$, it remains to verify

$$(7.19) \quad \#\{i \in \llbracket N'_1, N'_2 \rrbracket : q_i(t) \notin \mathcal{J}\} \leq 2(\log N)^{9/2}.$$

To show this, observe that if $i \in \llbracket N'_1, N'_2 \rrbracket$ satisfies $q_i(t) \notin \mathcal{J}$, then either $q_i(t) < q_{N'_1}(t)$ or $q_i(t) > q_{N'_2}(t)$. In the former case, we have

$$0 > q_i(t) - q_{N'_1}(t) \geq \alpha \cdot (i - N'_1) - |i - N'_1|^{1/2} \cdot (\log N)^2,$$

where the last inequality holds by our restriction to \mathbf{E}_2 . Hence $i < N'_1 + (\log N)^{9/2}$, and there are at most $(\log N)^{9/2}$ such indices in $\llbracket N'_1, N'_2 \rrbracket$. By similar reasoning, there are at most $(\log N)^{9/2}$ indices $i \in \llbracket N'_1, N'_2 \rrbracket$ such that $q_i(t) > q_{N'_2}(t)$. Summing these bounds yields (7.19) and thus (2.12).

As mentioned previously, the proof of (2.13) is entirely analogous (using (7.10) in place of (7.9)) upon observing that

$$\int_0^s j_k^{[m]}(s) ds = \sum_{i \leq k-1} \mathfrak{k}_i^{[m]}(0) - \sum_{i \leq k-1} \mathfrak{k}_i^{[m]}(t),$$

which holds by (2.6); integrating it over $s \in [0, t]$; and summing it over $i \in \llbracket N_1, k-1 \rrbracket$. \square

8. PROOF OF THE ASYMPTOTIC SCATTERING RELATION

8.1. Proof of Theorem 2.11. In this section we prove Theorem 2.11, which will follow from the following variant of it that replaces the $Q_t(i)$ -dependent sums in (2.15) with ones that depend on $\varphi_t(i)$. The latter will be shown in Section 8.3 below. We adopt Assumption 2.8 throughout.

Theorem 8.1. *There exists a constant $c > 0$ such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. Let $k \in \llbracket 1, N \rrbracket$ satisfy (2.14). Then, for any $t \in [0, T]$, we have*

$$(8.1) \quad \left| \lambda_k t - Q_k(t) + Q_k(0) - 2 \sum_{i: \varphi_t(i) < \varphi_t(k)} \log |\lambda_k - \lambda_i| + 2 \sum_{i: \varphi_0(i) < \varphi_0(k)} \log |\lambda_k - \lambda_i| \right| \leq 2(\log N)^{12}.$$

Proof of Theorem 2.11. Throughout this proof, we assume that $\alpha > 0$, as the proof when $\alpha < 0$ is entirely analogous; we also fix $k \in \llbracket 1, N \rrbracket$ satisfying (2.14). We begin by restricting to several events. Recalling Definition 3.13, first restrict to the event $\mathbf{E}_1 = \text{BND}_{\mathbf{L}(0)}(\log N) \cap \text{SEP}_{\mathbf{L}(0)}(e^{-(\log N)^2})$, which we may by Lemma 3.15, Lemma 3.18, and a union bound. Further restrict to the event \mathbf{E}_2 on which Theorem 8.1 holds; to the event \mathbf{E}_3 on which Lemma 7.2 holds; and to the event \mathbf{E}_4 on which Lemma 5.2 holds.

In view of (8.1) (and our restriction to \mathbf{E}_2), we must show that

$$2 \sum_{i=N_1}^{N_2} \left| (\mathbb{1}_{\varphi_t(i) < \varphi_t(k)} - \mathbb{1}_{\varphi_0(i) < \varphi_0(k)}) - (\mathbb{1}_{Q_i(t) < Q_k(t)} - \mathbb{1}_{Q_i(0) < Q_k(0)}) \right| \cdot \left| \log |\lambda_k - \lambda_i| \right| \leq (\log N)^{15} - 2(\log N)^{12}.$$

Since by our restriction to \mathbf{E}_1 we that $|\log |\lambda_k - \lambda_i|| \leq (\log N)^2$ for each $i \neq k$, it therefore suffices to show that

$$(8.2) \quad \sum_{i=N_1}^{N_2} \left| (\mathbb{1}_{\varphi_t(i) < \varphi_t(k)} - \mathbb{1}_{\varphi_0(i) < \varphi_0(k)}) - (\mathbb{1}_{Q_i(t) < Q_k(t)} - \mathbb{1}_{Q_i(0) < Q_k(0)}) \right| \leq (\log N)^8.$$

We first claim that the summand on the left side of (8.2) is equal to 0 if i satisfies $|\varphi_0(i) - \varphi_0(k)| > 2T(\log N)^5$. Indeed, fix such an index i , and assume $\varphi_0(i) < \varphi_0(k) - 2T(\log N)^5$, as the verification in the alternative case is entirely analogous. Then, (5.1) (with our restriction to \mathbf{E}_4) gives $|\varphi_t(i) - \varphi_0(i)| \leq T(\log N)^2$ and $|\varphi_t(k) - \varphi_0(k)| \leq T(\log N)^2$, so for sufficiently large N we have $\varphi_t(i) < \varphi_t(k) - T(\log N)^5$. As such, $\mathbb{1}_{\varphi_t(i) < \varphi_t(k)} = 1 = \mathbb{1}_{\varphi_0(i) < \varphi_0(k)}$. By (7.4) (and our restriction to \mathbf{E}_3) whose assumption holds by the above bounds, we have for each $s \in \{0, t\}$ that

$$Q_k(s) - Q_i(s) = q_{\varphi_s(k)} - q_{\varphi_s(i)} \geq \frac{\alpha}{2} \cdot (\varphi_s(k) - \varphi_s(i)) > 0.$$

So, it follows that $\mathbb{1}_{Q_i(t) < Q_k(t)} = 1 = \mathbb{1}_{Q_i(0) < Q_k(0)}$, meaning that the summand on the left side of (8.2) associated with i is equal to 0.

Therefore, we may restrict the sum on the left side of (8.2) to indices i satisfying $|\varphi_0(i) - \varphi_0(k)| \leq 2T(\log N)^5$. Fix such an index i ; we claim for each $s \in \{0, t\}$ that

$$(8.3) \quad \mathbb{1}_{\varphi_s(i) < \varphi_s(k)} = \mathbb{1}_{Q_i(s) < Q_k(s)}, \quad \text{unless } |\varphi_s(i) - \varphi_s(k)| \leq (\log N)^5,$$

which would imply (8.2) and thus the theorem. To confirm (8.3), fix $s \in \{0, t\}$, and assume that $|\varphi_s(i) - \varphi_s(k)| > (\log N)^5$; we will further suppose that $\varphi_s(i) < \varphi_s(k) - (\log N)^5$, as the proof in the alternative case is entirely analogous. By our assumptions (2.14) and $|\varphi_0(i) - \varphi_0(k)| \leq 2T(\log N)^5$, observe (by (5.1), using our restriction to \mathbf{E}_4) that $\varphi_s(i), \varphi_s(k) \in \llbracket N_1 + T(\log N)^5, N_2 - T(\log N)^5 \rrbracket$, so (7.3) holds (by our restriction to \mathbf{E}_4) with the (i, j) there equal to $(\varphi_s(i), \varphi_s(k))$ here. Hence,

$$Q_k(s) - Q_i(s) = q_{\varphi_k(s)}(s) - q_{\varphi_i(s)}(s) \geq \alpha \cdot (\varphi_k(s) - \varphi_i(s)) - |\varphi_k(s) - \varphi_i(s)|^{1/2} \cdot (\log N)^2 > 0,$$

where the last bound holds since $\varphi_k(s) - \varphi_i(s) \geq (\log N)^5$. Therefore, $\mathbb{1}_{Q_i(s) < Q_k(s)} = 1 = \mathbb{1}_{\varphi_s(i) < \varphi_s(k)}$, proving (8.3) and thus the theorem. \square

8.2. Proof of Theorem 8.1 if $T^2 \leq N$. In this section we establish the following case of Theorem 8.1, which assumes that $T^2 \leq N$ (and a slightly stronger condition (8.4) on $\varphi_0(k)$ than imposed in (2.14)) and addresses a fixed time $t \in [0, T]$. Recall we adopt Assumption 2.8 throughout.

Proposition 8.2. *Fix $t \in [0, T]$. There exists a constant $c > 0$ such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. Let $k \in \llbracket 1, N \rrbracket$ satisfy*

$$(8.4) \quad N_1 + T(\log N)^6 + N^{1/100} \leq \varphi_0(k) \leq N_2 - T(\log N)^6 - N^{1/100}.$$

If $T^2 \leq N$, then we have

$$(8.5) \quad \left| \lambda_k t - Q_k(t) + Q_k(0) - 2 \sum_{i: \varphi_t(i) < \varphi_t(k)} \log |\lambda_k - \lambda_i| + 2 \sum_{i: \varphi_0(i) < \varphi_0(k)} \log |\lambda_k - \lambda_i| \right| \leq 2(\log N)^{11}.$$

Proposition 8.2 is a quick consequence of the below two lemmas. In what follows, we denote

$$(8.6) \quad \mathfrak{C}(s) = \log \sum_{j=1}^N e^{-\lambda_j s} u_j(N_1; 0)^2.$$

Lemma 8.3. *Fix $t \in [0, T]$. There exists a constant $c > 0$ such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. For any $k \in \llbracket 1, N \rrbracket$ satisfying (8.4), we have*

$$(8.7) \quad \left| Q_k(t) - Q_k(0) + 2 \sum_{i: \varphi_t(i) < \varphi_t(k)} \log |\lambda_k - \lambda_i| - 2 \sum_{i: \varphi_0(i) < \varphi_0(k)} \log |\lambda_k - \lambda_i| - \lambda_k t - \mathfrak{C}(t) - q_{N_1}(t) + q_{N_1}(0) \right| \leq 2(\log N)^6.$$

Lemma 8.4. *Fix $t \in [0, T]$, and assume that $T^2 \leq N$. There exists a constant $c > 0$ such that*

$$\mathbb{P}[|q_{N_1}(0) - q_{N_1}(t) - \mathfrak{C}(t)| \leq (\log N)^{11}] \geq 1 - c^{-1}e^{-c(\log N)^2}.$$

Proof of Proposition 8.2. This follows from Lemma 8.3 and Lemma 8.4. \square

We now show Lemma 8.3 and Lemma 8.4.

Proof of Lemma 8.3. Fix $k \in \llbracket 1, N \rrbracket$ satisfying (8.4), and observe by Lemma 3.6 that

$$(8.8) \quad -2 \log |u_k(N_1; t)| = \lambda_k t - 2 \log |u_k(N_1; 0)| + \mathfrak{C}(t).$$

By Lemma 5.2, there is a constant $c_1 > 0$ and an event \mathbf{E}_1 with $\mathbb{P}[\mathbf{E}_1^c] \leq c_1^{-1} e^{-c_1(\log N)^2}$ so that, on \mathbf{E}_1 , we have $|\varphi_t(k) - \varphi_0(k)| \leq T(\log N)^2$. Observe on \mathbf{E}_1 that $N_1 + T(\log N)^5 + N^{1/100} \leq \varphi_s(k) \leq N_2 - T(\log N)^5 - N^{1/100}$ for each $s \in \{0, t\}$, by (8.4). We restrict to \mathbf{E}_1 in what follows.

By Proposition 6.1, there exists a constant $c_2 > 0$ and an event \mathbf{E}_2 with $\mathbb{P}[\mathbf{E}_2^c] \leq c_2^{-1} e^{-c_2(\log N)^2}$, such that the following holds on \mathbf{E}_2 . For any real number $s \in \{0, t\}$, we have

$$\left| 2 \log |u_k(N_1; s)| - 2 \sum_{i=N_1}^{\varphi_s(k)-1} \log L_{i,i+1}(s) + 2 \sum_{i:\varphi_s(i) < \varphi_s(k)} \log |\lambda_i - \lambda_k| \right| \leq 2(\log N)^6,$$

where we used that $N_1 + T(\log N)^5 + N^{1/100} \leq \varphi_s(k) \leq N_2 - T(\log N)^5 - N^{1/100}$. We further restrict to \mathbf{E}_2 below. Since Definition 2.1, (2.3), and Assumption 2.8 together imply that

$$2 \sum_{i=N_1}^{\varphi_s(k)-1} \log L_{i,i+1}(s) = 2 \sum_{i=N_1}^{\varphi_s(k)-1} \log a_i(s) = q_{N_1}(s) - q_{\varphi_s(k)}(s) = q_{N_1}(s) - Q_k(s),$$

we obtain by our restriction to $\mathbf{E}_1 \cap \mathbf{E}_2$ that, for each $s \in \{0, t\}$,

$$\left| 2 \log |u_k(N_1; s)| + Q_k(s) - q_{N_1}(s) + 2 \sum_{i:\varphi_s(i) < \varphi_s(k)} \log |\lambda_i - \lambda_k| \right| \leq (\log N)^6.$$

Subtracting this bound at $s = 0$ from it at $s = t$, and applying (8.8), then yields the lemma. \square

Proof of Lemma 8.4. We will establish this proposition by averaging an estimate on the left side of (8.7) over all $k \in \llbracket N_1, N_2 \rrbracket$. To that end, let \mathbf{E}_1 denote the event on which (8.7) holds for each $k \in \llbracket N_1, N_2 \rrbracket$ satisfying (8.4). By Lemma 8.3, there exists a constant $c_1 > 0$ such that $\mathbb{P}[\mathbf{E}_1^c] \leq c_1^{-1} e^{-c_1(\log N)^2}$. We will define additional events on which we can bound the left side of (8.7) for $k \in \llbracket N_1, N_2 \rrbracket$ not necessarily satisfying (8.4). Specifically, recalling Definition 3.13, set

$$\mathbf{E}_2 = \bigcap_{r \geq 0} \text{BND}_{\mathbf{L}(r)}(\log N) \cap \text{SEP}_{\mathbf{L}(0)}(e^{-(\log N)^2}); \quad \mathbf{E}_3 = \bigcap_{k=N_1}^{N_2} \{|\varphi_t(k) - \varphi_0(k)| \leq T(\log N)^2\}.$$

By Lemma 3.15 and Lemma 3.18, there exists a constant $c_2 > 0$ such that $\mathbb{P}[\mathbf{E}_2^c] \leq c_2^{-1} e^{-c_2(\log N)^2}$. Moreover, by Lemma 5.2, there exists a constant $c_3 > 0$ such that $\mathbb{P}[\mathbf{E}_3^c] \leq c_3^{-1} e^{-c_3(\log N)^2}$. We further define the event

$$\mathbf{E}_4 = \bigcap_{|i-j| \leq T(\log N)^2} \{|q_i(0) - q_j(0)| \leq T(\log N)^3\},$$

which by Lemma 3.12 and a union bound satisfies $\mathbb{P}[\mathbf{E}_4^c] \leq c_4^{-1} e^{-c_4(\log N)^2}$ for some constant $c_4 > 0$. Defining the event $\mathbf{E} = \mathbf{E}_1 \cap \mathbf{E}_2 \cap \mathbf{E}_3 \cap \mathbf{E}_4$, we may by a union bound restrict to \mathbf{E} in what follows.

Then, for any $k \in \llbracket N_1, N_2 \rrbracket$, we have

$$\begin{aligned}
(8.9) \quad & \left| Q_k(t) - Q_k(0) - q_{N_1}(t) + q_{N_1}(0) - \lambda_k t - \mathfrak{C}(t) \right. \\
& \left. + 2 \sum_{i: \varphi_t(i) < \varphi_t(k)} \log |\lambda_i - \lambda_k| - 2 \sum_{i: \varphi_0(i) < \varphi_0(k)} \log |\lambda_i - \lambda_k| \right| \\
& \leq |q_{\varphi_t(k)}(0) - q_{\varphi_0(k)}(0)| + |q_{\varphi_t(k)}(t) - q_{\varphi_t(k)}(0)| + |q_{N_1}(t) - q_{N_1}(0)| + T \log N + |\mathfrak{C}(t)| \\
& \quad + 2(\log N)^2 \cdot \#\{i \in \llbracket N_1, N_2 \rrbracket : \varphi_t(i) < \varphi_t(k), \varphi_0(i) > \varphi_0(k)\} \\
& \quad + 2(\log N)^2 \cdot \#\{i \in \llbracket N_1, N_2 \rrbracket : \varphi_t(i) > \varphi_t(k), \varphi_0(i) < \varphi_0(k)\},
\end{aligned}$$

where we used the definition $Q_k(s) = q_{\varphi_s(k)}(s)$ (from (2.9)) and the facts that $|\lambda_k| \leq \log N$ and that $|\log |\lambda_i - \lambda_k|| \leq (\log N)^2$ (both by our restriction to \mathbf{E}_2). We next bound the terms on the right side of (8.9). Observe by our restriction to \mathbf{E}_4 that

$$(8.10) \quad |q_{\varphi_t(k)}(0) - q_{\varphi_0(k)}(0)| \leq T(\log N)^3,$$

since $|\varphi_t(k) - \varphi_0(k)| \leq T(\log N)^2$, by our restriction to \mathbf{E}_3 . Additionally,

$$(8.11) \quad \max_{j \in \llbracket N_1, N_2 \rrbracket} |q_j(t) - q_j(0)| \leq \int_0^t |b_j(s)| ds \leq T \log N,$$

where in the first inequality we used (2.1) and (2.3); in the second we used the fact that $t \in [0, T]$ and our restriction to \mathbf{E}_2 . Moreover,

$$(8.12) \quad -2T \log N \leq -t \max_{\lambda \in \text{eig } \mathbf{L}} |\lambda| - \log N \leq \mathfrak{C}(t) \leq t \max_{\lambda \in \text{eig } \mathbf{L}} |\lambda| + \log N \leq 2T \log N,$$

where the first and fourth inequalities follow from our restriction to \mathbf{E}_2 ; the second from the definition (8.6) of $\mathfrak{C}(t)$, with the fact that there exists at least one index $j \in \llbracket 1, N \rrbracket$ for which $u_j(N_1; 0)^2 \geq N^{-1}$; and the third from the definition (8.6) of $\mathfrak{C}(t)$, with the fact that $u_j(N_1; 0)^2 \leq 1$ for each $j \in \llbracket 1, N \rrbracket$. We further have that

$$\begin{aligned}
(8.13) \quad & \#\{i \in \llbracket N_1, N_2 \rrbracket : \varphi_t(i) < \varphi_t(k), \varphi_0(i) > \varphi_0(k)\} \leq 2T(\log N)^2; \\
& \#\{i \in \llbracket N_1, N_2 \rrbracket : \varphi_t(i) > \varphi_t(k), \varphi_0(i) < \varphi_0(k)\} \leq 2T(\log N)^2.
\end{aligned}$$

Indeed, to verify the first bound in (8.13), observe (by our restriction to the event \mathbf{E}_3) that if $\varphi_t(i) < \varphi_t(k) - 2T(\log N)^2$ then $\varphi_0(i) \leq \varphi_t(i) + T(\log N)^2 < \varphi_t(k) - T(\log N)^2 \leq \varphi_0(k)$. In particular, we can only have $\varphi_t(i) < \varphi_t(k)$ and $\varphi_0(i) > \varphi_0(k)$ if $\varphi_t(i) \in \llbracket \varphi_t(k) - 2T(\log N)^2, \varphi_t(k) - 1 \rrbracket$, meaning that there are at most $2T(\log N)^2$ such indices i . This shows the first statement in (8.13); the second is confirmed entirely analogously.

Inserting (8.10), (8.11), (8.12), and (8.13) into (8.9), we obtain for any $k \in \llbracket N_1, N_2 \rrbracket$ that

$$\begin{aligned}
& \left| Q_k(t) - Q_k(0) - q_{N_1}(t) + q_{N_1}(0) - \lambda_k t - \mathfrak{C}(t) \right. \\
& \left. + 2 \sum_{i: \varphi_t(i) < \varphi_t(k)} \log |\lambda_i - \lambda_k| - 2 \sum_{i: \varphi_0(i) < \varphi_0(k)} \log |\lambda_i - \lambda_k| \right| \\
& \leq T(\log N)^3 + 5T \log N + 8T(\log N)^4 \leq 9T(\log N)^4.
\end{aligned}$$

Summing this inequality over $k \in \llbracket N_1, N_2 \rrbracket$ with $\varphi_0(k) \notin \llbracket N_1 + T(\log N)^6 + N^{1/100}, N_2 - T(\log N)^6 - N^{1/100} \rrbracket$ with (8.7) over $k \in \llbracket N_1, N_2 \rrbracket$ with $\varphi_0 \in \llbracket N_1 + T(\log N)^6 + N^{1/100}, N_2 - T(\log N)^6 - N^{1/100} \rrbracket$, we obtain

$$(8.14) \quad \left| \sum_{k=N_1}^{N_2} (q_k(t) - q_k(0)) - t \sum_{j=1}^N \lambda_j + 2 \sum_{k=N_1}^{N_2} \sum_{i: \varphi_t(i) < \varphi_t(k)} \log |\lambda_i - \lambda_k| - 2 \sum_{k=N_1}^{N_2} \sum_{i: \varphi_0(i) < \varphi_0(k)} \log |\lambda_i - \lambda_k| + N \cdot (q_{N_1}(0) - q_{N_1}(t) - \mathfrak{C}(t)) \right| \leq 2N(\log N)^6 + 18T^2(\log N)^{10} + 18TN^{1/100}(\log N)^4.$$

The difference between of the third and fourth terms on the left side of (8.14) is equal to 0, since both are equal to $\sum_{i \neq j} \log |\lambda_i - \lambda_j|$. The difference between the first and the second is also equal to 0, as

$$\sum_{k=N_1}^{N_2} (q_k(t) - q_k(0)) = \sum_{k=N_1}^{N_2} \int_0^t p_k(s) ds = \int_0^t \text{Tr } \mathbf{L}(s) ds = t \cdot \text{Tr } \mathbf{L}(0) = t \sum_{j=1}^N \lambda_j,$$

where the first statement follows from (2.1); the second from (2.3) and Definition 2.1; and the third and fourth from Lemma 2.2. It follows from (8.14) that

$$|q_{N_1}(0) - q_{N_1}(t) - \mathfrak{C}(t)| \leq 2(\log N)^6 + 18T^2N^{-1}(\log N)^{10} + 18TN^{-99/100}(\log N)^4 \leq (\log N)^{11},$$

where we used the fact that $T^2 \leq N$. This establishes the lemma. \square

8.3. Proof of Theorem 8.1. In this section we establish Theorem 8.1. We first prove the following variant of it that fixes the time $t \in [0, T]$. Recall we adopt Assumption 2.8 throughout.

Theorem 8.5. *Fix $t \in [0, T]$. There exists a constant $c > 0$ such that the following holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$. For any $k \in \llbracket 1, N \rrbracket$ satisfying (2.14), we have*

$$(8.15) \quad \left| \lambda_k t - Q_k(t) + Q_k(0) - 2 \sum_{i: \varphi_t(i) < \varphi_t(k)} \log |\lambda_k - \lambda_i| + 2 \sum_{i: \varphi_0(i) < \varphi_0(k)} \log |\lambda_k - \lambda_i| \right| \leq (\log N)^{12}.$$

Proof. By Proposition 8.2, (8.5) holds with probability at least $1 - c^{-1}e^{-c(\log N)^2}$ if $T^2 \leq N$. It suffices to show such a bound continues to hold for larger values of $T \leq N(\log N)^{-7}$ (and to weaken the constraint (8.4) on k to (2.14)).

To that end, we first apply (8.5) on a Toda lattice on a larger interval (at thermal equilibrium), and then use comparison estimates (such as Corollary 5.6 and Lemma 4.5) to approximate the original Toda lattice by the enlarged one. To implement this, first let $\tilde{N}_1 \leq \tilde{N}_2$ be integers satisfying

$$(8.16) \quad \tilde{N}_1 + T^2N \leq N_1 \leq N_2 \leq \tilde{N}_2 - T^2N, \quad \text{and} \quad \tilde{N} \leq N^5,$$

where $\tilde{N} = \tilde{N}_2 - \tilde{N}_1 + 1$. Let $(\tilde{\mathbf{a}}(s); \tilde{\mathbf{b}}(s)) \in \mathbb{R}^{\tilde{N}} \times \mathbb{R}^{\tilde{N}}$ denote the Flaschka variables for a Toda lattice on $\llbracket \tilde{N}_1, \tilde{N}_2 \rrbracket$; letting $\tilde{\mathbf{a}}(s) = (\tilde{a}_{\tilde{N}_1}(s), \tilde{a}_{\tilde{N}_1+1}(s), \dots, \tilde{a}_{\tilde{N}_2}(s))$ and $\tilde{\mathbf{b}}(s) = (\tilde{b}_{\tilde{N}_1}(s), \tilde{b}_{\tilde{N}_1+1}(s), \dots, \tilde{b}_{\tilde{N}_2}(s))$, they satisfy $\tilde{a}_{\tilde{N}_2}(s) = 0$, and (2.4) holds for each $(j, t) \in \llbracket \tilde{N}_1, \tilde{N}_2 \rrbracket \times \mathbb{R}$. We sample the initial data $(\tilde{\mathbf{a}}(0); \tilde{\mathbf{b}}(0))$ according to the thermal equilibrium $\mu_{\beta, \theta; \tilde{N}-1, \tilde{N}}$ of Definition 2.4; we couple $(\tilde{\mathbf{a}}(0); \tilde{\mathbf{b}}(0))$ with $(\mathbf{a}(0); \mathbf{b}(0))$ so that $(\tilde{a}_i(0), \tilde{b}_i(0)) = (a_i(0), b_i(0))$ for all $i \in \llbracket N_1, N_2 - 1 \rrbracket$.

For any $s \in \mathbb{R}$, denote the Lax matrix associated with $(\tilde{\mathbf{a}}(s); \tilde{\mathbf{b}}(s))$ (as in Definition 2.1) by $\tilde{\mathbf{L}}(s) = [\tilde{L}_{ij}(s)] \in \text{SymMat}_{\tilde{N} \times \tilde{N}}$. Set $\text{eig } \tilde{\mathbf{L}}(s) = (\tilde{\lambda}_1, \tilde{\lambda}_2, \dots, \tilde{\lambda}_{\tilde{N}})$, which does not depend on s (by Lemma 2.2). For each $s \in \mathbb{R}_{\geq 0}$, let $\tilde{\varphi}_s : \llbracket 1, \tilde{N} \rrbracket \rightarrow \llbracket \tilde{N}_1, \tilde{N}_2 \rrbracket$ denote a ζ -localization center bijection for $\tilde{\mathbf{L}}(s)$. Further let $(\tilde{\mathbf{p}}(s); \tilde{\mathbf{q}}(s)) \in \mathbb{R}^{\tilde{N}} \times \mathbb{R}^{\tilde{N}}$ denote the Toda state space variables associated with $(\tilde{\mathbf{a}}(s); \tilde{\mathbf{b}}(s))$, as in Section 2.1.1, where we have indexed $\tilde{\mathbf{p}}(s) = (\tilde{p}_{\tilde{N}_1}(s), \tilde{p}_{\tilde{N}_1+1}(s), \dots, \tilde{p}_{\tilde{N}_2}(s))$ and $\tilde{\mathbf{q}}(s) = (\tilde{q}_{\tilde{N}_1}(s), \tilde{q}_{\tilde{N}_1+1}(s), \dots, \tilde{q}_{\tilde{N}_2}(s))$. For each $s \in \mathbb{R}$ and $i \in \llbracket 1, \tilde{N} \rrbracket$, denote $\tilde{Q}_i(s) = \tilde{q}_{\tilde{\varphi}_s(i)}(s)$.

By (8.5) and the fact that $T^2 \leq \tilde{N} \leq N^5$, there exists a constant $c_1 > 0$ and an event \mathbf{E}_1 with $\mathbb{P}[\mathbf{E}_1^c] \leq c_1^{-1} e^{-c_1(\log N)^2}$ such that on \mathbf{E}_1 we have

$$(8.17) \quad \left| \tilde{\lambda}_m t - \tilde{Q}_m(t) + \tilde{Q}_m(0) - 2 \sum_{i: \tilde{\varphi}_t(i) < \tilde{\varphi}_t(m)} \log |\tilde{\lambda}_m - \tilde{\lambda}_i| + 2 \sum_{i: \tilde{\varphi}_0(i) < \tilde{\varphi}_0(m)} \log |\tilde{\lambda}_m - \tilde{\lambda}_i| \right| \leq 5^{12} (\log N)^{11},$$

for any $m \in \llbracket 1, \tilde{N} \rrbracket$ satisfying

$$(8.18) \quad \tilde{N}_1 + T(\log \tilde{N})^6 + \tilde{N}^{1/100} \leq \tilde{\varphi}_0(m) \leq \tilde{N}_2 - T(\log \tilde{N})^6 - \tilde{N}^{1/100}.$$

In what follows, we restrict to \mathbf{E}_1 . We must therefore approximate the parameters $(\tilde{Q}_j(s), \tilde{\lambda}_j, \tilde{\varphi}_s(j))$ by $(Q_j(s), \lambda_j, \varphi_s(j))$, which we will do using Lemma 4.5 and Corollary 5.6 (with the fact that $\mathbf{q}(0)$ and $\tilde{\mathbf{q}}(0)$ coincide near the origin).

To do so, we restrict to several additional events. Recalling Definition 3.13, restrict to the event

$$\mathbf{E}_2 = \text{SEP}_{\mathbf{L}(0)}(e^{-(\log N)^2}) \cap \text{SEP}_{\tilde{\mathbf{L}}(0)}(e^{-(\log N)^2}) \cap \bigcap_{r \geq 0} \text{BND}_{\mathbf{L}(r)}\left(\frac{\log N}{1600}\right) \cap \text{BND}_{\tilde{\mathbf{L}}(r)}\left(\frac{\log N}{1600}\right),$$

which we may by Lemma 3.15 and Lemma 3.18. Further restrict to the event \mathbf{E}_3 on which Lemma 7.2 holds, with the \mathbf{q} there equal to both \mathbf{q} here and $\tilde{\mathbf{q}}$ here. Additionally restrict to the event \mathbf{E}_4 on which Lemma 5.2 holds, with the $(\mathbf{L}; \varphi_j)$ there equal to both $(\mathbf{L}(0); \varphi_0(j))$ and $(\tilde{\mathbf{L}}(0); \tilde{\varphi}_0(j))$ here, and on which Proposition 2.9 holds, with the $\mathbf{L}(t)$ there equal to both $\mathbf{L}(t)$ and $\tilde{\mathbf{L}}(t)$ here.

Now, setting $K = T(\log N)^3/2$, we have by (the $A = \log N/400$ case of) Lemma 4.5 that

$$(8.19) \quad \sup_{t \in [0, T]} \max_{i \in \llbracket N_1+K, N_2-K \rrbracket} (|a_i(t) - \tilde{a}_i(t)| + |b_i(t) - \tilde{b}_i(t)|) \leq e^{-(\log N)^3/10}.$$

Next, by Proposition 4.4, there is a constant $c_1 > 0$; random matrices $\mathbf{M} = [M_{ij}] \in \text{SymMat}_{\llbracket N_1, N_2 \rrbracket}$ and $\tilde{\mathbf{M}} = [\tilde{M}_{ij}] \in \text{SymMat}_{\llbracket \tilde{N}_1, \tilde{N}_2 \rrbracket}$ with the same laws as $\mathbf{L}(0)$ and $\tilde{\mathbf{L}}(0)$, respectively; and an event \mathbf{E}_5 with $\mathbb{P}[\mathbf{E}_5^c] \leq c_1^{-1} e^{-c_1(\log N)^2}$, such that on \mathbf{E}_5 we have

$$(8.20) \quad \max_{i, j \in \llbracket N_1+K, N_2-K \rrbracket} |M_{ij} - L_{ij}(t)| \leq e^{-(\log N)^3/10}, \quad \max_{i, j \in \llbracket N_1+K, N_2-K \rrbracket} |\tilde{M}_{ij} - \tilde{L}_{ij}(t)| \leq e^{-(\log N)^3/10}.$$

Restricting to \mathbf{E}_5 , we may therefore (by (8.19) and (8.20)) further restrict to the event \mathbf{E}_6 on which Corollary 5.5 and Corollary 5.6 hold, with the $(\delta; \mathcal{D})$ there equal to $(2e^{-(\log N)^3/10}; \llbracket \tilde{N}_1, \tilde{N}_2 \rrbracket \setminus \llbracket N_1+K, N_2-K \rrbracket)$ here, and the $(\mathbf{L}, \tilde{\mathbf{L}})$ equal to any of $(\mathbf{L}(0), \tilde{\mathbf{L}}(0))$, $(\mathbf{M}, \mathbf{L}(t))$, $(\tilde{\mathbf{M}}, \tilde{\mathbf{L}}(t))$, and $(\tilde{\mathbf{M}}, \mathbf{M})$ here (where we view \mathbf{M} and $\mathbf{L}(0)$ as $\tilde{N} \times \tilde{N}$ matrices by setting $M_{ij} = L_{ij}(0) = 0$ if $(i, j) \in \llbracket \tilde{N}_1, \tilde{N}_2 \rrbracket \setminus \llbracket N_1, N_2 \rrbracket$).

Now, let $i \in \llbracket 1, N \rrbracket$ be any index satisfying

$$(8.21) \quad N_1 + 3T(\log \tilde{N})^3 \leq \varphi_0(i) \leq N_2 - 3T(\log \tilde{N})^3.$$

Our restriction to E_4 implies from (5.1) that $N_1 + 2T(\log \tilde{N})^3 \leq \varphi_t(i) \leq N_2 - 2T(\log \tilde{N})^3$. Recalling that $T \geq N^{1/2} \geq 250$ (as otherwise the theorem follows from Proposition 8.2), this will enable us to use our restriction to E_6 to apply Corollary 5.6 twice, first with the $(\mathbf{L}, \tilde{\mathbf{L}})$ there equal to $(\mathbf{M}; \mathbf{L}(t))$ here and then with them equal to $(\tilde{\mathbf{M}}; \mathbf{M})$, and also to apply Corollary 5.5 with the $(\mathbf{L}; \tilde{\mathbf{L}})$ there equal to $(\tilde{\mathbf{M}}; \tilde{\mathbf{L}}(t))$. The first yields a constant $c_2 > 0$ and an eigenvalue $\mu \in \text{eig } \mathbf{M}$ such that $|\mu - \lambda_i| \leq c_2^{-1} e^{-c_2(\log N)^3}$ and $\varphi_t(i)$ is an $N^{-1}\zeta$ -localization center for μ with respect to \mathbf{M} . The second yields an eigenvalue $\tilde{\mu} \in \text{eig } \tilde{\mathbf{M}}$ such that $|\mu - \tilde{\mu}| \leq c_2^{-1} e^{-c_2(\log N)^3}$ and $\varphi_t(i)$ is an $N^{-2}\zeta$ -localization center for $\tilde{\mu}$ with respect to $\tilde{\mathbf{M}}$. The third yields an index $\sigma(i) \in \llbracket 1, \tilde{N} \rrbracket$ such that $|\tilde{\mu} - \tilde{\lambda}_{\sigma(i)}| \leq c_2^{-1} e^{-c_2(\log N)^3}$ and $\varphi_t(i)$ is an $N^{-3}\zeta$ -localization center of $\tilde{\lambda}_{\sigma(i)}$ with respect to $\tilde{\mathbf{L}}(t)$. By Proposition 2.9 (and our restriction to E_4), we have $|\varphi_t(i) - \tilde{\varphi}_t(\sigma(i))| \leq (\log \tilde{N})^3$, and hence

$$(8.22) \quad \begin{aligned} |\lambda_i - \tilde{\lambda}_{\sigma(i)}| &\leq 3c_2^{-1} e^{-c_2(\log N)^3}; & |\varphi_t(i) - \tilde{\varphi}_t(\sigma(i))| &\leq (\log \tilde{N})^3; \\ & & |\varphi_0(i) - \tilde{\varphi}_0(\sigma(i))| &\leq (\log \tilde{N})^3, \end{aligned}$$

where the last bound follows again from similar reasoning to the second (taken at $t = 0$). Observe by our restriction to $E_2 \subseteq \text{SEP}_{\tilde{\mathbf{L}}(t)}(e^{-(\log N)^2})$ and the first inequality in (8.22) that the map from i to $\sigma(i)$ is injective.

Then, by our restriction to E_1 on which (8.17) holds (taking the m there to be $\sigma(k)$, which satisfies (8.18) by (2.14) and (8.22)), and the fact that $\llbracket N_1 + (\log N)^6, N_2 - (\log N)^6 \rrbracket \subseteq \llbracket N_1 + 3(\log \tilde{N})^3, N_2 - 3(\log \tilde{N})^3 \rrbracket \subseteq \llbracket \tilde{N}_1 + T(\log \tilde{N})^6 + \tilde{N}^{1/100}, \tilde{N}_2 - T(\log \tilde{N})^6 - \tilde{N}^{1/100} \rrbracket$ (by (8.16)), it suffices to show that

$$(8.23) \quad |\lambda_k - \tilde{\lambda}_{\sigma(k)}| \leq T^{-1}; \quad |Q_k(0) - \tilde{Q}_{\sigma(k)}(0)| \leq (\log N)^5; \quad |Q_k(t) - \tilde{Q}_{\sigma(k)}(t)| \leq (\log N)^5,$$

and

$$(8.24) \quad \left| \sum_{i=\tilde{N}_1}^{\tilde{N}_2} (\mathbb{1}_{\tilde{\varphi}_t(i) < \tilde{\varphi}_t(\sigma(k))} - \mathbb{1}_{\tilde{\varphi}_0(i) < \tilde{\varphi}_0(\sigma(k))}) \cdot \log |\tilde{\lambda}_{\sigma(k)} - \tilde{\lambda}_i| \right. \\ \left. - \sum_{i=N_1}^{N_2} (\mathbb{1}_{\varphi_t(i) < \varphi_t(k)} - \mathbb{1}_{\varphi_0(i) < \varphi_0(k)}) \cdot \log |\lambda_k - \lambda_i| \right| \leq (\log N)^9.$$

Taking $i = k$ in the first statement of (8.22) (using (2.14) to verify (8.21)) yields the first bound in (8.23). The proofs of the second and third are entirely analogous to each other, so we only verify the third. To do so, observe by (2.1), (2.3), and (8.19) that, for any $i \in \llbracket N_1 + K, N_2 - K \rrbracket$,

$$(8.25) \quad |q_i(t) - \tilde{q}_i(t)| \leq \int_0^t |b_i(s) - \tilde{b}_i(s)| ds \leq 2T \cdot e^{-(\log N)^3/10} \leq 1,$$

as $q_i(0) = \tilde{q}_i(0)$. Additionally, (7.3) implies for $i, j \in \llbracket N_1 + T(\log N)^3, N_2 - T(\log N)^3 \rrbracket$ that

$$(8.26) \quad |q_i(t) - q_j(t)| \leq \alpha \cdot |i - j| + |i - j|^{1/2} \cdot (\log N)^2,$$

by our restriction to E_3 . Combining this with (8.25), (8.22), and the definition (2.9) of Q_k yields

$$\begin{aligned} |Q_k(t) - \tilde{Q}_{\sigma(k)}(t)| &\leq |q_{\varphi_t(k)}(t) - q_{\tilde{\varphi}_t(\sigma(k))}(t)| + |\tilde{q}_{\varphi_t(k)}(t) - \tilde{q}_{\tilde{\varphi}_t(\sigma(k))}(t)| \\ &\leq \alpha \cdot |\varphi_t(k) - \tilde{\varphi}_t(\sigma(k))| + |\varphi_t(k) - \tilde{\varphi}_t(\sigma(k))|^{1/2} \cdot (\log N)^2 + 1 \leq (\log N)^5, \end{aligned}$$

which confirms the third bound in (8.23).

It therefore remains to verify (8.24). To that end, let

$$\begin{aligned}\mathcal{S} &= \{i \in \llbracket 1, N \rrbracket : \varphi_0(i) \in \llbracket N_1 + 2T(\log N)^6, N_2 - 2T(\log N)^6 \rrbracket\}; \\ \tilde{\mathcal{S}} &= \{i \in \llbracket 1, \tilde{N} \rrbracket : \tilde{\varphi}_0(i) \in \llbracket N_1 + 2T(\log N)^6, N_2 - 2T(\log N)^6 \rrbracket\},\end{aligned}$$

and $\sigma(\mathcal{S}) = \{\sigma(i) : i \in \mathcal{S}\}$. First suppose that $i \in \llbracket 1, N \rrbracket \setminus \mathcal{S}$. Then, since (5.1) (and our restriction to \mathbf{E}_4) implies that $|\varphi_t(i) - \varphi_0(i)| \leq T(\log N)^2$, we have from (2.14) that $\mathbb{1}_{\varphi_t(i) < \varphi_t(k)} = \mathbb{1}_{\varphi_0(i) < \varphi_0(k)}$. Hence, we may restrict the second sum in (8.24) to $i \in \mathcal{S}$. Similarly, we may restrict the first sum in (8.24) to $i \in \tilde{\mathcal{S}}$. Therefore, we have

$$(8.27) \quad \left| \sum_{i=\tilde{N}_1}^{\tilde{N}_2} (\mathbb{1}_{\tilde{\varphi}_t(i) < \tilde{\varphi}_t(\sigma(k))} - \mathbb{1}_{\tilde{\varphi}_0(i) < \tilde{\varphi}_0(\sigma(k))}) \cdot \log |\tilde{\lambda}_{\sigma(k)} - \tilde{\lambda}_i| - \sum_{i=N_1}^{N_2} (\mathbb{1}_{\varphi_t(i) < \varphi_t(k)} - \mathbb{1}_{\varphi_0(i) < \varphi_0(k)}) \cdot \log |\lambda_k - \lambda_i| \right| \leq A + B + C.$$

where

$$\begin{aligned}A &= \sum_{i \in \mathcal{S}} |\log |\tilde{\lambda}_{\sigma(k)} - \tilde{\lambda}_{\sigma(i)}| - \log |\lambda_k - \lambda_i||; \\ B &= \sum_{i \in \mathcal{S}} (|\log |\tilde{\lambda}_{\sigma(k)} - \tilde{\lambda}_{\sigma(i)}|| + |\log |\lambda_k - \lambda_i||) \\ &\quad \times \left| (\mathbb{1}_{\tilde{\varphi}_t(\sigma(i)) < \tilde{\varphi}_t(\sigma(k))} - \mathbb{1}_{\varphi_0(\sigma(i)) < \varphi_0(\sigma(k))}) - (\mathbb{1}_{\varphi_t(i) < \varphi_t(k)} - \mathbb{1}_{\varphi_0(i) < \varphi_0(k)}) \right|; \\ C &= \#((\tilde{\mathcal{S}} \cup \sigma(\mathcal{S})) \setminus (\tilde{\mathcal{S}} \cap \sigma(\mathcal{S}))) \cdot \max_{i \neq \sigma(k)} |\log |\tilde{\lambda}_{\sigma(k)} - \tilde{\lambda}_i||.\end{aligned}$$

By our restriction to the event \mathbf{E}_2 , we have

$$(8.28) \quad e^{-(\log N)^2} \leq |\lambda_i - \lambda_j| \leq 2 \log N, \quad \text{and} \quad e^{-(\log N)^2} \leq |\tilde{\lambda}_i - \tilde{\lambda}_j| \leq 2 \log N, \quad \text{whenever } i \neq j.$$

Together with the first bound in (8.22), this implies that $|\log |\tilde{\lambda}_{\sigma(k)} - \tilde{\lambda}_{\sigma(i)}| - \log |\lambda_k - \lambda_i|| \leq e^{-(\log N)^2}$ for each $i \in \mathcal{S}$, and so

$$(8.29) \quad A \leq N e^{-(\log N)^2} \leq 1.$$

By the second and third bounds in (8.22), that are at most $500(\log N)^3$ indices $i \in \mathcal{S}$ for which

$$(\mathbb{1}_{\tilde{\varphi}_t(\sigma(i)) < \tilde{\varphi}_t(\sigma(k))} - \mathbb{1}_{\varphi_0(\sigma(i)) < \varphi_0(\sigma(k))}) - (\mathbb{1}_{\varphi_t(i) < \varphi_t(k)} - \mathbb{1}_{\varphi_0(i) < \varphi_0(k)}) \neq 0.$$

Since this quantity is always bounded above by 2, (8.28) implies that

$$(8.30) \quad B \leq 2000(\log N)^5.$$

By the last inequality in (8.22) and the injectivity of σ , it is quickly verified that $\#((\tilde{\mathcal{S}} \cup \sigma(\mathcal{S})) \setminus (\tilde{\mathcal{S}} \cap \sigma(\mathcal{S}))) \leq 500(\log N)^3$, from which we deduce again by (8.28) that $C \leq 500(\log N)^5$. This, with (8.27), (8.29), and (8.30), yields (8.24) and thus the theorem. \square

Next, establish Theorem 8.1, which proves the bound in Theorem 8.5 uniformly in $t \in [0, T]$.

Proof of Theorem 8.1 (Outline). This theorem will follow from applying Theorem 8.5 on a mesh of times $\mathcal{T} \subset [0, T]$, and using continuity bounds to show that it continues to hold on all of $[0, T]$. Since the proof is similar to the discussions at the end of the proofs of Theorem 2.11 and Theorem 8.5, we only outline it.

Let \mathfrak{c} denote the constant c from Theorem 8.5 and, for any $t \in [0, T]$, let $\mathbf{E}(t)$ denote the event on which (8.15) holds for all $k \in \llbracket 1, N \rrbracket$ satisfying (2.14). Denoting $\mathfrak{c}' = \mathfrak{c}/2$, let $\mathcal{T} \subset [0, T]$ denote a $e^{-\mathfrak{c}'(\log N)^2}$ -mesh of $[0, T]$; by a union bound, we may restrict to the event $\mathbf{F}_1 = \bigcap_{s \in \mathcal{T}} \mathbf{E}(s)$. We further restrict to the event \mathbf{F}_2 on which Proposition 7.1 holds, with the \mathfrak{d} there equal to \mathfrak{c}' here; we additionally restrict to the intersection \mathbf{F}_3 of all of the events from the proof of Theorem 8.5.

Now fix $t \in [0, T]$, and let $s \in \mathcal{S}$ be such that $|s - t| \leq e^{-\mathfrak{c}'(\log N)^2}$. By our restriction to \mathbf{F}_1 , we have from (8.15) that

$$\left| \lambda_k s - Q_k(s) + Q_k(0) - 2 \sum_{i: \varphi_s(i) < \varphi_s(k)} \log |\lambda_k - \lambda_i| + 2 \sum_{i: \varphi_0(i) < \varphi_0(k)} \log |\lambda_k - \lambda_i| \right| \leq (\log N)^{12}.$$

Since $|\lambda_k| \cdot |s - t| \leq \log N \cdot e^{-\mathfrak{c}'(\log N)^2} \leq 1$ and $|\log |\lambda_k - \lambda_i|| \leq (\log N)^2$ for $i \neq k$ (both by our restriction to $\mathbf{BND}_{\mathbf{L}(0)}(\log N) \cap \mathbf{SEP}_{\mathbf{L}(0)}(e^{-(\log N)^2})$, from the event \mathbf{E}_2 in the proof of Theorem 8.5), it suffices to show that

$$(8.31) \quad |Q_k(t) - Q_k(s)| \leq (\log N)^{10}; \quad \sum_{i=N_1}^{N_2} |\mathbb{1}_{\varphi_t(i) < \varphi_t(k)} - \mathbb{1}_{\varphi_s(i) < \varphi_s(k)}| \leq (\log N)^8.$$

The first bound in (8.31) follows from the fact that

$$\begin{aligned} |Q_k(t) - Q_k(s)| &\leq |q_{\varphi_t(k)}(t) - q_{\varphi_s(k)}(t)| + |q_{\varphi_s(k)}(t) - q_{\varphi_s(k)}(s)| \\ &\leq \alpha \cdot |\varphi_t(k) - \varphi_s(k)| + |\varphi_t(k) - \varphi_s(k)|^{1/2} \cdot (\log N)^2 + |s - t| \cdot \max_{r \in [s, t]} |b_{\varphi_s(k)}(s)| \\ &\leq (\log N)^5. \end{aligned}$$

where the first statement follows from the definition (2.9) of Q_k ; the second follows from (8.26), (2.1), and (2.3); and the third follows from the fact that $|\varphi_t(k) - \varphi_s(k)| \leq (\log N)^3$ (by Proposition 7.1, as we restricted to \mathbf{F}_2), Definition 2.1, and our restriction to $\mathbf{F}_3 \subseteq \bigcap_{r \geq 0} \mathbf{BND}_{\mathbf{L}(r)}(\log N)$. The proof of the second bound in (8.31) is entirely analogous to that of (8.2) (using the fact that $|\varphi_t(j) - \varphi_s(j)| \leq (\log N)^3$ for $\varphi_s(j) \in \llbracket N_1 + T(\log N)^3, N_2 - T(\log N)^3 \rrbracket$, by Proposition 7.1 and our restriction to \mathbf{F}_2). This establishes the theorem. \square

APPENDIX A. PROOFS OF RESULTS FROM SECTION 3

A.1. Proofs of Lemma 3.11 and Lemma 3.12.

Proof of Lemma 3.11. Observe that the random variable $\mathfrak{r} \in \mathbb{R}$ has density $\mathbb{P}[\mathfrak{r} \in (r, r + dr)] = \beta^\theta \cdot \Gamma(\theta)^{-1} \cdot e^{-\theta r - \beta e^{-r}} dr$. Thus, denoting

$$(A.1) \quad F(\theta) = \int_{-\infty}^{\infty} e^{-\theta r - \beta e^{-r}} dr,$$

we have

$$(A.2) \quad F'(\theta) = - \int_{-\infty}^{\infty} r e^{-\theta r - \beta e^{-r}} dr = -F(\theta) \cdot \mathbb{E}[\mathfrak{r}].$$

Changing variables $u = \beta e^{-r}$ in (A.1) yields

$$F(\theta) = \beta^{-\theta} \int_0^\infty u^{\theta-1} e^{-u} du = \beta^{-\theta} \cdot \Gamma(\theta),$$

so the lemma follows from (A.2), together with the fact (recall (2.7)) that $\alpha = -F'(\theta) \cdot F(\theta)^{-1}$. \square

Proof of Lemma 3.12. We may assume that $i < j$; in what follows, we abbreviate $q_k = q_k(0)$ and $a_k = a_k(0)$ for each k . By (2.5), we have $q_{m+1} - q_m = -2 \log a_m$ for any $m \in \llbracket i+1, j \rrbracket$. Since $(\mathbf{a}; \mathbf{b})$ is sampled from the measure $\mu_{\beta, \theta; N-1, N}$ from Definition 2.4, all such a_m are mutually independent with the same law \mathbf{a} of density $\mathbb{P}[\mathbf{a} \in (a, a + da)] = 2\beta^\theta \cdot \Gamma(\theta)^{-1} \cdot a^{2\theta-1} e^{-\beta a^2} da$. Hence, denoting $\mathbf{r} = -2 \log \mathbf{a}$, each $q_{m+1} - q_m$ has law \mathbf{r} . By Lemma 3.11, we have $\alpha = \mathbb{E}[\mathbf{r}]$, so we find for any $u \in (0, 1/2)$ that

$$(A.3) \quad \mathbb{P}[q_j - q_i \geq \alpha(j-i) + R] \leq e^{-uR} \cdot \mathbb{E}[e^{u(q_j - q_i)} \cdot e^{\alpha u(j-i)}] = (e^{-uR/(j-i)} \cdot \mathbb{E}[e^{u(\mathbf{r}-\alpha)}])^{j-i},$$

where the first statement follows from a Markov bound, and the second from the mutual independence of the $(q_{m+1} - q_m)$ with the finiteness of the moment generating function $\mathbb{E}[e^{u\mathbf{r}}] = \mathbb{E}[\mathbf{a}^{-2u}] < \infty$ for $u \in (0, 1/2)$. Setting $u = \min\{cR/(j-i), c\}$ for some sufficiently small constant $c > 0$, and using the (quickly verified) fact that $e^{-uR/(j-i)} \cdot \mathbb{E}[e^{u(\mathbf{r}-\alpha)}] \leq e^{-cu^2}$ for this u , yields from (A.3) that

$$\mathbb{P}[q_j - q_i \geq \alpha(j-i) + R] \leq e^{-cu^2|i-j|} \leq e^{-c^3R^2/|j-i|} + e^{-c^3|i-j|}.$$

By similar reasoning, we have $\mathbb{P}[q_j - q_i \leq \alpha(j-i) - R] \leq e^{-c^3R^2/|i-j|} + e^{-c^3|i-j|}$, and so the lemma follows by a union bound. \square

A.2. Proofs of Results From Section 3.2 and Section 3.3.

Proof of Lemma 3.5. Let $\lambda \in \text{eig } \mathbf{L}(t)$ denote the eigenvalue of $\mathbf{L}(t)$ with maximal absolute value; if more than one exists, we select one arbitrarily. Then,

$$(A.4) \quad |\lambda| \leq \max_{i \in \mathcal{J}} \sum_{j \in \mathcal{J}} |L_{ij}(t')| \leq 2(A(t') + B(t')).$$

where the first inequality holds since $\lambda \in \text{eig } \mathbf{L}(t')$ (by Lemma 2.2), and the second inequality holds by Definition 2.1. Moreover,

$$(A.5) \quad |\lambda| \geq \max_{i \in \mathcal{J}} |L_{ii}(t)| = B(t); \quad |\lambda| \geq \frac{1}{2} \cdot \max_{i, j \in \mathcal{J}} |L_{ii}(t) + 2L_{ij}(t) + L_{jj}(t)| \geq A(t) - B(t),$$

where in both the first inequality follows from applying the min-max principle (to $\mathbf{L}(t)$ if $|\lambda| \in \text{eig } \mathbf{L}(t)$, and to $-\mathbf{L}(t)$ if $-|\lambda| \in \text{eig } \mathbf{L}(t)$), and the second inequality follows from Definition 2.1. Combining (A.4) and (A.5) yields $A(t) + B(t) \leq 3|\lambda| \leq 6(A(t') + B(t'))$, confirming the lemma. \square

Proof of Lemma 3.7. Set $\text{eig } \mathbf{A} = (\lambda_1, \lambda_2, \dots, \lambda_n)$ and $\text{eig } \mathbf{B} = (\mu_1, \mu_2, \dots, \mu_n)$. Further let (\mathbf{u}_j) and (\mathbf{v}_j) , for $j \in \llbracket 1, n \rrbracket$, denote orthonormal eigenbases for \mathbf{A} and \mathbf{B} , respectively. In this way, $\mathbf{u}_j = (u_j(i))_{i \in \mathcal{J}}$ is an eigenvector of \mathbf{A} with eigenvalue λ_j , and $\mathbf{v}_j = (v_j(i))_{i \in \mathcal{J}}$ is an eigenvector of \mathbf{B} with eigenvalue μ_j , for each $j \in \llbracket 1, n \rrbracket$. Let $k \in \llbracket 1, n \rrbracket$ be such that $\lambda_k = \lambda$; we may assume that the eigenbasis (\mathbf{u}_j) includes \mathbf{u} , in particular, that $\mathbf{u}_k = \mathbf{u}$. Then,

$$(A.6) \quad \text{Im } G_{\varphi\varphi}(z) = \sum_{j=1}^n u_j(\varphi)^2 \cdot \text{Im}(\lambda_j - z)^{-1} \geq u_k(\varphi)^2 \cdot \text{Im}(\lambda - z)^{-1} \geq \eta^{-1} \zeta^2.$$

where in the first statement we used (3.2); in the second we used the bound $\text{Im}(\lambda_j - z)^{-1} \geq 0$; and in the third we used the fact that $z = \lambda + i\eta$ and the second estimate in (3.5). It follows that

$$(A.7) \quad \left| \sum_{j=1}^n \frac{v_j(\varphi)^2}{\mu_j - z} \right| = |H_{\varphi\varphi}(z)| \geq |G_{\varphi\varphi}(z)| - \delta \geq \eta^{-1} \zeta^2 - \delta \geq (2\eta)^{-1} \zeta^2,$$

where the first statement follows from (3.2); the second from the third bound in (3.5); the third from (A.6); and the fourth from the first bound in (3.5).

Therefore, there exists $m \in \llbracket 1, n \rrbracket$ such that $|\mu_m - \lambda| \leq 3n\zeta^{-2}\eta$ and $|v_m(\varphi)| \geq (6n)^{-1/2}\zeta$. Indeed, assuming otherwise, we would have

$$(2\eta)^{-1}\zeta^2 \leq \left| \sum_{j=1}^n \frac{v_j(\varphi)^2}{\mu_j - z} \right| < n \cdot (3n\zeta^{-2}\eta)^{-1} + n \cdot \eta^{-1} \cdot ((6n)^{-1/2}\zeta)^2 = (2\eta)^{-1}\zeta^2,$$

where the first bound follows from (A.7) and the second from our assumption (with the fact that $z = \lambda + i\eta$). This is a contradiction, which establishes the lemma. \square

A.3. Proofs of Results From Section 3.4.

Proof of Lemma 3.8. For any $E \in \mathbb{R} \setminus \text{eig } \mathbf{M}$, denote the resolvent $\mathbf{G}(E) = [G_{ij}(E)] = (\mathbf{M} - E)^{-1}$. By (3.2) (with the fact by [15, Proposition 2.40(a)] that all eigenvalues of \mathbf{M} are mutually disjoint), we have

$$(A.8) \quad u_{N_1} \cdot u_{N_2} = \lim_{E \rightarrow \mu} (\mu - E) \cdot G_{N_1 N_2}(E).$$

To evaluate the right side of (A.8), we let $\mathbf{C}(E) = [C_{ij}(E)]$ denote the cofactor matrix of $\mathbf{M} - E \cdot \text{Id}$ and use the fact that $G_{N_1 N_2}(E) = (-1)^{N+1} \cdot C_{N_1 N_2}(E) \cdot (\det \mathbf{M} - E \cdot \text{Id})^{-1}$. Since removing the row of index N_1 and column of index N_2 from \mathbf{M} yields a lower triangular $(N-1) \times (N-1)$ matrix with diagonal entries $(M_{i,i+1})_{N_1 \leq i < N_2}$, we deduce that $C_{N_1 N_2}(E) = \prod_{i=N_1}^{N_2-1} M_{i,i+1}$. Hence,

$$(A.9) \quad \begin{aligned} G_{N_1 N_2}(E) &= (-1)^{N+1} \cdot C_{N_1 N_2}(E) \cdot (\det \mathbf{M} - E \cdot \text{Id})^{-1} \\ &= (-1)^{N+1} \cdot \prod_{i=N_1}^{N_2-1} M_{i,i+1} \cdot \prod_{\mu' \in \text{eig } \mathbf{M}} (\mu' - E)^{-1}. \end{aligned}$$

Combining (A.9) with (A.8) yields the lemma. \square

Proof of Lemma 3.9. Since $\mathbf{M} \cdot \mathbf{u} = \mu \cdot \mathbf{u}$ and \mathbf{M} is tridiagonal, we have for any index $k \in \llbracket N_1, N_2 - 1 \rrbracket$ that $M_{k,k+1} \cdot u_{k+1} = (\mu - M_{k,k}) \cdot u_k - M_{k,k-1} \cdot u_{k-1}$. Since $M_{k,k-1} = M_{k-1,k}$, this is equivalent to $\mathbf{S}_k(\mu) \cdot \mathbf{w}_k = \mathbf{w}_{k+1}$, so the lemma follows by induction on $j - i$. \square

Proof of Lemma 3.10. It quickly follows from the explicit forms (3.6) of \mathbf{S}_k and (3.7) of \mathbf{S}_K that there exist monic polynomials $P_{\ell-1}$, P_ℓ , $Q_{\ell-2}$, and $Q_{\ell-1}$ satisfying the following two properties. First, we have

$$\mathbf{S}_{\llbracket i, j \rrbracket}(E) = \begin{bmatrix} -Q_{\ell-2}(E) \cdot M_{i-1, i} \cdot \prod_{k=i}^{j-1} M_{k, k+1}^{-1} & P_{\ell-1}(E) \cdot \prod_{k=i}^{j-1} M_{k, k+1}^{-1} \\ -Q_{\ell-1}(E) \cdot M_{i-1, i} \cdot \prod_{k=i}^j M_{k, k+1}^{-1} & P_\ell(E) \cdot \prod_{k=i}^j M_{k, k+1}^{-1} \end{bmatrix}.$$

Second, we have that $\deg Q_{\ell-2} = \ell - 2$; that $\deg P_{\ell-1} = \deg Q_{\ell-1} = \ell - 1$; and that $\deg P_{\ell} = \ell$. Thus, it suffices to show that

$$(A.10) \quad \begin{aligned} P_{\ell}(E) &= \prod_{h=1}^{\ell} (E - \mu_h^{[i,j]}); & P_{\ell-1}(E) &= \prod_{h=1}^{\ell-1} (E - \mu_h^{[i,j-1]}); \\ Q_{\ell-1}(E) &= \prod_{h=1}^{\ell-1} (E - \mu_h^{[i+1,j]}); & Q_{\ell-2}(E) &= \prod_{h=1}^{\ell-2} (E - \mu_h^{[i+1,j-1]}). \end{aligned}$$

To that end, we locate the zeroes of P_{ℓ} . Fix any $\mu \in \text{eig } \mathbf{M}^{[i,j]}$, and let $\mathbf{v} = (v_i, v_{i+1}, \dots, v_j) \in \mathbb{R}^{\ell}$ denote an eigenvector of $\mathbf{M}^{[i,j]}$ with eigenvalue μ . Then setting $\mathbf{w}_k = (v_{k-1}, v_k)$ for each $k \in \llbracket i, j \rrbracket$, with $v_{i-1} = 0$, we have from Lemma 3.9 that $\mathbf{S}_{\llbracket i, j-1 \rrbracket}(\mu) \cdot \mathbf{w}_i = \mathbf{w}_j$. Furthermore, the second coordinate of $\mathbf{S}_j(\mu) \cdot \mathbf{w}_j \in \mathbb{R}^2$ is equal

$$M_{j,j+1}^{-1} \cdot ((\mu - M_{j,j})v_j - M_{j-1,j}v_{j-1}) = M_{j,j+1}^{-1} \cdot (\mu v_j - M_{j,j}v_j - M_{j,j-1}v_{j-1}) = 0,$$

where the first statement follows from the fact that \mathbf{M} is symmetric and the second from the fact that \mathbf{v} is an eigenvector of $\mathbf{M}^{[i,j]}$ with eigenvalue μ . Therefore, the second coordinate of $\mathbf{S}_{\llbracket i, j \rrbracket}(\mu) \cdot \mathbf{w}_i$ is equal to 0. Since the first coordinate of \mathbf{w}_i is equal to $v_{i-1} = 0$ (and its second coordinate is nonzero), this implies that the $(2, 2)$ -entry of $\mathbf{S}_{\llbracket i, j \rrbracket}(\mu)$ is equal to 0. Hence, $P_{\ell}(\mu) = 0$ for each $\mu \in \text{eig } \mathbf{M}^{[i,j]}$. Since $\deg P_{\ell} = \ell$ and P_{ℓ} is monic, this confirms the first statement in (A.10).

We can now quickly deduce the remaining equalities in (A.10). Indeed, the second statement in (A.10) follows from the first, together with the fact that the $(1, 2)$ -entry of $\mathbf{S}_{\llbracket i, j \rrbracket}(E)$ is equal to the $(2, 2)$ -entry of $\mathbf{S}_{\llbracket i, j-1 \rrbracket}(E)$ (by (3.6) and (3.7)). The third statement in (A.10) follows from the first, together with the fact that the $(2, 1)$ -entry of $\mathbf{S}_{\llbracket i, j \rrbracket}(E)$ is equal to the $(2, 2)$ -entry of $\mathbf{S}_{\llbracket i+1, j \rrbracket}(E)$ multiplied by $-M_{i,i+1}^{-1}M_{i-1,i}$ (again by (3.6) and (3.7)). The fourth statement in (A.10) follows from the third, again together with the fact that the $(1, 1)$ -entry of $\mathbf{S}_{\llbracket i, j \rrbracket}(E)$ is equal to the $(2, 1)$ -entry of $\mathbf{S}_{\llbracket i, j-1 \rrbracket}(E)$. This establishes the lemma. \square

A.4. Proofs of Results From Section 3.5.

Proof of Lemma 3.15. By Lemma 3.5 and Lemma 2.2, it suffices to verify the lemma at $t = 0$, that is, to show $\mathbb{P}[\text{BND}_{\mathbf{L}(0)}(A)] \geq 1 - c^{-1}Ne^{-cA^2}$. To that end, observe by Definition 2.4 (or Definition 3.3) and a union bound, there is a constant $C > 1$ such that

$$\mathbb{P} \left[\max_{a \in \mathbf{a}(0)} |a| + \max_{b \in \mathbf{b}(0)} |b| \geq \frac{A}{4} \right] \leq CNA^{2\theta} e^{-\beta A^2/2}.$$

By the deterministic bound (A.4), this yields $\mathbb{P}[\max_{\lambda \in \text{eig } \mathbf{L}(0)} |\lambda| \geq A] \leq CNA^{2\theta} e^{-\beta A^2/2}$. Together, these two estimates imply $\mathbb{P}[\text{BND}_{\mathbf{L}(0)}(A)] \geq 1 - 2CNA^{2\theta} e^{-\beta A^2/2}$, which (as mentioned above) implies the lemma. \square

Proof of Lemma 3.18. Denote $\text{eig } \mathbf{L} = (\lambda_1, \lambda_2, \dots, \lambda_N)$, and define the events

$$\mathbf{E} = \bigcap_{i=1}^{N-1} \{\lambda_i \geq \lambda_{i+1} + \delta\}; \quad \mathbf{F} = \left\{ \max_{1 \leq i \leq N} |\lambda_i| \leq 3N \right\}.$$

Let $\mathcal{J} = \llbracket -4\delta^{-1}N, 4\delta^{-1}N \rrbracket$. For each $j \in \mathcal{I}$, set $z_j = j\delta + \delta i$, and let

$$S = \sum_{j \in \mathcal{J}} \sum_{1 \leq i \neq k \leq N} \operatorname{Im}(\lambda_i - z_j)^{-1} \cdot \operatorname{Im}(\lambda_k - z_j)^{-1}.$$

Observe on $\mathbf{E}^{\mathbb{G}} \cap \mathbf{F}$ that there exists some $j \in \mathcal{J}$ and $i \in \llbracket 1, N-1 \rrbracket$ such that $\operatorname{Im}(\lambda_i - z_j)^{-1} \geq (2\delta)^{-1}$ and $\operatorname{Im}(\lambda_{i+1} - z_j)^{-1} \geq (2\delta)^{-1}$. Hence, on $\mathbf{E}^{\mathbb{G}} \cap \mathbf{F}$ we have $S \geq (2\delta)^{-2}$, so taking expectations yields

$$(A.11) \quad \mathbb{P}[\mathbf{E}^{\mathbb{G}} \cap \mathbf{F}] \leq (2\delta)^2 \cdot \mathbb{E}[S].$$

Now, set $\mathbf{G}(z_j) = [G_{ik}(z_j)] = (\mathbf{L} - z_j)^{-1}$ for each $j \in \mathcal{J}$. By [36, Equations (2.64) and (2.65)],¹⁰ there exists a constant $C_0 > 1$ such that

$$\mathbb{E}[S] = \sum_{j \in \mathcal{J}} \sum_{1 \leq i, k \leq N} \mathbb{E} \left[\det \begin{bmatrix} \operatorname{Im} G_{ii}(z_j) & \operatorname{Im} G_{ik}(z_j) \\ \operatorname{Im} G_{ik}(z_j) & \operatorname{Im} G_{kk}(z_j) \end{bmatrix} \right] \leq C_0 N^2 |\mathcal{J}|.$$

Together with (A.11) and the fact that $|\mathcal{J}| \leq 9\delta^{-1}N$, this yields

$$\mathbb{P}[\mathbf{E}^{\mathbb{G}} \cap \mathbf{F}] \leq 36C_0\delta N^3.$$

Since Lemma 3.15 yields constants $C_1 > 1 > c_1 > 0$ such that $\mathbb{P}[\mathbf{F}^{\mathbb{G}}] \leq C_1 e^{-c_1 N^2}$, this with a union bound yields the lemma. \square

APPENDIX B. HEURISTICS FOR EIGENVALUE VELOCITIES

Throughout this section, we adopt Assumption 2.8. For each $k \in \llbracket 1, N \rrbracket$ and $t \geq 0$, define $v_k(t)$ of Q_k by setting $Q_k(t) = Q_k(0) + t \cdot v_k(t)$. This quantity $v_k(t)$ can be thought of as the ‘‘velocity’’ of the eigenvalue λ_k under the Toda lattice. In this section, following the physics literature (for example, [16]), we explain how Theorem 2.11 can be used to heuristically derive the large T limit for $v_k(T)$, which will coincide with the predictions of [16, Equation (90)] and [45, Equation (6.19)]. The discussion in this section is not entirely rigorous, and we do not know of a direct way to mathematically justify it completely; the sequel work [1] will be devoted to establishing its output (when θ is sufficiently small) through another route. In what follows, we let $T \gg 1$ be large; abbreviate $v_k(T) = v_k$; and suppose for notational simplicity that $\alpha > 0$ (the case when $\alpha < 0$ is entirely analogous).

Applying the asymptotic scattering relation (2.15), we obtain with high probability that

$$(B.1) \quad \lambda_k \approx v_k + 2T^{-1} \sum_{i=1}^N (\mathbb{1}_{Q_i(0) < Q_k(0) + T(v_k - v_i)} - \mathbb{1}_{Q_i(0) < Q_k(0)}) \cdot \log |\lambda_k - \lambda_i|.$$

Any eigenvalue λ_i should (by Corollary 5.6, for example) approximately only depend on the entries of the Lax matrix \mathbf{L} with indices close to $\varphi_0(i)$. Since the entries of \mathbf{L} are independent under thermal equilibrium, this indicates that λ_i should be approximately independent from most of the other λ_j , and from $Q_i(0)$ (where the latter holds since the a -entries in \mathbf{L} depend on only the differences $q_{i+1} - q_i$). Now, let us assume that $v_k \approx v(\lambda_k)$ asymptotically only depends on λ_k (that it, it is approximately independent from the other $(\lambda_j)_{j \neq k}$).¹¹ By the above approximate independence, this suggests that $\mathbb{1}_{Q_i(0) < Q_k(0) + T(v_k - v_i)} - \mathbb{1}_{Q_i(0) < Q_k(0)}$ should behave as $\mathbb{1}_{Q_j(0) < Q_k(0) + T(v_k - v_i)} -$

¹⁰The results there are stated for a different random matrix, but it is quickly verified that its proofs apply for the one \mathbf{L} we study.

¹¹We are unaware of an (even heuritic) explanation for this, and so the proofs in [1] will proceed differently (based on regularizing the indicator functions), instead.

$\mathbb{1}_{Q_j(0) < Q_k(0)}$, for a uniformly random index $j \in \llbracket 1, N \rrbracket$. Therefore, the sum on the right side of (B.1) should approximate

$$(B.2) \quad \sum_{i=1}^N (\mathbb{1}_{Q_i(0) < Q_k(0) + T(v_k - v_i)} - \mathbb{1}_{Q_i(0) < Q_k(0)}) \cdot \log |\lambda_k - \lambda_i| \approx \sum_{i \neq k} \mathfrak{N}_i \cdot \log |\lambda_k - \lambda_i|,$$

where

$$\mathfrak{N}_i = N^{-1} \cdot (\#\{j : Q_j(0) < Q_k(0) + T(v_k - v_i)\} - \#\{j : Q_j(0) < Q_k(0)\}).$$

To approximate \mathfrak{N} , recall under thermal equilibrium that the $q_{i+1} - q_i$ are identically distributed under thermal equilibrium (due to (2.3), since the a_i are, by Corollary 4.8) and that $\mathbb{E}[q_{i+1} - q_i] = \alpha$ (by Lemma 3.11). Therefore, $\mathfrak{N}_i \approx N^{-1} \cdot \alpha^{-1} \cdot T(v_k - v_i)$, which upon insertion into (B.2) yields

$$(B.3) \quad \sum_{i=1}^N (\mathbb{1}_{Q_i(0) < Q_k(0)} - \mathbb{1}_{Q_i(0) < Q_k(0) + T(v_k - v_i)}) \cdot \log |\lambda_k - \lambda_i| \approx T(\alpha N)^{-1} \sum_{i \neq k} (v_k - v_i) \cdot \log |\lambda_k - \lambda_i|.$$

Now, the (λ_j) are eigenvalues of the random Lax matrix \mathbf{L} . The empirical spectral distribution of the latter is known (from [33, Lemma 4.3]) to converge to an explicit probability density $\varrho(x)dx$. Since $v_i = v(\lambda_i)$ is a function of λ_i , we expect the above sum to likely concentrate, namely,

$$(B.4) \quad (\alpha N)^{-1} \sum_{i \neq k} (v_k - v_i) \cdot \log |\lambda_k - \lambda_i| \approx \alpha^{-1} \int_{-\infty}^{\infty} (v(\lambda_k) - v(\lambda)) \cdot \log |\lambda_k - \lambda| \varrho(\lambda) d\lambda.$$

Together, the approximations (B.1), (B.3), and (B.4) yield

$$(B.5) \quad \lambda_k \approx v(\lambda_k) + 2\alpha^{-1} \int_{-\infty}^{\infty} (v(\lambda_k) - v(\lambda)) \cdot \log |\lambda_k - \lambda| \varrho(\lambda) d\lambda.$$

Replacing the approximation in (B.5) with an equality yields a linear system of equations for v that coincides with [16, Equation (90)] and [45, Equation (6.19)]. It was explained in [45, Equation (6.21)] that this system has a specific solution $v(\lambda) = v_{\text{eff}}(\lambda)$, given by [45, Equation (6.20)] (and [16, Equation (93)]); presumably, this solution v_{eff} is unique. This would then indicate $v_k \approx v_{\text{eff}}(\lambda_k)$.

REFERENCES

- [1] A. Aggarwal. In preparation.
- [2] M. Aizenman, J. H. Schenker, R. M. Friedrich, and D. Hundertmark. Finite-volume fractional-moment criteria for Anderson localization. volume 224, pages 219–253. 2001. Dedicated to Joel L. Lebowitz.
- [3] J. Avron and B. Simon. Almost periodic Schrödinger operators. II. The integrated density of states. *Duke Math. J.*, 50(1):369–391, 1983.
- [4] B. Bertini, M. Collura, J. De Nardis, and M. Fagotti. Transport in out-of-equilibrium XXZ chains: exact profiles of charges and currents. *Physical review letters*, 117(20):207201, 2016.
- [5] A. Bloch, F. Golse, T. Paul, and A. Uribe. Dispersionless Toda and Toeplitz operators. *Duke Math. J.*, 117(1):157–196, 2003.
- [6] C. Boldrighini, R. L. Dobrushin, and Y. M. Sukhov. One-dimensional hard rod caricature of hydrodynamics. *J. Statist. Phys.*, 31(3):577–616, 1983.
- [7] V. B. Bulchandani, X. Cao, and J. E. Moore. Kinetic theory of quantum and classical Toda lattices. *J. Phys. A*, 52(33):33LT01, 14, 2019.
- [8] M. J. Cantero, L. Moral, and L. Velázquez. Five-diagonal matrices and zeros of orthogonal polynomials on the unit circle. *Linear Algebra Appl.*, 362:29–56, 2003.
- [9] X. Cao, V. B. Bulchandani, and H. Spohn. The GGE averaged currents of the classical Toda chain. *J. Phys. A*, 52(49):495003, 12, 2019.

- [10] O. A. Castro-Alvaredo, B. Doyon, and T. Yoshimura. Emergent hydrodynamics in integrable quantum systems out of equilibrium. *Physical Review X*, 6(4):041065, 2016.
- [11] D. A. Croydon and M. Sasada. Generalized hydrodynamic limit for the box-ball system. *Comm. Math. Phys.*, 383(1):427–463, 2021.
- [12] P. Deift, S. Kamvissis, T. Kriecherbauer, and X. Zhou. The Toda rarefaction problem. *Comm. Pure Appl. Math.*, 49(1):35–83, 1996.
- [13] P. Deift and K. T.-R. McLaughlin. A continuum limit of the Toda lattice. *Mem. Amer. Math. Soc.*, 131(624):x+216, 1998.
- [14] P. Deift, T. Nanda, and C. Tomei. Ordinary differential equations and the symmetric eigenvalue problem. *SIAM J. Numer. Anal.*, 20(1):1–22, 1983.
- [15] P. A. Deift. *Orthogonal polynomials and random matrices: a Riemann-Hilbert approach*, volume 3 of *Courant Lecture Notes in Mathematics*. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 1999.
- [16] B. Doyon. Generalized hydrodynamics of the classical Toda system. *J. Math. Phys.*, 60(7):073302, 21, 2019.
- [17] B. A. Dubrovin, V. B. Matveev, and S. P. Novikov. Nonlinear equations of Korteweg-de Vries type, finite-band linear operators and Abelian varieties. *Uspehi Mat. Nauk*, 31(1(187)):55–136, 1976.
- [18] G. A. El. The thermodynamic limit of the Whitham equations. *Phys. Lett. A*, 311(4-5):374–383, 2003.
- [19] H. Flaschka. The Toda lattice. II. Existence of integrals. *Phys. Rev. B*, 9(4):1924, 1974.
- [20] M. Girotti, T. Grava, R. Jenkins, and K. D. T.-R. McLaughlin. Rigorous asymptotics of a KdV soliton gas. *Comm. Math. Phys.*, 384(2):733–784, 2021.
- [21] M. Girotti, T. Grava, R. Jenkins, K. T.-R. McLaughlin, and A. Minakov. Soliton versus the gas: Fredholm determinants, analysis, and the rapid oscillations behind the kinetic equation. *Comm. Pure Appl. Math.*, 76(11):3233–3299, 2023.
- [22] M. Girotti, T. Grava, K. McLaughlin, and J. Nanjudel. Law of large numbers and central limit theorem for random sets of solitons of the focusing nonlinear Schrödinger equation. Preprint, arXiv:2411.17036.
- [23] T. Grava, M. Gionni, G. Gubbiotti, and G. Mazzuca. Discrete integrable systems and random Lax matrices. *J. Stat. Phys.*, 190(1):Paper No. 10, 35, 2023.
- [24] A. Guionnet and R. Memin. Large deviations for Gibbs ensembles of the classical Toda chain. *Electron. J. Probab.*, 27:Paper No. 46, 29, 2022.
- [25] U. Islambekov, R. Sims, and G. Teschl. Lieb-Robinson bounds for the Toda lattice. *J. Stat. Phys.*, 148(3):440–479, 2012.
- [26] J. Q. K. Matetski and D. Remenik. Polynuclear growth and the Toda lattice. To appear in *J. Eur. Math. Soc.* Preprint, arXiv:2209.02643.
- [27] R. Killip and I. Nenciu. CMV: the unitary analogue of Jacobi matrices. *Comm. Pure Appl. Math.*, 60(8):1148–1188, 2007.
- [28] H. Krüger and G. Teschl. Long-time asymptotics of the Toda lattice for decaying initial data revisited. *Rev. Math. Phys.*, 21(1):61–109, 2009.
- [29] H. Kunz and B. Souillard. Sur le spectre des opérateurs aux différences finies aléatoires. *Comm. Math. Phys.*, 78(2):201–246, 1980/81.
- [30] O. E. Lanford, III, J. L. Lebowitz, and E. H. Lieb. Time evolution of infinite anharmonic systems. *J. Statist. Phys.*, 16(6):453–461, 1977.
- [31] E. H. Lieb and D. W. Robinson. The finite group velocity of quantum spin systems. *Comm. Math. Phys.*, 28:251–257, 1972.
- [32] S. Manakov. Complete integrability and stochastization of discrete dynamical systems. *Zh. Exp. Teor. Fiz.*, 67(2):543–555, 1974.
- [33] G. Mazzuca. On the mean density of states of some matrices related to the beta ensembles and an application to the Toda lattice. *J. Math. Phys.*, 63(4):Paper No. 043501, 13, 2022.
- [34] G. Mazzuca and R. Memin. CLT for β -Ensembles at High Temperature and for Integrable Systems: A Transfer Operator Approach. *Ann. Henri Poincaré*, 26(1):245–316, 2025.
- [35] H. P. McKean and E. Trubowitz. Hill’s operator and hyperelliptic function theory in the presence of infinitely many branch points. *Comm. Pure Appl. Math.*, 29(2):143–226, 1976.
- [36] N. Minami. Local fluctuation of the spectrum of a multidimensional Anderson tight binding model. *Comm. Math. Phys.*, 177(3):709–725, 1996.
- [37] S. A. Molchanov. Structure of the eigenfunctions of one-dimensional unordered structures. *Izv. Akad. Nauk SSSR Ser. Mat.*, 42(1):70–103, 214, 1978.

- [38] S. A. Molchanov. The local structure of the spectrum of a random one-dimensional Schrödinger operator. *Trudy Sem. Petrovsk.*, (8):195–210, 1982.
- [39] J. Moser. Finitely many mass points on the line under the influence of an exponential potential—an integrable system. In *Dynamical systems, theory and applications (Rencontres, Battelle Res. Inst., Seattle, Wash., 1974)*, volume Vol. 38 of *Lecture Notes in Phys.*, pages 467–497. Springer, Berlin-New York, 1975.
- [40] M. Opper. Analytical solution of the classical Bethe ansatz equation for the Toda chain. *Phys. Lett. A*, 112(5):201–203, 1985.
- [41] N. Saitoh. A transformation connecting the Toda lattice and the KdV equation. *J. Phys. Soc. Japan*, 49(1):409–416, 1980.
- [42] J. Schenker. Eigenvector localization for random band matrices with power law band width. *Comm. Math. Phys.*, 290(3):1065–1097, 2009.
- [43] B. Simon. *Orthogonal polynomials on the unit circle. Part 2*, volume 54, Part 2 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2005. Spectral theory.
- [44] H. Spohn. Generalized Gibbs ensembles of the classical Toda chain. *J. Stat. Phys.*, 180(1-6):4–22, 2020.
- [45] H. Spohn. *Hydrodynamic scales of integrable many-body systems*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2024.
- [46] D. Takahashi and J. Satsuma. A soliton cellular automaton. *J. Phys. Soc. Japan*, 59(10):3514–3519, 1990.
- [47] G. Teschl. *Jacobi operators and completely integrable nonlinear lattices*, volume 72 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2000.
- [48] D. J. Thouless. A relation between the density of states and range of localization for one dimensional random systems. *J. Phys. C: Solid State Phys.*
- [49] M. Toda. Vibration of a chain with nonlinear interaction. *Journal of the Physical Society of Japan*, 22(2):431–436, 1967.
- [50] S. Venakides, P. Deift, and R. Oba. The Toda shock problem. *Comm. Pure Appl. Math.*, 44(8-9):1171–1242, 1991.
- [51] V. E. Zakharov. Kinetic equation for solitons. *Sov. Phys. JETP*, 33(3):538–540, 1971.
- [52] V. E. Zakharov. Turbulence in integrable systems. *Stud. Appl. Math.*, 122(3):219–234, 2009.