Jordan-Holder Theorem for profinite groups and applications

Tamar Bar-On and Nikolay Nikolov

March 13, 2025

Abstract

We generalize the notions of composition series and composition factors for profinite groups, and prove a profinite version of the Jordan-Holder Theorem. We apply this to prove a Galois Theorem for infinite prosolvable extensions. In addition, we investigate the connection between the abstract and topological composition factors of a nonstrongly complete profinite group.

1 Composition factors and Jordan-Holder Theorem

The composition factors of a finite group are one of its most important invariants, and help translating structural questions in finite group theory to questions on the structure of finite simple groups - which have been classified.

The fundamental fact about composition factors and their multiplicity is that they do not depend on the composition series:

Theorem 1 (Jordan-Holder Theorem for finite groups). Let G be a finite group and let $\{e\} = G_n \trianglelefteq \cdots G_1 \trianglelefteq G_0 = G$ and $\{e\} = H_m \trianglelefteq \cdots H_1 \trianglelefteq H_0 = G$ be two composition series for G. Then for every finite simple group S, $|\{0 \le i \le n-1 : G_i/G_{i+1} \cong S\}| = |\{0 \le i \le m-1 : H_i/H_{i+1} \cong S\}|$. In particular m = n and the sets $\{G_i/G_{i+1}| \ 0 \le i \le n-1\}$ and $\{H_i/H_{i+1}| \ 0 \le i \le m-1\}$ are equal.

For a proof see for example [3]. For infinite groups the theorem no longer holds, as can be seen for example, by taking $G = \mathbb{Z}$ and considering the two composition series: $\mathbb{Z} \succeq p\mathbb{Z} \trianglerighteq p^2\mathbb{Z} \trianglerighteq \dots$ and $\mathbb{Z} \trianglerighteq q\mathbb{Z} \trianglerighteq q^2\mathbb{Z} \trianglerighteq \dots$ for $p \neq q$ primes.

Many generalizations to Jordan-Holder Theorem have been proven, as in [2] for well-ordered ascending T-invariant series, and the Jordan-Holder Theorem for composition factors of modules (see, for example, [4, Theorem 3.11]).

The purpose of this paper is to establish a generalization of the Jordan-Holder Theorem for profinite groups. More precisely, we define composition factors and composition series for profinite groups, and show that the composition factors of a profinite groups, as well as their multiplicity, are well-defined and independent on the choice of a composition series. Through this paper, unless stated otherwise (which will be relevant only in Section 3) subgroups in a profinite group are assumed to be closed and quotients are assumed to be topological quotients.

Our first observation is that for profinite groups, subnormal series are always replaced by *accessible series*:

Definition 2. [1, Chapter 8.3] Let G be a profinite group and $H \leq G$. An accessible series from G to H over some ordinal μ is a series $G = G_0 > ... > G_{\lambda} > ... > G_{\mu} = H$ which satisfies the following property:

- 1. For every $\alpha < \mu$, $G_{\alpha+1} \trianglelefteq G_{\alpha}$.
- 2. For every limit ordinal $\alpha < \mu$, $G_{\alpha} = \bigcap_{\beta < \alpha} G_{\beta}$.

If such a series exists we say that H is an *accessible subgroup*. In case $H = \{e\}$ we refer to an accessible series from G to $\{e\}$ simply as an accessible series for G.

Now we can define a *composition series*:

Definition 3. Let G be a profinite group. A composition series for G is an accessible series $G = G_0 > ... > G_{\lambda} > ... > G_{\mu} = \{e\}$ such that $G_{\lambda}/G_{\lambda+1}$ is a finite simple nontrivial group for every $\lambda < \mu$. The quotients $G_{\lambda}/G_{\lambda+1}$ for all $\lambda < \mu$ are called the *composition factors* of G.

Proposition 4. Every profinite group admits a composition series, and the composition factors are well defined. Moreover, the set of composition factors of G is precisely the set of all composition factors of all finite topological quotients of G.

Proof. First we show that every profinite group admits a composition series. Let $\eta = |G|$ and $\nu = 2^{\eta}$. We define a composition series over ν recursively, as follows. Assume that for all $\alpha < \beta \ G_{\alpha}$ has been defined. If $\beta = \gamma + 1$ for some γ then we define G_{β} to be a maximal open normal subgroup of G_{γ} in case $G_{\gamma} \neq \{e\}$ and to be $\{e\}$ otherwise. If β is a limit ordinal then we define $G_{\beta} = \bigcap_{\alpha < \beta} G_{\alpha}$. We claim that $G_{\nu} = \{e\}$. Indeed, otherwise we can use the axiom of choice to define a one-to-one map $\nu \to G$ by sending each α to an element in $G_{\alpha} \setminus G_{\alpha+1}$. Let μ be the first ordinal for which $G_{\mu} = \{e\}$, then $G = G_0 > \ldots > G_{\lambda} > \ldots > G_{\mu} = \{e\}$ is a composition series for G.

Now we prove that for every composition series $\{G_{\alpha}\}_{\alpha < \mu}$, the set

$$C_G := \{G_\alpha / G_{\alpha+1}\}_{\alpha < \mu}$$

equals the set of composition factors of all finite quotients of G, and that will imply that C_G does not depend on the choice of a composition series.

First we show the following: let $\{G_{\alpha}\}_{\alpha < \mu}$ be a composition series for G and A a finite image of G with some canonical projection $\varphi_A : G \to A$. Then the image series $\{\varphi_A(G_{\alpha})\}_{\alpha < \mu}$ is clearly a subnormal series in A which ends in $\{e\}$. Denote the different subgroups in this series by $A = A_0 \supseteq A_1 \supseteq \cdots A_n = \{e\}$.

For $0 < i \leq n$, let α be the first ordinal such that $\varphi_A(G_\alpha) = A_i$. We claim that α is a successor ordinal. Otherwise, for every $\beta < \alpha \ \varphi_A(G_\beta) \geq A_{i-1}$. Then by [1, Proposition 2.1.4]

$$\varphi_A(G_\alpha) = \varphi_A(\bigcap_{\beta < \alpha} G_\beta) = \bigcap_{\beta < \alpha} \varphi_A(G_\alpha) \ge A_{i-1}$$

Now let $\alpha = \beta + 1$. Then $\varphi_A(G_\beta) = A_{i-1}$ and since $G_\beta/G_{\beta+1}$ is simple, $G_\beta/G_{\beta+1} \cong A_{i-1}/A_i$. In particular, $A = A_0 \supseteq A_1 \supseteq \cdots A_n = \{e\}$ is a composition series and all its factors belongs to the set of the factors $\{G_\alpha/G_{\alpha+1} : \alpha < \mu\}$. We conclude that the set of the factors $\{G_\alpha/G_{\alpha+1} : \alpha < \mu\}$ contains the set of composition factors of all finite quotients of G.

Conversely, let $\alpha < \mu$. There is some finite quotient A of G in which $\varphi_A(G_\alpha) \neq \varphi_A(G_{\alpha+1})$. By the argument above, the different members of $\{\varphi_A(G_\alpha)\}_{\alpha < \mu}$ form a composition series $A = A_0 \supseteq A_1 \supseteq \cdots A_n = \{e\}$ of A. Hence if $\varphi_A(G_\alpha) = A_i$ then $\varphi(G_{\alpha+1}) = A_{i+1}$ and $G_\alpha/G_{\alpha+1} \cong A_i/A_{i+1}$, again from the simplicity of $G_\alpha/G_{\alpha+1}$. In particular, $G_\alpha/G_{\alpha+1}$ is a composition factor of A, and we are done.

Recall that a profinite group is called prosolvable if all its finite quotients are solvable.

Corollary 5. Let G be a profinite group. Then G is prosolvable if and only if all its composition factors are abelian.

Now we wish to prove the second part of the main goal of this paper:

Theorem 6. Let G be a profinite group and S a finite simple group. Let $G = G_0 > ... > G_{\alpha} > ... > G_{\mu} = \{e\}$ and $G = H_0 > ... > H_{\lambda} > ... > H_{\Delta} = \{e\}$ be two composition series for G. Let S be some finite simple group. Denote by $S_1 = \{\alpha < \mu : G_{\alpha}/G_{\alpha+1} \cong S\}$ and $S_2 = \{\lambda < \Delta : H_{\lambda}/H_{\lambda+1} \cong S\}$. Then $|S_1| = |S_2|$.

First we prove a stronger result in the case of S_1 being finite. Let S be a simple group. For a finite quotient A of G we denote by $n_{S,A}$ the number of appearances of S as a factor in some, and hence any, composition series of A.

Lemma 7. Let n be some natural number. Then $|S_1| = n$ if and only if

 $n = \max\{n_{S,A} \mid A \text{ is a finite quotient of } G\}$

Proof. As was shown in the proof of Proposition 4, the different image subgroups $\varphi_A(G_\alpha), \alpha < \mu$ induce a composition series $A = A_0 \supseteq \cdots A_m = \{e\}$ of A, for every finite image A of G. Moreover, there are $\alpha_0 < \ldots < \alpha_m < \mu$ such that $G_{\alpha_i}/G_{\alpha_i+1} \cong A_i/A_{i+1}$. Hence, $|\mathcal{S}_1| \ge n_{S,A}$ for every finite quotient A of G.

On the other hand, assume that l is a natural number and there are $\alpha_0 < \ldots < \alpha_l < \mu$ such that $G_{\alpha_i}/H_{\alpha_i+1} \cong S$ for all $0 \leq i \leq l$. There is some finite quotient A of G for which $\varphi_A(G_{\alpha_i}) \neq \varphi_A(G_{\alpha_i+1})$ for all $0 \leq i \leq l$. In particular, if we denote by $A = A_0 \supseteq \cdots A_m = \{e\}$ the induced composition series of A,

and $\varphi_A(G_{\alpha_i}) = A_{k_i}$ for some $0 \le k_i < m$ then $\varphi_A(G_{\alpha_i+1}) = A_{k_i+1}$. Moreover, $\varphi_A(G_{\alpha_i}) \ne \varphi_A(G_{\alpha_j})$ for all j > i. Thus, there are $0 \le k_1 < \ldots < k_l \le m$ such that $A_{k_i}/A_{k_i+1} \cong \varphi_A(G_{\alpha_i})/\varphi_A(G_{\alpha_i+1}) \cong S$ and we conclude that $\max\{n_{S,A} : A$ is a finite quotient of $G\} \ge n_{S,A} \ge |\mathcal{S}_1|$.

Before we can prove Theorem 6 in full generality we need some Lemmas.

Lemma 8. Let $A \neq B$ and H be subgroups of a given group G such that B is strictly contained in A and $A \cap H = B \cap H$. Then $AH \neq BH$.

Proof. Assume by contradiction that AH = BH. Let $a \in A \setminus B$, then $a \in BH$. Thus there exist $b \in B, h \in H$ such that a = bh. Since $B \leq A, b \in A \Rightarrow h \in A$. Thus $h \in A \cap H = B \cap H \Rightarrow h \in B$ and we get that $a \in B$. A contradiction. \Box

Lemma 9. Let G be a profinite group and $A, B \leq G$ such that B is strictly contained in A. Let $\{H_{\alpha} : \alpha < \mu\}$ be an accessible series from G to some accessible subgroup H and assume that $AH \neq BH$. Then there exists some α such that $(A \cap H_{\alpha})H_{\alpha+1} \neq (B \cap H_{\alpha})H_{\alpha+1}$.

Proof. Let $\beta < \mu$ be the first ordinal such that $AH_{\beta} \neq BH_{\beta}$. We claim that β is a successor ordinal. Otherwise, $H_{\beta} = \bigcap_{\alpha < \beta} H_{\alpha}$ and by [1, Proposition 2.1.4] $AH_{\beta} = A \bigcap_{\alpha < \beta} H_{\alpha} = \bigcap_{\alpha < \beta} AH_{\alpha} = \bigcap_{\alpha < \beta} BH_{\alpha} = B \bigcap_{\alpha < \beta} H_{\alpha} = BH_{\beta}$, a contradiction. Thus $\beta = \alpha + 1$ for some α . Notice that $(A \cap H_{\alpha})H_{\alpha+1} = AH_{\alpha+1} \cap H_{\alpha}$ and the same holds for B. Since $AH_{\alpha+1} \neq BH_{\alpha+1}$ and $AH_{\alpha+1}H_{\alpha} = AH_{\alpha} = BH_{\alpha} = BH_{\alpha} = BH_{\alpha+1}H_{\alpha}$, Putting $A' = AH_{\alpha+1}, B' = BH_{\alpha+1}, H = H_{\alpha}$, Lemma 8 applies the required.

Lemma 10. Let $\{G_{\alpha} : \alpha < \mu\}$ be a composition series for G, and H be an accessible subgroup of G. Then $G_{\alpha} \cap H$ is a composition series for H.

Proof. It is clear that this series is accessible. Fix an $\alpha < \mu$, we want to show that $(G_{\alpha} \cap H)/(G_{\alpha+1} \cap H)$ is either trivial or isomorphic to $G_{\alpha}/G_{\alpha+1}$. Let $G = H_0 \supseteq ... \supseteq H_{\beta} \supseteq ... \supseteq H_{\Delta} = H$ be an accessible series from G to H. The claim is now translated to $(G_{\alpha} \cap H_{\Delta})/(G_{\alpha+1} \cap H_{\Delta})$ is either trivial or isomorphic to $G_{\alpha}/G_{\alpha+1}$. We will prove by induction on β that in fact for every $\beta \leq \Delta$ $(G_{\alpha} \cap H_{\beta})/(G_{\alpha+1} \cap H_{\beta})$ is either trivial or isomorphic to $G_{\alpha}/G_{\alpha+1}$.

Obviously, for $\beta = 0$ the claim holds. Now let $\beta \leq \Delta$ and assume the claim holds for all $\gamma < \beta$. If $H_{\gamma} \cap G_{\alpha} = H_{\gamma} \cap G_{\alpha+1}$ for some $\gamma < \beta$ then the same holds for H_{β} . Now assume that $G_{\alpha} \cap H_{\gamma}/G_{\alpha+1} \cap H_{\gamma} \neq \{e\}$ for every $\gamma < \beta$.

First case: β is limit. By induction assumption, $(G_{\alpha} \cap H_{\gamma})/(G_{\alpha+1} \cap H_{\gamma}) \cong G_{\alpha}/G_{\alpha+1}$ for every $\gamma < \beta$. Recall that $G_{\alpha}/G_{\alpha+1}$ is finite. We get that the natural inclusion $(G_{\alpha} \cap H_{\gamma})/(G_{\alpha+1} \cap H_{\gamma}) \to G_{\alpha}/G_{\alpha+1}$ is in fact an epimorphism. Hence, $(G_{\alpha} \cap H_{\gamma})G_{\alpha+1} = G_{\alpha}$ for all $\gamma < \beta$. Using [1, Proposition 2.1.4] implies the same for H_{β} and we conclude that the natural inclusion $(G_{\alpha} \cap H_{\beta})/(G_{\alpha+1} \cap H_{\beta}) \to G_{\alpha}/G_{\alpha+1}$ is an isomorphism, as required.

Second case: $\beta = \gamma + 1$. Then $H_{\beta} \leq H_{\gamma}$ implies that $(H_{\beta} \cap G_{\alpha})/(H_{\beta} \cap G_{\alpha+1}) \leq (H_{\gamma} \cap G_{\alpha})/(H_{\gamma} \cap G_{\alpha+1}) \approx G_{\alpha}/G_{\alpha+1}$. The last isomorphism is due to the induction assumption. Now the claim follows by $G_{\alpha}/G_{\alpha+1}$ being simple. \Box

Now we are ready to prove the main theorem.

Proof of Proposition 6. Let $\{G_{\alpha}\}_{\alpha < \mu}$ and $\{H_{\beta}\}_{\beta < \nu}$ be two composition series. Let S be a simple group and $\alpha < \mu$ be some ordinal such that $G_{\alpha}/G_{\alpha+1} \cong S$. For more simplicity denote $G_{\alpha} = A, G_{\alpha+1} = B$. By Lemma 9 there is some $\beta < \nu$ such that $(A \cap H_{\beta})H_{\beta+1} \neq (B \cap H_{\beta})H_{\beta+1}$. By Lemma 10, $\{G_{\alpha} \cap H_{\beta} : \alpha < \mu\}$ is an composition series of H_{β} . Hence, as was explained in the proof of 4, its image is a composition series for $H_{\beta}/H_{\beta+1}$. In particular, $((A \cap H_{\beta})H_{\beta+1})/H_{\beta+1}$ and $((B \cap H_{\beta})H_{\beta+1})/H_{\beta+1}$ are subnormal subgroups of the simple group $H_{\beta}/H_{\beta+1}$. We conclude that $(A \cap H_{\beta})H_{\beta+1} = H_{\beta}$ and $(B \cap H_{\beta})H_{\beta+1} = H_{\beta+1}$. By the proof of Lemma 10, either $B \cap H_{\beta} = A \cap H_{\beta}$ or $(A \cap H_{\beta})/(B \cap H_{\beta}) \cong A/B$. The first option is impossible since $(A \cap H_{\beta})H_{\beta+1} \neq (B \cap H_{\beta})H_{\beta+1}$. Thus $(A \cap H_{\beta})/(B \cap H_{\beta}) \cong A/B$. Now $H_{\beta}/H_{\beta+1} = ((A \cap H_{\beta})H_{\beta+1})/(B \cap H_{\beta})H_{\beta+1}$. Look at the natural epimorphism $(A \cap H_{\beta})/(B \cap H_{\beta}) \Rightarrow ((A \cap H_{\beta})H_{\beta+1})/(B \cap H_{\beta})H_{\beta+1}$. Since $(A \cap H_{\beta})/(B \cap H_{\beta})$ is simple, this is in fact an isomorphism. We conclude that $H_{\beta}/H_{\beta+1} \cong S$.

Now let S_1 be the set of all $\alpha < \mu$ such that $G_{\alpha}/G_{\alpha+1} \cong S$ and S_2 be the set of all $\beta < \nu$ such that $H_{\beta}/H_{\beta+1} \cong S$. We define a map $S_1 \to S_2$ by choosing for every $\alpha \in S_1$ the first $\beta \in S_2$ such that $(G_{\alpha} \cap H_{\beta})H_{\beta+1} \neq (G_{\alpha+1} \cap H_{\beta})H_{\beta+1}$.

By Lemma 9 such β always exists. By the above computations, such β satisfies $H_{\beta}/H_{\beta+1} \cong S$ and thus $\beta \in S_2$. This map is one-to-one. Indeed, assume $\alpha < \gamma \in S_1$ and $(G_{\alpha} \cap H_{\beta})H_{\beta+1} \neq (G_{\alpha+1} \cap H_{\beta})H_{\beta+1}$.

In particular, as we computed above, $(G_{\alpha+1} \cap H_{\beta})H_{\beta+1} = H_{\beta+1}$. Since $\gamma \ge \alpha + 1$,

$$H_{\beta+1} \le (G_{\gamma} \cap H_{\beta})H_{\beta+1} \le (G_{\alpha+1} \cap H_{\beta})H_{\beta+1} = H_{\beta+1}$$

The same goes for $G_{\gamma+1}$, so $(G_{\gamma} \cap H_{\beta})H_{\beta+1} = (G_{\gamma+1} \cap H_{\beta})H_{\beta+1}$. In conclusion, $|\mathcal{S}_1| \leq |\mathcal{S}_2|$. By symmetry we conclude that $|\mathcal{S}_2| \leq |\mathcal{S}_1|$ and thus we have an equality.

We end this section by computing the length of a composition series. First we need to define the following invariant of closed subgroups.

Definition 11. Let G be a profinite group and H a closed non open subgroup of G. We define $\omega_0(G/H)$ to be the cardinality of the set of all open subgroups of G containing H. If $H = \{e\}$ then we simply write $\omega_0(G)$. The cardinal $w_0(G/H)$ is in fact equal to the cardinality of any set of open subgroups of Gwhose intersection is H and which is filtered from below. Indeed, assume \mathcal{A} is such a set. Let U be an open subgroup containing H. Since $\bigcap \mathcal{A} = H \subseteq U$ there are some open subgroups $O_1, ..., O_n$ in \mathcal{A} such that $O_1 \cap ... \cap O_n \subseteq U$. By the filtration property, there is some $O \in \mathcal{A}$ such that $O \leq U$. Hence using the axiom of choice we can define a map from the set of all open subgroups containing H to \mathcal{A} sending an open subgroup U to a open subgroup $O \in \mathcal{A}$ such that $O \leq U$. Since O is open, this map is finite to 1, so the cardinality of the set of all open subgroups of G containing H is less or equal then the cardinality of \mathcal{A} . The converse clearly holds, thus we have an equality.

5

Remark 12. The definition of $\omega_0(G/H)$ is different then the definition that was given in [1, Section 2.6]. However, one can show that these two definitions are equivalent.

Lemma 13. Let H be an infinite-indexed accessible subgroup of G. Assume that $G = H_0 > ... > H_{\lambda} > ... > H_{\mu} = H$ is an accessible series from G to H such that all quotients $H_{\lambda}/H_{\lambda+1}$ are finite nontrivial. Then $|\mu| = \omega_0(G/H)$.

Proof. Clearly $G/H \cong \lim_{\leftarrow H \leq U \leq_o G} G/U$. Here the inverse limit considered as an inverse limit of spaces, as the subgroups are not assumed to be normal. For every λ we choose some U such that $\varphi_{G/U}(H_{\lambda}) \neq \varphi_{G/U}(H_{\lambda+1})$ where $\varphi_{G/U}$ denoted the natural quotient map $G \to G/U$. Since the images $\varphi_{G/U}(H_{\lambda})$ form a chain of subsets in each G/U, which is a finite space, the map adjoining to each $\lambda < \mu$ such U is finite to 1. Hence, $|\mu| \leq \omega_0(G/H)$.

On the other hand, we will prove by transfinite induction that $\omega_0(G/H_\lambda) \leq |\lambda|$. The proof is very similar to the proof of [1, Theorem 2.6.4 (c)]. For $\lambda = 1$ it is obvious. Now assume that $\lambda = \gamma + 1$. Since $H_{\gamma+1}$ is an open subgroup of H_{γ} then there exists some $V \leq_o G$ such that $V \cap H_{\gamma} \leq H_{\gamma+1}$. It can be shown that $\{VH_{\gamma+1} \cap U\}_{H_{\gamma} \leq U \leq_o G}$ is a set of filtered from below open subgroups whose intersection equals $H_{\gamma+1}$. Indeed, the fact that it is filtered from below comes from the filtration property of the set $H_{\gamma} \leq U \leq_o G$. As for the intersection,

$$\bigcap_{H_{\gamma} \le U \le_{o} G} (VH_{\gamma+1}) \cap U = \bigcap_{H_{\gamma} \le U \le_{o} G} (V \cap U)H_{\gamma+1}$$

$$=H_{\gamma+1}\bigcap_{H_{\gamma}\leq U\leq_{o}G}(V\cap U)=H_{\gamma+1}(V\bigcap_{H_{\gamma}\leq U\leq_{o}G}U)=H_{\gamma+1}(V\cap H_{\gamma})=H_{\gamma+1}(V\cap H_{\gamma})$$

The second equality is [1, Proposition 2.1.4].

Thus $\omega_0(G/H_{\gamma+1}) \le \omega_0(G/H_{\gamma}) \le |\gamma| = |\gamma+1|.$

Now assume that λ is a limit cardinal. Let U be an open subgroup of G containing H_{λ} . We have $\bigcap_{\gamma < \lambda} (H_{\gamma} \cap (G \setminus U)) = H_{\lambda} \setminus U = \emptyset$ and from the compactness of $G \setminus U$ we deduce that $H_{\gamma} \setminus U = \emptyset$ for some $\gamma < \lambda$, i.e. $H_{\gamma} \leq U$. This shows

$$\{U: H_{\lambda} \le U \le_o G\} = \bigcup_{\gamma < \lambda} \{U: H_{\gamma} \le U \le_o G\},\$$

hence

$$|\{U: H_{\lambda} \le U \le_{o} G\}| \le \sum_{\gamma < \lambda} |\{U: H_{\gamma} \le U \le_{o} G\}| \le \sum_{\gamma < \lambda} |\gamma| \le |\lambda|.$$

2 Galois Theorem for infinite extensions

Now we present a generalization of Galois Theorem for infinite extensions, using composition series of profinite groups.

Definition 14. Let K/F be a separable extension.

- 1. We say that K is prosolvable if $\operatorname{Gal}(N/F)$ is prosolvable, for N being the Galois closure of K over F.
- 2. We say that K is solvable by radicals if there exists a field L containing K, an ordinal μ , and a series of field extensions $F = L_0 \leq ... \leq L_{\alpha} < ... \leq L_{\mu} = L$ which satisfying the following properties:
 - For every $\alpha < \mu$, $L_{\alpha+1}/L_{\alpha}$ is a radical extension, i.e, $L_{\alpha+1} = L_{\alpha}[\sqrt[n]{a}]$ for some natural number n and $a \in L_{\alpha}$.
 - For every $\alpha < \mu$ limit, $L_{\alpha} = \bigcup_{\beta < \alpha} L_{\beta}$.

Remark 15. Since a quotient of a prosolvable group is prosolvable, K/F is a prosolvable extension if and only if K is contained in some Galois extension N/F such that Gal(N/F) is prosolvable.

Theorem 16 (Galois Theorem for infinite extensions). Let K/F be a separable extension, then K is prosolvable if and only if it is solvable by radicals.

Proof. First assume that K/F is prosolvable. Let N be the Galois closure of K. By assumption, $\operatorname{Gal}(N/K)$ is prosolvable. By Corollary 5 $G = \operatorname{Gal}(N/F)$ admits a composition series $G = G_0 \geq \dots G_\alpha \geq \dots \geq G_\mu = \{e\}$ all its factors are cyclic. Let $N_\alpha = N^{G_\alpha}$ and L be the compositum of N with $F[\mu_\infty]$ where $F[\mu_\infty]$ denotes the field extension of F obtained by adjoining to F all n-th roots of unity for every natural number n. Obviously, $K \leq L$ and we have the following chain of Galois extensions: $F = L_0 \leq \dots \leq L_{n+1} = L_n [\sqrt[n!]{1}] \leq \dots \leq L_\omega = F[\mu_\infty] \leq L_{\omega+1} = L_\omega N_1 \leq \dots \leq L_{\omega+\alpha} = L_\omega N_\alpha \leq \dots \leq L_{\omega+\mu} = L_\omega N = L$. Obviously, for every n, L_{n+1}/L_n is radical. For every α , $L_{\omega+\alpha+1}/L_{\omega+\alpha}$ is cyclic over a field containing all roots of unity and thus radical too.

For the second direction assume that $K \leq N$ and there is a chain of radical extensions $F = N_0 \leq ... \leq N_\alpha \leq ... N_\mu = N$. Let $L = F[\mu_\infty]N$. Consider the following chain of field extensions: $F = L_0 \leq ... \leq L_{n+1} = L_n[\sqrt[n!]{1}] \leq ... \leq L_n$ $L_{\omega} = F[\mu_{\infty}] \le L_{\omega+1} = L_{\omega}N_1 \le \dots \le L_{\omega+\alpha} = L_{\omega}N_{\alpha} \le \dots \le L_{\omega+\mu} = L_{\omega}N = L.$ We define a new chain of field extensions $\{T_{\alpha}\}_{\alpha < \omega + \mu}$ as follows: Let $T_0 = F$. For every $\alpha < \omega + \mu$ limit define $T_{\alpha} = \bigcup_{\beta < \alpha} T_{\beta}$. For every $\alpha < \omega + \mu$, let a be such that $L_{\alpha+1} = L_{\alpha} [\sqrt[n]{a}]$ and define $T_{\alpha+1} = T_{\alpha} [\sqrt[n]{\sigma(a)} : \sigma \in \operatorname{Gal}(T_{\alpha}/F)]$. We claim that T_{α} is a Galois extension containing L_{α} . We prove it by transfinite induction. When $\alpha = 0$ this is clear. Assume that T_{α} containing L_{α} , then clearly $L_{\alpha+1} = L_{\alpha}[\sqrt[n]{a}] \subseteq T_{\alpha}[\sqrt[n]{\sigma(a)} : \sigma \in \operatorname{Gal}(T_{\alpha}/F)] = T_{\alpha+1}$. Moreover, $T_{\alpha+1}$ is Galois over F as the compositum of T_{α} with the splitting field over F of the polynomial $f_a(x^n)$, where f_a is the minimal polynomial of a over F. When α is a limit ordinal the claim is immediate. Hence $T_{\omega+\mu}$ is a Galois extension containing K. In addition, for every $\alpha < \omega + \mu$, $LT_{\alpha+1}/T_{\alpha}$ is abelian. Let $G_{\alpha} = \operatorname{Stab}_{\operatorname{Gal}(L/F)}(T_{\alpha})$, we get that G_{α} is an accessible series for $\operatorname{Gal}(L/F)$, $G_{\alpha+1}/G_{\alpha}$ is abelian and $\bigcap_{\alpha < \omega + \mu} G_{\alpha} = \{e\}$. We conclude from Corollary 5 that $\operatorname{Gal}(L/F)$ is prosolvable. **Definition 17.** Let K/F be a separable algebraic extension.

- 1. K/F is called *locally solvable* if every finite subextension L/F of K/F is solvable.
- 2. K/F is called *locally solvable by radicals* if every finite subextension L/F of K/F is solvable by radicals.

Remark 18. Observe that by Galois Theorem for finite extensions, K/F is locally solvable if and only if it is locally solvable by radicals.

Lemma 19. Let K/F be a separable algebraic extension, then K/F is locally solvable if and only if it is prosolvable. Consequently, K/F is locally solvable by radicals if and only if it is solvable by radicals.

Proof. First assume that K/F is prosolvable. Let N/F be the Galois closure of K and L/F be some finite subextension of K. Then L'/F, the Galois closure of L/F, is embedded into N/F and there is an epimorphism $\operatorname{Gal}(N/F) \to \operatorname{Gal}(L'/F)$. Hence, $\operatorname{Gal}(L'/F)$ is solvable. On the other hand, if N/F is the Galois closure of K/F, then $N = \bigcup L'/F$ the union of the Galois closures of all finite subextensions L/F inside a given separable closure of F. Thus $\operatorname{Gal}(N/F) = \lim_{\leftarrow} \operatorname{Gal}(L'/F)$ is prosolvable. □

Corollary 20. Any radical extension of F is locally radical. In other words, if a separable element x over F can be obtained by " μ steps" of extracting n'th roots for some ordinal μ and natural numbers n, then x can be obtained by finite steps of extracting n'th roots.

3 Topological vs. Abstract Composition Factors

In this section we study the connection between the composition factors of a profinite group and its profinite completion. Recall that the profinite completion of an abstract group G, which we denote by \hat{G} , is equal to the inverse limit of all its finite quotients with the natural projections between them. Then $\hat{G} = \lim_{N \leq f G} G/N$. Observe that the finite topological quotients of \hat{G} are precisely the finite abstract quotients of G. For more information about profinite completions see [1, Section 3.2].

This definition can in particular be applied for profinite groups G. Clearly, $G = \hat{G}$ if and only if all finite-index (normal) subgroups of G are open in G. Such groups are called *strongly complete* groups. In that case, properties of the complete system of finite abstract quotients of G are reflected in properties of the topological group G.

Although it seems a-priory unlikely to hold in general for all profinite groupsit turns out that some topological properties of a general profinite group are in fact properties of the complete system of its finite abstract quotients, or equivalently, are preserved under profinite completion. Some basic examples are being pro- π (prosolvable). By [1, Proposition 4.2.3 and Corollary 4.2.4], every abstract finite quotient of a pro- π (prosolvable) group is a finite π -group (solvable group). Here π denotes a set of primes and by (pro-) π groups we refer to (inverse limits of) finite groups whose orders are only divided by primes for π . In particular, the profinite completion of a pro- π (prosolvable) group is again a pro- π (prosolvable) group. Other interesting connections between a profinite group and its profinite completion can be found, for example, in [8], [9] and [10]. The importance of understanding those connections between *G* and \hat{G} lies in the fact that the profinite completion of a profinite group encodes, in some sense, the abstract group-theoretic properties of the group which are not immediate topological properties. Understanding these abstract group-theoretic properties of a profinite group helps us moving toward solution of the following question that was posed in [11]:

Question 21. Which abstract groups can be realized as Galois groups?

By the famous correspondence due to Krull and Leptin, this question is equivalent to: what abstract group can be given a profinite group-topology?

We say that a finite group is *anabelian* if all its composition factors are nonabelian. Let us say that a profinite group is anabelian if all its composition factors are non-abelian. By Proposition 4 this is equivalent to being an inverse limit of anabelian finite groups. Inspired by the prosolvable case we suggest the following question:

Question 22. Let G be an anabelian profinite group. Is \hat{G} an anabelian profinite group as well?

Some evidence for a positive answer might be found in the following proposition:

Proposition 23. Let G be an anabelian profinite group. Then G has no finite abstract abelian quotients.

Proof. Since G is anabelian it has no topological finite abelian quotients. Hence there is no proper open subgroup $[G,G] \leq U \leq_o G$. In particular there is no proper open subgroup $\overline{[G,G]} \leq U \leq_o G$. Recall that every closed normal subgroup of G is the intersection of all open normal subgroups containing it, see [1, Proposition 2.1.4 (d)]. We get that $\overline{[G,G]} = G$.

We will show a stronger result: in fact, G has no abelian quotients at all. Equivalently: G = [G, G]. By the preceding discussion it is enough to show that $[G, G] = \overline{[G, G]}$. For that we briefly recall some observations that were made in [5, Pg. 175].

Let $w = w(x_1, ..., x_k)$ be a group word. Denote by $G^{\{w\}} = \{w(g_1, ..., g_k)^{\pm 1} | g_1, ..., g_k \in G\}, (G^{\{w\}})^{*n} = \{s_1 \cdots s_n | s_1, ..., s_n \in G^{\{w\}}\}$ and $w(G) = \bigcup_{n \in \mathbb{N}} (G^{\{w\}})^{*n}$ the abstract subgroup that is generated by all substitutions of elements from G in $w(x_1, ..., x_k)$. Then w(G) is closed if and only if there exists some $n \in \mathbb{N}$ such that $w(G) = (G^{\{w\}})^{*n}$. Moreover, $w(G) = (G^{\{w\}})^{*n}$ if and only if $w(G/N) = (G/N^{\{w\}})^{*n}$ for every open normal subgroup $N \leq_o G$.

By [6, Theorem 2] there exists some constant D such that every element in an anabelian finite group is a product of D commutators. By definition of an anabelian profinite group, we are done.

It is worth mentioning that a profinite group might have finite abstract quotients that does not appear as topological quotients, as described in the following example from [5, Pg. 176].

Example 24. In [7, Lemma 2.2] the author constructed for every prime p and q a power of p a sequence of finite perfect groups K_n such that for each n, $K_n \cong P_n \rtimes SL_2(q)$ for P a finite p-group, and $\log |K_n| / \log |K_n^{\{w\}} \to \infty$ for $w = x^2$. Let $G = \prod_n K_n$. Then by [5, Pg. 176] $G = \overline{[G,G]}$ and thus has no topological finite abelian quotients. However, since $G^2 \leq G$, C_2 appears as an abstract finite quotient of G. One observes that C_2 does appear as a (topological) composition factor of G, since $C_2 \cong Z(SL_2(q))$.

This raises a question regarding the possible connection between the *abstract* and *topological* composition factors of a profinite group G. Let S be finite simple group. We say that S is an *abstract* (respectively *topological*) composition factor of G if it appears as a composition factor of some abstract (respectively topological) finite quotient of G.

Question 25. Does every abstract composition factor of a profinite group G also occur as a topological composition factor of G?

Since the abstract finite quotients of a profinite group are precisely the topological finite quotients of its profinite completion, by Proposition 4 Question 25 is equivalent to the following:

Question 25^{*}. Let G be a profinite group. Do G and \hat{G} have the same composition factors?

Although Question 25 remains widely open, we can prove some partial results. The first one says that a group with large topological composition factors cannot have an abstract finite image which is too small.

Proposition 26. There is a function $f : \mathbb{N} \to \mathbb{N}$ with the following property. Suppose that L is an abstract quotient of a profinite group G. Then G has a topological composition factor L_0 with $|L_0| < f(|L|)$.

Proof. Let q = |L|. By Proposition 10.1 of [5] there is are integers C = C(q) and m = m(q) with the following property:

Let Γ be a finite group with a semisimple normal subgroup N which is a product of simple groups, each of size at least C. For any g_1, \ldots, g_m in G we have

$$\{(g_1z_1)^q \cdots (g_mz_m)^q \mid z_i \in N, i = 1, \dots, m\} = g_1^q g_2^q \cdots g_m^q N.$$
(1)

Set $f = f(|L|) := \max\{C(q), q+1\}$ and suppose that every topological composition factor of G has size at least f. We claim that

$$G = \left\{ g_0 g_1^q g_2^q \cdots g_m^q \mid g_i \in G \text{ and } o(g_0) \in \hat{\mathbb{Z}} \text{ is coprime to } q \right\}.$$
(2)

Assuming this claim it follows that $G = G^q$ and since $L^q = \{1\}$ we conclude that L cannot be an abstract quotient of G, contradiction. Hence G must have a topological composition factor of size at least f.

It remains to prove the (2). The right hand side is a closed subset of G and thus it is enough to show that it is dense in G. Therefore we may assume from now on that G is a finite group. We argue by induction on |G| starting with the case when |G| = 1 when the equality (2) is trivially true. If G is simple we set N = G, otherwise we choose N to be a minimal normal subgroup of G. Let $x \in G$. By the induction hypothesis there are elements $g_0, \ldots, g_m \in G$ with order of g_0N in G/N coprime to q and such that $x \equiv g_0g_1^q \cdots g_m^q \mod N$. If N is semisimple then its simple factors are of size at least C(q) by assumption. In that case the existence of $z_1, \ldots, z_m \in N$ with $x = g_0(g_1z_1)^q \cdots (g_mz_m)^q$ follows directly from (1). If N is elementary abelian then it must be a p-group for some prime p > q and in particular |N| is coprime to q. In that case any element of $g_0N \subset G$ has order in G coprime to q and we can write $x = g'_0g_1^q \cdots, g_m^q$ where $g'_0 := x(g_1^q \cdots, g_m^q)^{-1} \in g_0N$ as required. The claim (2) is proved.

The next Proposition shows that if L is an abstract composition factor of G then the topological composition factors of G cannot be all smaller than L. Recall that for groups A, B we say that A is a section of B if $A \cong C/D$ where $D \leq C \leq B$ with D normal in C. It is easy to show that if H is a closed subgroup of a profinite group G then every topological composition factor of H occurs as a section of some topological composition factor of G.

Proposition 27. Let L be an abstract composition factor of a profinite group G. Then G has a topological composition factor L_0 such that L is a section of L_0 .

Proof. Let N be a normal subgroup of finite index in G such that L is a composition factor of G/N. Choose $a_1, \ldots, a_d \in G$ such that a_1N, \ldots, a_dN generate G/N and let $H := \langle a_1, \ldots, a_d \rangle \leq G$. Note that HN = G and hence $G/N \simeq H/(N \cap H)$. On the other hand H is a finitely generated profinite group and hence strongly complete by [5]. Therefore L occurs as a topological composition factor of H and hence occurs a section in some topological composition factor of G.

In particular since all proper subgroups of the alternating group A_5 are solvable we can deduce the following.

Corollary 28. Let G be a profinite group whose nonabelian topological composition factors are equal to A_5 . Then the nonabelian abstract composition factors of G must also be equal to A_5 .

References

[1] Ribes, L., Zalesskii, P. Profinite groups. Springer, 2010.

- [2] Birkhoff, G. Transfinite subgroup series. Bulletin of the American Mathematical Society, 40 (12): 847–850, 1934.
- [3] Baumslag, B. A simple way of proving the Jordan-Hölder-Schreier theorem. American Mathematical Monthly, 40 (12): 113 (10): 933–935, 2006.
- [4] Erdmann, K. & Holm, T. Simple Modules and the Jordan–Hölder Theorem. Algebras And Representation Theory. pp. 61-84 (2018)
- [5] Nikolov, N., Segal, D. On finitely generated profinite groups, I: strong completeness and uniform bounds. *Annals Of Mathematics*. pp. 171-238 (2007)
- [6] Nikolov, N. Verbal width in anabelian groups. Israel Journal Of Mathematics. 216 pp. 847-876 (2016)
- [7] Holt, D. Enumerating perfect groups. Journal Of The London Mathematical Society. 2, 67-78 (1989)
- [8] Bar-On, T. Profinite completion of free pro-C groups. Journal Of Algebra And Its Applications. pp. 2550338 (2024)
- [9] Bar-On, T. The profinite completion of a profinite projective group. Glasgow Mathematical Journal. 64, 499-503 (2022)
- [10] Bar-On, T. Cohomological properties of maximal pro-*p* Galois groups that are preserved under profinite completion. *Forum Mathematicum*. (2024)
- [11] Bar-On, T. & Nikolov, N. Groups of profinite type and profinite rigidity. Journal Of Group Theory. (2024)