

Schrödinger Bridges for Systems of Interacting Particles

Henri Orland¹

¹*Institut de Physique Théorique
CNRS, CEA
Université Paris-Saclay
France*

(Dated: March 13, 2025)

Abstract: A Schrödinger bridge is the most probable time-dependent probability distribution that connects an initial probability distribution w_i to a final one w_f . The problem has been solved and widely used for the case of simple Brownian evolution (non-interacting particles). It is related to the problem of entropy regularized Wasserstein optimal transport. In this article, we generalize Brownian bridges to systems of interacting particles. We derive some equations for the forward and backward single particle “wave-functions” which allow to compute the most probable evolution of the single-particle probability between the initial and final distributions.

In a seminal article published in 1931 [1–3], Schrödinger, noticing the analogy between his equation for the dynamics of quantum particles and the diffusion equation, which describes the statistics of independent particles performing Brownian motion, proposed and solved the following problem. Consider two sets of probability distributions $w_i(r)$ and $w_f(r)$. There is a finite probability that an ensemble of particles with distribution $w_i(r)$ at time 0, evolves under a Brownian motion to the distribution $w_f(r)$ at time t_f . How can one determine the probability distribution for the particles $w(r, t)$ at intermediate times t ? This is called the *Schrödinger bridge problem* and Schrödinger found an elegant solution to this problem. It is a question of paramount importance in non-equilibrium statistical physics, as it determines how an ensemble of particles in a given density configuration evolves into another one. For an extensive mathematical review, see [4].

The Schrödinger bridge problem has been shown to be equivalent to the entropy regularized optimal transport problem with Wasserstein distance [7]. Due to this fact, the Schrödinger bridge problem has found numerous applications in recent years in e.g. statistical physics [5], transition path theory [6], optimal transport [7], machine learning [8], computer vision [9], finances [10] etc. to name a few. The original problem is dealing with independent free (Brownian) particles. The goal of this article is to generalize the concept to the case of interacting particles.

Consider a system of N particles, interacting with a one-body potential $U(r)$ and a two-body potential $v(r - r')$. The initial coordinates of the particles at $t = 0$ are $\{r_1^i, \dots, r_N^i\}$ and their final coordinates at time t_f are $\{r_1^f, \dots, r_N^f\}$. These configurations define the probability distributions $w_i(r)$ and $w_f(r)$ by

$$w_i(r) = \frac{1}{N} \sum_{k=1}^N \delta(r - r_k^i) \quad (1)$$

$$w_f(r) = \frac{1}{N} \sum_{k=1}^N \delta(r - r_k^f) \quad (2)$$

and the corresponding particle densities by

$$\rho_i(r) = N w_i(r) \quad (3)$$

$$\rho_f(r) = N w_f(r) \quad (4)$$

The particles evolve under overdamped Langevin dynamics [11, 12] (Brownian dynamics)

$$\dot{r}_i = -\frac{1}{\gamma} \nabla_i W(\{r_i\}) + \eta_i(t) \quad (5)$$

where

$$W(\{r_i\}) = \sum_i U(r_i) + \frac{1}{2} \sum_{i \neq j} v(r_i - r_j) \quad (6)$$

and

$$\nabla_i W(\{r_i\}) = \nabla_i U(r_i) + \sum_{j \neq i} \nabla_i v(r_i - r_j) \quad (7)$$

The friction coefficient γ is related to the diffusion coefficient D and to the inverse temperature $\beta = 1/k_B T$, by the Einstein relation $\gamma = 1/\beta D$ and $\eta_i(t)$ is a random Gaussian force, with zero mean and correlation given by

$$\langle \eta_i(t) \eta_j(t') \rangle = 2D \delta(t - t') \quad (8)$$

The trajectories from any point of the initial distribution w_i to any point of the final distribution w_f are governed by the so-called *Brownian Bridge* equation [6, 13]. When searching for the transport of the distribution w_i to w_f , one must sum over all the trajectories that start from any point of w_i and ends up in any point of w_f . This involves some complex combinatorics, which was brilliantly solved by Schrödinger [1]. In this article, we tackle this problem in a completely different manner.

The probability distribution for the particles $P(\{r_i\}, t)$ satisfies the Fokker-Planck equation [11, 12]

$$\frac{dP(\{r_i\}, t)}{dt} = D \sum_i \nabla_i (\nabla_i P + \beta P \nabla_i W) \quad (9)$$

Note that the duration t_f of the process can be eliminated by rescaling the time t in the Langevin or Fokker-Planck equation, provided the diffusion constant and the temperature are rescaled properly. Most of the mathematical literature uses $t_f = 1$. Since, in this article, we emphasize the physical aspect of evolving particles, we shall keep the original time t_f .

Defining the function

$$\Phi(r_i, t) = e^{+\frac{\beta}{2} W(\{r_i\})} P(r_i, t) \quad (10)$$

it satisfies the Schrödinger equation [12, 14]

$$\frac{d\Phi(r_i, t)}{dt} = -H\Phi(r_i, t) \quad (11)$$

$$= D \sum_i \nabla_i^2 \Phi - D\beta^2 V_{eff}(r_i) \Phi(r_i, t) \quad (12)$$

where the Hamiltonian is given by

$$H = -D \sum_i \nabla_i^2 + D\beta^2 V_{eff}(\{r_i\}) \quad (13)$$

and the effective potential is given by

$$V_{eff}(\{r_i\}) = \frac{1}{4} \sum_i (\nabla_i W)^2 - \frac{k_B T}{2} \sum_i \nabla_i^2 W \quad (14)$$

which in terms of the potentials U and v takes the form

$$V_{eff}(\{r_i\}) = \frac{1}{4} \sum_i \left(\nabla_i U(r_i) + \sum_{j \neq i} \nabla_i v(r_i - r_j) \right)^2 - \frac{k_B T}{2} \sum_i \left(\nabla_i^2 U(r_i) + \sum_{j \neq i} \nabla_i^2 v(r_i - r_j) \right) \quad (15)$$

Note that the first term is a 3-body interaction, whereas the second is a 2-body one.

The probability that the set of particles starting at $\{r_1^i, \dots, r_N^i\}$ at time 0 ends in any permutation of the set $\{r_1^f, \dots, r_N^f\}$ at time t_f is given by

$$P(\{r_k^f\}, t_f | \{r_k^i\}, 0) = \sum_{p \in S_N} P(r_{p(1)}^f, \dots, r_{p(N)}^f, t_f | r_1^i, \dots, r_n^i, 0)$$

where the summation is over all permutations p in the symmetric group S_N .

Using the quantum mechanical Dirac notation [12, 15, 16], the probability for the system to evolve from the initial configuration to the final one is given by

$$P(\{r_k^f\}, t_f | \{r_k^i\}, 0) = \frac{1}{N!} e^{-\frac{\beta}{2}(W(\{r_k^f\}) - W(\{r_k^i\}))} \langle r_1^f, \dots, r_N^f | e^{-t_f H} | r_1^i, \dots, r_n^i \rangle \quad (16)$$

where the brackets $|r_1 \dots r_N\rangle = \sum_{p \in S_N} |r_{p(1)} \dots r_{p(N)}\rangle$ denote the symmetrized (bosonic) state summed over all permutations p of the symmetric group S_N . The factor $1/N!$ cancels one of the summations over permutations since the matrix element above comprises two such summations. In terms of the particle density distributions ρ_i and ρ_f , the pre-factor above can be written as

$$\begin{aligned} W(\{r_k^f\}) - W(\{r_k^i\}) &= \int dr U(r) (\rho_f(r) - \rho_i(r)) \\ &+ \frac{1}{2} \int dr dr' v(r - r') (\rho_f(r) \rho_f(r') - \rho_i(r) \rho_i(r')) \end{aligned} \quad (17)$$

In the following of this paper, we will heavily rely on coherent states [17]. We recall that coherent states constitute an overcomplete set of vectors in the Fock space of the system. The two fundamental identities that allow calculations with coherent states $|\phi\rangle$ are i) the scalar product

$$\langle \phi | \phi' \rangle = e^{\int dr \phi^*(r) \phi'(r)} \quad (18)$$

and ii) the decomposition of unity

$$\int d\phi^*(r) d\phi(r) e^{-\int dr \phi^*(r) \phi(r)} |\phi\rangle \langle \phi| = \mathbb{1} \quad (19)$$

where $\phi(r)$ and $\phi^*(r)$ are complex conjugate fields and $\mathbb{1}$ is the unit operator defined by $\langle r | \mathbb{1} | r' \rangle = \delta(r - r')$. In addition, we have

$$\langle r_1 \dots r_N | \phi \rangle = \phi(r_1) \dots \phi(r_N) \quad (20)$$

$$\langle \phi | r_1 \dots r_n \rangle = \phi^*(r_1) \dots \phi^*(r_n) \quad (21)$$

Using these identities, we have

$$\begin{aligned} \langle r_1^f, \dots, r_N^f | e^{-t_f H} | r_1^i, \dots, r_n^i \rangle &= \int d\phi_f^*(r) d\phi_f(r) \int d\phi_i^*(r) d\phi_i(r) e^{-\int dr (\phi_f^*(r) \phi_f(r) + \phi_i^*(r) \phi_i(r))} \\ &\times \phi_f(r_1^f) \dots \phi_f(r_N^f) \phi_i^*(r_1^i) \dots \phi_i^*(r_n^i) \langle \phi_f | e^{-t_f H} | \phi_i \rangle \end{aligned} \quad (22)$$

The coherent state matrix element above can be represented as a functional integral over complex fields [17] as

$$\begin{aligned} \langle \phi_f | e^{-t_f H} | \phi_i \rangle &= \int_{\phi(r,0)=\phi_i(r)}^{\phi^*(r,t_f)=\phi_f^*(r)} \mathcal{D}(\phi^*, \phi) \exp \left\{ \int dr \phi_f^*(r) \phi_f(r) \right. \\ &\left. - \int_0^{t_f} dt \int dr \left(\phi^*(r, t) \left(\frac{\partial \phi}{\partial t} - D \nabla^2 \right) \phi(r, t) \right) - \int_0^{t_f} dt A(\phi^*, \phi) \right\} \end{aligned} \quad (23)$$

where the interaction term is given by

$$\begin{aligned} A(\phi^*, \phi) &= D\beta^2 \int dr |\phi(r, t)|^2 \left(\frac{1}{4} \left(\nabla U(r) + \int dr' \nabla v(r - r') |\phi(r', t)|^2 \right)^2 \right. \\ &\left. - \frac{k_B T}{2} \left(\nabla^2 U(r) + \int dr' \nabla^2 v(r - r') |\phi(r', t)|^2 \right) \right) \end{aligned} \quad (24)$$

$$= D\beta^2 \int dr \rho(r, t) \left(\frac{1}{4} \left(\nabla U(r) + \int dr' \nabla v(r-r') \rho(r', t) \right)^2 \right) \quad (25)$$

$$- \frac{k_B T}{2} \left(\nabla^2 U(r) + \int dr' \nabla^2 v(r-r') \rho(r', t) \right) \quad (26)$$

where

$$\rho(r, t) = \phi^*(r, t) \phi(r, t) = |\phi(r, t)|^2$$

This term can be rewritten as

$$A(\phi^*, \phi) = D \int dr \rho(r, t) e^{\frac{\beta}{2} (U(r) + \int dr' v(r-r') \rho(r', t))} \nabla^2 e^{-\frac{\beta}{2} (U(r) + \int dr' v(r-r') \rho(r', t))} \quad (27)$$

Using the identity

$$\begin{aligned} \phi_f(r_1^f) \dots \phi_f(r_N^f) \phi_i^*(r_1^i) \dots \phi_i^*(r_N^i) &= \exp \left(\sum_k \left(\log \phi_f(r_k^f) + \log \phi_i^*(r_k^i) \right) \right) \\ &= \exp \left(\int dr (\rho_f(r) \log \phi_f(r) + \rho_i(r) \log \phi_i^*(r)) \right) \end{aligned} \quad (28)$$

we finally obtain the expression for the probability for the system to make a transition from the probability distribution $w_i(r)$ to $w_f(r)$ during time t_f :

$$\begin{aligned} P(w_f, t_f | w_i, 0) &= \frac{1}{N!} \exp \left(-\beta \int dr U(r) (\rho_f(r) - \rho_i(r)) \right. \\ &\quad \left. - \frac{\beta}{2} \int dr dr' v(r-r') (\rho_f(r) \rho_f(r') - \rho_i(r) \rho_i(r')) \right) \\ &\quad \times \int \mathcal{D}(\phi^*, \phi) \exp \left\{ - \int dr \phi^*(r, 0) \phi(r, 0) \right. \\ &\quad \left. - \int_0^{t_f} dt \int dr \left(\phi^*(r, t) \left(\frac{\partial \phi}{\partial t} - D \nabla^2 \right) \phi(r, t) \right) - \int_0^{t_f} dt A(\phi^*, \phi) \right. \\ &\quad \left. + \int dr (\rho_f(r) \log \phi(r, t_f) + \rho_i(r) \log \phi^*(r, 0)) \right\} \end{aligned} \quad (29)$$

where $A(\phi^*, \phi)$ is given by (24).

Note that the boundary conditions on the functional integral are not present anymore, since ϕ_f^*, ϕ_i are fully integrated over. To complement these equations, we note that the conditioned density $\rho_c(r, t)$ of particles at point r at time t is given by

$$\rho_c(r, t) = \left\langle |\phi(r, t)|^2 \right\rangle \quad (30)$$

where the bracket denotes the expectation value with the probability defined in eq. (29)

The one-body (non-interacting) case

We first consider the free case $v = 0$. In that case, there is no interaction term and the probability becomes

$$\begin{aligned}
P(w_f, t_f | w_i, 0) &= \frac{1}{N!} \exp \left(-\beta \int dr U(r) (\rho_f(r) - \rho_i(r)) \right) \\
&\times \int \mathcal{D}(\phi^*, \phi) \exp \left\{ - \int dr \phi^*(r, 0) \phi(r, 0) \right. \\
&- \int_0^{t_f} dt \int dr \left(\phi^*(r, t) \left(\frac{\partial \phi}{\partial t} - D \nabla^2 + \frac{D \beta^2}{4} \left((\nabla U(r))^2 - 2k_B T \nabla^2 U(r) \right) \right) \phi(r, t) \right) \\
&\left. + \int_0^{t_f} dr \left(\rho_f(r) \log \phi(r, t_f) + \rho_i(r) \log \phi^*(r, 0) \right) \right\}
\end{aligned} \tag{31}$$

Since this functional integral is Gaussian, the saddle-point approximation is exact. Note that although the fields ϕ and ϕ^* are complex conjugate in the functional integral, in the saddle-point approximation, they no longer need to be complex conjugate from each other, and they are to be treated as two independent complex fields.

The equations extremizing the exponent read:

$$\left(\frac{\partial}{\partial t} - D \nabla^2 + \frac{D \beta^2}{4} \left((\nabla U(r))^2 - 2k_B T \nabla^2 U(r) \right) \right) \phi(r, t) = 0 \tag{33}$$

$$\left(-\frac{\partial}{\partial t} - D \nabla^2 + \frac{D \beta^2}{4} \left((\nabla U(r))^2 - 2k_B T \nabla^2 U(r) \right) \right) \phi^*(r, t) = 0 \tag{34}$$

with the boundary conditions (obtained by extremizing the exponent w.r.t. the fields at 0 and t_f)

$$\phi(r, 0) = \frac{\rho_i(r)}{\phi^*(r, 0)} \tag{35}$$

$$\phi^*(r, t_f) = \frac{\rho_f(r)}{\phi(r, t_f)} \tag{36}$$

$$\tag{37}$$

Note that (34) is the time-reversed equation of (33). However, the boundary conditions above prevent the expression of $\phi^*(r, t)$ in terms of $\phi(r, t)$.

We define the Green's function $G(rt, r't')$ as

$$G(rt, r't') = \left(\frac{\partial}{\partial t} - D \nabla^2 + \frac{D \beta^2}{4} \left((\nabla U(r))^2 - 2k_B T \nabla^2 U(r) \right) \right)^{-1} (rt, r't') \tag{38}$$

solution of the equation

$$\left(\frac{\partial}{\partial t} - D \nabla^2 + \frac{D \beta^2}{4} \left((\nabla U(r))^2 - 2k_B T \nabla^2 U(r) \right) \right) G(rt, r't') = \delta(r - r') \delta(t - t') \tag{39}$$

In the absence of an external potential U , in space dimension d , this Green's function is

$$G_0(rt, r't') = \left(\frac{1}{4\pi D(t-t')} \right)^{d/2} e^{-\frac{(r-r')^2}{4D(t-t')}} \tag{40}$$

Another simple case when this Green's function can be computed exactly is that of the harmonic oscillator (Ornstein-Uhlenbeck process)

$$U(r) = \frac{1}{2} K r^2 \tag{41}$$

We introduce the notation

$$\phi_i(r) = \phi(r, 0) = \frac{\rho_i(r)}{\phi^*(r, 0)} \tag{42}$$

$$\phi_f^*(r) = \phi^*(r, t_f) = \frac{\rho_f(r)}{\phi(r, t_f)} \tag{43}$$

Eq. (33,34) with their boundary conditions can be solved as

$$\begin{aligned}\phi(r, t) &= \int dr' G(rt, r'0) \phi_i(r') \\ \phi^*(r, t) &= \int dr' \phi_f^*(r') G(r't_f, rt)\end{aligned}$$

which implies

$$\rho_f(r) = \phi_f^*(r) \int dr' G(rt_f, r'0) \phi_i(r') \quad (44)$$

$$\rho_i(r) = \phi_i(r) \int dr' \phi_f^*(r') G(r't_f, r0) \quad (45)$$

These are the standard Schrödinger bridge equations. They are the same equations as those of the entropy-regularized optimal transport problem [7], with the Wassestein cost function $(r-r')^2/4t_f$. Therefore, they can be solved efficiently by using the Sinkhorn fixed-point method [18]. Starting from a guess for $\phi_i(r)$, we obtain $\phi_f^*(r)$ from eq.(44). We plug it in (45) to obtain a new $\phi_i(r)$, and the procedure is iterated until convergence. Convergence is guaranteed and very fast ([19]).

From eq.(30) and since in the saddle-point approximation the expectation value is identical to the actual value, the conditioned particle density is given by

$$\begin{aligned}\rho_c(r, t) &= |\phi(r, t)|^2 \\ &= \int dr' dr'' \phi_f^*(r') G(r't_f, rt) G(rt, r''0) \phi_i(r'')\end{aligned}$$

which satisfies the boundary conditions $\rho_c(r, 0) = \rho_i(r)$ and $\rho_c(r, t_f) = \rho_f(r)$.

Finally, the transition probability is given by

$$\begin{aligned}P(w_f, t_f | w_i, 0) &= N! \exp\left(-\beta \int dr U(r) (\rho_f(r) - \rho_i(r))\right) \\ &\times \exp\left(N \int dr \left[w_i(r) \log \frac{w_i(r)}{\phi_i(r)} + w_f(r) \log \frac{w_f(r)}{\phi_f^*(r)} \right]\right)\end{aligned} \quad (46)$$

We recognize the Kullback-Leibler (KL) relative entropy [20] of the distributions ϕ_i and ϕ_f^* with respect to the distributions w_i and w_f . This expression shows the connection of Schrödinger bridges with large deviations theory and optimal transport.

The interacting case

We now consider the interacting case. For the sake of simplicity, we shall assume that $U = 0$. The expression for P is still given by eq.(29), however the expression (24) simplifies as

$$\begin{aligned}A(\phi^*, \phi) &= D\beta^2 \int dr \rho(r, t) \left(\frac{1}{4} \left(\int dr' \nabla v(r-r') \rho(r', t) \right)^2 - \frac{k_B T}{2} \int dr' \nabla^2 v(r-r') \rho(r', t) \right) \\ &= D \int dr \rho(r, t) e^{\frac{\beta}{2} \int dr' v(r-r') \rho(r', t)} \nabla^2 e^{-\frac{\beta}{2} \int dr' v(r-r') \rho(r', t)}\end{aligned} \quad (47)$$

where $\rho(r, t) = |\phi(r, t)|^2 = Nw(r, t)$, and $w(r, t)$ is the probability distribution at time t .

Due to the presence of implicit factors of N in A through $\rho(r, t)$, the saddle-point approximation is not exact. However, it provides the most probable probability distribution path connecting w_i and w_f .

The saddle point equations are

$$\left(\frac{\partial}{\partial t} - D\nabla^2 - D\beta \int dr' \nabla^2 v(r-r') |\phi(r', t)|^2 + \frac{D\beta^2}{4} \left(\int dr' \nabla v(r-r') |\phi(r', t)|^2 \right)^2 \right)$$

$$+ \frac{D\beta^2}{2} \int dr' dr'' |\phi(r', t)|^2 \nabla v(r' - r'') |\phi(r'', t)|^2 \nabla v(r' - r) \Big) \phi(r, t) = 0 \quad (48)$$

$$\left(-\frac{\partial}{\partial t} - D\nabla^2 - D\beta \int dr' \nabla^2 v(r - r') |\phi(r', t)|^2 + \frac{D\beta^2}{4} \left(\int dr' \nabla v(r - r') |\phi(r', t)|^2 \right)^2 \right. \\ \left. + \frac{D\beta^2}{2} \int dr' dr'' |\phi(r', t)|^2 \nabla v(r' - r'') |\phi(r'', t)|^2 \nabla v(r' - r) \right) \phi^*(r, t) = 0 \quad (49)$$

with the boundary conditions

$$\phi(r, 0) = \frac{\rho_i(r)}{\phi^*(r, 0)} \quad (50)$$

$$\phi^*(r, t_f) = \frac{\rho_f(r)}{\phi(r, t_f)} \quad (51)$$

As mentioned in the non-interacting case, (49) is the time-reversed equation of (48).

As before, the conditioned density of particles at r at time t is given by

$$\rho_c(r, t) = |\phi(r, t)|^2$$

As in the non-interacting case, these equations can be solved numerically by a Sinkhorn-like fixed-point method. One first guesses the function $\rho(r, t)$, with the condition $\rho(r, 0) = \rho_i(r)$ and $\rho(r, t_f) = \rho_f(r)$. For example, one can use $\rho(r, t) = \frac{t_f - t}{t_f} \rho_i(r) + \frac{t}{t_f} \rho_f(r)$ as an initial guess. One must also guess $\phi(r, 0)$. With this two functions, one can solve (48) to obtain $\phi(r, t_f)$. Using (51), we obtain $\phi^*(r, t_f)$ which is used as an initial condition for (49) to obtain $\phi^*(r, 0)$. Using (50), we get $\phi(r, 0)$ and we can recalculate the density ρ and iterate the procedure until convergence.

The above equations can be used to express the probability at the saddle-point as

$$P(w_f, t_f | w_i, 0) = N! \exp \left(-\frac{\beta}{2} \int dr dr' v(r - r') (\rho_f(r) \rho_f(r') - \rho_i(r) \rho_i(r')) \right) \\ \times \exp \left\{ -\frac{D\beta}{2} \int_0^{t_f} dt \int dr dr' \rho(r, t) \nabla^2 v(r - r') \rho(r', t) \right. \\ \left. + \frac{D\beta^2}{2} \int_0^{t_f} dt \int dr \rho(r, t) \left(\int \nabla v(r - r') \rho(r', t) \right)^2 \right. \\ \left. + N \int dr \left(w_f(r) \log \frac{w_f(r)}{\phi(r, t_f)} + w_i(r) \log \frac{w_i(r)}{\phi^*(r, 0)} \right) \right\}$$

Again, the KL relative entropy with respect to the original densities appears in this expression.

One-dimensional example

We now look at a specific one-dimensional example, namely

$$v(x) = g\delta(x)$$

The interaction term (24) becomes

$$A(\phi^*, \phi) = D\beta^2 \int dx \rho(x, t) \left(\frac{g^2}{4} \rho'^2(x, t) - \frac{g}{2} k_B T \rho''(x, t) \right) \quad (52)$$

Equations (48), (49) become

$$\left(\frac{\partial}{\partial t} - D \frac{\partial^2}{\partial x^2} - D\beta g \rho''(x, t) - \frac{D\beta^2 g^2}{4} \rho'^2(x, t) \right)$$

$$- \frac{D\beta^2 g^2}{2} \rho(x, t) \rho''(x, t) \Big) \phi(x, t) = 0 \quad (53)$$

$$\left(-\frac{\partial}{\partial t} - D \frac{\partial^2}{\partial x^2} - D\beta g \rho''(x, t) - \frac{D\beta^2 g^2}{4} \rho'^2(x, t) \right. \\ \left. - \frac{D\beta^2 g^2}{2} \rho(x, t) \rho''(x, t) \right) \phi^*(x, t) = 0 \quad (54)$$

with the boundary conditions

$$\phi(x, 0) = \frac{\rho_i(x)}{\phi^*(x, 0)} \quad (55)$$

$$\phi^*(x, t_f) = \frac{\rho_f(x)}{\phi(x, t_f)} \quad (56)$$

The corresponding probability is given by

$$P(w_f, t_f | w_i, 0) = N! \exp \left(-\frac{\beta g}{2} \int dx (\rho_f^2(x) - \rho_i^2(x)) \right) \quad (57)$$

$$\times \exp \left\{ -\frac{D\beta g}{2} \int_0^{t_f} dt \int dx \rho(x, t) \rho''(x, t) \right. \quad (58)$$

$$+ \frac{D\beta^2 g^2}{2} \int_0^{t_f} dt \int dx \rho(x, t) \rho'^2(x, t) \\ \left. + \int dx \left(w_f(x) \log \frac{w_f(x)}{\phi(x, t_f)} + w_i(x) \log \frac{w_i(x)}{\phi^*(x, 0)} \right) \right\} \quad (59)$$

To illustrate the method, we have solved numerically equations (53) and (54) for some typical values of g . We impose hard-wall boundary conditions. The initial and final distributions are identical Gaussians separated by a distance d

$$\rho_{i,f}(x) = \frac{N}{\sqrt{4\pi\sigma^2}} e^{-\frac{(x \mp d/2)^2}{2\sigma^2}} \quad (60)$$

The number of particles is $N = 100$, the distance $d = 10$, diffusion constant $D = T = 2$. We first study the case of no interaction $g = 0$ (a) and repulsive δ interaction $g = 0.03$ (b). The iteration method does not converge for larger values of g . Since the interaction energy of the system scales like $g\rho^2V$ where ρ is the average density of the system and V its volume, the natural scaling of g is $g \sim 1/\rho^2$. In our numerical example, $N = 100$ and the typical volume of the system is $V \sim 20$, which implies an average density $\rho \sim 5$. This means that g should be of the order 0.04. As can be seen in fig.(1), the effect of repulsive interaction is small, barely visible. This is expected.

The case of attractive interaction is more spectacular. In fig.(2), we plot the sequence of distributions between the two boundary distributions for $g = -0.1$.

As shown in fig.(2), the distribution remains much more concentrated than in the zero or repulsive case during the transition, due to the attractive interactions. In all cases, the fixed-point method does not converge for larger values of g .

In this article, we have generalized the concept of Schrödinger bridges to the case of interacting particles. We derived equations for the most probable distribution path between two boundary distributions. We have checked that these equations can be solved numerically by a generalization of the Sinkhorn algorithm, at least in some simple one-dimensional cases. This generalization should prove very useful in the non-equilibrium statistical physics of many-particle systems. Also, it would be interesting to adapt the above method to the study of bridges in the real-time Schrödinger equation, in order to study phenomena such as diffusion, echo, etc.. However, further studies are needed to devise reliable numerical methods to solve these equations for stronger couplings.




Figure 1: Snapshot of the distribution of particles between the initial (right) and final (left) distributions (thick black lines). The arrow denotes the direction of increasing time




Figure 2: $g = -0.1$ Attractive case

Acknowledgments

The author wishes to thank Prof. C.Leonard for introducing him to the problem of Schrödinger bridges and Prof. K. Mallick for illuminating discussions.

-
- [1] E.Schrödinger, *Sitzungsberichte der Preussischen Akademie der Wissenschaften*, Physikalisch-mathematische Klasse, **144**, 144 (1931) ([document](#))
 - [2] E.Schrödinger, *Sur la théorie relativiste de l'électron et l'interprétation de la mécanique quantique*, Annales de l'I.H.P., tome 2, 4 (1932), 269-310.
 - [3] R.Chetrite, P.Muratore-Ginanneschi and K.Schwieger, *Eur. Phys. J. H* (2021) 46:28. ([document](#))
 - [4] C.Leonard, *Discrete and Continuous Dynamical Systems*, **34**,4, April 2014. ([document](#))
 - [5] C.Leonard, *Journal of Functional Analysis*, **262** 4, 15 February 2012, 1879-1920. ([document](#))
 - [6] H.Orland, *J. Chem. Phys.* **134**, 174114 (2011). ([document](#))
 - [7] R.Otto and C.Villani, *Journal of Functional Analysis*, **173**, 2, 1 June 2000, 361-400. ([document](#))
 - [8] A.Genevay, G.Peyré, and M. Cuturi, *Learning generative models with Sinkhorn divergences* in International Conference on Artificial Intelligence and Statistics,1608-1617, PMLR, 2018. ([document](#))
 - [9] G.Peyré and M.Cuturi, *Computational Optimal Transport: With Applications to Data Science*, Foundations and Trends in Machine Learning: **11**:5-6, pp 355-607 (2019). ([document](#))
 - [10] M.Beiglböck, P.Henry-Labordère and F.Penkner, *Finance Stoch* **17**, 477-501 (2013). ([document](#))
 - [11] N.G.Van Kampen, *Stochastic Processes in Physics and Chemistry*. North-Holland (2007). ([document](#))
 - [12] R.Elber, D.E.Makarov and H.Orland, *Molecular Kinetics in Condensed Phases*, Wiley (2020). ([document](#))
 - [13] D. Revuz and M. Yor, *Continuous Martingales and Brownian Motion*, Springer-Verlag, Berlin, (1991). ([document](#))

- [14] R.Zwanzig, *Nonequilibrium Statistical Mechanics*, Oxford University Press (2001). ([document](#))
- [15] R.P.Feynman and A.R.Hibbs, *Quantum Mechanics and Path Integrals* McGraw-Hill (1965). ([document](#))
- [16] C.Cohen-Tannoudji, B.Diu and F.Laloe, *Quantum Mechanics* Wiley (1977). ([document](#))
- [17] J.W.Negele and H.Orland, *Quantum Many-Particle Systems* (Advanced Books Classics), Perseus Books (1998). ([document](#))
- [18] M.Cuturi in *Advances in Neural Information Processing Systems*, C.J. Burges, L. Bottou, M. Welling, Z. Ghahramani and K.Q. Weinberger ed., **26** (2013) ([document](#))
- [19] P.A.Knight, *The Sinkhorn-Knopp Algorithm: Convergence and Applications*, SIAM Journal on Matrix Analysis and Applications, **30** (1), 261-275 (2008). ([document](#))
- [20] S.Kullback and R.A.Leibler, *On information and sufficiency*, Annals of Mathematical Statistics. **22** (1): 79-86 (1951). ([document](#))