Song $\operatorname{He}^{a,b,c,e_1}$, Yi Li^{d,2}, Hao Ouyang^{c,3}, Yuan Sun^{d4}

^aInstitute of Fundamental Physics and Quantum Technology,

Ningbo University, Ningbo, Zhejiang 315211, China

^bSchool of Physical Science and Technology, Ningbo University, Ningbo, 315211, China

^cCenter for Theoretical Physics and College of Physics, Jilin University,

Changchun 130012, People's Republic of China

^dInstitute of Quantum Physics, School of Physics, Central South University,

Changsha 418003, China

^eMax Planck Institute for Gravitational Physics (Albert Einstein Institute),

Am Mühlenberg 1, 14476 Golm, Germany

Abstract

This review explores recent advances in the theory of $T\bar{T}$ deformation, an irrelevant yet solvable deformation of quantum field theories defined via the quadratic form of the energy-momentum tensor. It addresses classical and quantum aspects, highlighting significant developments across various fields, including field theory, holography, and string theory. Classically, $T\bar{T}$ deformation manifests through multiple geometric interpretations, notably random geometry, Jackiw-Teitelboim-like gravity, and uniform light-cone gauge frameworks. For quantum aspects, the deformation introduces notable features such as non-locality, UV-IR mixing, solvable renormalization structures, and intriguing modifications to correlation functions and entanglement properties. Furthermore, the paper examines the profound relationship between $T\bar{T}$ deformation and holography, particularly within the mixed boundary conditions/cutoff AdS holography proposal and holographic entanglement entropy. Connections to string theory through single-trace deformations and their holographic duals further reveal the deformed structure of the worldsheet. This review synthesizes recent developments and outlines potential directions for future research in the study of $T\bar{T}$ -like deformation.

¹hesong@nbu.edu.cn

²liyi@fudan.edu.cn

³haoouyang@jlu.edu.cn

⁴sunyuan@csu.edu.cn

Contents

1	Introduction	2
2	Basics of $T\bar{T}$ deformation	5
	2.1 The $T\bar{T}$ deformation	5
	2.2 Deformed Lagrangians	7
	2.3 The method of characteristics	9
	2.4 Geometric realization of $T\bar{T}$ deformation	11
	2.5 Uniform light-cone gauge approach	15
	2.6 $T\bar{T}$ -deformed conformal symmetry $\ldots \ldots \ldots$	17
3	$T\overline{T}$ deformation in quantum field theory	19
	3.1 Deformed spectrum and partition function	19
	3.2 Renormalization	21
	3.3 Correlation functions	22
	3.4 Correlation functions: further developments	32
	3.5 Quantum entanglement in $T\overline{T}$ deformed field theory	36
	3.6 $T\bar{T}$ flow at large central charge $\ldots \ldots \ldots$	37
4	$T\bar{T}$ deformation and holography	43
	4.1 Mixed boundary condition, cutoff and glue-on	43
	4.2 Holographic partition function and entanglement entropy	48
	4.3 Holographic correlation functions of the energy-momentum tensor $\ldots \ldots \ldots$	51
	4.4 Symmetries of boundary gravitons at a finite cutoff	53
	4.5 Further developments on holography	56
5	$T\bar{T}$ and string theory	59
	5.1 Little string theory perspective	64
	5.2 TsT and single trace $T\bar{T}$	66
6	Recent Developments and Perspectives	70
	6.1 Generalized $T\bar{T}$ -Like Deformations	70
	6.2 Some perspectives	75
A	Appendix: Useful integrals	77
	A.1 Dimensional regularization	77

1 Introduction

The study of deformations in quantum field theories (QFTs) has provided crucial insights into the structure and dynamics of physical models. A notable example is the $T\overline{T}$ deformation [1,2], first introduced by Zamolodchikov in 2004 [3], which has attracted significant attention due to its remarkable properties, such as solvability, integrability, and non-trivial modifications to the system's dynamics. This deformation, which couples the energy-momentum tensor to itself, offers a non-local yet solvable modification of two-dimensional quantum field theories while preserving key physical properties.

One of the primary motivations for reviewing energy-momentum tensor-induced deformations, such as the $T\overline{T}$ deformation, stems from their ability to provide new insights into nonlocality and ultraviolet (UV) completion of QFTs. Unlike most irrelevant deformations that typically lead to complex, non-renormalizable theories, the $T\overline{T}$ deformation is solvable (and integrable), suggesting that such deformations can be physically meaningful under certain conditions. As this area of research grows, we expect this review will help identify new connections and directions in the study of quantum gravity [4–11], holography [12–31], and string theory [32–35]. The new variant $\sqrt{T\overline{T}}$ of stress tensor flows inspired by the two-dimensional $T\overline{T}$ deformation has been proposed by [36–42]. More recently, the generic $T\overline{T}$ -like deformation has been reviewed in [43].

The study of these deformations spans multiple perspectives beyond standard solvability, integrability, and non-local properties often detailed in early review papers [44, 45], including but not limited to:

To understand how the graviton emerges from the field theory

We propose a thought experiment starting from the action of a massless boson in flat spacetime. According to Einstein's equations, this action necessarily induces an energy-momentum tensor, which must result in spacetime curvature. Simultaneously, the curvature alters the background metric of the massless boson's action or, equivalently, modifies the graviton. This modification leads to interactions between the massless boson and the graviton, inducing an interaction that couples the boson and graviton. This process demonstrates the transition from field theory on a fixed background to a gravitational theory coupled with matter field theory.

Studying field theories deformed by $T\overline{T}$ -like operators reveals a process remarkably similar to this thought experiment. Beginning with an undeformed field theory, coupling it to a $T\overline{T}$ -like deformation operator drives the flow of the field equations. Classically, the $T\overline{T}$ deformation in two dimensions corresponds to coupling undeformed field theories to flat-space Jackiw-Teitelboim-like gravity. This coupling transforms the original seed theory into a gravitational theory, offering a classical framework to explore the emergence of gravity from field theory. One can refer to [46] for higher dimensional realization. Although classical, this process provides valuable insights into how gravity may arise from field theory. For example, a free boson subjected to a $T\overline{T}$ deformation evolves into a non-critical string theory in a static gauge [2, 47]. Upon quantization, this theory inevitably produces gravitons, assuming the exclusion of unstable particles such as tachyons. This highlights the importance of studying $T\overline{T}$ -like deformations as they offer a mechanism for generating gravitons.

However, understanding this mechanism at the quantum level (for higher dimensional generalizations as in [46]) becomes a significant challenge, motivating a comprehensive review of $T\overline{T}$ -like deformations. While numerous technical issues remain resolved, this motivation underscores the critical role of energy-momentum tensor deformations in shedding light on the emergence of gravitational interactions.

Correlation function in the deformed field theories

Correlation functions are essential for understanding the properties of both undeformed and deformed theories. They are central in holography and black hole physics, particularly in quantum chaos and entanglement. Out-of-time-order correlation functions and K-complexity, key tools for diagnosing chaotic behavior in deformed field theories, highlight their significance. Similarly, the computation of entanglement entropy relies heavily on these functions. Constructing correlation functions is crucial for exploring the core features of deformed theories. Although $T\overline{T}$ is an irrelevant deformation, typically precluding a well-defined renormalized theory with a removable UV cutoff, the $T\overline{T}$ deformed theories is essential to understanding their non-local behavior. To consider correlation functions in nonlocal theory like $T\overline{T}$ deformed ones, the operator may be best defined in momentum space and thus is a non-local operator in coordinates space [49, 50].

Celestial holographical dual of the $T\overline{T}$ deformed field theory

At the classical level, a key observation is that the $T\overline{T}$ deformation of a massless free scalar field theory corresponds to the Nambu-Goto action in static gauge. The inherent non-locality of $T\overline{T}$ -deformed theories has been identified in studies of the effective dynamics of long relativistic strings [51, 52]. The connection between $T\overline{T}$ -deformed theories and string theory has been extensively explored in various works [35, 53–57]. Celestial holography provides a framework where quantum scattering amplitudes of massless particles in four-dimensional asymptotically Minkowskian spacetimes are reinterpreted as correlation functions on the celestial sphere at null infinity, defining celestial conformal field theories (CFTs) [58–60]. Based on the construction in [61], it can be proposed that a two-dimensional $T\overline{T}$ deformation of celestial CFTs may correspond to a four-dimensional gravitational theory, offering potential insights into the construction of a UV-complete formulation of General Relativity.

Duality and symmetry of $T\overline{T}$ -like deformed theories

The recent revival of interest in nonlinear electrodynamics (NED) has been significantly

driven by the exploration of $T\overline{T}$ -like deformations, particularly through the introduction of Modified Maxwell (ModMax) theory [62]. This theory, along with the discovery that both Born-Infeld and ModMax can be derived from the Maxwell theory via Lagrangian flows involving composite fields [63–65], has sparked a renewed focus on NED. A central theme in NED is the incorporation of interaction terms into Maxwell's theory while preserving the SO(2)duality symmetry, which deepens our understanding of these systems. Various methods have been proposed to construct self-dual theories of Maxwell's fields with SO(2) duality symmetry, including the Gaillard-Zumino approach [66, 67], the Hamiltonian method by Henneaux and Teitelboim [68], and the PST approach by Pasti, Sorokin, and Tonin [69–72]. These methods provide different frameworks for incorporating interaction terms while preserving duality. This review paper is motivated by the need to explore duality and symmetry in NED theories through $T\overline{T}$ -like deformations, which offer a novel approach for constructing effective Lagrangians that respect these duality symmetries, as demonstrated in previous studies of $T\overline{T}$ -like and marginal root- $T\overline{T}$ -like deformations in NED theories [39, 63] and scalar field theory [73].

This review consolidates the knowledge on the $T\overline{T}$ deformation and its generalized versions, spanning from their origins to recent developments. It aims to serve a broad audience, from students to experts. While several reviews already exist in the literature [43–45,74,75], this work focuses on recent theoretical advancements associated with the deformations induced by energymomentum tensor. We highlight their importance in quantum field theory, holography, and string theory, exploring their implications for spacetime structure, non-locality in QFTs, and the interplay between classical and quantum deformations. By synthesizing these developments, this review offers a helpful resource for researchers involved in the ongoing study of energymomentum tensor-induced deformations.

The structure of this review is organized as follows. Section 2 introduces the fundamental concepts of $T\overline{T}$ deformation, beginning with its definition, key properties, and the formulation of the deformed action. We also discuss various geometrical realizations, such as random geometry, Jackiw-Teitelboim gravity, and the uniform light-cone gauge approach. Section 3 delves into the quantum aspects of $T\overline{T}$ deformation, examining its impact on the spectrum, partition function, renormalization, correlation functions, and quantum entanglement. In Section 4, we investigate the relationship between $T\overline{T}$ deformation and holography, with a focus on the cutoff AdS/ $T\overline{T}$ CFT proposal, holographic partition functions, and entanglement entropy. Section 5 briefly introduces single-trace $T\overline{T}$ deformation in orbifold CFT and its holographic dual, in contrast to the double-trace version discussed in previous sections. Single-trace deformation is described in three aspects: worldsheet theory, bulk geometry, and boundary field theory. Furthermore, the deformed partition function, correlation functions, and non-locality implications from these quantities are reviewed. Finally, Section 6 reviews recent developments in generalized $T\overline{T}$ -like deformations and provides insights into potential future research directions.

2 Basics of $T\bar{T}$ deformation

This section explains the core concepts underlying the $T\overline{T}$ -deformation, beginning with its definition and proceeding to the deformed Lagrangians. We then explore various geometric realizations of the $T\overline{T}$ deformation, including random geometry, Jackiw-Teitelboim gravity, field-dependent coordinate transformations, and the uniform light-cone gauge. These geometric realizations collectively lead to a deeper understanding of how the deformation modifies the theory, making it non-local but solvable and preserving key properties, including integrability and symmetry. By analyzing these methods and studying these core concepts, we establish a foundational framework for understanding the effects and applications of the $T\overline{T}$ -deformation in quantum field theory, holography, string theory, and other aspects of theoretical physics.

2.1 The $T\bar{T}$ deformation

The $T\bar{T}$ deformation is a solvable irrelevant deformation of two-dimensional quantum field theories. It deforms the theory by the $T\bar{T}$ operator [3] ⁵

$$O_{T\bar{T}} = -\det\left[T^{\mu}{}_{\nu}\right] = -\frac{1}{2}\epsilon_{\mu\nu}\epsilon^{\rho\sigma}T^{\mu}{}_{\rho}T^{\nu}{}_{\sigma} = \frac{1}{2}\left(T^{\mu\nu}T_{\mu\nu} - T^{\mu}{}_{\mu}T^{\nu}{}_{\nu}\right).$$
(2.1)

For a CFT in a 2D flat space with complex coordinates z, \bar{z} and a flat local metric $ds^2 = dz d\bar{z}$, the energy-momentum tensor is traceless and the $T\bar{T}$ operator is reduced to $O_{T\bar{T}} = 4T_{zz}T_{\bar{z}\bar{z}} = \pi^{-2}T\bar{T}$, justifying the nomenclature. The deformed theory depends on a parameter λ of massdimension -2. When considering an infinitesimal deformation of a theory from λ to $\lambda + \delta\lambda$, the action is deformed as

$$S^{(\lambda+\delta\lambda)} = S^{(\lambda)} + \delta\lambda \int d^2x \sqrt{\pm\gamma} O_{T\bar{T}}^{(\lambda)}, \qquad (2.2)$$

where $\gamma_{\mu\nu}$ denotes the background metric with Euclidean or Lorentzian signature. Equivalently, the flow for theory can be described by the flow equation:

$$\frac{\partial S^{(\lambda)}}{\partial \lambda} = \int d^2 x \sqrt{\pm \gamma} O_{T\bar{T}}^{(\lambda)}.$$
(2.3)

 6 In the following, we will focus on $T\bar{T}$ deformation in flat spacetime.

The $T\bar{T}$ deformation is well-defined for two-dimensional quantum field theories in flat spacetime. The $T\bar{T}$ operator in flat spacetime can be defined by point splitting:

$$-\lim_{x \to y} \frac{1}{2} \epsilon_{\mu\nu} \epsilon_{\rho\sigma} T^{\mu\rho}(x) T^{\nu\sigma}(y), \qquad (2.4)$$

⁵Our convention for the stress energy tensor in Euclidean signature is $T_{\mu\nu} = \frac{2}{\sqrt{|\gamma|}} \frac{\delta S}{\delta \gamma^{\mu\nu}}$.

⁶The $T\bar{T}$ deformation parameter λ we used is related to other conventions or notations by: $\lambda = -t$ for tin [2] or $\lambda = -\mu$ for μ in [21], $\lambda = \frac{\mu}{4}$ for μ in [12], $\lambda = \pi^2 \lambda_{\text{KLM}}$ where λ_{KLM} is the deformation parameter denoted also by λ in [14] and $\lambda = 16t$ for t in [76].

where we use Euclidean Cartesian coordinates and the Levi-Civita symbol is defined as $\epsilon^{12} = \epsilon_{12} = 1$. Now we show that the $T\bar{T}$ operator is well-defined up to derivatives, following the derivation in [3]. Taking the derivative, we find

$$\frac{\partial}{\partial x^{\alpha}} \left(\frac{1}{2} \epsilon_{\mu\nu} \epsilon_{\rho\sigma} T^{\mu\rho}(x) T^{\nu\sigma}(y) \right)
= \frac{\partial}{\partial x^{\mu}} \left(\frac{1}{2} \epsilon_{\alpha\nu} \epsilon_{\rho\sigma} T^{\mu\rho}(x) T^{\nu\sigma}(y) \right) + \frac{\partial}{\partial x^{\nu}} \left(\frac{1}{2} \epsilon_{\mu\alpha} \epsilon_{\rho\sigma} T^{\mu\rho}(x) T^{\nu\sigma}(y) \right)$$

$$= \left(\frac{\partial}{\partial x^{\nu}} + \frac{\partial}{\partial y^{\nu}} \right) \left(\frac{1}{2} \epsilon_{\mu\alpha} \epsilon_{\rho\sigma} T^{\mu\rho}(x) T^{\nu\sigma}(y) \right),$$
(2.5)

where we have used the conservation law $\partial_{\mu}T^{\mu\nu} = 0$. The general form of the OPE of the operator in the bracket in the third line can be written as:

$$\frac{1}{2}\epsilon_{\mu\alpha}\epsilon_{\rho\sigma}T^{\mu\rho}(x)T^{\nu\sigma}(y) = \sum_{i}c^{i\nu}_{\alpha}(x-y)\mathcal{O}_{i}(y).$$
(2.6)

Therefore the OPE of (2.5) takes the form

$$\frac{\partial}{\partial x^{\alpha}} \left(\frac{1}{2} \epsilon_{\mu\nu} \epsilon_{\rho\sigma} T^{\mu\rho}(x) T^{\nu\sigma}(y) \right) = \sum_{i} c^{i\nu}_{\alpha} (x-y) \frac{\partial}{\partial y^{\nu}} \mathcal{O}_{i}(y).$$
(2.7)

Consider the OPE of the point-splitting operator:

$$\frac{1}{2}\epsilon_{\mu\nu}\epsilon_{\rho\sigma}T^{\mu\rho}(x)T^{\nu\sigma}(y) = \sum_{i}C^{i}(x-y)\mathcal{O}_{i}(y).$$
(2.8)

Taking the derivative, we find

$$\frac{\partial}{\partial x^{\alpha}} \left(\frac{1}{2} \epsilon_{\mu\nu} \epsilon_{\rho\sigma} T^{\mu\rho}(x) T^{\nu\sigma}(y) \right) = \sum_{i} \frac{\partial}{\partial x^{\alpha}} C^{i}(x-y) \mathcal{O}_{i}(y).$$
(2.9)

Comparing (2.7) with (2.9), one finds that any operator \mathcal{O}_i appearing in the OPE should be a derivative of some other operators or come with a constant coefficient C^i . Therefore, the $T\bar{T}$ operator can be defined up to total derivatives as

$$-\frac{1}{2}\epsilon_{\mu\nu}\epsilon_{\rho\sigma}T^{\mu\rho}(x)T^{\nu\sigma}(y) = O_{T\bar{T}}(y) + \text{derivatives}, \qquad (2.10)$$

and the $T\bar{T}$ deformation is well-defined because the Lagrangian is defined up to total derivatives. Computing the vacuum expectation value of (2.10), we find

$$\langle O_{T\bar{T}}(y)\rangle = \langle T_{12}(x)T_{12}(y)\rangle - \frac{1}{2}\langle T_{11}(x)T_{22}(y)\rangle - \frac{1}{2}\langle T_{22}(x)T_{11}(y)\rangle.$$
 (2.11)

The vacuum expectation values of the total derivatives vanish because the theory is invariant under translation. Taking x and y infinitely separated from each other, we obtain the factorization formula

$$\langle O_{T\bar{T}}(y)\rangle = \langle T_{12}(x)\rangle\langle T_{12}(y)\rangle - \frac{1}{2}\langle T_{11}(x)\rangle\langle T_{22}(y)\rangle - \frac{1}{2}\langle T_{22}(x)\rangle\langle T_{11}(y)\rangle.$$
(2.12)

The factorization formula is also valid for non-degenerate eigenstates of the momentum and energy operators [3]. In the rest of this section, we will focus on the classical aspects of the $T\bar{T}$ deformation.

TT deformation is also considered for curved spacetimes where the existence of the curvature and lack of symmetry can complicate the definition of quantum $T\bar{T}$ operator as what we did for the flat plane. In particular, the expectation value of the $T\bar{T}$ operator was studied in [19], showing the factorization fails without assuming a large c limit. Partition function was studied in [77] by computing path integrals of the 3D gravity theory corresponding to the $T\bar{T}$ deformation. Many geometric realizations of the $T\bar{T}$ deformation to be discussed in this section can be generalized to curved space more or less straightforwardly.

2.2 Deformed Lagrangians

In principle (2.3) determines the flow of the action under the $T\bar{T}$ deformation, but in general it's extremely difficult to solve the flow with the stress tensor defined as the functional derivative of the action with respect to the metric. When a Lagrangian describes the theory, the flow equation is reduced to

$$\frac{\partial \mathcal{L}}{\partial \lambda} = \frac{1}{2} \left(T^{\mu\nu} T_{\mu\nu} - T^{\mu}{}_{\mu} T^{\nu}{}_{\nu} \right), \qquad (2.13)$$

When the deformed Lagrangian depends algebraically on the metric, we have $T_{\mu\nu} = 2 \frac{\partial \mathcal{L}}{\partial \gamma^{\mu\nu}} - \gamma_{\mu\nu} \mathcal{L}$ and the flow equation of the Lagrangian takes the form

$$\frac{\partial \mathcal{L}}{\partial \lambda} = 2\gamma^{\mu\rho}\gamma^{\nu\sigma}\frac{\partial \mathcal{L}}{\partial \gamma^{\mu\nu}}\frac{\partial \mathcal{L}}{\partial \gamma^{\rho\sigma}} - 2(\gamma^{\mu\nu}\frac{\partial \mathcal{L}}{\partial \gamma^{\mu\nu}})^2 + 2\mathcal{L}\gamma^{\mu\nu}\frac{\partial \mathcal{L}}{\partial \gamma^{\mu\nu}} - \mathcal{L}^2.$$
(2.14)

If we further assume the Lagrangian depends on fields through scalar invariants of the fields and their derivatives, then the flow equation can be reduced to a partial differential equation, which in many cases can be solved by the "method of characteristics" with the initial condition given by the undeformed theory $\mathcal{L}(\lambda = 0)$. We will devote section 2.3 to the method of characteristics. Now, for simplicity, we focus on some simple examples in flat spacetimes in the Euclidean signature.

One method to solve the flow equation (2.13) is to consider the small λ expansion of the Lagrangian:

$$\mathcal{L}(\lambda) = \sum_{k=0}^{\infty} \lambda^k \mathcal{L}_k.$$
(2.15)

The energy-momentum tensors can also be expanded as

$$T^{\mu\nu}(\lambda) = \sum_{k=0}^{\infty} \lambda^k T_k^{\mu\nu}.$$
(2.16)

When $\mathcal{L}(\lambda)$ up to order λ^n is known, one can get $T^{\mu\nu}(\lambda)$ up to the same order. Therefore, \mathcal{L}_k can be solved recursively by substituting the expansions (2.15) and (2.16) into the flow equation. This method has been applied to compute deformed scalar Lagrangians in [2]. One can use Lorentz symmetries to simplify the flow equation for relativistic theory. This method has been used in [14,47]. Now, we consider the deformed theory of a free scalar ϕ as an example. Lorentz symmetries require that the deformed Lagrangian depend on ϕ only through the Lorentz invariant variable $X = \eta^{\mu\nu} \partial_{\mu} \phi \partial_{\nu} \phi$:

$$\mathcal{L} = \mathcal{L}(\lambda, X). \tag{2.17}$$

In this simple case, it is equivalent to using the canonical energy-momentum tensor

$$T^{\mu\nu} = \frac{\partial \mathcal{L}}{\partial(\partial_{\mu}\phi)} \partial^{\nu}\phi - \eta^{\mu\nu}\mathcal{L} = 2\partial_{X}\mathcal{L}\partial^{\mu}\phi\partial^{\nu}\phi - \eta^{\mu\nu}\mathcal{L}.$$
 (2.18)

Substituting (2.17) and (2.18) into the flow equation (2.13) we get

$$\partial_{\lambda} \mathcal{L} = X \partial_X \mathcal{L}^2 - \mathcal{L}^2, \qquad (2.19)$$

which is simply the Burgers' equation. The initial condition is given by

$$\mathcal{L}(0,X) = \frac{1}{2}X.$$
 (2.20)

Solving the Burgers' equation with the initial condition, we obtained the deformed Lagrangian:

$$\mathcal{L}(\lambda, X) = \frac{1 - \sqrt{1 - 2\lambda X}}{2\lambda}.$$
(2.21)

The deformed Lagrangian of a free massless scalar can be interpreted as a Nambu-Goto action in \mathbb{R}^3 in the static gauge, which reflects the non-locality of the $T\bar{T}$ deformation.

The closed form of many $T\bar{T}$ deformed theories, including interacting scalar fields, nonlinear σ models, WZW models, and massive Thirring model, can be found in [47]. Especially in fermion theories, the expansions of deformed Lagrangians in the deformation parameter usually terminate. $T\bar{T}$ deformed fermions are further explored in [78,79] and supersymmetric theories in [80–89]. When there are multiple Lorentz invariants in the theories, the flow equations for the Lagrangians are more complicated, and one often resorts to the method of characteristics, which has also been applied to study stress energy deformations beyond $T\bar{T}$ [90].

There are other methods to derive the deformed action based on geometric realizations of the $T\bar{T}$ deformation. One is the field-dependent coordinate transformation [4, 84, 91, 92], from which solutions [91, 93] and integrals of motion [94] can also be derived. Another powerful method is the uniform light-cone gauge approach proposed in [53, 54].

If the seed theory has a traceless energy-momentum tensor, following the flow equation of the Lagrangian (2.14), we have the trace relation

$$T^{\mu}_{\mu} = 2\lambda O_{T\bar{T}},\tag{2.22}$$

or in terms of the Lagrangian

$$\lambda \partial_{\lambda} \mathcal{L} - \gamma^{\mu\nu} \frac{\partial \mathcal{L}}{\partial \gamma^{\mu\nu}} + \mathcal{L} = 0.$$
 (2.23)

To prove the trace relation, we differentiate the left-hand side with respect to λ , using (2.14) we obtain

$$\partial_{\lambda} (\lambda \partial_{\lambda} \mathcal{L} - \gamma^{\alpha \beta} \frac{\partial \mathcal{L}}{\partial \gamma^{\alpha \beta}} + \mathcal{L}) = 4 \gamma^{\mu \rho} \gamma^{\nu \sigma} \frac{\partial}{\partial \gamma^{\mu \nu}} (\lambda \partial_{\lambda} \mathcal{L} - \gamma^{\alpha \beta} \frac{\partial \mathcal{L}}{\partial \gamma^{\alpha \beta}} + \mathcal{L}) \frac{\partial \mathcal{L}}{\partial \gamma^{\rho \sigma}} - 4 \gamma^{\mu \nu} \frac{\partial}{\partial \gamma^{\mu \nu}} (\lambda \partial_{\lambda} \mathcal{L} - \gamma^{\alpha \beta} \frac{\partial \mathcal{L}}{\partial \gamma^{\alpha \beta}} + \mathcal{L}) \gamma^{\rho \sigma} \frac{\partial \mathcal{L}}{\partial \gamma^{\rho \sigma}} + 2 (\lambda \partial_{\lambda} \mathcal{L} - \gamma^{\alpha \beta} \frac{\partial \mathcal{L}}{\partial \gamma^{\alpha \beta}} + \mathcal{L}) \gamma^{\mu \nu} \frac{\partial \mathcal{L}}{\partial \gamma^{\mu \nu}} + 2 \mathcal{L} \gamma^{\mu \nu} \frac{\partial}{\partial \gamma^{\mu \nu}} (\lambda \partial_{\lambda} \mathcal{L} - \gamma^{\alpha \beta} \frac{\partial \mathcal{L}}{\partial \gamma^{\alpha \beta}} + \mathcal{L}) - 2 \mathcal{L} (\lambda \partial_{\lambda} \mathcal{L} - \gamma^{\alpha \beta} \frac{\partial \mathcal{L}}{\partial \gamma^{\alpha \beta}} + \mathcal{L}).$$
(2.24)

(2.23) holds at $\lambda = 0$ by the assumption that the seed theory has a traceless energy-momentum tensor, then by the first-order PDE above, the relation holds along the $T\bar{T}$ flow. Our proof assumed the Lagrangian depends on the metric algebraically, a more general proof based on the gravity formulation of $T\bar{T}$ deformation can be found in [6].

2.3 The method of characteristics

This section is devoted to the method of characteristics.⁷ Although a bit of a digression, it's important for the derivation of deformed classical actions, deformed energy levels and the large $c T\bar{T}$ flow equations. The method of characteristics is most clearly motivated by the quasilinear partial differential equations

$$\vec{a}(\vec{x},u) \cdot \partial_{\vec{x}} u = c(\vec{x},u). \tag{2.25}$$

Suppose the integral surface is given by a graph

$$f(\vec{x}, u) = z(\vec{x}) - u = 0.$$
(2.26)

We have $df(\vec{a} \cdot \partial_{\vec{x}} + c\partial_u) = 0$ on the integral surface. So the vector field $\vec{a} \cdot \vec{\nabla} + c\partial_u$ is tangent to any integral surface, and integral surfaces are unions of integral curves along the vector field, a.k.a. characteristic curves. Explicitly, characteristic curves are obtained by solving the ODE

$$\frac{d\vec{x}}{ds} = \vec{a}(\vec{x}, u),$$

$$\frac{du}{ds} = c(\vec{x}, u).$$
(2.27)

Initial conditions specified as a codimension two surface Γ grows into an integral surface by flowing along characteristic curves. If the PDE is nonlinear in the partial derivatives, say, given by a generic function

$$F(\vec{x}, u, \partial_{\vec{x}}u) = 0, \qquad (2.28)$$

 $^{^7 \}mathrm{One}$ can refer to chapter 3 of [95].

we no longer have the simple geometrical picture described above. However, we can naturally promote $\partial_{\vec{x}} u$ to independent variables denoted by \vec{p} with the constraints $\partial_{\vec{x}} u = \vec{p}$ to hold on an integral surface, now a codimension n + 1 hypersurface. We want to find a vector field tangent to any integral surface. For a vector field to be tangent to an integral surface, we must have

$$\frac{\partial F}{\partial \vec{x}} \cdot \frac{d\vec{x}}{ds} + \frac{\partial F}{\partial u} \frac{du}{ds} + \frac{\partial F}{\partial \vec{p}} \cdot \frac{d\vec{p}}{ds} = 0,$$

$$\frac{du}{ds} - \vec{p} \cdot \frac{d\vec{x}}{ds} = 0,$$

$$\frac{\partial \vec{p}}{\partial \vec{x}} \frac{d\vec{x}}{ds} - \frac{d\vec{p}}{ds} = 0$$
(2.29)

with the first term of the last equation depending on the particular integral surface. To get rid of this term, we differentiate (2.28) with respect to \vec{x} for an integral surface

$$\frac{\partial F}{\partial \vec{x}} + \frac{\partial F}{\partial u}\vec{p} + \frac{\partial F}{\partial \vec{p}}\frac{\partial \vec{p}}{\partial \vec{x}} = 0.$$
(2.30)

We find the dependence on the particular integral surface can be eliminated if we set $\frac{d\vec{x}}{ds} = \frac{\partial F}{\partial \vec{p}}$. Then, we obtain the equation for characteristic curves of nonlinear first-order PDEs

$$\frac{d\vec{x}}{ds} = \frac{\partial F}{\partial \vec{p}},$$

$$\frac{du}{ds} = \vec{p} \cdot \frac{\partial F}{\partial \vec{p}},$$

$$\frac{d\vec{p}}{ds} = -\frac{\partial F}{\partial \vec{x}} - \frac{\partial F}{\partial u}\vec{p}.$$
(2.31)

In the context of $T\bar{T}$ deformation of classical Lagrangian, the flow equation (2.13) is a PDE in which $\frac{\partial \mathcal{L}}{\partial \lambda}$ appears on the left-hand side and nowhere else. So we have $\frac{d\lambda}{ds} = 1$ in the characteristic equation. Therefore, the $T\bar{T}$ deformation parameter λ can be taken as the parameter of the characteristic curves. Initial conditions are given by $\lambda = 0$ hypersurface, determined by the Lagrangian of the seed theory. The $T\bar{T}$ flow is the flow along characteristic curves [90].

Taking the interacting scalar for example,

$$\mathcal{L}^{(0)} = \frac{1}{2} g^{\mu\nu} \partial_{\mu} \phi \partial_{\nu} \phi + V(\phi).$$
(2.32)

We assume the deformed Lagrangian depends algebraically on ϕ , the scalar invariant $X = g^{\mu\nu}\partial_{\mu}\phi\partial_{\nu}\phi$ and, of course, the deformation parameter λ . Then the $T\bar{T}$ flow equation for the Lagrangian (2.14) is reduced to

$$\frac{\partial \mathcal{L}}{\partial \lambda} = 2\mathcal{L}X\frac{\partial \mathcal{L}}{\partial X} - \mathcal{L}^2$$
(2.33)

with the initial condition being

$$\mathcal{L}(\lambda = 0) = \mathcal{L}^{(0)} = \frac{1}{2}X + V(\phi).$$
(2.34)

Using the method of characteristics, we define

$$p_{\lambda} = \frac{\partial \mathcal{L}}{\partial \lambda}, \ p_X = \frac{\partial \mathcal{L}}{\partial X}, \ p_{\phi} = \frac{\partial \mathcal{L}}{\partial \phi}$$
 (2.35)

and rewrite (2.33) as

$$F(\lambda, X, \phi, \mathcal{L}, p_{\lambda}, p_{X}, p_{\phi}) = p_{\lambda} - 2\mathcal{L}Xp_{X} + \mathcal{L}^{2} = 0.$$
(2.36)

The equation for characteristic curves now takes the form

$$\frac{d\lambda}{ds} = 1, \ \frac{dX}{ds} = -2\mathcal{L}X, \ \frac{d\phi}{ds} = 0$$
$$\frac{d\mathcal{L}}{ds} = p_{\lambda} - 2\mathcal{L}Xp_X = -\mathcal{L}^2$$
$$\frac{dp_{\lambda}}{ds} = 2(Xp_X - \mathcal{L})p_{\lambda}, \ \frac{dp_X}{ds} = 2Xp_X, \ \frac{dp_{\phi}}{ds} = 2(Xp_X - \mathcal{L})p_{\phi}$$
(2.37)

From the first equation, we set $s = \lambda$. Integrating the fourth equation, we find

$$\mathcal{L} = \frac{\mathcal{L}_0}{1 + \lambda \mathcal{L}_0}.$$
(2.38)

Plugging into the second equation, we obtain

$$X = \frac{X_0}{(1 + \lambda \mathcal{L}_0)^2}.$$
 (2.39)

And by the third equation, we have

$$\phi = \phi_0. \tag{2.40}$$

For the seed theory we have $\mathcal{L}_0 = \frac{1}{2}X_0 + V(\phi_0)$. Substituting X_0, ϕ_0 by X, ϕ , we finally find the deformed Lagrangian

$$\mathcal{L}(\lambda, X, \phi) = \frac{1 + 2\lambda V(\phi) - \sqrt{1 - 2\lambda X(1 + \lambda V(\phi))}}{2\lambda (1 + \lambda V(\phi))}.$$
(2.41)

This was obtained as a series in [96] and later in a closed form in [47], which simplifies to our expression.

2.4 Geometric realization of $T\bar{T}$ deformation

Random geometry

Remarkable geometric interpretations exist for the $T\bar{T}$ deformation. It has been shown in [97] that $T\bar{T}$ deformation can be interpreted as coupling the original theory to a random geometry. The infinitesimal deformation of the action is:

$$S^{(\lambda+\delta\lambda)} = S^{(\lambda)} - \frac{1}{2}\delta\lambda \int d^2x \epsilon_{\mu\nu}\epsilon_{\rho\sigma}T^{\mu\rho}T^{\nu\sigma}.$$
 (2.42)

In the path integral formulation, one can use the Hubbard-Stratonovich transformation to rewrite the perturbation term as:

$$e^{\frac{\delta\lambda}{2}\int d^2x\epsilon_{\mu\nu}\epsilon_{\rho\sigma}T^{\mu\rho}T^{\nu\sigma}} = \int \mathcal{D}h \, e^{-\frac{1}{2\delta\lambda}\int\epsilon^{\mu\nu}\epsilon^{\rho\sigma}h_{\mu\rho}h_{\nu\sigma}d^2x - \int h_{\mu\nu}T^{\mu\nu}d^2x}.$$
(2.43)

The saddle point of $h_{\mu\nu}$ is of order $O(\delta\lambda)$:

$$h_{\mu\nu}^* = -\delta\lambda\epsilon_{\mu\rho}\epsilon_{\nu\sigma}T^{\rho\sigma} = \delta\lambda(T_{\mu\nu} - \delta_{\mu\nu}T_{\rho}^{\rho}).$$
(2.44)

According to the definition of the energy-momentum tensor, the last term can be viewed as an infinitesimal change in the metric $\gamma_{\mu\nu} = \delta_{\mu\nu} + h_{\mu\nu}$. The path integral of $h_{\mu\nu}$ appears to encompass all possible metrics infinitesimally close to $\delta_{\mu\nu}$. But we will show that one can restrict to flat metrics. The infinitesimal variation of the Ricci scalar can be computed as:

$$\delta R = \partial_{\mu} \partial_{\nu} h^{\mu\nu} - \partial^2 h^{\mu}_{\mu}. \tag{2.45}$$

At the saddle point (2.44), we find

$$\delta R = \partial_{\mu} \partial_{\nu} h^{*\mu\nu} - \partial^2 h^{*\mu}_{\mu} = 0, \qquad (2.46)$$

where we use the conservation of energy-momentum tensor $\partial_{\mu}T^{\mu\nu}$. Therefore, the metric is flat at the saddle point, and the variation can be restricted to be an infinitesimal diffeomorphism:

$$h_{\mu\nu}^* = \partial_\mu \alpha_\nu + \partial_\nu \alpha_\mu. \tag{2.47}$$

Flow equations for partition functions on a torus, a finite cylinder, and connected domains with boundaries can be derived using the random geometry interpretation.

Jackiw-Teitelboim-like Gravity

We begin by considering the 2-form curvature R_{ab} , expressed in terms of the Riemann curvature tensor R_{abcd} and the vierbein e^a . The relationship is given as:

$$R_{ab} = \frac{1}{2} R_{abcd} e^c \wedge e^d = \frac{1}{2} R_{abcd} g_{ac} g^{ac} g_{bd} g^{bd} e^c \wedge e^d = \frac{1}{2} R e_a \wedge e_b = \frac{1}{2} R e_{a\mu} e_{b\nu} dx^{\mu} \wedge dx^{\nu}.$$
 (2.48)

This formulation shows the curvature expressed in terms of the metric and the vierbein components. Now, comparing the two expressions for R_{ab} , we can derive the following relation:

$$Re_{a\mu}e_{b\nu} = \epsilon_{ab}(\partial_{\mu}\omega_{\nu} - \partial_{\nu}\omega_{\mu}), \qquad (2.49)$$

where ω_{μ} is the spin connection. Further, we obtain the relation in determinant form:

$$R\frac{1}{2}\epsilon^{\mu\nu}\epsilon^{ab}e_{a\mu}e_{b\nu} = \frac{1}{2}\epsilon^{\mu\nu}\epsilon^{ab}\epsilon_{ab}(\partial_{\mu}\omega_{\nu} - \partial_{\nu}\omega_{\mu}), \qquad (2.50)$$

which simplifies to:

$$R \det(e) = \epsilon^{\mu\nu} (\partial_{\mu}\omega_{\nu} - \partial_{\nu}\omega_{\mu}).$$
(2.51)

Finally, we find the expression of eR in terms of the spin connection ω_{μ}

$$eR = \epsilon^{\mu\nu} (\partial_{\mu}\omega_{\nu} - \partial_{\nu}\omega_{\mu}) = \epsilon^{\mu\nu} \partial_{\mu}\omega_{\nu} - \epsilon^{\nu\mu} \partial_{\mu}\omega_{\nu} = 2\epsilon^{\mu\nu} \partial_{\mu}\omega_{\nu}.$$
(2.52)

Next, we rewrite the Jackiw-Teitelboim-like gravity action, which is deformed by the TT operator, in the first-order formalism. The general action is

$$S_{GJT} = \int d^2x \ (eR\phi + eW(\phi))$$

= $\int d^2x \left(2\epsilon^{\mu\nu}\partial_{\mu}\omega_{\nu}\phi + \frac{1}{2}\epsilon^{\mu\nu}\epsilon_{ab}e^a{}_{\mu}e^b{}_{\nu}W(\phi) \right)$
= $\int d^2x \ \epsilon^{\mu\nu} \left(2\phi\partial_{\mu}\omega_{\nu} + \frac{W(\phi)}{2}\epsilon_{ab}e^a{}_{\mu}e^b{}_{\nu} - \sigma_a \left(\partial_{\mu}e^a{}_{\nu} + \omega_{\mu}\epsilon^a{}_{b}e^b{}_{\nu} \right) \right),$ (2.53)

where σ_a is a Lagrange multiplier that enforces the metricity constraint for the spin connection. For more details on the metricity constraint, refer to equation (16) in [5].

If we introduce the Newton constant G_N , the generalized Jackiw-Teitelboim (JT) action becomes:

$$S_{GJT} = -\frac{1}{16\pi G_N} \int d^2 x \,\epsilon^{\mu\nu} \left(2\phi \partial_\mu \omega_\nu + \frac{W(\phi)}{2} \epsilon_{ab} e^a{}_\mu e^b{}_\nu - \sigma_a \left(\partial_\mu e^a{}_\nu + \omega_\mu \epsilon^a{}_b e^b{}_\nu \right) \right). \tag{2.54}$$

We now consider the equations of motion derived from the generalized Jackiw-Teitelboimlike gravity action to demonstrate how this theory is related to $T\bar{T}$ -deformed conformal field theories. The $T\bar{T}$ deformation allows us to explore the effects on quantum field theories in two dimensions (2D), which have significant connections to holographic dualities and lowerdimensional gravitational solutions.

The equation of motion for the dilaton field ϕ is given by:

$$\epsilon^{\mu\nu} \left(2\partial_{\mu}\omega_{\nu} + \frac{\partial_{\phi}W(\phi)}{2} \epsilon_{ab} e^{a}{}_{\mu}e^{b}{}_{\nu} \right) = 0, \qquad (2.55)$$

where ω_{μ} is the spin connection, and $e^{a}{}_{\mu}$ are the components of the vierbein. Using the relations $eR = 2\epsilon^{\mu\nu}\partial_{\mu}\omega_{\nu}$ and $e = \frac{1}{2}\epsilon^{\mu\nu}\epsilon^{ab}e_{a\mu}e_{b\nu}$, we find that the scalar curvature satisfies

$$e\left(R[\omega] + \partial_{\phi}W(\phi)\right) = 0. \tag{2.56}$$

Assuming the potential function $W(\phi)$ takes the form

$$W(\phi) = -\phi\Lambda, \tag{2.57}$$

where Λ is a cosmological constant, we get

$$\partial_{\phi} W(\phi) = -\Lambda, \tag{2.58}$$

and the equation becomes

$$e\left(R[\omega] - \Lambda\right) = 0. \tag{2.59}$$

Substituting the equation of motion for ϕ into the action, the generalized Jackiw-Teitelboim action becomes

$$S_{GJT} = -\frac{1}{16\pi G_N} \int d^2 x \,\epsilon^{\mu\nu} \left(2\phi \partial_\mu \omega_\nu + \frac{W(\phi)}{2} \epsilon_{ab} e^a{}_\mu e^b{}_\nu - \sigma_a \left(\partial_\mu e^a{}_\nu + \omega_\mu \epsilon^a{}_b e^b{}_\nu \right) \right),$$

$$= -\frac{1}{16\pi G_N} \int d^2 x \left[-\phi \epsilon^{\mu\nu} \frac{\partial_\phi W(\phi)}{2} \epsilon_{ab} e^a{}_\mu e^b{}_\nu + \epsilon^{\mu\nu} \frac{W(\phi)}{2} \epsilon_{ab} e^a{}_\mu e^b{}_\nu - \epsilon^{\mu\nu} \left(\partial_\mu (\sigma_a e^a{}_\nu) - e^a{}_\nu \partial_\mu \sigma_a + \sigma_a \omega_\mu \epsilon^a{}_b e^b{}_\nu \right) \right],$$

$$= -\frac{1}{16\pi G_N} \int d^2 x \,\epsilon^{\mu\nu} \left[\frac{1}{2} \left(W(\phi) - \phi \partial_\phi W \right) \epsilon_{ab} e^a{}_\mu e^b{}_\nu + e^a{}_\nu \partial_\mu \sigma_a - \sigma_b \omega_\mu \epsilon^b{}_a e^a{}_\nu \right].$$
(2.60)

Thus, the $T\bar{T}$ -deformed action becomes

$$S_{T\bar{T}} = -\frac{1}{16\pi G_N} \int d^2 x \,\epsilon^{\mu\nu} \left(\frac{1}{2} \left(W - \phi \partial_\phi W \right) \epsilon_{ab} e^a{}_\mu e^b{}_\nu + \left(\partial_\mu \sigma_a + \omega_\mu \epsilon_a{}^b \sigma_b \right) e^a{}_\nu \right) + S_m \left(e^a_\mu, \Psi \right),$$
(2.61)

where $S_m(e^a_\mu,\Psi)$ represents the matter contribution.

Let us now introduce the notation $D_{\mu}\sigma_a = \partial_{\mu}\sigma_a + \omega_{\mu}\epsilon_a{}^b\sigma_b$ and $f = W - \phi\partial_{\phi}W$. The action then simplifies to

$$S_{T\bar{T}} = -\frac{1}{16\pi G_N} \int d^2 x \, \epsilon^{\mu\nu} \left(\frac{f}{2} \epsilon_{ab} e^a{}_{\mu} e^b{}_{\nu} + D_{\mu} \sigma_a e^a{}_{\nu} \right) + S_m \left(e^a_{\mu}, \Psi \right). \tag{2.62}$$

The action can also be written in a factorized form as

$$S_{T\bar{T}} = -\frac{1}{16\pi G_N} \int d^2x \, \frac{f}{2} \epsilon^{\mu\nu} \epsilon_{ab} \left(e^a{}_\mu - \frac{1}{f} \epsilon^{ac} D_\mu \sigma_c \right) \left(e^b{}_\nu - \frac{1}{f} \epsilon^{bd} D_\nu \sigma_d \right) + \frac{1}{16\pi G_N} \int d^2x \, \frac{1}{2f} \epsilon^{\mu\nu} \epsilon_{ab} \left(\epsilon^{ac} D_\mu \sigma_c \right) \left(\epsilon^{bd} D_\nu \sigma_d \right) + S_m.$$
(2.63)

In this sense, the $T\bar{T}$ -deformed action is classically equivalent to the undeformed theory coupled to two-dimensional generalized Jackiw-Teitelboim gravity.

Field-dependent coordinate transformation

The deformed theory is related to the original theory by a field-dependent coordinate transformation [4,91,92]. The deformed theory can be reformulated as

$$S_{\lambda} = \int d^2 w \mathcal{L}_{\lambda} = \int \frac{d^2 w}{\det \partial_{x^i} w^j} (\mathcal{L}_0(\phi(x(w))) + \lambda \det((T^{(\lambda=0)})^{\mu}_{\nu})), \qquad (2.64)$$

where x and w are coordinates for the original and deformed theory, respectively. They are related by a field-dependent coordinate transformation

$$dw^{\mu} = dx^{\mu} - \lambda \epsilon^{\mu\beta} \epsilon_{\alpha\nu} T^{\alpha}_{\ \beta} dx^{\nu}. \tag{2.65}$$

Now, we show how to use this approach to derive the deformed action using the example of a scalar with an arbitrary potential. The undeformed action is given by

$$S_0 = \int d^2 x \mathcal{L}_0 = \int dx^+ dx^- \left(-\partial_{x^+} \phi \partial_{x^-} \phi - V(\phi) \right).$$
(2.66)

where the light-cone coordinates are defined as $x^{\pm} = (x \pm t)/\sqrt{2}$. The canonical energymomentum tensor is

$$T^{\mu}_{\ \nu} = \left(\begin{array}{cc} V & -\partial_{x^{-}}\phi\partial_{x^{-}}\phi \\ -\partial_{x^{+}}\phi\partial_{x^{+}}\phi & V \end{array}\right).$$
(2.67)

We defined a field-dependent coordinate transformation with the Jacobian

$$\begin{pmatrix} \partial_{x^+}w^+ & \partial_{x^+}w^- \\ \partial_{x^-}w^+ & \partial_{x^-}w^- \end{pmatrix} = \begin{pmatrix} 1 - \lambda T^{x^-}_{x^-} & \lambda T^{x^-}_{x^+} \\ \lambda T^{x^+}_{x^-} & 1 - \lambda T^{x^+}_{x^+} \end{pmatrix}.$$
 (2.68)

Solving

$$\begin{pmatrix} \partial_{x^{+}}\phi\\ \partial_{x^{-}}\phi \end{pmatrix} = \begin{pmatrix} \partial_{x^{+}}w^{+} & \partial_{x^{+}}w^{-}\\ \partial_{x^{-}}w^{+} & \partial_{x^{-}}w^{-} \end{pmatrix} \begin{pmatrix} \partial_{w^{+}}\phi\\ \partial_{w^{-}}\phi \end{pmatrix},$$
(2.69)

we get

$$\partial_{x^+}\phi = \frac{\Omega - 1}{2\lambda\partial_{w^-}\phi}, \quad \partial_{x^-}\phi = \frac{\Omega - 1}{2\lambda\partial_{w^+}\phi}.$$
(2.70)

where

$$\Omega = \sqrt{1 + 4\lambda \partial_{w^+} \phi \partial_{w^-} \phi (1 - \lambda V)}.$$
(2.71)

The deformed action is given by

$$S_{\lambda} = \int d^2 w \mathcal{L}_{\lambda} = \int \frac{d^2 w}{\det \partial_{x^i} w^j} (\mathcal{L}_0(\phi(x(w))) + \lambda \det(T))$$

= $\int d^2 w \left(\frac{1 - \Omega - 2\lambda V}{2\lambda(1 - \lambda V)} \right),$ (2.72)

which is consistent with the result in [47].

2.5 Uniform light-cone gauge approach

This subsection reviews the relationship between string theories in the uniform light-cone gauge and $T\bar{T}$ deformation [53, 54]. We consider a string action

$$S = -\frac{1}{2} \int_{-r}^{r} d\sigma d\tau \,\gamma^{\alpha\beta} \partial_{\alpha} X^{M} \partial_{\beta} X^{N} G_{MN}, \qquad (2.73)$$

where the world-sheet coordinate σ ranges from -r to < r and the target space metric is given by ⁸

$$ds^{2} = G_{MN} dX^{M} dX^{N} = G_{tt} dt^{2} + 2G_{t\phi} dt d\phi + G_{\phi\phi} d\phi^{2} + G_{\mu\nu} dx^{\mu} dx^{\nu}.$$
 (2.74)

We assume that G_{MN} is independent of t and ϕ , and therefore, the shifts of t and ϕ are two isometries. We use μ , $\nu = 1, ..., n$ to denote indices of the transversal coordinates. It is convenient to use the action in the first-order form

$$S = \int_{-r}^{r} d\sigma d\tau \left(p_M \dot{X}^M + \frac{\gamma^{01}}{\gamma^{00}} C_1 + \frac{1}{2} \gamma^{00} C_2 \right).$$
(2.75)

⁸In this subsection, we use (t, x^{μ}, ϕ) and (σ, τ) to denote target space coordinates and world-sheet coordinates respectively.

where p_M are canonical momenta and the constraints $C_{1,2}$ are defined as

$$C_1 = p_M X'^M, \quad C_2 = G^{MN} p_M p_N + X'^M X'^N G_{MN}.$$
 (2.76)

We will use the conformal gauge $\gamma^{\alpha\beta} = \text{diag}(-1,1)$. The conserved charge corresponds to the invariance under the shift of t is the target space energy. If ϕ is an angular variable, the associated shift invariant conserved charge can be viewed as a target space angular momentum. The two conserved charges can be computed as

$$E = -\int_{-r}^{r} d\sigma p_t, \quad J = \int_{-r}^{r} d\sigma p_{\phi}.$$
 (2.77)

We define the light-cone target space coordinates and momenta as

$$x^{-} = \phi - t, \quad x^{+} = \frac{1}{2}(\phi + t) + \alpha x^{-}, \quad p_{+} = p_{\phi} + p_{t}, \quad p_{-} = \frac{1}{2}(p_{\phi} - p_{t}) - \alpha p_{+}, \quad (2.78)$$

where we introduce an arbitrary parameter α which will turn out to relate to the $T\bar{T}$ deformation λ . In light cone coordinates, the action and constraints can be written as

$$S = \int_{-r}^{r} d\sigma d\tau \left(p_{+} \dot{x}^{+} + p_{-} \dot{x}^{-} + p_{\mu} \dot{x}^{\mu} + \frac{\gamma^{01}}{\gamma^{00}} C_{1} + \frac{1}{2\gamma^{00}} C_{2} \right), \qquad (2.79)$$

$$C_1 = p_+ x'^+ + p_- x'^- + p_\mu x'^\mu, \tag{2.80}$$

$$C_{2} = G^{++}p_{+}^{2} + 2G^{-+}p_{-}p_{+} + G^{--}p_{-}^{2} + G_{++}(x'^{+})^{2} + 2G_{-+}x'^{-}x'^{+} + G_{--}(x'^{-})^{2} + 2\mathcal{H}_{x}, \quad (2.81)$$

$$\mathcal{H}_x = \frac{1}{2} (G^{\mu\nu} p_\mu p_\nu + G_{\mu\nu} x'^\mu x'^\nu). \tag{2.82}$$

The target space metric components in light cone coordinates are

$$\begin{aligned}
G_{++} &= G_{\phi\phi} + 2G_{t\phi} + G_{tt}, \\
G^{--} &= \frac{G_{++}}{\det G_{lc}}, \quad \det G_{lc} \equiv G_{tt}G_{t\phi} - G_{\phi\phi}^{2}, \\
G_{--} &= \left(\frac{1}{2} - \alpha\right)^{2}G_{\phi\phi} + \left(2\alpha^{2} - \frac{1}{2}\right)G_{t\phi} + \left(\frac{1}{2} + \alpha\right)^{2}G_{tt}, \quad G^{++} = \frac{G_{--}}{\det G_{lc}}, \\
G_{-+} &= \left(\frac{1}{2} - \alpha\right)G_{\phi\phi} - 2\alpha G_{t\phi} - \left(\alpha + \frac{1}{2}\right)G_{tt}, \quad G^{-+} = -\frac{G_{-+}}{\det G_{lc}}.
\end{aligned}$$
(2.83)

Now, we impose a uniform light-cone gauge, which is defined as

$$x^{+} = \tau + \pi (\frac{1}{2} + \alpha) \frac{m}{r} R_{\phi} \sigma, \quad p_{-} = 1,$$
 (2.84)

where $2\pi R_{\phi}$ is the period of ϕ and the integer m is the winding number. In the uniform light-cone gauge, $C_1 = 0$ can be solved by

$$x'^{-} = -\pi (\frac{1}{2} + \alpha) \frac{m}{r} R_{\phi} p_{+} - p_{\mu} x'^{\mu}.$$
(2.85)

This solution can be substituted into C_2 , and thus, one can solve $C_2 = 0$ for p_+ . Finally, the gauge-fixed action can be written as

$$S_{\alpha} = \int_{-r}^{r} d\sigma d\tau (p_{\mu} \dot{x}^{\mu} + p_{+}(p_{\mu}, x^{\mu}, x'^{\mu})).$$
(2.86)

The gauge-fixed action can describe $T\bar{T}$ deformed models with scalars, fermions, and chiral bosons by choosing appropriate target space metrics. For the example of n scalar fields with arbitrary potential, one can take

$$G_{tt} = -(1 + \frac{V}{2}), \quad G_{\phi\phi} = 1 - \frac{V}{2}, \quad G_{t\phi} = -\frac{V}{2}.$$
 (2.87)

We also set the winding number m = 0. Then, the gauge-fixed action becomes

$$S_{\alpha} = \int_{-r}^{r} d\sigma d\tau (p_{\mu} \dot{x}^{\mu} - \frac{1}{\alpha} + \frac{1}{2\tilde{\alpha}} + \frac{1}{2\tilde{\alpha}} \sqrt{1 - 4\tilde{\alpha}\mathcal{H}_{x} + 4\tilde{\alpha}^{2} (p_{\mu} x'^{\mu})^{2}}), \qquad (2.88)$$

with $\tilde{\alpha} = \alpha(1 + \alpha V)$. Integrating out the canonical momenta p_{μ} , we get

$$S_{\alpha} = \int_{-r}^{r} d\sigma d\tau \left(-\frac{1}{\alpha} + \frac{1}{2\tilde{\alpha}} + \frac{1}{2\tilde{\alpha}}\sqrt{1 + 2\tilde{\alpha}\left(\dot{x}^{2} - x^{\prime 2}\right) - 4\tilde{\alpha}^{2}\left(\dot{x}^{2}x^{\prime 2} - (\dot{x}x^{\prime})^{2}\right)}\right),\tag{2.89}$$

which agrees with the $T\bar{T}$ deformation of *n* scalar fields with arbitrary potential [63]. To match the convention in the previous subsections, we take $\alpha = -\lambda$ by comparing (2.89) with (2.72) for the n = 1 case.

2.6 $T\bar{T}$ -deformed conformal symmetry

The $T\bar{T}$ deformation is not only solvable but also preserves many important properties of the seed theory. This can usually be understood by the geometric realizations of $T\bar{T}$ deformation. Of particular interest to us, the conformal symmetry is preserved/deformed by $T\bar{T}$ deformation, taking a form of symmetry of field-dependent coordinate transformations as discussed in 2.4. The key observation is, while the energy-momentum tensor is no longer traceless, it still has only two independent components by the trace relation (2.22). Following [98], we work with flat space in light-cone coordinates U, V, and the trace relation takes the form

$$T_{UV} = 2\lambda (T_{UU}T_{VV} - T_{UV}^2)$$
(2.90)

It's convenient to parametrize the energy-momentum tensor by two functions $\mathcal{L}, \bar{\mathcal{L}}$

$$T_{UU} = \frac{\mathcal{L}}{2(1 - \lambda \mathcal{L}\bar{\mathcal{L}})}, \ T_{VV} = \frac{\bar{\mathcal{L}}}{2(1 - \lambda \mathcal{L}\bar{\mathcal{L}})}, \ T_{UV} = \frac{\lambda \mathcal{L}\bar{\mathcal{L}}}{2(1 - \lambda \mathcal{L}\bar{\mathcal{L}})}.$$
 (2.91)

The conservation equations of the energy-momentum tensor can be written as

$$(\partial_V + \lambda \bar{\mathcal{L}} \partial_U) \mathcal{L} = 0, \ (\partial_U + \lambda \mathcal{L} \partial_V) \bar{\mathcal{L}} = 0.$$
(2.92)

It's clear that for $\lambda = 0$, $\mathcal{L}, \overline{\mathcal{L}}$ are the holomorphic and anti-holomorphic components of the energy-momentum tensor of the seed CFT, and for a CFT the action is invariant under the coordinate transformation

$$U \to U + \epsilon f(U), \ V \to V - \epsilon \bar{f}(V).$$
 (2.93)

Now for a $T\bar{T}$ -deformed CFT, we assume the coordinate transformation is

$$U \to U + \epsilon f(u), \ V \to V - \epsilon \bar{f}(v).$$
 (2.94)

where u, v is a new set of coordinates. The variation of the action under the coordinate transformation is

$$\delta S = \epsilon \int dU dV \left[-f'(u) (T_{UU} \partial_V u + T_{VU} \partial_U u) + \bar{f}'(v) (T_V V \partial_U v + T_{UV} \partial_V v) \right].$$
(2.95)

The action is invariant if

$$\partial_V u + \lambda \bar{\mathcal{L}} \partial_U u = 0, \ \partial_U v + \lambda \mathcal{L} \partial_V v = 0.$$
(2.96)

The solution depends on \mathcal{L} and $\overline{\mathcal{L}}$; that is, the coordinate transformation that leaves the action invariant is field-dependent, in contrast to the field-independent two-dimensional conformal symmetry (2.93). The conservation equations take a much simpler form in the new coordinates

$$\partial_v \mathcal{L} = 0, \; \partial_u \bar{\mathcal{L}} = 0, \tag{2.97}$$

so the energy-momentum tensor is parametrized by two functions $\mathcal{L}(u), \bar{\mathcal{L}}(v)$, similar to the case of CFTs. The charge algebra and quantum aspects of the $T\bar{T}$ -deformed conformal symmetry were further studied in [50,98–101]. Given how the conformal symmetry determines CFTs up to conformal dimensions and OPE coefficients subject to bootstrap equations (see [102,103] for example), it's a very interesting question how much we can extract from the deformed conformal symmetry for $T\bar{T}$ -deformed CFTs. The first step is to find a set of operators that transform nicely under the deformed conformal symmetry. The definition of a primary operator is given in [104] for $J\bar{T}$ -deformed CFTs, while it remains an open question for $T\bar{T}$ deformation.

3 $T\overline{T}$ deformation in quantum field theory

The perturbative and non-perturbative studies of correlation functions in $T\overline{T}$ -deformed quantum field theories provide essential insights into the deformation's impact on fundamental properties of the theory. These analysis systematically addresses the induced non-locality and ultraviolet-infrared (UV-IR) mixing effects, clarifying the structure of observables and the renormalization procedures required to maintain the theory's consistency and predictive power. The solvability of TT-deformed theories further allows exact computations of correlation functions and energy spectra. Beyond perturbation theory, non-perturbative approaches, such as resurgence methods, have explored the partition function and spectral properties. Quantum entanglement entropy calculations in $T\overline{T}$ -deformed theories demonstrate modifications from the undeformed case, providing a novel probe of the deformation's non-local nature. Additionally, the large central charge limit simplifies the flow equations, offering a powerful tool for analyzing quantum corrections and non-perturbative effects. These studies also extend to black hole physics, where the deformation influences thermodynamic quantities, including entropy growth and microstate structure. This section systematically reviews these quantum aspects, emphasizing exact solvability, correlation functions, renormalization, entanglement properties, large central charge behavior, and recent developments, elucidating key characteristics and implications of the $T\overline{T}$ deformation.

3.1 Deformed spectrum and partition function

Following the conventions in section 2, Consider a $T\bar{T}$ deformed theory defined on a circle of circumference L. The deformed Hamiltonian and its eigenvalues in Euclidean signature are [12]

$$H_{int} = \int dx L_{int} = \lambda \int dx O_{T\bar{T}}, \qquad (3.1)$$

$$\frac{\partial E_n}{\partial \lambda} = \frac{\partial \langle n | H_{int} | n \rangle}{\partial \lambda} = \lambda \int dx \langle n | O_{T\bar{T}} | n \rangle = L \langle n | O_{T\bar{T}} | n \rangle, \tag{3.2}$$

where

$$\langle n|O_{T\bar{T}}|n\rangle = (\langle n|T^{x\tau}|n\rangle)^2 - \langle n|T^{xx}|n\rangle\langle n|T^{\tau\tau}|n\rangle.$$
(3.3)

By using the relation

$$\langle n|T^{\tau\tau}|n\rangle = \frac{E_n}{L}, \quad \langle n|T^{xx}|n\rangle = \frac{\partial E_n}{\partial L}, \quad \langle n|T^{\tau x}|n\rangle = \frac{iP_n}{L},$$
 (3.4)

one can obtain the so-called Burger's equation

$$\frac{\partial E_n}{\partial \lambda} + \frac{P_n^2}{L} + E_n \frac{\partial E_n}{\partial L} = 0, \qquad (3.5)$$

whose solution, i.e., the deformed spectrum is [1, 2, 12]

$$E_n(\lambda) = \frac{L}{2\lambda} \left(1 - \sqrt{1 - \frac{4\lambda}{L}} E_n^{(\text{CFT})} + \frac{4\lambda^2}{L^2} (P_n^{(\text{CFT})})^2 \right), \quad P_n(\lambda) = P_n^{(\text{CFT})}.$$
(3.6)

For positive λ , the highly excited states have complex energies, leading to a non-unitary deformed theory, while for negative spectrum, the spectrum exhibits Hagedorn growth. Notice that here the undeformed theory was chosen to be a CFT, with the energy and momentum being

$$E_n(0) \equiv E_n^{(\text{CFT})} = \frac{2\pi}{L} (L_0 + \bar{L}_0 - \frac{c}{12}), \quad P_n(0) \equiv P_n^{(\text{CFT})} = \frac{2\pi}{L} (L_0 - \bar{L}_0), \quad (3.7)$$

Letting $E_L = \frac{1}{2}(E+P), E_R = \frac{1}{2}(E-P)$, one have [105]

$$E_R(0) = E_R(\lambda) - \frac{2\lambda}{L} E_R(\lambda) E_L(\lambda),$$

$$E_L(0) = E_L(\lambda) - \frac{2\lambda}{L} E_R(\lambda) E_L(\lambda).$$
(3.8)

With periods of torus are $w_1 = L, w_2$, and the modular parameter $\tau = w_2/w_1$, the deformed partition function can be expressed as [97] [106]

$$Z(\tau, \bar{\tau}, \lambda) = \operatorname{tr}(e^{i\tau_1 LP - \tau_2 LH(\lambda)}) = \operatorname{tr}(e^{i\tau LE_L(\lambda) - i\bar{\tau} LE_R(\lambda)}).$$
(3.9)

The CFT partition function is

$$Z(\tau, \bar{\tau}, \lambda = 0) = \operatorname{tr}(q^{L_0 - c/24} \bar{q}^{\bar{L}_0 - c/24}).$$
(3.10)

Since $E_n(\lambda)L$ is dimensionless, it depends only on λ/L^2 , one has

$$L\frac{\partial E}{\partial L} + E = -2\lambda \frac{\partial E}{\partial \lambda}.$$
(3.11)

Combining (3.5) and (3.11), one can obtain the flow of partition function defined in (3.9)

$$L^{2}\partial_{\lambda}Z + \frac{2}{i}(\tau - \bar{\tau})\partial_{\tau}\partial_{\bar{\tau}}Z + 2i\lambda(\partial_{\tau} - \partial_{\bar{\tau}})\partial_{\lambda}Z - \frac{4i\lambda}{(\tau - \bar{\tau})}\partial_{\lambda}Z = 0.$$
(3.12)

The modular transformation property of the deformed partition function is [106] (see also [107] from holographic perspective)

$$Z\left(\frac{a\tau+b}{c\tau+d}, \frac{a\bar{\tau}+b}{c\bar{\tau}+d}, \frac{\lambda}{|c\tau+d|^2}\right) = Z(\tau, \bar{\tau}, \lambda).$$
(3.13)

Notice that $Z(\tau, \bar{\tau}, \lambda)$ does not have a closed form and instead can be expressed as a formal series in λ , with expansion coefficients computed via recursion relation. Based on this formal series, non-perturbative contribution to deformed partition function was recently investigated via resurgence method [108].

The $T\bar{T}$ -deformed partition function could also be derived from single-trace $T\bar{T}$ deformation (see section 5) by the argument of universality [34,109]. In [109] the deformed partition function reads

$$Z(\zeta,\bar{\zeta},\lambda) = \frac{\zeta_2}{2\lambda} \int_{\mathcal{H}_+} \frac{d^2\tau}{\tau_2^2} e^{-\frac{\pi}{2\lambda\tau_2}|\tau-\zeta|^2} Z_{\text{cft}}(\tau,\bar{\tau}).$$
(3.14)

Here \mathcal{H}_+ is the upper half-plane. (3.14) is consistent with the results obtained via different methods in [97] and [5]. The deformed spectrum (3.6) follows by substituting CFT partition function (3.10) into the RHS of (3.14).

The Cardy formula for CFT entropy

$$S = 2\pi \sqrt{\frac{c_R}{6} \left(L_0 - \frac{c_R}{24} \right)} + 2\pi \sqrt{\frac{c_L}{6} \left(\bar{L}_0 - \frac{c_L}{24} \right)}.$$
 (3.15)

For simplicity, consider the case $c_L = c_R, L_0 = \bar{L}_0$

$$S(E(0)) = 2\pi \sqrt{\frac{c}{3}E(0)}.$$
(3.16)

Since each state in the deformed theory is related to the state in the original CFT via (3.6), the entropy for deformed theory

$$S(E(\lambda)) = 2\pi \sqrt{\frac{c}{3}E(0)} = 2\pi \sqrt{\frac{c}{3}(E(\lambda) - \frac{4\lambda}{L}E(\lambda)^2)}.$$
(3.17)

For positive λ , the entropy exhibit Hagedorn growth $S = \beta_H E$ at high energy, where $\beta_H = 2\pi \sqrt{\frac{-4c\lambda}{3L}}$ is inverse Hagedorn temperature.

3.2 Renormalization

Renormalization of $T\bar{T}$ deformed theory is non-trivial since it is non-normalizable from standard quantum field theory. This means there are an infinite number of counterterms. However, since this kind of deformed is integrable, it leads to infinite constraints to fix the counterterms. Finally, it makes the usual perturbation method applicable in the $T\bar{T}$ deformed theories.

It is known that for a integrable 2D QFT, two-to-two S-matrix $S(\theta)$ of a single particle of mass m can be expressed by the CDD factor S_{α}

$$S(\theta) = \prod_{\alpha} S_{\alpha}(\theta), \quad S_{\alpha} = \frac{\sinh \theta - i \sin \alpha}{\sinh \theta + i \sin \alpha},$$
(3.18)

where $\theta = \theta_1 - \theta_2$ is the rapidity difference of two particles and α is a model-dependent real number.

In the $T\bar{T}$ deformed theory, the S-matrix can be obtained (by integrability) and is modified by picking up a phase factor [1]

$$S_{\lambda}(\theta) = S(\theta)S'(\theta), \quad S'(\theta) = e^{i\lambda m^{2}\sinh\theta}, \quad (3.19)$$

where $S(\theta)$ is the S-matrix of undeformed integrable system. The deformed S-matrix grows exponentially at large imaginary momentum, which indicates the deformed theory is not a local QFT. For example, consider a free massive scalar $S(\theta) = 1$, then $S_{\lambda}(\theta) = \exp(i\lambda m^2 \sinh \theta)$. One takes the limit $\theta \to \infty$ and $m \to 0$ for the massless case while keeping $s = (p_1 + p_2)^2$ fixed. Hence the $T\bar{T}$ -deformed S-matrix is

$$S_{\lambda}(\theta) = \exp(i\lambda s/2). \tag{3.20}$$

Alternatively, the S-matrix can be computed perturbatively in λ order by order, at least in some leading orders as did in [110].

To compute the correlation functions perturbatively by bare Lagrangian, one would usually come across UV divergence, which can be removed by introducing counterterms to the Lagrangian. It is ambiguous to determine these counterterms. However, as shown in [110] for the $T\bar{T}$ deformed free scalar, this ambiguity could be fixed by demanding the renormalized Lagrangian can reproduce S-matrix (3.19) or (3.20) order by order. For free massive scalar up to λ^2 -order, the renormalized Lagrangian turns out to be [110]

$$\mathcal{L} = 2\partial\phi\bar{\partial}\phi + \frac{1}{2}m^2\phi^2 - 4g(\partial\phi\bar{\partial}\phi)^2 + \frac{1}{4}hm^4\phi^4 + \dots$$
(3.21)

with

$$g = \lambda + \frac{\lambda^2 m^2}{8\pi} \left(10 \frac{\Lambda^2}{m^2} + 9 \log \frac{m^2}{\Lambda^2} \right),$$

$$h = \lambda + \frac{\lambda^2 m^2}{24\pi} \left(7 \frac{\Lambda^2}{m^2} + 39 \log \frac{m^2}{\Lambda^2} \right).$$
(3.22)

3.3 Correlation functions

Conformal perturbation theory

One of the most direct ways could be using conformal perturbation theory to study the correlation function in $T\bar{T}$ deformed CFT. In this method, one usually considers the correlation function of the undeformed operator in the deformed theory, such as the primary fields in CFT, namely

$$\langle \mathcal{O}_1 \mathcal{O}_2 \cdots \mathcal{O}_n \rangle_{\lambda} = \int [d\phi] \mathcal{O}_1 \mathcal{O}_2 \cdots \mathcal{O}_n e^{-S_{\lambda}[\phi]}$$
 (3.23)

with $\mathcal{O}_i \equiv \mathcal{O}_i(z_i, \bar{z}_i)$ being the primary field in the undeformed CFT. $S_{\lambda}[\phi]$ is the deformed action, and $\lambda = 0$ coreesponds to undeformed CFT. Therefore, in principle, one can expand the RHS in power of λ order by order perturbatively. Nevertheless, some operators indeed deformed under $T\bar{T}$ deformation. Such as the conserved current operator and stress tensor. This is because the EOM is deformed, to keep the conservation of the current or stress tensor under deformation, the functional form of these operators must be modified. Furthermore, it would be more important to define new operators that appear more natural in the deformed theory; for example, the symmetry of the deformed theory may be more manifest based on their operators. The conformal perturbation method was initially employed in [14] [111], where the firstorder two and three-point functions of undeformed primary operators were computed. In this method, one first expands the action in the power of λ as

$$S_{\lambda}[\phi] = S_{CFT}[\phi] + \lambda \int d^2 z T \bar{T} + \lambda^2 S_{(2)}[\phi] + \dots$$
(3.24)

where $S_{(2)}[\phi]$ is the second-order correction of action $S_{\lambda}[\phi]$. It then follows first-order correction of (3.23), which is

$$\langle \mathcal{O}_1 \mathcal{O}_2 \cdots \mathcal{O}_n \rangle^{(1)}_{\lambda} = \lambda \int d^2 z \left\langle T \bar{T}(z, \bar{z}) \mathcal{O}_1 \mathcal{O}_2 \cdots \mathcal{O}_n \right\rangle,$$
 (3.25)

where $\langle ... \rangle_{\lambda}^{(n)}$ represents the *n*-th correction of $T\bar{T}$ -deformed correlation function. And $\langle ... \rangle$ on the RHS denotes the CFT correlation function, which can be computed from conformal Ward identities

$$\langle T(z)\bar{T}(\bar{z})\mathcal{O}_{1}\mathcal{O}_{2}...\mathcal{O}_{n}\rangle = \sum_{i=1}^{n} \left(\frac{h_{i}}{(z-z_{i})^{2}} + \frac{1}{z-z_{i}}\partial_{z_{i}}\right) \left\langle \bar{T}(\bar{z})\mathcal{O}_{1}\mathcal{O}_{2}...\mathcal{O}_{n}\right\rangle,$$

$$\langle \bar{T}(\bar{z})\mathcal{O}_{1}\mathcal{O}_{2}...\mathcal{O}_{n}\rangle = \sum_{i=1}^{3} \left(\frac{\bar{h}_{1}}{(\bar{z}-\bar{z}_{i})^{2}} + \frac{1}{\bar{z}-\bar{z}_{i}}\partial_{\bar{z}_{i}}\right) \left\langle \mathcal{O}_{1}\mathcal{O}_{2}...\mathcal{O}_{n}\right\rangle.$$

$$(3.26)$$

By making use of explicit expression for conformal correlation functions, it turns out one can always arrive at the general form

$$\langle \mathcal{O}_1 \mathcal{O}_2 \cdots \mathcal{O}_n \rangle_{\lambda}^{(1)} = \lambda \langle \mathcal{O}_1 \mathcal{O}_2 \cdots \mathcal{O}_n \rangle \int d^2 z f_n(z, \bar{z}, \{z_i, \bar{z}_i\}),$$
 (3.27)

where the function f_n can be worked out for each n. The above integral is generally divergent and can usually be regularized by dimensional regularization. Consider for example, the cases n = 2 the CFT correlation functions are well-known

$$\langle \mathcal{O}_1 \mathcal{O}_2 \rangle = \frac{C_{12}}{z_{12}^{2h_1} \bar{z}_{12}^{2\bar{h}_1}},$$
(3.28)

where $z_{ij} = z_i - z_j$, and (h_i, \bar{h}_i) the conformal dimension. By plugging (3.28) and (3.26) into (3.25), using dimensional regularization, one obtains

$$\left\langle T(z)\bar{T}(z)\mathcal{O}\left(z_{1},\bar{z}_{1}\right)\mathcal{O}\left(z_{2},\bar{z}_{2}\right)\right\rangle_{\lambda}^{(1)}$$

$$= \lambda h\bar{h}\frac{8\pi}{z_{12}\bar{z}_{12}} \left(\frac{4}{\epsilon} + 2\log|z_{12}|^{2} + 2\log\pi + 2\gamma - 5\right)\left\langle \mathcal{O}\left(z_{1},\bar{z}_{1}\right)\mathcal{O}\left(z_{2},\bar{z}_{2}\right)\right\rangle$$

$$= \lambda h\bar{h}\frac{8\pi C_{12}}{z_{12}^{2h+1}\bar{z}_{12}^{2\bar{h}+1}} \left(\frac{4}{\epsilon} + 2\log|z_{12}|^{2} + 2\log\pi + 2\gamma - 5\right),$$

$$(3.29)$$

where $\epsilon \to 0$ is the parameter introduced in dimensional regularization, γ is Euler constant. By redefining ϵ , we can absorb the constants inside the bracket in the last line of (3.29). In addition, the results show that the first-order correction is universal in the sense that it depends only on the dimension and normalization of the operators. Notice that there are logarithmic divergences in this order. It was suggested that the logarithmic divergence appears in all order in λ [48], and this point will be discussed in a subsequent section. Also, the logarithmic divergence can be understood as a momentum-dependent correction to the conformal dimension [112]. Similar with two-point case, the three-point function first order correction can also be worked out, one can refer to, for example [113], for more details.

As for higher points cases $n \ge 4$, even though correlation functions in CFT are modeldependent and we don't have an explicit expression, one can still work out the integral in (3.25) by making use of the following form

$$\left\langle \mathcal{O}^{\dagger}(z_1, \bar{z}_1) \mathcal{O}(z_2, \bar{z}_2) \mathcal{O}^{\dagger}(z_3, \bar{z}_3) \mathcal{O}(z_4, \bar{z}_4) \right\rangle = \frac{G(\eta, \bar{\eta})}{z_{13}^{2h} z_{24}^{2h} \bar{z}_{13}^{2\bar{h}} z_{24}^{2\bar{h}}}$$
(3.30)

with the cross ratios

$$\eta = \frac{z_{12}z_{34}}{z_{13}z_{24}}, \quad \bar{\eta} = \frac{\bar{z}_{12}\bar{z}_{34}}{\bar{z}_{13}\bar{z}_{24}}.$$
(3.31)

Here, $G(\eta, \bar{\eta})$ is a model-dependent cross-ratio function related to conformal blocks. Then (3.27) turns out to take the form

$$\left\langle \mathcal{O}^{\dagger}\left(z_{1},\bar{z}_{1}\right)\mathcal{O}\left(z_{2},\bar{z}_{2}\right)\mathcal{O}^{\dagger}\left(z_{3},\bar{z}_{3}\right)\mathcal{O}\left(z_{4},\bar{z}_{4}\right)\right\rangle_{\lambda} \\ = \lambda \int d^{2}z \left\{ \left(\frac{hz_{13}^{2}}{\left(z-z_{1}\right)^{2}\left(z-z_{3}\right)^{2}} + \frac{hz_{24}^{2}}{\left(z-z_{2}\right)^{2}\left(z-z_{4}\right)^{2}} + \frac{z_{23}z_{14}}{\prod_{j=1}^{4}\left(z-z_{j}\right)}\frac{\eta\partial_{\eta}G(\eta,\bar{\eta})}{G(\eta,\bar{\eta})} \right) \right. \\ \left(\frac{\bar{h}\bar{z}_{13}^{2}}{\left(\bar{z}-\bar{z}_{1}\right)^{2}\left(\bar{z}-\bar{z}_{3}\right)^{2}} + \frac{\bar{h}\bar{z}_{24}^{2}}{\left(\bar{z}-\bar{z}_{2}\right)^{2}\left(\bar{z}-\bar{z}_{4}\right)^{2}} + \frac{\bar{z}_{23}\bar{z}_{14}}{\prod_{j=1}^{4}\left(\bar{z}-\bar{z}_{j}\right)}\frac{\eta\partial_{\bar{\eta}}G(\eta,\bar{\eta})}{G(\eta,\bar{\eta})} \right) \\ \left. - \eta\bar{\eta}\frac{z_{23}z_{14}\bar{z}_{23}\bar{z}_{14}}{\prod_{i}^{4}\left(z-z_{i}\right)\left(\bar{z}-\bar{z}_{i}\right)}\frac{\partial_{\eta}G(\eta,\bar{\eta})\partial_{\bar{\eta}}G(\eta,\bar{\eta})}{G(\eta,\bar{\eta})^{2}} + \eta\bar{\eta}\frac{z_{23}z_{14}\bar{z}_{23}\bar{z}_{14}}{\prod_{i}^{4}\left(z-z_{i}\right)\left(\bar{z}-\bar{z}_{i}\right)}\frac{\partial_{\eta}\partial_{\bar{\eta}}G(\eta,\bar{\eta})}{G(\eta,\bar{\eta})} \\ \times \left\langle \mathcal{O}^{\dagger}\left(z_{1},\bar{z}_{1}\right)\mathcal{O}\left(z_{2},\bar{z}_{2}\right)\mathcal{O}^{\dagger}\left(z_{3},\bar{z}_{3}\right)\mathcal{O}\left(z_{4},\bar{z}_{4}\right)\right\rangle.$$

$$(3.32)$$

The integral is worked out in [113] [114].

For the general local (non-primary) operator, by the conformal Ward identity, (3.25) can be written as [48]

$$\langle X \rangle^{(1)} = -\frac{1}{\pi} \left(\int d^2 x \sum_{m,n} \sum_{r,s \ge 1} \frac{1}{(z - z_m + \varepsilon)^r (\bar{z} - \bar{z}_n)^s} \right) \langle L_{r-2,m} \bar{L}_{s-2,n} X \rangle^{(0)}, \tag{3.33}$$

where $L_{r-2,m}$ is a shorthand notation for $L_{r-2}(z_m)$. Performing the integration, we obtain

$$\langle X \rangle^{(1)} = \langle \mathrm{d} X \rangle^{(0)}, \quad \mathrm{d} \equiv \sum_{m \neq n} \mathrm{d}_{z_m, z_n}$$

$$(3.34)$$

with

$$d_{z_m, z_n} \equiv \sum_{m \neq n} \left(\log(|z_{mn}|^2 / \varepsilon^2) \partial_{z_m} \partial_{\bar{z}_n} - \sum_{s \ge 2} \frac{1}{s - 1} \frac{\bar{L}_{s-2, n} \partial_{z_m}}{\bar{z}_{mn}^{s-1}} - \sum_{r \ge 2} \frac{1}{r - 1} \frac{L_{r-2, m} \partial_{\bar{z}_n}}{z_{nm}^{r-1}} \right).$$
(3.35)

Superconformal case

The $T\bar{T}$ deformation can be generalized to the case with supersymmetry presented, such as $\mathcal{N} = (0, 1)$ and extend SUSY with $\mathcal{N} = (1, 1), (2, 0), (2, 2)$ [80–84, 87]. In these studies, the supersymmetric versions of $T\bar{T}$ operator was constructed based on the supercurrent multiplets. It was shown that the deformation constructed in this way preserves both the solvability and the SUSY. In analogous to the bosonic case discussed above, we can study the correlation functions when the undeformed theory is supersymmetric and conformal. Working in Euclidean signature with $\mathcal{N} = (1, 1)$ superconformal symmetry, the variation of action under $T\bar{T}$ deformation can be constructed

$$\delta S = \lambda \int d^2 z T \bar{T}(z) = -\lambda \int d^2 z \int d\theta d\bar{\theta} J(Z) \bar{J}(\bar{Z}), \qquad (3.36)$$

where superspace coordinate is denoted by $Z = (z, \theta)$, the J(Z) is the superfield containing the stress tensor $J(Z) = \Theta(z) + \theta T(z)$. In parallel with (3.25) and (3.26), consider the first order correlation function of primary superfield $\Phi(Z, \overline{Z})$, with anomalous dimensions $(\Delta, \overline{\Delta})$

$$-\lambda \int d^2 z \int d\theta d\bar{\theta} \left\langle J(Z)\bar{J}(\bar{Z})\Phi\left(Z_1,\bar{Z}_1\right)\dots\Phi\left(Z_n,\bar{Z}_n\right)\right\rangle.$$
(3.37)

Here, the integrand can be computed via the $\mathcal{N} = (1, 1)$ superconformal Ward identity

$$\left\langle J\left(Z_{0}\right)\Phi_{1}\left(Z_{1},\bar{Z}_{1}\right)\dots\Phi_{n}\left(Z_{n},\bar{Z}_{n}\right)\right\rangle$$

$$=\sum_{i=1}^{n}\left(\frac{\theta_{0i}}{Z_{0i}}\partial_{z_{i}}+\frac{1}{2Z_{0i}}D_{i}+\Delta_{i}\frac{\theta_{0i}}{Z_{0i}^{2}}\right)\left\langle\Phi_{1}\left(Z_{1},\bar{Z}_{1}\right)\dots\Phi_{n}\left(Z_{n},\bar{Z}_{n}\right)\right\rangle,$$
(3.38)

and similar expressions for $\overline{J}(\overline{Z})$. Here *D* is superderivative and $Z_{12} = z_1 - z_2 - \theta_1 \theta_2$ and $\theta_{12} = \theta_1 - \theta_2$ the SUSY invariant distance. Using global $\operatorname{osp}(2|1)$ superconformal transformation, the undeformed two-point function can be fixed up to a constant c_{12} , similar to the bosonic case

$$\left\langle \Phi_1\left(Z_1, \bar{Z}_1\right) \Phi_2\left(Z_2, \bar{Z}_2\right) \right\rangle = c_{12} \frac{1}{Z_{12}^{2\Delta} \bar{Z}_{12}^{2\bar{\Delta}}}, \quad \Delta \equiv \Delta_1 = \Delta_2, \quad \bar{\Delta} \equiv \bar{\Delta}_1 = \bar{\Delta}_2. \tag{3.39}$$

From Ward identity and using dimensional regularization, (3.37) turns out to be [115]

$$\frac{1}{\left\langle \Phi_{1}\left(Z_{1},\bar{Z}_{1}\right)\Phi_{2}\left(Z_{2},\bar{Z}_{2}\right)\right\rangle}\int d^{2}zd\theta d\bar{\theta}\left\langle J(Z)\bar{J}(\bar{Z})\Phi_{1}\left(Z_{1},\bar{Z}_{1}\right)\Phi_{2}\left(Z_{2},\bar{Z}_{2}\right)\right\rangle
= -\frac{4\pi\Delta^{2}}{Z_{12}\bar{Z}_{12}}\left(-\frac{4}{\epsilon}+2\ln|z_{12}|^{2}+2\gamma+2\ln\pi-2\right).$$
(3.40)

By setting $\theta_{1,2} \to 0$

$$-\frac{4\pi\Delta^2}{|z_1|^2} \left(-\frac{4}{\epsilon} + 2\ln|z_{12}|^2 + 2\gamma + 2\ln\pi - 2\right), \qquad (3.41)$$

one recovers results for bosonic CFT (3.29). Note that (3.29) and (3.41) equal up to a constant inside the bracket, and this difference can be eliminated by redefining ϵ . So far, only the 2-point function case has been considered. However, for the general *n*-point function, the first-order correlation functions can also be worked out as the bosonic case.

Torus case

Consider first-order correlation functions in $T\bar{T}$ deformed CFT on a torus in the next step. In this case, we should instead perform the integral (3.25) on a torus T^2 , which is

$$\langle \mathcal{O}_1 \mathcal{O}_2 \cdots \mathcal{O}_n \rangle^{(1)}_{\lambda} = \lambda \int_{T^2} d^2 z \left\langle T \bar{T}(z, \bar{z}) \mathcal{O}_1 \mathcal{O}_2 \cdots \mathcal{O}_n \right\rangle_{T^2}.$$
 (3.42)

Here, the expectation value on the RHS $\langle ... \rangle_{T^2}$ denotes the CFT correlation functions on a torus ⁹, which can not be evaluated using the Ward identity (3.26) on a plane. It turns out that the torus CFT correlation function with an arbitrary number of T and \overline{T} insertion satisfies the following recursion relation [116] ¹⁰

$$\operatorname{tr}(T(w)[T(u_{1})...T(u_{n})\bar{T}(v_{1})...\bar{T}(v_{m})]Xq^{L_{0}-c/12})$$

$$=2\pi i \frac{\partial}{\partial \tau} \operatorname{tr}(T(u_{1})...T(u_{n})\bar{T}(v_{1})...\bar{T}(v_{m})Xq^{L_{0}-c/24})$$

$$=\sum_{i} h_{i} \Big(-\zeta'(w_{i}-w) + 2\eta_{1} \Big) \operatorname{tr}(T(u_{1})...T(u_{n})\bar{T}(v_{1})...\bar{T}(v_{m})Xq^{L_{0}-c/24})$$

$$+\sum_{i} (-\zeta(w_{i}-w) + 2\eta_{1}w_{i} - 2\eta_{1}w - \pi i) \partial_{w_{i}} \operatorname{tr}(T(u_{1})...T(u_{n})\bar{T}(v_{1})...\bar{T}(v_{m})e^{L_{0}-c/24})$$

$$+ \frac{c}{12} \sum_{j} P''(u_{j}-w) \operatorname{tr}(T(u_{1})...\hat{T}(u_{j})...T(u_{n})\bar{T}(v_{1})...\bar{T}(v_{m})Xq^{L_{0}-c/24})$$

$$+ \sum_{j} 2\Big(P(w-u_{j}) + 2\eta_{1}\Big) \operatorname{tr}(T(u_{1})...T(u_{n})\bar{T}(v_{1})...\bar{T}(v_{m})Xq^{L_{0}-c/24})$$

$$+ \sum_{j} \Big(\zeta(w-u_{j}) + 2\eta_{1}u_{j} - 2\eta_{1}w - \pi i\Big) \partial_{u_{i}} \operatorname{tr}(T(u_{1})...T(u_{n})\bar{T}(v_{1})...\bar{T}(v_{m})Xq^{L_{0}-c/24}) .$$

$$(3.44)$$

Here $X = \mathcal{O}_1 \mathcal{O}_2 \cdots \mathcal{O}_n$ is a string of primary operators, the Weierstrass P elliptic function and the Weierstrass ζ function is defined in [117]. From the recursion relation, one has

$$\langle T(w)\bar{T}(\bar{v})X\rangle$$

$$=2\pi i\partial_{\tau}\langle \bar{T}(\bar{v})X\rangle + 2\pi i(\partial_{\tau}\ln Z)\langle \bar{T}(\bar{v})X\rangle + \sum_{i}h_{i}\Big(-\zeta'(w_{i}-w)+2\eta_{1}\Big)\langle \bar{T}(\bar{v})X\rangle$$

$$+\sum_{i}\Big(-\zeta(w_{i}-w)+2\eta_{1}w_{i}-2\eta_{1}w-\pi i\Big)\partial_{w_{i}}\langle \bar{T}(\bar{v})X\rangle.$$
(3.45)

where the well-known result for a single T or \overline{T} insertion is given in [118]. By substituting (3.45) into (3.42), and performing the integral by some proper regularization scheme ¹¹, one could obtain the first-order correlation function on a torus [116].

$$\langle \dots \rangle = \frac{1}{Z} \operatorname{tr}(\dots), \tag{3.43}$$

where Z is the partition function.

¹¹The disk regularization method is used. we excise these singular points $v = w_i$ from the integral domain

$$\lambda \int_{T^2 - \sum_i D(w_i)} d^2 v \langle T(v) \bar{T}(\bar{v}) X \rangle, \qquad (3.46)$$

where $D(w_i)$ is a small disk centered at $v = w_i$.

⁹We will omit subscript T^2 in the following for simplicity.

 $^{^{10}\}mathrm{On}$ a torus, the correlation function is defined as

Partition function: pertubatively

In the last section, we consider $T\bar{T}$ -deformed correlation function on a torus. As a special case, one can consider the zero-point function, i.e., the deformed partition function. The first order correction can be obtained by simply setting X = 1 in (3.45)

$$\left\langle T\left(w_{1}\right)\bar{T}\left(\bar{w}_{2}\right)\right\rangle = -(2\pi i)^{2}\frac{1}{Z}\partial_{\tau}\partial_{\bar{\tau}}Z.$$
(3.47)

Thus

$$Z' = \int D\phi e^{-S + \lambda \int d^2 z T \bar{T}(z)} = Z \left(1 + \lambda \int d^2 z \langle T \bar{T} \rangle(z) \right) \dots = Z + \lambda (2\pi)^2 \tau_2 \partial_\tau \partial_{\bar{\tau}} Z + \dots$$
(3.48)

Note the integrand is in the intermediate steps is finite, hence no regularization is needed. This result is in good agreement with [106].

Next, consider the second-order correction of the deformed partition function. To this end, there are two main difficulties. The first one is that the flow of the stress operator in generic theory is necessary for higher-order computation. Fortunately, this problem was solved recently in [119]. Second, the second-order perturbation involves more complicated integrals; hence, a more intricate regularization method is needed. As a first attempt, one can start with the free boson

$$S = \frac{g}{2} \int_{T^2} d^2 x \partial_\mu \phi \partial^\mu \phi, \qquad (3.49)$$

then, the stress tensor up to first-order are

$$T^{(0)} = -2\pi g(\partial \phi)^2, \quad \bar{T}^{(0)} = -2\pi g(\bar{\partial}\phi)^2, \quad \Theta^{(0)} = 0.$$
 (3.50)

$$T^{(1)} = 8\pi g^2 (\partial \phi)^3 (\bar{\partial} \phi), \quad \bar{T}^{(1)} = 8\pi g^2 (\bar{\partial} \phi)^3 (\partial \phi), \quad \Theta^{(1)} = -4\pi g^2 (\partial \phi \bar{\partial} \phi)^2. \tag{3.51}$$

And up to the second-order, the lagrangian

$$\mathcal{L}^{(1)} = -\frac{1}{\pi^2} T^{(0)} \bar{T}^{(0)} = -4g^2 (\partial \phi \bar{\partial} \phi)^2, \qquad (3.52)$$

$$\mathcal{L}^{(2)} = -\frac{1}{\pi^2} \left(T^{(0)} \bar{T}^{(1)} + \bar{T}^{(0)} T^{(1)} \right) = 32g^3 (\partial \phi \bar{\partial} \phi)^3.$$
(3.53)

It follows that the second-order partition function is

$$\mathcal{Z}^{(2)} = \mathcal{Z}^{(0)} \int_{\mathrm{T}_{1}^{2}} \int_{\mathrm{T}_{2}^{2}} \left\langle \mathcal{L}^{(1)}(z_{1}, \bar{z}_{1}) \mathcal{L}^{(1)}(z_{2}, \bar{z}_{2}) \right\rangle - \mathcal{Z}^{(0)} \int_{\mathrm{T}^{2}} \mathcal{L}^{(2)}(z, \bar{z})$$

$$= 16g^{4} \mathcal{Z}^{(0)} \int_{\mathrm{T}_{1}^{2}} \int_{\mathrm{T}_{2}^{2}} \left\langle \left(\partial_{1} \phi \bar{\partial}_{1} \phi\right)^{2} \left(\partial_{2} \phi \bar{\partial}_{2} \phi\right)^{2} \right\rangle - 32g^{3} \mathcal{Z}^{(0)} \int_{\mathrm{T}^{2}} \left\langle \left(\partial \phi \bar{\partial} \phi\right)^{3} \right\rangle.$$
(3.54)

Similar to the plane case, the correlation function in the integrand is evaluated in CFT, which

can be computed using wick contraction and propagator for torus-free scalar [118]

$$\langle \phi(z_1, \bar{z}_1) \phi(z_2, \bar{z}_2) \rangle = (4\pi g)^{-1} \left(-\log \left| \frac{\vartheta_1(z_{12})}{\eta(\tau)} \right|^2 + 2\pi \frac{(\operatorname{Im}[z_{12}])^2}{\tau_2} \right),$$

$$\langle \partial \phi(z_1, \bar{z}_1) \partial \phi(z_2, \bar{z}_2) \rangle = (4\pi g)^{-1} \left(\frac{\pi}{\tau_2} - 2\eta_1 - P(z_{12}) \right),$$

$$\langle \partial \phi(z_1, \bar{z}_1) \partial \phi(z_1, \bar{z}_1) \rangle = \lim_{z_2 \to z_1} \left(\langle \partial \phi(z_1, \bar{z}_1) \partial \phi(z_2, \bar{z}_2) \rangle + \frac{1}{4\pi g z_{21}^2} \right) = (4\pi g)^{-1} \left(\frac{\pi}{\tau_2} - 2\eta_1 \right).$$

$$(3.55)$$

With the help of this correlation function, the 2nd order partition function can be worked out [120]

$$\mathcal{Z}^{(2)} = 16 \left(\tau_2^2 \partial_\tau^2 \partial_{\bar{\tau}}^2 + i \tau_2 \left(\partial_\tau^2 \partial_{\bar{\tau}} - \partial_{\bar{\tau}}^2 \partial_\tau \right) \right) \mathcal{Z}^{(0)}, \tag{3.56}$$

where minimal subtraction is performed to eliminate the divergent terms. Note (3.56) is consistent with [106]. In parallel with the bosonic case, one could also consider the partition function up to two orders for free fermion. For example, Majorana fermions with spin structure ν , Notice that different from the bosonic case, the deformed Lagrangian for free fermion truncates at the first-order It turns out that the second-order correction of the partition function is also given by

$$\mathcal{Z}_{\nu}^{(2)} = 16 \left(\tau_2^2 \partial_{\tau}^2 \partial_{\bar{\tau}}^2 + i\tau_2 \left(\partial_{\tau}^2 \partial_{\bar{\tau}} - \partial_{\bar{\tau}}^2 \partial_{\tau} \right) \right) \mathcal{Z}_{\nu}^{(0)}.$$
(3.57)

The correction to all orders of the partition function is universal in the sense that these corrections can be expressed in terms of the undeformed partition function [106], as shown here for first and second-order corrections for both free bosons and fermions.

KdV charge one-point function: pertubatively

Let us proceed with the perturbation method to study the deformation of the one-point function of the first KdV charge [121]. There are also studies on deformed KdV charges based on nonperturbation methods [122, 123]. Since the KdV charges are conserved, it will flow under $T\bar{T}$ deformation, similar to the stress tensor, as mentioned at the beginning of this section.

The left-moving KdV charges as P_s^{λ} is a integral of conserved currents $(T_{s+1}^{\lambda}, \Theta_{s-1}^{\lambda})$, with deformation parameter λ

$$P_s^{\lambda} = \frac{1}{2\pi} \int_0^L \left(\, \mathrm{d}z T_{s+1}^{\lambda} + \mathrm{d}\bar{z} \Theta_{s-1}^{\lambda} \right), \qquad (3.58)$$

For the first KdV charge, i.e., s = 1, is distinguished with others since it relates with Hamiltonian $H^{\lambda} = -\int_{0}^{L} dx T_{yy}^{\lambda}$ and momentum $P^{\lambda} = -i \int_{0}^{L} dx T_{xy}^{\lambda}$ of the system, as follows

$$P_1^{\lambda} = \frac{1}{2\pi} \int_0^L \left(\, \mathrm{d}z T^{\lambda} + \mathrm{d}\bar{z} \Theta^{\lambda} \right) \equiv -\frac{H^{\lambda} + P^{\lambda}}{2}. \tag{3.59}$$

The expectation value of P_1^{λ} in the deformed state $|n\rangle^{\lambda}$ thus reads

$${}^{\lambda}\left\langle n\left|P_{1}^{\lambda}\right|n\right\rangle^{\lambda}=-\frac{\mathcal{E}_{n}^{\lambda}+P_{n}^{\lambda}}{2},$$
(3.60)

where \mathcal{E}_n^{λ} and P_n^{λ} represent the energy and momentum of the state $|n\rangle^{\lambda}$, which can be read form deformed spectrum on cylinder

$$\mathcal{E}_{n}^{\lambda} = \frac{L}{2\lambda} \left(\sqrt{1 + \frac{4\lambda E_{n}}{L} + \frac{4\lambda^{2} \left(P_{n}\right)^{2}}{L^{2}}} - 1 \right), \quad P_{n}^{\lambda} = P_{n}, \tag{3.61}$$

where E_n and P_n are the energy and momentum of the undeformed state, respectively. Therefore

$${}^{\lambda}\left\langle n\left|P_{1}^{\lambda}\right|n\right\rangle^{\lambda} = \frac{L}{4\lambda}\left(1 - \sqrt{1 + \frac{4\lambda E_{n}}{L} + \frac{4\lambda^{2}\left(P_{n}\right)^{2}}{L^{2}}}\right) - \frac{P_{n}}{2}.$$
(3.62)

the exact result can be expanded in the power of λ to compare with the results obtained from perturbation theory. In what follows, this comparison will be performed in free CFT with ground state n = 0 case.

In undeformed case, for free bosons with periodic boundary condition $\phi(z+L) = \phi(z)$, and fermions with anti-periodic boundary condition $\psi(z+L) = -\psi(z)$, the vacuum energy and momentum are (n = 0)

$$E_0 = -\frac{\pi c}{6L}, \quad P_0 = 0, \tag{3.63}$$

while for periodic fermions

$$E_0 = \frac{\pi c}{3L}, \quad P_0 = 0, \tag{3.64}$$

Notice that for ground state n = 0, cylinder one-point function (3.62) can be computed from the corresponding one-point functions on the torus by taking the zero temperature limit

$$\lim_{\beta \to \infty} \left\langle \mathcal{O}^{\lambda} \right\rangle_{\text{tor.}}^{\lambda} \equiv \lim_{\beta \to \infty} \left\{ \text{Tr} \left[e^{-\beta H^{\lambda}} \right]^{-1} \cdot \text{Tr} \left[e^{-\beta H^{\lambda}} \mathcal{O}^{\lambda} \right] \right\} = \left\langle \mathcal{O}^{\lambda} \right\rangle_{0}^{\lambda}.$$
(3.65)

Given (3.65), in what follows, we will reproduce (3.62) for the ground state up to first-order in λ in the context of free boson and fermion perturbatively.

In Lagrangian path integral formalism, the one-point function on the LHS (3.65) equals

$$\left\langle \mathcal{O}^{\lambda} \right\rangle_{\text{tor.}}^{\lambda} = \frac{1}{\mathcal{Z}^{\lambda}} \int \mathcal{D}\phi \mathcal{O}^{\lambda} \exp\left\{-\int_{\mathrm{T}^{2}} \mathcal{L}^{\lambda}\right\}$$
 (3.66)

With the help of free propagators given previously and Wick contraction, one obtains for free boson case [120]

$$\langle T^{\lambda} \rangle_{\text{tor.FB}}^{\lambda} = \left(\eta_{1} - \frac{\pi}{2\tau_{2}} \right) + \lambda \cdot \left(\frac{2 |\eta_{1}|^{2}}{\pi} - \frac{1}{2\tau_{2}} (\eta_{1} + \bar{\eta}_{1}) + \left(\frac{2}{\pi} \tau_{2} \bar{\eta}_{1} - 1 \right) i \partial_{\tau} \eta_{1} \right) + O(\lambda^{2}),$$

$$\langle \Theta^{\lambda} \rangle_{\text{tor.FB}}^{\lambda} = \lambda \cdot \left(-\frac{|\eta_{1}|^{2}}{\pi} + \frac{2}{\tau_{2}} (\eta_{1} + \bar{\eta}_{1}) - \frac{3\pi}{4\tau_{2}^{2}} \right) + O(\lambda^{2}),$$

$$(3.67)$$

and for free Majorana fermions

$$\langle T^{\lambda} \rangle_{\nu; \text{ tor.MF}}^{\lambda} = -\frac{e_{\nu-1}}{4} + \lambda \cdot \left(\frac{|e_{\nu-1}|^2}{8\pi} + \frac{i\tau_2 \bar{e}_{\nu-1} \partial_{\tau} e_{\nu-1}}{8\pi} \right) + O\left(\lambda^2\right),$$

$$\langle \Theta^{\lambda} \rangle_{\nu; \text{ tor.MF}}^{\lambda} = \lambda \cdot \frac{-|e_{\nu-1}|^2}{16\pi} + O\left(\lambda^2\right), \quad \nu = 2, 3, 4.$$

$$(3.68)$$

where $\nu = 2, 3, 4$ correspond to the periodic(space)- antiperiodic(time), antiperiodic-periodic and antiperiodic-antiperiodic sectors respectively. Taking zero temperature limit $\tau_2 = \beta \rightarrow \infty$, ¹² one obtains the $T\bar{T}$ -flow of the KdV charge P_1 up to the first-order for three free theories

$$\langle P_1^{\lambda} \rangle_{o,\text{FB}}^{\lambda} = \frac{\pi}{12} + \lambda \cdot \frac{\pi^2}{72} + O\left(\lambda^2\right), \qquad \text{(periodic B.C.)}$$

$$\langle P_1^{\lambda} \rangle_{o,\text{MF}}^{\lambda} = \frac{\pi}{24} + \lambda \cdot \frac{\pi^2}{288} + O\left(\lambda^2\right), \qquad \text{(antiperiodic B.C.)} \qquad (3.69)$$

$$\langle P_1^{\lambda} \rangle_{o,\text{MF}}^{\lambda} = -\frac{\pi}{12} + \lambda \cdot \frac{\pi^2}{72} + O\left(\lambda^2\right). \qquad \text{(periodic B.C.)}$$

It's easy to check that the perturbative results (3.69) match the results come from the nonperturbative method (3.62).

Higher-order correlation functions

In the previous section, we discussed the first-order correlation function (3.25) of undeformed operators on the plane. A natural question is how to compute higher-order corrections. For example, the second-order takes the form

$$\langle \mathcal{O}_1 \mathcal{O}_2 \cdots \mathcal{O}_n \rangle_{\lambda}^{(2)} = \lambda^2 (\int d^2 z_1 d^2 z_2 \left\langle T \bar{T}(z_1, \bar{z}_1) T \bar{T}(z_2, \bar{z}_2) \mathcal{O}_1 \mathcal{O}_2 \cdots \mathcal{O}_n \right\rangle + \dots), \tag{3.70}$$

where the dots involves the term containing the second-order action $S_{(2)}$ in (3.24). Similar to the discussion above (3.53), the explicit expression for $S_{(2)}$ is unknown for general deformed CFT, so the traditional perturbation theory seems not to work for higher-order correlation functions. As an attempt to overcome this problem, a new method is developed to compute $\langle \mathcal{O}_1 \mathcal{O}_2 \cdots \mathcal{O}_n \rangle^{(2)}_{\lambda}$, which based on trace relation of $T\bar{T}$ deformation and conservation of stress tensor [124]. As a consistent check of the universal log² behavior at second order of two-point function can be produced by this method, agreed with prior analyses, the works of Cardy [48] and also [49, 112].

Non-perturbative two-point functions have been derived in three distinct frameworks: the resummation of leading logarithmic divergences at small momentum in [48], the JT-formalism analysis of large-momentum asymptotics in [49], and the worldsheet computation of TsT-deformed vertex operators in [112].¹³ While the results of [49] and [112] align precisely, they differ from [48] by a momentum-dependent factor.

¹²Here for simplicity the modular parameter of the torus is chosen to be pure imaginary $\tau = i\tau_2$.

 $^{^{13}}$ See (3.97) and (3.98) for the explicit expressions for two-point functions.

This discrepancy is addressed in [49], where a perturbative first-order correction yields agreement in the leading log (log¹) term. The analysis therein emphasizes that, within perturbation theory, only the leading logarithmic term is scheme-independent and thus appropriate for crosscomparison. Consistent with this observation, the result in [124] indicates that the subleading contributions to the first- and second-order corrections depend on the renormalization scale μ . Accordingly, a meaningful consistency check should focus on the leading logarithm.

Non-perturbartive method

For $T\overline{T}$ deformation with general parameter λ , it was found that on a 2D plane, the flow of the deformed correlation function can be written as the flow of each field [48]

$$\partial_{\lambda} \langle \prod_{n} \Phi_{n}(x_{n}) \rangle_{\lambda} = \sum_{n} \langle \partial_{\lambda} \Phi_{n}^{\lambda}(x_{n}) \prod_{m \neq n} \Phi_{m}^{\lambda}(x_{m}) \rangle$$
(3.71)

with

$$\partial_{\lambda}\Phi^{\lambda}(x) = 2\pi\epsilon^{ab}\epsilon^{ij}\int_{x}^{X} dx'_{j}T^{\lambda}_{ai}(x'+\varepsilon)\partial_{x^{b}}\Phi^{\lambda}(x).$$
(3.72)

The leading divergence on the RHS follows from OPE of T and Φ (3.72)

$$\partial_{\lambda} \Phi^{\lambda}(x) = -(\log |\varepsilon|) \nabla_x^2 \Phi^{\lambda}(x) + \dots$$
(3.73)

The all-orders analysis was performed in [48], and it turns out that the logarithmic divergence occurred at all orders, with leading terms of the form $(\lambda \log \Lambda)^N$ in each order. Resume all such logarithmic terms divergence and work in momentum space, which may be more convenient than the position space for non-local theory, yielding

$$\Phi^{\lambda}(k) = e^{-\lambda \log(\Lambda/\mu)k^2} \Phi^0(k).$$
(3.74)

It then follows that the renormalized field can be defined by canceling the logarithmic divergence

$$\widehat{\Phi}^{\lambda}(k) \equiv e^{\lambda \log(\Lambda/\mu)k^2} \Phi^{\lambda}(k), \qquad (3.75)$$

which is a non-local renormalization in contrast with the renormalization of local fields. Since the bare correlation functions of field Φ^{λ} are independent of scale μ , it follows the correlation functions satisfy Callan-Symanzik-like equation. For example the renormalized two-point function in momentum space $\hat{C}(k; \lambda, \mu)$ satisfies

$$\left[\mu\partial_{\mu} + k\partial_{k} - 2\lambda\partial_{\lambda} - 2\Delta\right]\widehat{C}(k;\lambda,\mu) = 0.$$
(3.76)

From (3.99), for undeformed case, i.e., CFT two-point function in momentum space is $\widehat{C}(k; \lambda = 0) \sim k^{2\Delta-2}$ ¹⁴ At fixed renormalization scale μ , the solution then follows

$$\widehat{C}(k;\lambda,\mu) = k^{2\Delta-2} (k/\mu)^{2\lambda k^2}.$$
(3.77)

¹⁴However, the two-point function reads differently $\widehat{C}(k; \lambda = 0) \sim k^{2\Delta - 2}$, as indicated above eq(4.25) in [48].

To match the correlation functions obtained from the dual string theory side, a more general solution to (3.76) was proposed by multiplying an addition factor $F(\lambda k^2)$ with F(0) = 1 to (3.77) [112].

3.4 Correlation functions: further developments

In the last subsection, we mainly focus on the conformal perturbation theory in computing deformed correlation functions. There have been much progress going beyond traditional perturbation methods, including the methods based on dynamical coordinate transformation [119] and flat JT gravity perspective [49, 125], random geometry [126], and point-splitting methods [48, 112, 126]. In this subsection, we will briefly review the first two methods. Let us begin with the former case. In [48], it was found that the field transformed under $T\bar{T}$ deformation as ¹⁵

$$\partial_{\lambda}\Phi^{\lambda}(x) = 2\pi\epsilon^{ab}\epsilon^{ij}\int_{x}^{X} dx'_{j}T^{\lambda}_{ai}(x'+\varepsilon)\partial_{x^{b}}\Phi^{\lambda}(x), \qquad (3.79)$$

where X is an arbitrary point, the integral path can be anyone, provided it does not intersect with the points where other fields are inserted inside the correlation function. ε is the pointsplitting regulator. (3.79) suggests that instead of deforming the field, one can deform the coordinate, i.e.,

$$\partial_{\lambda} \Phi^{\lambda}(x) = \partial_{x^{b}} \Phi^{\lambda}(x) \cdot \partial_{\lambda} x^{b}$$
(3.80)

with

$$\partial_{\lambda}x^{b} = 2\pi\epsilon^{ab}\epsilon^{ij}\int_{x}^{X}dx'_{j}T^{\lambda}_{ai}(x'+\varepsilon) = 2\pi\epsilon^{ab}\epsilon^{ij}\int_{x+\varepsilon}^{X}dx'_{j}T^{\lambda}_{ai}(x').$$
(3.81)

In component, this becomes

$$\partial_{\lambda}x^{z} = \partial_{\lambda}z = 4\pi \int_{x+\varepsilon}^{X} \left(dz'T_{\bar{z}\bar{z}}^{\lambda}(z') - d\bar{z}'T_{\bar{z}\bar{z}}^{\lambda}(z') \right),$$

$$\partial_{\lambda}x^{\bar{z}} = \partial_{\lambda}\bar{z} = 4\pi \int_{x+\varepsilon}^{X} \left(d\bar{z}'T_{z\bar{z}}^{\lambda}(z') - dz'T_{zz}^{\lambda}(z') \right),$$
(3.82)

which can also be derived from a random geometry perspective [97]. Put (3.82) equivalently into an integral form on a plane that is convenient for computing correlation function below $[119]^{16}$

$$\partial_{\lambda} x^{z} = \partial_{\lambda} z = 4\pi \int_{x+\varepsilon}^{X} \left(dz' T^{\lambda}_{\bar{z}z}(z') - d\bar{z}' T^{\lambda}_{\bar{z}\bar{z}}(z') \right)$$

$$= -2i \int dz' \wedge d\bar{z}' T^{\lambda}_{\bar{z}'\bar{z}'}(z') \left(\frac{1}{x-z'} - \frac{1}{X-z'} \right),$$
(3.84)

 15 Follow the conventions in [118]

$$e^{z\bar{z}} = -2i, \quad \epsilon_{z\bar{z}} = \frac{i}{2}, \quad g_{z\bar{z}} = \frac{1}{2}, \quad g^{z\bar{z}} = 2.$$
 (3.78)

¹⁶Here $dz \wedge d\bar{z} = -2idx \wedge dy$, and Stokes theorem

$$\int dz \wedge d\bar{z} (\partial_z F^z + \partial_{\bar{z}} F^{\bar{z}}) = \oint_C (F^z d\bar{z} - F^{\bar{z}} dz).$$
(3.83)

or equivalently,

$$z \longmapsto Z^{(\mu|\delta\mu)}(z,\bar{z}) = z + \frac{\delta\mu}{2\pi^2} \int_{\mathbb{R}^2} d^2x \frac{\bar{T}^{(\mu)}(x,\bar{x})}{z-x} + \mathcal{O}\left(\delta\mu^2\right).$$
(3.85)

This map is called dynamical coordinate transformation. Here the theory $\mathcal{T}^{(\mu)}$ is defined on the deformed space $Z^{(\mu|\delta\mu)}$ and the deformed theory $\mathcal{T}^{(\mu+\delta\mu)}$ is defined on undeformed space z (see also [4,5,91]). It was proposed that under $T\bar{T}$ -deformation, the theory deformed by λ_1 on space deformed by λ_2 is equivalent to the theory deformed by μ_1 on deformed space by μ_2 [48,119]. In the notation of [119], this equivalence can be expressed as

$$\mathcal{T}^{(\lambda_1)}\left[\mathbb{R}^2_{(\lambda_1|\lambda_2)}\right] = \mathcal{T}^{(\mu_1)}\left[\mathbb{R}^2_{(\mu_1|\mu_2)}\right], \quad \text{with} \quad \lambda_1 + \lambda_2 = \mu_1 + \mu_2, \tag{3.86}$$

where the coordinates on space $\mathbb{R}^2_{(\lambda_1|\lambda_2)}$ is denoted by $Z^{(\lambda_1|\lambda_2)}$. A special case is

$$\mathcal{T}^{(\mu)}\left[\mathbb{R}^{2}_{(\mu|0)}\right] = \mathcal{T}^{(0)}\left[\mathbb{R}^{2}_{(0|\mu)}\right],\tag{3.87}$$

which demonstrates that the $T\bar{T}$ -deformed CFT $\mathcal{T}^{(\mu)}$ on undeformed plane $\mathbb{R}^2_{(\mu|0)} \equiv \mathbb{R}^2$ (with coordinate (z, \bar{z})) is equivalent to undeformed CFT $\mathcal{T}^{(0)}$ on deformed space $\mathbb{R}^2_{(0|\mu)}$ (with coordinate (Z, \bar{Z})), the dynamic coordinate transformation in this case is

$$Z^{(\mu)}(z,\bar{z}) = z + \frac{\mu}{2\pi^2} \int_{\mathbb{R}^2} d^2 x \frac{\bar{T}^{(\mu)}(x,\bar{x})}{z-x},$$
(3.88)

Note the direct relation between infinitesimal version (3.85) and finite version (3.88) is checked for some particular case [119]. This equivalence relation (3.86) is proper both technically and conceptually. The flow of stress tensor can be inferred from (3.86), which is inaccessible in traditional perturbation expansion once the Lagrangian is unknown. Furthermore, (3.86) plays an important role in computing correlation functions in deformed theory. In analogy to quantum mechanics, there are two pictures, namely "Heisenberg" and "Schrödinger," to describe the deformed correlation function [48, 119]

$$\langle \mathcal{O}_{\Delta_1,\bar{\Delta}_1}^{(\lambda)}(z_1,\bar{z}_1)\cdots\mathcal{O}_{\Delta_n,\bar{\Delta}_n}^{(\lambda)}(z_n,\bar{z}_n)\rangle_0 = \langle \mathcal{O}_{\Delta_1,\bar{\Delta}_1}^{(0)}(z_1,\bar{z}_1)\cdots\mathcal{O}_{\Delta_n,\bar{\Delta}_n}^{(0)}(z_n,\bar{z}_n)\rangle_\lambda, \tag{3.89}$$

where $\langle ... \rangle_{\lambda}$ ($\langle ... \rangle_{0}$) denotes the correlation functions in the deformed (undeformed) theory. $\mathcal{O}_{\Delta_{1},\bar{\Delta}_{1}}^{(\lambda)}$ are deformed operator in the "Heisenberg" picture Note the deformed correlation function on the RHS can be converted to a CFT correlation function on the LHS. While the operator on LHS is related to the undeformed one via dynamic coordinates transformation

$$\mathcal{O}_{\Delta,\bar{\Delta}}^{(\lambda)}(z,\bar{z}) = \left(\det\frac{\partial x}{\partial X}\right)^{-\frac{\Delta+\bar{\Delta}}{2}} \mathcal{O}_{\Delta,\bar{\Delta}}^{(0)}(X)$$

$$= \left(\det\frac{\partial x}{\partial X}\right)^{-\frac{\Delta+\bar{\Delta}}{2}} \left(\mathcal{O}_{\Delta,\bar{\Delta}}^{(0)}(z) + \delta z \partial_z \mathcal{O}_{\Delta,\bar{\Delta}}^{(0)}(z) + o(\mu^2)\right),$$
(3.90)

where $x = (z, \bar{z})$, and $X = (Z, \bar{Z})$ defined in (3.88). Note the 2nd line is an expansion perturbatively in μ such that operators are defined in CFT and undeformed coordinates. Therefore, using (3.90), the LHS of (3.89) is expressed in terms of the CFT correlation function and can be evaluated by proper regularization. For example, the two-point function to first order

$$\langle \mathcal{O}_{\Delta_{1,\bar{\Delta}_{1}}}^{(\lambda)}(z_{1},\bar{z}_{1}) \mathcal{O}_{\Delta_{n,\bar{\Delta}_{n}}}^{(\lambda)}(z_{2},\bar{z}_{2}) \rangle_{0}$$

$$= \frac{1}{|z_{12}|^{4\Delta}} - \frac{\lambda\Delta}{\pi^{2} z_{12}^{2\Delta+1}} \int d^{2}x \frac{\langle \bar{T}(\bar{x}) \mathcal{O}_{\Delta}(\bar{z}_{1}) \mathcal{O}_{\Delta}(\bar{z}_{2}) \rangle_{0}}{z_{1} - x} + (z_{1} \leftrightarrow z_{2}) + \text{ c.c.}$$

$$= \frac{1}{|z_{12}|^{4\Delta}} - \frac{8\lambda\Delta^{2}}{\pi |z_{12}|^{4\Delta}} \left(\frac{\ln |z_{12}/\epsilon|^{2}}{|z_{12}|^{2}} - \frac{1}{|z_{12}|^{2}} - \frac{1}{2\epsilon} \left(\frac{1}{z_{12}} + \frac{1}{\bar{z}_{12}} \right) \right).$$

$$(3.91)$$

For the second order, the leading-log contribution can also be reproduced

$$\left\langle \mathcal{O}_{\Delta,\Delta}^{(\lambda)}(z_1,\bar{z}_1) \, \mathcal{O}_{\Delta,\Delta}^{(\lambda)}(z_2,\bar{z}_2) \right\rangle_0 = \dots + \frac{8\lambda^2}{\pi^2} \Delta^2 (2\Delta+1)^2 \frac{\ln^2 |z_{12}/\epsilon|^2}{|z_{12}|^{4(\Delta+1)}} + \dots \tag{3.92}$$

In summary, the dynamical coordinate transform offers a way of computing higher-order correlation functions, contrasting with the traditional perturbation theory.

Now, we turn to another method for deformed correlation functions based on flat JT gravity interpretation of $T\bar{T}$ deformation. As mentioned in the beginning of sec. 3.3, perturbatively, one usually considers the correlation function of undeformed theory. A question is, how can the appropriate operator be defined in the deformed theory? ¹⁷ It was suggested that the flat JT gravity interpretation of $T\bar{T}$ deformation could define the deformed operator as described below.

As introduced in the section 2, the action of $T\bar{T}$ deformed theory can be written as

$$S_{T\bar{T}}(\psi, e^a_{\alpha}, X^a) = S_0(\psi, e^a_{\alpha}) + S_{JT}(e^a_{\alpha}, X^a)$$

= $S_0(\psi, e^a_{\alpha}) + \frac{\Lambda}{2} \int d^2 \sigma \epsilon^{\alpha\beta} \epsilon_{ab} \left(\partial_{\alpha} X^a - e^a_{\alpha}\right) \left(\partial_{\beta} X^b - e^b_{\beta}\right),$ (3.93)

where X is the dynamical coordinates, which is the original space of the undeformed theory. σ^a is the coordinates of an auxiliary space. One can introduce local operators on X as a deformation of undeformed operators ¹⁸

$$\mathcal{O}(X_0) = \int d^2 \sigma \sqrt{g(\sigma)} O(\sigma) \delta(X(\sigma) - X_0).$$
(3.94)

Transforming into momentum space

$$\mathcal{O}(q_0) = \int d^2 \sigma \sqrt{g(\sigma)} O(\sigma) \exp\left(iq_0 \cdot X(\sigma)\right).$$
(3.95)

The corresponding correlation functions are then defined by the path integral

$$C(q_1, q_2) \equiv \langle \mathcal{O}(q_1) \mathcal{O}(q_2) \rangle = \frac{1}{Z_{T\bar{T}}} \int \frac{\mathcal{D}e\mathcal{D}X\mathcal{D}\psi}{V_{\text{diff}}} \mathcal{O}(q_1) \mathcal{O}(q_2) e^{-S_{T\bar{T}}}.$$
 (3.96)

¹⁷For discussion in this regard, see, for example, [127]. In Minkowski signature, two kinds of operators, namely undeformed and dressed ones, and their correlation functions were discussed in [128].

 $^{^{18}}$ For discussion on correlation functions of undeformed operators in gravity action formalism, see [129].

In $T\overline{T}$ -deformed theories, the two-point correlation function in momentum space in the large momentum limit is given by the expression [49]

$$\frac{C}{|\Lambda|} \frac{1}{\sin(\frac{q^2}{2|\Lambda|} + \pi\Delta)} \left(\frac{\mu}{2e}\right)^{2\Delta} \left(\frac{2\pi e|\Lambda|}{\mu|q|}\right)^{\frac{q^2}{\pi|\Lambda|} + 2\Delta},\tag{3.97}$$

which in contrast with Cardy's results [48], which is (3.77) but in a different notation

$$C_t(q) = C_0(q) \left(\frac{|q|}{\mu}\right)^{\frac{tq^2}{\pi}},$$
(3.98)

 $C_0(q)$ is the undeformed conformal field theory (CFT) correlation function, μ is a renormalization scale, and t is the deformation parameter. The correlation function (3.97) reveals the non-local nature of $T\overline{T}$ -deformations, ¹⁹ especially at large momenta , a key feature of these deformed theories. Additionally, renormalization in these models is momentum-dependent, contrasting with local quantum field theories. The function also exhibits ultraviolet (UV) divergences, which are regulated through techniques like point-splitting, further emphasizing the unique properties of $T\overline{T}$ -deformed models in locality and operator renormalization. This result highlights the deformation's impact on short-distance physics, differing significantly from conventional QFTs.

In [125], the two-point function (3.97) on plane was generalized to torus, based also on the JT gravity formulation of $T\bar{T}$ deformation. On a torus, at small momenta, $|q| \ll L/t$, the correlation function behaves similarly to the plane case, where $C_t(q) \propto |q|^{-\frac{tq^2}{\pi}}$, with t as the deformation parameter, q the momentum, and L the torus length scale. However, for large momenta, $|q| \gg L/t$, the geometry of the torus modifies the correlation function, yielding the expression

$$C_t(q) \propto \left(4\sqrt{\frac{tq^2}{\pi eL^3|T|^2}}\right)^{\frac{tq^2}{\pi}},\tag{3.100}$$

where T is the modular parameter of the torus. The main properties of the correlation function include UV-IR mixing, where the high-energy (UV) behavior is influenced by the global spacetime structure (IR), a characteristic feature of non-local theories. Furthermore, the deformation introduces momentum smearing, where operators carrying large momentum q are smeared over a distance scale proportional to t|q|, highlighting the non-local nature of the deformation. The torus geometry also affects the correlation function significantly. At the same time, small |q|behavior mimics that of the plane; at large |q|, the correlation function first decays and then

$$\int d^2x \frac{e^{i\mathbf{p}\cdot\mathbf{x}}}{|\mathbf{x}|^{2\Delta}} = \int dr d\theta r \frac{e^{ipr\cos\theta}}{r^{2\Delta}} = \int dr \frac{2\pi J_0(pr)}{r^{2\Delta-1}} = 2^{2-2\Delta} \pi \frac{\Gamma(1-\Delta)}{\Gamma(\Delta)} p^{2\Delta-2}.$$
(3.99)

¹⁹In momentum space, power-law behavior $|p|^n$ is a characteristic of local field theory. The two-point correlation function in CFT in momentum space is

grows, implying that the correlation function diverges at very large momenta. Overall, the correlation function in the $T\overline{T}$ -deformed theory on a torus reflects the distinct non-local properties induced by the deformation, deviating from the typical power-law decay found in traditional quantum field theories.

3.5 Quantum entanglement in $T\overline{T}$ deformed field theory

The investigation of entanglement entropy in $T\overline{T}$ -deformed quantum field theories has attracted considerable interest for its profound implications in quantum information and non-local field dynamics. As an irrelevant deformation, $T\overline{T}$ induces inherent non-locality and modifies infrared physics, providing a novel framework for probing deformed quantum systems. In twodimensional spacetime, the classical equivalence between $T\overline{T}$ -deformed theories and Jackiw-Teitelboim (JT) gravity-coupled systems has been extended to higher dimensions [46], prompting investigations into whether this correspondence persists at the quantum level.

A central challenge emerges from the divergent algebraic structures [130, 131] underlying quantum field theories and gravitational systems, particularly their consistent incorporation within $T\overline{T}$ -deformed frameworks. Entanglement entropy is a critical holography probe for analyzing deformation-induced geometric modifications and their dual gravitational manifestations. The intrinsic non-locality of $T\overline{T}$ deformation establishes direct connections to quantum gravity paradigms, where non-local effects play fundamental roles.

The $T\overline{T}$ -deformed conformal field theory (CFT) has emerged as a holographic dual to quantum gravity in finite spacetime regions. As proposed in [132] and supported by [12], this duality suggests that $T\overline{T}$ -deformed CFTs encode gravitational physics within bounded geometries. A key feature of this deformation is its introduction of a finite radial cutoff in AdS spacetimes, which modifies holographic prescriptions like the Ryu-Takayanagi formula [133]. Notably, the deformation reduces subsystem degrees of freedom, rescales energy scales, and can trigger phase transitions even for fixed subsystem sizes [133, 134]. Recent work extends these insights across diverse holographic settings. In [135], the replica sphere method was generalized to derive deformed entanglement entropy for arbitrary single intervals in both AdS and dS holography, revealing non-locality in timelike boundary theories under $T\overline{T}$ deformation. While multi-interval entanglement entropy in holographic CFTs typically sums single-interval contributions at large separations [136], the deformation introduces novel constraints. For 2D systems, an alternative interpretation frames $T\overline{T}$ -deformed CFTs as AdS₃ gravity with mixed boundary conditions [21], where coordinate transformations connect Bañados geometries to holographic entanglement entropy calculations [137], consistent with earlier results [138]. Progress in quantifying entanglement includes perturbative expressions for negativity in bipartite mixed states [139] and explorations of odd, reflected, and timeline entanglement entropies [140–142]. Notably, [143] recently proposed a holographic covariant entropy bound (HCEB) connecting bulk Cauchy

slices to boundary states via codimension-2 surfaces. In 3D AdS gravity with $T\overline{T}$ -deformed boundaries, HCEB matches extremal/marginal surface areas but exceeds trapped surface areas (e.g., inside black holes), enabling tensor network constructions for holographic entropy.

The behavior of the entanglement entropy in the deformed CFTs, especially in the context of the TT-deformations, has been extensively studied in recent years [132, 137, 138, 144, 145]. Recent studies on entanglement entropy in $T\overline{T}$ -deformed CFTs reveal key insights into the quantum structure of these systems. [138] explored leading-order corrections in vacuum states using perturbative methods within the twistor operator formalism. In generic 2D CFTs, [113] constructed the perturbative correlation functions to evaluate the entanglement entropy of the locally excited states and further demonstrated that the deformation preserves the maximal chaotic behavior of the holographic CFTs. It has been shown that pseudo entropy, like entanglement entropy, can be used to quantify the topological contributions of entanglement of excited states in two-dimensional rational CFTs [146–148]. More recently, the generalized entanglement entropy called pseudo-entanglement entropy has been proposed via the AdS/CFT correspondence and post-selection [149]. The pseudo-entanglement entropy corrected by the deformation has been investigated [150], which shows that the deformation does not change the characteristic behaviors of the quantum chaos [113]. In the massive quantum fields with $T\overline{T}$ [145], the leading correction to entanglement entropy in $T\overline{T}$ -deformed massive quantum field theories arises from the boundary of the entangling surface, with a finite correction for scalars and a log-square divergence for Dirac fermions. By applying a form factor bootstrap approach, the paper [151] calculates the $T\overline{T}$ -deformed entanglement entropy for integrable quantum field theories, revealing that the UV behavior remains similar to the undeformed case with a modified central charge.

3.6 $T\bar{T}$ flow at large central charge

In this subsection, we briefly switch gears to discuss the "more classical" aspects of $T\bar{T}$ -deformed quantum field theory. We review the flow equation of partition function and one-point function of stress tensor for $T\bar{T}$ -deformed CFTs in the large central charge limit. This flow equation leads to the important trace relation (2.22) that has been proved for classical field theories in previous subsections. Most importantly, it is closely related to the holographic interpretation of $T\bar{T}$ deformation for holographic CFTs, preparing for the discussion in the next section. Our derivation largely follows the original paper [21] with some technical and conceptual tweaks.

The large central charge limit is a limit of large degrees of freedom, similar to the large N limit of gauge theory. It was brought into the context of $T\bar{T}$ deformation in [76]

$$c$$
 large, λc comparable to the energy scales considered (3.101)

and its implication for correlation functions was discussed there. The key simplification of this

limit is that the connected correlation functions of the energy-momentum tensor all scale as the central charge c, so dominant contributions to a correlation function come from factorizations into the largest number of (nonvanishing) factors (large c factorization). This line of argument was well known for multitrace deformations, for example, in [152]. Consider a theory defined by the action $S^{[0]}[\phi, \gamma]$ with ϕ being the fundamental field in path integration and γ_{ij} the background metric, the partition function and generating functional for energy-momentum tensor are given by

$$Z^{[0]}[\gamma] = \int \mathcal{D}\phi e^{-S^{[0]}[\phi,\gamma]},$$

$$I^{[0]}[\gamma] = -\log Z^{[0]}[\gamma].$$
(3.102)

Connected correlation functions of the energy-momentum tensor are given by functional derivatives of the generating functional with respect to the background metric. If the theory is deformed by the $T\bar{T}$ deformation with infinitesimal deformation parameter λ , the partition function of the deformed theory is

$$Z[\gamma] = \int \mathcal{D}\phi e^{-S^{[0]}[\phi,\gamma] + \frac{1}{2}\lambda \int d^2x \sqrt{\gamma} \epsilon^{\mu\nu} \epsilon^{\rho\sigma} T^{[0]}_{\mu\rho} T^{[0]}_{\nu\sigma}}.$$
(3.103)

By doing a Hubbard-Stratonovich transformation ²⁰, that is, inserting the identity

$$1 = \sqrt{\det(\lambda M_{\gamma})} \int \mathcal{D}h e^{-\frac{1}{2}\lambda \int d^2x \sqrt{\gamma} \epsilon^{\mu\nu} \epsilon^{\rho\sigma} (h_{\mu\rho} + T^{[0]}_{\mu\rho})(h_{\nu\sigma} + T^{[0]}_{\nu\sigma})}.$$
 (3.104)

where M_{γ} denotes the operator $M_{\gamma}(f,g) = \int d^2x \sqrt{\gamma} \epsilon^{\alpha\gamma} \epsilon^{\beta\delta} f_{\alpha\beta} g_{\gamma\delta}$, we find

$$Z[\gamma] = \int \mathcal{D}\phi \sqrt{\det(\lambda M_{\gamma})} \int \mathcal{D}h e^{-S^{[0]}[\phi,\gamma_{\mu\nu}] - \frac{1}{2}\lambda \int d^{2}x \sqrt{\gamma} \epsilon^{\mu\rho} \epsilon^{\nu\sigma} h_{\mu\nu} h_{\rho\sigma} - \lambda \int d^{2}x \sqrt{\gamma} \epsilon^{\mu\rho} \epsilon^{\nu\sigma} T^{[0]}_{\mu\nu} h_{\rho\sigma}}$$

$$= \sqrt{\det(\lambda M_{\gamma})} \int \mathcal{D}h e^{-\frac{1}{2}\lambda \int d^{2}x \sqrt{\gamma} \epsilon^{\mu\rho} \epsilon^{\nu\sigma} h_{\mu\nu} h_{\rho\sigma}} \int \mathcal{D}\phi e^{-S^{[0]}[\phi,\gamma_{\mu\nu}] - \lambda \int d^{2}x \sqrt{\gamma} \epsilon^{\mu\rho} \epsilon^{\nu\sigma} T^{[0]}_{\mu\nu} h_{\rho\sigma}}$$

$$\approx \sqrt{\det(\lambda M_{\gamma})} \int \mathcal{D}h e^{-\frac{1}{2}\lambda \int d^{2}x \sqrt{\gamma} \epsilon^{\mu\rho} \epsilon^{\nu\sigma} h_{\mu\nu} h_{\rho\sigma}} \int \mathcal{D}\phi e^{-S^{[0]}[\phi,\gamma^{\mu\nu} + 2\lambda \epsilon^{\mu\rho} \epsilon^{\nu\sigma} h_{\rho\sigma}]}$$

$$= \sqrt{\det(\lambda M_{\gamma})} \int \mathcal{D}h e^{-\frac{1}{2}\lambda \int d^{2}x \sqrt{\gamma} \epsilon^{\mu\rho} \epsilon^{\nu\sigma} h_{\mu\nu} h_{\rho\sigma} - I^{[0]}[\gamma^{\mu\nu} + 2\lambda \epsilon^{\mu\rho} \epsilon^{\nu\sigma} h_{\rho\sigma}]}.$$
(3.105)

In the third line we only keep terms up to the first order in λ . In the large c limit, we approximate the path integral by the saddle point, determined by the equation

$$h_{\mu\nu} = -\langle T_{\mu\nu} \rangle^{[0]} [\gamma^{\alpha\beta} + 2\lambda \epsilon^{\rho\alpha} \epsilon^{\sigma\beta} h_{\rho\sigma}].$$
(3.106)

 $h_{\mu\nu}$ appears on the right-hand side in the background metric, making the equation a complicated functional equation, so there is no simple expression for the saddle point. It's thus

 $^{^{20}}$ Reviewed in previous sections, this transformation has been used in [97] to derive the flow equation of partition functions in spaces with simple geometries where considerable simplification of the path integral over h is possible.

convenient to introduce the background metric on the right-hand side as a new variable $\gamma^{[0]^{\mu\nu}}$, the "undeformed metric". Completing the path integral in (3.105), we find to first order in λ

$$I[\gamma_{\mu\nu}] = I^{[0]}[\gamma^{[0]}_{\mu\nu}] + \frac{1}{2}\lambda \int d^2x \sqrt{\gamma^{[0]}} \epsilon^{[0]\mu\rho} \epsilon^{[0]\nu\sigma} \langle T_{\mu\nu} \rangle^{[0]} \langle T_{\rho\sigma} \rangle^{[0]}.$$
(3.107)

In addition we have, by our definition

$$\gamma^{\mu\nu} = \gamma^{[0]^{\mu\nu}} + 2\lambda\epsilon^{[0]^{\rho\alpha}}\epsilon^{[0]^{\sigma\beta}} \langle T_{\rho\sigma} \rangle^{[0]}.$$
(3.108)

If we consider $T\bar{T}$ deformation with a continuously varying parameter λ , the infinitesimal equations above become differential equations

$$\partial_{\lambda} I^{[\lambda]}[\gamma_{\alpha\beta}^{[\lambda]}] = \frac{1}{2} \int d^2 x \sqrt{\gamma^{[\lambda]}} \epsilon^{[\lambda]^{\mu\rho}} \epsilon^{[\lambda]^{\nu\sigma}} \langle T_{\mu\nu} \rangle^{[\lambda]} \langle T_{\rho\sigma} \rangle^{[\lambda]} = -\frac{1}{2} \int d^2 x \sqrt{\gamma^{[\lambda]}} (\langle T_{\mu\nu} \rangle^{[\lambda]} \langle T^{\mu\nu} \rangle^{[\lambda]} - \langle T_{\rho}^{\rho} \rangle^{[\lambda]^2}) = -\int d^2 x \sqrt{\gamma^{[\lambda]}} \langle O_{T\bar{T}} \rangle^{[\lambda]}$$
(3.109)

and

$$\partial_{\lambda}\gamma^{[\lambda]^{\mu\nu}} = 2\epsilon^{[\lambda]^{\mu\rho}}\epsilon^{[\lambda]^{\nu\sigma}}\langle T_{\rho\sigma}\rangle^{[\lambda]} = -2\langle T^{\mu\nu}\rangle^{[\lambda]} + \langle T^{\rho}_{\rho}\rangle^{[\lambda]}\gamma^{[\lambda]^{\mu\nu}}$$
(3.110)

To derive the flow equation of the one-point function, we take a variation in the metric of (3.109) and obtain

$$\partial_{\lambda}(\sqrt{\gamma^{[\lambda]}}\langle T_{\mu\nu}\rangle^{[\lambda]}\delta\gamma^{[\lambda]\mu\nu}) = -2\delta(\sqrt{\gamma^{[\lambda]}}\langle O_{T\bar{T}}\rangle^{[\lambda]}).$$
(3.111)

It further simplifies to

$$\partial_{\lambda}(\sqrt{\gamma}\langle T_{\mu\nu}\rangle\rangle)\delta\gamma^{\mu\nu} + \sqrt{\gamma}\langle T_{\mu\nu}\rangle\delta(\partial_{\lambda}\gamma^{\mu\nu}) = \sqrt{\gamma} \Big[\big(\langle O_{T\bar{T}}\rangle\gamma_{\mu\nu} + 2\langle T_{\mu\rho}\rangle\langle T_{\nu}^{\rho}\rangle - 2\langle T_{\rho}^{\rho}\rangle\langle T_{\mu\nu}\rangle\big)\delta\gamma^{\mu\nu} - \langle T_{\mu\nu}\rangle\delta\big(2\langle T^{\mu\nu}\rangle - 2\gamma^{\mu\nu}\langle T_{\rho}^{\rho}\rangle\big) \Big]$$
(3.112)

where for simplicity we have omitted the notation for λ -dependence. Plugging (3.110) into the equation, we find

$$\partial_{\lambda}(\sqrt{\gamma}\langle T_{\mu\nu}\rangle) = \sqrt{\gamma} \left(\langle O_{T\bar{T}} \rangle \gamma_{\mu\nu} + 2\langle T_{\mu\rho} \rangle \langle T_{\nu}^{\rho} \rangle - 2\langle T_{\rho}^{\rho} \rangle \langle T_{\mu\nu} \rangle \right)$$
(3.113)

Introducing the trace-reversed energy-momentum tensor $\hat{T}_{\mu\nu} = T_{\mu\nu} - T^{\rho}_{\rho} \gamma_{\mu\nu}$, (3.110) and (3.113) are further simplified to

$$\partial_{\lambda}\gamma_{\mu\nu} = 2\langle \hat{T}_{\mu\nu} \rangle,$$

$$\partial_{\lambda}\langle \hat{T}_{\mu\nu} \rangle = \langle \hat{T}_{\mu\rho} \rangle \langle \hat{T}^{\rho}_{\nu} \rangle.$$
 (3.114)

Leaving the details of the derivation to the original paper [21], the solution to these equations was found to be

$$\gamma_{\mu\nu}^{[\lambda]} = \gamma_{\mu\nu}^{[0]} + 2\lambda \langle \hat{T}_{\mu\nu} \rangle^{[0]} + \lambda^2 \langle \hat{T}_{\mu\rho} \rangle^{[0]} \gamma^{[0]\rho\sigma} \langle \hat{T}_{\sigma\nu} \rangle^{[0]},$$

$$\langle \hat{T}_{\mu\nu} \rangle^{[\lambda]} = \langle \hat{T}_{\mu\nu} \rangle^{[0]} + \lambda \langle \hat{T}_{\mu\rho} \rangle^{[0]} \gamma^{[0]\rho\sigma} \langle \hat{T}_{\sigma\nu} \rangle^{[0]}.$$
 (3.115)

It was further shown that $\sqrt{\gamma} \langle O_{T\bar{T}} \rangle$ is constant along the flow [21], so for the generating functional we simply have

$$I^{[\lambda]}[\gamma^{[\lambda]}] = I^{[0]}[\gamma^{[0]}] - \lambda \int d^2x \sqrt{\gamma} \langle O_{T\bar{T}} \rangle, \qquad (3.116)$$

where $\sqrt{\gamma} \langle O_{T\bar{T}} \rangle$ can be taken at any "time" along the flow between 0 and λ . In addition, if the seed field theory is conformal, the energy-momentum tensor is traceless up to the Weyl anomaly

$$\langle T^{\rho}_{\rho} \rangle^{[0]} = \frac{c}{24\pi} R[\gamma^{[0]}]$$
 (3.117)

then we can prove the trace relation follows from the flow equation [21]

$$\langle T^{\rho}_{\rho} \rangle^{[\lambda]} = \frac{c}{24\pi} R[\gamma^{[\lambda]}] + 2\lambda \langle O_{T\bar{T}} \rangle^{[\lambda]}.$$
(3.118)

(3.111), known as the "variational principle", was the starting point of the derivation in [21] (also used in [153] for the $J\bar{T}$ deformation). As the general formalism for multi-trace deformation in [154, 155], it encapsulates the effects of multi-trace deformation in a compact form, in analogy to the canonical transformation in the classical field theory. With sources J^A dual to other operators O_A turned on, we generally have

$$\partial_{\lambda} \left(\frac{1}{2}\sqrt{\gamma^{[\lambda]}} \langle T_{\mu\nu} \rangle^{[\lambda]} \delta \gamma^{[\lambda]\mu\nu} + \sum_{A} \sqrt{\gamma^{[\lambda]}} \langle O_A \rangle^{[\lambda]} \delta J^{A[\lambda]} \right) = -\delta \left(\sqrt{\gamma^{[\lambda]}} \langle O_{T\bar{T}} \rangle^{[\lambda]}_{\text{factorized}}\right).$$
(3.119)

Since the variation of the action on the right-hand side does not involve O_A or J^A , we have

$$J^{A[\lambda]} = J^{A[0]}.$$

$$\sqrt{\gamma^{[\lambda]}} \langle O_A \rangle^{[\lambda]} = \sqrt{\gamma^{[0]}} \langle O_A \rangle^{[0]}.$$
 (3.120)

We would like to remark that, even though the metric is deformed as the $T\bar{T}$ deformation goes in the formulation of multi-trace deformation given above, there is no physical change of the background metric of field theories under the $T\bar{T}$ deformation. The flow of the metric is just a mathematical way to describe the flow of the generating functional as a functional of its argument, the metric. In fact, for double-trace deformations, we get the variation of the generating functional with the "correct" sign if we fix the source. That is, for the $T\bar{T}$ deformation we have

$$\partial_{\lambda} I^{[\lambda]}[\gamma] = \int d^2 x \sqrt{\gamma} \langle O_{T\bar{T}} \rangle^{[\lambda]}$$
(3.121)

if we fix the metric. The "variational principle" and the flow equations (3.110)(3.109)(3.113)are solutions to the functional differential equation (3.121) by the method of characteristics, c.f. our discussion in 2.3. While this flow equation looks simpler, the formulation with varying metrics has its merits for many purposes, for example, $\sqrt{\gamma} \langle O_{T\bar{T}} \rangle$ is constant along the flow, and the flow of metric takes a compelling physical meaning in holography to be discussed in the next section.

The large $c T\bar{T}$ flow equations derived above enable us to compute the partition function and correlation functions of a $T\bar{T}$ -deformed field theory if these were known for the seed theory. For example, the sphere partition function of a $T\bar{T}$ -deformed CFT was considered in [132] and [20](generalizing to higher dimensions). Since the sphere is maximally symmetric, the one-point function of the energy-momentum tensor should be proportional to the metric

$$\langle T_{\mu\nu} \rangle = \alpha \gamma_{\mu\nu} \tag{3.122}$$

and from the trace relation (3.118), the constant of proportionality is determined to be

$$\alpha = \frac{\sqrt{1 + \frac{\lambda c}{6\pi L^2}} - 1}{2\lambda} \tag{3.123}$$

where L is the radius of the sphere. The partition function satisfies the differential equation

$$L\frac{d}{dL}\log Z = \int d^2x \sqrt{\gamma} \langle T_{\rho}^{\rho} \rangle = 4\pi L^2 \frac{\sqrt{1 + \frac{\lambda c}{6\pi L^2} - 1}}{\lambda}.$$
(3.124)

In [132], the boundary condition $\log Z|_{L=0} = 0$ was chosen, and the sphere partition function was found to be

$$\log Z = \frac{c}{3} \log \left(\sqrt{\frac{6\pi L^2}{\lambda c}} + \sqrt{1 + \frac{6\pi L^2}{\lambda c}} \right) + \frac{2\pi L^2}{\lambda} \left(\sqrt{1 + \frac{\lambda c}{6\pi L^2}} - 1 \right).$$
(3.125)

Taking the face value, this expression does not have a proper CFT limit as $\lambda \to 0^{21}$. This is not surprising since there is no justification of the boundary condition $\log Z|_{L=0} = 0$ for a CFT, and we don't expect it to hold for a slightly $T\bar{T}$ -deformed CFT either. A more serious problem is that (3.125) doesn't satisfy the flow equation (3.121). To find the partition function with a proper CFT limit, a more straightforward computation can be done using the $T\bar{T}$ flow equations derived in this section [156]. In particular, (3.116) is convenient for the computation. For the second term on the right-hand side of (3.116), we find $\int d^2x \sqrt{\gamma} \langle O_{T\bar{T}} \rangle = -4\pi L^2 \alpha^2 =$ $-\frac{\pi L^2}{\lambda^2} \left(2 + \frac{\lambda c}{6\pi L^2} - 2\sqrt{1 + \frac{\lambda c}{6\pi L^2}}\right)$. For the first term on the right-hand side, the partition function of a CFT on a sphere with radius L_0 is given by

$$\log Z(L_0) = \frac{c}{3} \log \frac{L_0}{\epsilon}$$
(3.126)

where ϵ is a UV cutoff. The radius of the "undeformed metric" L_0 can be determined from the metric flow equation in (3.115)

$$L_0 = L \frac{1 + \sqrt{1 + \frac{\lambda c}{6\pi L^2}}}{2}.$$
 (3.127)

²¹Some people take the point of view that the deformation parameter λ also plays the role of effective UV cutoff, somewhat justifying this expression for the partition function. We think the UV cutoff and the deformation parameter are two different quantities.

Summing up every term, we find

$$\log Z = \frac{c}{3} \log \frac{L}{\epsilon} + \frac{c}{3} \log \frac{1 + \sqrt{1 + \frac{\lambda c}{6\pi L^2}}}{2} + \frac{2\pi L^2}{\lambda} (\sqrt{1 + \frac{\lambda c}{6\pi L^2}} - 1) - \frac{c}{6}.$$
 (3.128)

This expression of the sphere partition function satisfies the differential equation (3.124) as well as the flow equation (3.121). Still, now it has a proper CFT limit $\log Z = \frac{c}{3} \log \frac{L}{\epsilon}$ as $\lambda \to 0$. In addition, as λ goes to $+\infty$, the partition function diverges as $\lambda^{\frac{1}{2}}$ instead of becoming trivial in (3.125). As far as we know, this expression was first given in [15], it was also discussed in [157].

From the sphere partition function, one can compute the entanglement entropy of two antipodal points [132]. Using the replica trick, the entanglement entropy can be computed by

$$S = \left(1 - n\frac{d}{dn}\right)\log Z_n\Big|_{n=1} \tag{3.129}$$

where $\log Z_n$ is the partition function of the CFT on a replica sphere with metric

$$ds^{2} = L^{2} \left(d\theta^{2} + n^{2} \sin^{2} \theta d\phi^{2} \right).$$
(3.130)

The differentiation with respect to n yields the energy-momentum tensor one-point function

$$n\frac{d}{dn}\log Z_n = -\int dV \langle T_{\phi}^{\phi} \rangle_n.$$
(3.131)

Taking the limit $n \to 1$, we have $\lim_{n\to 1} \langle T^{\phi}_{\phi} \rangle_n = \frac{1}{2} \langle T^{\mu}_{\mu} \rangle$, so we finally obtain

$$S = (1 - \frac{L}{2}\frac{d}{dL})\log Z.$$
 (3.132)

Using (3.125), the entanglement entropy was computed to be

$$S = \frac{c}{3} \log\left(\sqrt{\frac{6\pi L^2}{\lambda c}} + \sqrt{1 + \frac{6\pi L^2}{\lambda c}}\right)$$
(3.133)

in [132]. If we use (3.128) instead, the entanglement entropy is found to be

$$S = \frac{c}{3}\log\frac{L}{\epsilon} + \frac{c}{3}\log\frac{1 + \sqrt{1 + \frac{\lambda c}{6\pi L^2}}}{2} - \frac{c}{6}.$$
 (3.134)

4 $T\bar{T}$ deformation and holography

For a holographic CFT, it's natural to ask what the holographic dual of its $T\bar{T}$ deformation is. It was well-understood that multi-trace deformations correspond to changing boundary conditions in the dual AdS space [154, 158–160]. A sharp proposal was given in [12] that for positive $T\bar{T}$ deformation parameter λ , the holographic dual is a Dirichlet cutoff in the AdS space. Evidences for this holographic proposal and works in this holographic setting include signal propagation speed [12], energy levels in a finite volume [12], holographic energy-momentum tensor correlation functions [14, 24], correspondence between the trace relation and Gauss-Codazzi equations [14, 17], partition functions [20, 105, 107, 132] and holographic entanglement entropy [132, 133, 136–138, 161–165]. Most evidences hinge on the $T\bar{T}$ flow of the partition function and the energy-momentum tensor. The crucial insight into the cutoff AdS holography proposal was given in [21], where the large $c T\bar{T}$ flow equation of the partition function and the one point function of the energy-momentum tensor was derived, as we have reviewed in Section 3.6. An explicit solution to the flow equation was obtained, translating to a mixed boundary condition for the AdS gravity. Furthermore, when the deformation parameter λ is positive, the mixed boundary condition coincides with a Dirichlet boundary condition on a cutoff surface. The cutoff location, determined by λ , moves into the bulk as λ increases. For negative λ , a glue-on AdS picture was later proposed in [157]. The cutoff AdS or glue-on AdS picture only works in the pure gravity sector or on the field theory side when the physical quantities being considered only involve the partition function or the energy-momentum tensor. A bulk cutoff for all fields requires double-trace deformation involving all operators in the boundary field theory [14, 18, 166, 167]. The main body of this section, the derivation of the mixed boundary and reduction to the cutoff or glue-on picture, is given as subsection 4.1. We then discuss the holographic partition function and entanglement entropy in 4.2. The computation of holographic correlation functions of the energy-momentum tensor is reviewed in 4.3, in which the trace relation plays a central role. The asymptotic symmetries of the boundary gravitons that satisfy the mixed boundary condition are discussed in subsection 4.4, realizing on the gravity side the TT-deformed conformal symmetry we discussed in subsection 2.6. Some generalizations and further developments are briefly mentioned in the last section 4.5, including $T\bar{T}$ in higher dimensions, constructing dS holography with $T\bar{T}$, and $T\bar{T}$ as a quantum theory of boundary gravitons at a finite cutoff (holography beyond large c).

4.1 Mixed boundary condition, cutoff and glue-on

The AdS/CFT holography is a statement of equivalence of a gravity theory in an AdS space and a conformal field theory on the conformal boundary of the AdS space [168–170]. Near the conformal boundary, the metric of an asymptotic AdS space takes the form ²²

$$ds^{2} = \frac{dr^{2}}{r^{2}} + \frac{1}{r^{2}}g_{ij}(x,r)dx^{i}dx^{j}$$
(4.1)

in the Fefferman-Graham coordinates [171, 172], where r is the holographic radial coordinate, and the conformal boundary is located at r = 0. The holography is encoded in the Gubser-Polyakov-Klebanov-Witten relation, that the partition function of conformal field theory with operator sources is equal to the gravity partition function with prescribed boundary conditions for bulk fields

$$\langle e^{\int JO} \rangle_{CFT} = Z_{\rm G}[\phi_0] \tag{4.2}$$

with field theory operators corresponding to bulk fields and sources identified as boundary conditions $J \sim \phi_0$. In the semi-classical limit, the gravity on-shell action becomes the generating functional of correlation functions of the conformal field theory. With this holographic prescription, we find an operator O with dimension Δ corresponds to a bulk field ϕ with mass M, satisfying the relation

$$\Delta(\Delta - d) = M^2 \tag{4.3}$$

where d is the dimension of the boundary (and the bulk has dimension d + 1). The near boundary solution to the on-shell equation of a bulk field, which is a second-order differential equation, takes the form

$$\phi = \phi^{(0)}(r^{d-\Delta} + \ldots) + \phi^{(2\Delta - d)}(r^{\Delta} + \ldots).$$
(4.4)

The coefficient $\phi^{(0)}$ corresponds to the source ϕ_0 , and $\phi^{(2\Delta-d)}$ corresponds to the one-point function $\langle O \rangle$. Specializing on the bulk metric and its dual operator, the energy-momentum tensor, we have the Fefferman-Graham expansion for the bulk metric

$$g_{ij} = g_{ij}^{(0)} + \ldots + g_{ij}^{(d)} r^d + \ldots$$
(4.5)

in which $g^{(0)}$ is identified with the field theory background metric γ and $g^{(d)}$ corresponds to the one-point function of energy-momentum tensor. For AdS₃/CFT₂, the expansion truncates as [173]

$$g_{ij}(x,r) = g_{ij}^{(0)}(x) + g_{ij}^{(2)}(x)r^2 + g_{ij}^{(4)}(x)r^4$$
(4.6)

and the Einstein's equation is reduced to one equation that determines $g^{(4)}$ in terms of $g^{(0)}$ and $q^{(2)}$:

$$g_{ij}^{(4)} = \frac{1}{4} g_{ik}^{(2)} g^{(0)kl} g_{lj}^{(2)}.$$
(4.7)

 $^{^{22}}$ Throughout this section, we set the AdS radius l to 1 for simplicity. One can recover the AdS radius in formulae by dimensional analysis.

and another two equations

$$\nabla^{(0)i} g_{ij}^{(2)} = \nabla^{(0)}_j g^{(2)i}_{\ i}, \tag{4.8}$$

$$g^{(2)i}_{\ i} = -\frac{1}{2}R[g^{(0)}]. \tag{4.9}$$

We renormalize the gravity action by introducing a cutoff $r = r_c$ and adding the Gibbons-Hawking boundary term and a counter-term [174–176]

$$I = I_{\rm EH} + I_{\rm GH} + I_{\rm CT} = -\frac{1}{16\pi G} \int_{r \ge r_c} dV (R - 2\Lambda) - \frac{1}{8\pi G} \int_{r=r_c} d\sigma K + \frac{1}{8\pi G} \int_{r=r_c} d\sigma. \quad (4.10)$$

As shown in [176], this renormalized gravity action, when put on-shell, has logarithm divergence as $r_c \rightarrow 0$, which is related to the Weyl anomaly [177]. The Weyl anomaly counter-term was proposed in [176]

$$I_{\rm WCT} = -\frac{1}{16\pi G} \log r_c \int_{r=r_c} d\sigma \hat{R}$$
(4.11)

to subtract this logarithm divergence, where \hat{R} is the scalar curvature of the induced metric. It is, in fact, a topological term since

$$\int_{r=r_c} d\sigma \hat{R} = 4\pi \chi = 4\pi (2 - 2g)$$
(4.12)

with χ being the Euler characteristic of the cutoff surface and g the genus. As a topological term, it doesn't affect holographic correlation functions. Functionally differentiating the gravity on-shell action with respect to the induced metric on the cutoff surface h_{ij} yields the Brown-York tensor [178] modified by the counter term I_{CT}

$$T_{ij}^{BY} = \frac{2}{\sqrt{h}} \frac{\delta I_{\text{on-shell}}}{\delta h^{ij}} = -\frac{1}{8\pi G} (K_{ij} - Kh_{ij} + h_{ij}).$$
(4.13)

Since $\gamma_{ij} = g_{ij}^{(0)} = \lim_{r_c \to 0} h_{ij} r_c^2$, we find the one-point function of energy-momentum tensor

$$\langle T_{ij} \rangle = \lim_{r_c \to 0} T_{ij}^{BY} = \frac{1}{8\pi G} (g_{ij}^{(2)} - g^{(0)kl} g_{kl}^{(2)} g_{ij}^{(0)}).$$
(4.14)

The two equations (4.8,4.9) translate to the conservation of the energy-momentum tensor and the holographic Weyl anomaly

$$\nabla^i \langle T_{ij} \rangle = 0, \tag{4.15}$$

$$\langle T \rangle = \frac{R}{16\pi G}.\tag{4.16}$$

Now, we consider the $T\bar{T}$ deformation of a holographic conformal field theory. For the seed theory, we have the holographic dictionary

$$\gamma_{ij}^{[0]} = g_{ij}^{(0)},$$

$$\langle \hat{T}_{ij} \rangle^{[0]} = \frac{1}{8\pi G} g_{ij}^{(2)}$$
(4.17)

with the Brown-Henneaux relation [179]

$$c = \frac{3}{2G} \tag{4.18}$$

and \hat{T}_{ij} denoting the trace-reversed energy-momentum tensor introduced in 3.6. By the $T\bar{T}$ flow equation (3.115) we find

$$\gamma_{ij}^{[\lambda]} = g_{ij}^{(0)} + \frac{\lambda}{4\pi G} g_{ij}^{(2)} + (\frac{\lambda}{4\pi G})^2 \frac{1}{4} g_{ik}^{(2)} g^{(0)kl} g_{lj}^{(2)} = g_{ij}^{(0)} + \frac{\lambda}{4\pi G} g_{ij}^{(2)} + (\frac{\lambda}{4\pi G})^2 g_{ij}^{(4)}.$$
(4.19)

As expected, the field theory metric corresponds to a mixed boundary condition for gravity. Furthermore, if λ is positive, then the mixed boundary can be viewed as a Dirichlet boundary condition on the cutoff surface $r = r_c$

$$\gamma_{ij}^{[\lambda]} = r_c^2 h_{ij}|_{r=r_c} \tag{4.20}$$

with the cutoff location r_c determined by λ

$$r_c^2 = \frac{\lambda}{4\pi G} = \frac{\lambda c}{6\pi}.$$
(4.21)

The λc combination in this dictionary makes the large c limit (3.101) manifest. Furthermore, from (3.115), the deformed one-point function of the trace-reversed energy-momentum tensor is found to be

$$\langle \hat{T}_{ij} \rangle^{[\lambda]} = \frac{1}{8\pi G} (g_{ij}^{(2)} + 2\frac{\lambda}{4\pi G} g_{ij}^{(4)}) = \frac{1}{8\pi G} (h_{ij} - K_{ij})|_{r=r_c}$$
(4.22)

with r_c given by (4.21), and the one-point function of the energy-momentum tensor itself is equal to the modified Brown-York tensor on the cutoff surface

$$\langle T_{ij} \rangle^{[\lambda]} = -\frac{1}{8\pi G} (K_{ij} - Kh_{ij} + h_{ij})|_{r=r_c} = T_{ij}^{\mathrm{BY}}|_{r=r_c}.$$
 (4.23)

Furthermore, from the contracted Gauss and Codazzi equations in an Einstein space with negative curvature (for example, see Appendix D in [180])²³

$$K^2 - K_{ij}K^{ij} = \hat{R} + 2 \tag{4.24}$$

$$\hat{\nabla}^i K_{ij} - \hat{\nabla}_j K = 0 \tag{4.25}$$

and expressing the extrinsic curvature in terms of the Brown-York tensor

$$K_{ij} = -8\pi G(T_{ij}^{\rm BY} - T^{\rm BY}h_{ij}) + h_{ij}$$
(4.26)

we find the Brown-York tensor is conserved

$$\hat{\nabla}^i T^{\rm BY}_{ij} = 0 \tag{4.27}$$

²³Here a hat symbol denotes geometrical quantities pertaining to the intrinsic geometry on the cutoff surface.

and satisfies

$$T^{\rm BY} = 4\pi G \left(T^{\rm BY}_{ij} T^{\rm BY} i^{j} - T^{\rm BY}^2 \right) + \frac{1}{16\pi G} \hat{R}[h].$$
(4.28)

These two equations translate to the conservation of the field theory energy-momentum tensor and the trace relation (3.118) [14,17]. We see the cutoff AdS picture works both for the metric and the energy-momentum tensor one-point function. In fact, the quickest way to see how $T\bar{T}$ emerges from the cutoff is to consider the flow of the gravity on-shell action as the cutoff moves into the bulk. We find

$$r_c \partial_{r_c} I_{\text{on-shell}} = \frac{1}{2} \int_{r=r_c} d\sigma T^{\text{BY}^{ij}} \mathcal{L}_n g_{ij} = \int_{r=r_c} d\sigma T^{\text{BY}^{ij}} K_{ij}$$
(4.29)

where $n = -r\partial_r$ is the unit normal of the cutoff surface. Expressing the extrinsic curvature in terms of the modified Brown-York tensor (4.26) and using the trace relation (4.28), we find

$$r_c \partial_{r_c} I_{\text{on-shell}} = -8\pi G \int_{r=r_c} d\sigma O_{T\bar{T}}^{\text{BY}} + \frac{1}{16\pi G} \int_{r=r_c} d\sigma \hat{R}[h]$$
(4.30)

where $O_{T\bar{T}}^{\text{BY}}$ means the combination of the $T\bar{T}$ form of the Brown-York tensor. With the Weyl anomaly counter-term (4.11), we rewrite the equation as

$$r_c \partial_{r_c} I'_{\text{on-shell}} = -8\pi G \int_{r=r_c} d\sigma O^{\text{BY}}_{T\bar{T}}$$

$$\tag{4.31}$$

with

$$I' = I + I_{WCT}.$$
 (4.32)

This flow equation of the gravity on-shell action as the cutoff moving into the bulk agrees with the $T\bar{T}$ flow equation of the generating functional (3.109) given the relation (4.21) between the AdS cutoff and the $T\bar{T}$ deformation parameter. A version of the statement for a finite displacement of the cutoff was also given in [92]. We see that even if we omit the Weyl anomaly counter-term for holographic CFT to subtract the logarithm divergence, this term emerges as the theory is $T\bar{T}$ -deformed, which can be written as

$$I_{\rm WCT} = -\frac{1}{16\pi G} \log \frac{r_c}{\epsilon} \int_{r=r_c} d\sigma \hat{R}$$
(4.33)

where ϵ is the CFT UV cutoff. Inclusion of this term is essential to match the $T\bar{T}$ flow equation as discussed in [92, 156]. We can obtain a holographic sphere partition function that agrees with (3.128), which has a proper CFT limit and satisfies the differential equation from the $T\bar{T}$ flow. The Weyl anomaly counter-term also gives us a generalization of the Ryu-Takayagani formula [181] for the holographic entanglement entropy, as we will see in (4.2).

The cutoff AdS picture only works for positive deformation parameter λ , as seen from (4.21). A glue-on AdS picture was proposed in [157], that for negative deformation parameter,

the boundary where the field theory lives moves beyond r = 0 to an AdS glued on the original AdS space²⁴. More precisely, we can use the holographic radial coordinate $\rho = r^2$ in which the bulk metric takes the form

$$ds^{2} = \frac{d\rho^{2}}{4\rho^{2}} + \frac{1}{\rho}g_{ij}(x,\rho)dx^{i}dx^{j}$$
(4.34)

Now, we allow ρ to take values in the whole real line, with $\rho > 0$ corresponding to the original AdS space and $\rho < 0$ corresponding to the AdS space glued on. The relation between the cutoff and the $T\bar{T}$ deformation parameter now reads

$$\rho_c = \frac{\lambda}{4\pi G} = \frac{\lambda c}{6\pi} \tag{4.35}$$

so a negative λ corresponds to cutoff at a negative ρ_c . This glue-on AdS proposal can be understood as a geometric realization of analytic continuation of the cutoff AdS holography to the region $\lambda < 0$, and various physics quantities, including the energy levels, the trace relation, partition functions and the entanglement entropy, were studied in [157, 184] to support this picture.

One must keep in mind that the cutoff or glue-on picture only works for the pure gravity sector. In particular, (4.20) does not hold in the presence of matter. The solution to Einstein's equation is much more complicated with matter, and the deformed metric will have no simple geometric meaning. In [21], a bulk geometry with a matter shell, BTZ outside, and thermal AdS inside was considered. If the perceived cutoff surface by (4.21) was inside the shell, the Brown-York tensor on the cutoff surface cannot reproduce the energy level of the $T\bar{T}$ -deformed field theory. The universal deformation of energy levels of the field theory should come from the universal asymptotics of the bulk metric near the conformal boundary. Therefore, in holography, we can't remove the part of AdS space outside the perceived cutoff. Holographic correlation functions of scalar operators were also computed in [14] with the boundary put at a finite cutoff. The results do not agree with the field theory unless the double-trace deformation of scalar operators is turned on. All in all, as is known before [166, 167] and discussed in the context of $T\bar{T}$ deformation in [18], we need double-trace deformation involving all operators to move the conformal boundary into the bulk in holography.

4.2 Holographic partition function and entanglement entropy

In this subsection, we consider the holographic partition function and entanglement entropy, which directly follow the holography analysis in the previous subsection. First, we compute the holographic sphere partition function. Working in the "global patch" of AdS space

$$ds^2 = d\varrho^2 + \sinh^2 \varrho d\Omega_2^2 \tag{4.36}$$

 $^{^{24}}$ As a powerful tool, the glue-on picture has been applied to construct the multiple AdS spacetime [182] and traversable wormholes wormhole [183].

the radial Fefferman-Graham coordinate r, which must be proportional to $e^{-\varrho}$, needs to be chosen such that at the cutoff $r = r_c$, the induced metric corresponds to a field theory metric of a sphere with radius L. Let the cutoff in terms of ϱ coordinate be ϱ_c , we must have

$$r_c \sinh \varrho_c = L. \tag{4.37}$$

The gravity on-shell action, with the Weyl anomaly counter-term, is evaluated to be


$$I_{\text{on-shell}}^{'} = -\frac{1}{2G} e^{-\varrho_c} \sinh \varrho_c - \frac{1}{2G} \varrho_c - \frac{1}{2G} \log \frac{r_c}{\epsilon} = -\frac{1}{2G} \frac{L^2}{r_c^2} \left(\sqrt{1 + \frac{r_c^2}{L^2}} - 1 \right) - \frac{1}{2G} \log \left(1 + \sqrt{1 + \frac{r_c^2}{L^2}}\right) - \frac{1}{2G} \log \frac{L}{\epsilon}.$$
(4.38)

It agrees with the $T\bar{T}$ -deformed CFT sphere partition function (3.128) up to a constant by the parameter map (4.18) and (4.21). Without the Weyl anomaly counter-term, the on-shell action will agree with (3.126) [20,132]. The same can be said for the holographic entanglement entropy, (3.133) agrees with the Ryu-Takayanagi formula [181] with the geodesic endpoints on the cutoff surface [132], as expected given that the Ryu-Takayanagi formula can be derived from the Einstein's gravity [185,186]. With the Weyl anomaly counter-term, a modification to the Ryu-Takayanagi formula has to be made.

The modification turns out to be simple and intuitive. Suppose we consider the entanglement entropy of m intervals, the *n*-replica manifold \mathcal{M}_n has genus $g_n = (n-1)(m-1)$. By the replica trick (3.129), the contribution of the Weyl anomaly counter-term to the entanglement entropy is $\frac{1}{2G}m\log\frac{r_c}{\epsilon}$, that is, each endpoint contributes $\frac{1}{4G}\log\frac{r_c}{\epsilon}$, which is exactly $\frac{1}{4G}$ times the distance from the endpoint to the cutoff ϵ near the conformal boundary. Therefore, we have the modified Ryu-Takayanagi formula

$$S = \frac{A}{4G} + \frac{2m}{4G}\log\frac{r_c}{\epsilon} = \frac{A}{4G} + \frac{A'}{4G}.$$
 (4.39)

The first term $\frac{A}{4G}$ is the Ryu-Takayanagi formula adapted to the cutoff AdS picture advocated in [132] and many subsequent works, with A being the length of the Ryu-Takayanagi surface, namely the geodesics connecting the points of the entangling surface located on the cutoff surface in the bulk. In the correction term, 2m is the area of the entangling surface, namely the number of points, explicitly retaining the area law of UV-divergence of the entanglement entropy [187]. A' is the length of the lines connecting the entangling surface to the conformal boundary. This picture of extending the Ryu-Takayanagi surface from the cutoff surface to the conformal boundary supports the perspective that the AdS space outside the perceived cutoff must be retained, see Figure 1. In our case, the geometry of the extension of the Ryu-Takayanagi surface is simple since the cutoff surface is a surface of constant Fefferman-Graham radial coordinates, and all points in the entangling surface are equidistant from the conformal boundary. In general, we may consider a state represented by a cutoff surface that



Conformal boundary Perceived cutoff

Figure 1: Ryu-Takayanagi surface extended to the conformal boundary

is not necessarily at a constant radial cutoff, which can be prepared by an inhomogeneous $T\bar{T}$ deformation. The extension of the Ryu-Takayanagi surface will be "less trivial".

We can check our modified Ryu-Takayanagi formula (4.39) for the perturbative computation of entanglement entropy to the first order in the deformation parameter λ as was done in [136,138]. Consider an interval of length L in a thermal state with inverse temperature β . The dual of the thermal state in the high-temperature limit is the BTZ black hole

$$ds^{2} = (\varrho^{2} - \varrho^{2}_{+})dt^{2} + \frac{d\varrho^{2}}{\varrho^{2} - \varrho^{2}_{+}} + \varrho^{2}dx^{2} = \frac{dr^{2}}{r^{2}} + (\frac{1}{r} - \frac{\varrho^{2}_{+}r}{4})^{2}dt^{2} + (\frac{1}{r} + \frac{\varrho^{2}_{+}r}{4})^{2}dx^{2}$$
(4.40)

with $r = \frac{2}{\varrho_+^2} \left(\varrho - \sqrt{\varrho^2 - \varrho_+^2} \right)$ (or $\varrho = \frac{\varrho_+}{2} \left(\frac{\varrho_+ r}{2} + \left(\frac{\varrho_+ r}{2} \right)^{-1} \right) = \frac{1}{r} + \frac{\varrho_+^2 r}{4}$). The geodesic distance between two points in the Euclidean AdS space takes a particularly simple form in the coordinates of flat space $\mathbb{R}_{1,3}$ where the Euclidean AdS space is embedded as a hyperboloid (as an "analytic continuation" of the apparent formula for the sphere)

$$d(X_1, X_2) = \cosh^{-1}(-X_1 \cdot X_2). \tag{4.41}$$

The map from the usual coordinates of the BTZ black hole to the embedding coordinates is

$$X^{0} = \frac{\varrho}{\varrho_{+}} \cosh(\varrho_{+}x)$$

$$X^{1} = \frac{\varrho}{\varrho_{+}} \sinh(\varrho_{+}x)$$

$$X^{2} = \sqrt{\frac{\varrho^{2}}{\varrho_{+}^{2}} - 1} \cos(\varrho_{+}t)$$

$$X^{3} = \sqrt{\frac{\varrho^{2}}{\varrho_{+}^{2}} - 1} \sin(\varrho_{+}t).$$
(4.42)

Then the length of the Ryu-Takayanagi surface for a single interval with coordinate difference Δx at cutoff ρ_c is found to be

$$A_{\rm BTZ} = \cosh^{-1}\left(1 + 2\frac{\varrho_c^2}{\varrho_+^2}\sinh^2\frac{\varrho_+\Delta x}{2}\right) = \cosh^{-1}\left(1 + \frac{1}{2}\left(\frac{\varrho_+r_c}{2} + \frac{2}{\varrho_+r_c}\right)^2\sinh^2\frac{\varrho_+\Delta x}{2}\right).$$
 (4.43)

Plugging in the length of the interval in the field theory metric $L = (1 + \frac{\varrho_+^2 r_c^2}{4})\Delta x$, and the inverse temperature $\beta = \frac{2\pi}{\varrho_+} (1 - \frac{\varrho_+^2 r_c^2}{4})$, we find

$$A_{\rm BTZ} = \cosh^{-1} \left(\frac{(4\pi^2 r_c^2 + \beta^2) \cosh \frac{2\pi L}{\sqrt{4\pi^2 r_c^2 + \beta^2}} - \beta^2}{4\pi^2 r_c^2} \right).$$
(4.44)

As $r_c \to 0$ we find

$$A_{\rm BTZ} = 2\log\frac{\beta\sinh\frac{\pi L}{\beta}}{\pi r_c} - \frac{4\pi^2 r_c^2}{\beta^2}\frac{\pi L}{\beta}\coth\frac{\pi L}{\beta} + \frac{2\pi^2 r_c^2}{\beta^2}(\coth^2\frac{\pi L}{\beta} + 1) + o(r_c^2).$$
(4.45)

The $\log r_c$ is canceled exactly by the correction term in (4.39), we find

$$A_{\rm BTZ} + A'_{\rm BTZ} = 2\log\frac{\beta\sinh\frac{\pi L}{\beta}}{\pi\epsilon} - \frac{4\pi^2 r_c^2}{\beta^2}\frac{\pi L}{\beta}\coth\frac{\pi L}{\beta} + \frac{2\pi^2 r_c^2}{\beta^2}(\coth^2\frac{\pi L}{\beta} + 1) + o(r_c^2). \quad (4.46)$$

In the high-temperature limit, the leading $T\bar{T}$ deformation of the holographic entanglement entropy is

$$\delta S = -\frac{\pi^2 r_c^2}{G\beta^2} \frac{\pi L}{\beta} \coth \frac{\pi L}{\beta}.$$
(4.47)

It was found to agree with the field theory perturbative computation in [138]

$$\delta S = -\frac{\lambda \pi^2 c^2 L}{9\beta^3} \coth \frac{\pi L}{\beta}.$$
(4.48)

by the parameter map (4.18) and (4.21). The same result was obtained in [137] using the Chern-Simons formulation of three-dimensional gravity. We remark that without the correction term in the Ryu-Takayanagi formula, r_c plays a dual role of the UV cutoff and the $T\bar{T}$ deformation parameter in the original paper [138], it's thus ambiguous in which part of the expression (4.45) r_c should be identified as the UV cutoff, and in which part identified as the $T\bar{T}$ deformation parameter. Our standpoint is that the UV cutoff and the $T\bar{T}$ deformation parameter are different quantities. We can repeat this computation for the entanglement entropy of the vacuum state in a circle as in [138], but we can't find agreement between both sides of holography for reasons not yet clear.

4.3 Holographic correlation functions of the energy-momentum tensor

Energy-momentum tensor correlation functions were computed as early evidence for the proposed cutoff AdS holography for $T\bar{T}$ -deformed CFTs [14]. It has been further studied on the field theory side in [24,76,126] and on the gravity side of the holography in [24,27,28,31,188,189]. Since three-dimensional gravity has no propagating degrees of freedom, the holographic computation can be reduced to the conservation and the trace relation of the Brown-York tensor on a finite cutoff surface. The Brown-York tensor, with three independent components, is determined by the three equations from the conservation and the trace relation up to constants that, in principle, can be further determined by the global geometry, physical considerations, or some mathematical tricks. In the previous subsection, we have shown that the one-point function corresponds to the Brown-York tensor (4.23) and the conservation and trace relation of the one-point function correspond to those of the Brown-York tensor, respectively. As a result, the holographic energy-momentum tensor correlation functions agree with the large c field theory computation based on the trace relation [14, 24]. The holographic correlation functions agree of the conservation equation is preserved.

$$\nabla^i \langle T_{ij} \rangle = 0, \tag{4.49}$$

and the trace relation

$$\langle T \rangle = \frac{R}{16\pi G} + 4\pi G r_c^2 (\langle T^{ij} \rangle \langle T_{ij} \rangle - \langle T \rangle^2).$$
(4.50)

for a given metric variation. Consider a $T\overline{T}$ -deformed CFT on a compactified plane with the metric

$$ds^2 = dz d\bar{z} \tag{4.51}$$

and a variation of the metric $\epsilon f_{ij}dx^i dx^j$, where ϵ is an infinitesimal parameter. The one-point function can be solve as a series in $\epsilon \langle T_{ij} \rangle = \epsilon \langle T_{ij} \rangle_1 + \epsilon^2 \langle T_{ij} \rangle_2 + \dots$ from (4.49) and (4.50). For the first order, we find

$$\langle T_{z\bar{z}} \rangle_1 = \frac{1}{16\pi G} \left(\partial_{\bar{z}}^2 f_{zz} - 2\partial_z \partial_{\bar{z}} f_{z\bar{z}} + \partial_z^2 f_{\bar{z}\bar{z}} \right)$$
(4.52)

and

$$\partial_{\bar{z}} \langle T_{zz} \rangle_1 + \partial_z \langle T_{z\bar{z}} \rangle_1 = 0$$

$$\partial_z \langle T_{\bar{z}\bar{z}} \rangle_1 + \partial_{\bar{z}} \langle T_{z\bar{z}} \rangle_1 = 0.$$
 (4.53)

The Green's function on the compactified plane is $\frac{1}{z-w}$ $(\frac{1}{\bar{z}-\bar{w}})$, that is

$$\frac{1}{\pi}\partial_{\bar{z}}\frac{1}{z-w} = \delta^{(2)}(z-w) \text{ and c.c..}$$
(4.54)

Using the Green's function, we find

$$\langle T_{zz}(z) \rangle_1 = \frac{1}{16\pi G} \Big(-\partial_z \partial_{\bar{z}} f_{zz}(z) + 2\partial_z^2 f_{z\bar{z}}(z) + \frac{6}{\pi} \int d^2 w \frac{f_{\bar{z}\bar{z}}(w)}{(z-w)^4} \Big)$$
and c.c. (4.55)

where we have ruled out a possible constant by the regularity at the infinity point of the compactified plane. Therefore, we obtain the non-vanishing two-point functions for distinct points

$$\langle T_{zz}(z)T_{zz}(w)\rangle^c = \frac{3}{16\pi^2 G} \frac{1}{(z-w)^4} \text{ and c.c..}$$
 (4.56)

We see the two-point functions are unaffected by the cutoff (or the $T\bar{T}$ deformation on the field theory side). For the second order, we find

$$\langle T_{z\bar{z}} \rangle_2 = 8\pi G r_c^2 (\langle T_{zz} \rangle_1 \langle T_{\bar{z}\bar{z}} \rangle_1 - \langle T_{z\bar{z}} \rangle_1^2) - (f_{zz} \langle T_{\bar{z}\bar{z}} \rangle_1 + f_{\bar{z}\bar{z}} \langle T_{zz} \rangle_1)$$

+ local quadratic functions of f and its derivatives. (4.57)

Plugging in $\langle T_{ij} \rangle_1$ we have obtained, the only non-vanishing three-point function involving $T_{z\bar{z}}$ is found to be

$$\langle T_{z\bar{z}}(z)T_{zz}(w)T_{\bar{z}\bar{z}}(v)\rangle^c = \frac{9r_c^2}{32\pi^3 G}\frac{1}{(z-w)^4(\bar{z}-\bar{v})^4}.$$
 (4.58)

Furthermore, from the conservation equation to the second order, we find

$$\partial_{\bar{z}} \langle T_{zz} \rangle_{2} + \partial_{z} \langle T_{z\bar{z}} \rangle_{2} = 2 \langle T_{z\bar{z}} \rangle_{1} (\partial_{\bar{z}} f_{zz} + \partial_{z} f_{z\bar{z}}) + 3 \langle T_{zz} \rangle_{1} \partial_{z} f_{\bar{z}\bar{z}} + \langle T_{\bar{z}\bar{z}} \rangle_{1} \partial_{z} f_{zz} + f_{z\bar{z}} (\partial_{\bar{z}} \langle T_{zz} \rangle_{1} + \partial_{z} \langle T_{z\bar{z}} \rangle_{1}) + 2 f_{zz} \partial_{\bar{z}} \langle T_{z\bar{z}} \rangle_{1} + 2 f_{\bar{z}\bar{z}} \partial_{z} \langle T_{zz} \rangle_{1} \text{ and c.c..}$$

$$(4.59)$$

Other non-vanishing three-point functions are found to be

$$\langle T_{zz}(z)T_{zz}(w)T_{zz}(v)\rangle^{c} = -\frac{3}{16\pi^{3}G}\frac{1}{(z-w)^{2}(w-v)^{2}(v-z)^{2}} \text{ and c.c.,} \langle T_{zz}(z)T_{\bar{z}\bar{z}}(w)T_{\bar{z}\bar{z}}(v)\rangle^{c} = \frac{3r_{c}^{2}}{8\pi^{3}G}(\frac{1}{(z-w)^{3}(\bar{w}-\bar{v})^{5}} + \frac{1}{(z-v)^{3}(\bar{v}-\bar{w})^{5}}) \text{ and c.c..}$$
(4.60)

Following this spirit, one can proceed to compute higher-point functions [24] or holographic correlation functions on other boundary spacetimes (with nontrivial topology), for example, the sphere [24], torus [27, 188], higher-genus surfaces [31], and non-orientable manifolds [189].

4.4 Symmetries of boundary gravitons at a finite cutoff

An important aspect of the AdS_3/CFT_2 correspondence is the symmetry, the Virasoro algebra emerges as the algebra of the boundary charges of the boundary gravitons of AdS_3 gravity [179]. It's thus a natural and important question how the symmetry evolves as the $T\bar{T}$ deformation is turned on. This question has been studied on the CFT side in [50, 98, 100, 101], and on the gravity side in [21, 190, 191]. Restricting to pure gravity, now the boundary gravitons are on a finite cutoff surface (or equivalently with mixed boundary conditions on the conformal boundary). We will study the symmetries, boundary charges, and the algebra of the boundary gravitons in this subsection, mainly following [190]. To this end, it's helpful first to recapitulate the canonical formalism for gravity.

Hamilton's equation and Darboux's theorem inspire the symplectic-geometrical description of classical mechanics [192, 193]. The phase space is a 2n-dimensional space with a symplectic form Ω , a non-degenerate closed 2-form. A symmetry is represented by a symplectomorphism, infinitesimally given by a vector field V that preserves the symplectic form

$$\mathcal{L}_V \Omega = 0. \tag{4.61}$$

By the Cartan formula $\mathcal{L}_V = di_V + i_V d$ and the Poincare lemma, the vector is locally generated by a "Hamiltonian" F

$$i_V \Omega = -dF \tag{4.62}$$

or in components

$$V^i = \Omega^{ij} \partial_j F. \tag{4.63}$$

Conversely, Hamiltonians generate symplectomorphisms. The Poisson bracket is defined as

$$\{F,G\} = \Omega^{ij}\partial_i F \partial_j G = -V_F(G) = V_G(F)$$
(4.64)

and the equation of motion

$$\frac{dF}{dt} = \{F, H\} = V_H(F) \tag{4.65}$$

is geometrically a flow along the vector field generated by the Hamiltonian H. The canonical formalism can be generalized to field theories in a covariant way [194]. In this case, the phase space is defined as the space of all on-shell solutions ϕ . A typical function is $\phi(x)$, the value of the on-shell solution ϕ at the point x, or more generally a functional $F[\phi]$. A tangent vector is defined as an infinitesimal variation of an on-shell solution $\delta\phi$ (in fact, a solution to the linearized equation); it acts on the functions naturally by computing the variation of the function by this variation of the on-shell solution. One forms are linear functionals on the tangent space, a typical one is $\delta\phi(x)$, namely the value of $\delta\phi$ at x. A k-form can be represented as a wedge product of 1-forms

$$A = \int dx_1 \dots dx_k a_{x_1,\dots,x_k}[\phi] \delta\phi(x_1) \dots \delta\phi(x_k).$$
(4.66)

The exterior derivative is defined by its action on a function

$$\delta F[\phi] = \int dx \frac{\delta F}{\delta \phi(x)} \delta \phi(x). \tag{4.67}$$

and extended to arbitrary k-forms by the usual algebraic properties of an exterior derivative. The symplectic form of the gravity theory was given in [194]

$$\Omega = i \int d\Sigma_i \sqrt{g} J^i \tag{4.68}$$

where the symplectic current is

$$J^{i} = \frac{1}{16\pi G} \Big[\delta\Gamma^{i}_{jk} (\delta g^{jk} + \frac{1}{2} g^{jk} \delta \log g) - \delta\Gamma^{j}_{kj} (\delta g^{ik} + \frac{1}{2} g^{ik} \delta \log g) \Big].$$
(4.69)

It can be shown that for a diffeomorphism ξ , the charge

$$Q[\xi] = i \int_{\partial \Sigma} d\sigma T_{ij} n^i \xi^j \tag{4.70}$$

generates the diffeomorphism in the sense that

$$i_{\delta_{\xi}\phi}\Omega = -\delta Q[\xi] \tag{4.71}$$

where $\delta_{\xi}\phi$ is a tangent vector, the variation of an on-shell solution by the diffeomorphism ξ .

Now we consider a boundary at a finite cutoff ρ_c with a cylinder topology and the metric

$$ds^2 = dt^2 + d\phi^2 = dw d\bar{w}.$$
(4.72)

The general Banados geometry that satisfies this boundary condition is

$$ds^{2} = \frac{d\rho^{2}}{4\rho^{2}} + \frac{\left[(1 - \rho\rho_{c}\mathcal{L}\bar{\mathcal{L}})dw + (\rho - \rho_{c})\bar{\mathcal{L}}d\bar{w}\right]\left[(1 - \rho\rho_{c}\mathcal{L}\bar{\mathcal{L}})d\bar{w} + (\rho - \rho_{c})\mathcal{L}dw\right]}{\rho(1 - \rho_{c}^{2}\mathcal{L}\bar{\mathcal{L}})}$$
(4.73)

with $\mathcal{L}, \bar{\mathcal{L}}$ satisfying

$$\partial_{\bar{w}}\mathcal{L} + \rho_c \bar{\mathcal{L}} \partial_w \mathcal{L} = 0, \; \partial_w \bar{\mathcal{L}} + \rho_c \mathcal{L} \partial_{\bar{w}} \bar{\mathcal{L}} = 0. \tag{4.74}$$

Symmetries are given by the boundary-preserving diffeomorphisms, which, to the order that can be used to compute the boundary charges, are found to be

$$\xi^{w} = f^{w} - \frac{1}{2} \partial_{w} f^{\rho} + O((\rho - \rho_{c})^{2})$$

$$\xi^{\bar{w}} = f^{\bar{w}} - \frac{1}{2} \partial_{\bar{w}} f^{\rho} + O((\rho - \rho_{c})^{2})$$

$$\xi^{\rho} = \rho f^{\rho}$$
(4.75)

with

$$f^{\rho} = \frac{1 - \rho_c^2 \mathcal{L}\bar{\mathcal{L}}}{1 + \rho_c^2 \mathcal{L}\bar{\mathcal{L}}} (\partial_w f^w + \partial_{\bar{w}} f^{\bar{w}})$$
(4.76)

These diffeomorphisms depend on \mathcal{L} and $\overline{\mathcal{L}}$. In other words, the symmetries are "statedependent" as compared to the case without $T\overline{T}$ -deformation. On a time slice, we decompose the diffeomorphism in Fourier modes

$$\xi_n = e^{-in\phi} \partial_w, \ \bar{\xi}_n = -e^{-in\phi} \partial_{\bar{w}} \tag{4.77}$$

and we define

$$Q_n = Q[\xi_n] = \frac{1}{8\pi G} \int d\phi \frac{(1 - \rho_c \bar{\mathcal{L}})\mathcal{L}}{1 - \rho_c^2 \mathcal{L}\bar{\mathcal{L}}} e^{-in\phi}, \ \bar{Q}_n = Q[\bar{\xi}_n] = \frac{1}{8\pi G} \int d\phi \frac{(1 - \rho_c \mathcal{L})\bar{\mathcal{L}}}{1 - \rho_c^2 \mathcal{L}\bar{\mathcal{L}}} e^{in\phi}.$$
 (4.78)

From (4.71) we have the relation

$$\{Q[\xi], Q[\chi]\} = -\delta_{\xi}Q[\chi] = \delta_{\chi}Q[\xi]$$
(4.79)

and using this relation, we can find the charge algebra [190]

$$i\{Q_m,Q_n\} = \frac{1}{8\pi G} \int_0^{2\pi} d\phi e^{-i(m+n)\phi} \Big[\frac{1}{2}m^3 + (m-n)\frac{(1-\rho_c q)q}{1-\rho_c(q+\bar{q})} - \frac{1}{2}mn(m-n)\rho_c \frac{\bar{q} - \frac{1}{2}\rho_c(q+\bar{q})^2}{(1-\rho_c(q+\bar{q}))^2} \Big]$$

and c.c.

$$i\{Q_m, \bar{Q}_n\} = \frac{1}{8\pi G} \int_0^{2\pi} d\phi e^{-i(m-n)\phi} \Big[\frac{\rho_c(m-n)q\bar{q}}{1-\rho_c(q+\bar{q})} + \frac{1}{2}\rho_c mn \frac{nq - m\bar{q} + \frac{\rho_c}{2}(m-n)(q+\bar{q})^2}{(1-\rho_c(q+\bar{q}))^2} \Big]$$
(4.80)

where

$$q = 4G \sum_{n} Q_{n} e^{-in\phi}, \ \bar{q} = 4G \sum_{n} \bar{Q}_{n} e^{in\phi}.$$
 (4.81)

This is a nonlinear generalization of the Virasoro algebra that reduces to it when $\rho_c = 0$.

4.5 Further developments on holography

There are many generalizations and further developments of the relation between $T\bar{T}$ deformation and holography, which we can't discuss in detail in this review. We mention some of them in this last subsection and refer the readers to the original paper.

Quantum boundary gravitons: holography beyond large c

Following the symmetry analysis, boundary gravitons at a finite cutoff were further studied in [190, 195, 196], focusing on quantization which can be interpreted as holography for $T\bar{T}$ deformed CFTs beyond large c. The theory of boundary gravitons on the conformal boundary is essentially a constrained free scalar. By $T\bar{T}$ deformation, it's turned into a Nambu-Goto action in the static gauge. One can quantize in the light-cone gauge, with some ambiguities settled by the trace relation. The energy levels [1] and energy-momentum tensor correlation functions [14] are reproduced to the first few orders of $\frac{1}{c}$ expansion.

$T\bar{T} + \Lambda^2$ and dS holography

The picture of a flow with a moving cutoff can be generalized to space with a positive cosmological constant, particularly for dS holography. For dS space, the contracted Gauss equation is

$$K^2 - K_{ij}K^{ij} = \hat{R} - 2 \tag{4.82}$$

while the Codazzi equation is unchanged. The trace relation of the modified Brown-York tensor now has an extra term, and as a consequence, the flow equation of the on-shell action as the cutoff moves also takes an extra term

$$r_c \partial_{r_c} I_{\text{on-shell}} = -8\pi G \int_{r=r_c} d\sigma O_{T\bar{T}}^{\text{BY}} + \frac{1}{16\pi G} \int_{r=r_c} d\sigma \hat{R}[h] - \frac{1}{4\pi G} \int_{r=r_c} d\sigma.$$
(4.83)

Keeping the same holographic dictionary, this translates to the flow equation of the field theory generating functional

$$\partial_{\lambda} I^{[\lambda]}[\gamma^{[\lambda]}] = -\int d^2x \sqrt{\gamma^{[\lambda]}} \langle O_{T\bar{T}} \rangle^{[\lambda]} - \frac{1}{2\lambda^2} \int d^2x \sqrt{\gamma^{[\lambda]}}.$$
(4.84)

With the extra term on the right-hand side, a λ -dependent two-dimensional cosmological constant term, the deformation was introduced by the name $T\bar{T} + \Lambda_2$ in [15] and subsequently studied in [163, 197–201].

Higher dimensional $T\bar{T}$ deformation from cutoff AdS

There have been proposals for generalizing $T\bar{T}$ deformation to higher dimensions from the cutoff AdS holography. They are based on the trace relation or the flow of the on-shell action [17,18]. In general, the Brown-York tensor modified by counter-terms takes the form

$$T_{ij}^{\rm BY} = -\frac{1}{8\pi G} (K_{ij} - Kh_{ij} + C_{ij})$$
(4.85)

where C_{ij} is the contribution from the local counter-term [176]. The contracted Gauss equation (4.24) in an Einstein space with negative curvature in general dimensions is

$$K^{2} - K_{ij}K^{ij} = \hat{R} + d(d-1).$$
(4.86)

In terms of the Brown-York tensor, it reads

$$(8\pi G)^{2} (T_{ij}^{\rm BY} T^{\rm BY^{ij}} - \frac{1}{d-1} T^{\rm BY^{2}}) + (C_{ij} C^{ij} - \frac{1}{d-1} C^{2}) + 16\pi G (T_{ij}^{\rm BY} C^{ij} - \frac{1}{d-1} T^{\rm BY} C) + \hat{R}[h] + d(d-1) = 0.$$

$$(4.87)$$

Using this relation, the flow equation for the on-shell action can be written as

$$r_{c}\partial_{r_{c}}I_{\text{on-shell}} = \frac{1}{2} \int_{r=r_{c}} d\sigma T^{\text{BY}^{ij}} \mathcal{L}_{n}g_{ij} = \int_{r=r_{c}} d\sigma T^{\text{BY}^{ij}}K_{ij}$$

$$= \int_{r=r_{c}} d\sigma \left[(-8\pi G)(T_{ij}^{\text{BY}}T^{\text{BY}^{ij}} - \frac{1}{d-1}T^{\text{BY}^{2}}) - (T_{ij}^{\text{BY}}C^{ij} - \frac{1}{d-1}T^{\text{BY}}C) \right]$$

$$= \int_{r=r_{c}} d\sigma \left[(-4\pi G)(T_{ij}^{\text{BY}}T^{\text{BY}^{ij}} - \frac{1}{d-1}T^{\text{BY}^{2}}) - \frac{1}{16\pi G}(C_{ij}C^{ij} - \frac{1}{d-1}C^{2} + \hat{R}[h] + d(d-1)) \right].$$

$$(4.88)$$

Now we can see the natural higher dimensional $T\bar{T}$ operator inspired by the cutoff AdS holography is

$$O_{T\bar{T}} = \frac{1}{2} (T_{ij} T^{ij} - \frac{1}{d-1} T^2)$$
(4.89)

Taking the leading counter-term to be $(d-1)h_{ij}$ as in [176] and using the dictionary

$$\langle T_{ij} \rangle = r_c^{2-d} T_{ij}^{\text{BY}}, \ \gamma_{ij} = r_c^2 h_{ij}|_{r=r_c},$$
(4.90)

we can map the Gauss equation in terms of the Brown-York tensor to the trace relation and the flow of the gravity on-shell action to the flow of the generating functional, up to contribution from counter-terms, with the relation between the deformation parameter and the cutoff

$$\lambda = \frac{8\pi G r_c^d}{d}.\tag{4.91}$$

5 $T\bar{T}$ and string theory

This section will briefly introduce the application of $T\bar{T}$ deformation in string holography, i.e., single trace $T\bar{T}$ deformation. In generic translational and rotational invariant CFT, one can define the $T\bar{T}$ deformation. In addition, we can define another deformation also induced by the bilinear form of energy-momentum tensors at the same time if the CFT is a symmetric product orbifold \mathcal{M}^p/S_p , where \mathcal{M} represents the seed CFT and S_p is permutation group. The deformed theory remains an orbifold with a different seed theory $(\mathcal{M}_{\lambda})^p/S_p$. \mathcal{M}^i_{λ} is $T\bar{T}$ deformation of \mathcal{M}^i induced by $T^i_{\lambda}\bar{T}^i_{\lambda}$. Here *i* is *i*th copy of the orbifold. Therefore, the deformation of \mathcal{M}^p/S_p is triggered by

$$\sum_{i=1}^{p} T_{\lambda}^{i} \bar{T}_{\lambda}^{i}.$$
(5.1)

This is the so-called single-trace $T\bar{T}$ deformation of \mathcal{M}^p/S_p , in contrast with the usual $T\bar{T}$ deformation (also called double-trace deformation). The latter is induced by

$$\sum_{i=1}^{p} T_{\lambda}^{i} \sum_{j=1}^{p} \bar{T}_{\lambda}^{j}.$$
(5.2)

The entropy for single-trace $T\bar{T}$ deformed orbifold CFT \mathcal{M}^p/S_p can be obtained from (3.17), which turns out to be [13]

$$S(\mathcal{M}) = pS_{\mathcal{M}}(E/p), \tag{5.3}$$

where the total energy and entropy is a sum of each copy $E = \sum_{i} E_{i}, S(\mathcal{M}) = \sum_{i} S_{\mathcal{M}}(E_{i}), S_{\mathcal{M}}$ is the entropy of seed theory, taking the form (3.17). (5.3) also exhibit Hagedorn behavior for negative λ in (3.17).

We are interested in single-trace deformation because the above orbifold is supposed to be holographically dual to long string sector of type IIB string theory in AdS_3 . Thus, it is natural to ask about the behavior of this kind of holographical duality when turning on the single-trace deformation.

For string on $AdS_3 \times N_7$, a single long string, which is a string configuration that can wrap around the conformal boundary of AdS_3 , is described by a sigma model on [202], see also [34,203], and recently [204]

$$\mathcal{M}_{6k}^{(L)} = R_{\phi} \times \mathcal{N}_7 \tag{5.4}$$

with central charge $c_{\mathcal{M}} = 6k$, and k is related to AdS radius. The theory on R_{ϕ} has a linear dilaton, with slope

$$Q = (k-1)\sqrt{\frac{2}{k}}.$$
 (5.5)

The long string coupling

 $g^L \sim e^{Q\phi}.\tag{5.6}$

Thus, the string is strongly coupled at the boundary $\phi \to \infty$ (ϕ can be considered a radial direction). The complete boundary theory dual to string on AdS₃ is unknown for generic k. For p long strings, it is described by the symmetric orbifold

$$\left(\mathcal{M}_{6k}^{(L)}\right)^p / S_p. \tag{5.7}$$

String on AdS₃

String propagate on AdS_3 with NS-NS fields have a tractable world-sheet description of the WZW model. The AdS_3 metric is ²⁵

$$ds^2 = l^2 (d\phi^2 + e^{2\phi} d\gamma d\bar{\gamma}) \tag{5.8}$$

and NS-NS ${\cal B}$ field

$$B = -\frac{1}{2}l^2 e^{2\phi} d\gamma \wedge d\bar{\gamma}.$$
(5.9)

The corresponding WZW model is based on group SL(2, R) with element g parametrized as

$$g = \begin{pmatrix} e^{\phi} & 0\\ 0 & e^{-\phi} \end{pmatrix} \begin{pmatrix} 1 & 0\\ \gamma & 1 \end{pmatrix} \begin{pmatrix} 1 & \bar{\gamma}\\ 0 & 1 \end{pmatrix} = \begin{pmatrix} e^{\phi} & \bar{\gamma}e^{\phi}\\ \gamma e^{\phi} & e^{-\phi} + \gamma \bar{\gamma}e^{\phi} \end{pmatrix}.$$
 (5.10)

The world-sheet action is written as

$$\mathcal{L} = \frac{k}{\pi} \int d^2 z (\partial \phi \bar{\partial} \phi + e^{2\phi} \bar{\partial} \gamma \partial \bar{\gamma}), \quad k = \frac{l^2}{\alpha'}.$$
(5.11)

which is SL(2,R) WZW model at level l^2/α' . (5.11) can be equivalently expressed in Wakamoto representation

$$\mathcal{L} = \frac{2l^2}{l_s^2} \left(\partial \phi \bar{\partial} \phi + e^{2\phi} \bar{\partial} \gamma \partial \bar{\gamma} \right) = \frac{2l^2}{l_s^2} \left(\partial \phi \bar{\partial} \phi + \beta \bar{\partial} \gamma + \bar{\beta} \partial \bar{\gamma} - e^{-2\phi} \beta \bar{\beta} \right), \tag{5.12}$$

where action is described by a free field ϕ and a holomorphic bosonic β , γ system with conformal dimension

$$h(\beta) = 1, h(\gamma) = 0,$$
 (5.13)

and two-point function

$$\langle \phi(z)\phi(0)\rangle = -\log|z|^2, \langle \beta(z)\gamma(0)\rangle = 1/z.$$
(5.14)

The SL(2,R) current

$$J^{3} = \beta \gamma + \frac{\alpha_{+}}{2} \partial \phi,$$

$$J^{+} = \beta \gamma^{2} + \alpha_{+} \gamma \partial \phi + k \partial \gamma,$$

$$J^{-} = \beta,$$

(5.15)

²⁵Euclidean AdS₃: H_3^+ =SL(2,C)/SU(2).

whose modes expansion $J^i(z) = \sum_{n=-\infty}^{\infty} z^{-n-1} J_n^i$ obey the SL(2,R) current algebra

$$\begin{split} [J_n^3, J_m^3] &= -\frac{1}{2} kn \delta(n+m), \\ [J_n^3, J_m^{\pm}] &= \pm J_{n+m}^{\pm}, \\ [J_n^+, J_m^-] &= kn \delta(n+m) - 2J_{m+n}^3. \end{split}$$
(5.16)

The representation of the current algebra can be found in [205], and the long string sector corresponds to vertex operators belonging to the continuous representation of the SL(2, R).

Taking quantum effect, the renormalized Lagrangian is

$$\mathcal{L}_{\rm ws} = \partial \phi \bar{\partial} \phi - \frac{2}{\alpha_+} \widehat{R} \phi + \beta \bar{\partial} \gamma + \bar{\beta} \partial \bar{\gamma} - \beta \bar{\beta} \exp\left(-\frac{2}{\alpha_+}\phi\right), \qquad (5.17)$$

To figure out which operator in the world-sheet theory duals to single-trace $T\bar{T}$ operator in the spacetime CFT, we need to investigate the vertex operator in the world-sheet WZW model. Two relevant kinds of vertex operators will be relevant: the current operator and the energy-momentum tensor.

Current operator

From the current (5.15), one introduces $[13]^{-26}$

$$J^{+}(x;z) = e^{-xJ_{0}^{-}}J^{+}(z)e^{xJ_{0}^{-}} = J^{+}(z) - 2xJ^{3}(z) + x^{2}J^{-}(z),$$

$$J^{3}(x;z) = e^{-xJ_{0}^{-}}J^{3}(z)e^{xJ_{0}^{-}} = J^{3}(z) - xJ^{-}(z) = -\frac{1}{2}\partial_{x}J^{+}(x;z),$$

$$J^{-}(x;z) = e^{-xJ_{0}^{-}}J^{-}(z)e^{xJ_{0}^{-}} = J^{-}(z) = \frac{1}{2}\partial_{x}^{2}J^{+}(x;z),$$
(5.19)

where z is world-sheet coordinates x can be regarded as spacetime coordinates where the dual CFT lives. It is convenient to compact the currents together as follows

$$J(x;z) \equiv -J^{+}(x;z) = 2xJ^{3}(z) - J^{+}(z) - x^{2}J^{-}(z), \qquad (5.20)$$

whose zero mode (on the worldsheet) is

$$J_0(x) = \int dz J(x, z).$$
 (5.21)

Note $J_0(x)$ is not related to J_0^{\pm} , but to J(x, z).

Another useful operator is

$$\Phi_h(x;z) = \frac{1}{\pi} \left(\frac{1}{|\gamma - x|^2 e^{\phi} + e^{-\phi}} \right)^{2h},$$
(5.22)

 26 For example, from the algebra (5.16), one can obtain [206]

$$e^{-xJ_0^-}J^+(z)e^{xJ_0^-}$$

= $J^+(z) + [-xJ_0^-, J^+(z)] + \frac{1}{2}[-xJ_0^-, [-xJ_0^-, J^+(z)]] + \cdots$ (5.18)
= $J^+(z) - 2xJ^3(z) + x^2J^-(z).$

which is an eigenfunction of Laplacian on AdS_3 . $\Phi_h(x; z)$ can be viewed as a primary of spacetime with dimension (h, h), and also primary of world-sheet with dimension (-h(h - 1)/(k - 2)). The transformation generated by the current for $\Phi_h(x; z)$ is

$$[J_{0}(x), \Phi_{h}(y \cdot \bar{y})] = \int dz \left[J(x, z), \Phi_{h}(y, \bar{y})\right]$$

$$= \int dz \left(2x \left[J^{3}(z), \Phi_{h}\right] - \left[J^{+}(z), \Phi_{h}\right] - x^{2} \left[J(z), \Phi_{h}\right]\right)$$

$$= 2x \left[J^{3}_{0}, \Phi_{h}\right] - \left[J^{+}_{0}, \Phi_{h}\right] - x^{2} \left[J^{-}_{0}, \Phi_{h}\right]$$

$$= 2x(-1) \left(y\partial_{y} + h\right) \Phi_{h} - (-1) \left(y^{2}\partial_{y} + 2hy\right) \Phi_{h} - x^{2}(-1)\partial_{y}\Phi_{h}$$

$$= (x - y)^{2}\partial_{y}\Phi_{h} + 2(y - x)h\Phi_{h}.$$

(5.23)

Spacetime energy-momentum tensor

From Sugawara construction, the world-sheet energy-momentum tensor can be expressed as 27

$$T^{\rm ws} = \frac{1}{2(k-2)} \left[J \partial_x^2 J - \frac{1}{2} \left(\partial_x J \right)^2 \right].$$
 (5.25)

Here J(x; z) has spacetime dimension (-1,0), x has spacetime dimension (-1,0), thus T^{ws} has spacetime dimension (0,0). $\Phi_{h=1}$ as spacetime dimension $(h,\bar{h}) = (1,1)$.

Thus, to construct (2,0) spacetime tensor, which is the right dimension for spacetime energymomentum tensor, one needs to combine $(\partial_x^2, J, \bar{J}, \Phi_1)$, taking BRST condition into account, it turns out that the spacetime stress tension could be constructed as

$$T(x) = \frac{1}{2k} \int d^2 z \left[J(x;z) \partial_x^2 \Phi_1 + 3 \partial_x J \partial_x \Phi_1 + 3 \partial_x^2 J \Phi_1 \right] \bar{J}(\bar{x};\bar{z})$$

$$= \frac{1}{2k} \int d^2 z \left(\partial_x J \partial_x \Phi_1 + 2 \partial_x^2 J \Phi_1 \right) \bar{J}(\bar{x};\bar{z}).$$
 (5.26)

In parallel, one can also obtain a similar expression for $\overline{T}(x)$. By taking the product of T and \overline{T} , one can obtain the $T\overline{T}$ operator with spacetime dimension (2,2) that triggers double-trace $T\overline{T}$ deformation. What interests us here is the operator corresponds to the single-trace $T\overline{T}$ operator, which turns out to be the operator of spacetime dimension (2,2), the vertex operator of the massive dilaton [13]

$$D(x) = \int d^2 z \left(\partial_x J \partial_x + 2 \partial_x^2 J \right) \left(\partial_{\bar{x}} \bar{J} \partial_{\bar{x}} + 2 \partial_{\bar{x}}^2 \bar{J} \right) \Phi_1.$$
(5.27)

This is a quasi-primary operator. This operator is supposed to trigger single-trace $T\bar{T}$ deformation in the spacetime theory

$$D(x) = A \sum_{i=1}^{p} T_i(x) \bar{T}_i(\bar{x}).$$
(5.28)

²⁷From (5.20), for J(x; z)

$$J\partial_x^2 J - \frac{1}{2}(\partial_x J)^2 = -2JJ^- - \frac{1}{2}\left(2J^3 - 2xJ^-\right)^2 = 2J^+J^- - 2(J^3)^2.$$
(5.24)

Integrating over spacetime, one has ²⁸

$$\int d^2x D(x,\bar{x}) = \int d^2x \int d^2z \partial_x^2 J \partial_{\bar{x}}^2 \bar{J} \Phi_1 = 4 \int d^2x \int d^2z J^-(z) \bar{J}^-(\bar{z}) \Phi_1(x;z)$$

$$\propto \int d^2z J^-(z) \bar{J}^-(\bar{z}),$$
(5.30)

which implies that the single-trace $T\bar{T}$ deformation corresponds to current-current $J^-\bar{J}^-$ deformation in the world-sheet, while the latter is a marginal deformation on the world-sheet classically.

Deformation of world-sheet WZW model

In the Wakimoto representation (5.17), the Lagrangian deformed by (5.30) can be expressed as $-\lambda \int d^2z J^-(z) \bar{J}^-(\bar{z}) = -\lambda \int d^2z \beta(z) \bar{\beta}(\bar{z})$. Thus the deformed world-sheet Lagrangian is

$$\mathcal{L}_{\rm ws} = \partial \phi \bar{\partial} \phi - \frac{2}{\alpha_+} \widehat{R} \phi + \beta \bar{\partial} \gamma + \bar{\beta} \partial \bar{\gamma} - \beta \bar{\beta} \left[\lambda + \exp\left(-\frac{2}{\alpha_+}\phi\right) \right].$$
(5.31)

After integrating over $\beta, \bar{\beta}$, yields

$$\mathcal{L}_{\rm ws} = \partial \phi \bar{\partial} \phi - \frac{2}{\alpha_+} \widehat{R} \phi + \frac{\bar{\partial} \gamma \partial \bar{\gamma}}{\lambda + \exp\left(-\frac{2}{\alpha_+}\phi\right)},\tag{5.32}$$

which is σ -model on a space interpolating between linear dilaton background in the UV region $\phi \to \infty$ and AdS₃ in the IR $\phi \to -\infty$. Note that $\lambda > 0$ is assumed in the above discussion. When $\lambda < 0$, the IR behavior remains AdS₃, while there is a curvature singularity located at certain value of ϕ . It was argued in [207] that to study single-trace $T\bar{T}$, both side of the singular wall should be included.

Deformation of symmetric orbifold CFT

An asymmetric orbifold like (5.4), based on the symmetric group S_N , is a particular kind of orbifold CFT. It is related to holographic CFT in many aspects [208]. As the same with generic orbifold CFT, the Hilbert space of symmetric orbifold 2D CFT contains untwist and twist sectors, where the latter can be derived by imposing modular invariance of the partition function. Following this spirit, the authors of [105] (see also [50]) figure out the modular invariant partition functions under single-trace $T\bar{T}$ in terms of generalized Heck operator denoted as T'_n ,

$$Z_N(\tau,\bar{\tau};\lambda) = \sum_{\{k_1,\dots,k_N\}} \frac{1}{\prod_{n=1}^N n^{k_n} k_n!} \prod_{n=1}^N (T'_n Z) (\tau,\bar{\tau};\lambda)^{k_n}, \quad \sum_{n=1}^N n k_n = N,$$
(5.33)

²⁸Note that ∂_x commute with $(\partial_{\bar{x}}\bar{J}\partial_{\bar{x}}+2\partial_{\bar{x}}^2\bar{J})$. And for example

$$\int \partial_x J \partial_x \cdot \left[\partial_{\bar{x}} \bar{J} \partial_{\bar{x}} \Phi_1\right] dz = \int \partial_x J \cdot \partial_{\bar{x}} \bar{J} \partial_x \partial_{\bar{x}} \Phi_1 d^2 z = \int \partial_x^2 J \cdot \partial_{\bar{x}}^2 \bar{J} \Phi_1 d^2 z.$$
(5.29)

which leads to the appearance of twisted sectors in the deformed theory. The deformed spectrum of single particle twisted states also follows from the above-deformed partition function, which (up to different conventions) matches the result (5.73) below derived in the bulk side. The generating function for partition function Z_N can be defined as

$$\mathcal{Z}(\tau,\bar{\tau};\lambda;p) := \sum_{N=0}^{\infty} p^N Z_N(\tau,\bar{\tau};\lambda) = \exp\left(\sum_{n=1}^{\infty} \frac{p^n}{n} \left(T'_n Z\right)(\tau,\bar{\tau};\lambda)\right).$$
(5.34)

The deformed partition function (5.33) matches the holographic calculation from bulk string theory side [209], suggested the single-trace $T\bar{T}$ deformed CFT is holographic dual to string in linear dilation background. Recently, an exact holographic duality was proposed, i.e., tensionless string/2D symmetric orbifold duality, with the seed CFT of the orbifold being a supersymmetric sigma model on \mathbb{T}^4 [210, 211]. This framework further investigated the single trace $T\bar{T}$ deformation in [212]. The deformed partition function of the orbifold is computed explicitly, including the spin structure for fermions, which matches the string partition function with current-current deformation on the background with the torus boundary. Therefore, it provides strong evidence that current-current deformation of tensionless strings on $\mathrm{AdS}_3 \times S^3 \times T^4$ is holographically dual to single-trace $T\bar{T}$ deformed \mathbb{T}^4 orbifold.

5.1 Little string theory perspective

As mentioned, the background implied by the lagrangian (5.32) interpolates between AdS₃ and linear dilaton. The latter are also known as little string theory (LST) background. LST is the decoupled theory of NS5 branes that are magnetically charged under the NS B field In type II string theory, k NS5-branes extending in $\mathbb{R}^{5,1}$ with coordinate x^{μ} , is described in string frame as

$$ds^{2} = dx_{\mu}dx^{\mu} + \left(1 + \frac{N\alpha'}{r^{2}}\right)dx^{i}dx^{i}, \quad \mu = 0, 1, ..., 5.$$

$$e^{2\Phi} = g_{s}^{2}\left(1 + \frac{N\alpha'}{r^{2}}\right)$$
(5.35)

with transverse direction $dx^i dx^i = dr^2 + r^2 d\Omega_3^2$. To decouple the dynamics on the fivebranes from the bulk, consider the limit $g_s \to 0, r/g_s$ fixed, one obtain the background

$$ds^{2} = dx_{\mu}dx^{\mu} + N\alpha' \left(d\sigma^{2} + d\Omega_{3}^{2} \right), \quad e^{2\Phi} = \frac{g_{s}^{2}N\alpha'}{r^{2}} \equiv e^{-2\sigma}.$$
 (5.36)

This background is holographically dual to LST on the fivebrane. The topology of the background (5.36) becomes

$$\mathbb{R}^{5,1} \times \mathbb{R}_{\phi} \times S^3. \tag{5.37}$$

Note the string coupling vanishes far away from brane $\sigma \to +\infty$ and divergent near the brane $\sigma \to -\infty$.

So far NS5-branes wrapped around $\mathbb{R}^{5,1}$ is considered, one can also let the k NS5-branes wrapped on compact $T^4 \times S^1$, then the geometry (5.37) becomes [13]

$$\mathbb{R}_t \times \mathbb{R}_\phi \times S^1 \times \mathcal{N} \quad \text{with} \quad \mathcal{N} = T^4 \times S^3. \tag{5.38}$$

String theory in this background is holographically dual to non-local, non-gravitational LST theory [213]. To make contact (5.38) with geometry induced by (5.32), one can add p F1 string around S^1 in (5.38), which will modify the IR geometry (near the brane region) while leaving the UV geometry intact. Meanwhile, the string coupling will grow when approaching the IR and saturate $g_s^2 \sim 1/p$. Thus, the perturbation theory applies to large p. Now the IR geometry becomes

$$AdS_3 \times T^4 \times S^3. \tag{5.39}$$

Therefore, we have obtained a geometry indicated by (5.32) starting from LST background [213]. Notice that the background with a black hole in the linear dilaton in the UV and BTZ in the IR arises in the $SL(2, \mathbb{R}) \times U(1)/U(1)$ coset model, which can be thought as finite temperature generalization of background implied by (5.32) [203]. In addition, the above geometry can also arise by null gauging of the CFT on $R^{1,1} \times AdS_3 \times \mathcal{N}$ [127,214].

As an example for the background, namely, local AdS in the IR and linear dilaton in the UV, let us consider type IIB string theory on $\mathbb{R}^{4,1} \times T^4 \times S^1$, k NS5-branes on $T^4 \times S^1$ and p fundamental strings on S^1 (smeared over the T^4), with n units of momentum along S^1 , on string frame [203, 207]

$$ds^{2} = \frac{1}{f_{1}} \left[-\frac{f}{f_{n}} dt^{2} + f_{n} \left(dx + \frac{r_{0}^{2} \sinh 2\alpha_{n}}{2f_{n}r^{2}} dt \right)^{2} \right] + f_{5} \left(\frac{1}{f} dr^{2} + r^{2} d\Omega_{3}^{2} \right) + \sum_{i=1}^{4} dx_{i}^{2},$$

$$e^{2\Phi} = g^{2} \frac{f_{5}}{f_{1}}, \quad H = dx \wedge dt \wedge d \left(\frac{r_{0}^{2} \sinh 2\alpha_{1}}{2f_{1}r^{2}} \right) + r_{0}^{2} \sinh 2\alpha_{5} d\Omega_{3}$$
(5.40)

with

$$f = 1 - \frac{r_0^2}{r^2}, \quad f_{1,5,n} = 1 + \frac{r_{1,5,n}^2}{r^2}, \quad r_{1,5,n}^2 = r_0^2 \sinh^2 \alpha_{1,5,n}.$$
 (5.41)

$$\sinh 2\alpha_1 = \frac{2l_s^2 p}{vX}, \quad \sinh 2\alpha_5 = \frac{2l_s^2 k}{r_0^2}, \quad \sinh 2\alpha_n = \frac{2l_s^4 n}{R^2 vX}, \quad X = \frac{r_0^2}{g^2}.$$
 (5.42)

Note that for p = 0, n = 0, (5.40) reduces to (non-extremal) NS5-branes

$$ds^{2} = \left[-fdt^{2} + dx^{2}\right] + f_{5}\left(\frac{1}{f}dr^{2} + r^{2}d\Omega_{3}^{2}\right) + \sum_{i=1}^{4} dx_{i}^{2},$$

$$e^{2\Phi} = g^{2}f_{5}, \quad H = r_{0}^{2}\sinh 2\alpha_{5}d\Omega_{3},$$
(5.43)

which further reduces to extremal NS5 when $r_0 = 0$.

To investigate the thermodynamics of the dual field, transform (5.40) to Einstein frame

$$ds_E^2 = e^{-\Phi/2} ds^2, (5.44)$$

in which frame we can compute ADM mass from tress tensor T_{00} as

$$M = \frac{V_{S_1} V_{T^4}}{8\pi G_N^{(10)}} \int d^4 y T_{00} = \frac{V_{S_1} V_{T^4}}{8\pi G_N^{(10)}} \int d^4 y (R_{00}^{(1)} - \frac{1}{2} \eta_{00} R^{(1)})$$

$$= \frac{RvX}{2l_s^4} \left(\cosh 2\alpha_1 + \cosh 2\alpha_5 + \cosh 2\alpha_n\right)$$
(5.45)

with the flat transverse coordinates $y^i, i = 1, ..., 4$ satisfying

$$dy^{i}dy^{i} = dr^{2} + r^{2}d\Omega_{3}^{2}.$$
(5.46)

By evaluating the area of the horizon at $r = r_0$, the entropy turns out to be

$$S = \frac{A}{4G_N^{(10)}} = \frac{2\pi R v X r_0}{l_s^4} \cosh \alpha_1 \cosh \alpha_5 \cosh \alpha_n, \qquad (5.47)$$

Next, taking the decoupling limit, the energy and entropy become

$$S = \frac{2\pi R v X r_0}{l_s^3} \sqrt{k} \cosh \alpha_1, \quad E = \frac{R v X}{2l_s^4} (e^{-2\alpha_1} + 1), \tag{5.48}$$

which satisfies

$$S = 2\pi \sqrt{\frac{kl_s^2 \mathcal{E}^2}{R^2} + 2kp\mathcal{E}}, \quad \mathcal{E} = ER,$$
(5.49)

which is consistent with the entropy (5.3) of single-trace $T\bar{T}$ deformed theory, provided one identifies the deformation parameter as $\lambda = l_s^2/R^2$. Therefore, it is suggested that the gravity in the above background is dual to single-trace $T\bar{T}$ deformed CFT. As one varies the λ , the energy \mathcal{E} will be changed, but not the number of states

$$S(\lambda) = 2\pi \sqrt{k\lambda \mathcal{E}(\lambda)^2 + 2kp\mathcal{E}(\lambda)} = S(0) = 2\pi \sqrt{2kp\mathcal{E}(0)}$$
(5.50)

It then follows the deformed spectrum

$$\frac{1}{p}\mathcal{E}(\lambda) = \frac{1}{\lambda}(-1 + \sqrt{1 + 2\lambda\mathcal{E}(0)/p}).$$
(5.51)

5.2 TsT and single trace TT

As introduced in the last subsection, the geometry dual to the single-trace $T\bar{T}$ deformation interpolated AdS₃ and linear dilaton, which is relevant in studying little string theory. More recently, it was suggested that the single-trace $T\bar{T}$ deformed CFT dual to long string sector on a background generated by TsT transformation of $AdS_3 \times T^4 \times S^3$. The latter duality is supported by matching the spectrum and thermodynamics [56]. Firstly, let us review the background of the TsT transformation. Following the convention in [56], the world-sheet action is

$$\tilde{S} = -\ell_s^{-2} \int d^2 z \left(\sqrt{-\eta} \eta^{ab} \tilde{G}_{\mu\nu} + \epsilon^{ab} \tilde{B}_{\mu\nu} \right) \partial_a \tilde{X}^{\mu} \partial_b \tilde{X}^{\nu} = \ell_s^{-2} \int d^2 z \tilde{M}_{\mu\nu} \partial \tilde{X}^{\mu} \bar{\partial} \tilde{X}^{\nu}, \qquad (5.52)$$

where $\tilde{M}_{\mu\nu} = \tilde{G}_{\mu\nu} + \tilde{B}_{\mu\nu}$ is the background fields [215]. The field $\tilde{G}_{\mu\nu}, \tilde{B}_{\mu\nu}$ and dilaton $\tilde{\Phi}$ is assumed to be independent of two coordinates denoted as $\tilde{X}^1, \tilde{X}^{\bar{2}}$. The associated Neother currents

$$j^a_{(n)} = \frac{\partial L}{\partial(\partial_a X^n)}, \quad n = 1, \bar{2}, \tag{5.53}$$

which will be useful in a moment.

Consider the TsT transformation of the background and coordinates, i.e., a T transformation along \tilde{X}^1 , followed by a shift of \tilde{x}^2

$$\tilde{x}^1 = x^1, \quad \tilde{x}^2 = x^2 - 2\tilde{\lambda}x^1/l_s^2,$$
(5.54)

and another T transformation along X^1 , which can be represented as ²⁹

$$(\tilde{G}_{\mu\nu}, \tilde{B}_{\mu\nu}, \tilde{\Phi}, \tilde{X}^{\mu}) \xrightarrow{T} (\tilde{g}_{\mu\nu}, \tilde{b}_{\mu\nu}, \tilde{\phi}, \tilde{x}^{\mu}) \xrightarrow{s} (g_{\mu\nu}, b_{\mu\nu}, \phi, x^{\mu}) \xrightarrow{T} (G_{\mu\nu}, B_{\mu\nu}, \Phi, X^{\mu}).$$
(5.56)

The T-duality and shift can be embedded in O(d, d) transformation (see, for example, [217]). The action of $g \in O(d, d)$ on background organized as

$$H(G,B)_{\hat{I}\hat{J}} = \begin{pmatrix} G - BG^{-1}B & BG^{-1} \\ -G^{-1}B & G^{-1} \end{pmatrix}_{\hat{I}\hat{J}}, \quad \hat{I}, \hat{J} = 1, \dots, 2d.$$
(5.57)

takes the form

$$H(G', B') = g^T H(G, B)g.$$
 (5.58)

Here O(d, d) element g is defined by obeying

$$g^{T}Lg = L$$
, with $L = \begin{pmatrix} 0 & \mathbb{I} \\ \mathbb{I} & 0 \end{pmatrix}$, $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. (5.59)

Beside (5.58), the background field M is transformed as [215]

$$M \to M' = [aM + b][cM + d]^{-1}.$$
 (5.60)

It turns out that a coordinate shift, or more generally, a diffeomorphism transformation of background, can be realized by

$$g = \begin{pmatrix} A & 0\\ 0 & (A^{-1})^t \end{pmatrix}, \tag{5.61}$$

such that

$$g^{t}H(G,B)g = H(A^{T}GA, A^{T}BA).$$
(5.62)

Note that for the shift in (5.56), one has

$$A_{\mu\nu} = \frac{\partial \tilde{X}^{\mu}}{\partial \tilde{x}^{\nu}} = \begin{pmatrix} 1 & 0\\ -2\tilde{\lambda}/l_s^2 & 1 \end{pmatrix}.$$
 (5.63)

$$\partial_a X^1 = -(\sqrt{-\eta \varepsilon_{ab} \eta^{cb} \tilde{g}_{1\mu} \partial_c \tilde{x}^{\mu}} + b_{1\mu} \partial_a \tilde{x}^{\mu}) = -\epsilon_{ab} J^o_{(1)}$$

$$\tilde{X}^i = \tilde{x}^i, \quad i \neq 1$$
(5.55)

²⁹The coordinate transformation corresponds T duality [216], taking the first T duality as an example, can be written $\partial_a \tilde{X}^1 = -(\sqrt{-\eta}\varepsilon_{ab}\eta^{cb}\tilde{a}_{1\mu}\partial_c \tilde{x}^{\mu} + \tilde{b}_{1\mu}\partial_a \tilde{x}^{\mu}) = -\epsilon_{ab}\tilde{J}^b_{(1)}$

As for T-duality along the k-th direction, the corresponding O(d, d) element is given by

$$g_{T_k} = \begin{pmatrix} \mathbb{I} - E_k & E_k \\ E_k & \mathbb{I} - E_k \end{pmatrix}.$$
(5.64)

 E_k the $d \times d$ -matrix with 1 in the (k, k)-entry and zero everywhere else. Then the metric and B-field transform according to the standard Buscher rules (k = 1)

$$\tilde{G}_{11} = \frac{1}{G_{11}} \quad G_{1i} = \frac{B_{1i}}{G_{11}}, \quad \tilde{G}_{ij} = G_{ij} - \frac{G_{1i}G_{1j} - B_{1i}B_{1j}}{G_{11}}.$$

$$\tilde{B}_{1i} = \frac{G_{1i}}{G_{11}}, i \neq 1; \quad \tilde{B}_{ij} = B_{ij} - \frac{G_{1i}B_{1j} - B_{1i}G_{1j}}{G_{11}}, i \neq 1.$$
(5.65)

Based on the above results, the background and after TsT transformation (5.56) is given by [56] [215] [218]

$$M = \tilde{M} \left(I + \frac{2\tilde{\lambda}}{\ell_s^2} \Gamma \tilde{M} \right)^{-1}, \quad \Phi_{\text{TsT}} = \tilde{\Phi} + \frac{1}{4} \log \left(\frac{\det G_{\mu\nu}}{\det \tilde{G}_{\mu\nu}} \right)$$
(5.66)

with $\Gamma_{\mu\nu} = \delta^1_{\mu} \delta^{\overline{2}}_{\nu} - \delta^{\overline{2}}_{\mu} \delta^1_{\nu}$. Here, the TsT transformation of the dilaton field is included. The action after TsT is

$$S = \ell_s^{-2} \int d^2 z M_{\mu\nu} \partial X^{\mu} \bar{\partial} X^{\nu}, \qquad (5.67)$$

Making use of (5.66), one has

$$\frac{\partial M}{\partial \tilde{\lambda}} = -2\ell_s^{-2}M\Gamma M. \tag{5.68}$$

Thus, one obtains the flow of deformed world-sheet action

$$\frac{\partial S}{\partial \tilde{\lambda}} = 2 \int \boldsymbol{j}_{(1)} \wedge \boldsymbol{j}_{(\overline{2})}$$
(5.69)

where $\mathbf{j}_{(1)}$ is one-form current related to two Noether currents (5.53) [56]. (5.69) implies that world-sheet action with TsT transformed background field corresponds to current-current deformation, which is reminiscent of (5.30), which dual to single-trace $T\bar{T}$ deformation. TsT transformed action is related to single-trace $T\bar{T}$ deformation on the boundary once proper coordinate $X^1, X^{(2)}$ is chosen. This holographic duality is further supported by matching the spectrum on world-sheet (5.67) and the deformed CFT side. The spectrum on the world sheet can be computed by relating the energy-momentum tensor before and after TsT. By repeated use coordinates transformation for T-duality (5.55) and shift (5.54), one can relate coordinates in TsT transformation [56] (see (5.56))³⁰

$$\partial_a \tilde{X}^n = \partial_a X^n - 2(\tilde{\lambda}/l_s^2) \epsilon^{nn'} \epsilon_{ab} J^b_{(n')}(X), \quad \epsilon^{12} = -1.$$
(5.72)

In [56], \tilde{X} is replaced by auxiliary fields \hat{X}^n to distinguish the non-trivial boundary condition for \hat{X}^n . With (5.72) and boundary conditions for auxiliary fields \hat{X}^n , one can relate the Virasoro constraint before and after TsT so that the deformed spectrum follows

$$E_L(0) = E_L(\lambda) - \frac{2\lambda\ell}{wk} E_L(\lambda) E_R(\lambda), \quad E_R(0) = E_R(\lambda) - \frac{2\lambda\ell}{wk} E_L(\lambda) E_R(\lambda), \quad (5.73)$$

in which for w = -1 is reminiscent of deformed spectrum (3.8) of double trace $T\bar{T}$ deformation. with deformation constant proportitional to λ . For w = -1, the deformed spectrum matches the spectrum of single-particle states in the untwisted sector of single-trace deformed orbifold theory. At the same time, for w < -1, it corresponds to Z_{-w} twisted sector of the deformed orbifold. Therefore, the spectrum on both sides of the duality match each other.

In addition to the spectrum, a further piece of evidence is provided by matching the deformed correlation functions [112] [50] [219], where the two-point function of the vertex operators on worldsheet after TsT is computed, both perturbative and non-perturbative results are presented. Meanwhile, in the dual single-trace $T\bar{T}$ orbifold, the two-point function of the twist operator, which is dual to vertex operator with a specific quantum number in the worldsheet, is calculated by conformal perturbation theory, as well as by the nonperturbative method (for untwisted sector correlation functions) developed in [48]. Consistent results from both sides were established. However, the matching of correlation functions of general primary operators in the twisted sector remains unsolved. In addition to the long string sector, the deformed correlation functions were also considered in the short string sector, which can not be described in symmetric orbifold [127, 203, 220].

 $^{30}(5.72)$ can be obtained by the following coordinate transformations

$$\partial_{a}\tilde{X}^{1} = -\left(\sqrt{-\eta}\varepsilon_{ab}\eta^{cb}\tilde{g}_{1\mu}\partial_{c}\tilde{x}^{\mu} + \tilde{b}_{1\mu}\partial_{a}\tilde{x}^{\mu}\right) = -\epsilon_{ab}\tilde{j}^{b}_{(1)}(\tilde{x}), \quad \tilde{X}^{2} = \tilde{x}^{2},$$

$$\tilde{g}_{1\mu}\partial\tilde{x}^{\mu} = \frac{\partial x^{\rho}}{\partial\tilde{x}^{1}}g_{\rho\mu}\partial x^{\mu} = g_{1\mu}\partial x^{\mu} + 2(\tilde{\lambda}/l_{s}^{2})g_{2\mu}\partial x^{\mu},$$

$$\partial_{a}X^{1} = -\left(\sqrt{-\eta}\varepsilon_{ab}\eta^{cb}g_{1\mu}\partial_{c}x^{\mu} + b_{1\mu}\partial_{a}x^{\mu}\right) = -\epsilon_{ab}j^{b}_{(1)}, \quad X^{2} = x^{2},$$
(5.70)

which corresponds to T-duality, shift, and another T-duality in (5.56), respectively. Also, the Buscher rules (5.65) for T-duality is used

$$m_{12} = \frac{M_{12}}{G_{11}}, \quad m_{22} = M_{22} - \frac{M_{12}M_{21}}{G_{11}}, \quad m_{i2} = M_{i2} - \frac{M_{i1}M_{12}}{G_{11}}.$$
 (5.71)

6 Recent Developments and Perspectives

In recent years, significant progress has been made in the study of $T\bar{T}$ -like deformation and its generalizations, providing new insights and establishing connections across field theory, holography, and string theory. Generalized $T\bar{T}$ -like operators broaden the original scope of the investigation. These developments enhance our understanding of non-locality and integrability and offer new perspectives on the relationship between classical geometric formulations and deformed field theories. This section briefly reviews recent advances and outlines potential directions for future research.

6.1 Generalized $T\overline{T}$ -Like Deformations

The generalization of the $T\bar{T}$ deformation can be explored from various perspectives, including spacetime symmetries, conserved currents, and dimensionality. Extensions to different settings often adapt the two-dimensional $T\bar{T}$ deformation framework to accommodate variations in symmetry, composite operators, and spacetime dimensions.

Beyond relativistic QFT

The $T\bar{T}$ deformation has been generalized to various systems beyond relativistic QFT and string theory. There are generalizations to non-relativistic QFTs [93,221–224], non-relativistic string theory [225], and many body systems such as interacting particles [226–228]. Classically, the $T\bar{T}$ -like deformations of free particles lead to novel classical integrable many-body systems [229,230]. Additionally, a lot of works have been focused on the $T\bar{T}$ deformation of integrable spin chains [209,231–233], where the $T\bar{T}$ deformation preserves integrability and the S-matrix is deformed by a CDD factor. Recently, $T\bar{T}$ deformation has also been studied in lattice theories in [234].

The analog of the $T\bar{T}$ deformation in one dimension has been introduced in [22,235], which can be defined through the flow equation of the Hamiltonian:

$$\partial_{\lambda}H = \frac{H^2}{1/2 - 2\lambda H},\tag{6.1}$$

which can be defined via the dimensional reduction of $T\bar{T}$ dimension in two dimensions. The $T\bar{T}$ deformation in one dimension can be viewed as a special case, a broad class solvable of deformation that maps the Hamiltonian H to a function f(H) of H. The deformed spectrum and partition function can be obtained from the undeformed theory. The geometrical realization of the above flow equation (6.1) has been explored in [236].

The one-dimensional analog of $T\bar{T}$ deformation also has interesting holographic interpretations. The two-dimensional Jackiw-Teitelboim gravity is holographically dual to a solvable one-dimensional quantum theory known as the Schwarzian theory [237–239]. The holographic duality between two-dimensional Jackiw-Teitelboim gravity at finite cutoff and $T\bar{T}$ deformed Schwarzian theory was studied in [7,22,235]. The genus expansion over two-dimensional surfaces of the partition function of the Jackiw-Teitelboim gravity can be computed by a particular double-scaled matrix integral [240]. $T\bar{T}$ deformation of matrices models are investigated in [26,241,242].

Other double current deformations

The $T\bar{T}$ deformation is a special case of double current deformations driven by bilinears of Noether currents of the original theory. Double current deformations play an important role in integrable models and string theory.

When there is a U(1) current in a two-dimensional CFT, one can define a solvable Lorentzbreaking $J\bar{T}$ deformation [243]. The holographic dual of the $J\bar{T}$ deformed CFT is related to mixed boundary conditions between the metric and a U(1) gauge field [153]. There are interesting relations between $J\bar{T}$ deformation and string theory [34,244,245]. Correlation functions and entanglement entropy of $J\bar{T}$ deformed CFTs were computed in [113]. Modular covariance of $J\bar{T}$ deformed CFTs was discussed in [246]. For theories with higher spin conserved currents, such as two-dimensional integrable field theories or CFTs, one can construct deformations using these currents [1,94,247,248].

Generic energy-momentum tensor induced deformation in various spacetime

Besides $T\overline{T}$, one can construct other Lorentz invariant operators from the energy-momentum tensor. One interesting example is the root- $T\overline{T}$ operator [36, 37, 40]:

$$R = \sqrt{\frac{1}{2}T^{\mu\nu}T_{\mu\nu} - \frac{1}{4}T^{\mu}{}_{\mu}T^{\nu}{}_{\nu}}.$$
(6.2)

The rigorous definition of the root- $T\bar{T}$ operator at the quantum level remains uncertain, and a recent proposal has been presented in [41]. Nevertheless, the deformation triggered by the root- $T\bar{T}$ operator has increased interest because it exhibits some nice properties at the classical level. The root- $T\bar{T}$ deformation commutes with the $T\bar{T}$ deformation. The classical integrability [42] of specific integrable models is preserved by the root- $T\bar{T}$ deformation. The root- $T\bar{T}$ deformed conformal field theories are related to the ultra-relativistic (BMS₃) field theories [249] [250]. The role of the root- $T\bar{T}$ deformed conformal field theories can be derived in [251,252]. The modular properties of root- $T\bar{T}$ deformed conformal field theories can be derived from the holographic description [107]. The geometric formulation of the combination of the $T\bar{T}$ and root- $T\bar{T}$ deformation has been presented in [46, 129].

We review some interesting details about the root- $T\bar{T}$ deformation. The root- $T\bar{T}$ deformed classical action of N real scalars can be computed exactly [36, 37, 40]. In the following, we mainly follow the derivation in [37]. To obtain the deformed Lagrangian, one needs to solve the flow equation

$$\frac{\partial \mathcal{L}_{\gamma}}{\partial \gamma} = R_{\gamma} = \sqrt{\frac{1}{2} T^{\mu\nu} T_{\mu\nu} - \frac{1}{4} T^{\mu}{}_{\mu} T^{\nu}{}_{\nu}}, \qquad (6.3)$$

with the initial condition

$$\mathcal{L}_0 = \frac{1}{2} \Big(G_{ij}(\phi) g^{\mu\nu} + B_{ij}(\phi) \varepsilon^{\mu\nu} \Big) \partial_\mu \phi^i \partial_\nu \phi^j - V(\phi).$$
(6.4)

We use γ to denote the dimensionless deformation parameter. G_{ij} , B_{ij} , and V represent the target space metric, Kalb-Ramond fields, and the potential, respectively. One can use the Hilbert energy-momentum tensor

$$T_{\mu\nu} = -2\frac{\partial \mathcal{L}}{\partial \gamma^{\mu\nu}} + \gamma_{\mu\nu}\mathcal{L}.$$
(6.5)

Now, we simplify the flow equation by using Lorentz invariant variables. We define matrices

$$(X_1)^{\nu}_{\mu} = G_{ij}\partial_{\mu}\phi^i\partial^{\nu}\phi^j, \quad (X_n)^{\nu}_{\mu} = (X_1^n)^{\nu}_{\mu}, \tag{6.6}$$

and traces $x_n = \text{tr}X_n$. In two dimension, $x_{n>2}$ algebraically depends on x_1 and x_2 . For example, we have

$$2x_3 = 3x_1x_2 - x_1^3, \quad 2x_4 = 2x_1^2x_2 - x_1^4 + x_2^2.$$
(6.7)

Then the energy-momentum tensor is given by

$$T_{\mu\nu} = -2(X_1)_{\mu\nu} \frac{\partial \mathcal{L}_{\gamma}}{\partial x_1} - 4(X_2)_{\mu\nu} \frac{\partial \mathcal{L}_{\gamma}}{\partial x_2} + \gamma_{\mu\nu} \mathcal{L}_{\gamma}.$$
(6.8)

The square of the root- $T\bar{T}$ operator is

$$R^{2} = \left(2x_{2} - x_{1}^{2}\right) \left(\frac{\partial \mathcal{L}_{\gamma}}{\partial x_{1}} + 2x_{1}\frac{\partial \mathcal{L}_{\gamma}}{\partial x_{2}}\right)^{2}.$$
(6.9)

Remarkably, it does not explicitly depend on \mathcal{L}_{γ} and depends on the derivatives of \mathcal{L}_{γ} through a perfect square. The flow equation can be simplified as

$$\frac{\partial \mathcal{L}_{\gamma}}{\partial \gamma} = \pm \sqrt{2x_2 - x_1^2} \left(\frac{\partial \mathcal{L}_{\gamma}}{\partial x_1} + 2x_1 \frac{\partial \mathcal{L}_{\gamma}}{\partial x_2} \right), \tag{6.10}$$

which can be solved by using the method of characteristics. The solution is

$$\mathcal{L}_{\gamma}(x_0, x_1, x_2) = \mathcal{L}_{\gamma=0}\left(x_0, x_1^{(\gamma)}, x_2^{(\gamma)}\right),$$
(6.11)

with

$$x_1^{(\gamma)} = \cosh(\gamma) x_1 \pm \sinh(\gamma) \sqrt{2x_2 - x_1^2},$$
 (6.12)

$$x_2^{(\gamma)} = \cosh(2\gamma)x_2 \pm \sinh(2\gamma)x_1\sqrt{2x_2 - x_1^2},\tag{6.13}$$

and x_0 denotes the terms independent of $x_{1,2}$. For the initial condition (6.4), we have

$$x_0 = \frac{1}{2} B_{ij}(\phi) \varepsilon^{\mu\nu} \partial_\mu \phi^i \partial_\nu \phi^j - V(\phi).$$
(6.14)

In the special case of free scalar with $x_0 = 0$ and $G_{ij} = \delta_{ij}$, the deformed Lagrangian is given by

$$\mathcal{L}_{\gamma} = \frac{\cosh \gamma}{2} \Phi_{\mu}{}^{\mu} \pm \frac{\sinh \gamma}{2} \sqrt{2 \Phi_{\mu}{}^{\nu} \Phi_{\nu}{}^{\mu} - (\Phi_{\mu}{}^{\mu})^2}.$$
 (6.15)

When there are two scalars, it can be obtained from the four-dimensional ModMax theory by dimensional reduction [36].

Like the $T\bar{T}$ deformation, the combination of the root- $T\bar{T}$ and $T\bar{T}$ deformation can be reformulated as coupling the undeformed theory to a gravity action [46, 129]:

$$S_{\gamma,\lambda}[\phi, e^a_{\mu}, f^a_{\mu}] = S_0[\phi, e^a_{\mu}] + S_{\text{grav}}[e^a_{\mu}, f^a_{\mu}], \qquad (6.16)$$

where S_0 denotes the action of the undeformed theory, and e^a_μ denotes an auxiliary dynamical zweibein. The physical background zweibein f^a_μ couples to e^a_μ through the gravity action $S_{\rm grav}$

$$S_{\text{grav}}[e^a_{\mu}, f^a_{\mu}] = \frac{1}{2\lambda} \int d^2x \det e\left(2 + y_1^2 - y_2 - 2y_1 \cosh\frac{\gamma}{2} + 2\sqrt{2y_2 - y_1^2} \sinh\frac{\gamma}{2}\right), \quad (6.17)$$

where

$$y_1 = f^a_\mu e^\mu_a, \quad y_2 = f^a_\mu e^\mu_b f^b_\nu e^\nu_a, \tag{6.18}$$

are two Lorentz invariant variables.

The explicit form is

$$S_{\text{grav}} = \frac{1}{\lambda} \int d^2 x \left(\left(e_1^2 f_2^1 - e_1^1 f_2^2 + e_2^1 f_1^2 - e_2^2 f_1^1 \right) \cosh \frac{\gamma}{2} - e_2^1 e_1^2 + e_2^2 e_1^1 - f_2^1 f_1^2 + f_1^1 f_2^2 + \sqrt{\left(e_1^1 e_2^2 - e_2^1 e_1^2 \right) \left(\frac{\left(-e_1^2 f_2^1 + e_1^1 f_2^2 - e_2^1 f_1^2 + e_2^2 f_1^1 \right)^2}{e_1^1 e_2^2 - e_2^1 e_1^2} + 4f_2^1 f_1^2 - 4f_2^2 f_1^1 \right) \sinh \frac{\gamma}{2} \right).$$

$$(6.19)$$

The deformed action can be obtained by solving the equation of motion for e^a_ν

$$\det e T_{0a}^{\nu} \equiv \frac{\delta S_0}{\delta e_{\nu}^a} = -\frac{\delta S_{\text{grav}}}{\delta e_{\nu}^a},\tag{6.20}$$

and plugging the solution back to (6.16).

The energy-momentum tensor of the deformed theory can be computed as

$$T^{\nu}_{\ \mu} \equiv \frac{1}{\det f} \frac{\delta S_{\gamma,\lambda}}{\delta f^a_{\nu}} f^a_{\mu} = \frac{1}{\det f} \frac{\delta S_{\text{grav}}}{\delta f^a_{\nu}} f^a_{\mu}.$$
(6.21)

)

Note that the result is the same whether we perform the extremization of the action with respect to e^a_{ν} before or after. One can verify the action satisfies the flow equations

$$\frac{\partial S_{\gamma,\lambda}}{\partial \lambda} = -\int d^2 x \det f \det(T^{\nu}_{\ \mu}) \\
= -\int d^2 x \det f \left(\frac{2\sqrt{2y_2 - y_1^2} \sinh\frac{\gamma}{2} - 2y_1 \cosh\frac{\gamma}{2} + y_1^2 - y_2 + 2}{\lambda^2 (y_1^2 - y_2)} \right), \quad (6.22)$$

$$\frac{\partial S_{\gamma,\lambda}}{\partial \gamma} = \int d^2 x \det f \sqrt{\frac{1}{2} T^{\mu}_{\nu} T^{\nu}_{\mu} - \frac{1}{4} (T^{\nu}_{\nu})^2} \\
= \int d^2 x \det f \left(\frac{\sqrt{2y_2 - y_1^2} \cosh\frac{\gamma}{2} - y_1 \sinh\frac{\gamma}{2}}{\lambda (y_1^2 - y_2)} \right). \quad (6.23)$$

Therefore the action (6.16) gives a gravitational description of the combination of the root- $T\bar{T}$ and $T\bar{T}$ deformations. Using a change of variables $y_1 = \alpha_1 + \alpha_2$ and $y_2 = \alpha_1^2 + \alpha_2^2$, the gravity action can be simplified as

$$S_{\text{grav}}[e^{a}_{\mu}, f^{a}_{\mu}] = \frac{1}{\lambda} \int d^{2}x \, \det e(\alpha_{1} - e^{\frac{\gamma}{2}})(\alpha_{2} - e^{-\frac{\gamma}{2}}).$$
(6.24)

The α parameters can be viewed as the eigenvalues of the matrix $e^{\mu}_{a} f^{a}_{\nu}$. This expression is useful for generalizing root- $T\bar{T}$ deformation to higher dimensions [46].

Energy-momentum tensor deformations in higher dimensions which generalize the $T\bar{T}$ deformation have been proposed in [17, 36, 47, 64, 90, 91, 97, 253–255]. Specific models in four dimensions include the Modified Maxwell theory [62] and its Born-Infeld-like extension [256] which have been studied as energy-momentum tensor flows [39,63,65]. There are also non-linear gauge theories as energy-momentum tensor flows in three [253], six and higher dimensions [257].

Inquiring whether general energy-momentum tensor deformations exhibit rich and interesting properties in various physical models is natural. Recent investigations have uncovered several significant features related to such deformations. It has been demonstrated in [258] that energy-momentum tensor deformations of the two-dimensional principal chiral models can be expressed through an auxiliary field formulation, which facilitates the proof of the classical integrability of these models. This result highlights the robustness of integrable structures in the presence of such deformations. Moreover, it has been shown that the non-Abelian T-duality of the principal chiral models commutes with arbitrary energy-momentum tensor deformations. This intriguing result is derived using the auxiliary field approach [259], indicating that the integrability properties are preserved under both operations. The applications of the auxiliary field formulation extend beyond the classical integrability framework and have been further explored in various contexts. For instance, [260–262] delve into more intricate properties of the auxiliary field formulation, providing deeper insights into its utility in both integrable and non-integrable systems. For four-dimensional electrodynamics, general energy-momentum tensor deformations preserve the electric-magnetic duality of Maxwell's theory. This preservation of duality under deformations is a remarkable result, as demonstrated in a series of works [38,39,64,65,263,264]. Furthermore, the relationship between duality symmetries and energy-momentum tensor deformations in *p*-form theories across various dimensions has been systematically investigated in [257], revealing the broad applicability of these deformations in different physical settings. Also, energy-momentum tensor deformations have a natural connection with modified gravity models. These deformations are pivotal in various modified gravity theories, providing a fertile ground for exploring non-trivial gravitational dynamics. Such connections have been extensively studied in [43, 46, 129, 265–268], shedding light on how energy-momentum tensor deformations can lead to modifications in gravitational theories, including the coupling of matter fields to modified geometries.

In summary, the study of general energy-momentum tensor deformations is an active area of research with implications across a wide range of physical models, including integrable systems, duality symmetries, and gravitational theories. Recent results suggest that these deformations provide a versatile framework for probing the deeper structures of classical and quantum field theories.

6.2 Some perspectives

Several promising directions can be pursued to understand various aspects of $T\overline{T}$ -like deformed theories regarding holography, UV completion, duality, etc.

Holography and $T\overline{T}$ -like deformation

The holographic interpretation of the $T\overline{T}$ deformation corresponds to asymptotically anti-de Sitter (AdS) geometries with modified boundary conditions or radial cutoffs [12, 14, 21]. Within this framework, bulk correlation functions are computed via holographic duality and mapped to boundary observables. In pure Einstein gravity, the boundary metric—determined by the Ricci flow equation [43, 267]—serves as the foundation for bulk reconstruction. Consistency with the AdS/CFT correspondence is ensured by verifying that the boundary energy-momentum tensor's correlation functions align with the standard holographic dictionary. Extending this to theories with matter fields involves imposing $T\overline{T}$ -compatible boundary conditions, after which the Gubser-Klebanov-Polyakov-Witten prescription enables systematic analysis of deformed holographic correlators. Recent developments further enhance this duality by emulating $T\overline{T}$ like deformations on hyperlattices using quantum circuits [269–271].

A discrete formulation of holography offers a powerful paradigm for studying $T\overline{T}$ -like deformations. Two key frameworks emerge:

- Holographic Tensor Networks: By discretizing bulk spacetime into interconnected tensors encoding local geometry and entanglement, this approach [272] provides a tractable representation of bulk fields. The resulting discrete structure simplifies computations of holographic correlation functions while preserving essential features of continuum physics.
- *p*-adic Conformal Field Theories: As a non-archimedean generalization of standard CFTs, *p*-adic frameworks directly relate deformation parameters to bulk geometric modifications [273]. This connection facilitates exploration of non-perturbative effects and insights into strongly coupled regimes [274].

These discrete frameworks not only enable controlled studies of $T\overline{T}$ -like deformations but also reveal novel deformation classes. For instance, holographic tensor networks [11,275–277] reinterpret bulk geometry as entanglement structures, while *p*-adic CFTs establish geometric dualities inaccessible to conventional continuum methods. The interplay between discrete bulk structures and boundary theories opens new avenues for understanding quantum field theories and their holographic duals, particularly in regimes where traditional continuum approaches face computational or conceptual challenges.

UV Completion and Nonlocality

Using bootstrap constraints to explore the possibility of UV completion or the nature of nonlocal interactions induced by the deformation [13, 17]. At the classical level, the deformed theory can be regarded as seed theory coupled with flat JT-like gravity. One can expect to check whether the equivalent relation can be held at the quantum level. The one direct check is to obtain finite entanglement entropy in the deformed theory due to UV completion and nonlocality. Whether the entanglement entropy can be regarded as the quantum extremal surface in the UV region, the Hilbert space of the deformed theory should satisfy the type II von Neumann algebra [130]. However, the entanglement entropy in the deformed theory is divergent regarding the perturbation theory approach. From the IR to UV, how to understand the $T\bar{T}$ deformation induces the dramatic changes in Hilbert space is an interesting open question.

Dualities and Symmetries

By leveraging deformation methods tied to the energy-momentum tensor, we can identify deformation theories that satisfy both duality constraints and integrability requirements. This approach enables the systematic construction of theories that preserve these dual principles. One can refer to some interesting examples [73, 278]. The investigation focuses on uncovering potential dualities or hidden symmetries in deformed theories, which may be constrained or revealed through bootstrap methods [207] or classical duality equations, such as those in [66, 67, 279–281]. For instance, three principal frameworks exist for constructing self-dual theories of Maxwell fields with SO(2) duality symmetry: Gaillard-Zumino approach [66, 67], refined by Gibbons and Rasheed [281]; non-covariant Hamiltonian formalism introduced by Henneaux and Teitelboim [68], extended by Deser et al. [282, 283]; PST method by Pasti, Sorokin, and Tonin [69–72]. These frameworks provide distinct pathways to incorporate interactions while retaining duality symmetry. Recent work in [259] further demonstrates that T-duality transformations commute with energy-momentum tensor deformations via an auxiliary field formulation [258, 261], offering a unified perspective on duality-preserving deformations.

Bootstrap $T\bar{T}$

The $T\overline{T}$ deformation leads to intriguing modifications of the spectrum and dynamics, often resulting in nonlocal theories with characteristics such as finite energy spectra and Hagedornlike behavior [2]. Despite its nontrivial nature, the solvability of $T\overline{T}$ -deformed theories provides a valuable framework for analytical studies, offering fertile ground to explore nonperturbative phenomena in quantum field theory.

A powerful tool in studying such theories has been the bootstrap approach, particularly the conformal bootstrap, which leverages symmetries and consistency conditions like crossing symmetry and unitarity to solve for operator dimensions and correlation functions in conformal field theories. However, extending these techniques to $T\overline{T}$ -deformed theories presents several challenges. First, the original conformal symmetry is broken, and we are left with a $T\overline{T}$ -deformed conformal symmetry consisting of field-dependent transformations. It's an open question of utilizing this complicated deformed symmetry to find constraints imposed on the theory. In particular, we don't know what operators can make the symmetry manifest. Second, the deformation introduces nonlocal interactions, complicating the definition and calculation of correlation functions. Traditional bootstrap methods rely on local operator product expansions (OPEs), but these may not hold in the context of nonlocal theories [106]. Finally, as an irrelevant deformation, the $T\overline{T}$ operator introduces ultraviolet divergences, making the theory sensitive to high-energy details. This sensitivity poses a significant challenge to the bootstrap approach, which often depends on infrared (IR) data and universal behavior [17]. Thus, while the $T\overline{T}$ deformation opens up intriguing avenues for nonperturbative study, its nonlocality, loss of conformal symmetry, and UV sensitivity introduce substantial challenges to applying traditional bootstrap techniques.

Acknowledgements

We would like to thank H. Babaei-Aghbolagh, Ronggen Cai, Bin Chen, Jingcheng Chang, Yi-Hong Gao, Miao He, Jue Hou, Ling-Yan Hung, Yunfeng Jiang, Yun-Ze Li, Yu-Xiao Liu, Zhangcheng Liu, Xin-Cheng Mao, Pujian Mao, Tommaso Morone, Hongfei Shu, Jia Tian, Roberto Tateo, Huajia Wang, Jun-Bao Wu, Wei Song, Hao-Yu Sun, Jia-Rui Sun, Stefan Theisen, Zhuo-Yu Xian, Yunfei Xie, Jiashi Yin, Yu-Xuan Zhang, Long Zhao and Zi-Xuan Zhao for valuable discussions related to this work at various stages. SH would like to appreciate the financial support from Ningbo University, the Max Planck Partner group, and the Natural Science Foundation of China Grants (No. 12475053, No. 12235016). HO is supported by the National Natural Science Foundation of China, Grant No. 12205115, and by the Science and Technology Development Plan Project of Jilin Province of China, Grant No. 20240101326JC. YS is supported by the National Natural Science Foundation of China, Reat No. 12105113.

A Appendix: Useful integrals

A.1 Dimensional regularization

In this appendix, we present the detailed computations of integrals appearing in the conformal perturbation theory in section 3.3, where the deformed correlation is investigated.

We will consider the plane case, namely, the deformed theory is defined on a 2D plane, while the torus case can be found in [116, 120]. The general integral on a plane can be expressed as [113]

$$\mathcal{I}_{a_1,\cdots,a_m,b_1,\cdots,b_n}(z_{i_1},\cdots,z_{i_m},\bar{z}_{j_1},\cdots,\bar{z}_{j_n}) \equiv \int \frac{d^2z}{(z-z_{i_1})^{a_1}\cdots(z-z_{i_m})^{a_m}(\bar{z}-\bar{z}_{j_1})^{b_1}\cdots(\bar{z}-\bar{z}_{j_n})^{b_n}}$$
(A.1)

Using Feynman parametrization and dimensional regularization, one can obtain the following basic integral [111]

$$\mathcal{I}_{11}(z_1, \bar{z}_2) = \int d^2 z \frac{1}{z_{01}\bar{z}_{02}} = -\pi \Big(-\frac{2}{\epsilon} + \ln|z_{12}|^2 + \gamma + \ln\pi \Big) + O(\epsilon), \tag{A.2}$$

with ϵ being an infinitesimal constant. Next example is $\mathcal{I}_{12}(z_1, \bar{z}_2)$

$$\int d^{2}z \frac{1}{z_{01}\bar{z}_{02}^{2}} = \int d^{2}z \frac{\bar{z}_{01}z_{02}^{2}}{|z_{01}|^{2}|z_{02}|^{4}}$$

$$= 2 \int_{0}^{1} du(1-u) \int d^{2}z \frac{\bar{z}_{01}z_{02}^{2}}{(u|z_{01}|^{2}+(1-u)|z_{02}|^{2})^{3}}$$

$$= 2 \int_{0}^{1} du(1-u) \int d^{2}y \frac{2uz_{12}|y|^{2}-(1-u)u^{2}z_{12}^{2}\bar{z}_{12}}{(|y|^{2}+(1-u)u|z_{12}|^{2})^{3}}$$

$$= 2z_{12} \int_{0}^{1} duu(1-u) \int d^{2}y \frac{2|y|^{2}-A^{2}}{(|y|^{2}+A^{2})^{3}}$$

$$= 2z_{12} \int_{0}^{1} duu(1-u)V_{d} \int d\rho\rho^{d-1} \frac{2\rho^{2}-A^{2}}{(\rho^{2}+A^{2})^{3}}$$

$$= 2z_{12} \int_{0}^{1} duu(1-u)V_{d}A^{-2} \frac{1}{4} = \frac{\pi}{\bar{z}_{12}},$$
(A.3)

with d = 2 and $V_d = 2\pi^{d/2}/\Gamma(d/2)$ is the area of (d-1)-sphere with unit radius, also we denote $A^2 = (1-u)u|z_{12}|^2$ and use the coordinates transformation

$$z = y + uz_1 + (1 - u)z_2, \quad z_{01} = y - (1 - u)z_{12}, \quad z_{02} = y + uz_{12}.$$
 (A.4)

Let us mention that the result in eq.(A.3) is consistent with eq.(A.2), i.e. they satisfy $\partial_{\bar{z}_2} \mathcal{I}_{11}(z_i, \bar{z}_j) = \mathcal{I}_{12}(z_i, \bar{z}_j).$

For $\mathcal{I}_{22}(z_1, \bar{z}_2)$ with $z_1 \neq z_2$, similarly we can obtain

$$\int d^{2}z \frac{1}{z_{01}^{2} \bar{z}_{02}^{2}} = \int d^{2}z \frac{\bar{z}_{01}^{2} z_{02}^{2}}{|z_{01}|^{4} |z_{02}|^{4}}$$

$$= 6 \int_{0}^{1} duu(1-u) \int d^{2}y \frac{(\bar{y} - (1-u)\bar{z}_{12})^{2}(y+uz_{12})^{2}}{(|y|^{2} + (1-u)u|z_{12}|^{2})^{4}}$$

$$= 6 \int_{0}^{1} duu(1-u) \int d^{2}y \frac{|y|^{4} - 4|y|^{2}u(1-u)|z_{12}|^{2} + (1-u)^{2}u^{2}|z_{12}|^{4}}{(|y|^{2} + (1-u)u|z_{12}|^{2})^{4}}$$

$$= 6 \int_{0}^{1} duu(1-u) \int d^{2}y \frac{|y|^{4} - 4|y|^{2}A^{2} + A^{4}}{(|y|^{2} + A^{2})^{4}} = 0.$$
(A.5)

In summary, by using dimensional regularization, we can obtain all the integrals needed, which

are [115]

$$\mathcal{I}_{11}(z_i, \bar{z}_j) = -\pi \left(-\frac{2}{\epsilon} + \ln|z_{ij}|^2 + \gamma + \ln\pi + \mathcal{O}(\epsilon)\right), \\
\mathcal{I}_{12}(z_i, \bar{z}_j) = \frac{\pi}{\bar{z}_{ij}}, \quad \mathcal{I}_{21}(z_i, \bar{z}_j) = -\frac{\pi}{z_{ij}}, \quad \mathcal{I}_{22}(z_i, \bar{z}_j) = 0, \\
\mathcal{I}_{13}(z_i, \bar{z}_j) = \frac{\pi}{(\bar{z}_{ij})^2}, \quad \mathcal{I}_{31}(z_i, \bar{z}_j) = \frac{\pi}{(z_{ij})^2}, \\
\mathcal{I}_{23}(z_i, \bar{z}_j) = \mathcal{I}_{32}(z_i, \bar{z}_j) = \mathcal{I}_{33}(z_i, \bar{z}_j) = 0.$$
(A.6)

References

- F. A. Smirnov and A. B. Zamolodchikov, On space of integrable quantum field theories, Nucl. Phys. B 915 (2017) 363–383, arXiv:1608.05499.
- [2] A. Cavaglià, S. Negro, I. M. Szécsényi, and R. Tateo, TT-deformed 2D Quantum Field Theories, JHEP 10 (2016) 112, arXiv:1608.05534.
- [3] A. B. Zamolodchikov, Expectation value of composite field T anti-T in two-dimensional quantum field theory, arXiv:hep-th/0401146.
- [4] S. Dubovsky, V. Gorbenko, and M. Mirbabayi, Asymptotic fragility, near AdS₂ holography and TT, JHEP 09 (2017) 136, arXiv:1706.06604.
- [5] S. Dubovsky, V. Gorbenko, and G. Hernández-Chifflet, TT partition function from topological gravity, JHEP 09 (2018) 158, arXiv:1805.07386.
- [6] A. J. Tolley, TT deformations, massive gravity and non-critical strings, JHEP 06 (2020) 050, arXiv:1911.06142.
- [7] L. V. Iliesiu, J. Kruthoff, G. J. Turiaci, and H. Verlinde, JT gravity at finite cutoff, SciPost Phys. 9 (2020) 023, arXiv:2004.07242.
- [8] S. Okumura and K. Yoshida, TT-deformation and Liouville gravity, Nucl. Phys. B 957 (2020) 115083, arXiv:2003.14148.
- [9] S. Ebert, C. Ferko, H.-Y. Sun, and Z. Sun, TT in JT Gravity and BF Gauge Theory, SciPost Phys. 13 no. 4, (2022) 096, arXiv:2205.07817.
- [10] A. Bhattacharyya, S. Ghosh, and S. Pal, Aspects of $T\bar{T} + J\bar{T}$ deformed 2D topological gravity : from partition function to late-time SFF, arXiv:2309.16658.
- [11] N. Bao, L.-Y. Hung, Y. Jiang, and Z. Liu, QG from SymQRG: AdS₃/CFT₂ Correspondence as Topological Symmetry-Preserving Quantum RG Flow, arXiv:2412.12045.

- [12] L. McGough, M. Mezei, and H. Verlinde, Moving the CFT into the bulk with TT, JHEP 04 (2018) 010, arXiv:1611.03470.
- [13] A. Giveon, N. Itzhaki, and D. Kutasov, TT and LST, JHEP 07 (2017) 122, arXiv:1701.05576.
- [14] P. Kraus, J. Liu, and D. Marolf, Cutoff AdS_3 versus the $T\overline{T}$ deformation, JHEP 07 (2018) 027, arXiv:1801.02714.
- [15] V. Gorbenko, E. Silverstein, and G. Torroba, dS/dS and TT, JHEP 03 (2019) 085, arXiv:1811.07965.
- [16] W. Cottrell and A. Hashimoto, Comments on $T\bar{T}$ double trace deformations and boundary conditions, Phys. Lett. B 789 (2019) 251–255, arXiv:1801.09708.
- [17] M. Taylor, TT deformations in general dimensions,
 Adv. Theor. Math. Phys. 27 no. 1, (2023) 37–63, arXiv:1805.10287.
- [18] T. Hartman, J. Kruthoff, E. Shaghoulian, and A. Tajdini, Holography at finite cutoff with a T² deformation, JHEP 03 (2019) 004, arXiv:1807.11401.
- [19] Y. Jiang, Expectation value of TT operator in curved spacetimes, JHEP 02 (2020) 094, arXiv:1903.07561.
- [20] P. Caputa, S. Datta, and V. Shyam, Sphere partition functions \& cut-off AdS, JHEP 05 (2019) 112, arXiv:1902.10893.
- [21] M. Guica and R. Monten, *TT* and the mirage of a bulk cutoff, SciPost Phys. 10 no. 2, (2021) 024, arXiv:1906.11251.
- [22] D. J. Gross, J. Kruthoff, A. Rolph, and E. Shaghoulian, TT in AdS₂ and Quantum Mechanics, Phys. Rev. D 101 no. 2, (2020) 026011, arXiv:1907.04873.
- [23] G. Jafari, A. Naseh, and H. Zolfi, Path Integral Optimization for $T\bar{T}$ Deformation, Phys. Rev. D 101 no. 2, (2020) 026007, arXiv:1909.02357.
- [24] Y. Li and Y. Zhou, Cutoff AdS_3 versus $T\overline{T}$ CFT₂ in the large central charge sector: correlators of energy-momentum tensor, JHEP 12 (2020) 168, arXiv:2005.01693.
- [25] L. Griguolo, R. Panerai, J. Papalini, and D. Seminara, Nonperturbative effects and resurgence in Jackiw-Teitelboim gravity at finite cutoff, Phys. Rev. D 105 no. 4, (2022) 046015, arXiv:2106.01375.

- [26] S. He, H. Ouyang, and Y. Sun, Note on TT deformed matrix models and JT supergravity duals, Eur. Phys. J. C 83 no. 10, (2023) 885, arXiv:2204.13636.
- [27] S. He, Y. Li, Y.-Z. Li, and Y. Zhang, Holographic torus correlators of stress tensor in AdS₃/CFT₂, JHEP 06 (2023) 116, arXiv:2303.13280.
- [28] S. He, Y.-Z. Li, and Y. Zhang, Holographic torus correlators in AdS₃ gravity coupled to scalar field, JHEP 05 (2024) 254, arXiv:2311.09636.
- [29] S. Fichet, E. Megias, and M. Quiros, Holography of linear dilaton spacetimes from the bottom up, Phys. Rev. D 109 no. 10, (2024) 106011, arXiv:2309.02489.
- [30] S. Ebert, Holographic Renormalization Group and Stress Tensor Operators. PhD thesis, UCLA, Los Angeles (main), UCLA, 2024. arXiv:2404.10190.
- [31] S. He, Y.-Z. Li, and Y. Xie, Holographic stress tensor correlators on higher genus Riemann surfaces, arXiv:2406.04042.
- [32] M. Baggio and A. Sfondrini, Strings on NS-NS Backgrounds as Integrable Deformations, Phys. Rev. D 98 no. 2, (2018) 021902, arXiv:1804.01998.
- [33] A. Dei and A. Sfondrini, Integrable S matrix, mirror TBA and spectrum for the stringy $AdS_3 \times S^3 \times S^3 \times S^1 WZW$ model, JHEP 02 (2019) 072, arXiv:1812.08195.
- [34] S. Chakraborty, A. Giveon, and D. Kutasov, TT, JT, TJ and String Theory,
 J. Phys. A 52 no. 38, (2019) 384003, arXiv:1905.00051.
- [35] N. Callebaut, J. Kruthoff, and H. Verlinde, TT deformed CFT as a non-critical string, JHEP 04 (2020) 084, arXiv:1910.13578.
- [36] R. Conti, J. Romano, and R. Tateo, Metric approach to a TT-like deformation in arbitrary dimensions, JHEP 09 (2022) 085, arXiv:2206.03415.
- [37] C. Ferko, A. Sfondrini, L. Smith, and G. Tartaglino-Mazzucchelli, Root-TT Deformations in Two-Dimensional Quantum Field Theories, Phys. Rev. Lett. 129 no. 20, (2022) 201604, arXiv:2206.10515.
- [38] C. Ferko, L. Smith, and G. Tartaglino-Mazzucchelli, Stress Tensor flows, birefringence in non-linear electrodynamics and supersymmetry, SciPost Phys. 15 no. 5, (2023) 198, arXiv:2301.10411.
- [39] H. Babaei-Aghbolagh, K. B. Velni, D. M. Yekta, and H. Mohammadzadeh, Emergence of non-linear electrodynamic theories from TT⁻-like deformations, Phys. Lett. B 829 (2022) 137079, arXiv:2202.11156.

- [40] H. Babaei-Aghbolagh, K. Babaei Velni, D. Mahdavian Yekta, and H. Mohammadzadeh, Marginal TT⁻-like deformation and modified Maxwell theories in two dimensions, Phys. Rev. D 106 no. 8, (2022) 086022, arXiv:2206.12677.
- [41] L. Hadasz and R. von Unge, Defining the root-TT⁻ operator, Phys. Rev. D 111 no. 4, (2025) 045024, arXiv:2405.17945.
- [42] R. Borsato, C. Ferko, and A. Sfondrini, Classical integrability of root-TT⁻ flows, Phys. Rev. D 107 no. 8, (2023) 086011, arXiv:2209.14274.
- [43] N. Brizio, T. Morone, and R. Tateo, Stress-energy tensor deformations, Ricci flows and black holes, arXiv:2408.06031.
- [44] A. Giveon, Comments on $T\overline{T}$, $J\overline{T}$ and String Theory, arXiv:1903.06883.
- [45] Y. Jiang, A pedagogical review on solvable irrelevant deformations of 2D quantum field theory, Commun. Theor. Phys. 73 no. 5, (2021) 057201, arXiv:1904.13376.
- [46] H. Babaei-Aghbolagh, S. He, T. Morone, H. Ouyang, and R. Tateo, Geometric formulation of generalized root- $T\bar{T}$ deformations, arXiv:2405.03465.
- [47] G. Bonelli, N. Doroud, and M. Zhu, *TT*-deformations in closed form, JHEP 06 (2018) 149, arXiv:1804.10967.
- [48] J. Cardy, TT deformation of correlation functions, JHEP 12 (2019) 160, arXiv:1907.03394.
- [49] O. Aharony and N. Barel, Correlation functions in TT-deformed Conformal Field Theories, JHEP 08 (2023) 035, arXiv:2304.14091.
- [50] S. B. Chakraborty, S. Georgescu, and M. Guica, States, symmetries and correlators of TT and JT symmetric orbifolds, SciPost Phys. 16 no. 1, (2024) 011, arXiv:2306.16454.
- [51] S. Dubovsky, R. Flauger, and V. Gorbenko, Solving the Simplest Theory of Quantum Gravity, JHEP 09 (2012) 133, arXiv:1205.6805.
- [52] S. Dubovsky, V. Gorbenko, and M. Mirbabayi, Natural Tuning: Towards A Proof of Concept, JHEP 09 (2013) 045, arXiv:1305.6939.
- [53] S. Frolov, TT Deformation and the Light-Cone Gauge,
 Proc. Steklov Inst. Math. 309 (2020) 107–126, arXiv:1905.07946.

- [54] S. Frolov, TT, JJ, JT and JT deformations, J. Phys. A 53 no. 2, 025401, arXiv:1907.12117.
- [55] A. Sfondrini and S. J. van Tongeren, TT deformations as TsT transformations, Phys. Rev. D 101 no. 6, (2020) 066022, arXiv:1908.09299.
- [56] L. Apolo, S. Detournay, and W. Song, TsT, $T\bar{T}$ and black strings, JHEP 06 (2020) 109, arXiv:1911.12359.
- [57] C. Esper and S. Frolov, TT deformations of non-relativistic models, JHEP 06 (2021) 101, arXiv:2102.12435.
- [58] S. Pasterski, S.-H. Shao, and A. Strominger, Flat Space Amplitudes and Conformal Symmetry of the Celestial Sphere, Phys. Rev. D 96 no. 6, (2017) 065026, arXiv:1701.00049.
- [59] S. Pasterski and S.-H. Shao, Conformal basis for flat space amplitudes, Phys. Rev. D 96 no. 6, (2017) 065022, arXiv:1705.01027.
- [60] S. Pasterski, S.-H. Shao, and A. Strominger, Gluon Amplitudes as 2d Conformal Correlators, Phys. Rev. D 96 no. 8, (2017) 085006, arXiv:1706.03917.
- [61] S. He, P. Mao, and X.-C. Mao, TT⁻ deformed soft theorem, Phys. Rev. D 107 no. 10, (2023) L101901, arXiv:2209.01953.
- [62] I. Bandos, K. Lechner, D. Sorokin, and P. K. Townsend, A non-linear duality-invariant conformal extension of Maxwell's equations, Phys. Rev. D 102 (2020) 121703, arXiv:2007.09092.
- [63] R. Conti, L. Iannella, S. Negro, and R. Tateo, Generalised Born-Infeld models, Lax operators and the TT perturbation, JHEP 11 (2018) 007, arXiv:1806.11515.
- [64] H. Babaei-Aghbolagh, K. Babaei Velni, D. M. Yekta, and H. Mohammadzadeh, TT-like flows in non-linear electrodynamic theories and S-duality, JHEP 04 (2021) 187, arXiv:2012.13636.
- [65] C. Ferko, L. Smith, and G. Tartaglino-Mazzucchelli, On Current-Squared Flows and ModMax Theories, SciPost Phys. 13 no. 2, (2022) 012, arXiv:2203.01085.
- [66] M. K. Gaillard and B. Zumino, Duality rotations for interacting fields, Nucl. Phys. B 193 (1981) 221–244.
- [67] M. K. Gaillard and B. Zumino, Nonlinear electromagnetic selfduality and legendre transformations,. [arXiv:hep-th/9712103 [hep-th]].

- [68] M. Henneaux and C. Teitelboim, Dynamics of chiral (self-dual) p-forms, Phys. Lett. B 206 (1988) 650.
- [69] P. Pasti, D. P. Sorokin, and M. Tonin, Duality symmetric actions with manifest space-time symmetries, Phys. Rev. D 52 (1995) R4277–R4281. [arXiv:hep-th/9506109 [hep-th]].
- [70] P. Pasti, D. P. Sorokin, and M. Tonin, Space-time symmetries in duality symmetric models,. [arXiv:hep-th/9509052 [hep-th]].
- [71] P. Pasti, D. P. Sorokin, and M. Tonin, On lorentz invariant actions for chiral p-forms, Phys. Rev. D 55 (1997) 6292–6298. [arXiv:hep-th/9611100 [hep-th]].
- [72] P. Pasti, D. P. Sorokin, and M. Tonin, Covariant action for a d = 11 five-brane with the chiral field, Phys. Lett. B 398 (1997) 41–46. [arXiv:hep-th/9701037 [hep-th]].
- [73] H. Babaei-Aghbolagh, S. He, and H. Ouyang, Generalized TT-like flows for scalar theories in two dimensions, arXiv:2501.14583.
- [74] M. Guica, TT deformations and holography, https://indico.cern.ch/event/857396/contributions/3706292/attachments/2036750/34103
- [75] L. Apolo, Single-trace $T\overline{T}$ deformations and string theory, https://lui-apolo.github.io/.
- [76] O. Aharony and T. Vaknin, The TT* deformation at large central charge, JHEP 05 (2018) 166, arXiv:1803.00100.
- [77] E. A. Mazenc, V. Shyam, and R. M. Soni, A TT Deformation for Curved Spacetimes from 3d Gravity, arXiv:1912.09179.
- [78] K.-S. Lee, P. Yi, and J. Yoon, TT-deformed fermionic theories revisited, JHEP 07 (2021) 217, arXiv:2104.09529.
- [79] A. Dey and A. Fortinsky, Perturbative renormalization of the TT-deformed free massive Dirac fermion, JHEP 12 (2021) 200, arXiv:2109.10525.
- [80] M. Baggio, A. Sfondrini, G. Tartaglino-Mazzucchelli, and H. Walsh, On TT deformations and supersymmetry, JHEP 06 (2019) 063, arXiv:1811.00533.
- [81] C.-K. Chang, C. Ferko, and S. Sethi, Supersymmetry and $T\overline{T}$ deformations, JHEP 04 (2019) 131, arXiv:1811.01895.

- [82] H. Jiang, A. Sfondrini, and G. Tartaglino-Mazzucchelli, $T\bar{T}$ deformations with $\mathcal{N} = (0, 2)$ supersymmetry, Phys. Rev. D 100 no. 4, (2019) 046017, arXiv:1904.04760.
- [83] C.-K. Chang, C. Ferko, S. Sethi, A. Sfondrini, and G. Tartaglino-Mazzucchelli, TT flows and (2,2) supersymmetry, Phys. Rev. D 101 no. 2, (2020) 026008, arXiv:1906.00467.
- [84] E. A. Coleman, J. Aguilera-Damia, D. Z. Freedman, and R. M. Soni, TT -deformed actions and (1,1) supersymmetry, JHEP 10 (2019) 080, arXiv:1906.05439.
- [85] H. Jiang and G. Tartaglino-Mazzucchelli, Supersymmetric $J\overline{T}$ and $T\overline{J}$ deformations, JHEP 05 (2020) 140, arXiv:1911.05631.
- [86] C. Ferko, H. Jiang, S. Sethi, and G. Tartaglino-Mazzucchelli, Non-linear supersymmetry and TT-like flows, JHEP 02 (2020) 016, arXiv:1910.01599.
- [87] S. Ebert, H.-Y. Sun, and Z. Sun, TT deformation in SCFTs and integrable supersymmetric theories, JHEP 09 (2021) 082, arXiv:2011.07618.
- [88] C. Ferko, Supersymmetry and Irrelevant Deformations. PhD thesis, Chicago U., Chicago U., 2021. arXiv:2112.14647.
- [89] K.-S. Lee and J. Yoon, TT⁻ deformation of N=(1,1) off-shell supersymmetry and partially broken supersymmetry, Phys. Rev. D 110 no. 2, (2024) 025001, arXiv:2306.08030.
- [90] J. Hou, $T\overline{T}$ flow as characteristic flows, JHEP 03 (2023) 243, arXiv:2208.05391.
- [91] R. Conti, S. Negro, and R. Tateo, The TT perturbation and its geometric interpretation, JHEP 02 (2019) 085, arXiv:1809.09593.
- [92] P. Caputa, S. Datta, Y. Jiang, and P. Kraus, Geometrizing TT, JHEP 03 (2021) 140, arXiv:2011.04664. [Erratum: JHEP 09, 110 (2022)].
- [93] P. Ceschin, R. Conti, and R. Tateo, TT-deformed nonlinear Schrödinger, JHEP 04 (2021) 121, arXiv:2012.12760.
- [94] R. Conti, S. Negro, and R. Tateo, Conserved currents and $T\overline{T}_s$ irrelevant deformations of 2D integrable field theories, JHEP 11 (2019) 120, arXiv:1904.09141.
- [95] L. C. Evans, Partial Differential Equations. American Mathematical Society, 2010.
- [96] A. Cavaglia, S. Negro, P. Szepietowski, and R. Tateo, TT-deformed 2D Quantum Field Theories, JHEP 2016 (2016) 112.

- [97] J. Cardy, The $T\overline{T}$ deformation of quantum field theory as random geometry, JHEP 10 (2018) 186, arXiv:1801.06895.
- [98] M. Guica and R. Monten, Infinite pseudo-conformal symmetries of classical $T\bar{T}$, $J\bar{T}$ and JT_a deformed CFTs, SciPost Phys. 11 no. 4, (2021) 078, arXiv:2011.05445.
- [99] G. Jorjadze and S. Theisen, Canonical maps and integrability in $T\bar{T}$ deformed 2d CFTs, arXiv:2001.03563.
- [100] S. Georgescu and M. Guica, Infinite TT-like symmetries of compactified LST, SciPost Phys. 16 no. 1, (2024) 006, arXiv:2212.09768.
- [101] M. Guica, R. Monten, and I. Tsiares, Classical and quantum symmetries of $T\bar{T}$ -deformed CFTs, arXiv:2212.14014.
- [102] D. Simmons-Duffin, The Conformal Bootstrap, in Theoretical Advanced Study Institute in Elementary Particle Physics: New Frontiers in Fields and Strings, pp. 1–74. 2017. arXiv:1602.07982.
- [103] D. Poland, S. Rychkov, and A. Vichi, The Conformal Bootstrap: Theory, Numerical Techniques, and Applications, Rev. Mod. Phys. 91 (2019) 015002, arXiv:1805.04405.
- [104] M. Guica, A definition of primary operators in JT-deformed CFTs, SciPost Phys. 13 no. 3, (2022) 045, arXiv:2112.14736.
- [105] L. Apolo, W. Song, and B. Yu, On the universal behavior of TT-deformed CFTs: single and double-trace partition functions at large c, JHEP 05 (2023) 210, arXiv:2301.04153.
- [106] S. Datta and Y. Jiang, $T\overline{T}$ deformed partition functions, JHEP 08 (2018) 106, arXiv:1806.07426.
- [107] J. Tian, T. Lai, and F. Omidi, Modular transformations of on-shell actions of (root-)TT deformed holographic CFTs, arXiv:2404.16354.
- [108] J. Gu, Y. Jiang, and H. Wang, Resurgence of TT-deformed Partition Function, arXiv:2410.19633.
- [109] A. Hashimoto and D. Kutasov, $T\overline{T}, J\overline{T}, T\overline{J}$ partition sums from string theory, JHEP 02 (2020) 080, arXiv:1907.07221.
- [110] V. Rosenhaus and M. Smolkin, Integrability and renormalization under $T\overline{T}$, Phys. Rev. D 102 no. 6, (2020) 065009, arXiv:1909.02640.

- [111] M. Guica, On correlation functions in JT-deformed CFTs,
 J. Phys. A 52 no. 18, (2019) 184003, arXiv:1902.01434.
- [112] W. Cui, H. Shu, W. Song, and J. Wang, Correlation functions in the $TsT/T\overline{T}$ correspondence, JHEP 04 (2024) 017, arXiv:2304.04684.
- [113] S. He and H. Shu, Correlation functions, entanglement and chaos in the $T\overline{T}/J\overline{T}$ -deformed CFTs, JHEP 02 (2020) 088, arXiv:1907.12603.
- [114] S. He, Note on higher-point correlation functions of the TT or JT deformed CFTs, Sci. China Phys. Mech. Astron. 64 no. 9, (2021) 291011, arXiv:2012.06202.
- [115] S. He, J.-R. Sun, and Y. Sun, The correlation function of (1,1) and (2,2) supersymmetric theories with $T\bar{T}$ deformation, JHEP 04 (2020) 100, arXiv:1912.11461.
- [116] S. He and Y. Sun, Correlation functions of CFTs on a torus with a $T\overline{T}$ deformation, Phys. Rev. D 102 no. 2, (2020) 026023, arXiv:2004.07486.
- [117] N. I. Akhiezer, Elements of the Theory of Elliptic Functions. American Mathematical Society, Providence, 1990.
- [118] P. Di Francesco, P. Mathieu, and D. Senechal, Conformal Field Theory. Graduate Texts in Contemporary Physics. Springer-Verlag, New York, 1997.
- [119] S. Hirano and M. Shigemori, Conformal field theory on TT-deformed space and correlators from dynamical coordinate transformations, JHEP 07 (2024) 190, arXiv:2402.08278.
- [120] S. He, Y. Sun, and Y.-X. Zhang, TT-flow effects on torus partition functions, JHEP 09 (2021) 061, arXiv:2011.02902.
- [121] V. V. Bazhanov, S. L. Lukyanov, and A. B. Zamolodchikov, Integrable structure of conformal field theory, quantum KdV theory and thermodynamic Bethe ansatz, Commun. Math. Phys. 177 (1996) 381–398, arXiv:hep-th/9412229.
- [122] B. Le Floch and M. Mezei, KdV charges in $T\bar{T}$ theories and new models with super-Hagedorn behavior, SciPost Phys. 7 no. 4, (2019) 043, arXiv:1907.02516.
- [123] M. Asrat, KdV charges and the generalized torus partition sum in $T\overline{T}$ deformation, Nucl. Phys. B 958 (2020) 115119, arXiv:2002.04824.
- [124] S. He, Y. Sun, and J. Yin, A systematic approach to correlators in $T\bar{T}$ deformed CFTs, arXiv:2310.20516.

- [125] N. Barel, Correlation Functions in TT-deformed Theories on the Torus, arXiv:2407.15090.
- [126] S. Hirano, T. Nakajima, and M. Shigemori, TT Deformation of stress-tensor correlators from random geometry, JHEP 04 (2021) 270, arXiv:2012.03972.
- [127] M. Asrat, A. Giveon, N. Itzhaki, and D. Kutasov, Holography Beyond AdS, Nucl. Phys. B 932 (2018) 241–253, arXiv:1711.02690.
- [128] J. Kruthoff and O. Parrikar, On the flow of states under $T\overline{T}$, arXiv:2006.03054.
- [129] E. Tsolakidis, Massive gravity generalization of $T\overline{T}$ deformations, arXiv:2405.07967.
- [130] E. Witten, Aps medal for exceptional achievement in research: Invited article on entanglement properties of quantum field theory, Rev. Mod. Phys. 90 no. 4, (2018) 045003, arXiv:1803.04993.
- [131] E. Witten, Why Does Quantum Field Theory in Curved Spacetime Make Sense? And What Happens to the Algebra of Observables in the Thermodynamic Limit?, pp. 241–284. Springer International Publishing, Cham, 2022. https://doi.org/10.1007/978-3-031-17523-7_11.
- [132] W. Donnelly and V. Shyam, Entanglement entropy and $T\overline{T}$ deformation, Phys. Rev. Lett. 121 no. 13, (2018) 131602, arXiv:1806.07444.
- [133] A. Banerjee, A. Bhattacharyya, and S. Chakraborty, Entanglement Entropy for TT deformed CFT in general dimensions, Nucl. Phys. B 948 (2019) 114775, arXiv:1904.00716.
- [134] T. Ota, Comments on holographic entanglements in cutoff AdS, arXiv:1904.06930.
- [135] J.-C. Chang, S. He, Y.-X. Liu, and L. Zhao, The holographic TT deformation of the entanglement entropy in (A)dS₃/CFT₂, arXiv:2409.08198.
- [136] H.-S. Jeong, K.-Y. Kim, and M. Nishida, Entanglement and Rényi entropy of multiple intervals in TT-deformed CFT and holography, Phys. Rev. D 100 no. 10, (2019) 106015, arXiv:1906.03894.
- [137] M. He and Y. Sun, Holographic entanglement entropy in TT-deformed AdS3, Nucl. Phys. B 990 (2023) 116190, arXiv:2301.04435.
- [138] B. Chen, L. Chen, and P.-X. Hao, Entanglement entropy in TT-deformed CFT, Phys. Rev. D 98 no. 8, (2018) 086025, arXiv:1807.08293.

- [139] D. Basu, Lavish, and B. Paul, Entanglement negativity in TT⁻-deformed CFT2s, Phys. Rev. D 107 no. 12, (2023) 126026, arXiv:2302.11435.
- [140] D. Basu, S. Biswas, A. Dey, B. Paul, and G. Sengupta, Odd entanglement entropy in TT⁻ deformed CFT2s and holography, Phys. Rev. D 108 no. 12, (2023) 126013, arXiv:2307.04832.
- [141] D. Basu and V. Raj, Reflected entropy and timelike entanglement in TT⁻-deformed CFT2s, Phys. Rev. D 110 no. 4, (2024) 046009, arXiv:2402.07253.
- [142] D. Basu and S. Biswas, Entanglement, TT and rotating black holes, arXiv:2410.06363.
- [143] R. M. Soni and A. C. Wall, A New Covariant Entropy Bound from Cauchy Slice Holography, arXiv:2407.16769.
- [144] S. He, Z.-C. Liu, and Y. Sun, Entanglement entropy and modular Hamiltonian of free fermion with deformations on a torus, JHEP 09 (2022) 247, arXiv:2207.06308.
- [145] S. Ashkenazi, S. Chakraborty, Z. Ma, and T. Shachar, Linear response of entanglement entropy to $T\overline{T}$ in massive QFTs, JHEP 04 (2023) 077, arXiv:2302.06688.
- [146] T. Nishioka, T. Takayanagi, and Y. Taki, Topological pseudo entropy, JHEP 09 (2021) 015, arXiv:2107.01797.
- [147] W.-z. Guo, S. He, and Y.-X. Zhang, On the real-time evolution of pseudo-entropy in 2d CFTs, JHEP 09 (2022) 094, arXiv:2206.11818.
- [148] S. He, J. Yang, Y.-X. Zhang, and Z.-X. Zhao, Pseudoentropy for descendant operators in two-dimensional conformal field theories, Phys. Rev. D 109 no. 2, (2024) 025014, arXiv:2301.04891.
- [149] Y. Nakata, T. Takayanagi, Y. Taki, K. Tamaoka, and Z. Wei, New holographic generalization of entanglement entropy, Phys. Rev. D 103 no. 2, (2021) 026005, arXiv:2005.13801.
- [150] S. He, J. Yang, Y.-X. Zhang, and Z.-X. Zhao, Pseudo entropy of primary operators in TT/JT-deformed CFTs, JHEP 09 (2023) 025, arXiv:2305.10984.
- [151] M. He, J. Hou, and Y. Jiang, TT-deformed entanglement entropy for IQFT, JHEP 03 (2024) 056, arXiv:2306.07784.
- [152] S. S. Gubser and I. R. Klebanov, A Universal result on central charges in the presence of double trace deformations, Nucl. Phys. B 656 (2003) 23-36, arXiv:hep-th/0212138.

- [153] A. Bzowski and M. Guica, The holographic interpretation of JT-deformed CFTs, JHEP 01 (2019) 198, arXiv:1803.09753.
- [154] S. Elitzur, A. Giveon, M. Porrati, and E. Rabinovici, Multitrace deformations of vector and adjoint theories and their holographic duals, JHEP 02 (2006) 006, arXiv:hep-th/0511061.
- [155] I. Papadimitriou, Multi-Trace Deformations in AdS/CFT: Exploring the Vacuum Structure of the Deformed CFT, JHEP 05 (2007) 075, arXiv:hep-th/0703152.
- [156] Y. Li, Comments on large central charge $T\bar{T}$ deformed conformal field theory and cutoff AdS holography, arXiv:2012.14414.
- [157] L. Apolo, P.-X. Hao, W.-X. Lai, and W. Song, Glue-on AdS holography for TT-deformed CFTs, JHEP 06 (2023) 117, arXiv:2303.04836.
- [158] E. Witten, Multitrace operators, boundary conditions, and AdS / CFT correspondence, arXiv:hep-th/0112258.
- [159] M. Berkooz, A. Sever, and A. Shomer, 'Double trace' deformations, boundary conditions and space-time singularities, JHEP 05 (2002) 034, arXiv:hep-th/0112264.
- [160] D. Marolf and S. F. Ross, Boundary Conditions and New Dualities: Vector Fields in AdS/CFT, JHEP 11 (2006) 085, arXiv:hep-th/0606113.
- [161] C. Murdia, Y. Nomura, P. Rath, and N. Salzetta, Comments on holographic entanglement entropy in TT deformed conformal field theories, Phys. Rev. D 100 no. 2, (2019) 026011, arXiv:1904.04408.
- [162] S. Grieninger, Entanglement entropy and $T\overline{T}$ deformations beyond antipodal points from holography, JHEP 11 (2019) 171, arXiv:1908.10372.
- [163] A. Lewkowycz, J. Liu, E. Silverstein, and G. Torroba, TT and EE, with implications for (A)dS subregion encodings, JHEP 04 (2020) 152, arXiv:1909.13808.
- [164] H. Geng, Some Information Theoretic Aspects of De-Sitter Holography, JHEP 02 (2020) 005, arXiv:1911.02644.
- [165] W. Donnelly, E. LePage, Y.-Y. Li, A. Pereira, and V. Shyam, Quantum corrections to finite radius holography and holographic entanglement entropy, JHEP 05 (2020) 006, arXiv:1909.11402.
- [166] I. Heemskerk and J. Polchinski, Holographic and Wilsonian Renormalization Groups, JHEP 06 (2011) 031, arXiv:1010.1264.

- [167] T. Faulkner, H. Liu, and M. Rangamani, Integrating out geometry: Holographic
 Wilsonian RG and the membrane paradigm, JHEP 08 (2011) 051, arXiv:1010.4036.
- [168] J. M. Maldacena, The Large N limit of superconformal field theories and supergravity, Adv. Theor. Math. Phys. 2 (1998) 231-252, arXiv:hep-th/9711200.
- [169] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, Gauge theory correlators from noncritical string theory, Phys. Lett. B 428 (1998) 105–114, arXiv:hep-th/9802109.
- [170] E. Witten, Anti-de Sitter space and holography, Adv. Theor. Math. Phys. 2 (1998) 253-291, arXiv:hep-th/9802150.
- [171] C. Fefferman and C. R. Graham, Conformal invariants, Astérisque S131 (1985) 95–116.
- [172] C. R. Graham and J. M. Lee, Einstein metrics with prescribed conformal infinity on the ball, Adv. Math. 87 no. 2, (1991) 186–225.
- [173] K. Skenderis and S. N. Solodukhin, Quantum effective action from the AdS / CFT correspondence, Phys. Lett. B 472 (2000) 316-322, arXiv:hep-th/9910023.
- [174] V. Balasubramanian and P. Kraus, A Stress tensor for Anti-de Sitter gravity, Commun. Math. Phys. 208 (1999) 413-428, arXiv:hep-th/9902121.
- [175] R. Emparan, C. V. Johnson, and R. C. Myers, Surface terms as counterterms in the AdS / CFT correspondence, Phys. Rev. D 60 (1999) 104001, arXiv:hep-th/9903238.
- [176] S. de Haro, S. N. Solodukhin, and K. Skenderis, Holographic reconstruction of space-time and renormalization in the AdS / CFT correspondence, Commun. Math. Phys. 217 (2001) 595–622, arXiv:hep-th/0002230.
- [177] M. Henningson and K. Skenderis, The Holographic Weyl anomaly, JHEP 07 (1998) 023, arXiv:hep-th/9806087.
- [178] J. D. Brown and J. W. York, Jr., Quasilocal energy and conserved charges derived from the gravitational action, Phys. Rev. D 47 (1993) 1407–1419, arXiv:gr-qc/9209012.
- [179] J. D. Brown and M. Henneaux, Central Charges in the Canonical Realization of Asymptotic Symmetries: An Example from Three-Dimensional Gravity, Commun. Math. Phys. 104 (1986) 207–226.
- [180] S. M. Carroll, Spacetime and Geometry: An Introduction to General Relativity. Cambridge University Press, 7, 2019.

- [181] S. Ryu and T. Takayanagi, Holographic derivation of entanglement entropy from AdS/CFT, Phys. Rev. Lett. 96 (2006) 181602, arXiv:hep-th/0603001.
- [182] T. Kawamoto, S.-M. Ruan, and T. Takayanagia, Gluing AdS/CFT, JHEP 07 (2023) 080, arXiv:2303.01247.
- [183] T. Kawamoto, R. Maeda, N. Nakamura, and T. Takayanagi, Traversable AdS Wormhole via Non-local Double Trace or Janus Deformation, arXiv:2502.03531.
- [184] L. Apolo, P.-X. Hao, W.-X. Lai, and W. Song, Extremal surfaces in glue-on AdS/TT holography, JHEP 01 (2024) 054, arXiv:2311.04883.
- [185] A. Lewkowycz and J. Maldacena, Generalized gravitational entropy, JHEP 08 (2013) 090, arXiv:1304.4926.
- [186] X. Dong, A. Lewkowycz, and M. Rangamani, Deriving covariant holographic entanglement, JHEP 11 (2016) 028, arXiv:1607.07506.
- [187] M. Srednicki, Entropy and area, Phys. Rev. Lett. 71 (1993) 666-669, arXiv:hep-th/9303048.
- [188] S. He, Y. Li, Y.-Z. Li, and Y. Zhang, Note on holographic torus stress tensor correlators in AdS₃ gravity, arXiv:2405.01255.
- [189] Y.-Z. Li, Y. Xie, and S. He, Holographic Correlators of Boundary/Crosscap CFTs in Two Dimensions, arXiv:2501.18386.
- [190] P. Kraus, R. Monten, and R. M. Myers, 3D Gravity in a Box, SciPost Phys. 11 (2021) 070, arXiv:2103.13398.
- [191] M. He, S. He, and Y.-h. Gao, Surface charges in Chern-Simons gravity with TT deformation, JHEP 03 (2022) 044, arXiv:2109.12885.
- [192] V. I. Arnold, Mathematical Methods of Classical Mechanics. Graduate Texts in Mathematics. Springer, 1989.
- [193] R. Abraham and J. E. Marsden, Foundations of Mechanics. Addison-Wesley, second ed., 1994.
- [194] C. Crnkovic and E. Witten, Covariant description of canonical formalism in geometrical theories., in Three Hundred Years of Gravitation, S. W. Hawking and W. Israel, eds., pp. 676–684. 9, 1986.

- [195] S. Ebert, E. Hijano, P. Kraus, R. Monten, and R. M. Myers, Field Theory of Interacting Boundary Gravitons, SciPost Phys. 13 no. 2, (2022) 038, arXiv:2201.01780.
- [196] P. Kraus, R. Monten, and K. Roumpedakis, Refining the cutoff 3d gravity/ $T\overline{T}$ correspondence, JHEP 10 (2022) 094, arXiv:2206.00674.
- [197] V. Shyam, $T\overline{T} + \Lambda_2$ deformed CFT on the stretched dS₃ horizon, JHEP 04 (2022) 052, arXiv:2106.10227.
- [198] E. Coleman, E. A. Mazenc, V. Shyam, E. Silverstein, R. M. Soni, G. Torroba, and S. Yang, De Sitter microstates from TT + Λ₂ and the Hawking-Page transition, JHEP 07 (2022) 140, arXiv:2110.14670.
- [199] G. Torroba, $T\overline{T} + \Lambda_2$ from a 2d gravity path integral, JHEP 01 (2023) 163, arXiv:2212.04512.
- [200] G. Batra, G. B. De Luca, E. Silverstein, G. Torroba, and S. Yang, Bulk-local dS₃ holography: the matter with $T\overline{T} + \Lambda_2$, JHEP 10 (2024) 072, arXiv:2403.01040.
- [201] S. E. Aguilar-Gutierrez, A. Svesko, and M. R. Visser, TT deformations from AdS₂ to dS₂, JHEP 01 (2025) 120, arXiv:2410.18257.
- [202] N. Seiberg and E. Witten, The D1 / D5 system and singular CFT, JHEP 04 (1999) 017, arXiv:hep-th/9903224.
- [203] S. Chakraborty, $\frac{\mathrm{SL}(2,\mathbb{R})\times\mathrm{U}(1)}{\mathrm{U}(1)}$ CFT, NS5+F1 system and single trace $T\overline{T}$, JHEP 03 (2021) 113, arXiv:2012.03995.
- [204] B. Knighton, Deriving the long-string CFT in AdS₃, arXiv:2410.16904.
- [205] J. M. Maldacena and H. Ooguri, Strings in AdS(3) and SL(2,R) WZW model 1.: The Spectrum, J. Math. Phys. 42 (2001) 2929–2960, arXiv:hep-th/0001053.
- [206] D. Kutasov and N. Seiberg, More comments on string theory on AdS(3), JHEP 04 (1999) 008, arXiv:hep-th/9903219.
- [207] S. Chakraborty, A. Giveon, and D. Kutasov, TT, black holes and negative strings, JHEP 09 (2020) 057, arXiv:2006.13249.
- [208] R. Argurio, A. Giveon, and A. Shomer, Superstrings on AdS(3) and symmetric products, JHEP 12 (2000) 003, arXiv:hep-th/0009242.
- [209] B. Pozsgay, Y. Jiang, and G. Takács, TT-deformation and long range spin chains, JHEP 03 (2020) 092, arXiv:1911.11118.

- [210] L. Eberhardt, M. R. Gaberdiel, and R. Gopakumar, The Worldsheet Dual of the Symmetric Product CFT, JHEP 04 (2019) 103, arXiv:1812.01007.
- [211] L. Eberhardt, M. R. Gaberdiel, and R. Gopakumar, Deriving the AdS₃/CFT₂ correspondence, JHEP 02 (2020) 136, arXiv:1911.00378.
- [212] A. Dei, B. Knighton, K. Naderi, and S. Sethi, Tensionless AdS_3/CFT_2 and Single Trace $T\overline{T}$, arXiv:2408.00823.
- [213] O. Aharony, M. Berkooz, D. Kutasov, and N. Seiberg, Linear dilatons, NS five-branes and holography, JHEP 10 (1998) 004, arXiv:hep-th/9808149.
- [214] A. Giveon, N. Itzhaki, and D. Kutasov, A solvable irrelevant deformation of AdS₃/CFT₂, JHEP 12 (2017) 155, arXiv:1707.05800.
- [215] A. Giveon, M. Porrati, and E. Rabinovici, Target space duality in string theory, Phys. Rept. 244 (1994) 77–202, arXiv:hep-th/9401139.
- [216] S. Frolov, Lax pair for strings in Lunin-Maldacena background, JHEP 05 (2005) 069, arXiv:hep-th/0503201.
- [217] D. Orlando, S. Reffert, Y. Sekiguchi, and K. Yoshida, O(d,d) transformations preserve classical integrability, Nucl. Phys. B 950 (2020) 114880, arXiv:1907.03759.
- [218] O. Lunin and J. M. Maldacena, Deforming field theories with U(1) x U(1) global symmetry and their gravity duals, JHEP 05 (2005) 033, arXiv:hep-th/0502086.
- [219] A. Giveon, 2pf in single-trace $T\overline{T}$ holography, JHEP 10 (2023) 112, arXiv:2309.15629.
- [220] G. Giribet, TT-deformations, AdS/CFT and correlation functions, JHEP 02 (2018) 114, arXiv:1711.02716.
- [221] J. Cardy, $T\overline{T}$ deformations of non-Lorentz invariant field theories, arXiv:1809.07849.
- [222] C. D. A. Blair, Non-relativistic duality and $T\bar{T}$ deformations, JHEP 07 (2020) 069, arXiv:2002.12413.
- [223] D. Hansen, Y. Jiang, and J. Xu, Geometrizing non-relativistic bilinear deformations, JHEP 04 (2021) 186, arXiv:2012.12290.
- [224] B. Chen, J. Hou, and J. Tian, Note on the nonrelativistic $T\overline{T}$ deformation, Phys. Rev. D 104 no. 2, (2021) 025004, arXiv:2012.14091.

- [225] S. Demulder, S. Driezen, B. Knighton, G. Oling, A. L. Retore, F. K. Seibold, A. Sfondrini, and Z. Yan, Exact approaches on the string worldsheet, arXiv:2312.12930.
- [226] J. Cardy and B. Doyon, $T\overline{T}$ deformations and the width of fundamental particles, JHEP 04 (2022) 136, arXiv:2010.15733.
- [227] Y. Jiang, TT-deformed 1d Bose gas, SciPost Phys. 12 no. 6, (2022) 191, arXiv:2011.00637.
- [228] D. Pavshinkin, TT deformation of the Calogero–Sutherland model via dimensional reduction, Theor. Math. Phys. 217 no. 2, (2023) 1726–1742, arXiv:2111.12080.
- [229] B. Doyon, F. Hübner, and T. Yoshimura, New Classical Integrable Systems from Generalized TT⁻-Deformations, Phys. Rev. Lett. 132 no. 25, (2024) 251602, arXiv:2311.06369.
- [230] B. Doyon, F. Hübner, and T. Yoshimura, Generalised TT-deformations of classical free particles, arXiv:2312.14855.
- [231] T. Bargheer, N. Beisert, and F. Loebbert, Boosting Nearest-Neighbour to Long-Range Integrable Spin Chains, J. Stat. Mech. 0811 (2008) L11001, arXiv:0807.5081.
- [232] E. Marchetto, A. Sfondrini, and Z. Yang, TT Deformations and Integrable Spin Chains, Phys. Rev. Lett. 124 no. 10, (2020) 100601, arXiv:1911.12315.
- [233] B. Pozsgay, T. Gombor, and A. Hutsalyuk, Integrable hard-rod deformation of the Heisenberg spin chains, Phys. Rev. E 104 no. 6, (2021) 064124, arXiv:2108.13724.
- [234] Y. Jiang, TT Deformation: A Lattice Approach, Symmetry 15 no. 12, (2023) 2212, arXiv:2312.12078.
- [235] D. J. Gross, J. Kruthoff, A. Rolph, and E. Shaghoulian, Hamiltonian deformations in quantum mechanics, $T\bar{T}$, and the SYK model, Phys. Rev. D 102 no. 4, (2020) 046019, arXiv:1912.06132.
- [236] S. He and Z.-Y. Xian, TT⁻ deformation on multiquantum mechanics and regenesis, Phys. Rev. D 106 no. 4, (2022) 046002, arXiv:2104.03852.
- [237] K. Jensen, Chaos in AdS₂ Holography, Phys. Rev. Lett. 117 no. 11, (2016) 111601, arXiv:1605.06098.

- [238] J. Maldacena, D. Stanford, and Z. Yang, Conformal symmetry and its breaking in two dimensional Nearly Anti-de-Sitter space, PTEP 2016 no. 12, (2016) 12C104, arXiv:1606.01857.
- [239] J. Engelsöy, T. G. Mertens, and H. Verlinde, An investigation of AdS₂ backreaction and holography, JHEP 07 (2016) 139, arXiv:1606.03438.
- [240] P. Saad, S. H. Shenker, and D. Stanford, JT gravity as a matrix integral, arXiv:1903.11115.
- [241] F. Rosso, TT deformation of random matrices,
 Phys. Rev. D 103 no. 12, (2021) 126017, arXiv:2012.11714.
- [242] S. Ebert, H.-Y. Sun, and Z. Sun, *TT*-deformed free energy of the Airy model, JHEP 08 (2022) 026, arXiv:2202.03454.
- [243] M. Guica, An integrable Lorentz-breaking deformation of two-dimensional CFTs, SciPost Phys. 5 no. 5, (2018) 048, arXiv:1710.08415.
- [244] L. Apolo and W. Song, Strings on warped AdS₃ via TJ deformations, JHEP 10 (2018) 165, arXiv:1806.10127.
- [245] S. Chakraborty, A. Giveon, and D. Kutasov, $J\overline{T}$ deformed CFT₂ and string theory, JHEP 10 (2018) 057, arXiv:1806.09667.
- [246] O. Aharony, S. Datta, A. Giveon, Y. Jiang, and D. Kutasov, Modular covariance and uniqueness of JT deformed CFTs, JHEP 01 (2019) 085, arXiv:1808.08978.
- [247] B. Le Floch and M. Mezei, Solving a family of $T\bar{T}$ -like theories, arXiv:1903.07606.
- [248] G. Hernández-Chifflet, S. Negro, and A. Sfondrini, Flow Equations for Generalized TT Deformations, Phys. Rev. Lett. 124 no. 20, (2020) 200601, arXiv:1911.12233.
- [249] P. Rodríguez, D. Tempo, and R. Troncoso, Mapping relativistic to ultra/non-relativistic conformal symmetries in 2D and finite $\sqrt{T\overline{T}}$ deformations, JHEP 11 (2021) 133, arXiv:2106.09750.
- [250] S. He and X.-C. Mao, Irrelevant and marginal deformed BMS field theories, JHEP 04 (2024) 138, arXiv:2401.09991.
- [251] S. Ebert, C. Ferko, and Z. Sun, Root-TT⁻ deformed boundary conditions in holography, Phys. Rev. D 107 no. 12, (2023) 126022, arXiv:2304.08723.

- [252] S. Ebert, C. Ferko, C. L. Martin, and G. Tartaglino-Mazzucchelli, Flows in the space of interacting chiral boson theories, Phys. Rev. D 110 no. 4, (2024) 046005, arXiv:2403.18242.
- [253] C. Ferko, Y. Hu, Z. Huang, K. Koutrolikos, and G. Tartaglino-Mazzucchelli, TT-like flows and 3d nonlinear supersymmetry, SciPost Phys. 16 no. 1, (2024) 038, arXiv:2302.10410.
- [254] C. Ferko, J. Hou, T. Morone, G. Tartaglino-Mazzucchelli, and R. Tateo, TT-like Flows of Yang-Mills Theories, arXiv:2409.18740.
- [255] C. D. A. Blair, J. Lahnsteiner, N. A. Obers, and Z. Yan, Matrix theory reloaded: a BPS road to holography, JHEP 02 (2025) 024, arXiv:2410.03591.
- [256] I. Bandos, K. Lechner, D. Sorokin, and P. K. Townsend, On p-form gauge theories and their conformal limits, JHEP 03 (2021) 022, arXiv:2012.09286.
- [257] C. Ferko, S. M. Kuzenko, K. Lechner, D. P. Sorokin, and G. Tartaglino-Mazzucchelli, Interacting chiral form field theories and TT-like flows in six and higher dimensions, JHEP 05 (2024) 320, arXiv:2402.06947.
- [258] C. Ferko and L. Smith, An Infinite Family of Integrable Sigma Models Using Auxiliary Fields, arXiv:2405.05899.
- [259] D. Bielli, C. Ferko, L. Smith, and G. Tartaglino-Mazzucchelli, T-Duality and TT-like Deformations of Sigma Models, arXiv:2407.11636.
- [260] O. Fukushima and K. Yoshida, 4D Chern-Simons theory with auxiliary fields, arXiv:2407.02204.
- [261] D. Bielli, C. Ferko, L. Smith, and G. Tartaglino-Mazzucchelli, Integrable Higher-Spin Deformations of Sigma Models from Auxiliary Fields, arXiv:2407.16338.
- [262] D. Bielli, C. Ferko, L. Smith, and G. Tartaglino-Mazzucchelli, Auxiliary Field Sigma Models and Yang-Baxter Deformations, arXiv:2408.09714.
- [263] H. Babaei-Aghbolagh, K. Babaei Velni, D. M. Yekta, and H. Mohammadzadeh, Manifestly SL(2, R) Duality-Symmetric Forms in ModMax Theory, JHEP 12 (2022) 147, arXiv:2210.13196.
- [264] C. Ferko, S. M. Kuzenko, L. Smith, and G. Tartaglino-Mazzucchelli, Duality-invariant nonlinear electrodynamics and stress tensor flows, Phys. Rev. D 108 no. 10, (2023) 106021, arXiv:2309.04253.

- [265] T. Flöss, D. Roest, and T. Westerdijk, Non-linear electrodynamics from massive gravity, JHEP 02 (2024) 194, arXiv:2308.04349.
- [266] T. Morone, S. Negro, and R. Tateo, Gravity and $T\bar{T}$ flows in higher dimensions, Nucl. Phys. B 1005 (2024) 116605, arXiv:2401.16400.
- [267] T. Morone and R. Tateo, Solutions to the Ricci Flow via Einstein Field Equations, arXiv:2411.10265.
- [268] F. Hao, X.-Y. Ran, and M. Yamada, Geometric realization via irrelevant deformations induced by the stress-energy tensor, arXiv:2410.02537.
- [269] Y.-H. Shi et al., Quantum simulation of Hawking radiation and curved spacetime with a superconducting on-chip black hole, Nature Commun. 14 no. 1, (2023) 3263, arXiv:2111.11092.
- [270] P. Basteiro, F. Dusel, J. Erdmenger, D. Herdt, H. Hinrichsen, R. Meyer, and M. Schrauth, Breitenlohner-Freedman Bound on Hyperbolic Tilings, Phys. Rev. Lett. 130 no. 9, (2023) 091604, arXiv:2205.05081.
- [271] J. Chen et al., AdS/CFT Correspondence in Hyperbolic Lattices, arXiv:2305.04862.
- [272] B. Swingle, Entanglement Renormalization and Holography, Phys. Rev. D 86 (2012) 065007, arXiv:0905.1317.
- [273] S. S. Gubser, J. Knaute, S. Parikh, A. Samberg, and P. Witaszczyk, *p*-adic AdS/CFT, Commun. Math. Phys. 352 no. 3, (2017) 1019–1059, arXiv:1605.01061.
- [274] L. Chen, X. Liu, and L.-Y. Hung, Emergent Einstein Equation in p-adic Conformal Field Theory Tensor Networks, Phys. Rev. Lett. 127 no. 22, (2021) 221602, arXiv:2102.12022.
- [275] L. Chen, K. Ji, H. Zhang, C. Shen, R. Wang, X. Zeng, and L.-Y. Hung, CFTD from TQFTD+1 via Holographic Tensor Network, and Precision Discretization of CFT2, Phys. Rev. X 14 no. 4, (2024) 041033, arXiv:2210.12127.
- [276] G. Cheng, L. Chen, Z.-C. Gu, and L.-Y. Hung, Precision reconstruction of rational CFT from exact fixed point tensor network, arXiv:2311.18005.
- [277] L.-Y. Hung and Y. Jiang, Building up quantum spacetimes with BCFT Legos, arXiv:2404.00877.

- [278] H. Babaei-Aghbolagh, S. He, and H. Ouyang, Generalized TT-like deformations in duality-invariant nonlinear electrodynamic theories, JHEP 09 (2024) 137, arXiv:2407.03698.
- [279] I. Bialynicki-Birula, Nonlinear electrodynamics: Variations on a theme by born and infeld, in Quantum Theory of Particles and Fields: Birthday Volume Dedicated to Jan Lopuszanski, B. Jancewicz and J. Lukierski, eds., pp. 31–48. World Scientific Publishing Co Pte Ltd, 1983.
- [280] I. Bialynicki-Birula, Field theory of photon dust, Acta Phys. Polon. B 23 (1992) 553–559.
- [281] G. W. Gibbons and D. A. Rasheed, Sl(2,r) invariance of nonlinear electrodynamics coupled to an axion and a dilaton, Phys. Lett. B 365 (1996) 46–50. [arXiv:hep-th/9509141 [hep-th]].
- [282] S. Deser, A. Gomberoff, M. Henneaux, and C. Teitelboim, Duality, selfduality, sources and charge quantization in abelian n form theories, Phys. Lett. B 400 (1997) 80–86. [arXiv:hep-th/9702184 [hep-th]].
- [283] S. Deser, A. Gomberoff, M. Henneaux, and C. Teitelboim, P-brane dyons and electric magnetic duality, Nucl. Phys. B 520 (1998) 179–204. [arXiv:hep-th/9712189 [hep-th]].