

# Error Bounds on the Universal Lindblad Equation in the Thermodynamic Limit

Teruhiro Ikeuchi and Takashi Mori

*Department of Physics, Keio University, Hiyoshi, Yokohama, 223-8522, Japan*

It is a central problem in various fields of physics to elucidate the behavior of quantum many-body systems subjected to bulk dissipation. In this context, several microscopic derivations of the Lindblad quantum master equation for many-body systems have been proposed so far. Among them, the universal Lindblad equation derived by Nathan and Rudner is fascinating because it has desired locality and its derivation seems to rely solely on the assumption that the bath correlation time is much shorter than the dissipation time, which is the case in the weak-coupling limit or the singular-coupling limit. However, it remains elusive whether errors due to several approximations in deriving the universal Lindblad equation keep small during the time evolution in the thermodynamic limit. Here, rigorous error bounds on the time evolution of a local quantity are given, and it is shown that, under the assumption of the accelerated dissipation in bulk-dissipated systems, those errors vanish in the weak-coupling limit or the singular-coupling limit after taking the thermodynamic limit.

## I. INTRODUCTION

The quantum Markov process with complete positivity is generally described by a quantum master equation of the Lindblad form [1, 2]. The Lindblad equation has widely been used to describe a small quantum system in contact with thermal reservoirs, especially in quantum optics [3, 4]. Recent interest is shifted to open quantum *many-body* systems in connection with quantum control utilizing dissipation [5–8], noisy quantum computation [9, 10], and nonequilibrium statistical physics [11–15]. The Lindblad equation is also a fundamental tool to study such a many-body system. Recent theoretical studies on open quantum many-body systems are largely triggered by experimental progress, which enables us to introduce tunable dissipation in many-body setups [5–8]. In addition to that, there is also a line of research based on the idea that weak dissipation can be utilized as a probe field to better understand a given quantum many-body system [16, 17].

New insights have been gained through recent theoretical studies on the many-body Lindblad dynamics. In particular, it is recognized that the operator scrambling dynamics [18, 19] has important implications for dissipative dynamics under bulk dissipation. The operator scrambling—or the operator spreading—implies that a local operator becomes highly nonlocal after quantum time evolution in the Heisenberg picture. In general, a nonlocal operator is more fragile to dissipation, and hence the dissipative decay is accelerated due to the operator growth [20], which was numerically confirmed in the decay of the Loschmidt echo fidelity that is faster than exponential [21]. As a consequence of such an amplification of the effect of bulk dissipation, we sometimes have a non-zero decay rate even in the dissipationless limit (the thermodynamic limit is taken first), which is known as the quantum Ruelle-Pollicott resonance [17, 22], or the anomalous relaxation [23, 24].

In this way, fundamental properties of many-body Lindblad equations have been investigated, but their theoretical grounds are not well established. It is highly

desired to give a microscopic derivation of the Lindblad equation and understand under what condition its use is justified. A standard derivation of the Lindblad equation involves using the secular approximation [25] (it is also referred to as the rotating-wave approximation), and the derived equation is known as the Davies equation [26]. The Davies equation is rigorously derived from the unitary time evolution of the system and the environment when the system of interest is finite and the system-bath interaction is infinitesimally weak. However, it is later recognized that the secular approximation is not justified and gives unphysical results in the many-body setup [27, 28]. We have to go beyond the secular approximation to derive the Lindblad equation for open quantum many-body systems.

So far, based on different approximations, various derivations of the Lindblad equation for many-body systems have been proposed [27, 29–35]. Among them, the universal Lindblad equation (ULE) derived by Nathan and Rudner [31, 32] has remarkable feature. Firstly, it has desired locality [31, 35] in contrast to the Davies equation. Secondly, the ULE appears to be solely derived from the assumption that the bath correlation time is much shorter than the dissipation time, which is the case in the weak-coupling limit (the system-bath coupling is infinitesimally weak) [26, 36] or in the singular-coupling limit (the bath correlation time is infinitesimally short) [2, 37, 38]. No assumption on the timescale of the intrinsic dynamics of the system of interest is explicitly used. Thus, it is expected that the derivation safely applies to many-body systems.

Although the existing derivation of the ULE given in Ref. [31] is physically reasonable, it is not rigorous and it is uncertain whether errors induced by several approximations needed to derive the ULE are negligible. Nathan and Rudner [31] evaluated rigorous error bounds on the approximations, but there are two problems:

- (i) Their error bounds diverge in the thermodynamic limit for bulk dissipation (i.e. each site of the system is coupled to its own bath). Therefore, it remains unclear whether those approximations are

justified in macroscopic systems.

- (ii) They evaluated the error *at the level of the equations of motion*. It is not proved that the error does not accumulate over time. Thus, it is uncertain whether the error keeps small during the time evolution.

The aim of this paper is to overcome these two difficulties. We consider a one-dimensional quantum spin system with local Hamiltonian in which each site is coupled to its own free-boson bath, and evaluate rigorous error bounds on the time evolution of a local quantity due to the approximations made in the derivation of the ULE. Under the assumption of the *accelerated dissipation*, which is expected to be a generic property of bulk-dissipated open quantum many-body systems [17, 20, 21], we show that the errors vanish *uniformly in time* when we take the weak-coupling limit or the singular-coupling limit *after taking the thermodynamic limit*. The ULE is therefore justified for arbitrarily long times in the thermodynamic limit.

To overcome the difficulty (i), we prove that the existing Lieb-Robinson bound [39] is applicable to the ULE. By using it, finite error bounds in the thermodynamic limit are obtained. To overcome the difficulty (ii), i.e. to obtain desired time-independent error bounds, we use the assumption of the accelerated dissipation. Roughly speaking, it states that the dissipative decay rate of a local operator in the Heisenberg picture increases with time and is eventually saturated at a value that is much larger than the initial decay rate. As we have already mentioned, it was argued that such an accelerated dissipation generically occurs due to the operator spreading [17, 20, 21]. This is not a mathematically proved fact, but is a reasonable assumption. We give its theoretical picture and numerical evidence under a local Lindblad dynamics.

The plan of the paper is as follows. Section II A introduces the microscopic setup. Section II B gives a summary of the main result presented in this paper. Section II C introduces theoretical methods used in deriving the main result: the Lieb-Robinson bound and the accelerated dissipation. The former is formulated as a rigorous inequality as is explained in Sec. III, whereas the latter is an assumption without mathematical proof. Section IV is devoted to a detailed account of the accelerated dissipation with an intuitive picture and numerical evidences. Section V reviews the microscopic derivation of the ULE following Nathan and Rudner [31]. Various approximations are introduced and the explicit expressions of their errors are given. Section VI derives error bounds on the approximations introduced in Sec. V. We see that the Lieb-Robinson bound and the accelerated dissipation lead to the main result presented in Sec. II B. We conclude with discussion in Sec. VII.




FIG. 1. Setup of a one-dimensional quantum many-body system under bulk dissipation. Each site  $i$  is coupled to an independent free-boson bath at the inverse temperature  $\beta_i$ .

## II. SUMMARY OF THE MAIN RESULT

Before going on to the theoretical detail, we summarize the main result presented in this paper. We first explain the theoretical setup and then our main result and key theoretical methods.

### A. Setup

Let us consider a one-dimensional quantum spin system defined on the set of sites  $\Lambda = \{1, 2, \dots, N\}$ . The result presented in this paper is straightforwardly extended to a  $d$ -dimensional system. The Hamiltonian of the system of interest is denoted by  $\hat{H}_S$ . We assume that  $\hat{H}_S$  is local in the sense that it is written as

$$\hat{H}_S = \sum_{i \in \Lambda} \hat{h}_i, \quad (1)$$

where  $\hat{h}_i$  is an operator acting nontrivially on sites near  $i$ . For simplicity, we assume that  $\{\hat{h}_i\}_{i \in \Lambda}$  are finite range: the diameter of the support of  $\hat{h}_i$  is not greater than a constant  $a$  for any  $i \in \Lambda$ .

In this paper, we focus on a static Hamiltonian, but the following discussion is extended to a time-dependent Hamiltonian  $\hat{H}_S(t)$  by making a slight modification.

We consider bulk dissipation, see Fig. 1 for a schematic picture. Each site  $i$  is coupled to its own bath consisting of free bosons, and hence the Hamiltonian of the baths is written as

$$\hat{H}_B = \sum_{i \in \Lambda} \hat{H}_B^{(i)}, \quad \hat{H}_B^{(i)} = \sum_k \omega_k \hat{b}_{i,k}^\dagger \hat{b}_{i,k}, \quad (2)$$

where  $\hat{b}_{i,k}^\dagger$  ( $\hat{b}_{i,k}$ ) is the creation (annihilation) operator of bosons interacting with the site  $i$ . The interaction between the system and the baths is expressed by

$$\hat{H}_I = \sum_{i \in \Lambda} \hat{A}_i \otimes \hat{B}_i, \quad (3)$$

where

$$\hat{B}_i = \sum_k \left( g_k \hat{b}_{i,k} + g_k^* \hat{b}_{i,k}^\dagger \right) \quad (4)$$

is the bath operator, which is assumed to be linear in  $\hat{b}_{i,k}$  and  $\hat{b}_{i,k}^\dagger$ , and  $\hat{A}_i$  is an Hermitian operator of the system that is normalized as  $\|\hat{A}_i\| = 1$  ( $\|\cdot\|$  is the operator norm). For simplicity, in the following, we assume that  $\hat{A}_i$  is a single-site operator, but it is straightforward to extend the result to an arbitrary local operator  $\hat{A}_i$  that acts nontrivially onto sites near  $i$ . It is also straightforward to multiple dissipation channels at each site  $i$ :  $\hat{H}_I = \sum_{i \in \Lambda} \sum_{m=1}^M \hat{A}_{i,m} \otimes \hat{B}_{i,m}$  as long as  $M$  is finite.

The bath correlation function in the state

$$\rho_B = \frac{\exp\left[-\sum_{i \in \Lambda} \beta_i \hat{H}_B^{(i)}\right]}{\text{Tr}_B \exp\left[-\sum_{i \in \Lambda} \beta_i \hat{H}_B^{(i)}\right]}, \quad (5)$$

where  $\beta_i$  is the inverse temperature of the  $i$ th bath, is defined by

$$\Phi_i(t) = \text{Tr}_B \left[ \hat{B}_i(t) \hat{B}_i(0) \rho_B \right], \quad (6)$$

where

$$\hat{B}_i(t) = e^{i\hat{H}_B t} \hat{B}_i e^{-i\hat{H}_B t}. \quad (7)$$

Without loss of generality, we can assume

$$\text{Tr}_B[\hat{B}_i \rho_B] = 0. \quad (8)$$

The Fourier transform of  $\Phi_i(t)$  is denoted by  $J_i(\omega)$ :

$$J_i(\omega) = \int_{-\infty}^{\infty} \Phi_i(t) e^{-i\omega t} dt. \quad (9)$$

In an infinitely large bath, which we consider in the following analysis,  $J_i(\omega)$  is a continuous function of  $\omega$  and  $\Phi_i(t)$  vanishes in the long-time limit.

Following Nathan and Rudner [31], we introduce a jump correlator  $g_i(t)$  as

$$g_i(t) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \sqrt{J_i(\omega)} e^{i\omega t}, \quad (10)$$

which is equivalent to

$$\Phi_i(t-t') = \int_{-\infty}^{\infty} ds g_i(t-s) g_i(s-t'). \quad (11)$$

We introduce two positive constants  $\gamma$  and  $\tau_B$  satisfying

$$|g_i(t)| \leq \frac{\sqrt{\gamma}}{2\tau_B} e^{-2|t|/\tau_B} \quad (12)$$

for arbitrary  $i \in \Lambda$  and  $t \in \mathbb{R}$ . From Eq. (12), we can derive the following inequality on  $\Phi_i(t)$ :

$$|\Phi_i(t)| \leq \frac{\gamma}{\tau_B} e^{-|t|/\tau_B} \quad \forall i \in \Lambda. \quad (13)$$

Physically,  $\gamma$  and  $\tau_B$  denote the strength of dissipation and the bath correlation time, respectively. It should be remarked that  $\tau_B$  should satisfy  $\tau_B \gtrsim \max_{i \in \Lambda} \beta_i$ , which

is a consequence of the Kubo-Martin-Schwinger condition [40]:  $J_i(\omega) = J_i(-\omega) e^{-\beta_i \omega}$ . The weak-coupling limit corresponds to the limit of  $\gamma \rightarrow +0$  with  $\tau_B$  held fixed, whereas the singular-coupling limit corresponds to the limit of  $\tau_B \rightarrow +0$  with  $\gamma$  held fixed.

Suppose a factorized initial state  $\rho_T(0) = \rho(0) \otimes \rho_B$ , where  $\rho_T(t)$  is the density matrix of the total system and  $\rho(t)$  is the reduced density matrix of the system of interest, which is defined as  $\rho(t) = \text{Tr}_B[\rho_T(t)]$ . We assume that  $\rho_B$  is given by Eq. (5), i.e., each bath is in thermal equilibrium but its temperature can depend on site  $i$ .

The density matrix of the total system obeys the Liouville-von Neumann equation

$$\frac{d\rho_T(t)}{dt} = -i[\hat{H}_T, \rho_T(t)]. \quad (14)$$

Formally,  $\rho_T(t)$  is expressed as

$$\rho_T(t) = \mathcal{U}(t) \rho_T(0), \quad (15)$$

where  $\mathcal{U}(t)(\cdot) = e^{i\hat{H}_T t}(\cdot)e^{-i\hat{H}_T t}$ . The exact reduced density matrix at time  $t$  is thus given by  $\rho(t) = \text{Tr}_B[\mathcal{U}(t)\rho(0) \otimes \rho_B]$ , which is in general non-Markovian. By using various approximations, all of which are expected to be valid as long as  $\gamma\tau_B \ll 1$ , Nathan and Rudner [31] derived the following ULE:

$$\begin{aligned} \frac{d\rho}{dt} &= -i[\hat{H}_S + \hat{\Delta}, \rho] + \sum_{i=1}^N \left( \hat{L}_i \rho \hat{L}_i^\dagger - \frac{1}{2} \{ \hat{L}_i^\dagger \hat{L}_i, \rho \} \right) \\ &=: \mathcal{L}\rho, \end{aligned} \quad (16)$$

where  $\mathcal{L}$  is the Liouvillian of the Lindblad form (it is also referred to as the Lindbladian). Here,

$$\hat{L}_i = \int_{-\infty}^{\infty} ds g_i(s) \hat{A}_i(-s) \quad (17)$$

is the jump operator at site  $i$  and  $\hat{\Delta} = \sum_{i \in \Lambda} \hat{\Delta}_i$ , where

$$\hat{\Delta}_i = \frac{1}{2i} \int_{-\infty}^{\infty} ds \int_{-\infty}^{\infty} ds' \text{sgn}(s-s') g(s) g(-s') \hat{A}_i(s) \hat{A}_i(s'), \quad (18)$$

is the modification of the system Hamiltonian due to the coupling to the heat bath, which is called the Lamb-shift Hamiltonian [25]. In both of Eqs. (17) and (18),  $\hat{A}_i(s) = e^{i\hat{H}_S s} \hat{A}_i e^{-i\hat{H}_S s}$ . In Sec. V, we briefly review the derivation of Eq. (16) following Ref. [31].

For later convenience, we decompose  $\mathcal{L}$  into the unitary time evolution under  $\hat{H}_S$  and the effect of the coupling to the environment:

$$\mathcal{L} = \mathcal{L}_S + \tilde{\mathcal{L}}, \quad (19)$$

where  $\mathcal{L}_S := -i[\hat{H}_S, \cdot]$ .  $\tilde{\mathcal{L}}$  is further decomposed into the Lamb-shift part  $\mathcal{L}_\Delta$  and the dissipative part  $\mathcal{D}$ :

$$\begin{cases} \tilde{\mathcal{L}} = \mathcal{L}_\Delta + \mathcal{D}, \\ \mathcal{L}_\Delta \rho = -i[\hat{\Delta}, \rho] = -i \sum_{i \in \Lambda} [\hat{\Delta}_i, \rho] =: \sum_{i \in \Lambda} \mathcal{L}_{\Delta, i}, \\ \mathcal{D} \rho = \sum_{i \in \Lambda} \left( \hat{L}_i \rho \hat{L}_i^\dagger - \frac{1}{2} \{ \hat{L}_i^\dagger \hat{L}_i, \rho \} \right) =: \sum_{i \in \Lambda} \mathcal{D}_i. \end{cases} \quad (20)$$

Let us introduce the adjoint  $\mathcal{L}^\dagger$  of  $\mathcal{L}$ , which is expressed as

$$\begin{aligned}\mathcal{L}^\dagger \hat{O} &= i[\hat{H}_S + \hat{\Delta}, \hat{O}] + \sum_{i=1}^N \left( \hat{L}_i^\dagger \hat{O} \hat{L}_i - \frac{1}{2} \{ \hat{L}_i^\dagger \hat{L}_i, \hat{O} \} \right) \\ &= \mathcal{L}_S^\dagger \hat{O} + \tilde{\mathcal{L}}^\dagger \hat{O} \\ &= \mathcal{L}_S^\dagger \hat{O} + \mathcal{L}_\Delta^\dagger \hat{O} + \mathcal{D}^\dagger \hat{O}.\end{aligned}\quad (21)$$

The superoperator  $\mathcal{L}^\dagger$  gives the time evolution of an operator in the Heisenberg picture:  $\text{Tr}_S[\hat{O} e^{\mathcal{L}t} \rho] = \text{Tr}_S[(e^{\mathcal{L}^\dagger t} \hat{O}) \rho]$ . It is noted that  $e^{\mathcal{L}t}$  is a unital map:  $\mathcal{L}^\dagger \hat{1} = 0$ , where  $\hat{1}$  is the identity operator.

## B. Main result

We consider the time evolution of a local operator  $\hat{O}_X$  that acts nontrivially to  $X \subset \Lambda$ , where  $|X|$  is finite ( $|X|$  denotes the number of elements of the set  $X$ ). Without loss of generality, we can put  $\|\hat{O}_X\| = 1$ .

We compare the exact time evolution of the expectation value  $\langle \hat{O}_X(t) \rangle := \text{Tr}_S[\hat{O}_X \rho(t)]$ , where  $\rho(t) = \text{Tr}_B[\mathcal{U}(t) \rho(0) \otimes \rho_B]$  is the exact reduced density matrix at time  $t$ , with the approximate time evolution obeying the ULE:  $\langle \hat{O}_X(t) \rangle_{\text{ULE}} := \text{Tr}_S[\hat{O}_X e^{\mathcal{L}t} \rho(0)]$ . We denote by  $\epsilon(t)$  the difference between the two:

$$\epsilon(t) := \left| \langle \hat{O}_X(t) \rangle - \langle \hat{O}_X(t) \rangle_{\text{ULE}} \right|, \quad (22)$$

which is interpreted as the error of the ULE.

What we are going to show is that  $\epsilon(t)$  is bounded from above by a constant  $\bar{\epsilon}$ , which depends on  $\gamma$  and  $\tau_B$  but neither on time  $t$  nor on the initial state  $\rho(0)$ , in the thermodynamic limit. Moreover,  $\bar{\epsilon}$  vanishes in the limit of  $\gamma \rightarrow +0$  or  $\tau_B \rightarrow +0$ .

By introducing dimensionless constants  $\tilde{\gamma} = \gamma/v$  and  $\tilde{\tau}_B = v\tau_B$ , where  $v$  denotes the Lieb-Robinson velocity (it will be defined in Sec. III), it turns out that  $\bar{\epsilon} = O(\tilde{\gamma}^{1/2} \tilde{\tau}_B)$  for small  $\gamma$  or small  $\tau_B$ , which precisely means that  $\bar{\epsilon} \sim \tilde{\gamma}^{1/2}$  for  $\tilde{\gamma} \ll 1$  when  $\tilde{\tau}_B$  is held fixed and  $\bar{\epsilon} \sim \tilde{\tau}_B$  for  $\tilde{\tau}_B \ll 1$  when  $\tilde{\gamma}$  is held fixed. Thus, our main result is summarized as follows:

$$\lim_{N \rightarrow \infty} \epsilon(t) \leq \bar{\epsilon} = O(\tilde{\gamma}^{1/2} \tilde{\tau}_B). \quad (23)$$

For a sufficiently small value of  $\gamma$  or  $\tau_B$ , the ULE is justified for arbitrarily long times even in the thermodynamic limit.

In a non-rigorous but physically sound derivation of the ULE [31] (see also Ref. [40] for a more informal derivation), it appears that the ULE is solely derived by the condition  $\gamma\tau_B \ll 1$ . However, the main result mentioned above implies that the condition of  $\gamma\tau_B \ll 1$  is not sufficient: a stronger condition  $\tilde{\gamma}^{1/2} \tilde{\tau}_B = v^{1/2} \gamma^{1/2} \tau_B \ll 1$  is required for small  $\tilde{\gamma}$ .




FIG. 2. Strategy to evaluate the error bound. Consider a local operator acting nontrivially onto  $X$ . After time evolution, the operator grows. To evaluate the error bound, we use the Lieb-Robinson bound for the contribution from outside the light cone  $X^c(t)$ , whereas we use the accelerated dissipation for the contribution from inside the light cone  $X(t)$ .

## C. Theoretical methods

We briefly explain key theoretical methods: the Lieb-Robinson bound and the accelerated dissipation.

The Lieb-Robinson bound is a mathematical bound on the speed of propagation of information in spin systems. It was originally proved for the unitary dynamics generated by a local Hamiltonian [41–44], and later it was extended to Markovian open quantum systems [39, 45, 46]. Poulin [45] proved the Lieb-Robinson bound for open quantum systems described by a strictly local Lindbladian. Nachtergaele *et al.* [39] extended it to exponentially decaying ones.

The ULE generator of Eq. (16) is, as it is, not written as a sum of exponentially decaying local terms, and hence it is not immediately obvious whether one can apply the result in Ref. [39] to the ULE. In Sec. III, we show that the Lieb-Robinson bound is valid for the ULE by showing that it can be further decomposed into exponentially decaying local terms.

The accelerated dissipation states that, under bulk dissipation, certain kinds of norms of a local operator decay at a rate increasing with time [17, 20, 21]. Eventually, the decay rate is saturated at a value much larger than the initial value. It occurs due to the combination of the operator spreading in the unitary time evolution [18, 19] and the presence of bulk dissipation. Although the accelerated dissipation is considered to be a generic phenomenon, it has not been given as a rigorous theorem.

In this paper, we *assume* an inequality, Eq. (39), representing the accelerated dissipation. We explain the statement of the accelerated dissipation and its intuitive picture in Sec. IV A, and verify it numerically in Sec. IV B. A large decay rate predicted by the accelerated dissipation is crucial to get an error bound that is uniform in time.

Figure 2 shows our strategy to evaluate the error bound. As we will see later in Sec. V, the error due to each approximation is expressed as a space-time sum-

mation. We decompose the spacetime into inside and outside of the light cone  $x = \pm pv|t|$ , where  $p > 2$  is a constant and  $v$  is the Lieb-Robinson velocity. Outside the light cone, we use the Lieb-Robinson bound. Inside the light cone, instead, we use the accelerated dissipation.

### III. LIEB-ROBINSON BOUND FOR THE ULE

Let us consider a time-dependent Lindbladian  $\mathcal{L}(t)$ , whose adjoint  $\mathcal{L}^\dagger(t)$  is written as a sum of local terms:

$$\mathcal{L}^\dagger(t) = \sum_{Z \subset \Lambda} \mathcal{L}_Z^\dagger(t), \quad (24)$$

where  $\mathcal{L}_Z^\dagger(t)$  is a superoperator with support  $Z$ , i.e.  $\mathcal{L}_Z^\dagger(t)\hat{O}_X = 0$  for any operator  $\hat{O}_X$  whose support  $X$  does not have intersection with  $Z$ :  $X \cap Z = \emptyset$ . For each  $Z$ ,  $\mathcal{L}_Z^\dagger(t)$  satisfies  $\mathcal{L}_Z^\dagger(t)\hat{1} = 0$ , but it is not necessary to assume that  $\mathcal{L}_Z^\dagger(t)$  is the adjoint of a Lindbladian in its own right.

We assume that the Lindbladian is exponentially decaying, which implies

$$\sum_{Z \supseteq \{i,j\}} \|\mathcal{L}_Z^\dagger(t)\|_{\text{cb}} \leq D e^{-\mu d(i,j)} \quad (25)$$

for any  $t$ , where  $\|\cdot\|_{\text{cb}}$  denotes the completely bounded norm [39, 47], and  $D$  and  $\mu$  are positive constants which are independent of the system size  $N$ . Here,  $d(i,j)$  is the distance between sites  $i$  and  $j$ :  $d(i,j) = |i-j|$  in the open boundary condition and  $d(i,j) = \min\{|i-j|, N-|i-j|\}$  in the periodic boundary condition.

For the operator  $\hat{O}_X$  whose support is  $X$ , let us define its dissipative time evolution in the Heisenberg picture as

$$\hat{O}_X^{\text{d}}(t) = \bar{\mathcal{T}} \exp\left(\int_0^t ds \mathcal{L}^\dagger(s)\right) \hat{O}_X, \quad (26)$$

where  $\bar{\mathcal{T}}$  is the anti-time-ordering operation. We here put the superscript ‘‘d’’ to distinguish the unitary time evolution  $\hat{O}_X(t) = e^{i\hat{H}st} \hat{O} e^{-i\hat{H}st}$ .

Under the conditions of Eqs. (24) and (25), the following Lieb-Robinson bound has been proved [39]:

$$\|[\hat{O}_X^{\text{d}}(t), \hat{O}_Y]\| \leq C \|\hat{O}_X\| \|\hat{O}_Y\| \min\{|X|, |Y|\} \times e^{-[d(X,Y)-vt]/\xi}, \quad (27)$$

where  $C$ ,  $v$ ,  $\xi$  are positive constants which depend on the lattice geometry and constants  $D$  and  $\mu$  in Eq. (25) but not on the system size  $N$ . Here,  $d(X,Y) = \min\{d(i,j) : i \in X, j \in Y\}$  is the distance between the regions  $X$  and  $Y$ .

It should be noted that the adjoint of the ULE Liouvillian, which is given by Eq. (21) with Eqs. (17) and (18), is *not* in the form of Eq. (24) satisfying Eq. (25). It is

therefore not obvious whether one can apply the Lieb-Robinson bound for the ULE. In the following, we show that the adjoint of the ULE Liouvillian can be brought into the form of Eq. (24) satisfying Eq. (25). It means that the Lieb-Robinson bound holds for Markovian time evolution obeying the ULE.

In the expression  $\mathcal{L}^\dagger = \mathcal{L}_S^\dagger + \mathcal{L}_\Delta^\dagger + \mathcal{D}^\dagger$ ,  $\mathcal{L}_S^\dagger$  is expressed in the desired form because  $\hat{H}_S$  is a local Hamiltonian by assumption. Therefore, the remaining task is to show that  $\mathcal{L}_\Delta^\dagger$  and  $\mathcal{D}^\dagger$  can be decomposed into exponentially decaying local terms.

The operators  $\hat{\Delta}_i$  and  $\hat{L}_i$  in the ULE are not strictly local because  $\hat{A}_i(t)$  in Eqs. (17) and (18) become non-local for large  $|t|$ . Considering the fact that the jump correlation function  $g(t)$  vanishes for  $|t| \gg \tau_B$ , it is expected that  $\hat{A}_i(t)$  in Eqs. (17) and (18) can be approximated by a strictly local operator  $\hat{A}_i^l(t)$ . Here, we define  $\hat{A}_i^l(t)$  as the restriction of  $\hat{A}_i(t)$  to the region  $R_{i,l} := \{j \in \Lambda : d(i,j) \leq l\}$ . More precisely, we define

$$\hat{A}_i^l(t) = \frac{1}{\text{Tr}_{R_{i,l}^c}[\hat{1}_{R_{i,l}^c}]}} \text{Tr}_{R_{i,l}^c}[\hat{A}_i(t)] \otimes \hat{1}_{R_{i,l}^c}, \quad (28)$$

where  $\hat{1}_{R_{i,l}^c}$  is the identity operator for the region  $R_{i,l}^c = \{j \in \Lambda : d(i,j) > l\}$ . The operator  $\hat{A}_i^l(t)$  is regarded as a local operator approximating  $\hat{A}_i(t)$ . By using the Lieb-Robinson bound for the unitary time evolution and recalling that  $\hat{A}_i$  is a single-site operator with  $\|\hat{A}_i\| = 1$ , we have

$$\|\hat{A}_i(t) - \hat{A}_i^l(t)\| \leq \min\{2, C e^{-(l-v|t|)/\xi}\}. \quad (29)$$

Let us define  $\hat{L}_i^l$  as the jump operator localized to the region  $R_{i,l}$  by replacing  $\hat{A}_i(-s)$  in Eq. (17) by  $\hat{A}_i^l(-s)$ . We also define  $\hat{K}_i^l := \hat{L}_i^l - \hat{L}_i^{l-1}$  for  $l = 0, 1, \dots$  with  $\hat{L}_i^{-1} := 0$ . By using Eqs. (12) and (29), it is shown that the operator norm of  $\hat{L}_i$  and  $\hat{L}_i - \hat{L}_i^l$  are bounded from above as

$$\|\hat{L}_i^l\| \leq \|\hat{L}_i\| \leq \int_{-\infty}^{\infty} ds |g_i(s)| \leq \frac{\sqrt{\gamma}}{2}, \quad (30)$$

and

$$\|\hat{L}_i - \hat{L}_i^l\| \leq \frac{\sqrt{\gamma}}{2} (C+2) e^{-l/\eta}, \quad (31)$$

where

$$\eta = \max\{2\xi, v\tau_B\}. \quad (32)$$

This inequality leads to the following bound:

$$\begin{aligned} \|\hat{K}_i^l\| &\leq \|\hat{L}_i - \hat{L}_i^l\| + \|\hat{L}_i - \hat{L}_i^{l-1}\| \\ &\leq \sqrt{\gamma} (C+2) e^{-(l-1)/\eta}. \end{aligned} \quad (33)$$

See Appendix A for the detail on the derivation of Eq. (31).

The dissipator  $\mathcal{D}^\dagger = \sum_{i \in \Lambda} \mathcal{D}_i^\dagger$  is further decomposed into local terms as  $\sum_{i \in \Lambda} \sum_l \mathcal{D}_{i,l}^\dagger$  with

$$\begin{aligned} \mathcal{D}_{i,l}^\dagger \hat{O} &= \hat{K}_i^{l\dagger} \hat{O} \hat{L}_i^l - \frac{1}{2} \{ \hat{K}_i^{l\dagger} \hat{L}_i^l, \hat{O} \} + \hat{L}_i^{l\dagger} \hat{O} \hat{K}_i^l - \frac{1}{2} \{ \hat{L}_i^{l\dagger} \hat{K}_i^l, \hat{O} \} \\ &\quad - \hat{K}_i^{l\dagger} \hat{O} \hat{K}_i^l + \frac{1}{2} \{ \hat{K}_i^{l\dagger} \hat{K}_i^l, \hat{O} \}. \end{aligned} \quad (34)$$

The completely bounded norm of  $\mathcal{D}_{i,l}^\dagger$  is evaluated as

$$\| \mathcal{D}_{i,l}^\dagger \|_{\text{cb}} \leq 4 \| \hat{L}_i^l \| \| \hat{K}_i^l \| + 2 \| \hat{K}_i^l \|^2. \quad (35)$$

By using Eqs. (30) and (33), we obtain

$$\| \mathcal{D}_{i,l}^\dagger \|_{\text{cb}} \leq 2\gamma(C+2)e^{-(l-1)/\eta} + \gamma(C+2)^2 e^{-2(l-1)/\eta}, \quad (36)$$

which shows that the dissipator of the ULE is exponentially decaying.

The Lamb-shift Hamiltonian  $\hat{\Delta}$  is also decomposed into local terms as  $\hat{\Delta} = \sum_{i \in \Lambda} \sum_l \hat{\Delta}_{i,l}$ , where

$$\begin{aligned} \hat{\Delta}_{i,l} &= \frac{1}{2i} \int_{-\infty}^{\infty} ds \int_{-\infty}^{\infty} ds' \text{sgn}(s-s') g_i(s) g_i(-s') \\ &\quad \times \left[ (\hat{A}_i^l(s) - \hat{A}_i^{l-1}(s)) \hat{A}_i^l(s') + \hat{A}_i^l(s) (\hat{A}_i^l(s') - \hat{A}_i^{l-1}(s')) \right]. \end{aligned} \quad (37)$$

Its operator norm is bounded as

$$\begin{aligned} \| \hat{\Delta}_{i,l} \| &\leq \frac{1}{2} \int_{-\infty}^{\infty} ds \int_{-\infty}^{\infty} ds' |g_i(s)| |g_i(-s')| \\ &\quad \times \left( \| \hat{A}_i^l(s) - \hat{A}_i^{l-1}(s) \| + \| \hat{A}_i^l(s') - \hat{A}_i^{l-1}(s') \| \right) \\ &\leq \frac{\gamma}{4\tau_B} \int_{-\infty}^{\infty} ds e^{-2|s|/\tau_B} \| \hat{A}_i^l(s) - \hat{A}_i^{l-1}(s) \| \\ &\leq \frac{\gamma}{2} (C+2) e^{-(l-1)/\eta}. \end{aligned} \quad (38)$$

This bound implies that the Lamb-shift Hamiltonian has exponentially decaying interactions.

In conclusion, the adjoint of the generator of the ULE can be written as a sum of exponentially decaying local terms. Therefore, we can apply the Lieb-Robinson bound of Eq. (27) to the time evolution obeying the ULE.

## IV. ACCELERATED DISSIPATION

### A. Statement and intuitive picture

Let us consider the dissipative time evolution of a local operator  $\hat{O}$  in the Heisenberg picture:  $\hat{O}^d(t) = e^{\mathcal{L}^\dagger t} \hat{O}$ . We assume that  $\mathcal{L}^\dagger$  is written in the form of Eq. (21) with  $\| \hat{L}_i \| \sim \sqrt{\gamma}$ . We also assume that each  $\hat{L}_i$  is approximated by a local operator acting sites near  $i$  (in the case of the ULE, Eq. (31) ensures this property).

Then, the statement of the accelerated dissipation is summarized as follows:

**Accelerated dissipation:** Under the setup explained above, there exist positive constants  $a$  and  $\zeta$  such that

$$\lim_{N \rightarrow \infty} \| [\hat{O}^d(t), \hat{A}(t')] \| \leq \zeta \| \hat{O} \| \| \hat{A} \| e^{-a\sqrt{\gamma}vt} \quad (39)$$

for any  $t > 0$  and  $t' \in \mathbb{R}$ , where  $\hat{A}$  is an arbitrary local operator and  $\hat{A}(t') = e^{i\hat{H}_S t'} \hat{A} e^{-i\hat{H}_S t'}$  with  $\hat{H}_S$  being a local Hamiltonian.

In the following, we explain an intuitive picture behind this statement. Under bulk dissipation of strength  $\gamma$ , the decay rate of an operator  $\hat{O}^d(t)$  is roughly proportional to  $\gamma \mathcal{S}(t)$ , where  $\mathcal{S}(t)$  is the average operator size of  $\hat{O}^d(t)$  (i.e. how many sites this operator acts on) [17, 20, 21]. If the operator  $\hat{O}$  has no overlap with all the powers of the Hamiltonian  $\hat{H}_S$ , i.e.  $\hat{O}$  has no diagonal elements in the energy basis,  $\mathcal{S}(t)$  roughly obeys the following equation [20, 21]:

$$\frac{d\mathcal{S}(t)}{dt} \sim v - 2\gamma\delta\mathcal{S}(t)^2, \quad (40)$$

where  $\delta\mathcal{S}(t)^2$  is the variance of the operator size. The first term of the right hand side implies that the operator size increases linearly  $\mathcal{S}(t) \sim vt$ , which is called operator spreading or operator growth [18, 19]. The second term shows that bulk dissipation decreases the operator size at a rate proportional to the variance of the operator size.

When there is no local conserved quantity at  $\gamma = 0$ , which can happen for a time-dependent Hamiltonian  $\hat{H}_S(t)$ , the operator size distribution evolves diffusively  $\delta\mathcal{S}(t) \sim \sqrt{t}$  [21], and thus the operator size reaches the peak value  $\mathcal{S}_{\text{peak}} \sim \tilde{\gamma}^{-1}$ , and then starts to decrease. In this case,  $\| \hat{O}^d(t) \| \sim e^{-\gamma vt^2}$  up to  $t \sim \gamma^{-1}$ , and then the operator eventually undergoes an exponential decay  $\| \hat{O}^d(t) \| \sim e^{-gt}$  for  $t \gg \tilde{\gamma}^{-1}$ . Here,  $g$  is the spectral gap of the Liouvillian, which, interestingly, converges to a non-zero value in the weak dissipation limit after taking the thermodynamic limit:  $\lim_{\gamma \rightarrow +0} \lim_{N \rightarrow \infty} g = \bar{g} > 0$ . This finite decay rate is related to the intrinsic relaxation of the system, which is referred to as the quantum Ruelle-Pollicott resonance [17, 22, 48–51]. It is thus expected that Eq. (39) holds for a time-dependent Hamiltonian with no local conserved quantity at  $\gamma = 0$ .

In contrast, when there is a local conserved quantity at  $\gamma = 0$  (the Hamiltonian  $\hat{H}_S$  is always conserved in a static system), the presence of slow hydrodynamic modes makes the operator-size distribution broad and leads to  $\delta\mathcal{S}(t) \sim \mathcal{S}(t)$ . As a result, from Eq. (40), the average operator size will be saturated:  $\mathcal{S}(t) \sim \tilde{\gamma}^{-1/2}$  for  $\tilde{\gamma}^{1/2} vt \gtrsim 1$ . As was already mentioned, the decay rate is roughly proportional to  $\gamma \mathcal{S}(t)$ . The above estimate of  $\mathcal{S}(t)$  for a static system thus implies the following behavior:

$$\| [\hat{O}^d(t), \hat{A}(t')] \| \lesssim e^{-a\sqrt{\gamma}vt}. \quad (41)$$

The decay rate for small  $\gamma$  is enhanced ( $\gamma \rightarrow \sqrt{\gamma}v$ ) due to operator spreading, which is the meaning of the accelerated decay.

When  $\hat{O}$  has an overlap with local conserved quantities (e.g. the system Hamiltonian), taking the commutator with a local operator  $\hat{A}(t')$  as in Eq. (41) is crucial for the accelerated dissipation. To investigate the effect of local conserved quantities, we decompose the operator  $\hat{O}$  into the diagonal part  $\hat{O}_{\text{diag}}$  and the off-diagonal part  $\hat{O}_{\text{off}}$  in the energy basis:  $\hat{O} = \hat{O}_{\text{diag}} + \hat{O}_{\text{off}}$ . We expect that Eq. (41) holds for the off-diagonal part:  $\|[\hat{O}_{\text{off}}^{\text{d}}(t), \hat{A}(t')]\| \lesssim e^{-a\sqrt{\gamma}vt}$ . While, the diagonal part does not spread because it is conserved under the unitary time evolution at  $\gamma = 0$ , and hence one might be tempted to consider that it does not show the accelerated decay.

We now argue that the accelerated decay actually occurs for the operator norm of the commutator  $\|[\hat{O}^{\text{d}}(t), \hat{A}(t')]\|$ . For simplicity, we assume that there is no local conserved quantity apart from  $\hat{H}_{\text{S}}$ , and the system of interest obeys the eigenstate thermalization hypothesis [52, 53], which implies

$$\hat{O}_{\text{diag}} = O_{\text{eq}}(\hat{H}_{\text{S}}/N) \quad (42)$$

up to an exponentially small error in  $N$ , where  $O_{\text{eq}}(\cdot)$  is a smooth function. Physically,  $O_{\text{eq}}(\varepsilon)$  corresponds to the equilibrium average of  $\hat{O}$  at the energy density  $\varepsilon$ . We then have  $\|[\hat{O}_{\text{diag}}, \hat{A}(t')]\| = \|[O_{\text{eq}}(\hat{H}_{\text{S}}/N), \hat{A}]\|$ , where we have used the fact that  $\hat{O}_{\text{diag}}$  commutes with  $\hat{H}_{\text{S}}$ . Performing the Taylor expansion of  $O_{\text{eq}}(\hat{H}_{\text{S}}/N)$  as

$$O_{\text{eq}}(\hat{H}_{\text{S}}/N) = \sum_{n=0}^{\infty} \frac{O_{\text{eq}}^{(n)}(0)}{n!} \left( \frac{\hat{H}_{\text{S}}}{N} \right)^n, \quad (43)$$

and using the inequality

$$\|[\hat{H}_{\text{S}}^n, \hat{A}]\| \leq n \|\hat{H}_{\text{S}}\|^{n-1} \|[\hat{H}_{\text{S}}, \hat{A}]\|, \quad (44)$$

we obtain

$$\|[O_{\text{eq}}(\hat{H}_{\text{S}}/N), \hat{A}]\| \lesssim \frac{\|[\hat{H}_{\text{S}}, \hat{A}]\|}{N}. \quad (45)$$

Here, we have assumed that the Taylor expansion of  $dO_{\text{eq}}(x)/dx$  around  $x = 0$  converges absolutely, and a non-essential constant factor is dropped in Eq. (45). When  $\hat{H}_{\text{S}}$  is a local Hamiltonian,  $\|[\hat{H}_{\text{S}}, \hat{A}]\|$  does not depend on the system size for any local operator  $\hat{A}$ , which implies

$$\|[\hat{O}_{\text{diag}}, \hat{A}(t')]\| \lesssim \frac{1}{N}, \quad (46)$$

and hence the contribution from the conserved quantity (i.e. the diagonal part) is negligible in the thermodynamic limit. It should be emphasized that the operator norm  $\|\hat{O}_{\text{diag}}\|$  does not vanish in the thermodynamic limit. Taking the commutator with a local operator is crucial in Eq. (39). In this way, the contribution from the overlaps with conserved quantities in Eq. (39) can be

dropped in the thermodynamic limit. It is thus expected that Eq. (39) also holds in a static Hamiltonian.

A similar argument shows that the diagonal part is also negligible in considering the Frobenius norm  $\|\hat{O}^{\text{d}}(t)\|_{\text{F}}$ , where  $\|\hat{O}\|_{\text{F}}^2 := \text{Tr}_{\text{S}}[\hat{O}^{\dagger}\hat{O}]/\text{Tr}_{\text{S}}[\hat{1}]$ . Without loss of generality, we can assume  $\text{Tr}_{\text{S}}[\hat{O}_{\text{diag}}] = 0$ . It is shown that  $\|\hat{O}_{\text{diag}}\|_{\text{F}} \propto 1/\sqrt{N} \rightarrow +0$  in the thermodynamic limit. That is why the accelerated decay has been observed in the Loschmidt echo fidelity [20, 21], which is equivalent to  $\|\hat{O}^{\text{d}}(t)\|_{\text{F}}^2$ .

## B. Numerical verification

To support the theoretical argument given above, we provide numerical results verifying Eq. (39). Let us consider the Ising model in a tilted field in the periodic boundary condition:

$$\hat{H}_{\text{S}} = - \sum_{i=1}^N (\hat{\sigma}_i^z \hat{\sigma}_{i+1}^z + h_x \hat{\sigma}_i^x + h_z \hat{\sigma}_i^z), \quad (47)$$

where  $\hat{\sigma}_i^{\alpha}$  ( $\alpha = x, y, z$ ) denote the Pauli matrices at site  $i$ . We set  $h_x = 0.9045$  and  $h_z = 0.8090$  following [54], for which the eigenstate thermalization hypothesis is numerically verified. Bulk dissipation is modeled by the two jump operators  $\hat{L}_i^+ = \sqrt{\gamma}\hat{\sigma}_i^+$  and  $\hat{L}_i^- = \sqrt{\gamma}\hat{\sigma}_i^-$  at each site  $i$ .

We choose  $\hat{O} = \hat{\sigma}_1^x$  and  $\hat{A} = \hat{\sigma}_1^z$ , and compute the norm appearing in Eq. (39). We decompose the norm into the diagonal part  $\|[\hat{O}_{\text{diag}}^{\text{d}}(t), \hat{A}]\|$  and the off-diagonal part  $\|[\hat{O}_{\text{off}}^{\text{d}}(t), \hat{A}]\|$ . According to the discussion in Sec. IV, we expect that the norm from the off-diagonal part undergoes an exponential decay with the decay rate proportional to  $\sqrt{\gamma}$ , and that the norm from the diagonal part decreases as the system size increases.

Our numerical results shown in Fig. 3 agree with those theoretical predictions. In Fig. 3 (a), the norm from the off-diagonal part is plotted as a function of  $\sqrt{\gamma}t$  for different values of  $\gamma$ . We see that the slope on a semi-log plot does not depend on  $\gamma$ , which implies  $\|[\hat{O}_{\text{off}}^{\text{d}}(t), \hat{A}]\| \sim e^{-a\sqrt{\gamma}t}$ , which is in accordance with Eq. (39). In Fig. 3 (b), the norm from the diagonal part is plotted for different system sizes. We see that the norm decreases as the system size increases, which agrees with the theoretical prediction  $\|[\hat{O}_{\text{diag}}^{\text{d}}(t), \hat{A}]\| \sim 1/N$ . It means that the effect of conserved quantities would be negligible for large system sizes.

Because of an exponential growth of the state space, numerical calculations are limited to relatively small systems ( $N \leq 12$ ). However, the consistency between numerical results and the theoretical argument presented in Sec. IV supports the validity of the accelerated dissipation in the form of Eq. (39).




FIG. 3. Contribution from (a) the off-diagonal part  $\|[\hat{O}_{\text{off}}^{\text{d}}(t), \hat{A}]\|$  and (b) the diagonal part  $\|[\hat{O}_{\text{diag}}^{\text{d}}(t), \hat{A}]\|$  to the accelerated dissipation in the bulk-dissipated Ising model in a tilted field. (a) The norm from the off-diagonal part decays exponentially, and the decay rate is proportional to  $\sqrt{\gamma}$ , which agrees with the inequality (39). (b) The norm from the diagonal part decreases as the system size increases, which is in accordance with the argument in Sec. IV.

## V. PRELIMINARY: DERIVATION OF THE ULE

In this section, we review the derivation of Eq. (16) following Ref. [31]. Through the derivation, we give the explicit expression of the error term for each approximation, which is used to evaluate the error bounds on the dynamics of a local quantity in Sec. VI.

Let us start with the Liouville-von Neumann equation for the total system, i.e., Eq. (14), and its formal solution given by Eq. (15) with a factorized initial state  $\rho_{\text{T}}(0) = \rho(0) \otimes \rho_{\text{B}}$ . In the interaction picture,  $\rho_{\text{T}}^{\text{I}}(t) = e^{i(\hat{H}_{\text{S}} + \hat{H}_{\text{B}})t} \rho_{\text{T}}(t) e^{-i(\hat{H}_{\text{S}} + \hat{H}_{\text{B}})t}$ , which is formally written as

$$\rho_{\text{T}}^{\text{I}}(t) = \mathcal{U}_{\text{I}}(t, 0) \rho_{\text{T}}(0), \quad \mathcal{U}_{\text{I}}(t, s) = \mathcal{T} e^{\int_s^t ds' \mathcal{L}_{\text{I}}(s')}, \quad (48)$$

where  $\mathcal{T}$  is the time-ordering operator and  $\mathcal{L}_{\text{I}}(t) = -i[\hat{H}_{\text{I}}(t), \cdot]$ . Here,  $\hat{H}_{\text{I}}(t)$  is the interaction Hamiltonian in the interaction picture, which is given by

$$\hat{H}_{\text{I}}(t) = \sum_{i \in \Lambda} \hat{A}_i(t) \otimes \hat{B}_i(t), \quad (49)$$

where  $\hat{A}_i(t) = e^{i\hat{H}_{\text{S}}t} \hat{A}_i e^{-i\hat{H}_{\text{S}}t}$  and  $\hat{B}_i(t) = e^{i\hat{H}_{\text{B}}t} \hat{B}_i e^{-i\hat{H}_{\text{B}}t}$ .

In the interaction picture, we have

$$\frac{d\rho_{\text{T}}^{\text{I}}(t)}{dt} = \mathcal{L}_{\text{I}}(t) \rho_{\text{T}}^{\text{I}}(t) = \mathcal{L}_{\text{I}}(t) \mathcal{U}_{\text{I}}(t, 0) \rho(0) \otimes \rho_{\text{B}}. \quad (50)$$

The reduced density matrix in the interaction picture is given by  $\rho^{\text{I}}(t) = \text{Tr}_{\text{B}}[\rho_{\text{T}}^{\text{I}}(t)]$ . By using Wick's theorem

for free bosons, we have

$$\begin{aligned} \frac{d\rho^{\text{I}}(t)}{dt} &= \text{Tr}_{\text{B}}[\mathcal{L}_{\text{I}}(t) \mathcal{U}^{\text{I}}(t, 0) \rho(0) \otimes \rho_{\text{B}}] \\ &= - \sum_{i \in \Lambda} \sum_{\nu = \pm 1} \nu \int_0^t ds \Phi_i(\nu(t-s)) \\ &\quad \times [\hat{A}_i(t), \text{Tr}_{\text{B}}[\mathcal{U}_{\text{I}}(t, s) \mathcal{A}_i^{\nu}(s) \mathcal{U}_{\text{I}}(s, 0) \rho(0) \otimes \rho_{\text{B}}]], \end{aligned} \quad (51)$$

where  $\mathcal{A}_i^{+1}(s)(\cdot) = \hat{A}_i(s)(\cdot)$  and  $\mathcal{A}_i^{-1}(s)(\cdot) = (\cdot) \hat{A}_i(s)$ . That is, we multiply  $\hat{A}_i(s)$  from left when  $\nu = 1$  and from right when  $\nu = -1$ .

Here,  $\mathcal{U}_{\text{I}}(t, s)$  is rewritten as

$$\mathcal{U}_{\text{I}}(t, s) = 1 + \int_s^t ds' \mathcal{L}_{\text{I}}(s') \mathcal{U}_{\text{I}}(s', s). \quad (52)$$

By substituting it into Eq. (51), we have

$$\begin{aligned} \frac{d\rho^{\text{I}}(t)}{dt} &= - \sum_{i \in \Lambda} \sum_{\nu = \pm 1} \nu \int_0^t ds \Phi_i(\nu(t-s)) \\ &\quad \times [\hat{A}_i(t), \mathcal{A}_i^{\nu}(s) \rho^{\text{I}}(s)] + \hat{\xi}_{\text{B}}^{\text{I}}(t), \end{aligned} \quad (53)$$

where

$$\begin{aligned} \hat{\xi}_{\text{B}}^{\text{I}}(t) &= i \sum_{i, j \in \Lambda} \sum_{\nu = \pm 1} \nu \int_0^t ds \Phi_i(\nu(t-s)) \int_s^t ds' \\ &\quad [\hat{A}_i(t), [\hat{A}_j(s'), \Psi_{ij}^{\nu}(s', s)]], \end{aligned} \quad (54)$$

where

$$\Psi_{ij}^{\nu}(s', s) = \text{Tr}_{\text{B}}[\hat{B}_j(s') \mathcal{U}_{\text{I}}(s', s) \mathcal{A}_i^{\nu}(s) \mathcal{U}_{\text{I}}(s, 0) \rho(0) \otimes \rho_{\text{B}}]. \quad (55)$$

Ignoring  $\hat{\xi}_{\text{B}}^{\text{I}}(t)$  corresponds to the Born approximation. Therefore,  $\hat{\xi}_{\text{B}}^{\text{I}}(t)$  expresses the error induced by the Born approximation in the interaction picture.

Next, we perform the Markov approximation: in Eq. (53), replace  $\rho^I(s)$  by  $\rho^I(t)$  and  $\int_0^t ds$  by  $\int_{-\infty}^t ds$ . As a result, we obtain

$$\frac{d\rho^I(t)}{dt} = - \sum_{i \in \Lambda} \sum_{\nu = \pm 1} \nu \int_0^\infty ds \Phi_i(\nu s) [\hat{A}_i(t), \mathcal{A}_i^\nu(t-s) \rho^I(t)] + \hat{\xi}_B^I(t) + \hat{\xi}_M^I(t), \quad (56)$$

where the error induced by the Markov approximation is given by

$$\begin{aligned} \hat{\xi}_M^I(t) &= \sum_{i \in \Lambda} \sum_{\nu = \pm 1} \nu \int_0^t ds \Phi_i(\nu s) \\ &\times [\hat{A}_i(t), \mathcal{A}_i^\nu(t-s) (\rho^I(t) - \rho^I(t-s))] \\ &+ \sum_{i \in \Lambda} \sum_{\nu = \pm 1} \nu \int_t^\infty ds \Phi_i(\nu s) [\hat{A}_i(t), \mathcal{A}_i^\nu(t-s) \rho^I(t)]. \end{aligned} \quad (57)$$

By introducing an operator

$$\hat{R}_i(t) = \int_0^\infty ds \Phi_i(s) \hat{A}_i(t-s), \quad (58)$$

and using  $\Phi_i(-t) = \Phi_i(t)^*$ , Eq. (56) is expressed as

$$\frac{d\rho^I}{dt} = - \sum_{i \in \Lambda} [\hat{A}_i(t), \hat{R}_i(t) \rho^I - \rho^I \hat{R}_i^\dagger(t)] + \hat{\xi}_{\text{BM}}^I(t), \quad (59)$$

where  $\hat{\xi}_{\text{BM}}^I(t) = \hat{\xi}_B^I(t) + \hat{\xi}_M^I(t)$ . Equation (59) is known as the Redfield equation [25] when we ignore  $\hat{\xi}_{\text{BM}}^I(t)$ .

In a textbook derivation of the Lindblad equation [25], the secular approximation is used, but it is in general not valid in many-body systems, as is mentioned in Introduction. Nathan and Rudner [31, 32] introduce the memory-dressing transformation to derive the ULE, which is explained below.

Let us first rewrite Eq. (59) by using the jump correlation function, see Eq. (11). By using a superoperator  $\mathcal{F}(t, s, s')$ , which is defined as

$$\begin{aligned} \mathcal{F}(t, s, s') \hat{O} &= \sum_{i \in \Lambda} \theta(t-s') \\ &\times \left\{ g_i(t-s) g_i(s-s') [\hat{A}_i(t), \hat{O} \hat{A}_i(s')] \right. \\ &\left. + g_i^*(t-s) g_i^*(s-s') [\hat{A}_i(s') \hat{O}, \hat{A}_i(t)] \right\}, \end{aligned} \quad (60)$$

Eq. (59) is expressed as follows:

$$\frac{d\rho^I}{dt} = \int_{-\infty}^\infty ds \int_{-\infty}^\infty ds' \mathcal{F}(t, s, s') \rho^I(t) + \hat{\xi}_{\text{BM}}^I(t). \quad (61)$$

A key observation by Nathan and Rudner is that if we replace  $\mathcal{F}(t, s, s')$  by  $\mathcal{F}(s, t, s')$ , the above equation becomes of the Lindblad form, which is nothing but the ULE, except for the error term. To make a connection

between the Redfield equation and the ULE, let us introduce the memory-dressing transformation  $\rho^I(t) \rightarrow \tilde{\rho}^I(t)$ , which is defined as

$$\tilde{\rho}^I(t) = [1 + \mathcal{M}(t)] \rho^I(t), \quad (62)$$

where

$$\mathcal{M}(t) = \int_t^\infty ds_1 \int_{-\infty}^t ds_2 \int_{-\infty}^\infty ds_3 [\mathcal{F}(s_1, s_2, s_3) - \mathcal{F}(s_2, s_1, s_3)]. \quad (63)$$

The transformed density matrix  $\tilde{\rho}^I(t)$  is referred to as the modified density matrix [31]. By differentiating Eq. (62) with respect to  $t$  and using Eqs. (61) and (63), we obtain

$$\frac{d\tilde{\rho}^I}{dt} = \int_{-\infty}^\infty ds \int_{-\infty}^\infty ds' \mathcal{F}(s, t, s') \tilde{\rho}^I(t) + \hat{\xi}_{\text{BM}}^I(t) + \hat{\xi}_1^I(t), \quad (64)$$

where

$$\hat{\xi}_1^I(t) = \mathcal{M}(t) \partial_t \rho^I(t). \quad (65)$$

Straightforward calculations show that the first term of the right hand side of Eq. (64) is nothing but that of the ULE, i.e. Eq. (16), in the interaction picture. The ULE Liouvillian in the interaction picture is given by

$$\tilde{\mathcal{L}}^I(t) = e^{-\mathcal{L}st} \mathcal{L} e^{\mathcal{L}st} - \mathcal{L}_S = e^{-\mathcal{L}st} \tilde{\mathcal{L}} e^{\mathcal{L}st}, \quad (66)$$

where recall that we write  $\mathcal{L} = \mathcal{L}_S + \tilde{\mathcal{L}}$ . We have

$$\tilde{\mathcal{L}}^I(t) = \int_{-\infty}^\infty ds \int_{-\infty}^\infty ds' \mathcal{F}(s, t, s'). \quad (67)$$

Equation (64) is thus written as

$$\frac{d\tilde{\rho}^I}{dt} = \tilde{\mathcal{L}}^I(t) \tilde{\rho}^I(t) + \hat{\xi}_{\text{BM}}^I(t) + \hat{\xi}_1^I(t) + \hat{\xi}_2^I(t), \quad (68)$$

where

$$\hat{\xi}_2^I(t) = \tilde{\mathcal{L}}^I(t) [\rho^I(t) - \tilde{\rho}^I(t)]. \quad (69)$$

The modified density matrix obeys the ULE when we ignore

$$\hat{\xi}_{\text{all}}^I(t) := \hat{\xi}_{\text{BM}}^I(t) + \hat{\xi}_1^I(t) + \hat{\xi}_2^I(t), \quad (70)$$

which is interpreted as the total correction to the ULE for the modified density matrix.

If the modified density matrix  $\tilde{\rho}^I(t)$  is close to the exact density matrix  $\rho^I(t)$  and  $\hat{\xi}_{\text{all}}^I$  is small enough,  $\rho^I(t)$  approximately obeys the ULE:  $d\rho^I(t)/dt \approx \mathcal{L}\rho^I(t)$  by going back to the Schrödinger picture.

In Ref. [31], the following upper bounds on the trace norm of  $\hat{\xi}_{\text{all}}^I(t)$  and that of  $\rho^I(t) - \tilde{\rho}^I(t)$  are obtained [55]:

$$\|\hat{\xi}_{\text{all}}^I(t)\|_1 \lesssim (N\gamma)^2 \tau_B \quad (71)$$

and

$$\|\rho^{\text{I}}(t) - \tilde{\rho}^{\text{I}}(t)\|_1 \lesssim N\gamma\tau_{\text{B}}, \quad (72)$$

respectively, where  $\|\cdot\|_1$  denotes the trace norm. Since  $\|\mathcal{L}^{\text{I}}(t)\rho^{\text{I}}(t)\|_1 \lesssim \gamma N$ , Eq. (71) implies that the correction term is actually small when  $N\gamma\tau_{\text{B}} \ll 1$ . Under the same condition, Eq. (72) shows that we can ignore the difference between the exact density matrix and the modified one.

However, there are two problems in the above evaluation as is already pointed out in Introduction. Firstly, if we are interested in many-body systems under bulk dissipation, the condition  $N\gamma\tau_{\text{B}} \ll 1$  is not satisfied in the thermodynamic limit ( $N \rightarrow \infty$ ). Secondly, Eq. (71) gives an error bound at the level of the equations of motion, but errors might be accumulated with time. It is important to exclude this possibility to ensure that the ULE can be used to predict long-time (i.e.  $t \gtrsim 1/\gamma$ ) evolution of the system.

In the next section, we give an error bound on  $\epsilon(t)$  that is given by Eq. (22). It turns out that  $\epsilon(t)$  keeps small during the time evolution in the thermodynamic limit when  $\gamma$  or  $\tau_{\text{B}}$  is small enough.

## VI. ERROR BOUNDS

In this section, we derive the main result summarized in Sec. II. First of all, we decompose the total error  $\epsilon(t)$  into three parts by using the triangle inequality:

$$\begin{aligned} \epsilon(t) &\leq \|\text{Tr}_{\text{S}}[\hat{O}_X(\rho(t) - \tilde{\rho}(t))]\| \\ &+ \|\text{Tr}_{\text{S}}[\hat{O}_X(\tilde{\rho}(t) - e^{\mathcal{L}t}\tilde{\rho}(0))]\| \\ &+ \|\text{Tr}_{\text{S}}[\hat{O}_X e^{\mathcal{L}t}(\rho(0) - \tilde{\rho}(0))]\| \end{aligned} \quad (73)$$

First term of the right hand side represents the error due to the difference between the exact density matrix  $\rho(t)$  and the modified one in the Schrödinger picture,  $\tilde{\rho}(t) = e^{-i\hat{H}st}\tilde{\rho}^{\text{I}}(t)e^{i\hat{H}st}$ . The second term represents the error due to the correction to the ULE for the modified density matrix, see Eq. (68). The last term represents the error due to the difference between the initial state of the exact density matrix and that of the modified one.

As for the second term of Eq. (73), by using the Schrödinger picture of Eq. (68), i.e.

$$\frac{d\tilde{\rho}(t)}{dt} = \mathcal{L}\tilde{\rho}(t) + \hat{\xi}_{\text{all}}(t), \quad (74)$$

where  $\hat{\xi}_{\text{all}}(t) = e^{-i\hat{H}st}\hat{\xi}_{\text{all}}^{\text{I}}(t)e^{i\hat{H}st}$ , we obtain

$$\tilde{\rho}(t) - e^{\mathcal{L}t}\tilde{\rho}(0) = \int_0^t dt' e^{\mathcal{L}t'} \hat{\xi}_{\text{all}}(t-t'). \quad (75)$$

By further decomposing  $\hat{\xi}_{\text{all}}(t)$  into  $\hat{\xi}_{\text{B}}(t) + \hat{\xi}_{\text{M}}(t) + \hat{\xi}_{\text{1}}(t) +$

$\hat{\xi}_{\text{2}}(t)$ , we have

$$\begin{aligned} &\|\text{Tr}_{\text{S}}[\hat{O}_X(\tilde{\rho}(t) - e^{\mathcal{L}t}\tilde{\rho}(0))]\| \\ &\leq \int_0^t dt' \left| \text{Tr}_{\text{S}}[\hat{O}_X e^{\mathcal{L}t'} \hat{\xi}_{\text{all}}(t-t')] \right| \\ &\leq \sum_{\alpha=\text{B},\text{M},\text{1},\text{2}} \int_0^t dt' \left| \text{Tr}_{\text{S}}[\hat{O}_X^{\text{d}}(t') \hat{\xi}_{\alpha}(t-t')] \right|, \end{aligned} \quad (76)$$

where recall that  $\hat{O}_X^{\text{d}}(t) = e^{\mathcal{L}^{\dagger}t}\hat{O}_X$ . In this way, we can decompose the total error into the error due to each approximation:  $\epsilon(t) \leq \epsilon_{\text{B}}(t) + \epsilon_{\text{M}}(t) + \epsilon_{\text{md}}(t)$ , where

(Born approximation)

$$\epsilon_{\text{B}}(t) = \int_0^t dt' \left| \text{Tr}_{\text{S}}[\hat{O}_X^{\text{d}}(t') \hat{\xi}_{\text{B}}(t-t')] \right| \quad (77)$$

(Markov approximation)

$$\epsilon_{\text{M}}(t) = \int_0^t dt' \left| \text{Tr}_{\text{S}}[\hat{O}_X^{\text{d}}(t') \hat{\xi}_{\text{M}}(t-t')] \right| \quad (78)$$

(Memory-dressing transformation)

$$\begin{aligned} \epsilon_{\text{md}}(t) &= \int_0^t dt' \sum_{\alpha=1,2} \left| \text{Tr}_{\text{S}}[\hat{O}_X^{\text{d}}(t') \hat{\xi}_{\alpha}(t-t')] \right| \\ &+ \left| \text{Tr}_{\text{S}}[\hat{O}_X(\rho(t) - \tilde{\rho}(t))] \right| + \left| \text{Tr}_{\text{S}}[\hat{O}_X^{\text{d}}(t)(\rho(0) - \tilde{\rho}(0))] \right| \end{aligned} \quad (79)$$

In the following, we evaluate each error separately by using theoretical methods explained in Sec. II C. It turns out that each error bound is at most  $O(\tilde{\gamma}^{1/2}\tilde{\tau}_{\text{B}})$  for small  $\gamma$  or small  $\tau_{\text{B}}$ .

### A. Born approximation

By using Eq. (54) and the cyclic property of the trace, and changing the integration variables as  $s \rightarrow t-t'-s$  and  $s' \rightarrow t-t'-s'$ , we obtain

$$\begin{aligned} \epsilon_{\text{B}}(t) &\leq \sum_{i,j \in \Lambda} \sum_{\nu=\pm 1} \int_0^t dt' \int_0^{t-t'} ds \int_0^s ds' |\Phi_i(\nu s)| \\ &\times \| [\hat{O}_X^{\text{d}}(t'), \hat{A}_i], \hat{A}_j(-s') \| \cdot \| \Psi_{ij}^{\nu}(t-t'-s', t-t'-s) \|_1. \end{aligned} \quad (80)$$

By using Wick's theorem, Eq. (13), and  $\|\hat{A}_i\| = 1$ , we have

$$\begin{aligned} &\| \Psi_{ij}^{\nu}(t-t'-s', t-t'-s) \|_1 \leq \\ &\sum_{\nu'=\pm 1} \int_{s'}^{t-t'} ds'' |\Phi_j(\nu'(s'-s''))| \\ &\leq \frac{2\gamma}{\tau_{\text{B}}} \int_{s'}^{\infty} ds'' e^{-|s''-s'|/\tau_{\text{B}}} = 2\gamma. \end{aligned} \quad (81)$$

By using Eq. (13) again,  $\epsilon_B(t)$  is bounded as

$$\epsilon_B(t) \leq \sum_{i,j \in \Lambda} \frac{4\gamma^2}{\tau_B} \int_0^t dt' \int_0^{t-t'} ds \int_0^s ds' e^{-s/\tau_B} \times \|\llbracket [\hat{O}_X^d(t'), \hat{A}_i], \hat{A}_j(-s') \rrbracket\|. \quad (82)$$

Since the right hand side is an increasing function of  $t$ , we obtain

$$\begin{aligned} \epsilon_B(t) &\leq \sum_{i,j \in \Lambda} \frac{4\gamma^2}{\tau_B} \int_0^\infty dt' \int_0^\infty ds \int_0^s ds' e^{-s/\tau_B} \\ &\quad \times \|\llbracket [\hat{O}_X^d(t'), \hat{A}_i], \hat{A}_j(-s') \rrbracket\| \\ &=: \bar{\epsilon}_B. \end{aligned} \quad (83)$$

It should be noted that this upper bound does not depend on  $t$ . Therefore, if it is shown that  $\bar{\epsilon}_B$  is sufficiently small, the Born approximation is justified for arbitrarily long times.

In the following analysis, we change the integration variable  $t'$  to  $t$  for notational simplicity. There is no confusion about it because  $\bar{\epsilon}_B$  does no longer depend on  $t$ .

We introduce a region  $X(t)$  defined as

$$X(t) = \{i \in \Lambda : d(X, i) \leq pvt\}, \quad (84)$$

where  $v$  is the Lieb-Robinson velocity and  $p$  is an arbitrary constant satisfying  $p > 2$  (e.g. we can put  $p = 3$ ). In the following analysis, the summation over  $i$  and  $j$  in Eq. (83) is decomposed into the three contributions: (i)  $i, j \in X(t)$ , (ii)  $i \in X^c(t)$  and  $d(X, i) \geq d(X, j)$ , and (iii)  $j \in X^c(t)$  and  $d(X, j) > d(X, i)$ . Correspondingly,  $\bar{\epsilon}_B = \bar{\epsilon}_B^{(i)} + \bar{\epsilon}_B^{(ii)} + \bar{\epsilon}_B^{(iii)}$ . Below, we evaluate each contribution. In the following analysis, it turns out that the case (i) gives the dominant contribution, and we have  $\bar{\epsilon}_B \lesssim \tilde{\gamma}^{1/2} \tilde{\tau}_B$ .

$$(i) \ i, j \in X(t)$$

In this case, we evaluate the double commutator in  $\bar{\epsilon}_B^{(i)}$  as

$$\|\llbracket [\hat{O}_X^d(t), \hat{A}_i], \hat{A}_j(-s') \rrbracket\| \leq 2\|\llbracket \hat{O}_X^d(t), \hat{A}_i \rrbracket\|. \quad (85)$$

By using the assumption of the accelerated dissipation, i.e. Eq. (39),

$$\bar{\epsilon}_B^{(i)} \leq 8\gamma^2 \tau_B \zeta \int_0^\infty dt (|X| + 2pvt)^2 e^{-a\tilde{\gamma}^{1/2}vt} \quad (86)$$

where we used  $\sum_{i,j \in X(t)} 1 = |X(t)|^2 \leq (|X| + 2pvt)^2$ . A straightforward calculation yields

$$\bar{\epsilon}_B^{(i)} \leq \left( \frac{64p^2}{a^3} \tilde{\gamma}^{1/2} + \frac{32p|X|}{a^2} \tilde{\gamma} + \frac{8|X|^2}{a} \tilde{\gamma}^{3/2} \right) \tilde{\tau}_B \zeta. \quad (87)$$

The dominant contribution to  $\bar{\epsilon}_B^{(i)}$  for small  $\tilde{\gamma}$  or small  $\tilde{\tau}_B$  is thus given by

$$\bar{\epsilon}_B^{(i)} \lesssim \tilde{\gamma}^{1/2} \tilde{\tau}_B. \quad (88)$$

It should be noted that if we used a milder assumption  $\|\hat{O}^d(t)\| \leq e^{-O(\gamma t)}$ , we would obtain  $\bar{\epsilon}_B^{(i)} \sim \tilde{\tau}_B/\tilde{\gamma}$ , which does not tend to zero in the weak-coupling limit  $\tilde{\gamma} \rightarrow +0$ . To obtain the error bound that vanishes in the weak-coupling limit, the assumption of the accelerated decay, i.e. Eq. (39), is needed.

$$(ii) \ i \in X^c(t) \text{ and } d(X, i) \geq d(X, j)$$

By using

$$\|\llbracket [\hat{O}_X^d(t), \hat{A}_i], \hat{A}_j(-s') \rrbracket\| \leq 2\|\llbracket \hat{O}_X^d(t), \hat{A}_i \rrbracket\| \leq 2C e^{-[d(X, i) - vt]/\xi}, \quad (89)$$

where the Lieb-Robinson bound (Eq. (27)) is used in the last inequality, we have

$$\bar{\epsilon}_B^{(ii)} \leq 8C\gamma^2 \tau_B \int_0^\infty dt \sum_{i \in X^c(t)} \sum_{j: d(X, j) \leq d(X, i)} e^{-[d(X, i) - vt]/\xi}. \quad (90)$$

By putting  $d(X, i) = r$ , we have  $\sum_{i \in X^c(t)} \approx 2 \int_{pvt}^\infty dr$  and  $\sum_{j: d(X, j) \leq d(X, i)} 1 = |X| + 2r$ . We thus have

$$\bar{\epsilon}_B^{(ii)} \leq 16C\gamma^2 \tau_B \int_0^\infty dt \int_{pvt}^\infty dr (|X| + 2r) e^{-(r - vt)/\xi}. \quad (91)$$

After carrying out the integrations, we obtain

$$\bar{\epsilon}_B^{(ii)} \leq 16C \frac{2(2p-1)\xi^3 + (p-1)\xi^2|X|}{(p-1)^2} \tilde{\gamma}^2 \tilde{\tau}_B, \quad (92)$$

which is of order  $\tilde{\gamma}^2 \tilde{\tau}_B$  and not dominant compared with  $\bar{\epsilon}_B^{(i)}$  for  $\tilde{\gamma} \ll 1$ .

$$(iii) \ j \in X^c(t) \text{ and } d(X, j) > d(X, i)$$

We introduce the region  $X_j$  defined as

$$X_j = \left\{ i \in \Lambda : d(X, i) \leq \frac{p-1}{p} d(X, j) \right\}. \quad (93)$$

We then introduce the operator  $\hat{O}_{X,j}^d(t)$  as the operator obtained by localizing  $\hat{O}_X^d(t)$  to the region  $X_j$ :

$$\hat{O}_{X,j}^d(t) = \frac{\text{Tr}_{X_j^c}[\hat{O}_X^d(t)] \otimes \hat{1}_{X_j^c}}{\text{Tr}_{X_j^c}[\hat{1}_{X_j^c}]}, \quad (94)$$

where  $X_j^c$  is the complement of  $X_j$ . Let us decompose  $\|[\hat{O}_X^d(t), \hat{A}_i, \hat{A}_j(-s')]\|$  in  $\bar{\epsilon}_B^{(iii)}$  as follows:

$$\begin{aligned} \|[\hat{O}_X^d(t), \hat{A}_i, \hat{A}_j(-s')]\| &\leq \|[\hat{O}_{X,j}^d(t), \hat{A}_i, \hat{A}_j(-s')]\| \\ &+ \|[\hat{O}_X^d(t) - \hat{O}_{X,j}^d(t), \hat{A}_i, \hat{A}_j(-s')]\|. \end{aligned} \quad (95)$$

By using the Lieb-Robinson bound, we have

$$\begin{aligned} \|[\hat{O}_{X,j}^d(t), \hat{A}_i, \hat{A}_j(-s')]\| &\leq \|[\hat{O}_{X,j}^d(t), \hat{A}_i]\| \min\{2, Ce^{-[d(X_j,j)-vs']/\xi}\} \\ &\leq 2I[i \in X_j] \min\{2, Ce^{-[d(X_j,j)-vs']/\xi}\} \\ &\leq 2I[i \in X_j] \min\{2, Ce^{-[\frac{1}{p}d(X,j)-vs']/\xi}\} \end{aligned} \quad (96)$$

where  $I[\cdot]$  is the indicator function, and

$$\begin{aligned} \|[\hat{O}_X^d(t) - \hat{O}_{X,j}^d(t), \hat{A}_i, \hat{A}_j(-s')]\| &\leq 4\|\hat{O}_X^d(t) - \hat{O}_{X,j}^d(t)\| \\ &\leq 4C|X|e^{-[\frac{p-1}{p}d(X,j)-vt]/\xi}. \end{aligned} \quad (97)$$

By substituting Eqs. (96) and (97) into Eq. (95), we obtain

$$\begin{aligned} \bar{\epsilon}_B^{(iii)} &\leq \frac{4\gamma^2}{\tau_B} \int_0^\infty dt \int_0^\infty ds \int_0^\infty ds' e^{-s/\tau_B} \sum_{j \in X^c(t)} \sum_{i: d(X,i) < d(X,j)} \\ &\times \left[ 2I[i \in X_j] \min\left\{2, Ce^{-[\frac{1}{p}d(X,j)-vs']/\xi}\right\} \right. \\ &\quad \left. + 4C|X|e^{-[\frac{p-1}{p}d(X,j)-vt]/\xi} \right]. \end{aligned} \quad (98)$$

As for  $\min\{2, Ce^{-[\frac{1}{p}d(X,j)-vs']/\xi}\}$ , we choose 2 when  $pvt < d(X,j) < pvs'$ , and choose  $Ce^{-[\frac{1}{p}d(X,j)-vs']/\xi}$  when  $d(X,j) \geq pvs'$ . As a result, we have

$$\bar{\epsilon}_B^{(iii)} \leq B_1 + B_2 + B_3, \quad (99)$$

where

$$B_1 = \frac{16\gamma^2}{\tau_B} \int_0^\infty dt \int_t^\infty ds \int_t^s ds' e^{-s/\tau_B} \times \sum_{j: pvt < d(X,j) < pvs'} \sum_{i \in X_j} 1, \quad (100)$$

$$B_2 = \frac{8C\gamma^2}{\tau_B} \int_0^\infty dt \int_0^\infty ds \int_0^s ds' e^{-s/\tau_B} \times \sum_{j: d(X,j) > pvs'} \sum_{i \in X_j} e^{-[\frac{d(X,j)}{p}-vs']/\xi}, \quad (101)$$

and

$$B_3 = \frac{16C|X|\gamma^2}{\tau_B} \int_0^\infty dt \int_0^\infty ds \int_0^s ds' e^{-s/\tau_B} \times \sum_{j: d(X,j) > pvt} \sum_{i: d(X,i) < d(X,j)} e^{-[\frac{p-1}{p}d(X,j)-vt]/\xi}. \quad (102)$$

By introducing  $r = d(X,j)$  and  $r' = d(X,i)$ , and replace the sum over  $i$  and  $j$  by integrations over  $r$  and  $r'$ , we can explicitly evaluate  $B_1$ ,  $B_2$ , and  $B_3$ . We omit tedious calculations, and just present the result:

$$\begin{aligned} B_1 &\approx 32\tilde{\gamma}^2\tilde{\tau}_B^3[p|X| + 4p(p-1)\tilde{\tau}_B], \\ B_2 &\approx 16C\tilde{\gamma}^2\tilde{\tau}_B[p|X|\xi(\xi + \tilde{\tau}_B) + 4p(p-1)\xi(\xi^2 + \xi\tilde{\tau}_B + \tilde{\tau}_B^2)], \\ B_3 &\approx \frac{32C|X|\xi^2p}{(p-1)(p-2)}\tilde{\gamma}^2\tilde{\tau}_B \left[ |X| + \frac{2p(2p-3)}{(p-1)(p-2)}\xi \right]. \end{aligned} \quad (103)$$

By collecting them,  $\bar{\epsilon}_B^{(iii)} = O(\tilde{\gamma}^2\tilde{\tau}_B)$  for small  $\tilde{\gamma}$  or  $\tilde{\tau}_B$ , which is not dominant compared with  $\bar{\epsilon}_B^{(i)}$ .

## B. Markov approximation

In the Schrödinger picture, Eq. (57) is expressed as

$$\hat{\xi}_M(t) = \hat{\xi}_M^{(1)}(t) + \hat{\xi}_M^{(2)}(t), \quad (104)$$

where

$$\begin{aligned} \hat{\xi}_M^{(1)} &= \sum_{i \in \Lambda} \sum_{\nu = \pm 1} \nu \int_0^t ds \Phi(\nu s) [\hat{A}_i, \mathcal{A}_i^\nu(-s) \\ &\quad \times e^{-i\hat{H}s}(\rho^I(t) - \rho^I(t-s))e^{i\hat{H}s}], \end{aligned} \quad (105)$$

and

$$\hat{\xi}_M^{(2)}(t) = \sum_{i \in \Lambda} \sum_{\nu = \pm 1} \nu \int_t^\infty ds \Phi(\nu s) [\hat{A}_i, \mathcal{A}_i^\nu(-s)\rho(t)]. \quad (106)$$

Let us start with the evaluation of the error caused by  $\hat{\xi}_M^{(2)}(t)$ , which is given by

$$\begin{aligned} \epsilon_M^{(2)}(t) &= \int_0^t dt' \left| \text{Tr}_S[\hat{O}_X^d(t')\hat{\xi}_M^{(2)}(t-t')] \right| \\ &\leq \sum_{i \in \Lambda} \sum_{\nu = \pm 1} \int_0^t dt' \int_{t-t'}^\infty ds \frac{\gamma}{\tau_B} e^{-s/\tau_B} \|\mathcal{A}_i^\nu(-s)[\hat{O}_X^d(t'), \hat{A}_i]\| \\ &\leq 2\gamma \sum_{i \in \Lambda} \int_0^t dt' e^{-(t-t')/\tau_B} \|[\hat{O}_X^d(t'), \hat{A}_i]\|. \end{aligned} \quad (107)$$

By using Eq. (39) (the accelerated dissipation) for  $i$  with  $d(X,i) \leq vt'$ , and by using Eq. (27) (the Lieb-Robinson bound) for  $i$  with  $d(X,i) > vt'$ , we find that it is bounded as

$$\epsilon_M^{(2)}(t) \leq \bar{\epsilon}_M^{(2)} := \frac{4\zeta}{a}\tilde{\gamma}^{1/2}\tilde{\tau}_B + 4(|X| + C\xi)\tilde{\gamma}\tilde{\tau}_B, \quad (108)$$

which is of order  $\tilde{\gamma}^{1/2}\tilde{\tau}_B$  for small  $\gamma$  or  $\tau_B$ .

Next, let us consider the error caused by  $\hat{\xi}_M^{(1)}(t)$ . In

Eq. (105),  $\rho^I(t) - \rho^I(t-s)$  is written as

$$\begin{aligned} \rho^I(t) - \rho^I(t-s) &= \int_{t-s}^t ds' \frac{d}{ds'} \rho^I(s') \\ &= \int_{t-s}^t ds' \frac{d}{ds'} \text{Tr}_B[\mathcal{U}_I(s', 0)\rho(0) \otimes \rho_B] \\ &= -i \sum_{j \in \Lambda} \int_{t-s}^t ds' [\hat{A}_j(s'), \text{Tr}_B(\hat{B}_j(s')\mathcal{U}_I(s', 0)\rho(0) \otimes \rho_B)]. \end{aligned} \quad (109)$$

By introducing

$$\hat{\Psi}_j(t, t-s') := -ie^{-i\hat{H}_S t} \text{Tr}_B[\hat{B}_j(s')\mathcal{U}_I(s', 0)\rho(0) \otimes \rho_B] e^{i\hat{H}_S t}, \quad (110)$$

we obtain

$$\begin{aligned} e^{-i\hat{H}_S t}(\rho^I(t) - \rho^I(t-s))e^{i\hat{H}_S t} \\ = \sum_{j \in \Lambda} \int_0^s ds' [\hat{A}_j(-s'), \hat{\Psi}_j(t, s')]. \end{aligned} \quad (111)$$

By using Wick's theorem,  $\hat{\Psi}_j(t, t-s')$  is rewritten as

$$\begin{aligned} e^{i\hat{H}_S t} \hat{\Psi}_j(t, t-s') e^{-i\hat{H}_S t} &= -i \int_0^{s'} ds'' \Phi_j(s' - s'') \sum_{\nu=\pm 1} \nu \\ &\times \text{Tr}_B[\mathcal{U}_I(s', s'') \mathcal{A}_j^\nu(s'') \mathcal{U}_I(s'', 0)\rho(0) \otimes \rho_B], \end{aligned} \quad (112)$$

and thus its trace norm is evaluated as

$$\begin{aligned} \|\hat{\Psi}_j(t, t-s')\|_1 &\leq \int_0^{s'} ds'' \frac{\gamma}{\tau_B} e^{-(s'-s'')/\tau_B} \times 2 \\ &\leq 2\gamma. \end{aligned} \quad (113)$$

By substituting Eq. (111) into Eq. (105), we have

$$\begin{aligned} \hat{\xi}_M^{(1)}(t) &= \sum_{ij \in \Lambda} \sum_{\nu=\pm 1} \nu \int_0^t ds \int_0^s ds' \Phi(\nu s) \\ &\times [\hat{A}_i, \mathcal{A}_i^\nu(-s) [\hat{A}_j(-s'), \hat{\Psi}_j(t, s')]]. \end{aligned} \quad (114)$$

Thus, the error caused by  $\hat{\xi}_M^{(1)}(t)$  is given by

$$\begin{aligned} \epsilon_M^{(1)}(t) &= \int_0^t dt' \left| \text{Tr}_S[\hat{O}_X^d(t') \hat{\xi}_M^{(1)}(t-t')] \right| \\ &\leq \sum_{ij \in \Lambda} \int_0^\infty dt' \int_0^\infty ds \int_0^s ds' \frac{\gamma}{\tau_B} e^{-s'/\tau_B} \sum_{\nu=\pm 1} \\ &\times \left| \text{Tr}_S \left\{ [\mathcal{A}_i^\nu(-s) [\hat{O}_X^d(t'), \hat{A}_i], \hat{A}_j(-s')] \hat{\Psi}_j(t, s') \right\} \right| \\ &\leq \frac{2\gamma^2}{\tau_B} \sum_{ij \in \Lambda} \int_0^\infty dt' \int_0^\infty ds \int_0^s ds' e^{-s'/\tau_B} \sum_{\nu=\pm 1} \\ &\times \|[\mathcal{A}_i^\nu(-s) [\hat{O}_X^d(t'), \hat{A}_i], \hat{A}_j(-s')]\| \\ &=: \bar{\epsilon}_M^{(1)}, \end{aligned} \quad (115)$$

where we have used Eq. (113). We notice that the expression of  $\bar{\epsilon}_M^{(1)}$  is similar to that of  $\bar{\epsilon}_B$  given in Eq. (83), except for the presence of  $\mathcal{A}_i^\nu(-s)$  in front of  $[\hat{O}_X^d(t'), \hat{A}_i]$ .

Similarly to the analysis of the Born approximation, let us decompose the summation over  $i$  and  $j$  in Eq. (115) into the three parts  $\bar{\epsilon}_M^{(1)} \leq \bar{\epsilon}_M^{(i)} + \bar{\epsilon}_M^{(ii)} + \bar{\epsilon}_M^{(iii)}$ : (i)  $i, j \in X(t)$ , (ii)  $i \in X^c(t)$  and  $d(X, i) \geq d(X, j)$ , and (iii)  $j \in X^c(t)$  and  $d(X, j) > d(X, i)$ . In the following, we evaluate each contribution separately.

(i)  $i, j \in X(t)$

In this case, we have

$$\begin{aligned} \bar{\epsilon}_M^{(i)} &\leq \frac{8\gamma^2}{\tau_B} \int_0^\infty dt \int_0^\infty ds \int_0^s ds' e^{-s'/\tau_B} \\ &\times \sum_{i, j \in X(t)} \|[\hat{O}_X^d(t), \hat{A}_i]\|. \end{aligned} \quad (116)$$

This is identical to the upper bound on  $\bar{\epsilon}_B^{(i)}$ , and Eq. (87) also holds for  $\bar{\epsilon}_M^{(i)}$ . We therefore conclude

$$\bar{\epsilon}_M^{(i)} = O(\tilde{\gamma}^{1/2} \tilde{\tau}_B). \quad (117)$$

(ii)  $i \in X^c(t)$  and  $d(X, i) \geq d(X, j)$

In this case, by using the simple bound  $\|[\mathcal{A}_i^\nu(-s) [\hat{O}_X^d(t), \hat{A}_i], \hat{A}_j(-s')]\| \leq 2\|[\hat{O}_X^d(t), \hat{A}_i]\|$  and using the Lieb-Robinson bound, we have

$$\bar{\epsilon}_M^{(ii)} \leq 8C\gamma^2 \tau_B \int_0^\infty dt \sum_{i \in X(t)} \sum_{j: d(X, j) \leq d(X, i)} e^{-[d(X, i) - vt]/\xi}. \quad (118)$$

This is identical to Eq. (90), and hence, following the same calculation, we conclude

$$\bar{\epsilon}_M^{(ii)} = O(\tilde{\gamma}^2 \tilde{\tau}_B). \quad (119)$$

(iii)  $j \in X^c(t)$  and  $d(X, j) > d(X, i)$

By using  $\|\hat{A}_i\| = 1$  and  $[\hat{A}\hat{B}, \hat{C}] = \hat{A}[\hat{B}, \hat{C}] + [\hat{A}, \hat{C}]\hat{B}$  for arbitrary operators  $\hat{A}$ ,  $\hat{B}$ , and  $\hat{C}$ , we have

$$\begin{aligned} \|[\mathcal{A}_i^\nu(-s) [\hat{O}_X^d(t), \hat{A}_i], \hat{A}_j(-s')]\| \\ \leq \|[[\hat{O}_X^d(t), \hat{A}_i], \hat{A}_j(-s')]\| \\ + \|[\hat{A}_i(-s), \hat{A}_j(-s')]\| \cdot \|[\hat{O}_X^d(t), \hat{A}_i]\|. \end{aligned} \quad (120)$$

By using this inequality, we obtain

$$\bar{\epsilon}_M^{(iii)} \leq \bar{\epsilon}_M^{(iii-1)} + \bar{\epsilon}_M^{(iii-2)}, \quad (121)$$

where

$$\begin{aligned} \bar{\epsilon}_M^{(iii-1)} &= \frac{4\gamma^2}{\tau_B} \int_0^\infty dt \int_0^\infty ds \int_0^s ds' e^{-s'/\tau_B} \\ &\times \sum_{j \in X^c(t)} \sum_{i: d(X, i) < d(X, j)} \|[[\hat{O}_X^d(t), \hat{A}_i], \hat{A}_j(-s')]\|, \end{aligned} \quad (122)$$

and

$$\begin{aligned} \bar{\epsilon}_M^{(\text{iii}-2)} &= \frac{4\gamma^2}{\tau_B} \int_0^\infty dt \int_0^\infty ds \int_0^s ds' e^{-s/\tau_B} \\ &\times \sum_{j \in X^c(t)} \sum_{i: d(X,i) < d(X,j)} \|\hat{A}_i(-s'), \hat{A}_j\| \cdot \|\hat{O}_X^d(t), \hat{A}_i\|. \end{aligned} \quad (123)$$

We find that  $\bar{\epsilon}_M^{(\text{iii}-1)}$  is identical to  $\bar{\epsilon}_B^{(\text{iii})}$ , and hence  $\bar{\epsilon}_M^{(\text{iii}-1)} = O(\tilde{\gamma}^2 \tilde{\tau}_B)$ .

Thus, the remaining problem is to evaluate  $\bar{\epsilon}_M^{(\text{iii}-2)}$ . The product  $\|\hat{A}_i(-s'), \hat{A}_j\| \cdot \|\hat{O}_X^d(t), \hat{A}_i\|$  in Eq. (123) is evaluated by using the Lieb-Robinson bound. We bound it as follows: when  $i \in X_j$ ,

$$\begin{aligned} \|\hat{A}_i(-s'), \hat{A}_j\| \cdot \|\hat{O}_X^d(t), \hat{A}_i\| &\leq 2 \min\{2, C e^{-[d(i,j)-vs']/\xi}\} \\ &\leq 2 \min\left\{2, C \exp\left[-\frac{1}{\xi} \left(\frac{d(X,j)}{p} - vs'\right)\right]\right\}, \end{aligned} \quad (124)$$

where we used  $d(i,j) \geq d(X,j) - d(X,i) \geq d(X,j)/p$ , and when  $i \in X_j^c$ ,

$$\begin{aligned} \|\hat{A}_i(-s'), \hat{A}_j\| \cdot \|\hat{O}_X^d(t), \hat{A}_i\| &\leq 2C e^{-[d(X,i)-vt]/\xi} \\ &\leq 2C \exp\left[-\frac{1}{\xi} \left(\frac{p-1}{p} d(X,j) - vt\right)\right], \end{aligned} \quad (125)$$

where we used  $d(X,i) > d(X,j)(p-1)/p$  (recall the definition of  $X_j$ , see Eq. (93)). By substituting these inequalities into Eq. (123), after calculations, we finally obtain

$$\begin{aligned} \bar{\epsilon}_M^{(\text{iii}-2)} &\leq \tilde{\gamma}^2 \tilde{\tau}_B \cdot 16Cp\xi \left[ \frac{p(2p-3)}{(p-1)^2(p-2)^2} \xi^2 \right. \\ &\quad \left. + |X|(\xi + \tilde{\tau}_B) + 4(p-1)(\xi^2 + \xi \tilde{\tau}_B + \tilde{\tau}_B^2) \right] \\ &\quad + \tilde{\gamma}^2 \tilde{\tau}_B^3 \cdot 32p[|X| + 4(p-1)\tilde{\tau}_B]. \end{aligned} \quad (126)$$

From this expression, we find that  $\bar{\epsilon}_M^{(\text{iii}-2)} = O(\tilde{\gamma}^2 \tilde{\tau}_B)$ .

In conclusion, we find  $\bar{\epsilon}_M^{(\text{iii})} \leq \bar{\epsilon}_M^{(\text{iii}-1)} + \bar{\epsilon}_M^{(\text{iii}-2)} = O(\tilde{\gamma}^2 \tilde{\tau}_B)$ .

### C. Memory-dressing transformation

By looking at Eq. (79), we see that  $\epsilon_{\text{md}}(t)$  is composed of four parts:

$$\begin{cases} \epsilon_{\text{md}}^{(1)}(t) = \int_0^t dt' \left| \text{Tr}_S[\hat{O}_X^d(t') \hat{\xi}_1(t-t')] \right|, \\ \epsilon_{\text{md}}^{(2)}(t) = \int_0^t dt' \left| \text{Tr}_S[\hat{O}_X^d(t') \hat{\xi}_2(t-t')] \right|, \\ \epsilon_{\text{md}}^{(3)}(t) = \left| \text{Tr}_S[\hat{O}_X(\rho(t) - \tilde{\rho}(t))] \right|, \\ \epsilon_{\text{md}}^{(4)}(t) = \left| \text{Tr}_S[\hat{O}_X^d(t)(\rho(0) - \tilde{\rho}(0))] \right|. \end{cases} \quad (127)$$

We evaluate each contribution.

By going back to the Schrödinger picture,  $\hat{\xi}_1(t) = e^{-i\hat{H}st} \hat{\xi}_1^I(t) e^{i\hat{H}st}$  is given by

$$\hat{\xi}_1(t) = \mathcal{M}(0) \left[ e^{-i\hat{H}st} \frac{\partial \rho^I(t)}{\partial t} e^{i\hat{H}st} \right], \quad (128)$$

where  $\mathcal{M}(t)$  is defined in Eq. (63), which is expressed as

$$\mathcal{M}(0) = \sum_{\nu=\pm 1} \int_0^\infty ds_1 \int_{-\infty}^0 ds_2 \int_{-\infty}^\infty ds_3 \nu \mathcal{F}(\nu s_1, \nu s_2, s_3). \quad (129)$$

By using  $\hat{\Psi}_j(t, s)$  defined in Eq. (110), we can write

$$e^{-i\hat{H}st} \frac{\partial \rho^I(t)}{\partial t} e^{i\hat{H}st} = \sum_{j \in \Lambda} [\hat{A}_j, \hat{\Psi}_j(t, 0)]. \quad (130)$$

By substituting Eqs. (129) and (130) into Eq. (128),  $\epsilon_{\text{md}}^{(1)}(t)$  is bounded from above as

$$\begin{aligned} \epsilon_{\text{md}}^{(1)}(t) &\leq 2 \sum_{\nu=\pm 1} \int_0^t dt' \int_0^\infty ds_1 \int_{-\infty}^0 ds_2 \int_{-\infty}^{\nu s_1} ds_3 \\ &\quad \times |g_i(\nu s_1 - \nu s_2)| \cdot |g_i(\nu s_2 - s_3)| \\ &\quad \times G(t', \nu s_1, s_3) \|\Psi_j(t-t', 0)\|_1, \end{aligned} \quad (131)$$

where

$$G(t, s, s') = \sum_{ij \in \Lambda} \|\hat{A}_i(s') [\hat{O}_X^d(t), \hat{A}_i(s)], \hat{A}_j\|. \quad (132)$$

By using Eqs. (12) and (113), we obtain

$$\begin{aligned} \epsilon_{\text{md}}^{(1)}(t) &\leq \frac{\gamma^2}{\tau_B} \sum_{\nu=\pm 1} \int_0^\infty dt' \int_0^\infty ds_1 \int_{-\infty}^0 ds_2 \int_{-\infty}^{\nu s_1} ds_3 \\ &\quad \times e^{-2(s_1-s_2)/\tau_B} e^{-2|\nu s_2-s_3|/\tau_B} G(t', \nu s_1, s_3) \\ &=: \bar{\epsilon}_{\text{md}}^{(1)}. \end{aligned} \quad (133)$$

By following similar calculations as in the previous subsections, an upper bound on  $\bar{\epsilon}_{\text{md}}^{(1)}$  is evaluated in Appendix B, where it turns out that

$$\bar{\epsilon}_{\text{md}}^{(1)} \lesssim \tilde{\gamma}^{1/2} \tilde{\tau}_B \quad (134)$$

for small  $\gamma$  or small  $\tau_B$ .

Next, we consider  $\epsilon_{\text{md}}^{(2)}(t)$ . By using  $\hat{\xi}_2(t) = -\tilde{\mathcal{L}}\mathcal{M}(0)\rho(t)$ , we obtain

$$\begin{aligned} \epsilon_{\text{md}}^{(2)}(t) &= \int_0^t dt' \left| \text{Tr}_S \left[ \left( \mathcal{M}^\dagger(0) \tilde{\mathcal{L}}^\dagger \hat{O}_X^d(t') \right) \rho(t) \right] \right| \\ &\leq \int_0^t dt' \|\mathcal{M}^\dagger(0) \tilde{\mathcal{L}}^\dagger \hat{O}_X^d(t')\|, \end{aligned} \quad (135)$$

where  $\mathcal{M}^\dagger(0)$  is the adjoint of  $\mathcal{M}(0)$ :  $\text{Tr}_S[\hat{X}_1 \mathcal{M}(0) \hat{X}_2] = \text{Tr}_S[(\mathcal{M}^\dagger(0) \hat{X}_1) \hat{X}_2]$  for arbitrary operators  $\hat{X}_1$  and  $\hat{X}_2$ . By using Eqs. (12), (60) and (63) combined with the

inequality  $\int_{-\infty}^s ds_3 e^{-2|s_2-s_3|/\tau_B} \leq \tau_B$  for an arbitrary  $s \in \mathbb{R}$ , we obtain the following simple bound on  $\epsilon_{\text{md}}^{(2)}(t)$ :

$$\begin{aligned} \epsilon_{\text{md}}^{(2)}(t) &\leq \frac{\gamma}{4} \sum_{\nu=\pm 1} \sum_{i,j \in \Lambda} \int_0^\infty dt' \int_0^\infty ds e^{-2s/\tau_B} \\ &\quad \times \|\llbracket \tilde{\mathcal{L}}_i^\dagger \hat{O}_X^d(t'), \hat{A}_j(\nu s) \rrbracket\| \\ &=: \bar{\epsilon}_{\text{md}}^{(2)}. \end{aligned} \quad (136)$$

where  $\tilde{\mathcal{L}}_i^\dagger = \mathcal{L}_{\Delta,i}^\dagger + \mathcal{D}_i^\dagger$ .

The evaluation of  $\bar{\epsilon}_{\text{md}}^{(2)}$  is provided in Appendix C. It turns out that the contribution from  $i, j \in X(t)$  is dominant, and we again obtain

$$\bar{\epsilon}_{\text{md}}^{(2)} \lesssim \tilde{\gamma}^{1/2} \tilde{\tau}_B. \quad (137)$$

Next, let us consider  $\epsilon_{\text{md}}^{(3)}(t)$ . It is written as

$$\epsilon_{\text{md}}^{(3)}(t) = \left| \text{Tr}_S[\hat{O}_X^d(t) \mathcal{M}(0) \rho(0)] \right| \leq \|\mathcal{M}^\dagger(0) \hat{O}_X^d(t)\|. \quad (138)$$

By using Eqs. (12) and (129), we have

$$\begin{aligned} \epsilon_{\text{md}}^{(3)}(t) &\leq \frac{\gamma}{2\tau_B^2} \sum_{\nu=\pm 1} \sum_{i \in \Lambda} \int_0^\infty ds_1 \int_{-\infty}^0 ds_2 \int_{-\infty}^{\nu s_1} ds_3 \\ &\quad \times e^{-2(s_1-s_2)/\tau_B} e^{-2|\nu s_2-s_3|/\tau_B} \|\llbracket \hat{O}_X^d(t), \hat{A}_i(\nu s_1) \rrbracket\|. \end{aligned} \quad (139)$$

Let us evaluate it for  $i \in X(t)$  and  $i \in X^c(t)$ . For  $i \in X(t)$ , we use Eq. (39). A straightforward calculation yields

$$\begin{aligned} &\sum_{\nu=\pm 1} \int_0^\infty ds_1 \int_{-\infty}^0 ds_2 \int_{-\infty}^{\nu s_1} ds_3 e^{-2(s_1-s_2)/\tau_B} e^{-2|\nu s_2-s_3|/\tau_B} \\ &= \tau_B^3/4. \end{aligned} \quad (140)$$

The contribution from  $i \in X(t)$  in the right hand side of Eq. (139) is thus given by

$$\frac{\gamma \tau_B}{8} (|X| + 2pvt) \zeta e^{-a\sqrt{\gamma}vt} \leq \frac{p\zeta}{4a} \tilde{\gamma}^{1/2} \tilde{\tau}_B + \frac{|X|}{8} \zeta \tilde{\gamma} \tilde{\tau}_B. \quad (141)$$

For  $i \in X^c(t)$ , we use the Lieb-Robinson bound, e.g. Eq. (27):

$$\begin{aligned} &\frac{\gamma}{2\tau_B^2} \sum_{\nu=\pm 1} \sum_{i \in X^c(t)} \int_0^\infty ds_1 \int_{-\infty}^0 ds_2 \int_{-\infty}^{\nu s_1} ds_3 \\ &\quad \times e^{-2(s_1-s_2)/\tau_B} e^{-2|\nu s_2-s_3|/\tau_B} \min\{2, C e^{-[d(X,i)-v(t+s_1)]/\xi}\}. \end{aligned} \quad (142)$$

For  $s_1 \leq (p-1)d(X,i)/pv$ , we use

$$\begin{aligned} \min\{2, C e^{-[d(X,i)-v(t+s_1)]/\xi}\} &\leq C e^{-[d(X,i)-v(t+s_1)]/\xi} \\ &\leq C e^{-[\frac{d(X,i)}{p}-vt]/\xi}, \end{aligned} \quad (143)$$

and for  $s_1 > (p-1)d(X,i)/pv$ , we use

$$\min\{2, C e^{-[d(X,i)-v(t+s_1)]/\xi}\} \leq 2. \quad (144)$$

We then obtain an upper bound

$$\frac{Cp\xi}{4} \tilde{\gamma} \tilde{\tau}_B + \frac{p}{4(p-1)} e^{-2(p-1)t/\tau_B} \tilde{\gamma} \tilde{\tau}_B^2 \quad (145)$$

from  $i \in X^c(t)$  in Eq. (139).

An upper bound on  $\epsilon_{\text{md}}^{(3)}(t)$  is given by the sum of Eqs. (141) and (145). We therefore find that  $\epsilon_{\text{md}}^{(3)}(t)$  behaves as

$$\epsilon_{\text{md}}^{(3)}(t) \lesssim \tilde{\gamma}^{1/2} \tilde{\tau}_B \quad (146)$$

for small  $\gamma$  or small  $\tau_B$ .

Finally, we consider  $\epsilon_{\text{md}}^{(4)}(t)$ . We find

$$\epsilon_{\text{md}}^{(4)}(t) = \left| \text{Tr}_S[\hat{O}_X^d(t) \mathcal{M}(0) \rho(0)] \right| \leq \|\mathcal{M}^\dagger(0) \hat{O}_X^d(t)\|, \quad (147)$$

which is identical to the right hand side of Eq. (138). We can therefore conclude that  $\epsilon_{\text{md}}^{(4)}(t)$  behaves as

$$\epsilon_{\text{md}}^{(4)}(t) \lesssim \tilde{\gamma}^{1/2} \tilde{\tau}_B \quad (148)$$

for small  $\gamma$  or small  $\tau_B$ .

## VII. CONCLUSION

The ULE describes dissipative time evolution of an open quantum many-body system. However, in its microscopic derivation, several approximations are introduced, and hence it has not been obvious at all whether the ULE gives accurate predictions, especially for the long-time behavior of macroscopically large quantum systems. In this paper, we evaluate errors on the time evolution of local quantities. For this purpose, we show that the Lieb-Robinson bound holds in an open system obeying the ULE, and introduce an assumption of the accelerated dissipation, which is physically motivated by recent works on the operator scrambling. The Lieb-Robinson bound and the accelerated dissipation are key tools to evaluate error bounds.

We find that the error is bounded from above by a quantity of order  $\tilde{\gamma}^{1/2} \tilde{\tau}_B$ , where  $\tilde{\gamma}$  is the dimensionless dissipation strength and  $\tilde{\tau}_B$  is the dimensionless correlation time of the environment. It is therefore concluded that either the weak-coupling limit  $\tilde{\gamma} \rightarrow +0$  or the singular-coupling limit  $\tilde{\tau}_B \rightarrow +0$  justify the use of the ULE for arbitrarily long times in the thermodynamic limit.

Dissipation usually destroys quantum coherence, and hence it is crucial to understand the effect of dissipation and develop techniques reducing environmental effects in quantum technologies [10]. Meanwhile, recent experimental progress [5–8] introduces a new perspective: engineered dissipation is rather useful to control and manipulate quantum states [56, 57]. The interplay between dissipation and interactions gives rise to rich nonequilibrium dynamics [13, 14] and new kinds of quantum phases [11, 58, 59]. It is important to develop new

theoretical tools and lay the foundation for them to tame dissipation in quantum many-body systems. This work takes a step forward in this direction.

Finally, we conclude with open problems. The accelerated dissipation is used as a key assumption in this work: it is yet to be formulated as a mathematical theorem. It is an important future problem to refine the notion of the accelerated dissipation as a fundamental tool to describe open quantum many-body systems. It is also a future work to justify the Lindblad equation in boundary dissipated quantum many-body systems. The setup of boundary dissipation is relevant for quantum transport in a nonequilibrium steady state [60, 61]. In such a system, we typically observe delayed relaxation in the sense that the relaxation time is much larger than the inverse of the spectral gap of the Liouvillian [62, 63]. The delayed relaxation is considered to be originated from the presence of bulk hydrodynamic modes, which do not decay until they reach the boundary. It is then nontrivial how to control errors stemming from the inside of the light cone in Fig. 2. We hope to address these problems in the future.

## ACKNOWLEDGMENTS

This work was supported by JSPS KAKENHI Grant No. JP21H05185 and by JST, PRESTO Grant No. JPMJPR2259.

### Appendix A: Derivation of Eq. (31)

The definition of the jump operator in the ULE is given in Eq. (17). By using it, we have

$$\|\hat{L}_i - \hat{L}_i^l\| \leq \int_{-\infty}^{\infty} ds |g_i(s)| \|\hat{A}_i(-s) - \hat{A}_i^l(-s)\|. \quad (\text{A1})$$

By using Eqs. (12) and (29), it is bounded as

$$\begin{aligned} \|\hat{L}_i - \hat{L}_i^l\| &\leq \int_{-\infty}^{\infty} ds \frac{\sqrt{\gamma}}{2\tau_B} e^{-2|s|/\tau_B} \min\{2, Ce^{-(l-v|s|)/\xi}\} \\ &= \frac{\sqrt{\gamma}}{\tau_B} \int_0^{\infty} ds e^{-2s/\tau_B} \min\{2, Ce^{-(l-vs)/\xi}\}. \end{aligned} \quad (\text{A2})$$

Here, we decompose the integration range as  $\int_0^{\infty} = \int_0^{l/2v} ds + \int_{l/2v}^{\infty} ds$ , where we use  $\min\{2, Ce^{-(l-vs)/\xi}\} \leq Ce^{-(l-vs)/\xi}$  in the former and  $\min\{2, Ce^{-(l-vs)/\xi}\} \leq 2$  in the latter. As a result,

$$\begin{aligned} \|\hat{L}_i - \hat{L}_i^l\| &\leq \frac{\sqrt{\gamma}}{\tau_B} \int_0^{l/2v} ds Ce^{-2s/\tau_B} e^{-(l-vs)/\xi} \\ &\quad + \frac{\sqrt{\gamma}}{\tau_B} \int_{l/2v}^{\infty} ds 2e^{-2s/\tau_B}. \end{aligned} \quad (\text{A3})$$

dqzzq In the first term of the right hand side,  $e^{-(l-vs)/\tau_B} \leq e^{-l/2\tau_B}$ , and hence

$$\begin{aligned} \|\hat{L}_i - \hat{L}_i^l\| &\leq \sqrt{\gamma} \left( \frac{C}{2} e^{-l/2\xi} + e^{-l/v\tau_B} \right) \\ &\leq \sqrt{\gamma} \left( 1 + \frac{C}{2} \right) \exp\left( -\frac{l}{\max\{2\xi, v\tau_B\}} \right), \end{aligned} \quad (\text{A4})$$

which is the desired inequality.

### Appendix B: Evaluation of $\bar{\epsilon}_{\text{md}}^{(1)}$

We evaluate an upper bound on

$$G(t, s, s') = \sum_{ij \in \Lambda} \|[\hat{A}_i(s') [\hat{O}_X^d(t), \hat{A}_i(s)], \hat{A}_j]\|, \quad (\text{B1})$$

which is used to evaluate  $\bar{\epsilon}_{\text{md}}^{(1)}$  given by Eq. (133). For a fixed  $t$ , the sum over  $i$  and  $j$  is decomposed into the three parts: (i)  $i, j \in X(t)$ , (ii)  $i \in X^c(t)$  and  $j$  with  $d(X, j) \leq d(X, i)$ , and (iii)  $j \in X^c(t)$  and  $i$  with  $d(X, i) \leq d(X, j)$ . Each contribution is denoted by  $G^{(i)}(t, s, s')$ ,  $G^{(ii)}(t, s, s')$ , and  $G^{(iii)}(t, s, s')$ , respectively.

(i)  $i, j \in X(t)$

In this case, we can simply evaluate  $G^{(i)}(t, s, s')$  as

$$G^{(i)}(t, s, s') \leq 2 \sum_{i, j \in X(t)} \|[\hat{O}_X^d(t), \hat{A}_i(s)]\|. \quad (\text{B2})$$

Here, we use the acceleration of dissipation, i.e. Eq. (39):

$$G^{(i)}(t, s, s') \leq 2\zeta(|X| + 2pvt)^2 e^{-a\tilde{\gamma}^{1/2}vt}. \quad (\text{B3})$$

By using Eq. (B3), we can evaluate the contribution to  $\bar{\epsilon}_{\text{md}}^{(1)}$  from  $i, j \in X(t)$ , which is denoted by  $\bar{\epsilon}_{\text{md}}^{(1-i)}$ . After an explicit calculation, we obtain

$$\begin{aligned} &\sum_{\nu=\pm 1} \int_0^{\infty} ds \int_{-\infty}^0 ds_2 \int_{-\infty}^{\nu s_1} ds_3 e^{-2(s_1-s_2)/\tau_B} e^{-2|\nu s_2-s_3|/\tau_B} \\ &= \frac{\tau_B^3}{4}. \end{aligned} \quad (\text{B4})$$

By using it and Eq. (B3) in Eq. (133), we find that

$$\bar{\epsilon}_{\text{md}}^{(1-i)} \leq \left( \frac{4p^2}{a^3} \tilde{\gamma}^{1/2} + \frac{2p|X|}{a^2} \tilde{\gamma} + \frac{|X|^2}{2a} \tilde{\gamma}^{3/2} \right) \tilde{\tau}_B \zeta, \quad (\text{B5})$$

which is of  $O(\tilde{\gamma}^{1/2} \tilde{\tau}_B)$ .

(ii)  $i \in X^c(t)$  and  $d(X, j) \leq d(X, i)$

In this case,  $G^{(ii)}(t, s, s')$  is evaluated as

$$G^{(ii)}(t, s, s') \leq 2 \sum_{i \in X^c(t)} \sum_{j: d(X, j) \leq d(X, i)} \|\hat{O}_X^d(t), \hat{A}_i(s)\|. \quad (\text{B6})$$

By putting  $d(X, i) = r$ , we have  $\sum_{i \in X^c(t)} \approx 2 \int_{pvt}^{\infty} dr$  and  $\sum_{j: d(X, j) \leq d(X, i)} = |X| + 2r$ . By using the Lieb-Robinson bound, we obtain

$$G^{(ii)}(t, s, s') \leq 4 \int_{pvt}^{\infty} dr (|X| + 2r) \min\{2, C e^{-[r-v(t+|s|)]/\xi}\}. \quad (\text{B7})$$

As for  $\min\{2, C e^{-[r-v(t+|s|)]/\xi}$ , when  $|s| \leq (p-2)t$ , we choose  $C e^{-[r-v(t+|s|)]/\xi}$ . When  $|s| > (p-2)t$ , we choose

$C e^{-[r-v(t+|s|)]/\xi}$  for  $r > pv|s|/(p-2)$ , and 2 for  $r \leq pv|s|/(p-2)$ . As a result, we obtain the following upper bound on  $G^{(ii)}(t, s, s')$ :

$$\begin{aligned} & 4I[|s| \leq (p-2)t] C \int_{pvt}^{\infty} dr (|X| + 2r) e^{-[r-v(t+|s|)]/\xi} \\ & + 4I[|s| > (p-2)t] \left[ \int_{pvt}^{\frac{p-2}{p}v|s|} dr 2(|X| + 2r) \right. \\ & \left. + \int_{\frac{p-2}{p}v|s|}^{\infty} dr (|X| + 2r) C e^{-[r-v(t+|s|)]/\xi} \right]. \quad (\text{B8}) \end{aligned}$$

By performing the integration over  $r$ , we obtain

$$\begin{aligned} G^{(ii)}(t, s, s') & \leq I[|s| \leq (p-2)t] \cdot 4C\xi(|X| + 2\xi + 2pvt)e^{-vt/\xi} \\ & + I[|s| > (p-2)t] \left\{ \frac{2p}{p-2}v|X| [ |s| - (p-2)t ] + 2 \left( \frac{p}{p-2} \right)^2 v^2 [ |s|^2 - (p-2)^2 t^2 ] + C\xi \left( |X| + 2\xi + \frac{2p}{p-2}v|s| \right) \right\}. \quad (\text{B9}) \end{aligned}$$

By substituting it into Eq. (133), after some calculations, we finally obtain  $\bar{\epsilon}_{\text{md}}^{(1-\text{ii})}$ , which corresponds to the contribution from  $i, j$  with  $i \in X^c(t)$  and  $d(X, j) \leq d(X, i)$ , as follows:

$$\begin{aligned} \bar{\epsilon}_{\text{md}}^{(1-\text{ii})} & \leq \tilde{\gamma}^2 \tilde{\tau}_{\text{B}} C \xi^2 [ |X| + 2(p+1)\xi ] \\ & + \frac{\tilde{\gamma}^2 \tilde{\tau}_{\text{B}}^2}{8(p-2)^3} [ C \xi (p-2)^2 (|X| + 2\xi) \\ & + p(p-2)(|X| + 2C\xi) \tilde{\tau}_{\text{B}} + 2p^2 \tilde{\tau}_{\text{B}}^2 ]. \quad (\text{B10}) \end{aligned}$$

For small  $\gamma$  or small  $\tau_{\text{B}}$ , we find  $\bar{\epsilon}_{\text{md}}^{(1-\text{ii})} \lesssim \tilde{\gamma}^2 \tilde{\tau}_{\text{B}}$ .

(iii)  $j \in X^c(t)$  and  $d(X, i) \leq d(X, j)$

By using  $[\hat{A}\hat{B}, \hat{C}] = \hat{A}[\hat{B}, \hat{C}] + [\hat{A}, \hat{C}]\hat{B}$  for arbitrary operators  $A, B$ , and  $C$ , we have

$$\begin{aligned} G^{(iii)}(t, s, s') & \leq \sum_{j \in X^c(t)} \sum_{i: d(X, i) \leq d(X, j)} \\ & \left( \|\hat{O}_X^d(t), \hat{A}_i(s)\| + \|\hat{A}_i(s'), \hat{A}_j\| \cdot \|\hat{O}_X^d(t), \hat{A}_i(s)\| \right). \quad (\text{B11}) \end{aligned}$$

By defining  $\hat{O}_X^d(t, s) := e^{i\hat{H}_S s} \hat{O}_X^d(t) e^{-i\hat{H}_S s}$ , we have

$$\begin{aligned} \|\hat{O}_X^d(t), \hat{A}_i(s)\| & = \|\hat{O}_X^d(t, s), \hat{A}_i\| \\ & \leq \|\hat{O}_{X,j}(t, -s), \hat{A}_i\| \\ & + 4\|\hat{O}_X^d(t, -s) - \hat{O}_{X,j}^d(t, -s)\|, \quad (\text{B12}) \end{aligned}$$

where  $\hat{O}_{X,j}(t, s)$  is the operator that is obtained by localizing  $\hat{O}_X^d(t, s)$  to the region  $X_j$  defined by Eq. (93); see Eq. (94). By using the Lieb-Robinson bound, we have

$$\begin{aligned} & \|\hat{O}_{X,j}(t, -s), \hat{A}_i\| \\ & \leq I[i \in X_j] \min\left\{2, C e^{-[\frac{d(X,j)}{p} - v|s|]/\xi}\right\}, \quad (\text{B13}) \end{aligned}$$

where we have used  $d(X_j, j) \geq d(X, j)/p$ , and

$$\begin{aligned} & \|\hat{O}_X^d(t, -s) - \hat{O}_{X,j}^d(t, -s)\| \\ & \leq \min\{2, C|X| e^{-[\frac{p-1}{p}d(X,j) - v(t+|s|)]/\xi}\}. \quad (\text{B14}) \end{aligned}$$

As for  $\|\hat{A}_i(s'), \hat{A}_j\| \cdot \|\hat{O}_X^d(t), \hat{A}_i(s)\|$  in Eq. (B11), the LR bound is used for  $\|\hat{A}_i(s'), \hat{A}_j\|$  when  $i \in X_j$ , and for  $\|\hat{O}_X^d(t), \hat{A}_i(s)\|$  when  $i \in X_j^c$ . We then have

$$\begin{aligned} & \|\hat{A}_i(s'), \hat{A}_j\| \cdot \|\hat{O}_X^d(t), \hat{A}_i(s)\| \\ & \leq 2I[i \in X_j] \min\{2, C e^{-[\frac{d(X,j)}{p} - v|s|]/\xi}\} \\ & + 2I[i \in X_j^c] \min\{2, C e^{-[\frac{p-1}{p}d(X,j) - v(t+|s|)]/\xi}\}, \quad (\text{B15}) \end{aligned}$$

where we used  $d(i, j) \geq d(X, j)/p$  for  $i \in X_j$  and  $d(X, i) \geq [(p-1)/p]d(X, j)$  for  $i \in X_j^c$ .

By substituting above results, i.e. Eqs. (B12) to (B15), into Eq. (B11), we obtain an upper bound on  $G^{(iii)}(t, s, s')$  and  $\bar{\epsilon}_{\text{md}}^{(1-\text{iii})}$  by using Eq. (133). Everything is finite, and we find

$$\bar{\epsilon}_{\text{md}}^{(1-\text{iii})} \lesssim \tilde{\gamma}^2 \tilde{\tau}_{\text{B}} \quad (\text{B16})$$

for small  $\gamma$  or small  $\tau_B$ .

### Appendix C: Evaluation of $\bar{\epsilon}_{\text{md}}^{(2)}$

We evaluate  $\bar{\epsilon}_{\text{md}}^{(2)}$ , which is defined by Eq. (136). As before, we decompose the sum over  $i, j \in \Lambda$  into the three parts: (i)  $i, j \in X(t)$ , (ii)  $i \in X^c(t)$  and  $d(X, j) \leq d(X, i)$ , and (iii)  $j \in X^c(t)$  and  $d(X, j) \leq d(X, i)$ . Each contribution is denoted by  $\bar{\epsilon}_{\text{md}}^{(2-i)}$ ,  $\bar{\epsilon}_{\text{md}}^{(2-ii)}$ , and  $\bar{\epsilon}_{\text{md}}^{(2-iii)}$ .

(i)  $i, j \in X(t)$

From Eq. (136), we have

$$\begin{aligned} \bar{\epsilon}_{\text{md}}^{(2-i)} &\leq \gamma \int_0^\infty dt \int_0^\infty ds e^{-2s/\tau_B} \sum_{i,j \in X(t)} \|\mathcal{L}_i^\dagger \hat{O}_X^d(t)\| \\ &= \frac{\gamma\tau_B}{2} \int_0^\infty dt \sum_{i \in X(t)} (|X| + 2pvt) \|\mathcal{L}_i^\dagger \hat{O}_X^d(t)\|. \end{aligned} \quad (\text{C1})$$

From the expression

$$\mathcal{L}_i^\dagger \hat{O}_X^d(t) = i[\hat{\Delta}_i, \hat{O}_X^d(t)] + \hat{L}_i^\dagger \hat{O}_X^d(t) \hat{L}_i - \frac{1}{2} \{\hat{L}_i^\dagger \hat{L}_i, \hat{O}_X^d(t)\}, \quad (\text{C2})$$

we obtain

$$\begin{aligned} \|\mathcal{L}_i^\dagger \hat{O}_X^d(t)\| &\leq \|[\hat{O}_X^d(t), \hat{\Delta}_i]\| + \frac{1}{2} \|\hat{L}_i^\dagger\| \cdot \|[\hat{O}_X^d(t), \hat{L}_i]\| \\ &\quad + \frac{1}{2} \|\hat{L}_i\| \cdot \|[\hat{O}_X^d(t), \hat{L}_i^\dagger]\|. \end{aligned} \quad (\text{C3})$$

By using  $\|\hat{L}_i\| = \|\hat{L}_i^\dagger\| \leq \sqrt{\gamma}/2$  and  $\|\hat{\Delta}_i\| \leq \gamma/8$ , and using the accelerated dissipation (39), we obtain

$$\begin{aligned} \bar{\epsilon}_{\text{md}}^{(2-i)} &\leq \frac{3}{16} \gamma^2 \int_0^\infty dt (|X| + 2pvt)^2 \\ &\quad \times (\eta e^{-a\tilde{\gamma}^{1/2}vt} + f(\tilde{\gamma}) e^{-b\tilde{\gamma}vt}), \end{aligned} \quad (\text{C4})$$

which is of  $O(\tilde{\gamma}^\alpha \tilde{\tau}_B)$  for small  $\gamma$  or small  $\tau_B$  (the same calculation as in Eq. (87)).

(ii)  $i \in X^c(t)$  and  $d(X, j) \leq d(X, i)$

Let us start with the expression

$$\begin{aligned} \bar{\epsilon}_{\text{md}}^{(2-ii)} &\leq \frac{\gamma\tau_B}{2} \int_0^\infty dt \sum_{i \in X^c(t)} \sum_{j: d(X, j) \leq d(X, i)} \\ &\quad \left( \|[\hat{O}_X^d(t), \hat{\Delta}_i]\| + \frac{\sqrt{\gamma}}{4} \|[\hat{O}_X^d(t), \hat{L}_i]\| + \frac{\sqrt{\gamma}}{4} \|[\hat{O}_X^d(t), \hat{L}_i^\dagger]\| \right). \end{aligned} \quad (\text{C5})$$

By putting  $l = d(X, i)/p$ , we do the following decompositions:

$$\begin{cases} \hat{\Delta}_i = \hat{\Delta}_i^l + (\hat{\Delta}_i - \hat{\Delta}_i^l), \\ \hat{L}_i = \hat{L}_i^l + (\hat{L}_i - \hat{L}_i^l), \end{cases} \quad (\text{C6})$$

where

$$\hat{\Delta}_i^l := \frac{1}{2i} \int_{-\infty}^\infty ds \int_{-\infty}^\infty ds' \text{sgn}(s-s') g_i(s) g_i(-s') \hat{A}_i^l(s) \hat{A}_i^l(s'). \quad (\text{C7})$$

The operators  $\hat{L}_i^l$  and  $\hat{A}_i^l$  were introduced in Sec. III: they are operators localized to the region  $R_{i,l} = \{j \in \Lambda : d(i, j) \leq l\}$ .

By substituting Eq. (C6) into Eq. (C5), we obtain

$$\begin{aligned} \bar{\epsilon}_{\text{md}}^{(2-ii)} &\leq \frac{\gamma\tau_B}{2} \int_0^\infty dt \sum_{i \in X^c(t)} \sum_{j: d(X, j) \leq d(X, i)} \\ &\quad \left( 2\|\hat{\Delta}_i - \hat{\Delta}_i^l\| + \|[\hat{O}_X^d(t), \hat{\Delta}_i^l]\| + \frac{\sqrt{\gamma}}{2} \|\hat{L}_i - \hat{L}_i^l\| \right. \\ &\quad \left. + \frac{\sqrt{\gamma}}{4} \|[\hat{O}_X^d(t), \hat{L}_i^l]\| + \frac{\sqrt{\gamma}}{4} \|[\hat{O}_X^d(t), \hat{L}_i^{\dagger l}]\| \right). \end{aligned} \quad (\text{C8})$$

As for  $\|\hat{\Delta}_i - \hat{\Delta}_i^l\|$ , by using the inequality  $\|\hat{A}_i(s) - \hat{A}_i^l(s)\| \leq \min\{2, C e^{-(l-|s|)/\xi}\}$ , which is obtained from the Lieb-Robinson bound, the operator norm is evaluated as

$$\begin{aligned} \|\hat{\Delta}_i - \hat{\Delta}_i^l\| &\leq \frac{\gamma}{8} (C e^{-l/2\xi} + 2e^{-l/v\tau_B}) \\ &\leq \frac{\gamma}{8} (C + 2) e^{-l/\eta}, \end{aligned} \quad (\text{C9})$$

where  $\eta = \max\{2\xi, v\tau_B\}$ . Next, as for  $\|[\hat{O}_X^d(t), \hat{\Delta}_i^l]\|$  in Eq. (C8), by using  $d(X, R_{i,l}) = d(X, i) - l = d(X, i)(p-1)/p$  and  $\|\hat{\Delta}_i^l\| \leq \gamma/8$ , the Lieb-Robinson bound yields

$$\|[\hat{O}_X^d(t), \hat{\Delta}_i^l]\| \leq \frac{\gamma C}{8} e^{-(\frac{p-1}{p}r - vt)/\xi}, \quad (\text{C10})$$

where  $r = d(X, i)$ . As for  $\|\hat{L}_i - \hat{L}_i^l\|$ , we can use Eq. (31):

$$\|\hat{L}_i - \hat{L}_i^l\| \leq \frac{\sqrt{\gamma}}{2} (C + 2) e^{-l/\eta}. \quad (\text{C11})$$

Finally, as for  $\|[\hat{O}_X^d(t), \hat{L}_i^l]\|$  and  $\|[\hat{O}_X^d(t), \hat{L}_i^{\dagger l}]\|$ , by using  $d(X, R_{i,l}) = d(X, i)(p-1)/p$  and  $\|\hat{L}_i\| \leq \sqrt{\gamma}/2$ , we have

$$\|[\hat{O}_X^d(t), \hat{L}_i^l]\| = \|[\hat{O}_X^d(t), \hat{L}_i^{\dagger l}]\| \leq \frac{\sqrt{\gamma} C}{2} e^{-(\frac{p-1}{p}r - vt)/\xi}. \quad (\text{C12})$$

By collecting them, Eq. (C8) is evaluated by approximating the sum over  $i$  by the integral:

$$\sum_{i \in X^c(t)} \sum_{j: d(X, j) \leq d(X, i)} \approx \int_{pvt}^\infty dr (|X| + 2r). \quad (\text{C13})$$

As a result, we find

$$\bar{\epsilon}_{\text{md}}^{(2-ii)} \lesssim \tilde{\gamma}^2 \tilde{\tau}_B \quad (\text{C14})$$

for small  $\gamma$  or small  $\tau_B$ .

(iii)  $j \in X^c(t)$  and  $d(X, j) \leq d(X, i)$

In Eq. (136), by using  $\hat{O}_{X,j}^d(t)$ , which is defined in Eq. (94), we have

$$\begin{aligned} \|\tilde{\mathcal{L}}_i^\dagger \hat{O}_X^d(t), \hat{A}_j(\nu s)\| &\leq \|\tilde{\mathcal{L}}_i^\dagger \hat{O}_{X,j}^d(t), \hat{A}_j(\nu s)\| \\ &\quad + 2\|\tilde{\mathcal{L}}_i^\dagger (\hat{O}_X^d(t) - \hat{O}_{X,j}^d(t))\|. \end{aligned} \quad (\text{C15})$$

We now decompose  $\tilde{\mathcal{L}}_i$  as

$$\tilde{\mathcal{L}}_i^\dagger = \sum_{l=0}^{\infty} \tilde{\mathcal{L}}_{i,l}^\dagger = \sum_{l'=0}^l \tilde{\mathcal{L}}_{i,l'}^\dagger + \sum_{l'=l+1}^{\infty} \tilde{\mathcal{L}}_{i,l'}^\dagger, \quad (\text{C16})$$

where  $l$  is chosen as

$$l = \left\lfloor \frac{d(X, j)}{4p} \right\rfloor, \quad (\text{C17})$$

and  $\tilde{\mathcal{L}}_{i,l}^\dagger$  is defined by

$$\tilde{\mathcal{L}}_{i,l}^\dagger \hat{O} = i[\hat{\Delta}_{i,l}, \hat{O}] + \mathcal{D}_{i,l} \hat{O} \quad (\text{C18})$$

for an arbitrary operator  $\hat{O}$ . See Eqs. (34) and (37) for the definition of  $\mathcal{D}_{i,l}$  and  $\hat{\Delta}_{i,l}$ , respectively. Equation (C15) is then bounded as

$$\begin{aligned} \|\tilde{\mathcal{L}}_i^\dagger \hat{O}_X^d(t), \hat{A}_j(\nu s)\| &\leq \left\| \left[ \sum_{l'=0}^l \tilde{\mathcal{L}}_{i,l'}^\dagger \hat{O}_{X,j}^d(t), \hat{A}_j(\nu s) \right] \right\| \\ &\quad + 2 \sum_{l'=l+1}^{\infty} \|\tilde{\mathcal{L}}_{i,l'}^\dagger\|_{\text{cb}} + 2\|\tilde{\mathcal{L}}_i^\dagger\|_{\text{cb}} \|\hat{O}_X^d(t) - \hat{O}_{X,j}^d(t)\|. \end{aligned} \quad (\text{C19})$$

If we choose  $l$  as in Eq. (C17), the distance between the support of  $\sum_{l'=0}^l \tilde{\mathcal{L}}_{i,l'}^\dagger \hat{O}_{X,j}^d(t)$  and the site  $j$  is greater than  $d(X, j)/2p$  (the former is  $R_{i,l} \cup X_j$  under the condition  $R_{i,l} \cap X_j \neq \emptyset$ ). Therefore, the Lieb-Robinson bound

yields

$$\begin{aligned} &\left\| \left[ \sum_{l'=0}^l \tilde{\mathcal{L}}_{i,l'}^\dagger \hat{O}_{X,j}^d(t), \hat{A}_j(\nu s) \right] \right\| \\ &\leq \left\| \sum_{l'=0}^l \tilde{\mathcal{L}}_{i,l'}^\dagger \hat{O}_{X,j}^d(t) \right\| \max\{2, C e^{-(\frac{d(X,j)}{2p} - \nu s)/\xi}\}. \end{aligned} \quad (\text{C20})$$

Since

$$\sum_{l'=0}^l \hat{O} = i[\hat{\Delta}_i^l, \hat{O}] + \hat{L}_i^{l\dagger} \hat{O} \hat{L}_i^l - \frac{1}{2} \{\hat{L}_i^{l\dagger} \hat{L}_i^l, \hat{O}\}, \quad (\text{C21})$$

we have

$$\left\| \sum_{l'=0}^l \tilde{\mathcal{L}}_{i,l'}^\dagger \hat{O}_{X,j}^d(t) \right\| \leq 2\|\hat{\Delta}_i^l\| + 2\|\hat{L}_i^l\|^2 \leq \frac{3}{4}\gamma, \quad (\text{C22})$$

where we have used  $\|\hat{\Delta}_i^l\| \leq \gamma/8$  and  $\|\hat{L}_i^l\| \leq \sqrt{\gamma}/2$ . By using Eqs. (36), (38) and (C17), we find that there is a constant  $D_\eta$  that depends solely on  $\eta$  and  $C$  such that

$$\sum_{l'=l+1}^{\infty} \|\tilde{\mathcal{L}}_{i,l'}\|_{\text{cb}} \leq \gamma D_\eta e^{-\eta d(X,j)/4p}. \quad (\text{C23})$$

As for  $\|\tilde{\mathcal{L}}_i\|_{\text{cb}} \|\hat{O}_X^d(t) - \hat{O}_{X,j}^d(t)\|$  in Eq. (C19),

$$\|\tilde{\mathcal{L}}_i\|_{\text{cb}} \leq 2\|\hat{\Delta}_i\| + 2\|\hat{L}_i\|^2 \leq \frac{3}{4}\gamma \quad (\text{C24})$$

and the Lieb-Robinson bound implies

$$\|\hat{O}_X^d(t) - \hat{O}_{X,j}^d(t)\| \leq C|X| e^{-[\frac{p-1}{p}d(X,j) - \nu t]/\xi}. \quad (\text{C25})$$

By collecting the above results, Eq. (C19) leads to

$$\begin{aligned} \|\tilde{\mathcal{L}}_i^\dagger \hat{O}_X^d(t), \hat{A}_j(\nu s)\| &\leq \frac{3}{4}\gamma \max\{2, C e^{-(\frac{d(X,j)}{2p} - \nu s)/\xi}\} \\ &\quad + \gamma D_\eta e^{-\eta d(X,j)/4p} + \frac{3}{2}\gamma C|X| e^{-[\frac{p-1}{p}d(X,j) - \nu t]/\xi}. \end{aligned} \quad (\text{C26})$$

By substituting it into Eq. (136) with the condition  $i \in X^c(t)$  and  $d(X, j) \leq d(X, i)$ , we can explicitly calculate an upper bound on  $\tilde{\epsilon}_{\text{md}}^{(2-\text{iii})}$ . We do not give lengthy calculations here, but it turns out that the upper bound is finite and

$$\tilde{\epsilon}_{\text{md}}^{(2-\text{iii})} \lesssim \tilde{\gamma}^2 \tilde{\tau}_B \quad (\text{C27})$$

for small  $\gamma$  or small  $\tau_B$ .

- 
- [1] G. Lindblad, On the generators of quantum dynamical semigroups, *Commun. Math. Phys.* **48**, 119 (1976).  
[2] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, Completely positive dynamical semigroups of N-level systems,

- J. Math. Phys.* **17**, 821 (1976).  
[3] H. Carmichael, *An Open Systems Approach to Quantum Optics* (Springer, Berlin, 1993).  
[4] W. H. Louisell, *Quantum Statistical Properties of Radia-*

- tion (Wiley-VCH, Berlin, 1990).
- [5] I. Bloch, J. Dalibard, and S. Nascimbène, Quantum simulations with ultracold quantum gases, *Nat. Phys.* **8**, 267 (2012).
  - [6] J. T. Barreiro, M. Müller, P. Schindler, D. Nigg, T. Monz, M. Chwalla, M. Hennrich, C. F. Roos, P. Zoller, and R. Blatt, An open-system quantum simulator with trapped ions, *Nature* **470**, 486–491 (2011).
  - [7] G. Barontini, R. Labouvie, F. Stubenrauch, A. Vogler, V. Guarrera, and H. Ott, Controlling the dynamics of an open many-body quantum system with localized dissipation, *Phys. Rev. Lett.* **110**, 035302 (2013).
  - [8] T. Tomita, S. Nakajima, I. Danshita, Y. Takasu, and Y. Takahashi, Observation of the Mott insulator to superfluid crossover of a driven-dissipative Bose-Hubbard system, *Sci. Adv.* **3**, e1701513 (2017).
  - [9] K. Bharti, A. Cervera-Lierta, T. H. Kyaw, T. Haug, S. Alperin-Lea, A. Anand, M. Degroote, H. Heimonen, J. S. Kottmann, T. Menke, W. K. Mok, S. Sim, L. C. Kwek, and A. Aspuru-Guzik, Noisy intermediate-scale quantum algorithms, *Rev. Mod. Phys.* **94**, 015004 (2022).
  - [10] Z. Cai, R. Babbush, S. C. Benjamin, S. Endo, W. J. Huggins, Y. Li, J. R. McClean, and T. E. O’Brien, Quantum error mitigation, *Rev. Mod. Phys.* **95**, 045005 (2023).
  - [11] S. Diehl, A. Tomadin, A. Micheli, R. Fazio, and P. Zoller, Dynamical Phase Transitions and Instabilities in Open Atomic Many-Body Systems, *Phys. Rev. Lett.* **105**, 015702 (2010).
  - [12] F. Minganti, A. Biella, N. Bartolo, and C. Ciuti, Spectral theory of Liouvillians for dissipative phase transitions, *Phys. Rev. A* **98**, 042118 (2018).
  - [13] Z. Cai and T. Barthel, Algebraic versus exponential decoherence in dissipative many-particle systems, *Phys. Rev. Lett.* **111**, 150403 (2013).
  - [14] R. Bouganne, M. Bosch Aguilera, A. Ghermaoui, J. Beugnon, and F. Gerbier, Anomalous decay of coherence in a dissipative many-body system, *Nat. Phys.* **16**, 21 (2020).
  - [15] T. Rakovszky, S. Gopalakrishnan, and C. von Keyserlingk, Defining stable phases of open quantum systems, *Phys. Rev. X* **14**, 041031 (2024).
  - [16] L. Pan, X. Chen, Y. Chen, and H. Zhai, Non-Hermitian linear response theory, *Nat. Phys.* **16**, 767 (2020).
  - [17] T. Mori, Liouvillian-gap analysis of open quantum many-body systems in the weak dissipation limit, *Phys. Rev. B* **109**, 064311 (2024).
  - [18] A. Nahum, J. Ruhman, and D. A. Huse, Dynamics of entanglement and transport in one-dimensional systems with quenched randomness, *Phys. Rev. B* **98**, 035118 (2018).
  - [19] C. W. Von Keyserlingk, T. Rakovszky, F. Pollmann, and S. L. Sondhi, Operator Hydrodynamics, OTOCs, and Entanglement Growth in Systems without Conservation Laws, *Phys. Rev. X* **8**, 021013 (2018).
  - [20] T. Shirai and T. Mori, Accelerated Decay due to Operator Spreading in Bulk-Dissipated Quantum Systems, *Phys. Rev. Lett.* **133**, 040201 (2024).
  - [21] T. Schuster and N. Y. Yao, Operator Growth in Open Quantum Systems, *Phys. Rev. Lett.* **131**, 160402 (2023).
  - [22] T. Prosen, Ruelle resonances in quantum many-body dynamics, *J. Phys. A: Math. Gen.* **35**, L737 (2002).
  - [23] A. M. García-García, L. Sá, J. J. Verbaarschot, and J. P. Zheng, Keldysh wormholes and anomalous relaxation in the dissipative Sachdev-Ye-Kitaev model, *Phys. Rev. D* **107**, 106006 (2023).
  - [24] T. Yoshimura and L. Sá, Robustness of Quantum Chaos and Anomalous Relaxation in Open Quantum Circuits, *Nat. Commun.* **15**, 9808 (2024).
  - [25] H. P. Breuer and F. Petruccione, *The theory of open quantum systems* (Oxford University Press, New York, 2002).
  - [26] E. B. Davies, Markovian Master Equations, *Commun. Math. Phys.* **39**, 91–110 (1974).
  - [27] H. Wichterich, M. J. Henrich, H. P. Breuer, J. Gemmer, and M. Michel, Modeling heat transport through completely positive maps, *Phys. Rev. E* **76**, 031115 (2007).
  - [28] T. Mori, Floquet States in Open Quantum Systems, *Annu. Rev. Condens. Matter Phys.* **14**, 35 (2023).
  - [29] B. Vacchini, Completely positive quantum dissipation, *Phys. Rev. Lett.* **84**, 1374 (2000).
  - [30] G. Schaller and T. Brandes, Preservation of positivity by dynamical coarse graining, *Phys. Rev. A* **78**, 022106 (2008).
  - [31] F. Nathan and M. S. Rudner, Universal Lindblad equation for open quantum systems, *Phys. Rev. B* **102**, 115109 (2020).
  - [32] F. Nathan and M. S. Rudner, Quantifying the accuracy of steady states obtained from the universal Lindblad equation, *Phys. Rev. B* **109**, 205140 (2024).
  - [33] C. Gneiting, Disorder-dressed quantum evolution, *Phys. Rev. B* **101**, 214203 (2020).
  - [34] T. Becker, L. N. Wu, and A. Eckardt, Lindbladian approximation beyond ultraweak coupling, *Phys. Rev. E* **104**, 014110 (2021).
  - [35] K. Shiraishi, M. Nakagawa, T. Mori, and M. Ueda, Quantum master equation for many-body systems: Derivation based on the Lieb-Robinson bound, [arXiv:2404.14067](https://arxiv.org/abs/2404.14067).
  - [36] L. van Hove, Quantum-mechanical perturbations giving rise to a statistical transport equation, *Physica* **21**, 517 (1954).
  - [37] K. Hepp and E. H. Lieb, Phase transitions in reservoir-driven open systems with applications to lasers and superconductors, *Helv. Phys. Acta* **46**, 573 (1973).
  - [38] P. F. Palmer, The singular coupling and weak coupling limits, *J. Math. Phys.* **18**, 527–529 (1977).
  - [39] B. Nachtergaele, A. Vershynina, and V. Zagrebnov, Lieb-Robinson bounds and existence of the thermodynamic limit for a class of irreversible quantum dynamics, *AMS Contemp. Math.* **552**, 161 (2011).
  - [40] T. Mori, Strong Markov dissipation in driven-dissipative quantum systems, *J. Stat. Phys.* **192**, 1 (2024).
  - [41] E. H. Lieb and D. W. Robinson, The finite group velocity of quantum spin systems, *Commun. Math. Phys.* **28**, 251–257 (1972).
  - [42] M. B. Hastings, Locality in quantum and Markov dynamics on lattices and networks, *Phys. Rev. Lett.* **93**, 140402 (2004).
  - [43] S. Bravyi, M. B. Hastings, and F. Verstraete, Lieb-Robinson bounds and the generation of correlations and topological quantum order, *Phys. Rev. Lett.* **97**, 050401 (2006), [arXiv:0603121](https://arxiv.org/abs/0603121) [quant-ph].
  - [44] B. Nachtergaele and R. Sims, Lieb-Robinson bounds and the exponential clustering theorem, *Commun. Math. Phys.* **265**, 119 (2006).
  - [45] D. Poulin, Lieb-Robinson Bound and Locality for General Markovian Quantum Dynamics, *Phys. Rev. Lett.* **104**, 190401 (2010).
  - [46] T. Barthel and M. Kliesch, Quasilocality and efficient

- simulation of Markovian quantum dynamics, *Phys. Rev. Lett.* **108**, 230504 (2012).
- [47] R. Sweke, J. Eisert, and M. Kastner, Lieb-Robinson bounds for open quantum systems with long-ranged interactions, *J. Phys. A Math. Theor.* **52**, 424003 (2019).
- [48] M. Žnidarič, Momentum-dependent quantum Ruelle-Pollicott resonances in translationally invariant many-body systems, *Phys. Rev. E* **110**, 054204 (2024).
- [49] J. A. Jacoby, D. A. Huse, and S. Gopalakrishnan, Spectral gaps of local quantum channels in the weak-dissipation limit, *Phys. Rev. B* **111**, 104303 (2025).
- [50] C. Zhang, L. Nie, and C. von Keyserlingk, Thermalization rates and quantum Ruelle-Pollicott resonances: insights from operator hydrodynamics, [arXiv:2409.17251](https://arxiv.org/abs/2409.17251).
- [51] T. Yoshimura and L. Sá, Theory of Irreversibility in Quantum Many-Body Systems, [arXiv:2501.06183](https://arxiv.org/abs/2501.06183).
- [52] L. D'Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics, *Adv. Phys.* **65**, 239 (2016).
- [53] T. Mori, T. N. Ikeda, E. Kaminishi, and M. Ueda, Thermalization and prethermalization in isolated quantum systems: A theoretical overview, *J. Phys. B At. Mol. Opt. Phys.* **51**, 112001 (2018).
- [54] H. Kim, T. N. Ikeda, and D. A. Huse, Testing whether all eigenstates obey the eigenstate thermalization hypothesis, *Phys. Rev. E* **90**, 052105 (2014).
- [55] Actually, Nathan and Rudner [31] considered the spectral norm, but we can repeat the same argument by using the trace norm without any change.
- [56] F. Verstraete, M. M. Wolf, and J. Ignacio Cirac, Quantum computation and quantum-state engineering driven by dissipation, *Nat. Phys.* **5**, 633 (2009).
- [57] P. M. Harrington, E. J. Mueller, and K. W. Murch, Engineered dissipation for quantum information science, *Nat. Rev. Phys.* **4**, 660 (2022).
- [58] J. Klinder, H. Keßler, M. Wolke, L. Mathey, and A. Hemmerich, Dynamical phase transition in the open Dicke model, *Proc. Natl. Acad. Sci. U. S. A.* **112**, 3290 (2015).
- [59] H. Keßler, P. Kongkhambut, C. Georges, L. Mathey, J. G. Cosme, and A. Hemmerich, Observation of a Dissipative Time Crystal, *Phys. Rev. Lett.* **127**, 043602 (2021).
- [60] T. Prosen and I. Pižorn, Quantum phase transition in a far-from-equilibrium steady state of an XY spin chain, *Phys. Rev. Lett.* **101**, 105701 (2008), [arXiv:0805.2878](https://arxiv.org/abs/0805.2878).
- [61] T. Prosen, Open XXZ spin chain: Nonequilibrium steady state and a strict bound on ballistic transport, *Phys. Rev. Lett.* **106**, 217206 (2011).
- [62] M. Žnidarič, Relaxation times of dissipative many-body quantum systems, *Phys. Rev. E* **92**, 042143 (2015).
- [63] T. Mori and T. Shirai, Resolving a Discrepancy between Liouvillian Gap and Relaxation Time in Boundary-Dissipated Quantum Many-Body Systems, *Phys. Rev. Lett.* **125**, 230604 (2020).