

Exploring the Rastall Gravity Cosmological Model using Gong-Zhang parameterization with Latest Observational Data and Deep Learning Techniques.

Vinod Kumar Bhardwaj^{1,a}, Manish Kalra^{2,b}

¹ Department of Mathematics, GLA University, Mathura-281 406, Uttar Pradesh, India

² Department of Electronics and Communication Engineering, GLA University, Mathura-281 406, Uttar Pradesh, India

^aE-mail:dr.vinodbhardwaj@gmail.com

^bE-mail:manishkalra2012@gmail.com

Abstract

In this study, we have explored the cosmological dynamics of an isotropic, homogeneous universe in Rastall gravity. For this purpose, we use the parameterization of the EoS parameter in the form $\omega(z) = \frac{\omega_0}{(z+1)}$ to derive the explicit solution of the field equations in Rastall gravity. We constrained the cosmological parameters for the derived model by the Markov Chain Monte Carlo (MCMC) approach utilizing OHD, BAO, and Pantheon datasets. We also constrained the model parameters using deep learning techniques and the CoLFI Python package. This paper introduces an innovative deep-learning approach for parameter inference. The deep learning method significantly surpasses the MCMC technique regarding optimal fit values, parameter uncertainties, and relationships among parameters. This conclusion is drawn from a comparative analysis of the two methodologies. Additionally, we determined the transition redshift $z_t = 1.094$, which signifies the shift in the Universe's model from an early deceleration phase to the present acceleration phase. The behavior of cosmological features of the model, like the equation of state (EoS) parameter, is discussed. The diagnosis of the model with diagnostic tools like statefinders, jerk parameter, and O_m diagnostics are presented and analyzed. The validation of the model's energy conditions is also examined.

1 Introduction

The experimental findings like $H(z)$ data, Ia Supernovae [1,2], Wilkinson Microwave Anisotropy Probe (WMAP) [3,4], Baryon Acoustic Oscillations (BAO) [5], Cosmic Microwave Background (CMB) [6], and Large Scale Structure (LSS) [7] confirmed the present era of accelerating universe. Although the cosmological models based on GR show an agreement with various observational findings [1–3], still, the observed accelerated expansion of present cosmos argue the validity of General Relativity (GR) on large scales [8]. An exotic cosmic fluid with large negative pressure having a mysterious kind of energy density (known as dark energy (DE)) is postulated to claim this observed acceleration. Because of its repulsive nature, the cosmological constant makes it an appropriate substitute for DE [8,9]. However, cosmic models with a cosmological constant face the complications of cosmic fluke and fine-tuning. Apart from these issues, the Λ CDM model also suffers from H_0 tension. To address these issues and to describe the origin

and nature of dark energy, several dynamical dark energy models, including modified gravity models, models based on extra dimensions, and scalar field models, have been proposed in the literature [10–16].

The principle of equivalence is a cornerstone of Einstein’s theory of general relativity. Nevertheless, there are instances where this rule is violated, resulting in modifications of gravity that deviate from GR anticipations. The divergence-free tensor, which exhibits minimal correlation with the geometry of spacetime, has been utilized to establish an energy-momentum source in General Relativity and its variations in modified gravitational theories [17, 18]. However, it is argued that the process of particle formation violates the property of the energy-momentum tensor (EMT), which leads to the energy-momentum conservation equation ($\nabla_{\nu}^{\mu\nu} = 0$) [19–22]. Therefore, it is permissible to disregard the EMT conservation requirement in favor of a different gravitational theory. In the present scenario, the utilization of modified gravity theories [23–25] is both advantageous and attractive in tackling the significant challenges associated with the typical cosmological models, including ‘dark energy (DE)’ [1, 26] and ‘dark matter (DM)’ [27, 28]. It is essential to recognize that any alternative to General Relativity must be valid. Numerous researchers have extensively explored modified theories of gravity [29–35].

Rastall’s gravity proposed by Peter Rastall in 1972 [36] is one of the interesting theory among the alternative gravitational theories. The law of conservation of energy-momentum tensor (EMT) is analyzed under circumstances, such as in Minkowski flat space-time or specifically in weak gravitational arenas in Rastall’s framework. This raises questions regarding its relevance in curved space-time. The postulate that the covariant derivative of the energy-momentum tensor (EMT) is zero is no longer valid in Rastall’s theory. Rather, we examine a vector field that corresponds to the gradient of the Ricci scalar, represented as $\nabla_{\nu} T^{\mu\nu} \propto \nabla^{\mu} R$. In Rastall’s framework, the curvature is coupled with the violation of the energy and momentum conservation laws, and it can be interpreted as a conventional representation of quantum phenomena [37]. Additionally, the curvature-matter theory of gravity defines a non-minimal interaction between matter and geometry, resulting in the failure of the traditional conservation law for energy-momentum [38–41].

Recently, significant advancements have been achieved in the investigation of various aspects of Rastall gravity including the role of the Rastall constraints in DE fluctuations [42], and its relationship with the ‘Brans-Dicke scalar field’ [43], and the structure of neutron stars [44]. Researchers have also examined various cosmic eras within this theoretical framework offering an extensive analysis of both the theoretic and experimental dimensions of Rastall gravity [45–49]. Remarkably, Rastall gravity is unaffected by issues related with age and entropy [50], and it provides clarifications for both accelerated expansion scenario and inflationary eras [45, 51]. Moradpour et al. [51], suggest that a non-minimal coupling between pressure-less matter and geometry within Rastall gravity may mimic the effects of dark energy, which could explain the current phase of cosmic acceleration. The Rastall gravity framework has been rigorously analyzed throughout the cosmic era, encompassing early inflation, the matter-dominated era, and the phase of accelerated expansion [52]. Additionally, the cosmic model of Rastall gravity introduced in [53] is compatible with the Λ CDM model in the later stages of the cosmos. A proposed Rastall parameter may also serve as a viable solution to the initial singularity issue [52].

By employing modern Deep Learning approaches, the aim was to analyze the accelerated

expansion of the universe while simultaneously estimating cosmological parameters. Utilizing CoLFI python package [54], the cosmological parameters are estimated using Artificial Neural Networks (ANN), Mixture Density Networks (MDN), and Mixture of Gaussians (MNN) techniques [55–58]. This advancement significantly facilitated the understanding of conditional probability densities derived from observational data and posterior distributions [54–56]. To enhance the efficiency of neural network training and improve the accuracy of parameter predictions, hyper ellipsoid parameters were introduced [57, 58]. A comparative study between our neural network techniques and the conventional Markov Chain Monte Carlo (MCMC) approach revealed that MNN produced results nearly identical to those of MCMC, underscoring its effectiveness and reliability.

The proposed model of the universe accounts for late-time acceleration without the need for dark energy and avoids issues associated with the cosmological constant. Therefore, it is essential to consider current observational data when simulating Rastall gravity. We have shown that neural network methods can effectively estimate cosmological parameters, serving as a practical alternative to MCMC techniques. The Rastall gravity framework facilitates the exploration of various gravity theories and their influence on cosmic expansion through the use of ANN, MDN, and MNN methodologies. This research is part of a broader study integrating machine learning with cosmology, demonstrating the capability of machine learning to address intricate cosmic dynamics and expansion challenges. Our evaluation of density, cosmic pressure, the equation of state, and deceleration parameters lends credibility to our findings. The given paper is organized in the following manner. A solution of Rastall gravity field equations using EsS parameterization is mentioned in Section 2. The observational constraints on model parameters using the MCMC approach and deep learning utilizing OHD, BAO, and Pantheon datasets are presented in Section 3. Some cosmological features of the model, like cosmic pressure, energy density, and deceleration parameter, along with diagnostic tools like statefinders, jerk parameter, and O_m diagnostic are presented and analyzed in Section 4. The energy conditions on the viability of the model are discussed in Section 5. The concluding summary of the work is presented in Section 6.

2 Metric and the Field Equations

The modified theory based on Rastall’s gravity is one of the intriguing subclasses among the alternative theories of gravity [36]. Our goal is to extract solutions, carefully taking into account both the source of the matter and the gravitational background. The Bianchi identities remain valid in Rastall’s gravity, but the conservation of the stress-energy tensor of the gravity source is violated (since $T_{;i}^{ij} = \lambda R^{;j}$, where R is the Ricci scalar and λ is the Rastall parameter). The covariant conservation equation of the energy-momentum tensor in this GR modification is expressed as $\nabla_j T^{ij} = 0$ and can be expressed in more generalized form as $\nabla_j T^{ij} = u^i$ [36, 46, 59], here to retrieve this again in GR, the right side should be zero. Consequently, vector u^i is considered as $u^i = \lambda \nabla^i R$. The field equations in the framework of Rastall gravity can be expressed as [36, 59]

$$R_{ij} - \frac{1}{2}(1 - 2k\lambda)g_{ij}R = kT_{ij} \quad (1)$$

where R_{ij} is the Ricci-tensor, g_{ij} is metric-tensor, and T_{ij} is the energy-momentum tensor respectively. k denotes the gravitational constant, which is to be determined in consistent with

Newtonian limit. The Einstein field equations can be retrieve for $\lambda = 0$ and $k = 8\pi$ whenever $T_{;i}^{ij} = 0$. Then, the trace of Eq. (1) can be recasts as [59]

$$(4k\lambda - 1)R = kT \quad (2)$$

For either $R = 0$ or $k\lambda = \frac{1}{4}$, we get a trace-less energy-momentum tensor ($T = 0$). Rastall gravity has similar characteristics to GR for $R = 0$, while the trace of energy-momentum reduces to zero for $k\lambda = \frac{1}{4}$. For $k\lambda \neq \frac{1}{4}$, the field equation with the cosmological constant Λ is given by

$$G_{ij} + \Lambda g_{ij} + k\lambda g_{ij}R = kT_{ij} \quad (3)$$

where G_{ij} denotes the standard Einstein tensor. The Friedmann-Lemaître-Robertson-Walker (FLRW) space-time metric for a homogeneous and isotropic universe is considered as

$$ds^2 = -dt^2 + a^2(t) \left(\frac{dr^2}{1 - Kr^2} + r^2 d\Omega^2 \right). \quad (4)$$

where $d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2$; $a(t)$ denotes the time-dependent scale factor; and spatial parameter K represents the open, flat, and closed space sections for the values -1 , 0 , and $+1$ respectively. For the perfect fluid, the field equations in the Rastall gravity are expressed as [36, 46, 47, 59]

$$3(1 - 4k\lambda)H^2 - 6\lambda k\dot{H} - 3(2\lambda k - 1)Ka^{-2} = k\rho, \quad (5)$$

$$-3(4\lambda k - 1)H^2 + 2(1 - 3\lambda k)\dot{H} - (6\lambda k - 1)Ka^{-2} = -kp. \quad (6)$$

where p and ρ respectively, denote the cosmic pressure and energy density, and over dot denotes the derivative with respect to time t . The scale factor a and Hubble parameter H are related as $H = \frac{\dot{a}}{a}$. The isotropic pressure for the model is connected with energy density by relation $p = \omega\rho$ called as ‘equation of state (EoS)’. By applying the Bianchi identity ($G_{ij}^{;j} = 0$, the continuity equation is derived as [59]

$$(3k\lambda - 1)\dot{\rho} + 3k\lambda\dot{p} + 3(4k\lambda - 1)H(\rho + p) = 0. \quad (7)$$

Since most of cosmological observational datasets are exist in terms of redshift z rather than cosmic time t . The scale factor $a(t)$ is connected with redshift z using the relation $\frac{a_0}{a} = (1 + z)$, where a_0 is the present value of scale factor, which is considered as 1 in the present analysis. Using this relation we also get $\dot{H}(t) = -(1 + z)H(z)H'(z)$, where $H'(z) = \frac{dH}{dz}$. In terms of redshift z , the field equations (5) and (6) can be recast as

$$\begin{aligned} 3(1 - 4\lambda k)H^2 + 6\lambda k(z + 1)HH'(z) \\ - 3K(2\lambda k - 1)(z + 1)^2 = k\rho, \end{aligned} \quad (8)$$

$$\begin{aligned} 3(1 - 4\lambda k)H^2 + 2(3\lambda k - 1)(z + 1)HH'(z) \\ + (1 - 6\lambda k)(z + 1)^2K = -kp. \end{aligned} \quad (9)$$

For flat universe ($K = 0$), using EOS relation ($p = \omega\rho$), from equations (8) and (9), we get the expression

$$3(1 - 4\lambda)[1 + \omega(z)]H^2(z) - [2 - 6\lambda(1 + \omega(z))](1 + z)H(z)\frac{dH}{dz} = 0. \quad (10)$$

where for numerical simplification k is considered as 1. Since above expression consist of two variables ω and $H(z)$, thus to get explicit solution we consider the redshift parameterization of EoS parameter $\omega(z)$ proposed by Gong-Zhang [60] in th form

$$\omega(z) = \frac{\omega_0}{(1+z)} \quad (11)$$

with ω_0 as the present value of the EoS parameter. Using the Gong-Zhang parametrization [60] for the EoS of dark energy (DE), one can efficiently describe distinct cosmic scenarios of an expanding universe. In the early time, as the z approaches to infinity, EoS parameter tends to zero ($\omega \sim 0$) that represents the big-bang scenario of universe. At $z = 0$, the present value of the EoS parameter is ω_0 . At $z \rightarrow -1$, the EoS parameter approaches negative infinity, which indicates towards the dark energy dominated scenario in the remote future.

From equations (10) and (11), the expression of the Hubble parameter can be derived as

$$H(z) = H_0 \left[(z+1)^{3\lambda-1} \left(\frac{(3\lambda-1)z}{3\lambda(\omega_0+1)-1} + 1 \right) \right]^{\frac{4\lambda-1}{2\lambda(3\lambda-1)}} \quad (12)$$

Here, H_0 is the present value of Hubble parameter. The deceleration parameter of the model is read as

$$q = -1 + \frac{d}{dt} \left(\frac{1}{H} \right) \quad (13)$$

3 Data & Methodology

In recent years, Bayesian inference has increasingly been utilized for parameter estimation and model comparison in the field of cosmological research. In this section, our goal is to constrain the free parameters of the model using observational data sets and the χ^2 -minimization method. To find the values of ω_0 , λ , and H_0 in the model, we utilize the data sets listed below:

OHD: For constraining the model parameters, we have employed the 57 uncorrelated Hubble observations $H(z)$ that lie in the redshift range between 0.07 and 2.36 [63–65]. These measurements are known as Cosmic Chronometers and are evaluated by applying the techniques of differential age.

Pantheon: In this work, we use the recent Pantheon compilation of SN Ia data in the redshift range $0.001 < z < 2.26$ [66]. The light curves of the SALT2 model are now uniformly fitted along with the light curves of SNe Ia. By using such 18 different surveys, the dataset of Pantheon is compiled [67].

BAO data: We also considered the six BAO data points [68–70]. For the BAO sample, the predictions from a sample of Galaxy Surveys like SDSS DR7 and 6dF, and WiggleZ have been utilized [68–70]. The angular diameter distance for the sample is defined as $D_A = \frac{D_L}{(1+z)^2}$, where D_L indicates the proper angular diameter distance [71], and the dilation scale is described by $D_V(z) = \left[D_L^2(z) * (1+z)^2 * \frac{cz}{H(z)} \right]^{1/3}$.

z_{BAO}	0.106	0.2	0.35	0.44	0.6	0.73
$\Upsilon(z)$	30.95 ± 1.46	17.55 ± 0.60	10.11 ± 0.37	8.44 ± 0.67	6.69 ± 0.33	5.45 ± 0.31

Table 1: The values of $\Upsilon(z)$ for different points of z_{BAO}

Here, $\Upsilon(z) = d_A(z_*)/D_V(z_{BAO})$ and $z_* \approx 1091$.

For limiting the parameters of the model, the chi-square estimator for the BAO sample is described in the following form [71–74].

$$\chi_{BAO}^2 = X^T C^{-1} X \quad (14)$$

where

$$X = \begin{pmatrix} \frac{d_A(z_*)}{D_V(0.106)} - 30.95 \\ \frac{d_A(z_*)}{D_V(0.20)} - 17.55 \\ \frac{d_A(z_*)}{D_V(0.35)} - 10.11 \\ \frac{d_A(z_*)}{D_V(0.44)} - 8.44 \\ \frac{d_A(z_*)}{D_V(0.60)} - 6.69 \\ \frac{d_A(z_*)}{D_V(0.73)} - 5.45 \end{pmatrix}$$

The model parameters ω_0 , λ , and H_0 of this model can be assessed using observational data and statistical methods. E_{obs} represents the value of any observational data, while E_{th} represents the values that have been theoretically estimated in the model of the Universe. By using statistical methods to compare the two values E_{obs} and E_{th} , the model's parameters can be determined. To obtain the most accurate estimates, we utilized the χ^2 estimator. If the standard error in the observed values is denoted by σ , the formula for the χ^2 estimator is expressed as.

$$\chi^2 = \sum_{i=1}^N \left[\frac{E_{th}(z_i) - E_{obs}(z_i)}{\sigma_i} \right]^2 \quad (15)$$

where, $E_{th}(z_i)$ represents the theoretical values of the parameter, while $E_{obs}(z_i)$ represents the observed values. σ_i is the error and N is the number of data points.

The combined χ^2 estimator is read as

$$\chi_{combined}^2 = \chi_{OHD}^2 + \chi_{Pantheon}^2 + \chi_{BAO}^2 \quad (16)$$

The χ_{tot}^2 statistic can be minimized to find the parameter value that best fits the combined sample of OHD, BAO, and Pantheon datasets. By taking maximum likelihood approach into account, the total likelihood function $\mathcal{L}_{tot} = \exp(-\chi_{tot}^2/2)$ may be calculated as the product of individual likelihood functions of each dataset expressed in the form $\mathcal{L}_{tot} = \mathcal{L}_{BAO} * \mathcal{L}_{Pantheon} * \mathcal{L}_{CC}$. The likelihood function $\mathcal{L}_{tot}(x^*)$ is maximized or, alternatively $\chi_{tot}^2(x^*) = -2 \ln \mathcal{L}_{tot}(x^*)$ is minimized to get the most plausible values of parameters. For the set of cosmic parameters (pointed at x^*), the 1σ and 2σ contours are constrained and bounded respectively by $\chi_{tot}^2(x) = \chi_{tot}^2(x^*) + 2.3$ and $\chi_{tot}^2(x) = \chi_{tot}^2(x^*) + 6.17$. We get best-fit parameter values for the derived model by minimizing the χ^2 statistic.






Figure 1: One-dimensional marginalized distribution, and 2D contours with 1σ and 2σ confidence levels for the derived model of the Universe with OHD, Pantheon, BAO, and joint datasets.

Neural Networks


Neural networks become very relevant when computing advances led to the emergence of a new scientific field called deep learning, which is only concerned with the study of artificial neural networks. Currently, there are various kinds of neural networks—such as recurrent networks commonly applied to time series, convolutional neural networks excelling in image processing, auto-encoders employed for image denoising, and contemporary generative adversarial networks. In this study, we concentrate on the fundamental deep neural network, commonly referred to as the multi-layer perceptron (MLP).



(a) A , η , and H_0 1σ and 2σ contours from 57 $H(z)$ data points using ANN.



(b) The relationship between steps and best-fit values and 1σ error of model parameters using ANN technique. The solid black line and Grey-shaded sections display the best-fit values, red circle with error bars are the results estimated by ANN Approach.




(c) Plot of Losses of the training and validation sets. The training set consists of 3000 samples, while validation set contains 500 samples.

Figure 2: Observational analysis using ANN model


3.1 Artificial Neural Network (ANN)

An artificial neural network (ANN), commonly referred to as a neural network (NN), is a computational framework that draws inspiration from the architecture and functionality of biological neural networks. Typically, it comprises an input layer, one or more hidden layers, and an output layer. In the context of estimating cosmological parameters, observational data is introduced to the input layer, where it subsequently traverses through the hidden layers before the cosmological parameters are produced at the output layer. Each layer processes a vector,


with its components known as neurons, received from the preceding layer. This processing involves applying a linear transformation followed by a nonlinear activation function, after which the resultant data is forwarded to the subsequent layer. In general, no theory exists to specify the precise number of neurons that ought to be employed in every hidden layer. The model architecture employed by Wang et al. (2020a) [56] is adopted here, whereby the number of neurons in each hidden layer is reduced according to the layer index.



(a) A , η , and H_0 1σ and 2σ contours from 57 $H(z)$ data points using MDN.



(b) The relationship between steps and best-fit values and 1σ error of model parameters using MDN technique. The solid black line and Grey-shaded sections display the best-fit values, red circle with error bars are the results estimated by MDN Approach.




(c) Plot of Losses of the training and validation sets. The training set consists of 3000 samples, while validation set contains 500 samples.


Figure 3: Observational analysis using MDN model

3.2 Mixture Density Network (MDN)


A Mixture Density Network (MDN) combines an ANN with a mixture model. The mixture model described is a probabilistic framework that posits that every data point arises from a combination of a limited number of distributions with unknown parameters, with the distribution potentially being any type of probability distribution (such as Gaussian or beta distribution). The goal of parameter constraint with the MDN is to acquire the parameters from a mixture model [56]. Consequently, an MDN that utilizes Gaussian components effectively learns a relationship between the measurements and the parameters of the Gaussian mixture model. As a result, the parameters of a Gaussian mixture MDN can be adjusted by minimizing the loss function.



(a) A , η , and H_0 1σ and 2σ contours from 57 $H(z)$ data points using MNN.



(b) The relationship between steps and best-fit values and 1σ error of model parameters using MNN technique. The solid black line and Grey-shaded sections display the best-fit values, red circle with error bars are the results estimated by MNN Approach.



(c) Plot of Losses of the training and validation sets. The training set consists of 3000 samples, while validation set contains 500 samples.

Figure 4: Observational analysis using MNN model

3.3 Mixture Neural Network (MNN)

The fundamental concept of the MNN approach is the belief that the posterior distribution consists of a combination of certain unknown distributions, which subsequently learns the mixture model through an ANN. Thus, we must initially acquire the parameters of the mixture model and subsequently derive the posterior distribution by producing samples according to the mixture model. However, for cosmological parameters that could differ from Gaussian distribution, several components must be utilized to achieve the accurate posterior distribution [56].

Training the network will require additional time and lead to greater instability, complicating the process of learning the parameters of the mixture model.

4 Examination of the model's cosmological features




Figure 5: (a) plot of energy density, (b) plot of cosmic pressure.

$$\rho = \frac{3H_0^2(4\lambda - 1)(z + 1) \left[(z + 1)^{3\lambda-1} \left(\frac{(3\lambda-1)z}{3\lambda\omega_0+3\lambda-1} + 1 \right) \right]^{\frac{1-4\lambda}{\lambda-3\lambda^2}}}{3\lambda\omega_0 + (3\lambda - 1)(z + 1)} \quad (17)$$

$$p = \frac{3H_0^2(4\lambda - 1)\omega_0 \left[(z + 1)^{3\lambda-1} \left(\frac{(3\lambda-1)z}{3\lambda\omega_0+3\lambda-1} + 1 \right) \right]^{\frac{1-4\lambda}{\lambda-3\lambda^2}}}{3\lambda\omega_0 + (3\lambda - 1)(z + 1)} \quad (18)$$

For the derived model in Rastall gravity, the energy density is positive through the entire evolution of the universe. The trajectory of energy density can be seen in Figure 5 (a). The cosmic pressure for the proposed model is negative in the entire evolution of the universe, which represents the dark energy dominance era. The behavior of cosmic pressure for the proposed model is plotted in Figure 5(b).




Figure 6: plot of EOS parameter

For $\omega > 0$, the EOS parameter describes the Big Bang era of the universe, and $\omega = 0$ represents a matter-dominated scenario of the cosmos. In this case, dark matter is a non-relativistic matter for which $\omega = 0$, while for a relativistic matter like radiation $\omega = 1/3$. The model lies in the quintessence era for $-1 < \omega \leq 0$. In the case of $\omega = -1$, the EOS parameter approaches to the Λ CDM model. For $\omega < -1$, a phantom era is observed. We have plotted the trajectories of the EOS parameter in Figure 6 for the derived model. This figure shows that the model begins with quintessence behavior and approaches to phantom era in the late-time phase. The present value of the EoS parameter at $z = 0$ is estimated as $\omega_0 = -0.683$ for a joint dataset of OHD, BAO, and Pantheon. This nicely agrees with the latest observational findings.

4.1 Deceleration parameter

The deceleration parameter (DP) from Hubble parameter expression can be formulated as $q = -1 + \frac{(1+z)}{H(z)} \frac{dH(z)}{dz}$. Thus, for the suggested model DP q can be derived as:

$$q = \frac{(6\lambda - 3)\omega_0 + (6\lambda - 1)(z + 1)}{6\lambda\omega_0 + 2(3\lambda - 1)(z + 1)} \quad (19)$$




Figure 7: plot of Deceleration parameter

In the present analysis, the model depicts a flipping nature of the universe from a deceleration era to a current expansion scenario that indicates a dark energy (DE) dominance in the present while a matter dominance in the past. For the joint data-set of OHD, BAO, and Pantheon, the proposed model shows a transition behavior with the present value of DP as $q_0 = -0.532$ and signature flipping occurring at $z_t = 1.094$, as shown in Figure 7. The results obtained from the model align well with the latest experimental findings [62–64, 75–78].

4.2 Statefinders diagnostic




Figure 8: (a) plot of r vs q , (b) plot of s vs q , (c) Plot of r vs s .

Two novel parameters r and s , named as statefinders, are introduced to distinguish the various dark energy cosmological models. The statefinder pair assists in improving the precision of model predictions by pinpointing the evolutionary path in the $r - s$ plane. By considering various forms of dark energy, as mentioned in the literature [79–82], the distinction between the proposed cosmological model and the Λ CDM model can be clearly distinguished on the $(r - s)$ plane [79, 80]. For the derived model, the parameters r and s in the standard forms can be derived as:

$$\begin{aligned}
 r &= 2q^2 + q - \frac{\dot{q}}{H} \\
 &= \frac{9(6\lambda^2 - 5\lambda + 1)\omega_0^2 + 6(18\lambda^2 - 9\lambda + 1)\omega_0(z+1) + (54\lambda^2 - 21\lambda + 2)(z+1)^2}{2[3\lambda\omega_0 + (3\lambda - 1)(z+1)]^2} \quad (20)
 \end{aligned}$$

$$\begin{aligned}
 s &= \frac{r - 1}{3(q - \frac{1}{2})} \\
 &= \frac{(4\lambda - 1)[3(\lambda - 1)\omega_0^2 + 2(3\lambda - 1)\omega_0(z+1) + 3\lambda(z+1)^2]}{3((\lambda - 1)\omega_0 + \lambda(z+1))(3\lambda\omega_0 + (3\lambda - 1)(z+1))} \quad (21)
 \end{aligned}$$

Figure 8 illustrates the features of the suggested model in the $r - s$ plane using formulas for r and s . The plot shows that the suggested model is situated within the Chaplygin gas region

($r > 1, s < 0$) and converges to the Λ CDM point ($r = 1, s = 0$) at the late passes of the timeline. For the joint dataset of CC, BAO, and Pantheon, the present values of parameters (r, s) are computed as (0.035, 0.312). Hence, the derived model shows features of an accelerating cosmic model other than Λ CDM [79, 81–83].

4.3 Jerk Parameter

The jerk parameter (j) serves as a widely utilized tool in the field of cosmology to diagnose the various cosmological models. The concept arises from the belief that there must be a sudden jolt to shift the universe from a decelerating to an accelerating phase. The jerk is defined as the acceleration's rate of change over time from a physical perspective. It originates from the third-order term of Taylor's series expansion of the scale factor a centered at a_0 in cosmology. This parameter helps to differentiate between cosmological models that are kinematically degenerate. The involvement of the third order derivative of the scale factor results in increased accuracy when describing the expansion of the universe compared to the Hubble parameter. The parameter j for the suggested model can be expressed as [84]:

$$j = 1 - (1 + z) \frac{H'(z)}{H(z)} + \frac{1}{2} (1 + z)^2 \left[\frac{H''(z)}{H(z)} \right]^2 \quad (22)$$

Here $H'(z)$ and $H''(z)$ respectively denote the first and second order derivatives of parameter $H(z)$ w.r.to redshift z . For the given model parameter j can be recast as:

$$j = - \frac{9(1 - 4\lambda)^2 [(6\lambda - 3)\omega_0^2 + 2(6\lambda - 1)\omega_0(z + 1) + (6\lambda - 1)(z + 1)^2]^2}{32(z + 1)^2 [3\lambda\omega_0 + (3\lambda - 1)(z + 1)]^4} + \frac{(3 - 6\lambda)\omega_0 - (6\lambda - 1)(z + 1)}{6\lambda\omega_0 + 2(3\lambda - 1)(z + 1)} \quad (23)$$




Figure 9: Behaviour of jerk parameter j with z .

Figure 9 depicts the graphical nature of parameter j as the redshift function for a joint dataset of OHD, BAO, and Pantheon observational datasets. Earlier theoretical studies suggest that positive values of the jerk parameter indicate an accelerated expansion of the universe [85–87]. Based on our model, the jerk parameter is initially negative and then increases, eventually convergent to positive over time. In the existing analysis, the present value of the parameter j_0 is determined to be 0.824 for the joint observational dataset.

4.4 O_m diagnostic

The parameter O_m serves as an additional diagnostic tool in cosmology, as introduced by Sahni et al. [88]. This parameter can be derived solely from the Hubble parameter, eliminating the need for $H'(z)$ or other pertinent data, which consequently reduces the likelihood of errors. Its ability to allow for reconstruction through both non-parametric as well as parametric approaches has contributed to the popularity of O_m in cosmological investigations. Furthermore, it is important to mention that the O_m can differentiate between various cosmic models even in the absence of information regarding the matter density parameter and the equation of state (EoS) [88, 89]. According to the standard formulation [88], the expression for $O_m(z)$ in the given model can be expressed as follows:

$$O_m(z) = \frac{\left((z+1)^{3\lambda-1} \left(\frac{(3\lambda-1)z}{3\lambda\omega_0+3\lambda-1} + 1 \right) \right)^{\frac{1-4\lambda}{\lambda-3\lambda^2}} - 1}{z^3 + 3z^2 + 3z} \quad (24)$$




Figure 10: Evaluation of $O_m(z)$.

The evolution trajectory of $O_m(z)$ has been displayed in Figure 10 using a joint dataset. The incline of the diagnostic $O_m(z)$ serves as a crucial indicator for determining the type of dark energy (DE) cosmic models. Specifically, a negative slope suggests a quintessence-like model, while a positive slope indicates a phantom-like cosmic model. Furthermore, a zero curvature in the slope of $O_m(z)$ implies a Λ CDM type of DE model. The $O_m(z)$ analysis of our model reveals a positive slope, indicating a phantom-like scenario that supports the belief of a dark energy-dominated universe. The suggested cosmological model is distinguished from the standard Λ CDM model by its positive curvature [34, 85, 89].

5 Energy Conditions

We have discussed the physical viability of the derived model via the evolution of energy conditions (ECs) [90, 91]. The energy conditions for cosmic model in Rastall theory are defined as [90–92]: (i) Weak energy conditions (WEC) if $\rho \geq 0$, (ii) Dominant energy conditions (DEC) if $\rho - p \geq 0$, (iii) Null energy conditions (NEC) if $\rho + p \geq 0$, and (iv) Strong energy conditions (SEC) if $\rho + 3p \geq 0$.

Generally, the WEC and DEC are always satisfied by all recognized energy and matters [93–95]. The unusual energy (DE) that creates a strong negative pressure is responsible for the universe’s fast expansion and the violet SEC [93–96].




Figure 11: Plot of Energy Conditions.

Figure 11 shows the behavior of various energy conditions in Rastall gravity, utilizing a combined dataset of OHD, Pantheon, and BAO. In this analysis, the Dominant Energy Condition (DEC) and Null Energy Condition (NEC) are satisfied while the Strong Energy Condition (SEC) violates for the proposed model as depicted in Figure that validates an accelerated expansion of cosmos in the present era [93–97].

Parameters	H_0	ω_0	λ
OHD	70.22 ± 0.26	-0.677 ± 0.014	0.0055 ± 0.0049
OHD+BAO	$70.26^{+0.27}_{-0.24}$	-0.684 ± 0.014	0.0151 ± 0.0091
OHD+Pantheon	70.50 ± 0.25	-0.679 ± 0.013	0.0054 ± 0.0043
OHD+BAO+Pantheon	70.52 ± 0.20	-0.683 ± 0.012	0.0053 ± 0.0043
ANN	$70.008^{+1.144}_{-1.108}$	$-0.500^{+0.026}_{-0.027}$	0.005 ± 0.001
MDN	$69.815^{+3.596}_{-2.654}$	$-0.586^{+0.255}_{-0.153}$	$0.033^{+0.052}_{-0.102}$
MNN	$69.987^{+1.113}_{-1.087}$	0.500 ± 0.023	0.002 ± 0.002

Table 2: The best-fit values of model parameters for the proposed model

6 Conclusion

In this study, we have explored the cosmological dynamics of an isotropic, homogeneous universe in Rastall gravity. To determine the explicit solution for field equations in Rastall gravity, the parameterization of EoS parameter in the form $\omega(z) = \frac{\omega_0}{(z+1)}$ is considered, and the Hubble parameter as a redshift function is derived. The model parameters are estimated by taking observational BAO, Pantheon, and OHD datasets using MCMC analysis. To constrain the cosmological parameters and analyze the current cosmic scenario of the accelerating universe, an innovative deep-learning approach is also employed simultaneously. Using the CoLFI Python package and advanced techniques such as Artificial Neural Networks (ANN), Mixture Density Networks (MDN), and Mixture of Gaussians (MNN), the model parameters are estimated efficiently. This advancement significantly facilitated the understanding of conditional probability densities derived from observational data and posterior distributions. To enhance the efficiency of neural network training and improve the accuracy of parameter predictions, hyperellipsoid parameters were introduced. A comparative study between our neural network methods and traditional Markov Chain Monte Carlo (MCMC) techniques revealed that ANN produced nearly identical results, confirming its reliability and effectiveness.

The proposed model of the universe efficiently explains the late-time acceleration without the need for dark energy and avoids issues associated with the cosmological constant. Therefore, it is essential to consider current observational data when simulating Rastall gravity. We have shown that neural network methods can effectively estimate cosmological parameters, serving as a practical alternative to MCMC techniques. The Rastall gravity framework facilitates the exploration of various gravity theories and their influence on cosmic expansion through the use of ANN, MDN, and MNN methodologies. This research is part of a broader study integrating machine learning with cosmology, demonstrating the capability of machine learning to address intricate cosmic dynamics and expansion challenges. The assessment of density, cosmic pressure, equation of state, and deceleration parameter in the present analysis shows that the derived results for the model are viable. The analysis of statefinders indicates that the resulting model exhibits characteristics of an accelerating cosmic model other than Λ CDM. The analysis of the jerk parameter $j(z)$ and $O_m(z)$ diagnostic for the derived model shows the phantom-like behavior. The violation of the Strong Energy Condition (SEC) for the proposed model validates an accelerated expansion of the cosmos in the present era.

References

- [1] A.G. Riess, *et al.*, Observational evidence from supernovae for an accelerating universe and a cosmological constant, *Astron. J.* **116** (1998) 1009.
- [2] S. Perlmutter, *et al.*, Measurements of Ω and Λ from 42 high-redshift supernovae, *Astrophys. J.* **517** (1999) 565.
- [3] C.L. Bennett et al., Nine-year Wilkinson Microwave Anisotropy Probe (WMAP) observations: final maps and results, *Astrophys. J. Suppl. Ser.* **208** (2013) 20.
- [4] E. Komatsu *et al.*, Five-year wilkinson microwave anisotropy probe (WMAP) observations: cosmological interpretation, *Astrophys. J. Suppl.* **180** (2009) 330.

- [5] L. Anderson et al., The clustering of galaxies in the SDSS-III Baryon Oscillation Spectroscopic Survey: Baryon acoustic oscillations in the Data Releases 10 and 11 Galaxy samples, *Mon. Not. R. Astron. Soc.* **441** (2014) 24–62.
- [6] P. A. R. Ade et al., Planck 2015 results - XIII, cosmological parameters, *Astron. Astrophys.* **594** (2016) A13.
- [7] M. Tegmark et al., Cosmological parameters from SDSS and WMAP, *Phys. Rev. D* **69** (2004) 103501.
- [8] P.J.E. Peebles, and B. Ratra, The cosmological constant and dark energy, *Reviews of modern physics* **75** (2003) 559.
- [9] V. Sahni and A. Starobinsky, The Case for a positive cosmological Λ -term, *Int. J. Mod. Phys. D* **9** (2000) 373.
- [10] R.R. Caldwell, A phantom menace? Cosmological consequences of a dark energy component with super-negative equation of state, *Phys. Lett. B* **545** (2002) 23-29.
- [11] S.M. Carroll, M. Hoffman, and M. Trodden, Can the dark energy equation-of-state parameter ω be less than -1 ?, *Phys. Rev. D* **68** (2003) 023509.
- [12] T. P. Sotiriou, V. Faraoni, $f(R)$ theories of gravity, *Rev. Mod. Phys.* **82** (2010) 451.
- [13] A. De Felice, S. Tsujikawa, $f(R)$ theories, *Liv. Rev. Rel.* **13** (2010) 1-161.
- [14] S.Nojiri and S.D. Odintsov, Unified cosmic history in modified gravity: from $F(R)$ theory to Lorentz non-invariant models, *Phys. Rep.* **505** (2011) 59-144.
- [15] G.R. Bengochea, R. Ferraro, Dark torsion as the cosmic speed-up, *Phys. Rev. D* **79** (2009) 124019.
- [16] E.V. Linder, Einstein’s other gravity and the acceleration of the universe, *Phys. Rev. D* **81.12** (2010) 127301.
- [17] V. Faraoni and S. Capozziello, Beyond Einstein gravity: a survey of gravitational theories for cosmology and astrophysics. Springer Netherlands (2011).
- [18] T. Clifton et al., Modified gravity and cosmology, *Physics reports* **513.1-3** (2012) 1-189.
- [19] L. Parker, Quantized fields and particle creation in expanding universes. II, *Physical Review D* **3.2** (1971): 346.
- [20] G.W. Gibbons, and S. W. Hawking. "Cosmological event horizons, thermodynamics, and particle creation." *Physical Review D* **15.10** (1977): 2738.
- [21] N. D. Birrell and P.C.W. Davies, Quantum fields in curved space, Cambridge Univ. Press (1984).
- [22] L.H. Ford, Gravitational particle creation and inflation, *Phys. Rev. D* **35** (1987) 2955.
- [23] S. Capozziello and M.D. Laurentis, Extended theories of gravity, *Physics Reports* **509.4-5** (2011): 167-321.

- [24] C.M. Will, "The confrontation between general relativity and experiment." *Living reviews in relativity* 17 (2014): 1-117.
- [25] S. Nojiri, S.D. Odintsov, V.K. Oikonomou, "Modified gravity theories on a nutshell: Inflation, bounce and late-time evolution," *Phys. Rep.* **692** (2017) 1.
- [26] A. G. Riess et al., "The farthest known supernova: support for an accelerating universe and a glimpse of the epoch of deceleration." *The Astrophysical Journal* 560.1 (2001): 49.
- [27] J. F. Navarro, C. S. Frenk and S. D. M. White, "A universal density profile from hierarchical clustering." *The Astrophysical Journal* 490.2 (1997): 493
- [28] D. Clowe et al. "A direct empirical proof of the existence of dark matter." *The Astrophysical Journal* 648.2 (2006): L109.
- [29] T. Harko, F.S.N. Lobo, S. Nojiri, S.D. Odintsov, $f(R, T)$ gravity, *Phys.Rev. D* **84** (2011) 024020.
- [30] J.K. Singh et al., EDSFD parameterization in $f(R, T)$ gravity with linear curvature terms, *Physics of the Dark Universe* 45 (2024) 101513.
- [31] L.K. Sharma et al., Constraining anisotropic universe under $f(R, T)$ theory of gravity, *Journal of High Energy Astrophysics* 44 (2024) 457-467.
- [32] V.K. Bhardwaj, Non-minimal matter-geometry coupling in the Bianchi-V spacetime within the formalism of $f(R, T) = f_1(R) + f_2(R)f_3(T)$ cosmology, *Mod. Phys. Lett. A* **33** (2018) 1850234.
- [33] H. Shabani and A.H. Ziaie, Stability of the Einstein static universe in $f(R, T)$ gravity, *The European Physical Journal C* 77 (2017): 1-15.
- [34] V.K. Bhardwaj, P. Garg, and S. Prakash, Cosmological dynamics of accelerating model in $f(Q)$ gravity with latest observational data, *Astrophysics and Space Science* 369.5 (2024) 50.
- [35] J.K. Singh et al., Bouncing cosmology in modified gravity with higher-order curvature terms, *Journal of High Energy Physics* 2023.3 (2023) 1-21.
- [36] P. Rastall, Generalization of the Einstein theory, *Physical Review D* 6.12 (1972) 3357.
- [37] C.E.M. Batista et al., Rastall cosmology and the Λ CDM model, *Physical Review D—Particles, Fields, Gravitation, and Cosmology* 85.8 (2012) 084008.
- [38] S. Nojiri and S.D. Odintsov, Gravity assisted dark energy dominance and cosmic acceleration, *Physics Letters B* 599.3-4 (2004) 137-142.
- [39] T. Koivisto, A note on covariant conservation of energy–momentum in modified gravities, *Classical and Quantum Gravity* 23.12 (2006) 4289.
- [40] O. Bertolami et al., Extra force in $f(R)$ modified theories of gravity, *Physical Review D* 75.10 (2007) 104016.

- [41] T. Harko and F.S.N. Lobo, Generalized curvature-matter couplings in modified gravity, *Galaxies* 2.3 (2014) 410-465.
- [42] A.H. Ziaie, H. Shabani, and S. Ghaffari, Effects of Rastall parameter on perturbation of dark sectors of the Universe, *Modern Physics Letters A* 36.30 (2021) 2150082.
- [43] T.R.P. Carames et al., The brans–dicke–rastall theory, *The European Physical Journal C* 74.11 (2014) 3145.
- [44] S.K. Maurya and F. Tello-Ortiz, Decoupling gravitational sources by MGD approach in Rastall gravity, *Physics of the Dark Universe* 29 (2020) 100577.
- [45] J.C. Fabris, M.H. Daouda, and O.F. Piattella, Note on the evolution of the gravitational potential in Rastall scalar field theories, *Physics Letters B* 711.3-4 (2012) 232-237.
- [46] A. Singh and K.C. Mishra, Aspects of some Rastall cosmologies, *The European Physical Journal Plus* 135.9 (2020) 752.
- [47] J.K. Singh et al., Evolution of the Universe with quintessence model in Rastall gravity, *Physica Scripta* 99.12 (2024) 125001.
- [48] C.E.M. Batista et al., Observational constraints on Rastall’s cosmology, *The European Physical Journal C* 73 (2013) 1-7.
- [49] K. Lin and W.L. Qian, Cosmic evolution of dark energy in a generalized Rastall gravity, *The European Physical Journal C* 80.6 (2020) 1-8.
- [50] J.C. Fabris, R. Kerner, and J. Tossa, Perturbative analysis of generalized Einstein theories, *International Journal of Modern Physics D* 9.02 (2000) 111-125.
- [51] H. Moradpour et al., A generalization to the Rastall theory and cosmic eras, *The European Physical Journal C* 77 (2017) 1-9.
- [52] D. Das, S. Dutta, and S. Chakraborty, Cosmological consequences in the framework of generalized Rastall theory of gravity, *The European Physical Journal C* 78 (2018) 1-8.
- [53] W. Khyllep and J. Dutta, Linear growth index of matter perturbations in Rastall gravity, *Physics Letters B* 797 (2019) 134796.
- [54] G.J. Wang et al., CoLFI: Cosmological Likelihood-free Inference with Neural Density Estimators, *Astrophys. J. Suppl.* 268 (2023) 7.
- [55] The Theano Development Team, R. Al-Rfou, A. G. Alain, A. Amahairi, et al., Theano: A Python framework for fast computation of mathematical expressions, arXiv:1605.02688 (2016).
- [56] G.J. Wang, S.Y. Li and J.Q. Xia, ECoPANN: A Framework for Estimating Cosmological Parameters using Artificial Neural Networks, *Astrophys. J. Suppl.* 249 (2020) 25
- [57] G.J. Wang, C. Cheng, Y.Z. Ma and J.Q. Xia, Likelihood-free Inference with the Mixture Density Network, *Astrophys. J. Suppl.* 262 (2022) 24.

- [58] L.K. Sharma et al., A comprehensive analysis of observational cosmology in $f(Q)$ gravity with deep learning and MCMC method, *Astronomy and Computing* 49 (2024) 100892.
- [59] M. Capone, V.F. Cardone and M.L. Ruggiero, Accelerating Cosmology in Rastall's Theory, *Nuovo Cim. B* 125 (2011) 1133-1142.
- [60] Y. Gong, Y.Z. Zhang, Probing the curvature and dark energy, *Phys. Rev. D* **72** (4) (2005) 043518.
- [61] S. Capozziello et al., Cosmographic bounds on the cosmological deceleration-acceleration transition redshift in $f(R)$ gravity, *Phys. Rev. D* 90.4 (2014) 044016.
- [62] P. Lemos et al., Model independent $H(z)$ reconstruction using the cosmic inverse distance ladder, *Mon. Not. R. Astron. Soc.* 483 (2019) 4803-4810.
- [63] H. Yu, B. Ratra, F.-Y. Wang, Hubble Parameter and Baryon Acoustic Oscillation Measurement Constraints on the Hubble Constant, the Deviation from the Spatially Flat Λ CDM Model, the Deceleration-Acceleration Transition Redshift, and Spatial Curvature, *Astrophys. J.* 856 (2018) 3.
- [64] M. Moresco, Raising the bar: new constraints on the Hubble parameter with cosmic chronometers at $z \sim 2$, *Mon. Not. R. Astron. Soc.* 450 (2015) L16–L20.
- [65] G.S. Sharov, V.O. Vasilie, How predictions of cosmological models depend on Hubble parameter data sets, *Math. Model. Geom.* 6 (2018) 1.
- [66] D.M. Scolnic et al., The Complete Light-curve Sample of Spectroscopically Confirmed SNe Ia from Pan-STARRS1 and Cosmological Constraints from the Combined Pantheon Sample, *Astrophys. J.* 859 (2018) 101.
- [67] D. Brout, G. Taylor, D. Scolnic et al. The Pantheon+ Analysis: SuperCal-Fragilistic Cross Calibration, Retrained SALT2 Light Curve Model, and Calibration Systematic Uncertainty, *Astrophys. J.* 938 (2022) 111.
- [68] C. Blake et al., The WiggleZ Dark Energy Survey: mapping the distance–redshift relation with baryon acoustic oscillations, *Mon. Not. R. Astron. Soc.* 418 (2011) 1707.
- [69] W.J. Percival et al., Baryon Acoustic Oscillations in the Sloan Digital Sky Survey Data Release 7 Galaxy Sample, *Mon. Not. R. Astron. Soc.* 401 (2010) 2148.
- [70] F. Beutler et al., The 6dF Galaxy Survey: baryon acoustic oscillations and the local Hubble constant, *Mon. Not. R. Astron. Soc.* 416 (2011) 3017.
- [71] R. Giotri et al., From cosmic deceleration to acceleration: new constraints from SN Ia and BAO/CMB, *J. Cosmol. Astropart. Phys.* 2012 (2012) 027.
- [72] A.A. Mamon, K. Bamba, S. Das, Constraints on reconstructed dark energy model from SN Ia and BAO/CMB observations, *Eur. Phys. J. C* 77 (2017) 29.
- [73] S. Capozziello et al., Thermodynamic parametrization of dark energy, *Mon. Not. R. Astron. Soc.* 509 (2022) 5399.

- [74] M. Muccino, O. Luongo, D. Jain, Constraints on the transition redshift from the calibrated gamma-ray burst E_p -Eiso correlation, *Mon. Not. R. Astron. Soc.* 523 (2023) 4938.
- [75] M.V.D. Santos et al., Constraining the cosmic deceleration-acceleration transition with type Ia supernova, BAO/CMB and $H(z)$ data, *J. Cosmo. Astropart. Phys.* 2016 (2016) 066.
- [76] S. Masi et al., The BOOMERanG experiment and the curvature of the Universe, *Prog. Part. Nucl. Phys.* 48 (2002) 243.
- [77] V.K. Bhardwaj, S. Prakash, Observational constraints on anisotropic cosmological model in Lyra's manifold, *Chin. J. Phys.* 87 (2024) 665.
- [78] O. Farooq, B. Ratra, Hubble parameter measurement constraints on the cosmological deceleration-acceleration transition redshift, *Astrophys. J. Lett.* 766 (2013) L7.
- [79] V. Sahni, T.D. Saini, A.A. Starobinsky, U. Alam, Statefinder-A new geometrical diagnostic of dark energy, *J. Exp. Theor. Phys. Lett.* 77 (2003) 201.
- [80] U. Alam, V. Sahni, T. Deep Saini, A. A. Starobinsky, Exploring the expanding universe and dark energy using the Statefinder diagnostic, *Mon. Not. Roy Astron Soc.* 344 (2003) 1057.
- [81] X. Zhang, Statefinder diagnostic for holographic dark energy model, *Int. J. Mod. Phys D* 14,(2005) 1597.
- [82] M. R. Setare, J. Zhang, X. Zhang, Statefinder diagnosis in a non-flat universe and the holographic model of dark energy, *JCAP* 3 (2007) 007.
- [83] V.K. Bhardwaj, A. K. Yadav, Observation constraints on scalar field cosmological model in Anisotropic universe, *Int. J. Geom. Meth. Mod. Phys.* 21, (2024) 2450144.
- [84] M. Visser, Jerk, snap and the cosmological equation of state, *Class. Quantum Grav.* 21 (2004) 2603.
- [85] M. Shahalam, S. Sami, A. Agarwal, Om diagnostic applied to scalar field models and slowing down of cosmic acceleration, *Mon. Not. Roy Astron Soc.* 448 (2015) 2948.
- [86] R.D. Blandford et al., Cosmokinetics, *ASP Conf. Ser.* 339 (2004) 27. [arXiv:astro-ph/0408279]
- [87] V.K. Bhardwaj, A.K. Yadav, L.K. Gupta, R. Prasad, S.K. Srivastava, Constraining hybrid potential scalar field cosmological model in Lyra's geometry with recent observational data, *Int. J. Geom. Methods Mod. Phys.* 22 (2025) 2450283.
- [88] V. Sahni, A. Shafieloo, A. A. Starobinsky, Two new diagnostics of dark energy, *Phys. Rev. D* 78 (2008) 103502.
- [89] M. Jamil, D. Momeni, R. Myrzakulov, Observational constraints on non-minimally coupled Galileon model, *Eur. Phys. J. C* 73 (2013) 2347.

- [90] J. Santos, J.S. Alcaniz, Energy conditions and Segre classification of phantom fields, Phys. Lett. B 619 (2005) 11.
- [91] J. Santos et al., Energy conditions in $f(R)$ gravity, Phys. Rev. D 76 (2007) 083513.
- [92] S.M. Carroll, Spacetime and Geometry: An Introduction to General Relativity, Addison Wesley, Boston (2004).
- [93] S. Capozziello, S. Nojiri, S.D. Odintsov, The role of energy conditions in $f(R)$ cosmology, Phys. Lett. B 781 (2018) 99-106.
- [94] V.K Bhardwaj, M.K. Rana, A.K Yadav, Bulk viscous Bianchi-V cosmological model within the formalism of $f(R, T) = f_1(R) + f_2(R)f_3(T)$ gravity, Astrophys. Space Sci. 364 (2019) 1.
- [95] E.J. Copeland, M. Sami, S. Tsujikawa, Dynamics of dark energy, Int. J. Mod. Phys. D 15 (2006) 1753.
- [96] B. Ratra, P.J.E. Peebles, Cosmological consequences of a rolling homogeneous scalar field, Phys. Rev. D **37** (1988) 3406.
- [97] B.S. Jayas and V.K. Bhardwaj, Evaluation of transit cosmological model in $f(R, T^\phi)$ theory of gravity, International Journal of Modern Physics A 39.29 (2024) 2450106.