QCD corrections to Higgs boson pair production and decay to the $b\bar{b}\tau^+\tau^-$ final state

Hai Tao Li^a, Zong-Guo Si^a, Jian Wang^{a,b}, Xiao Zhang^a, Dan Zhao^a

^aSchool of Physics, Shandong University, Jinan, Shandong 250100, China ^bCenter for High Energy Physics, Peking University, Beijing 100871, China

Abstract

We present a comprehensive investigation of next-to-leading order (NLO) quantum chromodynamics (QCD) corrections to Higgs boson pair production and subsequent decay into $b\bar{b}\tau^+\tau^-$. We adopt the narrow width approximation to separate the production and decay contributions, and employ the dipole subtraction method to address infrared divergences inherent in perturbative QCD corrections. After applying a set of typical experimental cuts, we show that NLO QCD corrections to the decay process induce a significant reduction of the fiducial cross section and reshape important kinematic distributions at the 13.6 TeV LHC, such as the invariant mass of the Higgs boson pair and the transverse momentum of the leading *b*-jet. We also investigate the dependence of the kinematic distributions on the Higgs self-coupling and provide the signal acceptance as a function of the Higgs self-coupling modifier with full QCD corrections.

1. Introduction

As the last elementary particle discovered at the LHC in 2012 [1, 2], the Higgs boson plays a crucial role in the Standard Model (SM) and various scenarios beyond the SM. Current measurements of its properties—including mass, width, spin, and CP property—are consistent with SM predictions [3–7]. The spontaneous electroweak symmetry breaking mechanism implies that the couplings of the Higgs boson with other particles are proportional to their masses, a relationship that has been experimentally validated by both the ATLAS and CMS collaborations [8, 9]. Despite these achievements, the self-couplings of the Higgs boson remain largely unconstrained, leaving an important gap in our understanding of the Higgs sector [10, 11].

Higgs boson pair production serves as a direct probe of the Higgs trilinear self-coupling λ_{HHH} . At the LHC, Higgs boson pairs are predominantly produced via gluon-gluon fusion (ggF) and vector boson fusion (VBF). Experimentally, the Higgs boson pair events have been searched for through various decay channels, including $b\bar{b}\gamma\gamma$, $b\bar{b}\tau^+\tau^-$, $b\bar{b}b\bar{b}$, $b\bar{b}ZZ$, and multilepton final states by CMS [12–17], as well as by ATLAS in the $b\bar{b}\gamma\gamma$, $b\bar{b}\tau^+\tau^-$, $b\bar{b}b\bar{b}$, $b\bar{b}WW^*$, $WW^*\gamma\gamma$ and WW^*WW^* decay channels [18–24]. Current experimental constraints on the Higgs self-coupling modifier $\kappa_{\lambda} \equiv \lambda_{\text{HHH}}/\lambda_{\text{HHH}}^{\text{SM}}$ from the single- and double-Higgs production lie in the range of (-0.4, 6.3) and (-1.2, 7.5) by ATLAS [25] and CMS [26], respectively, at 95% confidence level, assuming that the other couplings are the same as the SM values. Future runs of the LHC, particularly the High-Luminosity LHC (HL-LHC), are expected to greatly refine these measurements [27].

On the theoretical front, significant progress has been made in providing precision predictions on the production cross sections. For the ggF process, calculations have been performed up to QCD next-to-next-to-leading order (N³LO) [28–32] and next-to-next-to-next-to-leading logarithmic (N³LL) level [33–35] in the heavy topquark mass limit, as well as to QCD next-to-leading order (NLO) [36–39] and electroweak (EW) NLO [40–48] with full top-mass dependence. Similarly, the VBF production mode has been computed up to QCD N³LO [49– 53] and EW NLO [47, 54]. Given that the Higgs pair production rate is rather small, it is almost mandatory that experimentalists search for the signals with at least one of the Higgs bosons decaying into bottom quarks, which has the largest branching fraction. In such a case, the QCD corrections in $H \rightarrow b\bar{b}$ decay can not be neglected. Indeed, we have demonstrated that QCD NLO corrections to the decay process in $gg \rightarrow HH \rightarrow b\bar{b}\gamma\gamma$ result in substantial effects on both the inclusive and differential cross sections, which significantly exceed the N³LO QCD correction to production [55].

In this work, we extend the previous study to $gg \rightarrow HH \rightarrow b\bar{b}\tau^{+}\tau^{-}$, presenting a full calculation of NLO QCD corrections in both the production and decay processes. This channel is especially appealing because of its relatively large branching fraction, around 7.3%, which enhances the overall signal yield compared to other

decay modes, such as $b\bar{b}\gamma\gamma$. Moreover, despite the experimental challenges associated with τ lepton identification, the $b\bar{b}\tau^+\tau^-$ final state offers distinctive kinematic signatures that improve the discrimination ability between signal and backgrounds. Similar to the $b\bar{b}\gamma\gamma$ channel [55], we find that, after applying typical kinematical cuts, the QCD corrections to the decay lead to a significant reduction of the fiducial cross section as well as remarkable reshaping of key kinematic distributions. The signal acceptance is essential to extracting constraints on the total cross section from the observed events under kinematic cuts. We provide the signal acceptance in this process as a function of κ_{λ} .

The rest of this paper is organized as follows. In section 2 we present the framework for QCD corrections to the production and decay of the Higgs boson pair. The numerical results are discussed in section 3. We conclude in section 4.

2. Theoretical framework

We are going to calculate NLO QCD corrections to Higgs pair production and decay, $gg \rightarrow HH \rightarrow b\bar{b}\tau^{+}\tau^{-}$. Because the Higgs boson width ($\Gamma_{H} = 4.1 \text{ MeV}$) is so small compared to its mass ($m_{H} = 125 \text{ GeV}$) [56], the onshell production gives the dominant contribution to the cross section. Moreover, there is no interference between the corrections to the production and decay at NLO in QCD, since the Higgs boson is colorless. Therefore, the cross section can be written in the narrow width approximation as

$$\int d\sigma_{\rm pro+dec} = \int d\sigma_{\rm pro}(gg \to H_1H_2) \times \frac{\int d\Gamma_{H_1 \to b\bar{b}}}{\Gamma_H} \frac{\int d\Gamma_{H_2 \to \tau^+\tau^-}}{\Gamma_H} = \int d\sigma_{\rm pro}(gg \to HH) \times \frac{\int d\Gamma_{H_1 \to b\bar{b}}}{\Gamma_{H \to b\bar{b}}} \frac{\int d\Gamma_{H_2 \to \tau^+\tau^-}}{\Gamma_{H \to \tau^+\tau^-}} \times B_r \left(b\bar{b}\tau^+\tau^-\right),$$
(1)

where Γ_H is the total decay width and $\Gamma_{H_1 \to X}$ is the partial decay width of $H \to X$. In the first line, we have denoted the Higgs bosons decaying to $b\bar{b}$ and $\tau^+\tau^-$ as H_1 and H_2 , respectively, which are not considered as identital particles. In the second line, we adopt the conventional notation for the production of identical Higgs bosons. Consequently, $B_r(b\bar{b}\tau^+\tau^-)$ is equal to two times the product of branching ratios of $H \to b\bar{b}$ and $H \to \tau^+\tau^-$. Since they are stable against higher-order perturbative corrections, we consider them as constants in our calculation when the other parts of Eq. (1) are expanded in the strong coupling α_s .

From Eq. (1), the LO cross section of $gg \rightarrow HH \rightarrow b\bar{b}\tau^+\tau^-$ is given by

$$\int d\sigma_{\rm pro+dec}^{\rm LO} = \int d\sigma_{\rm pro}^{\rm LO} \times \frac{\int d\Gamma_{H_1 \to b\bar{b}}^{\rm LO}}{\Gamma_{H \to b\bar{b}}^{\rm LO}} \frac{\int d\Gamma_{H_2 \to \tau^+ \tau^-}}{\Gamma_{H \to \tau^+ \tau^-}} \times B_r \left(b\bar{b}\tau^+ \tau^- \right) \,. \tag{2}$$

We have added the superscript "LO" to denote the leading order expansion. The NLO cross section can be expressed as a sum of two contributions, i.e., the NLO corrections to the production process combined with LO decays and the NLO corrections to the decay process interfaced with the LO production,

$$\int d\sigma_{\rm pro+dec}^{\delta \rm NLO} = \int d\sigma_{\rm pro+dec}^{\delta \rm NLO^{\rm pro}} + \int d\sigma_{\rm pro+dec}^{\delta \rm NLO^{\rm dec}} , \qquad (3)$$

where

$$\int d\sigma_{\rm pro+dec}^{\delta\rm NLO\,pro} = \int d\sigma_{\rm pro}^{\delta\rm NLO} \times \frac{\int d\Gamma_{H_1 \to b\bar{b}}^{\rm LO}}{\Gamma_{H \to b\bar{b}}^{\rm LO}} \frac{\int d\Gamma_{H_2 \to \tau^+ \tau^-}}{\Gamma_{H \to \tau^+ \tau^-}} \times B_r \left(b\bar{b}\tau^+ \tau^- \right) \,, \tag{4}$$

and

$$\int d\sigma_{\rm pro+dec}^{\delta \rm NLO^{dec}} = \int d\sigma_{\rm pro}^{\rm LO} \times \frac{\int d\Gamma_{H_2 \to \tau^+ \tau^-}}{\Gamma_{H \to \tau^+ \tau^-}} \frac{\int d\Gamma_{H_1 \to b\bar{b}}^{\rm LO}}{\Gamma_{H \to b\bar{b}}^{\rm LO}} \left(\frac{\int d\Gamma_{H_1 \to b\bar{b}}^{\delta \rm NLO}}{\int d\Gamma_{H_1 \to b\bar{b}}^{\rm LO}} - \frac{\Gamma_{H \to b\bar{b}}^{\delta \rm NLO}}{\Gamma_{H \to b\bar{b}}^{\rm LO}} \right) \times B_r \left(b\bar{b}\tau^+\tau^- \right) \,. \tag{5}$$

The superscript " δ NLO" indicates the expansion at $O(\alpha_s)$. Note that we do not need to expand the decay width of $H \rightarrow \tau^+ \tau^-$ because it is not subject to NLO QCD corrections. Eq. (4) represents the NLO QCD correction to the production process, while Eq. (5) incorporates the NLO QCD correction to the decay process. The second term in the bracket of Eq. (5) comes from the QCD correction to the partial decay width $\Gamma_{H\rightarrow b\bar{b}}$ in the denominator

| | without decays | with decays but no cuts | | with decays and cuts | |
|---|-------------------------|---------------------------------------|----------------------------------|--------------------------|--------------------------------------|
| fb | | LO ^{dec} | $\delta \text{NLO}^{\text{dec}}$ | LO ^{dec} | $\delta \mathrm{NLO}^{\mathrm{dec}}$ |
| LO^{pro}_{∞} | $15.72^{+31\%}_{-22\%}$ | $1.148^{+31\%}_{-22\%}$ | 0 | $0.6876^{+31\%}_{-22\%}$ | $-0.0924^{+42\%}_{-28\%}$ |
| $LO_{m_t}^{pro}$ | $18.57^{+28\%}_{-20\%}$ | $1.357^{+28\%}_{-21\%}$ | 0 | $0.7765^{+27\%}_{-20\%}$ | $-0.1361^{+40\%}_{-27\%}$ |
| $\delta \mathrm{NLO}^{\mathrm{pro}}_{\infty}$ | $13.68^{+6\%}_{-7\%}$ | $0.9997^{+6\%}_{-7\%}$ | - | $0.5869^{+6\%}_{-7\%}$ | - |
| $\delta \text{NLO}_{m_t}^{\text{pro}}$ | $12.27^{+4\%}_{-8\%}$ | $0.8964^{+4\%}_{-8\%}$ | _ | $0.5057^{+4\%}_{-8\%}$ | _ |
| Full NLO result | | | | | |
| NLO∞ | $29.40^{+18\%}_{-15\%}$ | $2.148^{+18\%}_{-15\%}$ | | $1.182^{+15\%}_{-14\%}$ | |
| NLO _{mt} | $30.84^{+14\%}_{-13\%}$ | 2.253 ^{+14%} _{-13%} | | $1.146^{+10\%}_{-11\%}$ | |

Table 1: Inclusive and fiducial cross sections for Higgs boson pair production and decay at the 13.6 TeV LHC. The LO and NLO results are provided with values computed with and without decay effects. The subscript ∞ denotes the cross section calculated in the heavy top-quark mass limit, while the subscript m_t corresponds to the full SM result with complete top-quark mass dependence.

of Eq. (1). It will cancel with the first term if the latter is integrated over the full phase space. In practice, multiple kinematic cuts are applied and thus the decay process exhibits non-vanishing QCD corrections. This is the reason we keep the integration symbols $\int explicitly in above equations.$

Since ggF Higgs boson pair production is a loop-induced process in the SM, the virtual corrections involve two-loop calculations with multiple scales. In our calculation, we employ the numerical grid in POWHEG BOX [57, 58]. The real corrections for *HH* production come from the one-loop diagrams such as $gg \rightarrow HHg$, $qg \rightarrow HHq$, $q\bar{q} \rightarrow HHg$. These contributions are computed using the package OpenLoops [59, 60], interfaced with Collier/OneLOop to calculate the scalar integrals [61, 62].

All the amplitudes required for NLO QCD corrections to $H \rightarrow b\bar{b}$ are computed using FeynArts [63, 64] and FeynCalc [65–67] packages. The analytic results can be found in our previous paper [55]. In the calculation, we neglect the bottom quark mass in the final state but use finite bottom quark Yukawa coupling defined in the $\overline{\text{MS}}$ scheme.

All the virtual and real corrections contain infrared divergences, which are subtracted using the integrated and differential dipole terms [68, 69].

3. Numerical results

In numerical calculations, we adopt the following input parameters:

$$G_F = 1.1663787 \times 10^{-5} \text{ GeV}^{-2}$$
, $m_W = 80.399 \text{ GeV}$,
 $m_H = 125 \text{ GeV}$, $m_t = 173 \text{ GeV}$, $m_b(m_b) = 4.18 \text{ GeV}$,
 $m_\tau = 1.777 \text{ GeV}$. (6)

The Higgs boson decay branching ratios are chosen to be $R(H \rightarrow b\bar{b}) = 0.5824$ and $R(H \rightarrow \tau^+\tau^-) = 6.272 \times 10^{-2}$ [70–72]. We have used the parton distribution function set *PDF4LHC15_nlo_*100_*pdfas* [73] together with the associating strong coupling constant, α_s . The renormalization and factorization scales are set by default to $\mu_r = \mu_f = m_{HH}/2$. The theoretical uncertainties are estimated by varying μ_r and μ_f independently by a factor of two, excluding the cases of $\mu_r/\mu_f = 4$ and 1/4. To stabilize the numerical integration, we impose a technical cut on the Higgs boson pair transverse momentum, $p_T^{\min} = 0.1$ GeV, in the real corrections to the production, while a technical cut of $s_{ij}^{\min} \sim 10^{-5}$ GeV² is applied to the real emissions in the decay.

In this work, we do not consider the decay of τ leptons, since they can be reconstructed and identified using dedicated algorithms by the ATLAS [19, 74] and CMS [13, 75] collaborations. The *b*-jets are constructed using anti- k_t algorithm with a cone size R = 0.4 as implemented in the FastJet package [76]. To investigate QCD corrections to the cross sections with kinematic cuts, we select events that satisfy the following criteria:

$$p_T^{\tau} \ge 20 \text{ GeV} , \quad p_T^{\tau} \ge 20 \text{ GeV} , \quad |\eta^j| \le 2.4 ,$$

$$|\eta^{\tau}| \le 2.3 , \quad 90 \text{ GeV} \le m_{jj} \le 190 \text{ GeV} ,$$

$$R_{ll} > 0.3 , \quad R_{jl} > 0.5 , \quad R_{ji} > 0.4 , \qquad (7)$$

where R_{ll} represents the distance between the two τ leptons, R_{lj} denotes the separation between a *b*-jet and a τ lepton, and R_{jj} is the distance between two *b*-jets.

Table 1 shows the inclusive and fiducial cross sections for Higgs boson pair production and decay at the 13.6 TeV LHC. We have performed the calculation in both the full SM and the heavy top-quark mass limit. The results in the SM are higher than those in the heavy top-quark mass limit by 13% and 18% at LO with and without cuts on the decay products, respectively. However, the NLO QCD corrections in the production increase the cross section by 66% and 87% in the SM and large m_t limit, respectively. And the NLO QCD corrections in the decay decrease the cross section and are more pronounced in the SM, reaching -18%. Consequently, the NLO results after cuts in the SM is smaller than that in the heavy top-quark mass limit, although the difference is small, only 3%. The scale uncertainties are reduced by a factor of two in both the SM and large m_t limit.

From the table, we observe that the cross sections with decays but without cuts are just the production rates multiplied by the branching ratio, which serves as a check of our numerical program. Note that the cuts in Eq. (7) look rather loose. However, they cut down over 40% of the events at both LO and NLO. One of the reasons is that the subleading *b*-jet and τ lepton do not have large transverse momentum in most of the events.



Figure 1: Differential cross sections for Higgs boson pair production with QCD corrections. The black curves represent the LO predictions, while the blue dashed curves show the results with NLO corrections to production only. The red dotted curves display the outcome when full QCD corrections (production and decay) are included. The K-factors are defined by the ratio of NLO results over LO ones.

In Fig. 1, we display the kinematic distributions of the final states with NLO QCD corrections after cuts. In the Higgs boson pair invariant mass $m_{jj\tau\tau}$ distribution, which is reconstructed from the final-state *b*-jets and τ letpons, the QCD corrections in production are more significant in the smaller $m_{jj\tau\tau}$ region, reaching 80% at most. The QCD corrections in decay increase and decrease the distribution below and above $m_{jj\tau\tau} = 340$ GeV, respectively, and therefore shift the peak towards smaller value of $m_{jj\tau\tau}$. In addition, they reduce the peak height by 11%.

The *b*-jets are ordered according to their transverse momenta. The shape of the leading *b*-jet transverse momentum, denoted by p_T^{j1} , distribution is notably changed after including the NLO QCD corrections in decay, since both positive and negative corrections exist in different regions of p_T^{j1} . The subleading *b*-jet transverse momentum distribution is reduced by the QCD corrections in decay. The effect is most obvious around $p_T^{j2} = 80$ GeV. The QCD corrections to the rapidity distribution of the *b*-jets and τ leptons are almost flat over the whole regions. We show the case of the leading *b*-jet in Fig. 1.

The transverse momentum distributions of the τ leptons, shown in Fig. 1, also suffer from the suppression of QCD corrections in decay, especially in the peak regions.



Figure 2: Differential cross sections for Higgs boson pair production and decay at different values of κ_{λ} . The results of $\kappa_{\lambda} = 6$ have been rescaled by a factor of 1/10. The lower panels show the K-factors defined as NLO QCD results over the LO ones.

The kinematics of the signal events at different κ_{λ} plays a crucial role in determining the experimental constraints on the Higgs self-coupling, since the signal acceptance that is defined as the cross section passing the cuts over the total cross section is generally not a constant for different κ_{λ} [25, 77]. We show in Fig. 2 the kinematics at different values of κ_{λ} . The lower panels in Fig. 2 show that the QCD corrections are important to provide precise theoretical predictions for various distributions and do not appear as overall enhancement factors. It can be seen that the invariant mass $m_{jj\tau\tau}$ distribution varies dramatically as κ_{λ} changes from 1 to 6. Near the threshold of Higgs boson pair production (around 250 GeV), the SM cross section is almost vanishing due to the cancellation between the amplitudes with and without the Higgs self-coupling dependence. Choosing larger values of κ_{λ} breaks the cancellation and the cross section is remarkably enhanced in this region. Then the cancellation would happen in other regions of $m_{jj\tau\tau}$. This explains the dip around $m_{jj\tau\tau} = 360$ GeV in the curve for $\kappa_{\lambda} = 3$. Whenever the cancellation happens, QCD corrections increase the cross section tremendously, which can be observed by the large K-factor (larger than 3.6) in the lower panel.

The transverse momentum distributions of the *b*-jets and τ leptons at different values of κ_{λ} have very different shapes. The leading *b*-jets and leptons tend to be softer as the increasing of κ_{λ} , while the peak positions of the subleading *b*-jets and leptons move towards harder regions. In all these distributions, prominent QCD corrections can be seen, in particular, for large κ_{λ} .

Since the (differential) cross section is a quadratic function of κ_{λ} , the distributions for other values of κ_{λ} can be derived from the three curves in the plots, which correspond to $\kappa_{\lambda} = 1, 3$, and 6.

Based on the above calculation, we obtain the cross sections after the cuts (7) as functions of κ_{λ} ,

$$\sigma_{\rm LO}^{\rm cut} = 0.2028\kappa_{\lambda}^{2} - 1.010\kappa_{\lambda} + 1.583 \text{ fb},$$

$$\sigma_{\rm NLO}^{\rm cut} = 0.3703\kappa_{\lambda}^{2} - 1.740\kappa_{\lambda} + 2.516 \text{ fb},$$

$$\sigma_{\rm NLO/\delta NLO^{\rm dec}}^{\rm cut} = 0.4107\kappa_{\lambda}^{2} - 1.934\kappa_{\lambda} + 2.806 \text{ fb},$$
(8)

where the third equation represents the result without QCD corrections in decay. Comparing them to the total



Figure 3: The inverse of signal acceptance as a function of κ_{λ} .

cross sections without cuts

$$\sigma_{\rm LO} = 0.3797 \kappa_{\lambda}^2 - 1.851 \kappa_{\lambda} + 2.829 \text{ fb},$$

$$\sigma_{\rm NLO} = 0.7728 \kappa_{\lambda}^2 - 3.560 \kappa_{\lambda} + 5.037 \text{ fb},$$
 (9)

we see that a large amount of the events can not pass the cuts for a general κ_{λ} . The inverse of the signal acceptance, i.e., σ/σ^{cut} , multiplied with the cross section measured experimentally, provides information on the constraints of *HH* production total cross sections. We depict the inverse of the signal acceptance under the cuts Eq. (7) in Fig. 3. A dip occurs near $\kappa_{\lambda} = 2$, which is a feature also in the plot on the expected exclusion limit in the ATLAS analysis [25]. From Fig. 3, the allowed region of κ_{λ} would become wider after considering the QCD NLO corrections, especially those in the decay process.

4. Conclusion

Higgs boson pair production via gluon-gluon fusion provides a unique probe of the Higgs self-coupling, a critical parameter to test the Standard Model. In this work, we have presented a precise theoretical prediction for such a process and the subsequent decay to $b\bar{b}\tau^+\tau^-$, which possesses a large branching fraction and distinctive kinematic features. Our study reveals that, once typical experimental cuts are applied, the NLO QCD corrections to the decay process cause an approximate 18% reduction in the fiducial cross section compared to the LO prediction. Moreover, the kinematic distributions, particularly those of the invariant mass $m_{jj\tau\tau}$ and the leading *b*-jet transverse momentum, are remarkably reshaped by these corrections. We have further demonstrated that variations in the Higgs self-coupling modifier κ_{λ} lead to significant shifts in the kinematic distributions, underscoring the sensitivity of the $b\bar{b}\tau^+\tau^-$ channel to new physics effects, and that the QCD corrections in theoretical predictions to ensure precise extraction of the Higgs self-coupling from current and future experimental data.

Acknowledgements

This work was partly supported by the National Science Foundation of China under grant No. 12275156, No. 12321005, No. 12375076 and the Taishan Scholar Foundation of Shandong province (tsqn201909011).

References

- [1] ATLAS Collaboration, G. Aad et al., Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC, Phys. Lett. B 716 (2012) 1–29, [arXiv:1207.7214].
- [2] CMS Collaboration, S. Chatrchyan et al., Observation of a New Boson at a Mass of 125 GeV with the CMS Experiment at the LHC, Phys. Lett. B 716 (2012) 30–61, [arXiv:1207.7235].

- [3] CMS Collaboration, A. M. Sirunyan et al., A measurement of the Higgs boson mass in the diphoton decay channel, Phys. Lett. B 805 (2020) 135425, [arXiv:2002.06398].
- [4] ATLAS Collaboration, Measurement of the Higgs boson mass in the H → ZZ* → 4ℓ decay channel using 139 fb⁻¹ of √s = 13 TeV pp collisions recorded by the ATLAS detector at the LHC, Phys. Lett. B 843 (2023) 137880.
- [5] CMS Collaboration, A. Tumasyan et al., Measurement of the Higgs boson width and evidence of its off-shell contributions to ZZ production, Nature Phys. 18 (2022), no. 11 1329–1334, [arXiv:2202.06923].
- [6] ATLAS Collaboration, G. Aad et al., Study of the spin and parity of the Higgs boson in diboson decays with the ATLAS detector, Eur. Phys. J. C 75 (2015), no. 10 476, [arXiv:1506.05669]. [Erratum: Eur.Phys.J.C 76, 152 (2016)].
- [7] CMS Collaboration, V. Khachatryan et al., Constraints on the spin-parity and anomalous HVV couplings of the Higgs boson in proton collisions at 7 and 8 TeV, Phys. Rev. D 92 (2015), no. 1 012004, [arXiv:1411.3441].
- [8] ATLAS Collaboration, A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery, Nature 607 (2022), no. 7917 52–59, [arXiv:2207.00092]. [Erratum: Nature 612, E24 (2022)].
- [9] CMS Collaboration, A. Tumasyan et al., A portrait of the Higgs boson by the CMS experiment ten years after the discovery, Nature 607 (2022), no. 7917 60–68, [arXiv:2207.00043].
- [10] A. Adhikary, S. Banerjee, R. K. Barman, B. Bhattacherjee, and S. Niyogi, *Higgs Self-Coupling at the HL-LHC and HE-LHC, Springer Proc. Phys.* **277** (2022) 27–31.
- [11] J. de Blas et al., *Higgs Boson Studies at Future Particle Colliders*, *JHEP* **01** (2020) 139, [arXiv:1905.03764].
- [12] CMS Collaboration, A. M. Sirunyan et al., Search for nonresonant Higgs boson pair production in final states with two bottom quarks and two photons in proton-proton collisions at $\sqrt{s} = 13$ TeV, JHEP 03 (2021) 257, [arXiv:2011.12373].
- [13] CMS Collaboration, A. Tumasyan et al., Search for nonresonant Higgs boson pair production in final state with two bottom quarks and two tau leptons in proton-proton collisions at s=13 TeV, Phys. Lett. B 842 (2023) 137531, [arXiv:2206.09401].
- [14] CMS Collaboration, A. Tumasyan et al., Search for Higgs Boson Pair Production in the Four b Quark Final State in Proton-Proton Collisions at s=13 TeV, Phys. Rev. Lett. 129 (2022), no. 8 081802, [arXiv:2202.09617].
- [15] CMS Collaboration, A. Tumasyan et al., Search for Nonresonant Pair Production of Highly Energetic Higgs Bosons Decaying to Bottom Quarks, Phys. Rev. Lett. 131 (2023), no. 4 041803, [arXiv:2205.06667].
- [16] CMS Collaboration, A. Tumasyan et al., Search for Higgs boson pairs decaying to WW*WW*, WW* $\tau\tau$, and $\tau\tau\tau\tau$ in proton-proton collisions at $\sqrt{s} = 13$ TeV, JHEP **07** (2023) 095, [arXiv:2206.10268].
- [17] **CMS** Collaboration, A. Tumasyan et al., Search for nonresonant Higgs boson pair production in the four leptons plus twob jets final state in proton-proton collisions at $\sqrt{s} = 13$ TeV, JHEP **06** (2023) 130, [arXiv:2206.10657].
- [18] **ATLAS** Collaboration, G. Aad et al., Search for Higgs boson pair production in the two bottom quarks plus two photons final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, Phys. Rev. D 106 (2022), no. 5 052001, [arXiv:2112.11876].
- [19] **ATLAS** Collaboration, G. Aad et al., Search for resonant and non-resonant Higgs boson pair production in the $b\bar{b}\tau^+\tau^-$ decay channel using 13 TeV pp collision data from the ATLAS detector, JHEP **07** (2023) 040, [arXiv:2209.10910].

- [20] **ATLAS** Collaboration, G. Aad et al., Search for nonresonant pair production of Higgs bosons in the $b\bar{b}b\bar{b}$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, Phys. Rev. D 108 (2023), no. 5 052003, [arXiv:2301.03212].
- [21] **ATLAS** Collaboration, M. Aaboud et al., Search for Higgs boson pair production in the $b\bar{b}WW^*$ decay mode at $\sqrt{s} = 13$ TeV with the ATLAS detector, JHEP **04** (2019) 092, [arXiv:1811.04671].
- [22] **ATLAS** Collaboration, M. Aaboud et al., Search for Higgs boson pair production in the $\gamma\gamma WW^*$ channel using pp collision data recorded at $\sqrt{s} = 13$ TeV with the ATLAS detector, Eur. Phys. J. C **78** (2018), no. 12 1007, [arXiv:1807.08567].
- [23] **ATLAS** Collaboration, M. Aaboud et al., Search for Higgs boson pair production in the $WW^{(*)}WW^{(*)}$ decay channel using ATLAS data recorded at $\sqrt{s} = 13$ TeV, JHEP **05** (2019) 124, [arXiv:1811.11028].
- [24] **ATLAS** Collaboration, G. Aad et al., Search for the nonresonant production of Higgs boson pairs via gluon fusion and vector-boson fusion in the $b\bar{b}\tau^+\tau^-$ final state in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, Phys. Rev. D **110** (2024), no. 3 032012, [arXiv:2404.12660].
- [25] ATLAS Collaboration, G. Aad et al., Constraints on the Higgs boson self-coupling from single- and double-Higgs production with the ATLAS detector using pp collisions at s=13 TeV, Phys. Lett. B 843 (2023) 137745, [arXiv:2211.01216].
- [26] CMS Collaboration, Constraints on the Higgs boson self-coupling with combination of single and double Higgs boson production, .
- [27] **ATLAS** Collaboration, *Projected sensitivity of measurements of Higgs boson pair production with the ATLAS experiment at the HL-LHC*, .
- [28] S. Dawson, S. Dittmaier, and M. Spira, Neutral Higgs boson pair production at hadron colliders: QCD corrections, Phys. Rev. D 58 (1998) 115012, [hep-ph/9805244].
- [29] D. de Florian and J. Mazzitelli, *Higgs Boson Pair Production at Next-to-Next-to-Leading Order in QCD*, *Phys. Rev. Lett.* **111** (2013) 201801, [arXiv:1309.6594].
- [30] D. de Florian, M. Grazzini, C. Hanga, S. Kallweit, J. M. Lindert, P. Maierhöfer, J. Mazzitelli, and D. Rathlev, *Differential Higgs Boson Pair Production at Next-to-Next-to-Leading Order in QCD*, *JHEP* 09 (2016) 151, [arXiv:1606.09519].
- [31] L.-B. Chen, H. T. Li, H.-S. Shao, and J. Wang, *Higgs boson pair production via gluon fusion at N³LO in QCD*, *Phys. Lett. B* 803 (2020) 135292, [arXiv:1909.06808].
- [32] L.-B. Chen, H. T. Li, H.-S. Shao, and J. Wang, The gluon-fusion production of Higgs boson pair: N³LO QCD corrections and top-quark mass effects, JHEP 03 (2020) 072, [arXiv:1912.13001].
- [33] D. Y. Shao, C. S. Li, H. T. Li, and J. Wang, Threshold resummation effects in Higgs boson pair production at the LHC, JHEP 07 (2013) 169, [arXiv:1301.1245].
- [34] D. de Florian and J. Mazzitelli, *Higgs pair production at next-to-next-to-leading logarithmic accuracy at the LHC, JHEP* **09** (2015) 053, [arXiv:1505.07122].
- [35] A. H. Ajjath and H.-S. Shao, N³LO+N³LL QCD improved Higgs pair cross sections, JHEP **02** (2023) 067, [arXiv:2209.03914].
- [36] S. Borowka, N. Greiner, G. Heinrich, S. P. Jones, M. Kerner, J. Schlenk, U. Schubert, and T. Zirke, *Higgs Boson Pair Production in Gluon Fusion at Next-to-Leading Order with Full Top-Quark Mass Dependence, Phys. Rev. Lett.* **117** (2016), no. 1 012001, [arXiv:1604.06447]. [Erratum: Phys.Rev.Lett. 117, 079901 (2016)].
- [37] S. Borowka, N. Greiner, G. Heinrich, S. P. Jones, M. Kerner, J. Schlenk, and T. Zirke, Full top quark mass dependence in Higgs boson pair production at NLO, JHEP 10 (2016) 107, [arXiv:1608.04798].
- [38] J. Baglio, F. Campanario, S. Glaus, M. Mühlleitner, M. Spira, and J. Streicher, *Gluon fusion into Higgs pairs at NLO QCD and the top mass scheme, Eur. Phys. J. C* **79** (2019), no. 6 459, [arXiv:1811.05692].

- [39] J. Baglio, F. Campanario, S. Glaus, M. Mühlleitner, J. Ronca, M. Spira, and J. Streicher, *Higgs-Pair Production via Gluon Fusion at Hadron Colliders: NLO QCD Corrections*, JHEP 04 (2020) 181, [arXiv:2003.03227].
- [40] S. Borowka, C. Duhr, F. Maltoni, D. Pagani, A. Shivaji, and X. Zhao, Probing the scalar potential via double Higgs boson production at hadron colliders, JHEP 04 (2019) 016, [arXiv:1811.12366].
- [41] M. Mühlleitner, J. Schlenk, and M. Spira, *Top-Yukawa-induced corrections to Higgs pair production*, *JHEP* **10** (2022) 185, [arXiv:2207.02524].
- [42] J. Davies, G. Mishima, K. Schönwald, M. Steinhauser, and H. Zhang, *Higgs boson contribution to the leading two-loop Yukawa corrections to gg* → *HH*, *JHEP* **08** (2022) 259, [arXiv:2207.02587].
- [43] J. Davies, K. Schönwald, M. Steinhauser, and H. Zhang, *Next-to-leading order electroweak corrections to* $gg \rightarrow HH$ and $gg \rightarrow gH$ in the large- m_t limit, JHEP **10** (2023) 033, [arXiv:2308.01355].
- [44] H.-Y. Bi, L.-H. Huang, R.-J. Huang, Y.-Q. Ma, and H.-M. Yu, Electroweak Corrections to Double Higgs Production at the LHC, Phys. Rev. Lett. 132 (2024), no. 23 231802, [arXiv:2311.16963].
- [45] G. Heinrich, S. Jones, M. Kerner, T. Stone, and A. Vestner, *Electroweak corrections to Higgs boson pair production: the top-Yukawa and self-coupling contributions*, *JHEP* 11 (2024) 040, [arXiv:2407.04653].
- [46] H. Zhang, K. Schönwald, M. Steinhauser, and J. Davies, *Electroweak corrections to gg -> HH: Factorizable contributions*, PoS LL2024 (2024) 014, [arXiv:2407.05787].
- [47] H. T. Li, Z.-G. Si, J. Wang, X. Zhang, and D. Zhao, Improved constraints on Higgs boson self-couplings with quartic and cubic power dependencies of the cross section*, Chin. Phys. C 49 (2025), no. 2 023107, [arXiv:2407.14716].
- [48] M. Bonetti, P. Rendler, and W. J. Torres Bobadilla, Two-loop light-quark Electroweak corrections to Higgs boson pair production in gluon fusion, arXiv: 2503.16620.
- [49] R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, P. Torrielli, E. Vryonidou, and M. Zaro, *Higgs pair production at the LHC with NLO and parton-shower effects, Phys. Lett. B* 732 (2014) 142–149, [arXiv:1401.7340].
- [50] L.-S. Ling, R.-Y. Zhang, W.-G. Ma, L. Guo, W.-H. Li, and X.-Z. Li, NNLO QCD corrections to Higgs pair production via vector boson fusion at hadron colliders, Phys. Rev. D 89 (2014), no. 7 073001, [arXiv:1401.7754].
- [51] F. A. Dreyer and A. Karlberg, Vector-Boson Fusion Higgs Pair Production at N³LO, Phys. Rev. D 98 (2018), no. 11 114016, [arXiv:1811.07906].
- [52] F. A. Dreyer and A. Karlberg, Fully differential Vector-Boson Fusion Higgs Pair Production at Next-to-Next-to-Leading Order, Phys. Rev. D 99 (2019), no. 7 074028, [arXiv:1811.07918].
- [53] B. Jäger, A. Karlberg, and S. Reinhardt, *Precision tools for the simulation of double-Higgs production via vector-boson fusion*, arXiv:2502.09112.
- [54] F. A. Dreyer, A. Karlberg, J.-N. Lang, and M. Pellen, *Precise predictions for double-Higgs production via vector-boson fusion, Eur. Phys. J. C* 80 (2020), no. 11 1037, [arXiv:2005.13341].
- [55] H. T. Li, Z.-G. Si, J. Wang, X. Zhang, and D. Zhao, *Higgs boson pair production and decay at NLO in QCD: the b̄bγγ final state*, *JHEP* 04 (2024) 002, [arXiv:2402.00401].
- [56] Particle Data Group Collaboration, S. Navas et al., *Review of particle physics, Phys. Rev. D* 110 (2024), no. 3 030001.
- [57] S. Alioli, P. Nason, C. Oleari, and E. Re, A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX, JHEP 06 (2010) 043, [arXiv:1002.2581].
- [58] G. Heinrich, S. P. Jones, M. Kerner, and L. Scyboz, A non-linear EFT description of gg → HH at NLO interfaced to POWHEG, JHEP 10 (2020) 021, [arXiv:2006.16877].

- [59] OpenLoops 2 Collaboration, F. Buccioni, J.-N. Lang, J. M. Lindert, P. Maierhöfer, S. Pozzorini,
 H. Zhang, and M. F. Zoller, *OpenLoops 2, Eur. Phys. J. C* 79 (2019), no. 10 866, [arXiv:1907.13071].
- [60] F. Buccioni, S. Pozzorini, and M. Zoller, On-the-fly reduction of open loops, Eur. Phys. J. C 78 (2018), no. 1 70, [arXiv:1710.11452].
- [61] A. Denner, S. Dittmaier, and L. Hofer, Collier: a fortran-based Complex One-Loop Llbrary in Extended Regularizations, Comput. Phys. Commun. 212 (2017) 220–238, [arXiv:1604.06792].
- [62] A. van Hameren, OneLOop: For the evaluation of one-loop scalar functions, Comput. Phys. Commun. 182 (2011) 2427–2438, [arXiv:1007.4716].
- [63] J. Küblbeck, M. Böhm, and A. Denner, *Feyn arts computer-algebraic generation of feynman graphs and amplitudes, Computer Physics Communications* **60** (1990), no. 2 165–180.
- [64] T. Hahn, Generating Feynman diagrams and amplitudes with FeynArts 3, Comput. Phys. Commun. 140 (2001) 418–431, [hep-ph/0012260].
- [65] R. Mertig, M. Böhm, and A. Denner, Feyn calc computer-algebraic calculation of feynman amplitudes, Computer Physics Communications 64 (1991), no. 3 345–359.
- [66] V. Shtabovenko, R. Mertig, and F. Orellana, New Developments in FeynCalc 9.0, Comput. Phys. Commun. 207 (2016) 432–444, [arXiv:1601.01167].
- [67] V. Shtabovenko, R. Mertig, and F. Orellana, *FeynCalc 9.3: New features and improvements, Comput. Phys. Commun.* **256** (2020) 107478, [arXiv:2001.04407].
- [68] S. Catani and M. H. Seymour, A General algorithm for calculating jet cross-sections in NLO QCD, Nucl. Phys. B 485 (1997) 291–419, [hep-ph/9605323]. [Erratum: Nucl.Phys.B 510, 503–504 (1998)].
- [69] T. Gleisberg and F. Krauss, Automating dipole subtraction for QCD NLO calculations, Eur. Phys. J. C 53 (2008) 501–523, [arXiv:0709.2881].
- [70] LHC Higgs Cross Section Working Group Collaboration, D. de Florian et al., *Handbook of LHC Higgs Cross Sections: 4. Deciphering the Nature of the Higgs Sector*, arXiv:1610.07922.
- [71] HDECAY Collaboration, A. Djouadi, J. Kalinowski, M. Muehlleitner, and M. Spira, HDECAY: Twenty₊₊ years after, Comput. Phys. Commun. 238 (2019) 214–231, [arXiv:1801.09506].
- [72] A. Djouadi, J. Kalinowski, and M. Spira, HDECAY: A Program for Higgs boson decays in the standard model and its supersymmetric extension, Comput. Phys. Commun. 108 (1998) 56–74, [hep-ph/9704448].
- [73] A. Buckley, J. Ferrando, S. Lloyd, K. Nordström, B. Page, M. Rüfenacht, M. Schönherr, and G. Watt, LHAPDF6: parton density access in the LHC precision era, Eur. Phys. J. C 75 (2015) 132, [arXiv:1412.7420].
- [74] **ATLAS** Collaboration, G. Aad et al., *Identification and energy calibration of hadronically decaying tau leptons with the ATLAS experiment in pp collisions at* $\sqrt{s}=8$ *TeV, Eur. Phys. J. C* **75** (2015), no. 7 303, [arXiv:1412.7086].
- [75] **CMS** Collaboration, A. M. Sirunyan et al., *Performance of reconstruction and identification of* τ *leptons decaying to hadrons and* ν_{τ} *in pp collisions at* $\sqrt{s} = 13$ TeV, JINST **13** (2018), no. 10 P10005, [arXiv:1809.02816].
- [76] M. Cacciari, G. P. Salam, and G. Soyez, FastJet User Manual, Eur. Phys. J. C 72 (2012) 1896, [arXiv:1111.6097].
- [77] ATLAS Collaboration, Validation of signal Monte Carlo event generation in searches for Higgs boson pairs with the ATLAS detector, .