# Volume in the Extensive Thermodynamics of Black Holes<sup>\*</sup>

Réka Somogyfoki<sup>1,3,4</sup> and Péter Ván<sup>2,3,4</sup>

<sup>1</sup>ELTE Doctoral School of Physics, Eötvös Loránd University, Budapest, Hungary

<sup>2</sup>Department of Energy Engineering, Faculty of Mechanical Engineering, Budapest University of Technology and Economics, Budapest, Hungary <sup>3</sup>Department of Theoretical Physics, Institute of Particle and Nuclear

Physics, HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

<sup>4</sup>Montavid Thermodynamic Research Group, Budapest, Hungary

#### Received March 31, 2025

Since black holes lack a straightforward notion of geometrical volume due to their event horizon structure and coordinate dependence, various approaches have been proposed to introduce a meaningful geometric and thermodynamic volume. There is no volume in classical black hole thermodynamics, the nonextensivity of black holes is a consequence. Extensivity of such systems can be restored by introducing a new thermodynamic state variable, typically the volume, the mass or the number of particles. Smarr relations show that black hole thermodynamics is nonextensive. However, restored extensivity leads to a volume which corresponds to geometric concepts and is also meaningful from a physical point of view. The article examines various volume definitions in the context of Anti-de Sitter (AdS) black holes and their implications for phase transitions, focusing on the Hawking—Page transition in Kerr—AdS spacetime, including the Christodoulou—Rovelli volume.

### 1. Introduction

In 1973 Bekenstein formulated a supplementary version of the second law of thermodynamics, according to which the sum of the entropy of the black hole and the traditional entropy outside the black hole can never decrease [1]. In 1974 Stephen Hawking discovered that Hawking radiation can lead to the required equilibration, a mechanism for the zeroth law of thermodynamics [2, 3]. Therefore, black holes have a temperature that must

<sup>\*</sup> Presented in Prague, at VISEGRAD 2024 october

be inversely proportional to their mass. These discoveries made it possible to discuss the thermodynamic properties of black holes.

While the physics of black holes is based on a set of strict criteria from differential geometry, thermodynamics is a discipline that describes macroscopic systems and our usual viewpoint requires knowledge of a microscopic structure [4]. However, the no-hair theorem indicates the lack of internal structure analogous to thermodynamic universality, leading to a significant reassessment of the role of thermodynamics.

The paper first analyses the physical meaning and the mathematical realization of first-order homogeneity, the extensivity of entropy functions, then explores how these principles apply to black hole thermodynamics, particularly in defining a consistent notion of volume. The leading example is the technique introduced in thermodynamics of small systems by Hill [5].

## 2. Hawking-Page phase boundaries

In black hole thermodynamics, certain properties follow first-order homogeneous extensivity. The so-called Smarr relations emerge when the invariant mass is treated as a thermodynamic potential [6]. The identification of the mass with different thermodynamic potentials leads to distinct volume definitions (if such a concept is included), as discussed in the following section. A crucial aspect is thermodynamic stability, is a necessary condition for dynamic stability in non-equilibrium thermodynamics, particularly for time-dependent processes [7]. In case of black holes, dynamic stability against symmetric perturbations around an axis is ensured if and only if the black hole satisfies the analogue of the thermodynamic stability criterion for inhomogeneous systems [4].

In AdS spacetime, the energy-momentum tensor is related to the cosmological constant, which is proportional to the pressure. The expansion of the Universe manifests as a locally negative pressure, making the cosmological constant analogous to gas pressure.

Introducing pressure and volume reveals state-space regions in charged AdS black holes where the second derivative of the entropy is positive semidefinite, suggesting phase transitions. The Hawking—Page temperature marks a transition between stable black holes and thermal radiation, calculated as [3]:

$$T_{HP} = \sqrt{\frac{8p}{3\pi}},\tag{1}$$

for  $F(T_{HP}) = 0$  free energy, where p is pressure. Above  $T_{HP}$ , the system favors black hole formation over pure thermal radiation and black holes remain stable. Below  $T_{HP}$  their free energy turns into negative, causing evaporation.

Alternatively, AdS black hole thermodynamics can be formulated by treating the cosmological constant as pressure [8]. In this case, the black hole mass should be identified as enthalpy rather than the free energy. Also, the BH volume naturally emerges as its derivative with respect to pressure. Entropy aligns with the event horizon surface, ensuring that the second law holds. As positive-energy material crosses the event horizon, its mass and surface area increase, leading to entropy growth [9].

### 3. Volume

A black hole without volume is thermodynamically unstable. Introducing volume as an extensive quantity results in the Christodoulou — Rovelli volume concept [10]. In the U(V) parameter space, the entropy is concave, ensuring thermodynamic stability.

A major issue in black hole thermodynamics is the negative heat capacity problem: as a black hole loses mass, its temperature increases, leading to runaway evaporation. Biró et al. [11] introduced a volume concept that resolves this problematic behavior. Their definition, proportional to the fifth power of the radius (and mass), transforms entropy from a convex curve into a concave surface. Identifying black hole mass with enthalpy rather than internal energy incorporates volume and pressure naturally, leading to a completely thermodynamic formulation [12]:

$$H(S,p) = \frac{1}{2}\sqrt{\frac{S}{\pi}} + p\left(\frac{S}{\pi}\right)^{5/2}.$$
 (2)

From this, volume is derived as:

$$V = \frac{\partial H}{\partial p}\Big|_{S} = \left(\frac{S}{\pi}\right)^{5/2} = r_{S}^{5},\tag{3}$$

which confirms that volume is consistent with the Christodoulou—Rovelli volume, namely, that it is proportional to  $r_S^5$ .

### 4. Conclusion

The concept of volume in black hole thermodynamics challenges our conventional understanding of space, matter, and thermodynamic variables. Unlike ordinary systems, the theory black holes lacks a straightforward volume definition, necessitating alternative approaches. The Christodoulou — Rovelli volume is a compelling candidate, preserving thermodynamic consistency and providing a natural resolution to open questions such as negative heat capacity.



Fig. 1. The dimensionless temperature (T) expressed in terms of radius (R) in case of p = 0.15. The local minimum remarked with blue and the Hawking–Page temperature with green. Left: For Hawking and Page; Right: with the Christodoulou–Rovelli volume.

In this work we have shown that the volume concept can be consistently extended to AdS spacetimes and the value of the Hawking–Page temperature changes, in the new thermodynamic framework it becomes  $T_{HPnew} = 0.2067$ , instead of the classical  $T_{HP} = 0.3568$ . The different value of the Hawking–Page temperature and also the different stability properties may have observable consequences, suggesting that black holes in AdS spacetimes remain stable in different temperature regimes compared to the classical approach. Among the several possible cosmological implications, the change in the primordial black hole distribution is outstanding, since it may be detected by the Einstein Telescope [13].

There are several other theoretical consequences, offering a more refined framework for understanding black hole phase transitions like the Hawking—Page one. As black hole physics continues to evolve, refining these thermodynamic descriptions will be crucial in bridging the gap between classical gravity and theories of spacetime.

### 5. Acknowledgement

The work was supported by the grant NKFIH NKKP-Advanced 150038.

## REFERENCES

- [1] J.D. Bekenstein. Black holes and entropy. *Physical Review D*, 7(8):2333, 1973.
- [2] S.W. Hawking. Black hole explosions? Nature, 248(5443):30-31, 1974.
- [3] S.W. Hawking and D.N. Page. Thermodynamics of black holes in Anti-de Sitter space. Communications in Mathematical Physics, 87(4):577– 588, 1983.
- [4] R.M. Wald. General relativity. University of Chicago press, 2010.

- [5] T. L. Hill. Thermodynamics of small systems. Courier Corporation, 1994.
- [6] L. Smarr. Mass formula for kerr black holes. *Physical Review Letters*, 30(2):71, 1973.
- [7] T. Matolcsi. Ordinary thermodynamics. Akadémiai Kiadó Budapest, 2004.
- [8] E. Spallucci and A. Smailagic. Maxwell's equal area law and the Hawking-Page phase transition. *Journal of Gravity*, 2013, 2013.
- [9] T. S. Biró. Is there a temperature?: conceptual challenges at high energy, acceleration and complexity, volume 171. Springer Science & Business Media, 2011.
- [10] T.S. Biró, V.G. Czinner, H. Iguchi, and P. Ván. Black hole horizons can hide positive heat capacity. *Physics Letters B*, 782:228–231, 2018.
- [11] T.S. Biró, V.G. Czinner, H. Iguchi, and P. Ván. Volume dependent extension of Kerr-Newman black hole thermodynamics. *Physics Letters B*, 803:135344, 2020.
- [12] B.P. Dolan. Where is the PdV in the first law of black hole thermodynamics? *Open Questions in Cosmology*, 2012.
- [13] A. Abac et al. The science of the einstein telescope. arXiv preprint arXiv:2503.12263, 2025. Einstein Telescope Collaboration.