

Probing the general 2HDM with flavor-violation through $A \rightarrow ZH$

Wei-Shu Hou and Mohamed Krab

Department of Physics, National Taiwan University, Taipei 10617, Taiwan

We investigate the Large Hadron Collider (LHC) discovery prospects for a second Higgs doublet through $gg \rightarrow A$, followed by $A \rightarrow ZH$ weak decay, where A and H are exotic CP -odd and CP -even scalars of the general two Higgs doublet model (G2HDM) that has flavor-changing neutral Higgs (FCNH) couplings. The FCNH extra top Yukawa coupling ρ_{tc} and flavor-conserving ρ_{tt} coupling can drive electroweak baryogenesis (EWBG) to account for the disappearance of antimatter from the Universe. With dominant $H \rightarrow t\bar{c} + \bar{c}t$ decay, it gives rise to trilepton final states. By a phenomenological analysis, we show that the $A \rightarrow ZH \rightarrow \ell^+\ell^-t\bar{c}$ final state could be a promising probe of G2HDM at the LHC with flavor-violation.

Introduction.—The existence of a second Higgs doublet introduces extra Higgs bosons beyond the one discovered at the LHC [1, 2], and could provide the mechanism to explain the matter-antimatter asymmetry of the Universe. Its discovery would confirm the non-minimal structure of the Higgs sector and perhaps shed light on the mechanism behind the baryon asymmetry of the Universe (BAU).

We specifically advocate the general two Higgs doublet model (G2HDM), i.e. without the usual Z_2 symmetry. With two identical weak doublets, it allows for a second set of Yukawa couplings, other than fermion masses. Though it has not garnered much attention, this model has quite a few merits.

Quartic couplings at $\mathcal{O}(1)$ can provide [3] prerequisite first-order electroweak phase transition (EWPT), needed for EW baryogenesis (EWBG), while $\mathcal{O}(1)$ extra top Yukawa couplings ρ_{tc} and ρ_{tt} can each drive [4] EWBG to account for BAU, one of the biggest mysteries. Given this, one must confront the stringent electron electric dipole moment (eEDM) bounds of ACME [5] and JILA [6], where a spectacular cancellation mechanism [7] allows one to survive these bounds.

Without the Z_2 symmetry, G2HDM allows for flavor-changing neutral Higgs (FCNH) couplings, such as $t \rightarrow ch$ [8]. Interestingly, this decay still has not been seen, as it is *alignment* protected, by a rather small h - H mixing between the two CP -even scalars. However, FCNH couplings can lead to *alignment enhanced* processes such as $cg \rightarrow tH/tA \rightarrow tt\bar{c}, t\bar{t}$ [9], while processes such as $cg \rightarrow bH^+$ [10] and $\bar{b}g \rightarrow \bar{c}H^+$ [11], along with $c\bar{b} \rightarrow H^+$ [12] resonant production, are not subject to Cabibbo-Kobayashi-Maskawa (CKM) suppression as in the usual 2HDM type-II.

In this Letter, we suggest searching for $A \rightarrow ZH$ in $\ell^+\ell^-t\bar{c}$ final states. We propose a benchmark scenario and perform a detector level Monte Carlo analysis at the 14 TeV LHC. We show that searches for the $A \rightarrow ZH$ decay into the $\ell^+\ell^-t\bar{c}$ final state are promising: Run 2 data may already lead to discovery, thus providing a probe of G2HDM that has flavor-violation.

G2HDM.—In the Higgs basis where only one doublet gives mass, the most general CP -conserving scalar poten-

tial for two doublet Φ and Φ' is given by [13, 14]

$$V(\Phi, \Phi') = \mu_{11}^2 |\Phi|^2 + \mu_{22}^2 |\Phi'|^2 - (\mu_{12}^2 \Phi^\dagger \Phi' + \text{H.c.}) + \frac{\eta_1}{2} |\Phi|^4 + \frac{\eta_2}{2} |\Phi'|^4 + \eta_3 |\Phi|^2 |\Phi'|^2 + \eta_4 |\Phi^\dagger \Phi'|^2 + \left[\frac{\eta_5}{2} (\Phi^\dagger \Phi')^2 + (\eta_6 |\Phi|^2 + \eta_7 |\Phi'|^2) \Phi^\dagger \Phi' + \text{H.c.} \right], \quad (1)$$

where all mass and quartic coupling parameters are chosen to be *real*, Φ generates the vacuum expectation value v for EW symmetry breaking through a first minimization condition of $\mu_{11}^2 = -\frac{1}{2}\eta_1 v^2$ while $\langle \Phi' \rangle = 0$, hence $\mu_{22}^2 > 0$. A second minimization condition $\mu_{12}^2 = \frac{1}{2}\eta_6 v^2$ reduces the number of parameters to nine.

In the Higgs basis, the CP -odd scalar A and the charged Higgs H^\pm are mass eigenstates. The CP -even mass eigenstates h and H are obtained by diagonalizing the mass-squared matrix through a rotation characterized by the angle γ , denoted as $\beta - \alpha$ in 2HDM type-II convention, which satisfies $s_\gamma c_\gamma = \eta_6 v^2 / (m_H^2 - m_h^2)$ [13, 14], with $c_\gamma \equiv \cos \gamma$ ($s_\gamma \equiv \sin \gamma$). The emergent alignment phenomenon, that h closely resembles the SM Higgs boson [15], implies a rather small h - H mixing angle c_γ .

The general Yukawa couplings are [13, 14]

$$\mathcal{L}_Y = -\frac{1}{\sqrt{2}} \sum_{f=u,d,\ell} \bar{f}_i \left[(-\lambda_{ij}^f \sin \gamma + \rho_{ij}^f \cos \gamma) h + (\lambda_{ij}^f \cos \gamma + \rho_{ij}^f \sin \gamma) H - i \text{sgn}(Q_f) \rho_{ij}^f A \right] R f_j - \bar{u}_i [(V \rho^d)_{ij} R - (\rho^{u\dagger} V)_{ij} L] d_j H^+ - \bar{\nu}_i \rho_{ij}^\ell R \ell_j H^+ + \text{H.c.}, \quad (2)$$

where $i, j = 1, 2, 3$ are the generation indices, $\lambda_{ij}^f \equiv \delta_{ij} \sqrt{2} m_i^f / v$, $\text{sgn}(Q_f) = +1(-1)$ for $f = u$ ($f = d, \ell$), $L, R = (1 \mp \gamma_5)/2$, and V is the CKM matrix. The λ_{ij}^f matrices are real and diagonal, while ρ_{ij}^f are in general complex and non-diagonal. Here, we consider a CP -conserving model and take the ρ_{ij}^f matrices as real but not necessarily Hermitian.

In the *alignment* limit of $c_\gamma \rightarrow 0$, h couples diagonally as the SM Higgs boson, while H interacts through the extra Yukawa couplings ρ_{ij}^f . Thus, besides mass-mixing




FIG. 1. Branching ratios of the CP -odd scalar A as a function of m_A for $m_H = 200$ GeV, $\rho_{tt} = 0.3$ and $\rho_{tc} = 0.1$.

hierarchy protection [8] of FCNH couplings, *alignment* provides [14] further safeguard, such as for $t \rightarrow ch$, without relying on natural flavor conservation [16]. So enforcing the Z_2 symmetry may be overkill. The H^+ and A couplings are independent of γ .

In what follows, we drop the superscript f and set all $\rho_{ij} = 0$ for simplicity, except ρ_{tt} and ρ_{tc} . We will return to turning on other ρ_{ij} s, especially ρ_{bb} and $\rho_{\tau\tau}$, later in the paper. We set $c_\gamma = 0$, and fix $m_h = 125$ GeV. In Fig. 1, we plot the branching ratios of $A \rightarrow t\bar{c}$, $t\bar{t}$, ZH for $m_H = 200$ GeV, $\rho_{tt} = 0.3$ and $\rho_{tc} = 0.1$. We assume $m_{H^+} \simeq m_A$ (custodial symmetry), which is required by EW precision observables, implying that the decay $A \rightarrow W^-H^+$ is kinematically forbidden. We will return to $m_{H^+} \simeq m_H$ (twisted custodial [17]), where $A \rightarrow W^-H^+$ could be of similar magnitude to $A \rightarrow ZH$ later. We observe that $A \rightarrow ZH$ largely dominates over $A \rightarrow t\bar{c}$ and $A \rightarrow t\bar{t}$. Note that $A \rightarrow t\bar{c}$ dominates A decays for $m_A - m_H < m_Z$. The decay $A \rightarrow Zh$ is suppressed by c_γ . The subsequent decay of $H \rightarrow t\bar{c}$ is dominant, while $H \rightarrow W^+W^-$ and $H \rightarrow ZZ$ decays are c_γ -suppressed.

Constraints on G2HDM.—The model is subject to theoretical and experimental constraints. Higgs quartic couplings in Eq. (1) are required to fulfill positivity, tree-level unitarity, and perturbativity constraints, which are implemented in 2HDMC-1.8.0 [18]. G2DHM parameter space is further required to fulfill EW precision constraints through the oblique parameters S , T , and U [19] using the PDG values [20]. The code 2HDMC calculates the theoretical predictions for these observables. We refer to Ref. [11] for details on the parameter space scan of the G2HDM and constraints.

Flavor constraints on extra top Yukawa couplings ρ_{tt} and ρ_{tc} are relatively mild [21, 22], where it is found that [21, 22] B_q mixing ($q = d, s$) strongly constrains ρ_{ct} due to CKM $|V_{cq}/V_{tq}| \sim 25$ enhancement factor, hence ρ_{ct} must be tiny. Therefore, ρ_{ct} is turned off here. Following Ref. [22], we estimate the modifications




FIG. 2. Exclusion regions from $B \rightarrow X_s\gamma$, B_q mixing, and ATLAS $pp \rightarrow \bar{t}H^+b \rightarrow \bar{t}t\bar{b}b$ search in the m_{H^+} - $|\rho_{tt}|$ plane.

to the B_q - \bar{B}_q mixing amplitude and use the UTfit results [23]: $C_{B_d} = 1.09 \pm 0.09$ and $C_{B_s} = 1.10 \pm 0.06$, where $C_{B_q} \equiv M_{12}^q/M_{12}^{q, \text{SM}}$, while the phase ϕ_{B_q} is neglected in our study. We find the light blue region (extending to upper left) shown in Fig. 2 excluded at 2σ level. The B_q mixing constraint on ρ_{tc} is weaker [21]. Reinterpreting the limits from Ref. [21], we find the bound $|\rho_{tc}| \lesssim 1.7$ for $m_{H^+} = 500$ GeV.

The $B \rightarrow X_s\gamma$ constraint on m_{H^+} in G2HDM is not particularly strong [21, 22]. Ref. [22] finds that ρ_{bb} is more constrained than ρ_{tt} due to an enhancement factor of m_t/m_b , hence ρ_{bb} must be small. Setting $\rho_{bb} = 0$, we find the yellow exclusion region depicted in Fig. 2. The $B \rightarrow X_s\gamma$ constraint on ρ_{tc} is much weaker due to the small charm quark mass and the absence of any enhancement factor [22].

LHC indirect and direct measurement set limits on ρ_{tt} , but these are weaker [24] due to c_γ suppression [see Eq. (2)]. We further check these constraints using the new version of the code HiggsSignals [25], which incorporates the latest ATLAS and CMS Run 2 measurements, via HiggsTools [26]. ATLAS [27] and CMS [28] direct searches for $pp \rightarrow \bar{t}H^+(b)$ followed by $H^+ \rightarrow t\bar{b}$ (charge conjugate process implied) constrain ρ_{tt} as well. These searches set significant limits on $\sigma[pp \rightarrow \bar{t}H^+(b)] \times \mathcal{B}(H^+ \rightarrow t\bar{b})$. We reinterpret ATLAS limits [27] assuming $\mathcal{B}(H^+ \rightarrow t\bar{b}) = 100\%$, and obtain the light green exclusion region (extending upward) shown in Fig. 2. CMS limits [28] are also reinterpreted and found to be weaker; thus, only the ATLAS bounds are shown.

ATLAS [29] and CMS [30] have recently performed searches for exotic Higgs bosons with flavor-violating couplings. The analysis is interpreted in G2DHM with extra top Yukawa couplings ρ_{tt} , ρ_{tc} , and ρ_{tu} , which is a remarkable step toward searching for new Higgs bosons. The search results are used to constrain ρ_{tt} , where bounds on ρ_{tc} and ρ_{tu} are particularly stringent. For non-zero c_γ , LHC $t \rightarrow qh$ ($q = u, c$) searches [31, 32] set further limits on ρ_{tc} and ρ_{tu} . For $c_\gamma = 0.1$, we find that $|\rho_{tc}| \gtrsim 0.5$




FIG. 3. $P_T^{\ell_1}$ (left), H_T (middle), and $m_{\ell^+\ell^-}$ (right) distributions for background and signal processes.

($|\rho_{tu}| \gtrsim 0.38$) is excluded at 95% confidence level. The limit decreases for $c_\gamma < 0.1$ and disappears for $c_\gamma = 0$, which we assume in our analysis. Note that D - \bar{D} mixing and $B \rightarrow \tau\nu$ observables impose strong bounds on ρ_{tu} [21], requiring it to be very small; we set it to zero in our study.

Before discussing LHC search for $A \rightarrow ZH$ in $\ell^+\ell^-t\bar{c}$ final states at 14 TeV, we point out that the chosen benchmark model ($m_H = 200$ GeV, $|\rho_{tt}| = 0.3$, and $|\rho_{tc}| = 0.1$) evades all the constraints mentioned above.

LHC search for $A \rightarrow ZH$.— We now discuss the LHC search prospects for the $A \rightarrow ZH \rightarrow \ell^+\ell^-t\bar{c}$ channel, which leads to a final state consisting of three leptons plus at least two jets, with at least one b -tagged and one non- b -tagged, and missing transverse energy. We use the simple *cut and count* method and assume 14 TeV collision energy. We choose $m_A = 450$ (500) GeV as a benchmark point, denoted BP (BP'), where $\mathcal{B}(A \rightarrow ZH) \simeq 87$ (90)%. This benchmark could also be used to search for charged Higgs bosons via $H^+ \rightarrow W^+H$ [12]. Our signal can be searched for via $pp \rightarrow A \rightarrow ZH \rightarrow \ell^+\ell^-(t\bar{c} + \bar{t}c) + X$, with $t \rightarrow \ell^+\nu b$, where $\ell = e$ or μ . The dominant SM backgrounds are WZ , tZj , $t\bar{t}Z$, $t\bar{t}W$, and tZc , with $t\bar{t}h$ and $t\bar{t}\bar{t}$ are subdominant. Other backgrounds such as $t\bar{t}j$ and $t\bar{t}W$ are minor.

Signal and background samples are generated at leading order (LO) using `MadGraph5_aMC@NLO` [33]. These samples are passed to `Pythia-8.2` [34] for parton showering and hadronization, and then to `Delphes-3.5.0` [35], with a default ATLAS card for detector simulation. We consider one additional jet for WZ , $t\bar{t}Z$, and $t\bar{t}W$ backgrounds using the MLM matching scheme [36]. The tZj , tZc , $t\bar{t}h$, $t\bar{t}\bar{t}$ backgrounds and signal samples are estimated without additional jets.

We rescale the LO cross sections for our signal and the dominant background WZ to their respective next-to-next-to-leading-order values using K factors of 2.14 (2.09) for BP (BP'), and 2.07 (2.02) for W^-Z (W^+Z) [37]. Note that the signal K -factor is computed using the code `SusHi` [38]. The LO tZj , $t\bar{t}Z$, $t\bar{t}W^-$ ($t\bar{t}W^+$), $t\bar{t}h$, and $t\bar{t}\bar{t}$ backgrounds are adjusted to next-

to-leading-order (NLO) by factors of 1.44 [33], 1.40 [39], 1.54 (1.51) [39], 1.26 [39], and 2.04 [33], respectively. The tZc background is assumed to have the same NLO correction factor as tZj .

For event selection, we require the presence of exactly three charged leptons (samples with more than three leptons are vetoed), with $p_T^{\ell_1} > 80$, $p_T^{\ell_2} > 30$, and $p_T^{\ell_3} > 20$ GeV, and $|\eta_\ell| < 2.5$, together with at least two jets, of which at least one b -tagged and one non- b -tagged, with $P_T^j > 20$ GeV and $|\eta_j| < 2.5$, plus a missing transverse energy E_T^{miss} larger than 20 GeV. The scalar p_T sum of all jets and the three charged leptons (H_T) is required to be between 280 and 500 GeV. To further suppress the $t\bar{t}W$, $t\bar{t}h(\rightarrow WW^*)$, and $t\bar{t}\bar{t}$ backgrounds, the invariant mass $m_{\ell^+\ell^-}$ of all possible same-flavor, opposite-sign $\ell^+\ell^-$ pairs must lie within the mass window $70 < m_{\ell^+\ell^-} < 110$ GeV.

The $P_T^{\ell_1}$, H_T , and $m_{\ell^+\ell^-}$ distributions, before any selection cuts (only default cuts during event generation in `MadGraph5_aMC@NLO` are applied), are shown in Fig. 3. Signal and background cross sections, after selection cuts, are given in Table I. Note that the $t\bar{t}$ + jets process can also contribute if a jet is misidentified as a lepton (fake). We adopt a conservative misidentification rate of $\epsilon_{\text{fake}} = 10^{-4}$ [40, 41], and find the resulting cross section is below 0.001 after selection cuts. Thus, we neglect the fake contribution in our analysis.

Based on the results given in Table I, we determine

	Cross section
BP (BP')	0.87 (0.53)
WZ	0.81
tZj	0.36
$t\bar{t}Z$	0.17
$t\bar{t}W$	0.036
tZc	0.034
$t\bar{t}h$	0.002
$t\bar{t}\bar{t}$	< 0.001

TABLE I. Signal and background cross sections (in fb) at 14 TeV after selection cuts.

the statistical significance for our signal with $\mathcal{Z} = \sqrt{2}[(S+B)\ln(1+S/B) - S]$ [42], where S (B) is the number of signal (background) events. For 140 fb^{-1} , the significance for $A \rightarrow ZH \rightarrow \ell^+\ell^-t\bar{c}$ is $\sim 7.9\sigma$ ($\sim 5.0\sigma$) for BP (BP'). Thus, the available Run 2 data could lead to discovery. Assuming a conservative 10% systematic uncertainty in the background estimation, we find a significance of $\sim 4.4\sigma$ ($\sim 2.8\sigma$) for BP (BP').

Discussion and Summary. — ATLAS [29] and CMS [30] have recently conducted searches for G2HDM Higgs bosons, marking a remarkable step by exploring new channels beyond the conventional ones. G2HDM is a quite robust model, as it can drive [4] EWBG that could explain the observed matter-antimatter asymmetry of the Universe. It is found [43] that a strong first-order EWPT in 2HDM, as needed for EWBG, favors a rather heavy CP -odd Higgs boson with $m_A > 300 \text{ GeV}$, and a large mass splitting $m_A - m_H \gtrsim v$. Hence, the decay $A \rightarrow ZH$ is identified as a smoking gun signature of 2HDM scenarios with a strongly first-order EWPT, and searches for $A \rightarrow ZH$ in $\ell^+\ell^-b\bar{b}$ and $\ell^+\ell^-W^+W^-$ final states are found promising [43]. In addition, $A \rightarrow ZH$ can also be explored in the $\ell^+\ell^-t\bar{t}$ and $\nu\nu b\bar{b}$ final states [44]. In this Letter, we have shown that $A \rightarrow ZH$ in $\ell^+\ell^-t\bar{c}$ final states is very promising, and the current Run 2 data may be able to discover this exotic decay through the trilepton signal. A signature that could distinguish G2HDM from type-I(II) 2HDM, hence observing it would be a smoking gun for G2HDM.

We have assumed a scenario where $m_A \simeq m_{H^+}$ (custodial), but one can on other hand consider the $m_H \simeq m_{H^+}$ case (twisted custodial [17]), with $m_{H^+} < m_A$. For example, let us consider $m_H = m_{H^+} = 200 \text{ GeV}$ and $m_A = 500 \text{ GeV}$ (with $|\rho_{tt}| = 0.3$, $|\rho_{tc}| = 0.1$). In this benchmark scenario, the CP -odd Higgs boson A would decay into W^-H^+ , with a branching ratio of $\sim 65\%$, which dilutes the $A \rightarrow ZH$ decay, thereby altering the expected number of events in $\ell^+\ell^-t\bar{c}$ final states. In addition, the decay $H^+ \rightarrow c\bar{b}$ would be enhanced, suggesting $A \rightarrow W^-H^+ \rightarrow \ell^-\bar{\nu}c\bar{b}$ could serve as an alternative discovery channel. However, for Higgs masses above 200 GeV (e.g., $m_H \simeq m_{H^+} \simeq 300 \text{ GeV}$), the $H^+ \rightarrow t\bar{b}$ decay would dominate, making the $A \rightarrow W^-H^+ \rightarrow \ell^-\bar{\nu}t\bar{b}$ final state potentially more promising.

Away from the *alignment* limit, $A \rightarrow Zh$, $H \rightarrow W^+W^-$, and $H \rightarrow ZZ$ would be induced. For $c_\gamma = 0.1$, which agrees with SM Higgs measurements at 2σ level by **HiggsSignals**, the decay $A \rightarrow Zh$ remains negligible. However, $H \rightarrow W^+W^-$ (ZZ) is important and has a branching ratio of $\sim 35\%$ ($\sim 16\%$), diluting $H \rightarrow t\bar{c} + \bar{t}c$. For non-zero c_γ , $A \rightarrow ZH \rightarrow \ell^+\ell^-t\bar{c}$ and $A \rightarrow ZH \rightarrow \ell^+\ell^-W^+W^-$ would both lead to trilepton signal, which is itself another G2HDM bonus. Note that $A \rightarrow ZH \rightarrow \ell^+\ell^-ZZ$ is also possible, but it would require much more data for potential discovery.

So far we have only considered extra top Yukawa couplings ρ_{tt} and ρ_{tc} . Turning on ρ_{bb} ($\rho_{\tau\tau}$) would induce $H \rightarrow b\bar{b}$ ($\tau^+\tau^-$), and thus one can have $A \rightarrow ZH \rightarrow \ell^+\ell^-b\bar{b}$. However, flavor observables strongly constrain ρ_{bb} ($\rho_{\tau\tau}$), requiring $|\rho_{bb}| \lesssim \lambda_b$ ($|\rho_{\tau\tau}| \lesssim \lambda_\tau$) [21, 22]. For $\rho_{bb} \sim \rho_{\tau\tau} \sim 0.01$, $H \rightarrow b\bar{b}$ and $H \rightarrow \tau^+\tau^-$ are small, and their presence would slightly soften $H \rightarrow t\bar{c} + \bar{t}c$ but would not alter our conclusions. Note that non-zero ρ_{bb} and $\rho_{\tau\tau}$ would also induce $A \rightarrow b\bar{b}$ and $A \rightarrow \tau^+\tau^-$, but these decays are very small.

In summary, we suggest searching for $A \rightarrow ZH$ in $\ell^+\ell^-t\bar{c}$ final states, a smoking gun signature for G2HDM. We proposed a benchmark scenario and carried out a signal-to-background analysis at the 14 TeV LHC, and show that $A \rightarrow ZH$ is quite promising. Even current LHC Run 2 data may give potential discovery of this exotic decay into $\ell^+\ell^-t\bar{c}$ final states, thus providing a probe of G2HDM with flavor-violation.

Acknowledgments. — This work is supported by NSTC Grant No. 113-2639-M-002-006-ASP of Taiwan, and NTU Grants No. 113L86001 and No. 113L891801. MK acknowledges CERN TH Department for hospitality, where part of this work was carried out.

-
- [1] G. Aad *et al.* [ATLAS], Phys. Lett. B **716**, 1 (2012).
 - [2] S. Chatrchyan *et al.* [CMS], Phys. Lett. B **716**, 30 (2012).
 - [3] S. Kanemura, Y. Okada, and E. Senaha, Phys. Lett. B **606**, 361 (2005).
 - [4] K. Fuyuto, W.-S. Hou and E. Senaha, Phys. Lett. B **776**, 402 (2018).
 - [5] V. Andreev *et al.* (ACME Collaboration), Nature **562**, 355 (2018).
 - [6] T.S. Roussy, *et al.* Science **381**, 46 (2023).
 - [7] K. Fuyuto, W.-S. Hou and E. Senaha, Phys. Rev. D **101**, 011901(R) (2020).
 - [8] W.-S. Hou, Phys. Lett. B **296**, 179 (1992).
 - [9] M. Kohda, T. Modak, and W.-S. Hou, Phys. Lett. B **776**, 379 (2018).
 - [10] D.K. Ghosh, W.-S. Hou, and T. Modak, Phys. Rev. Lett. **125**, 221801 (2020).
 - [11] W.-S. Hou and M. Krab, Phys. Rev. D **110**, L011702 (2024).
 - [12] W.-S. Hou and M. Krab, Phys. Rev. D **111**, L031701 (2025).
 - [13] S. Davidson and H.E. Haber, Phys. Rev. D **72**, 035004 (2005).
 - [14] W.-S. Hou and M. Kikuchi, EPL **123**, 11001 (2018).
 - [15] G. Aad *et al.* [ATLAS and CMS], JHEP **08** (2016) 045.
 - [16] S. L. Glashow and S. Weinberg, Phys. Rev. D **15**, 1958 (1977).
 - [17] J.-M. Gérard and M. Herquet, Phys. Rev. Lett. **98**, 251802 (2007).
 - [18] D. Eriksson, J. Rathsmann and O. Stål, Comput. Phys. Commun. **181**, 189 (2010).
 - [19] W. Grimus, L. Lavoura, O. M. Ogreid and P. Osland, J. Phys. G **35**, 075001 (2008); Nucl. Phys. B **801**, 81 (2008).

- [20] S. Navas *et al.* [Particle Data Group], Phys. Rev. D **110**, 030001 (2024).
- [21] A. Crivellin, C. Greub and A. Kokulu, Phys. Rev. D **87**, 094031 (2013).
- [22] B. Altunkaynak, W.-S. Hou, C. Kao, M. Kohda and B. McCoy, Phys. Lett. B **751**, 135 (2015).
- [23] <http://www.utfit.org/UTfit/ResultsSummer2023NP>.
- [24] W.-S. Hou, M. Kohda and T. Modak, Phys. Rev. D **98**, 075007 (2018).
- [25] P. Bechtle, S. Heinemeyer, T. Klingl, T. Stefaniak, G. Weiglein and J. Wittbrodt, Eur. Phys. J. C **81**, 145 (2021).
- [26] H. Bahl, T. Biekötter, S. Heinemeyer, C. Li, S. Paasch, G. Weiglein and J. Wittbrodt, Comput. Phys. Commun. **291**, 108803 (2023).
- [27] G. Aad *et al.* [ATLAS], JHEP **06** (2021) 145.
- [28] A.M. Sirunyan *et al.* [CMS], JHEP **07** (2020) 126.
- [29] G. Aad *et al.* [ATLAS], JHEP **12** (2023) 081.
- [30] A. Hayrapetyan *et al.* [CMS], Phys. Lett. B **850**, 138478 (2024).
- [31] A. Tumasyan *et al.* [CMS], Phys. Rev. Lett. **129**, 032001 (2022); A. Hayrapetyan *et al.* [CMS], arXiv:2407.15172.
- [32] G. Aad *et al.* [ATLAS], Eur. Phys. J. C **84**, 757 (2024).
- [33] J. Alwall *et al.*, JHEP **07** (2014) 079.
- [34] T. Sjöstrand *et al.*, Comput. Phys. Commun. **191**, 159 (2015).
- [35] J. de Favereau *et al.* [DELPHES 3], JHEP **02** (2014) 057.
- [36] J. Alwall *et al.* Eur. Phys. J. C **53**, 473 (2008).
- [37] M. Grazzini, S. Kallweit, D. Rathlev and M. Wiesemann, Phys. Lett. B **761**, 179 (2016).
- [38] R. V. Harlander, S. Liebler and H. Mantler, Comput. Phys. Commun. **184**, 1605 (2013).
- [39] D. de Florian *et al.* [LHC Higgs Cross Section Working Group], arXiv:1610.07922.
- [40] G. Aad *et al.* [ATLAS], Eur. Phys. J. C **76**, 259 (2016).
- [41] E. Alvarez, D.A. Faroughy, J.F. Kamenik, R. Morales, and A. Szykman, Nucl. Phys. B **915**, 19 (2017).
- [42] G. Cowan, K. Cranmer, E. Gross and O. Vitells, Eur. Phys. J. C **71**, 1554 (2011).
- [43] G. C. Dorsch, S. J. Huber, K. Mimasu and J. M. No, Phys. Rev. Lett. **113**, 211802 (2014).
- [44] T. Biekötter, S. Heinemeyer, J. M. No, K. Radchenko, M. O. O. Romacho and G. Weiglein, JHEP **01** (2024) 107.