

A search for the three isomers of cyano-1,3-butadiene in TMC-1: Implications for bottom-up routes involving 1,3-butadiene^{*}

M. Agúndez¹, C. Cabezas¹, N. Marcelino^{2,3}, B. Tercero^{2,3}, R. Fuentetaja¹, P. de Vicente³, and J. Cernicharo¹

¹ Instituto de Física Fundamental, CSIC, Calle Serrano 123, E-28006 Madrid, Spain
e-mail: marcelino.agundez@csic.es, jose.cernicharo@csic.es

² Observatorio Astronómico Nacional, IGN, Calle Alfonso XII 3, E-28014 Madrid, Spain

³ Observatorio de Yebes, IGN, Cerro de la Palera s/n, E-19141 Yebes, Guadalajara, Spain

Received; accepted

ABSTRACT

The molecule 1,3-butadiene ($\text{CH}_2\text{CHCHCH}_2$) could play a key role in the synthesis of the cyclic molecules cyclopentadiene and benzene in cold dense clouds. Since 1,3-butadiene is non-polar, we searched for its cyano derivative, which exists in the form of three different polar isomers, in the cold dense cloud TMC-1. We used the most recent data obtained with the Yebes 40m telescope in the Q band (31.0-50.3 GHz) in the frame of the QUIJOTE project. We do not detect any of the two isomers of 1-cyano-1,3-butadiene, and derive 3σ upper limits to their column densities of $1.2 \times 10^{10} \text{ cm}^{-2}$ and $2.0 \times 10^{10} \text{ cm}^{-2}$ for *E*- and *Z*-1-cyano-1,3-butadiene, respectively. Our results are not consistent with those from Cooke et al. (2023), who determine a column density of $3.8 \times 10^{10} \text{ cm}^{-2}$ for *E*-1-cyano-1,3-butadiene in TMC-1 using GBT data and a line stack technique. At the current level of sensitivity of our data, there is tentative evidence for the presence of the third cyano derivative isomer, 2-cyano-1,3-butadiene, although a firm detection must await more sensitive data. We derive an upper limit to its column density of $3.1 \times 10^{10} \text{ cm}^{-2}$. This isomer cannot be formed in the reaction between CN and 1,3-butadiene, according to experimental and theoretical studies, and thus we speculate whether it could arise from neutral-neutral reactions like $\text{C}_2\text{H}_3 + \text{CH}_2\text{CHCN}$ and $\text{CH}_2\text{CCN} + \text{C}_2\text{H}_4$. From the upper limit on the abundance of 1-cyano-1,3-butadiene derived here, we estimate that the abundance of 1,3-butadiene in TMC-1 is below 10^{-11} - 10^{-10} relative to H_2 . The low abundance inferred for 1,3-butadiene makes it unlikely that it plays an important role in bottom-up routes to cyclopentadiene and benzene.

Key words. astrochemistry – line: identification – ISM: individual objects (TMC-1) – ISM: molecules – radio lines: ISM

1. Introduction

The hydrocarbon 1,3-butadiene ($\text{CH}_2\text{CHCHCH}_2$) is a very interesting molecule from an interstellar perspective because it shows a great reactivity with small hydrocarbon radicals, resulting in cyclization to form simple rings such as cyclopentadiene and benzene. The interest on the formation of these cycles in interstellar space is high because they are known or inferred to be abundant in cold dense clouds. Cyclopentadiene and its cyano- and ethynyl-derivatives (including the isomer of ethynyl cyclopentadiene fulvenallene) have been detected in TMC-1 (Cernicharo et al. 2021a,b, 2022a; McCarthy et al. 2021; Lee et al. 2021a), while in the case of benzene, its cyano and ethynyl derivatives have been detected as well (McGuire et al. 2018; Loru et al. 2023) and there is ample evidence based on chemical kinetics (Woon 2006; Goulay et al. 2006; Balucani et al. 1999, 2000; Landera et al. 2008; Trevitt et al. 2010; Jones et al. 2010; Cooke et al. 2020) that they are a proxy of the radio invisible molecule benzene. These mono-ringed structures are likely to open the door to form larger fused rings involving two or three cycles, several of which, such as indene, naphthalene, acenaphthylene, or pyrene, have been detected directly or through

their cyano derivatives in TMC-1 (Cernicharo et al. 2021a, 2024; Burkhardt et al. 2021a; McGuire et al. 2021; Wenzel et al. 2024). Moreover, some of these aromatic cycles have been also detected in other cold dense clouds, in addition to TMC-1 (Burkhardt et al. 2021b; Agúndez et al. 2023a).

Several theoretical and experimental studies have provided evidence that 1,3-butadiene can be a precursor of cycles under interstellar conditions. For example, the reaction of 1,3-butadiene with C_2H is a promising route to form benzene (Jones et al. 2011) and fulvene (Lockyear et al. 2015), while its reaction with CH can produce cyclopentadiene (He et al. 2020), which in turn can react with CH to produce benzene (Caster et al. 2019, 2021), and even 1,3-butadiene reacting with CH_3CC could form toluene (Thomas et al. 2019). The radical CH_3CC has not been for the moment observed in interstellar space but the radicals CH and C_2H are known to be abundant in cold dense clouds such as TMC-1 (Agúndez & Wakelam 2013), making the aforementioned reactions promising routes to form cyclopentadiene and benzene.

The validation of the hypothesis that 1,3-butadiene is a key intermediate in the bottom-up synthesis of large cycles relies on whether it is abundant or not in cold dense clouds. 1,3-butadiene has several rotamers depending on the internal rotation about the single C-C bond. The most stable form is the *s-trans*, while the *s-gauche* and *s-cis* lie 3.0 and 3.5 kcal mol⁻¹ (1500 and 1750 K), respectively, above the *s-trans*, according to the theoretical calculations by Feller & Craig (2009). The *s-trans* rotamer is

^{*} Based on observations carried out with the Yebes 40m telescope (projects 19A003, 20A014, 20D023, 21A011, 21D005, and 23A024). The 40m radio telescope at Yebes Observatory is operated by the Spanish Geographic Institute (IGN; Ministerio de Transportes, Movilidad y Agenda Urbana).




Fig. 1. Geometrical structure of the three cyano derivatives of 1,3-butadiene resulting from the substitution of one terminal H atom (*E* and *Z* isomers of 1-cyano-1,3-butadiene) and one internal H atom (2-cyano-1,3-butadiene).

non-polar and thus cannot be detected at radio wavelengths. The next rotamers are expected to be polar but they have proven to be very elusive to experimental characterization.

Given that the lowest energy rotamer of 1,3-butadiene is non-polar, its detection must rely on indirect methods. One possibility is to search for its protonated form, as done for other non-polar molecules, such as N_2 , CO_2 , C_3 , C_5 , $NCCN$, and NC_4N (Thaddeus & Turner 1975; Minh et al. 1991; Pety et al. 2012; Cernicharo et al. 2022b; Agúndez et al. 2015, 2023b). However, to our knowledge, the rotational spectrum of protonated 1,3-butadiene has not been measured in the laboratory, which prevents its radioastronomical search. An alternative indirect method is to search for the cyano substituted form. Indeed, several non-polar molecules, such as ethane, benzene, and naphthalene, are inferred to be present in cold dense clouds thanks to the detection of their cyano derivatives (Cernicharo et al. 2021c; McGuire et al. 2018, 2021). The hypothesis that cyano derivatives can be used as a proxy of 1,3-butadiene is validated because experimental and theoretical studies show that 1,3-butadiene reacts rapidly with the radical CN at low temperatures (Bullock & Cooper 1971; Butterfield et al. 1993; Morales et al. 2011; Gardez et al. 2012), forming the polar molecule 1-cyano-1,3-butadiene (Morales et al. 2011; Sun et al. 2014), which can thus be searched provided their rotational spectra are known.

The rotational spectrum of 1-cyano-1,3-butadiene has been recently measured in the laboratory (McCarthy et al. 2020; Zdanovskaia et al. 2021a), while that of 2-cyano-1,3-butadiene has been measured as well (Zdanovskaia et al. 2021a). Based on these laboratory data, Cooke et al. (2023) found statistical evidence of *E*-1-cyano-1,3-butadiene in TMC-1 using astronomical spectra obtained with the GBT telescope and a line stack technique. The statistical evidence of this isomer of cyano-1,3-butadiene is however modest, with a signal-to-noise ratio of 5.1σ . Given the importance of 1,3-butadiene as a potential precursor in bottom-up routes to synthesize large cycles in cold dense clouds, here we revisit the evidence for the presence of cyano-1,3-butadiene using independent data of TMC-1 obtained with the Yebes 40m telescope in the frame of the QUIJOTE (Q-band Ultrasensitive Inspection Journey to the Obscure TMC-1 Environment) project (Cernicharo et al. 2021d).

2. Astronomical observations

The astronomical observations presented here are part of the ongoing Yebes 40m Q-band line survey of TMC-1 QUIJOTE (Cernicharo et al. 2021d), which consists of a Q-band line survey (31.0-50.3 GHz) of the dark cloud TMC-1 at the position of the cyanopolyne peak ($\alpha_{J2000} = 4^h41^m41.9^s$ and $\delta_{J2000} = +25^\circ41'27.0''$). The observations are carried out using the frequency-switching observing technique, with a frequency throw of either 8 or 10 MHz. The full Q band is covered in one shot with a spectral resolution of 38.15 kHz in both polarizations using a 7 mm dual linear polarization receiver connected to a set of 2×8 fast Fourier transform spectrometers (Tercero et al. 2021). The intensity scale at the telescope is the antenna temperature, T_A^* , which has an estimated uncertainty due to calibration of 10%, which can be converted to main beam brightness temperature, T_{mb} , by dividing T_A^* by B_{eff}/F_{eff} , where the beam efficiency B_{eff} is given by the Ruze formula $B_{eff} = 0.797 \exp[-(\nu/71.1)^2]$, where ν is the frequency in GHz, and the forward efficiency F_{eff} is 0.97. The half power beam width (HPBW) can be approximated as $HPBW('') = 1763/\nu(\text{GHz})$. The latest dataset of QUIJOTE used here includes observations carried out between November 2019 and July 2024. The total on-source telescope time is 1509.2 h, of which 736.6 h correspond to a frequency throw of 8 MHz and 772.6 h to a throw of 10 MHz. The T_A^* rms noise level varies between 0.06 mK at 32 GHz and 0.18 mK at 49.5 GHz. The procedure used to reduce and analyze the data is described in Cernicharo et al. (2022a). The data were analyzed using the GILDAS software¹.

3. Spectroscopy of cyano-1,3-butadiene

The substitution of one hydrogen atom by a $-CN$ group in 1,3-butadiene can result in different isomers (see Fig. 1). If the substitution occurs in one of the four hydrogen atoms of the terminal carbon atoms, 1-cyano-1,3-butadiene is formed, while substitution in one of the two hydrogen atoms of the internal carbon atoms results in the 2-cyano-1,3-butadiene isomer. The most stable isomer is 1-cyano-1,3-butadiene, which

¹ <https://www.iram.fr/IRAMFR/GILDAS/>




Fig. 2. Spectra of TMC-1 in the Q band at the frequencies of the most favorable lines of *E*-1-cyano-1,3-butadiene. The noise level, as measured in a window of ± 8 MHz around the expected position of each line with the nominal spectral resolution of 38.15 kHz, is indicated by a gray horizontal band. The red lines correspond to the line intensities calculated adopting as column density for *E*-1-cyano-1,3-butadiene the 3σ upper limit of $1.2 \times 10^{10} \text{ cm}^{-2}$ determined here, while the blue dashed lines correspond to the line intensities calculated adopting as column density the value of $3.8 \times 10^{10} \text{ cm}^{-2}$ determined by Cooke et al. (2023). Our data is clearly inconsistent with such column density.

in turn has *E* and *Z* isomers, the *E* form being $0.3 \text{ kcal mol}^{-1}$ more stable than the *Z* form (Zdanovskaia et al. 2021a). The isomer 2-cyano-1,3-butadiene is 1 kcal mol^{-1} less stable than *E*-1-cyano-1,3-butadiene (Sun et al. 2014). These three isomers can exist in different conformations, the lowest in energy is *s-trans* for the *E* and *Z* forms of 1-cyano-1,3-butadiene and *syn* for 2-cyano-1,3-butadiene, all of them with sizable dipole moments of $\mu_a = 4.6 \text{ D}$ and $\mu_b = 1.0 \text{ D}$ for *E*-1-cyano-1,3-butadiene, $\mu_a = 3.6 \text{ D}$ and $\mu_b = 2.3 \text{ D}$ for *Z*-1-cyano-1,3-butadiene, and $\mu_a = 3.2 \text{ D}$ and $\mu_b = 2.3 \text{ D}$ for 2-cyano-1,3-butadiene (Zdanovskaia et al. 2021a,b).

The rotational spectra of the lowest *s-trans* conformers of the *E* and *Z* isomers of 1-cyano-1,3-butadiene have been characterized in the laboratory in the frequency ranges 6.5–26 GHz (McCarthy et al. 2020) and 130–375 GHz (Zdanovskaia et al.

2021a), while that of *syn*-2-cyano-1,3-butadiene has been measured in the laboratory by Zdanovskaia et al. (2021b) in the frequency range 130–360 GHz. Frequency predictions for the three cyano-1,3-butadiene isomers in the Q band were done using the molecular parameters derived from the rotational spectra analysis (Zdanovskaia et al. 2021a,b) employing the SPCAT program (Pickett 1991). In the case of the isomers of 1-cyano-1,3-butadiene, the availability of low-frequency data permitted to resolve the hyperfine structure due to the nitrogen nucleus, something that was not possible in the case of 2-cyano-1,3-butadiene. In this case, we estimated the ^{14}N nuclear quadrupole coupling constants through quantum chemical calculations at the B3LYP/cc-pVTZ level of theory. We used this method because it reproduces quite well the experimental values of χ_{aa} and χ_{bb} derived for the *E* and *Z* isomers of 1-cyano-1,3-butadiene (Mc-

Carthy et al. 2020; Zdanovskaia et al. 2021a). For the *E* isomer we calculate $\chi_{aa} = -3.53$ MHz and $\chi_{bb} = 1.40$ MHz, in good agreement with the experimental values of -3.33 and 1.28 MHz, respectively, while for the *Z* isomer we obtain $\chi_{aa} = -0.69$ MHz and $\chi_{bb} = -1.40$ MHz, in line with the experimental values of -0.36 and -1.65 MHz, respectively. For 2-cyano-1,3-butadiene we calculate $\chi_{aa} = -1.29$ MHz and $\chi_{bb} = -0.89$ MHz. For the three molecules, the frequency errors in the Q band are typically around 10-15 kHz for *a*- and *b*-type transitions with $K_a = 0, 1$. The hyperfine splitting for these lines is modest, on the order of a few tens of kHz.

4. Search for cyano-1,3-butadiene in TMC-1

We first searched for the isomer *E*-1-cyano-1,3-butadiene in our Yebes 40m data of TMC-1 because Cooke et al. (2023) announced its detection based on a line stack analysis of GBT data of the same cloud. In order to identify the most favorable lines of this molecule in the Q band we computed the line intensities at a rotational temperature of 9 K, which is the gas kinetic temperature in TMC-1 (Agúndez et al. 2023c). The most intense predicted lines correspond to *a*-type transitions with $K_a = 0$ and 1. We made a careful inspection of those lines in our QUIJOTE data. The $12_{0,12}-11_{0,11}$ transition at 33690.962 MHz coincides with the $J = 13-12$ line of HC_5NH^+ at 33690.975 MHz, which has a T_A^* intensity of 9.7 mK (Marcelino et al. 2020), while there is a clear line with a T_A^* intensity of 0.7 mK at the frequency of the $12_{1,12}-11_{1,11}$ transition, which is 33275.732 MHz. This line however cannot be due to *E*-1-cyano-1,3-butadiene because many other lines should be present with similar intensities, which is not the case. We do not have a clear assignment for this line, which for the moment remains unidentified. There are still many lines with T_A^* intensities below 1 mK in our QUIJOTE data that await identification.

In Figure 2 we show the spectra at the expected position of the most favorable lines of *E*-1-cyano-1,3-butadiene that lie in low-noise and relatively clean spectral regions. At the position of the $15_{0,15}-14_{0,14}$ and $16_{1,15}-15_{1,14}$ transitions the data show emission features with T_A^* intensities of 0.3 mK and 0.4 mK, respectively, which are probably real lines. However, their assignment to *E*-1-cyano-1,3-butadiene is inconsistent with the non-detection of other lines that should have comparable or even higher intensities. As for the rest of transitions shown in Fig. 2, for some of them there is a feature at the expected position of the line but it lies within the noise level of the spectrum. We must conclude that we do not have enough evidence for the presence of *E*-1-cyano-1,3-butadiene in TMC-1. The non-detection of the $13_{0,13}-12_{0,12}$ line at 36489.156 MHz provides the most stringent upper limit to the column density of this isomer of cyano-1,3-butadiene. Assuming a rotational temperature of 9 K and a line width of 0.6 km s^{-1} (Agúndez et al. 2023c), we derive a 3σ upper limit to the beam-averaged column density of *E*-1-cyano-1,3-butadiene in TMC-1 of $1.2 \times 10^{10} \text{ cm}^{-2}$, which is 3.2 times lower than the column density determined by Cooke et al. (2023). The calculated line profiles adopting as column density the 3σ upper limit derived here are shown in red in Fig. 2. If the column density of *E*-1-cyano-1,3-butadiene were $3.8 \times 10^{10} \text{ cm}^{-2}$, as derived by Cooke et al. (2023), the lines of this molecule should be clearly visible above the noise level in our sensitive Q band data (see blue dashed lines in Fig. 2). The calculated line profiles assume that the emission size of *E*-1-cyano-1,3-butadiene would be extended compared to the main beam of the Yebes 40m (35-57'' in the Q band) and GBT (20-94'' in the X, K and Ka bands) telescopes. Although we do not

have information on the true emission size of *E*-1-cyano-1,3-butadiene, if this species is indeed present in TMC-1, this is a reasonable assumption given the extended nature of the emission observed for several hydrocarbons and N-bearing molecules in TMC-1 (Pratap et al. 1997; Fossé et al. 2001; Cernicharo et al. 2023).

We also searched for the *Z* isomer of 1-cyano-1,3-butadiene in our TMC-1 data. As with the *E* isomer, for a rotational temperature of 9 K, the most favorable lines for its detection in the Q band correspond to *a*-type transitions with $K_a = 0$ and 1. In Fig. 3 we show spectra at the frequencies of some of these favorable lines that lie in uncontaminated and sensitive spectral regions. As in the case of the *E* isomer, for some of the transitions there is an emission feature around the expected position of the line. For example, in the case of the $10_{1,10}-9_{1,9}$ and $12_{0,12}-11_{0,11}$ transitions, there are emission features with T_A^* intensities of 0.3 mK and 0.5 mK, respectively, which are very likely real lines although they cannot be assigned to *Z*-1-cyano-1,3-butadiene because on the one hand they appear slightly shifted with respect to the expected position and on the other their intensities are inconsistent with the non-detection of other lines that should have comparable or even larger intensities. Here we conclude that *Z*-1-cyano-1,3-butadiene is not detected and derive a 3σ upper limit to its beam-averaged column density of $2.0 \times 10^{10} \text{ cm}^{-2}$.

The astronomical search for the third isomer of cyano-1,3-butadiene, which is 2-cyano-1,3-butadiene, was carried out similarly to that of the two other isomers. Again, the most favorable lines correspond to rotational transitions of *a*-type with quantum number $K_a = 0$ and 1. In this particular case, the transitions $J_{0,J} \rightarrow J-1_{0,J-1}$ and $J_{1,J} \rightarrow J-1_{1,J-1}$, which are predicted to be the brightest, tend to collapse to the same frequency as the quantum number J increases. Therefore, these pairs of lines stand out as the most favorable ones to search for 2-cyano-1,3-butadiene in the Q band. The transitions $7_{0,7}-6_{0,6}$ and $7_{1,7}-6_{1,6}$ are separated by 566 kHz and lie in a relatively noisy region of the spectrum (top-left panel in Fig. 4), while the transitions $8_{0,8}-7_{0,7}$ and $8_{1,8}-7_{1,7}$, which are separated by 94 kHz, coincide with a weak emission feature at the noise level seen as a shoulder in the red side of a $T_A^* \sim 0.5$ mK line (top-right panel in Fig. 4). The next two pairs of lines with $J = 9$ and $J = 10$ are separated by less than 15 kHz, which is smaller than the spectral resolution of 38.15 kHz of our TMC-1 data, and coincide with lines of T_A^* intensities of 0.5 mK that are seen above the noise level (two middle-right panels in Fig. 4). The next pair of lines, $11_{0,11}-10_{0,10}$ and $11_{1,11}-10_{1,10}$, lies again in a relatively noisy spectrum and no clear feature can be appreciated above the noise level (bottom-right panel in Fig. 4). Other transitions with $K_a = 1$ and 2 are not clearly seen at the current level of sensitivity of our data (remaining panels in Fig. 4). Therefore, the current evidence for the presence of 2-cyano-1,3-butadiene in TMC-1 comes from two lines that coincide with the pairs $9_{0,9}-8_{0,8}/9_{1,9}-8_{1,8}$ at 39364.05 MHz and $10_{0,10}-9_{0,9}/10_{1,10}-9_{1,9}$ at 43506.93 MHz. With just two lines and given the relatively high density of lines with T_A^* intensities below 1 mK, we must be cautious. We thus conclude that for the moment there is tentative evidence for the presence of 2-cyano-1,3-butadiene in TMC-1. We derive an upper limit to its column density of $3.1 \times 10^{10} \text{ cm}^{-2}$. The confirmation of this detection must await more sensitive data in the Q band. We note that, unlike in the cases of the *E* and *Z* isomers of 1-cyano-1,3-butadiene, the hyperfine splitting of the rotational transitions of 2-cyano-1,3-butadiene is somewhat uncertain because it is based on ab initio calculations and not in experimental measurements. The impact of the hyperfine splitting on the tentative detection is however likely to be small because the predicted splitting, which




Fig. 3. Spectra of TMC-1 in the Q band at the frequencies of the most favorable lines of Z-1-cyano-1,3-butadiene. The noise level, as measured in a window of ± 8 MHz around the expected position of each line with the nominal spectral resolution of 38.15 kHz, is indicated by a gray horizontal band. The red lines correspond to the line intensities calculated adopting as column density for Z-1-cyano-1,3-butadiene the 3σ upper limit of $2.0 \times 10^{10} \text{ cm}^{-2}$ determined here.

is similar to the experimentally determined ones for the two isomers of 1-cyano-1,3-butadiene, is comparable or smaller than the spectral resolution of the TMC-1 data.

5. Discussion

The non-detection of 1-cyano-1,3-butadiene and the tentative evidence of 2-cyano-1,3-butadiene can shed light on the abundance of 1,3-butadiene in TMC-1 and its potential role as intermediate in the synthesis of simple mono-ringed cyclic structures such as cyclopentadiene and benzene. To this regard it is interesting to have a look at the typical abundance ratios between a given molecule and its cyano derivative in TMC-1. There are various cases in which this abundance ratio has been observationally constrained. For example, the three $\text{C}_4\text{H}_3\text{N}$ isomers $\text{CH}_3\text{C}_3\text{N}$, CH_2CCHCN , and HCCCH_2CN are probably produced in the reactions of CN with CH_3CCH and CH_2CCH_2 (Carty et al. 2001; Balucani et al. 2000, 2002; Abeysekera et al. 2015). These three cyano derivatives together are 18 times less abundant than CH_3CCH , which is the polar hydrocarbon precursor detectable in TMC-1 (Marcelino et al. 2021; Cernicharo et al. 2022a). The abundance of propene is 90 times higher than the sum of the abundances of the five cyano derivatives detected in TMC-1 (Marcelino et al. 2007; Cernicharo et al. 2022c), which are likely formed in the reaction between propene and CN (Morales et al. 2010; Gu et al. 2008; Huang et al. 2009).

It is also known that maleonitrile (NC-CH=CH-CN) is 130 times less abundant than CH_2CHCN (Gratier et al. 2016; Agúndez et al. 2024), its likely precursor via the reaction with CN (Marchione et al. 2022). For other cyano derivatives observed in TMC-1, there is no experimental or theoretical evidence that the reaction with CN is the main source of them, although based on similar reactions it is a likely possibility. These cases comprise the cyano derivatives of vinyl acetylene, CHCCHCHCN and $\text{CH}_2\text{CHC}_3\text{N}$, which together account for 1/25 of the abundance of vinyl acetylene (Cernicharo et al. 2021c; Lee et al. 2021b), the cyanide $\text{CH}_2\text{CCHC}_3\text{N}$, which has an abundance 1/60 times that of CH_2CCHCCH (Shingledecker et al. 2021; Cernicharo et al. 2021e), the cyano derivatives of cyclopentadiene, which are 30 times less abundant than cyclopentadiene (Cernicharo et al. 2021a,b), and the cyano derivative of indene, which is present with an abundance 1/80 times of that of indene (Cernicharo et al. 2021a; Sita et al. 2022). It is also worth to note that the methyl cyanopolyynes $\text{CH}_3\text{C}_5\text{N}$ and $\text{CH}_3\text{C}_6\text{N}$ are 8-9 times less abundant than their potential precursors via reaction with CN, $\text{CH}_3\text{C}_4\text{H}$ and $\text{CH}_3\text{C}_6\text{H}$, respectively (Agúndez & Wakelam 2013; Siebert et al. 2022; Cernicharo et al. 2022a). In summary, the abundance ratios found in TMC-1 between a given precursor (via reaction with CN) and the resulting cyano derivative are in the range 8-130.

Based on the above observational information, we could estimate that 1,3-butadiene is present in TMC-1 with an abundance




Fig. 4. Spectra of TMC-1 in the Q band at the frequencies of the most favorable lines of 2-cyano-1,3-butadiene. The noise level, as measured in a window of ± 8 MHz around the expected position of each line with the nominal spectral resolution of 38.15 kHz, is indicated by a gray horizontal band. The red lines correspond to the line intensities calculated adopting a column density of 2-cyano-1,3-butadiene of $3.1 \times 10^{10} \text{ cm}^{-2}$.

8-130 times higher than that of 1-cyano-1,3-butadiene, which is the main product of the reaction between 1,3-butadiene and CN, according to crossed molecular beam experiments and theoretical calculations (Morales et al. 2011; Sun et al. 2014). Chemical model calculations can also provide some insights on the abundance ratio between 1,3-butadiene and 1-cyano-1,3-butadiene. In their chemical model, Cooke et al. (2023) predict a value of ~ 10 for this abundance ratio assuming that 1-cyano-1,3-butadiene is mostly destroyed through reactions with cations. We note that the abundance ratio of ~ 10 would increase if 1-cyano-1,3-butadiene is assumed to react rapidly with neutral atoms, such as C and H. In any case, it is encouraging that the findings from chemical models are consistent with the range of abundance ratios derived from observations. Since here we find that 1-cyano-1,3-butadiene is present with an abundance below 1.2×10^{-12} relative to H_2 , adopting the upper limit on the column density of

E-1-cyano-1,3-butadiene derived here and a column density of H_2 of 10^{22} cm^{-2} (Cernicharo & Guélin 1987), we can conclude that the abundance of 1,3-butadiene in TMC-1 is below 8-130 times that of 1-cyano-1,3-butadiene, i.e., approximately below 10^{-11} - 10^{-10} with respect to H_2 (see Fig. 5).

To shed further light on the abundance of 1,3-butadiene in cold dense clouds we carried out chemical modeling calculations using the latest release of the UMIST chemical network (Millar et al. 2024), which includes a reduced set of reactions involving 1,3-butadiene. We adopted physical conditions and elemental abundances typical of cold dense clouds, with the abundance of oxygen depleted to result in a C/O elemental abundance ratio of 1 (Agúndez & Wakelam 2013). There is evidence that oxygen could be depleted on dust grains in cold dense clouds (Jenkins 2009; Whittet 2010; Hincelin et al. 2011). This would lead to an increase in the gas-phase elemental C/O ratio and enhance




Fig. 5. Abundance of 1,3-butadiene calculated by the chemical model as a function of time (solid orange curve), estimated observational upper limit on the abundance of 1,3-butadiene in TMC-1 (horizontal orange band), and abundance observed (or inferred) for cyclopentadiene and benzene in TMC-1.

the abundances of C-bearing molecules (Agúndez & Wakelam 2013). The calculated abundance of 1,3-butadiene is rather low, it lies below 10^{-13} relative to H_2 at any time (see Fig. 5). According to the chemical model, 1,3-butadiene is mainly formed by the dissociative recombination with electrons of the cation $C_4H_7^+$, which arises mainly from the reaction $C_2H_4^+ + C_2H_4$, while it is mostly destroyed by reactions with neutral atoms, such as C and H, and abundant cations, such as HCO^+ . This chemical scheme reveals as insufficient to produce 1,3-butadiene with a high abundance. The non-detection of the two isomers of 1-cyano-1,3-butadiene is in line with a low abundance of 1,3-butadiene in TMC-1. The chemical model of Cooke et al. (2023) predicts an abundance of 1,3-butadiene somewhat higher than our model, although their maximum calculated abundance is also relatively low, a few times 10^{-12} relative to H_2 , which is within the conservative observational upper limit of $< 10^{-11}$ - 10^{-10} relative to H_2 estimated here.

Given that 1,3-butadiene seems to be less abundant than 10^{-11} - 10^{-10} relative to H_2 , it is unlikely that it plays a key role as precursor, via reactions with CH and C_2H (He et al. 2020; Jones et al. 2011), of cyclopentadiene and benzene, which are observed (or inferred in the case of benzene) to be present with rather large abundances, around 10^{-9} relative to H_2 (Cernicharo et al. 2021a,b, 2022a; see Fig. 5). In cold dense clouds, a precursor of a given molecule may be less abundant than the molecule itself, but this usually happens when the precursor is an unstable species, such as a radical or an ion. This is a general behavior predicted by chemical models, and in some cases also confirmed by observations. For example, in TMC-1 the cation H_2CCCH^+ is observed with a column density of $7.0 \times 10^{11} \text{ cm}^{-2}$ (Silva et al. 2023) and it is thought to be the precursor of H_2C_3 , which is found to be around three times more abundant (Cernicharo et al. 2022a). In this case, the precursor 1,3-butadiene is a stable molecule, and therefore, any species formed from it is unlikely to be more abundant. Note that, as discussed above, stable precursors reacting with CN produce cyano derivatives that are systematically less abundant. Although in each particular case, the relative abundances between a precursor and a product will depend on the particular reactivities of each of them, we find that given the stable nature of both the precursor 1,3-butadiene and the products cyclopentadiene and benzene, it is unlikely that

1,3-butadiene can act as precursor if it is less abundant than the putative products.

The non-detection of 1-cyano-1,3-butadiene implies that 1,3-butadiene is probably not an important precursor in the synthesis of the first mono-ringed cycles. However, if the tentative evidence for 2-cyano-1,3-butadiene is confirmed at some point in the future, it is not straightforward how could this isomer be formed. Given that experiments and theoretical calculations exclude that this isomer can be formed through the reaction of CN and 1,3-butadiene (Morales et al. 2011; Sun et al. 2014), other neutral-neutral reactions that could a priori form $CH_2C(CN)CHCH_2$ are



For these reactions to provide efficient routes to 2-cyano-1,3-butadiene in TMC-1, the reactants must be abundant enough and the reaction must be rapid and produce $CH_2C(CN)CHCH_2$. Given the relatively low abundance derived for 2-cyano-1,3-butadiene in TMC-1, $< 3.1 \times 10^{-12}$ relative to H_2 , the reactants involved in the two above reactions are probably abundant enough. CH_2CHCN and CH_2CCN have been detected in TMC-1 with abundances of 6.5×10^{-10} relative to H_2 (Gratier et al. 2016) and 2.5×10^{-11} (Cabezas et al. 2023), while the vinyl radical (C_2H_3) and ethylene (C_2H_4) are predicted by our chemical model to reach peak abundances relative to H_2 of $\sim 10^{-9}$ and $\sim 10^{-8}$, respectively. To our knowledge, the kinetics and product distribution of reactions (1) and (2) have not been investigated neither experimentally nor theoretically, and thus it is difficult to anticipate their expected behavior. Addition of vinyl radicals to multiple C–C bonds in hydrocarbons tends to have barriers (Ismail et al. 2007; Smith et al. 2020), although it is unknown whether reaction (1) would show a similar behavior. The kinetics of the radical CH_2CCN involved in reaction (2) is completely unexplored. Further studies on reactions (1) and (2) could shed light on their potential role in the synthesis of 2-cyano-1,3-butadiene.

6. Conclusions

We searched for the three isomers of cyano-1,3-butadiene in the cold dense cloud TMC-1 using the latest QUIJOTE data obtained with the Yebes 40m telescope. We do not find evidence for the *E* or *Z* isomers of 1-cyano-1,3-butadiene, and derive 3σ upper limits to their column densities of $1.2 \times 10^{10} \text{ cm}^{-2}$ and $2.0 \times 10^{10} \text{ cm}^{-2}$, respectively. Our 3σ upper limit on the column density of *E*-1-cyano-1,3-butadiene is 3.2 times lower than the column density determined by Cooke et al. (2023) using a line stack technique. We find tentative evidence for the presence of the third isomer, 2-cyano-1,3-butadiene, and derive an upper limit to its column density of $3.1 \times 10^{10} \text{ cm}^{-2}$. Since this isomer is unlikely to result from the reaction between CN and 1,3-butadiene, we suggest that it could arise from neutral-neutral reactions, such as $C_2H_3 + CH_2CHCN$ and $CH_2CCN + C_2H_4$. The upper limit imposed on the abundance of 1-cyano-1,3-butadiene allows us to estimate that the abundance of 1,3-butadiene in TMC-1 is below 10^{-11} - 10^{-10} relative to H_2 . This low abundance makes it unlikely that 1,3-butadiene plays an important role in bottom-up routes to the first mono-ringed cycles cyclopentadiene and benzene in cold dense clouds such as TMC-1. Therefore, other routes must be explored.

Acknowledgements. We acknowledge funding support from Spanish Ministerio de Ciencia, Innovación, y Universidades through grants PID2022-137980NB-I00 and PID2023-147545NB-I00.

References

- Abeyssekera, C., Joalland, B., Ariyasingha, N., et al. 2015, *J. Phys. Chem. Lett.*, 6, 1599
- Agúndez, M. & Wakelam, V. 2013, *Chem. Rev.*, 113, 8710
- Agúndez, M., Cernicharo, J., de Vicente, P., et al. 2015, *A&A*, 579, L10
- Agúndez, M., Marcelino, N., Tercero, B., & Cernicharo, J. 2023a, *A&A*, 677, L13
- Agúndez, M., Cabezas, C., Marcelino, N., et al. 2023b, *A&A*, 669, L1
- Agúndez, M., Marcelino, N., Tercero, B., et al. 2023c, *A&A*, 677, A106
- Agúndez, M., Bermúdez, C., Cabezas, C., et al. 2024, *A&A*, 688, L31
- Balucani, N., Asvany, O., Chang, A. H. H., et al. 1999, *J. Chem. Phys.*, 111, 7457
- Balucani, N., Asvany, O., Huang, L. C. L., et al. 2000, *ApJ*, 545, 892
- Balucani, N., Asvany, O., Kaiser, R. I., et al. 2002, *J. Phys. Chem. A*, 106, 4301
- Bullock, G. E. & Cooper, R. 1971, *Trans. Faraday Soc.*, 67, 3258
- Burkhardt, A. M., Lee, K. L. K., Changala, P. B., et al. 2021a, *ApJ*, 913, L18
- Burkhardt, A. M., Loomis, R. A., Shingledecker, C. N., et al. 2021b, *Nat. Astron.*, 5, 181
- Butterfield, M. T., Yu, T., & Lin, M. C. 1993, *Chem. Phys.*, 169, 129
- Cabezas, C., Tang, J., Agúndez, M., et al. 2023, *A&A*, 676, L5
- Carty, D., Le Page, V., Sims, I. R., & Smith, I. W. M. 2001, *Chem. Phys. Lett.*, 344, 310
- Caster, K. L., Donnellan, Z. N., Selby, T. M., & Goulay, F. 2019, *J. Phys. Chem.*, 123, 5692
- Caster, K. L., Selby, T. M., Osborn, D. L., et al. 2021, *J. Phys. Chem. A*, 125, 6927
- Cernicharo, J. & Guélin, M. 1987, 176, 299
- Cernicharo, J., Agúndez, M., Cabezas, C., et al. 2021a, *A&A*, 649, L15
- Cernicharo, J., Agúndez, M., Kaiser, R. I., et al. 2021b, *A&A*, 655, L1
- Cernicharo, J., Agúndez, M., Cabezas, C., et al. 2021c, *A&A*, 647, L2
- Cernicharo, J., Agúndez, M., Kaiser, R. I., et al. 2021d, *A&A*, 652, L9
- Cernicharo, J., Cabezas, C., Agúndez, M., et al. 2021e, *A&A*, 647, L3
- Cernicharo, J., Fuentetaja, R., Agúndez, M., et al. 2022a, *A&A*, 663, L9
- Cernicharo, J., Agúndez, M., Cabezas, C., et al. 2022b, *A&A*, 657, L16
- Cernicharo, J., Fuentetaja, R., Cabezas, C., et al. 2022c, *A&A*, 663, L5
- Cernicharo, J., Tercero, B., Marcelino, N., et al. 2023, *A&A*, 674, L4
- Cernicharo, J., Cabezas, C., Fuentetaja, R., et al. 2024, *A&A*, 690, L13
- Cooke, I. R., Gupta, D., Messinger, J. P., & Sims, I. R. 2020, *ApJ*, 891, L41
- Cooke, I. R., Xue, C., Changala, P. B., et al. 2023, *ApJ*, 948, 133
- Feller, D. & Craig, N. C. 2009, *J. Phys. Chem. A*, 113, 1601
- Fossé, D., Cernicharo, J., Gerin, M., & Cox, P. 2001, *ApJ*, 552, 168
- Gardez, A., Saidani, G., Biennier, L., et al. 2012, *Int. J. Chem. Kinet.*, 44, 753
- Goulay, F. & Leone, S. R. 2006, *J. Phys. Chem. A*, 110, 1875
- Gratier, P., Majumdar, L., Ohishi, M., et al. 2016, *ApJS*, 225, 25
- Gu, X. B., Zhang, F. T. & Kaiser, R. I. 2008, *J. Phys. Chem. A*, 112, 9607
- He, C., Zhao, L., Doddipatla, S., et al. 2020, *ChemPhysChem.*, 21, 1295
- Hincelin, U., Wakelam, V., Hersant, F., et al. 2011, *A&A*, 530, A61
- Huang, C. H., Kaiser, R. I. & Chang, H. H. 2009, *J. Phys. Chem. A*, 113, 12675
- Ismail, H., Goldsmith, F., Abel, P. R., et al. 2007, *J. Phys. Chem. A*, 111, 6843
- Jenkins, E. B. 2009, *ApJ*, 700, 1299
- Jones, B., Zhang, F., Maksyutenko, P., et al. 2010, *J. Phys. Chem. A*, 114, 5256
- Jones, B. M., Zhang, F., Kaiser, R. I., et al. 2011, *PNAS*, 108, 452
- Landera, A., Mebel, A. M., & Kaiser, R. I. 2008, *Chem. Phys. Lett.*, 459, 54
- Lee, K. L. K., Changala, P. B., Loomis, R. A., et al. 2021a, *ApJ*, 910, L2
- Lee, K. L. K., Loomis, R. A., Burkhardt, A. M., et al. 2021b, *ApJ*, 908, L11
- Lockyear, J. F., Fournier, M., Sims, I. R., et al. 2015, *Int. J. Mass Spectrom.*, 378, 232
- Loru, D., Cabezas, C., Cernicharo, J., et al. 2023, *A&A*, 677, A166
- Marcelino, N., Cernicharo, J., Agúndez, M., et al. 2007, *ApJ*, 665, L127
- Marcelino, N., Agúndez, M., Tercero, B., et al. 2020, *A&A*, 643, L6
- Marcelino, N., Tercero, B., Agúndez, M., & Cernicharo, J. 2021, *A&A*, 646, L9
- Marchione, D., Mancini, L., Liang, P., et al. 2022, *J. Phys. Chem. A*, 126, 3569
- McCarthy, M. C., Lee, K. L. K., Carroll, P. B., et al. 2020, *J. Phys. Chem. A*, 124, 5170
- McCarthy, M. C., Lee, K. L. K., Loomis, R. A., et al. 2021, *Nat. Astron.*, 5, 176
- McGuire, B. A., Burkhardt, A. M., Kalenskii, S., et al. 2018, *Science*, 359, 6372
- McGuire, B. A., Loomis, R. A., Burkhardt, A. M. et al. 2021, *Science*, 371, 1265
- Millar, T. J., Walsh, C., Van de Sande, M., & Markwick, A. J. 2024, *A&A*, 682, A109
- Minh, Y. C., Brewer, M. K., Irvine, W. M., et al. 1991, *A&A*, 244, 470
- Morales, S. B., Le Picard, S., Canosa, A., & Sims, I. R. 2010, *Faraday Discuss.*, 147, 155
- Morales, S. B., Bennett, C. J., Le Picard, S. D., et al. 2011, *ApJ*, 742, 26
- Pety, J., Gratier, P., Guzmán, V., et al. 2012, *A&A*, 548, A68
- Pickett, H. M. 1991, *J. Mol. Spectr.*, 148, 371
- Pratap, P., Dickens, J. E., Snell, R. L., et al. 1997, *ApJ*, 486, 862
- Shingledecker, C. N., Lee, K. L. K., Wandishin, J. T., et al. 2021, *A&A*, 652, L12
- Siebert, M. A., Lee, K. L. K., Remijan, A. J., et al. 2022, *ApJ*, 924, 21
- Silva, W. G. D. P., Cernicharo, J., Schlemmer, S., et al. 2023, *A&A*, 676, L1
- Sita, M. L., Changala, P. B., Xue, C., et al. 2022, *ApJ*, 938, L12
- Smith, M. C., Liu, G., Buras, Z. J., et al. 2020, *J. Phys. Chem. A*, 124, 2871
- Sun, B. J., Huang, C. H., Chen, S. Y., et al. 2014, *J. Phys. Chem. A*, 118, 7715
- Tercero, F., López-Pérez, J. A., Gallego, J. D., et al. 2021, *A&A*, 645, A37
- Thaddeus, P., & Turner, B. E. 1975, *ApJ*, 201, L25
- Thomas, A. M., He, C., Zhao, L., et al. 2019, *J. Phys. Chem. A*, 123, 4104
- Trevitt, A. J., Goulay, F., Taatjes, C. A., et al. 2010, *J. Phys. Chem. A*, 114, 1749
- Wenzel, G., Cooke, I. R., Changala, P. B., et al. 2024, *Science*, 386, 810
- Whittet, D. C. B. 2010, *ApJ*, 710, 1009
- Woon, D. E. 2006, *Chem. Phys.*, 331, 67
- Zdanovskaia, M. A., Dorman, P. M., Orr, V. L., et al. 2021a, *J. Am. Chem. Soc.*, 143, 9551
- Zdanovskaia, M. A., Esselman, B. J., Kougias, S. M., et al. 2021b, *Mol. Phys.*, 119, e1964629