

# Resonant ALP-Portal Dark Matter Annihilation as a Solution to the $B^\pm \rightarrow K^\pm \nu \bar{\nu}$ Excess

Kewen Ding,<sup>a</sup> Ying Li,<sup>a,\*</sup> Xuewen Liu,<sup>a,\*</sup> Yu Liu,<sup>b,\*</sup> Chih-Ting Lu,<sup>c,d,\*</sup> Bin Zhu<sup>a,\*</sup>

<sup>a</sup>*Department of Physics, Yantai University, Yantai 264005, China*

<sup>b</sup>*School of Physics, Henan Normal University, Xinxiang 453007, China*

<sup>c</sup>*Department of Physics and Institute of Theoretical Physics, Nanjing Normal University, Nanjing, 210023, China*

<sup>d</sup>*Nanjing Key Laboratory of Particle Physics and Astrophysics, Nanjing, 210023, China*

*E-mail:* [dlsc52000@s.ytu.edu.cn](mailto:dlsc52000@s.ytu.edu.cn), [liying@ytu.edu.cn](mailto:liying@ytu.edu.cn),  
[xuewenliu@ytu.edu.cn](mailto:xuewenliu@ytu.edu.cn), [liuyu2020@s.ytu.edu.cn](mailto:liuyu2020@s.ytu.edu.cn), [ctlun@nju.edu.cn](mailto:ctlun@nju.edu.cn),  
[zhubin@mail.nankai.edu.cn](mailto:zhubin@mail.nankai.edu.cn)

**ABSTRACT:** The Belle II collaboration recently reported a  $2.7\sigma$  excess in the rare decay  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$ , potentially signaling new physics. We propose an axion-like particle (ALP)-portal dark matter (DM) framework to explain this anomaly while satisfying the observed DM relic abundance. By invoking a resonant annihilation mechanism ( $m_a \sim 2m_\chi$ ), we demonstrate that the ALP-mediated interactions between the Standard Model and DM sectors simultaneously account for the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  anomaly and thermal freeze-out dynamics. Two distinct scenarios—long-lived ALPs decaying outside detectors (displaced diphotons) and ALPs decaying invisibly to DM pairs (missing energy)—are examined. While the displaced diphotons scenario is excluded by kaon decay bounds ( $K^\pm \rightarrow \pi^\pm + \text{inv.}$ ), the invisible decay channel remains unconstrained and aligns with Belle II’s missing energy signature. Using the coupled Boltzmann equation formalism, we rigorously incorporate early kinetic decoupling effects, revealing deviations up to a factor of 20 from traditional relic density predictions in resonance regions. For the missing energy scenario, the viable parameter space features ALP-SM and ALP-DM couplings:  $g_{aWW} \in (7.3 \times 10^{-5} - 1.1 \times 10^{-4}) \text{ GeV}^{-1}$  (from  $B^\pm \rightarrow K^\pm a$ ) and  $g_{a\chi\chi} \in (6.1 \times 10^{-5} - 6.0 \times 10^{-3}) \text{ GeV}^{-1}$  (for resonant annihilation), accommodating ALP masses  $m_a \in (0.6, 4.8) \text{ GeV}$ . Therefore, this work establishes the ALP portal as a viable bridge between the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  anomaly and thermal DM production, emphasizing precision calculations of thermal decoupling in resonance regimes.

---

\*Corresponding author

---

## Contents

|          |  |           |
|----------|--|-----------|
| <b>1</b> | <b>Introduction</b>  | <b>1</b>  |
| <b>2</b> | <b>ALP portal DM model and its annihilation mechanisms</b>   | <b>3</b>  |
| 2.1      | ALP portal DM model  | 3         |
| 2.2      | Thermal DM annihilation mechanisms   | 4         |
| <b>3</b> | <b>ALPs from exotic <math>B</math> meson decays and the <math>B^\pm \rightarrow K^\pm \nu \bar{\nu}</math> anomaly at Belle II</b> | <b>5</b>  |
| <b>4</b> | <b>Precision DM relic density in the resonance mechanism</b>   | <b>8</b>  |
| 4.1      | Coupled Boltzmann Equation   | 8         |
| 4.2      | EKD effect in the resonance mechanism  | 10        |
| <b>5</b> | <b>Combined Results</b>  | <b>12</b> |
| 5.1      | Displaced diphoton scenario  | 14        |
| 5.2      | Missing energy scenario  | 15        |
| <b>6</b> | <b>Conclusion</b>  | <b>17</b> |
| <b>A</b> | <b>Cross-sections for DM annihilation and scattering processes</b>   | <b>18</b> |

---

## 1 Introduction

Axion-like particles (ALPs) can be naturally generated through the breaking of a global  $U(1)$  symmetry [1, 2]. ALPs were originally proposed to solve the strong CP problem [3–6]. Later, they became compelling candidates for dark matter (DM) via the misalignment mechanism [7–9]. Additionally, ALPs could influence the dynamics of inflation and reheating [10, 11], as well as the generation of baryon asymmetry through CP-violating processes in the early Universe [12–14]. Moreover, ALPs can act as mediators between the visible and dark sectors, providing a portal to explore the dark sector, often referred to as the ALP portal [15]. Given their potential to address major unresolved problems and bridge various domains of physics from particle physics to cosmology and astrophysics, ALPs have become a highly attractive focus of modern theoretical and experimental research.

In this work, we focus on ALP-portal DM models. Since the couplings of ALPs to a DM pair and to pairs of Standard Model (SM) particles are naturally suppressed by a cutoff scale associated with global  $U(1)$  symmetry breaking, ALP-portal DM models allow for diverse production mechanisms, including the freeze-out [16, 17] and freeze-in [18–20] mechanisms. These mechanisms enable ALP-portal DM models to accommodate a broad range of coupling strengths and DM masses. Notably, when the ALP mass aligns with the

DM annihilation resonance, it significantly enhances the annihilation cross-section [21, 22]. When a pair of DM particles predominantly annihilates to photons, this feature provides a natural explanation for specific astrophysical signals, such as excesses observed in cosmic-ray or gamma-ray data, and allows precise tuning of DM properties [17, 23–26]. Furthermore, ALP-portal DM models exhibit rich experimental phenomenology, including searches in collider [27, 28] and beam-dump [19, 28, 29] experiments.

On the other hand, recently, Belle II reported the first evidence of the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  decay channel with a significance of  $3.5\sigma$  [30]. However, a  $2.7\sigma$  deviation from the SM prediction has also been announced in this process. This deviation may indicate the presence of new physics [31–63], and ALPs, as candidates for physics beyond the SM, could potentially explain the excess reported by Belle II [53, 57]. By further investigating the contributions of ALPs to this decay channel, we can evaluate whether they are responsible for the observed excess.

In the thermal freeze-out framework, several DM annihilation scenarios exist in the ALP-portal model, depending on the mass spectrum between the DM and the ALP, as well as the size of coupling constraints. In this study, we aim to explore the potential connection between the explanation of the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  excess and the DM relic abundance within the ALP-portal DM model. For the DM  $p$ -wave or forbidden annihilation scenario, the ALP predominantly decays into diphotons, and ALPs are required to be long-lived particles (LLPs) that decay outside the detectors, with the signature resulting in missing energy, which can explain this excess. In the DM  $s$ -wave annihilation scenario, despite the strict constraints imposed by the Cosmic Microwave Background and indirect detection experiments, these limitations can be effectively circumvented if DM annihilation during freeze-out is resonantly enhanced [64, 65]. In this case, the ALP can predominantly decay into a pair of DM particles, which directly accounts for this excess. Therefore, the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  excess observed at Belle II could provide a hint for the possible DM annihilation scenario in the early Universe.

Among these DM annihilation scenarios, special care must be taken in calculating the resonant  $s$ -wave annihilation. In this work, we employ the coupled Boltzmann equation (CBE) method, accurately accounting for the resonance mechanism, in which the early kinetic decoupling (EKD) effect has been taken into account. The EKD effect has been extensively studied in the literature [66–80] to calculate the DM relic density. In the context of the Boltzmann equation, it is generally imperative to consider the scattering collision terms for the resonant mechanism. The momentum transfer rate associated with elastic scattering processes, which are crucial for maintaining thermal equilibrium, is significantly suppressed relative to the annihilation rate. This suppression arises due to the small coupling constant required to meet the relic density criterion, absent any enhancement of the scattering rate. Consequently, kinetic equilibrium decouples prematurely and the outcomes derived from the CBE method diverge significantly from those obtained through the standard calculation formulated by Gondolo and Gelmini [81] (denoted as the ‘NBE’ method since this treatment involves solving the Boltzmann equations of number density)). Our investigation reveals that the observed relic abundance of DM can be produced within a precisely defined parameter space region. This region represents a promising avenue for

further exploration at the Belle II experiment, particularly in the context of explaining the excess observed in the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  decay mode.

The structure of the rest of this paper is organized as follows: In Sec. 2, we first review the ALP-portal DM model and its thermal annihilation mechanisms. In Sec. 3, we discuss exotic  $B$  meson decays that produce ALPs and explore their connection to the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  anomaly at Belle II. In Sec. 4, we present precise calculations of the DM relic density in the resonance annihilation scenario. In Sec. 5, we display our combined results for the parameter space that explains the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  excess and yields the correct DM relic density. Finally, we summarize our findings in Sec. 6.

## 2 ALP portal DM model and its annihilation mechanisms

In this section, we begin by reviewing the ALP portal DM model and the effective interactions of ALPs. Next, we discuss the decay patterns of ALPs. Finally, we explore various thermal DM production mechanisms within this model.

### 2.1 ALP portal DM model

For the interactions of a pseudoscalar ALP,  $a$ , with SM gauge bosons, we consider the Lagrangian

$$\mathcal{L}_{\text{EW}} = \frac{1}{2} \partial^\mu a \partial_\mu a - \frac{1}{2} m_a^2 a^2 - \frac{c_B}{4 f_a} a B^{\mu\nu} \tilde{B}_{\mu\nu} - \frac{c_W}{4 f_a} a W^{i,\mu\nu} \tilde{W}_{\mu\nu}^i, \quad (2.1)$$

where  $B^{\mu\nu}$  and  $W^{i,\mu\nu}$  denote the field strength tensors of  $U(1)_Y$  and  $SU(2)_L$  gauge groups before electroweak symmetry breaking (EWSB), respectively, and the dual field strength tensors are defined via  $\tilde{B}_{\mu\nu} = \frac{1}{2} \epsilon_{\mu\nu\rho\sigma} B^{\rho\sigma}$ . The parameters  $c_B, c_W$  are the corresponding Wilson coefficients for ALP-gauge boson interactions.  $m_a$  and  $f_a$  denote the ALP mass and decay constant, which are taken to be independent parameters.

After EWSB, the last two terms in Eq. (2.1) induce four different interactions between the ALP and SM gauge bosons:

$$\mathcal{L}_{\text{EW}} \supset -\frac{g_{a\gamma\gamma}}{4} a F_{\mu\nu} \tilde{F}^{\mu\nu} - \frac{g_{a\gamma Z}}{4} a F_{\mu\nu} \tilde{Z}^{\mu\nu} - \frac{g_{aZZ}}{4} a Z_{\mu\nu} \tilde{Z}^{\mu\nu} - \frac{g_{aWW}}{4} a W_{\mu\nu} \tilde{W}^{\mu\nu}, \quad (2.2)$$

where  $F_{\mu\nu}$ ,  $Z_{\mu\nu}$ , and  $W_{\mu\nu}$  are the field strength tensors of the photon,  $Z$ , and  $W^\pm$  bosons, respectively, and their duals have been defined as above. As pointed out in Ref. [82], the coupling of ALPs to  $W^\pm$  bosons can induce flavor-changing processes such as  $B \rightarrow Ka$  and offer the best sensitivity to ALPs with masses below 5 GeV. Therefore, the two most relevant couplings in this study are those of the ALP to a pair of photons and a pair of  $W$  bosons, which are expressed in terms of  $c_B$  and  $c_W$  [83–87],

$$g_{a\gamma\gamma} = \frac{c_B \cos^2 \theta_W + c_W \sin^2 \theta_W}{f_a}, \quad (2.3)$$


$$g_{aWW} = \frac{c_W}{f_a}, \quad (2.4)$$

where  $\theta_W$  denotes the Weinberg angle. In this study, we choose  $c_B = c_W$  as a benchmark model.

In addition to interactions between the ALP and gauge bosons, we also consider the ALP interacting with the dark sector via the ALP portal [16, 19, 88, 89],

$$\mathcal{L}_{\text{DM}} = \bar{\chi} i \partial_\mu \gamma^\mu \chi - m_\chi \bar{\chi} \chi - g_{a\chi\chi} \partial_\mu a \bar{\chi} \gamma^\mu \gamma^5 \chi \quad (2.5)$$

where  $\chi$  represents the Majorana fermionic dark matter, and  $g_{a\chi\chi}$  is the coupling of the ALP to a pair of DM particles, with dimensions of inverse mass.



**Figure 1.** Branching ratios for two ALP decay channels,  $a \rightarrow \gamma\gamma$  and  $a \rightarrow \chi\bar{\chi}$ . Here we take  $m_a = 1$  GeV,  $m_\chi = 0.475m_a$ ,  $g_{a\gamma\gamma} = 10^{-4}$  GeV $^{-1}$  as a benchmark point and vary the value of  $g_{a\chi\chi}$ .

For an  $\mathcal{O}(1)$  GeV ALP, it predominantly decays into either a pair of photons or a pair of DM particles, depending on the mass relation between the ALP and DM. The partial decay width for the process  $a \rightarrow \gamma\gamma$  is given by:

$$\Gamma_{a\gamma} = \frac{g_{a\gamma\gamma}^2 m_a^3}{64\pi}. \quad (2.6)$$

Additionally, if  $m_a > 2m_\chi$ , the partial decay width for  $a \rightarrow \chi\bar{\chi}$  must also be included:

$$\Gamma_{a\chi} = \frac{g_{a\chi\chi}^2 m_a m_\chi^2}{\pi} \sqrt{1 - \frac{4m_\chi^2}{m_a^2}}. \quad (2.7)$$

To illustrate the interplay between the two ALP decay channels,  $a \rightarrow \gamma\gamma$  and  $a \rightarrow \chi\bar{\chi}$ , for varying  $g_{a\chi\chi}$ , we fix  $m_a = 1$  GeV,  $m_\chi = 0.475m_a$ , and  $g_{a\gamma\gamma} = 10^{-4}$  GeV $^{-1}$  as a benchmark point, and present the results in Fig. 1. It is evident that when  $g_{a\chi\chi} \gtrsim 10^{-4}$  GeV $^{-1}$ , the decay channel  $a \rightarrow \chi\bar{\chi}$  dominates. Otherwise,  $a \rightarrow \gamma\gamma$  becomes the primary decay mode.

## 2.2 Thermal DM annihilation mechanisms

DM annihilations to photons in this model can be dominant by the  $s$ -wave contribution via  $a$ -resonance ( $\chi\bar{\chi} \rightarrow a \rightarrow \gamma\gamma$ ), or  $p$ -wave contribution ( $\chi\bar{\chi} \rightarrow aa$ ). In addition, some

different mass relations between  $\chi$  and  $a$  can alter the velocity dependence of the DM annihilation cross-section. For example, in the  $\chi\bar{\chi} \rightarrow aa$  cases, the main processes involve  $p$ -wave annihilation (velocity-squared suppression) and forbidden annihilation (velocity-exponential suppression). On the other hand, when  $2m_\chi \approx m_a$ , the decay channel  $a \rightarrow \chi\bar{\chi}$  opens and the resonant  $s$ -wave annihilation becomes dominant.

The possible annihilation mechanisms based on the relationships between  $m_\chi$  and  $m_a$  appear as what follows:

1.  $p$ -wave annihilation ( $m_a < m_\chi$ ): The dominant DM annihilation channel is  $\chi\bar{\chi} \rightarrow aa$  which is  $p$ -wave and then each ALP decays into diphoton.
2. forbidden annihilation ( $m_a \gtrsim m_\chi$ ): The annihilation process is also  $\chi\bar{\chi} \rightarrow aa$ , albeit with  $m_\chi \lesssim m_a$ . Essentially, forbidden annihilations cease when the kinetic energies of DM particles fall below threshold energies. Here each ALP also decays into diphoton.
3.  $s$ -wave annihilation ( $m_a > 2m_\chi$  or  $2m_\chi > m_a > m_\chi$ ): The annihilation process is  $\chi\bar{\chi} \rightarrow a \rightarrow \gamma\gamma$ , and both  $g_{a\gamma\gamma}$  and  $g_{a\chi\chi}$  must be sufficiently large to account for the DM relic abundance [16, 17]. Additionally, in this scenario, the ALP predominantly decays into either diphoton or a pair of DM particles, which can be probed in various accelerator [19, 27–29] and indirect detection [17, 23–26] experiments. Consequently, the parameter space for this scenario is already subject to significant constraints.
4. Resonance annihilation ( $m_a \sim 2m_\chi$ ): The annihilation cross-section for  $\chi\bar{\chi} \rightarrow \gamma\gamma$  can experience significant enhancement through an  $a$ -resonance. We focus on the scenario where DM predominantly couples to photons. Strong constraints on this scenario, imposed by the CMB and indirect detection experiments can be circumvented if DM annihilations during freeze-out are resonantly enhanced [64, 65]. In this context, we find that the observed DM relic abundance can be reproduced within a well-defined region of parameter space that is currently consistent with all experimental constraints. However, this parameter space can be explored in the coming years through invisible ALPs searches at the Belle II [88, 90]. To ensure accuracy in our analysis, we employ the coupled Boltzmann equation method [72, 78] to calculate the DM relic density, properly accounting for the resonance mechanism.

### 3 ALPs from exotic $B$ meson decays and the $B^\pm \rightarrow K^\pm \nu\bar{\nu}$ anomaly at Belle II

The ALPs are expected to be produced via flavor-changing processes at  $B$ -factories, specifically focusing on their generation during  $B$  meson decays. We examine the two-body decay process  $B^\pm \rightarrow K^\pm a$ . The partial decay width for this process can be written as [82, 91],

$$\Gamma(B^\pm \rightarrow K^\pm a) = \frac{g_{aWW}^2}{64\pi m_{B^\pm}^3} |C_{sb}|^2 F_K^2(m_a) (m_{B^\pm}^2 - m_{K^\pm}^2)^2 \lambda^{1/2}(m_{B^\pm}^2, m_{K^\pm}^2, m_a^2), \quad (3.1)$$

where  $m_{B^\pm}$ ,  $m_{K^\pm}$  represent the masses of the  $B^\pm$ ,  $K^\pm$  mesons, respectively, while  $|C_{sb}|^2$  denotes the square of the Wilson coefficient  $C_{sb}$ . The Wilson coefficient  $C_{sb}$  is expressed as:

$$C_{sb} = -\frac{3m_W^2}{16\pi^2 v^2} V_{ts}^* V_{tb} f\left(\frac{m_t^2}{m_W^2}\right), \quad (3.2)$$

where  $m_W$  is the mass of the  $W$ -boson,  $v$  is the vacuum expectation value of the Higgs field, and  $V_{ts}^*$  and  $V_{tb}$  are elements of the CKM matrix. The function  $f\left(\frac{m_t^2}{m_W^2}\right)$  depends on the ratio of the top quark mass  $m_t$  to the  $W$ -boson mass  $m_W$ .

We employ established theoretical frameworks to estimate the scalar form factor  $F_K(q)$  for  $K^\pm$ , which characterizes the influence of momentum transfer  $q^2$  on the decay process. This is expressed in the following form [92]:

$$F_K(q^2) = 0.33 \exp\left(\frac{0.63q^2}{m_{B^\pm}^2} - \frac{0.095q^4}{m_{B^\pm}^4} + \frac{0.591q^6}{m_{B^\pm}^6}\right), \quad (3.3)$$

providing a comprehensive description of the kinematic behavior and associated decay widths in the  $B$  meson decay process. Additionally, we introduce two key mathematical functions to represent the decay dynamics and their integrals. These include kinematic function:

$$\lambda(\alpha, \beta, \gamma) = \alpha^2 + \beta^2 + \gamma^2 - 2(\alpha\beta + \alpha\gamma + \beta\gamma), \quad (3.4)$$

and the integral function:

$$f(x) = x[1 + x(\log x - 1)]/(1 - x)^2, \quad (3.5)$$

which facilitate the analysis of the decay mechanisms.

The detection of ALPs in this process exhibits three distinct signatures, depending on the ALP decay length in the laboratory frame: (1) prompt diphoton, (2) displaced diphoton, (3) missing energy. The BaBar collaboration has already utilized the first two signatures, prompt and displaced diphotons, to search for ALPs, particularly in the mass range  $0.175 \text{ GeV} < m_a < 4.78 \text{ GeV}$ , where the coupling constant  $g_{aWW} \sim 10^{-5} \text{ GeV}^{-1}$  [93]. However, due to significant background contamination from  $\eta$  and  $\eta'$  mesons, the mass ranges  $0.45 \text{ GeV} < m_a < 0.63 \text{ GeV}$  and  $0.91 \text{ GeV} < m_a < 1.01 \text{ GeV}$  are excluded from this search. These regions are challenging because the  $\eta$  and  $\eta'$  mesons contribute substantially to the background, making it difficult to distinguish the ALP signal.

To overcome these limitations, a novel search method has been proposed for the Large Hadron Collider (LHC) experiment, which involves using photon-jets as a distinctive signature for exploring ALPs. This method, discussed in detail in Refs. [85, 94], relies on the fact that ALPs can decay into two collimated photons, forming a photon-jet. In high-energy collider environments like the LHC, this phenomenon becomes more prominent. By using this photon-jet feature, it is possible to mitigate background contamination from  $\eta$  and  $\eta'$  mesons in the excluded mass ranges, providing a more efficient experimental approach for ALP searches at the LHC. The combination of the photon-jet analysis at the LHC with existing  $B$ -factory studies could further expand the discovery potential for ALPs and improve constraints on their mass and coupling constants. This approach would enhance our ability to probe ALPs and offer valuable insights into new physics beyond the SM.

We also identify the mass range  $0.1 \text{ GeV} < m_a < 0.175 \text{ GeV}$  as a blind spot for ALP prompt decay to diphoton searches due to significant background contamination from  $\pi^0$  meson decays. This contamination complicates the separation of ALP signals from background noise, making it challenging to detect ALPs in this mass range using current experimental techniques. However, for the long-lived ALP in this blind spot, the ALPs decay length would be sufficiently large that it does not produce any visible decay products within the detector of  $B$  factories. As a result, the ALP would manifest as a missing energy signature. Therefore, we can employ missing energy searches to constrain the potential contribution of ALPs. Notably, this method can be applied to other ALP mass range when light ALPs become very long-lived. Specifically, we propose leveraging the existing  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  process to limit the presence of ALPs indirectly. Since the neutrino pair is electrically neutral and difficult to detect, this decay process appears as missing energy in the detectors. If the ALP decays outside the detector, its signature would resemble the missing energy feature of the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  process. By accurately measuring this process, we can place upper limits on the ALPs contribution.

Additionally, we consider a complementary scenario in which the ALP primarily decays into a pair of DM particles which makes the ALP decay products invisible to the detectors. As discussed above, distinguishing this scenario from the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  process in the SM is challenging. Therefore, precision measurements of the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  process can also place upper limits on the ALP contribution in this scenario.

Notably, the Belle II experiment has searched for the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  decay and observed a  $2.7\sigma$  deviation from the SM prediction of  $\text{BR}(B^\pm \rightarrow K^\pm \nu \bar{\nu}) = (5.58 \pm 0.37) \times 10^{-6}$  [95]. In contrast, the measured branching ratio is  $\text{BR}(B^\pm \rightarrow K^\pm \nu \bar{\nu}) = [2.3 \pm 0.5(\text{stat})_{-0.4}^{+0.5}(\text{syst})] \times 10^{-5}$  [30]. Although it is premature to assert whether this excess results from statistical fluctuations or is a genuine indication of new physics, it does represent the first evidence of the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  process with a significance of  $3.5\sigma$ . Therefore, further accumulation of events and improvements in measurements are urgently needed. It is also valuable to consider possible new physics explanations for this observed anomaly while awaiting the next update on the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  process. In fact, several proposals have already been put forward to explain this excess, including new  $U(1)$  extensions of the SM [31, 54], the SMEFT framework [33, 41, 43, 47, 48], light dark matter models [34, 35, 38–40, 45, 59, 60], ALPs [53, 57], and other possibilities [36, 37, 42, 44, 50, 52, 55, 56, 61–63].

In this study, we aim to use the ALP-portal DM model to explain the observed  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  excess from the Belle II experiment. As discussed in Sec. 2, ALPs predominantly decay into diphotons in both the DM  $p$ -wave annihilation and forbidden annihilation scenarios. In these scenarios, ALPs are required to be light, and the coupling  $g_{a\gamma\gamma}$  must be small enough to ensure that the ALP lifetime is long enough for them to travel outside the detectors, with the final state resulting in missing energy. Alternatively, in the DM  $s$ -wave annihilation scenario, ALPs can decay into diphotons and a pair of DM particles, as shown in Fig. 1, depending on the relative magnitudes of  $g_{a\gamma\gamma}$  and  $g_{a\chi\chi}$ . When the invisible decay channel of ALPs dominates, the lifetime of ALPs becomes less of a concern, as the final state already results in missing energy. By further investigating the contribution of ALPs to the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  excess in Sec. 5, we can assess whether they could be the source of this

observed excess.

## 4 Precision DM relic density in the resonance mechanism

As extensively discussed in the literature, the freeze-out dynamics of DM exhibit significant complexity within the resonance region of annihilation [66–80]. In this section, we provide a comprehensive review of the coupled Boltzmann equations, which play a crucial role in accurately calculating the relic density. Additionally, we explore in detail the early kinetic decoupling (EKD) effect that emerges in the resonant annihilation.

### 4.1 Coupled Boltzmann Equation

Keeping kinetic equilibrium during and even after the freeze-out period is a key assumption for standard relic density calculations. Nevertheless, this assumption is not invariably valid, especially in contexts where kinetic decoupling occurs prior to chemical decoupling. To investigate the DM relic density considering the impact of early kinetic decoupling, we should focus on the following Boltzmann equation for DM phase-space distribution [67, 71]:

$$E \left( \frac{\partial}{\partial t} - H \vec{p} \cdot \frac{\partial}{\partial \vec{p}} \right) f_\chi(t, \vec{p}) = C_{\text{ann.}}[f_\chi] + C_{\text{el.}}[f_\chi], \quad (4.1)$$

where  $E$  is the energy of the DM,  $H$  is the Hubble constant,  $\vec{p}$  is the momentum of DM, and  $f_\chi$  is the DM phase-space density. The collision terms,  $C_{\text{ann.}}$  and  $C_{\text{el.}}$ , account for the annihilation of DM particles into particles of the thermal bath and the elastic scattering processes between DM and SM particles, respectively. For two-body processes,

$$\begin{aligned} C_{\text{ann.}} &= \frac{1}{2g_\chi} \sum \int \frac{d^3 p'}{(2\pi)^3 2E_{p'}} \int \frac{d^3 k}{(2\pi)^3 2E_k} \int \frac{d^3 k'}{(2\pi)^3 2E_{k'}} (2\pi)^4 \delta^4(p + p' - k - k') \\ &\times \left( -|\mathcal{M}_{\chi\chi \rightarrow \mathcal{B}\mathcal{B}'}|^2 f_\chi(\vec{p}) f_\chi(\vec{p}') (1 \pm f_{\mathcal{B}}^{\text{eq}}(\vec{k})) (1 \pm f_{\mathcal{B}'}^{\text{eq}}(\vec{k}')) \right. \\ &\left. + |\mathcal{M}_{\mathcal{B}\mathcal{B}' \rightarrow \chi\chi}|^2 f_{\mathcal{B}}^{\text{eq}}(\vec{k}) f_{\mathcal{B}'}^{\text{eq}}(\vec{k}') (1 \pm f_\chi(\vec{p})) (1 \pm f_\chi(\vec{p}')) \right), \end{aligned} \quad (4.2)$$

$$\begin{aligned} C_{\text{el.}} &= \frac{1}{2g_\chi} \sum \int \frac{d^3 p'}{(2\pi)^3 2E_{p'}} \int \frac{d^3 k}{(2\pi)^3 2E_k} \int \frac{d^3 k'}{(2\pi)^3 2E_{k'}} (2\pi)^4 \delta^4(p + p' - k - k') \\ &\times \left( -|\mathcal{M}_{\chi\mathcal{B} \rightarrow \chi\mathcal{B}}|^2 f_\chi(\vec{p}) f_{\mathcal{B}}^{\text{eq}}(\vec{k}) (1 \pm f_\chi(\vec{p}')) (1 \pm f_{\mathcal{B}}^{\text{eq}}(\vec{k}')) \right. \\ &\left. + |\mathcal{M}_{\chi\mathcal{B}' \rightarrow \chi\mathcal{B}}|^2 f_\chi(\vec{p}') f_{\mathcal{B}'}^{\text{eq}}(\vec{k}') (1 \pm f_\chi(\vec{p})) (1 \pm f_{\mathcal{B}'}^{\text{eq}}(\vec{k}')) \right), \end{aligned} \quad (4.3)$$

where  $\mathcal{B}$  and  $\mathcal{B}'$  stand for particles in the thermal bath such as ALPs and photons,  $g_\chi$  is the number of internal degrees of freedom of DM, and  $f_{\mathcal{B}}^{\text{eq}}$  is given by the Fermi-Dirac or Bose-Einstein distribution depending on the spin of  $\mathcal{B}$ . The summation should be taken for all the internal degrees of freedom for all the particles. For the nonrelativistic DM,  $C_{\text{el.}}$  can be simplified as the Fokker-Planck operator [96–99]:

$$C_{\text{el.}} \simeq \frac{E}{2} \gamma(T) \left[ TE \partial_p^2 + \left( \frac{2TE}{p} + p + \frac{Tp}{E} \right) \partial_p + 3 \right] f_\chi. \quad (4.4)$$

where  $T$  is the temperature of the thermal bath and  $\partial_p \equiv \partial/\partial p$ . In the above equation, the momentum transfer rate  $\gamma(T)$  is given by (see also Ref. [100])

$$\gamma = \frac{1}{3g_\chi m_\chi T} \int \frac{d^3k}{(2\pi)^3} f_{\mathcal{B}}^\pm(E_k) [1 \mp f_{\mathcal{B}}^\pm(E_k)] \int_{-4k_{\text{cm}}^2}^0 dt(-t) \frac{d\sigma}{dt} v, \quad (4.5)$$

where  $f_{\mathcal{B}}^\pm(E_k)$  is Fermi-Dirac or Bose-Einstein distribution of bath particle,  $v$  is the relative velocity and  $t$  is the Mandelstam variable. The differential cross section can be expressed as  $(d\sigma/dt)v \equiv |\mathcal{M}|_{\chi\mathcal{B} \leftrightarrow \chi\mathcal{B}}^2 / (64\pi k E_k m_\chi^2)$ , and  $k_{\text{cm}}^2$  is given by

$$k_{\text{cm}}^2 = \frac{m_\chi^2(E_k^2 - m_{\mathcal{B}}^2)}{m_\chi^2 + m_{\mathcal{B}}^2 + 2m_\chi E_k}. \quad (4.6)$$

Here  $E_k$  is the energy of heat bath particle  $\mathcal{B}$ . Note that  $k_{\text{cm}}^2 \neq E_k^2 - m_{\mathcal{B}}^2 = |\vec{k}|^2$ . The integrals of the amplitudes of the elastic scattering processes are presented in Appendix A.

During the chemical decoupling, the scattering processes may not be frequent enough to maintain the kinetic equilibrium, which means that DM particles own a different temperature  $T_\chi$  from the thermal plasma in their following evolution. A common definition of the DM temperature is

$$T_\chi(T) = \frac{g_\chi}{3n_\chi} \int \frac{d^3p}{(2\pi)^3} \frac{\vec{p}^2}{E} f_\chi(\vec{p}) \equiv \frac{s^{2/3}}{m_\chi} y, \quad (4.7)$$

which is also a function of the thermal bath temperature  $T$ . In this definition  $n_\chi$  is the number density of the DM, and  $s$  is the entropy density. Here  $y$  is a dimensionless version in analogy to the DM yield  $Y (= n_\chi/s)$ .

To reach a suitable description of the DM temperature evolution and then explore the early kinetic decoupling (EKD) effect on the chemical decoupling process, we should consider the second moment of  $f_\chi$  as a dynamical degree of freedom. By integrating Eq. (4.1) with  $g_\chi \int \frac{d^3p}{(2\pi)^3} \frac{1}{E}$  and  $g_\chi \int \frac{d^3p}{(2\pi)^3} \frac{1}{E} \frac{\vec{p}^2}{E^2}$ , one obtains the zeroth and second moments of the Boltzmann equation, respectively. This leads to a relatively simple coupled system of Boltzmann differential equations,

$$\frac{Y'}{Y} = \frac{sY}{x\tilde{H}} \left[ \frac{Y_{\text{eq}}^2}{Y^2} \langle \sigma v \rangle_T - \langle \sigma v \rangle_{T_\chi} \right], \quad (4.8)$$

$$\begin{aligned} \frac{y'}{y} &= \frac{1}{x\tilde{H}} \langle C_{\text{el}} \rangle_2 + \frac{sY}{x\tilde{H}} \left[ \langle \sigma v \rangle_{T_\chi} - \langle \sigma v \rangle_{2,T_\chi} \right] \\ &+ \frac{sY}{x\tilde{H}} \frac{Y_{\text{eq}}^2}{Y^2} \left[ \frac{y_{\text{eq}}}{y} \langle \sigma v \rangle_{2,T} - \langle \sigma v \rangle_T \right] + 2(1-w) \frac{H}{x\tilde{H}}, \end{aligned} \quad (4.9)$$

where  $x$  is defined as usual  $x = m_\chi/T$ ,  $Y_{\text{eq}}(x) \equiv n_{\text{eq}}(T)/s$ , and

$$\tilde{H} \equiv H / [1 + (1/3)d(\log g_{\text{eff}}^s)/d(\log T)], \quad (4.10)$$

with  $g_{\text{eff}}^s$  being the entropy degrees of freedom of the background plasma.  $w(T_\chi) \equiv 1 - \langle p^4/E^3 \rangle_{T_\chi} / (6T_\chi)$ , with  $\langle p^4/E^3 \rangle = \frac{g_\chi}{n_\chi^{\text{eq}}(T_\chi)} \int \frac{d^3p}{(2\pi)^3} (\vec{p} \cdot \vec{p})^2 / E^3 e^{-\frac{E}{T_\chi}}$ . Note that the elastic

scattering term given in Eq. (4.4) does not contribute to the zeroth moment term. This is a natural consequence because the elastic scattering processes do not change the number density of DM.

The above compact form of the differential equations contains the following thermally averaged cross sections,

$$\langle C_{\text{el}} \rangle_2 \equiv \frac{g_\chi}{3nT_\chi} \int \frac{d^3p}{(2\pi)^3} \frac{p^2}{E^2} C_{\text{el}}, \quad (4.11)$$

$$\langle \sigma v \rangle_{T_\chi} \equiv \frac{g_\chi^2}{(n_\chi^{\text{eq}})^2} \int \frac{d^3p}{(2\pi)^3} \int \frac{d^3q}{(2\pi)^3} (\sigma v)_{\chi\chi \rightarrow BB'} f_\chi^{\text{eq}}(\vec{p}, T_\chi) f_\chi^{\text{eq}}(\vec{q}, T_\chi). \quad (4.12)$$


The thermal average  $\langle \sigma v \rangle_{2,T}$  is a variant of the commonly used thermal average  $\langle \sigma v \rangle_T$ , and is explicitly stated in Ref. [67] and introduced as

$$\langle \sigma v \rangle_{2,T_\chi} = \frac{g_\chi^2}{(n_\chi^{\text{eq}})^2 T_\chi} \int \frac{d^3p}{(2\pi)^3} \int \frac{d^3q}{(2\pi)^3} \frac{\vec{p} \cdot \vec{q}}{3E} (\sigma v)_{\chi\chi \rightarrow BB'} f_\chi^{\text{eq}}(\vec{p}, T_\chi) f_\chi^{\text{eq}}(\vec{q}, T_\chi). \quad (4.13)$$

For  $\langle \sigma v \rangle_T$  and  $\langle \sigma v \rangle_{2,T}$ , we replace  $T_\chi$  by  $T$  in  $\langle \sigma v \rangle_{T_\chi}$  and  $\langle \sigma v \rangle_{2,T_\chi}$ , respectively. Moreover,  $n_\chi^{\text{eq}}(T_\chi)$  is given by


$$n_\chi^{\text{eq}}(T_\chi) = g_\chi \int \frac{d^3p}{(2\pi)^3} f_\chi^{\text{eq}}(\vec{p}, T_\chi) = g_\chi \int \frac{d^3p}{(2\pi)^3} e^{-\frac{E_p}{T_\chi}}. \quad (4.14)$$

## 4.2 EKD effect in the resonance mechanism



**Figure 2.** The required value of the coupling  $g_{a\gamma\gamma}$  to achieve the correct relic density is shown in the figure. The red dashed line represents the results using the NBE method, which relies on the assumption of kinetic equilibrium throughout the entire freeze-out epoch. The blue line corresponds to the results derived from the CBE method. For illustration, we adopt  $m_a = 1$  GeV and  $g_{a\chi\chi} = 10^{-2}$  GeV $^{-1}$ .

Initially, we compute the relic density using both the standard treatment and the CBE treatment. We numerically solve Eqs. (4.8) and (4.9) for a given set of parameters  $(m_\chi, g_{a\gamma\gamma})$  and determine the resulting asymptotic value of  $Y_0$ . For illustration, we fix  $m_a = 1$  GeV and  $g_{a\chi\chi} = 10^{-2}$  GeV $^{-1}$ . The viable parameter space is determined by the matching the observed relic density reported by the Planck Collaboration, which is  $\Omega h^2 = 0.120 \pm 0.001$  [101]. The red dashed line in Fig. 2, corresponding to the NBE method, shows the contour in this plane that yields the correct relic density. In the same figure, we also display the required value of  $g_{a\gamma\gamma}$  obtained by solving the coupled system of Boltzmann equations, represented by the blue solid curve. Outside the resonance region, the coupled Boltzmann equations produce results identical to those of the standard approach, indicating that kinetic decoupling occurs much later than chemical decoupling and that the assumption of local thermal equilibrium during chemical freeze-out is satisfied. However, for DM masses within the resonance region, the CBE method yields significantly different results, suggesting that this assumption is violated. In the resonance region, the momentum transfer rate  $\gamma(T)$  is strongly suppressed compared to the annihilation rate. This suppression arises because the small coupling  $g_{a\gamma\gamma}$  required to satisfy the relic density condition does not benefit from a corresponding resonant enhancement of  $\gamma(T)$ , leading to the early decoupling of kinetic equilibrium.



**Figure 3.** The relic density ratio between CBE and NBE methods, where we require the parameters to match the observed abundance in the NBE approach, i.e.  $\Omega_{\text{NBE}} h^2 = 0.12$ . Here we set  $m_a = 1$  GeV.

To highlight the significance of the improved treatment of decoupling dynamics near the resonance region, we plot in Fig. 3 the ratio of the relic density obtained from the CBE method to that from the standard NBE approach. The parameters used here satisfy the requirement  $\Omega_{\text{NBE}} h^2 = 0.12$ . As shown in the figure, the CBE results differ significantly from the NBE results for the same parameters. First, as demonstrated by the benchmark points with  $r < 0.37$  and  $r > 0.55$ , for DM masses sufficiently far from the resonance, the phase-space distribution remains nearly Maxwellian in shape. For these cases, we observe,

as expected, excellent agreement in the evolution of  $Y(x)$  and  $y(x)$  between the CBE and the NBE approaches. However, in the resonance region, the ratio  $\Omega_{\text{CBE}}/\Omega_{\text{NBE}}$  exhibits a complex structure. For instance,  $\Omega_{\text{CBE}}/\Omega_{\text{NBE}} \simeq 20$  when  $r \simeq 0.41$ . This behavior can be understood by dividing the region into three distinct parts:

- Sub-resonance region ( $0.37 \lesssim r \lesssim 0.48$ )

EKD happens in this region, the DM phase-space density  $f_\chi$  begins to deviate from the equilibrium value of the thermal bath, requiring higher momenta to reach the resonance.

Consider an illustrative case where  $r = 0.45$ . In this scenario, resonant annihilation depletes  $f_\chi$  for momenta above the peak of the distribution [67]. This leads to a relative reduction in  $f_\chi$  compared to a thermal distribution at these momenta, thereby decreasing the DM velocity dispersion, often referred to as the “temperature”. This effect is clearly visible in Fig. 4, where the evolution of  $y_\chi$  shows a decline, as indicated by the blue curve.

Moreover, this behavior can be directly understood from Eq. (4.9). Specifically, in the sub-resonant regime, we find that  $\langle\sigma v\rangle_{T_\chi} < \langle\sigma v\rangle_{2,T_\chi}$ . This drives  $y_\chi$  to smaller values after decoupling. Additionally, the depletion in  $f_\chi$  reduces  $\langle\sigma v\rangle_{T_\chi}$ , leading to an earlier chemical decoupling. As a result, the relic density increases, as demonstrated in the left panel of Fig. 5.

- Resonance region ( $0.48 \lesssim r \lesssim 0.5$ )

When entering the resonant region, DM annihilation begins to deplete  $f_\chi$  below the peak of the would-be Maxwellian distribution. As a result, once equilibrium is no longer maintained (i.e. EKD), the depletion causes an increase in the velocity dispersion, in contrast to the decrease observed in the sub-resonant regime. This phenomenon is clearly illustrated by the orange and green curves in Fig. 4.


At the exact resonance point, the residual DM particles with higher velocities contribute to an enhanced cross section  $\langle\sigma v\rangle$ . This increase in  $\langle\sigma v\rangle$  reduces the DM abundance, as shown in the right panel of Fig. 5. This mechanism explains the formation of the narrow dip observed near the resonance position, as depicted in Fig. 3.

- Super-resonance region ( $0.5 \lesssim r \lesssim 0.55$ )


In the super-resonant regime, characterized by  $r \gtrsim 0.5$ , the condition  $\sqrt{s} \gtrsim m_a$  is inherently satisfied. Consequently, a resonantly enhanced annihilation rate can only occur within a highly confined region of phase space, where the relative DM momenta are nearly zero. This restriction implies that the influence of the resonance diminishes rapidly, rendering its effect practically negligible in this regime.

## 5 Combined Results

The determination of the DM relic density utilizing the CBE method enables a precise delineation of the parameter space, as discussed in the previous section. This parameter



**Figure 4.** The evolution of the DM ‘temperature’  $y_\chi$  with respect to  $x = m_\chi/T$ . The gray line represents the temperature of the thermal bath. The blue, orange, and green curves correspond to  $r \equiv m_\chi/m_a = 0.45, 0.49$ , and  $0.5$ , respectively. Here we take  $m_a = 1\text{GeV}$  for illustration.



**Figure 5.** The evolution of DM abundance  $Y_\chi$  with respect to  $x = m_\chi/T$ . Left panel: The blue lines depict the DM temperature evolution in the CBE approach, with benchmarks labeled in the plot. The gray dashed line represents the NBE approach. Right panel: The red line shows the DM yields in the CBE approach for different parameter settings. The gray dashed line corresponds to the yield in the NBE approach. The line styles follow the same convention as in the left panel. We observe significantly different evolutionary trajectories for varying mass ratios  $r \equiv m_\chi/m_a$ .

space represents a promising regime for further exploration, particularly in light of the observed excess in the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  decay mode. In this section, we will present the combined results that simultaneously account for the observed DM relic abundance and the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  excess.

### 5.1 Displaced diphoton scenario

As discussed in Sec. 2, ALPs predominantly undergo decay into diphotons in various scenarios, including DM  $p$ -wave annihilation, forbidden annihilation, and  $s$ -wave annihilation with certain coupling configurations (refer to Fig. 1). In these scenarios, ALPs are necessitated to possess low masses, and the coupling  $g_{a\gamma\gamma}$  must be sufficiently small to guarantee an ALP lifetime that permits them to traverse beyond detectors at the Belle II. This results in a signature characterized by missing energy.

To explain the observed  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  excess at the Belle II, we should determine the number of signal events that can generate invisible ALP signature at the detector for this scenario. This can be expressed as follows:

$$N(B^\pm \rightarrow K^\pm \cancel{E}) = N_B \times Br(B^\pm \rightarrow K^\pm a) \times Br(a \rightarrow \gamma\gamma) \times P_a(R). \quad (5.1)$$

Here,  $N_B$  denotes the number of B mesons produced at the Belle II. The probability that an ALP is detected at a distance  $R$  from the interaction point is given by


$$P_a(R) = \exp\left(-\frac{R}{L_a}\right), \quad L_a = \gamma\beta c\tau_a, \quad (5.2)$$

where  $\gamma$  represents the Lorentz factor,  $\beta$  denotes the velocity of the ALP, and  $\tau_a \simeq 1/\Gamma_{a \rightarrow \gamma\gamma}$  denotes the proper decay time of the ALP. In this scenario,  $m_a \ll m_{B^\pm}$ , and therefore the boost factor can be estimated as  $\gamma\beta \simeq m_{B^\pm}/(2m_a)$ .

The structure and geometry sizes of detectors are crucial in determining whether the signature of an ALP is a displaced diphoton or missing energy in Eq. (5.1). At the Belle II detector, for  $R > 150$  cm in radius from the interaction point, only the  $K_L$ -Muon (KLM) detector is covered [102]. The KLM detector is designed to detect  $K_L^0$  mesons, neutrons, and muons, and has negligible efficiency for detecting a photon pair [103]. Therefore, we use  $R > 150$  cm in Eq. (5.2) as a simple criterion for the long-lived ALP to be invisible in the Belle II experiment.

To estimate the bounds for the blind zone of  $100 \text{ MeV} < m_a < 175 \text{ MeV}$  and for even lighter ALPs, we employ Eq. (5.1) in conjunction with the Belle II (2023) measurement,  $Br(B^\pm \rightarrow K^\pm \nu \bar{\nu}) = (2.3 \pm 0.7) \times 10^{-5}$  [30]. The resulting bounds are illustrated as the blue bulk in Fig. 6. The shape of this new bound is influenced by the ALP production rate and its decay length in the lab frame. As the parameters  $g_{aWW}$  and  $m_a$  decrease, the decay length of the ALP in the lab frame increases significantly, but the production rate from  $B^\pm \rightarrow K^\pm a$  decreases considerably.

For further investigation, we incorporate existing constraints such as those derived from the  $K^\pm \rightarrow \pi^\pm + X_{\text{inv}}$  search conducted by the NA62 experiment [104] (represented by the green area) and the  $\pi^0 \rightarrow X_{\text{inv}}$  search by the same experiment [105] (depicted as the light blue region). Additionally, we present constraints stemming from searches of  $K_L \rightarrow \pi^0 a$  and



**Figure 6.** Signal region and experimental constraints in the ALP parameter space, assuming ALP decays into displaced diphotons. We apply the Belle II search for  $B \rightarrow K\nu\bar{\nu}$  [30] to constrain the decay  $B \rightarrow Ka$  (blue region). The parameter regions excluded by existing bounds:  $K^\pm \rightarrow \pi^\pm X_{inv}$  from NA62 experiment [104] (green region),  $\pi^0 \rightarrow X_{inv}$  from NA62 experiment [105] (light blue region), the LEP experiment [106] (upper gray region), and the searches of the long-lived particle decaying into a pair of photons by the FASER experiment [107] (gray region). Also the constraints from  $K_L \rightarrow \pi^0 a$ ,  $K^\pm \rightarrow \pi^\pm a$ , and other Beam Dump experiments [82] are included. The solid red line represents the required value of the coupling  $g_{a\gamma\gamma}$  to achieve the correct relic density in the CBE approach, while the dashed red line represents the corresponding value in the NBE approach. Here, we take  $r = 0.47$  and  $g_{a\chi\chi} = 10^{-3} \text{ GeV}^{-1}$  as a benchmark point.


$K^\pm \rightarrow \pi^\pm a$  processes, the LEP experiment [106], and various Beam Dump experiments [82]. Furthermore, we include the constraint imposed by the FASER experiment’s search for long-lived particles decaying into a pair of photons [107] (indicated by the gray area). Notably, our newly delineated parameter space in this specific scenario completely overlaps with the exclusion limits, suggesting that the scenario involving displaced diphoton final states of ALPs is unable to account for the observed excess.

Since we have set  $g_{a\gamma\gamma} = g_{aWW}$ , we are able to present the parameter space corresponding to the DM relic density within the same plot. We depict this parameter space using a solid (dashed) red curve to signify the results derived from the CBE (NBE) approach. For the purpose of illustration, we adopt  $r = 0.47$  and  $g_{a\chi\chi} = 10^{-3} \text{ GeV}^{-1}$  as our benchmark values. It is evident that the CBE results significantly diverge from those of the NBE, highlighting the need for a more precise treatment in our analysis.

## 5.2 Missing energy scenario

When the invisible decay channel of ALPs dominates in the DM  $s$ -wave annihilation scenario, the lifetime of ALPs becomes less of a concern, which directly accounts for this excess. The number of signal events can be expressed as follows:

$$N(B^\pm \rightarrow K^\pm \cancel{E}) = N_B \times Br(B^\pm \rightarrow K^\pm a) \times Br(a \rightarrow \text{inv.}). \quad (5.3)$$



**Figure 7.** Signal region and experimental constraints in the ALP parameter space assuming ALP decays to a pair of DM particles. The blue bulk represents the parameter space for  $B \rightarrow Ka$  determined by the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  measurement from Belle II [30]. We show existing constraints from SN 1987A [88] (gray region), BaBar  $\gamma + inv.$  ( $53 \text{ fb}^{-1}$ ) (green) searches [108]. The constraints are also derived from limits on ALP production in the decay of  $B \rightarrow Ka$  by BaBar data [82] (orange). Additionally, the expected sensitivity from Belle II  $\gamma + inv.$  ( $20 \text{ fb}^{-1}$ ) data [88] is shown by black dotted curve. The solid and dashed red (and light-red) curves have the same meaning as in Fig. 6.

Using the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  measurement from Belle II [30], we are able to ascertain the feasible parameter region for the decay of an ALP into a pair of DM particles, within the mass range where  $m_a$  is less than  $m_{B^\pm} - m_{K^\pm}$ . The outcomes of this analysis are represented in Fig. 7, highlighted in the blue region. Our findings indicate that the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  excess prefers a coupling strength  $g_{aWW}$  within the interval of  $(7.3 \times 10^{-5} \sim 1.1 \times 10^{-4}) \text{ GeV}^{-1}$  across the majority of the  $m_a$  spectrum.

For comparative analysis, the plot also incorporates existing constraints, namely those derived from SN 1987A [88], BaBar  $\gamma + inv.$  (with  $53 \text{ fb}^{-1}$ ) searches [108]. Notably, the constraint imposed by SN 1987A excludes the parameter space where  $m_a \lesssim 0.6 \text{ GeV}$ . BaBar searches of  $B \rightarrow K \nu \bar{\nu}$  [109] put strict constraints on the parameter space of ALP production by process  $B \rightarrow Ka$  as shown in the orange region [82]. Consequently, there remains limited room to accommodate the excess events. Of particular interest, we also include the expected sensitivity from Belle II searches of final state with  $\gamma + inv.$  ( $20 \text{ fb}^{-1}$ ) [88], showing that the most parameter space will be tested.

Regarding the DM relic density in this scenario, the parameter space is already subject to stringent constraints from CMB observations and indirect detection experiments. As a result, the DM production mechanism is strongly influenced by the resonance effects. Fig. 7 displays six curves (shown in (light-) red): the solid curves derived using the CBE method and the dashed ones using the NBE method, both of which satisfy the DM relic density criterion. For clarity, specific parameter combinations are selected:  $(r, g_{a\chi\chi}) =$

(0.475,  $10^{-3}$  GeV $^{-1}$ ), (0.477,  $10^{-3}$  GeV $^{-1}$ ), and (0.490,  $1.04 \times 10^{-5}$  GeV $^{-1}$ ). Notably, the CBE curves align closely with the blue region over a substantial range. In contrast, the NBE results deviate significantly from the CBE curves and fail to intersect with the viable region delineated by the Belle II measurement within this framework. Consequently, we conclude that the CBE method yields distinct and more accurate results compared to the traditional approach.

| BP | $m_a$ (GeV) | $g_{a\gamma\gamma}$ (GeV $^{-1}$ )  | $g_{a\chi\chi}$ (GeV $^{-1}$ ) |
|----|-------------|-------------------------------------|--------------------------------|
| 1  | 0.75        | Lower Bound: $7.398 \times 10^{-5}$ | $5.961 \times 10^{-3}$         |
|    |             | Upper Bound: $9.037 \times 10^{-5}$ | $6.930 \times 10^{-5}$         |
| 2  | 2.63        | Lower Bound: $7.345 \times 10^{-5}$ | $2.295 \times 10^{-3}$         |
|    |             | Upper Bound: $8.810 \times 10^{-5}$ | $1.280 \times 10^{-4}$         |
| 3  | 3.56        | Lower Bound: $7.360 \times 10^{-5}$ | $2.120 \times 10^{-3}$         |
|    |             | Upper Bound: $9.750 \times 10^{-5}$ | $8.400 \times 10^{-5}$         |
| 4  | 4.47        | Lower Bound: $8.429 \times 10^{-5}$ | $8.565 \times 10^{-4}$         |
|    |             | Upper Bound: $1.148 \times 10^{-4}$ | $6.100 \times 10^{-5}$         |

**Table 1.** Viable range of coupling  $g_{a\chi\chi}$  for the selected benchmark points (BPs). The table presents the ALP mass points with the allowed range of  $g_{a\gamma\gamma}$  selected as benchmarks from the four remaining mass intervals after the signal region is excluded by experimental limits. The lower and upper bounds of  $g_{a\gamma\gamma}$  are derived from the signal and experimental exclusion regions in Fig. 7. The corresponding values of  $g_{a\chi\chi}$  are computed where the relic density of DM is equal to 0.12 using the CBE method. Here  $r = 0.477$  is chosen.

To better interpret the influence of the coupling  $g_{a\chi\chi}$ , we select ALP mass points from the four remaining mass intervals that have not been ruled out by experimental limits within the signal region as benchmarks, as illustrated in Table 1. For each of these regions, the combination of signal region and experimental exclusions yields both a lower and an upper bound for  $g_{a\gamma\gamma}$ . We then use the CBE method to derive the corresponding values of  $g_{a\chi\chi}$  that ensure the DM has the observed relic density.

## 6 Conclusion

The  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  anomaly observed at Belle II provides a compelling opportunity to explore ALP-mediated dark matter scenarios. Within the resonant annihilation framework ( $m_a \sim 2m_\chi$ ), we demonstrate that ALP couplings to both the Standard Model ( $g_{aWW}(g_{a\gamma\gamma})$ ) and dark matter ( $g_{a\chi\chi}$ ) can simultaneously explain the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  excess and satisfy DM relic density constraints. Key advancements in this work include:

1. Distinct ALP decay signatures: Invisible decays ( $a \rightarrow \chi\bar{\chi}$ ) dominate for  $g_{a\chi\chi} \geq 10^{-4}$  GeV $^{-1}$ , producing missing energy signals consistent with the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  excess at the Belle II, while evading SN 1987A and BaBar  $B \rightarrow K \nu \bar{\nu}$  constraints for  $m_a > 0.6$  GeV. Notably, this parameter space can be further explored in the coming years through invisible ALPs searches at the Belle II. In contrast, the displaced diphoton

scenario for explaining the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  excess is ruled out by the  $K^\pm \rightarrow \pi^\pm a$  searches.

2. Resonant enhancement and early kinetic decoupling: Utilizing the CBE method, we find the discrepancies between traditional relic density calculations. In resonant regions ( $r = m_\chi/m_a \approx 0.4 - 0.5$ ), EKD effects affect the freeze-out process, amplifying relic densities by up to 20 times compared to standard treatments.
3. Viable parameter space: The interplay between  $g_{aWW} \in (7.3 \times 10^{-5} - 1.1 \times 10^{-4}) \text{ GeV}^{-1}$  and  $g_{a\chi\chi} \in (6.1 \times 10^{-5} - 6.0 \times 10^{-3}) \text{ GeV}^{-1}$  defines a testable regime for the not excluded ALP mass intervals in the range of  $m_a \in (0.6 - 4.8) \text{ GeV}$  that can simultaneously explain the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  excess at the Belle II and the observed dark matter relic abundance.

This work establishes the ALP portal as a robust framework for unifying the  $B^\pm \rightarrow K^\pm \nu \bar{\nu}$  anomaly and thermal dark matter production. Further studies (e.g., B-factories, beam-dump experiments, and astrophysical phenomenology) will provide deeper insights into this scenario.

## Acknowledgments

We are grateful to Shu-Yuan Guo for the valuable discussions. The work by X.L. is supported by the Project of Shandong Province Higher Educational Science and Technology Program under Grants No. 2022KJ271. Y.L. is supported in part by the National Natural Science Foundation of China (NNSFC) under the Grants No. 11925506, 12375089, 12435004, and the Natural Science Foundation of Shandong province under the Grant No. ZR2022ZD26. C.T.L. is supported by NNSFC under grant No. 12335005 and the Special funds for postdoctoral overseas recruitment, Ministry of Education of China. B.Z. is supported by NNSFC under grant Nos. 12275232

## A Cross-sections for DM annihilation and scattering processes

The cross-sections for DM annihilation process is

$$\sigma_{\chi\chi \rightarrow \gamma\gamma\nu} = \frac{g_{a\chi\chi}^2 g_{a\gamma\gamma}^2 m_\chi^2 s^{\frac{5}{2}} \sqrt{s(s-4m_\chi^2)}}{2\pi \sqrt{s-4m_\chi^2} (s-2m_\chi^2) (m_a^2 \Gamma_a^2 + (m_a^2 - s)^2)}, \quad (\text{A.1})$$

where  $s = E_{cm}^2$  is the square of the center-of-mass energy for DM annihilation. The decay width of ALP express as

$$\Gamma_a(m_a > 2m_\chi) = \frac{g_{a\chi\chi}^2 m_\chi^2 \sqrt{m_a^2 - 4m_\chi^2}}{\pi} + \frac{g_{a\gamma\gamma}^2 m_a^3}{64\pi}, \quad (\text{A.2})$$

and

$$\Gamma_a(m_a \leq 2m_\chi) = \frac{g_{a\gamma\gamma}^2 m_a^3}{64\pi}, \quad (\text{A.3})$$

depending on the mass relations of DM and ALP.

For relevant scattering processes we have

$$\int_{-4k_{\text{cm}}^2}^0 dt(-t)|\mathcal{M}_{\chi\gamma\rightarrow\chi\gamma}|^2 = 8g_{a\chi\chi}^2 g_{a\gamma\gamma}^2 m_\chi^2 \left( 4m_a^6 \left( \log(m_a^2) - \log\left(\frac{4k^2 m_\chi}{m_\chi+2\omega} + m_a^2\right) \right) \right. \\ \left. + \frac{4}{3}k^2 m_\chi \left( \frac{16k^4 m_\chi^2}{(m_\chi+2\omega)^3} - \frac{12k^2 m_a^2 m_\chi}{(m_\chi+2\omega)^2} \right) \right. \\ \left. + \frac{3m_a^6}{4k^2 m_\chi + m_a^2(m_\chi+2\omega)} + \frac{9m_a^4}{m_\chi+2\omega} \right), \quad (\text{A.4})$$

and

$$\int_{-4k_{\text{cm}}^2}^0 dt(-t)|\mathcal{M}_{\chi a\rightarrow\chi a}|^2 = -\frac{1}{(m_a^2+2m_\chi\omega)^2} 256g_{a\chi}^4 m_\chi^2 \left( -\frac{1}{3} \left( (m_a^2 + 2m_\chi\omega)(m_a^2 + m_\chi(m_\chi + 2\omega)) \right. \right. \\ \left. \left. \times \left( \frac{4k^2 m_\chi^2}{m_a^2+m_\chi^2+2m_\chi\omega} + m_a^2 - 2m_\chi\omega \right)^3 \right) \right. \\ \left. - \frac{m_a^4 m_\chi^2 (m_a^2 - 2m_\chi\omega)(m_a^2 + 2m_\chi\omega)^2}{\frac{4k^2 m_\chi^2}{m_a^2+m_\chi^2+2m_\chi\omega} + m_a^2 - 2m_\chi\omega} \right. \\ \left. - 2m_\chi^2 (m_a^8 - 2m_a^4 m_\chi^2 \omega^2 + 8m_a^2 m_\chi^3 \omega^3 + 8m_\chi^4 \omega^4) \right. \\ \left. \times \log\left(-\frac{-2m_\chi^2(\omega(m_\chi+2\omega)-2k^2)+m_a^4+m_a^2 m_\chi^2}{M_a^2+m_\chi(m_\chi+2\omega)}\right) \right. \\ \left. - (-m_a^6 + m_a^4 m_\chi(m_\chi - 2\omega) + 4m_a^2 m_\chi^2 \omega(m_\chi + \omega)) \right. \\ \left. + 2m_\chi^3 \omega^2 (3m_\chi + 4\omega) \left( \frac{4k^2 m_\chi^2}{m_a^2+m_\chi^2+2m_\chi\omega} + m_a^2 - 2m_\chi\omega \right)^2 \right. \\ \left. - (m_a^8 - 4m_a^6 m_\chi^2 - 8m_a^4 m_\chi^2 \omega^2 + 20m_a^2 m_\chi^4 \omega^2 \right. \\ \left. + 8m_\chi^4 \omega^3 (3m_\chi + 2\omega) \left( \frac{4k^2 m_\chi^2}{m_a^2+m_\chi^2+2m_\chi\omega} + m_a^2 - 2m_\chi\omega \right) \right. \\ \left. + \frac{1}{3} (m_a^{10} - m_a^8 m_\chi (5m_\chi + 2\omega) + 8m_a^6 m_\chi^2 \omega (4m_\chi - \omega) \right. \\ \left. + 2m_a^4 m_\chi^3 \omega^2 (27m_\chi + 8\omega) - 8m_a^2 m_\chi^4 \omega^3 (7m_\chi - 2\omega) \right. \\ \left. + 6m_\chi^2 (m_a^8 - 2m_a^4 m_\chi^2 \omega^2 + 8m_a^2 m_\chi^3 \omega^3 + 8m_\chi^4 \omega^4) \right. \\ \left. \times \log(2m_\chi\omega - m_a^2) - 8m_\chi^5 \omega^4 (11m_\chi + 4\omega) \right), \quad (\text{A.5})$$

where  $k$  and  $\omega$  respectively represent the four-momentum and the energy of the thermal bath particles involved in scattering with DM.

## References

- [1] P. W. Graham, I. G. Irastorza, S. K. Lamoreaux, A. Lindner, and K. A. van Bibber, [Ann. Rev. Nucl. Part. Sci.](#) **65**, 485 (2015), [arXiv:1602.00039 \[hep-ex\]](#).
- [2] K. Choi, S. H. Im, and C. Sub Shin, [Ann. Rev. Nucl. Part. Sci.](#) **71**, 225 (2021), [arXiv:2012.05029 \[hep-ph\]](#).
- [3] R. D. Peccei and H. R. Quinn, [Phys. Rev. Lett.](#) **38**, 1440 (1977).

- [4] S. Weinberg, *Phys. Rev. Lett.* **40**, 223 (1978).
- [5] F. Wilczek, *Phys. Rev. Lett.* **40**, 279 (1978).
- [6] J. E. Kim, *Phys. Rev. Lett.* **43**, 103 (1979).
- [7] J. Preskill, M. B. Wise, and F. Wilczek, *Phys. Lett. B* **120**, 127 (1983).
- [8] L. F. Abbott and P. Sikivie, *Phys. Lett. B* **120**, 133 (1983).
- [9] M. Dine and W. Fischler, *Phys. Lett. B* **120**, 137 (1983).
- [10] R. Blumenhagen and E. Plauschinn, *Phys. Lett. B* **736**, 482 (2014), [arXiv:1404.3542 \[hep-th\]](#) .
- [11] T. Kobayashi and L. Ubaldi, *JHEP* **09**, 052 (2020), [arXiv:2006.09389 \[hep-ph\]](#) .
- [12] E. I. Guendelman and D. A. Owen, *Phys. Lett. B* **276**, 108 (1992).
- [13] V. Domcke, Y. Ema, K. Mukaida, and M. Yamada, *JHEP* **08**, 096 (2020), [arXiv:2006.03148 \[hep-ph\]](#) .
- [14] R. T. Co, N. Fernandez, A. Ghalsasi, K. Harigaya, and J. Shelton, *Phys. Rev. D* **110**, 083534 (2024), [arXiv:2405.12268 \[hep-ph\]](#) .
- [15] Y. Nomura and J. Thaler, *Phys. Rev. D* **79**, 075008 (2009), [arXiv:0810.5397 \[hep-ph\]](#) .
- [16] A. Bharucha, F. Brümmer, N. Desai, and S. Mutzel, *JHEP* **02**, 141 (2023), [arXiv:2209.03932 \[hep-ph\]](#) .
- [17] S. Allen, A. Blackburn, O. Cardenas, Z. Messenger, N. H. Nguyen, and B. Shuve, *Phys. Rev. D* **110**, 095010 (2024), [arXiv:2405.02403 \[hep-ph\]](#) .
- [18] S. H. Im and K. S. Jeong, *Phys. Lett. B* **799**, 135044 (2019), [arXiv:1907.07383 \[hep-ph\]](#) .
- [19] D. K. Ghosh, A. Ghoshal, and S. Jeusun, *JHEP* **01**, 026 (2024), [arXiv:2305.09188 \[hep-ph\]](#) .
- [20] P. Arias, B. Diaz Saez, and J. Jaeckel, (2025), [arXiv:2501.17234 \[hep-ph\]](#) .
- [21] W.-L. Guo and Y.-L. Wu, *Phys. Rev. D* **79**, 055012 (2009), [arXiv:0901.1450 \[hep-ph\]](#) .
- [22] M. Laine, *JHEP* **01**, 157 (2023), [arXiv:2211.06008 \[hep-ph\]](#) .
- [23] H. M. Lee, M. Park, and W.-I. Park, *Phys. Rev. D* **86**, 103502 (2012), [arXiv:1205.4675 \[hep-ph\]](#) .
- [24] A. Ibarra, H. M. Lee, S. López Gehler, W.-I. Park, and M. Pato, *JCAP* **05**, 016 (2013), [Erratum: *JCAP* 03, E01 (2016)], [arXiv:1303.6632 \[hep-ph\]](#) .
- [25] T. Klangburam and C. Pongkitivanichkul, *JHEP* **10**, 145 (2024), [arXiv:2311.15681 \[hep-ph\]](#) .
- [26] M. Yang, Z.-Q. Guo, X.-Y. Luo, Z.-Q. Shen, Z.-Q. Xia, C.-T. Lu, Y.-L. S. Tsai, and Y.-Z. Fan, *JHEP* **10**, 094 (2024), [arXiv:2407.06815 \[hep-ph\]](#) .
- [27] S. Biswas, A. Chatterjee, E. Gabrielli, and B. Mele, *Phys. Rev. D* **100**, 115040 (2019), [arXiv:1906.10608 \[hep-ph\]](#) .
- [28] G. Armando, P. Panci, J. Weiss, and R. Ziegler, *Phys. Rev. D* **109**, 055029 (2024), [arXiv:2310.05827 \[hep-ph\]](#) .
- [29] K. Jodłowski, (2024), [arXiv:2411.19196 \[hep-ph\]](#) .
- [30] I. Adachi *et al.* (Belle-II), *Phys. Rev. D* **109**, 112006 (2024), [arXiv:2311.14647 \[hep-ex\]](#) .

- [31] P. Athron, R. Martinez, and C. Sierra, *JHEP* **02**, 121 (2024), [arXiv:2308.13426 \[hep-ph\]](#) .
- [32] R. Bause, H. Gisbert, and G. Hiller, *Phys. Rev. D* **109**, 015006 (2024), [arXiv:2309.00075 \[hep-ph\]](#) .
- [33] L. Allwicher, D. Becirevic, G. Piazza, S. Rosauero-Alcaraz, and O. Sumensari, *Phys. Lett. B* **848**, 138411 (2024), [arXiv:2309.02246 \[hep-ph\]](#) .
- [34] M. Abdughani and Y. Reyimuaji, *Phys. Rev. D* **110**, 055013 (2024), [arXiv:2309.03706 \[hep-ph\]](#) .
- [35] X.-G. He, X.-D. Ma, and G. Valencia, *Phys. Rev. D* **109**, 075019 (2024), [arXiv:2309.12741 \[hep-ph\]](#) .
- [36] A. Datta, D. Marfatia, and L. Mukherjee, *Phys. Rev. D* **109**, L031701 (2024), [arXiv:2310.15136 \[hep-ph\]](#) .
- [37] W. Altmannshofer, A. Crivellin, H. Haigh, G. Inguglia, and J. Martin Camalich, *Phys. Rev. D* **109**, 075008 (2024), [arXiv:2311.14629 \[hep-ph\]](#) .
- [38] D. McKeen, J. N. Ng, and D. Tuckler, *Phys. Rev. D* **109**, 075006 (2024), [arXiv:2312.00982 \[hep-ph\]](#) .
- [39] K. Fridell, M. Ghosh, T. Okui, and K. Tobioka, *Phys. Rev. D* **109**, 115006 (2024), [arXiv:2312.12507 \[hep-ph\]](#) .
- [40] S.-Y. Ho, J. Kim, and P. Ko, (2024), [arXiv:2401.10112 \[hep-ph\]](#) .
- [41] F.-Z. Chen, Q. Wen, and F. Xu, *Eur. Phys. J. C* **84**, 1012 (2024), [arXiv:2401.11552 \[hep-ph\]](#) .
- [42] E. Gabrielli, L. Marzola, K. Mürsepp, and M. Raidal, *Eur. Phys. J. C* **84**, 460 (2024), [arXiv:2402.05901 \[hep-ph\]](#) .
- [43] B.-F. Hou, X.-Q. Li, M. Shen, Y.-D. Yang, and X.-B. Yuan, *JHEP* **06**, 172 (2024), [arXiv:2402.19208 \[hep-ph\]](#) .
- [44] C.-H. Chen and C.-W. Chiang, *Phys. Rev. D* **110**, 075036 (2024), [arXiv:2403.02897 \[hep-ph\]](#) .
- [45] X.-G. He, X.-D. Ma, M. A. Schmidt, G. Valencia, and R. R. Volkas, *JHEP* **07**, 168 (2024), [arXiv:2403.12485 \[hep-ph\]](#) .
- [46] P. D. Bolton, S. Fajfer, J. F. Kamenik, and M. Novoa-Brunet, *Phys. Rev. D* **110**, 055001 (2024), [arXiv:2403.13887 \[hep-ph\]](#) .
- [47] D. Marzocca, M. Nardecchia, A. Stanzione, and C. Toni, *Eur. Phys. J. C* **84**, 1217 (2024), [arXiv:2404.06533 \[hep-ph\]](#) .
- [48] S. Rosauero-Alcaraz and L. P. S. Leal, *Eur. Phys. J. C* **84**, 795 (2024), [arXiv:2404.17440 \[hep-ph\]](#) .
- [49] C. S. Kim, D. Sahoo, and K. N. Vishnudath, *Eur. Phys. J. C* **84**, 882 (2024), [arXiv:2405.17341 \[hep-ph\]](#) .
- [50] C. Hati, J. Leite, N. Nath, and J. W. F. Valle, (2024), [arXiv:2408.00060 \[hep-ph\]](#) .
- [51] L. Allwicher, M. Bordone, G. Isidori, G. Piazza, and A. Stanzione, (2024), [arXiv:2410.21444 \[hep-ph\]](#) .
- [52] D. Bečirević, S. Fajfer, N. Košnik, and L. Pavičić, (2024), [arXiv:2410.23257 \[hep-ph\]](#) .

- [53] W. Altmannshofer and S. Roy, (2024), [arXiv:2411.06592 \[hep-ph\]](#) .
- [54] A. J. Buras and P. Stangl, (2024), [arXiv:2412.14254 \[hep-ph\]](#) .
- [55] Q.-Y. Hu, (2024), [arXiv:2412.19084 \[hep-ph\]](#) .
- [56] C.-Q. Zhang, J. Sun, Z.-P. Xing, and R.-L. Zhu, (2024), [arXiv:2501.00512 \[hep-ph\]](#) .
- [57] L. Calibbi, T. Li, L. Mukherjee, and M. A. Schmidt, (2025), [arXiv:2502.04900 \[hep-ph\]](#) .
- [58] J.-P. Lee, (2025), [arXiv:2502.06370 \[hep-ph\]](#) .
- [59] X.-G. He, X.-D. Ma, J. Tandean, and G. Valencia, (2025), [arXiv:2502.09603 \[hep-ph\]](#) .
- [60] A. Berezhnoy, W. Lucha, and D. Melikhov, (2025), [arXiv:2502.14313 \[hep-ph\]](#) .
- [61] P. D. Bolton, S. Fajfer, J. F. Kamenik, and M. Novoa-Brunet, (2025), [arXiv:2503.19025 \[hep-ph\]](#) .
- [62] T. M. Aliev, A. Elpe, I. Turan, and L. Selbuz, (2025), [arXiv:2503.22347 \[hep-ph\]](#) .
- [63] C.-H. Chen, C.-W. Chiang, and L. M. G. de la Vega, (2025), [arXiv:2503.22431 \[hep-ph\]](#) .
- [64] Y.-N. Wang, X.-C. Duan, T.-P. Tang, Z. Wang, and Y.-L. S. Tsai, (2025), [arXiv:2502.18263 \[hep-ph\]](#) .
- [65] G. Bélanger, S. Chakraborti, C. Delaunay, and M. Jomain, (2025), [arXiv:2503.08897 \[hep-ph\]](#) .
- [66] L. G. van den Aarssen, T. Bringmann, and Y. C. Goedecke, *Phys. Rev. D* **85**, 123512 (2012), [arXiv:1202.5456 \[hep-ph\]](#) .
- [67] T. Binder, T. Bringmann, M. Gustafsson, and A. Hryczuk, *Phys. Rev. D* **96**, 115010 (2017), [Erratum: *Phys.Rev.D* 101, 099901 (2020)], [arXiv:1706.07433 \[astro-ph.CO\]](#) .
- [68] F. Brümmer, *JHEP* **01**, 113 (2020), [arXiv:1910.01549 \[hep-ph\]](#) .
- [69] K. Ala-Mattinen and K. Kainulainen, *JCAP* **09**, 040 (2020), [arXiv:1912.02870 \[hep-ph\]](#) .
- [70] T. Abe, *Phys. Rev. D* **102**, 035018 (2020), [arXiv:2004.10041 \[hep-ph\]](#) .
- [71] T. Binder, T. Bringmann, M. Gustafsson, and A. Hryczuk, *Eur. Phys. J. C* **81**, 577 (2021), [arXiv:2103.01944 \[hep-ph\]](#) .
- [72] B. Zhu and X. Liu, *Sci. China Phys. Mech. Astron.* **65**, 231011 (2022), [arXiv:2104.03238 \[hep-ph\]](#) .
- [73] A. Hryczuk and M. Laletin, *JHEP* **06**, 026 (2021), [arXiv:2104.05684 \[hep-ph\]](#) .
- [74] T. Abe, *Phys. Rev. D* **104**, 035025 (2021), [arXiv:2106.01956 \[hep-ph\]](#) .
- [75] Y. Du, F. Huang, H.-L. Li, Y.-Z. Li, and J.-H. Yu, *JCAP* **04**, 012 (2022), [arXiv:2111.01267 \[hep-ph\]](#) .
- [76] K. Ala-Mattinen, M. Heikinheimo, K. Kainulainen, and K. Tuominen, *Phys. Rev. D* **105**, 123005 (2022), [arXiv:2201.06456 \[hep-ph\]](#) .
- [77] A. Hryczuk and M. Laletin, *Phys. Rev. D* **106**, 023007 (2022), [arXiv:2204.07078 \[hep-ph\]](#) .
- [78] Y. Liu, X. Liu, and B. Zhu, *Phys. Rev. D* **107**, 115009 (2023), [arXiv:2301.12199 \[hep-ph\]](#) .
- [79] A. Aboubrahim, M. Klasen, and L. P. Wiggering, *JCAP* **08**, 075 (2023), [arXiv:2306.07753 \[hep-ph\]](#) .

- [80] X.-C. Duan, R. Ramos, and Y.-L. S. Tsai, *Phys. Rev. D* **110**, 063535 (2024), [arXiv:2404.12019 \[hep-ph\]](#) .
- [81] P. Gondolo and G. Gelmini, *Nucl. Phys. B* **360**, 145 (1991).
- [82] E. Izaguirre, T. Lin, and B. Shuve, *Phys. Rev. Lett.* **118**, 111802 (2017), [arXiv:1611.09355 \[hep-ph\]](#) .
- [83] H. Georgi, D. B. Kaplan, and L. Randall, *Phys. Lett. B* **169**, 73 (1986).
- [84] I. Brivio, M. B. Gavela, L. Merlo, K. Mimasu, J. M. No, R. del Rey, and V. Sanz, *Eur. Phys. J. C* **77**, 572 (2017), [arXiv:1701.05379 \[hep-ph\]](#) .
- [85] J. Ren, D. Wang, L. Wu, J. M. Yang, and M. Zhang, *JHEP* **11**, 138 (2021), [arXiv:2106.07018 \[hep-ph\]](#) .
- [86] C.-T. Lu, X. Luo, and X. Wei, *Chin. Phys. C* **47**, 103102 (2023), [arXiv:2303.03110 \[hep-ph\]](#) .
- [87] K. Cheung and C. J. Ouseph, *Phys. Rev. D* **108**, 035003 (2023), [arXiv:2303.16514 \[hep-ph\]](#) .
- [88] M. J. Dolan, T. Ferber, C. Hearty, F. Kahlhoefer, and K. Schmidt-Hoberg, *JHEP* **12**, 094 (2017), [Erratum: *JHEP* 03, 190 (2021)], [arXiv:1709.00009 \[hep-ph\]](#) .
- [89] S. Gola, S. Mandal, and N. Sinha, *Int. J. Mod. Phys. A* **37**, 2250131 (2022), [arXiv:2106.00547 \[hep-ph\]](#) .
- [90] F. Acanfora, R. Franceschini, A. Mastroddi, and D. Redigolo, *JHEP* **11**, 156 (2024), [arXiv:2307.06369 \[hep-ph\]](#) .
- [91] T. Ferber, A. Filimonova, R. Schäfer, and S. Westhoff, *JHEP* **04**, 131 (2023), [arXiv:2201.06580 \[hep-ph\]](#) .
- [92] A. Abada and S. Nasri, *Phys. Rev. D* **85**, 075009 (2012), [arXiv:1201.1413 \[hep-ph\]](#) .
- [93] J. P. Lees et al. (BaBar), *Phys. Rev. Lett.* **128**, 131802 (2022), [arXiv:2111.01800 \[hep-ex\]](#) .
- [94] D. Wang, L. Wu, J. M. Yang, and M. Zhang, *Phys. Rev. D* **104**, 095016 (2021), [arXiv:2102.01532 \[hep-ph\]](#) .
- [95] W. G. Parrott, C. Bouchard, and C. T. H. Davies (HPQCD), *Phys. Rev. D* **107**, 014511 (2023), [Erratum: *Phys.Rev.D* 107, 119903 (2023)], [arXiv:2207.13371 \[hep-ph\]](#) .
- [96] T. Binder, L. Covi, A. Kamada, H. Murayama, T. Takahashi, and N. Yoshida, *JCAP* **11**, 043 (2016), [arXiv:1602.07624 \[hep-ph\]](#) .
- [97] E. Bertschinger, *Phys. Rev. D* **74**, 063509 (2006), [arXiv:astro-ph/0607319](#) .
- [98] T. Bringmann and S. Hofmann, *JCAP* **04**, 016 (2007), [Erratum: *JCAP* 03, E02 (2016)], [arXiv:hep-ph/0612238](#) .
- [99] T. Bringmann, *New J. Phys.* **11**, 105027 (2009), [arXiv:0903.0189 \[astro-ph.CO\]](#) .
- [100] P. Gondolo, J. Hisano, and K. Kadota, *Phys. Rev. D* **86**, 083523 (2012), [arXiv:1205.1914 \[hep-ph\]](#) .
- [101] N. Aghanim et al. (Planck), *Astron. Astrophys.* **641**, A6 (2020), [Erratum: *Astron.Astrophys.* 652, C4 (2021)], [arXiv:1807.06209 \[astro-ph.CO\]](#) .
- [102] M. Duerr, T. Ferber, C. Garcia-Cely, C. Hearty, and K. Schmidt-Hoberg, *JHEP* **04**, 146 (2021), [arXiv:2012.08595 \[hep-ph\]](#) .

- [103] W. Altmannshofer et al. (Belle-II), *PTEP* **2019**, 123C01 (2019), [Erratum: PTEP 2020, 029201 (2020)], [arXiv:1808.10567 \[hep-ex\]](#) .
- [104] E. Cortina Gil et al. (NA62), *JHEP* **06**, 093 (2021), [arXiv:2103.15389 \[hep-ex\]](#) .
- [105] E. Cortina Gil et al. (NA62), *JHEP* **02**, 201 (2021), [arXiv:2010.07644 \[hep-ex\]](#) .
- [106] J. Jaeckel and M. Spannowsky, *Phys. Lett. B* **753**, 482 (2016), [arXiv:1509.00476 \[hep-ph\]](#) .
- [107] R. Mammen Abraham et al. (FASER), *JHEP* **01**, 199 (2025), [arXiv:2410.10363 \[hep-ex\]](#) .
- [108] J. P. Lees et al. (BaBar), *Phys. Rev. Lett.* **119**, 131804 (2017), [arXiv:1702.03327 \[hep-ex\]](#) .
- [109] J. P. Lees et al. (BaBar), *Phys. Rev. D* **87**, 112005 (2013), [arXiv:1303.7465 \[hep-ex\]](#) .