

# Tensor Meson Pole contributions to the HLbL piece of $a_\mu^{\text{HLbL}}$ within R $\chi$ T

Emilio J. Estrada <sup>1</sup> and Pablo Roig <sup>1,2</sup>

<sup>1</sup>*Departamento de Física, Centro de Investigación y de Estudios Avanzados del  
Instituto Politécnico Nacional  
Apartado Postal 14-740, 07360 Ciudad de México, México*

<sup>2</sup>*IFIC, Universitat de València – CSIC, Catedrático José Beltrán 2, E-46980 Paterna,  
Spain*

## Abstract

We report our results for the tensor meson pole contributions to the Hadronic Light-by-Light piece of  $a_\mu$ , using Resonance Chiral Theory in the purely hadronic region. Given the differences between the dispersive and holographic results for it and the resulting discussion of the corresponding uncertainty estimate for the Hadronic Light-by-Light section of the  $g - 2$  theory initiative second White Paper, we consider timely to present an alternative evaluation. In our approach, in addition to the lightest tensor meson nonet, two vector meson resonance nonets are considered. We work in the chiral limit, where all parameters are determined by imposing short-distance QCD constraints, and the radiative tensor decay widths. Within this setting, only the form factor  $\mathcal{F}_1^T$  is non-vanishing, in agreement with the result put forward by the dispersive study. We obtain the following results for the different contributions (in units of  $10^{-11}$ ):  $a_\mu^{\text{a}_2\text{-pole}} = -1.09(10)_{\text{stat}}^{(+0.0)}_{(-0.11)}_{\text{syst}}$ ,  $a_\mu^{\text{f}_2\text{-pole}} = -3.4(3)_{\text{stat}}^{(+0.4)}_{(-0.0)}_{\text{syst}}$  and  $a_\mu^{\text{f}'_2\text{-pole}} = -0.046(14)_{\text{stat}}^{(+0.008)}_{(-0.000)}_{\text{syst}}$ , which add up to  $a_\mu^{(\text{a}_2+\text{f}_2+\text{f}'_2)\text{-pole}} = -4.5^{+0.5}_{-0.3}$ .

# 1 Introduction

The final measurement of the Muon  $g - 2$  Collaboration will be announced soon and its prediction in the Standard Model will be updated just before, marking a milestone in the exploration of the high-intensity frontier of particle physics.

The current experimental average is based on the legacy BNL result [1] and in the two measurements published so far by FNAL [2, 3], reaching an uncertainty of  $22 \times 10^{-11}$  [4]. The current SM prediction was published in the White Paper [5, 6], based on refs. [7–41]<sup>1</sup>, with an error of  $43 \times 10^{-11}$ . A reduction of this uncertainty at the level of  $\leq 16 \times 10^{-11}$  would be desirable to maximize the new physics implications of the final FNAL result [102].

Within this global effort, we will focus in this paper on a subleading contribution to the hadronic light-by-light (HLbL) piece of the muon anomaly ( $a_\mu = (g_\mu - 2)/2$ ) which seems puzzling at present.

A tension has recently appeared, between the determinations of the tensor meson contributions to the HLbL piece of  $a_\mu$  by the dispersive [103, 104] and holographic [105, 106] groups (see also the earlier holographic computation of ref. [107] and refs. [30–32, 108], which were used to estimate it in the first White Paper [5]). While these results basically agree on its magnitude, they differ in sign. This current puzzle motivated us to compute the dispersively defined lightest tensor meson pole contributions within Resonance Chiral Theory ( $R\chi T$ , see section 2) [109, 110], which bridges between the chiral and asymptotic QCD regimes.

The study of tensor mesons in  $R\chi T$  is scarce, as it contributes negligibly to chiral dynamics [111] (see also ref. [112]). We will benefit from the recent ref. [113], which particularly focused on understanding tensor meson masses and their di-pion and radiative decay widths. As first pointed out in ref. [103], the  $\mathcal{F}_1^T$  form factor (FF) is the leading effect at low energies, which is also true in the perturbative regime [114]. In accordance with these results, we find that the remaining four FFs vanish, within our approach. This is important, since one of such FFs,  $\mathcal{F}_3^T$ , appears responsible for the positive sign found in the holographic computation [105] of this contribution to  $a_\mu$ . Our description of the  $T\gamma\gamma$  interactions fulfills the short-distance QCD constraint of the tensor mesons transition form factors (TFFs) when both virtualities are asymptotic [115] and in the Brodsky-Lepage limit [116]. This requires a  $\sim 1/Q^4$  damping of the  $\mathcal{F}_1^T$  FF

---

<sup>1</sup>See also, e. g. refs. [42–101].

in the ultraviolet, which is achieved if at least two vector meson multiplets of resonances are accounted for. We describe our effective approach in section 2, impose high-energy constraints on the obtained FFs in section 3 and finally present our results –discussing the uncertainty estimation– for the  $a_\mu^{\text{HLbL: T-poles}}$  in section 4. We conclude in section 5.

## 2 Resonance Chiral Theory Lagrangian

Our theoretical framework will be a Lagrangian theory, that is briefly introduced in the following. In order to extend the applicability of Chiral Perturbation Theory [117–119] to larger energies, Resonance Chiral Theory ( $R\chi T$ ) [109, 110] includes the resonances as dynamical fields. Independence on the formalism used to describe the spin-one mesons is verified for the two-point functions and related observables, and the convenience of using antisymmetric tensors for the spin-one fields is explained [109, 110].  $R\chi T$  is built upon chiral symmetry for the pseudo-Goldstone bosons (pion, kaon and eta mesons) and unitary symmetry for the resonances. Its expansion parameter is the inverse of the number of colors of the QCD gauge group [120]. The number of relevant operators is limited by two observations. On the one hand, in  $R\chi T$  the number of resonance fields is typically restricted by the specific problem at hand. On the other, the inclusion of tensors of high order in the chiral counting conflicts with the requirements from short-distance QCD. Particularly, for order parameters of chiral symmetry breaking, the matching of  $R\chi T$  results to the corresponding operator product expansion (OPE) expressions has been pursued [109, 110, 121–127], rendering relations between Lagrangian parameters, thus increasing the predictive power of this framework.  $R\chi T$  has been employed successfully in the computation of a number of hadronic contributions to  $a_\mu$  [35, 48, 78, 82, 90, 126, 128–135].

We have considered the most general Lagrangian consistent with the symmetries of  $R\chi T$ , at leading order in the  $1/N_C$  expansion and in the chiral limit. In the minimal model, which includes pseudoscalar mesons, vector meson resonances, photons, tensor mesons, and their interactions, only 6 operators contribute – in this approximation– to the radiative tensor decays, which are collected in table 1. We have not included operators with additional derivatives that yield a short-distance behaviour at odds with short-distance QCD properties and the matching to OPE results. Their real coupling constants are named according to the interaction vertices they describe and are not fixed by symmetry requirements.

Coupling constant	Operator
$C_{T\gamma\gamma}$	$\langle T^{\mu\nu} f_{+\mu}^\alpha f_{+\alpha\nu} \rangle$
$C_{T\gamma V}$	$i \langle T^{\mu\nu} \{ f_{+\mu}^\alpha, V_{\alpha\nu} \} \rangle$
$C_{T\gamma V'}$	$i \langle T^{\mu\nu} \{ f_{+\mu}^\alpha, V'_{\alpha\nu} \} \rangle$
$C_{TVV}$	$\langle T^{\mu\nu} \{ V_\mu^\alpha, V_{\alpha\nu} \} \rangle$
$C_{TV'V'}$	$\langle T^{\mu\nu} \{ V'_\mu^\alpha, V'_{\alpha\nu} \} \rangle$
$C_{TVV'}$	$\langle T^{\mu\nu} \{ V_\mu^\alpha, V'_{\alpha\nu} \} \rangle$

Table 1: Relevant  $R\chi T$  Operators at leading order in the  $1/N_C$  expansion and in the chiral limit. We note that  $C_{T\gamma V^{(\prime)}}$  are dimensionless, while  $(C_{T\gamma\gamma})$   $C_{TV^{(\prime)}V^{(\prime)}}$  have (inverse) energy dimensions.

We recall the  $V \rightarrow \gamma$  terms (with trivial generalization for  $V \rightarrow V^{(\prime)}$ ), given in the original  $R\chi T$  Lagrangian [109], that are also needed to describe the tensor meson transition FFs,

$$\mathcal{L}^{V \rightarrow \gamma} = \frac{F_V}{2\sqrt{2}} \langle V_{\mu\nu} f_+^{\mu\nu} \rangle. \quad (2.1)$$

The tensor mesons,  $T^{\mu\nu}$ , are in the symmetric, on-shell traceless representation [111]:

$$T_{\mu\nu} = \begin{pmatrix} \frac{a_2^0}{\sqrt{2}} + \frac{f_2^8}{\sqrt{6}} + \frac{f_2^0}{\sqrt{3}} & a_2^+ & K_2^{*+} \\ a_2^- & -\frac{a_2^0}{\sqrt{2}} + \frac{f_2^8}{\sqrt{6}} + \frac{f_2^0}{\sqrt{3}} & K_2^{*0} \\ K_2^{*-} & \bar{K}_2^{*0} & -\frac{2f_2^8}{\sqrt{6}} + \frac{f_2^0}{\sqrt{3}} \end{pmatrix}_{\mu\nu}, \quad (2.2)$$

where the  $f_2^8$  and  $f_2^0$  octet and singlet states in flavor space, respectively, are related to the physical mass eigenstates via a single-angle mixing:

$$\begin{aligned} f_2^8 &= \sin \theta_T f_2 + \cos \theta_T f'_2, \\ f_2^0 &= \cos \theta_T f_2 - \sin \theta_T f'_2. \end{aligned} \quad (2.3)$$

The vector meson nonets are in the antisymmetric tensor representation [109]. Considering ideal mixing they read

$$V_{\mu\nu} = \begin{pmatrix} (\rho^0 + \omega)/\sqrt{2} & \rho^+ & K^{*+} \\ \rho^- & (-\rho^0 + \omega)/\sqrt{2} & K^{*0} \\ K^{*-} & \bar{K}^{*0} & \phi \end{pmatrix}_{\mu\nu}, \quad (2.4)$$

and similarly for  $V \rightarrow V'$ . The photon field is encoded in the  $f_+$  chiral tensor [136]:

$$f_+^{\mu\nu} = u F_L^{\mu\nu} u^\dagger + u^\dagger F_R^{\mu\nu} u, \quad F_{L(R)}^{\mu\nu} = -e Q F^{\mu\nu} + \dots, \quad (2.5)$$

where  $Q = \text{diag}(2/3, -1/3, -1/3)$ , and  $u = \exp\left(i\frac{\Phi}{\sqrt{2}F}\right)$  holds the information of the pseudo-Goldstone bosons ( $\pi$ , kaon and  $\eta$  mesons), which in this case do not play a role. Given these building blocks, we can construct the amplitude of the  $T \rightarrow \gamma\gamma$  decays and examine the associated transition FF (see Figure 1).




Figure 1: Tensor Meson transition FF, which gives the radiative decay with both photons on-shell.

We recall the matrix element of a massive tensor meson decaying into two virtual photons [103]

$$\langle \gamma^*(q_1, \lambda_1) \gamma^*(q_2, \lambda_2) | T(p, \lambda_T) \rangle = i(2\pi)^4 \delta^{(4)}(q_1 + q_2 - p) e^2 \varepsilon^{\lambda_1^*}(q_1) \varepsilon^{\lambda_2^*}(q_2) \varepsilon_{\alpha\beta}^{\lambda_T}(p) \mathcal{M}^{\mu\nu\alpha\beta}(q_1, q_2). \quad (2.6)$$

For the tensor mesons, the reduced amplitude (which -among other properties [103]- must be symmetric under  $q_1 \leftrightarrow q_2, \mu \leftrightarrow \nu$ ) can be split into 5 structures contributing to observables [115]:

$$\mathcal{M}^{\mu\nu\alpha\beta} = \sum_{i=5}^5 T_i^{\mu\nu\alpha\beta} \frac{1}{m_T^{n_i}} \mathcal{F}_i^T(q_1^2, q_2^2), \quad (2.7)$$

where  $n_1 = 1$  and  $n_{i \neq 1} = 3$ , and the tensor structures are:

$$\begin{aligned} T_1^{\mu\nu\alpha\beta} &= g^{\mu\alpha} P_{21}^{\nu\beta} + g^{\nu\alpha} P_{12}^{\mu\beta} + g^{\mu\beta} P_{21}^{\nu\alpha} + g^{\nu\beta} P_{12}^{\mu\alpha} + g^{\mu\nu} (q_1^\alpha q_2^\beta + q_1^\beta q_2^\alpha) - q_1 \cdot q_2 (g^{\mu\alpha} g^{\nu\beta} + g^{\nu\alpha} g^{\mu\beta}), \\ T_2^{\mu\nu\alpha\beta} &= (q_1^\alpha q_2^\beta + q_2^\alpha q_1^\beta) P_{12}^{\mu\nu}, \\ T_3^{\mu\nu\alpha\beta} &= P_{11}^{\mu\alpha} P_{22}^{\nu\beta} + P_{11}^{\mu\beta} P_{22}^{\nu\alpha}, \\ T_4^{\mu\nu\alpha\beta} &= P_{12}^{\mu\alpha} P_{22}^{\nu\beta} + P_{12}^{\mu\beta} P_{22}^{\nu\alpha}, \\ T_4^{\mu\nu\alpha\beta} &= P_{21}^{\nu\alpha} P_{11}^{\mu\beta} + P_{21}^{\nu\beta} P_{11}^{\mu\alpha}, \end{aligned} \quad (2.8)$$

where the  $P_{ij}^{\mu\nu}$  are defined as:

$$P_{ij}^{\mu\nu} = g^{\mu\nu} q_i \cdot q_j - q_i^\nu q_j^\mu. \quad (2.9)$$

In the model used in this work, only two derivatives appear in the  $T \rightarrow \gamma^*\gamma^*$  amplitude, resulting in just a single non-vanishing contribution, given by the  $\mathcal{F}_1^T$  FF, in

agreement with ref. [103]. This will determine that we get a negative result for the  $a_\mu^{\text{T-pole}}$  contribution, as in this reference, at variance with the holographic results, due to the  $\mathcal{F}_3^T$  contribution [105].

### 3 $\mathcal{F}_1^T$ and short-distance constraints

From the operators in table 1, we get the following transition FF for the 3 contributing neutral tensor mesons:

$$\begin{aligned} \mathcal{F}_1^{a_2}(q_1^2, q_2^2) = & -\frac{M_{a_2}}{2} \left[ \frac{\sqrt{2}}{3} C_{T\gamma\gamma} + C_{T\gamma V} \frac{4F_V}{3} (D_{M_V}^{-1}(q_1^2) + D_{M_V}^{-1}(q_2^2)) \right. \\ & \left. + C_{TVV} \frac{2\sqrt{2}F_V^2}{3} D_{M_V}^{-1}(q_1^2) D_{M_V}^{-1}(q_2^2) + C_{TVV'} \frac{2\sqrt{2}F_V F_{V'}}{3} D_{M_V}^{-1}(q_1^2) D_{M_{V'}}^{-1}(q_2^2) + V \leftrightarrow V' \right], \end{aligned} \quad (3.1)$$

$$\begin{aligned} \mathcal{F}_1^{f_2}(q_1^2, q_2^2) = & -\frac{M_{f_2}}{2} \left[ \frac{\sqrt{2}}{3\sqrt{3}} C_{T\gamma\gamma} (\sin \theta_T + 2\sqrt{2} \cos \theta_T) \right. \\ & + C_{T\gamma V} (\sin \theta_T + 11\sqrt{2} \cos \theta_T) \frac{2\sqrt{3}F_V}{27} (D_{M_V}^{-1}(q_1^2) + D_{M_V}^{-1}(q_2^2)) \\ & + C_{TVV} (\sin \theta_T + 2\sqrt{2} \cos \theta_T) \frac{2\sqrt{2}F_V^2}{3\sqrt{3}} D_{M_V}^{-1}(q_1^2) D_{M_V}^{-1}(q_2^2) \\ & \left. + C_{TVV'} (\sin \theta_T + 2\sqrt{2} \cos \theta_T) \frac{2\sqrt{2}F_V F_{V'}}{3\sqrt{3}} D_{M_V}^{-1}(q_1^2) D_{M_{V'}}^{-1}(q_2^2) + V \leftrightarrow V' \right], \end{aligned} \quad (3.2)$$

$$\mathcal{F}_1^{f'_2}(q_1^2, q_2^2) = \mathcal{F}_1^{f_2}(q_1^2, q_2^2)|_{\sin \theta_T \rightarrow \cos \theta_T, \cos \theta_T \rightarrow -\sin \theta_T, M_{f_2} \rightarrow M_{f'_2}}, \quad (3.3)$$

where  $D_R(q_i^2) = M_R^2 - q_i^2$ .

Ref. [115] derived the behavior of the  $\mathcal{F}_i^T$  form factors in the asymptotic region. Since in a model with finite resonances only a discrete amount of short-distance constraints (SDCs) can be imposed, we choose fixing the symmetric double virtual case and the single virtual  $1/Q^4$  behavior

$$\lim_{Q^2 \rightarrow \infty} \mathcal{F}_1^T(-Q^2, -Q^2) = -\frac{3F_T^{\text{eff}} M_T^3}{14Q^4}, \quad (3.4a)$$

$$\lim_{Q^2 \rightarrow \infty} \mathcal{F}_1^T(-Q^2, 0) \sim \frac{1}{Q^4}, \quad (3.4b)$$

where  $Q_i^2 = -q_i^2$ , and  $F_T^{\text{eff}} = 4 \sum_a C_a F_T^a = 79(8)$  MeV [103]. We note that the magnitude of  $\mathcal{F}_1^T$  is overestimated in this way, since for one finite and another asymptotic photon virtuality,  $\mathcal{F}_1^T$  should vanish as  $\sim 1/Q^4$ . However, with two multiplets of vector meson resonances (and also with three of them), we get  $\mathcal{F}_1^T \sim 1/Q^2$  in this mixed region, which we consider as a systematic error when computing  $a_\mu^{\text{HLbL: T-pole}}$ . The resulting FFs, after imposing the high-energy constraints mentioned above are:

$$\mathcal{F}_1^T(q_1^2, q_2^2) = c_T \frac{9 F_T^{\text{eff}} M_T^3 (M_V^4 - q_1^2 q_2^2) - 14\sqrt{2} C_{TVV} F_V^2 M_T (M_V^2 - M_{V'}^2)^2}{42 D_{M_V}(q_1^2) D_{M_V}(q_2^2) D_{M_{V'}}(q_1^2) D_{M_{V'}}(q_2^2)}, \quad (3.5)$$

where  $c_T$  is a flavor space rotation given by:  $c_{a_2} = 1$ ,  $c_{f_2} = \frac{\sin \theta_T + 2\sqrt{2} \cos \theta_T}{\sqrt{3}}$ ,  $c_{f'_2} = \frac{\cos \theta_T - 2\sqrt{2} \sin \theta_T}{\sqrt{3}}$ . The only free parameters left are the combination  $C_{TVV} F_V^2$  and the mixing angle  $\theta_T$ . These two parameters can be used to normalize the form factors to reproduce the radiative decay widths as was done in ref. [113]. We will take their chiral limit value for  $\theta_T$  and fix  $C_{TVV} F_V^2$  to the measured value of  $\Gamma(a_2 \rightarrow \gamma\gamma)$ . The corresponding uncertainty is negligible with respect to those discussed in the next section.

## 4 Form Factors and $a_\mu$ computation

For the rest of the parameters of eq. (3.5), we use the PDG input [4] for the tensor meson masses and their radiative decay widths. The effective masses of the vector meson multiplets are obtained from the study of  $a_\mu^{\text{P-pole}}$ , within  $R\chi T$ , in ref. [131] and  $F_T^{\text{eff}}$  is taken from ref. [137]. Thus, with all parameters fixed, we obtain a complete description for the tensor mesons  $\mathcal{F}_1^T$ , which are shown in Figure 2 for the single virtual and the symmetric double virtual case of  $a_2$ ,  $f_2$  and  $f'_2$ . There is scarce related experimental information, Belle [138] measured the single virtual form factors for the  $f_2(1270)$  tensor meson in the helicity basis. The change of basis is [115]:

$$\mathcal{F}_{\lambda=0}^T = \frac{Q^2}{\sqrt{6} M_T^2} \mathcal{F}_1^T(-Q^2, 0) - \frac{(M_T^2 + Q^2)^2}{2\sqrt{6} M_T^4} \mathcal{F}_2^T(-Q^2, 0) + \frac{Q^2}{\sqrt{6} M_T^2} \mathcal{F}_5^T(-Q^2, 0), \quad (4.1a)$$

$$\mathcal{F}_{\lambda=1}^T = \frac{\sqrt{Q^2}}{\sqrt{2} M_T} \mathcal{F}_1^T(-Q^2, 0) + \frac{\sqrt{Q^2} (M_T^2 - Q^2)}{2\sqrt{2} M_T^3} \mathcal{F}_5^T(-Q^2, 0), \quad (4.1b)$$

$$\mathcal{F}_{\lambda=2}^T = -\mathcal{F}_1^T(-Q^2, 0) + \frac{Q^2}{M_T^2} \mathcal{F}_5^T(-Q^2, 0). \quad (4.1c)$$

For our case, this is reduced as  $\mathcal{F}_{2,5}^T(-Q^2, 0) = 0$ . The comparison of these data with the Quark Model used in [103], the Hard-Wall model used in [105], and our results is shown in Figure 3. Our model exhibits good agreement with the experimental data.




Figure 2: Comparison between the simple quark model [114] used in the dispersive determination [103], the hQCD hard-wall model [105], the R $\chi$ T model used in this work and the  $C_2^0$  of eq. (4.4) for the single and symmetric double virtual form factor  $\mathcal{F}_1^T$ , for all 3 tensor mesons:  $a_2(1320)$ ,  $f_2(1270)$  and  $f_2'(1525)$ . Our one  $\sigma$  uncertainties are displayed by the green band.




Figure 3: Comparison between the simple quark model [114] used in ref. [103], the hard-wall model [105] and this work with the Belle data [138] in the helicity basis of eq. (4.1) for the  $f_2(1270)$  tensor meson, normalized by  $\mathcal{F}^{f_2}(0, 0) = \sqrt{\frac{5\Gamma_{T\gamma\gamma}}{\pi\alpha^2 M_T}}$ . Our one  $\sigma$  uncertainties are displayed by the green band.

It is important to remark that –by construction– these TFFs fulfill the doubly virtual asymptotic behavior given in ref. [115]. However, for arbitrary asymmetries –parametrized by  $w = \frac{q_1^2 - q_2^2}{q_1^2 + q_2^2}$ , so that  $\mathcal{F}_1^T(q_1^2, q_2^2) = \mathcal{F}_1^T(Q^2) f_1^T(\omega)$ –, for  $Q^2 = \frac{q_1^2 + q_2^2}{2} \rightarrow \infty$ , we get:

$$\mathcal{F}_1^T(q_1^2, q_2^2) \rightarrow \frac{4 \sum_a C_a F_T^a m_T^3}{Q^4} \left( -\frac{3}{14} (1 - w^2)^{-1} \right), \quad (4.2)$$

which is simpler than the ones obtained by refs. [115] and [105]. One reason for the lack of structure of our result is the truncation of the infinite tower of interactions to two vector and the lightest tensor meson multiplets. If the infinite tower of resonances were not integrated out, a logarithmic term would be expected, as in the quoted refs. The comparison between the asymmetry functions for the  $a_2$  –which is representative for all tensor mesons–, is shown in Figure 4, where it is clear that none of the 3 models that have




Figure 4: Comparison of the full asymmetry range of the asymptotic behavior of  $f_1^T(\omega)$  given by ref. [115] using the Light-Cone-Expansion, the Quark Model [114] used in ref. [103], the hard-wall model of ref. [105] and this work.

been used for computing the  $a_\mu^{\text{HLbL:T-poles}}$  fulfills the asymmetric asymptotic behavior predicted by ref. [115], being worse as the asymmetry approaches the single virtual case. However, noticeably, in the purely hadronic region where these contributions are computed, the high-energy behavior does not significantly influence the resulting contributions to  $a_\mu$  in any of the three considered approaches. The matching at  $Q_0 = 1.5$  GeV is shown in the left plot of Figure 5 where it can be seen, again, that the single virtual asymptotic behavior ( $\lambda = 0$ ) is not reproduced by any of the 3 models mentioned above. Near the symmetric ( $\lambda = 1$ ) asymptotic behavior, a better transition is achieved –as expected– for higher values of the matching scale  $Q_0 = 7.5$  GeV, as it can be appreciated in the right plot of Figure 5.

For the  $a_\mu$  computation, we use the master formula provided in ref. [139], using the optimized basis in ref. [87], which allows an evaluation of the tensor meson poles without kinematic singularities for specific cases, including the one where only  $\mathcal{F}_1^T$  is non-zero. In this way, we got the following results:

$$a_\mu^{\text{a2-pole}} = -1.09(10)_{\text{stat}}(^{+0.00}_{-0.11})_{\text{syst}} \times 10^{-11}, \quad (4.3a)$$




Figure 5: Transition from the purely hadronic to the asymptotic region as a function of the relative virtuality of both photons,  $\lambda$ , at the matching scale  $Q_0 = 1.5\text{GeV}$ (left) and  $5Q_0$ (right) for the 3 models considered and the Light-Cone-Expansion. Our one  $\sigma$  uncertainties are displayed by the green band.

$$a_\mu^{\text{f}_2\text{-pole}} = -3.4(3)_{\text{stat}}^{(+0.4)}_{(-0.0)} \text{syst} \times 10^{-11}, \quad (4.3b)$$

$$a_\mu^{\text{f}'_2\text{-pole}} = -0.046(14)_{\text{stat}}^{(+0.008)}_{(-0.000)} \text{syst} \times 10^{-11}, \quad (4.3c)$$

$$a_\mu^{a_2 + f_2 + f'_2\text{-pole}} = (-4.5^{+0.5}_{-0.3}) \times 10^{-11}. \quad (4.3d)$$

## 4.1 Error Computation

The statistical errors come from the propagation of the uncertainties on all parameters used to compute the  $a_\mu$  contributions, whereas the systematic ones come from two main sources: first concerning the specific value of  $M_T^3$  in eq. (3.4a); second, fulfilling all known SDCs being impossible in our setting.

### 4.1.1 Choice of $M_T$

The short-distance limits of the FFs can be imposed only to one of the three particles (either  $a_2$ ,  $f_2$  or  $f'_2$ ) which causes a factor of  $M_{T_{\text{choice}}}^2$  multiplying the asymptotic term of eq. (3.5). We used the respective mass for each tensor meson instead of choosing a common one for all 3. To compute the systematic error induced by this choice, we varied this parameter among 4 options: the 3 particle masses and their isospin average.

### 4.1.2 High-energy behavior

Our  $R\chi T$  result for the  $T$ -TFF, eq. (3.5), does fulfill the two constraints from eq. (3.4); however, the limiting case when only one of the  $Q_i$  is asymptotic and the other one is finite (but non-vanishing), should scale as  $1/Q^4$ , while it goes as  $1/Q^2$ . To estimate how this mismatch induces an error in our computation of  $a_\mu^{\text{HLbL:T-pole}}$ , we construct a Canterbury Approximant in the same fashion as in ref. [24]. For this purpose, the minimal choice is a  $C_2^0(Q_1^2, Q_2^2)$ :

$$\mathcal{F}_1^T(Q_1^2, Q_2^2) = \frac{\mathcal{F}_1^T(0, 0)}{1 + \beta_1^T(Q_1^2 + Q_2^2) + \beta_{1,1}^T Q_1^2 Q_2^2 + \beta_2^T(Q_1^4 + Q_2^4)}, \quad (4.4)$$

where we omit the terms with  $\beta_{1,2}^T$  and  $\beta_{2,2}^T$  coefficients, as they contribute to orders  $1/Q^6$  and  $1/Q^8$  in the asymptotic behavior. All single virtual, double virtual, and mixed asymptotic behaviors can be reproduced by the condition:

$$\beta_2^T = \frac{14\mathcal{F}_1^T(0, 0) - 3\beta_{1,1}^T F_T^{\text{eff}} M_T^3}{6F_T^{\text{eff}} M_T^3}. \quad (4.5)$$

The 2 remaining constants,  $\beta_1^T$  and  $\beta_{1,1}^T$ , will be fixed by matching the first two low-energy-expansion constants from our model and the considered  $C_2^0(Q_1^2, Q_2^2)$ , eq. (4.4):

$$\frac{\mathcal{F}_1^T(Q_1^2, Q_2^2)}{\mathcal{F}_1^T(0, 0)} \approx 1 - \frac{a_1^T}{M_T^2}(Q_1^2 + Q_2^2) + \frac{b_{1,1}^T}{M_T^4} Q_1^2 Q_2^2 + \dots, \quad (4.6)$$

leading to the following relations:

$$\beta_1^T = \frac{1}{M_\Lambda^2} = \frac{1}{M_V^2} + \frac{1}{M_{V'}^2}, \quad (4.7a)$$

$$\beta_{1,1}^T = \frac{1}{M_{\Lambda'}^4} = \frac{14\sqrt{2}C_{TVV}F_V^2(M_V^4 - M_{V'}^4)^2 - 9F_T^{\text{eff}} M_T^2 M_V^4 (M_V^4 + 2M_V^2 M_{V'}^2 + 2M_{V'}^4)}{M_V^4 M_{V'}^4 [14\sqrt{2}C_{TVV}F_V^2(M_V^2 - M_{V'}^2)^2 - 9F_T^{\text{eff}} M_T^2 M_V^4]}, \quad (4.7b)$$

where we introduced the two scales  $M_\Lambda \sim M_{\Lambda'} \sim M_\rho$ . We compute this error by comparing the  $a_\mu^{\text{T-pole}}$  value of our model to the results obtained in the  $C_2^0$  model, where both the asymptotic behavior and the normalization are correct. This indicates that the magnitude of  $a_\mu^{a_2-\text{pole}}$  is underestimated, while the contrary happens for the  $f_2$  and  $f'_2$  pole contributions. This gives the dominant (asymmetric) error of our results.

Logically, our result agrees in sign with the dispersive computation,  $a_\mu^{a_2+f_2+f'_2} = -2.5(3) \times 10^{-11}$  [103]. A comparison of our  $\mathcal{F}_1^T$  in  $R\chi T$  with the one of the simple quark model [114] used in the dispersive determination [103], Figure 2, shows us that in the

purely hadronic region ( $Q_i < 1.5\text{GeV}$ ) <sup>2</sup> the TFF from R $\chi$ T is larger than the quark model one, which explains why we get a larger magnitude for  $a_\mu$ . Noteworthy, they are in very nice agreement with the holographic ones, once the  $\mathcal{F}_3^T$  contribution (responsible for the sign change) has been discarded ,  $a_\mu^{a_2+f_2+f'_2} = -4.6(6) \times 10^{-11}$  [105] –which is clearly explained by the agreement of the  $\mathcal{F}_1^T$  between these two models in Figure 2 at both single virtual and symmetrically double virtual cases–. This result should be helpful for the update of the White Paper (to appear soon), which will include the revised  $a_\mu^{\text{T-pole}}$  contribution.

## 5 Conclusions

The FNAL Muon g-2 Collaboration will soon release its final measurement of this precision observable, with potentially crucial beyond the Standard Model implications. This has triggered and boosted the efforts of the Muon  $g - 2$  Theory Initiative, towards reducing the uncertainty of the Standard Model prediction, aiming to maximize the reach on new physics of this measurement. Within this global effort, not only the leading uncertainties, of hadronic origin, coming from the vacuum polarization piece, but also the subleading ones, coming from the light-by-light part, have been improved recently.

In this work we have focused on the tensor-pole contribution, in which the most recent evaluations, dispersive and holographic, differ in sign. We have studied it, within Resonance Chiral Theory, and found that only one form factor ( $\mathcal{F}_1^T$ ) is non-vanishing, in agreement with the expectations coming from the chiral expansion and from the quark model results. This makes this contribution negative (the  $\mathcal{F}_3^T$  contribution is responsible for its positiveness in holographic QCD). We have compared our  $\mathcal{F}_1^T$  to both computations and found closer agreement with the holographic form factor. Our results (in units of  $10^{-11}$ ):  $a_\mu^{\text{a}_2\text{-pole}} = -1.09(10)_{\text{stat}}^{(+0.0)}_{(-0.11)}_{\text{syst}}$ ,  $a_\mu^{\text{f}_2\text{-pole}} = -3.4(3)_{\text{stat}}^{(+0.4)}_{(-0.0)}_{\text{syst}}$  and  $a_\mu^{\text{f}'_2\text{-pole}} = -0.046(14)_{\text{stat}}^{(+0.008)}_{(-0.000)}_{\text{syst}}$ , adding up to  $a_\mu^{(\text{a}_2+\text{f}_2+\text{f}'_2)\text{-pole}} = (-4.5^{+0.5})_{-0.3}$ , agree very well with the holographic computation if  $\mathcal{F}_3^T$  is set to zero,  $-4.6(6) \times 10^{-11}$  [105], and exceed in magnitude the outcome of the dispersive analysis,  $-2.5(3) \times 10^{-11}$  [103].

---

<sup>2</sup>The contribution to  $a_\mu^{\text{T-pole}}$  is evaluated in this region, as it has become standard in the matching to the OPE results [23, 28, 140–145].

## Acknowledgements

The authors are very grateful to Antonio Rodríguez-Sánchez, Martin Hoferichter, and Anton Rebhan for enlightening feedback and insightful comments. The authors thank partial funding from SECIHTI (México).

## References

- [1] MUON G-2 collaboration, *Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL*, *Phys. Rev. D* **73** (2006) 072003 [[hep-ex/0602035](#)].
- [2] MUON G-2 collaboration, *Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm*, *Phys. Rev. Lett.* **126** (2021) 141801 [[2104.03281](#)].
- [3] MUON G-2 collaboration, *Measurement of the Positive Muon Anomalous Magnetic Moment to 0.20 ppm*, *Phys. Rev. Lett.* **131** (2023) 161802 [[2308.06230](#)].
- [4] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *Phys. Rev. D* **110** (2024) 030001.
- [5] T. Aoyama et al., *The anomalous magnetic moment of the muon in the Standard Model*, *Phys. Rept.* **887** (2020) 1 [[2006.04822](#)].
- [6] G. Colangelo et al., *Prospects for precise predictions of  $a_\mu$  in the Standard Model*, [2203.15810](#).
- [7] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *Reevaluation of the hadronic vacuum polarisation contributions to the Standard Model predictions of the muon  $g - 2$  and  $\alpha(m_Z^2)$  using newest hadronic cross-section data*, *Eur. Phys. J. C* **77** (2017) 827 [[1706.09436](#)].
- [8] A. Keshavarzi, D. Nomura and T. Teubner, *Muon  $g - 2$  and  $\alpha(M_Z^2)$ : a new data-based analysis*, *Phys. Rev. D* **97** (2018) 114025 [[1802.02995](#)].
- [9] G. Colangelo, M. Hoferichter and P. Stoffer, *Two-pion contribution to hadronic vacuum polarization*, *JHEP* **02** (2019) 006 [[1810.00007](#)].

- [10] M. Hoferichter, B.-L. Hoid and B. Kubis, *Three-pion contribution to hadronic vacuum polarization*, *JHEP* **08** (2019) 137 [[1907.01556](#)].
- [11] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *A new evaluation of the hadronic vacuum polarisation contributions to the muon anomalous magnetic moment and to  $\alpha(m_Z^2)$* , *Eur. Phys. J. C* **80** (2020) 241 [[1908.00921](#)].
- [12] A. Keshavarzi, D. Nomura and T. Teubner,  *$g - 2$  of charged leptons,  $\alpha(M_Z^2)$ , and the hyperfine splitting of muonium*, *Phys. Rev. D* **101** (2020) 014029 [[1911.00367](#)].
- [13] A. Kurz, T. Liu, P. Marquard and M. Steinhauser, *Hadronic contribution to the muon anomalous magnetic moment to next-to-next-to-leading order*, *Phys. Lett. B* **734** (2014) 144 [[1403.6400](#)].
- [14] FERMILAB LATTICE, LATTICE-HPQCD, MILC collaboration, *Strong-Isospin-Breaking Correction to the Muon Anomalous Magnetic Moment from Lattice QCD at the Physical Point*, *Phys. Rev. Lett.* **120** (2018) 152001 [[1710.11212](#)].
- [15] BUDAPEST-MARSEILLE-WUPPERTAL collaboration, *Hadronic vacuum polarization contribution to the anomalous magnetic moments of leptons from first principles*, *Phys. Rev. Lett.* **121** (2018) 022002 [[1711.04980](#)].
- [16] RBC, UKQCD collaboration, *Calculation of the hadronic vacuum polarization contribution to the muon anomalous magnetic moment*, *Phys. Rev. Lett.* **121** (2018) 022003 [[1801.07224](#)].
- [17] D. Giusti, V. Lubicz, G. Martinelli, F. Sanfilippo and S. Simula, *Electromagnetic and strong isospin-breaking corrections to the muon  $g - 2$  from Lattice QCD+QED*, *Phys. Rev. D* **99** (2019) 114502 [[1901.10462](#)].
- [18] PACS collaboration, *Hadronic vacuum polarization contribution to the muon  $g - 2$  with 2+1 flavor lattice QCD on a larger than  $(10 \text{ fm})^4$  lattice at the physical point*, *Phys. Rev. D* **100** (2019) 034517 [[1902.00885](#)].
- [19] FERMILAB LATTICE, LATTICE-HPQCD, MILC collaboration, *Hadronic-vacuum-polarization contribution to the muon's anomalous magnetic moment from four-flavor lattice QCD*, *Phys. Rev. D* **101** (2020) 034512 [[1902.04223](#)].

- [20] A. Gérardin, M. Cè, G. von Hippel, B. Hörz, H.B. Meyer, D. Mohler et al., *The leading hadronic contribution to  $(g - 2)_\mu$  from lattice QCD with  $N_f = 2 + 1$  flavours of  $O(a)$  improved Wilson quarks*, *Phys. Rev. D* **100** (2019) 014510 [[1904.03120](#)].
- [21] C. Aubin, T. Blum, C. Tu, M. Golterman, C. Jung and S. Peris, *Light quark vacuum polarization at the physical point and contribution to the muon  $g - 2$* , *Phys. Rev. D* **101** (2020) 014503 [[1905.09307](#)].
- [22] D. Giusti and S. Simula, *Lepton anomalous magnetic moments in Lattice QCD+QED*, *PoS LATTICE2019* (2019) 104 [[1910.03874](#)].
- [23] K. Melnikov and A. Vainshtein, *Hadronic light-by-light scattering contribution to the muon anomalous magnetic moment revisited*, *Phys. Rev. D* **70** (2004) 113006 [[hep-ph/0312226](#)].
- [24] P. Masjuan and P. Sánchez-Puertas, *Pseudoscalar-pole contribution to the  $(g_\mu - 2)$ : a rational approach*, *Phys. Rev. D* **95** (2017) 054026 [[1701.05829](#)].
- [25] G. Colangelo, M. Hoferichter, M. Procura and P. Stoffer, *Dispersion relation for hadronic light-by-light scattering: two-pion contributions*, *JHEP* **04** (2017) 161 [[1702.07347](#)].
- [26] M. Hoferichter, B.-L. Hoid, B. Kubis, S. Leupold and S.P. Schneider, *Dispersion relation for hadronic light-by-light scattering: pion pole*, *JHEP* **10** (2018) 141 [[1808.04823](#)].
- [27] A. Gérardin, H.B. Meyer and A. Nyffeler, *Lattice calculation of the pion transition form factor with  $N_f = 2 + 1$  Wilson quarks*, *Phys. Rev. D* **100** (2019) 034520 [[1903.09471](#)].
- [28] J. Bijnens, N. Hermansson-Truedsson and A. Rodríguez-Sánchez, *Short-distance constraints for the HLbL contribution to the muon anomalous magnetic moment*, *Phys. Lett. B* **798** (2019) 134994 [[1908.03331](#)].
- [29] G. Colangelo, F. Hagelstein, M. Hoferichter, L. Laub and P. Stoffer, *Longitudinal short-distance constraints for the hadronic light-by-light contribution to  $(g - 2)_\mu$  with large- $N_c$  Regge models*, *JHEP* **03** (2020) 101 [[1910.13432](#)].
- [30] V. Pauk and M. Vanderhaeghen, *Single meson contributions to the muon's anomalous magnetic moment*, *Eur. Phys. J. C* **74** (2014) 3008 [[1401.0832](#)].

- [31] I. Danilkin and M. Vanderhaeghen, *Light-by-light scattering sum rules in light of new data*, *Phys. Rev. D* **95** (2017) 014019 [[1611.04646](#)].
- [32] F. Jegerlehner, *The Anomalous Magnetic Moment of the Muon*, vol. 274, Springer, Cham (2017), [10.1007/978-3-319-63577-4](#).
- [33] M. Knecht, S. Narison, A. Rabemananjara and D. Rabetiarivony, *Scalar meson contributions to a  $\mu$  from hadronic light-by-light scattering*, *Phys. Lett. B* **787** (2018) 111 [[1808.03848](#)].
- [34] G. Eichmann, C.S. Fischer and R. Williams, *Kaon-box contribution to the anomalous magnetic moment of the muon*, *Phys. Rev. D* **101** (2020) 054015 [[1910.06795](#)].
- [35] P. Roig and P. Sánchez-Puertas, *Axial-vector exchange contribution to the hadronic light-by-light piece of the muon anomalous magnetic moment*, *Phys. Rev. D* **101** (2020) 074019 [[1910.02881](#)].
- [36] G. Colangelo, M. Hoferichter, A. Nyffeler, M. Passera and P. Stoffer, *Remarks on higher-order hadronic corrections to the muon  $g-2$* , *Phys. Lett. B* **735** (2014) 90 [[1403.7512](#)].
- [37] T. Blum, N. Christ, M. Hayakawa, T. Izubuchi, L. Jin, C. Jung et al., *Hadronic Light-by-Light Scattering Contribution to the Muon Anomalous Magnetic Moment from Lattice QCD*, *Phys. Rev. Lett.* **124** (2020) 132002 [[1911.08123](#)].
- [38] T. Aoyama, M. Hayakawa, T. Kinoshita and M. Nio, *Complete Tenth-Order QED Contribution to the Muon  $g-2$* , *Phys. Rev. Lett.* **109** (2012) 111808 [[1205.5370](#)].
- [39] T. Aoyama, T. Kinoshita and M. Nio, *Theory of the Anomalous Magnetic Moment of the Electron*, *Atoms* **7** (2019) 28.
- [40] A. Czarnecki, W.J. Marciano and A. Vainshtein, *Refinements in electroweak contributions to the muon anomalous magnetic moment*, *Phys. Rev. D* **67** (2003) 073006 [[hep-ph/0212229](#)].
- [41] C. Gnendiger, D. Stöckinger and H. Stöckinger-Kim, *The electroweak contributions to  $(g-2)_\mu$  after the Higgs boson mass measurement*, *Phys. Rev. D* **88** (2013) 053005 [[1306.5546](#)].

- [42] S. Borsanyi et al., *Leading hadronic contribution to the muon magnetic moment from lattice QCD*, *Nature* **593** (2021) 51 [[2002.12347](#)].
- [43] C. Lehner and A.S. Meyer, *Consistency of hadronic vacuum polarization between lattice QCD and the R-ratio*, *Phys. Rev. D* **101** (2020) 074515 [[2003.04177](#)].
- [44] M. Knecht, *On some short-distance properties of the fourth-rank hadronic vacuum polarization tensor and the anomalous magnetic moment of the muon*, *JHEP* **08** (2020) 056 [[2005.09929](#)].
- [45] P. Masjuan, P. Roig and P. Sánchez-Puertas, *The interplay of transverse degrees of freedom and axial-vector mesons with short-distance constraints in  $g - 2$* , *J. Phys. G* **49** (2022) 015002 [[2005.11761](#)].
- [46] J. Lüdtke and M. Procura, *Effects of longitudinal short-distance constraints on the hadronic light-by-light contribution to the muon  $g - 2$* , *Eur. Phys. J. C* **80** (2020) 1108 [[2006.00007](#)].
- [47] E.-H. Chao, A. Gérardin, J.R. Green, R.J. Hudspith and H.B. Meyer, *Hadronic light-by-light contribution to  $(g - 2)_\mu$  from lattice QCD with  $SU(3)$  flavor symmetry*, *Eur. Phys. J. C* **80** (2020) 869 [[2006.16224](#)].
- [48] J.A. Miranda and P. Roig, *New  $\tau$ -based evaluation of the hadronic contribution to the vacuum polarization piece of the muon anomalous magnetic moment*, *Phys. Rev. D* **102** (2020) 114017 [[2007.11019](#)].
- [49] B.-L. Hoid, M. Hoferichter and B. Kubis, *Hadronic vacuum polarization and vector-meson resonance parameters from  $e^+e^- \rightarrow \pi^0\gamma$* , *Eur. Phys. J. C* **80** (2020) 988 [[2007.12696](#)].
- [50] G. Colangelo, M. Hoferichter and P. Stoffer, *Constraints on the two-pion contribution to hadronic vacuum polarization*, *Phys. Lett. B* **814** (2021) 136073 [[2010.07943](#)].
- [51] E.-H. Chao, R.J. Hudspith, A. Gérardin, J.R. Green, H.B. Meyer and K. Ott nad, *Hadronic light-by-light contribution to  $(g - 2)_\mu$  from lattice QCD: a complete calculation*, *Eur. Phys. J. C* **81** (2021) 651 [[2104.02632](#)].
- [52] I. Danilkin, M. Hoferichter and P. Stoffer, *A dispersive estimate of scalar contributions to hadronic light-by-light scattering*, *Phys. Lett. B* **820** (2021) 136502 [[2105.01666](#)].

- [53] G. Colangelo, F. Hagelstein, M. Hoferichter, L. Laub and P. Stoffer, *Short-distance constraints for the longitudinal component of the hadronic light-by-light amplitude: an update*, *Eur. Phys. J. C* **81** (2021) 702 [[2106.13222](#)].
- [54] J. Leutgeb and A. Rebhan, *Hadronic light-by-light contribution to the muon g-2 from holographic QCD with massive pions*, *Phys. Rev. D* **104** (2021) 094017 [[2108.12345](#)].
- [55] G. Colangelo, M. Hoferichter, B. Kubis, M. Niehus and J.R. de Elvira, *Chiral extrapolation of hadronic vacuum polarization*, *Phys. Lett. B* **825** (2022) 136852 [[2110.05493](#)].
- [56] L. Cappiello, O. Catà and G. D'Ambrosio, *Scalar resonances in the hadronic light-by-light contribution to the muon (g-2)*, *Phys. Rev. D* **105** (2022) 056020 [[2110.05962](#)].
- [57] D. Giusti and S. Simula, *Window contributions to the muon hadronic vacuum polarization with twisted-mass fermions*, *PoS LATTICE2021* (2022) 189 [[2111.15329](#)].
- [58] M. Hoferichter and T. Teubner, *Mixed Leptonic and Hadronic Corrections to the Anomalous Magnetic Moment of the Muon*, *Phys. Rev. Lett.* **128** (2022) 112002 [[2112.06929](#)].
- [59] A. Miramontes, A. Bashir, K. Raya and P. Roig, *Pion and Kaon box contribution to  $a_\mu^{HLbL}$* , *Phys. Rev. D* **105** (2022) 074013 [[2112.13916](#)].
- [60] K. Raya, A. Bashir and P. Roig, *Contribution of neutral pseudoscalar mesons to  $a_\mu^{HLbL}$  within a Schwinger-Dyson equations approach to QCD*, *Phys. Rev. D* **101** (2020) 074021 [[1910.05960](#)].
- [61] D. Stamen, D. Hariharan, M. Hoferichter, B. Kubis and P. Stoffer, *Kaon electromagnetic form factors in dispersion theory*, *Eur. Phys. J. C* **82** (2022) 432 [[2202.11106](#)].
- [62] D. Boito, M. Golterman, K. Maltman and S. Peris, *Evaluation of the three-flavor quark-disconnected contribution to the muon anomalous magnetic moment from experimental data*, *Phys. Rev. D* **105** (2022) 093003 [[2203.05070](#)].
- [63] CHIQCD collaboration, *Muon g-2 with overlap valence fermions*, *Phys. Rev. D* **107** (2023) 034513 [[2204.01280](#)].

- [64] C. Aubin, T. Blum, M. Golterman and S. Peris, *Muon anomalous magnetic moment with staggered fermions: Is the lattice spacing small enough?*, *Phys. Rev. D* **106** (2022) 054503 [[2204.12256](#)].
- [65] G. Colangelo, A.X. El-Khadra, M. Hoferichter, A. Keshavarzi, C. Lehner, P. Stoffer et al., *Data-driven evaluations of Euclidean windows to scrutinize hadronic vacuum polarization*, *Phys. Lett. B* **833** (2022) 137313 [[2205.12963](#)].
- [66] M. Cè et al., *Window observable for the hadronic vacuum polarization contribution to the muon g-2 from lattice QCD*, *Phys. Rev. D* **106** (2022) 114502 [[2206.06582](#)].
- [67] EXTENDED TWISTED MASS collaboration, *Lattice calculation of the short and intermediate time-distance hadronic vacuum polarization contributions to the muon magnetic moment using twisted-mass fermions*, *Phys. Rev. D* **107** (2023) 074506 [[2206.15084](#)].
- [68] G. Colangelo, M. Hoferichter, B. Kubis and P. Stoffer, *Isospin-breaking effects in the two-pion contribution to hadronic vacuum polarization*, *JHEP* **10** (2022) 032 [[2208.08993](#)].
- [69] V. Biloshytskyi, E.-H. Chao, A. Gérardin, J.R. Green, F. Hagelstein, H.B. Meyer et al., *Forward light-by-light scattering and electromagnetic correction to hadronic vacuum polarization*, *JHEP* **03** (2023) 194 [[2209.02149](#)].
- [70] D. Boito, M. Golterman, K. Maltman and S. Peris, *Data-based determination of the isospin-limit light-quark-connected contribution to the anomalous magnetic moment of the muon*, *Phys. Rev. D* **107** (2023) 074001 [[2211.11055](#)].
- [71] J. Leutgeb, J. Mager and A. Rebhan, *Hadronic light-by-light contribution to the muon g-2 from holographic QCD with solved U(1) $A$  problem*, *Phys. Rev. D* **107** (2023) 054021 [[2211.16562](#)].
- [72] CMD-3 collaboration, *Measurement of the  $e+e-\rightarrow\pi^+\pi^-$  cross section from threshold to 1.2 GeV with the CMD-3 detector*, *Phys. Rev. D* **109** (2024) 112002 [[2302.08834](#)].
- [73] CMD-3 collaboration, *Measurement of the Pion Form Factor with CMD-3 Detector and its Implication to the Hadronic Contribution to Muon (g-2)*, *Phys. Rev. Lett.* **132** (2024) 231903 [[2309.12910](#)].

- [74] M. Hoferichter, B.-L. Hoid, B. Kubis and D. Schuh, *Isospin-breaking effects in the three-pion contribution to hadronic vacuum polarization*, *JHEP* **08** (2023) 208 [[2307.02546](#)].
- [75] P. Colangelo, F. Giannuzzi and S. Nicotri,  *$\pi 0, \eta, \eta'$  two-photon transition form factors in the holographic soft-wall model and contributions to  $(g-2)_\mu$* , *Phys. Lett. B* **840** (2023) 137878 [[2301.06456](#)].
- [76] FERMILAB LATTICE, HPQCD,, MILC collaboration, *Light-quark connected intermediate-window contributions to the muon  $g-2$  hadronic vacuum polarization from lattice QCD*, *Phys. Rev. D* **107** (2023) 114514 [[2301.08274](#)].
- [77] RBC, UKQCD collaboration, *Update of Euclidean windows of the hadronic vacuum polarization*, *Phys. Rev. D* **108** (2023) 054507 [[2301.08696](#)].
- [78] S.-J. Wang, Z. Fang and L.-Y. Dai, *Two body final states production in electron-positron annihilation and their contributions to  $(g-2)_\mu$* , *JHEP* **07** (2023) 037 [[2302.08859](#)].
- [79] J. Lüdtke, M. Procura and P. Stoffer, *Dispersion relations for hadronic light-by-light scattering in triangle kinematics*, *JHEP* **04** (2023) 125 [[2302.12264](#)].
- [80] R. Escribano, J.A. Miranda and P. Roig, *Radiative corrections to the  $\tau \rightarrow (P1P2) - v\tau$  ( $P1,2 = \pi, K$ ) decays*, *Phys. Rev. D* **109** (2024) 053003 [[2303.01362](#)].
- [81] RBC, UKQCD collaboration, *Hadronic light-by-light contribution to the muon anomaly from lattice QCD with infinite volume QED at physical pion mass*, *Phys. Rev. D* **111** (2025) 014501 [[2304.04423](#)].
- [82] P. Masjuan, A. Miranda and P. Roig,  *$\tau$  data-driven evaluation of Euclidean windows for the hadronic vacuum polarization*, *Phys. Lett. B* **850** (2024) 138492 [[2305.20005](#)].
- [83] G. Benton, D. Boito, M. Golterman, A. Keshavarzi, K. Maltman and S. Peris, *Data-Driven Determination of the Light-Quark Connected Component of the Intermediate-Window Contribution to the Muon  $g-2$* , *Phys. Rev. Lett.* **131** (2023) 251803 [[2306.16808](#)].

- [84] M. Hoferichter, G. Colangelo, B.-L. Hoid, B. Kubis, J.R. de Elvira, D. Schuh et al., *Phenomenological Estimate of Isospin Breaking in Hadronic Vacuum Polarization*, *Phys. Rev. Lett.* **131** (2023) 161905 [[2307.02532](#)].
- [85] M. Davier, Z. Fodor, A. Gerardin, L. Lellouch, B. Malaescu, F.M. Stokes et al., *Hadronic vacuum polarization: Comparing lattice QCD and data-driven results in systematically improvable ways*, *Phys. Rev. D* **109** (2024) 076019 [[2308.04221](#)].
- [86] M. Davier, A. Hoecker, A.-M. Lutz, B. Malaescu and Z. Zhang, *Tensions in  $e^+e^- \rightarrow \pi^+\pi^-(\gamma)$  measurements: the new landscape of data-driven hadronic vacuum polarization predictions for the muon  $g - 2$* , *Eur. Phys. J. C* **84** (2024) 721 [[2312.02053](#)].
- [87] M. Hoferichter, P. Stoffer and M. Zillinger, *An optimized basis for hadronic light-by-light scattering*, *JHEP* **04** (2024) 092 [[2402.14060](#)].
- [88] A. Boccaletti et al., *High precision calculation of the hadronic vacuum polarisation contribution to the muon anomaly*, [2407.10913](#).
- [89] S. Lahert, C. DeTar, A.X. El-Khadra, S. Gottlieb, A.S. Kronfeld and R.S. Van de Water, *The two-pion contribution to the hadronic vacuum polarization with staggered quarks*, [2409.00756](#).
- [90] G.L. Castro, A. Miranda and P. Roig, *Isospin breaking corrections in  $2\pi$  production in tau decays and  $e^+e^-$  annihilation: consequences for the muon  $g - 2$  and CVC tests*, [2411.07696](#).
- [91] M. Hoferichter, J. Lüdtke, L. Naterop, M. Procura and P. Stoffer, *An improved evaluation of the electroweak contribution to  $(g - 2)_\mu$* , [2503.04883](#).
- [92] A. Bazavov et al., *Hadronic vacuum polarization for the muon  $g - 2$  from lattice QCD: Long-distance and full light-quark connected contribution*, [2412.18491](#).
- [93] S. Holz, M. Hoferichter, B.-L. Hoid and B. Kubis, *Dispersion relation for hadronic light-by-light scattering:  $\eta$  and  $\eta'$  poles*, [2412.16281](#).
- [94] J. Leutgeb, J. Mager and A. Rebhan, *Superconnections in AdS/QCD and the hadronic light-by-light contribution to the muon  $g - 2$* , [2411.10432](#).
- [95] S. Holz, M. Hoferichter, B.-L. Hoid and B. Kubis, *A precision evaluation of the  $\eta$ - and  $\eta'$ -pole contributions to hadronic light-by-light scattering in the anomalous magnetic moment of the muon*, [2411.08098](#).

- [96] A.S. Miramontes, K. Raya, A. Bashir, P. Roig and G. Paredes-Torres, *Radially excited pion: electromagnetic form factor and the box contribution to the muon's  $g - 2$* , [2411.02218](#).
- [97] O. Deineka, I. Danilkin and M. Vanderhaeghen, *Dispersive estimate of the  $a_0(980)$  contribution to  $(g-2)\mu$* , *Phys. Rev. D* **111** (2025) 034009 [[2410.12894](#)].
- [98] E.J. Estrada, J.M. Márquez, D. Portillo-Sánchez and P. Roig, *Proton-box contribution to  $a_\mu^{\text{HLbL}}$* , [2411.07115](#).
- [99] S. Volkov, *Calculating the five-loop QED contribution to the electron anomalous magnetic moment: Graphs without lepton loops*, *Phys. Rev. D* **100** (2019) 096004 [[1909.08015](#)].
- [100] S. Volkov, *Calculation of the total 10th order QED contribution to the electron magnetic moment*, *Phys. Rev. D* **110** (2024) 036001 [[2404.00649](#)].
- [101] T. Aoyama, M. Hayakawa, A. Hirayama and M. Nio, *Verification of the tenth-order QED contribution to the anomalous magnetic moment of the electron from diagrams without fermion loops*, *Phys. Rev. D* **111** (2025) L031902 [[2412.06473](#)].
- [102] MUON G-2 collaboration, *Muon ( $g-2$ ) Technical Design Report*, [1501.06858](#).
- [103] M. Hoferichter, P. Stoffer and M. Zillinger, *Dispersion relation for hadronic light-by-light scattering: subleading contributions*, *JHEP* **02** (2025) 121 [[2412.00178](#)].
- [104] M. Hoferichter, P. Stoffer and M. Zillinger, *Complete Dispersive Evaluation of the Hadronic Light-by-Light Contribution to Muon  $g-2$* , *Phys. Rev. Lett.* **134** (2025) 061902 [[2412.00190](#)].
- [105] L. Cappiello, J. Leutgeb, J. Mager and A. Rebhan, *Tensor meson transition form factors in holographic QCD and the muon  $g - 2$* , [2501.09699](#).
- [106] J. Mager, L. Cappiello, J. Leutgeb and A. Rebhan, *Longitudinal short-distance constraints on hadronic light-by-light scattering and tensor meson contributions to the muon  $g - 2$* , [2501.19293](#).
- [107] P. Colangelo, F. Giannuzzi and S. Nicotri, *Hadronic light-by-light scattering contributions to  $(g-2)\mu$  from axial-vector and tensor mesons in the holographic soft-wall model*, *Phys. Rev. D* **109** (2024) 094036 [[2402.07579](#)].

- [108] V. Pascalutsa, V. Pauk and M. Vanderhaeghen, *Light-by-light scattering sum rules constraining meson transition form factors*, *Phys. Rev. D* **85** (2012) 116001 [[1204.0740](#)].
- [109] G. Ecker, J. Gasser, A. Pich and E. de Rafael, *The Role of Resonances in Chiral Perturbation Theory*, *Nucl. Phys. B* **321** (1989) 311.
- [110] G. Ecker, J. Gasser, H. Leutwyler, A. Pich and E. de Rafael, *Chiral Lagrangians for Massive Spin 1 Fields*, *Phys. Lett. B* **223** (1989) 425.
- [111] G. Ecker and C. Zauner, *Tensor meson exchange at low energies*, *Eur. Phys. J. C* **52** (2007) 315 [[0705.0624](#)].
- [112] S. Bellucci, J. Gasser and M.E. Sainio, *Low-energy photon-photon collisions to two loop order*, *Nucl. Phys. B* **423** (1994) 80 [[hep-ph/9401206](#)].
- [113] C. Chen, N.-Q. Cheng, L.-W. Yan, C.-G. Duan and Z.-H. Guo, *Revisiting the tensor-meson nonet in resonance chiral theory*, *Phys. Rev. D* **108** (2023) 014002 [[2302.11316](#)].
- [114] G.A. Schuler, F.A. Berends and R. van Gulik, *Meson photon transition form-factors and resonance cross-sections in e+ e- collisions*, *Nucl. Phys. B* **523** (1998) 423 [[hep-ph/9710462](#)].
- [115] M. Hoferichter and P. Stoffer, *Asymptotic behavior of meson transition form factors*, *JHEP* **05** (2020) 159 [[2004.06127](#)].
- [116] S.J. Brodsky and G.R. Farrar, *Scaling Laws at Large Transverse Momentum*, *Phys. Rev. Lett.* **31** (1973) 1153.
- [117] S. Weinberg, *Phenomenological Lagrangians*, *Physica A* **96** (1979) 327.
- [118] J. Gasser and H. Leutwyler, *Chiral Perturbation Theory to One Loop*, *Annals Phys.* **158** (1984) 142.
- [119] J. Gasser and H. Leutwyler, *Chiral Perturbation Theory: Expansions in the Mass of the Strange Quark*, *Nucl. Phys. B* **250** (1985) 465.
- [120] G. 't Hooft, *A Planar Diagram Theory for Strong Interactions*, *Nucl. Phys. B* **72** (1974) 461.

- [121] V. Cirigliano, G. Ecker, M. Eidemüller, A. Pich and J. Portolés, *The  $< VAP >$  Green function in the resonance region*, *Phys. Lett. B* **596** (2004) 96 [[hep-ph/0404004](#)].
- [122] V. Cirigliano, G. Ecker, M. Eidemüller, R. Kaiser, A. Pich and J. Portolés, *Towards a consistent estimate of the chiral low-energy constants*, *Nucl. Phys. B* **753** (2006) 139 [[hep-ph/0603205](#)].
- [123] V. Cirigliano, G. Ecker, M. Eidemüller, R. Kaiser, A. Pich and J. Portolés, *The  $< SPP >$  Green function and  $SU(3)$  breaking in  $K(l3)$  decays*, *JHEP* **04** (2005) 006 [[hep-ph/0503108](#)].
- [124] L.-Y. Dai, J. Fuentes-Martín and J. Portolés, *Scalar-involved three-point Green functions and their phenomenology*, *Phys. Rev. D* **99** (2019) 114015 [[1902.10411](#)].
- [125] P.D. Ruiz-Femenía, A. Pich and J. Portolés, *Odd intrinsic parity processes within the resonance effective theory of QCD*, *JHEP* **07** (2003) 003 [[hep-ph/0306157](#)].
- [126] K. Kampf and J. Novotny, *Resonance saturation in the odd-intrinsic parity sector of low-energy QCD*, *Phys. Rev. D* **84** (2011) 014036 [[1104.3137](#)].
- [127] P. Roig and J.J. Sanz Cillero, *Consistent high-energy constraints in the anomalous QCD sector*, *Phys. Lett. B* **733** (2014) 158 [[1312.6206](#)].
- [128] P. Roig, A. Guevara and G. López Castro,  *$VV'P$  form factors in resonance chiral theory and the  $\pi - \eta - \eta'$  light-by-light contribution to the muon  $g - 2$* , *Phys. Rev. D* **89** (2014) 073016 [[1401.4099](#)].
- [129] A. Guevara, P. Roig and J.J. Sanz-Cillero, *Pseudoscalar pole light-by-light contributions to the muon ( $g - 2$ ) in Resonance Chiral Theory*, *JHEP* **06** (2018) 160 [[1803.08099](#)].
- [130] T. Kadavy, K. Kampf and J. Novotny, *On the three-point order parameters of chiral symmetry breaking*, *JHEP* **03** (2023) 118 [[2206.02579](#)].
- [131] E.J. Estrada, S. Gonzàlez-Solís, A. Guevara and P. Roig, *Improved  $\pi^0$ ,  $\eta$ ,  $\eta'$  transition form factors in resonance chiral theory and their  $a_\mu^{HLbL}$  contribution*, *JHEP* **12** (2024) 203 [[2409.10503](#)].

- [132] A. Pich and J. Portolés, *The Vector form-factor of the pion from unitarity and analyticity: A Model independent approach*, *Phys. Rev. D* **63** (2001) 093005 [[hep-ph/0101194](#)].
- [133] V. Cirigliano, G. Ecker and H. Neufeld, *Isospin violation and the magnetic moment of the muon*, *Phys. Lett. B* **513** (2001) 361 [[hep-ph/0104267](#)].
- [134] V. Cirigliano, G. Ecker and H. Neufeld, *Radiative tau decay and the magnetic moment of the muon*, *JHEP* **08** (2002) 002 [[hep-ph/0207310](#)].
- [135] B.-H. Qin, W. Qin and L.-Y. Dai, *Study of electron-positron annihilation into  $KK^-\pi$  within resonance chiral theory*, *Phys. Rev. D* **111** (2025) 034025 [[2403.14294](#)].
- [136] J. Bijnens, G. Colangelo and G. Ecker, *The Mesonic chiral Lagrangian of order  $p^{**6}$* , *JHEP* **02** (1999) 020 [[hep-ph/9902437](#)].
- [137] V.M. Braun, N. Kivel, M. Strohmaier and A.A. Vladimirov, *Electroproduction of tensor mesons in QCD*, *JHEP* **06** (2016) 039 [[1603.09154](#)].
- [138] BELLE collaboration, *Study of  $\pi^0$  pair production in single-tag two-photon collisions*, *Phys. Rev. D* **93** (2016) 032003 [[1508.06757](#)].
- [139] G. Colangelo, M. Hoferichter, M. Procura and P. Stoffer, *Dispersion relation for hadronic light-by-light scattering: theoretical foundations*, *JHEP* **09** (2015) 074 [[1506.01386](#)].
- [140] A. Vainshtein, *Perturbative and nonperturbative renormalization of anomalous quark triangles*, *Phys. Lett. B* **569** (2003) 187 [[hep-ph/0212231](#)].
- [141] M. Knecht, S. Peris, M. Perrottet and E. de Rafael, *New nonrenormalization theorems for anomalous three point functions*, *JHEP* **03** (2004) 035 [[hep-ph/0311100](#)].
- [142] J. Bijnens, N. Hermansson-Truedsson, L. Laub and A. Rodríguez-Sánchez, *Short-distance HLbL contributions to the muon anomalous magnetic moment beyond perturbation theory*, *JHEP* **10** (2020) 203 [[2008.13487](#)].
- [143] J. Bijnens, N. Hermansson-Truedsson, L. Laub and A. Rodríguez-Sánchez, *The two-loop perturbative correction to the  $(g-2)_\mu$  HLbL at short distances*, *JHEP* **04** (2021) 240 [[2101.09169](#)].

- [144] J. Bijnens, N. Hermansson-Truedsson and A. Rodríguez-Sánchez, *Constraints on the hadronic light-by-light in the Melnikov-Vainshtein regime*, *JHEP* **02** (2023) 167 [[2211.17183](#)].
- [145] J. Bijnens, N. Hermansson-Truedsson and A. Rodríguez-Sánchez, *Constraints on the hadronic light-by-light tensor in corner kinematics for the muon  $g - 2$* , *JHEP* **03** (2025) 094 [[2411.09578](#)].