

**SOME NORM INEQUALITIES FOR COMMUTATORS
GENERATED BY THE RIESZ POTENTIALS ON
HOMOGENEOUS VARIABLE EXPONENT
HERZ-MORREY-HARDY SPACES**

FERİT GÜRBÜZ

ABSTRACT. In harmonic analysis, the studies of inequalities of classical operators (= singular, maximal, Riesz potentials etc.) in various function spaces have a very important place. The maturation of many topics in the field of harmonic analysis, as a result of various needs and developments to respond to the problems of the time, has also led to the emergence of many studies and works on these topics. In [3], under some conditions, the boundedness of Riesz potential on homogeneous variable exponent Herz-Morrey-Hardy spaces has been given. Inspired by the work of [3], in this work, by the atomic decompositions, we obtain the boundedness of commutators generated by the Riesz potentials on homogeneous variable exponent Herz-Morrey-Hardy spaces.

1. INTRODUCTION

In the last two decades it has become clear that classical function spaces are no longer adequate to solve a number of contemporary problems that arise naturally in various mathematical models of applied sciences. Therefore, it has become necessary to introduce and study new nonstandard function spaces from various points of view. Therefore, the study of variable exponent function spaces has attracted considerable attention and has made significant progress in various branches of mathematics, including real analysis, partial differential equations (PDEs), and applied mathematics. These spaces are an important generalization of classical function spaces and were developed to study problems involving variable exponent functions. These spaces eliminate several boundednesses of classical function spaces and allow generalized differential and integral operators to be considered in a broader context.

Herz spaces were initially introduced by Herz [4] in his paper to investigate the absolute convergence of Fourier transforms. Subsequently, these spaces have found significant applications in diverse branches of applied mathematics. Moreover, Herz [4] gave a new class of space are called the Herz spaces, which characterized certain properties of some classical functions. Since Fefferman and Stein [2] gave a real variable characterization of Hardy space, many important achievements have been made around Hardy space and its characterization as the main representative.

In recent years, many Herz type spaces have emerged. One of the important problems on Herz type spaces is boundedness of some integral operators. In this work, only Herz-Morrey-Hardy spaces (which will be defined in the next section)

2020 *Mathematics Subject Classification.* Primary 46E35; Secondary 42B25, 42B35.

Key words and phrases. Riesz potential, variable exponent, commutator, homogeneous Morrey-Herz-Hardy space, atomic decomposition.

will be discussed. In this context, Herz-Morrey-Hardy spaces are the generalized version of Herz-Hardy spaces. We will obtain some new results in these spaces. In 2015, Xu and Yang [9] first introduced Herz-Morrey-Hardy spaces with variable exponents, secondly gave their atomic decompositions. In 2016, Xu and Yang [10] gave the molecular decomposition of variable exponent Herz-Morrey-Hardy spaces. In 2024, by using the atomic decomposition for the domain Hardy space, Gürbüz [3] showed the boundedness of Riesz potential on homogeneous variable exponent Herz-Morrey-Hardy space. As a continuation of [3], we will discuss the commutator in this space in the third part of this article.

2. BASIC DEFINITIONS AND PREVIOUS RESULTS

Before giving the main result of this work, we first recall some elementary facts, notations and necessary definitions that will be needed.

Define $\mathcal{P}(\mathbb{R}^n)$ to be the set of $p(\cdot) : \mathbb{R}^n \rightarrow [1, \infty)$ when $1 \leq p_- := \operatorname{ess\,inf}_{x \in \mathbb{R}^n} p(x)$ and $p_+ := \operatorname{ess\,sup}_{x \in \mathbb{R}^n} p(x) < \infty$. The exponent $p'(\cdot)$ means the conjugate of $p(\cdot)$, that is $\frac{1}{p(x)} + \frac{1}{p'(x)} = 1$ holds. Let $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$. Variable exponent Lebesgue space $L^{p(\cdot)}(\mathbb{R}^n)$ is defined by

$$\|f\|_{L^{p(\cdot)}(\mathbb{R}^n)} := \inf \left\{ \eta > 0 : \int_{\mathbb{R}^n} \left(\frac{|f(x)|}{\eta} \right)^{p(x)} dx \leq 1 \right\} < \infty.$$

Let $f \in L_{loc}(\mathbb{R}^n)$. The Hardy-Littlewood maximal operator \mathcal{M} is defined by

$$\mathcal{M}f(x) := \sup_{r>0} \frac{1}{r^n} \int_{B(x,r)} |f(y)| dy, \quad \forall x \in \mathbb{R}^n,$$

where and follows $B(x, r) = \{y \in \mathbb{R}^n : |x - y| < r\}$ is the open ball centered at x with radius r and $L_{loc}(\mathbb{R}^n)$ is the collection of all locally integrable functions on \mathbb{R}^n . $\mathcal{B}(\mathbb{R}^n)$ is the set of $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ satisfying the condition that \mathcal{M} is bounded on $L^{p(\cdot)}(\mathbb{R}^n)$ as $p(\cdot) \in \mathcal{B}(\mathbb{R}^n)$ in [1]. We claim that the letter C stands for a positive constant that, which may vary from line to line. The expression $f \lesssim g$ means $f \leq Cg$, and $f \approx g$ means $f \lesssim g \lesssim f$. Throughout this work, for simplicity, we denote $L^{p(\cdot)}(\mathbb{R}^n)$ by $L^{p(\cdot)}$ and similarly $B(x, r)$ by B . We will use the following results:

Let $\forall x, y \in \mathbb{R}^n$ and $C > 0$. If $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ satisfies the requirement given below

$$(2.1) \quad |p(y) - p(x)| \leq \frac{-C}{\ln(|x - y|)}, \quad |x - y| \leq \frac{1}{2}$$

$$(2.2) \quad |p(y) - p(x)| \leq \frac{C}{\ln(e + |x|)}, \quad |x| \leq |y|,$$

then $p(\cdot) \in \mathcal{B}(\mathbb{R}^n)$ in [6].

For $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$, $f \in L^{p(\cdot)}(\mathbb{R}^n)$ and $g \in L^{p'(\cdot)}(\mathbb{R}^n)$, Hölder's inequality on variable exponent Lebesgue spaces holds in the form

$$(2.3) \quad \left| \int_{\mathbb{R}^n} f(x) g(x) dx \right| \leq \int_{\mathbb{R}^n} |f(x) g(x)| dx \leq r_p \|f\|_{L^{p(\cdot)}} \|g\|_{L^{p'(\cdot)}}, \quad r_p = 1 + \frac{1}{p_-} - \frac{1}{p_+},$$

see Theorem 2.1 in [7].

For $0 < \beta < n$, the Riesz potential

$$I^\beta f(x) = \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-\beta}} dy$$

has many applications in harmonic analysis. For example, suppose $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ satisfies (2.1) and (2.2). Then,

$$\|I^\beta f\|_{L^{p_2(\cdot)}} \lesssim \|f\|_{L^{p_1(\cdot)}}$$

is valid such that $0 < \beta < \frac{n}{p_+}$ and $\frac{1}{p_2(\cdot)} = \frac{1}{p_1(\cdot)} - \frac{\beta}{n}$ (see [1]). In 2024, Gürbüz [3] obtained the boundedness of the operator I^β on homogeneous variable exponent Herz-Morrey-Hardy spaces under some conditions. In addition, with $b \in L_{loc}(\mathbb{R}^n)$ ve $0 < \beta < n$, in this work, we can mainly focus on the the commutator generated by I^β and b is defined as follows:

$$\begin{aligned} [b, I^\beta] f(x) &= b(x) I^\beta f(x) - I^\beta (bf)(x) \\ (2.4) \quad &= \int_{\mathbb{R}^n} \frac{[b(x) - b(y)]}{|x-y|^{n-\beta}} f(y) dy. \end{aligned}$$

A locally integrable function b is said to belong to bounded mean oscillation space ($BMO(\mathbb{R}^n)$), if it satisfies

$$\|b\|_{BMO} := \sup_B \frac{1}{|B|} \int_B |b(y) - b_B| dy < \infty,$$

where B denotes the ball centered at $x \in \mathbb{R}^n$ and radius of $r > 0$, and b_B denotes the average of b on B , that is, $b_B := |B|^{-1} \int_B b(t) dt$.

Let $b \in BMO(\mathbb{R}^n)$, $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ and for $j, i \in \mathbb{Z}$ with $j > i$, we have the following inequalities:

$$\begin{aligned} C^{-1} \|b\|_{BMO} &\leq \sup_{B: \text{Ball}} \frac{1}{\|\chi_B\|_{L^{p(\cdot)}}} \|(b - b_B)\chi_B\|_{L^{p(\cdot)}} \leq C \|b\|_{BMO} \\ (2.5) \quad &\|(b - b_{B_i})\chi_{B_j}\|_{L^{p(\cdot)}} \leq C(j-i) \|b\|_{BMO} \|\chi_{B_j}\|_{L^{p(\cdot)}} \end{aligned}$$

(see [6]).

Assume that $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ satisfies (2.1) and (2.2), $0 < \beta < \frac{n}{p_+}$, and define $p_2(\cdot)$ as $\frac{1}{p_2(\cdot)} = \frac{1}{p_1(\cdot)} - \frac{\beta}{n}$. Then,

$$(2.6) \quad \|[b, I^\beta] f\|_{L^{p_2(\cdot)}} \lesssim \|b\|_{BMO} \|f\|_{L^{p_1(\cdot)}}$$

is satisfied (see [6]).

When $p(\cdot) \in \mathcal{B}(\mathbb{R}^n)$, then

$$(2.7) \quad \frac{\|\chi_S\|_{L^{p(\cdot)}}}{\|\chi_B\|_{L^{p(\cdot)}}} \leq C \left(\frac{|S|}{|B|} \right)^{\delta_1}, \frac{\|\chi_S\|_{L^{p'(\cdot)}(\mathbb{R}^n)}}{\|\chi_B\|_{L^{p'(\cdot)}(\mathbb{R}^n)}} \leq C \left(\frac{|S|}{|B|} \right)^{\delta_2}, S \subset B,$$

and

$$(2.8) \quad \|\chi_B\|_{L^{p(\cdot)}} \|\chi_B\|_{L^{p'(\cdot)}} \leq C |B|,$$

were proved in [5], respectively.

By (2.7) and (2.8), we can conclude that

$$(2.9) \quad \|\chi_i\|_{L^q(\cdot)} \leq \|\chi_{B_i}\|_{L^q(\cdot)}.$$

Let $l \in \mathbb{Z}$, $B_l := \{x \in \mathbb{R}^n : |x| \leq 2^l\}$, $D_l := B_l \setminus B_{l-1}$, $\chi_l := \chi_{D_l}$. Assume that the symbol \mathbb{N}_0 denotes the set of all nonnegative integers. For any $m \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$, we define

$$\tilde{\chi}_m := \begin{cases} \chi_{D_m} & , \quad m \geq 1 \\ \chi_{B_0} & , \quad m = 0 \end{cases}.$$

Now let's give the definition of the homogeneous variable exponent Herz-Morrey spaces.

Definition 1. Let $0 < q \leq \infty$, $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ and $0 \leq \lambda < \infty$. Let $\alpha(\cdot)$ also be a bounded real-valued measurable function on \mathbb{R}^n (that is, $\alpha(\cdot) \in L^\infty(\mathbb{R}^n)$). Then, the homogeneous variable exponent Herz-Morrey space $M\dot{K}_{p(\cdot),\lambda}^{\alpha(\cdot),q}(\mathbb{R}^n)$ is defined by

$$M\dot{K}_{p(\cdot),\lambda}^{\alpha(\cdot),q}(\mathbb{R}^n) := \left\{ f \in L_{loc}^{p(\cdot)}(\mathbb{R}^n \setminus \{0\}) : \|f\|_{M\dot{K}_{p(\cdot),\lambda}^{\alpha(\cdot),q}(\mathbb{R}^n)} < \infty \right\},$$

where

$$\|f\|_{M\dot{K}_{p(\cdot),\lambda}^{\alpha(\cdot),q}(\mathbb{R}^n)} := \sup_{L \in \mathbb{Z}} 2^{-L\lambda} \left(\sum_{l=-\infty}^L \|2^{l\alpha(\cdot)} f \chi_l\|_{L^{p(\cdot)}}^q \right)^{\frac{1}{q}}$$

and when $q = \infty$, there is the usual modification.

Now before giving the definition of homogeneous variable exponent Herz-Morrey-Hardy space $HM\dot{K}_{p(\cdot),\lambda}^{\alpha(\cdot),q}(\mathbb{R}^n)$, we need to recall some notations.

We assume that $\mathcal{S}(\mathbb{R}^n)$ be the Schwartz space of all rapidly decreasing infinitely differentiable functions on \mathbb{R}^n , and $\mathcal{S}'(\mathbb{R}^n)$ be the dual space of $\mathcal{S}(\mathbb{R}^n)$. Assume $G_N f$ is the grand maximal function of f defined by

$$G_N f(x) := \sup_{\phi \in \mathcal{A}_N} |\phi_{\nabla}^*(f)(x)|, \quad x \in \mathbb{R}^n,$$

where \mathcal{A}_N and ϕ_{∇}^* are defined in [9], respectively.

In 2015, Xu and Yang [9] introduced the homogeneous variable exponent Herz-Morrey-Hardy spaces $HM\dot{K}_{p(\cdot),\lambda}^{\alpha(\cdot),q}(\mathbb{R}^n)$ as follows:

Definition 2. Let $\alpha(\cdot) \in L^\infty(\mathbb{R}^n)$, $0 < q \leq \infty$, $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$, $0 \leq \lambda < \infty$ and $N > n + 1$. Then, the homogeneous variable exponent Herz-Morrey-Hardy space $HM\dot{K}_{p(\cdot),\lambda}^{\alpha(\cdot),q}(\mathbb{R}^n)$ is defined by

$$HM\dot{K}_{p(\cdot),\lambda}^{\alpha(\cdot),q}(\mathbb{R}^n) := \left\{ f \in \mathcal{S}'(\mathbb{R}^n) : \|f\|_{HM\dot{K}_{p(\cdot),\lambda}^{\alpha(\cdot),q}(\mathbb{R}^n)} := \left\| \begin{array}{l} f \\ \|G_N f\|_{M\dot{K}_{p(\cdot),\lambda}^{\alpha(\cdot),q}(\mathbb{R}^n)} < \infty \end{array} \right\| \right\}.$$

We can say that $G_N f(x) \leq CMf(x)$ ($C > 0$) for $x \in \mathbb{R}^n$ (see Proposition in Page 57 in [8]).

Now, we give the notion of atoms.

Definition 3. Let $p(\cdot) \in \mathcal{P}(\mathbb{R}^n)$, $\alpha(\cdot) \in L^\infty(\mathbb{R}^n) \cap \mathcal{P}_0^{\log}(\mathbb{R}^n) \cap \mathcal{P}_\infty^{\log}(\mathbb{R}^n)$, and nonnegative integer $s \geq [\alpha_r - n\delta_2]$; here $\alpha_r = \alpha(0)$, if $r < 1$, and $\alpha_r = \alpha_\infty$, if $r \geq 1$, $n\delta_2 \leq \alpha_r < \infty$ and δ_2 as in (2.7).

(i) If a function a on \mathbb{R}^n satisfies

$$\begin{aligned} (1) \quad & \text{supp } a \subset B(0, 2^r) \\ (2) \quad & \|a\|_{L^{p(\cdot)}} \leq |B(0, 2^r)|^{-\frac{\alpha r}{n}} \\ (3) \quad & \int_{\mathbb{R}^n} a(x) x^\beta dx = 0, \quad |\beta| \leq s, \end{aligned}$$

then it is called a central $(\alpha(\cdot), p(\cdot))$ -atom.

(ii) If a function a on \mathbb{R}^n satisfies (2) and (3) above and condition given below:

$$(1) \quad \text{supp } a \subset B(0, 2^r), \quad r \geq 1,$$

then it is called a central $(\alpha(\cdot), p(\cdot))$ -atom of restricted type, where follows $\mathcal{P}_0^{\log}(\mathbb{R}^n)$ and $\mathcal{P}_\infty^{\log}(\mathbb{R}^n)$ are the set of log-Hölder continuous functions at origin and the set of log-Hölder continuous functions at infinity, respectively; for their definitions see [3].

3. MAIN RESULT

The following theorem is our main result.

Theorem 1. Let $0 < q_1 \leq q_2 < \infty$, $0 \leq \lambda < \infty$, $0 < \beta < n$ and $\alpha(\cdot) \in L^\infty(\mathbb{R}^n) \cap \mathcal{P}_0^{\log}(\mathbb{R}^n) \cap \mathcal{P}_\infty^{\log}(\mathbb{R}^n)$ such that $2\lambda \leq \alpha(\cdot)$, $\beta - n\delta_2 < \alpha(0)$, $\beta - n\delta_2 < \alpha_\infty < \infty$ with $\delta_1, \delta_2 \in (0, 1)$ satisfying (2.7). Let also $[b, I^\beta]$ be defined as in (2.4). Assume that $p(\cdot), p_1(\cdot), p_2(\cdot) \in \mathcal{P}(\mathbb{R}^n)$ satisfy (2.1) and (2.2) and define $p_1(\cdot)$ by $\frac{1}{p_1(\cdot)} = \frac{\beta}{n} - \frac{1}{p_2(\cdot)}$ for $1 < p_1(\cdot) < p_2(\cdot) < \infty$. if $b \in \|b\|_{BMO}$, then the following inequality holds:

$$(3.1) \quad \|[b, I^\beta](f)\|_{M\dot{K}_{p_2(\cdot), \lambda}^{\alpha(\cdot), q_2}(\mathbb{R}^n)} \lesssim \|b\|_{BMO} \|f\|_{HM\dot{K}_{p_1(\cdot), \lambda}^{\alpha(\cdot), q_1}(\mathbb{R}^n)}$$

for all $f \in HM\dot{K}_{p_1(\cdot), \lambda}^{\alpha(\cdot), q_1}(\mathbb{R}^n)$.

Proof. Let $f \in HM\dot{K}_{p_1(\cdot), \lambda}^{\alpha(\cdot), q_1}(\mathbb{R}^n)$. By Theorem 2 in [3], take $a_l = a \cdot \chi_l = a \cdot \chi_{B_l}$ for each $l \in \mathbb{Z}$, we have

$$f = \sum_{l=0}^{\infty} \lambda_l a = \sum_{l=-\infty}^{\infty} \lambda_l a_l,$$

where each a_l is a central $(\alpha(\cdot), q_1(\cdot))$ -atom with support contained in B_l , and

$$\|f\|_{HM\dot{K}_{p_1(\cdot), \lambda}^{\alpha(\cdot), q_1}(\mathbb{R}^n)} \approx \inf_{L \in \mathbb{Z}} \sup 2^{-L\lambda} \left(\sum_{l=-\infty}^L |\lambda_l|^{q_1} \right)^{\frac{1}{q_1}}.$$

For convenience, we show

$$\Psi = \sup_{L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=-\infty}^L |\lambda_l|^{q_1}.$$

By (2.8) in [3], we have

$$\| [b, I^\beta] (f) \|_{MK_{p_2(\cdot), \lambda}^{\alpha(\cdot), q_2}(\mathbb{R}^n)}^{q_1} \approx \max \left\{ \begin{array}{l} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \left(\sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \| [b, I^\beta] (f) \chi_k \|_{L^{p_2(\cdot)}}^{q_1} \right), \\ \sup_{L > 0, L \in \mathbb{Z}} \left[\begin{array}{l} 2^{-L\lambda q_1} \left(\sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \| [b, I^\beta] (f) \chi_k \|_{L^{p_2(\cdot)}}^{q_1} \right) \\ + 2^{-L\lambda q_1} \left(\sum_{k=0}^L 2^{kq_1\alpha_\infty} \| [b, I^\beta] (f) \chi_k \|_{L^{p_2(\cdot)}}^{q_1} \right) \end{array} \right] \end{array} \right\} \\ \lesssim \{X, Y + Z\},$$

where

$$\begin{aligned} X &: = \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \left(\sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \| [b, I^\beta] (f) \chi_k \|_{L^{p_2(\cdot)}}^{q_1} \right) \\ Y &: = \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \| [b, I^\beta] (f) \chi_k \|_{L^{p_2(\cdot)}}^{q_1} \\ Z &: = \sup_{L > 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \left(\sum_{k=0}^L 2^{kq_1\alpha_\infty} \| [b, I^\beta] (f) \chi_k \|_{L^{p_2(\cdot)}}^{q_1} \right). \end{aligned}$$

To complete the proof of main result, we only need prove that $X, Y, Z \lesssim \Psi$. To do this we estimate X, Y, Z step by step. Indeed, we have

$$\begin{aligned} X &= \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \left(\sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \| [b, I^\beta] (f) \chi_k \|_{L^{p_2(\cdot)}}^{q_1} \right) \\ &\leq \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \left(\sum_{l=k}^{\infty} |\lambda_l| \| ([b, I^\beta] (a_l)) \chi_k \|_{L^{p_2(\cdot)}} \right)^{q_1} \\ &\quad + \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l| \| ([b, I^\beta] (a_l)) \chi_k \|_{L^{p_2(\cdot)}} \right)^{q_1} \\ &: = X_1 + X_2. \end{aligned}$$

$$\begin{aligned} Y &= \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \| [b, I^\beta] (f) \chi_k \|_{L^{p_2(\cdot)}}^{q_1} \\ &\leq \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=k}^{\infty} |\lambda_l| \| ([b, I^\beta] (a_l)) \chi_k \|_{L^{p_2(\cdot)}} \right)^{q_1} \\ &\quad + \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l| \| ([b, I^\beta] (a_l)) \chi_k \|_{L^{p_2(\cdot)}} \right)^{q_1} \\ &: = Y_1 + Y_2. \end{aligned}$$

$$\begin{aligned}
Z &= \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \left(\sum_{k=0}^L 2^{kq_1\alpha_\infty} \|[b, I^\beta](f)\chi_k\|_{L^{p_2(\cdot)}}^{q_1} \right) \\
&\lesssim \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \left(\sum_{l=k}^{\infty} |\lambda_l| \|[b, I^\beta](a_l)\chi_k\|_{L^{p_2(\cdot)}} \right)^{q_1} \\
&\quad + \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \left(\sum_{l=-\infty}^{k-1} |\lambda_l| \|[b, I^\beta](a_l)\chi_k\|_{L^{p_2(\cdot)}} \right)^{q_1} \\
&: = Z_1 + Z_2.
\end{aligned}$$

To proceed, we need a pointwise estimate for $[b, I^\beta]$ on F_l , where by (2.3), for any $l, k \in \mathbb{Z}$, with $l \leq k-1$ and $y \in F_l$, then $|x-y| \sim |x|$, $2|y| \leq |x|$, we obtain

$$\begin{aligned}
&|[b, I^\beta](a_j)\chi_k(x)| \\
&\leq \int_{F_l} \frac{|b(x) - b(y)|}{|x-y|^{n-\beta}} |a_l(y)| dy \cdot \chi_k(x) \\
&\lesssim 2^{k(\beta-n)} \left(|b(x) - b_{B_l}| \int_{F_l} |a_l(y)| dy + \int_{F_l} |b_{B_l} - b(y)| |a_l(y)| dy \right) \cdot \chi_k(x) \\
&\lesssim 2^{k(\beta-n)} \|a_l\|_{L^{p_1(\cdot)}} \\
(3.2) \quad &\times \left(|b(x) - b_{B_l}| \|\chi_l\|_{L^{p'_1(\cdot)}} + \|(b - b_{B_j})\chi_l\|_{L^{p'_1(\cdot)}} \right) \cdot \chi_k(x).
\end{aligned}$$

So applying (2.3), (2.5), (2.6), (2.7), (2.9) and (3.2), we have

$$\begin{aligned}
\|[b, I^\beta](a_l)\chi_k\|_{L^{p_2(\cdot)}} &\lesssim 2^{k(\beta-n)} \|a_l\|_{L^{p_1(\cdot)}} \\
&\quad \times \left(\|(b - b_{B_l})\chi_k\|_{L^{p_2(\cdot)}} \|\chi_l\|_{L^{p'_1(\cdot)}} + \|(b - b_{B_j})\chi_l\|_{L^{p'_1(\cdot)}} \|\chi_k\|_{L^{p_2(\cdot)}} \right) \\
&\lesssim 2^{k(\beta-n)} \|a_j\|_{L^{p_1(\cdot)}} \\
&\quad \times \left((k-l) \|b\|_{BMO} \|\chi_{B_k}\|_{L^{p_2(\cdot)}} \|\chi_l\|_{L^{p'_1(\cdot)}} + \|b\|_{BMO} \|\chi_{B_l}\|_{L^{p'_1(\cdot)}} \|\chi_k\|_{L^{p_2(\cdot)}} \right) \\
&\lesssim \|b\|_{BMO} 2^{k(\beta-n)} \|a_l\|_{L^{p_1(\cdot)}} (k-l) \|\chi_{B_k}\|_{L^{p_2(\cdot)}} \|\chi_l\|_{L^{p'_1(\cdot)}} \\
&\lesssim \|b\|_{BMO} 2^{k\beta} (k-l) \|a_l\|_{L^{p_1(\cdot)}} \|\chi_l\|_{L^{p'_1(\cdot)}} \|\chi_{B_k}\|_{L^{p'_2(\cdot)}}^{-1} \\
&\lesssim \|b\|_{BMO} 2^{k\beta} (k-l) \|a_l\|_{L^{p_1(\cdot)}} \|\chi_l\|_{L^{p'_1(\cdot)}} \|\chi_{B_k}\|_{L^{p'_2(\cdot)}}^{-1} \frac{\|\chi_{B_l}\|_{L^{p'_2(\cdot)}}}{\|\chi_{B_k}\|_{L^{p'_2(\cdot)}}} \\
&\lesssim \|b\|_{BMO} 2^{k\beta} (k-l) \|a_l\|_{L^{p_1(\cdot)}} \|\chi_{B_l}\|_{L^{p'_1(\cdot)}} \|\chi_{B_l}\|_{L^{p'_2(\cdot)}}^{-1} 2^{n\delta_2(l-k)} \\
&\lesssim \|b\|_{BMO} 2^{k\beta} (k-l) \|a_l\|_{L^{p_1(\cdot)}} 2^{(n-\beta)l} \|\chi_{B_l}\|_{L^{p_2(\cdot)}}^{-1} \|\chi_{B_l}\|_{L^{p'_2(\cdot)}}^{-1} 2^{n\delta_2(l-k)} \\
&\lesssim \|b\|_{BMO} 2^{(n\delta_2-\beta)(l-k)} (k-l) \|a_l\|_{L^{p_1(\cdot)}} \left(2^{-ln} \|\chi_{B_l}\|_{L^{p_2(\cdot)}} \|\chi_{B_l}\|_{L^{p'_2(\cdot)}} \right)^{-1} \\
&\lesssim \|b\|_{BMO} 2^{(\beta-n\delta_2)(k-l)} (k-l) \|a_l\|_{L^{p_1(\cdot)}} \\
&\lesssim \|b\|_{BMO} (k-l) 2^{(\beta-n\delta_2)(k-l)-\alpha_l l},
\end{aligned}$$

where in the sixth inequality we have used the following fact:

First, we know that

$$\chi_{B_l}(x) \lesssim 2^{-l\beta} [b, I^\beta](\chi_{B_l})(x).$$

Also, let $p_1(\cdot) \in \mathcal{P}(\mathbb{R}^n)$. and $0 < \beta < \frac{n}{(p_1)_+}$. Then, by (2.6) and using (2.8), we obtain

$$\begin{aligned} \|\chi_{B_l}\|_{L^{p_2(\cdot)}} &\lesssim 2^{-l\beta} \|[b, I^\beta](\chi_{B_l})\|_{L^{p_2(\cdot)}} \\ &\lesssim \|b\|_{BMO} 2^{-l\beta} \|\chi_{B_l}\|_{L^{p_1(\cdot)}} \\ &\lesssim \|b\|_{BMO} 2^{(n-\beta)l} \|\chi_{B_l}\|_{L^{p_1'(\cdot)}}^{-1}. \end{aligned}$$

Thus, we get

$$(3.3) \quad \|[b, I^\beta](a_l)\chi_k\|_{L^{p_2(\cdot)}} \lesssim \|b\|_{BMO} (k-l) 2^{(\beta-n\delta_2)(k-l)-\alpha_l l}.$$

To complete the proof, we consider X, Y, Z into two cases $0 < q_1 \leq 1$ and $1 < q_1 < \infty$.

Case 1: ($0 < q_1 \leq 1$). In this case, we always use (3.6) in [3] and the convergence of a geometric series and exchange order of summation and the convergence of geometric power series.

Firstly, since $[b, I^\beta]$ is bounded from $L^{p_1(\cdot)}$ to $L^{p_2(\cdot)}$ and using the condition (2) in Definition 3, then we get

$$(3.4) \quad \|[b, I^\beta](a_l)\chi_k\|_{L^{p_2(\cdot)}} \leq \|b\|_{BMO} \|a_l\|_{L^{p_1(\cdot)}} \leq \|b\|_{BMO} |B_l|^{-\frac{\alpha_l}{n}} = \|b\|_{BMO} 2^{-l\alpha_l}.$$

Thus, by (3.6) in [3] and (3.4), we obtain

$$\begin{aligned} X_1 &= \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda_{q_1}} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \left(\sum_{l=k}^{\infty} |\lambda_l| \|[b, I^\beta](a_l)\chi_k\|_{L^{p_2(\cdot)}} \right)^{q_1} \\ &\lesssim \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda_{q_1}} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \left(\sum_{l=k}^{\infty} |\lambda_l| \|b\|_{BMO} 2^{-l\alpha_l} \right)^{q_1} \\ &\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda_{q_1}} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \left(\sum_{l=k}^{-1} |\lambda_l|^{q_1} 2^{-l\alpha(0)q_1} + \sum_{l=0}^{\infty} |\lambda_l|^{q_1} 2^{-l\alpha_\infty q_1} \right) \\ &\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda_{q_1}} \sum_{k=-\infty}^L \sum_{l=k}^{-1} |\lambda_l|^{q_1} 2^{(k-l)\alpha(0)q_1} \\ &\quad + \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda_{q_1}} \sum_{k=-\infty}^L 2^{k\alpha(0)q_1} \sum_{l=0}^{\infty} |\lambda_l|^{q_1} 2^{-l\alpha_\infty q_1} \\ &\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda_{q_1}} \sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} \sum_{k=-\infty}^l 2^{(k-l)\alpha(0)q_1} \\ &\quad + \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} \sum_{l=0}^{\infty} 2^{-l\lambda_{q_1}} |\lambda_l|^{q_1} 2^{(\lambda-\alpha_\infty)lq_1} 2^{-L\lambda_{q_1}} \sum_{k=-\infty}^L 2^{k\alpha(0)q_1} \\ &\leq \|b\|_{BMO}^{q_1} \left(\sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda_{q_1}} \sum_{l=-\infty}^L |\lambda_l|^{q_1} + \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda_{q_1}} \sum_{l=L}^{-1} |\lambda_l|^{q_1} \sum_{k=-\infty}^l 2^{(k-l)\alpha(0)q_1} \right) \\ &\quad + \|b\|_{BMO}^{q_1} \Psi \sup_{L \leq 0, L \in \mathbb{Z}} \sum_{l=0}^{\infty} 2^{(\lambda-\alpha_\infty)lq_1} \sum_{k=-\infty}^L 2^{(\alpha(0)k-L\lambda)q_1}, \end{aligned}$$

which gives

$$\begin{aligned}
&\lesssim \|b\|_{BMO}^{q_1} \left(\Psi + \sup_{L \leq 0, L \in \mathbb{Z}} \sum_{l=L}^{-1} 2^{-l\lambda q_1} |\lambda_l|^{q_1} 2^{(l-L)\lambda q_1} \sum_{k=-\infty}^l 2^{(k-l)\alpha(0)q_1} + \Psi \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \left(\Psi + \Psi \sup_{L \leq 0, L \in \mathbb{Z}} \sum_{l=L}^{-1} 2^{(l-L)\lambda q_1} \sum_{k=-\infty}^l 2^{(k-l)\alpha(0)q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \Psi \quad (\alpha_\infty > \lambda).
\end{aligned}$$

Secondly, by (3.3), (3.6) in [3] and the hypothesis $\beta - n\delta_2 < \alpha(0)$, we get

$$\begin{aligned}
X_2 &= \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l| \|([b, I^\beta](a_l)) \chi_k\|_{L^{p_2(\cdot)}} \right)^{q_1} \\
&\lesssim \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} \|b\|_{BMO}^{q_1} (k-l)^{q_1} |\lambda_l|^{q_1} 2^{[(\beta-n\delta_2)(k-l)-\alpha_l]q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} (k-l)^{q_1} |\lambda_l|^{q_1} 2^{[(\beta-n\delta_2)(k-l)-\alpha(0)l]q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=-\infty}^L |\lambda_j|^{q_1} \sum_{k=l+1}^{-1} (k-l)^{q_1} 2^{(\beta-n\delta_2-\alpha(0))(k-l)q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \Psi.
\end{aligned}$$

Thirdly, by (3.4) and (3.6) in [3], we obtain that

$$\begin{aligned}
Y_1 &= \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=k}^{\infty} |\lambda_l| \|([b, I^\beta](a_l)) \chi_k\|_{L^{p_2(\cdot)}} \right)^{q_1} \\
&\lesssim \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=k}^{\infty} |\lambda_l| \|b\|_{BMO} 2^{-l\alpha_l} \right)^{q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=k}^{-1} |\lambda_l|^{q_1} 2^{-l\alpha(0)q_1} + \sum_{l=0}^{\infty} |\lambda_l|^{q_1} 2^{-l\alpha_\infty q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \left(\sum_{k=-\infty}^{-1} \sum_{l=k}^{-1} |\lambda_l|^{q_1} 2^{(k-l)\alpha(0)q_1} + \sum_{k=-\infty}^{-1} 2^{k\alpha(0)q_1} \sum_{l=0}^{\infty} |\lambda_l|^{q_1} 2^{-l\alpha_\infty q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \left(\sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} \sum_{k=\infty}^l 2^{(k-l)\alpha(0)q_1} + \sum_{l=0}^{\infty} |\lambda_l|^{q_1} 2^{-l\alpha_\infty q_1} \sum_{k=-\infty}^{-1} 2^{k\alpha(0)q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \left(\sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} + \sum_{l=0}^{\infty} |\lambda_l|^{q_1} 2^{-l\alpha_\infty q_1} 2^{-l\lambda q_1} \sum_{k=-\infty}^{-1} 2^{k\alpha(0)q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \left(\Psi + \Psi \sum_{i=-\infty}^l |\lambda_i|^{q_1} \sum_{l=0}^{\infty} 2^{(\lambda-\alpha_\infty)q_1 l} \sum_{k=-\infty}^l 2^{k\alpha(0)q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \Psi \quad (\alpha_\infty > \lambda).
\end{aligned}$$

Fourthly, by (3.3), (3.6) in [3] and the assumption $\beta - n\delta_2 < \alpha(0)$, we obtain that

$$\begin{aligned}
Y_2 &= \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l| \left\| ([b, I^\beta](a_l)) \chi_k \right\|_{L^{p_2(\cdot)}} \right)^{q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l|^{q_1} (k-l)^{q_1} 2^{[(\beta-n\delta_2)(k-l)-\alpha_l]q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \sum_{k=-\infty}^{-1} |\lambda_l|^{q_1} \sum_{k=l+1}^{-1} (k-l)^{q_1} 2^{(\beta-n\delta_2-\alpha(0))(k-l)q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \Psi.
\end{aligned}$$

Fifthly, similar to X_1 , using (3.4) and (3.6) in [3], we have

$$\begin{aligned}
Z_1 &= \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \left(\sum_{l=k}^{\infty} |\lambda_l| \left\| ([b, I^\beta](a_l)) \chi_k \right\|_{L^{p_2(\cdot)}} \right)^{q_1} \\
&\lesssim \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \sum_{l=k}^{\infty} |\lambda_l|^{q_1} \left\| ([b, I^\beta](a_l)) \chi_k \right\|_{L^{p_2(\cdot)}}^{q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \sum_{l=k}^{\infty} |\lambda_l|^{q_1} 2^{-l\alpha_l q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \sum_{l=k}^{\infty} |\lambda_l|^{q_1} 2^{-l\alpha_\infty q_1} \\
&= \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=0}^L |\lambda_l|^{q_1} \sum_{k=0}^l 2^{(k-l)q_1\alpha_\infty} \\
&\quad + \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=L}^{\infty} |\lambda_l|^{q_1} \sum_{k=0}^L 2^{(k-l)q_1\alpha_\infty} \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=0}^L |\lambda_l|^{q_1} \\
&\quad + \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} \sum_{l=L}^{\infty} 2^{(l-L)q_1\lambda} 2^{-l\lambda q_1} \sum_{i=-\infty}^l |\lambda_i|^{q_1} \sum_{k=0}^L 2^{(k-l)q_1\alpha_\infty},
\end{aligned}$$

which implies

$$\begin{aligned}
&\lesssim \|b\|_{BMO}^{q_1} \Psi + \|b\|_{BMO}^{q_1} \Psi \sup_{L>0, L \in \mathbb{Z}} \sum_{l=L}^{\infty} 2^{(l-L)\lambda q_1} 2^{(L-l)q_1\alpha_\infty} \\
&\lesssim \|b\|_{BMO}^{q_1} \Psi + \|b\|_{BMO}^{q_1} \Psi \sup_{L>0, L \in \mathbb{Z}} \sum_{l=L}^{\infty} 2^{(l-L)q_1(\lambda-\alpha_\infty)} \\
&\lesssim \|b\|_{BMO}^{q_1} \Psi \quad (\alpha_\infty > \lambda).
\end{aligned}$$

Finally, by (3.3), (3.6) in [3] and the assumptions that $\beta - n\delta_2 < \alpha(0)$ and $\beta - n\delta_2 < \alpha_\infty$, we get

$$\begin{aligned}
Z_2 &= \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1 \alpha_\infty} \left(\sum_{l=-\infty}^{k-1} |\lambda_l| \left\| ([b, I^\beta](a_l)) \chi_k \right\|_{L^{p_2(\cdot)}} \right)^{q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1 \alpha_\infty} \left(\sum_{l=-\infty}^{k-1} |\lambda_l|^{q_1} (k-l)^{q_1} 2^{[(\beta-n\delta_2)(k-l)-\alpha_l]q_1} \right) \\
&= \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1 \alpha_\infty} \left(\sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} (k-l)^{q_1} 2^{[(\beta-n\delta_2)(k-l)-\alpha(0)l]q_1} \right) \\
&\quad + \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1 \alpha_\infty} \left(\sum_{l=0}^{k-1} |\lambda_l|^{q_1} (k-l)^{q_1} 2^{[(\beta-n\delta_2)(k-l)-\alpha_\infty l]q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L (k-l)^{q_1} 2^{kq_1 [\alpha_\infty - (\beta-n\delta_2)]} \sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} 2^{[\beta-n\delta_2 - \alpha(0)]lq_1} \\
&\quad + \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L |\lambda_l|^{q_1} \sum_{k=l+1}^{\infty} (k-l)^{q_1} 2^{[(\beta-n\delta_2 - \alpha_\infty)(l-k)]q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \left(\sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} + \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=0}^{L-1} |\lambda_l|^{q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \Psi.
\end{aligned}$$

Case 2: ($1 < q_1 < \infty$). In this case, similar to Case 1, we always exchange order of summation and use the convergence of a geometric series, but use (2.3) instead of (3.6) in [3].

Indeed, for X_1 , let $\frac{1}{q_1} + \frac{1}{q'_1} = 1$. By (2.3) and (3.4), we have

$$\begin{aligned}
X_1 &= \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1 \alpha(0)} \left(\sum_{l=k}^{\infty} |\lambda_l| \left\| ([b, I^\beta](a_l)) \chi_k \right\|_{L^{p_2(\cdot)}} \right)^{q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1 \alpha(0)} \left(\sum_{l=k}^{\infty} |\lambda_l| 2^{-l\alpha_l} \right)^{q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L \left(\sum_{l=k}^{-1} |\lambda_l| 2^{(k-l)\alpha(0)} \right)^{q_1} \\
&\quad + \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{k\alpha(0)q_1} \left(\sum_{l=0}^{\infty} |\lambda_l| 2^{-l\alpha_\infty} \right)^{q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L \left(\sum_{l=k}^{-1} |\lambda_l|^{q_1} 2^{\frac{(k-l)\alpha(0)q_1}{2}} \right) \left(\sum_{l=k}^{-1} 2^{\frac{(k-l)\alpha(0)q'_1}{2}} \right)^{\frac{q_1}{q'_1}} \\
&\quad + \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1 \alpha(0)} \left(\sum_{l=0}^{\infty} |\lambda_l|^{q_1} 2^{\frac{-l\alpha_\infty q_1}{2}} \right) \left(\sum_{l=0}^{\infty} 2^{\frac{-l\alpha_\infty q'_1}{2}} \right)^{\frac{q_1}{q'_1}},
\end{aligned}$$

which gives

$$\begin{aligned}
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L \sum_{l=k}^{-1} |\lambda_l|^{q_1} 2^{\frac{(k-l)\alpha(0)q_1}{2}} \\
&\quad + \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \sum_{l=0}^{\infty} |\lambda_l|^{q_1} 2^{\frac{-l\alpha_{\infty}q_1}{2}} \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} \sum_{k=-\infty}^l 2^{\frac{(k-l)\alpha(0)q_1}{2}} \\
&\quad + \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} \sum_{l=0}^{\infty} 2^{-l\lambda q_1} |\lambda_l|^{q_1} 2^{(\lambda - \frac{\alpha_{\infty}}{2})lq_1} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{k\alpha(0)q_1} \\
&\leq \|b\|_{BMO}^{q_1} \left(\sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=-\infty}^L |\lambda_l|^{q_1} + \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=L}^{-1} |\lambda_l|^{q_1} \sum_{k=-\infty}^l 2^{\frac{(k-l)\alpha(0)q_1}{2}} \right) \\
&\quad + \|b\|_{BMO}^{q_1} \Psi \sup_{L \leq 0, L \in \mathbb{Z}} \sum_{l=0}^{\infty} 2^{(\lambda - \frac{\alpha_{\infty}}{2})lq_1} \sum_{k=-\infty}^L 2^{(\alpha(0)k - L\lambda)q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \left(\Psi + \sup_{L \leq 0, L \in \mathbb{Z}} \sum_{l=L}^{-1} 2^{-l\lambda q_1} |\lambda_l|^{q_1} 2^{(l-L)\lambda q_1} \sum_{k=-\infty}^l 2^{\frac{(k-l)\alpha(0)q_1}{2}} + \Psi \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \left(\Psi + \Psi \sup_{L \leq 0, L \in \mathbb{Z}} \sum_{l=L}^{-1} 2^{(l-L)\lambda q_1} \sum_{k=-\infty}^l 2^{\frac{(k-l)\alpha(0)q_1}{2}} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \Psi \quad (\alpha_{\infty} > 2\lambda).
\end{aligned}$$

For X_2 , let $\frac{1}{q_1} + \frac{1}{q_1'} = 1$. By (2.3), (3.3) and the assumption $\beta - n\delta_2 < \alpha(0)$, we get

$$\begin{aligned}
X_2 &= \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l| \left\| ([b, I^{\beta}](a_l)) \chi_k \right\|_{L^{p_2(\cdot)}} \right)^{q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l|^{q_1} (k-l)^{q_1} 2^{[(\beta - n\delta_2)(k-l) - \alpha_l]q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l|^{q_1} (k-l)^{q_1} 2^{[(\beta - n\delta_2)(k-l) - \alpha(0)l] \frac{q_1}{2}} \right) \\
&\quad \times \left(\sum_{l=-\infty}^{k-1} 2^{[(\beta - n\delta_2)(k-l) - \alpha(0)l] \frac{q_1'}{2}} \right)^{\frac{q_1}{q_1'}} \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=-\infty}^L 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l|^{q_1} (k-l)^{q_1} 2^{[(\beta - n\delta_2)(k-l) - \alpha(0)l] \frac{q_1}{2}} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \sup_{L \leq 0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=-\infty}^L |\lambda_l|^{q_1} \sum_{k=l+1}^{-1} 2^{[(\beta - n\delta_2 - \alpha(0))(k-l)] \frac{q_1}{2}} \\
&\lesssim \|b\|_{BMO}^{q_1} \Psi.
\end{aligned}$$

For Y_1 , let $\frac{1}{q_1} + \frac{1}{q'_1} = 1$. Then, applying (2.3) and (3.4), we know that

$$\begin{aligned}
Y_1 &= \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=k}^{\infty} |\lambda_l| \|([b, I^\beta](a_l)) \chi_k\|_{L^{p_2(\cdot)}} \right)^{q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=k}^{\infty} |\lambda_l| 2^{-l\alpha_l} \right)^{q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \left[\sum_{k=-\infty}^{-1} \left(\sum_{l=k}^{-1} |\lambda_l| 2^{(k-l)\alpha(0)} \right)^{q_1} + \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=0}^{\infty} |\lambda_l| 2^{-l\alpha_\infty} \right)^{q_1} \right] \\
&\lesssim \|b\|_{BMO}^{q_1} \sum_{k=-\infty}^{-1} \left(\sum_{l=k}^{-1} |\lambda_l|^{q_1} 2^{(k-l)\alpha(0)\frac{q_1}{2}} \right) \left(\sum_{l=k}^{-1} 2^{(k-l)\alpha(0)\frac{q'_1}{2}} \right)^{\frac{q_1}{q'_1}} \\
&\quad + \|b\|_{BMO}^{q_1} \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=0}^{\infty} |\lambda_l|^{q_1} 2^{(-l\alpha_\infty)\frac{q_1}{2}} \right) \left(\sum_{l=0}^{\infty} 2^{(-l\alpha_\infty)\frac{q'_1}{2}} \right)^{\frac{q_1}{q'_1}} \\
&\lesssim \|b\|_{BMO}^{q_1} \left(\sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} \sum_{k=-\infty}^l 2^{(k-l)\alpha(0)\frac{q_1}{2}} + \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \sum_{l=0}^{\infty} |\lambda_l|^{q_1} 2^{-l\alpha_\infty\frac{q_1}{2}} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \left(\sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} + \sum_{l=0}^{\infty} 2^{(\lambda - \frac{\alpha_\infty}{2})lq_1} 2^{-l\lambda q_1} \sum_{i=-\infty}^j |\lambda_i|^{q_1} \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \left(\Psi + \Psi \sum_{l=0}^{\infty} 2^{(\lambda - \frac{\alpha_\infty}{2})lq_1} \sum_{k=-\infty}^{-1} 2^{k\alpha(0)q_1} \right) \\
&\lesssim \|b\|_{BMO}^{q_1} \Psi \quad (\alpha_\infty > \lambda).
\end{aligned}$$

For Y_2 , let $\frac{1}{q_1} + \frac{1}{q'_1} = 1$. Then, (2.3), (3.3) and the assumption $\beta - n\delta_2 < \alpha(0)$, we have

$$\begin{aligned}
Y_2 &= \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l| \|([b, I^\beta](a_l)) \chi_k\|_{L^{p_2(\cdot)}} \right)^{q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l| (k-l) 2^{(\beta-n\delta_2)(k-l)-\alpha(0)l} \right)^{q_1} \\
&\lesssim \|b\|_{BMO}^{q_1} \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l|^{q_1} 2^{[(\beta-n\delta_2)(k-l)-\alpha(0)l]\frac{q_1}{2}} \right) \\
&\quad \times \left(\sum_{l=-\infty}^{k-1} (k-l) 2^{[(\beta-n\delta_2)(k-l)-\alpha(0)l]\frac{q'_1}{2}} \right)^{\frac{q_1}{q'_1}} \\
&\lesssim \|b\|_{BMO}^{q_1} \sum_{k=-\infty}^{-1} 2^{kq_1\alpha(0)} \left(\sum_{l=-\infty}^{k-1} |\lambda_l|^{q_1} 2^{[(\beta-n\delta_2)(k-l)-\alpha(0)l]\frac{q_1}{2}} \right),
\end{aligned}$$

which implies

$$\begin{aligned} &\lesssim \|b\|_{BMO}^{q_1} \sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} \sum_{k=l+1}^{-1} 2^{(\beta-n\delta_2-\alpha(0))(k-l)\frac{q_1}{2}} \\ &\lesssim \|b\|_{BMO}^{q_1} \Psi. \end{aligned}$$

For Z_1 , let $\frac{1}{q_1} + \frac{1}{q_1'} = 1$. Then, using (2.3) and (3.4), we have

$$\begin{aligned} Z_1 &= \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \left(\sum_{l=k}^{\infty} |\lambda_l| \left\| ([b, I^\beta](a_l)) \chi_k \right\|_{L^{p_2(\cdot)}} \right)^{q_1} \\ &\lesssim \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \left(\sum_{l=k}^{\infty} |\lambda_l|^{q_1} \left\| ([b, I^\beta](a_l)) \chi_k \right\|_{L^{p_2(\cdot)}}^{\frac{q_1}{2}} \right) \left(\sum_{l=k}^{\infty} \left\| ([b, I^\beta](a_l)) \chi_k \right\|_{L^{p_2(\cdot)}}^{\frac{q_1'}{2}} \right)^{\frac{q_1}{q_1'}} \\ &\lesssim \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \left(\sum_{l=k}^{\infty} |\lambda_l|^{q_1} \|a_l\|_{L^{p_1(\cdot)}}^{\frac{q_1}{2}} \right) \left(\sum_{l=k}^{\infty} \|a_l\|_{L^{p_1(\cdot)}}^{\frac{q_1'}{2}} \right)^{\frac{q_1}{q_1'}} \\ &\lesssim \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \left(\sum_{l=k}^{\infty} |\lambda_l|^{q_1} |B_l|^{-\frac{\alpha_l q_1}{(2^n)}} \right) \left(\sum_{l=k}^{\infty} |B_l|^{-\frac{\alpha_l q_1'}{(2^n)}} \right)^{\frac{q_1}{q_1'}} \\ &\lesssim \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{k\frac{q_1}{2}\alpha_\infty} \left(\sum_{l=k}^{\infty} |\lambda_l|^{q_1} |B_l|^{-\frac{\alpha_l q_1}{(2^n)}} \right) \\ &= \|b\|_{BMO}^{q_1} \left(\sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=0}^L |\lambda_l|^{q_1} \sum_{k=0}^l 2^{(k-l)\frac{q_1}{2}\alpha_\infty} + \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=L}^{\infty} |\lambda_l|^{q_1} \sum_{k=0}^L 2^{(k-l)\frac{q_1}{2}\alpha_\infty} \right) \\ &\lesssim \|b\|_{BMO}^{q_1} \left(\sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=0}^L |\lambda_l|^{q_1} + \sup_{L>0, L \in \mathbb{Z}} \sum_{l=L}^{\infty} 2^{(l-L)q_1\lambda} 2^{-l\lambda q_1} \sum_{i=-\infty}^l |\lambda_i|^{q_1} \sum_{k=0}^L 2^{(k-l)\frac{q_1}{2}\alpha_\infty} \right) \\ &\lesssim \|b\|_{BMO}^{q_1} \left(\Psi + \Psi \sup_{L>0, L \in \mathbb{Z}} \sum_{l=L}^{\infty} 2^{(l-L)\lambda q_1} 2^{(L-l)\frac{q_1}{2}\alpha_\infty} \right) \\ &\lesssim \|b\|_{BMO}^{q_1} \left(\Psi + \Psi \sup_{L>0, L \in \mathbb{Z}} \sum_{l=L}^{\infty} 2^{(l-L)\lambda q_1} (\lambda - \frac{\alpha_\infty}{2}) \right) \\ &\lesssim \|b\|_{BMO}^{q_1} \Psi \quad (\alpha_\infty > 2\lambda). \end{aligned}$$

Finally, we consider the term Z_2 . Now, let $\frac{1}{q_1} + \frac{1}{q_1'} = 1$. By (2.3), (3.3) and the conditions $\beta - n\delta_2 < \alpha(0)$ and $\beta - n\delta_2 < \alpha_\infty$, we obtain that

$$\begin{aligned} Z_2 &= \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \left(\sum_{l=-\infty}^{k-1} |\lambda_l| \left\| ([b, I^\beta](a_l)) \chi_k \right\|_{L^{p_2(\cdot)}} \right)^{q_1} \\ &\lesssim \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \left(\sum_{l=-\infty}^{k-1} |\lambda_l| (k-l) 2^{(\beta-n\delta_2)(k-l)-\alpha_l l} \right)^{q_1} \\ &\lesssim \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \left(\sum_{l=-\infty}^{-1} |\lambda_l| (k-l) 2^{(\beta-n\delta_2)(k-l)-\alpha(0)l} \right)^{q_1} \end{aligned}$$

$$\begin{aligned}
& + \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1\alpha_\infty} \left(\sum_{l=0}^{k-1} |\lambda_l| (k-l) 2^{(\beta-n\delta_2)(k-l)-\alpha_\infty l} \right)^{q_1} \\
\lesssim & \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L 2^{kq_1[\alpha_\infty - (\beta-n\delta_2)]} \left(\sum_{l=-\infty}^{-1} |\lambda_l| (k-l) 2^{l[(\beta-n\delta_2)-\alpha(0)]} \right)^{q_1} \\
& + \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L \left(\sum_{l=0}^{k-1} |\lambda_l| (k-l) 2^{(\beta-n\delta_2-\alpha_\infty)(k-l)} \right)^{q_1} \\
\lesssim & \|b\|_{BMO}^{q_1} \left(\sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} 2^{l[(\beta-n\delta_2)-\alpha(0)]l\frac{q_1}{2}} \right) \left(\sum_{l=-\infty}^{-1} (k-l)^{q_1} 2^{l[(\beta-n\delta_2)-\alpha(0)]l\frac{q_1}{2}} \right)^{\frac{q_1}{q_1-1}} \\
& + \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L \left(\sum_{l=0}^{k-1} |\lambda_l|^{q_1} (k-l)^{q_1} 2^{l[(\beta-n\delta_2-\alpha_\infty)(k-l)]\frac{q_1}{2}} \right) \\
& \times \left(\sum_{l=0}^{k-1} 2^{l[(\beta-n\delta_2-\alpha_\infty)(k-l)]\frac{q_1}{2}} \right)^{\frac{q_1}{q_1-1}} \\
\lesssim & \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} (k-l)^{q_1} 2^{l[(\beta-n\delta_2)-\alpha(0)]l\frac{q_1}{2}} \\
& + \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{k=0}^L \sum_{l=0}^{k-1} |\lambda_l|^{q_1} (k-l)^{q_1} 2^{l[(\beta-n\delta_2-\alpha_\infty)(k-l)]\frac{q_1}{2}} \\
\lesssim & \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} \\
& + \|b\|_{BMO}^{q_1} \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=0}^{L-1} |\lambda_l|^{q_1} \sum_{k=l+1}^L (k-l)^{q_1} 2^{l[(\beta-n\delta_2-\alpha_\infty)(k-l)]\frac{q_1}{2}} \\
\lesssim & \|b\|_{BMO}^{q_1} \left(\sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=-\infty}^{-1} |\lambda_l|^{q_1} + \sup_{L>0, L \in \mathbb{Z}} 2^{-L\lambda q_1} \sum_{l=0}^{L-1} |\lambda_l|^{q_1} \right) \\
\lesssim & \|b\|_{BMO}^{q_1} \Psi.
\end{aligned}$$

Then, by joining the above inequalities for X, Y and Z , we obtain (3.1). Thus, the proof is completed. \square

REFERENCES

- [1] C. Capone, D. Cruz-Urbe, A. Fiorenza. The fractional maximal operator and fractional integrals on variable L^p spaces. *Rev. Mat. Iberoam.* 23(3), 743–770 (2007).
- [2] C. Fefferman, E.M. Stein. H^p spaces of several variables. *Acta Math.* 129(3-4), 137–193 (1972). doi:10.1007/BF02392215
- [3] F. Gürbüz. Some inequalities for Riesz Potential on homogeneous variable exponent Herz-Morrey-Hardy spaces. (submitted.) arXiv:2411.13880v1 [math.FA] 21 Nov 2024 DOI: 10.48550/arXiv.2411.13880 (accepted for publication in the Springer Volume)
- [4] C.S. Herz. Lipschitz spaces and Bernstein's theorem on absolutely convergent Fourier transforms. *J. Math. Mech.* 18(4), 283–323 (1968/69).
- [5] M. Izuki. Boundedness of sublinear operators on Herz spaces with variable exponent and application to wavelet characterization. *Anal. Math.* 36(1), 33–50 (2010). doi: 10.1007/s10476-010-0102-8

- [6] M. Izuki. Boundedness of commutators on Herz spaces with variable exponent. *Rend. Circ. Mat. Palermo* (2) 59(2) (2010), 199-213.
- [7] O. Kováčik, J. Rákosník. On spaces $L^{p(x)}$ and $W^{k,p(x)}$. *Czechoslovak Math. J.* 41, 592-618 (1991). doi:10.21136/CMJ.1991.102493
- [8] E.M. Stein. *Harmonic Analysis*. Princeton University Press, Princeton, NJ, USA (1993).
- [9] J. Xu, X. Yang. Herz-Morrey-Hardy spaces with variable exponents and their applications. *J. Funct. Spaces*. 2015, Art. ID 160635, 1-19 (2015). doi: 10.1155/2015/160635
- [10] J. Xu, X. Yang. The molecular decomposition of Herz-Morrey-Hardy spaces with variable exponents and its application. *J. Math. Inequal.* 10, 977-1008 (2016). doi:10.7153/jmi-10-79

DEPARTMENT OF MATHEMATICS, KIRKLARELI UNIVERSITY, KIRKLARELI 39100, TÜRKİYE
Email address: feritgurbuz@klu.edu.tr