

MEAN OSCILLATION CONDITIONS FOR NONLINEAR EQUATION AND REGULARITY RESULTS

PETER HÄSTÖ, MIKYOUNG LEE, AND JIHOON OK

ABSTRACT. We consider general nonlinear elliptic equations of the form

$$\operatorname{div} A(x, Du) = 0 \quad \text{in } \Omega,$$

where $A : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfies a quasi-isotropic (p, q) -growth condition, which is equivalent to the point-wise uniform ellipticity of A . We establish sharp and comprehensive mean oscillation conditions on $A(x, \xi)$ with respect to the x variable to obtain C^1 - and $W^{1,s}$ -regularity results. The results provide new conditions even in the standard p -growth case with coefficient $\operatorname{div}(a(x)|Du|^{p-2}Du) = 0$. Also included are variable exponent growth with and without perturbation as well as borderline double-phase growth and double-phase growth with a coefficient.

1. INTRODUCTION

We study regularity properties of weak solutions to the following nonlinear, non-autonomous equation

$$(\operatorname{div} A) \quad \operatorname{div} A(x, Du) = 0$$

in a domain $\Omega \subset \mathbb{R}^n$, $n \geq 2$, where the nonlinearity $A : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfies the quasi-isotropic (p, q) -growth condition in Definition 1.1. In particular, we are interested in sharp conditions on A of mean oscillation type that imply desired regularity of the weak solution to $(\operatorname{div} A)$.

Let us start with reviewing relevant previous results. Regularity is well-known for linear equations with coefficient, $A(x, \xi) = M(x)\xi$, i.e.

$$(\Delta_{2,M}) \quad \operatorname{div}(M(x)Du) = 0 \quad \text{in } \Omega,$$

where the coefficient matrix $M : \Omega \rightarrow \mathbb{M}_n$ satisfies $L^{-1} \leq M(x)e \cdot e \leq L$ for some $L \geq 1$ and all $x \in \Omega$ and $e \in \partial B_1(0)$. We consider three implications in various settings:

$$\begin{aligned} (\Rightarrow C^{1,\alpha}) \quad & \omega(r) \lesssim r^\beta \quad \Longrightarrow \quad u \in C_{\text{loc}}^{1,\alpha}(\Omega), \\ (\Rightarrow W^{1,s}) \quad & \lim_{r \rightarrow 0^+} \omega(r) = 0 \quad \Longrightarrow \quad u \in W_{\text{loc}}^{1,s}(\Omega) \quad \forall s > 1, \\ (\Rightarrow C^1) \quad & \int_0^1 \omega(r) \frac{dr}{r} < \infty \quad \Longrightarrow \quad u \in C_{\text{loc}}^1(\Omega), \end{aligned}$$

where ω is chosen differently according to the setting. Note that $(\Rightarrow W^{1,s})$ includes the result $u \in C^{0,\alpha}$ for any $\alpha \in (0, 1)$ by the Sobolev embedding, as well as Calderón-Zygmund

Date: April 4, 2025.

2020 Mathematics Subject Classification. 35J62, 35B65, 35R05, 35J92, 46E35.

Key words and phrases. Mean oscillation, gradient continuity, Calderón-Zygmund estimate, non-standard growth, non-uniformly elliptic, quasi-isotropic, generalized Orlicz space.

type gradient estimates for the corresponding nonhomogeneous equations, as stated in Theorem 1.2. In the linear setting with ω equal to the point-wise continuity modulus

$$\omega_M(r) := \sup_{B_r \subset \Omega} \sup_{x, y \in B_r} |M(x) - M(y)|$$

the implications ($\Rightarrow C^{1,\alpha}$), ($\Rightarrow W^{1,s}$) and ($\Rightarrow C^1$) were proved by Caccioppoli/Schauder [18, 53, 54], Agmon–Douglis–Nirenberg [4] and Hartman–Wintner [36], respectively.

More recently, Di Fazio [30] and Dong–Kim [31] showed, respectively, that ($\Rightarrow W^{1,s}$) and ($\Rightarrow C^1$) remain true under the weaker assumption that ω is the modulus of mean continuity

$$\bar{\omega}_M(r) := \sup_{B_r \subset \Omega} \int_{B_r} \int_{B_r} |M(x) - M(y)| dx dy.$$

Note that the implication ($\Rightarrow C^{1,\alpha}$) is trivial for $\bar{\omega}_M$ since this modulus has an upper bound r^β if and only if ω_M does, by the Campanato embedding. We refer to [5, 14, 32, 40, 45] for more related results and the sharpness of mean oscillation type conditions. In short, mean oscillation type conditions are essentially the optimal ones in regularity theory for partial differential equations.

Many of the above implications have been generalized to the p -Laplace equations with coefficients of the form

$$(\Delta_{p,M}) \quad \operatorname{div} \left((M(x)Du \cdot Du)^{\frac{p-2}{2}} M(x)Du \right) = 0 \quad \text{in } \Omega, \quad 1 < p < \infty,$$

i.e. $A(x, \xi) = (M(x)\xi \cdot \xi)^{\frac{p-2}{2}} M(x)\xi$. Manfredi first proved ($\Rightarrow C^{1,\alpha}$) in [49]. Kuusi–Mingione [43] established the implication ($\Rightarrow C^1$) with the point-wise continuity modulus $\omega = \omega_M$, whereas Kinnunen–Zhou [42] established the implication ($\Rightarrow W^{1,s}$) with the mean continuity modulus $\omega = \bar{\omega}_M$. However, the implication ($\Rightarrow C^1$) with mean continuity modulus $\omega = \bar{\omega}_M$ has remained open for the p -Laplacian, even for scalar weights $M(x) = a(x)I_n$, where I_n is the identity matrix.

Research into equations with nonstandard growth conditions has exploded in the past 25 years. Nonstandard growth means that the growth of the equation strongly depends on the x variable, so that the equation does not satisfy the stronger global version of uniform ellipticity defined below. A model equation with nonstandard growth is the $p(x)$ -Laplace equation

$$(\Delta_{p(\cdot)}) \quad \operatorname{div} (|Du|^{p(x)-2} Du) = 0 \quad \text{in } \Omega, \quad 1 < p_1 \leq p(x) \leq p_2.$$

Here, ($\Rightarrow C^{1,\alpha}$) was first proved by Coscia–Mingione [22] and Acerbi–Mingione [1]. With $\omega(r) = \omega_p(r) \log \frac{1}{r}$ now including an extra logarithm, where ω_p is the point-wise modulus of continuity as before, the implications ($\Rightarrow W^{1,s}$) and ($\Rightarrow C^1$) were proved in [2, 17] and [51], respectively. However, to the best of our knowledge no results have been proved under mean continuity conditions for the $p(\cdot)$ -Laplacian. Similarly, for the borderline double phase problem

$$(\Delta_{\text{bdp}}) \quad \operatorname{div} (|Du|^{p-2} Du + a(x) \log(1 + |Du|) |Du|^{p-2} Du) = 0 \quad \text{in } \Omega,$$

where $1 < p < \infty$ and $0 \leq a(x) \leq L$, regularity results have been obtained under point-wise continuity assumptions [8, 9, 15] but no results are available when assuming only mean continuity.

In recent years, $C^{0,1}$ - and C^1 -regularity results under Sobolev–Lorentz type conditions have been developed. De Filippis and Mingione [25] proved that if $M(x) = a(x)I_n$ and $a \in W^{1,1}(\Omega)$ and $|Da|$ is in the Lorentz space $L^{n,1}(\Omega)$, then the weak solution to $(\Delta_{p,M})$ is locally Lipschitz. In fact, they considered a more general class of nonuniformly elliptic

Assumption type	Conclusion	$(\Delta_{2,M})$	$(\Delta_{p,M})$	$(\Delta_{p(\cdot)})$	(Δ_{bdp})	$(\text{div}A)$
Point-wise	$(\Rightarrow C^{1,\alpha})$	[18, 53, 54]	[49]	[1, 22]	[9]	[37, 38]
	$(\Rightarrow W^{1,s})$	[4]	[42]	[2, 17]	[15]	new
	$(\Rightarrow C^1)$	[36]	[43]	[51]	[8]	new
Mean	$(\Rightarrow W^{1,s})$	[30]	[42]	new	new	new
	$(\Rightarrow C^1)$	[31]*	new	new	new	new
Lorenz	$\Rightarrow C^1$	[25]	[25]	[6]	[7]	new

TABLE 1. A summary of previous results for different equations, assumptions and conclusions. The asterisk is explained in Remark 1.4.

equations. Baroni and Coscia [6, 7, 8] have obtained C^1 -regularity results for the (Δ_{bdp}) and $(\Delta_{p(\cdot)})$ under Lorenz-type assumptions on the coefficient and exponent. One of the advantages of our general framework is that also assumptions of Sobolev–Lorentz type are included as special cases of our Dini mean continuity condition, see Proposition 4.8.

The results mentioned so far are summarized in Table 1. In the table we see that results with mean continuity assumptions are especially lacking, although also other categories have gaps for more complicated types of examples. In particular, general equations of type $(\text{div}A)$ have almost no higher regularity results and neither do special cases not mentioned in the table like variable exponent double phase energies (e.g., [50, 57]).

In this paper, we fill in the blanks in the table by proving results with mean continuity assumptions for a general equation with the following fundamental growth and ellipticity conditions on the nonlinearity $A(x, \xi)$. For (aInc) and (aDec) , see Definition 2.1.

Definition 1.1. Let $1 < p < q$. We say that $A : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ with $A(x, \cdot) \in C^1(\mathbb{R}^n \setminus \{0\}, \mathbb{R}^n)$ satisfies *quasi-isotropic (p, q) -growth* condition if

- (i) $D_\xi A(x, \xi)$ satisfies $(\text{aInc})_{p-2}$ and $(\text{aDec})_{q-2}$ with constant $L \geq 1$ ((p, q) -growth);
- (ii) for every $x \in \Omega$ and $e, \xi, \xi' \in \mathbb{R}^n$ with $|e| = 1$ and $|\xi| = |\xi'| > 0$,

$$|D_\xi A(x, \xi')| \leq L D_\xi A(x, \xi) e \cdot e$$

for some $L \geq 1$ (quasi-isotropic ellipticity).

When $A(x, \xi) = D_\xi f(x, \xi)$ for some $f(x, \cdot) \in C^1(\mathbb{R}^n) \cap C^2(\mathbb{R}^n \setminus \{0\})$, the condition (ii) is equivalent to the following *point-wise uniform ellipticity* condition:

$$\mathcal{R}(x, t) := \frac{\sup\{\text{eigenvalues of } D_\xi A(x, \xi) : |\xi| = t\}}{\inf\{\text{eigenvalues of } D_\xi A(x, \xi) : |\xi| = t\}} \leq L \quad \text{for all } x \in \Omega \text{ and } t > 0.$$

Note that the stronger global version of uniform ellipticity condition:

$$\mathcal{R}_\Omega(t) := \frac{\sup\{\text{eigenvalues of } D_\xi A(x, \xi) : x \in \Omega, |\xi| = t\}}{\inf\{\text{eigenvalues of } D_\xi A(x, \xi) : x \in \Omega, |\xi| = t\}} \leq L \quad \text{for all } t > 0$$

plays an important role in regularity theory of partial differential equations and the calculus of variations. In particular, sharp $C^{1,\alpha}$ -regularity under oscillation type condition on A has been obtained in [37, 38]. We also refer to recent development in regularity theory for nonuniformly elliptic equations by De Filippis and Mingione [25, 26, 27, 28].

Another important model equation is the double phase equation

$$\text{div} (|Du|^{p-2} Du + a(x)|Du|^{q-2} Du) = 0 \quad \text{in } \Omega,$$

where $1 < p \leq q < \infty$ and $0 \leq a(x) \leq L$. For this equation, $C^{1,\alpha}$ - and $W^{1,s}$ -regularity results have been established in [10, 20, 21, 24] by assuming point-wise continuity $a \in C^{0,\alpha}(\Omega)$ and the inequality $\frac{q}{p} \leq 1 + \frac{\alpha}{n}$. Note that this energy is not included in the Table 1, since there is no gap for mean continuity results due to the Campanato embedding, see Example 4.3.

Now, we state our main results, Theorems 1.2 and 1.3, which provide sharp and comprehensive C^1 - and $W^{1,s}$ -regularity results for (point-wise) uniformly elliptic equations. Moreover, we will see in Section 4 that the theorems imply new results even in the special cases of Table 1. Define

$$A^{(-1)}(x, \xi) := |\xi|A(x, \xi).$$

The conditions on A in the following theorems are explained in Definitions 2.4 and 3.1.

We first have a Calderón–Zygmund type estimate for the following nonhomogeneous version of $(\operatorname{div}A)$:

$$(\operatorname{div}A; F) \quad \operatorname{div}A(x, Du) = \operatorname{div}A(x, F) \quad \text{in } \Omega,$$

where the given function $F \in L^1(\Omega)$ satisfies $|A^{(-1)}(\cdot, F)| \in L^1_{\operatorname{loc}}(\Omega)$. Note that $(\operatorname{div}A)$ is the special case when $F \equiv 0$ so the following result implies in particular the $W^{1,s}$ -regularity for the equation $(\operatorname{div}A)$, i.e. $(\Rightarrow W^{1,s})$.

Theorem 1.2. *Let $A : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfy the quasi-isotropic (p, q) -growth condition, and let $u \in W^{1,1}_{\operatorname{loc}}(\Omega)$ with $|A^{(-1)}(\cdot, Du)| \in L^1_{\operatorname{loc}}(\Omega)$ be a weak solution to $(\operatorname{div}A; F)$. If $A^{(-1)}$ satisfies $(\operatorname{VMA1})$ and $(\operatorname{SA1})$ and $|A^{(-1)}(\cdot, F)| \in L^s_{\operatorname{loc}}(\Omega)$ for some $s > 1$, then $|A^{(-1)}(\cdot, Du)| \in L^s_{\operatorname{loc}}(\Omega)$. Moreover, for any $\Omega' \Subset \Omega$ there exists a small $R_0 > 0$ depending on $n, p, q, L, s, \theta_1, \Omega'$ and Du such that*

$$\int_{B_r} |A^{(-1)}(x, Du)|^s dx \leq c \left(\int_{B_{2r}} |A^{(-1)}(x, Du)| dx \right)^s + c \int_{B_{2r}} |A^{(-1)}(x, F)|^s dx + c$$

for some $c = c(n, p, q, L, s) > 0$, whenever $r \leq R_0$ and $B_{2r} \subset \Omega'$.

Finally, we have the C^1 -result under Dini mean continuity assumption, which is the most delicate part of the paper and its central contribution.

Theorem 1.3. *Let $A : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfy the quasi-isotropic (p, q) -growth condition, and let $u \in W^{1,1}_{\operatorname{loc}}(\Omega)$ with $|A^{(-1)}(\cdot, Du)| \in L^1_{\operatorname{loc}}(\Omega)$ be a weak solution to $(\operatorname{div}A)$. If $A^{(-1)}$ satisfies $(\operatorname{DMA1})_\gamma$ for some $\gamma > 2$, then Du is continuous in Ω .*

Remark 1.4. The parameter γ in the previous theorem can be thought of as the power of the mean, as in $\omega_\gamma(r) = \sup_{B_r \subset \Omega} (\int_{B_r} \int_{B_r} |M(x) - M(y)|^\gamma dx dy)^{1/\gamma}$. In the linear case, Dong and Kim [31] were able to consider the case $\gamma = 1$, but we require the slightly stronger assumption $\gamma > 2$. For this reason we included an asterisk in Table 1 to indicate that our general result does not quite cover this one paper. See also Proposition 3.8.

Let us comment on the Dini mean oscillation version of $(\operatorname{A1})$, $(\operatorname{DMA1})$, and the vanishing mean oscillation version of $(\operatorname{A1})$, $(\operatorname{VMA1})$. According to Definition 3.1 for $G = A^{(-1)}$ we consider mean averages of the quantity

$$\tilde{\theta}(x, B_r) = \sup_{\xi \in D(B_r)} \frac{|A(x, \xi) - A_{B_r}(\xi)|}{|A_{B_r}(\xi)| + \omega(r)}.$$

The denominator depends on the choice of the ball B_r , which make it difficult to find the relation between fundamental point-wise $(\operatorname{A1})$ -type conditions in Definition 2.4 and

the mean oscillation type condition and proving properties of the function θ_γ defined in Definition 3.1. We investigate the mean oscillation type conditions in Section 3.

The proof of Theorem 1.3 consists of two major steps. The first step is to obtain a comparison estimate in L^1 -space between the gradients of the weak solutions to the original equation $(\operatorname{div}A)$ and its approximating autonomous equation in $(\operatorname{div}\tilde{A})$ (see Section 5). Note that the weak solution to $(\operatorname{div}\tilde{A})$ has good $C^{1,\alpha}$ -regularity estimates by Lemma 5.9. We emphasize that the comparison estimate in Lemma 5.10 is sharper than the one we obtained in [38]. Specially, the exponent of Θ in Lemma 5.10 is 2, whereas the approach used in the proof of [38, Lemma 6.2] yields a smaller exponent. This sharp estimate allows us to prove the C^1 -regularity under the (DMA1) condition. We also note that the approximation deriving the equation $(\operatorname{div}\tilde{A})$ is similar to the one in [38], but simpler since we apply the splicing technique only for large values of $|\xi| = t$, rather than for both large and small values. The second step is an iteration. We improve upon the iteration argument in [43], leading to Lemma 7.3, which can be applied for both the Lipschitz regularity and C^1 -regularity.

The proof of the Calderón–Zygmund estimate in Theorem 1.2 also involves two steps (see Section 6). The first step is, once again, a comparison. Since we are dealing with the non-homogeneous equation $(\operatorname{div}A; F)$, a new comparison estimate for the gradients of the weak solutions to $(\operatorname{div}A; F)$ and $(\operatorname{div}\tilde{A})$ is required, which is provided in Lemma 6.3. The next step is to prove the $W^{1,s}$ -estimate by estimating integrals of $\varphi(x, |Du|)$ over super-level sets. We follow the so-called maximal function free approach, introduced in [3]. Note that in this process, we must essentially use the (A1) condition of $A^{(-1)}$.

2. PRELIMINARIES AND NOTATION

Let Ω be a bounded domain in \mathbb{R}^n with $n \geq 2$. We denote by $B_r(x_0)$ the open ball with center $x_0 \in \mathbb{R}^n$ and radius $r > 0$. If the center is either clear or irrelevant, we simplify notation to $B_r = B_r(x_0)$. For a set $E \subset \mathbb{R}^n$, χ_E is the usual *characteristic function* of E such that $\chi_E(x) = 1$ if $x \in E$ and $\chi_E(x) = 0$ if $x \notin E$. We denote the Hölder conjugate exponent of $p \in [1, \infty]$ by $p' = \frac{p}{p-1}$. A generic constant denoted by $c > 0$ without subscript may vary between appearances.

Let $f, g : E \rightarrow \mathbb{R}$ be measurable in $E \subset \mathbb{R}^n$. We denote the average of f over E with $0 < |E| < \infty$ by $f_E := \int_E f dx := \frac{1}{|E|} \int_E f dx$. The notation $f \lesssim g$ means that there exists a constant $C > 0$ such that $f(y) \leq Cg(y)$ all $y \in E$ and $f \approx g$ means that $f \lesssim g \lesssim f$. When $E \subset \mathbb{R}$, we say that f is *almost increasing* on E with constant $L \geq 1$ if $f(s) \leq Lf(t)$ whenever $s, t \in E$ with $s \leq t$. If we can choose $L = 1$, we say that f is *increasing* on E . *Almost decreasing* and *decreasing* are defined similarly. *Modulus of continuity* refers to a concave and increasing function $\omega : [0, \infty) \rightarrow [0, \infty)$ with $\omega(0) = \lim_{r \rightarrow 0^+} \omega(r) = 0$.

We introduce fundamental conditions on the energy function $\varphi : \Omega \times [0, \infty) \rightarrow [0, \infty)$. We refer to [34, Chapter 2] for the following definitions and properties. We start with regularity with respect to the second variable, which are supposed to hold for all $x \in \Omega$ and a constant $L \geq 1$ independent of x .

Definition 2.1. Let $\varphi : \Omega \times [0, \infty) \rightarrow [0, \infty)$ and $\gamma \in \mathbb{R}$. We say that φ satisfies

- (aInc) $_\gamma$ if $t \mapsto t^{-\gamma}\varphi(x, t)$ is almost increasing on $(0, \infty)$ with constant L ;
- (Inc) $_\gamma$ if $t \mapsto t^{-\gamma}\varphi(x, t)$ is increasing on $(0, \infty)$;
- (aDec) $_\gamma$ if $t \mapsto t^{-\gamma}\varphi(x, t)$ is almost decreasing on $(0, \infty)$ with constant L ;

(Dec) $_\gamma$ if $t \mapsto t^{-\gamma}\varphi(x, t)$ is decreasing on $(0, \infty)$;

$$(A0) \quad L^{-1} \leq \varphi(x, 1) \leq L.$$

We say that φ satisfies (aInc) or (aDec) if it satisfies (aInc) $_\gamma$ or (aDec) $_\gamma$, respectively, for some $\gamma > 1$.

Furthermore, for a vector-valued function $G : \Omega \times \mathbb{R}^M \rightarrow \mathbb{R}^N$, we say that G satisfies (aInc) $_\gamma$ or (aDec) $_\gamma$ if $\varphi(x, t) := |G(x, te)|$ satisfies (aInc) $_\gamma$ or (aDec) $_\gamma$, respectively, with the constant L uniformly in $e \in \mathbb{R}^M$ with $|e| = 1$.

For $p, q > 0$, the conditions (aInc) $_p$ or (aDec) $_q$ on φ with constant $L \geq 1$ are equivalent to the following inequalities

$$\varphi(x, \lambda t) \leq L\lambda^p\varphi(x, t) \quad \text{or} \quad \varphi(x, \Lambda t) \leq L\Lambda^q\varphi(x, t), \quad \text{respectively,}$$

for all $(x, t) \in \Omega \times [0, \infty)$ and $0 \leq \lambda \leq 1 \leq \Lambda$. Additionally, (aInc) or (aDec) are equivalent to ∇_2 -condition or Δ_2 -condition, respectively. Although the definition of (A0) presented above slightly differs from that in [34], the two definitions coincide when φ satisfies (aDec). In the case $\varphi(x, \cdot) \in C^1((0, \infty))$, the conditions (Inc) $_p$ and (Dec) $_q$ for $0 < p \leq q$ are equivalent to

$$p \leq \frac{t\varphi'(x, t)}{\varphi(x, t)} \leq q \quad \text{for all } t \in (0, \infty).$$

Let us consider increasing functions $\varphi, \psi : [0, \infty) \rightarrow [0, \infty)$ such that φ satisfies (aInc) $_1$ and (aDec) $_q$ for some $q \geq 1$, and ψ satisfies (aDec) $_1$. Then there exist a convex function $\tilde{\varphi}$ and a concave function $\tilde{\psi}$ such that $\varphi \approx \tilde{\varphi}$ and $\psi \approx \tilde{\psi}$ from [34, Lemma 2.2.1]. In turn, Jensen's inequality for $\tilde{\varphi}$ and $\tilde{\psi}$ yields that

$$\varphi\left(\int_{\Omega} |f| dx\right) \lesssim \int_{\Omega} \varphi(|f|) dx \quad \text{and} \quad \int_{\Omega} \psi(|f|) dx \lesssim \psi\left(\int_{\Omega} |f| dx\right)$$

for every $f \in L^1(\Omega)$. Here, the implicit constants depend on L from (aInc) $_1$ and (aDec) $_q$ or (aDec) $_1$, based on the constants arising from the equivalence relation.

We define classes of Φ -functions and generalized Orlicz spaces, following [34]. Our primary focus is on convex functions relevant to minimization problems and associated PDEs; however, the class $\Phi_w(\Omega)$ is quite useful for approximating functionals.

Definition 2.2. Let $\varphi : \Omega \times [0, \infty] \rightarrow [0, \infty)$. Assume $x \mapsto \varphi(x, |f(x)|)$ is measurable for every measurable function f on Ω , $t \mapsto \varphi(x, t)$ is increasing for every $x \in \Omega$, and $\varphi(x, 0) = \lim_{t \rightarrow 0^+} \varphi(x, t) = 0$ and $\lim_{t \rightarrow \infty} \varphi(x, t) = \infty$ for every $x \in \Omega$. Then φ is said to be

- (1) a Φ -function, denoted $\varphi \in \Phi_w(\Omega)$, if it satisfies (aInc) $_1$;
- (2) a convex Φ -function, denoted $\varphi \in \Phi_c(\Omega)$, if $t \mapsto \varphi(x, t)$ is left-continuous and convex for every $x \in \Omega$.

The subsets of $\Phi_w(\Omega)$ and $\Phi_c(\Omega)$ consisting of functions without dependence on the first variable (i.e., $\varphi(x, t) = \varphi(t)$) are denoted by Φ_w and Φ_c , respectively.

Since convexity implies (Inc) $_1$, we see that $\Phi_c(\Omega) \subset \Phi_w(\Omega)$. Let us now consider $\varphi \in \Phi_w(\Omega)$. We define the (left-continuous) inverse function of φ with respect to t by

$$\varphi^{-1}(x, t) := \inf\{\tau \geq 0 : \varphi(x, \tau) \geq t\}.$$

If φ is strictly increasing and continuous in t , then φ^{-1} is the usual inverse function. We also define the conjugate function of φ by

$$\varphi^*(x, t) := \sup_{s \geq 0} (st - \varphi(x, s)).$$

From this definition, it follows that *Young's inequality*

$$ts \leq \varphi(x, t) + \varphi^*(x, s)$$

holds for all $s, t \geq 0$. If φ satisfies $(\mathbf{aInc})_p$ or $(\mathbf{aDec})_q$ for some $p, q > 1$, then φ^* satisfies $(\mathbf{aDec})_{p'}$ or $(\mathbf{aInc})_{q'}$, respectively [34, Proposition 2.4.9]. For simplicity, we write

$$\varphi_{B_r}^+(t) := \sup_{x \in B_r \cap \Omega} \varphi(x, t) \quad \text{and} \quad \varphi_{B_r}^-(t) := \inf_{x \in B_r \cap \Omega} \varphi(x, t).$$

If $\varphi \in \Phi_c(\Omega)$, then $(\varphi^*)^* = \varphi$ [29, Theorem 2.2.6] and that there exists an increasing and right-continuous function $\varphi' : \Omega \times [0, \infty) \rightarrow [0, \infty)$, the so-called (right-)derivative of φ , such that

$$\varphi(x, t) = \int_0^t \varphi'(x, s) ds.$$

We recall some results related to this φ' .

Proposition 2.3 (Proposition 3.6, [37]). *Let $\gamma > 0$ and $\varphi \in \Phi_c(\Omega)$.*

- (1) *If φ' satisfies $(\mathbf{Inc})_\gamma$, $(\mathbf{Dec})_\gamma$, $(\mathbf{aInc})_\gamma$ or $(\mathbf{aDec})_\gamma$, then φ satisfies $(\mathbf{Inc})_{\gamma+1}$, $(\mathbf{Dec})_{\gamma+1}$, $(\mathbf{aInc})_{\gamma+1}$ or $(\mathbf{aDec})_{\gamma+1}$, respectively, with the same constant $L \geq 1$.*
- (2) *If φ satisfies $(\mathbf{aDec})_\gamma$, then $(2^{\gamma+1}L)^{-1}t\varphi'(x, t) \leq \varphi(x, t) \leq t\varphi'(x, t)$.*
- (3) *If φ' satisfies $(\mathbf{A0})$ and $(\mathbf{aDec})_\gamma$ with constant $L \geq 1$, then φ also satisfies $(\mathbf{A0})$, with constant depending on L and γ .*
- (4) *$\varphi^*(x, \varphi'(x, t)) \leq t\varphi'(x, t)$.*

For $\varphi \in \Phi_w(\Omega)$, the *generalized Orlicz space* (also known as the *Musielak–Orlicz space*) is defined by

$$L^\varphi(\Omega) := \{f \in L_{\text{loc}}^1(\Omega) : \|f\|_{L^\varphi(\Omega)} < \infty\},$$

with the (Luxemburg) norm

$$\|f\|_{L^\varphi(\Omega)} := \inf \left\{ \lambda > 0 : \varrho_{L^\varphi(\Omega)}\left(\frac{f}{\lambda}\right) \leq 1 \right\}, \quad \text{where} \quad \varrho_{L^\varphi(\Omega)}(f) := \int_\Omega \varphi(x, |f|) dx.$$

We denote by $W^{1,\varphi}(\Omega)$ the set of functions $f \in W_{\text{loc}}^{1,1}(\Omega)$ with $\|f\|_{W^{1,\varphi}(\Omega)} := \|f\|_{L^\varphi(\Omega)} + \| |Df| \|_{L^\varphi(\Omega)} < \infty$. If φ satisfies (\mathbf{aDec}) , then we note that $f \in L^\varphi(\Omega)$ if and only if $\varrho_{L^\varphi(\Omega)}(f) < \infty$. The spaces $L^\varphi(\Omega)$ and $W^{1,\varphi}(\Omega)$ are reflexive Banach spaces when φ satisfies $(\mathbf{A0})$, (\mathbf{aInc}) and (\mathbf{aDec}) . We denote by $W_0^{1,\varphi}(\Omega)$ the closure of $C_0^\infty(\Omega)$ in $W^{1,\varphi}(\Omega)$. For more information on generalized Orlicz and Orlicz–Sobolev spaces, we refer to the monographs [19, 34] and also [29, Chapter 2].

In recent years, we have studied regularity theory for the general equations $(\mathbf{div}A)$ with quasi-isotropic (p, q) -growth condition [35, 37, 38, 39, 41]. In these papers, regularity conditions for the growth function $\varphi \in \Phi_w(\Omega)$ or a relevant function with respect to the space variable x is given in terms of point-wise oscillation. This is in contrast to (p, q) -growth approach, where usually the one assumes that $\frac{q}{p}$ is small, e.g. [12], although see also [23]. Let us recall these assumptions and the regularity results in [39]. We have made a slight alteration in that $\omega(r)|\xi|$ previously lacked the $|\xi|$; this does not affect which G satisfy the condition, but it does impact the functions ω and $\tilde{\omega}$, see Example 4.1.

Definition 2.4. Let $G : \Omega \times \mathbb{R}^M \rightarrow \mathbb{R}^N$ with $M, N \in \mathbb{N}$, $r \in (0, 1]$, and $\omega, \tilde{\omega} : [0, 1] \rightarrow [0, L]$ with $L > 0$. Consider the claim

$$|G(x, \xi) - G(y, \xi)| \leq \tilde{\omega}(r)(|G(y, \xi)| + \omega(r)|\xi|) \quad \text{when } |G(y, \xi)| \in [0, |B_r|^{-1}]$$

for all $x, y \in B_r$, $B_{2r} \subset \Omega$ and $\xi \in \mathbb{R}^M$. We say that G satisfies

- (A1) if the claim holds with $\omega = \tilde{\omega} \equiv L$;
- (SA1) if there exists a modulus of continuity ω such that the claim holds with $\tilde{\omega} \equiv L$;
- (VA1) if there exists a modulus of continuity $\omega = \tilde{\omega}$ such that the claim holds;
- (wVA1) if, for every $\varepsilon > 0$, G satisfies (VA1) with the range condition replaced by $|G(y, \xi)|^{1+\varepsilon} \in [0, |B_r|^{-1}]$, with moduli of continuity $\omega_\varepsilon := \omega = \tilde{\omega}$ depending on ε , but with a common L independent of ε .

We also use the definition for $\varphi : \Omega \times [0, \infty) \rightarrow [0, \infty)$ by interpreting $\varphi(x, \xi) = \varphi(x, |\xi|)$.

We refer to Section 8 of [39] for examples of functions satisfying these conditions. With these point-wise assumptions, we proved maximal regularity results in [37, 38, 39].

Theorem 2.5 (Theorem 1.2, [39]). *Let A satisfy the quasi-isotropic (p, q) -growth condition and let $u \in W_{\text{loc}}^{1,1}(\Omega)$ with $|A^{(-1)}(\cdot, Du)| \in L_{\text{loc}}^1(\Omega)$ be a local weak solution to $(\text{div} A)$.*

- (1) *If $A^{(-1)}$ satisfies (A1), then $u \in C_{\text{loc}}^{0,\alpha}(\Omega)$ for some $\alpha \in (0, 1)$ depending on n, p, q and L .*
- (2) *If $A^{(-1)}$ satisfies (wVA1), then $u \in C_{\text{loc}}^{0,\alpha}(\Omega)$ for every $\alpha \in (0, 1)$.*
- (3) *If $A^{(-1)}$ satisfies (wVA1) with Hölder-continuous ω_ε for every $\varepsilon > 0$, then $u \in C_{\text{loc}}^{1,\alpha}(\Omega)$ for some $\alpha \in (0, 1)$ depending on n, p, q and L .*

3. MEAN OSCILLATION CONDITIONS

We introduce the mean oscillation variants of the (A1)-condition, specifically, the *vanishing mean-A1 condition* (VMA1) and the *Dini mean-A1 condition* (DMA1).

Definition 3.1. Let $G : \Omega \times \mathbb{R}^M \rightarrow \mathbb{R}^N$ with $M, N \in \mathbb{N}$, $r \in (0, 1]$, $\omega : [0, 1] \rightarrow [0, 1]$, and

$$\tilde{\theta}(x, B_r) := \sup_{\xi \in D(B_r)} \frac{|G(x, \xi) - G_{B_r}(\xi)|}{|G_{B_r}(\xi)| + \omega(r)|\xi|} \quad \text{for } x \in B_r,$$

where $D(B_r) := \{\xi \in \mathbb{R}^M : |G_{B_r}(\xi)| \leq |B_r|^{-1}\}$. For $\gamma \geq 1$, we consider

$$\theta_\gamma(r) := \left(\sup_{B_{2r} \subset \Omega} \int_{B_r} \tilde{\theta}(x, B_r)^\gamma dx \right)^{\frac{1}{\gamma}}.$$

We say that G satisfies:

- (VMA1) if there exists a nondecreasing ω such that $\lim_{r \rightarrow 0^+} [\theta_1(r) + \omega(r)] = 0$.
- (DMA1) $_\gamma$ if there exists a nondecreasing ω such that $\int_0^1 [\theta_\gamma(r) + \omega(r)] \frac{dr}{r} < \infty$.

We refer to Section 4 for several examples of energies satisfying these conditions. There we show that all the cases from Table 1 are included as special cases. By Hölder's inequality, the (DMA1) $_\gamma$ is monotone in γ . Furthermore, if the case $\gamma = \infty$ is understood as a supremum, then θ_∞ is closely related to the quantity in the point-wise conditions of Definition 2.4.

Remark 3.2. In Definition 3.1, we consider the γ -mean of $\tilde{\theta}$ for some $\gamma \geq 1$. Proposition 3.8 implies that if we assume that θ is nondecreasing, then it suffices to consider the case $\gamma = 1$.

Remark 3.3. Similarly to the vanishing condition (VA1), we can also consider weak versions (wDMA1) and (wVMA1) of (DMA1) and (VMA1). However, we are not aware of any examples where these conditions would be needed. Thus we will for the sake of simplicity not consider them in what follows. Furthermore, we could define a point-wise condition (DA1) of Dini-type, but this is already obsolete since we directly handle the more general condition (DMA1).

We show that the (DMA1)₁ condition also implies the continuity of $G(x, \xi)$ in the x variable and the (SA1) condition. For this we need Spanne's result relating modulus of mean continuity with modulus of continuity:

Lemma 3.4 (Corollary 1 with its remark, [55]). *Assume that $f \in L^1(\Omega)$ satisfies the Dini mean continuity condition*

$$\int_0^1 \bar{\omega}_f(r) \frac{dr}{r} < \infty \quad \text{where} \quad \bar{\omega}_f(r) := \sup_{B_r \subset \Omega} \int_{B_r} |f(x) - (f)_{B_r}| dx.$$

Then f is continuous with modulus $\omega_f(r) = c \int_0^r \bar{\omega}_f(s) \frac{ds}{s}$ for some $c > 0$.

From this lemma we can get an intuition of how the point-wise and mean moduli are related. First, if $\bar{\omega}_f(r) \lesssim r^\beta$, then also $\omega_f(r) \lesssim r^\beta$, so there is no difference between the cases here. However, if $\bar{\omega}_f(r) = (\log \frac{1}{r})^{-\alpha}$, $\alpha > 1$, then $\omega_f(r) \approx (\log \frac{1}{r})^{1-\alpha}$, so there is a loss of a logarithm between the cases. In particular, if $\alpha \in (1, 2]$, then the point-wise modulus ω_f does not satisfy the Dini condition even though the mean oscillation modulus $\bar{\omega}_f$ does.

Proposition 3.5. *Let $G : \Omega \times \mathbb{R}^M \rightarrow \mathbb{R}^N$ with $M, N \in \mathbb{N}$ satisfy (A0), (aInc)_p and (aDec)_q for some $1 \leq p \leq q$, as well as (DMA1)₁.*

- (1) *For each $\xi \in \mathbb{R}^M$, $G(\cdot, \xi)$ is continuous.*
- (2) *G satisfies (SA1) with the same modulus of continuity ω and the relevant constant $L \geq 1$ depending on n, p, q, L , and θ_1 .*

Proof. Note that $G_U^+(\xi) := \sup_{x \in U} |G(x, \xi)|$ is finite by (A0) and (aDec)_q.

(1) Fix $\xi \in \mathbb{R}^M \setminus \{0\}$. Then there exists $r_0 > 0$ such that $G_\Omega^+(\xi) \leq |B_{r_0}|^{-1}$. For $r \in (0, r_0]$, from the definition of $\tilde{\theta}$ in Definition 3.1, $|G(x, \xi) - G_{B_r}(\xi)| \leq (G_\Omega^+(\xi) + |\xi|)\tilde{\theta}(x, B_r)$ for every $x \in B_r$ and $B_{2r} \subset \Omega$. Hence

$$\begin{aligned} \sup_{B_{2r} \subset \Omega} \int_{B_r} |G(x, \xi) - G_{B_r}(\xi)| dx &\leq (G_\Omega^+(\xi) + |\xi|) \sup_{B_{2r} \subset \Omega} \int_{B_r} \tilde{\theta}(x, B_r) dx \\ &= (G_\Omega^+(\xi) + |\xi|)\theta_1(r). \end{aligned}$$

Therefore, by Lemma 3.4, the (DMA1)₁ condition yields the continuity of $G(\cdot, \xi)$.

(2) Fix $B_{2r} \subset \Omega$ with small $r > 0$ to be determined later. Consider $y \in B_r$ and $\xi \in \mathbb{R}^n$ with $G_{B_r}^+(\xi) \leq |B_r|^{-1}$. Then for any ball $B_\rho \subset B_r$, since $|G_{B_\rho}(\xi)| \leq G_{B_r}^+(\xi) \leq |B_r|^{-1} \leq |B_\rho|^{-1}$, it follows from Definition 3.1 that

$$|G(x, \xi) - G_{B_\rho}(\xi)| \leq (|G_{B_\rho}(\xi)| + \omega(\rho)|\xi|)\tilde{\theta}(x, B_\rho) \lesssim (G_{B_r}^+(\xi) + \omega(r)|\xi|)\tilde{\theta}(x, B_\rho)$$

for any $x \in B_\rho$. Hence by Definition 3.1,

$$\sup_{B_\rho \subset B_r} \int_{B_\rho} |G(x, \xi) - G_{B_\rho}(\xi)| dx \lesssim (G_{B_r}^+(\xi) + \omega(r)|\xi|) \theta_1(\rho).$$

Therefore, in view of Lemma 3.4, the $(\text{DMA1})_1$ condition yields that

$$|G(x, \xi) - G(z, \xi)| \leq c_1 (G_{B_r}^+(\xi) + \omega(r)|\xi|) \int_0^r \theta_1(\rho) \frac{d\rho}{\rho}, \quad \text{for any } x, z \in B_r$$

for some constant $c_1 \geq 0$ depending on p, q, L and θ_1 . Choose $r \in (0, r_1]$, where r_1 is determined by $\int_0^{r_1} \theta(\rho) \frac{d\rho}{\rho} = \frac{1}{2c_1}$. Then

$$G_{B_r}^+(\xi) - G_{B_r}^-(\xi) \leq \sup_{x, z \in B_r} |G(x, \xi) - G(z, \xi)| \leq \frac{1}{2}(G_{B_r}^+(\xi) + \omega(r)|\xi|),$$

which implies that

$$G_{B_r}^+(\xi) \leq 2G_{B_r}^-(\xi) + \omega(r)|\xi|.$$

This implies the desired (SA1) -inequality, when $G_{B_r}^+(\xi) \leq |B_r|^{-1}$.

Suppose then that $G_{B_r}^-(\xi) \leq |B_r|^{-1}$ and let $s \in (0, 1]$ be the largest number with $G_{B_r}^+(s\xi) \leq |B_r|^{-1}$. The case $s = 1$ was handled above. If $s < 1$, then $G_{B_r}^+(s\xi) = |B_r|^{-1}$ and it follows from the earlier case that

$$|B_r|^{-1} = G_{B_r}^+(s\xi) \leq 2G_{B_r}^-(s\xi) + \omega(r)|s\xi|.$$

Since $r \leq 1$, it follows from (A0) that $|\xi| \gtrsim 1$. Then (A0) and $\omega(r) \leq 1$ yield $\omega(r)|s\xi| \lesssim G_{B_r}^-(s\xi)$. Using also $(\text{aDec})_q$, we find that

$$|B_r|^{-1} \leq 2G_{B_r}^-(s\xi) + \omega(r)|s\xi| \leq cG_{B_r}^-(s\xi) \leq c_2 s^p G_{B_r}^-(\xi) \leq c_2 s^p |B_r|^{-1}.$$

Hence $s \geq c_2^{1/p}$. Returning to the earlier case, we then conclude that

$$G_{B_r}^+(\xi) \leq Ls^{-q} G_{B_r}^+(s\xi) \leq Ls^{-q} (2G_{B_r}^-(s\xi) + \omega(r)|s\xi|) \leq 2Lc_2^{-q/p} (G_{B_r}^-(\xi) + \omega(r)|\xi|),$$

which implies that (SA1) -inequality with the correct assumption $G_{B_r}^-(\xi) \leq |B_r|^{-1}$ for $r \in (0, r_1]$. When $r \in (r_1, 1]$, we obtain the conclusion using a chain of a fixed number $\lceil \frac{1}{r_1} \rceil$ of balls of radius r_1 and the above argument. \square

The function θ in the above definition is not assumed to be increasing. This makes it harder to estimate geometric series based on θ , but the next lemma shows that it is possible.

Lemma 3.6. *Let $G : \Omega \times \mathbb{R}^M \rightarrow \mathbb{R}^N$ satisfy (A0) , $(\text{aInc})_p$ and $(\text{aDec})_q$ for some $1 \leq p \leq q$ and $(\text{DMA1})_\gamma$ for some $\gamma \geq 1$. There exists $\delta_1 = \delta_1(n, p, q, \theta_\gamma) \in (0, 1)$ such that for every $\delta \in (0, \delta_1)$ and $r \in (0, 1)$ with $|B_{2r}| \leq 1$,*

$$\sum_{k=0}^{\infty} \theta_\gamma(\delta^k r) \leq \frac{c}{\delta^{\frac{2n}{\gamma}+1} \log(\delta^{-1})} \int_0^r \theta_\gamma(\rho) \frac{d\rho}{\rho}$$

for some $c = c(n, p, q, \gamma, \theta_\gamma) > 0$.

Proof. Let $\tilde{r} := \delta^k r$ with $k \geq 1$ and $\rho \in (\delta\tilde{r}, \tilde{r})$. Fix $B_{2r} \subset \Omega$. Since G is satisfies (SA1) by Proposition 3.5(2),

$$|G(x, \xi)| \lesssim G_{B_{3\tilde{r}/2}}^-(\xi) + \omega(\tilde{r})$$

for any $x, y \in B_{3\tilde{r}/2}$ and $\xi \in \mathbb{R}^M$ with $G_{B_{3\tilde{r}/2}}^-(\xi) \leq |B_{\tilde{r}}|^{-1}$. Fix $\xi \in \mathbb{R}^M$ such that $|G_{B_{\tilde{r}}}(\xi)| \leq |B_{\tilde{r}}|^{-1}$. Then the preceding inequality yields for any $B_\rho(z) \subset B_{3\tilde{r}/2}$ with $z \in \overline{B_{\tilde{r}}}$ that

$$(3.7) \quad |G_{B_\rho(z)}(\xi)| \leq \tilde{L}(|G_{B_{\tilde{r}}}(\xi)| + \omega(\tilde{r})),$$

for some $\tilde{L} \geq 1$ depending on n, p, q and θ_1 . Therefore, since $\omega(r) \leq 1$,

$$|G_{B_\rho(z)}(\xi)| \leq \tilde{L} (|G_{B_{\tilde{r}}}(\xi)| + \omega(\tilde{r})) \leq (\tilde{L} + 1) |B_{\tilde{r}}|^{-1} \leq |B_\rho|^{-1}$$

provided that $\rho \leq \min\{(\tilde{L} + 1)^{-1/n}, 1/2\} r$. Thus $D(B_{\tilde{r}}) \subset D(B_\rho(z))$ for $z \in \overline{B_{\tilde{r}}}$.

Denote $H_\xi(x) := \frac{G(x, \xi)}{|G_{B_{\tilde{r}}}(\xi)| + \omega(r)|\xi|}$. We connect the integrals of H_ξ in the two balls by a chain argument. We cover $B_{\tilde{r}}$ with $c(n)\delta^{-n}$ balls B^i of radius $\frac{\rho}{2}$, whose centers are in $\overline{B_{\tilde{r}}}$, with the property that every $x, y \in B_{\tilde{r}}$ with $|x - y| \leq \frac{\rho}{4}$ belong to some ball in the cover. For $x \in B^i$ and $y \in B^j$ with $i \neq j$ and $|x - y| > \frac{\rho}{4}$ we use the estimate

$$\begin{aligned} |H_\xi(x) - H_\xi(y)| &\leq |H_\xi(x) - (H_\xi)_{B_{\rho/2}(x_0)}| + \sum_{k=0}^m |(H_\xi)_{B_{\rho/2}(x_{k-1})} - (H_\xi)_{B_{\rho/2}(x_k)}| \\ &\quad + |(H_\xi)_{B_{\rho/2}(x_m)} - H_\xi(y)| \end{aligned}$$

where $x_0 = x$, $x_m = y$ and $x_k \in [x, y] := \{tx + (1-t)y : t \in [0, 1]\}$ for $k = 1, \dots, m-1$ satisfy $\frac{\rho}{4} < |x_m - x_{m-1}| \leq \frac{\rho}{2}$ for $k = 1, 2, \dots, m$. Note that $m \leq 13\delta^{-1}$. Using $D(B_{\tilde{r}}) \subset D(B_\rho(x_k))$ and Hölder's inequality in the second step, we have

$$\begin{aligned} &\int_{B^i} \int_{B^j} \sup_{\xi \in D(B_{\tilde{r}})} |H_\xi(x) - H_\xi(y)|^\gamma dy dx \\ &\leq \int_{B^i} \sup_{\xi \in D(B_{\tilde{r}})} |H_\xi(x) - (H_\xi)_{B_{\rho/2}(x_0)}|^\gamma dx + \int_{B^j} \sup_{\xi \in D(B_{\tilde{r}})} |(H_\xi)_{B_{\rho/2}(x_{m_0})} - H_\xi(y)|^\gamma dy \\ &\quad + \sum_{k=1}^m \sup_{\xi \in D(B_{\tilde{r}})} |(H_\xi)_{B_{\rho/2}(x_{k-1})} - (H_\xi)_{B_{\rho/2}(x_k)}|^\gamma \\ &\lesssim \sum_{k=0}^m \int_{B_\rho(x_k)} \int_{B_\rho(x_k)} \sup_{\xi \in D(B_\rho(x_k))} |H_\xi(x) - H_\xi(y)|^\gamma dy dx. \end{aligned}$$

We can connect integrals over $\tilde{\theta}$ and H_ξ . We estimate

$$\int_{B_{\tilde{r}}} \tilde{\theta}(x, B_{\tilde{r}})^\gamma dx \approx \int_{B_{\tilde{r}}} \int_{B_{\tilde{r}}} \sup_{\xi \in D(B_{\tilde{r}})} |H_\xi(x) - H_\xi(y)|^\gamma dy dx$$

and, using also (3.7), we see that

$$\int_{B_\rho(x_k)} \int_{B_\rho(x_k)} \sup_{\xi \in D(B_\rho(x_k))} |H_\xi(x) - H_\xi(y)|^\gamma dy dx \lesssim \int_{B_\rho(x_k)} \tilde{\theta}(x, B_\rho(x_k))^\gamma dx.$$

Combining these with the estimate from the previous paragraph and $m \lesssim \frac{1}{\delta}$, we find that

$$\int_{B_{\tilde{r}}} \tilde{\theta}(x, B_{\tilde{r}})^\gamma dx \lesssim \sum_{k=0}^m \int_{B_\rho(x_k)} \tilde{\theta}(x, B_\rho(x_k))^\gamma dx \lesssim (\delta^{-1}\theta_\gamma(\rho))^\gamma.$$

Using this estimate for the second inequality, we conclude that

$$\begin{aligned} &\int_{B_{\tilde{r}}} \int_{B_{\tilde{r}}} \sup_{\xi \in D(B_{\tilde{r}})} \left(\frac{|G(x, \xi) - G(y, \xi)|}{|G_{B_{\tilde{r}}}(\xi)| + \omega(\tilde{r})|\xi|} \right)^\gamma dy dx \\ &\lesssim \delta^{-2n} \sum_{i,j} \int_{B^i} \int_{B^j} \sup_{\xi \in D(B_{\tilde{r}})} \left(\frac{|G(x, \xi) - G(y, \xi)|}{|G_{B_{\tilde{r}}}(\xi)| + \omega(\tilde{r})|\xi|} \right)^\gamma dy dx \lesssim \delta^{-2n-\gamma}\theta_\gamma(\rho)^\gamma. \end{aligned}$$

We have shown that $\theta_\gamma(\tilde{r}) \lesssim \delta^{-(2n/\gamma+1)}\theta_\gamma(\rho)$ for $\rho \in (\delta\tilde{r}, \sqrt{\delta\tilde{r}})$ with $\tilde{r} = \delta^k r$ and $\delta \leq \min\{(\tilde{L} + 1)^{-1/n}, 1/2\} =: \delta_1$. Thus we calculate

$$\int_{\delta^{k+1}r}^{\delta^k r} \theta_\gamma(\rho) \frac{d\rho}{\rho} \gtrsim \delta^{\frac{2n}{\gamma}+1} \theta_\gamma(\delta^k r) \int_{\delta^{k+1}r}^{\sqrt{\delta}\delta^k r} \frac{d\rho}{\rho} \approx \delta^{\frac{2n}{\gamma}+1} \log(\delta^{-1}) \theta_\gamma(\delta^k r).$$

Adding the inequalities over k yields the claim. \square

We derive some properties of the supremal counterpart θ_γ^* of θ_γ .

Proposition 3.8. *Let $G : \Omega \times \mathbb{R}^M \rightarrow \mathbb{R}^N$ satisfy (A0), (aInc) $_p$ and (aDec) $_q$ for some $1 \leq p \leq q$ as well as (SA1). With θ_γ from Definition 3.1 for non-decreasing ω and $\gamma \geq 1$ we define $\theta_\gamma^*(r) := \sup_{\rho \in (0, r]} \theta_\gamma(\rho)$. Then $\theta_\gamma^* \leq L_\gamma \theta_1^*$ for some $L_\gamma > 0$ depending on n, p, q and γ . We also note two consequences of this.*

- (1) If $\int_0^1 \theta_1^*(\rho) \frac{d\rho}{\rho} < \infty$, then (DMA1) $_\gamma$ holds for any $\gamma \geq 1$.
- (2) If G satisfies (SA1) and (VMA1), then $\lim_{r \rightarrow 0} \theta_\gamma^*(r) = 0$ for any $\gamma \geq 1$.

Proof. We follow ideas from the proof of the John–Nirenberg inequality. In Definition 3.1, the supremum is taken over balls, so we apply the Vitali covering lemma instead of the Calderón–Zygmund decomposition. Fix $r \in (0, 1]$ and abbreviate $\theta^* := \theta_1^*(r)$. It follows from the definition of θ^* that

$$(3.9) \quad \int_{B_\rho(y)} \tilde{\theta}(x, B_\rho(y)) dx \leq \theta^* \quad \text{for all } B_{2\rho}(y) \subset \Omega \text{ with } \rho \in (0, r].$$

Let $\delta \in (0, \frac{1}{10})$ be a small constant to be determined later, and $\beta_n > 1$ be a constant depending only on n such that $|B_\rho| \leq \beta_n |B_r \cap B_\rho(y)|$ for any $y \in B_r$ and $\rho \in (0, 2r]$. Denote $E(t, U) := \{x \in U \cap B_r : \tilde{\theta}(x, U) > t\}$.

We first estimate the measure of the set $E(\beta_n \delta^{-n} \theta^*, B_r)$. Observe that for every $y \in B_r$ and $\rho \in (\delta r, r]$,

$$\int_{B_r \cap B_\rho(y)} \tilde{\theta}(x, B_r) dx < \beta_n \delta^{-n} \int_{B_r} \tilde{\theta}(x, B_r) dx \leq \beta_n \delta^{-n} \theta^*,$$

and that for almost every $y \in E(\beta_n \delta^{-n} \theta^*, B_r)$

$$\lim_{\rho \rightarrow 0^+} \int_{B_r \cap B_\rho(y)} \tilde{\theta}(x, B_r) dx > \beta_n \delta^{-n} \theta^*.$$

Then by the continuity of the integral with respect to ρ , for almost every $y \in E(\beta_n \delta^{-n} \theta^*, B_r)$ one can find $\rho_y \in (0, \delta r]$ such that

$$\int_{B_r \cap B_{\rho_y}(y)} \tilde{\theta}(x, B_r) dx = \beta_n \delta^{-n} \theta^* \quad \text{and} \quad \int_{B_r \cap B_\rho(y)} \tilde{\theta}(x, B_r) dx < \beta_n \delta^{-n} \theta^* \quad \forall \rho \in (\rho_y, r].$$

Therefore, by the Vitali covering lemma, there exist $y_j^1 \in E(\beta_n \delta^{-n} \theta^*, B_r)$ and $\rho_j^1 \in (0, \delta r]$ for $j = 1, 2, \dots$ such that the balls $B_j^1 := B_{\rho_j^1}(y_j^1)$ are mutually disjoint,

$$\bigcup_{j=1}^{\infty} 5B_j^1 \supset E(\beta_n \delta^{-n} \theta^*, B_r) \setminus \mathcal{N}^0,$$

where $5B_j^1 := B_{5\rho_j^1}(y_j^1)$ and \mathcal{N}^0 is some measure zero set,

$$\int_{B_r \cap B_j^1} \tilde{\theta}(x, B_r) dx = \beta_n \delta^{-n} \theta^* \quad \text{and} \quad \int_{B_r \cap 5B_j^1} \tilde{\theta}(x, B_r) dx < \beta_n \delta^{-n} \theta^*;$$

the earlier setup is applicable, since $5\rho_j^1 \leq 5\delta r \leq r$. The equation in above together with (3.9) implies

$$\sum_{j=1}^{\infty} |B_j^1 \cap B_r| = \frac{\delta^n}{\beta_n \theta^*} \sum_{j=1}^{\infty} \int_{B_r \cap B_j^1} \tilde{\theta}(x, B_r) dx \leq \frac{\delta^n}{\beta_n \theta^*} \int_{B_r} \tilde{\theta}(x, B_r) dx \leq \frac{\delta^n}{\beta_n} |B_r|,$$

so that

$$|E(\beta_n \delta^{-n} \theta^*, B_r)| \leq \sum_{j=1}^{\infty} |5B_j^1 \cap B_r| \leq 5^n \sum_{j=1}^{\infty} |B_j^1| \leq \beta_n 5^n \sum_{j=1}^{\infty} |B_j^1 \cap B_r| \leq (5\delta)^n |B_r|.$$

We can use the same procedure in the ball $5B_j^1$ to conclude that

$$|E(\beta_n \delta^{-n} \theta^*, 5B_j^1)| \leq \sum_{l=1}^{\infty} |5B_{j,l}^1 \cap B_j^1| \leq (5\delta)^n |5B_j^1| = (5^2\delta)^n |B_j^1|,$$

where $B_{j,l}^1$, $l \in \mathbb{N}$, are mutually disjoint balls whose five-fold dilates cover B_j^1 . This holds in every B_j^1 , so we estimate

$$\begin{aligned} \sum_{j=1}^{\infty} |E(\beta_n \delta^{-n} \theta^*, 5B_j^1)| &\leq \sum_{j=1}^{\infty} \sum_{l=1}^{\infty} |5B_{j,l}^1 \cap B_j^1| \leq (5^2\delta)^n \beta_n \sum_{j=1}^{\infty} |B_j^1| \\ &\leq (5\delta)^n \beta_n \sum_{j=1}^{\infty} |5B_j^1 \cap B_r| \leq (5\delta)^{2n} \beta_n |B_r|. \end{aligned}$$

Denote by (B_j^2) the sequence of balls whose five-fold dilates cover the first generation balls B_j^1 . Repeating this process, for each $k \in \mathbb{N}$, we get a sequence of balls (B_j^k) in B_{2r} such that $\bigcup_{j=1}^{\infty} 5B_j^k \supset E(\beta_n \delta^{-n} \theta^*, B_r) \setminus \mathcal{N}^k$ for some measure zero set \mathcal{N}^k ,

$$(3.10) \quad \int_{5B_j^{k-1} \cap 5B_j^k} \tilde{\theta}(x, 5B_j^{k-1}) dx < \beta_n \delta^{-n} \theta^*,$$

where \tilde{j} is such that $B_{\tilde{j}}^{k-1}$ is the predecessor of B_j^k , and

$$(3.11) \quad \sum_{j=1}^{\infty} |E(\beta_n \delta^{-n} \theta^*, 5B_j^k)| \leq (5\delta)^{(k+1)n} \beta_n^k |B_r|.$$

We also note that the condition $\delta < \frac{1}{10}$ implies $B_j^k \subset B_{2r}$ for any k and j . We impose on δ the condition $(5\delta)^n \beta_n < 1$.

Note that, by the same argument as in the beginning of the proof of Lemma 3.6, we see from the (SA1) condition of G that $|G_{B_r}(\xi)| + \omega(r) \approx |G_{B_\rho(y)}(\xi)| + \omega(r)$ for every $B_\rho(y) \subset B_{3r/2}$ and $\xi \in D(B_\rho(y))$, and that $D(B_r) \subset D(B_\rho(y))$ for any $B_\rho(y) \subset B_{3r/2}$ with $y \in \overline{B_r}$ and $\rho \in (0, \delta r]$ by choosing sufficiently small δ , see the proof of the previous lemma. Now, fix $k \in \mathbb{N}$. Let $B^k := 5B_{j_k}^k$, $B^l := 5B_{j_l}^l$ be its predecessor at level $l = 1, 2, \dots, k-1$, ρ_l is the radius of B^l , and $B^0 = B_r$. Note that $|B^l| \leq \beta_n |B^l \cap B^{l-1}|$. Then for $x \in B^k$,

$$\begin{aligned} |G(x, \xi) - G_{B_r}(\xi)| &\leq |G(x, \xi) - G_{B^k \cap B^{k-1}}(\xi)| + |G_{B^k \cap B^{k-1}}(\xi) - G_{B^{k-1} \cap B^{k-2}}(\xi)| \\ &\quad + \dots + |G_{B^2 \cap B^1}(\xi) - G_{B^1 \cap B_r}(\xi)| + |G_{B^1 \cap B_r}(\xi) - G_{B_r}(\xi)| \\ &\leq |G(x, \xi) - G_{B^k}(\xi)| + \sum_{l=1}^k \left(|G_{B^k \cap B^{l-1}}(\xi) - G_{B^l}(\xi)| + |G_{B^l \cap B^{l-1}}(\xi) - G_{B^{l-1}}(\xi)| \right). \end{aligned}$$

Since ω is non-decreasing, we have $|G_{B_r}(\xi)| + \omega(r) \gtrsim |G_{B_\rho(y)}(\xi)| + \omega(\rho)$ from before. Hence, using the above observations, (3.9) and (3.10),

$$\begin{aligned}\tilde{\theta}(x, B_r) &= \sup_{\xi \in D(B_r)} \frac{|G(x, \xi) - G_{B_r}(\xi)|}{|G_{B_r}(\xi)| + \omega(r)|\xi|} \\ &\lesssim \tilde{\theta}(x, B^k) + \sum_{j=1}^k \left(\beta_n \int_{B^j} \tilde{\theta}(x, B^j) dx + \int_{B^j \cap B^{j-1}} \tilde{\theta}(x, B^{j-1}) dx \right) \\ &\lesssim \tilde{\theta}(x, B^k) + k\beta_n(\delta^{-n} + 1)\theta^*.\end{aligned}$$

Let $\alpha_1 \geq 1$ denote the implicit constant in the above estimate. Then it follows that $E(2\delta^{-n}\beta_n\alpha_1(k+1)\theta^*, B_r) \cap B^k \subset E(2\beta_n\delta^{-n}\theta^*, B^k)$ for each ball $B^k = 5B_j^k$, $j \in \mathbb{N}$. Therefore, by the covering property and (3.11), we obtain

$$|E(2\beta_n\delta^{-n}\alpha_1(k+1)\theta^*, B_r)| \leq \sum_{j=1}^{\infty} |E(2\beta_n\delta^{-n}\theta^*, 5B_j^k)| \leq (5\delta)^{(k+1)n} \beta_n^k |B_r|.$$

Finally, when $2\beta_n\delta^{-n}\alpha_1(k+1)\theta^* \leq t < 2\beta_n\delta^{-n}\alpha_1(k+2)\theta^*$ for some $k \in \mathbb{N}$, we find that

$$\begin{aligned}|\{x \in B_r : \tilde{\theta}(x, B_r) > t\}| &= |E(t, B_r)| \leq (5\delta)^{n(k+1)} \beta_n^k |B_r| \\ &\leq (5\delta)^n \exp\left(-\underbrace{\frac{\delta^n}{6\beta_n\alpha_1} \ln\left(\frac{1}{(5\delta)^n \beta_n}\right)}_{=:\alpha_2} \frac{t}{\theta^*}\right) |B_r|.\end{aligned}$$

As in the John–Nirenberg inequality, this implies that

$$\begin{aligned}\int_{B_r} \tilde{\theta}(x, B_r)^\gamma dx &= \frac{1}{|B_r|} \int_0^\infty \gamma t^{\gamma-1} |\{x \in B_r : \tilde{\theta}(x, B_r) > t\}| dt \\ &\leq \gamma(5\delta)^n \int_0^\infty t^{\gamma-1} \exp\left(-\frac{\alpha_2}{\theta^*} t\right) dt = \gamma(5\delta)^n \Gamma(\gamma) \left(\frac{\theta^*}{\alpha_2}\right)^\gamma,\end{aligned}$$

where Γ is the gamma function. Thus $\theta_\gamma \leq L_\gamma \theta_1^*$. Claims (1) and (2) follow directly from this inequality. \square

4. EXAMPLES

In this section we consider what the Dini mean oscillation condition $(DMA1)_\gamma$ and the vanishing mean oscillation condition $(VMA1)$ means for specific model energies. The first example shows how they are related to $(VA1)$.

Example 4.1 (Point-wise conditions). Let $G : \Omega \times \mathbb{R}^M \rightarrow \mathbb{R}^N$ satisfy $(VA1)$ with $\tilde{\omega} = \omega = \omega_V$. Fix a ball $B_{2r} \subset \Omega$ and $\xi \in \mathbb{R}^M$ with $|G_{B_r}(\xi)| \leq |B_r|^{-1}$. Choosing $y \in B_r$ such that $G_{B_r}(\xi) = G(y, \xi)$, we see from $(VA1)$ that

$$|G(x, \xi) - G_{B_r}(\xi)| \leq \omega_V(r)(|G_{B_r}(\xi)| + \omega_V(r)|\xi|)$$

for every $x \in B_r$. Thus

$$\tilde{\theta}(x, B_r) = \sup_{\xi \in D(B_r)} \frac{|G(x, \xi) - G_{B_r}(\xi)|}{|G_{B_r}(\xi)| + \omega_V(r)|\xi|} \leq \omega_V(r),$$

so that $\theta_\gamma(r) \leq c\omega_V(r)$ for every $\gamma \geq 1$ and so G satisfies $(VMA1)$.

Moreover, if ω_V satisfies the Dini condition: $\int_0^1 \omega_V(r) \frac{dr}{r} < \infty$, then G satisfies $(DMA1)_\gamma$ for any $\gamma \geq 1$. In particular, if G satisfies $(VA1)$ with $\omega(r) = \tilde{\omega}(r) = r^\alpha$ for some $\alpha > 0$, which yields the $C^{1,\alpha}$ -regularity (Theorem 2.5), then G satisfies $(DMA1)_\gamma$ for any $\gamma \geq 1$.

If we consider the old version of (VA1) without the $|\xi|$ as described before Definition 2.4, then we can obtain similar conclusions based on the estimate

$$|G(x, \xi) - G_{B_r}(\xi)| \leq \omega_V(r) (|G_{B_r}(\xi)| + \omega_V(r)) \leq c\omega_V(r)^{\frac{1}{p'}} (|G_{B_r}(\xi)| + \omega_V(r)^{\frac{1}{p'}} |\xi|),$$

provided G satisfies (aInc) $_p$ with $p > 1$ (the inequality is proved like Proposition 3.6, [38], so we omit the details). Again, we find that G satisfies (VMA1) but now the modulus of continuity is $\omega_V(r)^{1/p'}$. The corresponding Dini condition is $\int_0^1 \omega_V(r)^{1/p'} \frac{dr}{r} < \infty$. This means that the old version does not give as precise control of the decay, which is the reason for the change to the new version of (VA1).

We next look at several special cases of the nonlinearity $A(x, \xi)$ in (divA) to see what the DMO condition entails. We will use the BMO-seminorm

$$\|f\|_{\text{BMO},r} := \sup_{0 < \rho \leq r} \sup_{B_\rho \subset \mathbb{R}^n} \int_{B_\rho} |f(x) - (f)_{B_\rho}| dx.$$

With the second supremum, this seminorm is increasing in r , so it corresponds to θ^* in Proposition 3.8; we could also consider a version where the supremum is taken only over balls of radius r . We say that f satisfies VMO (vanishing mean oscillation) if $\|f\|_{\text{BMO},r} \rightarrow 0$ when $r \rightarrow 0$, and DMO (Dini mean oscillation) if $\int_0^1 \|f\|_{\text{BMO},r} \frac{dr}{r} < \infty$.

Example 4.2 (Orlicz with coefficient). Let $\varphi \in \Phi_w$ and $a : \Omega \rightarrow [L^{-1}, L]$ for some $L \geq 1$. Define

$$A(x, \xi) = D_\xi[a(x)\varphi(|\xi|)] = a(x) \frac{\varphi'(|\xi|)}{|\xi|} \xi.$$

Since $(A^{(-1)})_{B_r}(\xi) = (a)_{B_r} \varphi'(|\xi|) \xi$ and $a \geq L^{-1}$,

$$\tilde{\theta}(x, B_r) \leq \frac{|A^{(-1)}(x, \xi) - (A^{(-1)})_{B_r}(\xi)|}{|(A^{(-1)})_{B_r}(\xi)|} \leq L|a(x) - (a)_{B_r}|, \quad \xi \in \mathbb{R}^n,$$

and, for any $\gamma \geq 1$,

$$\int_{B_r} \tilde{\theta}(x, B_r)^\gamma dx \lesssim \int_{B_r} |a(x) - (a)_{B_r}|^\gamma dx \lesssim \|a\|_{\text{BMO},r}^\gamma.$$

Therefore, $\theta_\gamma(r) \lesssim \|a\|_{\text{BMO},r}^\gamma$ and so $A^{(-1)}$ satisfies (DMA1) $_\gamma$ for any $\gamma \geq 1$ or (VMA1) if a satisfies DMO or VMO, respectively.

We next consider double phase energies. In the following example, the mean continuity does not give us anything new compared to the point-wise continuity modulus.

Example 4.3 (Double phase). Let $\varphi(x, t) = t^p + a(x)t^q$, where $1 < p \leq q$ and $a : \Omega \rightarrow [0, L]$, and define

$$A(x, \xi) := D_\xi[\varphi(x, |\xi|)] = p|\xi|^{p-2}\xi + qa(x)|\xi|^{q-2}\xi.$$

Then for $\varepsilon \geq 0$,

$$\begin{aligned} \tilde{\theta}_\varepsilon(x, B_r) &:= \sup_{\xi \in D_\varepsilon(B_r)} \frac{|A^{(-1)}(x, \xi) - (A^{(-1)})_{B_r}(\xi)|}{|(A^{(-1)})_{B_r}(\xi)|} \leq \sup_{\xi \in D_\varepsilon(B_r)} \frac{q|a(x) - (a)_{B_r}| |\xi|^q}{p|\xi|^p} \\ &\lesssim |a(x) - (a)_{B_r}| r^{-\frac{n(q-p)}{p(1+\varepsilon)}}, \end{aligned}$$

where $D_\varepsilon(B_r) := \{\xi \in \mathbb{R}^n : |(A^{(-1)})_{B_r}(\xi)|^{1+\varepsilon} \leq |B_r|^{-1}\} \subset \{\xi \in \mathbb{R}^n : |\xi|^{p(1+\varepsilon)} \leq |B_r|^{-1}\}$. Hence

$$\int_{B_r} \tilde{\theta}_\varepsilon(x, B_r)^\gamma dx \lesssim r^{-\frac{\gamma n(q-p)}{p(1+\varepsilon)}} \int_{B_r} |a(x) - (a)_{B_r}|^\gamma dx \lesssim \left(r^{-\frac{n(q-p)}{p(1+\varepsilon)}} \|a\|_{\text{BMO},r} \right)^\gamma.$$

Now in order for this to be useful, we need that $\|a\|_{\text{BMO},r} \lesssim r^{\frac{n(q-p)}{p(1+\varepsilon)}}$ for some $\varepsilon \geq 0$. But then a must be $\frac{n(q-p)}{p(1+\varepsilon)}$ -Hölder continuous by the Campanato-embedding, so a has point-wise continuity modulus and the mean continuity becomes irrelevant.

In the next example, we exchange the coefficients 1 and $a(x)$ of $|\xi|^p$ and $|\xi|^q$ in the double phase problem. The end result has standard q -growth, but illustrates the role of ω in Definition 3.1 which is related to small values of $|\xi|$.

Example 4.4. Let $\varphi(x, t) = a(x)t^p + t^q$, where $1 < p \leq q$ and $a : \Omega \rightarrow [0, L]$, and define

$$A(x, \xi) := D_\xi[\varphi(x, |\xi|)] = pa(x)|\xi|^{p-2}\xi + q|\xi|^{q-2}\xi.$$

Then we calculate, for $\xi \in \mathbb{R}^n$,

$$\begin{aligned} \frac{|A(x, \xi) - A_{B_r}(\xi)|}{|A_{B_r}(\xi)| + \omega(r)} &= \frac{p|a(x) - (a)_{B_r}| |\xi|^{p-1}}{p(a)_{B_r}|\xi|^{p-1} + q|\xi|^{q-1} + \omega(r)} \\ &\approx |a(x) - (a)_{B_r}| \min \left\{ \frac{1}{(a)_{B_r}}, |\xi|^{p-q}, \frac{|\xi|^{p-1}}{\omega(r)} \right\}. \end{aligned}$$

Since t^{p-q} is decreasing and $\frac{t^{p-1}}{\omega(r)}$ is increasing in t , the largest value of $\min\{t^{p-q}, \frac{t^{p-1}}{\omega(r)}\}$ occurs when $t^{q-1} = \omega(r)$ and equals $\omega(r)^{(p-q)/(q-1)}$. Thus

$$\left(\int_{B_r} \tilde{\theta}(x, B_r)^\gamma dx \right)^{1/\gamma} \lesssim \omega(r)^{\frac{p-q}{q-1}} \left(\int_{B_r} |a(x) - (a)_{B_r}|^\gamma dx \right)^{1/\gamma} \approx \omega(r)^{\frac{p-q}{q-1}} \|a\|_{\text{BMO},r}.$$

Thus $\theta_\gamma(r) + \omega(r) \lesssim \omega(r)^{\frac{p-q}{q-1}} \|a\|_{\text{BMO},r} + \omega(r) \leq \|a\|_{\text{BMO},r}^{\frac{q-1}{2q-p-1}}$, where the last estimate is obtained by the minimizing choice of $\omega(r)$. Therefore, $A^{(-1)}$ is $(\text{DMA1})_\gamma$ for any $\gamma \geq 1$ if the coefficient a is $\frac{q-1}{2q-p-1}$ -DMO.

Next, we consider nontrivial examples whose mean oscillation conditions are weaker than the corresponding point-wise ones. For $f : \Omega \rightarrow \mathbb{R}$, we say that f is log-VMO or log-DMO if

$$\lim_{r \rightarrow 0^+} \|f\|_{\text{BMO},r} \log \frac{1}{r} = 0 \quad \text{or} \quad \int_0^1 \|f\|_{\text{BMO},r} \log \left(\frac{1}{r} \right) \frac{dr}{r} < \infty,$$

respectively. If $\|f\|_{\text{BMO},r}$ is replaced by the point-wise continuity modulus ω_f , then we get the stronger vanishing log-Hölder continuity and the log-Dini continuity. By Lemma 3.4, the log-DMO condition implies the log-Hölder continuity $\omega_f(r) \lesssim (\log \frac{1}{r})^{-1}$. However, the log-VMO does not imply the log-Hölder continuity.

Example 4.5 (Borderline double phase). Let $\varphi(x, t) = t^p + a(x) \log(1+t)t^p$, where $1 < p < \infty$ and $a : \Omega \rightarrow [0, L]$, and define

$$A(x, \xi) := D_\xi[\varphi(x, |\xi|)] = \left\{ p + a(x) \left(p \log(1 + |\xi|) + \frac{|\xi|}{1 + |\xi|} \right) \right\} |\xi|^{p-2} \xi.$$

For any $\xi \in D(B_r)$, since $|\xi|^p \leq |B_r|^{-1}$,

$$\begin{aligned} \frac{|A(x, \xi) - A_{B_r}(\xi)|}{|A_{B_r}(\xi)|} &= \frac{|a(x) - (a)_{B_r}| \{p \log(1 + |\xi|) + \frac{|\xi|}{1+|\xi|}\}}{p + (a)_{B_r} \{p \log(1 + |\xi|) + \frac{|\xi|}{1+|\xi|}\}} \\ &\lesssim |a(x) - (a)_{B_r}| \log(e + |\xi|) \lesssim |a(x) - (a)_{B_r}| \log \frac{1}{r}. \end{aligned}$$

This implies, for $\gamma \geq 1$, that

$$\int_{B_r} \tilde{\theta}(r)^\gamma dx \lesssim (\|a\|_{\text{BMO}, r} \log \frac{1}{r})^\gamma.$$

Therefore, $A^{(-1)}$ satisfies **(DMA1) $_\gamma$** for any $\gamma \geq 1$ or **(VMA1)** if a is log-DMO or log-VMO, respectively.

The Dini mean oscillation condition is inspired by the coefficient case. Consequently, it is more difficult to obtain results when the variability is not in a coefficient, as the next result illustrates.

Proposition 4.6 (Variable exponent). *Let $\varphi(x, t) := \frac{1}{p(x)} t^{p(x)}$, where $p : \Omega \rightarrow [p^-, p^+]$ with $1 < p^- \leq p^+$, and define*

$$A(x, \xi) := D_\xi[\varphi(x, |\xi|)] = |\xi|^{p(x)-2} \xi.$$

If p is log-VMO, then $A^{(-1)}$ satisfies Definition 3.1 for any $\gamma \geq 1$ with $\theta(r) = \|p\|_{\text{BMO}, r} \log \frac{1}{r}$ and $\omega(r) = r^{p^- - 1}$.

Proof. Let $r \in (0, 1]$. Note that $D(B_r) = \{\xi \in \mathbb{R}^n : |\xi|^{p(x)} \leq |B_r|^{-1}\} \subset \{\xi \in \mathbb{R}^n : |\xi| \in [0, r^{-n}]\}$. If $|\xi| \in [0, r^2]$,

$$\frac{|A^{(-1)}(x, \xi) - A_{B_r}^{(-1)}(\xi)|}{|A_{B_r}^{(-1)}(\xi)| + \omega(r) |\xi|} = \frac{|A(x, \xi) - A_{B_r}(\xi)|}{|A_{B_r}(\xi)| + r^{p^- - 1}} \leq 2r^{p^- - 1}.$$

If $|\xi| = 1$, $|A(x, \xi) - A_{B_r}(\xi)| = 0$ and our estimate will be trivial. Suppose $|\xi| \in [r^2, r^{-n}] \setminus \{1\}$ and define

$$\bar{p} := \log_{|\xi|} \int_{B_r} |\xi|^{p(y)} dy \in [p_{B_r}^-, p_{B_r}^+].$$

From $|\xi|^{p(x)-p(y)} \leq \max\{r^{-n|p(x)-p(y)|}, r^{-2|p(x)-p(y)|}\} = r^{-n|p(x)-p(y)|}$ it follows that

$$|\xi|^{p(x)-\bar{p}} = \int_{B_r} |\xi|^{p(x)-p(y)} dy \leq \int_{B_r} r^{-n|p(x)-p(y)|} dy = \int_{B_r} \exp(n|p(x) - p(y)| \log \frac{1}{r}) dy.$$

An analogous argument gives a lower bound with integrand $\exp(-n|p(x) - p(y)| \log \frac{1}{r})$. Furthermore, $1 - e^{-s} \leq e^s - 1$ when $s \geq 0$. Thus we obtain that

$$\begin{aligned} \frac{|A(x, \xi) - A_{B_r}(\xi)|}{|A_{B_r}(\xi)| + r^{p^- - 1}} &\leq \frac{||\xi|^{p(x)-1} - |\xi|^{\bar{p}-1}|}{|\xi|^{\bar{p}-1}} = ||\xi|^{p(x)-\bar{p}} - 1| \\ &\leq \int_{B_r} [\exp(n|p(x) - p(y)| \log \frac{1}{r}) - 1] dy. \end{aligned}$$

Combining the above results and Hölder's inequality, we have shown that

$$\int_{B_r} \tilde{\theta}(x, B_r)^\gamma dx \lesssim \int_{B_r} \int_{B_r} [\exp(n|p(x) - p(y)| \log \frac{1}{r}) - 1]^\gamma dy dx + r^{\gamma(p^- - 1)}.$$

By differentiation it follows that $e^{ab} - 1 \leq ae^b$ when $b > 0$ and $a \in [0, 1 - \frac{\ln b}{b}]$. In particular, this holds for all $a \in [0, 1 - \frac{1}{e}]$. It follows that

$$\begin{aligned} & \int_{B_r} \int_{B_r} \left[\exp(n|p(x) - p(y)| \log \frac{1}{r}) - 1 \right]^\gamma dy dx \\ & \leq (c_n n \gamma \|p\|_{\text{BMO}_r} \log \frac{1}{r})^\gamma \int_{B_r} \int_{B_r} \exp\left(\frac{|p(x) - p(y)|}{c_n \|p\|_{\text{BMO}_r}}\right) dy dx \\ & \lesssim (c_n n \gamma \|p\|_{\text{BMO}_r} \log \frac{1}{r})^\gamma \int_{B_r} \exp\left(\frac{|p(x) - p_{B_r}|}{c_n \|p\|_{\text{BMO}_r}}\right) dx, \end{aligned}$$

where $c_n > 1$ is the constant from the John–Nirenberg lemma depending on n and $r > 0$ is so small that $c_n n \gamma \log \frac{1}{r} \|p\|_{\text{BMO}_r} \leq 1 - \frac{1}{e}$ by the log-VMO assumption. The John–Nirenberg lemma ensures that the integral on the right-hand side is finite. Thus we have shown that

$$\left(\int_{B_r} \tilde{\theta}(x, B_r)^\gamma dx \right)^{\frac{1}{\gamma}} \lesssim \log \frac{1}{r} \|p\|_{\text{BMO}_r} + r^{p^- - 1}$$

for all sufficiently small r . For $r \geq r_1$, the estimate still holds since $\tilde{\theta}$ is bounded by the constant $cr_1^{p^- - p^+}$. \square

The definition of log-DMO and log-VMO combined with the previous result gives the following example.

Example 4.7 (Variable exponent). In the setting of Proposition 4.6, $A^{(-1)}$ satisfies $(\text{DMA1})_\gamma$ for any $\gamma \geq 1$ or (VMA1) if p is log-DMO or log-VMO, respectively.

Finally, we show that also the Lorentz-type restriction is a special case of our condition. Interestingly, this proof does not hold if we include the supremum with respect to the radius in the definition of θ_γ , i.e. use θ_γ^* from Proposition 3.8. This shows the importance of having the optimization in our condition. We define the *non-increasing rearrangement* f_E^* by

$$f_E^*(t) := \inf\{s \geq 0 \mid \mu_E(s) \leq t\} \quad \text{where} \quad \mu_E(s) := |\{x \in E \mid |f(x)| > s\}|.$$

Note that f_E^* is a kind of inverse of the distribution function μ_E . The definition directly implies that $(|f|^\gamma)_\Omega^* = (f_\Omega^*)^\gamma$. The *Lorentz space* $L^{n,1}(\Omega)$ is defined by the condition $\int_0^\infty t^{1/n} f_\Omega^*(t) \frac{dt}{t} < \infty$.

Proposition 4.8 (Orlicz with Lorentz coefficient). *Let $A(x, \xi) = a(x) \frac{\psi'(|\xi|)}{|\xi|} \xi$, where $a : \Omega \rightarrow [L^{-1}, L]$ with $L \geq 1$. If $a \in W^{1,1}(\Omega)$ with $|Da| \in L^{n,1}(\Omega)$, then $A^{(-1)}$ satisfies $(\text{DMA1})_\gamma$ for any $\gamma \geq 1$.*

Proof. We follow the argument in [43, Section 2.3]. We consider $\gamma > n'$; the case $\gamma \leq n'$ follows from this by Hölder's inequality. For $B_{2r} \subset \Omega$ and $\tilde{\gamma} := \frac{\gamma n}{n+\gamma} \in (1, n)$, by the Sobolev–Poincaré inequality,

$$\begin{aligned} \left(\int_{B_r} |a - (a)_{B_r}|^\gamma dx \right)^{\frac{1}{\gamma}} & \leq cr \left(\int_{B_r} |Da|^{\tilde{\gamma}} dx \right)^{\frac{1}{\tilde{\gamma}}} \leq cr \left(\int_0^{|B_r|} (|Da|^{\tilde{\gamma}})_{B_r}^*(t) dt \right)^{\frac{1}{\tilde{\gamma}}} \\ & \leq cr \left(\int_0^{|B_r|} (|Da|^{\tilde{\gamma}})_\Omega^*(t) dt \right)^{\frac{1}{\tilde{\gamma}}} \end{aligned}$$

where we used that $f_{U_2}^* \leq f_{U_1}^*$ if $U_2 \subset U_1$. Note that the right hand side is independent of the center of B_r . Taking supremum over $B_r \subset \Omega$ and changing variables $\tau = |B_r| = |B_1|r^n$, we find that

$$\int_0^1 \theta_\gamma(r) \frac{dr}{r} \leq c \int_0^1 \left(\int_0^{|B_r|} (|Da|^{\tilde{\gamma}})_\Omega^*(t) dt \right)^{\frac{1}{\tilde{\gamma}}} dr \leq c \int_0^\infty \tau^{\frac{1}{n}-1} \left(\int_0^\tau (|Da|^{\tilde{\gamma}})_\Omega^*(t) dt \right)^{\frac{1}{\tilde{\gamma}}} d\tau.$$

We recall a Hardy-type inequality for the quasinorm case $\beta < 1$ (see [56, Theorem 3] with $p = q$ and $w(\tau) = v(\tau) = \tau^{\alpha-1}$):

$$\int_0^\infty \tau^{\alpha-1} \left(\int_0^\tau f(t) dt \right)^\beta d\tau \leq c(\alpha, \beta) \int_0^\infty t^{\alpha-1} f(t)^\beta dt, \quad 0 < \alpha < \beta < 1,$$

where f is a non-negative, non-increasing function. Using this inequality with $\beta = \frac{1}{\gamma} \in (\frac{1}{n}, 1)$ and $\alpha = \frac{1}{n}$, we obtain

$$\int_0^1 \theta_\gamma(r) \frac{dr}{r} \lesssim \int_0^\infty t^{\frac{1}{n}-1} (|Da|^{\tilde{\gamma}})_\Omega^*(t)^{\frac{1}{\tilde{\gamma}}} dt = \int_0^\infty t^{\frac{1}{n}} (|Da|)_\Omega^*(t) \frac{dt}{t} < \infty,$$

by $(|f|^\gamma)_\Omega^* = (f_\Omega^*)^\gamma$ and the definition of the Lorentz space. \square

We can combine the previous two propositions to cover the variable exponent with coefficient and regularity given by Lorentz spaces as considered by Baroni [6].

Example 4.9 (Variable exponent with Lorentz conditions). Let $A(x, \xi) = \frac{a(x)}{p(x)} |\xi|^{p(x)}$, where $a : \Omega \rightarrow [L^{-1}, L]$ with $L \geq 1$. If $a, p \in W^{1,1}(\Omega)$ with $|Da|, |Dp| \in L^{n,1}(\Omega)$ and $\|Dp\|_{L^{n,1}(B_r)} \log \frac{1}{r} \leq c$, then we can show that $A^{(-1)}$ satisfies **(DMA1) $_\gamma$** for any $\gamma \geq 1$.

To reach this conclusion, we estimate $|A_{B_r}(\xi)| \geq L^{-1} |\xi|^{\bar{p}-1}$ and

$$|a(x) |\xi|^{p(x)-1} - a(y) |\xi|^{p(y)-1}| \leq L \left| |\xi|^{p(x)-1} - |\xi|^{p(y)-1} \right| + |a(x) - a(y)| |\xi|^{p(y)-1}.$$

Since $L^{n,1}(\Omega) \hookrightarrow L_{\text{loc}}^\infty(\Omega)$, p is log-Hölder continuous and so $|\xi|^{p(y)-\bar{p}} \leq c$ [6, (1.10)]. Then we handle $|a(x) - a(y)|$ in the second term as in Proposition 4.8 using $|Da| \in L^{n,1}(\Omega)$. For the first term we arrive as in Proposition 4.6 at an estimate

$$\log \frac{1}{r} \int_{B_r} |p(x) - (p)_{B_r}| dx,$$

from which the integral is estimated as in Proposition 4.8. The details are left to the interested reader.

5. COMPARISON ESTIMATES

In this section, we always assume that $A : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfies the quasi-isotropic (p, q) -growth condition in Definition 1.1.

5.1. Growth functions and approximating energies. We start with recalling the properties of a so-called growth function φ of A .

Proposition 5.1 (Proposition 3.3, [38]). *There exists $\varphi \in \Phi_c(\Omega)$ with $\varphi'(x, \cdot) \in C([0, \infty))$ such that φ' is **(A0)**, **(Inc) $_{p-1}$** and **(Dec) $_{q_1-1}$** for some $q_1 \geq q$ and that*

$$(5.2) \quad \bar{L}^{-1} (|A(x, \xi)| + |\xi| |D_\xi A(x, \xi)|) \leq \varphi'(x, |\xi|) \leq \bar{L} |\xi| |D_\xi A(x, \xi)| e \cdot e$$

for every $x \in \Omega$ and $\xi, e \in \mathbb{R}^n$ with $\xi \neq 0$ and $|e| = 1$. Here, q_1 and $\bar{L} \geq 1$ depend on n, p, q and L . We call this φ the growth function of A .

By the monotonicity of the (aDec)-condition, we can replace our original q by q_1 from the proposition above and assume without loss of generality that $q_1 = q$.

From growth and ellipticity inequalities of growth functions in Proposition 5.1, a standard calculation yields the following monotonicity and coercivity/growth properties:

$$(A(x, \xi_1) - A(x, \xi_2)) \cdot (\xi_1 - \xi_2) \gtrsim \frac{\varphi'(|\xi_1| + |\xi_2|)}{|\xi_1| + |\xi_2|} |\xi_1 - \xi_2|^2$$

and

$$c_*^{-1} \varphi(x, |\xi|) \leq \xi \cdot A(x, \xi) \leq |\xi| |A(x, \xi)| \leq c_* \varphi(x, |\xi|)$$

for some $c_* = c_*(n, p, q, L) \geq 1$. In particular, this implies that the function space associated with the local weak solutions to (divA) is the Sobolev space $W_{\text{loc}}^{1, \varphi}(\Omega)$, and that φ satisfies (A1) or (SA1) if and only if $A^{(-1)}$ does.

We state the higher integrability result for the weak solutions to (divA; F) which requires the (A1) condition of $A^{(-1)}$. The homogeneous case, when $F \equiv 0$, can be found in [39, Theorem 4.1]; see also [37, Lemma 4.7] and references therein. Essentially the same argument can be applied to the nonhomogeneous case, so it is not repeated here.

Lemma 5.3. *Suppose that $A^{(-1)}$ satisfies (A1). Let $u \in W_{\text{loc}}^{1, \varphi}(\Omega)$ be a weak solution to (divA; F) with $\varphi(\cdot, |F|) \in L_{\text{loc}}^s(\Omega)$ for some $s > 1$. There exists $\sigma = \sigma(n, p, q, L, s) > 0$ such that $\varphi(\cdot, |Du|) \in L_{\text{loc}}^{1+\sigma}(\Omega)$ and*

$$\int_{B_r} \varphi(x, |Du|)^{1+\sigma} dx \leq c \left\{ \varphi_{B_{2r}}^- \left(\int_{B_{2r}} |Du| dx \right)^{1+\sigma} + \int_{B_{2r}} \varphi(x, |F|)^{1+\sigma} dx + 1 \right\}$$

for some $c = c(n, p, q, L, s) \geq 1$, whenever $B_{2r} \Subset \Omega$ and $\varrho_{L\varphi(B_{2r})}(|Du|) \leq 1$.

We establish an approximating autonomous problem for (divA). Denote the averages of $A(\cdot, \xi)$, $\varphi(\cdot, t)$ and $\varphi'(\cdot, t)$ over $B_r \subset \Omega$ by A_{B_r} , φ_{B_r} and φ'_{B_r} , and set

$$(5.4) \quad t_0 := 2(\varphi_{B_r})^{-1}(c_* |B_r|^{-1}),$$

where $c_* \geq 1$ is as above, and

$$\psi'(t) := \begin{cases} \varphi'_{B_r}(t) & \text{if } t \leq t_0, \\ \varphi'_{B_r}(t_0) \left(\frac{t}{t_0}\right)^{p-1} & \text{if } t_0 < t. \end{cases}$$

Note that φ'_{B_r} is the same as the derivative of φ_{B_r} and ψ' is continuous since φ' satisfies (Inc) $_{p-1}$ and (Dec) $_{q-1}$. We consider

$$\psi(t) := \int_0^t \psi'(s) ds.$$

We see that $\psi = \varphi_{B_r}$ in $[0, t_0]$ and, since φ'_{B_r} satisfies (Inc) $_{p-1}$, $\psi \leq \varphi_{B_r}$ in $[t_0, \infty)$. Fix $\eta \in C_0^\infty(\mathbb{R})$ with $\eta \geq 0$, $\text{supp } \eta \subset (0, 1)$ and $\|\eta\|_1 = 1$. We define

$$\tilde{\varphi}(t) := \int_0^\infty \psi(s) \eta_{rt}(s-t) ds \quad \text{where} \quad \eta_r(t) := \frac{1}{r} \eta\left(\frac{t}{r}\right).$$

The construction of $\tilde{\varphi}$ is analogous to the one in [37, Section 5] with $t_1 = 0$ and $t_2 = t_0$, except that here we use the average φ_{B_r} instead of the function at the center-point, $\varphi(x_0, \cdot)$. Therefore, we have the following analogue of Proposition 5.10 in [37].

Proposition 5.5. *Suppose that $A^{(-1)}$ satisfies (A1). The following hold with $c \geq 1$ depending only on n, p, q and L :*

- (1) $\psi(t) \leq \tilde{\varphi}(t) \leq (1 + cr)\psi(t) \leq c\psi(t)$ for all $t > 0$.

- (2) $\tilde{\varphi} \in C^1([0, \infty)) \cap C^2((0, \infty))$ satisfies (A0), (Inc) $_p$ and (Dec) $_q$ while $\tilde{\varphi}'$ satisfies (A0), (Inc) $_{p-1}$ and (Dec) $_{q-1}$.
(3) $\tilde{\varphi}(t) \leq c(\varphi(x, t) + 1)$ for all $(x, t) \in B_r \times [0, \infty)$.
(4) Let

$$\tilde{\zeta}(x, t) := \begin{cases} \varphi(x, \tilde{\varphi}^{-1}(1))t, & 0 \leq t < 1, \\ \varphi(x, \tilde{\varphi}^{-1}(t)), & t \geq 1, \end{cases}$$

and $\zeta(x, t) := \tilde{\zeta}(x, t)^{1+\tilde{\sigma}}$ for $\tilde{\sigma} > 0$. Then $\zeta \in \Phi_w(B_r)$ satisfies (A0), (aInc) $_{1+\tilde{\sigma}}$ and (aDec) $_{q(1+\tilde{\sigma})/p}$ with relevant constants depending on n, p, q, L and $\tilde{\sigma}$.

Proof. The proofs of (1) – (3) can be obtained from [37, Proposition 5.10] when we replace $\varphi'(x_0, t)$ by $\varphi'_{B_r}(t)$.

Let us prove (4) by following the proof of [37, Proposition 5.12]. Note that $\zeta \in \Phi_w(B_r)$ is clear once we show (aInc) $_1$. As φ and $\tilde{\varphi}$ satisfy (A0), so does $\tilde{\zeta}$ and thus $\tilde{\zeta}(x, t) \approx t$ for $t \in [0, 1)$. Now we prove that $\tilde{\zeta}$ satisfies (aInc) $_1$ and (aDec) $_{q/p}$, hence ζ satisfies (aInc) $_{1+\tilde{\sigma}}$ and (aDec) $_{q(1+\tilde{\sigma})/p}$. For $t \in [1, \tilde{\varphi}(t_0)]$, since $\varphi_{B_r}^-(\tilde{\varphi}^{-1}(t)) \lesssim \varphi_{B_r}(t_0) \lesssim |B_r|^{-1}$, the (A1) condition of φ yields $\varphi(x, \tilde{\varphi}^{-1}(t)) \approx \varphi_{B_r}(\tilde{\varphi}^{-1}(t)) = \psi(\tilde{\varphi}^{-1}(t)) \approx t$. Therefore $\tilde{\zeta}(x, t) \approx t$ in $[0, \tilde{\varphi}(t_0)]$. For $t \in [\tilde{\varphi}(t_0), \infty)$, setting $s := \tilde{\varphi}^{-1}(t)$,

$$\frac{\tilde{\zeta}(x, t)}{t} = \frac{\varphi(x, \tilde{\varphi}^{-1}(t))}{t} \approx \frac{\varphi(x, s)}{\psi(s)} = \frac{t_0^{p-1}}{\varphi'_{B_r}(t_0)} \frac{\varphi(x, s)}{s^p}$$

and, similarly,

$$\frac{\tilde{\zeta}(x, t)}{t^{q/p}} \approx \left(\frac{t_0^{p-1}}{\varphi'_{B_r}(t_0)} \right)^{q/p} \frac{\varphi(x, s)}{s^q}.$$

Therefore, (aInc) $_p$ and (aDec) $_q$ of φ imply (aInc) $_1$ and (aDec) $_{q/p}$ of $\tilde{\zeta}$. Finally, we show that ζ satisfies (A1). Let $B_\rho \subset B_r$, and assume that $\zeta_{B_\rho}^-(t) \leq |B_\rho|^{-1}$. Then

$$\tilde{\zeta}_{B_\rho}^-(t) = \varphi_{B_\rho}^-(\tilde{\varphi}^{-1}(t)) \leq |B_\rho|^{-1/(1+\tilde{\sigma})} \leq |B_\rho|^{-1}.$$

Therefore, (A1) of φ implies that

$$\zeta_{B_\rho}^+(t) = [\varphi_{B_\rho}^+(\tilde{\varphi}^{-1}(t))]^{1+\tilde{\sigma}} \lesssim [\varphi_{B_\rho}^-(\tilde{\varphi}^{-1}(t))]^{1+\tilde{\sigma}} = \zeta_{B_\rho}^-(t)$$

and so ζ satisfies (A1). \square

From the nonlinearity A , we define an autonomous function $\tilde{A} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ as

$$\tilde{A}(\xi) := \eta_1(|\xi|)A_{B_r}(\xi) + \eta_2(|\xi|) \frac{\varphi'_{B_r}(t_0)}{t_0^{p-1}} |\xi|^{p-2} \xi,$$

where $\eta_1 \in C^\infty([0, \infty))$ is such that $0 \leq \eta \leq 1$, $\eta \equiv 1$ on $[0, t_0]$, $\eta \equiv 0$ on $[2t_0, \infty)$, $-2/t_0 \leq \eta'_1 \leq 0$, and $\eta_2 \in C^\infty([0, \infty))$ is such that $0 \leq \eta_2 \leq 1$, $\eta_2 \equiv 0$ on $[0, t_0/2]$, $\eta_2 \equiv 1$ on $[t_0, \infty)$ and $0 \leq \eta'_2 \leq 4/t_0$. Then by the same computations as in [39, Lemma 5.2] with $t_1 = 0$, $t_2 = t_0$ and replacing η_2 and η_3 by the functions η_1 and η_2 defined above, we can show that $\tilde{\varphi}$ defined above is a growth function of \tilde{A} , i.e.

$$(5.6) \quad \tilde{L}^{-1}(|\tilde{A}(\xi)| + |\xi| |D\tilde{A}(\xi)|) \leq \tilde{\varphi}'(|\xi|) \leq \tilde{L} |\xi| |D\tilde{A}(\xi)| e \cdot e$$

for some $\tilde{L} \geq 1$ and every $\xi, e \in \mathbb{R}^n$ with $\xi \neq 0$ and $|e| = 1$.

5.2. Approximation and comparison estimates. Let $u \in W_{\text{loc}}^{1,\varphi}(\Omega)$ be a local weak solution to $(\text{div}A)$. Fix $\Omega' \Subset \Omega$. Then, by Lemma 5.3, $\varphi(\cdot, |Du|) \in L^{1+\sigma}(\Omega')$. Assume that r satisfies

$$(5.7) \quad r \in (0, \frac{1}{2}] \quad \text{and} \quad |B_{2r}| \leq 2^{-\frac{2(1+\sigma)}{\sigma}} \left(\int_{\Omega'} \varphi(x, |Du|)^{1+\sigma} dx + 1 \right)^{-\frac{2+\sigma}{\sigma}} \leq \frac{1}{2}.$$

By Hölder's inequality this gives

$$\int_{B_{2r}} \varphi(x, |Du|)^{1+\frac{\sigma}{2}} dx \leq |B_{2r}|^{\frac{\sigma}{2(1+\sigma)}} \left(\int_{\Omega'} \varphi(x, |Du|)^{1+\sigma} dx \right)^{\frac{2+\sigma}{2(1+\sigma)}} \leq \frac{1}{2}$$

so that

$$\varrho_{L^\varphi(B_{2r})}(|Du|) \leq \int_{B_{2r}} \varphi(x, |Du|)^{1+\frac{\sigma}{2}} dx + |B_r| \leq 1.$$

We fix $B_{2r} \Subset \Omega'$, and let $\tilde{u} \in u + W_0^{1,\tilde{\varphi}}(B_r)$ be the unique weak solution to

$$(\text{div}\tilde{A}) \quad \text{div}\tilde{A}(D\tilde{u}) = 0$$

in $B_r \subset \Omega$. The solutions u and \tilde{u} of $(\text{div}A)$ and $(\text{div}\tilde{A})$ satisfy the following estimates.

Lemma 5.8. *Suppose that $A^{(-1)}$ satisfies (A1) and r satisfies (5.7) and let $\sigma > 0$ be from Lemma 5.3. There exists $C_0 \geq 1$ depending only on n, p, q and L such that*

- (1) $\int_{B_r} \tilde{\varphi}(|D\tilde{u}|) dx \leq C_0 \int_{B_r} \tilde{\varphi}(|Du|) dx,$
- (2) $\left(\int_{B_r} \varphi(x, |Du|)^{1+\sigma} dx \right)^{\frac{1}{1+\sigma}} \leq C_0 \tilde{\varphi} \left(\int_{B_{2r}} |Du| dx \right) + C_0,$
- (3) $\int_{B_r} |D\tilde{u}| dx \leq C_0 \int_{B_{2r}} |Du| dx + C_0,$
- (4) if $\varrho_{L^{\varphi^{1+\tilde{\sigma}}}(B_r)}(|Du|) \leq M_0 < \infty$ for $\tilde{\sigma} > 0$, then

$$\int_{B_r} \varphi(x, |D\tilde{u}|)^{1+\tilde{\sigma}} dx \leq C_0 (M_0^{\frac{q}{p}-1} + 1) \left(\int_{B_{2r}} \varphi(x, |Du|)^{1+\tilde{\sigma}} dx + 1 \right),$$

where the constant C_0 depends also on $\tilde{\sigma}$.

Proof. Testing $(\text{div}\tilde{A})$ by $\tilde{u} - u \in W_0^{1,\tilde{\varphi}}(B_{2r})$ and using that $\tilde{\varphi} \approx |\tilde{A}^{(-1)}|$ by (5.6) and Young's inequality, we obtain (1). Note that this part does not require (A1).

Let $t_1 := \int_{B_{2r}} |Du| dx$. Since $\varrho_{L^\varphi(B_{2r})}(|Du|) \leq 1$ by the choice of r , we conclude from Jensen's inequality and (A1) of φ that

$$t_1 \lesssim (\varphi_{B_{2r}}^-)^{-1} \left(\int_{B_{2r}} \varphi_{B_{2r}}^- (|Du|) dx \right) \lesssim \varphi_{B_r}^- (|B_{2r}|^{-1}) \lesssim t_0.$$

Hence, by the definition of ψ and Proposition 5.5(1), $\varphi_{B_{2r}}^-(t_1) \leq \varphi_{B_r}(t_1) \approx \psi(t_1) \approx \tilde{\varphi}(t_1)$. Then, (2) follows from Lemma 5.3, when $F \equiv 0$, and the fact that $\varphi_{B_{2r}}^-(t_1) \lesssim \tilde{\varphi}(t_1)$. Moreover, using the two estimates we have shown and Proposition 5.5(3), we also obtain

$$\tilde{\varphi} \left(\int_{B_r} |D\tilde{u}| dx \right) \lesssim \int_{B_r} \tilde{\varphi}(|Du|) dx \lesssim \int_{B_r} \varphi(x, |Du|) dx + 1 \lesssim \tilde{\varphi} \left(\int_{B_{2r}} |Du| dx + 1 \right)$$

which implies (3). Finally, (4) follows the Calderón–Zygmund estimate in [38, Lemma 4.15] with φ and θ replaced by $\tilde{\varphi}$ and ζ given in Proposition 5.5(4):

$$\begin{aligned} \int_{B_r} \varphi(x, |D\tilde{u}|)^{1+\tilde{\sigma}} dx &\lesssim \int_{B_r} \tilde{\zeta}(x, \tilde{\varphi}(|D\tilde{u}|))^{1+\tilde{\sigma}} dx + 1 \\ &\lesssim (M_0^{\frac{q}{p}-1} + 1) \left(\int_{B_r} \tilde{\zeta}(x, \tilde{\varphi}(|Du|))^{1+\tilde{\sigma}} dx + 1 \right) \\ &\lesssim (M_0^{\frac{q}{p}-1} + 1) \left(\int_{B_r} \varphi(x, |Du|)^{1+\tilde{\sigma}} dx + 1 \right). \quad \square \end{aligned}$$

The next result is a supremum estimate and an excess decay estimate for the derivative $D\tilde{u}$ of the solution to $(\operatorname{div}\tilde{A})$ in L^1 -space, which can essentially be found in [47] (see also [48]). Although the paper [47] proved the excess decay estimate in $L^{\tilde{\varphi}}$ -space, a minor modification yields the desired estimate in L^1 -space. More details of the proof of the lemma are provided in Appendix A.

Lemma 5.9. *Let $\tilde{A} : \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfy the quasi-isotropic (p, q) -growth condition and $\tilde{\varphi} \in C^1([0, \infty)) \cap C^2((0, \infty))$ be its growth function satisfying (5.6). Let $\tilde{u} \in W^{1, \tilde{\varphi}}(B_r)$ be a weak solution to $(\operatorname{div}\tilde{A})$. There exist $\alpha \in (0, 1)$ and $C_1 \geq 1$ depending on n, p, q and \tilde{L} such that, for any $B_\nu(y) \subset B_\rho(y) \subset B_r$,*

$$\|D\tilde{u}\|_{L^\infty(B_{\rho/2}(y))} \leq C_1 \int_{B_\rho(y)} |D\tilde{u}| dx,$$

and

$$\int_{B_\nu(y)} |D\tilde{u} - (D\tilde{u})_{B_{\tilde{\rho}}(y)}| dx \leq \operatorname{osc}_{B_\nu(y)} D\tilde{u} \leq C_1 \left(\frac{\nu}{\rho}\right)^\alpha \int_{B_\rho(y)} |D\tilde{u} - (D\tilde{u})_{B_\rho(y)}| dx.$$

Now we derive comparison estimates between the gradients of u and \tilde{u} , in terms of the mean oscillation conditions in Definition 3.1. The following first one corresponds to Lemma 3.1 in [6]. Note that the resulting estimates are almost the same. However, the proof of next lemma requires more delicate analysis since we consider general structure. A similar inequality was proved in [37, Lemma 6.2], but with the much worse power $\omega^{p/q}$, even though there ω was a point-wise modulus of continuity, not mean continuity like here.

Lemma 5.10. *In the setting of Lemma 5.8 with $\varrho_{L^{\varphi^{\gamma/(\gamma-2)}}(B_r)}(|Du|) \leq 1$ for some $\gamma > 2$,*

$$\int_{B_r} \frac{\tilde{\varphi}'(|Du| + |D\tilde{u}|)}{|Du| + |D\tilde{u}|} |Du - D\tilde{u}|^2 dx \leq c \Theta(r)^2 \left(\int_{B_r} \varphi(x, |Du|)^{\frac{\gamma}{\gamma-2}} dx + 1 \right)^{\frac{\gamma-2}{\gamma}}$$

for some $c = c(n, p, q, L, \gamma, C_\gamma^*) \geq 1$, where $\Theta(r) := \omega(r) + \theta_\gamma(r) + r^{\alpha_1}$ with $\alpha_1 = \frac{4n}{\gamma(\gamma-2)}$.

Proof. Denote $\Psi := \frac{\tilde{\varphi}'(|Du| + |D\tilde{u}|)}{|Du| + |D\tilde{u}|} |Du - D\tilde{u}|^2$. By the monotonicity of \tilde{A} and testing with $u - \tilde{u} \in W_0^{1, \varphi}(B_r) \subset W_0^{1, \tilde{\varphi}}(B_r)$ in the weak formulations of $(\operatorname{div}A)$ and $(\operatorname{div}\tilde{A})$, we have

$$\begin{aligned} \int_{B_r} \Psi dx &\leq c_0 \int_{B_r} (\tilde{A}(Du) - \tilde{A}(D\tilde{u})) \cdot (Du - D\tilde{u}) dx \\ &= c_0 \int_{B_r} (\tilde{A}(Du) - A(x, Du)) \cdot (Du - D\tilde{u}) dx \\ &\leq c_0 \int_{B_r} \underbrace{|A(x, Du) - \tilde{A}(Du)|}_{=:\Delta} |Du - D\tilde{u}| dx. \end{aligned}$$

In the set $E := \{x \in B_r : |A_{B_r}^{(-1)}(Du(x))| \leq |B_r|^{-1}\}$, since $\varphi_{B_r}(|Du|) \leq c_* |A_{B_r}^{(-1)}(Du)| \leq c_* |B_r|^{-1}$, it follows from (5.4) that $|Du| \leq \frac{t_0}{2}$ and so $\tilde{A}(Du) = A_{B_r}(Du)$. Hence by the definition of $\tilde{\theta}$ (Definition 3.1 with $G = A^{(-1)}$) and the fact $\varphi_{B_r}(|Du|) \approx \psi(|Du|) \leq \tilde{\varphi}(|Du|)$ from Proposition 5.5(1), we conclude that

$$\begin{aligned} |\tilde{A}(Du) - A(x, Du)| &\leq \tilde{\theta}(x, B_r)(|A_{B_r}(Du)| + \omega(r)) \\ &\lesssim \tilde{\theta}(x, B_r) \tilde{\varphi}(|Du| + |D\tilde{u}|)^{\frac{1}{2}} \left[\frac{\tilde{\varphi}'(|Du| + |D\tilde{u}|)}{|Du| + |D\tilde{u}|} \right]^{\frac{1}{2}} + \tilde{\theta}(x, B_r) \omega(r). \end{aligned}$$

It then follows from the Cauchy–Schwartz inequality and $\tilde{\theta}(x, B_r) \omega(r) \leq \tilde{\theta}(x, B_r)^2 + \omega(r)^2$ that

$$\Delta \chi_E \leq c \tilde{\theta}(x, B_r)^2 \tilde{\varphi}(|Du| + |D\tilde{u}|) + \frac{1}{2c_0} \Psi + (\tilde{\theta}(x, B_r)^2 + \omega(r)^2) |Du - D\tilde{u}|.$$

Integrating the previous inequality over B_r , we obtain that

$$\int_{B_r} \Delta \chi_E dx \leq \int_{B_r} [\tilde{\theta}(x, B_r)^2 + \omega(r)^2] \tilde{\varphi}(|Du| + |D\tilde{u}| + 1) dx + \frac{1}{2c_0} \int_{B_r} \Psi dx.$$

The first integral is estimated using Hölder's inequality, the definition of θ_γ , Proposition 5.5(3) and Lemma 5.8(4) with $\varrho_{L^{\varphi^\gamma/(\gamma-2)}(B_r)}(|Du|) \leq 1$:

$$\begin{aligned} &\int_{B_r} \left[\tilde{\theta}(x, B_r)^2 + \omega(r)^2 \right] \tilde{\varphi}(|Du| + |D\tilde{u}| + 1) dx \\ &\leq \left\{ \left(\int_{B_r} \tilde{\theta}(x, B_r)^\gamma dx \right)^{\frac{2}{\gamma}} + \omega(r)^2 \right\} \left(\int_{B_r} \tilde{\varphi}(|Du| + |D\tilde{u}| + 1)^{\frac{\gamma}{\gamma-2}} dx \right)^{\frac{\gamma-2}{\gamma}} \\ &\leq c (\theta_\gamma(r)^2 + \omega(r)^2) \left(\int_{B_r} \varphi(x, |Du|)^{\frac{\gamma}{\gamma-2}} dx + 1 \right)^{\frac{\gamma-2}{\gamma}}. \end{aligned}$$

Hence we obtain that

$$\int_{B_r} \Delta \chi_E dx \leq c (\theta_\gamma(r)^2 + \omega(r)^2) \left(\int_{B_r} \varphi(x, |Du|)^{\frac{\gamma}{\gamma-2}} dx + 1 \right)^{\frac{\gamma-2}{\gamma}} + \frac{1}{2c_0} \int_{B_r} \Psi dx.$$

In the set $B_r \setminus E$, $\varphi_{B_r}(|Du|) \gtrsim |B_r|^{-1}$. This implies $\varphi_{B_r}^-(|Du|) \gtrsim |B_r|^{-1}$ since if $\varphi_{B_r}^-(|Du|) \leq |B_r|^{-1}$, by (A1) of φ , $|B_r|^{-1} \lesssim \varphi_{B_r}(|Du|) \lesssim \varphi_{B_r}^-(|Du|) + 1 \lesssim \varphi_{B_r}^-(|Du|)$. In the last estimate we used the fact that $1 \lesssim |Du| \lesssim \varphi_{B_r}^-(|Du|)$ in $B_r \setminus E$. Furthermore, by Proposition 5.5(1)&(3) and (A0),

$$\tilde{\varphi}'(|Du|) \approx \frac{\tilde{\varphi}(|Du|)}{|Du|} \lesssim \frac{\varphi(x, |Du|) + 1}{|Du|} \approx \varphi'(x, |Du|).$$

Using the growth conditions (5.2) and (5.6), the above inequalities and Young's inequality twice, we conclude that

$$\begin{aligned} \Delta \chi_{B_r \setminus E} &\lesssim [|B_r| \varphi^-(|Du|)]^{\tilde{\sigma}} \varphi'(x, |Du|) (|Du| + |D\tilde{u}|) \\ &\lesssim r^{n\tilde{\sigma}} (\varphi(x, |Du|)^{1+\tilde{\sigma}} + \varphi(x, |Du|)^{\tilde{\sigma}} \varphi(x, |D\tilde{u}|)) \\ &\lesssim r^{n\tilde{\sigma}} (\varphi(x, |Du|)^{1+\tilde{\sigma}} + \varphi(x, |D\tilde{u}|)^{1+\tilde{\sigma}}), \end{aligned}$$

where $\tilde{\sigma} := \frac{2}{\gamma-2}$. From Lemma 5.8(4) with $\varrho_{L\varphi^{\gamma/(\gamma-2)}(B_r)}(|Du|) \leq 1$ it follows that

$$\begin{aligned} \int_{B_r} \Delta \chi_{B_r \setminus E} dx &\lesssim r^{n\tilde{\sigma}} \left(\int_{B_r} [\varphi(x, |Du|)^{1+\tilde{\sigma}} + \varphi(x, |D\tilde{u}|)^{1+\tilde{\sigma}}] dx \right)^{\frac{1}{1+\tilde{\sigma}} + \frac{\tilde{\sigma}}{1+\tilde{\sigma}}} \\ &\lesssim r^{n(\tilde{\sigma} - \frac{\tilde{\sigma}}{1+\tilde{\sigma}})} \left(\int_{B_r} \varphi(x, |Du|)^{1+\tilde{\sigma}} dx + 1 \right)^{\frac{1}{1+\tilde{\sigma}}}. \end{aligned}$$

Since $1 + \tilde{\sigma} = \frac{\gamma}{\gamma-2}$, we obtain the desired inequality by combining the estimates in E and $B_r \setminus E$ and absorbing the integral over Ψ into the left-hand side. \square

Applying the same approach as in [37, Corollary 6.3], we obtain the following rough comparison estimates from the previous lemma with an unwanted exponent $\frac{1}{q}$ on Θ .

Lemma 5.11. *In the setting of Lemma 5.10, we further assume that*

$$\left(\int_{B_r} \varphi(x, |Du|)^{\frac{\gamma}{\gamma-2}} dx \right)^{\frac{\gamma-2}{\gamma}} \leq C_\gamma^* \tilde{\varphi} \left(\int_{B_{2r}} |Du| dx + 1 \right),$$

for some $C_\gamma^* > 0$. Then there exists $C_2 = C_2(n, p, q, L, \gamma, C_\gamma^*) \geq 1$ such that

$$\int_{B_r} |Du - D\tilde{u}| dx \leq \tilde{\varphi}^{-1} \left(\int_{B_r} \tilde{\varphi}(|Du - D\tilde{u}|) dx \right) \leq C_2 \Theta(r)^{\frac{1}{q}} \left(\int_{B_{2r}} |Du| dx + 1 \right).$$

The next result is a sharper version of the previous lemma with better exponent of Θ . This is analogous to Lemma 3.5 in [6], but much simpler since we consider a single homogeneous equation.

Lemma 5.12. *In the setting of Lemma 5.11, there exists $C_3 = C_3(n, p, q, L, \gamma, C_\gamma^*) \geq 1$ such that if*

$$\frac{\lambda}{\Lambda} \leq \inf_{B_\rho} |D\tilde{u}| \quad \text{and} \quad \int_{B_{2r}} |Du| dx \leq \lambda$$

for some $\lambda, \Lambda \geq 1$ and $B_\rho \subset B_r$, then

$$\int_{B_\rho} |Du - D\tilde{u}| dx \leq C_3 \Lambda^{q-1} \left(\frac{r}{\rho} \right)^n \Theta(r) \lambda.$$

Proof. By the Cauchy–Schwartz inequality,

$$\int_{B_\rho} |Du - D\tilde{u}| dx \leq \Theta(r) \int_{B_\rho} (|Du| + |D\tilde{u}|) dx + \frac{1}{\Theta(r)} \int_{B_\rho} \frac{|Du - D\tilde{u}|^2}{|Du| + |D\tilde{u}|} dx.$$

By Lemma 5.8(3), the first integral on the right-hand side is bounded by $c(\frac{r}{\rho})^n \lambda$. For the second integral, we use that $\tilde{\varphi}'$ is increasing and satisfies $(\text{Dec})_{q-1}$ and that $\frac{\lambda}{\Lambda} \leq |D\tilde{u}| + |Du|$ in B_ρ , to estimate by Lemma 5.10 and the assumption of Lemma 5.11 that

$$\begin{aligned} \left(\frac{\rho}{r} \right)^n \int_{B_\rho} \frac{|Du - D\tilde{u}|^2}{|Du| + |D\tilde{u}|} dx &\leq \frac{1}{\tilde{\varphi}'(\lambda/\Lambda)} \int_{B_r} \frac{\tilde{\varphi}'(|Du| + |D\tilde{u}|)}{|Du| + |D\tilde{u}|} |Du - D\tilde{u}|^2 dx \\ &\lesssim \Lambda^{q-1} \Theta(r)^2 \frac{1}{\tilde{\varphi}'(\lambda)} \tilde{\varphi} \left(\int_{B_{2r}} |Du| dx \right) \\ &\approx \Lambda^{q-1} \Theta(r)^2 \lambda. \end{aligned} \quad \square$$

6. CALDERÓN–ZYGmund ESTIMATES

In this section, we prove the Calderón–Zygmund estimates of Theorem 1.2. We assume throughout that $A : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfies the quasi-isotropic (p, q) -growth condition in Definition 1.1. Let $u \in W_{\text{loc}}^{1,\varphi}(\Omega)$ be a weak solution to $(\operatorname{div} A; F)$. We start with deriving comparison estimates. Note that by Lemma 5.3, $\varphi(\cdot, |Du|) \in L_{\text{loc}}^{1+\sigma}(\Omega)$. Fix $B_{2r} \subset \Omega' \Subset \Omega$ and let $u_r \in u + W_0^{1,\varphi}(B_{2r})$ be the unique weak solution to $(\operatorname{div} A)$ in B_{2r} . Then we have the following energy and higher integrability estimates for Du_r .

Lemma 6.1. *In the setting of Proposition 5.1,*

$$\int_{B_{2r}} \varphi(x, |Du_r|) dx \leq c \int_{B_{2r}} \varphi(x, |Du|) dx.$$

Suppose also that $A^{(-1)}$ satisfies (A1). Then there exist $C_4 \geq 1$ and $\sigma_1 \in (0, \sigma)$ depending on n, p, q and L such that

$$\int_{B_{2r}} \varphi(x, |Du_r|)^{1+\sigma_1} dx \leq C_4 \left(\int_{B_{2r}} \varphi(x, |Du|)^{1+\sigma_1} dx + 1 \right).$$

Proof. The proofs of the estimates are quite standard; hence we only briefly sketch them. By testing $(\operatorname{div} A)$ with $u - u_r \in W_0^{1,\varphi}(B_{2r})$ and using the coercivity property of growth functions and Young's inequality we obtain the first claim. We next prove the second estimate. From Lemma 5.3,

$$\int_{B_r} \varphi(x, |Du_r|)^{1+\sigma} dx \leq c \left\{ \left(\int_{B_{2r}} \varphi(x, |Du_r|) dx \right)^{1+\sigma} + 1 \right\}.$$

Using a similar approach as in the proof of Lemma 5.3 (see, e.g., [16, Lemma 3.5]), we also find the following higher integrability on the boundary:

$$\begin{aligned} & \int_{B_r(y) \cap B_{2r}} \varphi(x, |Du_r|)^{1+\sigma_1} dx \\ & \leq c \left\{ \left(\int_{B_{2r}(y) \cap B_{2r}} \varphi(x, |Du_r|) dx \right)^{1+\sigma_1} + \int_{B_{2r}(y) \cap B_{2r}} \varphi(x, |Du|)^{1+\sigma_1} dx + 1 \right\}, \end{aligned}$$

where $y \in \partial B_{2r}$. The standard covering argument with the previous estimates implies the second claim. \square

Now, suppose that $r > 0$ satisfies the following stronger version of (5.7):

$$(6.2) \quad r \leq \frac{1}{2} \quad \text{and} \quad |B_{2r}| \leq \left\{ 4^{\frac{1+\sigma_1}{\sigma_1}} C_4 \left(\int_{\Omega'} \varphi(x, |Du|)^{1+\sigma_1} dx + 1 \right)^{\frac{2+\sigma_1}{\sigma_1}} \right\}^{-1} \leq \frac{1}{4C_4}.$$

Analogously as (5.7), this and Lemma 6.1 imply that

$$\varrho_{L\varphi(B_{2r})}(|Du|) \leq 1 \quad \text{and} \quad \int_{B_{2r}} \varphi(x, |Du_r|)^{1+\frac{\sigma_1}{2}} dx \leq \frac{1}{2}.$$

Moreover, Lemma 5.8(2) for u_r and $\sigma_1 \in (0, \sigma)$ imply that

$$\left(\int_{B_r} \varphi(x, |Du_r|)^{1+\frac{\sigma_1}{2}} dx \right)^{\frac{1}{1+\sigma_1/2}} \leq c\tilde{\varphi} \left(\int_{B_{2r}} |Du_r| dx + 1 \right)$$

for some $c > 0$ depending on n, p, q and L . Therefore, u_r satisfies the inequalities required in Lemma 5.11 with $\gamma = 2 + \frac{4}{\sigma_1}$. Let $\tilde{u}_r \in u_r + W_0^{1,\tilde{\varphi}}(B_r)$ be the unique weak solution to

($\operatorname{div} \tilde{A}$) where \tilde{A} and $\tilde{\varphi}$ are defined in Section 5.2. Then Lemma 5.11 gives the following comparison estimates:

Lemma 6.3. *In the setting of Lemma 5.8,*

$$\int_{B_r} \varphi(x, |D\tilde{u}_r|) dx \leq c \left(\int_{B_{2r}} \varphi(x, |Du|) dx + 1 \right),$$

and, for any $\varepsilon \in (0, 1)$,

$$\int_{B_r} \varphi(x, |Du - D\tilde{u}_r|) dx \leq c(\varepsilon + \Theta(r)^{\alpha_2}) \left(\int_{B_{2r}} \varphi(x, |Du|) dx + 1 \right) + \varepsilon^{-\frac{p+1}{p}} \int_{B_{2r}} \varphi(x, |F|) dx$$

for some $c > 0$ and $\alpha_2 \in (0, 1)$ depending on n, p, q and L .

Proof. We first obtain a comparison estimate between Du and Du_r . We take $u - u_r \in W_0^{1,\varphi}(B_{2r})$ as a test function in the weak formulations of ($\operatorname{div} A; F$) and ($\operatorname{div} A$) to discover that

$$\int_{B_{2r}} (A(x, Du) - A(x, Du_r)) \cdot (Du - Du_r) dx = \int_{B_{2r}} A(x, F) \cdot (Du - Du_r) dx.$$

By [37, Proposition 3.8(3)], the monotonicity and coercivity properties of growth functions, the equation above, and Lemma 6.1, we see that

$$\begin{aligned} & \int_{B_{2r}} \varphi(x, |Du - Du_r|) dx \\ & \lesssim \varepsilon \int_{B_{2r}} \varphi(x, |Du|) + \varphi(x, |Du_r|) dx + \varepsilon^{-1} \int_{B_{2r}} \frac{\varphi'(x, |Du| + |Du_r|)}{|Du| + |Du_r|} |Du - Du_r|^2 dx \\ & \lesssim \varepsilon \int_{B_{2r}} \varphi(x, |Du|) dx + \varepsilon^{-1} \int_{B_{2r}} \varphi'(x, |F|) |Du - Du_r| dx. \end{aligned}$$

Then, using Young's inequality we have

$$(6.4) \quad \int_{B_{2r}} \varphi(x, |Du - Du_r|) dx \lesssim \varepsilon \int_{B_{2r}} \varphi(x, |Du|) dx + \varepsilon^{-\frac{p+1}{p}} \int_{B_{2r}} \varphi(x, |F|) dx.$$

Let $\tau > 0$ be the solution of $\frac{\tau}{1+\sigma_1/2} + q(1-\tau) = 1$. By Hölder inequality with exponents $\frac{1+\sigma_1/2}{\tau}$ and $\frac{1}{q(1-\tau)}$ applied to $\varphi = \varphi^\tau \varphi^{1-\tau}$, we derive that

$$\begin{aligned} & \int_{B_r} \varphi(x, |Du_r - D\tilde{u}_r|) dx \\ & \leq \left(\int_{B_r} \varphi(x, |Du_r - D\tilde{u}_r|)^{1+\frac{\sigma_1}{2}} dx \right)^{\frac{\tau}{1+\sigma_1/2}} \left(\int_{B_r} \varphi(x, |Du_r - D\tilde{u}_r|)^{\frac{1}{q}} dx \right)^{q(1-\tau)}. \end{aligned}$$

By Lemma 5.8(4) and Lemma 5.3 for u_r with $\sigma = \sigma_1$ we have

$$\begin{aligned} \left(\int_{B_r} \varphi(x, |Du_r - D\tilde{u}_r|)^{1+\frac{\sigma_1}{2}} dx \right)^{\frac{1}{1+\sigma_1/2}} & \lesssim \left(\int_{B_r} \varphi(x, |Du_r|)^{1+\frac{\sigma_1}{2}} dx \right)^{\frac{1}{1+\sigma_1/2}} + 1 \\ & \lesssim \int_{B_{2r}} \varphi(x, |Du_r|) dx + 1. \end{aligned}$$

Moreover, from by Jensen's inequality, Lemma 5.11 for u with $\gamma = 2 + \frac{4}{\sigma_1}$ and (A1) with the fact that $\int_{B_{2r}} |Du_r| dx \lesssim (\varphi_{B_{2r}}^-)^{-1}(|B_{2r}|^{-1})$, we obtain that

$$\begin{aligned} \left(\int_{B_r} \varphi(x, |Du_r - D\tilde{u}_r|)^{\frac{1}{q}} dx \right)^q &\leq \left(\int_{B_r} \varphi_{B_r}^+ (|Du_r - D\tilde{u}_r|)^{\frac{1}{q}} dx \right)^q \\ &\lesssim \varphi_{B_r}^+ \left(\int_{B_r} |Du_r - D\tilde{u}_r| dx \right) \lesssim \Theta(r)^{\frac{p}{q}} \varphi_{B_r}^+ \left(\int_{B_{2r}} |Du_r| dx + 1 \right) \\ &\lesssim \Theta(r)^{\frac{p}{q}} \varphi_{B_{2r}}^- \left(\int_{B_{2r}} |Du_r| dx + 1 \right) \lesssim \Theta(r)^{\frac{p}{q}} \left(\int_{B_{2r}} \varphi(x, |Du_r|) dx + 1 \right). \end{aligned}$$

Combining the previous three estimates yields

$$\int_{B_r} \varphi(x, |Du_r - D\tilde{u}_r|) dx \lesssim \Theta(r)^{(1-\tau)\frac{p}{q}} \left(\int_{B_{2r}} \varphi(x, |Du_r|) dx + 1 \right).$$

We obtain the desired estimates from this, (6.4) and Lemma 6.1. \square

Now, we are ready to prove the Calderón–Zygmund estimate.

Proof of Theorem 1.2. Since we have obtained the relevant comparison estimates, we can prove the theorem by using the well-known argument mentioned in Section 1. Therefore, we will provide a sketch of the proof and refer to, for instance, our recent paper [44] for details. We first observe that by Proposition 3.8,

$$\lim_{r \rightarrow 0^+} \theta_\gamma(r) = 0 \quad \text{for } \gamma = 2 + \frac{4}{\sigma_1}.$$

Let $\delta \in (0, 1)$ be a small constant to be determined later, and $B_{2r} \subset \Omega' \Subset \Omega$, where $r > 0$ is a small number which satisfies (6.2) and $\Theta(r)^{\alpha_2} \leq \varepsilon := \delta^{\frac{p}{2p+1}p}$ with notation from Lemma 6.3. Set

$$(6.5) \quad \lambda_0 := \int_{B_{2r}} \varphi(x, |Du|) dx + \frac{1}{\delta} \int_{B_{2r}} \varphi(x, |F|) dx + 1$$

and for $\rho \in (0, 2r]$ and $\lambda > 0$

$$E(\lambda, \rho) = E(\lambda, B_\rho) := \{x \in B_\rho : \varphi(x, |Du|) > \lambda\}.$$

By the Vitali covering lemma, for a given $\lambda > \alpha\lambda_0$, where $\alpha := (\frac{20}{\tau_2 - \tau_1})^n$, and $1 \leq \tau_1 < \tau_2 \leq 2$, there exists a disjoint family of balls $\{B_{\rho_i}(y^i)\}_{i=1}^\infty$ with $y^i \in E(\lambda, \tau_1 r)$ and $\rho_i \in (0, \frac{(\tau_2 - \tau_1)r}{10})$ such that

$$\begin{aligned} E(\lambda, \tau_1 r) &\subset \bigcup_{i=1}^\infty B_{5\rho_i}(y^i), \\ \int_{B_{\rho_i}(y^i)} \varphi(x, |Du|) dx + \frac{1}{\delta} \int_{B_{\rho_i}(y^i)} \varphi(x, |F|) dx &= \lambda, \end{aligned}$$

and, for any $\rho \in (\rho_i, (\tau_2 - \tau_1)r]$,

$$\int_{B_\rho(y^i)} \varphi(x, |Du|) dx + \frac{1}{\delta} \int_{B_\rho(y^i)} \varphi(x, |F|) dx < \lambda.$$

Note that the previous inequality can be applied when $\rho = 10\rho_i$ and that the penultimate display yields

$$(6.6) \quad \frac{\lambda}{2} |B_{\rho_i}(y^i)| \leq \int_{B_{\rho_i}(y^i) \cap \{\varphi(x, |Du|) > \frac{\lambda}{4}\}} \varphi(x, |Du|) dx + \frac{1}{\delta} \int_{B_{\rho_i}(y^i) \cap \{\varphi(x, |F|) > \frac{\delta\lambda}{4}\}} \varphi(x, |F|) dx.$$

We use a two-step approximation approach in order to be able to apply our earlier results. For each i , we consider the unique solution $h_r^i \in u + W_0^{1,\varphi}(B_{10\rho_i}(y^i))$ to $(\operatorname{div}A)$ in $B_{10\rho_i}(y^i)$ and the unique weak solution $\tilde{u}_r^i \in h_r^i + W_0^{1,\tilde{\varphi}}(B_{5\rho_i}(y^i))$ to $(\operatorname{div}\tilde{A})$ in $B_{5\rho_i}(y^i)$. Then by Lemma 6.3 and $\Theta(r)^{\alpha_2} \leq \varepsilon$, we have

$$(6.7) \quad \int_{B_{5\rho_i}(y^i)} \varphi(x, |Du - D\tilde{u}_r^i|) dx \lesssim \varepsilon\lambda.$$

Moreover, by Lemma 5.8(3), Jensen's inequality, Lemma 6.1 and $\varrho_{L^\varphi(B_{2r})}(|Du|) \leq 1$ we conclude that

$$\begin{aligned} \varphi_{B_{10\rho_i}(y^i)}^- \left(\int_{B_{5\rho_i}(y^i)} |D\tilde{u}_r^i| dx \right) &\lesssim \varphi_{B_{10\rho_i}(y^i)}^- \left(\int_{B_{10\rho_i}(y^i)} |Dh_r^i| dx + 1 \right) \\ &\lesssim \int_{B_{10\rho_i}(y^i)} \varphi(x, |Dh_r^i|) dx + 1 \lesssim |B_{10\rho_i}|^{-1}. \end{aligned}$$

Hence by Lemma 5.9, (A1) and Jensen's inequality, we have

$$\begin{aligned} \varphi(y, \|D\tilde{u}_r^i\|_{L^\infty(B_{5\rho_i}(y^i))}) &\leq c\varphi\left(y, \int_{B_{10\rho_i}(y^i)} |D\tilde{u}_r^i| dx\right) \\ &\leq c\varphi_{B_{5\rho_i}(y^i)}^- \left(\int_{B_{10\rho_i}(y^i)} |D\tilde{u}_r^i| dx + 1 \right) \leq c \int_{B_{10\rho_i}(y^i)} \varphi(x, |D\tilde{u}_r^i|) dx + 1 \leq c_0\lambda \end{aligned}$$

for some constant $c_0 \geq 1$.

Set $K := 2^q c_0$ and let $y \in B_{5\rho_i}(y^i)$ with $\varphi(y, |Du|) > K\lambda$. Then

$$\varphi(y, |D\tilde{u}_r^i|) \leq c_0\lambda = 2^{-q}K\lambda < 2^{-q}\varphi(y, |Du|) \leq \varphi(y, \frac{1}{2}|Du|).$$

Therefore $|D\tilde{u}_r^i| \leq \frac{1}{2}|Du|$ so that $|Du| \leq 2|Du - D\tilde{u}_r^i|$, which implies that

$$\varphi(y, |Du|) \leq 2^q \varphi(y, |Du - D\tilde{u}_r^i|).$$

By (6.6) and (6.7), we then derive that

$$\begin{aligned} &\int_{B_{5\rho_i}(y^i) \cap E(K\lambda, \tau_1 r)} \varphi(x, |Du|) dx \\ &\leq 2^q \int_{B_{5\rho_i}(y^i)} \varphi(x, |Du - D\tilde{u}_r^i|) dx \leq c\varepsilon\lambda |B_{\rho_i}(y^i)| \\ &\leq c\varepsilon \left(\int_{B_{\rho_i}(y^i) \cap \{\varphi(x, |Du|) > \frac{\lambda}{4}\}} \varphi(x, |Du|) dx + \frac{1}{\delta} \int_{B_{\rho_i}(y^i) \cap \{\varphi(x, |F|) > \frac{\delta\lambda}{4}\}} \varphi(x, |F|) dx \right). \end{aligned}$$

Note that the balls $B_{\rho_i}(y^i)$ are mutually disjoint and

$$E(K\lambda, \tau_1 r) \subset E(\lambda, \tau_1 r) \subset \bigcup_{i=1}^{\infty} B_{5\rho_i}(y^i) \subset B_{\tau_2 r}.$$

Thus

$$\begin{aligned} \int_{E(K\lambda, \tau_1 r)} \varphi(x, |Du|) dx &\leq \sum_{i=1}^{\infty} \int_{B_{5\rho_i}(y^i) \cap E(K\lambda, \tau_1 r)} \varphi(x, |Du|) dx \\ &\leq c\varepsilon \left(\int_{B_{\tau_2 r} \cap \{\varphi(x, |Du|) > \frac{\lambda}{4}\}} \varphi(x, |Du|) dx + \frac{1}{\delta} \int_{B_{\tau_2 r} \cap \{\varphi(x, |F|) > \frac{\delta\lambda}{4}\}} \varphi(x, |F|) dx \right) \end{aligned}$$

for some constant $c > 0$.

We next apply a truncation argument to complete the proof. For $k \geq \lambda$, let us define $\varphi_k(x, |Du|) := \min \{\varphi(x, |Du|), k\}$, and consider the super-level set of φ_k

$$E_k(\lambda, \rho) := \{y \in B_\rho : \varphi_k(y, |Du|) > \lambda\} \quad \text{for } \lambda, \rho > 0.$$

Note that $E_k(\lambda, \rho) = E(\lambda, \rho)$ if $\lambda \leq k$ and $E_k(\lambda, \rho) = \emptyset$ if $\lambda > k$. Hence we can continue the earlier estimate for any $\lambda \geq \alpha\lambda_0$,

$$\int_{E_k(K\lambda, \tau_1 r)} \varphi(x, |Du|) dx \leq c\varepsilon \left(\int_{E_k(\frac{\lambda}{4}, \tau_2 r)} \varphi(x, |Du|) dx + \frac{1}{\delta} \int_{B_{\tau_2 r} \cap \{\varphi(x, |F|) > \frac{\delta\lambda}{4}\}} \varphi(x, |F|) dx \right).$$

Then we multiply both sides by λ^{s-2} , $s \in (1, \infty)$, and integrate with respect to λ over $(\alpha\lambda_0, \infty)$ to discover that

$$\begin{aligned} I_0 &:= \int_{\alpha\lambda_0}^{\infty} \lambda^{s-2} \int_{E_k(K\lambda, \tau_1 r)} \varphi(x, |Du|) dx d\lambda \\ &\leq c\varepsilon \left(\int_{\alpha\lambda_0}^{\infty} \lambda^{s-2} \int_{E_k(\frac{\lambda}{4}, \tau_2 r)} \varphi(x, |Du|) dx d\lambda \right. \\ &\quad \left. + \int_{\alpha\lambda_0}^{\infty} \lambda^{s-2} \int_{B_{\tau_2 r} \cap \{\frac{\varphi(x, |F|)}{\delta} > \frac{\lambda}{4}\}} \frac{\varphi(x, |F|)}{\delta} dx d\lambda \right) \\ &=: c\varepsilon(I_1 + I_2). \end{aligned}$$

By Fubini's theorem, we then derive that

$$I_0 = \frac{1}{s-1} \int_{E_k(K\alpha\lambda_0, \tau_1 r)} \varphi(x, |Du|) \left[\left(\frac{\varphi_k(x, |Du|)}{K} \right)^{s-1} - (\alpha\lambda_0)^{s-1} \right] dx,$$

$$\begin{aligned} I_1 &= \int_{E_k(\frac{\alpha\lambda_0}{4}, \tau_2 r)} \varphi(x, |Du|) \left(\int_{\alpha\lambda_0}^{4\varphi_k(x, |Du|)} \lambda^{s-2} d\lambda \right) dx \\ &\leq \frac{4^{s-1}}{s-1} \int_{B_{\tau_2 r}} \varphi(x, |Du|) \varphi_k(x, |Du|)^{s-1} dx, \end{aligned}$$

and, similarly,

$$I_2 \leq \frac{4^{s-1}}{s-1} \int_{B_{\tau_2 r}} \left[\frac{\varphi(x, |F|)}{\delta} \right]^s dx.$$

With these expressions for I_0 , I_1 , I_2 , we continue our earlier estimate as

$$\begin{aligned} &\int_{B_{\tau_1 r}} \varphi(x, |Du|) \varphi_k(x, |Du|)^{s-1} dx \\ &\leq \int_{E_k(K\alpha\lambda_0, \tau_1 r)} \varphi(x, |Du|) \varphi_k(x, |Du|)^{s-1} dx + (K\alpha\lambda_0)^{s-1} \int_{B_{\tau_1 r}} \varphi(x, |Du|) dx \\ &\leq c_1\varepsilon \int_{B_{\tau_2 r}} \varphi(x, |Du|) \varphi_k(x, |Du|)^{s-1} dx + c(\alpha\lambda_0)^{s-1} \int_{B_{2r}} \varphi(x, |Du|) dx \\ &\quad + c\varepsilon \int_{B_{2r}} \left[\frac{\varphi(x, |F|)}{\delta} \right]^s dx \end{aligned}$$

for some $c_1 = c_1(n, p, q, L, L_1) \geq 0$. At this stage, we fix $\varepsilon \in (0, 1)$ such that $c_1\varepsilon \leq \frac{1}{2}$, which also determines δ and r . From the definition of α we finally have

$$\begin{aligned} \int_{B_{r_1 r}} \varphi(x, |Du|) \varphi_k(x, |Du|)^{s-1} dx &\leq \frac{1}{2} \int_{B_{r_2 r}} \varphi(x, |Du|) \varphi_k(x, |Du|)^{s-1} dx \\ &\quad + \frac{c\lambda_0^{s-1}}{(\tau_2 - \tau_1)^{n(s-1)}} \int_{B_{2r}} \varphi(x, |Du|) dx + c \int_{B_{2r}} \varphi(x, |F|)^s dx. \end{aligned}$$

Then a standard iteration argument for $1 \leq \tau_1 < \tau_2 \leq 2$ yields

$$\int_{B_r} \varphi(x, |Du|) \varphi_k(x, |Du|)^{s-1} dx \leq c\lambda_0^{s-1} \int_{B_{2r}} \varphi(x, |Du|) dx + c \int_{B_{2r}} \varphi(x, |F|)^s dx.$$

Therefore, from Lebesgue's monotone convergence theorem, Hölder's inequality and Young's inequality together with the definition of λ_0 in (6.5), we derive that

$$\begin{aligned} \int_{B_r} \varphi(x, |Du|)^s dx &= \lim_{k \rightarrow \infty} \int_{B_r} \varphi(x, |Du|) \varphi_k(x, |Du|)^{s-1} dx \\ &\leq c\lambda_0^{s-1} \int_{B_{2r}} \varphi(x, |Du|) dx + c \int_{B_{2r}} \varphi(x, |F|)^s dx \\ &\leq c \left(\int_{B_{2r}} \varphi(x, |Du|) dx \right)^s + c \int_{B_{2r}} \varphi(x, |F|)^s dx + c, \end{aligned}$$

which together with the growth condition of growth functions implies the desired estimates. \square

7. CONTINUITY OF THE DERIVATIVE

We next prove Theorem 1.3. In this section we always suppose that $A : \Omega \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfies the quasi-isotropic (p, q) -growth condition and $A^{(-1)}$ satisfies the $(DMA1)_\gamma$ condition for some $\gamma > 2$. Then $A^{(-1)}$ also satisfies the $(VMA1)$ condition and, by Proposition 3.5, the $(SA1)$ condition, and hence the $(A1)$ condition, with relevant constant L depending on n, p, q, L, L_1 and θ_γ . By Theorem 1.2 with $F \equiv 0$, $\varphi(\cdot, |Du|) \in L^s_{\text{loc}}(\Omega)$ for any $s > 1$. Moreover, for $\Omega' \Subset \Omega$ there exists $R_1 > 0$ depending on $n, p, q, L, \gamma, \theta_\gamma, Du$ and Ω' and such that for any $B_{2r} \subset \Omega'$ with $r \leq R_1$, the condition (5.7) holds,

$$\varrho_{L\varphi^{\gamma/(\gamma-2)}(B_{2r})}(|Du|) = \int_{B_{2r}} \varphi(x, |Du|)^{\frac{\gamma}{\gamma-2}} dx \leq 1,$$

and, by Theorem 1.2 with $F \equiv 0$, $s = \frac{\gamma}{\gamma-2}$ and scaled balls,

$$\left(\int_{B_r} \varphi(x, |Du|)^{\frac{\gamma}{\gamma-2}} dx \right)^{\frac{\gamma-2}{\gamma}} \leq c \left(\int_{B_{3r/2}} \varphi(x, |Du|) dx + 1 \right) \leq C_\gamma^* \tilde{\varphi} \left(\int_{B_{2r}} |Du| dx + 1 \right),$$

where the last inequality follows from Lemma 5.8(2), with scaled balls, and the constant $C_\gamma^* > 0$ depends on n, p, q, L and γ . Therefore, we obtain the necessary inequalities in the assumptions to apply Lemmas 5.11 and 5.12.

Let $u \in W_{\text{loc}}^{1,\varphi}(\Omega)$ be a weak solution to $(\text{div} A)$. Fix $B_{2r}(x_0) \subset \Omega'$ with $r \leq R_1$. For $\delta \in (0, \frac{1}{4})$ to be determined later and $k \in \mathbb{N} \cup \{0\}$, set

$$\Theta_k := \Theta(\delta^k r), \quad B_k := B_{\delta^k r}(x_0), \quad \text{and} \quad E_k := \int_{B_k} |Du - (Du)_{B_k}| dx,$$

where Θ is defined in Lemma 5.10. Suppose that

$$\delta \leq \delta_1,$$

where δ_1 is given in Lemma 3.6. Then this lemma implies that

$$\sum_{k=0}^{\infty} \Theta_k = \sum_{k=0}^{\infty} \Theta(\delta^k r) \leq \frac{c}{\delta^{2n/\gamma+1} \log \delta^{-1}} \int_0^r \Theta(\rho) \frac{d\rho}{\rho} < \infty.$$

For each $k \in \mathbb{N}$, we construct a function $\tilde{\varphi}$ in the ball B_k as in Section 5 and let $\tilde{u}_k \in u + W_0^{1,\tilde{\varphi}}(B_k)$ be the weak solution to $(\operatorname{div} \tilde{A})$ in B_k . We will use the named constants C_0 , C_1 , C_2 and C_3 from Lemmas 5.8, 5.9, 5.11 and 5.12. The following results hold assuming only the inequalities from these four results.

Let us first record an estimate for E_{k+1} with $k \geq j$. We integrate the inequality

$$|Du - (Du)_{B_{k+1}}| \leq |Du - D\tilde{u}_j| + |D\tilde{u}_j - (D\tilde{u}_j)_{B_{k+1}}| + |(D\tilde{u}_j)_{B_{k+1}} - (Du)_{B_{k+1}}|$$

over B_{k+1} , then use Lemma 5.9 with $(B_\rho, B_\rho) \leftarrow (B_{k+1}, B_k)$ and choose δ small so that $C_1 \delta^\alpha \leq \frac{1}{32}$, to conclude that

$$\begin{aligned} E_{k+1} &\leq \int_{B_{k+1}} |D\tilde{u}_j - (D\tilde{u}_j)_{B_{k+1}}| dx + 2 \int_{B_{k+1}} |Du - D\tilde{u}_j| dx \\ &\leq C_1 \delta^\alpha \int_{B_k} |D\tilde{u}_j - (D\tilde{u}_j)_{B_k}| dx + 2\delta^{-n} \int_{B_k} |Du - D\tilde{u}_j| dx \\ (7.1) \quad &\leq 2C_1 \delta^\alpha \int_{B_k} |D\tilde{u}_j - (Du)_{B_k}| dx + 2\delta^{-n} \int_{B_k} |Du - D\tilde{u}_j| dx \\ &\leq \frac{1}{16} E_k + 4\delta^{-n} \int_{B_k} |Du - D\tilde{u}_j| dx. \end{aligned}$$

Lemma 7.2. *In the setting fixed in the beginning of the section, let $\delta \in (0, \delta_1]$, $\varepsilon \in (0, \frac{1}{4})$ and $k \in \mathbb{N}$ satisfy*

$$\max\{C_2 \delta^{-n} \Theta_k^{1/q}, 32C_0 C_1 \delta^\alpha\} \leq \varepsilon,$$

where $\alpha \in (0, 1)$ is from Lemma 5.9. If

$$\int_{2B_k} |Du| dx \leq \lambda \quad \text{and} \quad \int_{2B_{k+1}} |Du| dx \geq \varepsilon \lambda,$$

for some $\lambda \geq 1$, then

$$E_{k+2} \leq \frac{1}{16} E_{k+1} + 4^q C_3 \varepsilon^{-(q-1)} \delta^{-2n} \Theta_k \lambda.$$

Proof. We use the second inequality in the assumption and Lemma 5.11 in B_k to conclude that

$$\varepsilon \lambda \leq \int_{2B_{k+1}} |Du - D\tilde{u}_k| dx + \int_{2B_{k+1}} |D\tilde{u}_k| dx \leq 2^{1-n} C_2 \delta^{-n} \Theta_k^{1/q} \lambda + \int_{2B_{k+1}} |D\tilde{u}_k| dx.$$

By the assumption $C_2 \delta^{-n} \Theta_k^{1/q} \leq \varepsilon$, the first term on the right is at most $\frac{\varepsilon \lambda}{2}$ and can be absorbed into the left-hand side. By Lemma 5.9 with $(B_\rho, B_r) \leftarrow (2B_{k+1}, \tilde{B}_k)$, we have

$$\inf_{2B_{k+1}} |D\tilde{u}_k| \geq \int_{2B_{k+1}} |D\tilde{u}_k| dx - \operatorname{osc}_{2B_{k+1}} |D\tilde{u}_k| \geq \frac{\varepsilon \lambda}{2} - 2^{1+\alpha} C_1 \delta^\alpha \int_{B_k} |D\tilde{u}_k| dx \geq \frac{\varepsilon \lambda}{4},$$

where in the last step we used Lemma 5.8(3) with $\tilde{u} = \tilde{u}_k$ and the assumption $32C_0C_1\delta^\alpha \leq \varepsilon$. Therefore, applying Lemma 5.12 with $(\tilde{u}, \tilde{C}, B_r, B_\rho) \leftarrow (\tilde{u}_k, \frac{4}{\varepsilon}, B_k, B_{k+1})$, we estimate

$$\int_{B_{k+1}} |Du - D\tilde{u}_k| dx \leq C_3 \left(\frac{4}{\varepsilon}\right)^{q-1} \delta^{-n} \Theta_k \lambda.$$

Note that $C_1\delta^\alpha \leq \frac{1}{32}$ follows from our assumption so we can use (7.1) with $(k, j) \leftarrow (k+1, k)$ and the previous estimate to conclude that

$$E_{k+2} \leq \frac{1}{16} E_{k+1} + 4\delta^{-n} \int_{B_{k+1}} |Du - D\tilde{u}_k| dx \leq \frac{1}{16} E_{k+1} + 4^q C_3 \varepsilon^{-(q-1)} \delta^{-2n} \Theta_k \lambda. \quad \square$$

Lemma 7.3. *In the setting fixed in the section, let $\delta \in (0, \delta_1]$, $\varepsilon_0 := 2^{-n-2}$, $\varepsilon \in (0, \varepsilon_0]$ and $j \in \mathbb{N}$ satisfy the conditions*

$$\max\{64C_2\delta^{-2n} \sup_{k \geq j} \Theta_k^{1/q}, 32C_0C_1\delta^\alpha\} \leq \varepsilon_0\varepsilon \quad \text{and} \quad \sum_{k=j}^{\infty} \Theta_k \leq \frac{\delta^{3n}\varepsilon_0\varepsilon^q}{8 \cdot 4^q C_3}.$$

For $m \geq j+1$, we assume that

$$(7.4) \quad \frac{1}{\delta^n \varepsilon} E_j + \int_{2B_j} |Du| dx \leq 2\varepsilon_0 \lambda$$

and

$$(7.5) \quad \int_{2B_k} |Du| dx \geq \varepsilon \lambda \quad \text{for every } k \in \{j+1, \dots, m-1\} \quad \text{when } m \geq j+2,$$

for some $\lambda \geq 1$. Then

$$E_m \leq \frac{\delta^n \varepsilon_0 \varepsilon}{2} \lambda, \quad \int_{B_{m+1}} |Du - (Du)_{B_j}| dx \leq 3\varepsilon_0 \varepsilon \lambda \quad \text{and} \quad \int_{2B_m} |Du| dx \leq \lambda.$$

Proof. We first note that (7.1) with $k = j$, (7.4) and Lemma 5.11 in B_j imply that

$$E_{j+1} \leq \frac{1}{16} E_j + 4\delta^{-n} \int_{B_j} |Du - D\tilde{u}_j| dx \leq \frac{\delta^n \varepsilon_0 \varepsilon}{8} \lambda + 4C_2 \delta^{-n} \Theta_j^{1/q} \left(\int_{2B_j} |Du| dx + 1 \right).$$

By assumption $4C_2\delta^{-n}\Theta_j^{1/q} \leq \frac{1}{16}\delta^n\varepsilon_0\varepsilon$ and $\int_{2B_j} |Du| dx + 1 \leq \frac{3}{2}\lambda$ so that $E_{j+1} \leq \frac{1}{4}\delta^n\varepsilon_0\varepsilon\lambda$. Moreover, by the assumption (7.4),

$$\begin{aligned} \int_{2B_{j+1}} |Du| dx &\leq \int_{2B_{j+1}} |Du - (Du)_{B_j}| dx + |(Du)_{B_j}| \leq (2\delta)^{-n} E_j + 2^n \int_{2B_j} |Du| dx \\ &\leq 2^n \cdot 2\varepsilon_0 \lambda \leq \lambda. \end{aligned}$$

Using also the previous estimate for E_{j+1} , we further obtain that

$$\begin{aligned} \int_{B_{j+2}} |Du - (Du)_{B_j}| dx &\leq \int_{B_{j+2}} |Du - (Du)_{B_{j+1}}| dx + |(Du)_{B_{j+1}} - (Du)_{B_j}| \\ &\leq \delta^{-n} (E_{j+1} + E_j) \leq \left(\frac{1}{4} + 2\right) \varepsilon_0 \lambda \leq 3\varepsilon_0 \lambda. \end{aligned}$$

Thus we have proved all the claims in the case $m = j+1$.

We prove the claim by induction when $m \geq j+2$; specifically, we assume that the claim holds up to $m-1$ and show it for m . By the induction assumption $\int_{2B_k} |Du| dx \leq \lambda$ when $k \leq m-1$. The assumption (7.5) gives $\int_{2B_{k+1}} |Du| dx \geq \varepsilon \lambda$ for $k \in \{j, \dots, m-2\}$. Thus we can use Lemma 7.2 to conclude that

$$2E_{k+2} - E_{k+1} \leq c' \Theta_k \lambda$$

for $k \in \{j, \dots, m-2\}$ and $c' := 2 \cdot 4^q C_3 \varepsilon^{-(q-1)} \delta^{-2n}$. Therefore

$$(7.6) \quad \sum_{k=j+1}^m E_k = \sum_{k=j}^{m-2} (2E_{k+2} - E_{k+1}) + 2E_{j+1} - E_m \leq c' \lambda \sum_{k=j}^{m-2} \Theta_k + 2E_{j+1} \leq \frac{1}{2} \delta^n \varepsilon_0 \varepsilon \lambda$$

since $\sum_{k=j}^{\infty} \Theta_k \leq \frac{\delta^n \varepsilon_0 \varepsilon}{4c'}$ by assumption and $E_{j+1} \leq \frac{\delta^n \varepsilon_0 \varepsilon}{4} \lambda$ as shown above. In particular, the first claim $E_m \leq \frac{1}{2} \delta^n \varepsilon_0 \varepsilon \lambda$ is proved. We write

$$Du - (Du)_{B_j} = [Du - (Du)_{B_m}] + [(Du)_{B_m} - (Du)_{B_{m-1}}] + \dots + [(Du)_{B_{j+1}} - (Du)_{B_j}].$$

Integrating this over B_{m+1} and using (7.4) and (7.6) to estimate E_k , we observe that

$$\int_{B_{m+1}} |Du - (Du)_{B_j}| dx \leq \sum_{k=j}^m \int_{B_{k+1}} |Du - (Du)_{B_k}| dx \leq \delta^{-n} \sum_{k=j}^m E_k \leq 3\varepsilon_0 \varepsilon \lambda.$$

This proves the second claim. Similarly, we prove the inequality $\int_{2B_m} |Du - (Du)_{B_j}| dx \leq (2\delta)^{-n} \sum_{k=j}^{m-1} E_k \leq \frac{3\varepsilon_0 \varepsilon}{2^n} \lambda$. Using this and the estimate $\int_{2B_j} |Du| dx \leq \frac{3}{2} \varepsilon_0 \lambda = 3 \cdot 2^{-n-3} \lambda$, we obtain

$$\int_{2B_m} |Du| dx \leq \int_{2B_m} |Du - (Du)_{B_j}| dx + 2^n \int_{2B_j} |Du| dx \leq \frac{3\varepsilon_0 \varepsilon}{2^n} \lambda + \frac{1}{8} \lambda \leq \lambda.$$

This concludes the proof of the third claim in the induction step, so the claim holds for any number of steps. \square

Proposition 7.7. *In the setting fixed in the beginning of the section, the gradient Du is locally essentially bounded. Moreover,*

$$\|Du\|_{L^\infty(B_r)} \leq c \left(\int_{B_{2r}} |Du| dx + 1 \right)$$

for some $c = c(n, p, q, \gamma, L, L_1, L_\gamma, \theta_\gamma) \geq 1$ and any $B_{2r} \subset \Omega'$ with $r \in (0, R_1]$.

Proof. Let $\varepsilon := \varepsilon_0 = 2^{-n-2}$ and choose $\delta := \min\{\delta_1, (\frac{\varepsilon_0^2}{32C_0C_1})^{1/\alpha}\}$. Let j_0 satisfy the conditions of Lemma 7.3 with $j = j_0$; such j_0 exists since the series $\sum \Theta_k$ is convergent.

Now we fix a Lebesgue point $x_0 \in B_r$ of Du and consider balls $B_k := B_{\delta^k r}(x_0)$, $k \in \mathbb{N}$. For $k \geq j_0$, we define

$$F_k := \frac{1}{\delta^n \varepsilon_0} E_k + \int_{2B_k} |Du| dx \quad \text{and} \quad \lambda := \frac{1}{2\varepsilon_0} F_{j_0} + 1.$$

Since $\lambda \leq F_{j_0} \lesssim \delta^{-j_0} \int_{B_{2r}} |Du| dx$ and both δ and j_0 depend only on the parameters and can be included in the constant c , we complete the proof by showing that $|Du(x_0)| \leq \lambda$.

Since x_0 is a Lebesgue point,

$$|Du(x_0)| \leq \liminf_{k \rightarrow \infty} \int_{2B_k} |Du| dx.$$

The claim follows if $F_k \leq 2\varepsilon_0 \lambda$ for infinitely many k . If this is not the case we let $j \geq j_0$ be the largest index with $F_j \leq 2\varepsilon_0 \lambda$; note that such index exists as $F_{j_0} \leq 2\varepsilon_0 \lambda$ by the choice of λ . Then $F_k \geq 2\varepsilon_0 \lambda$ for every $k > j$. Now Lemma 7.3 with $m = j+1$ implies that $E_{j+1} \leq \frac{1}{2} \delta^n \varepsilon_0^2 \lambda$. Therefore,

$$\int_{2B_{j+1}} |Du| dx = F_{j+1} - \delta^{-n} \varepsilon_0^{-1} E_{j+1} \geq 2\varepsilon_0 \lambda - \frac{1}{2} \varepsilon_0 \lambda \geq \varepsilon_0 \lambda.$$

From this we conclude that (7.5) is satisfied for $m = j + 2$, which by the lemma in turn implies that $E_{j+2} \leq \frac{1}{2} \delta^n \varepsilon_0^2 \lambda$. This in turn implies $E_{j+3} \leq \frac{1}{2} \delta^n \varepsilon_0 \lambda$, and so on. Thus (7.5) holds for every $k > j$ and so Lemma 7.3 implies that $\int_{2B_m} |Du| dx \leq \lambda$ for every $m \geq j$, so that $|Du(x_0)| \leq \lambda$. \square

Now, we are ready to prove the continuity of the derivative Du .

Proof of Theorem 1.3. We continue in the setting fixed in the beginning of the section and note that Proposition 7.7 implies that Du is bounded in Ω' . Let $\lambda := \varepsilon_0^{-1} \|Du\|_{L^\infty(\Omega')} + 1$. For $\varepsilon \in (0, \varepsilon_0]$, choose $\delta := \min\{\delta_1, (\frac{\varepsilon_0 \varepsilon}{32C_0C_1})^{1/\alpha}\}$ and then j_0 so large that the conditions of Lemma 7.3 are satisfied when $j = j_0$. We observe that $E_{j_0} \leq \lambda$ by the definition of λ and $E_{k+1} \leq \frac{1}{16} E_k + \frac{1}{8} \delta^n \varepsilon_0 \varepsilon \lambda$ for any $k \geq j_0$ (see beginning of the proof of Lemma 5.11). Thus we may assume by increasing j_0 if necessary that $E_j \leq \delta^n \varepsilon_0 \varepsilon \lambda$ for all $j \geq j_0$. Then assumption (7.4) of Lemma 7.3 holds for any $j \geq j_0$.

Fix $\Omega'' \Subset \Omega'$ and assume by increasing j_0 if necessary that $2B_{j_0}(x) \subset \Omega'$ for every $x \in \Omega''$. Consider Lebesgue points $x, y \in \Omega''$ of Du so close to one another that $|B_j(x) \setminus B_j(y)| \leq \varepsilon |B_j|$ for some $j \geq j_0$. Suppose that

$$|(Du)_{B_j} - (Du)_{B_{m+1}}| \leq (6 + 2^{n+1}) \varepsilon \lambda$$

for all $m \geq j + 2$ (this is proved below). Then

$$\begin{aligned} |Du(x) - Du(y)| &\leq \lim_{m \rightarrow \infty} |(Du)_{B_j(x)} - (Du)_{B_{m+1}(x)}| + \lim_{m \rightarrow \infty} |(Du)_{B_j(y)} - (Du)_{B_{m+1}(y)}| \\ &\quad + |(Du)_{B_j(x)} - (Du)_{B_j(y)}| \\ &\leq (6 + 2^{n+1}) \varepsilon \lambda + 2 |B_j(x) \setminus B_j(y)| |B_j|^{-1} \|Du\|_{L^\infty(\Omega')} \leq 2^{n+2} \varepsilon \lambda. \end{aligned}$$

This implies the uniform continuity of Du in Ω'' .

It remains to prove the inequality $|(Du)_{B_j} - (Du)_{B_{m+1}}| \leq (6 + 2^{n+1}) \varepsilon \lambda$. Let us set

$$K := \{k \in \mathbb{N} : F_k < 2\varepsilon \lambda\},$$

where F_k is defined in the proof of Proposition 7.7. If $K \cap [j + 1, m - 1] = \emptyset$, then

$$\int_{2B_k} |Du| dx = F_k - \delta^{-n} \varepsilon_0^{-1} E_k \geq 2\varepsilon \lambda - \varepsilon \lambda \geq \varepsilon \lambda \quad \text{for all } j + 1 \leq k \leq m - 1.$$

Therefore, the second assumption (7.5) of Lemma 7.3 holds and so, from the second estimate in Lemma 7.3, the desired inequality follows. Otherwise, let $j' = \max(K \cap [j + 1, m - 1])$ and $m' = \min(K \cap [j + 1, m - 1])$. Then applying Lemma 7.3 when $m = m' - 1 > j$, we conclude that

$$|(Du)_{B_j} - (Du)_{B_{m'}}| \leq 3\varepsilon_0 \varepsilon \lambda,$$

which together with the fact that $m' \in K$ implies

$$|(Du)_{B_j}| \leq |(Du)_{B_j} - (Du)_{B_{m'}}| + |(Du)_{B_{m'}}| \leq 3\varepsilon_0 \varepsilon \lambda + F_{m'} \leq (3 + 2^n) \varepsilon \lambda.$$

Similarly, applying the second estimate in Lemma 7.3 when $j = j'$, we also obtain $|(Du)_{B_{j'}} - (Du)_{B_m}| \leq \varepsilon_0 \varepsilon \lambda$ and so $|(Du)_{B_m}| \leq (3 + 2^n) \varepsilon \lambda$. Therefore, $|(Du)_{B_j} - (Du)_{B_m}| \leq |(Du)_{B_j}| + |(Du)_{B_m}| \leq (6 + 2^{n+1}) \varepsilon \lambda$, as claimed. \square

APPENDIX A. EXCESS DECAY ESTIMATES FOR AUTONOMOUS EQUATIONS

We provide more details of the proof of Lemma 5.9.

Sketch of the proof of Lemma 5.9. Without loss of generality we may assume that $1 < p \leq 2 \leq q$ and, by a smoothing argument, $\tilde{\varphi} \in C^2((0, \infty))$ so that

$$(A.1) \quad p \leq \frac{t\tilde{\varphi}''(t)}{\tilde{\varphi}'(t)} + 1 \leq q, \quad \text{for all } t > 0.$$

We further consider the weak solutions $u_\varepsilon \in W^{1,\varphi_\varepsilon}(B_r)$, $\varepsilon \in (0, 1)$, to the approximate non-degenerate equations

$$(A.2) \quad \operatorname{div} A_\varepsilon(Du_\varepsilon) = 0 \quad \text{in } B_r, \quad \text{and} \quad u_\varepsilon = \tilde{u} \quad \text{on } \partial B_r,$$

where

$$A_\varepsilon(\xi) := \tilde{A}\left(\frac{\varepsilon}{\varepsilon+|\xi|}\xi\right) \frac{|\xi|}{\varepsilon+|\xi|} \quad \text{and} \quad \varphi_\varepsilon(t) = \int_0^t \varphi'_\varepsilon(s) ds \quad \text{with} \quad \varphi'_\varepsilon(t) := \frac{\tilde{\varphi}'(\varepsilon+t)}{\varepsilon+t} t.$$

Note that φ_ε is a growth function of A_ε which satisfies the inequality (A.1). Following the argument in the proof of [52, Lemma 4.3] with modification for the elliptic setting, one can show that Du_ε converges to $D\tilde{u}$ in $L^{\tilde{\varphi}}(B_r)$. Therefore, it is enough to derive the desired estimates for the equation (A.2) with the constant C_1 independent of $\varepsilon \in (0, 1)$.

We revisit the proofs of [47, Lemma 5.1] and [46, Lemma 1]. By setting $g(t) := \varphi'_\varepsilon(t)$ and $F(t) := t^{-1}g(t) = t^{-1}\varphi'_\varepsilon(t)$, we observe from properties of growth functions and (A.1) that all the assumptions in [46, Lemma 1] hold, except for $F(t) \geq \tilde{\varepsilon}$; in [47, Lemma 5.1], all assumption hold except for $g(t) \geq \tilde{\varepsilon}t$. We point out that these two missing assumptions are equivalent and are only used in the step proving the twice differentiability of the weak solution u_ε to (A.2). See [46, p. 1205], specifically just below their equation (1.7), for the comment on the assumption $F(t) \geq \tilde{\varepsilon}$ in their equation (1.3b). In particular, the quantity $\tilde{\varepsilon} > 0$ is not used in the regularity estimate.

Since the equation (A.2) is non-degenerate and uniformly elliptic, we deduce from, e.g., [47, Theorem 1.7] that the gradient of its weak solution is locally bounded. We also refer to [11, 13, 25] for local Lipschitz regularity for very general non-uniformly elliptic problems. Once we know the local boundedness of the gradient of the weak solution u_ε , we do not need to consider large values of t , hence the above assumption can be replaced by $\varphi'_\varepsilon(t) = g(t) \geq \tilde{\varepsilon}t$ for $t \in [0, M]$, where M is any positive number and $\tilde{\varepsilon}$ depends on M . Note that this holds as follows:

$$\varphi'_\varepsilon(t) = \frac{\varphi'_0(\varepsilon+t)}{\varepsilon+t} t \geq \frac{\varphi'_0(\varepsilon)}{\varepsilon+M} t \quad \text{for } t \in [0, M].$$

Therefore, the problem (A.2) satisfies all conditions in [47, Lemma 5.1] and [46, Lemma 1]. We follow the proof of [47, Lemma 5.1]. Fix $B_\rho(y) \subset B_r$ and $B_\nu = B_\nu(y)$ with $0 < \nu \leq \rho$. Then we first have that

$$\sup_{B_{\rho/2}} \varphi_\varepsilon(|Du_\varepsilon|) \leq c \int_{B_{3\rho/2}} \varphi_\varepsilon(|Du_\varepsilon|) dx \leq c\varphi_\varepsilon \left(\int_{B_\rho} |Du_\varepsilon| dx \right),$$

Note that the second inequality is a reverse Hölder type inequality like the first inequality in Lemma 5.8, and the case $\varepsilon = 0$ can be obtained as $\varepsilon \rightarrow 0^+$. This implies the first claim of the lemma.

We next prove the excess decay estimate. For simplicity, we write $w := u_\varepsilon$, $\varepsilon > 0$, $D_i w := \frac{dw}{dx_i}$, $i = 1, 2, \dots, n$, and set

$$M(\nu) := \max_{1 \leq i \leq n} \left(\sup_{B_\nu} |D_i w| \right) \quad \text{and} \quad I(\nu) := \int_{B_\nu} |Dw - (Dw)_{B_\nu}| dx, \quad \nu \in (0, \rho].$$

Then, as in the proof [46, Lemma 1], there exist $\mu, \eta \in (0, 1)$ depending only on n, p, q and \tilde{L} such that if

$$(A.3) \quad |\{D_i w < \frac{M(\nu)}{2}\} \cap B_\nu| \leq \mu |B_\nu| \quad \text{for some } i = 1, 2, \dots, n$$

or

$$(A.4) \quad |\{D_i w > -\frac{M(\nu)}{2}\} \cap B_\nu| \leq \mu |B_\nu| \quad \text{for some } i = 1, 2, \dots, n,$$

then

$$|Dw| > \frac{3}{8} M(\nu) \quad \text{in } B_{\nu/2},$$

while if neither (A.3) nor (A.4) holds, then

$$M(\frac{\nu}{2}) \leq \eta M(\nu) \quad \text{in } B_{\nu/2}.$$

Denote $\nu_j := 2^{-j}\rho$ for $j \in \mathbb{N} \cup \{0\}$. We first assume that either (A.3) or (A.4) holds for $\nu = \nu_j$ with some j . Then $\frac{DA_\varepsilon(Dw)}{\varphi_\varepsilon''(M(\nu_j))}$ is uniformly elliptic in $B_{\nu_j/2}$ such that $\tilde{L}|\xi|^2 \geq (\frac{DA_\varepsilon(Dw)}{\varphi_\varepsilon''(M(\nu_j))}\xi) \cdot \xi \geq \tilde{L}^{-1}(\frac{3}{8})^{q-2}|\xi|^2$. By differentiation of (A.2), $h_i := D_i w - (D_i w)_{B_{\nu_j}}$ is a weak solution to

$$\operatorname{div} \left(\frac{DA_\varepsilon(Dw)}{\varphi_\varepsilon''(M(\nu_j))} Dh_k \right) = 0 \quad \text{in } B_{\nu_j/2}$$

for every $i = 1, 2, \dots, n$. Therefore by a standard oscillation decay estimate from De Giorgi's theorem for linear elliptic equations (see, for instance, [33, Theorem 2.4]), there exists $\alpha_1 \in (0, 1)$ such that, for every $\nu \in (0, \frac{\nu_j}{4})$,

$$(A.5) \quad \begin{aligned} I(\nu) &\leq \sup_{1 \leq i \leq n} \left(\operatorname{osc}_{B_\nu} h_i \right) \leq c \left(\frac{\nu}{\nu_j} \right)^{\alpha_1} \sup_{1 \leq i \leq n} \left(\operatorname{osc}_{B_{\nu_j/4}} h_i \right) \leq c \left(\frac{\nu}{\nu_j} \right)^{\alpha_1} \sup_{1 \leq i \leq n} \left(\sup_{B_{\nu_j/4}} |h_i| \right) \\ &\leq c \left(\frac{\nu}{\nu_j} \right)^{\alpha_1} \sup_{1 \leq i \leq n} \int_{B_{\nu_j/2}} |h_i| dx \leq c \left(\frac{\nu}{\nu_j} \right)^{\alpha_1} I(\nu_j). \end{aligned}$$

Let $j_1 \geq j_0$, where $j_0 \in \mathbb{N}$ is so large so that $2C_1\eta^{j_0-1} \leq \frac{1}{2}$, and suppose that neither (A.3) nor (A.4) holds for any ν_j with $j = 0, 1, \dots, j_1$. In the case $|(Dw)_{B_\rho}| \leq 2M(\nu_{j_0})$, using $M(\frac{\nu_j}{2}) \leq \eta M(\nu_j)$ and the already proved first inequality of the lemma, we then have

$$\begin{aligned} M(\nu_{j_0}) &\leq \eta^{j_0-1} M(\frac{\rho}{2}) \leq C_1 \eta^{j_0-1} |(Dw)|_{B_\rho} \leq C_1 \eta^{j_0-1} (I(\rho) + |(Dw)_{B_\rho}|) \\ &\leq C_1 \eta^{j_0-1} (I(\rho) + 2M(\nu_{j_0})). \end{aligned}$$

Since $2C_1\eta^{j_0-1} \leq \frac{1}{2}$, this yields that $M(\nu_{j_0}) \leq \frac{1}{2}I(\rho)$. Therefore, for every $j = j_0 + 1, j_0 + 2, \dots, j_1$,

$$I(\nu_j) \leq 2M(\nu_j) \leq 2\eta^{j-j_0} M(\nu_{j_0}) \leq \frac{1}{\eta^{j_0}} \eta^j I(\rho),$$

which implies that there exists $\alpha_2 \in (0, 1)$ and $c > 0$ such that

$$(A.6) \quad I(\nu) \leq c \left(\frac{\nu}{\rho} \right)^{\alpha_2} I(\rho) \quad \text{for every } \nu \in (2^{-j_1}\rho, \rho).$$

Next we consider the case $|(Dw)_{B_\rho}| > 2M(\nu_{j_0})$. Then, since $|Dw - (Dw)_{B_\rho}| \geq |(Dw)_{B_\rho}| - |Dw| \geq M(\nu_{j_0})$ in $B_{\nu_{j_0}}$, we obtain for $j = j_0 + 1, \dots, j_1$ that

$$I(\nu_j) \leq 2M(\nu_j) \leq 2\eta^{j-j_0} M(\nu_{j_0}) \leq 2\eta^{j-j_0} \int_{B_{\nu_{j_0}}} |Dw - (Dw)_{B_\rho}| dx \leq \frac{2^{1+nj_0}}{\eta^{j_0}} \eta^j I(\rho),$$

which again implies (A.6).

Combining (A.5) and (A.6), we obtain the second inequality from the claim of the lemma for $w = u_\varepsilon$ with $\alpha = \min\{\alpha_1, \alpha_2\}$. Finally passing $\varepsilon \rightarrow 0^+$ implies the second claim for \tilde{u} . \square

ACKNOWLEDGMENT

M. Lee was supported by NRF grant funded by MSIT (NRF-2022R1F1A1063032). J. Ok was supported by NRF grant funded by MSIT (NRF-2022R1C1C1004523).

REFERENCES

1. E. Acerbi and G. Mingione: *Regularity results for a class of functionals with non-standard growth*, Arch. Ration. Mech. Anal. 156 (2001), no. 2, 121–140.
2. E. Acerbi and G. Mingione: *Gradient estimates for the $p(x)$ -Laplacean system*, J. Reine Angew. Math. 584 (2005), 117–148.
3. E. Acerbi and G. Mingione: *Gradient estimates for a class of parabolic systems*, Duke Math. J. 136 (2) (2007), 285–320.
4. S. Agmon, A. Douglis and L. Nirenberg, *Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions. I*, Comm. Pure Appl. Math. 12 (1959), 623–727.
5. A. Ancona: *Elliptic operators, conormal derivatives and positive parts of functions. With an appendix by Haïm Brezis*, J. Funct. Anal. 257 (2009), no. 7, 2124–2158.
6. P. Baroni: *Gradient continuity for $p(x)$ -Laplacian systems under minimal conditions on the exponent*, J. Differential Equations 367 (2023), 415–450.
7. P. Baroni: *A new condition ensuring gradient continuity for minimizers of non-autonomous functionals with mild phase transition*, Calc. Var. Partial Differential Equations 64 (2025), no. 4, Paper No. 112.
8. P. Baroni and A. Coscia: *Gradient regularity for non-autonomous functionals with Dini or non-Dini continuous coefficients*, Electron. J. Differential Equations (2022), article 80.
9. P. Baroni, M. Colombo, G. Mingione: *Non-autonomous functionals, borderline cases and related function classes*, St. Petersburg Math. J. 27 (2016), no. 3, 347–379.
10. P. Baroni, M. Colombo, G. Mingione: *Regularity for general functionals with double phase*, Calc. Var. Partial Differential Equations 57 (2018), no. 2, Paper No. 62, 48 pp.
11. L. Beck and G. Mingione: *Lipschitz bounds and nonuniform ellipticity*, Comm. Pure Appl. Math. 73 (2020), no. 5, 944–1034.
12. P. Bella and M. Schäffner: *On the regularity of minimizers for scalar integral functionals with growth*, Anal. PDE 13 (2020), 2241–2257.
13. P. Bousquet and L. Brasco: *Global Lipschitz continuity for minima of degenerate problems*, Math. Ann. 366 (2016), no. 3–4, 1403–1450.
14. H. Brezis: *On a conjecture of J. Serrin*, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. 19 (2008), no. 4, 335–338.
15. S. Byun and J. Oh: *Global gradient estimates for the borderline case of double phase problems with BMO coefficients in nonsmooth domains*, J. Differential Equations 263 (2017), no. 2, 1643–1693.
16. S. Byun and J. Ok, *On $W^{1,q(\cdot)}$ -estimates for elliptic equations of $p(x)$ -Laplacian type*, J. Math. Pures Appl. (9) 106 (2016), no. 3, 512–545.
17. S. Byun, J. Ok and S. Ryu, *Global gradient estimates for elliptic equations of $p(x)$ -Laplacian type with BMO nonlinearity*, J. Reine Angew. Math. 715 (2016), 1–38.
18. R. Caccioppoli, *Sulle equazioni ellittiche a derivate parziali con n variabili indipendenti*, Atti Accad. Naz. Lincei, Rend. Lincei, Mat. Appl. 19 (1934) 83–89.

19. I. Chlebicka, P. Gwiazda, A. Świerczewska-Gwiazda and A. Wróblewska-Kamińska: *Partial Differential Equations in Anisotropic Musielak-Orlicz Spaces*, Springer Monographs in Mathematics, Springer, Cham, 2021.
20. M. Colombo and G. Mingione, *Regularity for double phase variational problems*, Arch. Ration. Mech. Anal. 215 (2015), no. 2, 443–496.
21. M. Colombo and G. Mingione, *Calderón-Zygmund estimates and non-uniformly elliptic operators*, J. Funct. Anal. 270 (2016), no. 4, 1416–1478.
22. A. Coscia and G. Mingione: *Hölder continuity of the gradient of $p(x)$ -harmonic mappings*, C. R. Acad. Sci. Paris Sér. I Math. 328 (1999), no. 4, 363–368.
23. G. Cupini and P. Marcellini: *Global boundedness of weak solutions to a class of nonuniformly elliptic equations*, Math. Ann., to appear.
24. C. De Filippis and G. Mingione: *A borderline case of Calderón-Zygmund estimates for nonuniformly elliptic problems*, Algebra i Analiz 31 (2019), no. 3, 82–115; reprinted in St. Petersburg Math. J. 31 (2020), no. 3, 455–477.
25. C. De Filippis and G. Mingione: *Lipschitz bounds and nonautonomous integrals*, Arch. Ration. Mech. Anal. 242 (2021), no. 2, 973–1057.
26. C. De Filippis and G. Mingione: *Regularity for double phase problems at nearly linear growth*, Arch. Ration. Mech. Anal. 247 (2023), no. 5, article 85.
27. C. De Filippis and G. Mingione: *Nonuniformly elliptic Schauder theory*, Invent. Math. 234 (2023), no. 3, 1109–1196.
28. C. De Filippis and G. Mingione: *The sharp growth rate in nonuniformly elliptic Schauder theory*, Duke Math. J., to appear. arXiv:2401.07160.
29. L. Diening, P. Harjulehto, P. Hästö and M. Růžička: *Lebesgue and Sobolev Spaces with Variable Exponents*, Lecture Notes in Mathematics, vol. 2017, Springer, Heidelberg, 2011.
30. G. Di Fazio, *L^p estimates for divergence form elliptic equations with discontinuous coefficients*, Boll. Un. Mat. Ital. A (7) 10 (1996), no. 2, 409–420.
31. H. Dong and S. Kim, *On C^1 , C^2 , and weak type- $(1, 1)$ estimates for linear elliptic operators*, Comm. Partial Differential Equations 42 (2017), no. 3, 417–435.
32. H. Dong and S. Kim, *On C^1 , C^2 , and weak type- $(1, 1)$ estimates for linear elliptic operators: Part II*, Math. Ann. 370 (2018), no. 1-2, 447–489.
33. Q. Han and F. Lin: *Elliptic partial differential equations. Second edition*, Courant Lecture Notes in Mathematics, 1. Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2011.
34. P. Harjulehto and P. Hästö: *Orlicz spaces and generalized Orlicz spaces*, Lecture Notes in Mathematics, vol. 2236, Springer, Cham, 2019.
35. P. Harjulehto, P. Hästö and M. Lee: *Hölder continuity of ω -minimizers of functionals with generalized Orlicz growth*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 22 (2021), no. 2, 549–582.
36. P. Hartman and A. Wintner, *On uniform Dini conditions in the theory of linear partial differential equations of elliptic type*, Amer. J. Math. 77 (1955), 329–354.
37. P. Hästö and J. Ok: *Maximal regularity for local minimizers of non-autonomous functionals*, J. Eur. Math. Soc. (JEMS) 24 (2022), no. 4, 1285–1334.
38. P. Hästö and J. Ok: *Regularity theory for non-autonomous partial differential equations without Uhlenbeck structure*, Arch. Ration. Mech. Anal. 245 (2022), no. 3, 1401–1436.
39. P. Hästö and J. Ok: *Regularity theory for non-autonomous problems with a priori assumptions*, Calc. Var. Partial Differential Equations 62 (2023), article 251.
40. T. Jin, V. Maz'ya and J. Van Schaftingen: *Pathological solutions to elliptic problems in divergence form with continuous coefficients*, C. R. Math. Acad. Sci. Paris 347 (2009), no. 13-14, 773–778.
41. A. Karppinen and M. Lee: *Hölder continuity of the minimizer of an obstacle problem with generalized Orlicz growth*, Int. Math. Res. Not. IMRN 2022, no. 19, 15313–15354.
42. J. Kinnunen and S. Zhou: *A local estimate for nonlinear equations with discontinuous coefficients*, Comm. Partial Differential Equations 24 (1999), no. 11-12, 2043–2068.
43. T. Kuusi and G. Mingione: *A nonlinear Stein theorem*, Calc. Var. Partial Differential Equations 51 (2014), no. 1-2, 45–86.
44. M. Lee, J. Ok and J. Pyo: *Calderón-Zygmund estimates for nonlinear equations of differential forms with BMO coefficients*, J. Differential Equations 397(2024) 262–288.

45. Y. Li: *On the C^1 regularity of solutions to divergence form elliptic systems with Dini-continuous coefficients*, Chinese Ann. Math. Ser. B 38 (2017), no. 2, 489–496.
46. G. M. Lieberman: *Boundary regularity for solutions of degenerate parabolic equations*, Nonlinear Anal. 14 (1990), no. 6, 501–524.
47. G. M. Lieberman: *The natural generalization of the natural conditions of Ladyzhenskaya and Ural'tseva for elliptic equations*, Comm. Partial Differential Equations 16 (1991), no. 2-3, 311–361.
48. G. M. Lieberman: *On the natural generalization of the natural conditions of Ladyzhenskaya and Ural'tseva*, Banach Center Publ., 27, Part 1, 2 Polish Academy of Sciences, Institute of Mathematics, Warsaw, 1992, 295–308.
49. J. J. Manfredi: *Regularity for minima of functionals with p -growth*, J. Differential Equations 76 (1988), no. 2, 203–212.
50. T. Ohno and T. Shimomura: *Boundedness of maximal operators and Sobolev inequalities on Musielak-Orlicz spaces over unbounded metric measure spaces*, Bul. Sci. Math. 199 (2025), article 103546.
51. J. Ok, *Gradient continuity for $p(\cdot)$ -Laplace systems*, Nonlinear Anal. 141 (2016), 139–166.
52. J. Ok, G. Scilla and B. Stroffolini: *Regularity theory for parabolic systems with Uhlenbeck structure*, J. Math. Pures Appl. 182 (2024), 116–163.
53. J. Schauder: *Über lineare elliptische Differentialgleichungen zweiter Ordnung*, Math. Z. 38 (1934), no. 1, 257–282.
54. J. Schauder: *Numerische Abschätzungen in elliptischen linearen Differentialgleichungen*, Stud. Math. 5 (1937) 34–42.
55. S. Spanne: *Some function spaces defined using the mean oscillation over cubes*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3) 19 (1965), 593–608.
56. V. D. Stepanov: *The weighted Hardy's inequality for nonincreasing functions*, Trans. Amer. Math. Soc. 338 (1993), no. 1, 173–186.
57. F. Weisz, G. Xie and D. Yang: *Boundedness of dyadic maximal operators on Musielak-Orlicz type spaces and its applications*, J. Geom. Anal. 35 (2025), article 81.

DEPARTMENT OF MATHEMATICS AND STATISTICS, FI-00014 UNIVERSITY OF HELSINKI, FINLAND
 AND DEPARTMENT OF MATHEMATICS AND STATISTICS, FI-20014 UNIVERSITY OF TURKU, FINLAND
Email address: peter.hasto@helsinki.fi

DEPARTMENT OF MATHEMATICS AND INSTITUTE OF MATHEMATICAL SCIENCE, PUSAN NATIONAL
 UNIVERSITY, BUSAN 46241, REPUBLIC OF KOREA
Email address: mikyounglee@pusan.ac.kr

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 04107, REPUBLIC OF KOREA
Email address: jihoonok@sogang.ac.kr