Tilted dipolar bosons in the quasi-2D regime: from liquid stripes to droplets.

J. Sánchez-Baena^{1,*}

¹Departament de Física, Universitat Politècnica de Catalunya, Campus Nord B4-B5, 08034 Barcelona, Spain

(Dated: April 4, 2025)

We characterize a system of tilted dipoles in a quasi two-dimensional (flattened) geometry and in the thermodynamic limit. We consider a finite trapping in the z-axis achievable in current experiments. We compute the phase diagram of the system at its equilibrium density for high tilting angles, where it becomes self-bound, and a striped liquid state emerges. To characterize the system, we perform a variational calculation, which is benchmarked with the solution of the extended Gross-Pitaevskii equation. We connect the phenomenology in the thermodynamic limit to the physics of the finite-size system, provide parameters for the realization of potentially supersolid striped states and study the critical number for dipolar droplet formation. Our results are helpful to guide potential experiments in the study of dipolar atoms in quasi two-dimensional geometries in the dipole-dominated regime.

I. INTRODUCTION

Ultracold quantum gases with dipolar interactions have risen to prominence in the last decade as a platform to study a plethora of exotic phenomena in a controlled environment. The interplay between the anisotropy and long-range behaviour of the dipole-dipole interaction (DDI) is responsible for the emergence of dipolar droplets [1-5], which constitute an ultra-dilute liquid state, supersolids [6–14], which combine two apparently mutually exclusive properties like global phase coherence and spatial periodicity, or anomalous thermal behaviour [14–17], in the form of the promotion of supersolidity by increasing the temperature. Recently, many efforts are being directed towards the realization of ultracold atom experiments in a quasi two-dimensional geometry, in a regime where physical phenomena exclusive to the two-dimensional realm can be accessed. This includes the measurement of the Berezinskii-Kosterlitz-Thouless (BKT) transition for a system of ⁸⁷Rb atoms [18], the realization of a Bose glass in a quasicrystalline optical lattice [19], or the direct probing of a 1D-2D crossover for caesium atoms [20], to name a few. In the case of dipolar systems, quasi two-dimensional systems have the added interest of studying the interplay between the unique properties of the DDI and a dimensional constrain. Due to the anisotropy of the DDI, limiting the motion of particles in one direction introduces a new variable with respect to the three-dimensional case: the polarization direction of the dipoles. Moreover, quantum fluctuations, which play a crucial role in dipolar systems by arresting the mean-field collapse [21–23] depend significantly on dimensionality [24, 25]. Because of this, experimental activity targeting these systems has emerged recently. In particular, a bilayer of Dy atoms with a layer width on the sub-50 nm scale has been realized [26], and the BKT transition has been experimentally accessed for a dipolar system, where the wave function of atoms along the

tightly trapped direction approximately corresponds that of the harmonic oscillator ground state [27]. In light of this experimental activity, it is necessary to provide theoretical calculations pointing at the relevant parameters for the observation of novel physical phenomena exclusive to the quasi two-dimensional regime.

Several theoretical works have been produced along the years with the goal of characterizing dipolar systems in flattened geometries under different specific conditions. These include the study of the ground state, collapse instability and excitations of the gas phase at the mean field level [28–31], including finite temperature [32], and the characterization of the stripe phase for bosons [30, 33, 34], fermions [35, 36] and mixtures [37], where a spin-stripe phase emerges. The system has also been studied in the 3D limit [38–41] and in the presence of a two-dimensional lattice, both at zero and finite temperature [42]. In particular, the results in the 3D limit show the importance to modify the tilting angle of the dipoles to reduce the necessary density for the experimental realization of the stripe phase, which is excessively large in the case of dipoles perpendicular to the plane of motion, and thus induces important three-body losses [39]. Through this approach, striped arrangements of tilted dipoles have been observed in experiments [43], with the number of stripes being controllable with the trap aspect ratio. In the strict two-dimensional case, quantum Monte Carlo methods have been employed to study the phases of the system for the case of a single layer [44–48], a bilayer [49] and a multilayer [50] configuration.

Recently, a theoretical work brought the attention into the liquid properties of the system of tilted dipoles in the fully two-dimensional case [51], where the emergence of an exotic, striped liquid phase in the thermodynamic limit is shown. Nevertheless, the strict two-dimensional limit, which is achieved when $l \ll a$, with l the oscillator length in the z-axis and a the scattering length, lies away from the state of the art experiments, which typically fulfill l > a. This has important consequences on relating the phenomenology of the system with the s-wave scattering length, which is of crucial importance in experiments, while also impacting greatly the critical number

^{*} juan.sanchez.baena@upc.edu



FIG. 1. Schematic representation of the system of dipoles in a flattened geometry. Left: isodensity surfaces in the homogeneous liquid phase. The different colors correspond to surfaces where $n = 0.5n_{\text{peak}}$ (red), $n = 0.8n_{\text{peak}}$ (light blue) and $n = n_{\text{peak}}$ (dark blue), with n the 3D density and n_{peak} the peak density. Center: isodensity surface ($n = 0.5n_{\text{peak}}$ (dark blue)) of the system in the liquid stripe phase. Right: graphical representation of a polarized dipole (thick black arrow) with tilting angle α .

for the formation of droplets, as we shall illustrate below.

In this work, we perform a variational calculation to obtain the phase diagram of an infinite system of tilted dipoles in a flattened geometry at the equilibrium density. In specific cases, we benchmark our variational results with the solution of the extended Gross-Pitaevskii equation (eGPE), which we also employ to perform calculations of a finite size system under a box trap. This paper is organized as follows. In Sec. II, we present the variational theory and the eGPE. In Sec. III, we present our results for both the thermodynamic limit (Sec. III A) and the finite size system (Sec. III B) while in Sec. IV we summarize the main conclusions of our work.

II. THEORY

We study a system governed by the energy functional

$$E[\psi] = \int d\mathbf{r} \psi^*(\mathbf{r}) \left[-\frac{\hbar^2 \nabla^2}{2m} + U(\mathbf{r}) \right] \psi(\mathbf{r}) + \frac{1}{2} \int \int d\mathbf{r} d\mathbf{r}' |\psi(\mathbf{r}')|^2 V(\mathbf{r} - \mathbf{r}') |\psi(\mathbf{r}')|^2 + \int d\mathbf{r} \epsilon_{\text{BMF}} \left(|\psi(\mathbf{r})|^2 \right)$$
(1)

where $U(\mathbf{r}) = \frac{1}{2}m\omega^2 z^2$ for the infinite system and $U(\mathbf{r}) = \frac{1}{2}m\omega^2 z^2 + U_{\text{box}}(x,y)$ for the finite one with $U_{\text{box}}(x,y)$ a box trap. $V(\mathbf{r} - \mathbf{r}')$ is the contact plus dipole-dipole interaction, i.e.

$$V(\mathbf{r} - \mathbf{r}') = \frac{4\pi\hbar^2 a}{m} \delta\left(\mathbf{r} - \mathbf{r}'\right) + \frac{C_{\rm dd}}{4\pi} \frac{\left(1 - 3\cos^2\theta_{\alpha}\right)}{\left|\mathbf{r} - \mathbf{r}'\right|^3} ,$$
(2)

with a the s-wave scattering length and θ_{α} the angle between the vector $\mathbf{r} - \mathbf{r}'$ and the polarization axis of the dipoles. In the calculations, we change this direction within the x-z plane, and define α as the angle between the polarization direction and the z-axis (see Fig. 1). Here $C_{\rm dd} = \mu_0 \mu^2$ for magnetic dipoles. We restrict ourselves to the regime where the collisions between particles can be treated as a three-dimensional process, meaning that l > a with $l = \sqrt{\frac{\hbar}{m\omega}}$, which is necessary for the model interaction of Eq. 2 to be valid. We define characteristic energy and length scales given by $r_0 = 12\pi a_{\rm dd}$ and $E_0 = \hbar^2/(mr_0^2)$ and present our results in these units.

The quantity $\epsilon_{\text{BMF}} \left(n = |\psi(\mathbf{r})|^2 \right)$ is an energy functional accounting for beyond mean-field (BMF) effects. In order to compute it, we follow Ref. [25], where the BMF energy density is obtained for an homogeneous system with periodic boundary conditions on the z-axis, thus accounting approximately for the discretized excitations due to finite trapping effects. Remarkably, the scaling with the density depends on its value, since lower densities only populate the lowest excitation along the z-axis (meaning that the system is in the quasi two-dimensional limit) while higher densities yield the 3D result. In practice, if we denote by n_{lim} the limiting value of the density where the scaling changes, the BMF energy density can be fitted to [25]

$$\epsilon_{\rm BMF}^{\rm low}(n) = C_1 n^2 + C_2 n^3 + C_3 n^{5/2} \qquad n < n_{\rm lim} \qquad (3)$$

$$\epsilon_{\rm BMF}^{\rm high}(n) = C_4 n^{5/2} \qquad n > n_{\rm lim} \qquad (4)$$

The term $C_3 n^{5/2}$ is included in Eq. 3 with respect to the result of Ref. [25] to improve the accuracy of the fit, while Eq. 4 shows the well-known density scaling in the 3D limit [21, 22]. In the calculation for $\epsilon_{\rm BMF}$, we set the value of the length of the z-axis to $L_z = 4l$, with lthe harmonic oscillator length of the trapping, and find $n_{\rm lim}$ numerically (see the Appendix A). This density is an important parameter, as it helps to determine whether the system is effectively two or three-dimensional. When evaluating Eq. 1, we have to choose one of the two density functionals from Eqs. 3 and 4. The choice is determined by the peak density of the converged ground state. We employ Eq. 3 if $n_{\rm peak} \leq n_{\rm lim}$ and Eq. 4 if $n_{\rm peak} \gg n_{\rm lim}$. It can be seen numerically that the energy functional of Eq. 4 lays quantitatively very close (less than a few % away) to the standard LHY energy density correction in three dimensions given in Refs. [21, 22].

At high enough values of the tilting and for a tight confinement along the z-axis, the system is expected to host either an unmodulated state, or a striped state [34, 51]. In order to find the lowest energy state, we perform a variational minimization of Eq. 1 for two different ansatz. These are the senoidal ansatz

$$\psi_{\rm S}(\mathbf{r}) = \sqrt{\frac{n_{\rm 2D}}{\sqrt{\pi}\sigma_z}} \sqrt{1 + A\cos(ky)} e^{-\frac{1}{2}\left(\frac{z}{\sigma_z}\right)^2} , \quad (5)$$

with A, k and σ_z variational parameters, and the gaussian ansatz

$$\psi_{\rm G}(\mathbf{r}) = \sum_{j=-\infty}^{\infty} \int d\mathbf{r}' \sqrt{\frac{n_{\rm 2D} L_y}{\pi \sigma_z \sigma_y}} e^{-\frac{1}{2} \left[\left(\frac{y'}{\sigma_y} \right)^2 + \left(\frac{z'}{\sigma_z} \right)^2 \right]} \\ \times \delta(y + jL_y - y') , \qquad (6)$$

where the variational parameters are L_y , σ_y and σ_z . In both expressions, L_y is the length of the unit cell in the y-axis ($L_y = 2\pi/k$ for the senoidal ansatz), $n_{2\rm D} = \frac{N}{L_x L_y}$ is the 2D density and the normalization ensures that $N = \int_{-L_x/2}^{L_x/2} \int_{-L_y/2}^{L_y/2} \int_{-\infty}^{\infty} d\mathbf{r} |\psi_{\mathrm{S,G}}(\mathbf{r})|^2$. The senoidal ansatz converges to an unmodulated state in the limit $A \to 0$, meaning that with these wave functions we can account for an unmodulated state, a phase coherent, striped state and a highly incoherent striped state.

We insert Eqs. 5 and 6 into Eq. 1 to obtain expressions for the energy of the system. In a general way, we split the contributions of the energy into

$$E_{\rm S,G} = E_1 + E_{\rm V} + E_{\rm BMF} \tag{7}$$

$$E_{1} = \int d\mathbf{r}\psi^{*}(\mathbf{r}) \left[-\frac{\hbar^{2}\nabla^{2}}{2m} + U(\mathbf{r}) \right] \psi_{\mathrm{S,G}}(\mathbf{r})$$
(8)

$$E_{\rm V} = \frac{1}{2} \int \int d\mathbf{r} d\mathbf{r}' |\psi_{\rm S,G}(\mathbf{r}')|^2 V(\mathbf{r} - \mathbf{r}') |\psi_{\rm S,G}(\mathbf{r}')|^2$$
(9)

$$E_{\rm BMF} = \int d\mathbf{r} \epsilon_{\rm BMF} \left(\left| \psi_{\rm S,G}(\mathbf{r}) \right|^2 \right) \ . \tag{10}$$

where the subindex "S, G" of Eq. 7 refers to the senoidal and gaussian ansatz, respectively. The contributions to the energy per particle E/N for the senoidal ansatz are given by

$$E_1/N = \frac{\hbar^2 A^2 k^2}{8m} \frac{1}{2\pi} \int_{-\pi}^{\pi} d\xi \frac{\sin^2(\xi)}{1 + A\cos\xi} + \frac{\hbar^2}{4m\sigma_z^2} + \frac{m\omega^2 \sigma_z^2}{4}$$
(11)

$$E_{\rm V}/N = \frac{n_{\rm 2D}}{2} \left(\tilde{V}(0,0) + \frac{A^2}{2} \tilde{V}(0,k) \right)$$
(12)

$$E_{\rm BMF}^{\rm high}/N = C_4 n_{\rm 2D}^{3/2} \sqrt{\frac{2}{5\sigma_z^3 \pi^{3/2}} \frac{1}{2\pi} \int_{-\pi}^{\pi} d\xi \left(1 + A\cos\xi\right)^{5/2}}$$
(13)

$$E_{\rm BMF}^{\rm low}/N = C_1 \frac{1}{\sqrt{2\pi}\sigma_z} n_{2\rm D} \frac{1}{2\pi} \int_{-\pi}^{\pi} d\xi \left(1 + A\cos\xi\right)^2 + C_2 \frac{1}{\sqrt{3\pi}\sigma_z^2} n_{2\rm D}^2 \frac{1}{2\pi} \int_{-\pi}^{\pi} d\xi \left(1 + A\cos\xi\right)^3 + C_3 n_{2\rm D}^{3/2} \sqrt{\frac{2}{5\sigma_z^3 \pi^{3/2}}} \frac{1}{2\pi} \int_{-\pi}^{\pi} d\xi \left(1 + A\cos\xi\right)^{5/2} , \qquad (14)$$

where the term $\tilde{V}(k_x, k_y)$ of Eq. 12 is [33]

$$\tilde{V}(k_x, k_y) = \frac{2\sqrt{2\pi}\hbar^2 a}{m\sigma_z} - \frac{2\sqrt{2\pi}\hbar^2 a_{\rm dd}}{m\sigma_z} + \frac{6\hbar^2 a_{\rm dd}}{m} \int_{-\infty}^{\infty} dk_z \frac{k_x^2}{k^2} \exp\left(-\frac{k_z^2 \sigma_z^2}{2}\right) \sin^2 \alpha + \frac{6\hbar^2 a_{\rm dd}}{m} \int_{-\infty}^{\infty} dk_z \frac{k_z^2}{k^2} \exp\left(-\frac{k_z^2 \sigma_z^2}{2}\right) \cos^2 \alpha .$$
(15)

Here, $a_{\rm dd} = mC_{\rm dd}/(12\pi\hbar^2)$ is the dipole length. In much the same way, the contributions of E/N for the gaussian ansatz are

$$E_1/N = \frac{\hbar^2}{2m} \left(\frac{1}{2\sigma_y^2} + \frac{1}{2\sigma_z^2} \right) + \frac{m\omega^2 \sigma_z^2}{4}$$
(16)

$$E_{\rm V}/N = \frac{n_{\rm 2D}}{2} \left(\frac{L_y}{\sqrt{\pi}\sigma_y}\right)^2 \left(A_0^2 \tilde{V}(0,0) + \sum_{j=1}^{\infty} \frac{A_j^2}{2} \tilde{V}(0,jk)\right)$$
(17)

$$E_{\rm BMF}^{\rm high}/N = \frac{2C_4}{5} \left(\frac{n_{\rm 2D}L_y}{\pi\sigma_y\sigma_z}\right)^{3/2} \tag{18}$$

$$E_{\rm BMF}^{\rm low}/N = \frac{C_1}{2} \frac{n_{\rm 2D}L_y}{\pi\sigma_y\sigma_z} + \frac{C_2}{3} \left(\frac{n_{\rm 2D}L_y}{\pi\sigma_y\sigma_z}\right)^2 + \frac{2C_3}{5} \left(\frac{n_{\rm 2D}L_y}{\pi\sigma_y\sigma_z}\right)^{3/2} .$$
(19)

where we have assumed that $L_y \gg \sigma_y$. The coefficients A_j from Eq. 17 are obtained from the Fourier series of the density and given by

$$A_j = f_j \frac{\sqrt{\pi}\sigma_y}{L_y} \exp\left(-\frac{(2\pi j\sigma_y)^2}{4L_y^2}\right) , \qquad (20)$$

where $f_0 = 1$ and $f_j = 2$ for j > 0. The variational minimization of the energy per particle for each ansatz is performed numerically via a Simulated Annealing algorithm.

Performing a functional minimization of Eq. 1 with respect to $\psi^*(\mathbf{r})$ with the constrain $N = \int d\mathbf{r} |\psi(\mathbf{r})|^2$ we obtain the zero temperature, extended Gross-Pitaevskii equation (eGPE), which is given by

$$\mu\psi(\mathbf{r}) = \left[-\frac{\hbar^2 \nabla^2}{2m} + U(\mathbf{r}) + \int d\mathbf{r}' V(\mathbf{r} - \mathbf{r}') |\psi(\mathbf{r}')|^2 + \frac{\partial\epsilon_{\rm BMF}}{\partial n} \Big|_{n = |\psi(\mathbf{r})|^2} \right] \psi(\mathbf{r}) .$$
(21)

We use the eGPE to benchmark our variational calculations in the thermodynamic limit and to study the finite size system under a box trap in the x-y plane.

III. RESULTS

A. Infinite system

We start by discussing the results for the infinite system. In this case, the trapping potential in Eq. 1 is set to $U(\mathbf{r}) = \frac{1}{2}m\omega^2 z^2$, while the system extends indefinitely in the *x-y* plane. Unless specified otherwise, we set the trapping strength along the *z*-axis to $\omega = 2\pi \times 906$ Hz in all calculations, which in our units corresponds to $\hbar\omega/E_0 = 1$. This trapping strength is close to the one employed in the recent experiment [27], where the BKT transition was probed for a dipolar system. For a given value of the *s*-wave scattering length *a*, tilting α and 2D density n_{2D} , we can minimize the energy per particle E/N for each of the two ansatz of Eqs. 5 and 6. In particular, we are interested in the state of the system at the



FIG. 2. Energy per particle for the senoidal (top, see Eq. 5) and gaussian (bottom, see Eq. 6) ansatz, respectively. The value of the scattering length is set to $a/a_{\rm dd} = 0.7$. We have subtracted the energy of the non-interacting system, $E_{\rm NI}/(NE_0) = \hbar\omega/(2E_0)$.

equilibrium 2D density $n_{2D}^{\text{eq.}}$, i.e. the 2D density at which the energy per particle is minimized. We illustrate the different qualitative behaviour of the energy per particle as a function of n_{2D} for the two ansatz in Fig. 2 for two different sets of parameters $\{a/a_{dd}, \alpha\} = \{0.7, 1.23\}$ and $\{0.7, 1.35\}$. From the figure, we see that E_G/N increases monotonously with the density in both cases and the minimum value is achieved for $n_{2D} \to 0$, with N, $L_x L_y \to \infty$. This is because the gaussian ansatz describes an array of incoherent stripes, spaced by a distance L_y . Since the inter-stripe energy is globally repulsive, the minimum energy is achieved for infinitely spaced stripes, charac-



FIG. 3. Phase diagram of the infinite dipolar system in a flattened geometry at the equilibrium 2D density.

terized by a given ratio of particles per length along the x-axis, N/L_x . Moreover, in the range of values of L_y where the inter-stripe interactions are neglectable, $E_{\rm G}/N$ stays essentially unchanged for $n_{2D}L_y = \text{constant}$. The gaussian ansatz describes a striped system that can correspond to either a gas $(E_{\rm G}/N > E_{\rm NI}/N)$ or a liquid $(E_{\rm G}/N < E_{\rm NI}/N)$, where $E_{\rm NI}/(NE_0) = \hbar\omega/(2E_0)$ is the energy per particle of the non-interacting system. This is because if $E_{\rm G}/N > E_{\rm NI}/N$, a finite system expands indefinitely to reduce its energy while if $E_{\rm G}/N < E_{\rm NI}/N$, the system prefers to bunch into stripes to do so. Moving on to the results for the senoidal ansatz, $E_{\rm S}/N$ shows two different qualitative trends for the parameters chosen: either it increases monotonously with n_{2D} or it shows a minimum at a given 2D density. As known from thermodynamics, the former trend corresponds to a gas (expanding) state, while the latter corresponds to a liquid.

For each set of parameters $\{a, \alpha\}$ we perform the variational energy minimization for the senoidal and gaussian ansatz and look for the lowest energy state at the equilibrium 2D density. From these calculations, we obtain the phase diagram of the system, which is shown in Fig. 3. The system shows three distinct phases: a gas phase, which is characterized by a vanishing $n_{2D}^{eq.}$ and absence of order, an unmodulated liquid phase, which has a finite equilibrium 2D density at which $E/N < \hbar\omega/2$ with absent order and finally a liquid, modulated phase, where $E/N < \hbar \omega/2$ for $n_{2D}^{\text{eq.}}$ and the system shows modulations. We observe there is a maximum value of the scattering length for the emergence of modulations at the equilibrium density, which lies around $a/a_{\rm dd} \simeq 0.76$ for our parameters of choice, or $a \simeq 99.4a_0$ for ¹⁶⁴Dy atoms. This phenomenology is consistent to the one observed in the 2D case [51]. However, the comparison between the results of Fig. 3 with the phase diagram provided in Ref. [51] shows relevant qualitative differences drawn from the finite trapping strength along the z-axis. While in the strict 2D case the system can transition directly from the gas phase to the modulated liquid, we find that the phase boundary of the gas phase is always



FIG. 4. 2D equilibrium density (top) and contrast of the density $(C = \frac{n_{\max} - n_{\min}}{n_{\max} + n_{\min}})$ (center, bottom) as a function of the tilting angle. The scattering length is set to $a/a_{\rm dd} = 0.7$ in the top and bottom figures while it is set to $a/a_{\rm dd} = 0.6$ in the center figure.

connected to the unmodulated liquid for our parameters of choice. The difference stems from the dimensional nature of the modulated phase, since in all cases it has a 3D peak density significantly larger than the threshold density $n_{\rm lim}r_0^3 \simeq 0.06$ at which excitations along the z axis get significantly populated (see Sec. II). This means that this state is three-dimensional.

Another major qualitative difference between our phase diagram and that of Ref. [51] is the fact that the liquid phases in the fully 2D case appear only when the system hosts a two-body bound state. In contrast, we find the liquid states with a model potential that does not host any two-body bound state, meaning that the self-bound nature of the system arises due to many-body effects. This has important consequences in the formation of self-bound droplet states in the finite size system.



FIG. 5. Column densities $n(y) = \int dx dz |\psi(\mathbf{r})|^2$ (top) and $n(z) = \int dx dy |\psi(\mathbf{r})|^2$ (bottom) from the ground state solution of the eGPE (Eq. 21) and the gaussian ansatz at the variational energy minimum (Eq. 6). The scattering length and tilting angle are $a/a_{\rm dd} = 0.7$ and $\alpha = 1.325$, respectively, while the 2D density is set to $n_{\rm 2D}r_0^2 = 20$.

While in the 2D case of Ref. [51] any many-body state in the liquid phase supports the formation of a self-bound droplet, in our case there is a varying critical number of particles $N_{\rm crit} > 2$ for droplet formation. We study the formation of dipolar clusters in the finite system in Sec. III B.

We have analyzed the transitions between the different regions of the phase diagram of Fig. 3 for two values of the scattering length, $a/a_{\rm dd} = 0.6$ and 0.7. To do so, we have looked at the equilibrium 2D density $n_{2D}^{\text{eq.}}$ in the gasliquid transition and at the contrast of the 3D density, $C = \frac{n_{\max} - n_{\min}}{n_{\max} + n_{\min}}$ in the transition between the modulated and unmodulated liquid regions. Results are shown in Fig. 4. For both values of the scattering length, the equilibrium density 2D density changes continuously from $n_{2D}^{\text{eq.}} = 0$ in the gas phase to $n_{2D}^{\text{eq.}} \neq 0$ in the liquid phase, meaning that the liquid state emerges continuously. For this reason, we only report the case with $a/a_{\rm dd} = 0.7$. In regards to the transition in and out of the striped region, the contrast can change either abruptly $(a/a_{dd} = 0.6)$ or continuously $(a/a_{dd} = 0.7)$ when entering the striped region by increasing the tilting angle, as shown in the figure. This is because, for $a/a_{dd} = 0.6$, the gaussian ansatz becomes energetically favorable at the equilibrium density over an unmodulated state described by the senoidal ansatz. In contrast, for $a/a_{dd} = 0.7$, the senoidal ansatz, for which modulations arise continuously, is preferable to the gaussian one over a larger range of densities. Ex-



FIG. 6. Energy per particle for the senoidal (see Eq. 5) and gaussian (see Eq. 6) ansatz as a function of the 2D density for $\alpha = 1.35$ (top, striped liquid region) and $\alpha = 1.42$ (bottom, unmodulated liquid region). The value of the scattering length is set to $a/a_{\rm dd} = 0.7$. The red squares indicate the values at which the senoidal ansatz shows a density modulation in the variational energy minimum. We have subtracted the energy of the non-interacting system, $E_{\rm NI}/(NE_0) = \hbar\omega/(2E_0)$.

iting the modulated region by increasing the tilting at constant scattering length causes to contrast to abruptly vanish for both cases, and hence it is not shown in the figure.

In order to evaluate the effect of increasing the trapping strength in our results, we perform a variational calculation for $a/a_{dd} = 0.7$ using a trapping strength of $\omega' = 2\pi \times 90600$ Hz, with an associated harmonic oscillator length of l = 26 nm, around the order of magnitude of that employed in the experiment of Ref. [26]. Such an increase of the harmonic oscillator frequency has an important effect on n_{lim} , which increases by almost two orders of magnitude $(n_{\rm lim}r_0^2 \simeq 3)$. The results indicate that for the trapping strength ω' , the system only hosts the gas and unmodulated liquid phases at the equilibrium density, with the transition happening at $\alpha = 1.285$. This corresponds to the behaviour of the phase diagram at larger values of $a/a_{\rm dd}$ for ω , which implies that, effectively, the modulated liquid phase is pushed to lower values of the scattering length.

In order to benchmark our variational calculations, we run simulations to solve the eGPE of Eq. 21 and compare the outcome with the variational results in two cases. Firstly, we set $a/a_{\rm dd} = 0.6$, $\alpha = 1.325$, $n_{\rm 2D}r_0^2 = 20$



FIG. 7. Integrated density $n(x, y) = \int dz |\psi(\mathbf{r})|^2$ for the ground state of the flattened dipolar system in a box trap. The tilting angle and the lengths of the box trap are $\alpha = 1.35$, $L_x/r_0 = 41.7$, $L_y/r_0 = 61.5$ (top) and $\alpha = 1.42$, $L_x/r_0 = 26.3$, $L_y/r_0 = 96.3$ (bottom) while, in both cases, $a/a_{\rm dd} = 0.7$, N = 30000.



FIG. 8. Maximum 2D density at which a striped state appears as the variational energy minimum in the infinite system vs tilting angle. We set $a/a_{\rm dd} = 0.7$. The values of the tilting are chosen within the striped liquid region of Fig. 2.

and compute the ground state wave function and energy of the system. The eGPE yields a striped state with $E/N = 4.003 \times 10^{-2} E_0$ and a stripe spacing of $L_y/r_0 =$ 19 while the variational calculation yields a striped gaussian state with $E/N = 5.627 \times 10^{-2} E_0$ with $L_y/r_0 = 18.5$, providing reasonable agreement. We show a comparison of the column densities $n(y) = \int dx dz |\psi(\mathbf{r})|^2$ and $n(z) = \int dx dy |\psi(\mathbf{r})|^2$ obtained with both methods in Fig. 5. Secondly, we perform eGPE simulations for $a/a_{\rm dd} = 0.6$, $\alpha = 1.185$, 1.22 at $n_{\rm 2D}r_0^2 = 0.56$ and find results compatible with the variational calculations of Fig. 3, mainly an unmodulated, gas state at $\alpha = 1.185$ and a modulated, liquid state at $\alpha = 1.22$.

B. Finite system

We now turn our attention to the finite system, where we consider a trapping potential of the form $U(\mathbf{r}) = \frac{1}{2}m\omega^2 z^2 + U_{\text{box}}(x, y)$. As in the infinite case, we set the trapping strength along the z-axis to $\omega = 2\pi \times 906$ Hz in all calculations. The box trap potential is characterized by parameters L_x and L_y and is given by

$$U_{\text{box}}(x,y) = \begin{cases} 0 & \text{if } |x| < \frac{L_x}{2} \text{ and } |y| < \frac{L_y}{2} \\ \infty & \text{if } |x| > \frac{L_x}{2} \text{ and } |y| > \frac{L_y}{2} \end{cases}$$
(22)

In order to obtain the ground state of the system, we solve the eGPE given in Eq. 21 by simulating imaginary time evolution. All the states computed for the finite system fulfill $n_{\text{peak}} \gg n_{\text{lim}}$, meaning that we employ the beyond mean-field energy functional of Eq. 4. We first focus on the realization of the striped liquid phase in experimental conditions. In this sense, the results of Fig. 3 are particularly useful. This is illustrated in Fig. 6, where we show the behaviour of E/N for the infinite system for two points in the liquid phase: $\{a, \alpha\} = \{0.7, 1.35\} \pmod{100}$ ulated liquid) and $\{a, \alpha\} = \{0.7, 1.42\}$ (unmodulated liquid). Notice that in the unmodulated liquid region, the system's energetically preferable state is an unmodulated liquid for $n_{2D} \ge n_{2D}^{\text{eq.}}$. This means that, when experimentally realizing the finite system under the box trapping, the system is unmodulated for $n_{2D} \ge n_{2D}^{eq.}$ while it converges to an unmodulated state also for $n_{2D} < n_{2D}^{eq.}$, since particles minimize their energy by bunching and producing a state with 2D density as close to n_{2D}^{eq} as possible. On the other hand, this does not happen in the modulated liquid region. To illustrate this phenomenology, we perform simulations at the two aforementioned points, $\{a, \alpha\} = \{0.7, 1.35\}$ and $\{0.7, 1.42\}$ setting N = 30000and $\{L_x/r_0, L_y/r_0\} = \{41.7, 61.5\}$ in the former case and $\{L_x/r_0, L_y/r_0\} = \{26.3, 96.3\}$ in the latter. The lengths of the box trap are chosen to fix a 2D density where the variational calculation in the infinite case leads to density modulations, with L_y chosen to be commensurate with their spatial period. We show the results in Fig. 7. As can be seen from the results, we see the emergence of a striped state for $\{a, \alpha\} = \{0.7, 1.35\}$ but not so for $\{0.7, 1.42\}$, where the atoms bunch into an unmodulated state with higher 2D density to minimize their energy. Remarkably, in this case, the atoms do not occupy the full volume of the trap. In view of these results, it is thus important to characterize the maximum 2D density at which stripes emerge in the infinite system within the modulated liquid region, since setting an experiment with



FIG. 9. Potentially supersolid states: integrated densities $n(x, y) = \int dz |\psi(\mathbf{r})|^2$ (top) and column densities $n(y) = \int dx dz |\psi(\mathbf{r})|^2$ (bottom) for the ground state of the flattened dipolar system in a box trap. The parameters are N = 10000, $\alpha = 1.3$, $L_x/r_0 = 45.6$, $L_y/r_0 = 45.6$ (left), N = 28000, $\alpha = 1.32$, $L_x/r_0 = 54.3$, $L_y/r_0 = 54.3$ (center) and N = 40000, $\alpha = 1.35$, $L_x/r_0 = 41.1$, $L_y/r_0 = 61.5$ (right). In all cases $a/a_{\rm dd} = 0.7$.

higher 2D density would destroy modulations. Thus, we report this quantity in Fig. 8 for $a/a_{dd} = 0.7$.

Next, we move on to the discussion on supersolidity. In qualitative terms, a modulated state should feature a cloud of atoms surrounding the density modulations in order for global phase coherence to exist, and thus for supersolidity to emerge. We show in Fig. 9 the integrated density n(x, y) of three examples of potentially supersolid states, which involve parameters currently achievable in experiments. One concern for the realization of such states is the peak density, as a high value can induce excessive three-body losses. In all the states reported in Fig. 9, the peak 3D density is of the order $n_{\text{peak}}r_0^3 \sim \mathcal{O}(10)$, which means $n_{\text{peak}} \sim 5 \times 10^{14} \text{ cm}^{-3}$. This value lays within experimental reach [52], such that three-body losses do not translate into an excessively short life-time [39]. These results may thus guide potential future experiments in the realization of supersolid stripes, which were not observed in the experiment of Ref. [43].

Finally, we address the formation of dipolar droplets. The appearance of a liquid phase in the thermodynamic limit (see Fig. 3) is indicative of the emergence of selfbound liquid droplets in the finite size system. In the absence of a trap in the x-y plane, dipoles minimize their energy by forming a droplet elongated along the x-axis (which can be understood as a single, finite stripe) [43]. However, and distinctly to what happens in the strict 2D regime, there exist a minimum number of particles $N_{\rm crit.} > 2$ necessary for the emergence of a many-body bound state. We refer to this quantity as the critical number $N_{\rm crit.}$. We show in Fig. 10 $N_{\rm crit.}$ as a function of the tilting angle α for $a/a_{\rm dd} = 0.7$, as well as a sample of the column density n(x, y). For technical reasons, in these calculations we have replaced the box trap defined in Eq. 22 by a weak harmonic trap in the *x-y* plane $U(x,y) = \frac{1}{2}m(\omega_x^2 x^2 + \omega_y^2 y^2)$. The harmonic frequencies are $\hbar\omega_x/E_0 = 2.20 \times 10^{-6}$ and $\hbar\omega_y/E_0 =$ 8.83×10^{-6} for N > 1750 and $\hbar \omega_x / E_0 = 3.91 \times 10^{-6}$ and $\hbar\omega_u/E_0 = 3.53 \times 10^{-5}$ for $N \leq 1750$. These trapping frequencies have been chosen such that the characteristic length scale of the trap is significantly larger than the size of the self-bound droplets, such that the contribution to the energy by the trapping term is neglectable. It must be remarked that we are only considering the formation of stable droplets, for which $E/N < \hbar\omega/2$, although metastable droplets for which $E/N > \hbar\omega/2$ may arise for slightly lower tilting angles/particle numbers. The solution of the eGPE yields a minimum tilting angle for stable droplet formation of $\alpha = 1.262$ for the parameters considered, which lays close to the tilting for the liquidgas phase transition for the infinite system (see Fig. 3), $\alpha = 1.251$. The slight discrepancy between both values most likely arises from the variational approximation employed in the calculations of the infinite system. The realization of these self-bound droplets in experiments with flattened geometries is also feasible, since the peak density fulfills $n_{\text{peak}} < 10^{15} \text{ cm}^{-3}$ for $\alpha < 1.36$, meaning that three-body losses should be manageable in the regime $\alpha \in (1.262, 1.36)$.



FIG. 10. Top: critical number for stable droplet formation vs tilting angle for $a/a_{\rm dd} = 0.7$. In all cases where a droplet is formed, the length scales of the harmonic trap in the x-y plane are significantly larger than the size of the droplet. Bottom: integrated density $n(x, y) = \int dz |\psi(\mathbf{r})|^2$ for a selfbound droplet of N = 10000 atoms at $\alpha = 1.28$.

IV. CONCLUSIONS

In conclusion, we have computed the phase diagram at the 2D equilibrium density of a system of tilted dipolar bosons under a realistic harmonic confinement along the z-axis. We have done so by performing a variational calculation, which we have benchmarked with the solution of the extended Gross-Pitaevskii equation. We have included the effect of beyond mean-field corrections accounting for the discretization of excitation energies along the trapping direction. The diagram hosts three distinct phases in the thermodynamic limit: a gas phase, an unmodulated liquid phase and a striped, modulated liquid phase. Remarkably, and distinctly to the phenomenology in the strict 2D limit, the liquid phases emerge even if the two-body interaction does not host a two-body bound state. We have characterized the transitions between these phases, and have sketched the effect of increasing the trapping strength. We have also performed simulations for the finite system, and have connected the phenomenology in the thermodynamic limit



FIG. 11. Beyond mean-field correction to the chemical potential obtained from the solution of the Bogoliubov de-Gennes equations (black dots) as a function of the 3D density. The solid and dashed lines correspond to the fits of Eqs. A2 and A3, respectively. The labels "Low density region" and "High density region" make reference to the superscripts "low" and "high" of Eqs. A2 and A3. The parameters are $a/a_{\rm dd} = 0.7$ and $\alpha = 1.325$.

with the results. We have illustrated several cases within experimental reach where the realization of potentially supersolid striped states should be possible, and have also characterized the formation of self-bound, stable liquid droplets for a realistic value of the scattering length of $a = 91.56a_0$ for ¹⁶⁴Dy atoms, and for realistic particle numbers.

Our results should serve to motivate and guide potential future experiments in the study of dipolar systems in flattened geometries, in the quasi-2D limit, especially in regards to the realization of liquid phases and supersolid striped states. The proper quantification of the superfluid properties of such states is a matter for future work, where exact quantum Monte Carlo methods could be applied to exactly obtain the superfluid fraction across stripes. Moreover, the role played by dimensionality in thermal effects should also be more extensively characterized in order to check if the surprising and counterintuitive promotion of supersolidity by heating seen in three dimensions [15–17] is robust to dimensionality.

We acknowledge financial support from Ministerio de Ciencia e Innovación MCIN/AEI/10.13039/501100011033 (Spain) under Grant No. PID2023-147469NB-C21 and from AGAUR-Generalitat de Catalunya Grant No. 2021-SGR-01411.

Appendix A: Determination of n_{lim} and the fitting constants of Eqs. 3 and 4

In order to determine the limiting 3D density $n_{\rm lim}$ from Eqs. 3 and 4 in the main text, we numerically compute the beyond mean-field correction to the chemical potential, $\mu_{\rm BMF}$, which can be obtained from the solution of the Bogoliubov de-Gennes equations in the usual manner. The only remmarks are the use of the potential in Fourier space employed in Ref. [25] and the fact that the excitations along the z-axis are discrete. Since this quantity is ultraviolet divergent, as it is well known, we employ the standard renormalization procedure used in three-dimensions, analogously to the renormalization of the energy density performed in Ref. [25]. We then choose $n_{\rm lim}$ such that the ratio $\mu_{\rm BMF}/\mu_{\rm 3D} < 1.3$ for $n > n_{\rm lim}$ for all the calculations considered in this work. Here, $\mu_{\rm 3D}$ is the usual beyond mean-field correction for a dipolar system in three-dimensions, i.e.

$$\mu_{\rm 3D} = \frac{32g\sqrt{a^3}}{3\sqrt{\pi}} \mathcal{Q}_5\left(\frac{a_{\rm dd}}{a}\right) n^{3/2},\tag{A1}$$

where $g = 4\pi\hbar^2 a/m$ and and $Q_5(\varepsilon_{\rm dd}) = \frac{1}{2} \int_0^{\pi} d\alpha \sin \alpha \left[1 + \varepsilon_{\rm dd} (3\cos^2 \alpha - 1)\right]^{5/2}$.. The con-

- Holger Kadau, Matthias Schmitt, Matthias Wenzel, Clarissa Wink, Thomas Maier, Igor Ferrier-Barbut, and Tilman Pfau, "Observing the rosensweig instability of a quantum ferrofluid," Nature 530, 194–197 (2016).
- [2] Matthias Schmitt, Matthias Wenzel, Fabian Böttcher, Igor Ferrier-Barbut, and Tilman Pfau, "Self-bound droplets of a dilute magnetic quantum liquid," Nature 539, 259–262 (2016).
- [3] Igor Ferrier-Barbut, Holger Kadau, Matthias Schmitt, Matthias Wenzel, and Tilman Pfau, "Observation of quantum droplets in a strongly dipolar bose gas," Phys. Rev. Lett. **116**, 215301 (2016).
- [4] L. Chomaz, S. Baier, D. Petter, M. J. Mark, F. Wächtler, L. Santos, and F. Ferlaino, "Quantum-fluctuation-driven crossover from a dilute bose-einstein condensate to a macrodroplet in a dipolar quantum fluid," Phys. Rev. X 6, 041039 (2016).
- [5] Fabian Böttcher, Matthias Wenzel, Jan-Niklas Schmidt, Mingyang Guo, Tim Langen, Igor Ferrier-Barbut, Tilman Pfau, Raúl Bombín, Joan Sánchez-Baena, Jordi Boronat, and Ferran Mazzanti, "Dilute dipolar quantum droplets beyond the extended gross-pitaevskii equation," Phys. Rev. Res. 1, 033088 (2019).
- [6] L. Tanzi, E. Lucioni, F. Famà, J. Catani, A. Fioretti, C. Gabbanini, R. N. Bisset, L. Santos, and G. Modugno, "Observation of a dipolar quantum gas with metastable supersolid properties," Phys. Rev. Lett. **122**, 130405 (2019).
- [7] Fabian Böttcher, Jan-Niklas Schmidt, Matthias Wenzel, Jens Hertkorn, Mingyang Guo, Tim Langen, and Tilman Pfau, "Transient supersolid properties in an array of dipolar quantum droplets," Phys. Rev. X 9, 011051 (2019).
- [8] L. Chomaz, D. Petter, P. Ilzhöfer, G. Natale, A. Trautmann, C. Politi, G. Durastante, R. M. W. van Bijnen, A. Patscheider, M. Sohmen, M. J. Mark, and F. Ferlaino, "Long-lived and transient supersolid behaviors in dipolar quantum gases," Phys. Rev. X 9, 021012 (2019).
- [9] L. Tanzi, S. M. Roccuzzo, E. Lucioni, F. Famà,

stants of Eqs. 3 and 4 are then obtained by fitting the correction of the chemical potential to

$$\mu_{\rm BMF}^{\rm low}(n) = \tilde{C}_1 n + \tilde{C}_2 n^2 + \tilde{C}_3 n^{3/2} \qquad n < n_{\rm lim} \quad (A2)$$

$$\mu_{\rm BMF}^{\rm high}(n) = \tilde{C}_4 n^{3/2} \qquad n > n_{\rm lim} \ .$$
 (A3)

Then, using the relation $\frac{\partial \epsilon_{\text{BMF}}}{\partial n}\Big|_V = \mu_{\text{BMF}}$ we obtain $C_1 = \tilde{C}_1/2, C_2 = \tilde{C}_2/3$ and $C_{3,4} = 2\tilde{C}_{3,4}/5$. We show in Fig. 11, the ratio $\mu_{\text{BMF}}/\mu_{3\text{D}}$ as a function of the density for $a/a_{\text{dd}} = 0.7, \alpha = 1.325, \hbar\omega/E_0 = 1$, together with the fits of Eqs. A2 and A3. For these parameters, we find $n_{\text{lim}}r_0^3 = 0.06$.

A. Fioretti, C. Gabbanini, G. Modugno, A. Recati, and S. Stringari, "Supersolid symmetry breaking from compressional oscillations in a dipolar quantum gas," Nature 574, 382–385 (2019).

- [10] Mingyang Guo, Fabian Böttcher, Jens Hertkorn, Jan-Niklas Schmidt, Matthias Wenzel, Hans Peter Büchler, Tim Langen, and Tilman Pfau, "The low-energy goldstone mode in a trapped dipolar supersolid," Nature 574, 386–389 (2019).
- [11] L. Tanzi, J. G. Maloberti, G. Biagioni, A. Fioretti, C. Gabbanini, and G. Modugno, "Evidence of superfluidity in a dipolar supersolid from nonclassical rotational inertia," Science **371**, 1162–1165 (2021).
- [12] Matthew A. Norcia, Claudia Politi, Lauritz Klaus, Elena Poli, Maximilian Sohmen, Manfred J. Mark, Russell N. Bisset, Luis Santos, and Francesca Ferlaino, "Twodimensional supersolidity in a dipolar quantum gas," Nature **596**, 357–361 (2021).
- [13] Giulio Biagioni, Nicolò Antolini, Aitor Alaña, Michele Modugno, Andrea Fioretti, Carlo Gabbanini, Luca Tanzi, and Giovanni Modugno, "Dimensional crossover in the superfluid-supersolid quantum phase transition," Phys. Rev. X 12, 021019 (2022).
- [14] Maximilian Sohmen, Claudia Politi, Lauritz Klaus, Lauriane Chomaz, Manfred J. Mark, Matthew A. Norcia, and Francesca Ferlaino, "Birth, life, and death of a dipolar supersolid," Phys. Rev. Lett. **126**, 233401 (2021).
- [15] J. Sánchez-Baena, C. Politi, F. Maucher, F. Ferlaino, and T. Pohl, "Heating a dipolar quantum fluid into a solid," Nature Communications 14, 1868 (2023).
- [16] J. Sánchez-Baena, T. Pohl, and F. Maucher, "Superfluidsupersolid phase transition of elongated dipolar boseeinstein condensates at finite temperatures," Phys. Rev. Res. 6, 023183 (2024).
- [17] Liang-Jun He, Juan Sanchez-Baena, Fabian Maucher, and Yong-Chang Zhang, "Accessing elusive twodimensional phases of dipolar bose-einstein condensates by finite temperature," (2024), arXiv:2410.19260 [condmat.quant-gas].

- [18] S. Sunami, V. P. Singh, D. Garrick, A. Beregi, A. J. Barker, K. Luksch, E. Bentine, L. Mathey, and C. J. Foot, "Observation of the berezinskii-kosterlitz-thouless transition in a two-dimensional bose gas via matter-wave interferometry," Phys. Rev. Lett. **128**, 250402 (2022).
- [19] Jr-Chiun Yu, Shaurya Bhave, Lee Reeve, Bo Song, and Ulrich Schneider, "Observing the two-dimensional bose glass in an optical quasicrystal," Nature 633, 338–343 (2024).
- [20] Yanliang Guo, Hepeng Yao, Satwik Ramanjanappa, Sudipta Dhar, Milena Horvath, Lorenzo Pizzino, Thierry Giamarchi, Manuele Landini, and Hanns-Christoph Nägerl, "Observation of the 2d–1d crossover in strongly interacting ultracold bosons," Nature Physics 20, 934– 938 (2024).
- [21] Aristeu R P Lima and Axel Pelster, "Quantum fluctuations in dipolar Bose gases," Physical Review A 84, 041604–4 (2011).
- [22] A. R. P. Lima and A. Pelster, "Beyond mean-field lowlying excitations of dipolar bose gases," Phys. Rev. A 86, 063609 (2012).
- [23] F Wächtler and L Santos, "Ground-state properties and elementary excitations of quantum droplets in dipolar Bose-Einstein condensates," Physical Review A 94, 043618–7 (2016).
- [24] D. Edler, C. Mishra, F. Wächtler, R. Nath, S. Sinha, and L. Santos, "Quantum fluctuations in quasi-onedimensional dipolar bose-einstein condensates," Phys. Rev. Lett. **119**, 050403 (2017).
- [25] Paweł Zin, Maciej Pylak, Tomasz Wasak, Krzysztof Jachymski, and Zbigniew Idziaszek, "Quantum droplets in a dipolar bose gas at a dimensional crossover," Journal of Physics B: Atomic, Molecular and Optical Physics 54, 165302 (2021).
- [26] Li Du, Pierre Barral, Michael Cantara, Julius de Hond, Yu-Kun Lu, and Wolfgang Ketterle, "Atomic physics on a 50-nm scale: Realization of a bilayer system of dipolar atoms," Science **384**, 546–551 (2024), https://www.science.org/doi/pdf/10.1126/science.adh3023.
- [27] Yifei He, Ziting Chen, Haoting Zhen, Mingchen Huang, Mithilesh K Parit, and Gyu-Boong Jo, "Exploring the berezinskii-kosterlitz-thouless transition in a twodimensional dipolar bose gas," (2024), arXiv:2403.18683 [cond-mat.quant-gas].
- [28] Uwe R. Fischer, "Stability of quasi-two-dimensional boseeinstein condensates with dominant dipole-dipole interactions," Phys. Rev. A 73, 031602 (2006).
- [29] Abdelâali Boudjemâa and G. V. Shlyapnikov, "Twodimensional dipolar bose gas with the roton-maxon excitation spectrum," Phys. Rev. A 87, 025601 (2013).
- [30] Chinmayee Mishra and Rejish Nath, "Dipolar condensates with tilted dipoles in a pancake-shaped confinement," Phys. Rev. A 94, 033633 (2016).
- [31] D Baillie and P B Blakie, "A general theory of flattened dipolar condensates," New Journal of Physics 17, 033028 (2015).
- [32] Pengtao Shen and Khandker F. Quader, "Finitetemperature instabilities of a two-dimensional dipolar bose gas at arbitrary tilt angle," Phys. Rev. A 103, 043317 (2021).
- [33] A. K. Fedorov, I. L. Kurbakov, Y. E. Shchadilova, and Yu. E. Lozovik, "Two-dimensional bose gas of tilted dipoles: Roton instability and condensate depletion," Phys. Rev. A **90**, 043616 (2014).

- [34] A. N. Aleksandrova, I. L. Kurbakov, A. K. Fedorov, and Yu. E. Lozovik, "Density-wave-type supersolid of twodimensional tilted dipolar bosons," Phys. Rev. A 109, 063326 (2024).
- [35] F. M. Marchetti and M. M. Parish, "Density-wave phases of dipolar fermions in a bilayer," Phys. Rev. B 87, 045110 (2013).
- [36] J. K. Block and G. M. Bruun, "Properties of the densitywave phase of a two-dimensional dipolar fermi gas," Phys. Rev. B 90, 155102 (2014).
- [37] Au-Chen Lee, D. Baillie, and P. B. Blakie, "Excitations and phase ordering of the spin-stripe phase of a binary dipolar condensate," Phys. Rev. A 109, 023323 (2024).
- [38] Yong-Chang Zhang, Fabian Maucher, and Thomas Pohl, "Supersolidity around a critical point in dipolar boseeinstein condensates," Phys. Rev. Lett. **123**, 015301 (2019).
- [39] Yong-Chang Zhang, Thomas Pohl, and Fabian Maucher, "Phases of supersolids in confined dipolar bose-einstein condensates," Phys. Rev. A 104, 013310 (2021).
- [40] Yong-Chang Zhang, Thomas Pohl, and Fabian Maucher, "Metastable patterns in one- and two-component dipolar bose-einstein condensates," Phys. Rev. Res. 6, 023023 (2024).
- [41] Daniel Lima, Matheus Grossklags, Vinicius Zampronio, Fabio Cinti, and Alejandro Mendoza-Coto, "Supersolid dipolar phases in planar geometry: effects of tilted polarization," (2025), arXiv:2501.09641 [cond-mat.quantgas].
- [42] C Zhang, A Safavi-Naini, Ana Maria Rey, and B Capogrosso-Sansone, "Equilibrium phases of tilted dipolar lattice bosons," New Journal of Physics 17, 123014 (2015).
- [43] Matthias Wenzel, Fabian Böttcher, Tim Langen, Igor Ferrier-Barbut, and Tilman Pfau, "Striped states in a many-body system of tilted dipoles," Phys. Rev. A 96, 053630 (2017).
- [44] A. Macia, F. Mazzanti, and J. Boronat, "Ground state properties and excitation spectrum of a two dimensional gas of bosonic dipoles," The European Physical Journal D 66, 301 (2012).
- [45] A. Macia, D. Hufnagl, F. Mazzanti, J. Boronat, and R. E. Zillich, "Excitations and stripe phase formation in a two-dimensional dipolar bose gas with tilted polarization," Phys. Rev. Lett. **109**, 235307 (2012).
- [46] A. Macia, J. Boronat, and F. Mazzanti, "Phase diagram of dipolar bosons in two dimensions with tilted polarization," Phys. Rev. A 90, 061601 (2014).
- [47] R. Bombin, J. Boronat, and F. Mazzanti, "Dipolar bose supersolid stripes," Phys. Rev. Lett. 119, 250402 (2017).
- [48] Raúl Bombín, Ferran Mazzanti, and Jordi Boronat, "Berezinskii-kosterlitz-thouless transition in twodimensional dipolar stripes," Phys. Rev. A 100, 063614 (2019).
- [49] G. Guijarro, G. E. Astrakharchik, and J. Boronat, "Ultradilute quantum liquid of dipolar atoms in a bilayer," Phys. Rev. Lett. **128**, 063401 (2022).
- [50] G. Guijarro, G. E. Astrakharchik, G. Morigi, and J. Boronat, "Self-assembled chains and solids of dipolar atoms in a multilayer," Phys. Rev. Lett. 133, 233402 (2024).
- [51] Clemens Staudinger, Diana Hufnagl, Ferran Mazzanti, and Robert E. Zillich, "Striped dilute liquid of dipolar bosons in two dimensions," Phys. Rev. A 108, 033303

[52] Holger Kadau, Matthias Schmitt, Matthias Wenzel, Clarissa Wink, Thomas Maier, Igor Ferrier-Barbut, and Tilman Pfau, "Observing the Rosensweig instability of a quantum ferrofluid," Nature **530**, 194–197 (2016).