

Characterizing direct and indirect causal effects when outcomes are dependent due to treatment spillover and outcome spillover

Subhankar Bhadra*

Department of Statistics, The Pennsylvania State University
and

Michael Schweinberger

Department of Statistics, The Pennsylvania State University

Abstract

We provide novel insight into causal inference when both treatment spillover and outcome spillover occur in connected populations, by taking advantage of recent advances in statistical network analysis. Scenarios with treatment spillover and outcome spillover are challenging, because both forms of spillover affect outcomes and therefore treatment spillover and outcome spillover are intertwined, and outcomes are dependent conditional on treatments by virtue of outcome spillover. As a result, the direct and indirect causal effects arising from spillover have remained black boxes: While the direct and indirect causal effects can be identified, it is unknown how these causal effects explicitly depend on the effects of treatment, treatment spillover, and outcome spillover. We make three contributions, facilitated by low-rank random interference graphs. First, we provide novel insight into direct and indirect causal effects by disentangling the contributions of treatment, treatment spillover, and outcome spillover. Second, we provide scalable estimators of direct and indirect causal effects. Third, we establish rates of convergence for estimators of direct and indirect causal effects. These are the first convergence rates in scenarios in which treatment spillover and outcome spillover are intertwined and outcomes are dependent conditional on treatments, and the interference graph is sparse or dense.

Keywords: Causal Inference, Markov Random Field, Random Dot Product Graph, Random Interference Graph

*The authors acknowledge support by ARO award W911NF-21-1-0335.

1 Introduction

At the heart of science is the question of cause and effect (Pearl, 2009; Imbens and Rubin, 2015; Ding and Li, 2018). Statistical approaches to learning cause and effect have witnessed a surge of interest, driven by applications in science and technology (e.g., artificial intelligence). The study of cause and effect involving economic, social, and health-related phenomena is intricate, because the units of interest may be connected, and treating a subset of units can affect the outcomes of both treated and untreated units due to spillover among connected units (e.g., Hudgens and Halloran, 2008; Toulis and Kao, 2013; van der Laan, 2014; Aronow and Samii, 2017; Choi, 2017; Ogburn et al., 2020; Sävje et al., 2021; Tchetgen Tchetgen et al., 2021; Forastiere et al., 2022; Li and Wager, 2022; Ogburn et al., 2024). Understanding spillover is imperative in real-world applications, as demonstrated by the following illustrative examples:

- **Example 1:** If advertisers target teenagers on social media with advertisements featuring designer clothes (treatments), then some of the targeted teenagers may purchase the advertised designer clothes (outcomes). In addition, friends of targeted teenagers may observe them purchasing and wearing designer clothes, and may therefore decide to purchase designer clothes themselves (spillover).
- **Example 2:** To study whether financial incentives encourage couples to have more children, a government conducts a pilot study by sampling couples and providing them with financial incentives to have more children (treatments). Some of the sampled couples may decide to have more children (outcomes). Spillover occurs when unsampled couples observe the benefits that sampled couples with newborn children derive from parenthood and decide to have children themselves.

In both examples, the treatment of a unit can have a direct causal effect on its own outcome, in addition to an indirect causal effect on the outcomes of other units due to spillover. To estimate the total causal effect of treatments, one needs to quantify the direct causal effect of treatments on outcomes along with the indirect causal effect of treatments on outcomes due to spillover. Two forms of spillover can be distinguished: treatment spillover and outcome spillover.

Treatment spillover. The bulk of research on causal inference under interference assumes that the outcome Y_i of unit i is affected by its own treatment W_i and the treatments W_j of other units j (e.g., Aronow and Samii, 2017; Jagadeesan et al., 2020; Sävje et al., 2021; Li and Wager, 2022; Yu et al., 2022). In such scenarios, the outcome Y_i of unit i is not affected by the outcomes Y_j of other units j , so the outcomes Y_1, \dots, Y_N of $N \geq 2$ units are independent conditional on the treatment assignments W_1, \dots, W_N .

Outcome spillover. Outcome spillover occurs when the outcome Y_i of unit i is affected by the outcomes Y_j of other units j (e.g., Ogburn et al., 2020; Tchetgen Tchetgen et al., 2021; Clark and Handcock, 2024; Bhattacharya and Sen, 2024; Ogburn et al., 2024). Outcome spillover implies that the outcome Y_i of unit i depends on the outcomes Y_j of other units j , so the outcomes Y_1, \dots, Y_N are not independent conditional on the treatment assignments W_1, \dots, W_N .

Importance of outcome spillover. While the conditional independence of outcomes Y_1, \dots, Y_N given treatment assignments W_1, \dots, W_N under treatment spillover is convenient, the ignorance-of-outcomes assumption underlying treatment spillover may be violated in the study of economic, social, and health-related phenomena: e.g., why would rational human beings make choices without taking into account the outcomes of other human beings who made the same choices?

Capturing treatment and outcome spillover. A reasonable model should accommodate both treatment spillover and outcome spillover, and should provide insight into how treatment spillover and outcome spillover affect the causal effect of treatment assignments on outcomes. A simple but non-trivial starting point is to assume that the outcomes Y_1, \dots, Y_N conditional on the treatment assignments W_1, \dots, W_N are generated by Markov random fields, which can be viewed as generalizations of linear and generalized linear models (GLMs) for dependent outcomes. Linear and generalized linear models and Markov random fields are widely used tools in data science and have been used to capture outcome spillover in the recent works of Tchetgen Tchetgen et al. (2021), Bhattacharya and Sen (2024), Clark and Handcock (2024), and Ogburn et al. (2020). For example, if the outcomes Y_1, \dots, Y_N are real-valued, treatment spillover and outcome spillover can be captured by Gaussian Markov random fields, which imply that the conditional distributions of outcomes Y_i can be represented by linear models of the form

$$\begin{aligned} \mathbb{E}[Y_i \mid (\mathbf{Y}_{-i}, \mathbf{W}, \mathbf{Z}) = (\mathbf{y}_{-i}, \mathbf{w}, \mathbf{z})] &:= \beta w_i + \gamma c_{N,i,1}(\mathbf{z}) \sum_{j=1}^N w_j z_{i,j} \\ &+ \delta c_{N,i,2}(\mathbf{z}) \sum_{j=1}^N y_j z_{i,j}, \end{aligned} \tag{1}$$

where $N \geq 2$ is the number of units, $\mathbf{Y} \in \mathbb{R}^N$ is the N -vector of outcomes, $\mathbf{Y}_{-i} \in \mathbb{R}^{N-1}$ is the $(N-1)$ -vector of outcomes excluding the outcome $Y_i \in \mathbb{R}$ of unit i , $\mathbf{W} \in \{0, 1\}^N$ is the N -vector of treatment assignments, and $\mathbf{Z} \in \{0, 1\}^{\binom{N}{2}}$ is a $N \times N$ matrix of connection indicators. The quantities $c_{N,i,1}(\mathbf{z})$ and $c_{N,i,2}(\mathbf{z})$ are scaling constants that may depend on the number of units N , unit i , and connections \mathbf{z} (see Section 2), while β , γ , and δ are parameters free from N :

- $\beta \in \mathbb{R}$ captures the effect of the treatment assignment w_i of unit i on its outcome Y_i ;

- $\gamma \in \mathbb{R}$ quantifies the effect of the treatment assignments w_j of other units j connected to i ($z_{i,j} = 1$) on the outcome Y_i of unit i (treatment spillover);
- $\delta \in \mathbb{R}$ quantifies the effect of the outcomes y_j of other units j connected to i ($z_{i,j} = 1$) on the outcome Y_i of unit i (outcome spillover).

Gaussian Markov random fields capture the causal effect of the treatment assignments W_i on the outcomes Y_i of units i and the outcomes Y_j of other units j who are either directly or indirectly connected to i , in addition to correlations (not causation) among the outcomes Y_1, \dots, Y_N arising from outcome spillover ($\delta \neq 0$).

Insight into causal effects. In scenarios in which there is either treatment spillover or outcome spillover or both, two related questions arise:

1. While the direct and indirect causal effect arising from treatment spillover and outcome spillover can be identified (i.e., can be expressed as functions of the data-generating probability model), it is natural to ask: How do these causal effects depend on the direct effect of treatment β , the indirect effect of treatment via treatment spillover γ , and the effect of outcome spillover δ ?
2. How can the effects of treatment, treatment spillover, and outcome spillover be disentangled, and what is the added value of outcome spillover?

Despite the apparent simplicity of models of the form (1), answers to the two questions posed above are lacking. In the absence of outcome spillover ($\delta = 0$), Hu et al. (2022) characterized the direct and indirect causal effects as explicit mathematical functions of β and γ . In the presence of outcome spillover ($\delta \neq 0$), such characterizations are unavailable and more challenging, because treatment spillover and outcome spillover both affect outcomes and are therefore intertwined. In addition, the outcomes Y_1, \dots, Y_N are

dependent conditional on the treatment assignments W_1, \dots, W_N when there is outcome spillover. While outcome spillover has been studied by Tchetgen Tchetgen et al. (2021), Clark and Handcock (2024), Bhattacharya and Sen (2024), and Ogburn et al. (2020), these works do not characterize the direct and indirect causal effects and do not disentangle the contributions of treatment, treatment spillover, and outcome spillover.

Contributions. We provide insight into causal effects by leveraging recent advances in statistical network analysis (Kolaczyk, 2017). First, we characterize the direct and indirect causal effects as explicit mathematical functions of the direct effect of treatment β , the indirect effect of treatment via treatment spillover γ , and the effect of outcome spillover δ with a view to answering the two questions posed above. Second, we take advantage of the characterization of the direct and indirect causal effects to construct scalable plug-in estimators of these causal effects based on low-rank interference graphs. Third, we provide rates of convergence for estimators of the direct and indirect causal effects. To the best of our knowledge, these are the first convergence rates for estimators of direct and indirect causal effects under outcome spillover: While some works offer consistency results and convergence rates for estimators of parameters (e.g., Tchetgen Tchetgen et al., 2021; Bhattacharya and Sen, 2024) and consistency results for estimators of causal effects (e.g., Bhattacharya and Sen, 2024), none of them provides convergence rates for estimators of causal effects under outcome spillover. For example, Tchetgen Tchetgen et al. (2021) report consistency results and convergence rates for estimators of parameters under sparse interference graphs (see regularity conditions 1 and 2 and Theorem 2 in the supplement of Tchetgen Tchetgen et al., 2021), but do not provide theoretical guarantees for estimators of direct and indirect causal effects. Bhattacharya and Sen (2024) report consistency results for estimators of direct and indirect causal effects, without offering convergence rates for

estimators of direct and indirect causal effects. Other works focus on treatment spillover without considering outcome spillover (e.g., Li and Wager, 2022).

Structure. The statistical and causal framework are described in Sections 2 and 3, respectively. We characterize the direct and indirect causal effects as explicit mathematical functions of β , γ , and δ in Section 4, provide estimators of causal effects along with rates of convergence in Section 5, introduce scalable plug-in estimators based on low-rank interference graphs in Section 6, and present simulation results in Section 7.

Notation. We denote the ℓ_1 -, ℓ_2 -, and ℓ_∞ -norm of vectors $\mathbf{a} \in \mathbb{R}^d$ ($d \geq 1$) by $\|\mathbf{a}\|_1$, $\|\mathbf{a}\|_2$, and $\|\mathbf{a}\|_\infty$, respectively, and the matrix norms of matrices $\mathbf{A} \in \mathbb{R}^{d_1 \times d_2}$ ($d_1, d_2 \geq 1$) induced by the ℓ_1 -, ℓ_2 -, and ℓ_∞ -norm by $\|\mathbf{A}\|_1$, $\|\mathbf{A}\|_2$, and $\|\mathbf{A}\|_\infty$, respectively. The Frobenius norm of matrices $\mathbf{A} \in \mathbb{R}^{d_1 \times d_2}$ is denoted by $\|\mathbf{A}\|_F$. The eigenvalues of symmetric matrices $\mathbf{A} \in \mathbb{R}^{d \times d}$ are denoted by $\lambda_1(\mathbf{A}) \geq \dots \geq \lambda_d(\mathbf{A})$. The vector $\mathbf{1}_d \in \mathbb{R}^d$ is the d -vector of 1's and $\mathbf{I}_d \in \mathbb{R}^{d \times d}$ is the identity matrix of order d . The function $\mathbb{I}(\cdot)$ is an indicator function, which is 1 if its argument is true and is 0 otherwise. The matrix $\text{diag}(r_1, \dots, r_d) \in \mathbb{R}^{d \times d}$ is the $d \times d$ diagonal matrix with diagonal elements $r_1, \dots, r_d \in \mathbb{R}$. For two symmetric matrices \mathbf{A} and \mathbf{B} , $\mathbf{A} \succeq \mathbf{B}$ indicates that $\mathbf{A} - \mathbf{B}$ is positive semi-definite. For two sequences of numbers $\{a_N\}$ and $\{b_N\}$, $a_N = O(b_N)$ implies that there exists a constant $C > 0$ such that $a_N \leq C b_N$ for all large N , whereas $a_N = o(b_N)$ implies that $a_N/b_N \rightarrow 0$ as $N \rightarrow \infty$. Probability measures and expectations are denoted by \mathbb{P} and \mathbb{E} , respectively. All theoretical results are proved in the supplement.

2 Statistical framework

We consider a connected population with $N \geq 2$ units. Each unit i

- is either assigned to a control group ($W_i = 0$) or a treatment group ($W_i = 1$);

- has an outcome $Y_i \in \mathbb{R}$;
- has undirected connections $Z_{i,j} \in \{0, 1\}$ to other units j , where $Z_{i,j} = 1$ indicates that i and j are connected and $Z_{i,j} = 0$ otherwise.

The undirected connections $Z_{i,j}$ between units i and j are symmetric in the sense that $Z_{i,j} = Z_{j,i}$, and represent the channels that allow spillover from i to j or from j to i . Self-connections are not of interest for the purpose of studying spillover, and are excluded by defining $Z_{i,i} := 0$. The collections of treatment assignments W_i , outcomes Y_i , and connections $Z_{i,j}$ are denoted by \mathbf{W} , \mathbf{Y} , and \mathbf{Z} , respectively, and $\mathbf{Y}_{-i} \in \mathbb{R}^{N-1}$ is identical to $\mathbf{Y} \in \mathbb{R}^N$ excluding $Y_i \in \mathbb{R}$.

To characterize the direct and indirect causal effects, we make assumptions about the probability laws of \mathbf{W} , \mathbf{Z} , and $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$.

Population Probability Model for \mathbf{W} . We assume that the treatment assignments W_i of units i are independent Bernoulli(π) random variables, where $\pi \in (0, 1)$ is the probability that unit i is assigned to the treatment group. Bernoulli models are standard models for treatment assignments (e.g., Sävje et al., 2021). The treatment assignment probability π may be known or unknown. For the purpose of characterizing and estimating causal effects in Sections 4, 5, and 6, it is immaterial whether π is known or unknown.

Population Probability Model for \mathbf{Z} . We assume that the connections \mathbf{Z} are generated by a random dot product graph model (Tang et al., 2013; Athreya et al., 2021). In other words, the connections $Z_{i,j}$ are independent Bernoulli($P_{i,j}$) random variables with connection probabilities $P_{i,j} := \rho_N \boldsymbol{\alpha}_i^\top \boldsymbol{\alpha}_j$, where $\rho_N \in [0, 1]$ is a sparsity parameter and $\boldsymbol{\alpha} := (\boldsymbol{\alpha}_1, \dots, \boldsymbol{\alpha}_N)^\top \in \mathbb{R}^{N \times K}$ is a matrix satisfying $\boldsymbol{\alpha}_i^\top \boldsymbol{\alpha}_j \in [0, 1]$ for all $i < j$. The sparsity parameter ρ_N may depend on N . The vectors $\boldsymbol{\alpha}_i \in \mathbb{R}^K$ and $\boldsymbol{\alpha}_j \in \mathbb{R}^K$ capture unobserved heterogeneity in the propensities of units i and j to be connected. The dimension K of

the vectors $\boldsymbol{\alpha}_i$ is assumed to be a constant, independent of N , which imposes a low-rank structure on the interference graph. To address the identifiability issue arising from the fact that ρ_N can be multiplied by a constant $C \neq 0$ and $\boldsymbol{\alpha}$ can be divided by \sqrt{C} without changing $\mathbf{P} := \rho_N \boldsymbol{\alpha} \boldsymbol{\alpha}^\top$, we impose the constraint $\max_{1 \leq i \leq N, 1 \leq j \leq K} \alpha_{i,j} = 1$. Random dot product graphs include stochastic block models, degree-corrected block models, and mixed membership models as special cases (Athreya et al., 2021). Such models are popular and facilitate the characterization of direct and indirect causal effects.

Population Probability Model for $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$. In line with Tchetgen Tchetgen et al. (2021), Clark and Handcock (2024), Bhattacharya and Sen (2024), and Ogburn et al. (2020), we assume that $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ is generated by a Markov random field. To facilitate theoretical insight in scenarios in which treatment spillover and outcome spillover are intertwined and outcomes $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ are dependent due to outcome spillover, we assume that $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ is generated by a Gaussian Markov random field. In general, Markov random fields can be specified by either specifying the joint probability law of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ or the conditional probability law of $Y_i \mid (\mathbf{Y}_{-i}, \mathbf{W}, \mathbf{Z}) = (\mathbf{y}_{-i}, \mathbf{w}, \mathbf{z})$, as is well-known in the literature in spatial statistics (Besag, 1974) and graphical models (Lauritzen, 1996; Rue and Held, 2005). We specify a Gaussian Markov random field by specifying the conditional probability law of outcomes $Y_i \mid (\mathbf{Y}_{-i}, \mathbf{W}, \mathbf{Z}) = (\mathbf{y}_{-i}, \mathbf{w}, \mathbf{z})$ as Gaussian, with conditional means

$$\begin{aligned} \mathbb{E}[Y_i \mid (\mathbf{Y}_{-i}, \mathbf{W}, \mathbf{Z}) = (\mathbf{y}_{-i}, \mathbf{w}, \mathbf{z})] &:= \beta w_i + \gamma c_{N,i,1}(\mathbf{z}) \sum_{j=1}^N w_j z_{i,j} \\ &+ \delta c_{N,i,2}(\mathbf{z}) \sum_{j=1}^N y_j z_{i,j} \end{aligned} \tag{2}$$

and conditional variances κ^2 , where β , γ , δ , and κ^2 are parameters free from N , whereas

$c_{N,i,1}(\mathbf{z})$ and $c_{N,i,2}(\mathbf{z})$ are scaling constants. A conventional choice of the scaling constants is $c_{N,i,1}(\mathbf{z}) := 1/\sum_{l=1}^N z_{i,l}$ (e.g., Tchetgen Tchetgen et al., 2021), while the scaling constants $c_{N,i,2}(\mathbf{z})$ are chosen so that the conditional distribution of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ is multivariate Gaussian: see Lemma 1 below.

Three remarks are in place. First, the intercept is set to 0, because the proportion $\gamma c_{N,i,1}(\mathbf{z}) \sum_{j=1}^N w_j z_{i,j} = \gamma \sum_{j=1}^N w_j z_{i,j} / \sum_{l=1}^N z_{i,l}$ tends to the constant $\gamma \pi$ provided $\sum_{l=1}^N z_{i,l} \rightarrow \infty$ as $N \rightarrow \infty$. Such identifiability issues are common in linear-in-means models (see, e.g., Hayes and Levin, 2024). That said, for the purpose of studying causal effects, the intercept can be assumed to be 0 without loss of generality, because the causal effects of interest are sums of differences between potential outcomes (see Section 3), and the intercept does not contribute to the causal effects when the potential outcomes are generated by linear models of the form (2). Second, if there is outcome spillover ($\delta \neq 0$), the outcomes Y_i of units i depend on the outcomes y_j of other units j connected to i , so that outcomes $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ are dependent conditional on treatment assignments and connections. Third, Gaussian Markov random fields help study the causal effect of treatments W_1, \dots, W_N on outcomes Y_1, \dots, Y_N , in addition to capturing correlations among outcomes Y_1, \dots, Y_N due to outcome spillover ($\delta \neq 0$); note that the correlations among outcomes Y_1, \dots, Y_N do not admit causal interpretations, which would require additional data (e.g., temporal observations of outcomes: van der Laan, 2014; Eckles et al., 2017).

The Gaussian distributions of $Y_i \mid (\mathbf{Y}_{-i}, \mathbf{W}, \mathbf{Z}) = (\mathbf{y}_{-i}, \mathbf{w}, \mathbf{z})$ imply that the joint distribution of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ is multivariate Gaussian.

Lemma 1. *Assume that the conditional distributions of $Y_i \mid (\mathbf{Y}_{-i}, \mathbf{W}, \mathbf{Z}) = (\mathbf{y}_{-i}, \mathbf{w}, \mathbf{z})$ are Gaussian with means $\mathbb{E}[Y_i \mid (\mathbf{Y}_{-i}, \mathbf{W}, \mathbf{Z}) = (\mathbf{y}_{-i}, \mathbf{w}, \mathbf{z})]$ defined in (2) and variances $\kappa^2 \in (0, \infty)$, with scaling constants $c_{N,i,1}(\mathbf{z}) = 1/\sum_{l=1}^N z_{i,l}$ and $c_{N,i,2}(\mathbf{z}) = c_N(\mathbf{z})$ ($i = 1, \dots, N$).*

If $(\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})$ is positive definite, then $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ is multivariate Gaussian

- with mean vector $\boldsymbol{\mu}(\mathbf{w}, \mathbf{z}) := (\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})^{-1} \mathbf{G}(\mathbf{w}, \mathbf{z}) \boldsymbol{\vartheta}$,
- and precision matrix $\boldsymbol{\Omega}(\mathbf{w}, \mathbf{z}) := (1/\kappa^2) (\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})$,

where $\boldsymbol{\vartheta} := (\beta, \gamma)^\top \in \mathbb{R}^2$ and $\mathbf{G}(\mathbf{w}, \mathbf{z})$ is a $N \times 2$ matrix with i -th row

$$(\mathbf{G}(\mathbf{w}, \mathbf{z}))_{i,\cdot} := \left(w_i, \frac{\sum_{j=1}^N w_i z_{i,j}}{\sum_{j=1}^N z_{i,j}} \right), \quad i = 1, \dots, N.$$

If $|\delta| < 1$ and $c_N(\mathbf{Z}) \leq 1/\|\mathbf{Z}\|_2$ almost surely, then the multivariate Gaussian distribution of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ is non-singular.

We henceforth assume that the conditions of Lemma 1 hold, so that the distribution of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ is multivariate Gaussian and its conditional independence structure can be represented by a Gaussian Markov random field.

Remarks: scaling constants $c_{N,i,1}(\mathbf{z})$ and $c_{N,i,2}(\mathbf{z})$. The choice of the scaling constants $c_{N,i,1}(\mathbf{z}) := 1/\sum_{l=1}^N z_{i,l}$ is conventional (e.g., Tchetgen Tchetgen et al., 2021). To facilitate the characterization of direct and indirect causal effects, we choose the scaling constants $c_{N,i,2}(\mathbf{z})$ so that the conditional distribution of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ is multivariate Gaussian. By Lemma 1, $c_{N,i,2}(\mathbf{z}) \leq 1/\|\mathbf{z}\|_2$ ensures that $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ is multivariate Gaussian. Since \mathbf{z} is symmetric and $\|\mathbf{z}\|_2 \leq \sqrt{\|\mathbf{z}\|_1 \|\mathbf{z}\|_\infty} = \|\mathbf{z}\|_\infty$, the condition $c_{N,i,2}(\mathbf{z}) \leq 1/\|\mathbf{z}\|_2$ is satisfied when $c_{N,i,2}(\mathbf{z}) := 1/\max_{1 \leq i \leq N} \sum_{l=1}^N z_{i,l}$, in which case both $c_{N,i,1}(\mathbf{z})$ and $c_{N,i,2}(\mathbf{z})$ are functions of degrees $\sum_{l=1}^N z_{i,l}$.

3 Causal framework

In line with Rubin’s causal framework (Imbens and Rubin, 2015), we posit the existence of potential outcomes $Y_i(\mathbf{w})$ for each possible treatment combination $\mathbf{w} \in \{0, 1\}^N$ of the N units and assume that, conditional on the treatment assignments $\mathbf{W} = \mathbf{w}$, we observe outcomes $Y_i(\mathbf{w})$ ($i = 1, \dots, N$).

Ignorability. We assume that the treatment assignments \mathbf{W} are independent of the set of all potential outcomes $Y_1(\mathbf{w}), \dots, Y_N(\mathbf{w})$ under all possible treatment assignments $\mathbf{w} \in \{0, 1\}^N$ and connections \mathbf{Z} among units:

$$\mathbf{W} \perp\!\!\!\perp \{Y_1(\mathbf{w}), \dots, Y_N(\mathbf{w}) : \mathbf{w} \in \{0, 1\}^N\}, \mathbf{Z}. \quad (3)$$

The so-called ignorability assumption (3) is a standard assumption in causal inference, which facilitates the identification of causal effects and can be enforced in experiments (Rosenbaum and Rubin, 1983). In non-experimental settings, the ignorability assumption (3) may not be satisfied, but a form of conditional ignorability (conditional on covariates) may be satisfied (Rosenbaum and Rubin, 1983).

Definition of causal effects. In line with Li and Wager (2022), we define the total causal effect as $\tau_T := \tau_D + \tau_I$, where τ_D is the direct causal effect and τ_I is the indirect causal effect of treatments on outcomes:

$$\begin{aligned} \tau_D &:= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \mathbb{E}[Y_i(W_i = 1, \mathbf{W}_{-i}) - Y_i(W_i = 0, \mathbf{W}_{-i})] \\ \tau_I &:= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \sum_{j=1: j \neq i}^N \mathbb{E}[Y_j(W_i = 1, \mathbf{W}_{-i}) - Y_j(W_i = 0, \mathbf{W}_{-i})]. \end{aligned}$$

The expectation \mathbb{E} is with respect to the joint probability distribution of $(\mathbf{Y}, \mathbf{W}, \mathbf{Z})$. The direct causal effect captures the average effect of the treatment assigned to a unit on its own outcome, whereas the indirect causal effect captures the average effect of the treatment assigned to a unit on the outcomes of other units.

4 Characterizing causal effects

We provide insight into causal inference under interference by characterizing the direct and indirect causal effects τ_D and τ_I as explicit mathematical functions of the direct effect of treatment β , the indirect effect of treatment via treatment spillover γ , and the effect of outcome spillover δ . These characterizations help disentangle treatment spillover and outcome spillover, in addition to quantifying the added value of outcome spillover. Throughout, $\boldsymbol{\theta} := (\beta, \gamma, \delta)^\top$ is the parameter vector of primary interest, and $\boldsymbol{\Theta} := \mathbb{R}^2 \times (-1, +1)$ is its parameter space.

We begin with a general characterization of causal effects, without making assumptions about the model generating the interference graph \mathbf{Z} .

Proposition 1. *Assume that dependent outcomes $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ are generated by a Gaussian Markov random field with parameter vector $\boldsymbol{\theta} \in \boldsymbol{\Theta}$. Define $\mathbf{M}(\delta, \mathbf{Z}) := (\mathbf{I}_N - \delta \mathbf{c}_N(\mathbf{Z}) \mathbf{Z})^{-1}$. Then, without assumptions about the model generating \mathbf{Z} , the direct and total causal effects can be characterized as follows:*

(a) *The direct causal effect is*

$$\begin{aligned} \tau_D &= \beta \lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E}[\text{trace}(\mathbf{M}(\delta, \mathbf{Z}))] \\ &+ \gamma \lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E} \left[\sum_{i=1}^N \sum_{j=1}^N \frac{M_{i,j}(\delta, \mathbf{Z}) Z_{i,j}}{N_j} \right], \end{aligned} \tag{4}$$

where $N_j := \sum_{l=1}^N Z_{j,l}$, provided these limits exist.

(b) The total causal effect is

$$\tau_T \quad := \quad \tau_D + \tau_I \quad = \quad (\beta + \gamma) \lim_{N \rightarrow \infty} \frac{1}{N} \mathbf{1}_N^\top \mathbb{E}[\mathbf{M}(\delta, \mathbf{Z})] \mathbf{1}_N, \quad (5)$$

provided this limit exists.

To characterize the limits in (4) and (5) in Proposition 1 as explicit mathematical functions of β , γ , and δ , we make standard sparsity assumptions about the random dot product graph model generating the interference graph \mathbf{Z} .

Condition C.1. *There exist constants $D > 1$, $p_{\min} > 0$, and $\alpha_{\min} > 0$, which do not depend on N , such that $\rho_N \geq D (\log N)/N$, $\min_{1 \leq i \leq N} \sum_{j=1: j \neq i}^N P_{i,j} \geq p_{\min} N \rho_N$, and $\lambda_K(\boldsymbol{\alpha}^\top \boldsymbol{\alpha}) \geq \alpha_{\min}^2 N$, where $\lambda_K(\boldsymbol{\alpha}^\top \boldsymbol{\alpha})$ is the smallest eigenvalue of $\boldsymbol{\alpha}^\top \boldsymbol{\alpha}$.*

The condition $\rho_N \geq D (\log N)/N$ controls the global sparsity of the interference graph in terms of the expected number of connections, whereas the condition $\min_{1 \leq i \leq N} \sum_{j=1: j \neq i}^N P_{i,j} \geq p_{\min} N \rho_N$ controls the local sparsity of the interference graph in terms of the expected degrees of units. Taken together, these two conditions require the expected degrees of units to grow at least at rate $\log N$, which is a mild sparsity assumption. Sparsity assumptions are common in the literature on network models (Lei and Rinaldo, 2015), and ensure that the network is not too sparse and the information in the data about the estimands of interest is not too low.

Under Condition C.1, the direct and indirect causal effects τ_D and τ_I can be characterized as explicit mathematical functions of β , γ , and δ .

Theorem 1. *Assume that dependent outcomes $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ are generated by a Gaussian Markov random field with parameter vector $\boldsymbol{\theta} \in \boldsymbol{\Theta}$ and scaling constant $c_N(\mathbf{Z}) := (1 / \|\mathbf{Z}\|_2) \mathbb{I}(\|\mathbf{Z}\|_2 > 0)$, and the random dot product graph model generating the interference graph \mathbf{Z} satisfies Condition C.1. Then:*

(a) *The direct causal effect is $\tau_D = \beta$.*

(b) *The indirect causal effect is $\tau_I = \gamma + (\beta + \gamma) a(\delta)$, where*

$$a(\delta) := \delta \lim_{N \rightarrow \infty} \frac{1}{N} (\mathbf{1}_N^\top \boldsymbol{\alpha}) (\|\boldsymbol{\alpha}\|_2^2 \mathbf{I}_K - \delta \boldsymbol{\alpha}^\top \boldsymbol{\alpha})^{-1} (\boldsymbol{\alpha}^\top \mathbf{1}_N),$$

provided this limit exists. If $K = 1$, then

$$a(\delta) := \frac{\delta}{1 - \delta} \lim_{N \rightarrow \infty} \frac{(\mathbf{1}_N^\top \boldsymbol{\alpha})^2}{N \|\boldsymbol{\alpha}\|_2^2}.$$

(c) The total causal effect is $\tau_T = \beta + \gamma + (\beta + \gamma) a(\delta)$.

Theorem 1 characterizes the direct, indirect, and total causal effects τ_D , τ_I , and τ_T as explicit mathematical functions of β , γ , δ , and $a(\delta)$. We can characterize $a(\delta)$ as an explicit mathematical function of δ in special cases of random dot product graph models: e.g., rank-one models (Corollary 1) and degree-corrected block models (Corollary 2). To stress that τ_D , τ_I , and τ_T are functions of $\boldsymbol{\theta}$, we henceforth write $\tau_D(\boldsymbol{\theta})$, $\tau_I(\boldsymbol{\theta})$, and $\tau_T(\boldsymbol{\theta})$.

Corollary 1. *Assume that the conditions of Theorem 1 are satisfied and $Z_{i,j} \stackrel{\text{ind}}{\sim} \text{Bernoulli}(P_{i,j})$ with $P_{i,j} := \rho_N \alpha_i \alpha_j$ ($\rho_N \in [0, 1]$, $\alpha_i \in [0, 1]$, $\alpha_j \in [0, 1]$). If $\alpha_1, \dots, \alpha_N$ are independent and identically distributed random variables with first and second moment $\nu_1 := \mathbb{E}(\alpha_1)$ and $\nu_2 := \mathbb{E}(\alpha_1^2)$, the indirect causal effect is*

$$\tau_I(\boldsymbol{\theta}) = \gamma + (\beta + \gamma) \frac{\delta}{1 - \delta} \frac{\nu_1^2}{\nu_2}.$$

The result in Corollary 1 extends to degree-corrected block models (Karrer and Newman, 2011). Degree-corrected block models are popular models for detecting communities in networks. Assume that a population of N units is partitioned into $K \geq 2$ communities and $Z_{i,j} \stackrel{\text{ind}}{\sim} \text{Bernoulli}(P_{i,j})$, where $P_{i,j} := \rho_N \eta_i \eta_j B_{c_i, c_j}$, $\rho_N \in [0, 1]$, $\eta_i \in [0, 1]$, $\eta_j \in [0, 1]$, $\mathbf{B} := (B_{k,l})_{1 \leq k, l \leq K} \in [0, 1]^{K \times K}$ is positive definite, and c_i is the community of unit i . For ease of presentation, assume that all K communities have the same size n and K is fixed.

Corollary 2. *Assume that the conditions of Theorem 1 are satisfied and $Z_{i,j} \stackrel{\text{ind}}{\sim} \text{Bernoulli}(P_{i,j})$ with $P_{i,j} := \rho_N \eta_i \eta_j B_{c_i, c_j}$, where η_1, \dots, η_N are independent and identically distributed random variables with first and second moment $\nu_1 := \mathbb{E}(\eta_1)$ and $\nu_2 := \mathbb{E}(\eta_1^2)$. If \mathbf{B} is of the form $\mathbf{B} := (p - q) \mathbf{I}_K + q \mathbf{1}_K \mathbf{1}_K^\top$ and $0 \leq q < p \leq 1$, the indirect causal effect is*

$$\tau_I(\boldsymbol{\theta}) = \gamma + (\beta + \gamma) \frac{\delta}{1 - \delta} \frac{\nu_1^2}{\nu_2}.$$

We interpret the results in Theorem 1 and Corollaries 1 and 2 in Section 4.1. We then compare these results to the literature in Section 4.2 and describe the computational and statistical advantages of characterizing causal effects.

4.1 Disentangling contributions to causal effects

Theorem 1 and Corollaries 1 and 2 provide insight into the direct, indirect, and total causal effects $\tau_D(\boldsymbol{\theta})$, $\tau_I(\boldsymbol{\theta})$, and $\tau_T(\boldsymbol{\theta})$ by disentangling the direct effect of treatment β , the indirect effect of treatment via treatment spillover γ , and the effect of outcome spillover δ , in addition to characterizing the added value of outcome spillover. To demonstrate, note that the direct causal effect $\tau_D(\boldsymbol{\theta}) = \beta$ aligns with expectation, whereas the indirect causal effect $\tau_I(\boldsymbol{\theta}) = \gamma + (\beta + \gamma) a(\delta)$ and the total causal effect $\tau_T(\boldsymbol{\theta}) = \beta + \gamma + (\beta + \gamma) a(\delta)$ require a more careful interpretation.

Vanishing total causal effect. The total causal effect $\tau_T(\boldsymbol{\theta})$ can vanish even when the direct causal effect $\tau_D(\boldsymbol{\theta})$ does not vanish: If $\tau_D(\boldsymbol{\theta}) = \beta \neq 0$ and $\gamma = -\beta$, the indirect effect of treatment spillover γ cancels the direct effect of treatment β , so that $\tau_T(\boldsymbol{\theta}) = 0$.

Vanishing indirect causal effect. The indirect causal effect $\tau_I(\boldsymbol{\theta})$ vanishes either when there is no spillover ($\gamma = 0$ and $\delta = 0$) or when the indirect effect of treatment

spillover γ cancels the direct effect of treatment β , in the sense that

$$\gamma = -\beta \frac{a(\delta)}{1 + a(\delta)}; \quad (6)$$

note that the proof of Theorem 1 establishes that $a(\delta) \neq -1$. A special case of (6) is $\beta = 0$ (which implies $\gamma = 0$), in which case $\tau_I(\boldsymbol{\theta}) = 0$ regardless of whether $\delta = 0$ or $\delta \neq 0$. In other words, if the direct effect of treatment and the indirect effect of treatment spillover vanish ($\tau_D(\boldsymbol{\theta}) = \beta = 0$ and $\gamma = 0$), the indirect and total causal effects $\tau_I(\boldsymbol{\theta})$ and $\tau_T(\boldsymbol{\theta})$ vanish, even when there is outcome spillover ($\delta \neq 0$). The reason is that $\tau_I(\boldsymbol{\theta})$ and $\tau_T(\boldsymbol{\theta})$ capture the indirect and total causal effects of treatments on outcomes, and (2) reveals that the treatment does not have any effect on outcomes when there is neither a direct effect of treatment ($\beta = 0$) nor an indirect effect of treatment via treatment spillover ($\gamma = 0$).

Indirect causal effect, without direct causal effect. An indirect causal effect $\tau_I(\boldsymbol{\theta}) \neq 0$ can arise when there is no direct causal effect: If $\tau_D(\boldsymbol{\theta}) = \beta = 0$, there is no direct causal effect, but there is an indirect causal effect $\tau_I(\boldsymbol{\theta}) = \gamma(1 + a(\delta)) \neq 0$ as long as there is treatment spillover ($\gamma \neq 0$); recall that $a(\delta) \neq -1$.

Treatment spillover, without outcome spillover. If there is treatment spillover ($\gamma \neq 0$) without outcome spillover ($\delta = 0$), the indirect causal effect is $\tau_I(\boldsymbol{\theta}) = \gamma$.

Added value of outcome spillover. If there is outcome spillover ($\delta \neq 0$) in addition to treatment spillover ($\gamma \neq 0$), the indirect causal effect is $\tau_I(\boldsymbol{\theta}) = \gamma + (\beta + \gamma)a(\delta)$ and the total causal effect is $\tau_T(\boldsymbol{\theta}) = \beta + \gamma + (\beta + \gamma)a(\delta)$. In such scenarios, $(\beta + \gamma)a(\delta)$ can be interpreted as the value added by outcome spillover to $\tau_I(\boldsymbol{\theta})$ and $\tau_T(\boldsymbol{\theta})$, because $\tau_I(\boldsymbol{\theta}) = \gamma$ and $\tau_T(\boldsymbol{\theta}) = \beta + \gamma$ when $\delta = 0$.

Treatment and outcome spillover are intertwined. The added value of outcome spillover $(\beta + \gamma)a(\delta)$ reveals that the greater the direct effect of treatment β and the indirect

effect of treatment via treatment spillover γ are, the greater is the effect of outcome spillover δ on the indirect and total causal effects $\tau_I(\boldsymbol{\theta})$ and $\tau_T(\boldsymbol{\theta})$. The fact that $\beta + \gamma$ magnifies the effect of δ underscores that the effects of treatment β , treatment spillover γ , and outcome spillover δ are intertwined, and that characterizing causal effects is critical to disentangling the effects of treatment, treatment spillover, and outcome spillover.

4.2 Advantages of characterizing causal effects

The literature on causal inference under interference has not characterized the direct, indirect, and total causal effects $\tau_D(\boldsymbol{\theta})$, $\tau_I(\boldsymbol{\theta})$, and $\tau_T(\boldsymbol{\theta})$ when there is both treatment spillover and outcome spillover ($\gamma \neq 0$ and $\delta \neq 0$). While Hu et al. (2022) characterized causal effects in the absence of outcome spillover ($\delta = 0$), characterizing them in the presence of outcome spillover ($\delta \neq 0$) is more challenging, because treatment spillover and outcome spillover both affect outcomes and are therefore intertwined, and outcomes are dependent conditional on treatments. The characterization of $\tau_D(\boldsymbol{\theta})$, $\tau_I(\boldsymbol{\theta})$, and $\tau_T(\boldsymbol{\theta})$ in Theorem 1 and Corollaries 1 and 2 comes with at least three benefits, in addition to providing insight:

1. **Estimating sources of causal effects:** Without characterizing the direct, indirect, and total causal effects $\tau_D(\boldsymbol{\theta})$, $\tau_I(\boldsymbol{\theta})$, and $\tau_T(\boldsymbol{\theta})$ and disentangling the contributions of treatment β , treatment spillover γ , and outcome spillover δ , it is challenging to estimate the contributions of these effects to $\tau_D(\boldsymbol{\theta})$, $\tau_I(\boldsymbol{\theta})$, and $\tau_T(\boldsymbol{\theta})$. For example, while one can approximate the indirect and total causal effects $\tau_I(\boldsymbol{\theta})$ and $\tau_T(\boldsymbol{\theta})$ by Markov chain Monte Carlo simulations along the lines of Tchetgen Tchetgen et al. (2021), it is not evident how one can estimate the value added by outcome spillover to $\tau_I(\boldsymbol{\theta})$ and $\tau_T(\boldsymbol{\theta})$. As a result, the literature on causal inference under interference has not been able to estimate the added value of outcome spillover. By contrast, by

characterizing $\tau_I(\boldsymbol{\theta})$ and $\tau_T(\boldsymbol{\theta})$ as explicit mathematical functions of $\boldsymbol{\theta}$ and revealing that the added value of outcome spillover is $(\beta + \gamma) a(\delta)$, one can estimate the added value of outcome spillover $(\beta + \gamma) a(\delta)$ by estimating β , γ , δ , and $a(\delta)$.

2. **Transportability:** Characterizing the direct, indirect, and total causal effects $\tau_D(\boldsymbol{\theta})$, $\tau_I(\boldsymbol{\theta})$, and $\tau_T(\boldsymbol{\theta})$ as explicit mathematical functions of $\boldsymbol{\theta}$ helps generalize causal conclusions from one population to other populations (Pearl and Bareinboim, 2014). For example, suppose that an experiment is conducted in Los Angeles to assess the total causal effect $\tau_T^{LA}(\boldsymbol{\theta})$ of economic, financial, or political interventions in Los Angeles, and suppose that New York City is interested in estimating the total causal effect $\tau_T^{NY}(\boldsymbol{\theta})$ of these interventions in New York City, without conducting a time-consuming and expensive experiment. If there is reason to believe that Los Angeles and New York City share the same treatment, treatment spillover, and outcome spillover effects β , γ , and δ while the interference graph that enables spillover in New York City differs from the one in Los Angeles, then researchers could

- (a) collect network data in New York City, without conducting an experiment in New York City;
- (b) estimate the function $a(\delta)$ based on the network data collected in New York City, using the spectral estimator $\hat{a}(\delta)$ described in Section 5;
- (c) report $\tau_T^{NY}(\hat{\boldsymbol{\theta}}) = \hat{\beta} + \hat{\gamma} + (\hat{\beta} + \hat{\gamma}) \hat{a}(\hat{\delta})$ as an educated guess of the total causal effect in New York City, where $\hat{\boldsymbol{\theta}} := (\hat{\beta}, \hat{\gamma}, \hat{\delta})^\top$ is the maximum likelihood estimator of $\boldsymbol{\theta} := (\beta, \gamma, \delta)^\top$ based on the Los Angeles experiment, whereas $\hat{a}(\hat{\delta})$ is the spectral estimator of $a(\hat{\delta})$ based on the New York City network data.

In other words, transportability helps estimate the total causal effect $\tau_T^{NY}(\boldsymbol{\theta})$ of inter-

ventions in New York City, without conducting an experiment in New York City. The characterization of the total causal effect $\tau_T(\boldsymbol{\theta})$ as an explicit mathematical function of $\boldsymbol{\theta}$ facilitates transportability.

3. **Theoretical guarantees:** In addition to constructing estimators of direct and indirect causal effects $\tau_D(\boldsymbol{\theta})$ and $\tau_I(\boldsymbol{\theta})$ and the added value of outcome spillover $(\beta + \gamma) a(\delta)$, characterizing causal effects facilitates theoretical guarantees for estimators of $\tau_D(\boldsymbol{\theta})$, $\tau_I(\boldsymbol{\theta})$, and $(\beta + \gamma) a(\delta)$.

We introduce estimators of causal effects along with rates of convergence in Section 5, and propose scalable plug-in estimators based on low-rank interference graphs in Section 6.

5 Estimators of causal effects and rates of convergence

The characterization of the direct and indirect causal effects $\tau_D(\boldsymbol{\theta}^*)$ and $\tau_I(\boldsymbol{\theta}^*)$ as explicit mathematical functions of the data-generating parameter vector $\boldsymbol{\theta}^*$ and the function $a(\delta^*)$ paves the way for plug-in estimators of $\tau_D(\boldsymbol{\theta}^*)$ and $\tau_I(\boldsymbol{\theta}^*)$ using estimators of $\boldsymbol{\theta}^*$ and $a(\delta^*)$.

We estimate $\boldsymbol{\theta}^*$ by maximizing the likelihood, which is based on the multivariate Gaussian distribution of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ by Lemma 1. The function $a(\delta^*)$ can be estimated by spectral methods, and δ^* can be replaced by its maximum likelihood estimator $\hat{\delta}$. Let $\hat{\mathbf{U}}$ be a $N \times K$ matrix containing the K leading eigenvectors of \mathbf{Z} , let $\hat{\lambda}_1 \geq \dots \geq \hat{\lambda}_K$ be the corresponding leading eigenvalues, and define $\mathbf{T} := \sqrt{\rho_N} \boldsymbol{\alpha}$. By the definition of the function $a(\delta^*)$ and the matrix \mathbf{T} , the function $a(\delta^*)$ can be expressed in terms of \mathbf{T} as

$$a(\delta^*) := \delta^* \lim_{N \rightarrow \infty} \frac{1}{N} (\mathbf{1}_N^\top \mathbf{T}) (\|\mathbf{T}\|_2^2 \mathbf{I}_K - \delta \mathbf{T}^\top \mathbf{T})^{-1} (\mathbf{T}^\top \mathbf{1}_N).$$

Since \mathbf{T} can be estimated (up to an orthogonal transformation) by

$$\hat{\mathbf{T}} := \hat{\mathbf{U}} \operatorname{diag} \left(\sqrt{|\hat{\lambda}_1|}, \dots, \sqrt{|\hat{\lambda}_K|} \right), \quad (7)$$

we can estimate $a(\delta^*)$ by the plug-in estimator

$$\hat{a}(\hat{\delta}) := \frac{\hat{\delta}}{N} (\mathbf{1}_N^\top \hat{\mathbf{T}}) \left(\|\hat{\mathbf{T}}\|_2^2 \mathbf{I}_K - \hat{\delta} \hat{\mathbf{T}}^\top \hat{\mathbf{T}} \right)^{-1} (\hat{\mathbf{T}}^\top \mathbf{1}_N). \quad (8)$$

Upon estimating $\boldsymbol{\theta}^* := (\beta^*, \gamma^*, \delta^*)^\top$ by the maximum likelihood estimator $\hat{\boldsymbol{\theta}} := (\hat{\beta}, \hat{\gamma}, \hat{\delta})^\top$ and estimating $a(\hat{\delta})$ by the spectral estimator $\hat{a}(\hat{\delta})$, we can estimate the direct and indirect causal effects $\tau_D(\boldsymbol{\theta}^*)$ and $\tau_I(\boldsymbol{\theta}^*)$ by

$$\begin{aligned} \tau_D(\hat{\boldsymbol{\theta}}) &:= \hat{\beta} \\ \tau_I(\hat{\boldsymbol{\theta}}) &:= \hat{\gamma} + (\hat{\beta} + \hat{\gamma}) \hat{a}(\hat{\delta}). \end{aligned}$$

To obtain rates of convergence for $\tau_D(\hat{\boldsymbol{\theta}})$ and $\tau_I(\hat{\boldsymbol{\theta}})$, we need three additional conditions.

Condition C.2. *The data-generating parameter vector $\boldsymbol{\theta}^*$ satisfies $\beta^* + \gamma^* \neq 0$ and $\delta^* \in (-1, +1)$, and the scaling constant $c_N(\mathbf{Z})$ is $c_N(\mathbf{Z}) := (1/\|\mathbf{Z}\|_2) \mathbb{I}(\|\mathbf{Z}\|_2 > 0)$.*

Condition C.3. *There exists a constant $p^* \in (0, 1)$, which does not depend on N , such that $(\mathbf{1}_N^\top \mathbf{u})^2/N \leq p^*$ for all unit vectors \mathbf{u} in the column space of \mathbf{P} .*

Condition C.4. *There exists a constant $C(\delta^*) > 0$, which depends on $\delta^* \in (-1, +1)$ but does not depend on N , such that $|a_N(\delta^*) - a(\delta^*)| \leq C(\delta^*) \sqrt{(\log N)/(N\rho_N)}$ for all large enough N , where $a_N(\delta^*)$ is the finite-population approximation of $a(\delta^*)$ defined by*

$$a_N(\delta^*) := \frac{\delta^*}{N} (\mathbf{1}_N^\top \mathbf{T}) \left(\|\mathbf{T}\|_2^2 \mathbf{I}_K - \delta^* \mathbf{T}^\top \mathbf{T} \right)^{-1} (\mathbf{T}^\top \mathbf{1}_N).$$

Condition C.2 rules out the special case $\beta^* + \gamma^* = 0$, in which case the total causal effect $\tau_T(\theta^*)$ vanishes by Theorem 1, and ensures that the conditional distribution of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ is multivariate Gaussian by Lemma 1. Condition C.3 implies that the constant vector is not too close to the column space of \mathbf{P} . If the rank of \mathbf{P} is $K = 1$, Condition C.3 is satisfied if $(\mathbf{1}_N^\top \boldsymbol{\alpha})^2 / (N \|\boldsymbol{\alpha}\|_2^2)$ converges to a real number less than 1. Corollary 1 shows that when $\alpha_1, \dots, \alpha_N$ are independent and identically distributed random variables, then $(\mathbf{1}_N^\top \boldsymbol{\alpha})^2 / (N \|\boldsymbol{\alpha}\|_2^2)$ converges to $\nu_1^2 / \nu_2 = [\mathbb{E}(\alpha_1)]^2 / \mathbb{E}(\alpha_1^2)$ almost surely. The limit ν_1^2 / ν_2 is less than 1 provided that the α_i 's are not constant, that is, as long as the interference graph \mathbf{Z} exhibits degree heterogeneity. Degree heterogeneity is a reasonable requirement, because many real-world networks exhibit degree heterogeneity. Condition C.4 helps bound $\widehat{a}(\widehat{\delta}) - a(\delta^*)$, which can be decomposed into three parts: $\widehat{a}(\widehat{\delta}) - a_N(\widehat{\delta})$, $a_N(\widehat{\delta}) - a_N(\delta^*)$, and $a_N(\delta^*) - a(\delta^*)$. Condition C.4 controls the third part, $a_N(\delta^*) - a(\delta^*)$. We can verify Condition C.4 in special cases, such as rank-one models of the interference graph \mathbf{Z} , as demonstrated by Proposition 2:

Proposition 2. *Assume that $Z_{i,j} \stackrel{\text{ind}}{\sim} \text{Bernoulli}(P_{i,j})$ and $P_{i,j} := \rho_N \alpha_i \alpha_j$ ($\rho_N \in [0, 1]$, $\alpha_i \in [0, 1]$, $\alpha_j \in [0, 1]$) satisfies Condition C.1. Let $\alpha_1, \dots, \alpha_N$ be independent and identically distributed random variables with first and second moment $\nu_1 := \mathbb{E}(\alpha_1)$ and $\nu_2 := \mathbb{E}(\alpha_1^2)$ and Subgaussian tails: that is, for any $t > 0$,*

$$\mathbb{P}(|\alpha_i - \nu_1| > t) \leq 2 \exp\left(-\frac{t^2}{2\sigma^2}\right), \quad \sigma^2 \in (0, \infty), \quad i = 1, \dots, N.$$

Then Condition C.4 is satisfied with high probability, in the sense that

$$|a_N(\delta^*) - a(\delta^*)| \leq C(\delta^*) \sqrt{\frac{\log N}{N}}$$

holds with probability at least $1 - 4/N^2$, provided N is large enough.

Under Conditions C.1–C.4, we obtain convergence rates for the estimators $\tau_D(\hat{\boldsymbol{\theta}})$ and $\tau_I(\hat{\boldsymbol{\theta}})$ of the direct and indirect causal effects $\tau_D(\boldsymbol{\theta}^*)$ and $\tau_I(\boldsymbol{\theta}^*)$. These are the first convergence rates for estimators of direct and indirect causal effects under both treatment spillover and outcome spillover based on a single observation of dependent outcomes $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$, as discussed in Section 1.

Theorem 2. *Assume that dependent outcomes $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ are generated by a Gaussian Markov random field with parameters $\boldsymbol{\theta}^* \in \boldsymbol{\Theta}$ and $\kappa^{*2} \in (0, \infty)$, and that Conditions C.1–C.4 are satisfied. Then there exist constants $C_D(\boldsymbol{\theta}^*, \kappa^{*2}) > 0$ and $C_I(\boldsymbol{\theta}^*, \kappa^{*2}) > 0$, which depend on $\boldsymbol{\theta}^*$ and κ^{*2} but do not depend on N , and a constant $C > 0$, independent of N , such that*

$$\begin{aligned} |\tau_D(\hat{\boldsymbol{\theta}}) - \tau_D(\boldsymbol{\theta}^*)| &\leq C_D(\boldsymbol{\theta}^*, \kappa^{*2}) \sqrt{\frac{\log N}{N}} \\ |\tau_I(\hat{\boldsymbol{\theta}}) - \tau_I(\boldsymbol{\theta}^*)| &\leq C_I(\boldsymbol{\theta}^*, \kappa^{*2}) \sqrt{\frac{\log N}{N \rho_N}} \end{aligned}$$

with probability at least $1 - C/N^2$, provided N is large enough.

Theorem 2 shows that the convergence rate of the estimator $\tau_D(\hat{\boldsymbol{\theta}})$ of the direct causal effect $\tau_D(\boldsymbol{\theta}^*)$ depends on N , whereas the convergence rate of the estimator $\tau_I(\hat{\boldsymbol{\theta}})$ of the indirect causal effect $\tau_I(\boldsymbol{\theta}^*)$ depends on both N and ρ_N . The sparsity parameter ρ_N controls the expected number of channels available for treatment and outcome spillover: The higher ρ_N is, the higher is expected number of channels available for spillover, which helps estimate the indirect causal effect $\tau_I(\boldsymbol{\theta}^*)$ arising from treatment and outcome spillover. As discussed above, the convergence rates in Theorem 2 are the first convergence rates for estimators of direct and indirect causal effects in scenarios with both treatment and outcome spillover.

6 Scalable estimators of causal effects

The plug-in estimators $\tau_D(\hat{\boldsymbol{\theta}})$ and $\tau_I(\hat{\boldsymbol{\theta}})$ of the direct and indirect causal effects $\tau_D(\boldsymbol{\theta}^*)$ and $\tau_I(\boldsymbol{\theta}^*)$ introduced in Section 5 are based on the maximum likelihood estimator $\hat{\boldsymbol{\theta}}$ of $\boldsymbol{\theta}^*$. We introduce here more scalable plug-in estimators by approximating the maximum likelihood estimator $\hat{\boldsymbol{\theta}}$. The approximate maximum likelihood estimator is based on approximations of the likelihood leveraging low-rank interference graphs. As a case in point, we consider rank-one approximations of the likelihood, which assume that the rank of $\mathbb{E}(\mathbf{Z}) = \mathbf{P}$ is $K = 1$. Higher-rank approximations of the likelihood are possible, but come at a computational cost. The simulation results in Section 7 demonstrate that rank-one approximations perform well, which suggests that the added value of higher-rank approximations is limited.

The computational bottleneck of likelihood inference is the precision matrix $\boldsymbol{\Omega}(\mathbf{w}, \mathbf{z}) := (1/\kappa^2)(\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})$ of the multivariate Gaussian distribution of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$. If N is large, computing the determinant and inverse of the $N \times N$ matrix $\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z}$ is inconvenient. We propose low-rank approximations of the likelihood to avoid the need to compute the determinant and inverse of $\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z}$, taking advantage of low-rank interference graphs \mathbf{Z} . To introduce low-rank approximations of the likelihood, let $\mathbf{T} := \sqrt{\rho_N} \boldsymbol{\alpha}$ and $c_N(\mathbf{Z}) := (1/\|\mathbf{Z}\|_2) \mathbb{I}(\|\mathbf{Z}\|_2 > 0)$. Since the probability mass of smooth functions of \mathbf{Z} concentrates around the expectations of those functions (Vershynin, 2018), the Sherman-Morrison formula suggests the approximations

$$(\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} \approx (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1} = \mathbf{I}_N + \frac{\delta \mathbf{T} \mathbf{T}^\top}{(1 - \delta) \mathbf{T}^\top \mathbf{T}} \quad (9)$$

$$\det(\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z}) \approx \det(\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P}) = \det\left(\mathbf{I}_N - \frac{\delta \mathbf{T} \mathbf{T}^\top}{\mathbf{T}^\top \mathbf{T}}\right) = 1 - \delta. \quad (10)$$

The matrix \mathbf{T} can be estimated by the spectral estimator stated in Equation (7). If \mathbf{T} is

replaced by $\widehat{\mathbf{T}}$ in Equations (9) and (10), it is straightforward to compute the inner and outer product of $\widehat{\mathbf{T}}$ and update the inverse of $\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P}$ whenever δ is updated. The resulting approximation of the loglikelihood is

$$\begin{aligned} \tilde{\ell}(\boldsymbol{\theta}, \kappa^2) &:= -\frac{N}{2} \log(\kappa^2) + \frac{1}{2} \log(1 - \delta) - \frac{1}{2\kappa^2} \boldsymbol{\vartheta}^\top \mathbf{G}(\mathbf{w}, \mathbf{z})^\top \mathbf{G}(\mathbf{w}, \mathbf{z}) \boldsymbol{\vartheta} - \frac{1}{2\kappa^2} \mathbf{y}^\top \mathbf{y} \\ &- \frac{1}{2\kappa^2} \frac{\delta (\boldsymbol{\vartheta}^\top \mathbf{G}(\mathbf{w}, \mathbf{z})^\top \mathbf{T})^2}{(1 - \delta) \mathbf{T}^\top \mathbf{T}} + \frac{c_N(\mathbf{z}) \delta}{2\kappa^2} \mathbf{y}^\top \mathbf{z} \mathbf{y} + \frac{\boldsymbol{\vartheta}^\top}{\kappa^2} \mathbf{G}(\mathbf{w}, \mathbf{z})^\top \mathbf{y}, \end{aligned}$$

where $\boldsymbol{\vartheta} := (\beta, \gamma)^\top$. We maximize $\tilde{\ell}(\boldsymbol{\theta}, \kappa^2)$ by iterating three steps:

- **Step 1:** Given $\boldsymbol{\vartheta}_{(t)}$ and $\kappa_{(t)}^2$, update δ subject to the constraint that $\delta \in (-1, +1)$:

$$\delta_{(t+1)} = 1 - \frac{\frac{1}{2\kappa_{(t)}^2} \frac{(\boldsymbol{\vartheta}_{(t)}^\top \mathbf{G}(\mathbf{w}, \mathbf{z})^\top \mathbf{T})^2}{\mathbf{T}^\top \mathbf{T}}}{\sqrt{\frac{1}{4} + \frac{c_N(\mathbf{z})}{\kappa_{(t)}^4} \frac{(\boldsymbol{\vartheta}_{(t)}^\top \mathbf{G}(\mathbf{w}, \mathbf{z})^\top \mathbf{T})^2}{\mathbf{T}^\top \mathbf{T}} \mathbf{y}^\top \mathbf{z} \mathbf{y} - \frac{1}{2}}},$$

provided $\mathbf{y}^\top \mathbf{z} \mathbf{y} > 0$.

- **Step 2:** Given $\delta_{(t+1)}$, update $\boldsymbol{\vartheta}$ by

$$\boldsymbol{\vartheta}_{(t+1)} = \left(\mathbf{G}(\mathbf{w}, \mathbf{z})^\top \left(\mathbf{I}_N + \frac{\delta_{(t+1)} \mathbf{T} \mathbf{T}^\top}{(1 - \delta_{(t+1)}) \mathbf{T}^\top \mathbf{T}} \right) \mathbf{G}(\mathbf{w}, \mathbf{z}) \right)^{-1} \mathbf{G}(\mathbf{w}, \mathbf{z})^\top \mathbf{y}.$$

- **Step 3:** Given $\boldsymbol{\vartheta}_{(t+1)}$ and $\delta_{(t+1)}$, update κ^2 by

$$\begin{aligned} \kappa_{(t+1)}^2 &= \frac{1}{N} \left(\|\mathbf{y} - \mathbf{G}(\mathbf{w}, \mathbf{z}) \boldsymbol{\vartheta}_{(t+1)}\|_2^2 + \frac{\delta_{(t+1)} (\boldsymbol{\vartheta}_{(t+1)}^\top \mathbf{G}(\mathbf{w}, \mathbf{z})^\top \mathbf{T})^2}{(1 - \delta_{(t+1)}) \mathbf{T}^\top \mathbf{T}} \right. \\ &\quad \left. - c_N(\mathbf{z}) \delta_{(t+1)} \mathbf{y}^\top \mathbf{z} \mathbf{y} \right). \end{aligned}$$

In Step 1, the denominator of the ratio on the right-hand side can vanish when $\mathbf{y}^\top \mathbf{z} \mathbf{y}$ is

too small, in which case a solution for δ may not exist. It can be shown that the event that $\mathbf{Y}^\top \mathbf{Z} \mathbf{Y}$ is too small is a rare event (i.e., it occurs with a low probability). In the rare event that $\mathbf{Y}^\top \mathbf{Z} \mathbf{Y}$ is too small, one can use maximum likelihood estimators instead of approximate maximum likelihood estimators.

7 Simulation results

We shed light onto the finite-sample behavior of the plug-in estimators $\tau_D(\hat{\boldsymbol{\theta}})$ and $\tau_I(\hat{\boldsymbol{\theta}})$ introduced in Section 6 using simulations. Throughout, $\hat{\boldsymbol{\theta}}$ is the approximate maximum likelihood estimator of the data-generating parameter vector $\boldsymbol{\theta}^*$ introduced in Section 6 based on rank-one approximations of the likelihood, and $\tau_D(\hat{\boldsymbol{\theta}})$ and $\tau_I(\hat{\boldsymbol{\theta}})$ are the corresponding plug-in estimators of $\tau_D(\boldsymbol{\theta}^*)$ and $\tau_I(\boldsymbol{\theta}^*)$. We compare $\tau_D(\hat{\boldsymbol{\theta}})$ and $\tau_I(\hat{\boldsymbol{\theta}})$ to the Hajek-estimator $\hat{\tau}_D^{\text{HAJ}}$ of the direct causal effect and the PC-balancing estimator $\hat{\tau}_I^{\text{PC}}$ of the indirect causal effect introduced and defined in Li and Wager (2022). While Li and Wager (2022) focus on scenarios with treatment spillover but without outcome spillover ($\delta = 0$), the estimators of Li and Wager (2022) provide the best benchmark, because Li and Wager (2022) consider real-valued outcomes $Y_i \in \mathbb{R}$ and a random interference graph \mathbf{Z} , which dovetails with our assumptions; note that direct comparisons with other works are problematic, because all of them make assumptions that are incompatible with our assumptions: e.g., Tchetgen Tchetgen et al. (2021) assume that the conditional densities of Y_i are bounded away from 0 by a constant $C > 0$, which rules out real-valued outcomes $Y_i \in \mathbb{R}$ with Gaussian distributions (see regularity condition 3 in the supplement of Tchetgen Tchetgen et al., 2021); Bhattacharya and Sen (2024) assume that $Y_i \in [-1, +1]$; and Clark and Handcock (2024) assume that $Y_i \in \{0, 1\}$.

We generate data from the joint probability model proposed in Section 2 as follows.

First, we generate treatment assignments $W_i \stackrel{\text{iid}}{\sim} \text{Bernoulli}(\pi)$ using $\pi = 2/5$, and then generate the interference graph \mathbf{Z} from a rank-one random dot product graph model with $\mathbb{E}(Z_{i,j}) = P_{i,j} = \rho_N \alpha_i \alpha_j$ and $\alpha_i \stackrel{\text{iid}}{\sim} \text{Beta}(1, 3)$ ($i = 1, \dots, N$), which induces long-tailed distributions that emulate the degree distributions of many real-world networks and implies that $\nu_1^2/\nu_2 = 0.625$. The sparsity parameter is $\rho_N = 1$ in the dense-graph scenario and $\rho_N = N^{-2/5}$ in the sparse-graph scenario. Conditional on $(\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$, the outcomes \mathbf{Y} are generated by a Gaussian Markov random field with mean vector $\boldsymbol{\mu}(\mathbf{w}, \mathbf{z})$ and precision matrix $\boldsymbol{\Omega}(\mathbf{w}, \mathbf{z})$ defined in Section 2. We consider three scenarios:

- **Scenario A:** treatment spillover, without outcome spillover: $\tau_D(\boldsymbol{\theta}^*) = \beta^* = 1$, $\gamma^* = 1$, $\delta^* = 0$, and $\tau_I(\boldsymbol{\theta}^*) = 1$.
- **Scenario B:** outcome spillover, without treatment spillover: $\tau_D(\boldsymbol{\theta}^*) = \beta^* = 1$, $\gamma^* = 0$, $\delta^* = 1/2$, and $\tau_I(\boldsymbol{\theta}^*) = 0.625$.
- **Scenario C:** both treatment spillover and outcome spillover: $\tau_D(\boldsymbol{\theta}^*) = \beta^* = 1$, $\gamma^* = 1$, $\delta^* = 1/2$, and $\tau_I(\boldsymbol{\theta}^*) = 2.25$.

The conditional variance is $\kappa^{*2} = 1$ in all scenarios. We generate 500 replications in each scenario. To compare the performance of estimators, we report the mean and standard deviation (S.D.) of the estimators. The simulation results are presented in Tables 1–4. The proposed estimator of the direct causal effect $\tau_D(\hat{\boldsymbol{\theta}})$ and the Hajek estimator $\hat{\tau}_D^{\text{HAJ}}$ of Li and Wager (2022) can recover the true direct causal effect in all scenarios, and the standard deviations of both estimators are similar. By contrast, the proposed estimator of the indirect causal effect $\tau_I(\hat{\boldsymbol{\theta}})$ outperforms the PC-balancing estimator $\hat{\tau}_I^{\text{PC}}$ of Li and Wager (2022) in all scenarios, both in terms of mean and variance. We compare the estimators $\tau_I(\hat{\boldsymbol{\theta}})$ and $\hat{\tau}_I^{\text{PC}}$ in scenarios A, B, and C in more detail below.

Table 1: Simulation results in scenario A: treatment spillover, without outcome spillover.

N	ρ_N	$\tau_D(\boldsymbol{\theta}^*)$	$\tau_D(\hat{\boldsymbol{\theta}}) \pm \text{S.D.}$	$\hat{\tau}_D^{\text{HAJ}} \pm \text{S.D.}$	$\tau_I(\boldsymbol{\theta}^*)$	$\tau_I(\hat{\boldsymbol{\theta}}) \pm \text{S.D.}$	$\hat{\tau}_I^{\text{PC}} \pm \text{S.D.}$
500	1	1	0.99 ± 0.09	0.99 ± 0.10	1	1.00 ± 0.14	0.71 ± 0.64
1000	1	1	1.00 ± 0.06	1.00 ± 0.06	1	0.99 ± 0.10	0.69 ± 0.65
2000	1	1	1.00 ± 0.04	1.00 ± 0.04	1	1.00 ± 0.07	0.74 ± 0.66
500	$N^{-\frac{2}{5}}$	1	0.95 ± 0.08	0.99 ± 0.10	1	1.10 ± 0.12	0.77 ± 0.23
1000	$N^{-\frac{2}{5}}$	1	0.97 ± 0.06	1.00 ± 0.07	1	1.06 ± 0.08	0.84 ± 0.19
2000	$N^{-\frac{2}{5}}$	1	0.98 ± 0.04	1.00 ± 0.05	1	1.05 ± 0.06	0.89 ± 0.16

Scenario A: treatment spillover, without outcome spillover. According to Table 1, the performance of the estimator $\hat{\tau}_I^{\text{PC}}$ is better when the interference graph is sparse ($\rho_N = N^{-\frac{2}{5}}$) than dense ($\rho_N = 1$), which aligns with the theoretical results of Li and Wager (2022, Theorem 6): For $\hat{\tau}_I^{\text{PC}}$ to be consistent, Li and Wager (2022) require $\rho_N = o(1)$. By comparison, $\tau_I(\hat{\boldsymbol{\theta}})$ is consistent in both sparse- and dense-graph settings according to Theorem 2. According to Table 1, $\tau_I(\hat{\boldsymbol{\theta}})$ outperforms $\hat{\tau}_I^{\text{PC}}$ in terms of mean and variance for small and large N , although it performs better in dense- than sparse-graph settings, which dovetails with Theorem 2: The less sparse the interference graph is, the more channels are available for spillover, which helps estimate the indirect causal effect $\tau_I(\boldsymbol{\theta}^*)$.

Table 2: Simulation results in scenario B: outcome spillover, without treatment spillover.

N	ρ_N	$\tau_D(\boldsymbol{\theta}^*)$	$\tau_D(\hat{\boldsymbol{\theta}}) \pm \text{S.D.}$	$\hat{\tau}_D^{\text{HAJ}} \pm \text{S.D.}$	$\tau_I(\boldsymbol{\theta}^*)$	$\tau_I(\hat{\boldsymbol{\theta}}) \pm \text{S.D.}$	$\hat{\tau}_I^{\text{PC}} \pm \text{S.D.}$
500	1	1	0.99 ± 0.09	1.00 ± 0.10	0.625	0.63 ± 0.16	0.09 ± 0.62
1000	1	1	1.00 ± 0.06	1.00 ± 0.06	0.625	0.62 ± 0.11	0.07 ± 0.62
2000	1	1	1.00 ± 0.04	1.00 ± 0.05	0.625	0.63 ± 0.08	0.11 ± 0.64
500	$N^{-\frac{2}{5}}$	1	0.97 ± 0.08	1.02 ± 0.10	0.625	0.57 ± 0.14	0.24 ± 0.22
1000	$N^{-\frac{2}{5}}$	1	0.98 ± 0.06	1.02 ± 0.06	0.625	0.59 ± 0.10	0.26 ± 0.18
2000	$N^{-\frac{2}{5}}$	1	0.98 ± 0.04	1.01 ± 0.05	0.625	0.62 ± 0.07	0.28 ± 0.16

Scenario B: outcome spillover, without treatment spillover. According to Table 2, the estimator $\hat{\tau}_I^{\text{PC}}$ exhibits a strong bias when there is outcome spillover ($\delta^* \neq 0$). Worse, $\hat{\tau}_I^{\text{PC}}$ suggests that there is no indirect causal effect at all, because the estimates

of $\tau_I(\boldsymbol{\theta}^*) = \gamma^* + (\beta^* + \gamma^*) a(\delta^*) = 0.625$ cluster around 0 (though the variance is large). The bias of $\hat{\tau}_I^{\text{PC}}$ stems from misspecification: It estimates the effect of treatment spillover $\gamma^* = 0$, while omitting the value $(\beta^* + \gamma^*) a(\delta^*) = 0.625$ added by outcome spillover to the indirect causal effect $\tau_I(\boldsymbol{\theta}^*)$. By contrast, the estimator $\tau_I(\hat{\boldsymbol{\theta}})$ is close to the true value of $\tau_I(\boldsymbol{\theta}^*) = 0.625$, and its variance is small.

Table 3: Simulation results in scenario C: both treatment spillover and outcome spillover.

N	ρ_N	$\tau_D(\boldsymbol{\theta}^*)$	$\tau_D(\hat{\boldsymbol{\theta}}) \pm \text{S.D.}$	$\hat{\tau}_D^{\text{HAJ}} \pm \text{S.D.}$	$\tau_I(\boldsymbol{\theta}^*)$	$\tau_I(\hat{\boldsymbol{\theta}}) \pm \text{S.D.}$	$\hat{\tau}_I^{\text{PC}} \pm \text{S.D.}$
500	1	1	1.00 ± 0.09	1.00 ± 0.10	2.25	2.23 ± 0.16	0.95 ± 0.64
1000	1	1	1.00 ± 0.06	1.00 ± 0.07	2.25	2.24 ± 0.11	0.94 ± 0.65
2000	1	1	1.00 ± 0.04	1.00 ± 0.05	2.25	2.26 ± 0.08	0.99 ± 0.66
500	$N^{-\frac{2}{5}}$	1	1.02 ± 0.08	1.08 ± 0.11	2.25	2.02 ± 0.15	1.06 ± 0.28
1000	$N^{-\frac{2}{5}}$	1	1.02 ± 0.06	1.07 ± 0.07	2.25	2.08 ± 0.10	1.15 ± 0.22
2000	$N^{-\frac{2}{5}}$	1	1.01 ± 0.04	1.04 ± 0.05	2.25	2.16 ± 0.07	1.21 ± 0.18

Scenario C: both treatment spillover and outcome spillover. If there is both treatment spillover and outcome spillover, it is natural to ask whether we can separate the effect of treatment spillover captured by γ^* from the effect of outcome spillover captured by δ^* , and whether we can estimate the added value of outcome spillover $(\beta^* + \gamma^*) a(\delta^*)$. According to Tables 3 and 4, the answer is affirmative: The means of the estimators of the effect of treatment spillover γ^* and the effect of outcome spillover δ^* are close to the truth, and the variances of the estimators decrease as N increases. In addition, the value added by outcome spillover $(\beta^* + \gamma^*) a(\delta^*)$ to the indirect causal effect $\tau_I(\boldsymbol{\theta}^*) = \gamma^* + (\beta^* + \gamma^*) a(\delta^*)$ and the total causal effect $\tau_T(\boldsymbol{\theta}^*) = \beta^* + \gamma^* + (\beta^* + \gamma^*) a(\delta^*)$ is likewise estimated well.

Variance of estimators. In all scenarios, the variance of $\tau_I(\hat{\boldsymbol{\theta}})$ is much lower than the variance of $\hat{\tau}_I^{\text{PC}}$ for any given N . The superior performance of $\tau_I(\hat{\boldsymbol{\theta}})$ compared with $\hat{\tau}_I^{\text{PC}}$ in terms of variance stems from the fact that $\hat{\tau}_I^{\text{PC}}$ is a nonparametric estimator and may require more data than the parametric estimator $\tau_I(\hat{\boldsymbol{\theta}})$, which exploits additional knowledge

Table 4: Simulation results in scenario C: both treatment spillover and outcome spillover.

N	ρ_N	γ^*	$\hat{\gamma} \pm \text{S.D.}$	δ^*	$\hat{\delta} \pm \text{S.D.}$	$(\beta^* + \gamma^*) a(\delta^*)$	$(\hat{\beta} + \hat{\gamma}) \hat{a}(\hat{\delta}) \pm \text{S.D.}$
500	1	1	1.02 ± 0.18	0.5	0.50 ± 0.05	1.25	1.22 ± 0.16
1000	1	1	1.01 ± 0.12	0.5	0.50 ± 0.03	1.25	1.22 ± 0.11
2000	1	1	1.00 ± 0.10	0.5	0.50 ± 0.02	1.25	1.25 ± 0.08
500	$N^{-\frac{2}{5}}$	1	1.06 ± 0.12	0.5	0.53 ± 0.04	1.25	0.95 ± 0.12
1000	$N^{-\frac{2}{5}}$	1	1.03 ± 0.09	0.5	0.52 ± 0.03	1.25	1.06 ± 0.09
2000	$N^{-\frac{2}{5}}$	1	1.02 ± 0.18	0.5	0.50 ± 0.05	1.25	1.14 ± 0.06

about the structure of the data-generating model compared with $\hat{\tau}_I^{\text{PC}}$.

8 Conclusion

We provide insight into the direct and indirect causal effects arising from treatment and outcome spillover, by characterizing the direct and indirect causal effects as explicit mathematical functions of the direct effect of treatment, the indirect effect of treatment via treatment spillover, and the effect of outcome spillover. Characterizing the direct and indirect causal effects helps disentangle and estimate the contributions of treatment, treatment spillover, and outcome spillover, in addition to facilitating transportability across populations, as discussed in Sections 4.1 and 4.2. To characterize direct and indirect causal effects, we have used a simple but non-trivial starting point: Gaussian Markov random fields, which can be viewed as generalizations of linear models for dependent outcomes. Two interesting directions for future research are non-Gaussian Markov random fields and semiparametric extensions.

References

- Aronow, P. M. and C. Samii (2017). Estimating average causal effects under general interference, with application to a social network experiment. *The Annals of Applied Statistics* 11, 1912 – 1947.
- Athreya, A., M. Tang, Y. Park, and C. E. Priebe (2021). On estimation and inference in latent structure random graphs. *Statistical Science* 36, 68–88.
- Bapat, R. B. (2012). *Linear algebra and linear models*. Springer Science & Business Media.
- Besag, J. (1974). Spatial interaction and the statistical analysis of lattice systems. *Journal of the Royal Statistical Society, Series B* 36, 192–225.
- Bhattacharya, S. and S. Sen (2024). Causal effect estimation under network interference with mean-field methods. *Available from: arXiv:2407.19613*.
- Brown, L. (1986). *Fundamentals of Statistical Exponential Families: With Applications in Statistical Decision Theory*. Hayworth, CA, USA: Institute of Mathematical Statistics.
- Choi, D. (2017). Estimation of monotone treatment effects in network experiments. *Journal of the American Statistical Association* 112(519), 1147–1155.
- Clark, D. A. and M. S. Handcock (2024). Causal inference over stochastic networks. *Journal of the Royal Statistical Society, Series A (Statistics in Society)* 187, 772–795.
- Ding, P. and F. Li (2018). Causal inference: A missing data perspective. *Statistical Science* 33, 214–237.
- Eckles, D., B. Karrer, and J. Ugander (2017). Design and analysis of experiments in networks: Reducing bias from interference. *Journal of Causal Inference* 5, 20150021.

- Forastiere, L., F. Mealli, A. Wu, and E. M. Airolidi (2022). Estimating causal effects under network interference with bayesian generalized propensity scores. *Journal of Machine Learning Research* 23(289), 1–61.
- Hayes, A. and K. Levin (2024). Peer effects in the linear-in-means model may be inestimable even when identified. *arXiv preprint arXiv:2410.10772*.
- Hu, Y., S. Li, and S. Wager (2022). Average direct and indirect causal effects under interference. *Biometrika* 109(4), 1165–1172.
- Hudgens, M. G. and M. E. Halloran (2008). Toward causal inference with interference. *Journal of the American Statistical Association* 103, 832–842.
- Imbens, G. W. and D. B. Rubin (2015). *Causal Inference for Statistics, Social, and Biomedical Sciences: An Introduction*. Cambridge University Press.
- Jagadeesan, R., N. S. Pillai, and A. Volfovsky (2020). Designs for estimating the treatment effect in networks with interference. *The Annals of Statistics* 48, 679 – 712.
- Karrer, B. and M. E. J. Newman (2011). Stochastic blockmodels and community structure in networks. *Physical Review E* 83, 016107.
- Kolaczyk, E. D. (2017). *Topics at the Frontier of Statistics and Network Analysis: (Re)Visiting the Foundations*. Cambridge University Press.
- Lauritzen, S. (1996). *Graphical Models*. Oxford, UK: Oxford University Press.
- Lei, J. and A. Rinaldo (2015). Consistency of spectral clustering in stochastic block models. *The Annals of Statistics* 43, 215–237.
- Li, S. and S. Wager (2022). Random graph asymptotics for treatment effect estimation under network interference. *The Annals of Statistics* 50, 2334 – 2358.

- Magnus, J. R. and H. Neudecker (1988). *Matrix Differential Calculus with Applications in Statistics and Econometrics*. New York: John Wiley & Sons.
- Ogburn, E. L., I. Shpitser, and Y. Lee (2020). Causal inference, social networks and chain graphs. *Journal of the Royal Statistical Society, Series A* 183, 1659–1676.
- Ogburn, E. L., O. Sofrygin, I. Diaz, and M. J. Van der Laan (2024). Causal inference for social network data. *Journal of the American Statistical Association* 119, 597–611.
- Pearl, J. (2009). *Causality: Models, Reasoning and Inference* (2 ed.). Cambridge: Cambridge University Press.
- Pearl, J. and E. Bareinboim (2014). External validity: From do-calculus to transportability across populations. *Statistical Science* 29, 579–595.
- Rosenbaum, P. R. and D. B. Rubin (1983). The central role of the propensity score in observational studies for causal effects. *Biometrika* 70(1), 41–55.
- Rudelson, M. and R. Vershynin (2013). Hanson-Wright inequality and sub-gaussian concentration. *Electronic Communications in Probability* 18, 1–9.
- Rue, H. and L. Held (2005). *Gaussian Markov random fields: theory and applications*. CRC press.
- Sävje, F., P. Aronow, and M. Hudgens (2021). Average treatment effects in the presence of unknown interference. *The Annals of Statistics* 49, 673.
- Schweinberger, M. and J. R. Stewart (2020). Concentration and consistency results for canonical and curved exponential-family models of random graphs. *The Annals of Statistics* 48, 374–396.

- Stewart, J. R. and M. Schweinberger (2024). Pseudo-likelihood-based M -estimators for random graphs with dependent edges and parameter vectors of increasing dimension. *Available from: arXiv:2012.07167*.
- Tang, M., D. L. Sussman, and C. E. Priebe (2013). Universally consistent vertex classification for latent positions graphs. *The Annals of Statistics* 41, 1406–1430.
- Tchetgen Tchetgen, E. J., I. R. Fulcher, and I. Shpitser (2021). Auto-G-computation of causal effects on a network. *Journal of the American Statistical Association* 116, 833–844.
- Toulis, P. and E. Kao (2013). Estimation of causal peer influence effects. In S. Dasgupta and D. McAllester (Eds.), *Proceedings of the 30th International Conference on Machine Learning*, Volume 28, pp. 1489–1497.
- van der Laan, M. J. (2014). Causal inference for a population of causally connected units. *Journal of Causal Inference* 2, 13–74.
- Vershynin, R. (2018). *High-Dimensional Probability. An Introduction with Applications in Data Science*. Cambridge, UK: Cambridge University Press.
- Xie, F. (2024). Entrywise limit theorems for eigenvectors of signal-plus-noise matrix models with weak signals. *Bernoulli* 30, 388 – 418.
- Yu, C. L., E. M. Airoldi, C. Borgs, and J. T. Chayes (2022). Estimating the total treatment effect in randomized experiments with unknown network structure. *Proceedings of the National Academy of Sciences* 119(44), e2208975119.
- Yu, Y., T. Wang, and R. J. Samworth (2015). A useful variant of the Davis–Kahan theorem for statisticians. *Biometrika* 102(2), 315–323.

SUPPLEMENTARY MATERIAL

Characterizing Direct and Indirect Causal Effects When Outcomes are Dependent due to Treatment Spillover and Outcome Spillover

9	Characterizing direct and indirect causal effects	2
9.1	Proof of Proposition 1	2
9.2	Proof of Theorem 1	4
9.3	Proof of Corollaries 1 and 2	7
10	Estimating direct and indirect causal effects	11
10.1	Estimating parameters	11
10.2	Proof of Proposition 2	16
10.3	Proof of Proposition 3	18
10.4	Proof of Proposition 4	27
10.5	Proof of Proposition 5	36
10.6	Proof of Theorem 2	39
11	Auxiliary results	45
11.1	Lemma 1	45
11.2	Lemma 2	47
11.3	Lemma 3	61
11.4	Lemma 4	66
11.5	Lemma 5	66

9 Characterizing direct and indirect causal effects

9.1 Proof of Proposition 1

We start with characterizing the direct causal effect τ_D :

$$\begin{aligned}\tau_D &:= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \mathbb{E}[Y_i(1; \mathbf{W}_{-i}) - Y_i(0; \mathbf{W}_{-i})] \\ &= \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \mathbb{E}[\mu_i(1; \mathbf{W}_{-i}, \mathbf{Z}) - \mu_i(0; \mathbf{W}_{-i}, \mathbf{Z})],\end{aligned}\tag{9.1}$$

noting that the probability law of the potential outcomes $\{\mathbf{Y}(\mathbf{w}) : \mathbf{w} \in \{0, 1\}^N\}$ is specified by the probability law of $(\mathbf{Y}, \mathbf{W}, \mathbf{Z})$ under the ignorability assumption (3).

According to Lemma 1, we can express the mean vector as

$$\boldsymbol{\mu}(\mathbf{w}, \mathbf{z}) = (\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})^{-1} \mathbf{G}(\mathbf{w}, \mathbf{z}) \boldsymbol{\vartheta} = \mathbf{M}(\delta, \mathbf{z}) \mathbf{G}(\mathbf{w}, \mathbf{z}) \boldsymbol{\vartheta},$$

where $\mathbf{M}(\delta, \mathbf{z}) := (\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})^{-1}$. Plugging the value of $\boldsymbol{\mu}(\mathbf{w}, \mathbf{z})$ into (9.1), we obtain

$$\mathbb{E}[\mu_i(1; \mathbf{W}_{-i}, \mathbf{Z}) - \mu_i(0; \mathbf{W}_{-i}, \mathbf{Z})] = (\mathbf{M}(\delta, \mathbf{Z}))_{i,\cdot}^\top \mathbb{E}[(\mathbf{G}(1; \mathbf{W}_{-i}, \mathbf{Z}) - \mathbf{G}(0; \mathbf{W}_{-i}, \mathbf{Z})) \boldsymbol{\vartheta}].$$

For any fixed i , the j -th element of $(\mathbf{G}(1; \mathbf{W}_{-i}, \mathbf{Z}) - \mathbf{G}(0; \mathbf{W}_{-i}, \mathbf{Z})) \boldsymbol{\vartheta}$ is

$$\begin{aligned}& \left(\beta W_j + \gamma \frac{\sum_{k=1}^N W_k Z_{j,k}}{\sum_{k=1}^N Z_{j,k}} \right) \Big|_{W_i=1} - \left(\beta W_j + \gamma \frac{\sum_{k=1}^N W_k Z_{j,k}}{\sum_{k=1}^N Z_{j,k}} \right) \Big|_{W_i=0} \\ &= \begin{cases} \beta & \text{if } i = j \\ \gamma \frac{Z_{i,j}}{N_j} & \text{if } i \neq j, \end{cases}\end{aligned}$$

so that

$$\mathbb{E}[\mu_i(1; \mathbf{W}_{-i}, \mathbf{Z}) - \mu_i(0; \mathbf{W}_{-i}, \mathbf{Z})] = \beta \mathbb{E}[M_{i,i}(\delta, \mathbf{Z})] + \gamma \mathbb{E} \left[\sum_{j=1: j \neq i}^N \frac{M_{i,j}(\delta, \mathbf{Z}) Z_{i,j}}{N_j} \right].$$

Therefore, the direct causal effect is

$$\tau_D = \beta \lim_{N \rightarrow \infty} \left(\frac{1}{N} \mathbb{E}[\text{trace}(\mathbf{M}(\delta, \mathbf{Z}))] \right) + \gamma \lim_{N \rightarrow \infty} \mathbb{E} \left[\frac{1}{N} \sum_{i=1}^N \sum_{j=1}^N \frac{M_{i,j}(\delta, \mathbf{Z}) Z_{i,j}}{N_j} \right],$$

provided these limits exist.

We obtain the indirect causal effect τ_I from the total causal effect $\tau_T := \tau_D + \tau_I$. Hu et al. (2022) showed that the total causal effect is

$$\tau_T = \lim_{N \rightarrow \infty} \mathbb{E}[\tau_{T,N}],$$

where

$$\tau_{T,N} := \left. \frac{d}{d\pi'} \left(\frac{1}{N} \sum_{i=1}^N \mathbb{E}_{\mathbb{P}_{\mathbf{W}, \pi'}}[Y_i] \right) \right|_{\pi' = \pi},$$

where $\mathbb{P}_{\mathbf{W}, \pi'}$ is the probability law of independent and identically distributed random variables W_1, \dots, W_N following Bernoulli(π') with $\pi' \in (0, 1)$. The expression for $\tau_{T,N}$ is obtained conditional on potential outcomes $\mathbf{Y}(\cdot) := \{Y_1(\mathbf{w}), \dots, Y_N(\mathbf{w}) : \mathbf{w} \in \{0, 1\}^N\}$ and connections \mathbf{Z} , and hence does not depend on distributional assumptions regarding $\mathbf{Y}(\cdot)$ and \mathbf{Z} . Among other things, it does not require conditional independence of potential outcomes $Y_1(\mathbf{w}), \dots, Y_N(\mathbf{w}) \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$.

Therefore, the total causal effect is

$$\tau_T = \lim_{N \rightarrow \infty} \mathbb{E}[\tau_{T,N}] = \lim_{N \rightarrow \infty} \mathbb{E}_{\mathbb{P}_{\mathbf{Y}(\cdot), \mathbf{Z}}} \left. \frac{d}{d\pi'} \left(\frac{1}{N} \sum_{i=1}^N \mathbb{E}_{\mathbb{P}_{\mathbf{W}, \pi'}}[Y_i] \right) \right|_{\pi' = \pi}$$

$$\begin{aligned}
&= \lim_{N \rightarrow \infty} \frac{d}{d\pi'} \left(\frac{1}{N} \sum_{i=1}^N \mathbb{E}_{\mathbb{P}_{Y(\cdot), \mathbf{Z}}} \mathbb{E}_{\mathbb{P}_{W, \pi'}} [Y_i] \right) \Big|_{\pi'=\pi} \\
&= \lim_{N \rightarrow \infty} \frac{d}{d\pi'} \left(\frac{1}{N} \sum_{i=1}^N \mathbb{E}_{\mathbb{P}_{\mathbf{Z}}} \mathbb{E}_{\mathbb{P}_{W, \pi'}} [\mathbf{e}_i^\top \mathbf{M}(\delta, \mathbf{Z}) \mathbf{G}(\mathbf{W}, \mathbf{Z}) \boldsymbol{\vartheta}] \right) \Big|_{\pi'=\pi} \\
&= \lim_{N \rightarrow \infty} \frac{d}{d\pi'} \left(\frac{1}{N} \mathbb{E}_{\mathbb{P}_{\mathbf{Z}}} \mathbb{E}_{\mathbb{P}_{W, \pi'}} [\mathbf{1}_N^\top \mathbf{M}(\delta, \mathbf{Z}) \mathbf{G}(\mathbf{W}, \mathbf{Z}) \boldsymbol{\vartheta}] \right) \Big|_{\pi'=\pi} \\
&= \lim_{N \rightarrow \infty} \frac{d}{d\pi'} \left(\frac{1}{N} \mathbb{E}_{\mathbb{P}_{\mathbf{Z}}} [\mathbf{1}_N^\top \mathbf{M}(\delta, \mathbf{Z}) (\beta \pi' \mathbf{1}_N + \gamma \pi' \mathbf{1}_N)] \right) \Big|_{\pi'=\pi} \\
&= \lim_{N \rightarrow \infty} \frac{d}{d\pi'} \left(\frac{1}{N} (\beta \pi' + \gamma \pi') \mathbb{E}_{\mathbb{P}_{\mathbf{Z}}} [\mathbf{1}_N^\top \mathbf{M}(\delta, \mathbf{Z}) \mathbf{1}_N] \right) \Big|_{\pi'=\pi} \\
&= (\beta + \gamma) \lim_{N \rightarrow \infty} \left(\frac{1}{N} \mathbb{E}_{\mathbb{P}_{\mathbf{Z}}} [\mathbf{1}_N^\top \mathbf{M}(\delta, \mathbf{Z}) \mathbf{1}_N] \right),
\end{aligned}$$

provided this limit exists.

9.2 Proof of Theorem 1

Define $\mathbf{M}(\delta, \mathbf{Z}) := (\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1}$ and $\mathbf{M}(\delta, \mathbf{P}) := (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1}$, where $c_N(\mathbf{Z}) := (1 / \|\mathbf{Z}\|_2) \mathbb{I}(\|\mathbf{Z}\|_2 > 0)$ and $c_N(\mathbf{P}) := 1 / \|\mathbf{P}\|_2$.

By Proposition 1, the direct effect and total causal effects are

$$\begin{aligned}
\tau_D &= \beta \lim_{N \rightarrow \infty} \left(\frac{1}{N} \mathbb{E}[\text{trace}(\mathbf{M}(\delta, \mathbf{Z}))] \right) + \gamma \lim_{N \rightarrow \infty} \mathbb{E} \left[\frac{1}{N} \sum_{i=1}^N \sum_{j=1}^N \frac{M_{i,j}(\delta, \mathbf{Z}) Z_{i,j}}{N_j} \right] \\
\tau_T &= (\beta + \gamma) \lim_{N \rightarrow \infty} \left(\frac{1}{N} \mathbf{1}_N^\top \mathbb{E}[\mathbf{M}(\delta, \mathbf{Z})] \mathbf{1}_N \right).
\end{aligned}$$

We first show that the second term in the above expansion of τ_D vanishes:

$$\begin{aligned}
&\frac{1}{N} \left| \sum_{i=1}^N \sum_{j=1}^N \frac{M_{i,j}(\delta, \mathbf{Z}) Z_{i,j}}{N_j} \right| = \frac{1}{N} \left| \sum_{j=1}^N \frac{1}{N_j} \sum_{i=1}^N M_{i,j}(\delta, \mathbf{Z}) Z_{i,j} \right| \\
&\leq \frac{1}{N} \sum_{j=1}^N \frac{1}{N_j} \left| \sum_{i=1}^N M_{i,j}(\delta, \mathbf{Z}) Z_{i,j} \right|
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{N} \sum_{j=1}^N \frac{1}{N_j} \sqrt{N_j \sum_{i=1}^N M_{i,j}(\delta, \mathbf{Z})^2} \text{ by the Cauchy-Schwarz inequality} \\
&= \frac{1}{N} \sum_{j=1}^N \frac{1}{\sqrt{N_j}} \sqrt{\sum_{i=1}^N M_{i,j}(\delta, \mathbf{Z})^2} \\
&\leq \frac{1}{N \sqrt{\min_{1 \leq j \leq N} N_j}} \sum_{j=1}^N \sqrt{\sum_{i=1}^N M_{i,j}(\delta, \mathbf{Z})^2} \\
&\leq \frac{1}{N \sqrt{\min_{1 \leq j \leq N} N_j}} \sum_{j=1}^N \max \left\{ 1, \sum_{i=1}^N M_{i,j}(\delta, \mathbf{Z})^2 \right\} \\
&\quad \text{since } x \leq \max\{1, x^2\} \text{ for all } x > 0 \\
&\leq \frac{1}{\sqrt{\min_{1 \leq j \leq N} N_j}} \max \left\{ 1, \frac{\|\mathbf{M}(\delta, \mathbf{Z})\|_F^2}{N} \right\} \\
&\leq \frac{1}{\sqrt{\min_{1 \leq j \leq N} N_j}} \max \{1, \|\mathbf{M}(\delta, \mathbf{Z})\|_2^2\}.
\end{aligned}$$

Since $|\delta| < 1$ and $c_N(\mathbf{Z}) \|\mathbf{Z}\|_2 \leq 1$, we have $\|\mathbf{M}(\delta, \mathbf{Z})\|_2 \leq 1/(1 - |\delta|)$. Thus,

$$\frac{1}{N} \left| \sum_{i=1}^N \sum_{j=1}^N \frac{M_{i,j}(\delta, \mathbf{Z}) Z_{i,j}}{N_j} \right| \leq \frac{1}{\sqrt{\min_{1 \leq j \leq N} N_j}} \frac{1}{(1 - |\delta|)^2}. \quad (9.2)$$

By convention, $Z_{i,j}/N_j := 0$ when $N_j = 0$. Leveraging part (a) of Lemma 3, we have

$$\mathbb{P} \left(\frac{1}{N} \left| \sum_{i=1}^N \sum_{j=1}^N \frac{M_{i,j}(\delta, \mathbf{Z}) Z_{i,j}}{N_j} \right| \leq \frac{1}{(1 - |\delta|)^2 \sqrt{d_1} N \rho_N} \right) \geq 1 - \frac{1}{N^2}, \quad (9.3)$$

where $d_1 := p_{\min}(1 - \sqrt{6/p_{\min} D})$. Combining (9.2) and (9.3), we obtain

$$\begin{aligned}
&\frac{1}{N} \mathbb{E} \left[\left| \sum_{i=1}^N \sum_{j=1}^N \frac{M_{i,j}(\delta, \mathbf{Z}) Z_{i,j}}{N_j} \right| \right] \\
&\leq \frac{1}{(1 - |\delta|)^2 \sqrt{d_1} N \rho_N} \left(1 - \frac{1}{N^2} \right) + \frac{1}{(1 - |\delta|)^2} \frac{1}{N^2} \rightarrow 0
\end{aligned}$$

as $N \rightarrow \infty$, since $N\rho_N \rightarrow \infty$ by Condition C.1. Part (b) of Lemma 3 implies that

$$\begin{aligned}\lim_{N \rightarrow \infty} \left(\frac{1}{N} \mathbb{E}[\text{trace}(\mathbf{M}(\delta, \mathbf{Z}))] \right) &= \lim_{N \rightarrow \infty} \left(\frac{1}{N} \text{trace}(\mathbf{M}(\delta, \mathbf{P})) \right) \\ \lim_{N \rightarrow \infty} \left(\frac{1}{N} \mathbf{1}_N^\top \mathbb{E}[\mathbf{M}(\delta, \mathbf{Z})] \mathbf{1}_N \right) &= \lim_{N \rightarrow \infty} \left(\frac{1}{N} \mathbf{1}_N^\top \mathbf{M}(\delta, \mathbf{P}) \mathbf{1}_N \right).\end{aligned}$$

Therefore,

$$\tau_D = \beta \lim_{N \rightarrow \infty} \left(\frac{1}{N} \text{trace}(\mathbf{M}(\delta, \mathbf{P})) \right) \text{ and } \tau_T = (\beta + \gamma) \lim_{N \rightarrow \infty} \left(\frac{1}{N} \mathbf{1}_N^\top \mathbf{M}(\delta, \mathbf{P}) \mathbf{1}_N \right).$$

We show that the first limit is 1. So the direct causal effect is

$$\begin{aligned}\tau_D &= \beta \lim_{N \rightarrow \infty} \left(\frac{1}{N} \text{trace}(\mathbf{M}(\delta, \mathbf{P})) \right) = \beta \lim_{N \rightarrow \infty} \left(\frac{1}{N} \sum_{i=1}^N \lambda_i(\mathbf{M}(\delta, \mathbf{P})) \right) \\ &= \beta \lim_{N \rightarrow \infty} \left(\frac{1}{N} \sum_{i=1}^N \frac{1}{1 - \delta c_N(\mathbf{P}) \lambda_i(\mathbf{P})} \right) \\ &= \beta \lim_{N \rightarrow \infty} \left(1 + \frac{1}{N} \sum_{i=1}^N \frac{\delta c_N(\mathbf{P}) \lambda_i(\mathbf{P})}{1 - \delta c_N(\mathbf{P}) \lambda_i(\mathbf{P})} \right) \\ &= \beta \left(1 + \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^K \frac{\delta c_N(\mathbf{P}) \lambda_i(\mathbf{P})}{1 - \delta c_N(\mathbf{P}) \lambda_i(\mathbf{P})} \right) \text{ since } \mathbf{P} \text{ has rank } K \\ &= \beta.\end{aligned}$$

The limit in the last step is 0 because $|\delta c_N(\mathbf{P}) \lambda_i(\mathbf{P}) / (1 - \delta c_N(\mathbf{P}) \lambda_i(\mathbf{P}))| \leq |\delta| / (1 - |\delta|)$ and K is finite. To compute the second limit, we use the Sherman–Morrison formula to obtain

$$\mathbf{M}(\delta, \mathbf{P}) = (\mathbf{I}_N - \delta c_N(\mathbf{P}) \rho_N \boldsymbol{\alpha} \boldsymbol{\alpha}^\top)^{-1} = \mathbf{I}_N + \delta \boldsymbol{\alpha} (\|\boldsymbol{\alpha}\|_2^2 \mathbf{I}_K - \delta \boldsymbol{\alpha}^\top \boldsymbol{\alpha})^{-1} \boldsymbol{\alpha}^\top,$$

noting that $c_N(\mathbf{P}) \rho_N = 1 / \|\boldsymbol{\alpha}\|_2^2$. The total causal effect is

$$\begin{aligned}
\tau_T &= (\beta + \gamma) \lim_{N \rightarrow \infty} \left(\frac{1}{N} \mathbf{1}_N^\top \mathbf{M}(\delta, \mathbf{P}) \mathbf{1}_N \right) \\
&= (\beta + \gamma) \lim_{N \rightarrow \infty} \left(\frac{1}{N} \mathbf{1}_N^\top \left(\mathbf{I}_N + \delta \boldsymbol{\alpha} (\|\boldsymbol{\alpha}\|_2^2 \mathbf{I}_K - \delta \boldsymbol{\alpha}^\top \boldsymbol{\alpha})^{-1} \boldsymbol{\alpha}^\top \right) \mathbf{1}_N \right) \\
&= (\beta + \gamma) \lim_{N \rightarrow \infty} \left(1 + \frac{\delta}{N} (\mathbf{1}_N^\top \boldsymbol{\alpha}) (\|\boldsymbol{\alpha}\|_2^2 \mathbf{I}_K - \delta \boldsymbol{\alpha}^\top \boldsymbol{\alpha})^{-1} (\boldsymbol{\alpha}^\top \mathbf{1}_N) \right) \\
&= (\beta + \gamma) (1 + a(\delta)),
\end{aligned}$$

where

$$a(\delta) := \lim_{N \rightarrow \infty} \frac{\delta}{N} (\mathbf{1}_N^\top \boldsymbol{\alpha}) (\|\boldsymbol{\alpha}\|_2^2 \mathbf{I}_K - \delta \boldsymbol{\alpha}^\top \boldsymbol{\alpha})^{-1} (\boldsymbol{\alpha}^\top \mathbf{1}_N),$$

provided this limit exists. As a consequence, the indirect causal effect is

$$\tau_I = \tau_T - \tau_D = \gamma + (\beta + \gamma) a(\delta).$$

9.3 Proof of Corollaries 1 and 2

Corollary 1 is a special case of Corollary 2. We therefore prove Corollary 2.

Let $\Xi \in \{0, 1\}^{N \times K}$ be the membership matrix corresponding to the DCBM, that is, for $1 \leq i \leq N, 1 \leq k \leq K$,

$$\Xi_{i,k} := \begin{cases} 1 & \text{if the } i\text{-th unit belongs to the } k\text{-th community} \\ 0 & \text{otherwise.} \end{cases}$$

Then the edge probability matrix is given by

$$\mathbf{P} = \rho_N \text{diag}(\boldsymbol{\eta}) \Xi \mathbf{B} \Xi^\top \text{diag}(\boldsymbol{\eta}). \quad (9.4)$$

By definition,

$$a(\delta) := \delta \lim_{N \rightarrow \infty} \frac{1}{N} (\mathbf{1}_N^\top \boldsymbol{\alpha}) (\|\boldsymbol{\alpha}\|_2^2 \mathbf{I}_K - \delta \boldsymbol{\alpha}^\top \boldsymbol{\alpha})^{-1} (\boldsymbol{\alpha}^\top \mathbf{1}_N).$$

Let $\mathbf{P} = \mathbf{U} \boldsymbol{\Lambda} \mathbf{U}^\top$ be the eigen decomposition of \mathbf{P} , where $\boldsymbol{\Lambda} = \text{diag}(\lambda_1(\mathbf{P}), \lambda_2(\mathbf{P}), \dots, \lambda_K(\mathbf{P}))$, is the diagonal matrix consisting of the non-zero eigenvalues of \mathbf{P} . We can write

$$\mathbf{P} = \rho_N \boldsymbol{\alpha} \boldsymbol{\alpha}^\top, \text{ where } \sqrt{\rho_N} \boldsymbol{\alpha} = \mathbf{U} \boldsymbol{\Lambda}^{\frac{1}{2}}.$$

Plugging in $\boldsymbol{\alpha} = (1 / \sqrt{\rho_N}) \mathbf{U} \boldsymbol{\Lambda}^{\frac{1}{2}}$, and using the fact that $\mathbf{U}^\top \mathbf{U} = \mathbf{I}_K$, we simplify $a(\delta)$ as

$$\begin{aligned} a(\delta) &= \delta \lim_{N \rightarrow \infty} \frac{1}{N} \mathbf{1}_N^\top \mathbf{U} \boldsymbol{\Lambda}^{\frac{1}{2}} (\lambda_1 \mathbf{I}_K - \delta \boldsymbol{\Lambda})^{-1} \boldsymbol{\Lambda}^{\frac{1}{2}} \mathbf{U}^\top \mathbf{1}_N \\ &= \delta \lim_{N \rightarrow \infty} \frac{1}{N} \mathbf{1}_N^\top \mathbf{U} \tilde{\boldsymbol{\Lambda}} \mathbf{U}^\top \mathbf{1}_N, \end{aligned} \tag{9.5}$$

where $\tilde{\boldsymbol{\Lambda}}$ is the diagonal matrix with diagonal elements $\lambda_k(\mathbf{P}) / (\lambda_1(\mathbf{P}) - \delta \lambda_k(\mathbf{P}))$, $1 \leq k \leq K$.

Now, using the structure of the probability matrix for the DCBM in (9.4), it is possible to characterize \mathbf{U} and $\boldsymbol{\Lambda}$ following Lei and Rinaldo (2015). Let $\boldsymbol{\eta}_{(k)}$ be the subvector of $\boldsymbol{\eta}$ containing only units in the k -th community, for all $1 \leq k \leq K$. Define

$$s_k = \|\boldsymbol{\eta}_{(k)}\|_2, \quad \mathbf{s} = (s_1, s_2, \dots, s_K).$$

Then \mathbf{U} can expressed as

$$\mathbf{U} = \text{diag}(\boldsymbol{\eta}) \boldsymbol{\Xi} \text{diag}(\mathbf{s})^{-1} \mathbf{O}, \tag{9.6}$$

where \mathbf{O} is a $K \times K$ orthogonal matrix such that

$$\tilde{\mathbf{B}} := \rho_N \text{diag}(\mathbf{s}) \mathbf{B} \text{diag}(\mathbf{s}) = \mathbf{O} \mathbf{\Lambda} \mathbf{O}^\top.$$

We calculate that

$$\mathbf{1}_N^\top \text{diag}(\boldsymbol{\eta}) \Xi \text{diag}(\mathbf{s})^{-1} = \left(\frac{\mathbf{1}_n^\top \boldsymbol{\eta}_{(1)}}{\|\boldsymbol{\eta}_{(1)}\|_2}, \frac{\mathbf{1}_n^\top \boldsymbol{\eta}_{(2)}}{\|\boldsymbol{\eta}_{(2)}\|_2}, \dots, \frac{\mathbf{1}_n^\top \boldsymbol{\eta}_{(K)}}{\|\boldsymbol{\eta}_{(K)}\|_2} \right) =: \mathbf{v}^\top. \quad (9.7)$$

Plugging (9.6) and (9.7) into (9.5), we obtain

$$a(\delta) = \delta \lim_{N \rightarrow \infty} \frac{1}{N} \mathbf{v}^\top \mathbf{O} \tilde{\mathbf{\Lambda}} \mathbf{O}^\top \mathbf{v} = \delta \lim_{N \rightarrow \infty} \frac{1}{K} \left(\frac{\mathbf{v}}{\sqrt{n}} \right)^\top \mathbf{O} \tilde{\mathbf{\Lambda}} \mathbf{O}^\top \left(\frac{\mathbf{v}}{\sqrt{n}} \right). \quad (9.8)$$

Since η_1, \dots, η_N are independent and identically distributed, the strong law of large numbers implies that

$$\frac{\mathbf{1}_n^\top \boldsymbol{\eta}_{(k)}}{\sqrt{n} \|\boldsymbol{\eta}_{(k)}\|_2} = \frac{\sum_{i: c_i=k} \eta_i}{\sqrt{n} \sum_{i: c_i=k} \eta_i^2} \xrightarrow{\text{a.s.}} \frac{\nu_1}{\sqrt{\nu_2}} \text{ for all } 1 \leq k \leq K,$$

and hence

$$\frac{\mathbf{v}}{\sqrt{n}} \xrightarrow{\text{a.s.}} \frac{\nu_1}{\sqrt{\nu_2}} \mathbf{1}_K. \quad (9.9)$$

Next, observe that since $\tilde{\mathbf{\Lambda}} := \mathbf{\Lambda}^{\frac{1}{2}} (\lambda_1 \mathbf{I}_K - \delta \mathbf{\Lambda})^{-1} \mathbf{\Lambda}^{\frac{1}{2}}$ and $\tilde{\mathbf{B}} := \mathbf{O} \mathbf{\Lambda} \mathbf{O}^\top$, we have

$$\mathbf{O} \tilde{\mathbf{\Lambda}} \mathbf{O}^\top = \tilde{\mathbf{B}}^{\frac{1}{2}} (\|\tilde{\mathbf{B}}\|_2 \mathbf{I}_K - \delta \tilde{\mathbf{B}})^{-1} \tilde{\mathbf{B}}^{\frac{1}{2}}.$$

By strong law

$$\frac{1}{\rho_N} \frac{\tilde{\mathbf{B}}}{n \nu_2} = \frac{\text{diag}(\mathbf{s})}{\sqrt{n} \nu_2} \mathbf{B} \frac{\text{diag}(\mathbf{s})}{\sqrt{n} \nu_2} \xrightarrow{\text{a.s.}} \mathbf{B}.$$

Noting that $\mathbf{O} \tilde{\mathbf{\Lambda}} \mathbf{O}^\top$ does not change if $\tilde{\mathbf{B}}$ is scaled by a constant, we have

$$\mathbf{O} \tilde{\mathbf{\Lambda}} \mathbf{O}^\top \xrightarrow{\text{a.s.}} \mathbf{B}^{\frac{1}{2}} (\|\mathbf{B}\|_2 \mathbf{I}_K - \delta \mathbf{B})^{-1} \mathbf{B}^{\frac{1}{2}}. \quad (9.10)$$

From (9.8), (9.9) and (9.10), we conclude that $a(\delta)$ exists almost surely, and is given by

$$a(\delta) = \delta \frac{\nu_1^2}{\nu_2} \frac{1}{K} \mathbf{1}_K^\top \mathbf{B}^{\frac{1}{2}} (\|\mathbf{B}\|_2 \mathbf{I}_K - \delta \mathbf{B})^{-1} \mathbf{B}^{\frac{1}{2}} \mathbf{1}_K.$$

We consider the case where $\mathbf{B} = (p - q) \mathbf{I}_K + q \mathbf{1}_K \mathbf{1}_K^\top$, $0 \leq q < p \leq 1$. Then, some elementary matrix calculations yield that the eigenvalues of \mathbf{B} :

$$\lambda_j(\mathbf{B}) = \begin{cases} p + (K - 1)q & j = 1 \\ p - q & j = 2, \dots, K. \end{cases}$$

Moreover, $\mathbf{1}_K$ is an eigenvector of \mathbf{B} corresponding to the eigenvalue $\lambda_1(\mathbf{B})$, which is unique.

Next, we note that \mathbf{B} and $\mathbf{B}^{\frac{1}{2}} (\|\mathbf{B}\|_2 \mathbf{I}_K - \delta \mathbf{B})^{-1} \mathbf{B}^{\frac{1}{2}}$ have the same set of eigenvectors, and if λ is eigenvalue of \mathbf{B} , then $\lambda / (\|\mathbf{B}\|_2 - \delta \lambda)$ is the corresponding eigenvalue of $\mathbf{B}^{\frac{1}{2}} (\|\mathbf{B}\|_2 - \delta \mathbf{B})^{-1} \mathbf{B}^{\frac{1}{2}}$. Hence, $\mathbf{1}_K$ is an eigenvector of $\mathbf{B}^{\frac{1}{2}} (\|\mathbf{B}\|_2 - \delta \mathbf{B})^{-1} \mathbf{B}^{\frac{1}{2}}$, and the corresponding eigenvalue is

$$\frac{\lambda_1(\mathbf{B})}{\|\mathbf{B}\|_2 - \delta \lambda_1(\mathbf{B})} = \frac{1}{1 - \delta},$$

since $\|\mathbf{B}\|_2 = \lambda_1(\mathbf{B})$ by definition. Therefore,

$$\begin{aligned} a(\delta) &= \delta \frac{\nu_1^2}{\nu_2} \frac{1}{K} \mathbf{1}_K^\top \mathbf{B}^{\frac{1}{2}} (\|\mathbf{B}\|_2 \mathbf{I}_K - \delta \mathbf{B})^{-1} \mathbf{B}^{\frac{1}{2}} \mathbf{1}_K \\ &= \frac{\delta}{1 - \delta} \frac{\nu_1^2}{\nu_2} \frac{1}{K} \mathbf{1}_K^\top \mathbf{1}_K \\ &= \frac{\delta}{1 - \delta} \frac{\nu_1^2}{\nu_2}. \end{aligned}$$

10 Estimating direct and indirect causal effects

The specification of the Gaussian Markov random field in Section 2 implies that the resulting probability density function belongs to a regular exponential family of probability densities in the sense of Brown (1986, p. 2). Plugging the values of $\boldsymbol{\mu}(\mathbf{w}, \mathbf{z})$ and $\boldsymbol{\Omega}(\mathbf{w}, \mathbf{z})$ into the probability density function of the multivariate Gaussian, the conditional probability density of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ can be written as

$$f_{\mathbf{y}|\mathbf{w},\mathbf{z}}(\mathbf{y} \mid \mathbf{w}, \mathbf{z}) = \psi(\boldsymbol{\theta}, \kappa^2) b(\mathbf{y}) \exp(-\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2)^\top t(\mathbf{y})), \quad (10.1)$$

where $\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2) := (\beta/\kappa^2, \gamma/\kappa^2, \delta/\kappa^2, 1/\kappa^2)^\top$ and

$$\begin{aligned} \psi(\boldsymbol{\theta}, \kappa^2) &:= \sqrt{\frac{\det(\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})}{(2\pi\kappa^2)^N}} \\ &\times \exp\left(-\frac{1}{2\kappa^2} \boldsymbol{\vartheta}^\top \mathbf{G}(\mathbf{w}, \mathbf{z})^\top (\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})^{-1} \mathbf{G}(\mathbf{w}, \mathbf{z}) \boldsymbol{\vartheta}\right) \\ t(\mathbf{y}) &:= \left(\mathbf{y}^\top \mathbf{G}(\mathbf{w}, \mathbf{z}), \frac{c_N(\mathbf{z})}{2} \mathbf{y}^\top \mathbf{z} \mathbf{y}, -\frac{\mathbf{y}^\top \mathbf{y}}{2}\right)^\top \\ b(\mathbf{y}) &:= 1. \end{aligned}$$

We will use the exponential family form above in the proof of Lemma 2 and Proposition 4.

10.1 Estimating parameters

10.1.1 Estimating $\boldsymbol{\theta}$ and κ^2

We state theoretical guarantees for maximum likelihood estimators $\hat{\boldsymbol{\theta}}$ and $\hat{\kappa}^2$ of $\boldsymbol{\theta}$ and κ^2 , which pave the way for proving Theorem 2.

The loglikelihood of $\boldsymbol{\theta}$ and κ^2 is

$$\begin{aligned}
\ell(\boldsymbol{\theta}, \kappa^2) &= -\frac{N}{2} \log(\kappa^2) + \frac{1}{2} \log(\det(\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})) - \frac{1}{2\kappa^2} \mathbf{y}^\top \mathbf{y} \\
&\quad - \frac{1}{2\kappa^2} \boldsymbol{\vartheta}^\top \mathbf{G}(\mathbf{w}, \mathbf{z})^\top (\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})^{-1} \mathbf{G}(\mathbf{w}, \mathbf{z}) \boldsymbol{\vartheta} \\
&\quad + \frac{\delta c_N(\mathbf{z})}{2\kappa^2} \mathbf{y}^\top \mathbf{z} \mathbf{y} + \frac{\boldsymbol{\vartheta}^\top}{\kappa^2} \mathbf{G}(\mathbf{w}, \mathbf{z})^\top \mathbf{y}.
\end{aligned} \tag{10.2}$$

The set of maximum likelihood estimators can be expressed as

$$(\widehat{\boldsymbol{\theta}}, \widehat{\kappa}^2) := \{(\boldsymbol{\theta}, \kappa^2) \in \boldsymbol{\Theta} \times (0, \infty) : \|\nabla_{\boldsymbol{\theta}, \kappa^2} \ell(\boldsymbol{\theta}, \kappa^2)\|_2 = 0\}.$$

To establish convergence rates for $\widehat{\boldsymbol{\theta}}$ along with $\tau_D(\widehat{\boldsymbol{\theta}})$ and $\tau_I(\widehat{\boldsymbol{\theta}})$, we first derive convergence rates for the natural parameter vector $\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2)$ of the multivariate Gaussian, which is defined by

$$\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2) := (\beta/\kappa^2, \gamma/\kappa^2, \delta/\kappa^2, 1/\kappa^2)^\top.$$

We then obtain convergence rates for $\widehat{\boldsymbol{\theta}}$ based on $\boldsymbol{\eta}(\widehat{\boldsymbol{\theta}}, \widehat{\kappa}^2)$. To obtain convergence rates for $\widehat{\boldsymbol{\theta}}$ and $\boldsymbol{\eta}(\widehat{\boldsymbol{\theta}}, \widehat{\kappa}^2)$, note that consistency is possible on a subset of well-behaved $(\mathbf{w}, \mathbf{z}) \in \{0, 1\}^N \times \{0, 1\}^{\binom{N}{2}}$, but it is not possible for all $(\mathbf{w}, \mathbf{z}) \in \{0, 1\}^N \times \{0, 1\}^{\binom{N}{2}}$: e.g., when none of the units receives treatment and hence \mathbf{w} is an N -dimensional vector of 0's, the model parameters β and γ do not show up in $\ell(\boldsymbol{\theta}, \kappa^2)$.

To introduce a subset of well-behaved (\mathbf{w}, \mathbf{z}) , let $\mathbf{H}(\mathbf{w}, \mathbf{z})$ be the projection matrix onto the column space of $\mathbf{G}(\mathbf{w}, \mathbf{z})$. Define the following subsets of $(\mathbf{w}, \mathbf{z}) \in \{0, 1\}^N \times \{0, 1\}^{\binom{N}{2}}$:

$$\begin{aligned}
\mathcal{A}_{1,N} &:= \left\{ (\mathbf{w}, \mathbf{z}) : \sum_j z_{i,j} \geq d_1 N \rho_N \text{ for all } i, \|\mathbf{z} - \mathbf{P}\|_2 \leq l \sqrt{N \rho_N} \right\} \\
\mathcal{A}_{2,N} &:= \left\{ (\mathbf{w}, \mathbf{z}) : \left| \frac{\sum_j w_j z_{i,j}}{\sum_j z_{i,j}} - \pi \right| \leq \sqrt{\frac{3 \log N}{\sum_j z_{i,j}}} \text{ for all } i, \left| \frac{\sum_j w_j}{N} - \pi \right| \leq \sqrt{\frac{2 \log N}{N}} \right\}
\end{aligned}$$

$$\mathcal{A}_{3,N} := \left\{ (\mathbf{w}, \mathbf{z}) : \|(\mathbf{I}_N - \mathbf{H}(\mathbf{w}, \mathbf{z})) \mathbf{P} \mathbf{v}\|_2 \geq \frac{1 - \sqrt{(1 + p^*)/2}}{2} \|\mathbf{P} \mathbf{v}\|_2 \text{ for all } \mathbf{v} \in \mathbb{R}^N \right\}$$

$$\mathcal{A}_{4,N} := \left\{ (\mathbf{w}, \mathbf{z}) : \left\| \mathbf{z} \mathbf{G}(\mathbf{w}, \mathbf{z}) \begin{pmatrix} a \\ b \end{pmatrix} \right\|_2 \geq \frac{|a + b| \pi p_{\min}}{4} \sqrt{N} N \rho_N \right\},$$

where $d_1 := p_{\min} (1 - \sqrt{6/p_{\min} D})$, $p^* \in (0, 1)$, $l > 0$, and $a, b \in \mathbb{R}$ are constants.

Proposition 3 proves that the event $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_N := \bigcap_{k=1}^4 \mathcal{A}_{k,N}$ occurs with high probability provided N is large enough.

Proposition 3. Let $W_i \stackrel{\text{iid}}{\sim} \text{Bernoulli}(\pi)$ with $\pi \in (0, 1)$ and $Z_{i,j} \stackrel{\text{iid}}{\sim} \text{Bernoulli}(P_{i,j})$ with $P_{i,j} := \rho_N \boldsymbol{\alpha}_i^\top \boldsymbol{\alpha}_j$ satisfy Condition C.1. If there exists a constant $p^* \in (0, 1)$ such that $(\mathbf{1}_N^\top \mathbf{u})^2 / N \leq p^*$ for all unit vectors \mathbf{u} in the column space of \mathbf{P} and the constant D in Condition C.1 satisfies $D \geq 96 (1 + 2\sqrt{2})^2 / (p_{\min} (1 - \sqrt{(1 + p^*)/2})^2 \pi^3 (1 - \pi))$, then

$$\mathbb{P}((\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N} \cap \mathcal{A}_{2,N} \cap \mathcal{A}_{3,N}) \geq 1 - \frac{\tilde{C}_2}{N^2}$$

provided N is large enough, where $\tilde{C}_2 > 0$ is a constant independent of N . If, in addition, $a + b \neq 0$ and $D \geq 96 (|a| + 2\sqrt{(a^2 + b^2)/p_{\min}})^2 / ((a + b)^2 \pi^2 p_{\min}^2)$, then

$$\mathbb{P}((\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_N) \geq 1 - \frac{\tilde{C}_2}{N^2}.$$

Proposition 3 is proved in Supplement 10.3. We establish convergence rates for $\boldsymbol{\eta}(\hat{\boldsymbol{\theta}}, \hat{\kappa}^2)$ in the event $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_N$, which occurs with high probability according to Proposition 3. The following lemma bounds the spectral norm of the inverse Fisher information matrix,

defined by

$$\mathcal{I}(\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2)) := -\mathbb{E}_{\boldsymbol{\theta}, \kappa^2} \left[\nabla_{\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2)}^2 \ell(\boldsymbol{\theta}, \kappa^2) \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z}) \right].$$

Lemma 2. Assume that $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z})$ is generated by a Gaussian Markov random field with parameters $\boldsymbol{\theta} \in \boldsymbol{\Theta}$ and $\kappa^2 \in (0, \infty)$, $W_i \stackrel{\text{iid}}{\sim} \text{Bernoulli}(\pi)$ with $\pi \in (0, 1)$, and $Z_{i,j} \stackrel{\text{iid}}{\sim} \text{Bernoulli}(P_{i,j})$ with $P_{i,j} := \rho_N \boldsymbol{\alpha}_i^\top \boldsymbol{\alpha}_j$ subject to Condition C.1. If $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_N$, the constant D in Condition C.1 satisfies $D \geq 1536 (1 + 2\pi)^2 / (p_{\min} \pi^6 (1 - \pi)^2)$, and $\beta + \gamma \neq 0$, then $\mathcal{I}(\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2))$ is invertible at $\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2)$, and there exists a constant $L(\boldsymbol{\theta}, \kappa^2) > 0$, which depends on $\boldsymbol{\theta}$ and κ^2 but does not depend on N , such that

$$\|\mathcal{I}(\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2))^{-1}\|_2 \leq L(\boldsymbol{\theta}, \kappa^2) \frac{1}{c_N(\mathbf{Z})^2 N (N \rho_N)^2},$$

where

$$\begin{aligned} L(\boldsymbol{\theta}, \kappa^2) &:= \frac{Q_1(\boldsymbol{\theta}, \kappa^2)}{Q_2(\boldsymbol{\theta}, \kappa^2)} \\ Q_1(\boldsymbol{\theta}, \kappa^2) &:= \left[\frac{\kappa^2}{1 + |\delta|} \left(\sqrt{2} + \frac{2\sqrt{2} \|\boldsymbol{\vartheta}\|_2}{1 - |\delta|} \right)^2 + \frac{\kappa^4}{(1 - |\delta|)^2} \right]^3 \\ Q_2(\boldsymbol{\theta}, \kappa^2) &:= \frac{\kappa^{10} (\beta + \gamma)^2 p_{\min}^2}{(1 + |\delta|)^7} \frac{\pi^5 (1 - \pi)}{8192} (1 - \sqrt{(1 + p^*)/2})^2. \end{aligned}$$

Lemma 2 is proved in Supplement 11.2. Lemma 2 helps establish convergence rates. To state them, let $\boldsymbol{\theta}^* := (\beta^*, \gamma^*, \delta^*)$ and κ^{*2} be the data-generating parameters, and assume that $\beta^* + \gamma^* \neq 0$ and $|\delta^*| < 1$. It can be shown that, for all

$$\epsilon^* \in \left(0, \frac{1}{\kappa^{*2}} \min \left\{ |\beta^* + \gamma^*|, \frac{1 - |\delta^*|}{3} \right\} \right), \quad (10.3)$$

we have $L(\boldsymbol{\theta}, \kappa^2) < \infty$ for all $\boldsymbol{\theta} \in \boldsymbol{\Theta}$ and all $\kappa^2 \in (0, \infty)$ for which $\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2) \in \mathcal{B}(\boldsymbol{\eta}(\boldsymbol{\theta}^*, \kappa^{*2}), \epsilon^*)$, and the Fisher information matrix $\mathcal{I}(\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2))$ is invertible at $\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2)$. The following result establishes convergence rates for $\boldsymbol{\eta}(\widehat{\boldsymbol{\theta}}, \widehat{\kappa}^2)$ by bounding

$$\Lambda_N(\boldsymbol{\theta}^*, \kappa^{*2}) := \sup_{\boldsymbol{\eta} \in \mathcal{B}(\boldsymbol{\eta}(\boldsymbol{\theta}^*, \kappa^{*2}), \epsilon^*)} \|\mathcal{I}(\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2))^{-1}\|_2.$$

We then obtain the convergence rates for $\widehat{\boldsymbol{\theta}}$ based on $\boldsymbol{\eta}(\widehat{\boldsymbol{\theta}}, \widehat{\kappa}^2)$.

Proposition 4. Let the assumptions of Lemma 2 be satisfied and consider the event $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_N$. If $c_N(\mathbf{Z})^2 (N\rho_N)^2 \sqrt{N/\log N} \rightarrow \infty$ as $N \rightarrow \infty$, there exist constants $C_1(\boldsymbol{\theta}^*, \kappa^{*2}) > 0$ and $C_2(\boldsymbol{\theta}^*, \kappa^{*2}) > 0$, which depend on $\boldsymbol{\theta}^*$ and κ^{*2} but do not depend on N , such that

$$\begin{aligned} \|\boldsymbol{\eta}(\widehat{\boldsymbol{\theta}}, \widehat{\kappa}^2) - \boldsymbol{\eta}(\boldsymbol{\theta}^*, \kappa^{*2})\|_2 &\leq C_1(\boldsymbol{\theta}^*, \kappa^{*2}) \frac{1}{c_N(\mathbf{Z})^2 (N\rho_N)^2} \sqrt{\frac{\log N}{N}} \\ \|\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_2 &\leq C_2(\boldsymbol{\theta}^*, \kappa^{*2}) \frac{1}{c_N(\mathbf{Z})^2 (N\rho_N)^2} \sqrt{\frac{\log N}{N}} \end{aligned}$$

with probability at least $1 - 8/N^2$, provided N is large enough.

Proposition 4 is proved in Supplement 10.4.

10.1.2 Estimating $a(\delta)$

We state theoretical guarantees for the estimator $\widehat{a}(\delta)$ of $a(\delta)$ to pave the way for proving Theorem 2. For any δ such that $|\delta| < 1$, define

$$a_N(\delta) := \frac{\delta}{N} (\mathbf{1}_N^\top \mathbf{T}) (\|\mathbf{T}\|_2^2 \mathbf{I}_K - \delta \mathbf{T}^\top \mathbf{T})^{-1} (\mathbf{T}^\top \mathbf{1}_N),$$

so that $a(\delta) = \lim_{N \rightarrow \infty} a_N(\delta)$. We have the following result.

Proposition 5. Assume that Condition C.1 holds and that $|\delta| < 1$. Then there exists a constant $C_a(\delta) > 0$, which is a non-decreasing function of $|\delta|$ that does not depend on N , such that

$$|\widehat{a}(\delta) - a_N(\delta)| \leq C_a(\delta) g(\mathbf{Z}),$$

and there exists a constant $\widetilde{C}_1 > 0$ such that

$$\mathbb{P}\left(g(\mathbf{Z}) \leq \sqrt{\frac{\log N}{N\rho_N}}\right) \geq 1 - \frac{\widetilde{C}_1}{N^2},$$

where $g : \{0, 1\}^{\binom{N}{2}} \mapsto (0, \infty)$ is a function of interference graph $\mathbf{Z} \in \{0, 1\}^{\binom{N}{2}}$.

Proposition 5 is proved in Supplement 10.5.

10.2 Proof of Proposition 2

Based on Corollary 1, we know that for the rank-one case,

$$a_N(\delta^\star) = \frac{\delta^\star}{1 - \delta^\star} \frac{(\sum_{i=1}^N \alpha_i)^2}{N \sum_{i=1}^N \alpha_i^2} \quad \text{and} \quad a(\delta^\star) = \frac{\delta^\star}{1 - \delta^\star} \frac{\nu_1^2}{\nu_2}.$$

Since $|\delta^\star| < 1$,

$$\begin{aligned} |a_N(\delta^\star) - a(\delta^\star)| &= \frac{|\delta^\star|}{|1 - \delta^\star|} \left| \frac{(\sum_{i=1}^N \alpha_i)^2}{N \sum_{i=1}^N \alpha_i^2} - \frac{\nu_1^2}{\nu_2} \right| \\ &\leq \frac{1}{1 - |\delta^\star|} \left| \frac{\sum_{i=1}^N \alpha_i}{\sqrt{N \sum_{i=1}^N \alpha_i^2}} - \frac{\nu_1}{\sqrt{\nu_2}} \right| \left| \frac{\sum_{i=1}^N \alpha_i}{\sqrt{N \sum_{i=1}^N \alpha_i^2}} + \frac{\nu_1}{\sqrt{\nu_2}} \right| \\ &\leq \frac{2}{1 - |\delta^\star|} \left| \frac{\sum_{i=1}^N \alpha_i}{\sqrt{N \sum_{i=1}^N \alpha_i^2}} - \frac{\nu_1}{\sqrt{\nu_2}} \right|, \end{aligned} \tag{10.4}$$

noting that $\sum_{i=1}^N \alpha_i / \sqrt{N \sum_{i=1}^N \alpha_i^2}$ and $\nu_1 / \sqrt{\nu_2}$ are bounded above by 1.

$$\begin{aligned}
& \left| \frac{\sum_{i=1}^N \alpha_i}{\sqrt{N \sum_{i=1}^N \alpha_i^2}} - \frac{\nu_1}{\sqrt{\nu_2}} \right| \\
& \leq \left| \frac{\sum_{i=1}^N (\alpha_i - \nu_1)}{\sqrt{N \sum_{i=1}^N \alpha_i^2}} \right| + \left| \frac{\sqrt{N} \nu_1}{\sqrt{\sum_{i=1}^N \alpha_i^2}} - \frac{\nu_1}{\sqrt{\nu_2}} \right| \\
& = \frac{|\sum_{i=1}^N (\alpha_i - \nu_1)|}{\sqrt{N \sum_{i=1}^N \alpha_i^2}} + \nu_1 \left| \frac{N \nu_2 - \sum_{i=1}^N \alpha_i^2}{\sqrt{\sum_{i=1}^N \alpha_i^2} \sqrt{\nu_2} \left(\sqrt{N \nu_2} + \sqrt{\sum_{i=1}^N \alpha_i^2} \right)} \right| \\
& = \frac{|\sum_{i=1}^N (\alpha_i - \nu_1)/N|}{\sqrt{\sum_{i=1}^N \alpha_i^2/N}} + \frac{\nu_1 |\sum_{i=1}^N (\alpha_i^2 - \nu_2)/N|}{\sqrt{\sum_{i=1}^N \alpha_i^2/N} \sqrt{\nu_2} \left(\nu_2 + \sqrt{\sum_{i=1}^N \alpha_i^2/N} \right)}.
\end{aligned} \tag{10.5}$$

Since the random variables $\alpha_1, \dots, \alpha_N$ are Subgaussian with parameter σ^2 , we have

$$\mathbb{P} \left(\left| \frac{1}{N} \sum_{i=1}^N \alpha_i - \nu_1 \right| > t \right) \leq 2 \exp \left(-\frac{N t^2}{2 \sigma^2} \right). \tag{10.6}$$

The assumption that the random variables $\alpha_1, \dots, \alpha_N$ are Subgaussian implies that the random variables $\alpha_1^2, \dots, \alpha_N^2$ are Subexponential with parameters α^2 and λ (depending on σ^2), so that

$$\mathbb{P}(|\alpha_i^2 - \nu_2| > t) \leq 2 \exp \left(-\frac{1}{2} \min \left\{ \frac{t^2}{\alpha^2}, \frac{t}{\lambda} \right\} \right), \quad i = 1, \dots, N.$$

This in turn implies that

$$\mathbb{P} \left(\left| \frac{1}{N} \sum_{i=1}^N \alpha_i^2 - \nu_2 \right| > t \right) \leq 2 \exp \left(-\frac{N}{2} \min \left\{ \frac{t^2}{\alpha^2}, \frac{t}{\lambda} \right\} \right). \tag{10.7}$$

From (10.6) and (10.7), we conclude that there exists a constant $\tilde{C} > 0$ such that

$$\left| \frac{1}{N} \sum_{i=1}^N \alpha_i - \nu_1 \right|, \quad \left| \frac{1}{N} \sum_{i=1}^N \alpha_i^2 - \nu_2 \right| \leq \tilde{C} \sqrt{\frac{\log N}{N}},$$

with probability at least $1 - 4/N^2$. Incorporating this result into (10.5), we obtain

$$\begin{aligned} & \left| \frac{\sum_{i=1}^N \alpha_i}{\sqrt{N \sum_{i=1}^N \alpha_i^2}} - \frac{\nu_1}{\sqrt{\nu_2}} \right| \\ & \leq \frac{\tilde{C} \sqrt{\log N / N}}{\sqrt{\nu_2 - \tilde{C} \sqrt{\log N / N}}} + \frac{\nu_1 \tilde{C} \sqrt{\log N / N}}{\sqrt{\nu_2 - \tilde{C} \sqrt{\log N / N}} \nu_2^{3/2}}, \end{aligned} \tag{10.8}$$

with probability at least $1 - 4/N^2$.

Combining (10.4) and (10.8), we conclude that there exists a constant $C(\delta^*) > 0$ such that for all sufficiently large N ,

$$|a_N(\delta^*) - a(\delta^*)| \leq C(\delta^*) \sqrt{\frac{\log N}{N}},$$

with probability at least $1 - 4/N^2$.

10.3 Proof of Proposition 3

Bounding $\mathbb{P}((\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N})$. Lemma 3 implies that $\min_{1 \leq i \leq N} \sum_{j=1}^N Z_{i,j} > d_1 N \rho_N$ with probability at least $1 - 1/N^2$. Also, we showed in the proof of Lemma 3 (see Equation (11.47)) that under Condition C.1, there exists a constant $l > 0$ such that $\|\mathbf{Z} - \mathbf{P}\|_2 \leq l \sqrt{N \rho_N}$ with probability at least $1 - 1/N^2$.

Therefore,

$$\mathbb{P}((\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N}) \geq 1 - \frac{2}{N^2}.$$

Bounding $\mathbb{P}((\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{2,N})$. We invoke the Hoeffding's inequality to obtain

$$\mathbb{P}_{\mathbf{W}} \left(\left| \frac{\sum_{j=1}^N W_j Z_{i,j}}{\sum_{j=1}^N Z_{i,j}} - \pi \right| \leq \frac{t}{\sum_{j=1}^N Z_{i,j}} \right) \geq 1 - 2 \exp \left(-\frac{t^2}{\sum_{j=1}^N Z_{i,j}} \right).$$

Choosing $t := \sqrt{3 \log N \sum_{j=1}^N Z_{i,j}}$, and taking a union bound over all $i \in \{1, \dots, N\}$, we obtain

$$\mathbb{P} \left(\bigcap_{i=1}^N \left\{ \left| \frac{\sum_{j=1}^N W_j Z_{i,j}}{\sum_{j=1}^N Z_{i,j}} - \pi \right| \leq \sqrt{\frac{3 \log N}{\sum_{j=1}^N Z_{i,j}}} \right\} \right) \geq 1 - \frac{2}{N^2}.$$

Hoeffding's inequality also implies that

$$\mathbb{P} \left(\left| \frac{\sum_{j=1}^N W_j}{N} - \pi \right| \leq \sqrt{\frac{2 \log N}{N}} \right) \geq 1 - \frac{2}{N^2}.$$

Combining these two results gives

$$\mathbb{P}((\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{2,N}) \geq 1 - \frac{4}{N^2}.$$

Bounding $\mathbb{P}((\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{3,N})$. The matrix $\mathbf{H}(\mathbf{W}, \mathbf{Z})$ is defined as the projection matrix onto the column space of $\mathbf{G}(\mathbf{W}, \mathbf{Z})$, where the i -th row of $\mathbf{G}(\mathbf{W}, \mathbf{Z})$ is given by $(W_i, \sum_{j=1}^N W_j Z_{i,j} / \sum_{j=1}^N Z_{i,j})$. Let us define a new matrix $\overline{\mathbf{G}}(\mathbf{W})$ such that its i -th row is (W_i, π) . Then

$$\|\mathbf{G}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{G}}(\mathbf{W})\|_F^2 = \sum_{i=1}^N \left(\frac{\sum_{j=1}^N W_j Z_{i,j}}{\sum_{j=1}^N Z_{i,j}} - \pi \right)^2 \leq \frac{3 \log N}{d_1 \rho_N}, \quad (10.9)$$

provided $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N} \cap \mathcal{A}_{2,N}$. Let $\overline{\mathbf{H}}(\mathbf{W})$ be the projection matrix onto the column space of $\overline{\mathbf{G}}(\mathbf{W})$. To bound the probability of event $\mathcal{A}_{3,N}$, we first show that if $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N} \cap \mathcal{A}_{2,N}$, the projection matrices $\mathbf{H}(\mathbf{W}, \mathbf{Z})$ and $\overline{\mathbf{H}}(\mathbf{W})$ are close in terms of the

spectral norm.

Bounding $\|H(\mathbf{W}, \mathbf{Z}) - \overline{H}(\mathbf{W})\|_2$. The Davis-Kahan theorem (Theorem 1, Yu et al. (2015)) gives

$$\|H(\mathbf{W}, \mathbf{Z}) - \overline{H}(\mathbf{W})\|_2 \leq \frac{\|G(\mathbf{W}, \mathbf{Z}) G(\mathbf{W}, \mathbf{Z})^\top - \overline{G}(\mathbf{W}) \overline{G}(\mathbf{W})^\top\|_F}{\lambda_{\min}(\overline{G}(\mathbf{W})^\top \overline{G}(\mathbf{W}))}. \quad (10.10)$$

The numerator on the right-hand side above,

$$\begin{aligned} & \|G(\mathbf{W}, \mathbf{Z}) G(\mathbf{W}, \mathbf{Z})^\top - \overline{G}(\mathbf{W}) \overline{G}(\mathbf{W})^\top\|_F \\ & \leq \|G(\mathbf{W}, \mathbf{Z}) - \overline{G}(\mathbf{W})\|_F^2 + 2 \|\overline{G}(\mathbf{W})\|_F \|G(\mathbf{W}, \mathbf{Z}) - \overline{G}(\mathbf{W})\|_F \\ & \leq \frac{3 \log N}{d_1 \rho_N} + 2 \sqrt{2N} \sqrt{\frac{3 \log N}{d_1 \rho_N}}, \end{aligned}$$

leveraging (10.9) and noting that $\|\overline{G}(\mathbf{W})\|_F \leq \sqrt{2N}$. Since $N\rho_N \geq D \log N$ by Condition C.1, we obtain

$$\|G(\mathbf{W}, \mathbf{Z}) G(\mathbf{W}, \mathbf{Z})^\top - \overline{G}(\mathbf{W}) \overline{G}(\mathbf{W})^\top\|_F \leq N \left(\frac{3}{d_1 D} + 2\sqrt{2} \sqrt{\frac{3}{d_1 D}} \right).$$

Recall that $d_1 := p_{\min}(1 - \sqrt{6/p_{\min}D})$. The condition $D \geq 96(1 + 2\sqrt{2})^2/p_{\min}((1 - \sqrt{(1+p^*)/2})^2 \pi^3(1 - \pi))$ implies that $D > 24/p_{\min}$, so that we have $d_1 > p_{\min}/2$. Hence,

$$\begin{aligned} & \|G(\mathbf{W}, \mathbf{Z}) G(\mathbf{W}, \mathbf{Z})^\top - \overline{G}(\mathbf{W}) \overline{G}(\mathbf{W})^\top\|_F \\ & \leq N \left(\frac{6}{p_{\min} D} + 2\sqrt{2} \sqrt{\frac{6}{p_{\min} D}} \right) \leq N(1 + 2\sqrt{2}) \sqrt{\frac{6}{p_{\min} D}}. \end{aligned} \quad (10.11)$$

To bound the denominator on the right-hand side of (10.10), note that

$$\overline{\mathbf{G}}(\mathbf{W})^\top \overline{\mathbf{G}}(\mathbf{W}) = \begin{pmatrix} \sum_{i=1}^N W_i & \pi \sum_{i=1}^N W_i \\ \pi \sum_{i=1}^N W_i & N\pi^2 \end{pmatrix},$$

which implies,

$$\begin{aligned} \lambda_{\min}(\overline{\mathbf{G}}(\mathbf{W})^\top \overline{\mathbf{G}}(\mathbf{W})) &\geq \sqrt{\det(\overline{\mathbf{G}}(\mathbf{W})^\top \overline{\mathbf{G}}(\mathbf{W}))} \\ &= \sqrt{\pi^2 \sum_{i=1}^N W_i \left(N - \sum_{i=1}^N W_i \right)}. \end{aligned}$$

If $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{2,N}$, then we have

$$\begin{aligned} &\lambda_{\min}(\overline{\mathbf{G}}(\mathbf{W})^\top \overline{\mathbf{G}}(\mathbf{W})) \\ &\geq \sqrt{\pi^2 (N - \sqrt{2N \log N}) (N(1 - \pi) - \sqrt{2N \log N})} \\ &= \sqrt{\pi^2 N^2 \left(\pi - \sqrt{\frac{2 \log N}{N}} \right) \left((1 - \pi) - \sqrt{\frac{2 \log N}{N}} \right)} \\ &\geq N\pi \sqrt{\frac{\pi(1 - \pi)}{4}} \end{aligned} \tag{10.12}$$

for all $N \geq N_1$, choosing N_1 so that $\sqrt{2 \log N/N} \leq \min\{\pi, 1 - \pi\}/2$ for all $N \geq N_1$.

Combining (10.10), (10.11) and (10.12), we obtain that if $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N} \cap \mathcal{A}_{2,N}$, then

$$\begin{aligned} \|\mathbf{H}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{H}}(\mathbf{W})\|_2 &\leq N(1 + 2\sqrt{2}) \sqrt{\frac{6}{p_{\min} D}} \cdot \frac{1}{N\pi} \sqrt{\frac{4}{\pi(1 - \pi)}} \\ &\leq \frac{2(1 + 2\sqrt{2})}{\pi \sqrt{\pi(1 - \pi)}} \sqrt{\frac{6}{p_{\min} D}}. \end{aligned} \tag{10.13}$$

Now, we proceed to establish a lower bound on $\|(\mathbf{I}_N - \mathbf{H}(\mathbf{W}, \mathbf{Z})) \mathbf{P} \mathbf{v}\|_2$:

$$\begin{aligned}
& \|(\mathbf{I}_N - \mathbf{H}(\mathbf{W}, \mathbf{Z})) \mathbf{P} \mathbf{v}\|_2 \\
& \geq \|(\mathbf{I}_N - \overline{\mathbf{H}}(\mathbf{W})) \mathbf{P} \mathbf{v}\|_2 - \|(\mathbf{H}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{H}}(\mathbf{W})) \mathbf{P} \mathbf{v}\|_2 \\
& \geq \|\mathbf{P} \mathbf{v}\|_2 - \|\overline{\mathbf{H}}(\mathbf{W}) \mathbf{P} \mathbf{v}\|_2 - \|(\mathbf{H}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{H}}(\mathbf{W})) \mathbf{P} \mathbf{v}\|_2.
\end{aligned} \tag{10.14}$$

Bounding $\|\overline{\mathbf{H}}(\mathbf{W}) \mathbf{P} \mathbf{v}\|_2$. $\overline{\mathbf{H}}(\mathbf{W})$ is the projection matrix onto the column space of $\overline{\mathbf{G}}(\mathbf{W})$, that is, the space spanned by the vectors $\mathbf{1}_N$ and \mathbf{W} . Using Gram-Schmidt orthogonalization, we obtain that the column space of $\overline{\mathbf{G}}(\mathbf{W})$ is also spanned by the orthogonal vectors $\mathbf{1}_N$ and \mathbf{U} , where

$$\mathbf{U} := \left(\mathbf{I}_N - \frac{\mathbf{1}_N \mathbf{1}_N^\top}{N} \right) \mathbf{W}.$$

So, we can write the projection matrix $\overline{\mathbf{H}}(\mathbf{W})$ as the sum of rank-one projection matrices corresponding to $\mathbf{1}_N$ and \mathbf{U} ,

$$\overline{\mathbf{H}}(\mathbf{W}) = \frac{\mathbf{1}_N \mathbf{1}_N^\top}{N} + \frac{\mathbf{U} \mathbf{U}^\top}{\mathbf{U}^\top \mathbf{U}}.$$

Then

$$\begin{aligned}
\|\overline{\mathbf{H}}(\mathbf{W}) \mathbf{P} \mathbf{v}\|_2^2 &= \left\| \frac{\mathbf{1}_N \mathbf{1}_N^\top}{N} \mathbf{P} \mathbf{v} \right\|_2^2 + \left\| \frac{\mathbf{U} \mathbf{U}^\top}{\mathbf{U}^\top \mathbf{U}} \mathbf{P} \mathbf{v} \right\|_2^2 \\
&= \frac{((\mathbf{P} \mathbf{v})^\top \mathbf{1}_N)^2}{N} + \frac{((\mathbf{P} \mathbf{v})^\top \mathbf{U})^2}{\mathbf{U}^\top \mathbf{U}}.
\end{aligned} \tag{10.15}$$

By assumption, there exists a constant $p^* \in (0, 1)$ such that the first term on the right-hand side of above,

$$\frac{((\mathbf{P} \mathbf{v})^\top \mathbf{1}_N)^2}{N} \leq p^* \|\mathbf{P} \mathbf{v}\|_2^2. \tag{10.16}$$

For the second term, note that the denominator,

$$\mathbf{U}^\top \mathbf{U} = \mathbf{W} \left(\mathbf{I}_N - \frac{\mathbf{1}_N \mathbf{1}_N^\top}{N} \right) \mathbf{W} = \frac{\sum_{i=1}^N W_i (N - \sum_{i=1}^N W_i)}{N} \geq \frac{N \pi (1 - \pi)}{4} \quad (10.17)$$

for all $N \geq N_1$, provided that $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{2,N}$, as shown in (10.12). Also, the numerator $(\mathbf{P} \mathbf{v})^\top \mathbf{U}$ is linear in \mathbf{W} , with coefficients $(\mathbf{I}_N - (\mathbf{1}_N \mathbf{1}_N^\top)/N) \mathbf{P} \mathbf{v} = \mathbf{P} \mathbf{v} - \overline{\mathbf{P} \mathbf{v}} \mathbf{1}_N$, where $\overline{\mathbf{P} \mathbf{v}}$ is the mean of $(\mathbf{P} \mathbf{v})_i$'s. The coefficients are bounded above in magnitude by $|(\mathbf{P} \mathbf{v})_i - \overline{\mathbf{P} \mathbf{v}}|$. We compute

$$\mathbb{E}[(\mathbf{P} \mathbf{v})^\top \mathbf{U}] = (\mathbf{P} \mathbf{v})^\top \left(\mathbf{I}_N - \frac{\mathbf{1}_N \mathbf{1}_N^\top}{N} \right) \mathbb{E}[\mathbf{W}] = (\mathbf{P} \mathbf{v})^\top \left(\mathbf{I}_N - \frac{\mathbf{1}_N \mathbf{1}_N^\top}{N} \right) \pi \mathbf{1}_N = 0.$$

So, by applying Hoeffding's inequality, we obtain

$$\mathbb{P} \left(|(\mathbf{P} \mathbf{v})^\top \mathbf{U}| \leq \sqrt{2 \left(\sum_{i=1}^N ((\mathbf{P} \mathbf{v})_i - \overline{\mathbf{P} \mathbf{v}})^2 \right) \log N} \right) \geq 1 - \frac{2}{N^2}. \quad (10.18)$$

Combining (10.15), (10.16), (10.17) and (10.18), we obtain that with probability at least $1 - 6/N^2$ (recall that we also required $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{2,N}$),

$$\begin{aligned} \|\overline{\mathbf{H}}(\mathbf{W}) \mathbf{P} \mathbf{v}\|_2^2 &\leq p^* \|\mathbf{P} \mathbf{v}\|_2^2 + \frac{8 \log N}{N \pi (1 - \pi)} \left(\sum_{i=1}^N ((\mathbf{P} \mathbf{v})_i - \overline{\mathbf{P} \mathbf{v}})^2 \right) \\ &\leq p^* \|\mathbf{P} \mathbf{v}\|_2^2 + \frac{8 \log N}{N \pi (1 - \pi)} \|\mathbf{P} \mathbf{v}\|_2^2 \\ &\leq (p^* + (1 - p^*)/2) \|\mathbf{P} \mathbf{v}\|_2^2 \\ &\leq \frac{1 + p^*}{2} \|\mathbf{P} \mathbf{v}\|_2^2. \end{aligned} \quad (10.19)$$

for all $N \geq N_2$, where N_2 ensures that

$$\frac{8 \log N}{N \pi (1 - \pi)} \leq \frac{1 - p^*}{2}$$

for all $N \geq N_2$.

Bounding $\|(\mathbf{H}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{H}}(\mathbf{W})) \mathbf{P} \mathbf{v}\|_2$. We showed in (10.13) that if $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N} \cap \mathcal{A}_{2,N}$, then

$$\|\mathbf{H}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{H}}(\mathbf{W})\|_2 \leq \frac{2(1 + 2\sqrt{2})}{\pi \sqrt{\pi(1 - \pi)}} \sqrt{\frac{6}{p_{\min} D}}.$$

This implies,

$$\|(\mathbf{H}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{H}}(\mathbf{W})) \mathbf{P} \mathbf{v}\|_2 \leq \frac{2(1 + 2\sqrt{2})}{\pi \sqrt{\pi(1 - \pi)}} \sqrt{\frac{6}{p_{\min} D}} \|\mathbf{P} \mathbf{v}\|_2.$$

Therefore, for any $\epsilon_1 > 0$, if $D \geq 24(1 + 2\sqrt{2})^2 / (\epsilon_1^2 p_{\min} \pi^3 (1 - \pi))$, then

$$\mathbb{P}(\|(\mathbf{H}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{H}}(\mathbf{W})) \mathbf{P} \mathbf{v}\|_2 \leq \epsilon_1 \|\mathbf{P} \mathbf{v}\|_2) \geq 1 - \frac{6}{N^2}. \quad (10.20)$$

Combining (10.14), (10.19) and (10.20), we obtain that with probability at least $1 - 12/N^2$,

$$\begin{aligned} \|(\mathbf{I}_N - \mathbf{H}(\mathbf{W}, \mathbf{Z})) \mathbf{P} \mathbf{v}\|_2 &\geq (1 - \sqrt{(1 + p^*)/2} - \epsilon_1) \|\mathbf{P} \mathbf{v}\|_2 \\ &\geq \frac{1 - \sqrt{(1 + p^*)/2}}{2} \|\mathbf{P} \mathbf{v}\|_2, \end{aligned} \quad (10.21)$$

choosing $\epsilon_1 := (1 - \sqrt{(1 + p^*)/2})/2$. Therefore,

$$\mathbb{P}((\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{3,N}) \geq 1 - \frac{12}{N^2}.$$

Bounding $\mathbb{P}((\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{4,N})$. Define $c := a + b \neq 0$ and $\mathbf{q} := \begin{pmatrix} a \\ b \end{pmatrix}$.

$$\begin{aligned} & \|\mathbf{Z} \mathbf{G}(\mathbf{W}, \mathbf{Z}) \mathbf{q}\|_2 \\ & \geq \|\mathbf{Z} \overline{\mathbf{G}}(\mathbf{W}) \mathbf{q}\|_2 - \|\mathbf{Z} (\mathbf{G}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{G}}(\mathbf{W})) \mathbf{q}\|_2 \\ & = \left\| \mathbf{Z} \overline{\mathbf{G}}(\mathbf{W}) \begin{pmatrix} a \\ c-a \end{pmatrix} \right\|_2 - \|\mathbf{Z} (\mathbf{G}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{G}}(\mathbf{W})) \mathbf{q}\|_2 \\ & = |c| \pi \left\| \mathbf{Z} \overline{\mathbf{G}}(\mathbf{W}) \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\|_2 - \left\| \mathbf{Z} \overline{\mathbf{G}}(\mathbf{W}) \begin{pmatrix} a \\ -a \end{pmatrix} \right\|_2 - \|\mathbf{Z} (\mathbf{G}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{G}}(\mathbf{W})) \mathbf{q}\|_2. \end{aligned}$$

Recalling that the second column of $\overline{\mathbf{G}}(\mathbf{W})$ is $\mathbf{1}_N$, we have

$$\begin{aligned} \|\mathbf{Z} \mathbf{G}(\mathbf{W}, \mathbf{Z}) \mathbf{q}\|_2 & \geq |c| \pi \|\mathbf{Z} \mathbf{1}_N\|_2 - |a| \left\| \mathbf{Z} \overline{\mathbf{G}}(\mathbf{W}) \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\|_2 \\ & \quad - \|\mathbf{Z} (\mathbf{G}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{G}}(\mathbf{W})) \mathbf{q}\|_2. \end{aligned} \tag{10.22}$$

We show that the first term on the right-hand side of (10.22) dominates the second and the third term.

Concerning the first term, note that $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N}$ implies that

$$\|\mathbf{Z} \mathbf{1}_N\|_2 = \sqrt{\sum_{i=1}^N \left(\sum_{j=1}^N Z_{i,j} \right)^2} \geq d_1 \sqrt{N} N \rho_N. \quad (10.23)$$

The second term is bounded above by

$$\begin{aligned} \left\| \mathbf{Z} \overline{\mathbf{G}}(\mathbf{W}) \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\|_2 &= \sqrt{\sum_{i=1}^N \left(\sum_{j=1}^N W_j Z_{i,j} - \pi \sum_{j=1}^N Z_{i,j} \right)^2} \\ &\leq \sqrt{3 \log N \sum_{i=1}^N \sum_{j=1}^N Z_{i,j}} \\ &\leq \sqrt{3 N \log N \|\mathbf{Z}\|_2}, \end{aligned} \quad (10.24)$$

where the first inequality holds if $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{2,N}$.

The third term is bounded above by

$$\begin{aligned} \|\mathbf{Z} (\mathbf{G}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{G}}(\mathbf{W})) \mathbf{q}\|_2 &= \|\mathbf{q}\|_2 \|\mathbf{Z}\|_2 \|\mathbf{G}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{G}}(\mathbf{W})\|_2 \\ &\leq \|\mathbf{q}\|_2 \sqrt{\frac{3 \log N}{d_1 \rho_N}} \|\mathbf{Z}\|_2, \end{aligned} \quad (10.25)$$

incorporating (10.9), which holds if $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N} \cap \mathcal{A}_{2,N}$.

Now, if $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N}$,

$$\|\mathbf{Z}\|_2 \leq \|\mathbf{P}\|_2 + \|\mathbf{Z} - \mathbf{P}\|_2 \leq N \rho_N + l \sqrt{N \rho_N} \leq 2 N \rho_N, \quad (10.26)$$

for all sufficiently large N since $N \rho_N \rightarrow \infty$ by Condition C.1.

Collecting (10.23), (10.24), (10.25) and (10.26), we have that with probability at least

$$1 - 6/N^2,$$

$$\begin{aligned}
|c| \pi \|\mathbf{Z} \mathbf{1}_N\|_2 &\geq |c| \pi d_1 \sqrt{N} N \rho_N, \\
\left\| \mathbf{Z} \overline{\mathbf{G}}(\mathbf{W}) \begin{pmatrix} 1 \\ -1 \end{pmatrix} \right\|_2 &\leq \sqrt{6 N^2 \rho_N \log N} \leq \sqrt{\frac{6}{D}} \sqrt{N} N \rho_N, \\
\|\mathbf{Z} (\mathbf{G}(\mathbf{W}, \mathbf{Z}) - \overline{\mathbf{G}}(\mathbf{W})) \mathbf{q}\|_2 &\leq 2 \|\mathbf{q}\|_2 \sqrt{\frac{3 \rho_N \log N}{d_1}} N \leq 2 \|\mathbf{q}\|_2 \sqrt{\frac{3}{d_1 D}} \sqrt{N} N \rho_N,
\end{aligned} \tag{10.27}$$

applying the inequality $N \rho_N \geq D \log N$ from Condition C.1.

Finally, recall that $d_1 > p_{\min}/2$ because $D > 24/p_{\min}$. Thus, combining (10.22) and (10.27), we obtain that with probability at least $1 - 6/N^2$,

$$\begin{aligned}
\|\mathbf{Z} \mathbf{G}(\mathbf{W}, \mathbf{Z}) \mathbf{q}\|_2 &\geq \left(\frac{|c| \pi p_{\min}}{2} - |a| \sqrt{\frac{6}{D}} - 2 \|\mathbf{q}\|_2 \sqrt{\frac{6}{p_{\min} D}} \right) \sqrt{N} N \rho_N \\
&\geq \frac{|c| \pi p_{\min}}{4} \sqrt{N} N \rho_N,
\end{aligned}$$

provided that $D \geq 96 (|a| + 2 \|\mathbf{q}\|_2 / \sqrt{p_{\min}})^2 / (c^2 \pi^2 p_{\min}^2)$.

Therefore,

$$\mathbb{P}((\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{4,N}) \geq 1 - \frac{6}{N^2}.$$

10.4 Proof of Proposition 4

The parameter vector of primary interest is $\boldsymbol{\theta} := (\beta, \gamma, \delta)^\top$. To establish convergence rates for the maximum likelihood estimator $\hat{\boldsymbol{\theta}}$ of the data-generating parameter vector $\boldsymbol{\theta}^*$, we take advantage of recent advances in statistical exponential-family theory (Stewart and Schweinberger, 2024). Since multivariate Gaussians are statistical exponential families, there are two parameterizations that are more convenient to work with than the $\boldsymbol{\theta}$ -parameterization: the

natural and mean-value parameterization. The natural parameter vector of the multivariate Gaussian is defined by

$$\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2) := (\beta/\kappa^2, \gamma/\kappa^2, \delta/\kappa^2, 1/\kappa^2)^\top.$$

We henceforth suppress the notational dependence of $\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2)$ on $(\boldsymbol{\theta}, \kappa^2)$ and write $\boldsymbol{\eta}$ instead of $\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2)$, and define $\hat{\boldsymbol{\eta}} := \boldsymbol{\eta}(\hat{\boldsymbol{\theta}}, \hat{\kappa}^2)$ and $\boldsymbol{\eta}^* := \boldsymbol{\eta}(\boldsymbol{\theta}^*, \kappa^{*2})$. In accordance, we write $\Lambda_N(\boldsymbol{\eta}^*) := \Lambda_N(\boldsymbol{\theta}^*, \kappa^{*2})$. The mean-value parameter vector of the multivariate Gaussian is defined by

$$\boldsymbol{\mu}(\boldsymbol{\eta}) := \mathbb{E}_{\boldsymbol{\eta}} t(\mathbf{Y}),$$

where $t(\mathbf{Y})$ is the vector of sufficient statistics defined at the beginning of Supplement 10. Let \mathbb{N} be the natural parameter space and $\mathbb{M} := \boldsymbol{\mu}(\mathbb{N})$ be the mean-value parameter space of the statistical exponential family; note that the symbol \mathbb{N} should not be confused with the set of natural numbers. The map $\boldsymbol{\mu} : \mathbb{N} \mapsto \mathbb{M}$ is a homeomorphism (Brown, 1986, Theorem 3.6, p. 74), so the inverse map $\boldsymbol{\mu}^{-1} : \mathbb{M} \mapsto \mathbb{N}$ exists and $\boldsymbol{\mu}$ and $\boldsymbol{\mu}^{-1}$ are continuous, one-to-one, and onto. Throughout,

$$\mathcal{B}(\mathbf{c}, \rho) := \{\mathbf{v} \in \mathbb{R}^4 : \|\mathbf{v} - \mathbf{c}\|_2 \leq \rho\}$$

denotes a ℓ_2 -ball in \mathbb{R}^4 with center $\mathbf{c} \in \mathbb{R}^4$ and radius $\rho \in (0, \infty)$. We first establish convergence rates for $\hat{\boldsymbol{\eta}}$ and then establish convergence rates for $\hat{\boldsymbol{\theta}}$. The convergence rates for $\hat{\boldsymbol{\eta}}$ are based on the proof of Theorem 1 of Stewart and Schweinberger (2024), but we consider real-valued outcomes $Y_i \in \mathbb{R}$ rather than binary connections $Z_{i,j} \in \{0, 1\}$, and we extend the proof of Theorem 1 of Stewart and Schweinberger (2024) from natural parameters to continuous functions of natural parameters using ideas from Section 3.1 of

Schweinberger and Stewart (2020).

Idea of proof. Let

$$\epsilon^* := \frac{1}{\kappa^{*2}} \min \left\{ |\beta^* + \gamma^*|, \frac{1 - |\delta^*|}{3} \right\} \quad (10.28)$$

and consider any $\epsilon \in (0, \epsilon^*)$. We will show at the end of the proof that the restriction to the subset $\mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*) \subset \mathbb{N}$ is legitimate, because $\hat{\boldsymbol{\eta}}$ falls into $\mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*) \subset \mathbb{N}$ with high probability, provided that N is large enough.

Define

$$\varphi(\epsilon) := \sup \{ \varphi \in (0, \infty) : \mathcal{B}(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \varphi) \subseteq \boldsymbol{\mu}(\mathcal{B}(\boldsymbol{\eta}^*, \epsilon)) \}. \quad (10.29)$$

By the definition of $\varphi(\epsilon)$, $\mathcal{B}(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \varphi(\epsilon))$ is the largest ball contained in the image $\boldsymbol{\mu}(\mathcal{B}(\boldsymbol{\eta}^*, \epsilon))$ of $\mathcal{B}(\boldsymbol{\eta}^*, \epsilon)$, which implies that

$$\varphi(\epsilon) = \sup \{ \varphi \in (0, \infty) : \boldsymbol{\mu}^{-1}(\mathcal{B}(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \varphi)) \subseteq \mathcal{B}(\boldsymbol{\eta}^*, \epsilon) \}.$$

The definition of $\varphi(\epsilon)$ implies the following fact:

$$\text{If } \varphi \in (0, \infty) \text{ guarantees that } \boldsymbol{\mu}^{-1}(\mathcal{B}(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \varphi)) \subseteq \mathcal{B}(\boldsymbol{\eta}^*, \epsilon), \quad (10.30)$$

$$\text{then } \varphi \leq \varphi(\epsilon).$$

The key of the proof of Proposition 4 is to determine which real numbers $\varphi \in (0, \infty)$ guarantee that all elements $\boldsymbol{\mu}' \in \mathcal{B}(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \varphi)$ map to elements $\boldsymbol{\eta}' \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon)$. We will

prove the following result:

$$\text{All } \varphi \leq \frac{\epsilon}{\sup_{\boldsymbol{\eta} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*)} \|\mathcal{I}(\boldsymbol{\eta})^{-1}\|_2} \text{ ensure that } \boldsymbol{\mu}^{-1}(\mathcal{B}(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \varphi)) \subseteq \mathcal{B}(\boldsymbol{\eta}^*, \epsilon),$$

which implies that

$$\boldsymbol{\mu}^{-1} \left(\mathcal{B} \left(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \frac{\epsilon}{\sup_{\boldsymbol{\eta} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*)} \|\mathcal{I}(\boldsymbol{\eta})^{-1}\|_2} \right) \right) \subseteq \mathcal{B}(\boldsymbol{\eta}^*, \epsilon). \quad (10.31)$$

In light of Equation (10.30), Equation (10.31) implies that

$$\frac{\epsilon}{\sup_{\boldsymbol{\eta} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*)} \|\mathcal{I}(\boldsymbol{\eta})^{-1}\|_2} \leq \varphi(\epsilon). \quad (10.32)$$

To prove Equation (10.31), note that $\mathcal{B}(\boldsymbol{\eta}^*, \epsilon) \subset \mathbb{N}$ implies that $\boldsymbol{\mu}(\mathcal{B}(\boldsymbol{\eta}^*, \epsilon)) \subset \mathbb{M}$, and the maximum likelihood estimator $\hat{\boldsymbol{\eta}}$ exists, is unique, and solves $\boldsymbol{\mu}(\hat{\boldsymbol{\eta}}) = t(\mathbf{Y})$ in the event $t(\mathbf{Y}) \in \boldsymbol{\mu}(\mathcal{B}(\boldsymbol{\eta}^*, \epsilon)) \subset \mathbb{M}$ (Brown, 1986, Theorem 5.5, p. 148). Thus, there exist constants $B_1(\boldsymbol{\eta}^*) > 0$ and $B_2(\boldsymbol{\eta}^*) > 0$ such that the probability of event $\hat{\boldsymbol{\eta}} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon)$ is bounded below by

$$\begin{aligned} & \mathbb{P}_{\mathbf{y}|\mathbf{w}, \mathbf{z}}(\hat{\boldsymbol{\eta}} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon)) \\ &= \mathbb{P}_{\mathbf{y}|\mathbf{w}, \mathbf{z}}(t(\mathbf{Y}) \in \boldsymbol{\mu}(\mathcal{B}(\boldsymbol{\eta}^*, \epsilon))) \\ &\geq \mathbb{P}_{\mathbf{y}|\mathbf{w}, \mathbf{z}}(t(\mathbf{Y}) \in \mathcal{B}(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \varphi(\epsilon))) \\ &\geq 1 - 8 \exp \left(- \min \left\{ \frac{B_1(\boldsymbol{\eta}^*) \varphi(\epsilon)^2}{N}, B_2(\boldsymbol{\eta}^*) \varphi(\epsilon) \right\} \right), \end{aligned} \quad (10.33)$$

where

- the second line follows from the fact that $\boldsymbol{\mu} : \boldsymbol{\eta} \mapsto \mathbb{M}$ is a homeomorphism (Brown, 1986, Theorem 3.6, p. 74) and $\hat{\boldsymbol{\eta}}$ exists, is unique, and solves $\boldsymbol{\mu}(\hat{\boldsymbol{\eta}}) = t(\mathbf{Y})$ in the event $t(\mathbf{Y}) \in \boldsymbol{\mu}(\mathcal{B}(\boldsymbol{\eta}^*, \epsilon)) \subset \mathbb{M}$ (Brown, 1986, Theorem 5.5, p. 148);

- the third line follows from the fact that $\mathcal{B}(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \varphi(\epsilon)) \subseteq \boldsymbol{\mu}(\mathcal{B}(\boldsymbol{\eta}^*, \epsilon))$, which holds by definition of $\varphi(\epsilon)$ in Equation (10.29);
- the fourth line follows from $\boldsymbol{\mu}(\hat{\boldsymbol{\eta}}) = t(\mathbf{Y})$ in the event $t(\mathbf{Y}) \in \boldsymbol{\mu}(\mathcal{B}(\boldsymbol{\eta}^*, \epsilon)) \subset \mathbb{M}$ (Brown, 1986, Theorem 5.5, p. 148) along with Lemma 5 in Supplement 11.5.

We bound the right-hand side of Equation (10.33) by bounding $\varphi(\epsilon)$.

Bounding the right-hand side of Equation (10.33). Consider an arbitrary $\varphi \in (0, \infty)$ and an arbitrary $\boldsymbol{\mu}' \in \mathcal{B}(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \varphi)$, and let $\boldsymbol{\eta}' := \boldsymbol{\mu}^{-1}(\boldsymbol{\mu}')$ be the parameter vector corresponding to $\boldsymbol{\mu}'$; note that there is one and only one $\boldsymbol{\eta}'$ corresponding to $\boldsymbol{\mu}'$, because $\boldsymbol{\mu}$ and $\boldsymbol{\mu}^{-1}$ are one-to-one. We will determine which real numbers $\varphi \in (0, \infty)$ satisfy $\varphi \leq \varphi(\epsilon)$ by using the multivariate mean-value theorem. The multivariate mean-value theorem implies that there exists a real number $\lambda \in (0, 1)$ and a parameter vector

$$\boldsymbol{\eta}'' := \lambda \boldsymbol{\eta}^* + (1 - \lambda) \boldsymbol{\eta}' \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*)$$

such that

$$\boldsymbol{\mu}(\boldsymbol{\eta}') - \boldsymbol{\mu}(\boldsymbol{\eta}^*) = \mathcal{I}(\boldsymbol{\eta}'') (\boldsymbol{\eta}' - \boldsymbol{\eta}^*). \quad (10.34)$$

By Lemma 2 and its discussion, $\mathcal{I}(\boldsymbol{\eta})$ is invertible for all $\boldsymbol{\eta} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*)$, including $\boldsymbol{\eta}'' \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*)$. Since $\mathcal{I}(\boldsymbol{\eta}'')$ is invertible, Equation (10.34) implies that

$$\boldsymbol{\eta}' - \boldsymbol{\eta}^* = \mathcal{I}(\boldsymbol{\eta}'')^{-1} (\boldsymbol{\mu}(\boldsymbol{\eta}') - \boldsymbol{\mu}(\boldsymbol{\eta}^*)), \quad (10.35)$$

which in turn implies that

$$\|\boldsymbol{\eta}' - \boldsymbol{\eta}^*\|_2 \leq \|\mathcal{I}(\boldsymbol{\eta}'')^{-1}\|_2 \|\boldsymbol{\mu}(\boldsymbol{\eta}') - \boldsymbol{\mu}(\boldsymbol{\eta}^*)\|_2. \quad (10.36)$$

To obtain an upper bound on the right-hand side of Equation (10.36), observe that

$$\|\boldsymbol{\eta}' - \boldsymbol{\eta}^*\|_2 \leq \|\mathcal{I}(\boldsymbol{\eta}'')^{-1}\|_2 \|\boldsymbol{\mu}(\boldsymbol{\eta}') - \boldsymbol{\mu}(\boldsymbol{\eta}^*)\|_2 < \|\mathcal{I}(\boldsymbol{\eta}'')^{-1}\|_2 \varphi, \quad (10.37)$$

because $\boldsymbol{\mu}(\boldsymbol{\eta}') = \boldsymbol{\mu}'$ and $\boldsymbol{\mu}' \in \mathcal{B}(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \varphi)$ implies that $\|\boldsymbol{\mu}' - \boldsymbol{\mu}(\boldsymbol{\eta}^*)\|_2 < \varphi$. To establish Equation (10.31), we want to bound the right-hand side of Equation (10.37) from above by ϵ , that is, we want to determine which real numbers $\varphi \in (0, \infty)$ ensure that

$$\|\boldsymbol{\eta}' - \boldsymbol{\eta}^*\|_2 \leq \|\mathcal{I}(\boldsymbol{\eta}'')^{-1}\|_2 \|\boldsymbol{\mu}(\boldsymbol{\eta}') - \boldsymbol{\mu}(\boldsymbol{\eta}^*)\|_2 < \|\mathcal{I}(\boldsymbol{\eta}'')^{-1}\|_2 \varphi \leq \epsilon. \quad (10.38)$$

It is evident that the inequality in Equation (10.38) holds as long as

$$\varphi \leq \frac{\epsilon}{\sup_{\boldsymbol{\eta} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*)} \|\mathcal{I}(\boldsymbol{\eta})^{-1}\|_2}. \quad (10.39)$$

Since $\boldsymbol{\mu}'$ was arbitrary, Equations (10.38) and (10.39) prove that each element

$$\boldsymbol{\mu}' \in \mathcal{B}\left(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \frac{\epsilon}{\sup_{\boldsymbol{\eta} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*)} \|\mathcal{I}(\boldsymbol{\eta})^{-1}\|_2}\right)$$

pulls back to an element

$$\boldsymbol{\eta}' \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon),$$

which establishes

$$\boldsymbol{\mu}^{-1}\left(\mathcal{B}\left(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \frac{\epsilon}{\sup_{\boldsymbol{\eta} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*)} \|\mathcal{I}(\boldsymbol{\eta})^{-1}\|_2}\right)\right) \subseteq \mathcal{B}(\boldsymbol{\eta}^*, \epsilon). \quad (10.40)$$

To relate ϵ to $\varphi(\epsilon)$, recall that the definition of $\varphi(\epsilon)$ in Equation (10.29) implies the following

fact:

$$\text{If } \varphi \in (0, \infty) \text{ guarantees that } \boldsymbol{\mu}^{-1}(\mathcal{B}(\boldsymbol{\mu}(\boldsymbol{\eta}^*), \varphi)) \subseteq \mathcal{B}(\boldsymbol{\eta}^*, \epsilon), \quad (10.41)$$

$$\text{then } \varphi \leq \varphi(\epsilon).$$

Combining Equations (10.40) and (10.41) hence establishes

$$\frac{\epsilon}{\sup_{\boldsymbol{\eta} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*)} \|\mathcal{I}(\boldsymbol{\eta})^{-1}\|_2} \leq \varphi(\epsilon). \quad (10.42)$$

Armed with Equation (10.42), we revisit Equation (10.33) to obtain

$$\begin{aligned} \mathbb{P}(\widehat{\boldsymbol{\eta}} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon)) &\geq 1 - 8 \exp \left(- \min \left\{ \frac{B_1(\boldsymbol{\eta}^*) \varphi(\epsilon)^2}{N}, B_2(\boldsymbol{\eta}^*) \varphi(\epsilon) \right\} \right) \\ &\geq 1 - 8 \exp \left(- \min \left\{ \frac{B_1(\boldsymbol{\eta}^*) \epsilon^2}{\Lambda_N(\boldsymbol{\eta}^*)^2 N}, \frac{B_2(\boldsymbol{\eta}^*) \epsilon}{\Lambda_N(\boldsymbol{\eta}^*)} \right\} \right), \end{aligned}$$

where

$$\Lambda_N(\boldsymbol{\eta}^*) := \sup_{\boldsymbol{\eta} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*)} \|\mathcal{I}(\boldsymbol{\eta})^{-1}\|_2.$$

Choose

$$\epsilon := \sqrt{\frac{2}{B_1(\boldsymbol{\eta}^*)}} \Phi_N(\boldsymbol{\eta}^*), \quad (10.43)$$

where

$$\Phi_N(\boldsymbol{\eta}^*) := \Lambda_N(\boldsymbol{\eta}^*) \sqrt{N \log N},$$

and note that $\epsilon \in (0, \epsilon^*)$ for all large enough N by Equation (10.45). Then there exists an integer $N_1 \geq 1$ such that, for all $N \geq N_1$,

$$\mathbb{P} \left(\widehat{\boldsymbol{\eta}} \in \mathcal{B} \left(\boldsymbol{\eta}^*, \sqrt{\frac{2}{B_1(\boldsymbol{\eta}^*)}} \Phi_N(\boldsymbol{\eta}^*) \right) \right) \geq 1 - \frac{8}{N^2}, \quad (10.44)$$

because

$$\min \left\{ \frac{B_1(\boldsymbol{\eta}^*) \epsilon^2}{\Lambda_N(\boldsymbol{\eta}^*)^2 N}, \frac{B_2(\boldsymbol{\eta}^*) \epsilon}{\Lambda_N(\boldsymbol{\eta}^*)} \right\} = 2 \log N \quad \text{for all } N \geq N_1$$

by the choice of ϵ in Equation (10.43).

Showing that the restriction to the subset $\mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*) \subset \mathbb{N}$ is legitimate. To complete the proof, we show that the focus on the subset $\mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*) \subset \mathbb{N}$ is legitimate.

First, by Lemma 2, if $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_N$, then there exists a constant $L(\boldsymbol{\eta}^*) < \infty$, which depends on $\boldsymbol{\eta}^*$ but does not depend on N , such that

$$\Lambda_N(\boldsymbol{\eta}^*) \leq L(\boldsymbol{\eta}^*) \frac{1}{c_N(\mathbf{Z})^2 N (N \rho_N)^2}.$$

Therefore,

$$\Phi_N(\boldsymbol{\eta}^*) \leq L(\boldsymbol{\eta}^*) \frac{1}{c_N(\mathbf{Z})^2 (N \rho_N)^2} \sqrt{\frac{\log N}{N}} \rightarrow 0 \quad \text{as } N \rightarrow \infty, \quad (10.45)$$

because we have assumed that $c_N(\mathbf{Z})^2 (N \rho_N)^2 \sqrt{N / \log N} \rightarrow \infty$ as $N \rightarrow \infty$. The above result implies that there exists an integer $N_2 \geq 1$ such that $\epsilon := \sqrt{2/B_1(\boldsymbol{\eta}^*)} \Phi_N(\boldsymbol{\eta}^*) < \epsilon^*$ for all $N \geq N_2$.

Second, the fact that $\sqrt{2/B_1(\boldsymbol{\eta}^*)} \Phi_N(\boldsymbol{\eta}^*) < \epsilon^*$ for all $N \geq N_2$ implies that

$$\mathcal{B}(\boldsymbol{\eta}^*, \sqrt{2/B_1(\boldsymbol{\eta}^*)} \Phi_N(\boldsymbol{\eta}^*)) \subset \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*). \quad (10.46)$$

Combining (10.44) and (10.46) shows that, for all $N \geq \max\{N_1, N_2\}$,

$$\mathbb{P}(\hat{\boldsymbol{\eta}} \in \mathcal{B}(\boldsymbol{\eta}^*, \epsilon^*)) \geq \mathbb{P} \left(\hat{\boldsymbol{\eta}} \in \mathcal{B} \left(\boldsymbol{\eta}^*, C_1(\boldsymbol{\eta}^*) \frac{1}{c_N(\mathbf{Z})^2 (N \rho_N)^2} \sqrt{\frac{\log N}{N}} \right) \right) \geq 1 - \frac{8}{N^2}.$$

where $C_1(\boldsymbol{\eta}^*) := \sqrt{2/B_1(\boldsymbol{\eta}^*)} L(\boldsymbol{\eta}^*)$.

Convergence rates for $\hat{\boldsymbol{\theta}}$ based on $\hat{\boldsymbol{\eta}}$. We obtain convergence rates for the maximum likelihood estimator $\hat{\boldsymbol{\theta}}$ of $\boldsymbol{\theta}^*$ based on the convergence rates for the estimator $\hat{\boldsymbol{\eta}} := \boldsymbol{\eta}(\hat{\boldsymbol{\theta}}, \hat{\kappa}^2)$ of $\boldsymbol{\eta}^* := \boldsymbol{\eta}(\boldsymbol{\theta}^*, \kappa^{*2})$ by noting that

$$\begin{aligned}
\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_2 &= \left\| \frac{1}{\hat{\kappa}^2} \hat{\boldsymbol{\theta}} - \frac{1}{\hat{\kappa}^2} \boldsymbol{\theta}^* \right\|_2 \hat{\kappa}^2 \\
&= \left\| \left(\frac{1}{\hat{\kappa}^2} \hat{\boldsymbol{\theta}} - \frac{1}{\kappa^{*2}} \boldsymbol{\theta}^* \right) + \left(\frac{1}{\kappa^{*2}} \boldsymbol{\theta}^* - \frac{1}{\hat{\kappa}^2} \boldsymbol{\theta}^* \right) \right\|_2 \hat{\kappa}^2 \\
&\leq \left[\left\| \boldsymbol{\eta}(\hat{\boldsymbol{\theta}}, \hat{\kappa}^2) - \boldsymbol{\eta}(\boldsymbol{\theta}^*, \kappa^{*2}) \right\|_2 + \left| \frac{1}{\kappa^{*2}} - \frac{1}{\hat{\kappa}^2} \right| \|\boldsymbol{\theta}^*\|_2 \right] \hat{\kappa}^2 \\
&\leq C_1(\boldsymbol{\eta}^*) (1 + \|\boldsymbol{\theta}^*\|_2) \hat{\kappa}^2 \frac{1}{c_N(\mathbf{Z})^2 (N\rho_N)^2} \sqrt{\frac{\log N}{N}}
\end{aligned}$$

with probability at least $1 - 8/N^2$, provided that $N > \max\{N_1, N_2\}$. To deal with the random term $\hat{\kappa}^2$ in the upper bound on $\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_2$, note that

$$\begin{aligned}
\hat{\kappa}^2 &= \frac{1}{\frac{1}{\kappa^{*2}} + \left(\frac{1}{\hat{\kappa}^2} - \frac{1}{\kappa^{*2}} \right)} \\
&\leq \frac{1}{\frac{1}{\kappa^{*2}} - C_1(\boldsymbol{\eta}^*) \frac{1}{c_N(\mathbf{Z})^2 (N\rho_N)^2} \sqrt{\frac{\log N}{N}}} \leq 2\kappa^{*2}
\end{aligned}$$

with probability at least $1 - 8/N^2$, provided that $N > \max\{N_1, N_2\}$. Combining the above results proves that, for all $N > \max\{N_1, N_2\}$, the maximum likelihood estimator $\hat{\boldsymbol{\theta}}$ of $\boldsymbol{\theta}^*$ satisfies

$$\begin{aligned}
\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_2 &\leq C_1(\boldsymbol{\eta}^*) (1 + \|\boldsymbol{\theta}^*\|_2) \hat{\kappa}^2 \frac{1}{c_N(\mathbf{Z})^2 (N\rho_N)^2} \sqrt{\frac{\log N}{N}} \\
&\leq C_2(\boldsymbol{\eta}^*) \frac{1}{c_N(\mathbf{Z})^2 (N\rho_N)^2} \sqrt{\frac{\log N}{N}}.
\end{aligned}$$

with probability at least $1 - 8/N^2$, where

$$C_2(\boldsymbol{\eta}^*) := 2 C_1(\boldsymbol{\eta}^*) (1 + \|\boldsymbol{\theta}^*\|_2) \kappa^{*2}.$$

10.5 Proof of Proposition 5

According to Corollary 4.1 of Xie (2024), we have that under Condition C.1, there exists constants $C_T > 0$, $\tilde{C}_1 > 0$, and an orthogonal matrix $\mathbf{O} \in \mathbb{R}^{K \times K}$ such that

$$\mathbb{P} \left(\|\hat{\mathbf{T}} - \mathbf{T} \mathbf{O}\|_2 \leq C_T \sqrt{\log N} \right) \geq 1 - \frac{\tilde{C}_1}{N^2}. \quad (10.47)$$

As a result, we show that $\hat{\mathbf{T}}/\|\hat{\mathbf{T}}\|_2$ is also close to $\mathbf{T} \mathbf{O}/\|\mathbf{T} \mathbf{O}\|_2$ with high probability, which will be useful to derive the convergence rate of $\hat{a}(\delta)$.

$$\begin{aligned} \left\| \frac{\hat{\mathbf{T}}}{\|\hat{\mathbf{T}}\|_2} - \frac{\mathbf{T} \mathbf{O}}{\|\mathbf{T} \mathbf{O}\|_2} \right\|_2 &\leq \left\| \frac{\hat{\mathbf{T}}}{\|\hat{\mathbf{T}}\|_2} - \frac{\hat{\mathbf{T}}}{\|\mathbf{T} \mathbf{O}\|_2} \right\|_2 + \left\| \frac{\hat{\mathbf{T}}}{\|\mathbf{T} \mathbf{O}\|_2} - \frac{\mathbf{T} \mathbf{O}}{\|\mathbf{T} \mathbf{O}\|_2} \right\|_2 \\ &= \|\hat{\mathbf{T}}\|_2 \left| \frac{1}{\|\hat{\mathbf{T}}\|_2} - \frac{1}{\|\mathbf{T} \mathbf{O}\|_2} \right| + \frac{\|\hat{\mathbf{T}} - \mathbf{T} \mathbf{O}\|_2}{\|\mathbf{T} \mathbf{O}\|_2} \\ &= \frac{|\|\hat{\mathbf{T}}\|_2 - \|\mathbf{T} \mathbf{O}\|_2|}{\|\mathbf{T} \mathbf{O}\|_2} + \frac{\|\hat{\mathbf{T}} - \mathbf{T} \mathbf{O}\|_2}{\|\mathbf{T} \mathbf{O}\|_2} \\ &\leq 2 \frac{\|\hat{\mathbf{T}} - \mathbf{T} \mathbf{O}\|_2}{\|\mathbf{T} \mathbf{O}\|_2}. \end{aligned}$$

Moreover, by Condition C.1, we have that $\|\mathbf{T} \mathbf{O}\|_2 = \|\mathbf{T}\|_2 = \sqrt{\rho_N} \|\boldsymbol{\alpha}\|_2 \geq \alpha_{\min} \sqrt{N \rho_N}$.

Therefore,

$$\mathbb{P} \left(\left\| \frac{\hat{\mathbf{T}}}{\|\hat{\mathbf{T}}\|_2} - \frac{\mathbf{T} \mathbf{O}}{\|\mathbf{T} \mathbf{O}\|_2} \right\|_2 \leq \frac{2 C_T}{\alpha_{\min}} \sqrt{\frac{\log N}{N \rho_N}} \right) \geq 1 - \frac{\tilde{C}_1}{N^2}. \quad (10.48)$$

Now, we derive the convergence rate of $\hat{a}(\delta)$. Note that in the expression of $a_N(\delta)$, we

can replace \mathbf{T} by $\mathbf{T}\mathbf{O}$ for any orthogonal matrix $\mathbf{O} \in \mathbb{R}^{K \times K}$. So, we have

$$\begin{aligned}
& \widehat{a}(\delta) - a_N(\delta) \\
&= \frac{\delta}{N} (\mathbf{1}_N^\top \widehat{\mathbf{T}}) (\|\widehat{\mathbf{T}}\|_2^2 \mathbf{I}_K - \delta \widehat{\mathbf{T}}^\top \widehat{\mathbf{T}})^{-1} (\widehat{\mathbf{T}}^\top \mathbf{1}_N) \\
&- \frac{\delta}{N} (\mathbf{1}_N^\top \mathbf{T}\mathbf{O}) (\|\mathbf{T}\mathbf{O}\|_2^2 \mathbf{I}_K - \delta (\mathbf{T}\mathbf{O})^\top (\mathbf{T}\mathbf{O}))^{-1} ((\mathbf{T}\mathbf{O})^\top \mathbf{1}_N) \\
&= \frac{\delta}{N \|\widehat{\mathbf{T}}\|_2^2} (\mathbf{1}_N^\top \widehat{\mathbf{T}}) \mathbf{J}_1^{-1} (\widehat{\mathbf{T}}^\top \mathbf{1}_N) - \frac{\delta}{N \|\mathbf{T}\mathbf{O}\|_2^2} (\mathbf{1}_N^\top \mathbf{T}\mathbf{O}) \mathbf{J}_2^{-1} ((\mathbf{T}\mathbf{O})^\top \mathbf{1}_N) \\
&\quad \text{where } \mathbf{J}_1 := \mathbf{I}_K - \delta \widehat{\mathbf{T}}^\top \widehat{\mathbf{T}} / \|\widehat{\mathbf{T}}\|_2^2, \mathbf{J}_2 := \mathbf{I}_K - \delta (\mathbf{T}\mathbf{O})^\top (\mathbf{T}\mathbf{O}) / \|\mathbf{T}\mathbf{O}\|_2^2 \\
&= \frac{\delta}{N \|\widehat{\mathbf{T}}\|_2^2} (\mathbf{1}_N^\top \widehat{\mathbf{T}}) (\mathbf{J}_1^{-1} - \mathbf{J}_2^{-1}) (\widehat{\mathbf{T}}^\top \mathbf{1}_N) + \frac{\delta}{N \|\widehat{\mathbf{T}}\|_2^2} (\mathbf{1}_N^\top \widehat{\mathbf{T}}) \mathbf{J}_2^{-1} (\widehat{\mathbf{T}}^\top \mathbf{1}_N) \\
&- \frac{\delta}{N \|\mathbf{T}\mathbf{O}\|_2^2} (\mathbf{1}_N^\top \mathbf{T}\mathbf{O}) \mathbf{J}_2^{-1} ((\mathbf{T}\mathbf{O})^\top \mathbf{1}_N).
\end{aligned}$$

Bounding $|\widehat{a}(\delta) - a_N(\delta)|$. Using sub-multiplicativity of the operator norm, we obtain

$$\begin{aligned}
& |\widehat{a}(\delta) - a_N(\delta)| \\
&\leq \|\mathbf{J}_1^{-1} - \mathbf{J}_2^{-1}\|_2 + \frac{1}{N} \left| \frac{(\mathbf{1}_N^\top \widehat{\mathbf{T}}) \mathbf{J}_2^{-1} (\widehat{\mathbf{T}}^\top \mathbf{1}_N)}{\|\widehat{\mathbf{T}}\|_2^2} - \frac{(\mathbf{1}_N^\top \mathbf{T}\mathbf{O}) \mathbf{J}_2^{-1} ((\mathbf{T}\mathbf{O})^\top \mathbf{1}_N)}{\|\mathbf{T}\mathbf{O}\|_2^2} \right|. \tag{10.49}
\end{aligned}$$

We establish bounds on the two terms on the right-hand side above.

Bounding $\|\mathbf{J}_1^{-1} - \mathbf{J}_2^{-1}\|_2$. Since $|\delta| < 1$, \mathbf{J}_1 and \mathbf{J}_2 are positive definite, and

$$\|\mathbf{J}_1^{-1}\|_2 \leq 1/(1 - |\delta|), \quad \|\mathbf{J}_2^{-1}\|_2 \leq 1/(1 - |\delta|). \tag{10.50}$$

Hence,

$$\|\mathbf{J}_1^{-1} - \mathbf{J}_2^{-1}\|_2 = \|\mathbf{J}_1^{-1}(\mathbf{J}_2 - \mathbf{J}_1)\mathbf{J}_2^{-1}\|_2$$

$$\begin{aligned}
&\leq \|J_1^{-1}\|_2 \|J_2 - J_1\|_2 \|J_2^{-1}\|_2 \leq \frac{1}{(1-|\delta|)^2} \|J_2 - J_1\|_2 \\
&= \frac{1}{(1-|\delta|)^2} \left\| \frac{\hat{T}^\top \hat{T}}{\|\hat{T}\|_2^2} - \frac{(TO)^\top (TO)}{\|TO\|_2^2} \right\|_2 \\
&\leq \frac{1}{(1-|\delta|)^2} \left(\left\| \frac{\hat{T}}{\|\hat{T}\|_2} - \frac{TO}{\|TO\|_2} \right\|_2^2 + 2 \left\| \frac{\hat{T}}{\|\hat{T}\|_2} - \frac{TO}{\|TO\|_2} \right\|_2 \right).
\end{aligned}$$

$$\text{Bounding } \frac{1}{N} \left| \frac{(\mathbf{1}_N^\top \hat{T}) J_2^{-1} (\hat{T}^\top \mathbf{1}_N)}{\|\hat{T}\|_2^2} - \frac{(\mathbf{1}_N^\top TO) J_2^{-1} ((TO)^\top \mathbf{1}_N)}{\|TO\|_2^2} \right|.$$

$$\begin{aligned}
&\frac{1}{N} \left| \frac{(\mathbf{1}_N^\top \hat{T}) J_2^{-1} (\hat{T}^\top \mathbf{1}_N)}{\|\hat{T}\|_2^2} - \frac{(\mathbf{1}_N^\top TO) J_2^{-1} ((TO)^\top \mathbf{1}_N)}{\|TO\|_2^2} \right| \\
&= \frac{1}{N} \left| \left\| \frac{J_2^{-\frac{1}{2}} \hat{T}^\top \mathbf{1}_N}{\|\hat{T}\|_2} \right\|_2^2 - \left\| \frac{J_2^{-\frac{1}{2}} (TO)^\top \mathbf{1}_N}{\|TO\|_2} \right\|_2^2 \right| \\
&= \frac{1}{N} \left(\left\| \frac{J_2^{-\frac{1}{2}} \hat{T}^\top \mathbf{1}_N}{\|\hat{T}\|_2} \right\|_2 + \left\| \frac{J_2^{-\frac{1}{2}} (TO)^\top \mathbf{1}_N}{\|TO\|_2} \right\|_2 \right) \\
&\times \left| \left\| \frac{J_2^{-\frac{1}{2}} \hat{T}^\top \mathbf{1}_N}{\|\hat{T}\|_2} \right\|_2 - \left\| \frac{J_2^{-\frac{1}{2}} (TO)^\top \mathbf{1}_N}{\|TO\|_2} \right\|_2 \right| \\
&\leq \frac{1}{N} (2 \|J_2^{-\frac{1}{2}}\|_2 \sqrt{N}) \left\| J_2^{-\frac{1}{2}} \left(\frac{\hat{T}}{\|\hat{T}\|_2} - \frac{TO}{\|TO\|_2} \right)^\top \mathbf{1}_N \right\|_2 \\
&\leq 2 \|J_2^{-1}\|_2 \left\| \frac{\hat{T}}{\|\hat{T}\|_2} - \frac{TO}{\|TO\|_2} \right\|_2 \\
&\leq \frac{2}{1-|\delta|} \left\| \frac{\hat{T}}{\|\hat{T}\|_2} - \frac{TO}{\|TO\|_2} \right\|_2,
\end{aligned}$$

incorporating (10.50) in the final step above.

Conclusion. Combining terms, we obtain that for any δ such that $|\delta| < 1$,

$$|\hat{a}(\delta) - a_N(\delta)| \leq \tilde{C}_a(\delta) C_0 g(\mathbf{Z}), \tag{10.51}$$

where

$$\begin{aligned}\tilde{C}_a(\delta) &:= \frac{1}{(1 - |\delta|)^2}, \\ g(\mathbf{Z}) &:= \frac{1}{C_0} \left(\left\| \frac{\hat{\mathbf{T}}}{\|\hat{\mathbf{T}}\|_2} - \frac{\mathbf{T}\mathbf{O}}{\|\mathbf{T}\mathbf{O}\|_2} \right\|_2^2 + 4 \left\| \frac{\hat{\mathbf{T}}}{\|\hat{\mathbf{T}}\|_2} - \frac{\mathbf{T}\mathbf{O}}{\|\mathbf{T}\mathbf{O}\|_2} \right\|_2 \right),\end{aligned}$$

and $C_0 > 0$ is a proportionality constant chosen appropriately. Leveraging (10.48), we obtain that with probability at least $1 - \tilde{C}_1/N^2$,

$$g(\mathbf{Z}) \leq \frac{1}{C_0} \left(\frac{4 C_T^2 \log N}{\alpha_{\min}^2 N \rho_N} + \frac{8 C_T}{\alpha_{\min}} \sqrt{\frac{\log N}{N \rho_N}} \right).$$

Noting that $\sqrt{\log N / (N \rho_N)}$ is the dominating term above due to Condition C.1, we can choose C_0 such that

$$\mathbb{P} \left(g(\mathbf{Z}) \leq \sqrt{\frac{\log N}{N \rho_N}} \right) \geq 1 - \frac{\tilde{C}_1}{N^2}. \quad (10.52)$$

10.6 Proof of Theorem 2

Throughout, we suppress the notational dependence of $\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2)$ on $(\boldsymbol{\theta}, \kappa^2)$ and write $\boldsymbol{\eta}$ instead of $\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2)$, and define $\hat{\boldsymbol{\eta}} := \boldsymbol{\eta}(\hat{\boldsymbol{\theta}}, \hat{\kappa}^2)$ and $\boldsymbol{\eta}^\star := \boldsymbol{\eta}(\boldsymbol{\theta}^\star, \kappa^{\star 2})$. Conditional on the event $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_N$ defined in Supplement 10.1.1,

$$\mathbb{P}_{\mathbf{y}|\mathbf{W}, \mathbf{Z}} \left(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^\star\|_2 \leq C_2(\boldsymbol{\theta}^\star, \kappa^{\star 2}) \sqrt{\frac{\log N}{N}} \frac{1}{c_N(\mathbf{Z})^2 (N \rho_N)^2} \right) \geq 1 - \frac{8}{N^2}$$

by Proposition 4. The event $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_N$ occurs with probability at least

$$\mathbb{P}((\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_N) \geq 1 - \frac{\tilde{C}_2}{N^2}$$

by Proposition 3. Taken together, Propositions 3 and 4 imply that

$$\begin{aligned} & \mathbb{P} \left(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_2 \leq C_2(\boldsymbol{\theta}^*, \kappa^{\star 2}) \sqrt{\frac{\log N}{N}} \frac{1}{c_N(\mathbf{Z})^2 (N\rho_N)^2} \right) \\ & \geq \left(1 - \frac{8}{N^2}\right) \left(1 - \frac{\tilde{C}_2}{N^2}\right) \geq 1 - \frac{8 + \tilde{C}_2}{N^2}. \end{aligned}$$

Since $c_N(\mathbf{Z}) := (1 / (\|\mathbf{Z}\|_2) \mathbb{I}(\|\mathbf{Z}\|_2 > 0))$, (10.26) implies that

$$c_N(\mathbf{Z}) N\rho_N = \frac{N\rho_N}{\|\mathbf{Z}\|_2 \mathbb{I}(\|\mathbf{Z}\|_2 > 0)} \geq \frac{1}{2}$$

in the event $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N}$. Therefore,

$$\mathbb{P} \left(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_2 \leq 4 C_2(\boldsymbol{\theta}^*, \kappa^{\star 2}) \sqrt{\frac{\log N}{N}} \right) \geq 1 - \frac{8 + \tilde{C}_2}{N^2}. \quad (10.53)$$

Armed with these results, we establish convergence rates for the direct and indirect causal effects.

Bounding $|\tau_D(\hat{\boldsymbol{\theta}}) - \tau_D(\boldsymbol{\theta}^*)|$. Bounding the statistical error of the direct causal effect is straightforward, because

$$|\tau_D(\hat{\boldsymbol{\theta}}) - \tau_D(\boldsymbol{\theta}^*)| = |\hat{\beta} - \beta^*| \leq \|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_2 \leq C_D(\boldsymbol{\theta}^*, \kappa^{\star 2}) \sqrt{\frac{\log N}{N}}$$

with probability at least $1 - (8 + \tilde{C}_2) / N^2$, where $C_D(\boldsymbol{\theta}^*, \kappa^{\star 2}) := 4 C_2(\boldsymbol{\theta}^*, \kappa^{\star 2}) > 0$.

Bounding $|\tau_I(\widehat{\boldsymbol{\theta}}) - \tau_I(\boldsymbol{\theta}^*)|$. Write

$$\begin{aligned}
& |\tau_I(\widehat{\boldsymbol{\theta}}) - \tau_I(\boldsymbol{\theta}^*)| \\
&= |(\widehat{\gamma} - \gamma^*) + (\widehat{a}(\widehat{\delta}) - a(\delta^*))(\widehat{\beta} + \widehat{\gamma}) + a(\delta^*)((\widehat{\beta} + \widehat{\gamma}) - (\beta^* + \gamma^*))| \\
&\leq |\widehat{\gamma} - \gamma^*| + |\widehat{a}(\widehat{\delta}) - a(\delta^*)| |(\widehat{\beta} + \widehat{\gamma}) - (\beta^* + \gamma^*)| \\
&+ |\widehat{a}(\widehat{\delta}) - a(\delta^*)| |\beta^* + \gamma^*| + |a(\delta^*)| |(\widehat{\beta} + \widehat{\gamma}) - (\beta^* + \gamma^*)|.
\end{aligned} \tag{10.54}$$

We bound the terms on the right-hand side of Equation (10.54) one by one.

Bounding $|\widehat{\gamma} - \gamma^*|$.

$$|\widehat{\gamma} - \gamma^*| \leq \|\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_2. \tag{10.55}$$

Bounding $|(\widehat{\beta} + \widehat{\gamma}) - (\beta^* + \gamma^*)|$.

$$|(\widehat{\beta} + \widehat{\gamma}) - (\beta^* + \gamma^*)| \leq \sqrt{2} \|\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_2. \tag{10.56}$$

Bounding $|\widehat{a}(\widehat{\delta}) - a(\delta^*)|$. We next establish bounds for $|\widehat{a}(\widehat{\delta}) - a(\delta^*)|$ by breaking up $\widehat{a}(\widehat{\delta}) - a(\delta^*)$ into three parts:

- $\widehat{a}(\widehat{\delta}) - a_N(\widehat{\delta})$
- $a_N(\widehat{\delta}) - a_N(\delta^*)$
- $a_N(\delta^*) - a(\delta^*)$.

Bounding $|\widehat{a}(\widehat{\delta}) - a_N(\widehat{\delta})|$. By Proposition 5, for any δ satisfying $|\delta| < 1$,

$$|\widehat{a}(\delta) - a_N(\delta)| \leq C_a(\delta) g(\mathbf{Z}), \tag{10.57}$$

where $g(\mathbf{Z})$ is a function of \mathbf{Z} such that

$$\mathbb{P}\left(g(\mathbf{Z}) \leq \sqrt{\frac{\log N}{N\rho_N}}\right) \geq 1 - \frac{\tilde{C}_1}{N^2}, \quad (10.58)$$

and $C_a(\delta) > 0$ is a non-decreasing function of $|\delta|$ that does not depend on N . We next show that $|\hat{\delta}| \leq (2 + |\delta^*|)/3 < 1$ with high probability. First, observe that

$$\begin{aligned} |\hat{\delta}| &\leq \hat{\kappa}^2 \left(\left| \frac{\hat{\delta}}{\hat{\kappa}^2} - \frac{\delta^*}{\kappa^{*2}} \right| + \frac{|\delta^*|}{\kappa^{*2}} \right) \\ &= \frac{\left| \hat{\delta}/\hat{\kappa}^2 - \delta^*/\kappa^{*2} \right| + |\delta^*|/\kappa^{*2}}{1/\hat{\kappa}^2 - 1/\kappa^{*2} + 1/\kappa^{*2}} \\ &\leq \frac{\left| \hat{\delta}/\hat{\kappa}^2 - \delta^*/\kappa^{*2} \right| + |\delta^*|/\kappa^{*2}}{1/\kappa^{*2} - |1/\hat{\kappa}^2 - 1/\kappa^{*2}|} \\ &\leq \frac{\|\hat{\boldsymbol{\eta}} - \boldsymbol{\eta}^*\|_2 + |\delta^*|/\kappa^{*2}}{1/\kappa^{*2} - \|\hat{\boldsymbol{\eta}} - \boldsymbol{\eta}^*\|_2}. \end{aligned} \quad (10.59)$$

According to Proposition 4, we know that

$$\mathbb{P}(\|\hat{\boldsymbol{\eta}} - \boldsymbol{\eta}^*\|_2 < \epsilon^*) \geq 1 - \frac{8 + \tilde{C}_2}{N^2}. \quad (10.60)$$

Since $\epsilon^* \leq (1 - |\delta^*|)/3\kappa^{*2}$,

$$\frac{\|\hat{\boldsymbol{\eta}} - \boldsymbol{\eta}^*\|_2 + |\delta^*|/\kappa^{*2}}{1/\kappa^{*2} - \|\hat{\boldsymbol{\eta}} - \boldsymbol{\eta}^*\|_2} \leq \frac{(1 - |\delta^*|)/3\kappa^{*2} + |\delta^*|/\kappa^{*2}}{1/\kappa^{*2} - (1 - |\delta^*|)/3\kappa^{*2}} = \frac{1 + 2|\delta^*|}{2 + |\delta^*|} \leq \frac{2 + |\delta^*|}{3}.$$

We obtain

$$\mathbb{P}(|\hat{\delta}| \leq (2 + |\delta^*|)/3) \geq 1 - \frac{8 + \tilde{C}_2}{N^2}. \quad (10.61)$$

By combining (10.57), (10.58), and (10.61), we obtain

$$\begin{aligned}
& \mathbb{P} \left(|\widehat{a}(\widehat{\delta}) - a_N(\widehat{\delta})| \leq C_a \left(\frac{2 + |\delta^*|}{3} \right) \sqrt{\frac{\log N}{N\rho_N}} \right) \\
& \geq \mathbb{P} \left(|\widehat{a}(\widehat{\delta}) - a_N(\widehat{\delta})| \leq C_a \left(\frac{2 + |\delta^*|}{3} \right) \sqrt{\frac{\log N}{N\rho_N}} \cap |\widehat{\delta}| \leq \frac{2 + |\delta^*|}{3} \right) \\
& \geq \mathbb{P} \left(|\widehat{a}(\widehat{\delta}) - a_N(\widehat{\delta})| \leq C_a(\widehat{\delta}) \sqrt{\frac{\log N}{N\rho_N}} \cap |\widehat{\delta}| \leq \frac{2 + |\delta^*|}{3} \right) \tag{10.62} \\
& \geq \mathbb{P} \left(g(\mathbf{Z}) \leq \sqrt{\frac{\log N}{N\rho_N}} \cap |\widehat{\delta}| \leq \frac{2 + |\delta^*|}{3} \right) \\
& \geq 1 - \frac{8 + \widetilde{C}_2 + \widetilde{C}_1}{N^2}.
\end{aligned}$$

The second inequality uses the fact that $C_a(\delta)$ is a non-decreasing function of $|\delta|$ and $|\widehat{\delta}| \leq (2 + |\delta^*|)/3$ with probability at least $1 - (8 + \widetilde{C}_2)/N^2$ by (10.61). The third inequality uses (10.57) and (10.58).

Bounding $|a_N(\widehat{\delta}) - a_N(\delta^*)|$. By the mean-value theorem, there exists a real number $\delta_m \in (\delta^*, \widehat{\delta})$ such that

$$a_N(\widehat{\delta}) - a_N(\delta^*) = a'_N(\delta_m) (\widehat{\delta} - \delta^*), \tag{10.63}$$

where $a'_N(\delta_m)$ is the derivative of $a_N(\delta)$ evaluated at $\delta = \delta_m$:

$$\begin{aligned}
& a'_N(\delta_m) \\
& = \left. \frac{d}{d\delta} \frac{\delta}{N} (\mathbf{1}_N^\top \mathbf{T}) (\|\mathbf{T}\|_2^2 \mathbf{I}_K - \delta \mathbf{T}^\top \mathbf{T})^{-1} (\mathbf{T}^\top \mathbf{1}_N) \right|_{\delta=\delta_m} \\
& = \frac{\delta_m}{N} (\mathbf{1}_N^\top \mathbf{T}) (\|\mathbf{T}\|_2^2 \mathbf{I}_K - \delta_m \mathbf{T}^\top \mathbf{T})^{-1} \mathbf{T}^\top \mathbf{T} (\|\mathbf{T}\|_2^2 \mathbf{I}_K - \delta_m \mathbf{T}^\top \mathbf{T})^{-1} (\mathbf{T}^\top \mathbf{1}_N) \\
& + \frac{1}{N} (\mathbf{1}_N^\top \mathbf{T}) (\|\mathbf{T}\|_2^2 \mathbf{I}_K - \delta_m \mathbf{T}^\top \mathbf{T})^{-1} (\mathbf{T}^\top \mathbf{1}_N).
\end{aligned}$$

If $|\delta_m| \leq (2 + |\delta^*|)/3$, then—by applying the operator norm on the right-hand side—we

obtain

$$|a'_N(\delta_m)| \leq \frac{1}{(1 - (2 + |\delta^*|)/3)^2} + \frac{1}{1 - (2 + |\delta^*|)/3} \leq \frac{18}{(1 - |\delta^*|)^2}.$$

Since $\delta_m \in (\delta^*, \widehat{\delta})$ and $\mathbb{P}(|\widehat{\delta}| \leq (2 + |\delta^*|)/3) \geq 1 - (8 + \widetilde{C}_2)/N^2$ by (10.61), we obtain

$$\mathbb{P}(|\delta_m| \leq (2 + |\delta^*|)/3) \geq 1 - \frac{8 + \widetilde{C}_2}{N^2}$$

and hence

$$\mathbb{P}\left(|a'_N(\delta_m)| \leq \frac{18}{(1 - |\delta^*|)^2}\right) \geq 1 - \frac{8 + \widetilde{C}_2}{N^2}. \quad (10.64)$$

By combining (10.63), (10.64), and (10.53), we obtain

$$\mathbb{P}\left(|a_N(\widehat{\delta}) - a_N(\delta^*)| \leq \frac{72 C_2(\boldsymbol{\theta}^*, \kappa^{\star 2})}{(1 - |\delta^*|)^2} \sqrt{\frac{\log N}{N}}\right) \geq 1 - \frac{16 + 2 \widetilde{C}_2}{N^2}. \quad (10.65)$$

Bounding $|a_N(\delta^*) - a(\delta^*)|$. By Condition C.4,

$$|a_N(\delta^*) - a(\delta^*)| \leq C(\delta^*) \sqrt{\frac{\log N}{N \rho_N}}, \quad (10.66)$$

where $C(\delta^*) > 0$ depends on δ^* but does not depend on N .

Conclusion: $|\widehat{a}(\widehat{\delta}) - a(\delta^*)|$. Combining (10.62), (10.65), and (10.66) gives

$$\begin{aligned} \mathbb{P}\left(|\widehat{a}(\widehat{\delta}) - a(\delta^*)| \leq \left(C_a \left(\frac{2 + |\delta^*|}{3}\right) + \frac{72 C_2(\boldsymbol{\theta}^*, \kappa^{\star 2})}{(1 - |\delta^*|)^2} + C(\delta^*)\right) \sqrt{\frac{\log N}{N \rho_N}}\right) \\ \geq 1 - \frac{24 + 3 \widetilde{C}_2 + \widetilde{C}_1}{N^2}. \end{aligned} \quad (10.67)$$

Bounding $|a(\delta^*)|$. We know that $a(\delta^*) := \lim_{N \rightarrow \infty} a_N(\delta^*)$, hence

$$|a_N(\delta^*)| = \left| \frac{\delta^*}{N} (\mathbf{1}_N^\top \mathbf{T}) (\|\mathbf{T}\|_2^2 \mathbf{I}_K - \delta^* \mathbf{T}^\top \mathbf{T})^{-1} (\mathbf{T}^\top \mathbf{1}_N) \right| \leq \frac{1}{1 - |\delta^*|},$$

where the inequality is obtained by applying the operator norm, and using the fact that $|\delta^*| < 1$ by assumption. We conclude that

$$|a(\delta^*)| \leq \frac{1}{1 - |\delta^*|}. \quad (10.68)$$

Conclusion: $|\tau_I(\hat{\boldsymbol{\theta}}) - \tau_I(\boldsymbol{\theta}^*)|$. By combining (10.54), (10.55), (10.56), (10.67), (10.68), and (10.53), we conclude that there exists a constant $C_I(\boldsymbol{\theta}^*, \kappa^{*2}) > 0$, which depends on $\boldsymbol{\theta}^*$ and κ^{*2} but does not depend on N , such that

$$\mathbb{P} \left(|\tau_I(\hat{\boldsymbol{\theta}}) - \tau_I(\boldsymbol{\theta}^*)| \leq C_I(\boldsymbol{\theta}^*, \kappa^{*2}) \sqrt{\frac{\log N}{N \rho_N}} \right) \geq 1 - \frac{32 + 4\tilde{C}_2 + \tilde{C}_1}{N^2}.$$

11 Auxiliary results

11.1 Lemma 1

Proof of Lemma 1. By assumption,

$$\mathbb{E}[Y_i \mid (\mathbf{Y}_{-i}, \mathbf{W}, \mathbf{Z}) = (\mathbf{y}_{-i}, \mathbf{w}, \mathbf{z})] = \beta w_i + \gamma \frac{\sum_j w_j z_{i,j}}{\sum_j z_{i,j}} + \delta c_N(\mathbf{z}) \sum_{j=1}^N y_j z_{i,j}. \quad (11.1)$$

Let $\mu_i(\mathbf{w}, \mathbf{z}) := \mathbb{E}[Y_i \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})]$. Upon taking expectations on both sides above, we obtain

$$\mu_i(\mathbf{w}, \mathbf{z}) = \beta w_i + \gamma \frac{\sum_j w_j z_{i,j}}{\sum_j z_{i,j}} + \delta c_N(\mathbf{z}) \sum_{j=1}^N \mu_j(\mathbf{w}, \mathbf{z}) z_{i,j}. \quad (11.2)$$

Subtracting (11.2) from (11.1), we obtain

$$\mathbb{E}[Y_i \mid (\mathbf{Y}_{-i}, \mathbf{W}, \mathbf{Z}) = (\mathbf{y}_{-i}, \mathbf{w}, \mathbf{z})] = \mu_i(\mathbf{w}, \mathbf{z}) + \delta c_N(\mathbf{z}) \sum_{j=1}^N (y_j - \mu_j(\mathbf{w}, \mathbf{z})) z_{i,j}. \quad (11.3)$$

We have also assumed that

$$\text{Var}[Y_i \mid (\mathbf{Y}_{-i}, \mathbf{W}, \mathbf{Z}) = (\mathbf{y}_{-i}, \mathbf{w}, \mathbf{z})] = \kappa^2. \quad (11.4)$$

Upon invoking Theorem 2.6 of (Rue and Held, 2005), we can conclude from (11.3) and (11.4) that if $(\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})$ is positive definite, then $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z}) = (\mathbf{w}, \mathbf{z})$ is multivariate Gaussian with mean vector $\boldsymbol{\mu}(\mathbf{w}, \mathbf{z})$ and precision matrix

$$\boldsymbol{\Omega}(\mathbf{w}, \mathbf{z}) := \frac{1}{\kappa^2} (\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z}). \quad (11.5)$$

To obtain the expression for $\boldsymbol{\mu}(\mathbf{w}, \mathbf{z})$, note that (11.2) can be written in matrix form as

$$\boldsymbol{\mu}(\mathbf{w}, \mathbf{z}) = \mathbf{G}(\mathbf{w}, \mathbf{z}) \boldsymbol{\vartheta} + \delta c_N(\mathbf{z}) \mathbf{z} \boldsymbol{\mu}(\mathbf{w}, \mathbf{z}), \quad (11.6)$$

where $\boldsymbol{\vartheta} := (\beta, \gamma) \in \mathbb{R}^2$ and $\mathbf{G}(\mathbf{w}, \mathbf{z})$ is a $N \times 2$ matrix with i -th row

$$(\mathbf{G}(\mathbf{w}, \mathbf{z}))_{i,\cdot} := \left(w_i, \frac{\sum_j w_j z_{i,j}}{\sum_j z_{i,j}} \right), \quad i = 1, \dots, N.$$

If $|\delta| < 1$ and $c_N(\mathbf{z}) \|\mathbf{z}\|_2 \leq 1$, then $(\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})$ is positive definite, and we have

$$\boldsymbol{\mu}(\mathbf{w}, \mathbf{z}) := (\mathbf{I}_N - \delta c_N(\mathbf{z}) \mathbf{z})^{-1} \mathbf{G}(\mathbf{w}, \mathbf{z}) \boldsymbol{\vartheta}. \quad (11.7)$$

11.2 Lemma 2

Proof of Lemma 2. Recall that the natural parameter vector is defined as

$$\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2) := \left(\frac{\beta}{\kappa^2}, \frac{\gamma}{\kappa^2}, \frac{\delta}{\kappa^2}, \frac{1}{\kappa^2} \right)^\top = (\eta_1, \eta_2, \eta_3, \eta_4)^\top, \text{ say.} \quad (11.8)$$

Also, define $\boldsymbol{\phi} := (\eta_1, \eta_2)^\top$. We suppress the notational dependence of $\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2)$ on $(\boldsymbol{\theta}, \kappa^2)$ and write $\boldsymbol{\eta}$ instead of $\boldsymbol{\eta}(\boldsymbol{\theta}, \kappa^2)$.

The loglikelihood function in (10.2) can be expressed in terms of the natural parameters as

$$\begin{aligned} \ell(\boldsymbol{\eta}) &= \frac{1}{2} \log(\det(\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})) - \frac{\eta_4}{2} \mathbf{Y}^\top \mathbf{Y} \\ &\quad - \frac{1}{2} \boldsymbol{\phi}^\top \mathbf{G}(\mathbf{W}, \mathbf{Z})^\top (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{G}(\mathbf{W}, \mathbf{Z}) \boldsymbol{\phi} \\ &\quad + \frac{\eta_3 c_N(\mathbf{Z})}{2} \mathbf{Y}^\top \mathbf{Z} \mathbf{Y} + \boldsymbol{\phi}^\top \mathbf{G}(\mathbf{W}, \mathbf{Z})^\top \mathbf{Y}. \end{aligned} \quad (11.9)$$

From Theorems 1 and 2 of Magnus and Neudecker (1988, Chapter 8), we obtain

$$\begin{aligned} \frac{d}{d\eta_3} \det(\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z}) &= -c_N(\mathbf{Z}) \det(\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z}) \\ &\quad \times \text{trace}((\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{Z}), \\ \frac{d}{d\eta_4} \det(\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z}) &= \det(\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z}) \\ &\quad \times \text{trace}((\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1}), \\ \frac{d}{d\eta_3} (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} &= c_N(\mathbf{Z}) (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{Z} (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1}, \end{aligned}$$

$$\frac{d}{d\eta_4} (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} = - (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-2}.$$

Applying the above equations to the loglikelihood function (11.9), we obtain

$$\begin{aligned} -\nabla_{\phi}^2 \ell(\eta) &= \mathbf{G}(\mathbf{W}, \mathbf{Z})^{\top} (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{G}(\mathbf{W}, \mathbf{Z}) \\ -\nabla_{\eta_3}^2 \ell(\eta) &= c_N(\mathbf{Z})^2 \boldsymbol{\mu}(\mathbf{W}, \mathbf{Z})^{\top} \mathbf{Z} (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{Z} \boldsymbol{\mu}(\mathbf{W}, \mathbf{Z}) \\ &\quad + \frac{c_N(\mathbf{Z})^2}{2} \text{trace}(((\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{Z})^2) \\ -\nabla_{\eta_4}^2 \ell(\eta) &= \boldsymbol{\mu}(\mathbf{W}, \mathbf{Z})^{\top} (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} \boldsymbol{\mu}(\mathbf{W}, \mathbf{Z}) \\ &\quad + \frac{1}{2} \text{trace}((\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-2}) \\ -\nabla_{\phi} \nabla_{\eta_3} \ell(\eta) &= c_N(\mathbf{Z}) \mathbf{G}(\mathbf{W}, \mathbf{Z})^{\top} (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{Z} \boldsymbol{\mu}(\mathbf{W}, \mathbf{Z}) \\ -\nabla_{\phi} \nabla_{\eta_4} \ell(\eta) &= -\mathbf{G}(\mathbf{W}, \mathbf{Z})^{\top} (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} \boldsymbol{\mu}(\mathbf{W}, \mathbf{Z}). \\ -\nabla_{\eta_3} \nabla_{\eta_4} \ell(\eta) &= -c_N(\mathbf{Z}) \boldsymbol{\mu}(\mathbf{W}, \mathbf{Z})^{\top} \mathbf{Z} (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} \boldsymbol{\mu}(\mathbf{W}, \mathbf{Z}) \\ &\quad - \frac{c_N(\mathbf{Z})}{2} \text{trace}((\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{Z} (\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1}). \end{aligned}$$

Observe that $(\eta_4 \mathbf{I}_N - \eta_3 c_N(\mathbf{Z}) \mathbf{Z})^{-1}$ is the covariance matrix $\boldsymbol{\Sigma}(\mathbf{W}, \mathbf{Z}) := \boldsymbol{\Omega}(\mathbf{W}, \mathbf{Z})^{-1}$ of the multivariate Gaussian distribution of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z})$. To simplify notation, we write henceforth \mathbf{G} , $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$ instead of $\mathbf{G}(\mathbf{W}, \mathbf{Z})$, $\boldsymbol{\mu}(\mathbf{W}, \mathbf{Z})$ and $\boldsymbol{\Sigma}(\mathbf{W}, \mathbf{Z})$, respectively. Then,

we have the information matrix as,

$$\begin{aligned}
\mathcal{I}(\boldsymbol{\eta}) &= \begin{pmatrix} \mathbf{G} \\ c_N(\mathbf{Z}) \mathbf{Z} \boldsymbol{\mu} \\ -\boldsymbol{\mu} \end{pmatrix}^\top \boldsymbol{\Sigma} \begin{pmatrix} \mathbf{G} & c_N(\mathbf{Z}) \mathbf{Z} \boldsymbol{\mu} & -\boldsymbol{\mu} \end{pmatrix} \\
&+ \frac{1}{2} \begin{pmatrix} \mathbf{0}_{2 \times 2} & \mathbf{0}_{2 \times 1} & \mathbf{0}_{2 \times 1} \\ \mathbf{0}_{1 \times 2} & c_N(\mathbf{Z})^2 \text{trace}((\boldsymbol{\Sigma} \mathbf{Z})^2) & -c_N(\mathbf{Z}) \text{trace}(\boldsymbol{\Sigma} \mathbf{Z} \boldsymbol{\Sigma}) \\ \mathbf{0}_{1 \times 2} & -c_N(\mathbf{Z}) \text{trace}(\boldsymbol{\Sigma} \mathbf{Z} \boldsymbol{\Sigma}) & \text{trace}(\boldsymbol{\Sigma}^2) \end{pmatrix} \quad (11.10) \\
&=: \mathcal{I}_1(\boldsymbol{\eta}) + \mathcal{I}_2(\boldsymbol{\eta}),
\end{aligned}$$

where $\mathbf{0}_{d_1 \times d_2}$ stands for the $d_1 \times d_2$ matrix of 0's.

Define

$$\tilde{\mathcal{I}}_1(\boldsymbol{\eta}) := \begin{pmatrix} \mathbf{G} \\ c_N(\mathbf{Z}) \mathbf{Z} \boldsymbol{\mu} \\ -\boldsymbol{\mu} \end{pmatrix}^\top \begin{pmatrix} \mathbf{G} & c_N(\mathbf{Z}) \mathbf{Z} \boldsymbol{\mu} & -\boldsymbol{\mu} \end{pmatrix}.$$

Since $c_N(\mathbf{Z}) \|\mathbf{Z}\|_2 \leq 1$, the smallest eigenvalue of $\boldsymbol{\Sigma}$ is at least $\kappa^2/(1 + |\delta|)$, which implies,

$$\mathcal{I}_1(\boldsymbol{\eta}) \succeq \frac{\kappa^2}{1 + |\delta|} \tilde{\mathcal{I}}_1(\boldsymbol{\eta}),$$

and hence,

$$\mathcal{I}(\boldsymbol{\eta}) \succeq \frac{\kappa^2}{1 + |\delta|} \tilde{\mathcal{I}}_1(\boldsymbol{\eta}) + \mathcal{I}_2(\boldsymbol{\eta}) =: \tilde{\mathcal{I}}(\boldsymbol{\eta}), \text{ say.}$$

Then

$$\|\mathcal{I}(\boldsymbol{\eta})^{-1}\|_2 = \frac{1}{\lambda_{\min}(\mathcal{I}(\boldsymbol{\eta}))} \leq \frac{1}{\lambda_{\min}(\tilde{\mathcal{I}}(\boldsymbol{\eta}))}. \quad (11.11)$$

To derive a lower bound on $\lambda_{\min}(\tilde{\mathcal{I}}(\boldsymbol{\eta}))$, we first prove an upper and lower bound on the

determinant $\det(\tilde{\mathcal{I}}(\boldsymbol{\eta}))$, and use the fact that $\det(\tilde{\mathcal{I}}(\boldsymbol{\eta}))$ is the product of its eigenvalues.

Upper bound on $\det(\tilde{\mathcal{I}}(\boldsymbol{\eta}))$.

$$\det(\tilde{\mathcal{I}}(\boldsymbol{\eta})) = \prod_{i=1}^4 \lambda_i(\tilde{\mathcal{I}}(\boldsymbol{\eta})) \leq \|\tilde{\mathcal{I}}(\boldsymbol{\eta})\|_2^3 \lambda_{\min}(\tilde{\mathcal{I}}(\boldsymbol{\eta})), \quad (11.12)$$

since $\lambda_1(\tilde{\mathcal{I}}(\boldsymbol{\eta})) = \|\tilde{\mathcal{I}}(\boldsymbol{\eta})\|_2$ and $\lambda_4(\tilde{\mathcal{I}}(\boldsymbol{\eta}))$ is the smallest eigenvalue of $\tilde{\mathcal{I}}(\boldsymbol{\eta})$.

$$\begin{aligned} \|\tilde{\mathcal{I}}(\boldsymbol{\eta})\|_2 &\leq \frac{\kappa^2}{1+|\delta|} \|\tilde{\mathcal{I}}_1(\boldsymbol{\eta})\|_2 + \|\mathcal{I}_2(\boldsymbol{\eta})\|_2 \\ &\leq \frac{\kappa^2}{1+|\delta|} \left\| \begin{pmatrix} \mathbf{G} & c_N(\mathbf{Z}) \mathbf{Z} \boldsymbol{\mu} & -\boldsymbol{\mu} \end{pmatrix} \right\|_2^2 + \|\mathcal{I}_2(\boldsymbol{\eta})\|_F \\ &\leq \frac{\kappa^2}{1+|\delta|} (\|\mathbf{G}\|_2 + c_N(\mathbf{Z}) \|\mathbf{Z} \boldsymbol{\mu}\|_2 + \|\boldsymbol{\mu}\|_2)^2 \\ &\quad + \max\{c_N(\mathbf{Z})^2 |\text{trace}((\boldsymbol{\Sigma} \mathbf{Z})^2)|, c_N(\mathbf{Z}) |\text{trace}(\boldsymbol{\Sigma} \mathbf{Z} \boldsymbol{\Sigma})|, \text{trace}(\boldsymbol{\Sigma}^2)\}. \end{aligned} \quad (11.13)$$

Note that, $\|\mathbf{G}\|_2 \leq \sqrt{2N}$, since all the elements of \mathbf{G} lie within 0 and 1. Also,

$$\begin{aligned} c_N(\mathbf{Z}) \|\mathbf{Z} \boldsymbol{\mu}\|_2 &\leq c_N(\mathbf{Z}) \|\mathbf{Z}\|_2 \|\boldsymbol{\mu}\|_2 \leq \|\boldsymbol{\mu}\|_2, \text{ and,} \\ \|\boldsymbol{\mu}\|_2 &\leq \|(\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1}\|_2 \|\mathbf{G}\|_2 \|\boldsymbol{\vartheta}\|_2 \leq \frac{\sqrt{2N} \|\boldsymbol{\vartheta}\|_2}{1-|\delta|}, \end{aligned} \quad (11.14)$$

since $c_N(\mathbf{Z}) \|\mathbf{Z}\|_2 \leq 1$ and $|\delta| < 1$. So, the first term on the right-hand side of (11.13) is bounded above by $K_1(\boldsymbol{\theta}, \kappa^2) N$, where $K_1(\boldsymbol{\theta}, \kappa^2) := \frac{\kappa^2}{1+|\delta|} \left(\sqrt{2} + \frac{2\sqrt{2} \|\boldsymbol{\vartheta}\|_2}{1-|\delta|} \right)^2$. For the

second term, compute

$$\begin{aligned}
c_N(\mathbf{Z})^2 |\text{trace}((\mathbf{\Sigma} \mathbf{Z})^2)| &\leq N c_N(\mathbf{Z})^2 \|\mathbf{\Sigma} \mathbf{Z}\|_2^2 \leq N c_N(\mathbf{Z})^2 \|\mathbf{Z}\|_2^2 \|\mathbf{\Sigma}\|_2^2 \\
&\leq N \frac{\kappa^4}{(1-|\delta|)^2} \\
c_N(\mathbf{Z}) |\text{trace}(\mathbf{\Sigma} \mathbf{Z} \mathbf{\Sigma})| &\leq N c_N(\mathbf{Z}) \|\mathbf{\Sigma} \mathbf{Z} \mathbf{\Sigma}\|_2 \leq N c_N(\mathbf{Z}) \|\mathbf{Z}\|_2 \|\mathbf{\Sigma}\|_2^2 \quad (11.15) \\
&\leq N \frac{\kappa^4}{(1-|\delta|)^2} \\
\text{trace}(\mathbf{\Sigma}^2) &\leq N \|\mathbf{\Sigma}\|_2^2 \leq N \frac{\kappa^4}{(1-|\delta|)^2},
\end{aligned}$$

using the fact that $\|\mathbf{\Sigma}\|_2 \leq \kappa^2/(1-|\delta|)$ because $c_N(\mathbf{Z}) \|\mathbf{Z}\|_2 \leq 1$ and $|\delta| < 1$. So, the second term on the right-hand side of (11.13) is bounded above by $\frac{\kappa^4}{(1-|\delta|)^2} N$. Therefore,

$$\begin{aligned}
\det(\tilde{\mathcal{I}}(\boldsymbol{\eta})) &\leq \left(K_1(\boldsymbol{\theta}, \kappa^2) + \frac{\kappa^4}{(1-|\delta|)^2} \right)^3 N^3 \lambda_{\min}(\tilde{\mathcal{I}}(\boldsymbol{\eta})) \\
&=: K_2(\boldsymbol{\theta}, \kappa^2) N^3 \lambda_{\min}(\tilde{\mathcal{I}}(\boldsymbol{\eta})). \quad (11.16)
\end{aligned}$$

Lower bound on $\det(\tilde{\mathcal{I}}(\boldsymbol{\eta}))$.

$$\begin{aligned}
&\tilde{\mathcal{I}}(\boldsymbol{\eta}) \\
&= \frac{\kappa^2}{1+|\delta|} \begin{pmatrix} \mathbf{G}^\top \mathbf{G} & c_N(\mathbf{Z}) \mathbf{G}^\top \mathbf{Z} \boldsymbol{\mu} & -\mathbf{G}^\top \boldsymbol{\mu} \\ c_N(\mathbf{Z}) \boldsymbol{\mu}^\top \mathbf{Z} \mathbf{G} & c_N(\mathbf{Z})^2 \boldsymbol{\mu}^\top \mathbf{Z}^2 \boldsymbol{\mu} & -c_N(\mathbf{Z}) \boldsymbol{\mu}^\top \mathbf{Z} \boldsymbol{\mu} \\ -\boldsymbol{\mu}^\top \mathbf{G} & -c_N(\mathbf{Z}) \boldsymbol{\mu}^\top \mathbf{Z} \boldsymbol{\mu} & \boldsymbol{\mu}^\top \boldsymbol{\mu} \end{pmatrix} \\
&+ \frac{1}{2} \begin{pmatrix} \mathbf{0}_{2 \times 2} & \mathbf{0}_{2 \times 1} & \mathbf{0}_{2 \times 1} \\ \mathbf{0}_{1 \times 2} & c_N(\mathbf{Z})^2 \text{trace}((\mathbf{\Sigma} \mathbf{Z})^2) & -c_N(\mathbf{Z}) \text{trace}(\mathbf{\Sigma} \mathbf{Z} \mathbf{\Sigma}) \\ \mathbf{0}_{1 \times 2} & -c_N(\mathbf{Z}) \text{trace}(\mathbf{\Sigma} \mathbf{Z} \mathbf{\Sigma}) & \text{trace}(\mathbf{\Sigma}^2) \end{pmatrix}
\end{aligned}$$

$$= \frac{\kappa^2}{1 + |\delta|} \begin{pmatrix} \mathbf{G}^\top \mathbf{G} & c_N(\mathbf{Z}) \mathbf{G}^\top \mathbf{Z} \boldsymbol{\mu} & -\mathbf{G}^\top \boldsymbol{\mu} \\ c_N(\mathbf{Z}) \boldsymbol{\mu}^\top \mathbf{Z} \mathbf{G} & c_N(\mathbf{Z})^2 \boldsymbol{\mu}^\top \mathbf{Z}^2 \boldsymbol{\mu} + U_{11} & -c_N(\mathbf{Z}) \boldsymbol{\mu}^\top \mathbf{Z} \boldsymbol{\mu} + U_{12} \\ -\boldsymbol{\mu}^\top \mathbf{G} & -c_N(\mathbf{Z}) \boldsymbol{\mu}^\top \mathbf{Z} \boldsymbol{\mu} + U_{12} & \boldsymbol{\mu}^\top \boldsymbol{\mu} + U_{22} \end{pmatrix},$$

where,

$$U_{11} := \frac{(1 + |\delta|) c_N(\mathbf{Z})^2}{2 \kappa^2} \text{trace}((\boldsymbol{\Sigma} \mathbf{Z})^2), \quad U_{12} := -\frac{(1 + |\delta|) c_N(\mathbf{Z})}{2 \kappa^2} \text{trace}(\boldsymbol{\Sigma} \mathbf{Z} \boldsymbol{\Sigma}),$$

$$U_{22} := \frac{1 + |\delta|}{2 \kappa^2} \text{trace}(\boldsymbol{\Sigma}^2).$$

By the formula of the determinant of partitioned matrices (see 1.3, Chapter 3 of Bapat (2012)),

$$\det(\tilde{\mathcal{I}}(\boldsymbol{\eta})) = \frac{\kappa^8}{(1 + |\delta|)^4} \det(\mathbf{G}^\top \mathbf{G}) \det(\mathbf{Q}), \quad (11.17)$$

where

$$\begin{aligned} \mathbf{Q} &:= \begin{pmatrix} c_N(\mathbf{Z})^2 \boldsymbol{\mu}^\top \mathbf{Z}^2 \boldsymbol{\mu} + U_{11} & -c_N(\mathbf{Z}) \boldsymbol{\mu}^\top \mathbf{Z} \boldsymbol{\mu} + U_{12} \\ -c_N(\mathbf{Z}) \boldsymbol{\mu}^\top \mathbf{Z} \boldsymbol{\mu} + U_{12} & \boldsymbol{\mu}^\top \boldsymbol{\mu} + U_{22} \end{pmatrix} \\ &- \begin{pmatrix} c_N(\mathbf{Z}) \boldsymbol{\mu}^\top \mathbf{Z} \mathbf{G} \\ -\boldsymbol{\mu}^\top \mathbf{G} \end{pmatrix}^\top (\mathbf{G}^\top \mathbf{G})^{-1} \begin{pmatrix} c_N(\mathbf{Z}) \mathbf{G}^\top \mathbf{Z} \boldsymbol{\mu} & -\mathbf{G}^\top \boldsymbol{\mu} \end{pmatrix} \\ &= \begin{pmatrix} c_N(\mathbf{Z})^2 \boldsymbol{\mu}^\top \mathbf{Z} (\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu} + U_{11} & -c_N(\mathbf{Z}) \boldsymbol{\mu}^\top (\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu} + U_{12} \\ -c_N(\mathbf{Z}) \boldsymbol{\mu}^\top (\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu} + U_{12} & \boldsymbol{\mu}^\top (\mathbf{I}_N - \mathbf{H}) \boldsymbol{\mu} + U_{22} \end{pmatrix}, \end{aligned}$$

and $\mathbf{H} := \mathbf{G}(\mathbf{G}^\top \mathbf{G})^{-1} \mathbf{G}^\top$.

Lower bound on $\det(\mathbf{G}^\top \mathbf{G})$. Let $N_i := \sum_{j=1}^N Z_{i,j}$ and $F_i := \sum_{j=1}^N W_j Z_{i,j} / N_i$ for all

$i = 1, \dots, N$. Then,

$$\mathbf{G}^\top \mathbf{G} = \begin{pmatrix} \sum_{i=1}^N W_i & \sum_{i=1}^N W_i F_i \\ \sum_{i=1}^N W_i F_i & \sum_{i=1}^N F_i^2 \end{pmatrix},$$

and,

$$\det(\mathbf{G}^\top \mathbf{G}) = \sum_{i=1}^N W_i \sum_{i=1}^N F_i^2 - \left(\sum_{i=1}^N W_i F_i \right)^2. \quad (11.18)$$

We show that $\det(\mathbf{G}^\top \mathbf{G})$ is of order N^2 . First, using the fact that $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N} \cap \mathcal{A}_{2,N}$

and $N\rho_N \geq D \log N$, we obtain that $|F_i - \pi| \leq \sqrt{3/d_1 D}$ for all i . So,

$$\sum_{i=1}^N F_i^2 = \sum_{i=1}^N (F_i - \pi)^2 + 2\pi \sum_{i=1}^N (F_i - \pi) + N\pi^2 = N\pi^2 + R_1,$$

where $|R_1| \leq (3/d_1 D)N + 2\pi \sqrt{3/d_1 D}N$. In addition,

$$\sum_{i=1}^N W_i F_i = \sum_{i=1}^N W_i (F_i - \pi) + \pi \sum_{i=1}^N W_i = \pi \sum_{i=1}^N W_i + R_2,$$

where $|R_2| \leq \sqrt{3/d_1 D}N$.

Therefore,

$$\begin{aligned} & \sum_{i=1}^N W_i \sum_{i=1}^N F_i^2 - \left(\sum_{i=1}^N W_i F_i \right)^2 \\ &= \left(\sum_{i=1}^N W_i \right) (N\pi^2 + R_1) - \left(\pi \sum_{i=1}^N W_i + R_2 \right)^2 \\ &= N\pi^2 \sum_{i=1}^N W_i + R_1 \sum_{i=1}^N W_i - \pi^2 \left(\sum_{i=1}^N W_i \right)^2 - 2\pi R_2 \sum_{i=1}^N W_i - R_2^2 \\ &= \pi^2 \sum_{i=1}^N W_i \left(N - \sum_{i=1}^N W_i \right) + R_3, \end{aligned} \quad (11.19)$$

where

$$|R_3| \leq 2(1+2\pi)N^2 \max\left\{\frac{3}{d_1 D}, \sqrt{\frac{3}{d_1 D}}\right\}.$$

We showed in (10.12) that if $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{2,N}$,

$$\sum_{i=1}^N W_i \left(N - \sum_{i=1}^N W_i \right) \geq \frac{N^2 \pi (1 - \pi)}{4} \quad (11.20)$$

for all sufficiently large N .

Combining (11.18), (11.19) and (11.20), we obtain

$$\det(\mathbf{G}^\top \mathbf{G}) \geq \frac{N^2 \pi^3 (1 - \pi)}{4} - 2(1+2\pi)N^2 \max\left\{\frac{3}{d_1 D}, \sqrt{\frac{3}{d_1 D}}\right\}.$$

Recall that $d_1 := p_{\min} (1 - \sqrt{6/p_{\min} D})$. Applying the condition $D > 1536 (1+2\pi)^2 / p_{\min} \pi^6 (1 - \pi)^2$, we obtain

$$\det(\mathbf{G}^\top \mathbf{G}) \geq \frac{N^2 \pi^3 (1 - \pi)}{8}. \quad (11.21)$$

Lower bound on $\det(\mathbf{Q})$. Define

$$V_{11} := c_N(\mathbf{Z})^2 \boldsymbol{\mu}^\top \mathbf{Z}(\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu}, \quad V_{12} := -c_N(\mathbf{Z}) \boldsymbol{\mu}^\top (\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu},$$

$$V_{22} := \boldsymbol{\mu}^\top (\mathbf{I}_N - \mathbf{H}) \boldsymbol{\mu}.$$

Then, we can write the matrix \mathbf{Q} as

$$\mathbf{Q} = \begin{pmatrix} V_{11} + U_{11} & V_{12} + U_{12} \\ V_{11} + U_{12} & V_{22} + U_{22} \end{pmatrix}, \quad (11.22)$$

and,

$$\begin{aligned}
\det(\mathbf{Q}) &= (V_{11} + U_{11})(V_{22} + U_{22}) - (V_{12} + U_{12})^2 \\
&= (V_{11} V_{22} - V_{12}^2) + V_{11} U_{22} + U_{11} (V_{22} + U_{22}) - 2 V_{12} U_{12} - U_{12}^2.
\end{aligned} \tag{11.23}$$

We first show that $V_{11} V_{22} - V_{12}^2 = 0$. From definition,

$$\begin{aligned}
&V_{11} V_{22} - V_{12}^2 \\
&= c_N(\mathbf{Z})^2 (\boldsymbol{\mu}^\top \mathbf{Z} (\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu} \boldsymbol{\mu}^\top (\mathbf{I}_N - \mathbf{H}) \boldsymbol{\mu} - (\boldsymbol{\mu}^\top (\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu})^2).
\end{aligned}$$

Now,

$$\begin{aligned}
\mathbf{Z} \boldsymbol{\mu} &= \mathbf{Z} (\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{G} \boldsymbol{\vartheta} \\
&= \frac{1}{c_N(\mathbf{Z}) \delta} (\delta c_N(\mathbf{Z}) \mathbf{Z} - \mathbf{I}_N + \mathbf{I}_N) (\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{G} \boldsymbol{\vartheta} \\
&= \frac{1}{c_N(\mathbf{Z}) \delta} ((\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} - \mathbf{I}_N) \mathbf{G} \boldsymbol{\vartheta} \\
&= \frac{1}{c_N(\mathbf{Z}) \delta} (\boldsymbol{\mu} - \mathbf{G} \boldsymbol{\vartheta}).
\end{aligned} \tag{11.24}$$

Thereby, noting that $(\mathbf{I}_N - \mathbf{H}) \mathbf{G}$ is $\mathbf{0}_{N \times 2}$, we obtain

$$(\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu} = \frac{1}{c_N(\mathbf{Z}) \delta} (\mathbf{I}_N - \mathbf{H}) \boldsymbol{\mu},$$

which implies,

$$V_{11} V_{22} - V_{12}^2 = \frac{1}{\delta^2} (\boldsymbol{\mu}^\top (\mathbf{I}_N - \mathbf{H}) \boldsymbol{\mu} \cdot \boldsymbol{\mu}^\top (\mathbf{I}_N - \mathbf{H}) \boldsymbol{\mu} - (\boldsymbol{\mu}^\top (\mathbf{I}_N - \mathbf{H}) \boldsymbol{\mu})^2) = 0. \tag{11.25}$$

Note that, in the above derivation, we assumed $\delta \neq 0$. If $\delta = 0$, then $(\mathbf{I}_N - \mathbf{H}) \boldsymbol{\mu} =$

$(\mathbf{I}_N - \mathbf{H}) \mathbf{G} \boldsymbol{\vartheta} = 0$, which implies $V_{11} V_{22} - V_{12}^2 = 0$. So, from (11.23), we obtain

$$\det(\mathbf{Q}) = V_{11} U_{22} + U_{11} (V_{22} + U_{22}) - 2 V_{12} U_{12} - U_{12}^2. \quad (11.26)$$

Upper bounds on $|V_{11}|$, $|V_{12}|$ and $|V_{22}|$. Based on our previous calculations, we argue that $|V_{11}|$, $|V_{12}|$ and $|V_{22}|$ are $O(N)$. First, noting that $\|\mathbf{I}_N - \mathbf{H}\|_2 \leq 1$, we obtain

$$|V_{11}| \leq c_N(\mathbf{Z})^2 \boldsymbol{\mu}^\top \mathbf{Z}^2 \boldsymbol{\mu}, \quad |V_{12}| \leq c_N(\mathbf{Z}) |\boldsymbol{\mu}^\top \mathbf{Z} \boldsymbol{\mu}|, \quad |V_{22}| \leq \boldsymbol{\mu}^\top \boldsymbol{\mu}.$$

Observe that the upper bounds on $|V_{11}|$, $|V_{12}|$ and $|V_{22}|$ are elements (in absolute value) of $\tilde{\mathcal{I}}_1(\boldsymbol{\eta})$. We showed in (11.14) that $\|\tilde{\mathcal{I}}_1(\boldsymbol{\eta})\|_2$ is $O(N)$, which implies, the elements of $\tilde{\mathcal{I}}_1(\boldsymbol{\eta})$ in absolute value has to be $O(N)$. Therefore, $|V_{11}|$, $|V_{12}|$ and $|V_{22}|$ are also $O(N)$.

Upper bounds on $|U_{11}|$, $|U_{12}|$ and $|U_{22}|$. We show that

$$\begin{aligned} |U_{11}| &= o(c_N(\mathbf{Z})^2 N (N \rho_N)^2), \quad |U_{12}| = o(c_N(\mathbf{Z}) N (N \rho_N)), \\ |U_{22}| &= O(N). \end{aligned} \quad (11.27)$$

$$\begin{aligned} |U_{11}| &= \frac{(1 + |\delta|) c_N(\mathbf{Z})^2}{2 \kappa^2} |\text{trace}((\boldsymbol{\Sigma} \mathbf{Z})^2)| \\ &= \frac{(1 + |\delta|) c_N(\mathbf{Z})^2}{2 \kappa^2} |\text{trace}((\boldsymbol{\Sigma} (\mathbf{Z} - \mathbf{P}) + \boldsymbol{\Sigma} \mathbf{P})^2)| \\ &= \frac{(1 + |\delta|) c_N(\mathbf{Z})^2}{2 \kappa^2} |\text{trace}((\boldsymbol{\Sigma} (\mathbf{Z} - \mathbf{P}))^2 + 2 \boldsymbol{\Sigma} (\mathbf{Z} - \mathbf{P}) \boldsymbol{\Sigma} \mathbf{P} + (\boldsymbol{\Sigma} \mathbf{P})^2)| \\ &\leq \frac{(1 + |\delta|) c_N(\mathbf{Z})^2}{2 \kappa^2} (N \|\boldsymbol{\Sigma}\|_2^2 \|\mathbf{Z} - \mathbf{P}\|_2^2 + 2 N \|\boldsymbol{\Sigma}\|_2^2 \|\mathbf{Z} - \mathbf{P}\|_2 \|\mathbf{P}\|_2 \\ &\quad + \text{trace}(\boldsymbol{\Sigma} \mathbf{P})^2) \\ |U_{12}| &= \frac{(1 + |\delta|) c_N(\mathbf{Z})}{2 \kappa^2} |\text{trace}(\boldsymbol{\Sigma} \mathbf{Z} \boldsymbol{\Sigma})| \end{aligned}$$

$$\begin{aligned}
&= \frac{(1 + |\delta|) c_N(\mathbf{Z})}{2 \kappa^2} |\text{trace}(\mathbf{\Sigma} (\mathbf{Z} - \mathbf{P}) \mathbf{\Sigma} + \mathbf{\Sigma} \mathbf{P} \mathbf{\Sigma})| \\
&\leq \frac{(1 + |\delta|) c_N(\mathbf{Z})}{2 \kappa^2} (N \|\mathbf{\Sigma}\|_2^2 \|\mathbf{Z} - \mathbf{P}\|_2 + \text{trace}(\mathbf{\Sigma} \mathbf{P} \mathbf{\Sigma})) \\
|U_{22}| &= \frac{(1 + |\delta|)}{2 \kappa^2} |\text{trace}(\mathbf{\Sigma}^2)|.
\end{aligned}$$

Observe that

$$\begin{aligned}
|\text{trace}((\mathbf{\Sigma} \mathbf{P} \mathbf{\Sigma} \mathbf{P}))| &\leq \|\mathbf{\Sigma}\|_2^2 \|\mathbf{P}\|_2 \text{trace}(\mathbf{P}) \leq K \|\mathbf{\Sigma}\|_2^2 \|\mathbf{P}\|_2^2, \\
|\text{trace}(\mathbf{\Sigma} \mathbf{P} \mathbf{\Sigma})| &\leq \|\mathbf{\Sigma}\|_2^2 \text{trace}(\mathbf{P}) \leq K \|\mathbf{\Sigma}\|_2^2 \|\mathbf{P}\|_2 \\
|\text{trace}(\mathbf{\Sigma}^2)| &\leq N \|\mathbf{\Sigma}\|_2^2.
\end{aligned}$$

The inequalities above are obtained using the result that for matrices \mathbf{A} and \mathbf{B} where \mathbf{B} is positive definite, $\text{trace}(\mathbf{A} \mathbf{B}) \leq \|\mathbf{A}\|_2 \text{trace}(\mathbf{B})$. In our case, \mathbf{P} is positive definite and has rank K .

Now, $\|\mathbf{Z} - \mathbf{P}\|_2 \leq l \sqrt{N \rho_N}$ since $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N}$, $\|\mathbf{P}\|_2 \leq \|\mathbf{P}\|_F \leq N \rho_N$, and $\|\mathbf{\Sigma}\|_2 \leq \kappa^2 / (1 - |\delta|)$. We leverage these results to obtain

$$\begin{aligned}
|U_{11}| &\leq \frac{(1 + |\delta|) c_N(\mathbf{Z})^2}{2 \kappa^2} \left(\frac{\kappa^4}{(1 - |\delta|)^2} (l^2 N (N \rho_N) + 2 l N (N \rho_N)^{\frac{3}{2}} + K (N \rho_N)^2) \right) \\
|U_{12}| &\leq \frac{(1 + |\delta|) c_N(\mathbf{Z})}{2 \kappa^2} \left(\frac{\kappa^4}{(1 - |\delta|)^2} (l N \sqrt{N \rho_N} + K (N \rho_N)) \right) \\
|U_{22}| &\leq \frac{(1 + |\delta|) c_N(\mathbf{Z})}{2 \kappa^2} N \frac{\kappa^4}{(1 - |\delta|)^2}.
\end{aligned}$$

So, we conclude that (11.27) holds since $N \rho_N \rightarrow \infty$ due to Condition C.1. This result, together with the fact that $|V_{12}|$ and $|V_{22}|$ are $O(N)$, gives that in (11.26), the terms $|U_{11} (V_{22} + U_{22})|$ and U_{12}^2 are $o(c_N(\mathbf{Z})^2 N^2 (N \rho_N)^2)$. We would like to show that the term $|V_{12} U_{12}|$ also has the same order of magnitude, but for that, we need to prove a finer bound

on $|V_{12}|$ instead of $O(N)$.

Upper bound on $|V_{12} U_{12}|$.

$$\begin{aligned} |V_{12}| &\leq c_N(\mathbf{Z}) \|\mathbf{Z}\|_2 \|\boldsymbol{\mu}\|_2^2 \leq c_N(\mathbf{Z}) (\|\mathbf{Z} - \mathbf{P}\|_2 + \|\mathbf{P}\|_2) \|\boldsymbol{\mu}\|_2^2 \\ &= O(c_N(\mathbf{Z}) N (N\rho_N)), \end{aligned}$$

since $\|\mathbf{Z} - \mathbf{P}\|_2 \leq l\sqrt{N\rho_N}$, $\|\mathbf{P}\|_2 \leq N\rho_N$, and $\|\boldsymbol{\mu}\|_2^2 = O(N)$ from (11.14). This proves that $V_{12} U_{12}$ is also $o(c_N(\mathbf{Z})^2 N^2 (N\rho_N)^2)$.

So, from (11.26), we obtain

$$\det(\mathbf{Q}) \geq V_{11} U_{22} - o(c_N(\mathbf{Z})^2 N^2 (N\rho_N)^2). \quad (11.28)$$

Lower bounds on V_{11} and U_{22} . We show that both V_{11} and U_{22} has a lower bound of order N .

$$V_{11} := c_N(\mathbf{Z})^2 \boldsymbol{\mu}^\top \mathbf{Z} (\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu} := c_N(\mathbf{Z})^2 \|(\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu}\|_2^2. \quad (11.29)$$

We derive a lower bound on $\|(\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu}\|_2$.

$$\|(\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu}\|_2 \geq \|(\mathbf{I}_N - \mathbf{H}) \mathbf{P} \boldsymbol{\mu}\|_2 - \|(\mathbf{Z} - \mathbf{P}) \boldsymbol{\mu}\|_2,$$

using the fact that $\|\mathbf{I}_N - \mathbf{H}\|_2 \leq 1$. Now, since $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{3,N}$,

$$\begin{aligned} \|(\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu}\|_2 &\geq \frac{1 - \sqrt{(1+p^*)/2}}{2} \|\mathbf{P} \boldsymbol{\mu}\|_2 - \|(\mathbf{Z} - \mathbf{P}) \boldsymbol{\mu}\|_2 \\ &\geq \frac{1 - \sqrt{(1+p^*)/2}}{2} \|\mathbf{Z} \boldsymbol{\mu}\|_2 - \frac{3 - \sqrt{(1+p^*)/2}}{2} \|(\mathbf{Z} - \mathbf{P}) \boldsymbol{\mu}\|_2 \\ &\geq \frac{1 - \sqrt{(1+p^*)/2}}{2} \|\mathbf{Z} \boldsymbol{\mu}\|_2 - \frac{3}{2} \|\mathbf{Z} - \mathbf{P}\|_2 \|\boldsymbol{\mu}\|_2. \end{aligned}$$

Also, since $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{1,N}$, we have

$$\|(\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu}\|_2 \geq \frac{1 - \sqrt{(1+p^*)/2}}{2} \|\mathbf{Z} \boldsymbol{\mu}\|_2 - \frac{3l}{2} \sqrt{N\rho_N} \|\boldsymbol{\mu}\|_2. \quad (11.30)$$

Hence, we can obtain a lower bound on $\|(\mathbf{I}_N - \mathbf{H}) \mathbf{Z} \boldsymbol{\mu}\|_2$ by deriving a lower bound on $\|\mathbf{Z} \boldsymbol{\mu}\|_2$ and an upper bound on $\|\boldsymbol{\mu}\|_2$.

$$\begin{aligned} \|\mathbf{Z} \boldsymbol{\mu}\|_2^2 &= \boldsymbol{\vartheta}^\top \mathbf{G}^\top (\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{Z}^2 (\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{G} \boldsymbol{\vartheta} \\ &= \boldsymbol{\vartheta}^\top \mathbf{G}^\top \mathbf{Z} (\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-2} \mathbf{Z} \mathbf{G} \boldsymbol{\vartheta}, \end{aligned}$$

since $(\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1}$ and \mathbf{Z} are commutative (11.31)

$$\begin{aligned} &\geq \frac{1}{(1+|\delta|)^2} \boldsymbol{\vartheta}^\top \mathbf{G}^\top \mathbf{Z}^2 \mathbf{G} \boldsymbol{\vartheta} \\ &\geq \frac{(\beta + \gamma)^2 \pi^2 p_{\min}^2 N (N\rho_N)^2}{16(1+|\delta|)^2}. \end{aligned}$$

The first inequality holds because $c_N(\mathbf{Z}) \|\mathbf{Z}\|_2 \leq 1$ almost surely, and the second inequality holds because $(\mathbf{W}, \mathbf{Z}) \in \mathcal{A}_{4,N}$. We also showed in (11.14) that

$$\|\boldsymbol{\mu}\|_2 \leq \frac{\sqrt{2N} \|\boldsymbol{\vartheta}\|_2}{1 - |\delta|}. \quad (11.32)$$

Combining (11.30), (11.31) and (11.32), we obtain

$$\begin{aligned}
& \|(\mathbf{I}_N - \mathbf{H})\mathbf{Z}\boldsymbol{\mu}\|_2 \\
& \geq \frac{1 - \sqrt{(1 + p^*)/2}}{2} \frac{|\beta + \gamma| \pi p_{\min}}{4(1 + |\delta|)} \sqrt{N} N \rho_N - \frac{3l}{\sqrt{2}} \frac{\|\boldsymbol{\vartheta}\|_2}{1 - |\delta|} \sqrt{N^2 \rho_N} \\
& \geq \frac{1 - \sqrt{(1 + p^*)/2}}{2} \frac{|\beta + \gamma| \pi p_{\min}}{8(1 + |\delta|)} \sqrt{N} N \rho_N,
\end{aligned}$$

for all sufficiently large N , because $N \rho_N \rightarrow \infty$ by Condition C.1. Leveraging this in (11.29), we obtain

$$V_{11} \geq (1 - \sqrt{(1 + p^*)/2})^2 \frac{(\beta + \gamma)^2 \pi^2 p_{\min}^2}{256(1 + |\delta|)^2} c_N(\mathbf{Z})^2 N (N \rho_N)^2. \quad (11.33)$$

Since the eigenvalues of $\boldsymbol{\Sigma}$ are at least $\kappa^2 / (1 + |\delta|)$, we have

$$U_{22} = \frac{1 + |\delta|}{2 \kappa^2} \text{trace}(\boldsymbol{\Sigma}^2) \geq \frac{1 + |\delta|}{2 \kappa^2} N \frac{\kappa^4}{(1 + |\delta|)^2} = \frac{\kappa^2}{2(1 + |\delta|)} N. \quad (11.34)$$

Combining (11.28), (11.33) and (11.34), we obtain

$$\begin{aligned}
\det(\mathbf{Q}) & \geq (1 - \sqrt{(1 + p^*)/2})^2 \frac{(\beta + \gamma)^2 \pi^2 p_{\min}^2}{256(1 + |\delta|)^2} \frac{\kappa^2}{2(1 + |\delta|)} c_N(\mathbf{Z})^2 N^2 (N \rho_N)^2 \\
& - o(c_N(\mathbf{Z})^2 N^2 (N \rho_N)^2) \\
& \geq K_3(\boldsymbol{\theta}, \kappa^2) c_N(\mathbf{Z})^2 N^2 (N \rho_N)^2,
\end{aligned} \quad (11.35)$$

where $K_3(\boldsymbol{\theta}, \kappa^2) > 0$ is a constant that depends on $\boldsymbol{\theta}$ and κ^2 .

Combining (11.17), (11.21) and (11.35) yields

$$\begin{aligned} \det(\tilde{\mathcal{I}}(\boldsymbol{\eta})) &\geq \frac{\kappa^8}{(1+|\delta|)^4} \frac{N^2 \pi^3 (1-\pi)}{8} K_3(\boldsymbol{\theta}, \kappa^2) c_N(\mathbf{Z})^2 N^2 (N\rho_N)^2 \\ &=: K_4(\boldsymbol{\theta}, \kappa^2) c_N(\mathbf{Z})^2 N^4 (N\rho_N)^2. \end{aligned} \quad (11.36)$$

Lower bound on $\lambda_{\min}(\tilde{\mathcal{I}}(\boldsymbol{\eta}))$. From the lower and upper bound on $\det(\tilde{\mathcal{I}}(\boldsymbol{\eta}))$ derived in (11.16) and (11.36) respectively, we obtain

$$\lambda_{\min}(\tilde{\mathcal{I}}(\boldsymbol{\eta})) \geq \frac{K_4(\boldsymbol{\theta}, \kappa^2)}{K_2(\boldsymbol{\theta}, \kappa^2)} c_N(\mathbf{Z})^2 N (N\rho_N)^2. \quad (11.37)$$

Leveraging (11.11), we conclude

$$\|\mathcal{I}(\boldsymbol{\eta})^{-1}\|_2 \leq \frac{K_2(\boldsymbol{\theta}, \kappa^2)}{K_4(\boldsymbol{\theta}, \kappa^2)} \frac{1}{c_N(\mathbf{Z})^2 N (N\rho_N)^2},$$

where

$$\begin{aligned} K_2(\boldsymbol{\theta}, \kappa^2) &:= \left(\frac{\kappa^2}{1+|\delta|} \left(\sqrt{2} + \frac{2\sqrt{2} \|\boldsymbol{\vartheta}\|_2}{1-|\delta|} \right)^2 + \frac{\kappa^4}{(1-|\delta|)^2} \right)^3 \\ K_4(\boldsymbol{\theta}, \kappa^2) &:= \frac{\kappa^8}{(1+|\delta|)^4} \frac{\pi^3 (1-\pi)}{8} (1 - \sqrt{(1+p^*)/2})^2 \frac{(\beta + \gamma)^2 \pi^2 p_{\min}^2}{256 (1+|\delta|)^2} \frac{\kappa^2}{4(1+|\delta|)}. \end{aligned}$$

11.3 Lemma 3

Lemma 3. Assume that Condition C.1 holds. Then:

(a) If the constant D in Condition C.1 satisfies $D > 6/p_{\min}$, then

$$\mathbb{P} \left(\min_{1 \leq i \leq N} \sum_{j=1}^N Z_{i,j} > p_{\min} \left(1 - \sqrt{\frac{6}{p_{\min} D}} \right) N\rho_N \right) \geq 1 - \frac{1}{N^2}.$$

(b) The limits of the following terms vanish:

$$\lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E}[\|\text{trace}((\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} - (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1})\|] = 0, \quad (11.38)$$

$$\lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E}[\|\mathbf{1}_N^\top ((\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} - (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1}) \mathbf{1}_N\|] = 0, \quad (11.39)$$

assuming that $c_N(\mathbf{Z}) := (1 / \|\mathbf{Z}\|_2) \mathbb{I}(\|\mathbf{Z}\|_2 > 0)$, $c_N(\mathbf{P}) := 1 / \|\mathbf{P}\|_2$, and $|\delta| < 1$.

Proof of Lemma 3. $Z_{i,j} \stackrel{\text{ind}}{\sim} \text{Bernoulli}(P_{i,j})$ implies that $N_i := \sum_{j=1: j \neq i}^N Z_{i,j}$ is a sum of $N - 1$ independent Bernoulli variables with expectation $\sum_{j=1: j \neq i}^N P_{i,j}$. By a Chernoff bound, for any $t \in (0, 1)$,

$$\mathbb{P}\left(N_i \geq (1 - t) \sum_{j=1: j \neq i}^N P_{i,j}\right) \geq 1 - \exp\left(-\frac{t^2 \sum_{j=1: j \neq i}^N P_{i,j}}{2}\right).$$

From Condition C.1, we have $\sum_{j=1: j \neq i}^N P_{i,j} > p_{\min} N \rho_N$, so that

$$\mathbb{P}(N_i > (1 - t) p_{\min} N \rho_N) \geq 1 - \exp\left(-\frac{t^2 p_{\min} N \rho_N}{2}\right).$$

Choosing

$$t := \sqrt{\frac{6 \log N}{p_{\min} N \rho_N}}$$

and using the fact that $N \rho_N \geq D \log N$, we obtain

$$\mathbb{P}\left(N_i > p_{\min} (1 - \sqrt{6 / p_{\min} D}) N \rho_N\right) \geq 1 - \frac{1}{N^3}.$$

Taking a union bound over all $i \in \{1, \dots, N\}$ establishes (a).

We proceed with (b). In Equation (11.38),

$$\begin{aligned} & \frac{1}{N} |\text{trace}((\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} - (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1})| \\ & \leq \quad \|(\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} - (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1}\|_2, \end{aligned} \quad (11.40)$$

since for any symmetric matrix $\mathbf{A} \in \mathbb{R}^{N \times N}$, $\text{trace}(\mathbf{A}) = \sum_{i=1}^N \lambda_i(\mathbf{A})$ and $|\lambda_i(\mathbf{A})| \leq \|\mathbf{A}\|_2$.

In Equation (11.39),

$$\begin{aligned} & \frac{1}{N} |\mathbf{1}_N^\top ((\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} - (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1}) \mathbf{1}_N| \\ & \leq \quad \|(\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} - (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1}\|_2, \end{aligned} \quad (11.41)$$

which follows from the definition of the $\|\cdot\|_2$ - norm. Therefore, we focus on the term

$$\|(\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} - (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1}\|_2.$$

$$\begin{aligned} & \|(\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} - (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1}\|_2 \\ & = \quad \|(\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1} (\delta c_N(\mathbf{Z}) \mathbf{Z} - \delta c_N(\mathbf{P}) \mathbf{P}) (\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1}\|_2 \\ & \quad \text{since } \mathbf{A}^{-1} - \mathbf{B}^{-1} = \mathbf{B}^{-1}(\mathbf{B} - \mathbf{A}) \mathbf{A}^{-1} \\ & \leq \quad \|(\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1}\|_2 \|\delta c_N(\mathbf{Z}) \mathbf{Z} - \delta c_N(\mathbf{P}) \mathbf{P}\|_2 \|(\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1}\|_2. \end{aligned} \quad (11.42)$$

We bound each of three terms from above.

Since $|\delta| < 1$ and $c_N(\mathbf{Z}) \|\mathbf{Z}\|_2 \leq 1$,

$$\|(\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1}\|_2 = \max_{1 \leq i \leq N} \frac{1}{1 - \delta c_N(\mathbf{Z}) \lambda_i(\mathbf{Z})} \leq \frac{1}{1 - |\delta|}. \quad (11.43)$$

Along the same lines, we can show that

$$\|(\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1}\|_2 \leq \frac{1}{1 - |\delta|}. \quad (11.44)$$

To bound $\|\delta c_N(\mathbf{Z}) \mathbf{Z} - \delta c_N(\mathbf{P}) \mathbf{P}\|_2$ from above, first note that

$$\|\delta c_N(\mathbf{Z}) \mathbf{Z} - \delta c_N(\mathbf{P}) \mathbf{P}\|_2 \leq |\delta| c_N(\mathbf{Z}) \|\mathbf{Z}\|_2 + |\delta| c_N(\mathbf{P}) \|\mathbf{P}\|_2 \leq 2. \quad (11.45)$$

Next, we obtain an upper bound on $\|\delta c_N(\mathbf{Z}) \mathbf{Z} - \delta c_N(\mathbf{P}) \mathbf{P}\|_2$ that holds with high probability:

$$\begin{aligned} \|\delta c_N(\mathbf{Z}) \mathbf{Z} - \delta c_N(\mathbf{P}) \mathbf{P}\|_2 &= \|(\delta c_N(\mathbf{Z}) - \delta c_N(\mathbf{P})) \mathbf{Z} + \delta c_N(\mathbf{P}) (\mathbf{Z} - \mathbf{P})\|_2 \\ &\leq |\delta c_N(\mathbf{Z}) - \delta c_N(\mathbf{P})| \|\mathbf{Z}\|_2 + |\delta c_N(\mathbf{P})| \|\mathbf{Z} - \mathbf{P}\|_2 \\ &\leq \frac{|\delta| |\|\mathbf{Z}\|_2 - \|\mathbf{P}\|_2|}{\|\mathbf{Z}\|_2 \|\mathbf{P}\|_2} \|\mathbf{Z}\|_2 \mathbb{I}(\|\mathbf{Z}\|_2 > 0) + \frac{|\delta|}{\|\mathbf{P}\|_2} \|\mathbf{Z} - \mathbf{P}\|_2 \\ &\leq \frac{2 \|\mathbf{Z} - \mathbf{P}\|_2}{\|\mathbf{P}\|_2}. \end{aligned}$$

In the denominator, we have

$$\|\mathbf{P}\|_2 = \rho_N \|\boldsymbol{\alpha} \boldsymbol{\alpha}^\top\|_2 = \rho_N \|\boldsymbol{\alpha}\|_2^2 \geq (\alpha_{\min})^2 N \rho_N, \quad (11.46)$$

applying Condition C.1. In the numerator, using the fact that \mathbf{Z} is generated by an undirected Bernoulli random graph, we invoke Theorem 5.2 of Lei and Rinaldo (2015) to bound $\|\mathbf{Z} - \mathbf{P}\|_2$ from above. Since $N \max_{1 \leq i < j \leq N} P_{i,j} \leq N \rho_N$ and $N \rho_N \geq D \log N$, there

exists a constant $l > 0$ such that

$$\mathbb{P}(\|\mathbf{Z} - \mathbb{E}[\mathbf{Z}]\|_2 \leq (l/2) \sqrt{N\rho_N}) \geq 1 - \frac{1}{N^2}.$$

Now, we note that $\mathbf{P} = \mathbb{E}[\mathbf{Z}] + \text{diag}(\mathbf{P})$. So

$$\|\mathbf{Z} - \mathbf{P}\|_2 \leq \|\mathbf{Z} - \mathbb{E}[\mathbf{Z}]\|_2 + \|\text{diag}(\mathbf{P})\|_2 \leq \|\mathbf{Z} - \mathbb{E}[\mathbf{Z}]\|_2 + 1.$$

Since $N\rho_N > 1$ by Condition C.1, we conclude that

$$\mathbb{P}(\|\mathbf{Z} - \mathbf{P}\|_2 \leq l \sqrt{N\rho_N}) \geq 1 - \frac{1}{N^2}. \quad (11.47)$$

Thus,

$$\mathbb{P}\left(\|\delta c_N(\mathbf{Z}) \mathbf{Z} - \delta c_N(\mathbf{P}) \mathbf{P}\|_2 \leq \frac{2l}{\alpha_{\min}^2 \sqrt{N\rho_N}}\right) \geq 1 - \frac{1}{N^2}. \quad (11.48)$$

Combining (11.42), (11.43), (11.44), (11.45), and (11.48), we obtain

$$\begin{aligned} & \left\| (\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} - (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1} \right\|_2 \leq \frac{2}{(1 - |\delta|)^2} \\ \mathbb{P}\left(\left\| (\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} - (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1} \right\|_2 \leq \frac{2l}{\alpha_{\min}^2 \sqrt{N\rho_N} (1 - |\delta|)^2}\right) & \geq 1 - \frac{1}{N^2}. \end{aligned}$$

Therefore,

$$\begin{aligned} & \mathbb{E} \left[\left\| (\mathbf{I}_N - \delta c_N(\mathbf{Z}) \mathbf{Z})^{-1} - (\mathbf{I}_N - \delta c_N(\mathbf{P}) \mathbf{P})^{-1} \right\|_2 \right] \\ & \leq \frac{2l}{\alpha_{\min}^2 \sqrt{N\rho_N} (1 - |\delta|)^2} \left(1 - \frac{1}{N^2} \right) + \frac{2}{(1 - |\delta|)^2} \frac{1}{N^2} \rightarrow 0 \text{ as } N \rightarrow \infty, \end{aligned}$$

because $N\rho_N \rightarrow \infty$ by Condition C.1. The above result, along with (11.40) and (11.41),

completes the proof of (b).

11.4 Lemma 4

Lemma 4. (Theorem 5.2.15 of Vershynin (2018, p. 109)) *Suppose that \mathbf{X} is a N -dimensional random vector with a density (dominated by the Lebesgue measure on the Borel sets of \mathbb{R}^N) of the form*

$$f_{\mathbf{X}}(\mathbf{x}) = \exp(-U(\mathbf{x})),$$

where $U : \mathbb{R}^N \mapsto \mathbb{R}$ is a twice differentiable function with respect to \mathbf{x} for all $\mathbf{x} \in \mathbb{R}^N$. Assume that there exists $\lambda > 0$ such that the smallest eigenvalue of $\nabla_{\mathbf{x}}^2 U(\mathbf{x})$ is bounded below by λ for all $\mathbf{x} \in \mathbb{R}^N$. Then, for any Lipschitz function $g : \mathbb{R}^N \mapsto \mathbb{R}$ and any $t > 0$,

$$\mathbb{P}_{\mathbf{X}}(|g(\mathbf{X}) - \mathbb{E}[g(\mathbf{X})]| > t) \leq 2 \exp\left(-\frac{v_1 \lambda t^2}{\|g\|_{Lip}^2}\right),$$

where $\|g\|_{Lip}$ is the Lipschitz constant of g and $v_1 > 0$ is a constant, independent of N .

11.5 Lemma 5

Lemma 5. Suppose that the conditional distribution of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z})$ is generated by a Gaussian Markov random field with parameters $\boldsymbol{\theta}^* := (\beta^*, \gamma^*, \delta^*) \in \boldsymbol{\Theta}$ and $\kappa^{*2} \in (0, \infty)$. Then, for all $t > 0$,

$$\begin{aligned} & \mathbb{P}_{\mathbf{y}|\mathbf{w},\mathbf{z}}\left(\|t(\mathbf{Y}) - \mathbb{E}_{\mathbb{P}_{\mathbf{y}|\mathbf{w},\mathbf{z}}}[t(\mathbf{Y})]\|_2 \geq t\right) \\ & \leq 8 \exp\left(-\min\left\{\frac{B_1(\boldsymbol{\theta}^*, \kappa^{*2}) t^2}{N}, B_2(\boldsymbol{\theta}^*, \kappa^{*2}) t\right\}\right), \end{aligned}$$

where $B_1(\boldsymbol{\theta}^*, \kappa^{\star 2}), B_2(\boldsymbol{\theta}^*, \kappa^{\star 2}) > 0$ are constants that depend on $(\boldsymbol{\theta}^*, \kappa^{\star 2})$.

Proof of Lemma 5. Recall that

$$t(\mathbf{Y}) := (\mathbf{Y}^\top \mathbf{G}(\mathbf{W}, \mathbf{Z}), c_N(\mathbf{Z}) \mathbf{Y}^\top \mathbf{Z} \mathbf{Y} / 2, -\mathbf{Y}^\top \mathbf{Y} / 2)^\top. \quad (11.49)$$

The first two elements of $t(\mathbf{Y})$, $\mathbf{G}(\mathbf{W}, \mathbf{Z})^\top \mathbf{Y}$, are linear in \mathbf{Y} . Therefore, $\mathbf{G}(\mathbf{W}, \mathbf{Z})^\top \mathbf{Y}$ is elementwise Lipschitz as a function of \mathbf{Y} , with Lipschitz coefficients bounded above by

$$\max_{1 \leq k \leq 2} \|\mathbf{G}_{\cdot, k}(\mathbf{W}, \mathbf{Z})\|_2 \leq \sqrt{N},$$

because all elements of $\mathbf{G}(\mathbf{W}, \mathbf{Z})$ are between 0 and 1. We invoke Lemma 4 to show that $\mathbf{G}_{\cdot, k}(\mathbf{W}, \mathbf{Z})^\top \mathbf{Y}$ concentrates around its expectation. Recall from (10.1) that the joint density of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z})$ is

$$f_{\mathbf{y}|\mathbf{w}, \mathbf{z}}(\mathbf{y} \mid \mathbf{w}, \mathbf{z}) = \psi(\boldsymbol{\theta}) \exp \left(-\frac{1}{2\kappa^{\star 2}} \mathbf{y}^\top \mathbf{y} + \frac{c_N(\mathbf{z}) \delta^*}{2\kappa^{\star 2}} \mathbf{y}^\top \mathbf{z} \mathbf{y} + \frac{\boldsymbol{\vartheta}^{\star \top}}{\kappa^{\star 2}} \mathbf{G}(\mathbf{w}, \mathbf{z})^\top \mathbf{y} \right).$$

If we write $f_{\mathbf{y}|\mathbf{w}, \mathbf{z}}(\mathbf{y} \mid \mathbf{w}, \mathbf{z}) = \exp(-U(\mathbf{y}, \mathbf{w}, \mathbf{z}))$, then

$$\nabla_{\mathbf{y}}^2 U(\mathbf{y}, \mathbf{w}, \mathbf{z}) = \frac{1}{\kappa^{\star 2}} (\mathbf{I}_N - \delta^* c_N(\mathbf{z}) \mathbf{z}).$$

Since $c_N(\mathbf{Z}) \|\mathbf{Z}\|_2 \leq 1$ and $|\delta^*| < 1$, the smallest eigenvalue of $\nabla_{\mathbf{y}}^2 U(\mathbf{y}, \mathbf{w}, \mathbf{z})$ is at least $(1 - |\delta^*|)/\kappa^{\star 2}$. So, for any $t > 0$,

$$\begin{aligned} & \mathbb{P}_{\mathbf{y}|\mathbf{w}, \mathbf{z}}(|\mathbf{G}_{\cdot, k}(\mathbf{W}, \mathbf{Z})^\top \mathbf{Y} - \mathbb{E}_{\mathbb{P}_{\mathbf{y}|\mathbf{w}, \mathbf{z}}}[\mathbf{G}_{\cdot, k}(\mathbf{W}, \mathbf{Z})^\top \mathbf{Y}]| > t) \\ & \leq 2 \exp \left(-\frac{v_1 (1 - |\delta^*|) t^2}{N \kappa^{\star 2}} \right), \end{aligned} \quad (11.50)$$

where $k = 1, 2$, and $v_1 > 0$ is a constant, independent of N .

The third and fourth elements of $t(\mathbf{Y})$, $c_N(\mathbf{Z}) \mathbf{Y}^\top \mathbf{Z} \mathbf{Y} / 2$ and $-\mathbf{Y}^\top \mathbf{Y} / 2$, are quadratic in \mathbf{Y} , and are not Lipschitz-continuous. Let $\boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})$ and $\boldsymbol{\Sigma}^*(\mathbf{W}, \mathbf{Z})$ be the conditional mean and the conditional covariance matrix of $\mathbf{Y} \mid (\mathbf{W}, \mathbf{Z})$ respectively. We can express $c_N(\mathbf{Z}) \mathbf{Y}^\top \mathbf{Z} \mathbf{Y} / 2$ and $-\mathbf{Y}^\top \mathbf{Y} / 2$ as a function of independent normal variables by considering the following transformation:

$$\mathbf{R} := \mathbf{R}(\mathbf{W}) = \boldsymbol{\Sigma}^*(\mathbf{W}, \mathbf{Z})^{-\frac{1}{2}} (\mathbf{Y}(\mathbf{W}) - \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})).$$

Conditional on (\mathbf{W}, \mathbf{Z}) , \mathbf{R} follows multivariate Gaussian with mean $\mathbf{0}$ and covariance matrix \mathbf{I}_N . Also note that, $\boldsymbol{\Omega}^*(\mathbf{W}, \mathbf{Z})^{\frac{1}{2}} = \boldsymbol{\Sigma}^*(\mathbf{W}, \mathbf{Z})^{-\frac{1}{2}}$. We derive

$$\begin{aligned} \frac{c_N(\mathbf{Z})}{2} \mathbf{Y}^\top \mathbf{Z} \mathbf{Y} &= \frac{c_N(\mathbf{Z})}{2} (\mathbf{Y} - \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z}))^\top \mathbf{Z} (\mathbf{Y} - \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})) \\ &\quad + c_N(\mathbf{Z}) \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \mathbf{Z} \mathbf{Y} - \frac{c_N(\mathbf{Z})}{2} \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \mathbf{Z} \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z}) \\ &= \frac{c_N(\mathbf{Z})}{2} \mathbf{R}^\top \boldsymbol{\Omega}^*(\mathbf{W}, \mathbf{Z})^{-\frac{1}{2}} \mathbf{Z} \boldsymbol{\Omega}^*(\mathbf{W}, \mathbf{Z})^{-\frac{1}{2}} \mathbf{R} \\ &\quad + c_N(\mathbf{Z}) \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \mathbf{Y} - \frac{c_N(\mathbf{Z})}{2} \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \mathbf{Z} \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z}). \\ -\frac{1}{2} \mathbf{Y}^\top \mathbf{Y} &= -\frac{1}{2} (\mathbf{Y} - \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z}))^\top (\mathbf{Y} - \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})) - \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \mathbf{Y} \\ &\quad + \frac{1}{2} \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z}) \\ &= -\frac{1}{2} \mathbf{R}^\top \boldsymbol{\Omega}^*(\mathbf{W}, \mathbf{Z})^{-1} \mathbf{R} - \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \mathbf{Y} \\ &\quad + \frac{1}{2} \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z}). \end{aligned}$$

The first term in both the expansions above are quadratic in \mathbf{R} . Note that $\mathbf{R} := (R_1, \dots, R_N)$

is a random vector of independent components R_i such that $\mathbb{E}_{\mathbb{P}_{\mathbf{y}|\mathbf{W}, \mathbf{Z}}}[R_i \mid \mathbf{W}, \mathbf{Z}] = 0$ and its

Subgaussian norm is 1. Hence, we can apply the Hanson-Wright inequality (Rudelson and Vershynin,

2013, Theorem 1.1) to obtain that, for any matrix \mathbf{A} and any scalar $t > 0$,

$$\begin{aligned} & \mathbb{P}_{y|W,Z} \left(|\mathbf{R}^\top \mathbf{A} \mathbf{R} - \mathbb{E}_{\mathbb{P}_{y|W,Z}}[\mathbf{R}^\top \mathbf{A} \mathbf{R}]| > \frac{t}{2} \right) \\ & \leq 2 \exp \left(-v_2 \min \left\{ \frac{t^2}{4 \|\mathbf{A}\|_F^2}, \frac{t}{2 \|\mathbf{A}\|_2} \right\} \right), \end{aligned} \quad (11.51)$$

where $v_2 > 0$ is a constant, independent of N .

To derive concentration bound for $c_N(\mathbf{Z}) \mathbf{Y}^\top \mathbf{Z} \mathbf{Y} / 2$, choose

$$\mathbf{A} := \frac{c_N(\mathbf{Z})}{2} \boldsymbol{\Omega}^*(\mathbf{W}, \mathbf{Z})^{-\frac{1}{2}} \mathbf{Z} \boldsymbol{\Omega}^*(\mathbf{W}, \mathbf{Z})^{-\frac{1}{2}},$$

which results in

$$\begin{aligned} \|\mathbf{A}\|_F & \leq \sqrt{N} \|\mathbf{A}\|_2 = \frac{c_N(\mathbf{Z}) \sqrt{N}}{2} \|\boldsymbol{\Omega}^*(\mathbf{W}, \mathbf{Z})^{-\frac{1}{2}} \mathbf{Z} \boldsymbol{\Omega}^*(\mathbf{W}, \mathbf{Z})^{-\frac{1}{2}}\|_2 \\ & \leq \frac{c_N(\mathbf{Z}) \sqrt{N}}{2} \|\mathbf{Z}\|_2 \|\boldsymbol{\Omega}^*(\mathbf{W}, \mathbf{Z})^{-\frac{1}{2}}\|_2^2 \leq \frac{\sqrt{N} \kappa^{*2}}{2(1 - |\delta^*|)}, \end{aligned}$$

using the fact that $\|\boldsymbol{\Omega}^*(\mathbf{W}, \mathbf{Z})^{-1}\|_2 \leq \kappa^{*2} / (1 - |\delta^*|)$ because $c_N(\mathbf{Z}) \|\mathbf{Z}\|_2 \leq 1$ and $|\delta^*| < 1$.

Then, (11.51) implies that

$$\begin{aligned} & \mathbb{P}_{y|W,Z} \left(|\mathbf{R}^\top \mathbf{A} \mathbf{R} - \mathbb{E}_{\mathbb{P}_{y|W,Z}}[\mathbf{R}^\top \mathbf{A} \mathbf{R}]| > \frac{t}{2} \right) \\ & \leq 2 \exp \left(-v_2 \min \left\{ \frac{(1 - |\delta^*|)^2 t^2}{N \kappa^{*4}}, \frac{(1 - |\delta^*|) t}{\kappa^{*2}} \right\} \right). \end{aligned} \quad (11.52)$$

The second term in the expansion of $c_N(\mathbf{Z}) \mathbf{Y}^\top \mathbf{Z} \mathbf{Y} / 2$ is linear in \mathbf{Y} , and its Lipschitz

coefficient is bounded above by

$$\begin{aligned}
& c_N(\mathbf{Z}) \left\| \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \mathbf{Z} \right\|_2 \\
&= c_N(\mathbf{Z}) \left\| \boldsymbol{\vartheta}^{*\top} \mathbf{G}(\mathbf{W}, \mathbf{Z})^\top (\mathbf{I}_N - \delta^* c_N(\mathbf{Z}) \mathbf{Z})^{-1} \mathbf{Z} \right\|_2 \\
&\leq c_N(\mathbf{Z}) \left\| \boldsymbol{\vartheta}^* \right\|_2 \left\| \mathbf{G}(\mathbf{W}, \mathbf{Z}) \right\|_2 \left\| (\mathbf{I}_N - c_N(\mathbf{Z}) \delta^* \mathbf{Z})^{-1} \right\|_2 \left\| \mathbf{Z} \right\|_2 \\
&\leq \frac{\sqrt{2N} \left\| \boldsymbol{\vartheta}^* \right\|_2}{1 - |\delta^*|},
\end{aligned} \tag{11.53}$$

since $\left\| (\mathbf{I}_N - c_N(\mathbf{Z}) \delta^* \mathbf{Z})^{-1} \right\|_2 \leq 1/(1 - |\delta^*|)$ and $\left\| \mathbf{G}(\mathbf{W}, \mathbf{Z}) \right\|_2 \leq \sqrt{2N}$. We apply Lemma 4 to obtain

$$\begin{aligned}
& \mathbb{P}_{\mathbf{y}|\mathbf{w},\mathbf{z}} \left(c_N(\mathbf{Z}) \left| \boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \mathbf{Z} \mathbf{Y} - \mathbb{E}_{\mathbb{P}_{\mathbf{y}|\mathbf{w},\mathbf{z}}} [\boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \mathbf{Z} \mathbf{Y}] \right| > \frac{t}{2} \right) \\
&\leq 2 \exp \left(-\frac{v_1 (1 - |\delta^*|)^3 t^2}{8N \left\| \boldsymbol{\vartheta}^* \right\|_2^2 \kappa^{*2}} \right).
\end{aligned} \tag{11.54}$$

Finally, the third term in the expansion of $c_N(\mathbf{Z}) \mathbf{Y}^\top \mathbf{Z} \mathbf{Y} / 2$ is constant with respect to \mathbf{Y} . Therefore, collecting terms

$$\begin{aligned}
& \mathbb{P}_{\mathbf{y}|\mathbf{w},\mathbf{z}} \left(\frac{c_N(\mathbf{Z})}{2} \left| \mathbf{Y}^\top \mathbf{Z} \mathbf{Y} - \mathbb{E}_{\mathbb{P}_{\mathbf{y}|\mathbf{w},\mathbf{z}}} [\mathbf{Y}^\top \mathbf{Z} \mathbf{Y}] \right| > t \right) \\
&\leq 2 \exp \left(-v_2 \min \left\{ \frac{(1 - |\delta^*|)^2 t^2}{N \kappa^{*4}}, \frac{(1 - |\delta^*|) t}{\kappa^{*2}} \right\} \right) \\
&+ 2 \exp \left(-\frac{v_1 (1 - |\delta^*|)^3 t^2}{8N \left\| \boldsymbol{\vartheta}^* \right\|_2^2 \kappa^{*2}} \right).
\end{aligned} \tag{11.55}$$

We take a similar approach to derive concentration bound for $-\frac{1}{2} \mathbf{Y}^\top \mathbf{Y}$. Choosing

$$\mathbf{A} := \frac{1}{2} \boldsymbol{\Omega}^*(\mathbf{W}, \mathbf{Z})^{-1},$$

results in

$$\|\mathbf{A}\|_F \leq \sqrt{N} \|\mathbf{A}\|_2 = \frac{\sqrt{N}}{2} \|\boldsymbol{\Omega}^*(\mathbf{W}, \mathbf{Z})^{-1}\|_2 \leq \frac{\sqrt{N} \kappa^{\star 2}}{2(1 - |\delta|^\star)}.$$

Plugging this into (11.51), we obtain

$$\begin{aligned} & \mathbb{P}_{\mathbf{y}|\mathbf{W}, \mathbf{Z}} \left(|\mathbf{R}^\top \mathbf{A} \mathbf{R} - \mathbb{E}_{\mathbb{P}_{\mathbf{y}|\mathbf{W}, \mathbf{Z}}}[\mathbf{R}^\top \mathbf{A} \mathbf{R}]| > \frac{t}{2} \right) \\ & \leq 2 \exp \left(-v_2 \min \left\{ \frac{(1 - |\delta|^\star)^2 t^2}{N \kappa^{\star 4}}, \frac{(1 - |\delta|^\star) t}{\kappa^{\star 2}} \right\} \right). \end{aligned} \quad (11.56)$$

The second term in the expansion of $-\mathbf{Y}^\top \mathbf{Y}/2$ is likewise linear in \mathbf{Y} , and its Lipschitz coefficient is bounded above by

$$\|\boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top\|_2 \leq \frac{\sqrt{2N} \|\boldsymbol{\vartheta}^*\|_2}{\kappa^{\star 2} (1 - |\delta|^\star)},$$

following the steps as in (11.53). Again, applying Lemma 4, we obtain

$$\begin{aligned} & \mathbb{P}_{\mathbf{y}|\mathbf{W}, \mathbf{Z}} \left(|\boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \mathbf{Y} - \mathbb{E}_{\mathbb{P}_{\mathbf{y}|\mathbf{W}, \mathbf{Z}}}[\boldsymbol{\mu}^*(\mathbf{W}, \mathbf{Z})^\top \mathbf{Y}]| > \frac{t}{2} \right) \\ & \leq 2 \exp \left(-\frac{v_1 (1 - |\delta|^\star)^3 t^2}{8 N \|\boldsymbol{\vartheta}^*\|_2^2 \kappa^{\star 2}} \right). \end{aligned} \quad (11.57)$$

The third term in the expansion of $-\mathbf{Y}^\top \mathbf{Y}/2$ is constant with respect to \mathbf{Y} . Therefore, collecting terms

$$\begin{aligned} & \mathbb{P}_{\mathbf{y}|\mathbf{W}, \mathbf{Z}} \left(\frac{1}{2} |\mathbf{Y}^\top \mathbf{Y} - \mathbb{E}_{\mathbb{P}_{\mathbf{y}|\mathbf{W}, \mathbf{Z}}}[\mathbf{Y}^\top \mathbf{Y}]| > t \right) \\ & \leq 2 \exp \left(-v_2 \min \left\{ \frac{(1 - |\delta|^\star)^2 t^2}{N \kappa^{\star 4}}, \frac{(1 - |\delta|^\star) t}{\kappa^{\star 2}} \right\} \right) \\ & + 2 \exp \left(-\frac{v_1 (1 - |\delta|^\star)^3 t^2}{8 N \|\boldsymbol{\vartheta}^*\|_2^2 \kappa^{\star 2}} \right). \end{aligned} \quad (11.58)$$

From (11.50), (11.55) and (11.58), we conclude that

$$\begin{aligned} & \mathbb{P}_{y|W,Z} \left(\| t(\mathbf{Y}) - \mathbb{E}_{\mathbb{P}_{y|W,Z}}[t(\mathbf{Y})] \|_2 \geq t \right) \\ & \leq 8 \exp \left(- \min \left\{ \frac{B_1(\boldsymbol{\theta}^*, \kappa^{*2}) t^2}{N}, B_2(\boldsymbol{\theta}^*, \kappa^{*2}) t \right\} \right), \end{aligned}$$

where $B_1(\boldsymbol{\theta}^*, \kappa^{*2})$, $B_2(\boldsymbol{\theta}^*, \kappa^{*2})$ are constants that depend on $(\boldsymbol{\theta}^*, \kappa^{*2})$.