

Thermodynamic phase transition and Joule-Thomson expansion of a quantum corrected black hole in anti-de Sitter spacetime

Rui-Bo Wang ^{1,*} Lei You ^{1,†} Shi-Jie Ma ^{1,‡} Jian-Bo Deng ^{1,§} and Xian-Ru Hu ^{1,¶}

¹*School of Physical Science and Technology, Lanzhou University, Lanzhou, Gansu 730000, China*
(Dated: April 10, 2025)

The thermodynamics in the extended phase space of a quantum corrected black hole (BH) proposed recently is presented in this work. Our study shows that the phase transition behavior of the BH is analogous to that of conventional Schwarzschild BH in anti-de Sitter (AdS) space; however, a critical temperature exists such that when the BH temperature exceeds this critical value, the small BH phase and the large BH phase become separated, and no phase transition occurs. Due to the introduction of the quantum parameter ξ , the BH equation of state splits into two branches. One branch reduces to the Schwarzschild-AdS case as $\xi \rightarrow 0$, with its phase transition pressure lower than the critical pressure; another branch's phase transition pressure is greater than the critical pressure. We further investigate the temperature curve and the isobaric heat capacity of the BH. The results show that the $T - r_+$ phase transition and heat capacity behavior are very similar to those of the Schwarzschild-AdS BH, with the small BH phase exhibiting a negative heat capacity (unstable phase) and the large BH phase exhibiting a positive heat capacity (stable phase). Furthermore, owing to the corrected first law, the Gibbs free energy curve becomes smooth near the phase transition point. Moreover, we examine the Joule-Thomson expansion of the BH. The constant mass expansion process of the BH is divided into two stages: in the earlier stage, the BH pressure increases until it reaches a maximum; in the later stage, the pressure gradually decreases. Consequently, the inversion curve is also divided into two branches: one corresponding to the inversion point that may be experienced during the earlier stage and the other corresponding to the inversion point during the later stage. In addition, each expansion stage has a minimum inversion mass, below which any BH (in each respective stage) has no inversion point.

Keywords: Quantum Corrected Black Hole, Black Hole Thermodynamics, Joule-Thomson Expansion.

arXiv:2504.06907v1 [gr-qc] 9 Apr 2025

* Rui-Bo Wang: wangrb2021@lzu.edu.cn

† Lei You: youl21@lzu.edu.cn

‡ Shi-Jie Ma: 220220939731@lzu.edu.cn

§ Jian-Bo Deng (corresponding author): dengjb@lzu.edu.cn

¶ Xian-Ru Hu: huxianru@lzu.edu.cn

I. INTRODUCTION

Due to the inconsistency between general relativity (GR) and quantum theory, as well as issues such as singularities [1–5], quantum gravity has long been a subject of extensive interest. One approach to achieve spacetime quantization is non-commutative geometry [6–13], which is characterized by the commutation relation among spacetime coordinate operators $[x^\mu, x^\nu] = i\theta^{\mu\nu}$, where $\theta^{\mu\nu}$ is a anti-symmetric constant matrix. A. Smilagic’s research shows that noncommutative effects eliminate the point-like mass distribution [2, 3], a phenomenon that may circumvent the BH singularity problem. Inspired by this, P. Nicolini and K. Nozari respectively proposed that the point mass distribution model (represented by the Dirac delta function) can be replaced by either a Gaussian distribution or a Lorentzian distribution [14, 15]. In addition, effective field theory (EFT) and loop quantum gravity (LQG) are also candidate theories for quantum gravity. In Ref. [16] the authors analyze low-energy one-loop quantum corrections to the Schwarzschild geometry. In their approach, general relativity is treated as a quantum field theory (QFT) within the framework of EFT, implying that at low energies the degrees of freedom organize themselves as quantum fields governed by a local Lagrangian [17]. Similarly, LQG has received significant attention [18–25]. In LQG, the minimal fundamental units of spacetime are suggested as a series of discrete, elementary loops. One interesting result is that by studying the gravitational collapse of spherically symmetric dust matter in LQG, researchers have constructed a quantum corrected Schwarzschild BH and found that it can be interpreted as the formation of a white hole [26–28].

The BH discussed in this paper was proposed in a recent study that aimed to address the longstanding issue of general covariance in quantum gravity models. It is suggested that the classical GR could be modified by constructing a Hamiltonian constraints formulation [29–31]. In this approach, an issue arose: How to keep the general diffeomorphism covariance [32]. Recently, this problem is effectively addressed in the framework of spherically symmetric vacuum gravity [33]. In Zhang’s team’s work [33], general covariance is rigorously defined as a set of constraint equations that ensure the general covariance of the spacetime theory. Based on this formulation, they derived the equations of motion for the effective Hamiltonian constraint, which depends on a quantum parameter ξ . Furthermore, they provided two candidate forms for the effective Hamiltonian constraint, each corresponding to a quantum corrected Schwarzschild spacetime. In this work, we investigate the thermodynamics of one such static, spherically symmetric quantum corrected BH.

The thermodynamics of AdS BHs in the extended phase space is a novel and highly intriguing subject. Within this framework, the cosmological constant term in Einstein equation is interpreted as the energy-momentum tensor of a special “vacuum static ideal fluid”, and it is assumed that the BH achieves phase equilibrium with the vacuum fluid, thereby acquiring pressure [34]. To ensure that the BH exhibits a positive pressure, the cosmological constant is required to be negative, implying that the BH resides in an AdS spacetime. Consequently, the AdS BH is effectively regarded as a unique thermodynamic system, which has attracted extensive discussion in the academic community. In 1983, Hawking first investigated the thermodynamics of Schwarzschild-AdS BHs, revealing the existence of phase transitions within these systems [35]. This pioneering work is widely regarded as the inception of the field of AdS BH thermodynamics. Numerous studies have explored the thermodynamics of AdS BHs [36–49]. The thermodynamic properties of BHs are influenced by several additional parameters, such as electric charge [50–53], nonlinear magnetic charge [54–56], dark matter [57, 58], nonlinear electrodynamics models [59–65], noncommutative geometry [66–69], lower or higher-dimensional gravity theories [70–73], and various other modified gravity models [74]. Many BHs exhibit thermodynamic properties analogous to those of Van der Waals system, implying a profound connection between gravitational theory, quantum physics and thermodynamics, and thereby revealing a richer array of physical properties for these enigmatic objects.

The structure of this paper is organized as follows. In Sect. II, we introduce the quantum corrected BH that is the focus of this paper. Under the assumption that this quantum gravity decouples from the cosmological constant, we derive the BH metric in AdS spacetime and compute its thermodynamic functions as well as the corrected first law. Sect. III will provide a detailed investigation into the thermodynamic phase transitions and critical phenomena of the BH. The isobaric heat capacity and the Gibbs free energy of the BH are researched in Sect. IV and Sect. V respectively. In Sect. VI, we conduct a detailed investigation of the constant mass expansion of the BH and present its inversion curve. Finally, the summary and outlook of this work will be presented in Sect. VII. Our study will demonstrate the influence of the quantum correction parameter on the thermodynamics of conventional Schwarzschild–AdS BH, thereby enriching the fruits of research on quantum BH thermodynamics and providing a valuable reference for subsequent investigations.

For computational convenience, natural units $G = \hbar = c = k_B = 1$ are adopted throughout this work.

II. THERMODYNAMIC FUNCTIONS AND THE FIRST LAW

The BH considered in this paper is a quantum corrected BH, described by a static spherically symmetric spacetime [33]

$$ds^2 = -f(r) dt^2 + \frac{1}{f(r)} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2, \quad (2.1)$$

with $f(r)$ being

$$f(r) = 1 - \frac{2M}{r} \left(1 + \frac{\xi^2}{r^2} \left(1 - \frac{2M}{r} \right) \right), \quad (2.2)$$

where ξ is a quantum parameter, which is considered to be on the order of the Planck length ℓ_p . This BH corresponds to an equivalent energy distribution

$$T_{t(\text{BH})}^t = -\frac{\xi^2 (r - 2M)(r - 6M)}{8\pi r^6}. \quad (2.3)$$

Assuming that this quantum gravity decouples from the cosmological constant Λ , the Einstein equation with Λ is

$$R_\mu^\nu - \frac{1}{2} \delta_\mu^\nu R = 8\pi \left(T_{\mu(\text{BH})}^\nu + T_{\mu(\Lambda)}^\nu \right), \quad (2.4)$$

where $T_{\mu(\Lambda)}^\nu = -\frac{\Lambda}{8\pi} \delta_\mu^\nu$ represents an equivalent energy-momentum tensor corresponding to the contribution of the cosmological constant term. Thus the metric of this quantum corrected BH in AdS spacetime is obtained as

$$f(r) = 1 - \frac{2M}{r} \left(1 + \frac{\xi^2}{r^2} \left(1 - \frac{2M}{r} \right) \right) - \frac{\Lambda r^2}{3}. \quad (2.5)$$

The BH solution satisfies $f(r_+) = 0$, where r_+ is the event horizon radius. The mass of BH could be solved

$$M = \frac{2r_+ \xi^2 + r_+^3 (1 \pm \sqrt{\alpha})}{4\xi^2}, \quad (2.6)$$

where

$$\alpha = 1 + \frac{4}{3} \Lambda \xi^2. \quad (2.7)$$

Here, we discard the positive-sign branch because this branch does not reduce to the Schwarzschild-AdS case as $\xi \rightarrow 0$, but instead diverges to infinity. It is required that $\alpha \geq 0$, which gives

$$\Lambda \geq \Lambda_{min} = -\frac{3}{4\xi^2}. \quad (2.8)$$

On the other hand, the existence of BH requires $M > 0, r_+ > 0, \Lambda < 0$ (AdS spacetime), which reduces this parameter constraint

$$\frac{r_+}{2} < M < \frac{1}{2} \left(r_+ + \frac{r_+^3}{\xi^2} \right), \quad (2.9)$$

or

$$\frac{-3^{\frac{2}{3}} \xi^{\frac{4}{3}} + \xi^{\frac{2}{3}} 3^{\frac{1}{3}} \beta^{\frac{2}{3}}}{3\beta^{\frac{1}{3}}} < r_+ < 2M. \quad (2.10)$$

where $\beta = 9M + \sqrt{81M^2 + 3\xi^2}$.

The BH's temperature is defined by its surface gravity

$$T = \frac{\partial_r f}{4\pi} \Big|_{r_+} = \frac{\sqrt{\alpha}}{4\pi r_+} + \frac{3r_+ (\sqrt{\alpha} - \alpha)}{8\pi \xi^2} \quad (2.11)$$




FIG. 1. Condition of BH's existence in (r_+, M) coordinate. The orange region signifies the parameter range in which BH solutions exist, while the remaining areas indicate the absence of BH solutions.

In the extended phase space, the energy-momentum tensor $T_{\mu(\Lambda)}^\nu$ contributed by the cosmological constant is regarded as an equivalent "vacuum static ideal fluid". It is assumed that the BH reaches a phase equilibrium with this fluid, acquiring its pressure. The pressure of the BH is

$$P = -\frac{\Lambda}{8\pi}. \quad (2.12)$$

Eq. 2.8 gives the maximum pressure of BH

$$P_{max} = -\frac{\Lambda_{min}}{8\pi} = \frac{3}{32\pi\xi^2}. \quad (2.13)$$

According to the studies in references, when the energy-momentum tensor outside the BH event horizon explicitly includes the BH's mass M , the first law of BH thermodynamics takes a corrected form [55, 75–77]

$$WdM = TdS + VdP, \quad (2.14)$$

where correction function W is

$$W = 1 + \int_{r_+}^{\infty} 4\pi r^2 \frac{\partial T_t^t}{\partial M} dr = \sqrt{\alpha}. \quad (2.15)$$

The entropy of the BH is the Bekenstein-Hawking function [78, 79]

$$S = \int \frac{W}{T} dM = \int \frac{W}{T} \frac{\partial M}{\partial r_+} dr_+ = \pi r_+^2. \quad (2.16)$$

The BH's thermodynamic volume is

$$V = W \left(\frac{\partial M}{\partial P} \right)_S = \frac{4\pi r_+^3}{3}. \quad (2.17)$$

III. THERMODYNAMIC PHASE TRANSITION

The equation of state of BH is solved from Eq. 2.11

$$P_{\pm} = \frac{T}{4r_+} - \frac{1}{16\pi r_+^2} + \frac{3}{64\pi\xi^2} - \frac{\xi^2}{48\pi r_+^4} \pm \left(\frac{\sqrt{9r_+^4 + 12r_+^2(1 - 8\pi r_+ T)\xi^2 + 4\xi^4}}{96\pi r_+^4} + \frac{\sqrt{9r_+^4 + 12r_+^2(1 - 8\pi r_+ T)\xi^2 + 4\xi^4}}{64\pi r_+^2 \xi^2} \right). \quad (3.1)$$

It is evident that the $P - r_+$ diagram of the BH consists of two branches, corresponding respectively to the positive-sign branch P_+ (Fig. 2) and negative-sign branch P_- in the above function (Fig. 3).





FIG. 2. BH's $P_+ - r_+$ diagram. Black point "critic" is the critical point. We set $\xi = 1$.

FIG. 3. BH's $P_- - r_+$ diagram. Black point "critic" is the critical point. We set $\xi = 1$.

In particular, there exists a critical temperature

$$T_c = \frac{1}{3\sqrt{2}\pi\xi}. \quad (3.2)$$

When $T > T_c$, the small BH phase ($r_+ < r_1(T)$) and the large BH phase ($r_+ > r_2(T)$) become distinctly separated, and no phase transition occurs. $r_1(T)$ and $r_2(T)$ are the smaller and larger positive real roots of equation with parameter T

$$9r_+^4 + 12r_+^2\xi^2 - 96\pi r_+^3 T\xi^2 + 4\xi^4 = 0, \quad (3.3)$$

respectively.

When $T = T_c$, the small BH-large BH phase transition occurs at the critical radius

$$r_{+c} = \sqrt{2}\xi. \quad (3.4)$$

At the critical point, BH reaches its critical pressure

$$P_c = \frac{5}{96\pi\xi^2} = \frac{5}{9}P_{max}. \quad (3.5)$$

This critical point gives a dimensionless constant

$$\rho_c = \frac{2P_c r_{+c}}{T_c} = \frac{5}{8}. \quad (3.6)$$

So this quantum corrected BH shows a larger critical ratio than Van der Waals system ($\frac{3}{8}$) [52]. Furthermore, please notice that the equation of state P_- shown in Fig. 3 will reduce to the Schwarzschild-AdS case in the limit $\xi \rightarrow 0$. In

Fig. 3, during an isothermal expansion process, the pressure of the small BH increases rather than decreases, signifying that it corresponds to an unstable phase. In contrast, the large BH exhibits decreasing pressure with increasing radius. However, the situation in the BH phase diagram presented in Fig. 2 is completely opposite, where the small BH corresponds to the stable phase, while the large BH corresponds to the unstable phase. This characteristic arises due to the quantum parameter ξ , and this feature does not occur in the ordinary Schwarzschild-AdS BH. The BH's $P_- - r_+$ diagram for different quantum parameters ξ is shown in Fig. 4.




FIG. 4. BH's $P_- - r_+$ diagram for different quantum parameters ξ . Black line corresponds to the Schwarzschild-AdS BH ($\xi = 0$). We set $T = T_c$ ($\xi = 1$).

Specifically, when $P_- = 0$, the horizon radius of the BH is independent of the size of the quantum parameter ξ

$$r_+|_{P=0} = \frac{1}{4\pi T}, \quad (3.7)$$

which is also the result of Schwarzschild-AdS BH.

The phase transition occurs at

$$\frac{\partial P}{\partial r_+} = 0. \quad (3.8)$$

The variation of the BH phase transition pressure P_{pt} with respect to the phase transition temperature T_{pt} is illustrated in Fig. 5. Thus, it is apparent that the critical temperature represents the maximum phase transition temperature. Moreover, if we consider only the BH pressure as described in Fig. 3, then the critical pressure will also correspond to the maximum phase transition pressure. This characteristic is analogous to that observed in the Van der Waals system.

The BH's $T - r_+$ diagram is shown in Fig. 6 and 7. It is evident that the BH exhibits a small BH-large BH phase transition similar to that of the Schwarzschild-AdS BH. The BH pressure initially increases with the horizon radius (corresponding to the small BH phase) and subsequently decreases (corresponding to the large BH phase). This phase transition, in contrast to that observed in Van der Waals fluids, occurs instantaneously at the maximum pressure point.

In particular, phase separation does not occur in BH's $T - r_+$ diagram. It can be verified that $T > 0$ always holds. When $P = P_{max}$, the BH is an extremal BH, consistently maintaining a zero temperature $T \equiv 0$. For any given pressure, the BH temperature always undergoes a small BH-large BH phase transition. The temperature of the small BH decreases as the horizon radius increases, whereas the large BH exhibits the opposite behavior.

The phase transition temperature T_{pt} satisfies

$$\frac{\partial T}{\partial r_+} = 0. \quad (3.9)$$




FIG. 5. Phase transition points in $T - P$ coordinate. Point (T_c, P_c) is the critical point. We set $\xi = 1$.




FIG. 6. BH's $T - r_+$ curve corresponding to the isobaric process. We set $\xi = 1$.




FIG. 7. BH's $T - r_+$ diagram for different quantum parameters ξ . Black line ($\xi = 0$) represents the Schwarzschild-AdS BH. We set $\Lambda = -0.01$.

Using this condition, one could prove that at $P = P_c$, T_{pt} has its maximum

$$T_{pt}^{max} = T_c. \quad (3.10)$$

This also, from the perspective of the $T - r_+$ diagram, demonstrates that BHs with temperatures exceeding the critical temperature cannot undergo phase transitions.

IV. HEAT CAPACITY

According to

$$S = \pi \left(\frac{3V}{4\pi} \right)^{\frac{2}{3}}, \quad (4.1)$$

the BH's isochoric heat capacity is zero

$$C_V = T \left(\frac{\partial S}{\partial T} \right)_V = 0. \quad (4.2)$$

The BH's isobaric heat capacity is

$$C_p = T \left(\frac{\partial S}{\partial T} \right)_P = \frac{2\pi r_+^2 \left(24P\pi r_+^4 - \xi^2 + r_+^2 \sqrt{9 - 96P\pi\xi^2} \right)}{-3r_+^2 + 24P\pi r_+^4 + \xi^2}. \quad (4.3)$$




FIG. 8. Isobaric heat capacity for different quantum parameters ξ . Black line represents the Schwarzschild-AdS BH. We set $P = 0.001$.




FIG. 9. Isobaric heat capacity for different pressure. We set $\xi = 1$.

The heat capacity as a function of the horizon radius is presented in Fig. 8 and Fig. 9. In comparison with Fig. 6, it is readily evident that the small BH-satisfying $\partial T/\partial r_+ < 0$ -exhibits a negative heat capacity, corresponding to an unstable phase, whereas the large BH-described by $\partial T/\partial r_+ > 0$ -displays a positive heat capacity, indicative of a stable phase. At the phase transition point, the heat capacity diverges to infinity. Moreover, when the horizon radius is sufficiently small, the heat capacity exhibits asymptotic behavior

$$C_p = -2\pi r_+^2 + \mathcal{O}(r_+)^4. \quad (4.4)$$

In the limit as $r_+ \rightarrow \infty$, the BH's isobaric heat capacity exhibits a quadratic divergence

$$C_p = 2\pi r_+^2 + \frac{3 + \sqrt{9 - 96P\pi\xi^2}}{12P} + \mathcal{O}(r_+)^{-2}. \quad (4.5)$$

V. GIBBS FREE ENERGY

The BH's Gibbs free energy is

$$G = M - TS = \frac{r_+}{2} - 4P\pi r_+^3 - \frac{1}{12}r_+ \sqrt{9 - 96P\pi\xi^2} + \frac{5r_+^3 \left(3 - \sqrt{9 - 96P\pi\xi^2} \right)}{24\xi^2}. \quad (5.1)$$




FIG. 10. Gibbs free energy of the BH for different pressures. The black points in the figure are phase transition points. We set $\xi = 1$.

One could calculate the differential form of G

$$dG = (W^{-1} - 1) T dS - S dT + W^{-1} V dP, \quad (5.2)$$

which gives

$$\left(\frac{\partial G}{\partial T} \right)_P = -S + (W^{-1} - 1) C_p. \quad (5.3)$$

Phase transition satisfies $C_p = \infty$, so in the BH's $G - T$ diagram, the points at which $\partial G / \partial T = \infty$ are exactly the phase transition points.

Specifically, when $P < P_c$, there exists a point at which

$$\frac{\partial G}{\partial T} = 0, \quad (5.4)$$

shown as orange line ($P = 0.5P_c$) and lightblue line ($P = 0.75P_c$) in Fig. 10.

Fig. 11 displays the Gibbs free energy of the BH at a fixed pressure for various values of the quantum parameter ξ . It is evident that when $\xi = 0$, the $G - T$ curve of the BH reverts to that of the Schwarzschild-AdS case. As ξ increases, the $G - T$ curve near the phase transition point becomes progressively smoother. Moreover, when ξ is further increased such that the pressure exceeds the critical pressure, the $G - T$ curve of the BH deviates significantly from the Schwarzschild-AdS scenario.

VI. JOULE-THOMSON EXPANSION

The Joule-Thomson process of BHs essentially refers to their constant mass expansion process $dM = 0$. In many AdS BHs, the Joule-Thomson process exhibits striking similarities: during the constant mass expansion, the pressure continuously decreases while the temperature changes accordingly. The inversion curve partitions the $T - P$ plane into distinct cooling and heating regions, and as the BH mass increases, the BH traverses increasingly broader ranges of pressure and temperature [53, 67, 69, 72, 73, 77, 80–87].




FIG. 11. Gibbs free energy of the BH for different quantum parameters ξ . We set $P = P_c$ ($\xi = 1.5$).

For the convenience of studying the constant mass expansion of the BH, we first rewrite the BH temperature and pressure in terms of the horizon radius and BH mass as parameters

$$P = -\frac{3(r_+ - 2M)(r_+^3 - 2M\xi^2 + r_+\xi^2)}{8\pi r_+^6}, \quad (6.1)$$

$$T = \frac{-3(r_+^3 - 4M\xi^2 + 2r_+\xi^2)^2 + (3r_+^3 + 2r_+\xi^2)|r_+^3 - 4M\xi^2 + 2r_+\xi^2|}{8\pi r_+^5 \xi^2}. \quad (6.2)$$

The Joule-Thomson coefficient is

$$\mu_{JT} = \left(\frac{\partial T}{\partial P}\right)_M = \frac{r_+(3r_+^6 - 8r_+^3(r_+ - 3M)\xi^2 + 4(8M - 3r_+)r_+\xi^4 - 3(r_+^3 + 20M\xi^2 - 6r_+\xi^2)|r_+^3 + 2(r_+ - 2M)\xi^2|)}{6(r_+ - 3M)\xi^2|(r_+^3 - 4M\xi^2 + 2r_+\xi^2)|}. \quad (6.3)$$

By setting $\mu_{JT} = 0$, one gets the inversion mass M_i

$$M_i^1 = \frac{9r_i^3 + 28r_i\xi^2 + \sqrt{171r_i^6 + 204r_i^4\xi^2 + 64r_i^2\xi^4}}{60\xi^2}, \quad r_i > \frac{\sqrt{2 + \sqrt{22}}}{3}\xi, \quad (6.4)$$

$$M_i^2 = \frac{3r_i^3 + 20r_i\xi^2 + \sqrt{9r_i^6 + 60r_i^4\xi^2 + 40r_i^2\xi^4}}{60\xi^2}, \quad r_i > \frac{\sqrt{2}}{2}\xi, \quad (6.5)$$

where r_i is the inversion radius. Thus, the inversion curve of the BH is divided into two branches; for a given inversion radius, M_i^1 induces the first branch (IC_1), while M_i^2 induces the second branch (IC_2).




FIG. 12. BH's inversion curves. We set $\xi = 1$.

IC_1 and IC_2 are exactly tangent at their minimum points T_i^{min} , and aside from the tangency point, $T_i^1 < T_i^2$ is identically satisfied. Moreover, the inversion pressure and inversion temperature at this tangency point exactly coincide with the BH's critical pressure and critical temperature. That is $T_i^{min} = T_c$.

In the limit as $r_+ \rightarrow \infty$, IC_1 and IC_2 both exhibit a maximum inversion pressure

$$P_i^{1max} = \frac{3 + 6\sqrt{19}}{400\pi\xi^2}, \quad P_i^{2max} = \frac{3}{50\pi\xi^2}. \quad (6.6)$$

This results in two constants that are independent of the quantum parameter ξ

$$\frac{P_i^{1max}}{P_c} = \frac{6}{125} (3 + 6\sqrt{19}), \quad \frac{P_i^{2max}}{P_c} = \frac{144}{125}. \quad (6.7)$$

Fig. 13 depicts the constant-mass expansion process of the BH. In the figure, the arrows indicate the direction of increasing horizon radius r_+ , representing the expansion process of the BH. It is evident that the constant-mass expansion of the BH can be divided into two distinct stages. In the earlier stage, the pressure increases until reaching the BH's maximum pressure P_{max} ; in the later stage, the pressure decreases. The black dots in the figure denote the inversion points experienced by the BH in each stage.

It should be noted that, because the BH pressure increases rather than decreases during the pre-expansion stage, using the sign of μ_{JT} to differentiate between cooling and heating processes is no longer appropriate. More accurately, the sign in $(\partial T/\partial V)_M$ (or equivalently, $(\partial T/\partial r_+)_M$) should be employed to distinguish between the cooling process and the heating process.

$$\left(\frac{\partial T}{\partial V}\right)_M = \begin{cases} < 0, & \text{cooling point,} \\ = 0, & \text{inversion point,} \\ > 0, & \text{heating point.} \end{cases} \quad (6.8)$$




FIG. 13. Joule-Thomson expansion of the BH. We set $\xi = 1$, $M = 5$.

Thus, as depicted in Fig. 13, during both the earlier stage and the later stage, the BH initially undergoes a heating phase before transitioning into a cooling phase at the inversion point.

Moreover, considering that for a given inversion radius r_i , $M_i^1 > M_i^2$ always holds, and $M_i^1(r_i)$, $M_i^2(r_i)$ are both increasing functions, for a constant mass process, the BH necessarily first passes through the inversion point on IC_1 , followed by that on IC_2 . In other words, IC_1 precisely corresponds to the inversion points experienced by BHs in the earlier stage, whereas IC_2 pertains to the inversion points attainable by BHs in the later stage. In addition, since a given point in the (P, T) plane may correspond to several distinct BH phases, the inversion curve does not, by itself, partition the (P, T) plane into separate cooling and heating regions.

The earlier and later stages of the BH are connected at the point of maximum pressure P_{max} , at which the horizon radius r_+ is

$$r_{+c}^{JT} = \frac{6^{\frac{2}{3}}\xi^{\frac{4}{3}} - 6^{\frac{1}{3}}\xi^{\frac{2}{3}}\gamma^{\frac{2}{3}}}{3\gamma^{\frac{1}{3}}}, \quad (6.9)$$

where $\gamma = \sqrt{81M^2 + 6\xi^2} - 9M$.

Fig. 14 and Fig. 15 respectively depict the $T - P$ diagrams for BHs with varying masses during the earlier stage and later stage of expansion. In the left panel, the inversion curve is IC_1 , while in the right panel, the inversion curve is IC_2 . It is evident that, for sufficiently small BH masses, no inversion point exists. By setting $P_i = 0$, the minimum inversion mass M_i^{min} for each case can be determined

$$M_{ie}^{min} = \frac{\sqrt{770 + 187\sqrt{22}}}{54}\xi, \quad M_{il}^{min} = \frac{1}{2\sqrt{2}}\xi. \quad (6.10)$$

The minimum inversion mass also could be derived from Eq. 6.4 and Eq. 6.5 directly.

$$M_{ie}^{min} = M_i^1 \left(\frac{\sqrt{2 + \sqrt{22}}}{3}\xi \right), \quad M_{il}^{min} = M_i^2 \left(\frac{\sqrt{2}}{2}\xi \right). \quad (6.11)$$

In both of the aforementioned figures, the first constant mass curve in each does not exhibit an inversion point, which is because

$$0.6 < M_{ie}^{min} \approx 0.75, \quad 0.3 < M_{il}^{min} \approx 0.35. \quad (6.12)$$




FIG. 14. Earlier stage of BH's constant mass expansion. The points circled by small squares in the figure denote the inversion points. We set $\xi = 1$.




FIG. 15. Later stage of BH's constant mass expansion. The points circled by small squares in the figure denote the inversion points. We set $\xi = 1$.




FIG. 16. Earlier stage of constant mass expansion for BH with M_i^{min} . The points circled by small squares in the figure denote the inversion points. We set $\xi = 1$.




FIG. 17. Later stage of constant mass expansion for BH with M_{il}^{min} . The points circled by small squares in the figure denote the inversion points. We set $\xi = 1$.

As seen in Fig. 16 and Fig. 17, the BHs with M_i^{min} reach their critical points at $P = 0$.

VII. CONCLUSION AND OUTLOOK

In this article, we investigated the thermodynamics in the extended phase space of a quantum corrected BH proposed recently. Our analysis demonstrates that the phase transition behavior of the BH is analogous to that of conventional Schwarzschild-AdS BH. In particular, there exists a critical temperature T_c such that for any BH with a temperature

exceeding T_c , the small BH phase and the large BH phase become separated, and no phase transition occurs. Due to the effect of the quantum parameter ξ , the BH equation of state $P(T, r_+)$ bifurcates into two branches with completely opposite stability properties. One branch converges to the Schwarzschild-AdS BH as $\xi \rightarrow 0$, and it is associated with a phase transition pressure lower than the critical pressure P_c , whereas the other branch exhibits a phase transition pressure greater than P_c , which is not observed in conventional Schwarzschild-AdS BHs.

The temperature and heat capacity C_p of the BH closely resemble those of the Schwarzschild-AdS BH. Specifically, the small BH phase exhibits a negative heat capacity, indicative of an unstable configuration, whereas the large BH phase shows a positive heat capacity, corresponding to a stable state. Moreover, the BH does not exhibit any region of negative temperature. Moreover, owing to the corrected first law, the Gibbs free energy curve $G - T$ in the vicinity of the BH phase transition point becomes smooth, thereby deviating from the behavior observed in conventional Schwarzschild-AdS BHs.

In addition, we conducted a detailed study of the Joule-Thomson expansion process of the BH. Our analysis reveals that the constant mass expansion process of the BH has two stages. In the earlier stage, the BH pressure increases with expansion until it reaches its maximum value; subsequently, during the later stage, the pressure gradually decreases. Consequently, the inversion curve of the BH splits into two branches, corresponding to the inversion points that may be experienced in the earlier and later stages, respectively. Moreover, each expansion stage is associated with a minimum inversion mass M_i^{min} , below which any BH undergoing a constant mass expansion process (in either stage) will not exhibit an inversion point.

It is our hope that these findings could offer novel insights into the interplay between quantum corrections and classical BH thermodynamics, and serve as a valuable reference for future research.

In this work, our study is based on the assumption that the quantum gravity theory responsible for generating this quantum corrected BH decouples from the cosmological constant. This assumption, also adopted in Ref. [77], is generally regarded as valid since the cosmological constant, being a constant, typically does not interact with other fields. Nevertheless, some modified gravity models suggest that the cosmological constant may be linked to other interactions. For instance, in Ref. [88], the authors constructed a gravastar in non-commutative BTZ geometry, finding the non-commutativity parameter plays a role of the cosmological constant for gravastar formation and stability. If this theory is correct, the cosmological constant might be suggested as a non-commutative effect. In the context of Lorentz symmetry breaking theories, researchers have found that the choice of the self-interaction potential in the Kalb-Ramond field differs between the cases of a zero and a nonzero cosmological constant [89]. It is foreseeable that the cosmological constant may no longer be merely a constant term, but instead could couple with a variety of modified gravity models. In addition, although the quantum parameter ξ is presumed to be extremely small, its precise value remains undetermined. One approach is to estimate ξ by comparing the observed BH shadow radius with the theoretical value predicted for a Schwarzschild BH [32, 90, 91]. However, this method can only constrain ξ to be on the order of the Schwarzschild radius, which is evidently much larger than the Planck length ℓ_p . Therefore, seeking alternative methods to constrain this quantum parameter more precisely represents an interesting and worthwhile topic for future research.

CONFLICTS OF INTEREST

The authors declare that there are no conflicts of interest regarding the publication of this paper.

ACKNOWLEDGMENTS

We want to thank School of Physical Science and Technology, Lanzhou University.

DATA AVAILABILITY

This work is a theoretical study. No data was used or generated during this research.

[1] Jan Ambjørn, Jerzy Jurkiewicz, and Renate Loll, “Dynamically triangulating Lorentzian quantum gravity,” *Nuclear Physics B* **610**, 347–382 (2001).

- [2] Joseph Polchinski, *String theory, volume 1: An introduction to the bosonic string*, Vol. 1 (Cambridge university press Cambridge, 2005).
- [3] Abhay Ashtekar, Beverly K Berger, James Isenberg, and Malcolm MacCallum, *General relativity and gravitation: a centennial perspective* (Cambridge University Press, 2015).
- [4] Sumati Surya, “The causal set approach to quantum gravity,” *Living Reviews in Relativity* **22**, 1–75 (2019).
- [5] Roger Penrose, “Gravitational collapse and space-time singularities,” *Physical Review Letters* **14**, 57 (1965).
- [6] Markus B Fröb, Albert Much, and Kyriakos Papadopoulos, “Noncommutative geometry from perturbative quantum gravity,” *Physical Review D* **107**, 064041 (2023).
- [7] Anais Smailagic and Euro Spallucci, “Feynman path integral on the non-commutative plane,” *Journal of Physics A: Mathematical and General* **36**, L467 (2003).
- [8] Anais Smailagic and Euro Spallucci, “UV divergence-free QFT on noncommutative plane,” *Journal of Physics A: Mathematical and General* **36**, L517 (2003).
- [9] Sergio Doplicher, Klaus Fredenhagen, and John E Roberts, “The quantum structure of spacetime at the Planck scale and quantum fields,” *Communications in Mathematical Physics* **172**, 187–220 (1995).
- [10] Nathan Seiberg and Edward Witten, “String theory and noncommutative geometry,” *Journal of High Energy Physics* **1999**, 032 (1999).
- [11] Fedele Lizzi and Richard J Szabo, “Noncommutative geometry and spacetime gauge symmetries of string theory,” *Chaos, Solitons & Fractals* **10**, 445–458 (1999).
- [12] Edward Witten, “Non-commutative geometry and string field theory,” *Nuclear Physics B* **268**, 253–294 (1986).
- [13] Narges Heidari, Hassan Hassanabadi, AA Araújo Filho, and John Kriz, “Exploring non-commutativity as a perturbation in the Schwarzschild black hole: quasinormal modes, scattering, and shadows,” *The European Physical Journal C* **84**, 566 (2024).
- [14] Kourosh Nozari and S Hamid Mehdipour, “Hawking radiation as quantum tunneling from a noncommutative Schwarzschild black hole,” *Classical and Quantum Gravity* **25**, 175015 (2008).
- [15] Piero Nicolini, “Noncommutative black holes, the final appeal to quantum gravity: a review,” *International Journal of Modern Physics A* **24**, 1229–1308 (2009).
- [16] Emmanuele Battista, “Quantum Schwarzschild geometry in effective field theory models of gravity,” *Physical Review D* **109**, 026004 (2024).
- [17] John F. Donoghue, “Quantum General Relativity and Effective Field Theory,” in *Handbook of Quantum Gravity*, edited by Cosimo Bambi, Leonardo Modesto, and Ilya Shapiro (Springer Nature Singapore, Singapore, 2023) pp. 1–24.
- [18] Leonardo Modesto, “Disappearance of the black hole singularity in loop quantum gravity,” *Physical Review D* **70**, 124009 (2004).
- [19] Alejandro Corichi and Parampreet Singh, “Loop quantization of the Schwarzschild interior revisited,” *Classical and Quantum Gravity* **33**, 055006 (2016).
- [20] Cong Zhang, Yongge Ma, Shupeng Song, and Xiangdong Zhang, “Loop quantum Schwarzschild interior and black hole remnant,” *Physical Review D* **102**, 041502 (2020).
- [21] Sabine Hossenfelder, Leonardo Modesto, and Isabeau Prémont-Schwarz, “Model for nonsingular black hole collapse and evaporation,” *Physical Review D* **81**, 044036 (2010).
- [22] Wen-Cong Gan, Nilton O Santos, Fu-Wen Shu, and Anzhong Wang, “Properties of the spherically symmetric polymer black holes,” *Physical Review D* **102**, 124030 (2020).
- [23] Viqar Husain, Jarod George Kelly, Robert Santacruz, and Edward Wilson-Ewing, “Quantum gravity of dust collapse: shock waves from black holes,” *Physical Review Letters* **128**, 121301 (2022).
- [24] Abhay Ashtekar, Javier Olmedo, and Parampreet Singh, “Quantum transfiguration of Kruskal black holes,” *Physical review letters* **121**, 241301 (2018).
- [25] Rodolfo Gambini and Jorge Pullin, “Black holes in loop quantum gravity: the complete space-time,” *Physical review letters* **101**, 161301 (2008).
- [26] Jerzy Lewandowski, Yongge Ma, Jinsong Yang, and Cong Zhang, “Quantum oppenheimer-snyder and swiss cheese models,” *Physical Review Letters* **130**, 101501 (2023).
- [27] Cong Zhang, Yongge Ma, and Jinsong Yang, “Black hole image encoding quantum gravity information,” *Physical Review D* **108**, 104004 (2023).
- [28] Jinsong Yang, Cong Zhang, and Yongge Ma, “Shadow and stability of quantum-corrected black holes,” *The European Physical Journal C* **83**, 619 (2023).
- [29] Thomas Thiemann, *Modern canonical quantum general relativity* (Cambridge University Press, 2008).
- [30] Abhay Ashtekar and Jerzy Lewandowski, “Background independent quantum gravity: a status report,” *Classical and Quantum Gravity* **21**, R53 (2004).
- [31] Andrea Addazi, J Alvarez-Muniz, R Alves Batista, G Amelino-Camelia, V Antonelli, M Arzano, M Asorey, J-L Atteia, S Bahamonde, F Bajardi, *et al.*, “Quantum gravity phenomenology at the dawn of the multi-messenger era—A review,” *Progress in Particle and Nuclear Physics* **125**, 103948 (2022).
- [32] Yu-Heng Shu and Jia-Hui Huang, “Circular orbits and thin accretion disk around a quantum corrected black hole,” *Physics Letters B* , 139411 (2025).
- [33] Cong Zhang, Jerzy Lewandowski, Yongge Ma, and Jinsong Yang, “Black holes and covariance in effective quantum gravity,” *arXiv preprint arXiv:2407.10168* (2024).
- [34] David Kastor, Sourya Ray, and Jennie Traschen, “Enthalpy and the mechanics of AdS black holes,” *Classical and Quantum Gravity* **26**, 195011 (2009).

- [35] Stephen W Hawking and Don N Page, “Thermodynamics of black holes in anti-de Sitter space,” *Communications in Mathematical Physics* **87**, 577–588 (1983).
- [36] Natacha Altamirano, David Kubizňák, Robert B Mann, and Zeinab Sherkatghanad, “Thermodynamics of rotating black holes and black rings: phase transitions and thermodynamic volume,” *Galaxies* **2**, 89–159 (2014).
- [37] MH Dehghani, “Thermodynamics of rotating charged black strings and (A) dS/CFT correspondence,” *Physical Review D* **66**, 044006 (2002).
- [38] Í DD Carvalho, G Alencar, and CR Muniz, “Thermodynamics of static and stationary black holes in Einstein–Gauss–Bonnet gravity with dark matter,” *Physics of the Dark Universe* **42**, 101290 (2023).
- [39] Seyed Hossein Hendi and Ali Dehghani, “Criticality and extended phase space thermodynamics of AdS black holes in higher curvature massive gravity,” *The European Physical Journal C* **79**, 227 (2019).
- [40] Di Wu and Shuang-Qing Wu, “Topological classes of thermodynamics of rotating AdS black holes,” *Physical Review D* **107**, 084002 (2023).
- [41] Robert M Wald, “The thermodynamics of black holes,” *Living reviews in relativity* **4**, 1–44 (2001).
- [42] Jacob D Bekenstein, “Statistical black-hole thermodynamics,” *Physical Review D* **12**, 3077 (1975).
- [43] Rong-Gen Cai, Ya-Peng Hu, Qi-Yuan Pan, and Yun-Long Zhang, “Thermodynamics of black holes in massive gravity,” *Physical Review D* **91**, 024032 (2015).
- [44] Jacob D Bekenstein, “Generalized second law of thermodynamics in black-hole physics,” *Physical Review D* **9**, 3292 (1974).
- [45] David Kubizňák, Robert B Mann, and Mae Teo, “Black hole chemistry: thermodynamics with Lambda,” *Classical and Quantum Gravity* **34**, 063001 (2017).
- [46] Robert M Wald, ““ernst theorem” and black hole thermodynamics,” *Physical Review D* **56**, 6467 (1997).
- [47] Brian P Dolan, “The cosmological constant and black-hole thermodynamic potentials,” *Classical and Quantum Gravity* **28**, 125020 (2011).
- [48] Yu Tian and Xiao-Ning Wu, “Thermodynamics of black holes from equipartition of energy and holography,” *Physical Review D—Particles, Fields, Gravitation, and Cosmology* **81**, 104013 (2010).
- [49] Peng Wang, Houwen Wu, Haitang Yang, and Feiyu Yao, “Extended phase space thermodynamics for black holes in a cavity,” *Journal of High Energy Physics* **2020**, 1–19 (2020).
- [50] Andrew Chamblin, Roberto Emparan, Clifford V Johnson, and Robert C Myers, “Charged AdS black holes and catastrophic holography,” *Physical Review D* **60**, 064018 (1999).
- [51] Andrew Chamblin, Roberto Emparan, Clifford V Johnson, and Robert C Myers, “Holography, thermodynamics, and fluctuations of charged AdS black holes,” *Physical Review D* **60**, 104026 (1999).
- [52] David Kubizňák and Robert B Mann, “P- V criticality of charged AdS black holes,” *Journal of High Energy Physics* **2012**, 1–25 (2012).
- [53] Özgür Ökcü and Ekrem Aydiner, “Joule–Thomson expansion of the charged AdS black holes,” *The European Physical Journal C* **77**, 1–7 (2017).
- [54] Cong Li, Chao Fang, Miao He, Jiacheng Ding, and Jianbo Deng, “Thermodynamics of the Bardeen black hole in anti-de Sitter space,” *Modern Physics Letters A* **34**, 1950336 (2019).
- [55] Dharm Veer Singh and Sanjay Siwach, “Thermodynamics and Pv criticality of Bardeen-AdS black hole in 4D Einstein-Gauss-Bonnet gravity,” *Physics Letters B* **808**, 135658 (2020).
- [56] Shan-Ping Wu and Shao-Wen Wei, “Thermodynamics and phase transition of Bardeen-AdS-class black holes,” arXiv preprint arXiv:2407.19702[gr-qc;hep-th] (2024. doi: <https://doi.org/10.48550/arXiv.2407.19702>).
- [57] Shi-Jie Ma, Rui-Bo Wang, Jian-Bo Deng, and Xian-Ru Hu, “Euler–Heisenberg black hole surrounded by perfect fluid dark matter,” *The European Physical Journal C* **84**, 595 (2024).
- [58] He-Xu Zhang, Yuan Chen, Tian-Chi Ma, Peng-Zhang He, and Jian-Bo Deng, “Bardeen black hole surrounded by perfect fluid dark matter,” *Chinese Physics C* **45**, 055103 (2021).
- [59] Hernan A Gonzalez, Mokhtar Hassaine, and Cristian Martinez, “Thermodynamics of charged black holes with a nonlinear electrodynamics source,” *Physical Review D—Particles, Fields, Gravitation, and Cosmology* **80**, 104008 (2009).
- [60] Heng Dai, Zixu Zhao, and Shuhang Zhang, “Thermodynamic phase transition of Euler-Heisenberg-AdS black hole on free energy landscape,” *Nuclear Physics B* **991**, 116219 (2023).
- [61] MB Tataryn and MM Stetsko, “Thermodynamics of a static electric-magnetic black hole in Einstein-Born-Infeld-AdS theory with different horizon geometries,” *General Relativity and Gravitation* **53**, 72 (2021).
- [62] Sharmanthie Fernando, “Thermodynamics of Born-Infeld–anti-de Sitter black holes in the grand canonical ensemble,” *Physical Review D—Particles, Fields, Gravitation, and Cosmology* **74**, 104032 (2006).
- [63] Xu Ye, Zi-Qing Chen, Ming-Da Li, and Shao-Wen Wei, “QED effects on phase transition and Ruppeiner geometry of Euler-Heisenberg-AdS black holes,” *Chinese Physics C* **46**, 115102 (2022).
- [64] Guan-Ru Li, Sen Guo, and En-Wei Liang, “High-order QED correction impacts on phase transition of the Euler-Heisenberg AdS black hole,” *Physical Review D* **106**, 064011 (2022).
- [65] SI Kruglov, “Magnetic black holes with generalized ModMax model of nonlinear electrodynamics,” *International Journal of Modern Physics D* **31**, 2250025 (2022).
- [66] Piero Nicolini, Anais Smailagic, and Euro Spallucci, “Noncommutative geometry inspired Schwarzschild black hole,” *Physics Letters B* **632**, 547–551 (2006).
- [67] Rui-Bo Wang, Shi-Jie Ma, Lei You, Jian-Bo Deng, and Xian-Ru Hu, “Thermodynamics of Schwarzschild-AdS black hole in non-commutative geometry,” *Chinese Physics C* (2025).

- [68] Baoyu Tan, “Thermodynamics of high order correction for Schwarzschild-AdS black hole in non-commutative geometry,” *Nuclear Physics B*, 116868 (2025).
- [69] Rui-Bo Wang, Lei You, Shi-Jie Ma, Jian-Bo Deng, and Xian-Ru Hu, “Thermodynamic properties and Joule-Thomson expansion of AdS black hole with Gaussian distribution in non-commutative geometry,” arXiv preprint arXiv:2503.12363 (2025).
- [70] Daniel Grumiller and Robert McNees, “Thermodynamics of black holes in two (and higher) dimensions,” *Journal of High Energy Physics* **2007**, 074 (2007).
- [71] Chikun Ding, Yu Shi, Jun Chen, Yuebing Zhou, and Changqing Liu, “High dimensional AdS-like black hole and phase transition in Einstein-bumblebee gravity,” *Chinese Physics C* **47**, 045102 (2023).
- [72] Jing Liang, Benrong Mu, and Peng Wang, “Joule-Thomson expansion of lower-dimensional black holes,” *Physical Review D* **104**, 124003 (2021).
- [73] Jie-Xiong Mo, Gu-Qiang Li, Shan-Quan Lan, and Xiao-Bao Xu, “Joule-Thomson expansion of d-dimensional charged AdS black holes,” *Physical Review D* **98**, 124032 (2018).
- [74] Rong-Gen Cai, Li-Ming Cao, and Nobuyoshi Ohta, “Thermodynamics of black holes in Hořava–Lifshitz gravity,” *Physics Letters B* **679**, 504–509 (2009).
- [75] Meng-Sen Ma and Ren Zhao, “Corrected form of the first law of thermodynamics for regular black holes,” *Classical and Quantum Gravity* **31**, 245014 (2014).
- [76] Manuel E Rodrigues, Marcos V de S. Silva, and Henrique A Vieira, “Bardeen-Kiselev black hole with a cosmological constant,” *Physical Review D* **105**, 084043 (2022).
- [77] Rui-Bo Wang, Shi-Jie Ma, Lei You, Yu-Cheng Tang, Yu-Hang Feng, Xian-Ru Hu, and Jian-Bo Deng, “Thermodynamics of AdS-Schwarzschild-like black hole in loop quantum gravity,” *The European Physical Journal C* **84**, 1161 (2024).
- [78] JD Bekenstein, “Black holes and the second law,” *Lettere al Nuovo Cimento* (1971-1985) **4**, 737–740 (1972).
- [79] Stephen W Hawking, “Particle creation by black holes,” *Communications in mathematical physics* **43**, 199–220 (1975).
- [80] Sen Guo, Jin Pu, Qing-Quan Jiang, and Xiao-Tao Zu, “Joule-Thomson expansion of the regular (Bardeen)-AdS black hole,” *Chinese Physics C* **44**, 035102 (2020).
- [81] Cong Li, Pengzhang He, Ping Li, and Jian-Bo Deng, “Joule–Thomson expansion of the Bardeen-AdS black holes,” *General Relativity and Gravitation* **52**, 1–10 (2020).
- [82] Özgür Ökcü and Ekrem Aydıner, “Joule–Thomson expansion of Kerr–AdS black holes,” *The European Physical Journal C* **78**, 1–6 (2018).
- [83] Shan-Quan Lan, “Joule-Thomson expansion of charged Gauss-Bonnet black holes in AdS space,” *Physical Review D* **98**, 084014 (2018).
- [84] Shihao Bi, Minghao Du, Jun Tao, and Feiyu Yao, “Joule-Thomson expansion of Born-Infeld AdS black holes,” *Chinese Physics C* **45**, 025109 (2021).
- [85] Adolfo Cisterna, Shi-Qian Hu, and Xiao-Mei Kuang, “Joule-Thomson expansion in AdS black holes with momentum relaxation,” *Physics Letters B* **797**, 134883 (2019).
- [86] Jing-Tong Xing, Yuan Meng, and Xiao-Mei Kuang, “Joule-Thomson expansion for hairy black holes,” *Physics Letters B* **820**, 136604 (2021).
- [87] Jing Liang, Wei Lin, and Benrong Mu, “Joule–Thomson expansion of the torus-like black hole,” *The European Physical Journal Plus* **136**, 1169 (2021).
- [88] ATN Silva, MA Anacleto, and L Casarini, “Thin-shell gravastar in a noncommutative BTZ geometry,” *Physics of the Dark Universe* **44**, 101479 (2024).
- [89] Ke Yang, Yue-Zhe Chen, Zheng-Qiao Duan, and Ju-Ying Zhao, “Static and spherically symmetric black holes in gravity with a background Kalb-Ramond field,” *Physical Review D* **108**, 124004 (2023).
- [90] Yiyang Wang, Amnish Vachher, Qiang Wu, Tao Zhu, and Sushant G Ghosh, “Strong gravitational lensing by static black holes in effective quantum gravity,” *The European Physical Journal C* **85**, 1–18 (2025).
- [91] RA Konoplya and OS Stashko, “Probing the effective quantum gravity via quasinormal modes and shadows of black holes,” arXiv preprint arXiv:2408.02578 (2024).