

Adaptivity Gaps for Stochastic Probing with Subadditive Functions

Jian Li Yinchun Liu Yiran Zhang

Institute for Interdisciplinary Information Sciences, Tsinghua University, China.
lapordge@gmail.com, liuyinch23@mails.tsinghua.edu.cn, zhangyir22@mails.tsinghua.edu.cn

Abstract

In this paper, we study the stochastic probing problem under a *general monotone norm* objective. We are given a ground set $U = [n] = \{1, 2, \dots, n\}$, where each element i is associated with an independent nonnegative random variable X_i (with a known distribution). We may probe these elements adaptively, and upon probing an element i , its value X_i is realized. The sequence of probed elements must satisfy a prefix-closed feasibility constraint \mathcal{F} , such as a matroid, an orienteering constraint, or any other downward-closed constraint. We also have a monotone norm function $f : \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}$. Let $P \subseteq U$ be the set of probed elements. Then the reward is $f(X_P)$, where X_P is an n -dimensional vector whose i -th coordinate equals the realized value of X_i if $i \in P$ (i.e., element i is probed), and 0 otherwise. Our objective is to design a probing strategy that maximizes the expected reward $\mathbb{E}[f(X_P)]$.

We study the *adaptivity gap* of the problem, defined as the ratio between the expected reward of an optimal adaptive strategy and that of an optimal non-adaptive strategy. A small adaptivity gap allows us to focus on designing non-adaptive strategies, which are typically simpler to represent and analyze. Establishing tight adaptivity gaps is a central challenge in stochastic combinatorial optimization and has been studied extensively for stochastic probing problems with various objective functions. In this paper, we resolve a central open problem in this line of research, posed in [GNS17, KMS24], by proving that the adaptivity gap for stochastic probing with general monotone norms is bounded by $O(\log^2 n)$. With a refined analysis, we can further strengthen the bound to $O(\log r \log n / \log \log n)$ where r is the maximum length of a sequence in the feasibility constraint ($2 \leq r \leq n$). As a by-product, we obtain an asymptotically *tight* adaptivity gap $\Theta(\log n / \log \log n)$ for Bernoulli stochastic probing with binary-XOS objectives, matching the lower bound in [GNS17]. We also obtain an $O(\log^3 n)$ upper bound for Bernoulli stochastic probing with general subadditive objectives. Furthermore, for *monotone symmetric norms*, we prove that the adaptivity gap can be bounded by $O(1)$, answering an open question posed in [PRS23] and improving upon their $O(\log n)$ upper bound.

arXiv:2504.15547v1 [cs.DS] 22 Apr 2025

Contents

1	Introduction	3
1.1	Other Related Work	5
1.2	Organization of the Paper	5
1.3	Proof Overview	6
1.3.1	Step 1: Reduction to Bernoulli Setting	6
1.3.2	A Natural Non-adaptive Strategy	7
1.3.3	Step 2: Reduction to a Special XOS Norm	8
1.3.4	Warm Up: A Simplified Setting	8
1.3.5	Step 3: The Main Proof Ideas	9
2	Preliminaries	13
2.1	Mathematical Preliminaries and Notations	13
2.2	Bernoulli Stochastic Probing	14
3	Adaptivity Gap for Bernoulli Stochastic Probing	16
3.1	Reduction to a Special XOS Norm	16
3.2	Main Algorithm to Categorize Outcomes (ℓ, R)	18
3.3	Proof for Theorem 3.3	24
4	Tight Bounds for Binary-XOS	27
4.1	Technical Inequalities	27
4.2	Proof for Theorem 4.1	28
5	Adaptivity Gap for Symmetric Norm	32
6	Generalization to non-Bernoulli Case and Other Objectives	36
7	Conclusion and Future Directions	37
	Appendices	38
A	Basic Lemmas for Stochastic Probing	38
B	Missing Proof of Lemma 4.4	40
C	Missing Proof for Section 6	43

1 Introduction

In this paper, we investigate the *stochastic probing* problem, a fundamental framework in the algorithmic study of optimization under uncertainty. This problem has been widely explored in the presence of various combinatorial constraints and objective functions (see, e.g., [GN13, GNS16, GNS17, FLX18, BSZ19, PRS23, KMS24]). Further discussions can be found in Section 1.1.

Stochastic probing has numerous applications across different domains, including Bayesian mechanism design [GN13], online learning [CHL⁺16], influence maximization [GNS17], robot path planning [GNS16, GNS17] and database management [LPRY08]. In this work, we focus on a general stochastic probing problem characterized by a monotone norm objective function and general combinatorial constraints.

Definition 1.1 (Stochastic Probing). We are given a ground universe $U = [n] = \{1, 2, \dots, n\}$ of elements, where each element i is associated with an independent nonnegative random variable $X_i \in \mathbb{R}_{\geq 0}$ (with a known distribution that may vary across elements). The exact values of the X_i 's are initially unknown. We are allowed to *probe* the elements adaptively, and upon probing an element, we observe the realized value of the corresponding random variable. In addition, we are provided with an objective function $f : \mathbb{R}_{\geq 0}^U \rightarrow \mathbb{R}_{\geq 0}$ and a *prefix-closed* feasibility constraint $\mathcal{F} \subseteq U^*$,¹ such that probing is only allowed along sequences in \mathcal{F} (which captures combinatorial constraints like matroids, orienteering, or any arbitrary downward-closed constraint). Let $P \subseteq U$ denote the set of probed items. The reward is then given by $f(X_P)$, where $X_P \in \mathbb{R}_{\geq 0}^U$ is defined by setting its i -th coordinate to X_i if i is probed (i.e., if $i \in P$) and to 0 otherwise. Our objective is to design a probing strategy that maximizes the expected reward $\mathbb{E}[f(X_P)]$.

We also examine a special case where each random variable X_i follows a Bernoulli distribution.

Definition 1.2 (Bernoulli Stochastic Probing). In the stochastic probing problem, each random variable X_i is 1 with probability p_i and 0 with probability $1 - p_i$ for each $i \in U$. If we probe element i and X_i is realized to 1, we say that element i is *active*. The objective function f simplifies to a set function $f : 2^U \rightarrow \mathbb{R}_{\geq 0}$. Let $P \subseteq U$ denote the set of probed items, and $\text{Ac}(P) \subseteq P$ be the set of active elements in P . The reward in this case is $f(\text{Ac}(P))$. All other settings are the same as Definition 1.1, and our goal is also to find a probing strategy to maximize the expected reward $\mathbb{E}[f(\text{Ac}(P))]$.

In this work, we consider general objective functions, including monotone norms, subadditive functions, fractionally subadditive (XOS) functions, and binary-XOS functions (a subclass of XOS functions where weight vectors have 0/1 coordinates, see [BV22]). The detailed definitions and relationships among these function classes are provided in Section 2.1.

In general, the optimal strategy for stochastic probing is *adaptive*, meaning that elements are probed sequentially, with each decision based on previously observed outcomes. This decision process can be represented as an exponentially large tree, making it conceptually challenging and computationally expensive.

To address these difficulties, researchers often consider *non-adaptive* strategies, which commit in advance to a probing sequence $P \in \mathcal{F}$ without utilizing observed realizations. The expected reward in this case is $\mathbb{E}[f(X_P)]$, where X_P contains the ‘revealed’ values in P with unprobed entries set to zero. Non-adaptive strategies significantly simplify computation and storage by eliminating the need to maintain a large decision tree.

To evaluate the effectiveness of non-adaptive strategies, we define the *adaptivity gap* as the ratio between the expected reward of the optimal adaptive strategy and that of the best non-adaptive strategy.² A small adaptivity gap justifies the use of non-adaptive strategies, which are typically easier to analyze. Consequently, proving tight adaptivity gaps has been a central challenge in stochastic optimization, with prior studies covering problems such as stochastic knapsack [DGV04], stochastic covering [GV06, AAK19], stochastic matching [BGL⁺12], stochastic orienteering [BN14], stochastic TSP [JLLS20], and the general stochastic probing problems with various objective functions [GNS16, GNS17, PRS23, KMS24].

¹ U^* denotes the set of all sequences of distinct elements in U of length at most n . We use $|F|$ to denote the length of the sequence F for $F \in U^*$. \mathcal{F} is *prefix-closed* means that if a sequence $F \in \mathcal{F}$, then all its prefixes are also included in \mathcal{F} .

²It is known that one can restrict attention to deterministic adaptive strategies, which can be modeled as decision trees; see Section 2.2 for further discussion.

We first review prior results on the adaptivity gap in stochastic probing. The Bernoulli stochastic probing problem with general feasibility constraints was introduced in [GN13] and its adaptivity gap was systematically studied by Gupta and Nagarajan [GNS16], who show a constant adaptivity gap for prefix-closed constraints when f is a linear or weighted coverage function. Gupta, Nagarajan, and Singla [GNS17] then extended these results by establishing constant adaptivity gaps for both monotone and non-monotone submodular functions. In the same paper, they considered the more general *subadditive functions*,¹ and conjectured that the adaptivity gap for all subadditive functions is poly-logarithmic in the size of the ground set (Conjecture 1). Recall that a set function $f : 2^U \rightarrow \mathbb{R}_{\geq 0}$ is subadditive if $f(X \cup Y) \leq f(X) + f(Y)$ for all $X, Y \in 2^U$. The class of subadditive functions is very general and includes submodular functions and XOS functions as special cases. They also showed that the adaptivity gap for XOS functions is $O(\log W)$, where W denotes the width of the XOS representation. It is known that a subadditive function can be approximated by an XOS function within a logarithmic factor [Dob07, BR11]. Therefore, to resolve Conjecture 1, it suffices to provide a poly-logarithmic upper bound for XOS functions (for the Bernoulli case). However, there are subadditive functions (even submodular functions) that require exponentially large width in XOS representation [BH11, BDF⁺12]. For lower bound, the authors also show that there are instances where the adaptivity gap is at least $\Omega(\log W / \log \log W)$ for $W = \Theta(n)$.

The conjecture has attracted significant attention in recent years. Patton et al. [PRS23] proved the conjecture for symmetric norm objectives and general non-negative random variables (not necessarily Bernoulli). In particular, they showed the adaptivity gap for symmetric norms is $O(\log n)$, and posed the open question that whether the true adaptivity gap is sub-logarithmic or even a constant (Conjecture 2). Kesselheim et al. [KMS24] approached the problem from the notion of p -supermodularity for norms. In particular, a norm is p -submodular if its p th power is supermodular. They showed that for p -supermodular norms, the adaptivity gap is at most $O(p)$. Motivated by the conjecture of [GNS17], Kesselheim et al. [KMS24] extended the conjecture to monotone norms,² and conjectured that the adaptivity gap for stochastic probing is polylogn for any monotone norm (Conjecture 3).

In this paper, we resolve all three conjectures mentioned above. For the Bernoulli stochastic probing problem, we prove the following theorem, thus answering Conjecture 1 [GNS17] affirmatively.

Theorem 1.1 (See also Corollary 6.4). *The adaptivity gap for Bernoulli stochastic probing with any subadditive objective f is upper bounded by $O(\log^3 n)$.*

For the (general) stochastic probing problem with general monotone norms, we have the following theorem, confirming Conjecture 3 [KMS24] affirmatively:

Theorem 1.2 (See also Corollary 6.5). *The adaptivity gap for stochastic probing with any monotone norm f (not necessarily symmetric) is upper bounded by $O\left(\frac{\log r \log n}{\log \log n}\right)$, where $2 \leq r \leq n$ is an upper bound for the maximal length of a sequence in \mathcal{F} .*

In view of the lower bound $\Omega(\log n / \log \log n)$ established in [GNS17], we reduce the gap between the upper and lower bounds to a factor of $O(\log r)$. Moreover, as a by-product of our proof for Theorem 1.2, we establish that the adaptivity gap for Bernoulli stochastic probing with binary-XOS objectives is upper bounded by $O(\log n / \log \log n)$. Notably, in the lower bound instance in [GNS17], the objective function f is indeed a binary-XOS objective. Therefore, our upper bound is asymptotically tight, as it matches the existing lower bound.

Theorem 1.3 (See also Theorem 4.3). *The adaptivity gap for Bernoulli stochastic probing with binary-XOS objective is $\Theta\left(\frac{\log n}{\log \log n}\right)$.*

Finally, we improve the result in [PRS23] by reducing the upper bound of the adaptivity gap for monotone symmetric norms from $O(\log n)$ to $O(1)$, thereby answering one of the open problems posed in that work (Conjecture 2).

¹Normally, the objective f is always supposed to be **monotone** and **normalized** ($f(\emptyset) = 0$ or $f(\mathbf{0}) = 0$) unless stated otherwise, see also Section 2.1. We usually omit the two conditions for clarity.

²A monotone norm can be also be written as the max over linear functions, hence can be seen as an extension of XOS set functions to the domain $\mathbb{R}_{\geq 0}^U$

Theorem 1.4 (See also Corollary 6.6). *The adaptivity gap for stochastic probing with monotone symmetric norm is $O(1)$.*

1.1 Other Related Work

Stochastic probing is a versatile framework that encompasses many well-studied problems in stochastic combinatorial optimization. For instance, stochastic matching [CIK⁺09, Ada11, BGL⁺12]—with applications in kidney exchanges [RSU05, AR12] and online dating—considers a random graph G where probing an edge e reveals the outcome of an independent Bernoulli variable X_e , indicating whether $e \in E(G)$. Here, the objective function f is the size of the maximum matching, and the feasibility constraints are the valid matchings on the vertex set. Beyond matching, stochastic probing has been applied to Bayesian mechanism design [GN13], robot path planning [GNS16, GNS17], orienteering [GM09, GKNR12, BN14], and stochastic set cover problems in databases [LPRY08, DHK14].

The study of adaptivity in the Bernoulli stochastic probing problem with arbitrary prefix-closed constraints \mathcal{F} was initiated by Gupta et al. [GNS17]. They showed that even with a simple cardinality constraint, there exists a monotone function $f : 2^U \rightarrow \mathbb{R}_{\geq 0}$ for which the adaptivity gap is at least $2^{\Omega(\sqrt{n})}$, implying that further structural assumptions are necessary for tighter bounds. They further proved that for non-negative submodular functions $f : 2^U \rightarrow \mathbb{R}_{\geq 0}$, the gap is bounded by 3 (or 40 when f is non-monotone), with Bradac et al. [BSZ19] later tightening the bound to 2 for monotone submodular functions.

For specific constraints, Asadpour and Nazerzadeh [AN16] established that the adaptivity gap for monotone submodular functions under a single matroid constraint is exactly $\frac{e}{e-1}$. In addition, building on [GNS16], Bradac et al. [BSZ19] demonstrated that the adaptivity gap for weighted rank functions of an intersection of k matroids or a k -extendible system (see [Mes06, CCPV11]) lies between k and $O(k \log k)$.

Recent research has focused on general objectives such as ordered norms [BSS18, CS18] and symmetric norms [CS19a, CS19b, IS20, DLR23, ABB⁺23, KMS24]. Approximation algorithms for norm minimization, initiated by Chakrabarty and Swamy [CS19a], have been applied to problems including generalized load balancing [DLR23, ABP23], stochastic optimization [IS20], online algorithms, and stochastic probing [PRS23], as well as parameterized algorithms [ABB⁺23]. A key technique in these works is to approximate general norms by classical ones (e.g., ordered, Top- k , Orlicz, or super/submodular norms; see [KMS23, KMS24]), which is particularly effective for analyzing the adaptivity gap.

In particular, Patton et al. [PRS23] showed that any monotone symmetric norm can be approximated within a factor of $O(\log n)$ by a submodular norm, yielding an adaptivity gap of $O(\log n)$. Kesselheim et al. [KMS24] introduced p -supermodular norms and established an $O(p)$ adaptivity gap for this class, further showing that any monotone symmetric norm can be approximated by an $O(\log n)$ -supermodular norm, leading again to an $O(\log n)$ gap.

A summary of these results, along with a comparison of our findings on adaptivity gaps for stochastic probing with prefix-closed constraints, is provided in Table 1.

1.2 Organization of the Paper

The remainder of the paper is organized as follows. In Section 1.3, we provide a detailed overview of our proof. In Section 2, we present the necessary background and preliminaries. In Section 3, we prove our main theorem (Theorem 3.1), showing that the adaptivity gap for Bernoulli stochastic probing with any XOS objective is upper bounded by $O(\log r \log n)$, where $2 \leq r \leq n$ is an upper bound for the maximum length of a sequence in \mathcal{F} . In Section 4, through more refined analysis, we further improve this bound to $O\left(\log r \cdot \frac{\log n}{\log \log n}\right)$ (see Theorem 4.2) and also obtain the tight bound $\Theta\left(\frac{\log n}{\log \log n}\right)$ of the adaptivity gap for Bernoulli stochastic probing with binary-XOS objective. In Section 5, we establish that the adaptivity gap for Bernoulli stochastic probing with any symmetric norm objective (defined in Definition 5.1) is $O(1)$ (see Theorem 5.4). Finally, in Section 6, we invoke a well-known result that any subadditive objective can be approximated by an XOS function within logarithmic factors to obtain poly-logarithmic adaptivity gap for Bernoulli stochastic probing with any subadditive objective (see Corollary 6.4). In the same section, we present a reduction from the setting of general random variables to the Bernoulli setting, which leads to a poly-logarithmic adaptivity gap for (general) stochastic probing with any monotone norm (see Corollary 6.5) and $O(1)$ gap for (general) stochastic probing with symmetric monotone norm (see Corollary 6.6).

Variables X_i	Objective Class f^*	Adaptivity Gap
Bernoulli	Weighted rank of k -extendible system	$[k, O(k \log k)]$ [BSZ19]
Bernoulli	Submodular	≤ 3 [GNS17] (= 2 in [BSZ19])
Bernoulli	Non-monotone submodular	≤ 40 [GNS17]
Bernoulli	XOS of width W	$O(\log W)$ [GNS17]
Bernoulli	binary-XOS	$\Omega\left(\frac{\log n}{\log \log n}\right)$ [GNS17]
Bernoulli	binary-XOS	$O\left(\frac{\log n}{\log \log n}\right)$ (Theorem 1.3)
Bernoulli	XOS	$O\left(\frac{\log r \log n}{\log \log n}\right)^\dagger$ (Theorem 4.2)
Bernoulli	Subadditive	$O(\log^3 n)$ (Theorem 1.1)
General	Symmetric norm	$O(\log n)$ [PRS23, KMS24]
General	Symmetric norm	$O(1)$ (Theorem 1.4)
General	p -supermodular norm	$O(p)$ [KMS24]
General	General norm	$O\left(\frac{\log r \log n}{\log \log n}\right)$ (Theorem 1.2)

* All objectives are assumed to be **monotone** and **normalized** ($f(\emptyset) = 0$ or $f(\mathbf{0}) = 0$) unless stated otherwise.

$\dagger 2 \leq r \leq n$ denotes an upper bound for the maximal length of a sequence in feasibility constraint \mathcal{F} .

Table 1: Adaptivity gaps for stochastic probing with prefix-closed constraints

1.3 Proof Overview

In this section, we provide an overview of the proof for Theorem 1.2, the main result of this paper. For a roadmap of our proof, please refer to Fig. 1. For simplicity, we present only the main ideas that establish an $O(\log^2 n)$ upper bound on the adaptivity gap for any monotone norm objective.




Figure 1: Roadmap for Theorem 1.2.

1.3.1 Step 1: Reduction to Bernoulli Setting

We first reduce general random variables to Bernoulli random variables. The idea of the reduction is standard and already appears implicitly in [Sin18, GNS17]. Suppose that we aim to prove that the adaptivity gap is bounded by the function $\Lambda(n) \geq 2$. We denote the expected reward for the optimal (adaptive) strategy by OPT . Without loss of generality, we assume $\text{OPT} = 1$ by scaling.

We say that an outcome x_i of X_i is λ -large if $f(x_i) \geq \lambda \cdot \text{OPT}$.¹ Otherwise, we say it is λ -small. Then, for any probed sequence $P \subseteq [n]$, we can decompose the random variable X_P into two parts: $Y_{P,\lambda}$ only

¹ $f(x_i)$ is a shorthand for $f(0, 0, \dots, 0, x_i, 0, 0, \dots, 0)$.

contains λ -large outcomes (with placeholder zero if an outcome is λ -small), and $Z_{P,\lambda} = X_P - Y_{P,\lambda}$ contains the remaining λ -small outcomes. We also call a random variable λ -small if all of its possible outcomes are λ -small. An analogue of the work in [GNS17] shows that, if $\mathbb{E}[f(Y_{P,\lambda})] \geq \text{OPT}/2$, then the adaptivity gap is at most $O(1 + 1/\lambda)$, see also Lemma A.2. Therefore, we can always assume $\mathbb{E}[f(Z_{P,\lambda})] \geq \text{OPT}/2$ using subadditivity of f , if we set $\lambda = \Lambda(n)^{-1}$. Hence, we can replace X_i by the random variables $Z_{i,\lambda}$ with only λ -small outcomes. Since $\text{OPT} = 1$, by the above procedure, we can assume that all outcomes of X_i are $1 \geq 2\Lambda(n)^{-1}$ -small. We can also ignore any outcome x_i of X_i such that $f(x_i) \leq 1/(2n)$ since their total contribution is at most $1/2$. Now, we round the outcome of X_i to the nearest negative power of 2. Hence, each random variable X_i has at most $2 \log n$ different outcomes, which implies that each node in the decision tree (corresponding to an adaptive strategy) has at most $O(\log n)$ children. We can replace the current decision tree by a binary decision tree, which corresponds to a Bernoulli stochastic probing problem with a monotone norm objective. The details can be found in Appendix C. Moreover, it is well known that the restriction of any monotone norm on 2^U defines an XOS function, see Section 2.1 for details. Hence, we obtain the following theorem.

Theorem 1.5 (See also Theorem 6.2). *Suppose the adaptivity gap for Bernoulli stochastic probing with XOS objective is $\Lambda_{\text{xos}}(n)$, then the adaptivity gap for stochastic probing with any general monotone norm is $O(\Lambda_{\text{xos}}(n \log 8n))$.*

1.3.2 A Natural Non-adaptive Strategy

In the following, it suffices to consider the Bernoulli setting. We first need to define some notations regarding the decision tree for an adaptive strategy, and then introduce the construction of the non-adaptive strategy.

Adaptive Strategies. We denote the optimal adaptive strategy by `ALGTREE`, which can be represented by a rooted binary tree \mathcal{T} , where each internal node v represents an element $i \in U$, denoted as $\text{elt}(v) = i$. We also say that the node v **represents** the element i . If i is *active* (with probability p_i given in the definition of Bernoulli setting), the **Yes** arc is followed; otherwise, the **No** arc is taken. We require that no element can be represented by more than one node on any root-to-leaf path in the tree, so the length of each root-leaf path is trivially bounded by n . Each root-to-leaf path must be feasible, i.e., the element sequence must be contained in \mathcal{F} . The leaves indicate the termination of the probing process but do not represent elements in U .

Each leaf ℓ is associated with a unique path P_ℓ , which consists of all nodes visited from the root to ℓ . For a node v , we call an ancestor u of v its **active ancestor** if the **Yes** arc from u is taken on the path from root to v . Further, we call a node of **depth** k if it has k active ancestors. An illustration is shown in Fig. 4. The set of active ancestors of ℓ on P_ℓ is denoted as A_ℓ . Nodes in one single A_ℓ must have different depth. A node belongs to A_ℓ if its corresponding element is active when probed in the adaptive strategy `ALGTREE`. For any subset $S \subseteq P_\ell$, we denote the corresponding set of elements as $\text{elt}(S) = \{\text{elt}(v) \in U : v \in S\}$.

The tree induces a probability distribution $\pi_{\mathcal{T}}$ over its leaves. Given a monotone function f , the expected reward of `ALGTREE` is:

$$\text{Adap}(\mathcal{T}, f) := \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(\text{elt}(A_\ell))].$$

A Natural Non-adaptive Strategy. We consider a natural non-adaptive strategy which is derived from `ALGTREE` (with tree \mathcal{T}). The non-adaptive strategy first selects a leaf $\ell \leftarrow \pi_{\mathcal{T}}$, probes all elements on the path P_ℓ , and selects the active elements. Clearly, this non-adaptive strategy satisfies the feasibility constraint. We denote such strategy as `ALGPATH`. The same non-adaptive strategy has been studied in [GNS16, GNS17, BSZ19]. Note that the realization of the elements during the probing process of `ALGPATH` is independent of that in `ALGTREE` (e.g., a node $v \in A_\ell$ may be inactive in `ALGPATH`). To capture the randomness of the realizations in the probing process of `ALGPATH`, we define \mathbb{P}_R , which is a distribution of the random subset R of nodes on \mathcal{T} . In particular, R is obtained as follows: for each element e , with independent probability p_e , all nodes v that $\text{elt}(v) = e$ are included in R . In other words, each node $v \in P_\ell$ is included in R with probability $p_{\text{elt}(v)}$, which means that its corresponding element is active in `ALGPATH`. Hence, we can describe the non-adaptive strategy `ALGPATH` as first sampling a leaf $\ell \leftarrow \pi_{\mathcal{T}}$, then collecting all elements in $R \cap P_\ell$, where $R \sim \mathbb{P}_R$. Therefore, the expected reward of `ALGPATH` is:

$$\text{Nadp}(\mathcal{T}, f) := \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f(\text{elt}(R \cap P_\ell))] \right].$$

By a similar argument as in Section 1.3.1, we can assume without loss of generality that there is no "large" elements (those e with $f(e) \geq \Lambda(n)^{-1} \text{Adap}(\mathcal{T}, f)$). Hence, by scaling the function f , it suffices to focus on the case where $\text{Adap}(\mathcal{T}, f) = \Lambda(n)$ and $f(e_j) \leq 1, \forall j \in U$, and we need to show that $\text{Nadp}(\mathcal{T}, f) \geq 1$.

1.3.3 Step 2: Reduction to a Special XOS Norm

Now, we would like to further simplify the XOS norm to a special family of XOS norms. We first recall the XOS norm as $f(S) = \max_{i \in [W]} \left\{ \sum_{j \in S} w_j^{(i)} \right\}$, where W is the width of the XOS norm and $w_j^{(i)} \geq 0$ for all i, j . Since $f(e_j) \leq 1$, we have all $w_j^{(i)} \leq 1$ for all $i \in [W], j \in U$. Next, by rounding each weight $w_j^{(i)}$ to a negative power of 2 and ignoring those less than $1/(2n)$, we are left with an instance with only $O(\log n)$ possible weights. By picking the one with the largest contribution (suppose it is the one with weight 2^{-j_0}), by losing a factor of at most $O(\log n)$, it suffices to deal with the case where all weights are 2^{-j_0} or zero.

Through the above manipulations (see the full details in Lemma 3.2), we arrive at a special family of XOS norms as follows: for each ℓ , there is a subset $A'_\ell \subseteq A_\ell$, and the special XOS norm objective (with respect to the subsets A'_ℓ) is

$$f_{\text{xos}}(S) = \max_{\ell} \{ |\text{elt}(A'_\ell) \cap S| \}.$$

Suppose that we can prove the following statement: when $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|] = \Omega\left(\frac{\Lambda(n)}{\log n}\right)$, we always have

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \geq \frac{\log n}{\Lambda(n)} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|].$$

Then, we can use $f \geq 2^{-j_0} f_{\text{xos}}$ and $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|] = \Omega\left(2^{j_0} \frac{\Lambda(n)}{\log n}\right)$ (by the maximality of the contribution of weight 2^{-j_0}) to prove that:

$$\begin{aligned} \text{Nadp}(\mathcal{T}, f) &\geq 2^{-j_0} \text{Nadp}(\mathcal{T}, f_{\text{xos}}) = 2^{-j_0} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \\ &\geq \frac{\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|]}{2^{j_0}} \cdot \frac{\log n}{\Lambda(n)} = \Omega(1). \end{aligned}$$

Thus, we can conclude that the adaptivity gap is $O(\Lambda(n))$. We encapsulate the above argument as follows:

Lemma 1.6 (See also Lemma 3.2). *Suppose that for any decision tree \mathcal{T} and any choice of subsets $A'_\ell \subseteq A_\ell$ satisfying $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|] \geq 20 \log n$, the following condition holds:*

$$\text{Nadp}(\mathcal{T}, f_{\text{xos}}) = \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \geq \frac{1}{20 \log n} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|].$$

Then, the adaptivity gap of Bernoulli stochastic probing with XOS Objective is upper bounded by $O(\log^2 n)$.

1.3.4 Warm Up: A Simplified Setting

From the previous sections, we can see that it suffices to prove the condition of Lemma 1.6 holds. To begin with, we consider the following simplified case, where the number of leaves of \mathcal{T} is $O(n)$. The simplified case motivates the most technical part of the proof, presented in Section 1.3.5.

When the number of leaves of \mathcal{T} is $O(n)$, we can directly see the lower bound

$$f_{\text{xos}}(\text{elt}(R \cap P_\ell)) \geq |\text{elt}(R \cap P_\ell) \cap \text{elt}(A'_\ell)| = |R \cap A'_\ell| \tag{1.1}$$

for any leaf ℓ . Then, by definition of the distribution $\pi_{\mathcal{T}}$, p_ℓ is the probability of reaching the leaf ℓ , which is the product of the probability for each node on P_ℓ taking the corresponding Yes or No arc. Therefore, using AM-GM inequality, we can see that

$$p_\ell = \prod_{u \in A_\ell} p_{\text{elt}(u)} \cdot \prod_{u \in P_\ell \setminus (A_\ell \cup \{\ell\})} (1 - p_{\text{elt}(u)}) \leq \prod_{u \in A'_\ell} p_{\text{elt}(u)} \leq \left(\frac{\sum_{u \in A'_\ell} p_{\text{elt}(u)}}{|A'_\ell|} \right)^{|A'_\ell|}.$$

Moreover, it is clear that the $\mathbb{E}_{R \sim \mathbb{P}_R} |R \cap A'_\ell| = \sum_{u \in A'_\ell} p_{\text{elt}(u)}$. Putting together, we have

$$\begin{aligned} \text{Nadp}(\mathcal{T}, f_{\text{xos}}) &= \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \geq \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [|R \cap A'_\ell|] \right] \\ &\geq \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[|A'_\ell| p_\ell^{1/|A'_\ell|} \right] = \sum_{\ell} |A'_\ell| p_\ell^{1+1/|A'_\ell|}. \end{aligned}$$

Now, suppose that $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} |A'_\ell| = \sum_{\ell} p_\ell |A'_\ell| \geq 10 \log n$. We only need to consider the ‘‘good’’ leaves ℓ such that $|A'_\ell| \geq \log n$ and $p_\ell \geq 1/n^2$, since the contribution of other leaves at most $\log n + O(n)/n^2$ (using the number of leaves bounded by $O(n)$), and is at most half of $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} |A'_\ell|$. Now, for a good leaf ℓ , we have $p_\ell^{1/|A'_\ell|} \geq n^{-2/\log n} = \Omega(1)$. Therefore, we can conclude

$$\text{Nadp}(\mathcal{T}, f_{\text{xos}}) \geq \sum_{\ell \text{ good}} |A'_\ell| p_\ell^{1+1/|A'_\ell|} \geq \Omega(1) \cdot \sum_{\ell \text{ good}} |A'_\ell| p_\ell \geq \Omega(1) \cdot \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|].$$

Therefore, the condition for Lemma 1.6 indeed holds for this special case (with even better factor).

However, for the general setting, the above proof does not work anymore since the probability p_ℓ of each leaf can be extremely small. In fact, simply lower bounding $f_{\text{xos}}(\text{elt}(R \cap P_\ell))$ by $|R \cap A'_\ell|$ as in (1.1) is no longer sufficient in the general case. Indeed, one can consider the example where $A_\ell = A'_\ell$ and \mathcal{T} is a complete binary rooted tree of height h , where each node has probability λ to be active. Simple calculation shows that $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} |A_\ell| = \lambda h$ and $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} |R \cap A_\ell| = \lambda^2 h$. Therefore, in the case that $\lambda h = \log^k n$, and $\lambda = \log^{-2k} n$, where $k = \omega(1)$, the above proof does not yield a poly-logarithmic ratio (the number of nodes is $2^h \gg n$ in this example). Therefore, it is crucial to consider the structure of the tree and consider leaf ℓ' other than ℓ itself to make $|\text{elt}(R \cap P_\ell) \cap \text{elt}(A'_{\ell'})|$ as large as possible (thus improving lower bound of (1.1)).

Although the above proof does not extend to the general case, there are still some key insights in it. One key observation is that we can view the set $R \cap A'_\ell$ as the output of the following algorithm: on input (ℓ, R) , we start at leaf ℓ and climb up the path P_ℓ , while we greedily grab nodes in A'_ℓ that turns out to be in R and eventually obtain and output the whole set $R \cap A'_\ell$. This motivates us to construct an algorithm that starts from leaf ℓ , climbs along P_ℓ and try to select a subset of nodes in $R \cap P_\ell$ that is guaranteed to be contained within $A'_{\ell'}$ for some other leaf ℓ' . In fact, in the main proof (ideas presented below in Section 1.3.5), we design an (inefficient) algorithm on input ℓ and each possible outcome of R , starting from leaf ℓ and change the leaf ℓ' greedily to make the intersection $|\text{elt}(R \cap P_\ell) \cap \text{elt}(A'_{\ell'})|$ larger. We prove a tailored *tail bound* to prove the probability that the algorithm terminates with small intersections is exponentially small.

1.3.5 Step 3: The Main Proof Ideas

In this section, we sketch the main idea of the proof. To avoid confusion, we recall that the word *active ancestor* refers to the active nodes in ALGTREE, while we use $v \in R$ to indicate that $\text{elt}(v)$ is realized *active* in the probing of ALGPATH. We only consider the case $A_\ell = A'_\ell$ for simplicity: extending to the general case follows essentially the same idea but with more technical details. Recalling the form of Lemma 1.6, in this section, we aim to prove that under condition $\mathbb{E}_\ell [|A_\ell|] \geq 20 \log n$, it holds that

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \geq \frac{1}{20 \log n} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A_\ell|], \quad (1.2)$$

where $f_{\text{xos}}(S) = \max_{\ell} \{|\text{elt}(A_\ell) \cap S|\}$.

As alluded in Section 1.3.4 above, for each (ℓ, R) , we want to find a leaf ℓ_0 to make $|\text{elt}(R \cap P_\ell) \cap \text{elt}(A_{\ell_0})|$ as large as possible. The intuition is to use a *greedy* algorithm to find this ℓ_0 . We describe the algorithm as follows. The input of the algorithm is (ℓ, R) . We iterate from the leaf ℓ to the root in P_ℓ , while maintaining a sequence $B = (e_1, \dots, e_m)$, where e_i are elements in U . B is initialized as an empty sequence. At each node $v \in P_\ell$, we examine whether the following two conditions hold:

- (1) $v \in R$.
- (2) There is a leaf ℓ' of v such that $v \in A_{\ell'}$, and for each element e_i in the current sequence B , there is a descendant of v (not allowed to be v itself) in $A_{\ell'}$ that *represents* e_i .




Figure 2: In the decision tree, red edges indicate Yes arcs, and blue edges indicate No arcs. We suppose nodes labeled with the same letters (but with different number of primes) represent the same element: $\text{elt}(v_1) = \text{elt}(v'_1) = \text{elt}(v''_1) = \text{elt}(v'''_1)$, $\text{elt}(v_2) = \text{elt}(v'_2) = \text{elt}(v''_2)$, etc. v_4 is the root of the whole tree. Suppose that the specific outcome R_0 includes all green nodes in R_0 , but excludes all white nodes. The algorithm with input (ℓ, R_0) first visits ℓ and next discovers that $v_1 \in A_\ell$ satisfies both conditions. Thus, $B \leftarrow \{\text{elt}(v_1)\}$. After moving up one step without updating, the algorithm detects $v_2 \in R_0 \cap A_{\ell_1}$ has a descendant v'_1 such that v_1, v'_1 represent the same element. Then $B \leftarrow \{\text{elt}(v_1), \text{elt}(v_2)\}$. One should note that we do not require $v_2 \in A_\ell$, so we are considering all nodes in P_ℓ from the leaf to the root, not just the nodes in A_ℓ . After two steps, $v_3 \in R_0 \cap A_{\ell_2}$ is detected, and $B \leftarrow \{\text{elt}(v_1), \text{elt}(v_2), \text{elt}(v_3)\}$. The same thing happens for $v_4 \in R_0 \cap A_{\ell_3}$, where $B \leftarrow \{\text{elt}(v_1), \text{elt}(v_2), \text{elt}(v_3), \text{elt}(v_4)\}$, and the algorithm terminates with output $B(\ell, R_0) := B = \{\text{elt}(v_1), \text{elt}(v_2), \text{elt}(v_3), \text{elt}(v_4)\}$.

If both conditions hold, we append $\text{elt}(v)$ to the end of sequence B . Otherwise, we do nothing. Then we continue to consider the parent of v . Finally, the algorithm outputs B , denoted as $B(\ell, R)$. We use $|B(\ell, R)|$ to denote the number of elements in the sequence $B(\ell, R)$. An example of the algorithm is illustrated in Fig. 2, where only a portion of the binary tree is drawn for clarity.

Recall that different nodes in P_ℓ represents distinct elements. Therefore, due to the two conditions in the algorithm, $|\text{elt}(R \cap P_\ell) \cap \text{elt}(A_{\ell'})| \geq |B|$ always holds throughout the algorithm, where ℓ' is the descendant leaf of v that satisfies condition (2) and incurs the last update for B . Thus, upon termination, the reward for the non-adaptive strategy is at least the output $|B(\ell, R)|$. Hence, we just need to prove the following lower bound

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [|B(\ell, R)|] \stackrel{?}{\geq} \mathbb{E}_{\ell} |A_\ell| / 20 \log n.$$

The idea to prove this result is to bound the tail of the probability that $|B(\ell, R)|$ is at most $0.1|A_\ell|/\log n$ over $\ell \leftarrow \pi_{\mathcal{T}}$ and $R \sim \mathbb{P}_R$. In particular, for any $0 \leq s \leq n$, and any possible sequence B_0 , we can prove that if $|B_0| \leq 0.1s/\log n$ (here, $|B_0|$ denotes the number of elements in the sequence B_0),

$$\Pr_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [B(\ell, R) = B_0, |A_\ell| = s] \leq 2^{-0.9s}. \quad (1.3)$$

Before presenting the proof of (1.3), we first show how (1.3) leads to (1.2). Notice that for each fixed size $m = |B_0|$, there are at most n^m different B_0 's, since elements are in U . We consider the total contribution of (ℓ, R) such that $|A_\ell| > 10 \log n |B(\ell, R)|$ inside $\mathbb{E}_{\ell} [|A_\ell|]$ by summing all possible B_0 and $|A_\ell|$. When n is sufficiently large,

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [\mathbf{1}_{|A_\ell| > 10 \log n \cdot |B(\ell, R)|} |A_\ell|] \leq \sum_{m \geq 0, s > 10m \log n} 2^{-0.9s} \cdot n^m \cdot s \leq \sum_{s \geq 1} 2^{-0.8s} \cdot s^3 = O(1).$$

Using the condition that $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A_{\ell}|] \geq 20 \log n$,

$$\begin{aligned} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [|B(\ell, R)|] &\geq \frac{1}{10 \log n} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [\mathbf{1}_{|A_{\ell}| \leq 10 \log n \cdot |B(\ell, R)|} |A_{\ell}|] \\ &\geq \frac{\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A_{\ell}|] - O(1)}{10 \log n} \geq \frac{\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A_{\ell}|]}{20 \log n}, \end{aligned}$$

which confirms the condition of Lemma 1.6 as promised.

Proof of Inequality (1.3): Now, the only remaining piece of the proof is to prove the tail probability (1.3), which is the most technical part of the whole proof. In the following, we fix the algorithm output B_0 and only consider the leaves ℓ that $|A_{\ell}| = s$. Let $S(B_0)$ be the set of all leaves ℓ such that there exists some $R \sim \mathbb{P}_R$ satisfying $B(\ell, R) = B_0$. Note that the set S does not depend on the specific choice of R .

In the following, we would like to identify nodes u in the tree with the following property (the property is independent of R): if a leaf ℓ which is a descendant of u and the outcome R satisfy $B(\ell, R) = B_0$, then it is guaranteed that u not in R (this is a consequence of u being an **impossible** node, a notion formally defined later). We collect all such nodes to form a set T . We can see that we have the inclusion of the event (over ℓ and R)

$$[B(\ell, R) = B_0] \subseteq [\ell \in S(B_0)] \wedge [(P_{\ell} \cap T) \cap R = \emptyset]$$

(because the latter event is a consequence of the former). We will show that if the length of B_0 is small enough compared to s , the number of impossible nodes in each A_{ℓ} (for $\ell \in S(B_0), |A_{\ell}| = s$) is large, hence the probability that $(P_{\ell} \cap T) \cap R = \emptyset$ is negligible. Therefore, by the inclusion of the event, the probability (over ℓ, R) of $B(\ell, R) = B_0$ is also small.

Recall that in one single execution with input (ℓ, R) , when the iterator reaches a node u , whether u updates B depends on two conditions: (1) $u \in R$; (2) there is a leaf ℓ' of u such that $u \in A_{\ell'}$, and for each element e_i in sequence B , there is a (strict) descendant of u in $A_{\ell'}$ that *represents* e_i . The following observation is crucial to our analysis:

Observation 1.1. *For any descendant leaf ℓ of u and any outcome R that $B(\ell, R) = B_0$, in such execution with input (ℓ, R) , the maintained sequence B must be a prefix of B_0 at the time the iterator visits u (independent of ℓ and R). Specifically, the prefix B contains all elements e_i in B_0 excluding those represented by the ancestors of u (including u).*

From the above observation, whether condition (2) is satisfied does not depend on the particular input R or leaf ℓ , as long as ℓ is a descendant of u and $B(\ell, R) = B_0$. We then call node u an **impossible** node (for fixed B_0) if $\text{elt}(u) \notin B_0$ and condition (2) is satisfied. As a direct consequence, in any execution such that ℓ is a descendant of u and $B(\ell, R) = B_0$, u must not trigger an update (since $\text{elt}(u) \notin B_0$) and hence u must not be in R since it already satisfies condition (2). We conclude that for any impossible node u (for B_0), if $B(\ell, R) = B_0$ and ℓ is a descendant of u , then $u \notin R$. We note that the conclusion holds for *any* descendant leaf ℓ of u , no matter ℓ is in u 's Yes or No subtrees (note that if ℓ is in the No subtree of u , $u \notin A_{\ell}$).

An illustration is drawn in Fig. 3. The main portion of the decision tree is essentially the same as Fig. 2, where nodes of the same letters stand for the same nodes. In addition, some irrelevant parts are abbreviated, while other crucial parts are emphasized.

Now, we need to show that impossible nodes are not rare. From the example in Fig. 3, one may guess that for each $\ell \in S(B_0)$, any node u in A_{ℓ} that $\text{elt}(u) \notin B_0$ is an impossible node. In fact, this is the case. We consider the specific input R_0 and ℓ such that $B(\ell, R_0) = B_0$, whose existence is guaranteed by $\ell \in S(B_0)$. When the iterator approaches u , we consider the node v below u in P_{ℓ} that incurs the latest update ($v \neq u$ because $\text{elt}(u) \notin B_0$). The node v must have a descendant leaf ℓ' that satisfies condition (2). Note that ℓ' is also a descendant of u and hence $u \in A_{\ell'}$ because ℓ and ℓ' share the same active ancestors above v . Since condition (2) does not depend on the input R or leaf ℓ , u is indeed an impossible node. Therefore, for each $\ell \in S(B_0)$, the number of impossible nodes in A_{ℓ} is at least $|A_{\ell}| - |B_0|$, counting the nodes mentioned above.

Now we present the crucial ‘tail’ bound.

Lemma 1.7 (See also Lemma 3.9). *Let S be a subset of leaves in the tree \mathcal{T} . There is a subset T of nodes in \mathcal{T} , which does not contain any leaf of \mathcal{T} . There exists an integer $h \in \mathbb{N}$ such that for each $\ell \in S$, $|A_{\ell} \cap T| \geq h$.*

sequence y_j (the depth of the last $j \log n$ -th node in A'_ℓ) to control the positions of nodes in A'_ℓ . At the same time, the nodes in A_ℓ for $\ell \in S(B_0)$ that represent elements not in B_0 are no longer ensured to be impossible nodes, since they may not be in A'_ℓ . Thus, the arguments presented in the current subsection do not work directly. However, these problems can be remedied by refining the definition of sequence B , which is technically more involved.

2 Preliminaries

2.1 Mathematical Preliminaries and Notations

This subsection introduces basic mathematical definitions and results. Throughout the paper, all objective and functions (whose domain can be 2^U or $\mathbb{R}_{\geq 0}^U$) are assumed to be **monotone** ($f(\mathbf{u}) \leq f(\mathbf{v})$ if $\mathbf{u} \leq \mathbf{v}$, or $f(S) \leq f(T)$ if $S \subseteq T$) and **normalized** ($f(\emptyset) = 0$ or $f(\mathbf{0}) = 0$). We denote the ground set by $U = [n]$. Throughout this paper, \log denotes the logarithm to base 2.

Definition 2.1 ((Monotone) Norm Objective). A function $f : \mathbb{R}_{\geq 0}^U \rightarrow \mathbb{R}_{\geq 0}$ is a *norm objective*, if the following holds:

- (1) $f(\mathbf{0}) = 0$;
- (2) $f(\mathbf{u}) \leq f(\mathbf{v})$ for any $\mathbf{u} \leq \mathbf{v}$, that is, $u_i \leq v_i, \forall 1 \leq i \leq n$, where $\mathbf{u} = (u_i)_{i \in U}$ and $\mathbf{v} = (v_i)_{i \in U} \in \mathbb{R}_{\geq 0}^U$.
- (3) $f(\mathbf{u} + \mathbf{v}) \leq f(\mathbf{u}) + f(\mathbf{v})$ for all $\mathbf{u}, \mathbf{v} \in \mathbb{R}_{\geq 0}^U$;
- (4) $f(\theta \mathbf{v}) = \theta f(\mathbf{v})$ for all $\mathbf{v} \in \mathbb{R}_{\geq 0}^U, \theta \in \mathbb{R}_{\geq 0}$.

Definition 2.2 ((Monotone) Symmetric Norm). A norm $f : \mathbb{R}_{\geq 0}^U \rightarrow \mathbb{R}_{\geq 0}$ is a *symmetric norm* if and only if f remains unchanged under any permutation of its coordinates, i.e., $f(\mathbf{v}) = f((v_{\pi(i)})_{i \in U})$, for any permutation $\pi : U \rightarrow U$ and $\mathbf{v} = (v_i)_{i \in U} \in \mathbb{R}_{\geq 0}^U$.

Definition 2.3 (XOS (set) Function). A set function $f : 2^U \rightarrow \mathbb{R}_{\geq 0}$ is XOS (max-of-sums)¹ if there exist non-negative weight vectors $\mathbf{w}^{(1)}, \mathbf{w}^{(2)}, \dots, \mathbf{w}^{(W)} \in \mathbb{R}_{\geq 0}^U$, such that $f(S) = \max_{i \in [W]} \{\sum_{j \in S} w_j^{(i)}\}, \forall S \subseteq U$.

Definition 2.4 (binary-XOS (set) Function). A set function $f : 2^U \rightarrow \mathbb{R}_{\geq 0}$ is binary-XOS if there exist weight vectors $\mathbf{w}^{(1)}, \mathbf{w}^{(2)}, \dots, \mathbf{w}^{(W)} \in \{0, 1\}^U$, such that $f(S) = \max_{i \in [W]} \{\sum_{j \in S} w_j^{(i)}\}, \forall S \subseteq U$.

Definition 2.5 (Subadditive (set) Objective). A function $f : 2^U \rightarrow \mathbb{R}_{\geq 0}$ is a *subadditive objective*, if and only if the following holds:

- (1) $f(\emptyset) = 0$.
- (2) $f(S) \leq f(T)$ for all $S \subseteq T \subseteq U$.
- (3) $f(S \cup T) \leq f(S) + f(T)$ for all $S, T \subseteq U$.

It is clear that a binary-XOS set function is a special case of an XOS set function, and every XOS function is subadditive. Furthermore, Feige [Fei06] established that a set function is (monotone) fractionally subadditive if and only if it belongs to the XOS class, where a set function $f : 2^U \rightarrow \mathbb{R}_{\geq 0}$ is defined as (monotone) fractionally subadditive if it satisfies $f(T) \leq \sum_i \alpha_i f(S_i)$ for any collection of sets that $\mathbf{1}_T \leq \sum_i \alpha_i \mathbf{1}_{S_i}$ and $\alpha_i \geq 0$. Here, $\mathbf{1}_T$ denotes the indicator function of the subset $T \subseteq U$.

We will also use the well-known Chernoff bound:

Fact 2.1 (Chernoff Bound, see [MU05]). X_1, X_2, \dots, X_n is a sequence of independent random variables. Each X_i is a Bernoulli random variable with parameter $p_i \in [0, 1]$, i.e., $X_i \sim B(p_i)$. Let $X = \sum_{i=1}^n X_i$, and the expected value $\mu = \mathbb{E}[X] = \sum_{i=1}^n p_i$. Then, for any $0 < \epsilon < 1$,

$$\Pr[X < (1 - \epsilon)\mu] \leq \exp\left(-\frac{\epsilon^2 \mu}{2}\right).$$

¹Note that here we implicitly require f to be monotone and normalized.

2.2 Bernoulli Stochastic Probing

Recall that we have defined Bernoulli stochastic probing in Definition 1.2. In this section, we introduce basic definitions and ideas, following the framework of [GNS17].

Definition 2.6 (Bernoulli stochastic probing with XOS (binary-XOS, subadditive, resp.) Objective). In this special case of the stochastic probing problem, each random variable X_i follows a Bernoulli distribution: $X_i = 1$ with probability p_i and $X_i = 0$ with probability $1 - p_i$ for all $i \in U$. The objective function f simplifies to an XOS (binary-XOS, subadditive, resp.) set function $f : 2^U \rightarrow \mathbb{R}_{\geq 0}$. Let $P \subseteq U$ denote the set of probed items, and $\text{Ac}(P) \subseteq P$ be the set of active elements in P . The reward in this case is given by $f(\text{Ac}(P))$. All other settings are the same as in Definition 1.1, and our goal is also to find a probing strategy to maximize the expected reward $\mathbb{E}[f(\text{Ac}(P))]$.

In the Bernoulli setting, the objective degenerates into a set function: $f(S)$ for $S \subseteq U = [n]$ is defined as $f(\mathbf{v})$ where \mathbf{v} is an n -dimensional vector where the i -th coordinate is 0 if $i \notin S$ and is 1 otherwise. We can always assume without loss of generality that $p_i > 0$ for all i . If $p_i = 0$, we can remove the corresponding element from the ground set U and modify the norm accordingly by inserting a zero placeholder at the corresponding position. This modification does not affect the reward or any properties of the objective function f .

We denote the supremum of the adaptivity gap for *Bernoulli Stochastic Probing with any XOS Objective* and any *prefix-closed* feasibility constraint \mathcal{F} with $\max_{F \in \mathcal{F}} |F| \leq r$ as $\Lambda_{\text{XOS}}(n, r)$. We also denote the supremum of the adaptivity gap for *Bernoulli Stochastic Probing with any binary-XOS Objective* and any *prefix-closed* feasibility constraint \mathcal{F} as $\Lambda_{\text{bXOS}}(n)$.

Now we formally introduce the decision tree model of an adaptive strategy.

Adaptive Strategies. An adaptive strategy tree, denoted by \mathcal{T} , is a binary rooted tree where each internal node v represents an element $i \in U$, denoted by $\text{elt}(v) = i$. Every internal node has two outgoing arcs: the **Yes** arc, which leads to the next node if the element $i = \text{elt}(v)$ is active (with probability p_i) when probed, and the **No** arc, which leads to a different node if i is not active (with probability $1 - p_i$). We require that no element can be represented by more than one node on any root-to-leaf path in the tree. Additionally, any root-to-leaf path in \mathcal{T} must satisfy the feasibility constraint \mathcal{F} , meaning it represents a valid sequence of probes for both adaptive and non-adaptive strategies. Each leaf ℓ in \mathcal{T} is associated with a unique root-to-leaf path, denoted by P_ℓ , where the elements probed along this path are given by $\text{elt}(P_\ell)$. For any subset $S \subseteq P_\ell$, we denote the distinct elements represented by the nodes in S as $\text{elt}(S) = \{\text{elt}(v) \in U : v \in S\}$. Let A_ℓ represent the active nodes along the path P_ℓ , i.e., the nodes on P_ℓ where the **Yes** arc is taken. It is important to note that the leaves of the tree do not represent elements in U ; rather, they simply signal the termination of the probing process.

The tree \mathcal{T} naturally gives us a probability distribution $\pi_{\mathcal{T}}$ over its leaves: start at the root, at each node v , follow the **Yes** branch with probability $p_{\text{elt}(v)}$ and the **No** branch otherwise, and finally reach a leaf upon termination.

Given a monotone function f and a tree \mathcal{T} , the associated adaptive strategy probes elements following a root-to-leaf path, observes the realized values until we reach a leaf ℓ , and receives the reward $f(\text{elt}(A_\ell))$. Let $\text{Adap}(\mathcal{T}, f)$ denote the expected reward obtained this way; it can be written compactly as

$$\text{Adap}(\mathcal{T}, f) := \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(\text{elt}(A_\ell))].$$

In fact, any deterministic adaptive strategy can be modeled by such a decision tree model. Moreover, it can be easily proven that the reward of any *randomized* adaptive strategy is bounded by the reward of *some* deterministic adaptive strategy. Since the cardinality of a decision tree is finite, the *optimal* adaptive strategy exists. With the central theme to upper bound the gap between *optimal* adaptive strategies and *optimal* non-adaptive ones in mind, we will only consider deterministic adaptive strategies and the decision tree model described above. We denote the optimal adaptive strategy by **ALGTREE**.

A Natural Non-adaptive Strategy. We can also define a *natural* non-adaptive algorithm given the optimal adaptive strategy **ALGTREE** (with tree \mathcal{T}): just pick a leaf $\ell \leftarrow \pi_{\mathcal{T}}$ from the distribution given by \mathcal{T} , probe all elements on that path, and choose the active elements to get the reward. Clearly, such

strategy satisfies feasibility constraint. We denote this non-adaptive strategy as ALGPATH. We note that the same non-adaptive strategy has been studied in [GNS16, GNS17, BSZ19]. Since the objective f is always assumed to be monotone throughout the paper, the reward is actually the function value of the set of *all* active elements. To capture the randomness of the realizations in the probing process of ALGPATH, we define \mathbb{P}_R , which is a distribution of the random subset R of nodes on \mathcal{T} . A node u is in the random subset R , if the element $\text{elt}(u)$ is active in the probing process of ALGPATH. Specifically, for each element e , with independent probability p_e , all nodes v such that $\text{elt}(v) = e$ are included in R . Otherwise, with probability $1 - p_e$, these nodes are not included.

We denote the expected reward of ALGPATH by $\text{Nadp}(\mathcal{T}, f)$, and we can see that

$$\text{Nadp}(\mathcal{T}, f) := \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f(\text{elt}(R \cap P_{\ell}))] \right].$$

Another way to view the distribution $R \cap P_{\ell}$ is to first sample $\ell \leftarrow \pi_{\mathcal{T}}$, then sample R *only* on the path, with each node $v \in P_{\ell}$ included of independent probability $p_{\text{elt}(v)}$.

Bounding the Adaptivity Gap. A common technique for bounding the adaptivity gap, employed in [GNS16, GNS17, BSZ19], is to bound the ratio between the *optimal* adaptive strategy ALGTREE and its corresponding non-adaptive strategy ALGPATH. Formally, for the optimal tree \mathcal{T}^* , if we can show that the following holds for the function $\Lambda(n)$ when n is sufficiently large:

$$\Lambda(n) \cdot \text{Nadp}(\mathcal{T}^*, f) \geq \text{Adap}(\mathcal{T}^*, f),$$

then the true adaptivity gap of the problem is at most $\Lambda(n)$.

Throughout the paper, we also need the following definitions for trees.

Definition 2.7 (Depth). We say a node has depth k if there are k Yes arcs on the path from the root to this node.

Definition 2.8 (Active Ancestor). For a node v , we call an ancestor u of v its *active ancestor* if the Yes arc from u is taken on the path from root to v . Thus, a node of depth k has k active ancestors. Also recall that for a leaf ℓ of \mathcal{T} , A_{ℓ} is exactly the set of active ancestors of ℓ .

In the following, we will use the definition of depth and active ancestors extensively. It is useful to think of the original tree \mathcal{T} as being layered by depth, and forms a new layered tree, see Fig. 4.




Figure 4: A binary decision tree of height 3 is transformed to a layered tree of depth 3. Note that A is an active ancestor of B and C , but not for D . The active ancestor set $A_C = \{\text{root}, A\}$.

It is clear from definition that different nodes in A_{ℓ} have distinct depths. This fact will be frequently used in the algorithm and proofs in Section 3.2 and Section 3.3.

Now, we introduce some tools for the adaptivity gap problem in the Bernoulli setting. As usual, we denote $\text{OPT} = \text{Adap}(\mathcal{T}^*, f)$, the expected reward for ALGTREE.

Definition 2.9 (λ -large (small) elements). For any $\lambda > 0$, an element e is λ -large if $f(e) \geq \lambda \cdot \text{OPT}$ and λ -small otherwise, where $f(e)$ means the f value when only e is active.

The following lemma already appears implicitly in [GNS17], and the proof is deferred to Appendix A.

Lemma 2.2. *For any $\lambda > 0$, suppose that the adaptivity gap in the Bernoulli setting is upper bounded by $\Lambda(n) \geq 1$ for a certain subadditive objective f and a feasibility constraint \mathcal{F} , under the condition that every element is λ -small with respect to f . Then, the adaptivity gap in the Bernoulli setting for general instances (with general random variables, the same f , and \mathcal{F}) is at most $\max\{2\Lambda(n), 2 + 4/\lambda\}$.*

3 Adaptivity Gap for Bernoulli Stochastic Probing

In this section, we analyze *Bernoulli stochastic probing with XOS objective*, as defined in Definition 2.6. We follow the notations introduced in Section 2.2. Furthermore, recall that $2 \leq r \leq n$ denotes an upper bound on the maximal length of a sequence in \mathcal{F} . Intuitively, r serves as an upper bound on the maximum cardinality of elements that a strategy can probe. Consequently, r also bounds the length of any root-to-leaf path in any adaptive strategy tree \mathcal{T} . Formally, for any leaf ℓ , $|A_\ell| \leq |P_\ell| \leq r$.

The primary result of this section is the following theorem:

Theorem 3.1. *For $2 \leq r \leq n$, the adaptivity gap of Bernoulli Stochastic Probing with any XOS Objective under any prefix-closed \mathcal{F} with $\max_{F \in \mathcal{F}} |F| \leq r$ is $\Lambda_{\text{xos}}(n, r) = O(\log r \log n)$.*

3.1 Reduction to a Special XOS Norm

In this subsection, we present a reduction from the original arbitrary XOS Objective to a specific norm that is more amenable to analysis. Recall the definition of the active ancestor set A_ℓ in Definition 2.8. For each leaf ℓ , we define a subset $A'_\ell \subseteq A_\ell$ (in the proof of Lemma 3.2). Using these subsets, we define a new norm f_{xos} as follows:

$$f_{\text{xos}}(S) = \max_{\ell} \{|\text{elt}(A'_\ell) \cap S|\}, \quad (3.1)$$

for any subset $S \subseteq U = [n]$.

We establish the following lemma:

Lemma 3.2. *Let $\Lambda(n) \geq 2$ be an increasing function of n . Suppose that for any decision tree \mathcal{T} and any choice of subsets $A'_\ell \subseteq A_\ell$ satisfying*

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|] \geq \Lambda(n),$$

the following condition holds:

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \geq \frac{1}{\Lambda(n)} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|].$$

Then the following two results hold:

- (a) *For $r \geq 2$, $\Lambda_{\text{xos}}(n, r)$ is upper bounded by $O(\Lambda(n) \log r)$.*
- (b) *$\Lambda_{\text{bxos}}(n)$ is upper bounded by $O(\Lambda(n))$.*

Proof. (a) We first consider $\Lambda_{\text{xos}}(n, r)$ and the case where f is an XOS (set) function. From Lemma 2.2, we only need to address the case where all elements are $\lambda = 1/(5\Lambda(n) \log r)$ -small, since $1 + \lambda^{-1} = O(\Lambda(n) \log r)$. We further assume that $\text{OPT} = \text{Adap}(\mathcal{T}, f) = 5\Lambda(n) \log r$ by scaling f , and \mathcal{T} is the optimal decision tree.

Let $f(S) = \max_{i \in [W]} \left\{ \sum_{j \in S} w_j^{(i)} \right\}$, where $w_j^{(i)} \geq 0$ for all $i \in [W], j \in U$. Since all elements are $1/(\Lambda(n) \log r)$ -small, $f(\{j\}) = \max_{i \in [W]} w_j^{(i)} < 1$ implies $w_j^{(i)} < 1$ for all i, j . We note that this XOS function f is the objective function of the stochastic probing problem, which is different from $f_{\text{xos}}(S)$ defined in Eq. (3.1).

For each leaf ℓ , let $I(\ell)$ be the index $i \in [W]$ such that the linear function $w^{(i)}$ attains the value $f(\text{elt}(A_\ell))$ at the set $\text{elt}(A_\ell)$, that is, $\arg \max_{i \in [W]} \left\{ \sum_{j \in \text{elt}(A_\ell)} w_j^{(i)} \right\}$ (breaking ties arbitrarily). Now, for each ℓ and integer

$1 \leq j \leq \log r + 1$, we define $A_{\ell,j} = \left\{ u : u \in A_\ell, 2^{-j} \leq w_{\text{elt}(u)}^{(I(\ell))} < 2^{-j+1} \right\}$. We define an additional $A_{\ell,0}$ to be $A_\ell \setminus \bigcup_{j=1}^{\lfloor \log r \rfloor + 1} A_{\ell,j}$. It is clear that for any node $u \in A_{\ell,0}$, $w_{\text{elt}(u)}^{(I(\ell))} \leq 1/r$.

Therefore, the reward for the adaptive strategy ALGTREE is

$$5\Lambda(n) \log r = \text{OPT} = \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(\text{elt}(A_\ell))] = \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\sum_{u \in A_\ell} w_{\text{elt}(u)}^{(I(\ell))} \right] \leq \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\sum_{j=1}^{\lfloor \log r \rfloor + 1} \frac{|A_{\ell,j}|}{2^{j-1}} + \frac{|A_{\ell,0}|}{r} \right]$$

Using the assumption on the maximal length of a sequence in \mathcal{F} , which implies that $|A_{\ell,0}| \leq |A_\ell| \leq r$, we obtain

$$5\Lambda(n) \log r \leq \sum_{j=1}^{\lfloor \log r \rfloor + 1} \frac{1}{2^{j-1}} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A_{\ell,j}|] + \frac{r}{r} \leq \sum_{j=1}^{\lfloor \log r \rfloor + 1} \frac{1}{2^{j-1}} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A_{\ell,j}|] + 1.$$

Thus, there exists $1 \leq j_0 \leq \log r + 1$ such that $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A_{\ell,j_0}|] / 2^{j_0-1} \geq 4\Lambda(n) \log r / (\log r + 1)$ by the averaging principle. Thus,

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A_{\ell,j_0}|] \geq \frac{4 \cdot 2^{j_0-1} \Lambda(n) \log r}{\log r + 1} \geq 2^{j_0} \Lambda(n) \geq \Lambda(n).$$

Consider the f_{xos} that is defined by the subsets $A'_\ell = A_{\ell,j_0}$ for each ℓ . Then by the lemma condition,

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \geq \frac{1}{\Lambda(n)} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|] = \frac{1}{\Lambda(n)} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A_{\ell,j_0}|] \geq 2^{j_0}.$$

Notice that for each set $S \subseteq U$, when $f_{\text{xos}}(S) = s$ for some nonnegative integer s , there is a leaf ℓ such that $|S \cap \text{elt}(A_{\ell,j_0})| \geq s$. Thus,

$$f(S) \geq \sum_{j \in S} w_j^{(I(\ell))} \geq \sum_{j \in S \cap \text{elt}(A_{\ell,j_0})} w_j^{(I(\ell))} = \sum_{u \in A_{\ell,j_0}, \text{elt}(u) \in S} w_{\text{elt}(u)}^{(I(\ell))} \geq \frac{|S \cap \text{elt}(A_{\ell,j_0})|}{2^{j_0}} = \frac{f_{\text{xos}}(S)}{2^{j_0}},$$

where in the last inequality we use $w_{\text{elt}(u)}^{(I(\ell))} \geq 2^{-j_0}$ when $u \in A_{\ell,j_0}$, $1 \leq j_0 \leq \log r + 1$. Therefore, combining the above two inequalities, the reward for the non-adaptive strategy ALGPATH is

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f(\text{elt}(R \cap P_\ell))] \right] \geq \frac{1}{2^{j_0}} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \geq 1.$$

Thus, the adaptivity gap is upper bounded by $O(\Lambda(n) \log r)$.

(b) Next, we consider the case when f is a binary-XOS (set) function. From Lemma 2.2, we only need to address the case where all elements are $\lambda = 1/\Lambda(n)$ -small, since $1 + \lambda^{-1} = O(\Lambda(n))$. We further assume that $\text{OPT} = \text{Adap}(\mathcal{T}, f)$, and \mathcal{T} is the optimal decision tree.

Let $f(S) = \max_{i \in [W]} \left\{ \sum_{j \in S} w_j^{(i)} \right\}$, where $w^{(i)} \in \{0, 1\}^U$ for all $i \in [W], j \in U$. Since all elements are $1/\Lambda(n)$ -small, $f(\{j\}) = \max_{i \in [W]} w_j^{(i)} < \frac{\text{OPT}}{\Lambda(n)}$ shows that, either f is the zero function (trivial), or $\text{OPT} > \Lambda(n)$.

Similarly, for each leaf ℓ , let $I(\ell)$ be the index $i \in [W]$ such that the linear function $w^{(i)}$ attains the value $f(\text{elt}(A_\ell))$ at the set $\text{elt}(A_\ell)$, that is, $\arg \max_{i \in [W]} \left\{ \sum_{j \in \text{elt}(A_\ell)} w_j^{(i)} \right\}$ (breaking ties arbitrarily). Then, we directly

set $A'_\ell = \{u \in A_\ell : w_{\text{elt}(u)}^{(I(\ell))} = 1\}$. Since different nodes in A_ℓ represent distinct elements in U ,

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|] = \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(\text{elt}(A_\ell))] = \text{OPT} > \Lambda(n).$$

Then, by the lemma condition, we obtain

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \geq \frac{1}{\Lambda(n)} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|] = \frac{\text{OPT}}{\Lambda(n)}.$$

However, we have the following lower bound of the reward of the path P_ℓ :

$$f(\text{elt}(R \cap P_\ell)) = \max_{i \in [W]} \left\{ \sum_{j \in \text{elt}(R \cap P_\ell)} w_j^{(i)} \right\} \geq \max_{\ell'} \left\{ \sum_{j \in \text{elt}(R \cap P_\ell)} w_j^{(I(\ell'))} \right\} = \max_{\ell'} \{ |\text{elt}(R \cap P_\ell) \cap \text{elt}(A'_{\ell'})| \},$$

where clearly the last term is exactly $f_{\text{xos}}(\text{elt}(R \cap P_\ell))$. Hence, we obtain $\text{Nadp}(\mathcal{T}, f)$ as

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f(\text{elt}(R \cap P_\ell))] \right] \geq \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \geq \frac{\text{OPT}}{\Lambda(n)},$$

which shows that $\Lambda_{\text{bxos}}(n) = O(\Lambda(n))$. \square

The remaining task is to show the premise of Lemma 3.2 holds. This is highly nontrivial, and we make it a theorem as follows, and prove it in the next two subsections.

Theorem 3.3. *Let A'_ℓ be a subset of A_ℓ for each leaf $\ell \in \mathcal{T}$. Assume that*

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|] \geq 200 \log n.$$

Then for $n \geq 100$,

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \geq \frac{1}{200 \log n} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|].$$

We can see that taking $\Lambda(n) = 200 \log n$ in Lemma 3.2, and Theorem 3.3 completes the proof of Theorem 3.1.

3.2 Main Algorithm to Categorize Outcomes (ℓ, R)

In this section, we present a novel *greedy* algorithm on the decision tree \mathcal{T} . We will use the definitions of **depth** in Definition 2.7 and **active ancestors** in Definition 2.8 introduced in Section 2.2 to further exploit the structure of the tree. To prove Theorem 3.3, the intuition is to show that for each ℓ, R , one can find a leaf ℓ_0 such that $|\text{elt}(R \cap P_\ell) \cap \text{elt}(A'_{\ell_0})|$ is large. As explained in Section 1.3.5, our goal is to design an algorithm with input (ℓ, R) and output a sequence $B(\ell, R)$ such that $f_{\text{xos}}(\text{elt}(R \cap P_\ell)) = |\text{elt}(R \cap P_\ell) \cap \text{elt}(A'_{\ell_0})| \geq |B(\ell, R)| - 1$, where the subtraction of 1 is only due to technical reasons. Here, the length $|B(\ell, R)|$ does not refer to the standard notion of length and will be defined explicitly later. Thus, we can relate $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))]]$ to $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} |B(\ell, R)| - 1$. Moreover, for a fixed output sequence $B(\ell, R) = B_0$ and a non-negative integer $s \geq 0$ such that $|B_0| \leq s / \log n$, the design of the algorithm ensures a small upper bound on the probability $\Pr_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [B(\ell, R) = B_0, |A'_\ell| = s]$.

In this section, we describe the core algorithm with a fixed parameter $K \in \mathbb{N}_+$ and establish its basic properties, including the proof that $f_{\text{xos}}(\text{elt}(R \cap P_\ell)) \geq |B(\ell, R)| - 1$. In Section 3.3, we provide an upper bound on the probability $\Pr_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [B(\ell, R) = B_0, |A'_\ell| = s]$, for each fixed B_0 and $s \geq 0$. Combining the two sections, we obtain the main theorem. In this section, we let $K = \Theta(\log n)$.

First, we formally define the set of possible outputs for (ℓ, R) . We define \mathfrak{B} as the set of sequences of the form $(m; s; d_1, \dots, d_m; e_1, \dots, e_m; y_1, \dots, y_{\lfloor \frac{s}{K} \rfloor})$, satisfying the following conditions:

- $0 \leq s \leq n, m \geq 0$;
- $n \geq d_1 > d_2 > \dots > d_m \geq 0, n \geq y_1 \geq y_2 \geq \dots \geq y_{\lfloor \frac{s}{K} \rfloor} \geq 0$;
- e_2, \dots, e_m are distinct elements in $U = [n]$, with $e_1 = 0$ serving as a placeholder.

We refer to such a sequence $B \in \mathfrak{B}$ as a **label**. In our context, m is called the *length* of the label (the notation $|B(\ell, R)|$ mentioned above denotes the length m of the label $B(\ell, R)$, analogous to the notation $|\cdot|$ used in Section 1.3.5). The main difference from Section 1.3.5 is that, since we consider general A'_ℓ , we need the depths d_i and an additional sequence y_j to control the depths of nodes in A'_ℓ . This information ensures that the cardinality of **impossible** nodes (as introduced in Section 1.3.5) can be lower bounded by a quantity related to m and s , see Lemma 3.6.

In the following, we introduce the notion of a *prefix* of a label in \mathfrak{B} . The *prefix* of length $1 \leq a \leq m$ of a label $B = (m; s; d_1, \dots, d_m; e_1, \dots, e_m; y_1, \dots, y_{\lfloor \frac{s}{K} \rfloor})$ is defined to be

$$\text{Pref}_a(B) = (a; s; d_1, \dots, d_a; e_1, \dots, e_a; y_1, \dots, y_{\lfloor \frac{s}{K} \rfloor}).$$

Additionally, the *prefix* of length 0 of any label is defined as (0) for uniformity. It is straightforward to verify that \mathfrak{B} is *prefix-closed* (a prefix of a label is also in \mathfrak{B}).

We first outline the ideas behind Definition 3.1 and Algorithm 1 (see below). Algorithm 1 serves as the main algorithm that, given an input pair (ℓ, R) , produces a label $B(\ell, R) \in \mathfrak{B}$ as output. Definition 3.1 provides the preparatory framework for Algorithm 1 by defining m sets of nodes L_1, L_2, \dots, L_m (called the *level sets*) for a given label B of length m .

Intuitively, the a -th level set L_a consists of the nodes that can potentially trigger the a -th update of the maintained sequence. Roughly speaking, a node u is in $L_a(B)$ if u is at depth d_a , $\text{elt}(u) = e_a$, and there exists a leaf ℓ (subject to additional conditions) such that $u \in A'_\ell$ and the unique node in A'_ℓ at depth d_i represents e_i for every $1 \leq i \leq a$. Moreover, if multiple leaves satisfy these conditions, one must be uniquely chosen to resolve ambiguity; our convention is to select the ‘leftmost’ leaf (solely for tie-breaking) and define the set $\mathcal{A}_u(B)$, which we call the *special-ancestor set* of u with respect to B , as the set of nodes in A'_ℓ with depths less than d_a for that particular ℓ (note that all such nodes are active ancestor of u).

Specifically, the notion of ‘leftmost’ is determined by the dictionary order of the root-to-leaf path in the binary tree \mathcal{T} , where at each internal node, No path is ordered to be larger than Yes path.

Then, based on the definition of the level set $L_a(B)$, Algorithm 1 implements the greedy algorithm which, on input (ℓ, R) , maintains a label B , progresses upward along P_ℓ , and at each node encountered u seeks a label B' of length $a + 1$ (where a denotes the current length of B) that satisfies $u \in R \cap L_{a+1}(B')$, then updates B to B' . Note that Algorithm 1 is essentially the same as the algorithm presented in Section 1.3.5, albeit with additional details.

Now, let us present the formal description of Definition 3.1, with an example illustrated in the first item of Fig. 5. Recall that in a single $A'_\ell \subseteq P_\ell$, different nodes have different depths and represent distinct elements.

Definition 3.1 (Level sets $L_a(B)$ and Special-Ancestor sets $\mathcal{A}_u(B)$).

Let $B = (m; s; d_1, \dots, d_m; e_1, \dots, e_m; y_1, \dots, y_{\lfloor \frac{s}{K} \rfloor}) \in \mathfrak{B}$ be a label with length $m \geq 1$. Given a positive integer $1 \leq a \leq m$, we define $L_a(B)$ and $\mathcal{A}_u(B)$ for $u \in L_a(B)$ as follows.

- (a) When $a = 1$, a node u belongs to $L_1(B)$ if and only if it is a leaf at depth d_1 , satisfies $|A'_u| = s$, and for each $1 \leq i \leq \lfloor \frac{s}{K} \rfloor$, the depth of the iK -th node from the bottom of A'_u (i.e., counted upward from the leaf toward the root) is equal to y_i . We define $\mathcal{A}_u(B)$ to be the set of nodes in A'_u that are at depths less than d_1 .
- (b) When $a \geq 2$, a node u belongs to $L_a(B)$ if and only if it is at depth d_a , satisfies $\text{elt}(u) = e_a$, and there exists a leaf $\ell \in L_1(B)$ such that the unique node in A'_ℓ at depth d_i represents e_i for all $2 \leq i \leq a$ (which implies $u \in A'_\ell$). Among all such leaves, we select the leftmost one (for tie-breaking), and define $\mathcal{A}_u(B)$ to be the set of nodes in A'_ℓ that lie at depths less than d_a .

In both cases, any node $v \in \mathcal{A}_u(B)$ is of depth less than d_a , the depth of u . Moreover, $v \in \mathcal{A}_u(B) \subseteq A'_\ell$ shows that v is an active ancestor of ℓ , thus on the path from root to ℓ , the Yes arc from v is taken. At the same time, when $a \geq 2$, $u \in A'_\ell$ must be a descendant of v (also true for $a = 1$). Therefore, we conclude:

Observation 3.1. *For any $1 \leq a \leq m$, the special ancestor set $\mathcal{A}_u(B)$ is a subset of active ancestors of u .*

The following observation is immediate from the definition, since membership of a node in $L_a(B)$ depends only on m, s, y_j (for $1 \leq j \leq \lfloor \frac{s}{K} \rfloor$) and d_i, e_i (for $i \leq a$); furthermore, the tie-breaking process depends solely on the tree structure.

Observation 3.2. *For any two labels $B_1, B_2 \in \mathfrak{B}$ of lengths m and m' , respectively, and for any $1 \leq a \leq \min\{m, m'\}$, the level sets $L_a(B_1) = L_a(B_2)$, provided that $\text{Pref}_a(B_1) = \text{Pref}_a(B_2)$. Moreover, for any $u \in L_a(B_1) = L_a(B_2)$, the special ancestor sets $\mathcal{A}_u(B_1) = \mathcal{A}_u(B_2)$.*

The following observation will be used in the description of Algorithm 1.

Observation 3.3. For any two distinct labels $B_1, B_2 \in \mathfrak{B}$, both of length $m \geq 1$, if $\text{Pref}_{m-1}(B_1) = \text{Pref}_{m-1}(B_2)$, then $L_m(B_1) \cap L_m(B_2) = \emptyset$.

Proof. Assume, for contradiction, that $u \in L_m(B_1) \cap L_m(B_2)$. When $m = 1$, the contradiction is immediate since any node $u \in L_1(B)$, with $B = (m; s; d_i; e_i; y_j)$, must be a leaf with $s = |A'_u|$, the quantity y_i being the depth of the last iK -th node in A'_u , and d_1 the depth of u . Moreover, by definition, e_1 is fixed as 0, hence the label B is uniquely determined by u , which gives $B_1 = B_2$.

For $m \geq 2$, Definition 3.1(b) implies that if $u \in L_m(B)$ for $B = (m; s; d_i; e_i; y_j)$, then d_m must be equal to the depth of u , and $e_m = \text{elt}(u)$. Since $\text{Pref}_{m-1}(B_1) = \text{Pref}_{m-1}(B_2)$, it follows that $B_1 = B_2$, contradicting the assumption. Thus, the result is established. \square

We now proceed to present a detailed description of Algorithm 1. The goal of the algorithm is to classify each pair (ℓ, R) using appropriate labels from the set \mathfrak{B} . Initially, the algorithm sets the iterator u to the leaf ℓ and initializes the maintained label B to (0) , with the length variable a set to 0.

The core functionality of the algorithm is implemented through a while loop. Within each iteration, the algorithm examines whether there exists a label $B_1 \in \mathfrak{B}$ that satisfies three conditions: it has length $a + 1$, its prefix matches the current label (i.e., $B = \text{Pref}_a(B_1)$), and the current node belongs to the corresponding level set (i.e., $u \in L_{a+1}(B_1)$).

When such a label exists and the additional condition that either $u \in R$ or u is the starting leaf is met, the algorithm identifies the unique label B_1 (whose uniqueness is guaranteed by Observation 3.3). It then updates B to B_1 , increments a by 1 and advances u to its parent node in the tree. If then u becomes the root of \mathcal{T} , then the loop terminates. Upon termination, the algorithm returns the final label B .

It should be noted that Algorithm 1 is essentially identical to the algorithm presented in Section 1.3.5, with the interpretation that nodes in the level sets $L_{a+1}(B_1)$ represent those capable of extending the length of the maintained label B .

The pseudocode of the algorithm is defined as follows:

Algorithm 1: Categorizing (ℓ, R)

Input: (ℓ, R)
Output: $B = (m; s; d_1; \dots, d_m; e_1, \dots, e_m; y_1, \dots, y_{\lfloor \frac{m}{K} \rfloor}) \in \mathfrak{B}$

- 1 $u \leftarrow \ell; B \leftarrow (0); a \leftarrow 0$ \triangleright Initialization of output
- 2 **while** the depth of $u \geq 0$ **do**
- 3 **if** $\exists B_1 \in \mathfrak{B}$ of length $a + 1$, s.t. $B = \text{Pref}_a(B_1)$, $u \in L_{a+1}(B_1)$ **then**
- 4 **if** $u \in R$ **or** $u = \ell$ **then**
- 5 **find** such unique B_1 ; \triangleright Uniqueness by Observation 3.3
- 6 $B \leftarrow B_1; a \leftarrow a + 1$
- 7 $u \leftarrow u.\text{parent}$; \triangleright Climbing up the path P_ℓ . If u is already the root of \mathcal{T} , jump out while loop
- 8 **return** B ;

By the algorithm, it is clear that any (ℓ, R) is given a unique label. As mentioned above, we denote this label by $B(\ell, R)$. We also need certain entries of the label. To stress the dependency of the input (ℓ, R) , the length of the label is denoted as $m(\ell, R)$, the second entry as $s(\ell, R)$, etc.

Before proceeding, we need to examine further the properties of the output label B produced by Algorithm 1. It is also worth mentioning that $u = \ell$ always updates the label B to length 1, where d_1 is the depth of ℓ , $e_1 = 0$, each y_j is the depth of the jK -th node in A'_ℓ (counted from ℓ), and $s = |A'_\ell|$. These statistics never change after such initialization. Moreover, one can observe the *greedy* nature of this algorithm. Specifically, when traversing upward along P_ℓ , the algorithm always processes lower nodes before higher ones. When conditions Line 3 and Line 4 are satisfied, the maintained label B is irrevocably updated. Consequently, when we consider a fixed label B_0 , the fact that an outcome (ℓ, R) satisfies $B(\ell, R) = B_0$ already imposes specific constraints on R . This concept is illustrated in Fig. 5. The following lemma formalizes these properties:

Lemma 3.4. For each (ℓ, R) pair, assume $B(\ell, R) = B_0 = (m; s; d_i; e_i; y_j) \in \mathfrak{B}$. Then the following holds for each $1 \leq a \leq m$:

- (a) $\ell \in L_1(B_0)$ and it updates B from length 0 to 1, hence the length m is always at least 1.
- (b) For $a \geq 2$, there exists a unique node v_a of depth d_a in P_ℓ , $\text{elt}(v_a) = e_a$, such that $v_a \in R \cap L_a(B_0)$ and the label B is updated from length $(a - 1)$ to a at node v_a .
- (c) There does not exist any node $z \in R \cap P_\ell$ satisfying the two conditions:
- (1) z is a descendant of v_{a+1} (ignore this condition when $a = m$);
 - (2) $\exists B_1 \in \mathfrak{B}$ of length $a + 1$ such that $\text{Pref}_a(B_0) = \text{Pref}_a(B_1)$ and $z \in L_{a+1}(B_1)$.

In particular, condition $z \in P_\ell \cap \mathcal{A}_v(B_0)$ for some $v \in L_a(B_0)$ is **sufficient** to ensure condition (2).

Proof. (a) For any leaf ℓ , we define a label $B_1 = (1; s; d_1; e_1; y_j)_{1 \leq j \leq \lfloor \frac{s}{K} \rfloor}$, where d_1 is the depth of ℓ , $e_1 = 0$, each y_j is the depth of the last jK -th node in A'_ℓ , and $s = |A'_\ell|$. We can verify that $B_1 \in \mathfrak{B}$, as $0 \leq d_1 \leq n$, $e_1 = 0$, and $m = 1 \geq 0$. It follows directly from Definition 3.1 that $\ell \in L_1(B_1)$. Therefore, when $u = \ell$, the conditions in Line 3 are satisfied, causing B to be updated by B_1 from length 0 to 1. It is evident from the algorithm that the length of B never decreases during updates. Therefore, $\ell \in L_1(B_1) = L_1(B_0)$ by Observation 3.2, since $B_1 = \text{Pref}_1(B_0)$.

- (b) From part (a), we know that the first update of B always occurs at the leaf ℓ . We observe from Line 3 that the transition of B from length $(a - 1)$ to length a corresponds to a label $B_1 \in \mathfrak{B}$ of length a , where $u \in R \cap L_a(B_1)$ when $a \geq 2$. Let us denote this node u as v_a . Since the length of B only increases throughout the algorithm, we have $B_1 = \text{Pref}_a(B_0)$, which by Observation 3.2 implies $L_a(B_1) = L_a(B_0)$. According to Definition 3.1, if $v_a \in L_a(B_1) = L_a(B_0)$, then the depth of v_a equals d_a , and $\text{elt}(v_a) = e_a$, matching the assertion. The uniqueness of v_a follows from the fact that nodes in P_ℓ represent distinct elements. Furthermore, even for different outcomes R_0 and R_1 such that $B(\ell, R_0) = B_0 = B(\ell, R_1)$, the corresponding nodes v_a remain identical for this same reason.

- (c) Let us assume, for contradiction, that there exists a node z satisfying the two conditions. Then, assume $B_1 = (a + 1; s; d_1, \dots, d_a, d'_{a+1}; e_i; y_j)$. Then, $z \in L_{a+1}(B_1)$ shows the depth of z is d'_{a+1} by Definition 3.1, which should be strictly less than d_a since $B_1 \in \mathfrak{B}$. When examining the execution of Algorithm 1, we find a contradiction as follows. Between the updates of B from length $a - 1$ to a at node v_a , and then from length a to $a + 1$ at node v_{a+1} (or maintaining length m till termination if $a = m$), no intermediate node should trigger an update. However, the node z would appear in the iterator u after processing v_a ($d'_{a+1} < d_a$) but before reaching v_{a+1} (it is a descendant of v_{a+1}). At this point, B equals $\text{Pref}_a(B_0)$. $z \in R$ matches the conditions for updating B in Line 3, forcing an update of B , which gives a contradiction. Therefore, no such node z can exist.

Now we show that $z \in \mathcal{A}_v(B_0)$ for some $v \in L_a(B_0)$ suffices to ensure condition (2). We define d'_{a+1} as the depth of z , $v \in L_a(B_0)$ and $e'_{a+1} = \text{elt}(z)$. Then we define:

$$B_1 = (a + 1; s; d_1 \dots, d_a, d'_{a+1}; e_1, \dots, e_a, e'_{a+1}; y_1, \dots, y_{\lfloor \frac{s}{K} \rfloor}).$$

We now verify that $B_1 \in \mathfrak{B}$. By Observation 3.1, the special ancestor set $\mathcal{A}_v(B_0)$ only contains nodes of depth less than d_a . Thus, $0 \leq d'_{a+1} < d_a$. Moreover, from part (b), the nodes $v_i \in P_\ell$ ensure that $\text{elt}(v_i) = e_i$ differ from $\text{elt}(z) = e'_{a+1}$, for $2 \leq i \leq a$, as they appear in a same P_ℓ . Thus, $B_1 \in \mathfrak{B}$. Given that $z \in \mathcal{A}_v(B_0)$, there must exist a leaf $\ell \in L_1(B_0) = L_1(B_1)$ (since $\text{Pref}_a(B_0) = \text{Pref}_a(B_1)$), such that the unique node at depth d_i in A'_ℓ represents e_i for $2 \leq i \leq a$, and $z \in \mathcal{A}_v(B_0) \subseteq A'_\ell$. Since the depth of z is d'_{a+1} and $\text{elt}(z) = e'_{a+1}$, the leaf ℓ satisfies the conditions in Definition 3.1 that ensures $z \in L_{a+1}(B_1)$. Hence, condition (2) is satisfied since $\text{Pref}_a(B_0) = \text{Pref}_a(B_1)$. □

From the above lemma, we can relate $f_{\text{xos}}(\text{elt}(R \cap P_\ell))$, $|A'_\ell|$ with the entries of $B(\ell, R)$.

Lemma 3.5. *For any (ℓ, R) , we have $f_{\text{xos}}(\text{elt}(R \cap P_\ell)) \geq m(\ell, R) - 1$, $|A'_\ell| = s(\ell, R)$.*

Proof. Assume that the label of (ℓ, R) is $B_0 = (m; s; d_1, \dots, d_m; e_1, \dots, e_m; y_1, \dots, y_{\lfloor \frac{s}{K} \rfloor})$, and $m \geq 2$. Then, by Lemma 3.4 (b), nodes v_a are of depth d_a and $v_a \in R \cap P_\ell \cap L_a(B_0)$ for any $2 \leq a \leq m$. From Definition 3.1 (b), $v_a \in L_a(B_0)$ implies $\text{elt}(v_a) = e_a$. In particular, $v_m \in L_m(B_0)$ shows that there exists a leaf $\ell_0 \in L_1(B_0)$




Figure 5: In the decision tree, red edges indicate Yes arcs, and blue edges indicate No arcs. Nodes with the same depth are connected using dashed lines, and their depth is labeled on them. Nodes higher than v_4 and in the Yes subtree of v_4 are not drawn. Let us assume that K is large and $|A'_\ell| = 4K$ and the depth of the K -th, $2K$ -th, $3K$ -th, $4K$ -th nodes of A'_ℓ (counted from ℓ) have depths $4K + 10, 3K + 10, 2K + 10, K + 10$ for all leaves $\ell = \ell_0, \ell', \ell'', \ell'''$. Moreover, we assume $v_2 \in A'_{\ell_0}, v_2' \in A'_{\ell'}, v_2'' \in A'_{\ell''}, v_2''' \in A'_{\ell'''}, v_3 \in A'_{\ell'}, A'_{\ell''}$, but $v_3 \notin A'_{\ell_0}, A'_{\ell'''}$. Regarding elements, $\text{elt}(v_2) = \text{elt}(v_2') = \text{elt}(v_2'') = \text{elt}(v_2''')$. Consider the label $B_0 = (4; 4K; 6K, 6K - 1, 6K - 3, 6K - 4; 0, \text{elt}(v_2), \text{elt}(v_3), \text{elt}(v_4); 4K + 10, 3K + 10, 2K + 10, K + 10)$.

- All leaves $\ell_0, \ell', \ell'', \ell'''$ are in $L_1(B_0)$, and all v_2, v_2', v_2'', v_2''' are in $L_2(B_0)$. v_3 is in $L_3(B_0)$, but note that the satisfying ℓ in the definition can only be ℓ' or ℓ'' , and hence $\mathcal{A}_{v_3}(B_0)$ is defined to be the nodes of depth less than $6K - 3$ in $A'_{\ell'}$ (as ℓ' is on the ‘left’ of ℓ''').
- Suppose that the specific outcome R_0 includes all green nodes, but does not include white nodes and the nodes above v_4 . The algorithm with input (ℓ_0, R_0) updates B at ℓ_0, v_2, v_3, v_4 , and hence $B(\ell_0, R_0) = B_0$ (note that $v_3 \in L_3(B_0)$ can trigger an update, even though $v_3 \notin A'_{\ell_0}$).
- Note that u is a descendant of $v_3 = v_3(\ell_0)$, and $u \in \mathcal{A}_{v_2'}(B_0)$ for $v_2' \in L_2(B_0)$. These facts imply the existence of the label $B_1 = (3; 4K; 6K, 6K - 1, 6K - 2; 0, \text{elt}(v_2), \text{elt}(u); 4K + 10, 3K + 10, 2K + 10, K + 10)$ that $\text{Pref}_2(B_1) = \text{Pref}_2(B_0)$ by Lemma 3.4 (c). This means that $u \in \text{Imp}(\ell_0, B_0)$. Intuitively $u \notin R_0$, since otherwise, it can update B at u before v_3 .

such that the unique node of depth d_i in A'_{ℓ_0} represents e_i , for any $2 \leq i \leq m$. This already means that $e_a \in \text{elt}(R \cap P_\ell) \cap \text{elt}(A'_{\ell_0}), \forall 2 \leq a \leq m$. Hence, the conclusion $f_{\text{xos}}(\text{elt}(R \cap P_\ell)) = \max_{\ell_1} |\text{elt}(A'_{\ell_1}) \cap \text{elt}(R \cap P_\ell)| \geq |\text{elt}(A'_{\ell_0}) \cap \text{elt}(R \cap P_\ell)| \geq m - 1$ follows. For the second assertion, Lemma 3.4 (a) implies that $\ell \in L_1(B_0)$ and $|A'_\ell| = s = s(\ell, R)$. \square

We now proceed to derive consequences of Lemma 3.4. We first define the subset $S(B_0)$ of leaves as in Section 1.3.5.

Definition 3.2 (Generating Set $S(B_0)$ of B_0 and the a -th Triggering Node $v_a(\ell)$).

For a label $B_0 = (m, s, d_i, e_i, y_j) \in \mathfrak{B}$, we define the generating set of B_0 , denoted as $S(B_0)$, as the set of leaves ℓ for which there exists some outcome R such that $B(\ell, R) = B_0$, i.e.,

$$S(B_0) = \{\ell \text{ is a leaf of } \mathcal{T} : \exists R \text{ such that } B(\ell, R) = B_0\}.$$

For any leaf $\ell \in S(B_0)$, Lemma 3.4 (b) ensures the existence of a unique node v_a of depth d_a , $\text{elt}(v_a) = e_a$, such that $v_a \in P_\ell \cap L_a(B_0)$ for each $2 \leq a \leq m$. v_a triggers the a -th update of the label, and we call

it the a -th triggering node of ℓ . We make the dependency on ℓ explicit, and denote it as $v_a(\ell)$, but omit its dependency on B_0 , since we fix B_0 in the rest of this section. For notational convenience, we define $v_1(\ell) := \ell \in L_1(B_0)$. Note that the definitions of $S(B_0)$ and $v_a(\ell)$ do **not** depend on the specific choice of R .

Part (c) of Lemma 3.4 tells us that when $B(\ell, R) = B_0$, there are constraints on the active nodes (in ALGPATH) along P_ℓ . In detail, there does not exist a node in $R \cap P_\ell$ that satisfies the following two conditions: it is a descendant of $v_{a+1}(\ell)$; it is in $L_{a+1}(B_1)$ for some $B_1 \in \mathfrak{B}$ of length $a + 1$ such that $\text{Pref}_a(B_0) = \text{Pref}_a(B_1)$. Therefore, for each $\ell \in S(B_0)$, we define the subset of P_ℓ that cannot be active (i.e. $\notin R$) as follows (the condition on $v_{m+1}(\ell)$ when $a = m$ is ignored):

$$\text{Imp}(\ell, B_0) := \bigcup_{a=1}^m \{u \in P_\ell \cap L_{a+1}(B_1) : B_1 \in \mathfrak{B}, \text{Pref}_a(B_1) = \text{Pref}_a(B_0), u \text{ is a descendant of } v_{a+1}(\ell)\}. \quad (3.2)$$

Notice that $\text{Imp}(\ell, B_0)$ is defined with respect to only ℓ and B_0 , and does not depend on R , since after fixing $\ell \in S(B_0)$ and B_0 , the sets $L_a(B_0)$, $\mathcal{A}_v(B_0)$ and the nodes $v_a(\ell)$ are the same for any R that satisfies $B(\ell, R) = B_0$ as mentioned in Lemma 3.4 (b).

As discussed in the beginning of this section, we can view nodes in $L_a(B_0)$ as nodes that can potentially trigger an update in Section 1.3.5. Therefore, the set $\text{Imp}(\ell, B_0)$ defined here corresponds to the notion of **impossible** nodes defined there. In the following lemma, we investigate the properties of $\text{Imp}(\ell, B_0)$ and provide a lower bound on the cardinality of $\text{Imp}(\ell, B_0) \cap A_\ell$, as we did in Section 1.3.5.

Lemma 3.6. *For a fixed label $B_0 = (m; s; d_i; e_i; y_j) \in \mathfrak{B}$, we define $\text{Imp}(\ell, B_0)$ for $\ell \in S(B_0)$ as above. Then $\text{Imp}(\ell, B_0)$ does not contain any leaf of the original tree \mathcal{T} . Moreover, $|\text{Imp}(\ell, B_0) \cap A_\ell| \geq s - mK$.*

Proof. First, for any $u \in \text{Imp}(\ell, B_0)$ in the a -th union, the depth of $u \in P_\ell \cap L_{a+1}(B_1)$ is the $(a + 1)$ -th entry of the depth sequence of B_1 , which is strictly smaller than its a -th entry d_a . Thus, u is an ancestor of $v_a(\ell) \neq u$ (they are both in P_ℓ), hence not a leaf of \mathcal{T} .

Second, we consider the following subset of $\text{Imp}(\ell, B_0)$, where $v_a(\ell)$ is defined for $\ell \in S(B_0)$ in Definition 3.2 (in the following, we abbreviate $v_a(\ell)$ to v_a , and the special-ancestor set of v_a , i.e., $\mathcal{A}_{v_a}(B_0)$, to \mathcal{A}_{v_a} , since the leaf and label are fixed):

$$\text{Imp} = \bigcup_{a=1}^m \{u \in \mathcal{A}_{v_a} : \text{depth of } u \in (d_{a+1}, d_a)\}.$$

We first prove $\text{Imp} \subseteq A_\ell$. By Observation 3.1, any node $u \in \mathcal{A}_{v_a}$ is an active ancestor of $v_a \in P_\ell$. In particular, $u \in P_\ell$ and on the path from root to ℓ , the Yes arc from u is taken, and hence $u \in A_\ell$. Now, we take $v = v_a \in L_a(B_0)$ in the last sentence of Lemma 3.4 (c) shows that $\text{Imp} \subseteq \text{Imp}(\ell, B_0)$ (note that $v_{a+1}, u \in P_\ell$, so the depth of $u > d_{a+1}$ implies that it is a descendant of v_{a+1}).

Next, we lower bound $|\text{Imp}(\ell, B_0) \cap A_\ell|$ using $|\text{Imp}|$. By Lemma 3.4 (a), $\ell \in S(B_0)$ shows $v_1 := \ell \in L_1(B_0)$, hence d_1 is the depth of ℓ . Moreover, d_a is strictly decreasing since $B_0 \in \mathfrak{B}$. For any $1 \leq a \leq m$, we consider the node $v_a \in L_a(B_0)$. Moreover, we denote the ‘leftmost’ leaf for $v_a \in L_a(B_0)$ in Definition 3.1 as $\ell_a \in L_1(B_0)$. Therefore, the depth of the last jK -th node in $A'_{\ell_a} \supseteq \mathcal{A}_{v_a}$ is y_j , for $1 \leq j \leq \lfloor \frac{s}{K} \rfloor$.

Now, all $\lfloor \frac{s}{K} \rfloor$ indices from $1 \leq j \leq \lfloor \frac{s}{K} \rfloor$ of y_j divide the interval $[y_{\lfloor \frac{s}{K} \rfloor}, d_1)$ into $\lfloor \frac{s}{K} \rfloor$ intervals $[y_j, y_{j-1})$ (denote $y_0 = d_1$). However, there are at most $m - 1$ indices of d_i , $2 \leq i \leq m$ in $[y_{\lfloor \frac{s}{K} \rfloor}, d_1)$. Therefore, there are at least $\lfloor \frac{s}{K} \rfloor - m + 1$ indices $1 \leq j \leq \lfloor \frac{s}{K} \rfloor$, such that the interval $[y_j, y_{j-1})$ contains no d_a , for any $2 \leq a \leq m$. See Figure 6 for an example. We define $I(B_0)$ as the union of such intervals $[y_j, y_{j-1})$. For each of such intervals, there must be some interval (d_{a+1}, d_a) containing the entire interval $[y_j, y_{j-1})$, $1 \leq a \leq m$, as $(d_{m+1} := -1, d_1) \supseteq [y_{\lfloor \frac{s}{K} \rfloor}, d_1)$. Now, the cardinality of nodes in \mathcal{A}_{v_a} of depth in $[y_j, y_{j-1})$ is exactly K , since the depth of the jK -th, $(j - 1)K$ -th node in A'_{ℓ_a} (counted from the bottom ℓ_a) is y_j, y_{j-1} , and \mathcal{A}_{v_a} is defined to be all nodes in A'_{ℓ_a} with depth less than $d_a \geq y_{j-1}$. All such K nodes are entirely contained in the subset $\{u \in \mathcal{A}_{v_a} : \text{depth of } u \in (d_{a+1}, d_a)\}$. Summing up over all intervals in $I(B_0)$ (all intervals $[y_j, y_{j-1})$ are disjoint), we obtain $|\text{Imp}(\ell, B_0) \cap A_\ell| \geq |\text{Imp}| \geq K(\lfloor \frac{s}{K} \rfloor - m + 1) \geq s - mK$. \square




Figure 6: $[y_2, y_1)$ and $[y_5, y_4)$ do not contain any d_i , hence contained in (d_3, d_2) and $(-1, d_4)$ respectively. \mathcal{A}_{v_2} is defined to be nodes in A'_{ℓ_2} with depth less than $d_2 \geq y_1$, hence contain all K nodes at depth $[y_2, y_1)$. Similarly, A'_{ℓ_4} contain all K nodes at depth $[y_5, y_4)$.

Remark 3.7. In the proof, we have defined $I(B_0)$ as the union of intervals $[y_j, y_{j-1})$ for $1 \leq j \leq \lfloor \frac{s}{K} \rfloor$ such that no d_a ($2 \leq a \leq m$) is in the interval. Then the nodes in $\text{Imp}(\ell, B_0) \cap A_\ell$ whose depths are in $I(B_0)$ are at least $|I(B_0)| \geq s - mK$. This fact is not immediately used in this section, but will be used in Section 4.

We define the subset $T(B_0)$ of nodes in \mathcal{T} as $T(B_0) = \bigcup_{\ell \in S(B_0)} \text{Imp}(\ell, B_0)$. The following lemma captures the constraints on the nodes P_ℓ to be in R . This result is similar to the one mentioned in Section 1.3.5: if $B(\ell, R) = B_0$, then any **impossible** node above ℓ can not be in R .

Lemma 3.8. Fix a label $B_0 \in \mathfrak{B}$. For any $\ell \in S(B_0), R$ where $B(\ell, R) = B_0$, we have $R \cap P_\ell \cap T(B_0) = \emptyset$.

Proof. Assume that $B(\ell, R) = B_0$. Also, assume that there is a $u \in T(B_0)$ such that $u \in R \cap P_\ell$. By the definition of $T(B_0)$, $u \in \text{Imp}(\ell', B_0)$ for some $\ell' \in S(B_0)$. Expanding further Definition 3.2, there exists $1 \leq a \leq m$, $u \in P_{\ell'}$ is a descendant of $v_{a+1}(\ell')$ and $u \in L_{a+1}(B_1)$ for some $B_1 \in \mathfrak{B}$, $\text{Pref}_a(B_1) = \text{Pref}_a(B_0)$. To derive a contradiction, we show that u is also a descendant $v_{a+1}(\ell)$. Thus, we arrive at a contradiction to Lemma 3.4 (c) via taking $z = u$. Note that the result holds trivially when $a = m$, where conditions for $v_{m+1}(\ell)$ and $v_{m+1}(\ell')$ are ignored.

Hence, we will now assume by contradiction that $a < m$, and u is not a descendant of $v_{a+1}(\ell)$. Since $u, v_{a+1}(\ell) \in P_\ell$, u must be an ancestor of $v_{a+1}(\ell)$ (including $v_{a+1}(\ell)$ itself). However, this implies that $v_{a+1}(\ell)$ is a descendant of u , and also a descendant of $v_{a+1}(\ell')$. Therefore, $v_{a+1}(\ell') \in P_\ell$, which implies $v_{a+1}(\ell) = v_{a+1}(\ell')$, because the two nodes represent the same element e_{a+1} in P_ℓ . This is a contradiction. Then, the conclusion follows. \square

3.3 Proof for Theorem 3.3

In this section, we prove Theorem 3.3. Recall that we need to lower bound $\mathbb{E}[f_{\text{xos}}(\text{elt}(R \cap P_\ell))]$ for the f_{xos} defined in the beginning of this section. We set $K = \lceil 25 \log n \rceil + 25$ in this subsection.

First, we need to prove the following lemma:

Lemma 3.9. Let S be a subset of leaves in the tree \mathcal{T} . There is a subset T of nodes in \mathcal{T} , not containing any leaf of \mathcal{T} . There exists an integer $h \in \mathbb{N}$, such that for each $\ell \in S$, $|A_\ell \cap T| \geq h$. Then the following inequality holds:

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbf{1}_{\ell \in S} \Pr_{R \sim \mathbb{P}_R} [R \cap P_\ell \cap T = \emptyset] \right] \leq 2^{-h}.$$

Here, $\mathbf{1}_E$ denotes the indicator function of the event E .

Proof. The result is trivial for $h = 0$, as the right-hand side is 1. Therefore, in the following, we may assume that $h \geq 1$. We use induction on the height (the length of the longest root-leaf path, not the depth) of \mathcal{T} to prove the desired statement.

Assume now that the claim holds for trees of smaller height. For the current tree \mathcal{T} , let \mathcal{T}_1 and \mathcal{T}_2 denote the subtrees corresponding to the Yes and No arcs from the root r_0 , respectively. Let λ be the probability that the root is active (if r_0 has only one child, we set $\lambda = 0$ or 1 accordingly). Then, the path distribution $\pi_{\mathcal{T}}$ assigns probability λ to the Yes subtree \mathcal{T}_1 and $1 - \lambda$ to the No subtree \mathcal{T}_2 , preserving the distributions $\pi_{\mathcal{T}_1}$ and $\pi_{\mathcal{T}_2}$.

Now, we treat the following two cases separately. We denote

$$E_1 := \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_1}} \left[\mathbf{1}_{\ell \in S} \Pr_{R \sim \mathbb{P}_R} [R \cap P_\ell \cap T = \emptyset] \right], E_2 := \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_2}} \left[\mathbf{1}_{\ell \in S} \Pr_{R \sim \mathbb{P}_R} [R \cap P_\ell \cap T = \emptyset] \right].$$

Case 1. $r_0 \in T$.

In this case, we see that for leaves ℓ in $S \cap \mathcal{T}_1$, there are at least $h - 1$ active ancestors of it inside $T \cap \mathcal{T}_1$. Moreover, we denote the corresponding root-leaf path in \mathcal{T}_1 or \mathcal{T}_2 as $P'_\ell = P_\ell \setminus \{r_0\}$, then the corresponding relation $\Pr_{R \sim \mathbb{P}_R} [R \cap P_\ell \cap T = \emptyset] = (1 - \lambda) \Pr_{R \sim \mathbb{P}_R} [R \cap P'_\ell \cap T = \emptyset]$ holds, as the root r_0 has an independent probability of $1 - \lambda$ to be active. Therefore, by the induction hypothesis for \mathcal{T}_1 ,

$$E_1 = (1 - \lambda) \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_1}} \left[\mathbf{1}_{\ell \in S} \Pr_{R \sim \mathbb{P}_R} [R \cap P'_\ell \cap T = \emptyset] \right] \leq (1 - \lambda) 2^{-h+1}.$$

Similarly, for the **No** subtree \mathcal{T}_2 , each leaf has at least h active ancestors inside $T \cap \mathcal{T}_2$, since the root is not an active ancestor for leaves in the **No** subtree. Hence,

$$E_2 = (1 - \lambda) \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_2}} \left[\mathbf{1}_{\ell \in S} \Pr_{R \sim \mathbb{P}_R} [R \cap P'_\ell \cap T = \emptyset] \right] \leq (1 - \lambda) 2^{-h}.$$

Putting together,

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbf{1}_{\ell \in S} \Pr_{R \sim \mathbb{P}_R} [R \cap P_\ell \cap T = \emptyset] \right] = \lambda E_1 + (1 - \lambda) E_2 \leq 2^{-h} (2\lambda(1 - \lambda) + (1 - \lambda)^2) \leq 2^{-h}.$$

Case 2. $r_0 \notin T$.

In this case, we see that for leaves ℓ in both subtrees $S \cap \mathcal{T}_1$ and $S \cap \mathcal{T}_2$, there are at least h active ancestors of it inside $T \cap \mathcal{T}_1$ and $T \cap \mathcal{T}_2$, since $r_0 \notin T$. Moreover, we denote the corresponding root-leaf path in \mathcal{T}_1 or \mathcal{T}_2 as $P'_\ell = P_\ell \setminus \{r_0\}$, then the corresponding relation $\Pr_{R \sim \mathbb{P}_R} [R \cap P_\ell \cap T = \emptyset] = \Pr_{R \sim \mathbb{P}_R} [R \cap P'_\ell \cap T = \emptyset]$ holds, as the root $r_0 \notin T$. Therefore, by the induction hypothesis on \mathcal{T}_1 and \mathcal{T}_2 , we obtain $E_1, E_2 \leq 2^{-h}$. Putting together, $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [\mathbf{1}_{\ell \in S} \Pr_{R \sim \mathbb{P}_R} [R \cap P_\ell \cap T = \emptyset]] = \lambda E_1 + (1 - \lambda) E_2 \leq 2^{-h}$.

In conclusion, the result holds for all binary trees \mathcal{T} . \square

Lemma 3.10. For a fixed label $B_0 = (m; s; d_i; e_i; y_j) \in \mathfrak{B}$, the probability that the label of a random leaf and outcome (ℓ, R) is B_0 is at most 2^{mK-s} , i.e.,

$$\Pr_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [B(\ell, R) = B_0] \leq 2^{mK-s}.$$

Proof. The result is trivial when $s \leq mK$, so we assume $s > mK$. Define $S = S(B_0)$ as in Definition 3.2 and $T = T(B_0)$ as in the discussion preceding Lemma 3.8. Recall that $S(B_0) = \{\ell \text{ is a leaf of } \mathcal{T} : \exists R \text{ such that } B(\ell, R) = B_0\}$, and $T(B_0)$ is the union of all $\text{Imp}(\ell, B_0)$ for $\ell \in S(B_0)$.

It is clear from Lemma 3.6 that T does not contain any leaf of \mathcal{T} . Moreover, for any $\ell \in S$, we have $|\text{Imp}(\ell, B_0) \cap A_\ell| \geq h := s - mK > 0$. Hence, by Lemma 3.9,

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbf{1}_{\ell \in S} \Pr_{R \sim \mathbb{P}_R} [R \cap P_\ell \cap T = \emptyset] \right] \leq 2^{-h} = 2^{mK-s}.$$

By Lemma 3.8 and the definition of S , for any (ℓ, R) , the condition $B(\ell, R) = B_0$ implies that $R \cap P_\ell \cap T = \emptyset$ and $\ell \in S$. As a result,

$$\Pr_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [B(\ell, R) = B_0] \leq \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbf{1}_{\ell \in S} \Pr_{R \sim \mathbb{P}_R} [R \cap P_\ell \cap T = \emptyset] \right] \leq 2^{mK-s}.$$

\square

Now we are ready to prove Theorem 3.3 (formally stated in Section 3.1). Recall that we aim to prove the following:

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \geq \frac{1}{200 \log n} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|].$$

Here, f_{xos} is defined as

$$f_{\text{xos}}(S) = \max_{\ell} \{|\text{elt}(A'_\ell) \cap S|\},$$

where A'_ℓ is a subset of the active ancestors of ℓ , and $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|] \geq 200 \log n$.

Proof of Theorem 3.3. Before proceeding, we first give an upper bound on the number of labels in \mathfrak{B} with $s \geq 2mK$. For a fixed $1 \leq s \leq n$, there are at most $s/2K \leq n$ choices for $1 \leq m \leq s/2K$, and for each m , there are at most $(n+1)^m \times n^m \times (n+1)^{s/K} \leq (2n)^{2s/K}$ possible combinations of d_i, e_i, y_j .

From Lemma 3.5, we can express the term $\text{Adap} := \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_{\ell}|]$ as $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [\mathbb{E}_{R \sim \mathbb{P}_R} [s(\ell, R)]]$. Since for any outcome of $R \cap P_{\ell}$, the label is unique,

$$s(\ell, R) = \sum_{B_0 \in \mathfrak{B}} s(B_0) \cdot \mathbf{1}_{B(\ell, R) = B_0}.$$

Plugging in this formula and exchanging the order of summation, we obtain

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_{\ell}|] = \sum_{B_0 \in \mathfrak{B}} s(B_0) \cdot \Pr_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [B(\ell, R) = B_0].$$

We use $(m; s; *)$ to denote a label of the form $(m; s; d_i; e_i; y_j)$. Now we split the sum over $B_0 = (m; s; *) \in \mathfrak{B}$ into two cases: $s \geq 2mK$ and $s < 2mK$. For the former, we apply Lemma 3.10 to bound the probability for each $s \geq 1$, and then sum over all such terms. For the latter, we bound the contribution using the length of the random variable $m(\ell, R)$.

In detail, we have the bound for the case $s \geq 2mK$:

$$\begin{aligned} \sum_{\substack{B_0 = (m; s; *) \in \mathfrak{B} \\ 1 \leq m \leq n, s \geq 2mK}} s \Pr_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [B(\ell, R) = B_0] &\leq \sum_{\substack{B_0 = (m; s; *) \in \mathfrak{B} \\ 1 \leq m \leq n, s \geq 2mK}} s \cdot 2^{mK-s} \\ &\leq \sum_{s=2K}^{+\infty} s \cdot 2^{-s/2} \sum_{m=1}^{\lfloor s/2K \rfloor} 2^{-K(s/2K-m)} \sum_{B_0 = (m; s; *) \in \mathfrak{B}} 1 \\ &\leq \sum_{s=2K}^{+\infty} s \cdot 2^{-s/2} \cdot 2 \cdot (2n)^{2s/K} \leq \sum_{s=2K}^{+\infty} (2n)^{1+2s/K} \cdot 2^{-s/2}. \end{aligned}$$

We also have the bound for the case $s < 2mK$:

$$\begin{aligned} \sum_{\substack{B_0 = (m; s; *) \in \mathfrak{B} \\ 1 \leq m \leq n, s < 2mK}} s \Pr_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [B(\ell, R) = B_0] &\leq 2K \sum_{\substack{B_0 = (m; s; *) \in \mathfrak{B} \\ m, s}} m \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\Pr_{R \sim \mathbb{P}_R} [B(\ell, R) = B_0] \right] \\ &= 2K + 2K \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [m(\ell, R) - 1] \right], \end{aligned}$$

where the last equality splits m into $(m-1) + 1$, and uses the identity $1 = \sum_{\substack{B_0 = (m; s; *) \in \mathfrak{B} \\ m, s}} \mathbf{1}_{B(\ell, R) = B_0}$.

Finally, by Lemma 3.5, we have

$$\text{Nadp} := \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_{\ell}))] \right] \geq \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [m(\ell, R) - 1] \right].$$

Moreover, for $K = \lceil 25 \log n \rceil + 25$, when $s \geq 2K$, we have:

$$(1 + \log n)(1 + 2s/K) - s/2 \leq 1 + \log n + \frac{2s}{25} - \frac{s}{2} \leq \frac{s}{50} + \frac{2s}{25} - \frac{s}{2} \leq -\frac{s}{4}.$$

Combining all the expressions, we obtain that for $K = \lceil 25 \log n \rceil + 25$,

$$\begin{aligned} \text{Adap} &= \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_{\ell}|] = \sum_{\substack{B_0 = (m; s; *) \in \mathfrak{B} \\ 1 \leq m \leq n, s \geq 2mK}} s \Pr_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [B(\ell, R) = B_0] + \sum_{\substack{B_0 = (m; s; *) \in \mathfrak{B} \\ 1 \leq m \leq n, s < 2mK}} s \Pr_{\ell \leftarrow \pi_{\mathcal{T}}, R \sim \mathbb{P}_R} [B(\ell, R) = B_0] \\ &\leq \sum_{s=2K}^{+\infty} (2n)^{1+2s/K} \cdot 2^{-s/2} + 2K + 2K \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [m(\ell, R) - 1] \right] \\ &= \sum_{s=2K}^{+\infty} 2^{(1+\log n)(1+2s/K)-s/2} + 2K + 2K \cdot \text{Nadp} \leq \sum_{s=2K}^{+\infty} 2^{-s/4} + 2K(1 + \text{Nadp}) \leq 1 + 2K(1 + \text{Nadp}). \end{aligned}$$

Assuming that $\text{Adap} = \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|] \geq 200 \log n$, we conclude that

$$\text{Nadp} \geq \frac{1}{2K} (\text{Adap} - 2K - 1) \geq \frac{\text{Adap} - 1}{90 \log n} - 1 \geq \frac{\text{Adap}}{200 \log n},$$

as desired, provided that $n \geq 100$. \square

4 Tight Bounds for Binary-XOS

In this section, we improve upon the result in Section 3. In light of Lemma 3.2, we only need to consider the special norm f_{xos} . Specifically, we define a subset $A'_\ell \subseteq A_\ell$ for each leaf ℓ . As usual, we define a new norm with respect to A'_ℓ as $f_{\text{xos}}(S) = \max_{\ell} \{|\text{elt}(A'_\ell) \cap S|\}$. Our improvement to Theorem 3.3 is summarized in the following theorem:

Theorem 4.1. *Let A'_ℓ be a subset of A_ℓ for every leaf $\ell \in \mathcal{T}$, and define $K = 50 \lceil \frac{\log n}{\log \log n} \rceil + 50$. Assume that $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|] \geq 6K$. Then, when n is sufficiently large,*

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right] \geq \frac{1}{6K} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_\ell|].$$

By combining Lemma 3.2 and setting $\Lambda(n) = 6K = O\left(\frac{\log n}{\log \log n}\right)$, we obtain the desired bound:

Theorem 4.2. *For $2 \leq r \leq n$, the adaptivity gap of Bernoulli stochastic probing with XOS objective under any prefix-closed \mathcal{F} with $\max_{F \in \mathcal{F}} |F| \leq r$ is $\Lambda_{\text{xos}}(n, r) = O\left(\log r \frac{\log n}{\log \log n}\right)$.*

Using the same lemma, we obtain an upper bound of $O\left(\frac{\log n}{\log \log n}\right)$ for the binary-XOS objective. On the other hand, the lower bound construction in [GNS17] demonstrates that there exists an instance of Bernoulli stochastic probing with binary-XOS objective for which the adaptivity gap is $\Omega\left(\frac{\log n}{\log \log n}\right)$. This example shows that our adaptivity gap upper bound for the binary-XOS objective is *tight* up to a constant factor. Therefore, we conclude the following:

Theorem 4.3. *The adaptivity gap of Bernoulli stochastic probing with binary-XOS objective is $\Theta\left(\frac{\log n}{\log \log n}\right)$.*

The proof of Theorem 4.1 relies heavily on the results in Sections 3.2 and 3.3, using essentially the same notation and algorithms. However, the analysis is significantly more technical and intricate.

4.1 Technical Inequalities

In this section, we introduce a technical function $g(h, p)$ defined as

$$g(h, p) = p \cdot \exp(-0.1 h p^{1/h}),$$

where $h \in \mathbb{N}_+$ and $p \in [0, 1]$. For completeness, we define $g(0, p) = p$ for $p \in [0, 1]$. This is a natural extension since $\lim_{h \rightarrow 0^+} g(h, p) = p \cdot \exp(-0.1 \cdot 0) = p$. We now list some useful properties of g . Fix $h \in \mathbb{N}_+$. Since $0.1 p^{1/h} \leq 0.1 < 1$ for $p \leq 1$, we have:

$$\frac{\partial g}{\partial p}(h, p) = \exp(-0.1 h p^{1/h}) (1 - 0.1 p^{1/h}) > 0. \quad (4.1)$$

Furthermore,

$$\frac{\partial^2 g}{\partial p^2}(h, p) = -0.1 p^{1/h-1} \exp(-0.1 h p^{1/h}) \left(1 + \frac{1}{h} - 0.1 p^{1/h}\right) \leq 0. \quad (4.2)$$

These equations show that, for fixed h , the function g is increasing and concave on $[0, 1]$.

The following result is analogous to Lemma 3.9, but with a more careful treatment when the proportion of S is given by p . In fact, it is a stronger result (up to a constant in h when setting $x_\ell = 1$), since

$$g(h, p) = p \exp(-0.1h p^{1/h}) = \exp\left(h\left(\log p^{1/h} - 0.1p^{1/h}\right)\right) \leq \exp(-0.1h),$$

where the inequality uses $\log x \leq 0.1(x - 1)$ for $x = p^{1/h} \in (0, 1]$. The looseness of Lemma 3.9 for $p < 2^{-h}$ is evident, as the left-hand side is trivially bounded by p . The tightness of Lemma 4.4 when $p = n^{-O(m)}$ is what allows us to strengthen the main result from Theorem 3.3 to Theorem 4.1.

Lemma 4.4. *Let S be a subset of leaves in the tree \mathcal{T} . Suppose there exists a subset T of nodes in \mathcal{T} , not containing any leaves, and an integer $h \in \mathbb{N}$ such that for every $\ell \in S$, $|A_\ell \cap T| \geq h$. Furthermore, assume that each $\ell \in S$ is assigned a weight $x_\ell \in [0, 1]$ such that $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}}[x_\ell \mathbf{1}_{\ell \in S}] = p$. Then,*

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[x_\ell \mathbf{1}_{\ell \in S} \cdot \Pr_{R \sim \mathbb{P}_R} [R \cap P_\ell \cap T = \emptyset] \right] \leq g(h, p).$$

The proof follows the same strategy as in Lemma 3.9: we analyze separately the cases where the root $r_0 \in T$ and $r_0 \notin T$. For the former, we use a corresponding technical inequality (see Lemma B.1); for the latter, the concavity in Eq. (4.2) suffices. The complete proof is deferred to Appendix B.

4.2 Proof for Theorem 4.1

We now present the improved proof of Theorem 3.3. Although this proof employs the same Definition 3.1, Algorithm 1, and Lemma 3.5 as before, we refine the construction of the set T , choose a different parameter K , and perform a more detailed analysis using conditional probability, which improves the adaptivity gap by a $\log \log n$ factor.

To begin, we introduce a refined set T' for a fixed label B_0 , which differs from the one defined in the discussion before Lemma 3.8.

Fix a label $B_0 = (m; s; d_i; e_i; y_j) \in \mathfrak{B}$. We consider all intervals $I_j(B_0) = [y_j, y_{j-1})$, where $1 \leq j \leq \lfloor \frac{s}{K} \rfloor$ and additionally $y_0 := d_1$. Let $J(B_0)$ be the set of all j such that $d_i \in I_j(B_0)$ for some $2 \leq i \leq m$. We define $I(B_0) = \bigcup_{j \in J(B_0), 1 \leq j \leq \lfloor \frac{s}{K} \rfloor} I_j(B_0)$. Therefore, $d_i \notin I(B_0)$, for any $1 \leq i \leq m$.

Recall the definition of $\text{Imp}(\ell, B_0)$ in Eq. (3.2), $S(B_0) = \{\ell \text{ is a leaf of } \mathcal{T} : \exists R, B(\ell, R) = B_0\}$ in Definition 3.2, and $T(B_0)$ as the union of all $\text{Imp}(\ell, B_0)$ for $\ell \in S(B_0)$. For each label $B_0 \in \mathfrak{B}$, we define $T'(B_0)$ to be the set of nodes in $T(B_0)$ whose depth lies in $I(B_0)$. Formally,

$$T'(B_0) = \bigcup_{\ell \in S(B_0)} \{u : u \in \text{Imp}(\ell, B_0), \text{depth of } u \in I(B_0)\}.$$

Clearly, $T'(B_0)$ is a subset of $T(B_0)$, and every node in it has depth in $I(B_0)$. Thus, we obtain:

Lemma 4.5. *Fix a label $B_0 \in \mathfrak{B}$. Define $T'(B_0)$ as above. Then $T'(B_0)$ does not contain any leaf of \mathcal{T} , and for every $\ell \in S(B_0)$, there are at least $(s - mK)$ active ancestors in $T'(B_0)$.*

Proof. Since $T'(B_0) \subseteq T(B_0)$ and by Lemma 3.6, $T'(B_0)$ contains no leaves. Furthermore, by Remark 3.7, for any $\ell \in S(B_0)$, the nodes in $\text{Imp}(\ell, B_0)$ whose depths lie in $I(B_0)$ contribute at least $(s - mK)$ active ancestors. \square

We now recall some consequences of the assumption that $\ell \in S(B_0)$. From Lemma 3.4 (a), $\ell \in L_1(B_0)$ implies $s = |A'_\ell|$, and y_j is the depth of the jK -th node of A'_ℓ (counted from ℓ).

By part (b) of the same lemma, for $2 \leq a \leq m$, there exists a unique node $v_a = v_a(\ell)$ of depth d_a , representing e_a , in $P_\ell \cap L_a(B_0)$ (see also Definition 3.2). Moreover, Lemma 3.4 (c) states that if $B(\ell, R) = B_0$, then $v_a(\ell)$ must be the ‘lowest’ active (i.e., in R) node in P_ℓ and satisfy the condition in Line 3 of Algorithm 1; that is, there exists $B_1 \in \mathfrak{B}$ of length a such that $\text{Pref}_{a-1}(B_1) = \text{Pref}_{a-1}(B_0)$ and $u \in L_a(B_1)$ (here ‘lowest’ refers to the descendant-ancestor order in the tree and should not be confused with the notion of ‘depth’).

We now define an event on R , denoted $D(\ell, B_0)$, for each label $B_0 \in \mathfrak{B}$ and each leaf $\ell \in S(B_0)$ as follows. The event $D(\ell, B_0)$ occurs if:

- For every $2 \leq a \leq m$, $v_a(\ell) \in R$.
- For every node $u \in P_\ell \cap (T(B_0) \setminus T'(B_0))$, it holds that $u \notin R$.

The following lemma summarizes key properties of the event $D(\ell, B_0)$ for fixed ℓ and B_0 . In particular, part (c) asserts that if we fix ℓ and group the labels with respect to certain technical conditions, then any outcome R cannot satisfy two distinct D events in the same group simultaneously. This reflects the fact that $v_a(\ell)$ is the unique ‘lowest’ node meeting the condition in Line 3 of Algorithm 1, as mentioned above.

Lemma 4.6. *Fix a label B_0 and a leaf $\ell \in S(B_0)$. Define the event $D(\ell, B_0)$ as above. Then:*

(a) *The following event inclusion holds:*

$$[B(R \cap P_\ell) = B_0] \subseteq D(\ell, B_0) \cap [R \cap P_\ell \cap T'(B_0) = \emptyset].$$

(b) *The event $D(\ell, B_0)$ is independent of the event $[R \cap P_\ell \cap T'(B_0) = \emptyset]$.*

(c) *Let $|A'_\ell| = s$, and fix a subset $J \subseteq [\lfloor \frac{s}{K} \rfloor]$. Let \mathfrak{B}' denote the set of all labels B' such that $J(B') = J$ and $\ell \in S(B')$. Then,*

$$\sum_{B' \in \mathfrak{B}'} \Pr[D(\ell, B')] \leq 1.$$

Proof. Since we assume $\ell \in S(B_0)$, the nodes $v_a(\ell)$ (for $2 \leq a \leq m$, where m is the length of B_0) are uniquely determined by B_0 . We work within a single path P_ℓ , so any two nodes in P_ℓ are in a descendant-ancestor relation, and two nodes represent the same element if and only if they are identical. For clarity, we abbreviate $v_a(\ell)$ to v_a , though these nodes may differ with respect to different labels.

(a) This inclusion follows directly from part (b) of Lemma 3.4 and from Lemma 3.8:

$$[B(R \cap P_\ell) = B_0] \subseteq \left([R \cap P_\ell \cap T(B_0) = \emptyset] \wedge \bigwedge_{2 \leq a \leq m} [v_a \in R] \right) = D(\ell, B_0) \cap [R \cap P_\ell \cap T'(B_0) = \emptyset].$$

(b) It is clear that $D(\ell, B_0)$ or $[R \cap P_\ell \cap T'(B_0) = \emptyset]$ are just conditions on certain nodes in P_ℓ being in R or not. The nodes involved in $D(\ell, B_0)$ are either $v_a(\ell)$, which is at depth $d_a \notin I(B_0)$, or in $T(B_0) \setminus T'(B_0)$, hence disjoint from $T'(B_0)$. Therefore, nodes involved in $D(\ell, B_0)$ and $[R \cap P_\ell \cap T'(B_0) = \emptyset]$ are disjoint. Since different nodes in P_ℓ represents distinct elements, the two events are independent.

(c) We show that the events $D(\ell, B')$ for all $B' \in \mathfrak{B}'$ are mutually disjoint—that is, any outcome R can satisfy at most one such event.

Suppose, for contradiction, that there exist two labels $B^1, B^2 \in \mathfrak{B}'$ such that $R \in D(\ell, B^1) \cap D(\ell, B^2)$. From Lemma 3.4(a), $\ell \in S(B^1) \cap S(B^2)$ implies $\ell \in L_1(B^1) \cap L_1(B^2)$. Thus, in both labels, $s = |A'_\ell|$, and y_j is the depth of the jK -th node of A'_ℓ (counted from ℓ). We write $B^1 = (m^1; s; d_i^1; e_i^1; y_j)$ and $B^2 = (m^2; s; d_i^2; e_i^2; y_j)$.

Let $a \geq 2$ be the first index such that $e_a^1 \neq e_a^2$ (note that $e_1^1 = e_1^2 = 0$, and we allow $a = \min\{m^1, m^2\} + 1$). By Definition 3.2 and the assumption that $\ell \in S(B^1) \cap S(B^2)$, assume without loss of generality that the corresponding node $v_a^1 := v_a^1(\ell) \in L_a(B^1)$ is an ancestor of $v_a^2 := v_a^2(\ell) \in L_a(B^2)$, and $v_a^1 \neq v_a^2$ (if either node does not exist due to $a = \min\{m^1, m^2\} + 1$, we define the non-existent one to be v_a^1 , a virtual node).

Since $e_i^1 = e_i^2$ for all $2 \leq i < a$, it follows that $v_i^1 = v_i^2$, and hence $d_i^1 = d_i^2$. Consequently, $\text{Pref}_{a-1}(B^1) = \text{Pref}_{a-1}(B^2)$. Furthermore, since $B^1, B^2 \in \mathfrak{B}'$, we have $J(B^1) = J(B^2) = J$, so $I(B^1) = I(B^2)$, as all y_j values are equal.

Because $d_a^2 \notin I(B^2) = I(B^1)$, the node v_a^2 at depth d_a^2 is not contained in $T'(B^1)$. On the other hand, the label $\text{Pref}_a(B^2)$ (of length a) satisfies $\text{Pref}_{a-1}(\text{Pref}_a(B^2)) = \text{Pref}_{a-1}(B^1)$ and $v_a^2 \in P_\ell \cap L_a(B^2)$. Then v_a^2 satisfies all conditions in the $(a-1)$ -th union in Eq. (3.2) for B^1 , since v_a^2 is a descendant of v_a^1 (this still holds even when $a = \min\{m^1, m^2\} + 1$).

Thus, $v_a^2 \in \text{Imp}(\ell, B^1) \subseteq T(B^1)$. Since $v_a^2 \notin T'(B^1)$, it follows that $v_a^2 \in P_\ell \cap (T(B^1) \setminus T'(B^1))$, and therefore $v_a^2 \notin R$ because $D(\ell, B^1)$ occurs. But this contradicts the assumption that $D(\ell, B^2)$ occurs, which would imply $v_a^2 \in R$.

□

We now proceed with the proof of the main theorem Theorem 4.1. Intuitively, in Lemma 3.9 we "consume" the condition $R \cap P_\ell \cap T(B_0) = \emptyset$, yet the set $T(B_0) \cap A_\ell$ can be much larger than $(s - mK)$. In this proof, we only "consume" the condition $R \cap P_\ell \cap T'(B_0) = \emptyset$ and instead employ Lemma 4.4 in place of Lemma 3.9. By leveraging the concavity and monotonicity of $g(h, \cdot)$, we can combine conditional probabilities to obtain the desired bound.

Proof of Theorem 4.1. First, using Lemma 3.5, we can express

$$\text{Adap} := \mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} [|A'_\ell|] = \sum_{B_0=(m;s;*) \in \mathfrak{B}} s \cdot \mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} \left[\Pr_{R \sim \mathbb{P}_R} [B(\ell, R) = B_0] \right],$$

where $(m; s; *)$ denotes a sequence of the form $(m; s; d_i; e_i; y_j)$.

We split the sum over $B_0 = (m; s; *) \in \mathfrak{B}$ into two cases: $s \geq 2mK$ and $s < 2mK$. We bound the latter case using Lemma 3.5:

$$\begin{aligned} \sum_{\substack{B_0=(m;s;*) \in \mathfrak{B} \\ 1 \leq m \leq n, s < 2mK}} s \mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}, R \sim \mathbb{P}_R} [B(\ell, R) = B_0] &\leq 2K \sum_{B_0=(m;s;*) \in \mathfrak{B}} m \mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} \left[\Pr_{R \sim \mathbb{P}_R} [B(\ell, R) = B_0] \right] \\ &= 2K + 2K \mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [m(\ell, R) - 1] \right] \\ &\leq 2K + 2K \mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))] \right]. \end{aligned} \quad (4.3)$$

We denote the target quantity $\text{Nadp} := \mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} [\mathbb{E}_{R \sim \mathbb{P}_R} [f_{\text{xos}}(\text{elt}(R \cap P_\ell))]]$.

Now, we aim to bound the remaining term where $s \geq 2mK$.

For each fixed label $B_0 = (m; s; *)$ with $s \geq 2mK$, we define $T'(B_0)$ and $S(B_0)$ as in the discussion preceding Lemma 4.5. Moreover, for each $\ell \in S(B_0)$, we define $x_\ell \in [0, 1]$ to be the probability $\Pr_{R \sim \mathbb{P}_R} [D(\ell, B_0)]$ that the event $D(\ell, B_0)$ occurs.

From Lemma 4.6 (a) and the definition of $S(B_0)$, we have the inclusion of events:

$$[B(R \cap P_\ell) = B_0] \subseteq D(\ell, B_0) \cap [R \cap P_\ell \cap T' = \emptyset] \cap [\ell \in S(B_0)].$$

Taking expectations, we obtain:

$$\mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} \left[\Pr_{R \sim \mathbb{P}_R} [B(\ell, R) = B_0] \right] \leq \mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} \left[\mathbf{1}_{\ell \in S(B_0)} \cdot \Pr_{R \sim \mathbb{P}_R} [D(\ell, B_0) \cap [R \cap P_\ell \cap T' = \emptyset]] \right].$$

Furthermore, from Lemma 4.6 (b), for fixed B_0 and $\ell \in S(B_0)$, the event $D(\ell, B_0)$ is independent of $[R \cap P_\ell \cap T' = \emptyset]$. Therefore,

$$\mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} \left[\Pr_{R \sim \mathbb{P}_R} [B(\ell, R) = B_0] \right] = \mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} \left[x_\ell \cdot \mathbf{1}_{\ell \in S(B_0)} \cdot \Pr_{R \sim \mathbb{P}_R} [R \cap P_\ell \cap T' = \emptyset] \right].$$

This is exactly the form required by Lemma 4.4. Hence, using Lemma 4.5, we denote $p_{B_0} := \mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} [x_\ell \cdot \mathbf{1}_{\ell \in S(B_0)}]$ with $h := K \lfloor \frac{s}{2K} \rfloor \leq s - mK$ (since $s \geq 2mK$), taking $T = T'(B_0)$, $S = S(B_0)$, we apply Lemma 4.4 to get:

$$\mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} \left[\Pr_{R \sim \mathbb{P}_R} [B(\ell, R) = B_0] \right] \leq g \left(K \lfloor \frac{s}{2K} \rfloor, p_{B_0} \right).$$

Let $s_0 := \lfloor \frac{s}{2K} \rfloor$. Now, we use the fact that the function $g(h, p)$ is concave for fixed $h \geq 1$ and $p \in [0, 1]$, as shown via Jensen's inequality (see Eq. (4.2)). We then sum over all labels with a fixed s_0 (i.e., we sum over all s such that $2Ks_0 \leq s < 2K(s_0 + 1)$ and m satisfying $1 \leq m \leq s_0$), and define

$$q_{s_0} = \sum_{s=2Ks_0}^{2K(s_0+1)-1} \sum_{\substack{B_0=(m;s;*) \in \mathfrak{B} \\ 1 \leq m \leq s_0}} [p_{B_0}] = \sum_{s=2Ks_0}^{2K(s_0+1)-1} \sum_{\substack{B_0=(m;s;*) \in \mathfrak{B} \\ 1 \leq m \leq s_0}} \mathbb{E}_{\ell \leftarrow \pi_\mathcal{T}} [x_\ell \mathbf{1}_{\ell \in S(B_0)}].$$

Furthermore, we denote by r_{s_0} the number of labels satisfying $B_0 = (m; s; *) \in \mathfrak{B}$ with $1 \leq m \leq s_0$ and $2Ks_0 \leq s < 2K(s_0 + 1)$. We can bound r_{s_0} as follows.

There are at most s_0 possibilities for m and $2K$ possibilities for s . For each m, s , there are at most $(n+1)^m \times n^m \times (n+1)^{s/K} \leq (2n)^{2s/K}$ possibilities for d_i, e_i, y_j . Therefore,

$$r_s \leq 2Ks_0(2n)^{2s/K} \leq (2n)^{2s/K+1} \leq (2n)^{4s_0+5}.$$

Applying Jensen's inequality, we obtain:

$$\sum_{s=2Ks_0}^{2K(s_0+1)-1} \sum_{\substack{B_0=(m;s;*) \in \mathfrak{B} \\ 1 \leq m \leq s_0}} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\Pr_{R \sim \mathbb{P}_R} [B(\ell, R) = B_0] \right] \leq \sum_{s=2Ks_0}^{2K(s_0+1)-1} \sum_{\substack{B_0=(m;s;*) \in \mathfrak{B} \\ 1 \leq m \leq s_0}} g(s_0K, p_{B_0}) \leq r_{s_0} g\left(s_0K, \frac{q_{s_0}}{r_{s_0}}\right). \quad (4.4)$$

Note that the quantity $r_{s_0} g\left(s_0K, \frac{q_{s_0}}{r_{s_0}}\right) = q_{s_0} \exp\left(-0.1s_0K \left(\frac{q_{s_0}}{r_{s_0}}\right)^{1/(s_0K)}\right)$ is an increasing function of r_{s_0} for fixed q_{s_0} . Hence,

$$r_{s_0} g\left(s_0K, \frac{q_{s_0}}{r_{s_0}}\right) \leq (2n)^{4s_0+5} g\left(s_0K, \frac{q_{s_0}}{(2n)^{4s_0+5}}\right) = q_{s_0} \exp\left(-0.1s_0K \left(\frac{q_{s_0}}{(2n)^{4s_0+5}}\right)^{\frac{1}{s_0K}}\right). \quad (4.5)$$

To bound q_{s_0} , we expand its definition:

$$q_{s_0} = \sum_{\substack{B_0=(m;s;*) \in \mathfrak{B} \\ 1 \leq m \leq s_0 \\ 2s_0K \leq s < 2(s_0+1)K}} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [\mathbf{1}_{\ell \in S(B_0)} \Pr_{R \sim \mathbb{P}_R} [D(\ell, B_0)]] \leq \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \sum_{\substack{B_0=(*;s;*) \in \mathfrak{B} \\ 2s_0K \leq s < 2(s_0+1)K}} [\mathbf{1}_{\ell \in S(B_0)} \Pr_{R \sim \mathbb{P}_R} [D(\ell, B_0)]],$$

where the last inequality follows by exchanging summation and relaxing the range of m .

For fixed ℓ , note that $\ell \in S(B_0)$ implies $s = |A'_\ell|$ by Lemma 3.4 (a). Thus, if ℓ contributes to the above sum, then $2s_0K \leq |A'_\ell| < 2(s_0+1)K$.

Now we group all labels $B_0 = (*; s; *)$ and $\ell \in S(B_0)$ with respect to $J(B_0) \subseteq [\lfloor \frac{s}{K} \rfloor]$. For a fixed group, Lemma 4.6 (c) ensures that the sum of $\Pr[D(\ell, B_0)]$ over all such labels B_0 is at most 1.

Therefore,

$$\begin{aligned} q_{s_0} &= \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \sum_{\substack{B_0=(m;s;*) \in \mathfrak{B} \\ 2s_0K \leq s < 2(s_0+1)K}} [\mathbf{1}_{\ell \in S(B_0)} \Pr_{R \sim \mathbb{P}_R} [D(\ell, B_0)]] \\ &\leq \max_{\substack{\ell: |A'_\ell|=s \\ 2s_0K \leq s < 2(s_0+1)K}} \sum_{J \subseteq [\lfloor \frac{s}{K} \rfloor]} \sum_{B_0: J(B_0)=J, \ell \in S(B_0)} \Pr[D(\ell, B_0)] \\ &\leq \max_{2s_0K \leq s < 2(s_0+1)K} \sum_{J \subseteq [\lfloor \frac{s}{K} \rfloor]} 1 \leq 2^{2s_0+1}. \end{aligned}$$

Combining Eq. (4.4), Eq. (4.5), the monotonicity of $g(s_0K, p)$ in p (from Lemma B.1), and the bound $q_{s_0} \leq 2^{2s_0+1}$, we get (for $s_0 \geq 1, n > 100$):

$$\begin{aligned} \sum_{s=2Ks_0}^{2K(s_0+1)-1} \sum_{\substack{B_0=(m;s;*) \in \mathfrak{B} \\ 1 \leq m \leq s_0}} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\Pr_{R \sim \mathbb{P}_R} [B(\ell, R) = B_0] \right] &\leq q_{s_0} \exp\left(-0.1s_0K \left(\frac{q_{s_0}}{(2n)^{4s_0+5}}\right)^{\frac{1}{s_0K}}\right) \\ &\leq 2^{2s_0+1} \exp\left(-0.1s_0K n^{-10/K}\right). \end{aligned}$$

Therefore, summing over all s_0 (when n is large enough and $s \geq 2K$):

$$\sum_{\substack{B_0=(m;s;*) \in \mathfrak{B} \\ 1 \leq m \leq r, s \geq 2mK}} s \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\Pr_{R \sim \mathbb{P}_R} [B(\ell, R) = B_0] \right] \leq \sum_{s_0=1}^{+\infty} 2K(s_0+1) \cdot 2^{2s_0+1} \exp(-0.1s_0K n^{-10/K}).$$

Since $K = 50 \lceil \frac{\log n}{\log \log n} \rceil + 50$, we have $n^{10/K} = 2^{\frac{10 \log n}{K}} \leq 2^{\frac{\log \log n}{5}} = \log^{1/5} n \leq K/40$ for sufficiently large n . Hence,

$$\begin{aligned} \sum_{s_0=1}^{+\infty} 2K(s_0+1) \cdot 2^{2s_0+1} \exp(-0.1s_0Kn^{-10/K}) &\leq \sum_{s_0=1}^{+\infty} 2K(s_0+1) \cdot 2^{2s_0+1} \exp(-4s_0) \\ &\leq 2K \sum_{s_0=1}^{+\infty} (s_0+1) \cdot 2^{2s_0+1-5s_0} \leq 2K. \end{aligned}$$

Combining this with Eq. (4.3) and the partition strategy of $s \geq 2mK$ and $s < 2mK$, we conclude:

$$\text{Adap} \leq 2K + 2K \cdot \text{Nadp} + 2K \iff \text{Nadp} \geq \frac{\text{Adap}}{2K} - 2.$$

Thus, when $\text{Adap} = \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [|A'_{\ell}|] \geq 6K$, it follows that $\text{Nadp} \geq \frac{\text{Adap}}{2K} - 2 \geq \frac{\text{Adap}}{6K}$. The result follows. \square

5 Adaptivity Gap for Symmetric Norm

We first introduce the Bernoulli setting for monotone symmetric norms.

Definition 5.1 (Bernoulli Stochastic Probing with Symmetric Norm Objective). In this special case of the Stochastic Probing problem, each random variable X_i takes value $c_i \in \mathbb{R}_{\geq 0}$ with probability p_i , and 0 with probability $1 - p_i$, for all $i \in U$. Also, the objective function f is a *monotone symmetric norm*. Other settings are the same as Definition 1.1.

It is important to emphasize that the definition here is slightly different from Definition 2.6 (in which the Bernoulli random variable is 0/1). In Definition 5.1, the random variable may take values $c_i \in \mathbb{R}_{\geq 0}$, which is crucial for our analysis. The reason that we do not absorb c_i into the objective function f , as doing so would destroy the symmetry property of f .

Without loss of generality, we may assume that $p_i, c_i > 0$ for all i . We denote by $f(S)$ (for $S \subseteq U = [n]$) the value $f(\mathbf{v})$, where the n -dimensional vector \mathbf{v} has its i -th coordinate equal to c_i if $i \in S$ and 0 otherwise, as we did in Definition 2.6.

Let $\Lambda_{\text{sym}}(n)$ denote the supremum of the adaptivity gap for *Bernoulli stochastic probing with any symmetric norm objective* and any *prefix-closed* feasibility constraint \mathcal{F} . We show in this section that the adaptivity gap for this problem is bounded by a constant, namely, $\Lambda_{\text{sym}}(n) = O(1)$ (with an explicit upper bound of 2050).

In the remainder of this section, we adopt the binary decision tree model and use the notation for **active ancestors** $A_{\ell} \subseteq P_{\ell}$ and the random variable $R \sim \mathbb{P}_R$ introduced in Section 2.2. For simplicity, we write $f(A_{\ell})$ instead of $f(\text{elt}(A_{\ell}))$, as our analysis here avoids the complicated method of changing the maximizer leaf employed in Section 3 and Section 4.

In view of Lemma 2.2 (whose proof in Appendix A applies to this setting as well), by setting $\lambda = 4^{-4}$ and possibly incurring a factor of 2 loss in the adaptivity gap, we may assume that all elements are 4^{-4} -small. By scaling f by a constant, we normalize the optimal reward (the optimal binary decision tree denoted by \mathcal{T}_0) to

$$\text{OPT} = \text{Adap}(\mathcal{T}_0, f) = \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(A_{\ell})] = 1.$$

Furthermore, we assume that $f(1, 0, 0, \dots, 0) = 1$. Indeed, if $f(1, 0, 0, \dots, 0) = \frac{1}{a} > 0$, we can simply divide each c_i by a and define a new function $f_1(\mathbf{v}) = f(a\mathbf{v})$ to replace f . This transformation preserves the λ -smallness, the adaptivity gap, and the value of OPT. Consequently, we have $f(x, 0, 0, \dots, 0) = xf(1, 0, 0, \dots, 0) = x$, which we denote by $f(x)$. Now, for any internal node $u \in \mathcal{T}_0$, we round each $c_{\text{elt}(u)} = f(c_{\text{elt}(u)}) \leq 4^{-4}$ to the largest negative power of 4 strictly smaller than $c_{\text{elt}(u)}$. After rounding, each $c_{\text{elt}(u)} \leq 4^{-5}$. Using linearity of the norm objective, the final reward OPT_1 is still at least $1/4$.

Next, we modify the optimal decision tree \mathcal{T}_0 slightly into another decision tree \mathcal{T} with some additional requirements, as in the following lemma.

Lemma 5.1. *There exists a decision tree \mathcal{T} satisfying the following requirements:*

1. *Each root-leaf path is a subsequence of some root-leaf path in \mathcal{T}_0 . Thus, the corresponding element sequence satisfies the feasibility constraint \mathcal{F} ;*
2. *For any leaf $\ell \in \mathcal{T}$, $f(A_\ell) \leq 2$;*
3. $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(A_\ell)] \geq 1/8$.

Proof. In the original adaptive strategy, the algorithm starts at the root of \mathcal{T}_0 , follows the decision tree based on the probing outcomes, and eventually terminates at a leaf ℓ , obtaining the set A_ℓ . In our modified strategy, the algorithm still follows \mathcal{T}_0 , but before probing any node u , it checks whether A_u (the set of active ancestors of u) satisfies $f(A_u) \geq 1$. If it does, the algorithm terminates at node u . Looking into the decision tree model, the new decision tree \mathcal{T} is obtained by replacing, for each node u in \mathcal{T}_0 where $f(A_u) \geq 1$ but $f(A_v) < 1$ (v is the parent of u), the entire subtree \mathcal{T}_u rooted at u with a single leaf u . Clearly, no termination signs in the new strategy have an ancestor-descendant relationship in \mathcal{T}_0 .

We now argue that the reward for the new decision tree is at least $1/8$. To see this, consider any execution of the new strategy terminating at a node u in the original tree \mathcal{T}_0 . By the subsequence property (see the formal definition in Lemma A.1; the precise statement is not critical for understanding), the expected additional reward after u is at most 1; otherwise, a better adaptive strategy could be constructed by first following the tree to u and then applying the adaptive strategy at the subtree \mathcal{T}_u . Formally, if $\pi_{\mathcal{T}_u}$ denotes the original distribution of leaves $\pi_{\mathcal{T}}$ conditioned on $\ell \in \mathcal{T}_u$, then

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_u}} [f(A_\ell \setminus A_u)] \leq \text{OPT}_1 = 1.$$

Hence, when $f(A_u) \geq 1$, by subadditivity we have

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_u}} [f(A_\ell)] \leq \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_u}} [f(A_\ell \setminus A_u)] + f(A_u) \leq 1 + f(A_u) \leq 2f(A_u).$$

When $f(A_u) < 1$, then u must be a leaf in the original tree, and we have

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_u}} [f(A_\ell)] = f(A_u) \leq 2f(A_u).$$

Now, we take expectation with respect to the leaf distribution $u \leftarrow \pi_{\mathcal{T}}$ (note that now u is a leaf in $\pi_{\mathcal{T}}$), we obtain

$$\text{OPT}_1 = \mathbb{E}_{u \leftarrow \pi_{\mathcal{T}}} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_u}} [f(A_\ell)] \leq 2 \mathbb{E}_{u \leftarrow \pi_{\mathcal{T}}} [f(A_u)],$$

hence $\mathbb{E}_{u \leftarrow \pi_{\mathcal{T}}} [f(A_u)] \geq \text{OPT}_1/2 \geq 1/8$.

Moreover, in the new tree \mathcal{T} , we have $f(A_u) \leq 2$ for every leaf u . To see this, let v be the parent of u . Then,

$$f(A_u) \leq f(A_v) + f(u) \leq 1 + c_{\text{elt}(u)} \leq 1 + 4^{-5} \leq 2,$$

where we have used the fact that after rounding, each $c_{\text{elt}(u)} \leq 4^{-5}$ for any internal node $u \in \mathcal{T} \subseteq \mathcal{T}_0$. \square

By the above lemma, it suffices to lower bound the non-adaptive reward $\text{Nadp}(\mathcal{T}, f)$ for the modified decision tree \mathcal{T} , using the natural non-adaptive strategy ALGPATH discussed in Section 2.2.

For each internal node $u \in \mathcal{T}$, if $c_{\text{elt}(u)} = 4^{-k}$, we place u into a set Q_k , and refer to u as being in class k . Note that no leaf is included in any Q_k , as leaves do not represent any elements. Recall that after rounding, $c_{\text{elt}(u)} \leq 4^{-5}$ for each internal node $u \in \mathcal{T}$. Therefore, in the current tree, only classes Q_k with $k \geq 5$ are nonempty.

For each leaf ℓ , let $D_\ell \subseteq A_\ell$ be the union of the sets $A_\ell \cap Q_k$ for all $k \geq 5$ satisfying the condition $|A_\ell \cap Q_k| \geq 2^k$. Thus, since each node is in exactly one Q_k , we have the following relation:

$$D_\ell = \bigcup_{k: |A_\ell \cap Q_k| \geq 2^k} A_\ell \cap Q_k, \quad A_\ell \setminus D_\ell = \bigcup_{\substack{k \geq 5 \\ |A_\ell \cap Q_k| < 2^k}} A_\ell \cap Q_k.$$

Thus, the contribution from $A_\ell \setminus D_\ell$ to the optimal reward is small, leaving a remaining contribution that is at least a constant:

$$\begin{aligned}
\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(A_\ell \cap D_\ell)] &\geq \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(A_\ell) - f(A_\ell \setminus D_\ell)] \\
&= \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(A_\ell)] - \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(A_\ell \setminus D_\ell)] \\
&\stackrel{(\star)}{\geq} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(A_\ell)] - \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\sum_{k=5}^{\infty} 2^k f(4^{-k}) \right] \\
&= 1/8 - 1/16 = 1/16.
\end{aligned} \tag{5.1}$$

Note that the inequality marked (\star) uses the fact that f is subadditive and symmetric. This bound implies that for each ℓ , we need only consider the classes in which a sufficiently large number of nodes in A_ℓ appear.

Now, we introduce a technical definition.

Definition 5.2. For each $k \geq 5$, we call a leaf ℓ a **k -bad leaf** if $|A_\ell \cap Q_k| \geq 2^k$, but the following holds:

$$\sum_{e \in \text{elt}(P_\ell \cap Q_k)} p_e = \mathbb{E}_{R \sim \mathbb{P}_R} [|R \cap P_\ell \cap Q_k|] \leq \frac{1}{8} |A_\ell \cap Q_k|.$$

Intuitively, a **k -bad leaf** is one for which the non-adaptive algorithm is unlikely to achieve a comparable reward from class Q_k . However, we are able to show that the proportion of **k -bad leaves** (with respect to the distribution $\pi_{\mathcal{T}}$) is small. The following lemma, whose proof is analogous to those of Lemma 3.9 and Lemma 4.4 (by induction on the height of the tree), establishes this bound.

Lemma 5.2. *Let S be a subset of leaves in the tree \mathcal{T} , and let $T \subseteq \mathcal{T}$ be a subset of nodes that does not include any leaf. Suppose that for every $\ell \in S$, the number of active ancestors of ℓ in T is at least h , i.e., $|A_\ell \cap T| \geq h$. Furthermore, assume there exists a real number $q \in \mathbb{R}$ such that for every $\ell \in S$, the following inequality holds:*

$$|A_\ell \cap T| \geq 8 \sum_{e \in \text{elt}(P_\ell \cap T)} p_e + q.$$

Then, the following bound holds:

$$\Pr_{\ell \leftarrow \pi_{\mathcal{T}}} [\ell \in S] \leq e^{-h-q}.$$

Proof. We prove the claim by induction on the height of \mathcal{T} .

When the height of \mathcal{T} is 1, $h = 0, q \leq 0$, since T does not contain any leaf. Hence, the base case is trivial. Assume now that the claim holds for trees of smaller height. For the current tree \mathcal{T} , let \mathcal{T}_1 and \mathcal{T}_2 denote the subtrees corresponding to the **Yes** and **No** arcs from the root r_0 , respectively. Let λ be the probability that the root is active (if r_0 has only one child, we set $\lambda = 0$ or 1 accordingly). Then, the path distribution $\pi_{\mathcal{T}}$ assigns probability λ to the **Yes** subtree \mathcal{T}_1 and $1 - \lambda$ to the **No** subtree \mathcal{T}_2 . Conditioning on the **Yes** subtree (**No** subtree resp.), $\pi_{\mathcal{T}}$ has the same distribution of leaves as in $\pi_{\mathcal{T}_1}$ ($\pi_{\mathcal{T}_2}$ resp.). Denote $\Pr_{\ell \leftarrow \pi_{\mathcal{T}_1}} [\ell \in S] = u$ and $\Pr_{\ell \leftarrow \pi_{\mathcal{T}_2}} [\ell \in S] = v$. Then, we have $\Pr_{\ell \leftarrow \pi_{\mathcal{T}}} [\ell \in S] = \lambda u + (1 - \lambda)v$.

We now consider two cases.

Case 1. $r_0 \notin T$.

In this case, the two subtrees inherit essentially the same conditions (h, q) as the original tree because $r_0 \notin T$. Thus, by the induction hypothesis, $u, v \leq e^{-h-q}$, so that $\Pr_{\ell \leftarrow \pi_{\mathcal{T}}} [\ell \in S] = \lambda u + (1 - \lambda)v \leq e^{-h-q}$.

Case 2. $r_0 \in T$.

In this case, for leaves ℓ in $S \cap \mathcal{T}_1$, there are at least $h - 1$ active ancestors in $T \cap \mathcal{T}_1$. We denote $P'_\ell = P_\ell \setminus \{r_0\}$ and $A'_\ell = A_\ell \setminus \{r_0\}$ for $\ell \in S \cap \mathcal{T}_1$. Then, for $\ell \in S \cap \mathcal{T}_1$,

$$|A'_\ell \cap T| \geq 8 \sum_{e \in \text{elt}(P'_\ell \cap T)} p_e + 8\lambda - 1 + q.$$

Applying the induction hypothesis on \mathcal{T}_1 , we deduce $u \leq e^{-h+2-8\lambda-q}$. Similarly, for the **No** subtree, every leaf has at least h active ancestors in $T \cap \mathcal{T}_2$, and

$$|A'_\ell \cap T| \geq 8 \sum_{e \in \text{elt}(P'_\ell \cap T)} p_e + 8\lambda + q.$$

By the induction hypothesis on \mathcal{T}_2 , we obtain $v \leq e^{-h-8\lambda-q}$. Combining these estimates and applying Fact 2 to the inequality marked (\star) , we have

$$\Pr_{\ell \leftarrow \pi_{\mathcal{T}}} [\ell \in S] = \lambda u + (1 - \lambda)v \leq e^{-h-q-8\lambda} \cdot (\lambda e^2 + (1 - \lambda)) \leq e^{-h-q-8\lambda} \cdot e^{(e^2-1)\lambda} \stackrel{(\star)}{\leq} e^{-h-q},$$

which completes the proof. \square

Thus, for each $k \geq 5$, if we take the set of **k -bad leaves** as S and the nodes in class k as $T = Q_k$, then by choosing $h = 2^k$ and $q = 0$, the proportion of **k -bad leaves** is at most e^{-2^k} . For each ℓ , let

$$G_\ell = \bigcup_{\ell \text{ is a } k\text{-bad leaf}} (A_\ell \cap Q_k).$$

Note that for any leaf ℓ , since f is monotone, $f(G_\ell) \leq f(A_\ell) \leq 2$. Therefore,

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(G_\ell)] \leq 2 \Pr_{\ell \leftarrow \pi_{\mathcal{T}}} [G_\ell \neq \emptyset] \leq 2 \sum_{k=5}^{+\infty} \Pr_{\ell \leftarrow \pi_{\mathcal{T}}} [\ell \text{ is a } k\text{-bad leaf}] \leq 2 \sum_{k=5}^{+\infty} 2^{-2^k} \leq 2 \sum_{k=5}^{+\infty} 2^{-27-k} \leq 2^{-30}. \quad (5.2)$$

We now state the following lemma, which shows that if ℓ is not a **k -bad leaf**, then we can always obtain comparable non-adaptive reward.

Lemma 5.3. *For any fixed leaf ℓ , we have that*

$$\mathbb{E}_{R \sim \mathcal{P}_R} [f(R \cap P_\ell \cap D_\ell)] \geq \frac{1}{64} f((A_\ell \cap D_\ell) \setminus G_\ell).$$

Proof. For each $k \geq 5$ such that $b_{\ell,k} := |A_\ell \cap Q_k| \geq 2^k$ and ℓ is not a **k -bad leaf**, we include such k in a set $K_\ell \subseteq \mathbb{N}$. For each $k \in K_\ell$, consider the probability over R that $|R \cap P_\ell \cap Q_k|$ is at least $b_{\ell,k}/32$. Observe that $|R \cap P_\ell \cap Q_k|$ is the sum of $t_{\ell,k} = |P_\ell \cap Q_k| \geq b_{\ell,k}$ Bernoulli random variables $Y_1, \dots, Y_{t_{\ell,k}}$, each corresponding to an element in $\text{elt}(P_\ell \cap Q_k)$. Since $\mathbb{E}[\sum_{i=1}^{t_{\ell,k}} Y_i] = \mathbb{E}[|R \cap P_\ell \cap Q_k|] \geq b_{\ell,k}/8$ (as ℓ is not a **k -bad leaf**), the Chernoff bound (see Fact 2.1) with $\epsilon = 3/4$ implies that

$$\Pr \left[\sum_{i=1}^{t_{\ell,k}} Y_i \leq \frac{b_{\ell,k}}{32} \right] \leq \exp \left(-\frac{t_{\ell,k}}{16} \right) \leq \exp \left(-\frac{b_{\ell,k}}{16} \right) \leq \exp(-2^{k-4}).$$

By the union bound, the probability that there exists such a k is at most $\sum_{k=5}^{\infty} \exp(-2^{k-4}) \leq \frac{1}{2}$. Therefore, we have that

$$\Pr_R \left[|R \cap P_\ell \cap Q_k| \geq \frac{b_{\ell,k}}{32}, \forall k \in K_\ell \right] \geq \frac{1}{2}.$$

Next, we observe that

$$(A_\ell \cap D_\ell) \setminus G_\ell = \bigcup_{k \in K_\ell} (A_\ell \cap Q_k).$$

Since this set is a subset of P_ℓ and each node represents a distinct element, its corresponding vector has exactly $b_{\ell,k}$ copies of 4^{-k} for each $k \in K_\ell$. By the symmetry and subadditivity of f , the f -value of this vector is at most 32 times the f -value of a vector with $\lceil b_{\ell,k}/32 \rceil$ copies of 4^{-k} . We have shown that, with probability at least 1/2, $|R \cap P_\ell \cap Q_k| \geq \lceil b_{\ell,k}/32 \rceil$ for every $k \in K_\ell$. For such R , it follows that

$$f(R \cap P_\ell \cap D_\ell) \geq \frac{1}{32} f((A_\ell \cap D_\ell) \setminus G_\ell).$$

Averaging over all R (and using the fact that $f \geq 0$), the lemma follows. \square

Finally, we are in a position to lower bound $\text{Nadp}(\mathcal{T}, f)$ by a constant.

Theorem 5.4. *The adaptivity gap for Bernoulli Stochastic Probing with Symmetric Norm Objective is $O(1)$.*

Proof. By Lemma 5.3, Eq. (5.1) and Eq. (5.2), we have

$$\begin{aligned}
\text{Nadp}(\mathcal{T}, f) &= \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f(R \cap P_{\ell})] \right] \\
&\geq \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[\mathbb{E}_{R \sim \mathbb{P}_R} [f(R \cap P_{\ell} \cap D_{\ell})] \right] \\
&\geq \frac{1}{64} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f((A_{\ell} \cap D_{\ell}) \setminus G_{\ell})] \\
&\geq \frac{1}{64} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(A_{\ell} \cap D_{\ell})] - \frac{1}{64} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [f(G_{\ell})] \\
&\geq \frac{1}{1024} - \frac{1}{64 \cdot 2^{30}} \geq \frac{1}{1025}.
\end{aligned}$$

This completes the proof, as the original $\text{OPT} = 1$. \square

6 Generalization to non-Bernoulli Case and Other Objectives

In this section, we generalize our results on Bernoulli Stochastic Probing to general random variables based on the previous results for Bernoulli variables.

We state the following three theorems, where n denotes the cardinality of the ground set, and $2 \leq r \leq n$ denotes an upper bound for the maximal length of a sequence in \mathcal{F} (the same as in Section 3.1). Recall the adaptivity gap functions $\Lambda_{\text{xos}}(n, r)$ and $\Lambda_{\text{sym}}(n)$ for the two Bernoulli settings defined in Definition 2.6 and Definition 5.1.

Theorem 6.1 ([GNS17]). *The adaptivity gap for Bernoulli stochastic probing with any subadditive objective under any prefix-closed \mathcal{F} with $\max_{F \in \mathcal{F}} |F| \leq r$ is $O(\Lambda_{\text{xos}}(n, r) \log n)$.*

Theorem 6.2. *The adaptivity gap for stochastic probing with any monotone norm under any prefix-closed \mathcal{F} with $\max_{F \in \mathcal{F}} |F| \leq r$ is $O(\Lambda_{\text{xos}}(n \log 8r, r \log 8r))$.*

Theorem 6.3. *The adaptivity gap for stochastic probing with monotone symmetric norm is $O(\Lambda_{\text{sym}}(n \log 8n))$.*

The first theorem is a direct consequence of the relationship between (monotone, normalized) subadditive objectives and XOS functions, and already appear informally in [GNS17]. The proofs of the other two theorems are very similar to the classical methods in [Sin18] but in slightly different settings. All three proofs are deferred to Appendix C.

Combining the three theorems with the upper bounds of adaptivity gap for Bernoulli settings: Theorem 4.2 and Theorem 5.4, we obtain the promised results for general random variables, subadditive objectives and monotone symmetric norms.

Corollary 6.4. *For $2 \leq r \leq n$, the adaptivity gap for Bernoulli stochastic probing with any subadditive objective under any prefix-closed \mathcal{F} with $\max_{F \in \mathcal{F}} |F| \leq r$ is upper bounded by*

$$O\left(\log r \frac{\log^2 n}{\log \log n}\right) = O(\log^3 n).$$

Corollary 6.5. *For $2 \leq r \leq n$, the adaptivity gap for stochastic probing with any monotone norm under any prefix-closed \mathcal{F} with $\max_{F \in \mathcal{F}} |F| \leq r$ is upper bounded by*

$$O\left(\log r \frac{\log n}{\log \log n}\right).$$

Corollary 6.6. *The adaptivity gap for stochastic probing with monotone symmetric norm is $O(1)$.*

7 Conclusion and Future Directions

In this paper, we investigated the stochastic probing problem under a general monotone norm and subadditive objectives. First, we resolved a central open problem posed in [GNS17, KMS24] by establishing that the adaptivity gap for stochastic probing with general monotone norms is bounded by $O(\log^2 n)$, which can be further refined to $O\left(\log r \cdot \frac{\log n}{\log \log n}\right)$ where r is the maximum number of elements that can be probed under the feasibility constraint. As a by-product, we obtain an asymptotically tight adaptivity gap $\Theta\left(\frac{\log n}{\log \log n}\right)$ for Bernoulli stochastic probing with binary-XOS objectives. We also derived an $O(\log^3 n)$ upper bound for Bernoulli stochastic probing with general subadditive objectives. Second, for the special case of monotone symmetric norms, we prove that the adaptivity gap is $O(1)$ (in fact, bounded by 2050), answering an open question raised in [PRS23] and improving upon their $O(\log n)$ bound.

Several intriguing open questions remain. Notably, while our upper bound for general monotone norms shows an $O(\log r)$ gap compared to the best known lower bound of $\Omega\left(\frac{\log n}{\log \log n}\right)$, bridging this gap likely requires deeper understanding of the non-binary weight vectors in the XOS representation of the objective function f , representing an important direction for future research. In addition, refining the constant for monotone symmetric norms to obtain a tight bound remains an appealing challenge. Finally, extending our techniques to more general stochastic optimization settings and further exploring algorithmic applications of our insight in stochastic combinatorial optimization remain interesting further directions.

Acknowledgments

We thank Yichuan Wang for valuable feedback on an earlier draft of this manuscript. We also thank Sahil Singla for detailed and insightful discussions, which simplified the presentation of the overview.

Appendix

A Basic Lemmas for Stochastic Probing

In this section, we introduce several key tools for analyzing the general *stochastic probing* problem, as defined in Definition 1.1. The lemmas and proofs here closely resemble those in [GNS17]. However, unlike [GNS17], which focuses on the case where all random variables X_i follow Bernoulli distributions, we extend the analysis to handle multi-valued random variables and present the arguments using our own notation. These results play an important role in the proof of Lemma 2.2 and are also applied in Appendix C (i.e., the arguments given in Section 6).

Throughout this section, we only assume that the objective function f satisfies the standard properties of being normalized, monotone, and subadditive, as described in conditions (1), (2), and (3) of Definition 2.1. We also assume that the feasibility constraint is *prefix-closed*. In our model, the random variables X_i are allowed to take on multiple values. Let OPT denote the expected reward achieved by the optimal adaptive strategy.

Definition A.1 (λ -large (small) outcomes and variables). For any $\lambda > 0$, we call an outcome x_i of X_i λ -large if and only if $f(0, 0, \dots, 0, x_i, 0, 0, \dots, 0) \geq \lambda \cdot \text{OPT}$, and λ -small otherwise. If every possible outcome of X_i is λ -large (or λ -small), then we say that X_i is a λ -large (or λ -small) variable.

Definition A.2 (λ -large decomposition). For any $\lambda > 0$ and any coordinate $i \in [n]$, define the random variable $Y_{i,\lambda}$ by

$$Y_{i,\lambda} = \begin{cases} X_i, & \text{if } X_i \text{ is a } \lambda\text{-large outcome,} \\ 0, & \text{otherwise.} \end{cases}$$

Similarly, define $Z_{i,\lambda}$ by replacing “large” with “small” in the above definition.

We begin by proving a subsequence property, which is analogous to the subtree property in [GNS17, Assumption 5.3]. In adaptive strategies, the decision of which element to probe next can depend on the outcomes observed so far. Suppose that during the execution of an adaptive strategy we have already probed a sequence $P_1 \in \mathcal{F}$ (recall that \mathcal{F} is prefix-closed) and obtained the outcome X_{P_1} (that is, the n -dimensional vector whose i th coordinate equals X_i if $i \in P_1$ and 0 otherwise). Then, the remaining probing process produces a distribution of final outcomes, which we denote by \mathcal{D}_{P_1} . For any $X_P \leftarrow \mathcal{D}_{P_1}$, the difference $X_P - X_{P_1}$ represents the outcomes obtained after probing P_1 .

Lemma A.1 (Subsequence Property). *For any adaptive strategy and any sequence $P_1 \in \mathcal{F}$ probed during its execution, we have*

$$\mathbb{E}_{X_P \leftarrow \mathcal{D}_{P_1}} [f(X_P - X_{P_1})] \leq \text{OPT}.$$

Proof. Consider an alternative adaptive strategy that first probes the sequence $P_1 \in \mathcal{F}$ and receives X_{P_1} , and then follows the original adaptive strategy. Since the overall strategy still satisfies the feasibility constraint, its expected reward is at least $\mathbb{E}_{X_P \leftarrow \mathcal{D}_{P_1}} [f(X_P)]$. By the monotonicity of f , we have $f(X_P) \geq f(X_P - X_{P_1})$, so that the expected reward is at least $\mathbb{E}_{X_P \leftarrow \mathcal{D}_{P_1}} [f(X_P - X_{P_1})]$. Since OPT is the reward of the optimal adaptive strategy, the claim follows. \square

For any probed sequence $P \subseteq U = [n]$, the λ -large decomposition of X_P is given by $X_P = Y_{P,\lambda} + Z_{P,\lambda}$, where $Y_{P,\lambda}$ and $Z_{P,\lambda}$ consist of the outcomes $Y_{i,\lambda}$ and $Z_{i,\lambda}$, respectively, for each coordinate $i \in P$ (with coordinates for $i \notin P$ set to 0).

Lemma A.2. *Assume that the adaptive strategy is optimal. If $\mathbb{E}[f(Y_{P,\lambda})] \geq \text{OPT}/2$, then there exists a non-adaptive algorithm whose expected reward is at least $\frac{\text{OPT}}{2(1+1/\lambda)}$.*

Proof. We begin by constructing an intermediate strategy derived from the optimal adaptive strategy. This new strategy stops probing as soon as it encounters a λ -large outcome. Let OPT' denote the expected reward from the (at most one) λ -large outcome under this strategy. Let OPT_0 be the expected contribution from

any additional λ -large outcomes that would be probed in the original optimal strategy *after* the first one is found.

By construction, every λ -large outcome seen in any execution of the optimal strategy is either the first such outcome (counted in OPT'), or a later one (counted in OPT_0). Since the objective f is subadditive, we have: $\text{OPT}' + \text{OPT}_0 \geq \mathbb{E}[f(Y_{P,\lambda})] \geq \text{OPT}/2$.

Next, consider one execution of this intermediate strategy. There are two cases:

Case 1. The strategy encounters a λ -large outcome and terminates. By definition of λ -large, this outcome has f -value at least $\lambda \cdot \text{OPT}$. Now imagine continuing to probe as in the original optimal strategy. By the subsequence property (Lemma A.1), the expected total reward of these further probes is at most OPT . Thus, the expected value contributed by any additional λ -large outcomes is also at most OPT by monotonicity.

Case 2. The strategy finds no λ -large outcome and continues to the end. In this case, the expected contribution by further λ -large outcomes is zero.

Thus, in either case, the intermediate strategy achieves at least a λ fraction of the expected value contributed by later λ -large outcomes in the original strategy. Taking expectation over all executions gives: $\lambda \cdot \text{OPT}_0 \leq \text{OPT}'$. Combining with $\text{OPT}' + \text{OPT}_0 \geq \text{OPT}/2$, we derive $\text{OPT}' \geq \frac{\text{OPT}}{2(1+1/\lambda)}$.

Now, we construct a *non-adaptive* strategy that matches the reward of the intermediate adaptive strategy. This strategy simulates the behavior of the original optimal strategy. Whenever the adaptive strategy probes an element, the non-adaptive strategy samples an element by itself from the same known distribution *conditioned* on the outcome being λ -small. It then probes the same element as the adaptive strategy does upon receiving this (unreal) conditioned outcome sampled by the non-adaptive strategy itself.

This strategy is clearly feasible and non-adaptive. We show, by induction on the height of the decision tree for the intermediate strategy, that it achieves at least the same expected reward as the intermediate strategy.

The base case is trivial when the decision tree is of height 1 (the outcome of the probed element has same distribution in both strategies). For the general case, suppose the root element gives a λ -large outcome with probability p , and a λ -small outcome with probability $1 - p$. The intermediate strategy achieves expected reward A and terminates with probability p , and with probability $1 - p$, continues probing based on the received λ -small outcome, say distributed with probabilities $\{q_i\}$ (so that $\sum q_i = 1 - p$).

For the non-adaptive strategy, it occurs with probability p (independent of the strategy) that the first probe gives a λ -large outcome. Under this condition, by monotonicity, the expected reward is at least the f value of only the λ -large outcome for the first probe, which is exactly A .

There is also another $1 - p$ probability (independent of the strategy) that the first probe gives a λ -small outcome. Hence, conditioning on such event, the probability that the strategy chooses the i -th element to probe is exactly $q_i / (\sum q_i) \times (1 - p) = q_i$, the same as the probability of the intermediate adaptive strategy chooses the i -th element to probe. Hence, we use induction on the sub-execution of both strategies. Together with the p probability discussed above, we conclude that the reward of the non-adaptive strategy is at least that of the intermediate strategy.

Hence, the non-adaptive strategy achieves expected reward at least $\text{OPT}' \geq \frac{\text{OPT}}{2(1+1/\lambda)}$, as desired. \square

We now present the proof of Lemma 2.2.

Proof. Note that the Bernoulli settings for XOS, subadditive, and binary-XOS objectives described in Definition 2.6, as well as the Bernoulli setting for symmetric norm objectives in Definition 5.1, can both be understood as cases where the random variables X_i take values in the set $\{0, c_i\}$ for some fixed constant c_i .

By applying Lemma A.2, either the adaptivity gap is at most $2(1 + 1/\lambda)$, or we are in the case where

$$\mathbb{E}[f(Z_{P,\lambda})] \geq \mathbb{E}[f(X_P)] - \mathbb{E}[f(Y_{P,\lambda})] \geq \text{OPT} - \text{OPT}/2 = \text{OPT}/2,$$

which follows from the subadditivity of f .

In this second case, we analyze the adaptivity gap under the modified random variables $Z_{i,\lambda}$. Since for each coordinate $X_P \geq Z_{P,\lambda}$, the best non-adaptive strategy on $Z_{P,\lambda}$ cannot outperform the best non-adaptive strategy on X_P . Therefore, the adaptivity gap for the $Z_{i,\lambda}$ variables is at most twice the original gap.

Moreover, since the optimal reward is at least $\mathbb{E}[f(Z_{P,\lambda})] \geq \text{OPT}/2$, the random variables $Z_{P,\lambda}$ can be considered 2λ -small. The final result then follows by substituting λ with 2λ . \square

B Missing Proof of Lemma 4.4

This section provides the whole proof for Lemma 4.4. Recall the function $g(h, p) = p \cdot \exp(-0.1hp^{\frac{1}{h}})$ where $h \in \mathbb{N}_+$, and $p \in [0, 1]$. Additionally, we define $g(0, p) = p$, where $p \in [0, 1]$. Eq. (4.1) and Eq. (4.2) shows that g is increasing and concave on $[0, 1]$ when fixing $h \in \mathbb{N}$.

We first prove the following technical inequality lemma.

Lemma B.1. *For any reals $p, q, \lambda \in [0, 1]$, positive integer $h \geq 1$, the following inequality holds:*

$$g(h, \lambda p + (1 - \lambda)q) \geq (1 - \lambda)[\lambda g(h - 1, p) + (1 - \lambda)g(h, q)]. \quad (\text{B.1})$$

Proof. The idea is that, first we use concavity of function $g(h, \cdot)$ as well as the slackness in the coefficient (say, the coefficient $(1 - \lambda)$ outside the squared brackets in Eq. (B.1)) to rule out most of the non-tight case. The remaining possibility indicates that λ is relatively small, and p is larger than a constant multiple of q . In such case, we directly use the derivative of $g(h, \cdot)$ to approximate $g(h, \lambda p + (1 - \lambda)q) - g(h, q)$. In either case, we need to upper bound $g(h - 1, p)$ by $g(h, \cdot)$ using the so-called Young's inequality bellow:

Fact 1 (Young's Inequality, see [You12]). *If $a, b \geq 0$ are non-negative real numbers, and $x, y > 1$ are real numbers such that $\frac{1}{x} + \frac{1}{y} = 1$, then $ab \leq a^x/x + b^y/y$.*

The inequality can be equivalently stated in another form, by substituting $x = \frac{u+v}{u}$, $y = \frac{u+v}{v}$, and $a \leftarrow a^{u+v}$, $b \leftarrow b^{u+v}$: if $a, b \geq 0$, and $u, v > 0$, then $(u + v)(ab)^{\frac{1}{u+v}} \leq ua^{\frac{1}{u}} + vb^{\frac{1}{v}}$.

We may also use the following two fundamental inequalities related to the natural logarithm e .

Fact 2. $e^x \geq x + 1, \forall x \in \mathbb{R}$. $(1 + \frac{1}{x})^x \leq e, \forall x \in \mathbb{R}_+$.

Now we treat the following two cases respectively, as mentioned ahead.

Case 1. $(0.2 - 1.2\lambda)p \leq (1 - \lambda)q$.

By Young's inequality (Fact 1), $(h - 1)p^{\frac{1}{h-1}} + 1 \geq hp^{\frac{1}{h}}$ for $h \geq 2$. Then, for such $h \geq 2$

$$g(h - 1, p) = p \exp(-0.1(h - 1)p^{\frac{1}{h-1}}) \leq p \exp(-0.1hp^{\frac{1}{h}} + 0.1) \leq 1.2 \exp(-0.1hp^{\frac{1}{h}}) = 1.2g(h, p).$$

Note that the above also holds for $h = 1$, as the first inequality $0 \leq 0.1 - 0.1p$ still holds for $p \in [0, 1]$.

Now we utilize the property that $g(h, \cdot)$ is a concave function, and $g(h, 0) = 0$ for $h \in \mathbb{N}_+$. Then the following holds for any real $a, b \geq 0$ satisfying $a + b \leq 1$:

$$ag(h, p) + bg(h, q) = ag(h, p) + bg(h, q) + (1 - a - b)g(h, 0) \leq g(h, ap + bq + (1 - a - b) \cdot 0) = g(h, ap + bq).$$

Taking $a = 1.2\lambda(1 - \lambda)$, $b = (1 - \lambda)^2$, $a + b = (1 - \lambda)(1 + 0.2\lambda) \leq (1 - \lambda)(1 + \lambda) \leq 1$ in the above inequality,

$$\begin{aligned} (1 - \lambda)[\lambda g(h - 1, p) + (1 - \lambda)g(h, q)] &\leq 1.2\lambda(1 - \lambda)g(h, p) + (1 - \lambda)^2g(h, q) \\ &\leq g(h, 1.2\lambda(1 - \lambda)p + (1 - \lambda)^2q) \\ &= g(h, \lambda p + (1 - \lambda)q + \lambda[(0.2 - 1.2\lambda)p - (1 - \lambda)q]) \\ &\leq g(h, \lambda p + (1 - \lambda)q), \end{aligned}$$

where the last inequality uses the monotonicity of $g(h, \cdot)$ and the condition given in this case.

Case 2. $(0.2 - 1.2\lambda)p > (1 - \lambda)q$.

Note that the condition already implies $0.2 - 1.2\lambda > 0$, or $\lambda < \frac{1}{6}$, and $p > \frac{1 - \lambda}{0.2 - 1.2\lambda}q \geq 5q \geq 0$. We see that $\lambda p + (1 - \lambda)q \geq (1 + 4\lambda)q \geq q$, as well as $\lambda p + (1 - \lambda)q < (0.2 - 0.2\lambda)p \leq 0.2p$. Putting together,

$$q \leq \lambda p + (1 - \lambda)q < 0.2p. \quad (\text{B.2})$$

Since $g(h, \cdot)$ is concave when we fix h , the partial derivative $\frac{\partial g}{\partial p}(h, \cdot)$ with respect to the second variable is monotonically decreasing. In the following, we use $g'(\cdot)$ to denote the partial derivative $\frac{\partial g}{\partial p}(h, \cdot)$. Therefore,

$$g(h, \lambda p + (1 - \lambda)q) - g(h, q) = \int_q^{\lambda p + (1 - \lambda)q} g'(h, x) dx \geq \lambda(p - q)g'(h, \lambda p + (1 - \lambda)q).$$

Thus, by dividing λ in the target inequality Eq. (B.1), we only need to prove that

$$(p - q)g'(h, \lambda p + (1 - \lambda)q) + (2 - \lambda)g(h, q) \geq (1 - \lambda)g(h - 1, p).$$

For clarity, we denote $\lambda p + (1 - \lambda)q = r$. Using the derivative formula in Eq. (4.1), we have

$$g'(h, r) = \exp(-0.1hr^{\frac{1}{h}})(1 - 0.1r^{\frac{1}{h}}) \geq \exp(-0.1hr^{\frac{1}{h}}) \exp(-0.2r^{\frac{1}{h}}) = \exp(-0.1(h + 2)r^{\frac{1}{h}}),$$

where the first inequality uses a small inequality that $1 - x \geq \exp(-2x)$, when $x = 0.1r^{\frac{1}{h}} \in [0, 0.1]$. One can check that it indeed holds since $e^{2x} \geq 1 + 2x \geq \frac{1}{1-x}$, for $x \leq \frac{1}{2}$. Therefore, the target inequality can be relaxed further to be

$$(p - q) \exp(-0.1(h + 2)r^{\frac{1}{h}}) + (2 - \lambda)g(h, q) \geq (1 - \lambda)g(h - 1, p). \quad (\text{B.3})$$

Now we bound the right-hand side. Similar as in the previous case, we use Young's inequality (Fact 1) and obtain for $h \geq 2$ (using the implicit fact that $p > 0$ mentioned in the first line in the current case),

$$(h - 1)p^{\frac{1}{h-1}} + e^2 \frac{r}{p} \geq (h - 1)p^{\frac{1}{h-1}} + \left(\frac{h + 2}{h}\right)^h \frac{r}{p} \geq hp^{\frac{1}{h}} \times \frac{h + 2}{h} \left(\frac{r}{p}\right)^{\frac{1}{h}} = (h + 2)r^{\frac{1}{h}},$$

where the first inequality is given by the second inequality in Fact 2 together with the substitution $x = \frac{h}{2}$.

In the following, we still need another small inequality that $\exp(0.1e^2x) \leq 1 + x, \forall x \leq 0.2$. One can also check it indeed holds since $\exp(-0.1e^2x) \geq \exp(-0.8x) \geq 1 - 0.8x \geq \frac{1}{1+x}$, for $x \leq 0.2$. Therefore, by Eq. (B.2), taking $x = \frac{r}{p} = \frac{\lambda p + (1 - \lambda)q}{p} < 0.2$ in the above inequality, we obtain for $h \geq 2$:

$$\begin{aligned} g(h - 1, p) &= p \exp(-0.1(h - 1)p^{\frac{1}{h-1}}) \leq p \exp\left(-0.1(h + 2)r^{\frac{1}{h}} + 0.1e^2 \frac{r}{p}\right) \\ &\leq p \exp\left(-0.1(h + 2)r^{\frac{1}{h}}\right) \left(1 + \frac{r}{p}\right) \\ &= ((1 + \lambda)p + (1 - \lambda)q) \exp\left(-0.1(h + 2)r^{\frac{1}{h}}\right). \end{aligned}$$

Note that the above also holds for $h = 1$, as the first inequality $0 \leq -0.3r + 0.1e^2 \frac{r}{p} = (0.1e^2 - 0.3p) \frac{r}{p}$ still holds for $p \in [0, 1]$. Thus, in either case, the right-hand side of Eq. (B.3) can be upper bounded as:

$$(1 - \lambda)g(h - 1, p) \leq ((1 - \lambda^2)p + (1 - \lambda)^2q) \exp\left(-0.1(h + 2)r^{\frac{1}{h}}\right) \leq (p + (1 - \lambda)q) \exp\left(-0.1(h + 2)r^{\frac{1}{h}}\right).$$

Plugging in the upper bound above in Eq. (B.3) and dividing $2 - \lambda$ in both sides, we are left to prove

$$g(h, q) \geq q \exp\left(-0.1(h + 2)r^{\frac{1}{h}}\right).$$

By expressing $g(h, q)$ by definition and using the monotonicity of $\exp(\cdot)$, such inequality is equivalent to $hq^{\frac{1}{h}} \leq (h + 2)r^{\frac{1}{h}}$. But this is obvious, as we've already mentioned in Eq. (B.2) that $r = \lambda p + (1 - \lambda)q \geq q$. Thus, $hq^{\frac{1}{h}} \leq hr^{\frac{1}{h}} \leq (h + 2)r^{\frac{1}{h}}$, as desired. \square

Now we prove Lemma 4.4. Recall the setting as follows: S is a subset of leaves in the tree \mathcal{T} . There is a subset T of nodes in \mathcal{T} , not containing any leaf of \mathcal{T} . There exists an integer $h \in \mathbb{N}$, such that for each $\ell \in S$, $|A_\ell \cap T| \geq h$. Moreover, each $\ell \in S$ is assigned a real weight $x_\ell \in [0, 1]$, we denote $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}}[x_\ell \mathbf{1}_{\ell \in S}] = p$. We need to prove the following inequality:

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[x_\ell \mathbf{1}_{\ell \in S} \cdot \Pr_{R \sim \mathbb{P}_R} [R \cap P_\ell \cap T = \emptyset] \right] \leq g(h, p).$$

Proof of Lemma 4.4. First, the result is trivially true for $h = 0$, since

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[x_{\ell} \mathbf{1}_{\ell \in S} \cdot \Pr_{R \sim \mathbb{P}_R} [R \cap P_{\ell} \cap T = \emptyset] \right] \leq \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [x_{\ell} \mathbf{1}_{\ell \in S}] = p = g(0, p).$$

Therefore, we may assume $h \geq 1$ for the rest of the proof. We proceed by induction on the height of \mathcal{T} .

When the height of \mathcal{T} is 1, then h can only be 0, since T does not contain any leaf. Hence, the base case is already covered. Assume the statement holds for trees of height less than that of \mathcal{T} . Let the root of \mathcal{T} be r_0 . Let \mathcal{T}_1 and \mathcal{T}_2 be the subtrees rooted at the **Yes** and **No** children of r_0 , respectively. Denote the probability for the root to be active is $\lambda \in [0, 1]$ (note that we can also handle the case where r_0 has only one child by setting $\lambda = 0$ or 1). It is clear that the path distribution $\pi_{\mathcal{T}}$ is λ probability to **Yes** subtree \mathcal{T}_1 and $1 - \lambda$ to **No** subtree \mathcal{T}_2 , then preserves the distribution given by $\pi_{\mathcal{T}_1}$ and $\pi_{\mathcal{T}_2}$. Denote $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_1}} [x_{\ell} \mathbf{1}_{\ell \in S}] = u$, and $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_2}} [x_{\ell} \mathbf{1}_{\ell \in S}] = v$. Then $\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} [x_{\ell} \mathbf{1}_{\ell \in S}] = p = \lambda u + (1 - \lambda)v$.

Now we treat the following two cases respectively, as in Lemma 3.9.

Let $E_1 := \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_1}} [x_{\ell} \mathbf{1}_{\ell \in S} \cdot \Pr_{R \sim \mathbb{P}_R} [R \cap P_{\ell} \cap T = \emptyset]]$, $E_2 := \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_2}} [x_{\ell} \mathbf{1}_{\ell \in S} \cdot \Pr_{R \sim \mathbb{P}_R} [R \cap P_{\ell} \cap T = \emptyset]]$.

Case 1. $r_0 \in T$.

In this case, for leaves ℓ in $S \cap \mathcal{T}_1$, there are at least $h - 1$ active ancestors of it inside $T \cap \mathcal{T}_1$. Moreover, we denote the corresponding root-leaf path in \mathcal{T}_1 or \mathcal{T}_2 as $P'_{\ell} = P_{\ell} \setminus \{r_0\}$, then the corresponding relation $\Pr_{R \sim \mathbb{P}_R} [R \cap P_{\ell} \cap T = \emptyset] = (1 - \lambda) \Pr_{R \sim \mathbb{P}_R} [R \cap P'_{\ell} \cap T = \emptyset]$ holds, as the root r_0 has an independent probability of $1 - \lambda$ to be active. Therefore, by the induction hypothesis on \mathcal{T}_1 , we obtain:

$$E_1 = (1 - \lambda) \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_1}} \left[x_{\ell} \mathbf{1}_{\ell \in S} \cdot \Pr_{R \sim \mathbb{P}_R} [R \cap P'_{\ell} \cap T = \emptyset] \right] \leq (1 - \lambda)g(h - 1, u).$$

Similarly, for the **No** subtree, each leaf has at least h active ancestors inside $T \cap \mathcal{T}_2$, since the root is not an active ancestor for leaves in the **No** subtree. Hence,

$$E_2 = (1 - \lambda) \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}_2}} \left[x_{\ell} \mathbf{1}_{\ell \in S} \cdot \Pr_{R \sim \mathbb{P}_R} [R \cap P_{\ell} \cap T = \emptyset] \right] \leq (1 - \lambda)g(h, v).$$

Putting together,

$$\mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[x_{\ell} \mathbf{1}_{\ell \in S} \cdot \Pr_{R \sim \mathbb{P}_R} [R \cap P_{\ell} \cap T = \emptyset] \right] = \lambda E_1 + (1 - \lambda)E_2 \leq (1 - \lambda)[\lambda g(h - 1, u) + (1 - \lambda)g(h, v)].$$

Since $h \geq 1$, and $u, v, \lambda \in [0, 1]$ being expectation of a random variable in $[0, 1]$, we use Lemma B.1 and concludes:

$$(1 - \lambda)[\lambda g(h - 1, u) + (1 - \lambda)g(h, v)] \leq g(h, \lambda u + (1 - \lambda)v) = g(h, p).$$

Case 2. $r_0 \notin T$.

In this case, for leaves ℓ in both subtrees $S \cap \mathcal{T}_1$ and $S \cap \mathcal{T}_2$, there are at least h active ancestors of it inside $T \cap \mathcal{T}_1$ and $T \cap \mathcal{T}_2$, since $r_0 \notin T$. Moreover, we denote the corresponding root-leaf path in \mathcal{T}_1 or \mathcal{T}_2 as $P'_{\ell} = P_{\ell} \setminus \{r_0\}$, then the corresponding relation $\Pr_{R \sim \mathbb{P}_R} [R \cap P_{\ell} \cap T = \emptyset] = \Pr_{R \sim \mathbb{P}_R} [R \cap P'_{\ell} \cap T = \emptyset]$ holds, as the root $r_0 \notin T$. Therefore, by the induction hypothesis on \mathcal{T}_1 and \mathcal{T}_2 , we obtain $E_1 \leq g(h, u)$ and $E_2 \leq g(h, v)$. Putting together,

$$\begin{aligned} \mathbb{E}_{\ell \leftarrow \pi_{\mathcal{T}}} \left[x_{\ell} \mathbf{1}_{\ell \in S} \cdot \Pr_{R \sim \mathbb{P}_R} [R \cap P_{\ell} \cap T = \emptyset] \right] &= \lambda E_1 + (1 - \lambda)E_2 \leq \lambda g(h, u) + (1 - \lambda)g(h, v) \\ &\leq g(h, \lambda u + (1 - \lambda)v) = g(h, p), \end{aligned}$$

where the last inequality uses the concavity of $g(h, \cdot)$ as in Eq. (4.2). \square

C Missing Proof for Section 6

Recall that n represents the size of the ground set, and r is an upper bound on the length of a sequence in \mathcal{F} . We first present the proof of Theorem 6.1, the idea has already appeared implicitly in [GNS17].

Proof of Theorem 6.1. We consider the Bernoulli setting with a subadditive objective function $f : 2^U \rightarrow \mathbb{R}_{\geq 0}$. It is a known result that any monotone and normalized ($f(\emptyset) = 0$) subadditive set function can be approximated within a factor of $\log |U| = O(\log n)$ by an XOS function [DFF21, BR11]. Consequently, the adaptivity gap for *Bernoulli Stochastic Probing with Subadditive Objective* is $O(\Lambda_{\text{xos}}(n, r) \log n)$. \square

For the remaining two theorems, both settings require that the objective f is a monotone norm. At the same time, we consider general random variables. We define $f(x_i) = f(0, 0, \dots, 0, x_i, 0, \dots, 0)$ and likewise treat $f(X_i)$ as the corresponding random variable. Since f is a norm, we can assume that $f(0, 0, \dots, 0, 1, 0, \dots, 0) > 0$, since otherwise $f(x_i) = 0$ for any x_i . Then, we can remove the element from U and modify the norm accordingly by inserting a zero placeholder at the corresponding position.

We first establish the following lemma.

Lemma C.1. *Suppose the adaptivity gap for general random variables X_i and a monotone norm objective f under a feasibility constraint \mathcal{F} is $\Lambda \geq 1$, provided that each X_i takes value either 0 or in a geometric progression of ratio 2 and length at most $\log 4r$. Then, the adaptivity gap for general random variables with the same f and \mathcal{F} is at most 8Λ .*

Proof. By appropriately scaling the objective f , we may assume without loss of generality that the optimal adaptive reward is $\text{OPT} = 1$ in both cases. As explained in Section 1.3.1, we apply the 1-large decomposition introduced in Definition A.2: $X_P = Y_{P,1} + Z_{P,1}$. Under the optimal adaptive strategy. If $\mathbb{E}[f(Y_{P,1})] \geq 1/2$, Lemma A.2 implies an adaptivity gap of at most 4. Otherwise, we consider the 1-small random variables $Z_{P,1}$. Since the original optimal strategy ensures $\mathbb{E}[f(Z_{P,1})] \geq 1/2$, and the non-adaptive strategy cannot exceed this value due to monotonicity, the adaptivity gap for the original variables is at most twice that for $Z_{P,1}$. Reusing the notation, we assume that the current random variables X_i satisfy $f(X_i) \leq 1$.

We can further simplify X_i by setting $H_i = X_i$ if $f(X_i) \leq 1/(4r)$ and $H_i = 0$ otherwise. This gives $\mathbb{E}[f(H_P)] \leq r \cdot 1/(4r) \leq 1/4$. Thus, replacing X_i with H_i reduces the expected adaptive reward by at most $1/4$, and the non-adaptive reward does not increase. So, the adaptivity gap increases by at most a factor of 2. Now, all $f(X_i)$ are either zero or within $[1/(4r), 1]$, and the optimal reward is at least $1/4$.

Furthermore, for any possible outcome x_i of X_i , we scale x_i by a suitable constant $\theta \in [1/2, 1]$ to make $f(\theta x_i) = \theta f(x_i)$ the maximum negative power of 2 strictly below the original $f(x_i)$. Now, each $f(X_i)$ can take at most $\log 8r$ possible values, namely 0 and negative powers of 2 contained in the interval $[1/(4r), 1/2]$. Due to the homogeneity of the norm f , it follows that X_i itself assumes 0 or values in a geometric progression of ratio 2 and length at most $\log 4r$ (note that we are using $f(0, 0, \dots, 0, 1, 0, \dots, 0) > 0$). Again, the adaptive reward decreases by at most a factor of 2 by monotonicity, and the non-adaptive reward does not increase. Thus, the adaptivity gap increases by another factor of 2.

Combining the three reductions, the adaptivity gap increases by at most a factor of 8. \square

By incurring only a constant factor loss in the approximation, we may now assume that all X_i satisfy the condition of Lemma C.1. We can represent the optimal adaptive strategy using a decision tree \mathcal{T} , where each node corresponds to an element with at most $\log 8r$ children, each representing an outcome of X_i .

For each i , let X_i take values $0 = c_{i,0} < c_{i,1} < \dots < c_{i,k_i}$ ($c_{i,j}$, $1 \leq j \leq k_i$ forms a geometric progression of ratio 2) with probabilities $p_{i,0}, \dots, p_{i,k_i}$ (summing to 1, and $p_{i,0}$ can be zero). Following the approach in [Sin18], we introduce k_i new elements $e_{i,1}, \dots, e_{i,k_i}$. For $1 \leq j \leq k_i$, we associate $e_{i,j}$ with a Bernoulli random variable $R_{i,j}$ with activation probability $\frac{p_{i,j}}{\sum_{h=0}^j p_{i,h}}$. It takes value in $0, 1$ in the proof of Theorem 6.2, and $c_{i,j}$ in the proof of Theorem 6.3.

We construct a new decision tree \mathcal{T}' using these Bernoulli variables. At each node u with $\text{elt}(u) = i$ and subtrees T_0, \dots, T_{k_i} , we replace u with k_i nodes that represent elements $e_{i,k_i}, \dots, e_{i,1}$ respectively. For each j , the node represents $e_{i,j}$ proceeds to T_j if active or to the node represents $e_{i,j-1}$ otherwise. If none of $e_{i,j}$ is active, the path leads to T_0 . Notice that the replacement is from root to leaf, so the replacement procedure finally terminates. An example is illustrated in Fig. 7.




Figure 7: Illustration of How to Change the Tree

We can define a Bernoulli stochastic probing instance according to this decision tree. The elements are all $e_{i,j}$, with Bernoulli random variables $R_{i,j}$. In addition, we define the new feasibility constraint \mathcal{F}' to contain all prefixes of element sequences that correspond to a root-leaf path for \mathcal{T}' . The new functions f are defined in the proof sections of the two theorems, as they differ slightly in their definitions.

The following lemma demonstrates that the leaf distributions according to the two trees \mathcal{T} and \mathcal{T}' are essentially the same:

Lemma C.2. *For each $1 \leq i \leq n$ and $1 \leq j \leq k_i$, the probability that $e_{i,j}$ is active while all $e_{i,j'}$ with $j' > j$ are inactive is exactly $p_{i,j}$.*

Proof. This follows from direct computation of probability:

$$\begin{aligned}
& \frac{p_{i,j}}{p_{i,0} + p_{i,1} + \cdots + p_{i,j}} \prod_{j'=j+1}^{k_i} \left(1 - \frac{p_{i,j'}}{p_{i,0} + p_{i,1} + \cdots + p_{i,j'}} \right) \\
&= \frac{p_{i,j}}{p_{i,0} + p_{i,1} + \cdots + p_{i,j}} \prod_{j'=j+1}^{k_i} \frac{p_{i,0} + p_{i,1} + \cdots + p_{i,j'-1}}{p_{i,0} + p_{i,1} + \cdots + p_{i,j'}} \\
&= \frac{p_{i,j}}{p_{i,0} + p_{i,1} + \cdots + p_{i,k_i}} = p_{i,j}.
\end{aligned}$$

□

Note that we have defined a new tree \mathcal{T}' , a new feasibility constraint \mathcal{F}' , new elements $e_{i,j}$ and their associated Bernoulli random variables $R_{i,j}$. The current number of elements $e_{i,j}$ is at most $n \log 8r$, and the maximal length of a sequence in the feasibility constraint \mathcal{F}' is at most the height of the new decision tree, which is clearly bounded by $r \log 8r$. Let U' be the set of all $e_{i,j}$.

Now, based on the original objective f , we should define a new objective f' in the new setting. We show that the adaptivity gap for the new instances is at least that of the original instance.

First, we show the construction for the monotone norm and prove Theorem 6.2.

Proof of Theorem 6.2. We now define an objective $f': \mathbb{R}_{\geq 0}^{U'} \rightarrow \mathbb{R}_{\geq 0}$:

$$f'((x_{ij})_{i \in U, j \in [k_i]}) = f\left(\left(\max_{j \in [k_i]} c_{i,j} x_{i,j}\right)_{i \in U}\right).$$

Clearly, f' is a monotone norm on $\mathbb{R}_{\geq 0}^{U'}$. Thus, its restriction to a set function (as Definition 2.6 does) is fractionally subadditive by Jensen's inequality and therefore can be written as an XOS (set) function, see [Fei06].

Now, we first analyze the optimal reward for the new problem. Clearly, a valid adaptive strategy is to probe elements according to the probed result, following the decision tree \mathcal{T}' . Given a path P' for \mathcal{T}' and all the realized R_{ij} along the path, the reward $f'(R_{P'})$ is equal to the value $f(X_P)$, where P is a root-leaf path in the original tree \mathcal{T} that corresponds to P' under the transformation. By Lemma C.2, the leaf distribution of \mathcal{T} and \mathcal{T}' is the same. Thus, we conclude that the expected reward following \mathcal{T}' with objective f' is the same as the original optimal reward. Therefore, the *optimal* adaptive reward is at least the original.

For the non-adaptive strategy in the new Bernoulli instance, it is clear that the feasibility constraint \mathcal{F}' requires the algorithm to probe a prefix of any root-leaf path (in each branch). Therefore, the best non-adaptive strategy must correspond to a root-leaf path P' in \mathcal{T} (although in general, non-adaptive strategy, such as ALGPATH is randomized, a standard averaging argument shows that there must exist an optimal deterministic non-adaptive strategy). We compare the non-adaptive reward for P' to that of P , the path in the original tree \mathcal{T} that corresponds to P' under the transformation. We prove that for each coordinate i , the random variable $\max_{t \leq j \leq k_i} c_{i,j} R_{i,j}$ is dominated¹ by the original variable X_i , where t means that the path P' leaves $e_{i,t}$ and goes to T_t . It is clear that $\max_{t \leq j \leq k_i} c_{i,j} R_{i,j}$ takes the value $c_{i,j}$ if and only if $R_{i,j} = 1$ and $R_{i,j}$ for larger j are not active. By Lemma C.2, the probability is exactly $p_{i,j}$, the same as the random variable X_i , for $t \leq j \leq k_i$. For the remaining probability $\sum_{j=0}^{t-1} p_{i,j-1}$, the random variable $\max_{t \leq j \leq k_i} c_{i,j} R_{i,j}$ takes zero, while $X_i \geq 0$, hence the assertion of dominance holds.

Therefore, since dominated (multidimensional) random variables have a smaller f value for monotone norm f , we conclude that the best non-adaptive strategy in the new instance is at most that of the original instance. Therefore, the adaptivity gap of the worst case for Bernoulli setting is at least the adaptivity gap for the underlying instance. Hence, the adaptivity gap is at most $\Lambda_{\text{XOS}}(n \log 8r, r \log 8r)$. \square

Finally, we prove Theorem 6.3.

Proof of Theorem 6.3. We now define an objective $g' : \mathbb{R}_{\geq 0}^{U'} \rightarrow \mathbb{R}_{\geq 0}$ as follows:

$$g'((x_{ij})_{i \in U, j \in [k_i]}) = f(\text{Top}_n(x_{ij})),$$

where Top_n stands for the n largest values (break tie arbitrarily) in x_{ij} . Clearly, g' is a monotone symmetric norm. The same proof as above shows that the expected gain following \mathcal{T}' with objective g' is the same as the original optimal reward. Therefore, the *optimal* adaptive reward is at least the original.

For the non-adaptive strategy in the new instance, we use the norm f' in the proof of Theorem 6.2:

$$f'((x_{ij})_{i \in U, j \in [k_i]}) = f\left(\left(\max_{j \in [k_i]} x_{i,j}\right)_{i \in U}\right).$$

We consider P , the path in the original tree \mathcal{T} that corresponds to P' under the transformation. The same proof shows that the non-adaptive reward for P with norm f (note that the random variables R_{ij} take value $\{0, c_{i,j}\}$, so we change f' correspondingly) is at least that for P' with norm f' . We now prove that the non-adaptive reward for P' with norm f' is at least half the same path with norm g' . Indeed, it suffices to prove for any input $x_{i,j} \in \{0, c_{i,j}\}$, $2f'(x_{ij}) \geq g'(x_{ij})$. Among the n largest elements in x_{ij} , we assume that for each i , $x_{i,t_i} = c_{i,t_i}$ is taken, where $t_i \in [0, k_i]$ is the largest possible (if no $x_{i,j}$ appears in the n largest, or its value is zero, set $t_i = 0$). Then, by definition of f' , we have $f'(x_{ij}) \geq f(x_{i,t_i}) = f(c_{i,t_i})$. On the other hand, we can partition $x_{i,j}$ appearing in $\text{Top}_n(x_{ij})$ for each i with respect to the distance $t_i - j$. Therefore, by symmetry, subadditivity and monotonicity of f (treat $x_{i,t_i-j} = 0$ if $t_i - j \leq 0$),

$$g'(\text{Top}_n(x_{ij})) \leq \sum_{j=0}^{\infty} f(x_{i,t_i-j}) \leq \sum_{j=0}^{\infty} f(c_{i,t_i-j}) \leq \sum_{j=0}^{\infty} 2^{-j} f(c_{i,t_i}) = 2f(c_{i,t_i}) \leq 2f'(x_{ij}).$$

Therefore, the adaptivity gap of the worst case for Bernoulli setting is at least half the adaptivity gap for the underlying instance. Hence, the adaptivity gap for general random variables is at most $2\Lambda_{\text{sym}}(n \log 8n)$, using $r = \max_{F \in \mathcal{F}} \{2, \max |F|\} \leq n$ in Lemma C.1. \square

¹A (non-negative) random variable X is dominated by another random variable Y if and only if for any $z \geq 0$, $\Pr[X \geq z] \leq \Pr[Y \geq z]$.

References

- [AAK19] Arpit Agarwal, Sepehr Assadi, and Sanjeev Khanna. Stochastic submodular cover with limited adaptivity. In *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 323–342. SIAM, 2019.
- [ABB⁺23] Fateme Abbasi, Sandip Banerjee, Jaroslaw Byrka, Parinya Halermsook, Ameet Gaddekar, Kamyar Khodamoradi, Daniel Marx, Roohani Sharma, and Joachim Spoerhase. Parameterized approximation schemes for clustering with general norm objectives. In *2023 IEEE 64th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 1377–1399. IEEE, 2023.
- [ABP23] Nikhil Ayyadevara, Nikhil Bansal, and Milind Prabhu. On minimizing generalized makespan on unrelated machines. *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*, page 21, 2023.
- [Ada11] Marek Adamczyk. Improved analysis of the greedy algorithm for stochastic matching. *Information Processing Letters*, 111(15):731–737, 2011.
- [AN16] Arash Asadpour and Hamid Nazerzadeh. Maximizing stochastic monotone submodular functions. *Management Science*, 62(8):2374–2391, 2016.
- [AR12] Itai Ashlagi and Alvin E. Roth. New challenges in multihospital kidney exchange. *American Economic Review*, 102(3):354–59, May 2012.
- [BDF⁺12] Ashwinkumar Badanidiyuru, Shahar Dobzinski, Hu Fu, Robert Kleinberg, Noam Nisan, and Tim Roughgarden. Sketching valuation functions. In *Proceedings of the Twenty-Third Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA ’12, page 1025–1035, USA, 2012. Society for Industrial and Applied Mathematics.
- [BGL⁺12] Nikhil Bansal, Anupam Gupta, Jian Li, Julián Mestre, Viswanath Nagarajan, and Atri Rudra. When lp is the cure for your matching woes: Improved bounds for stochastic matchings. *Algorithmica*, 63(4):733–762, August 2012.
- [BH11] Maria-Florina Balcan and Nicholas J.A. Harvey. Learning submodular functions. In *Proceedings of the Forty-Third Annual ACM Symposium on Theory of Computing*, STOC ’11, page 793–802, New York, NY, USA, 2011. Association for Computing Machinery.
- [BN14] Nikhil Bansal and Viswanath Nagarajan. On the adaptivity gap of stochastic orienteering. In Jon Lee and Jens Vygen, editors, *Integer Programming and Combinatorial Optimization*, pages 114–125, Cham, 2014. Springer International Publishing.
- [BR11] Kshipra Bhawalkar and Tim Roughgarden. Welfare guarantees for combinatorial auctions with item bidding. In *Proceedings of the Twenty-Second Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA ’11, page 700–709, USA, 2011. Society for Industrial and Applied Mathematics.
- [BSS18] Jaroslaw Byrka, Krzysztof Sornat, and Joachim Spoerhase. Constant-factor approximation for ordered k -median. In *Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing*, pages 620–631, 2018.
- [BSZ19] Domagoj Bradac, Sahil Singla, and Goran Zuzic. (near) optimal adaptivity gaps for stochastic multi-value probing. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques, APPROX/RANDOM 2019*, Leibniz International Proceedings in Informatics, LIPIcs, September 2019.
- [BV22] Siddharth Barman and Paritosh Verma. Approximating nash social welfare under binary xos and binary subadditive valuations. In Michal Feldman, Hu Fu, and Inbal Talgam-Cohen, editors, *Web and Internet Economics*, pages 373–390, Cham, 2022. Springer International Publishing.

- [CCPV11] Gruia Calinescu, Chandra Chekuri, Martin Pál, and Jan Vondrák. Maximizing a monotone submodular function subject to a matroid constraint. *SIAM Journal on Computing*, 40(6):1740–1766, 2011.
- [CHL⁺16] Wei Chen, Wei Hu, Fu Li, Jian Li, Yu Liu, and Pinyan Lu. Combinatorial multi-armed bandit with general reward functions. *Advances in Neural Information Processing Systems*, 29, 2016.
- [CIK⁺09] Ning Chen, Nicole Immorlica, Anna R. Karlin, Mohammad Mahdian, and Atri Rudra. Approximating matches made in heaven. In *Proceedings of the 36th International Colloquium on Automata, Languages and Programming: Part I, ICALP '09*, page 266–278, Berlin, Heidelberg, 2009. Springer-Verlag.
- [CS18] Deeparnab Chakrabarty and Chaitanya Swamy. Interpolating between k -median and k -center: Approximation algorithms for ordered k -median. In *45th International Colloquium on Automata, Languages, and Programming*, volume 107 of *LIPICs*, pages 29:1–29:14, 2018.
- [CS19a] Deeparnab Chakrabarty and Chaitanya Swamy. Approximation algorithms for minimum norm and ordered optimization problems. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing*, pages 126–137, 2019.
- [CS19b] Deeparnab Chakrabarty and Chaitanya Swamy. Simpler and better algorithms for minimum-norm load balancing. In *27th Annual European Symposium on Algorithms*, volume 144 of *LIPICs*, pages 27:1–27:12, 2019.
- [DFF21] Shahar Dobzinski, Uriel Feige, and Michal Feldman. Are gross substitutes a substitute for submodular valuations? In *Proceedings of the 22nd ACM Conference on Economics and Computation, EC '21*, page 390–408, New York, NY, USA, 2021. Association for Computing Machinery.
- [DGV04] Brian C Dean, Michel X Goemans, and Jan Vondrak. Approximating the stochastic knapsack problem: The benefit of adaptivity. In *Proceedings of the 45th Annual IEEE Symposium on Foundations of Computer Science*, pages 208–217, 2004.
- [DHK14] Amol Deshpande, Lisa Hellerstein, and Devorah Kletenik. Approximation algorithms for stochastic boolean function evaluation and stochastic submodular set cover. In *Proceedings of the 25th Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2014*, Proceedings of the Annual ACM-SIAM Symposium on Discrete Algorithms, pages 1453–1467. Association for Computing Machinery, 2014. 25th Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2014 ; Conference date: 05-01-2014 Through 07-01-2014.
- [DLR23] Shichuan Deng, Jian Li, and Yuval Rabani. Generalized unrelated machine scheduling problem. In *Proceedings of the 2023 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 2898–2916. SIAM, 2023.
- [Dob07] Shahar Dobzinski. Two randomized mechanisms for combinatorial auctions. In *Proceedings of the 10th International Workshop on Approximation and the 11th International Workshop on Randomization, and Combinatorial Optimization. Algorithms and Techniques, APPROX '07/RANDOM '07*, page 89–103, Berlin, Heidelberg, 2007. Springer-Verlag.
- [Fei06] Uriel Feige. On maximizing welfare when utility functions are subadditive. In *Proceedings of the Thirty-Eighth Annual ACM Symposium on Theory of Computing, STOC '06*, page 41–50, New York, NY, USA, 2006. Association for Computing Machinery.
- [FLX18] Hao Fu, Jian Li, and Pan Xu. A ptas for a class of stochastic dynamic programs. In *45th International Colloquium on Automata, Languages, and Programming, ICALP 2018*, page 56. Schloss Dagstuhl-Leibniz-Zentrum für Informatik GmbH, Dagstuhl Publishing, 2018.

- [GKNR12] Anupam Gupta, Ravishankar Krishnaswamy, Viswanath Nagarajan, and R. Ravi. Approximation algorithms for stochastic orienteering. In *Proceedings of the Twenty-Third Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '12, page 1522–1538, USA, 2012. Society for Industrial and Applied Mathematics.
- [GM09] Sudipto Guha and Kamesh Munagala. Multi-armed bandits with metric switching costs. In *Automata, Languages and Programming*, pages 496–507, Berlin, Heidelberg, 2009. Springer Berlin Heidelberg.
- [GN13] Anupam Gupta and Viswanath Nagarajan. A stochastic probing problem with applications. In *Integer Programming and Combinatorial Optimization*, pages 205–216, Berlin, Heidelberg, 2013. Springer Berlin Heidelberg.
- [GNS16] Anupam Gupta, Viswanath Nagarajan, and Sahil Singla. Algorithms and adaptivity gaps for stochastic probing. In *Proceedings of the 2016 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1731–1747, 2016.
- [GNS17] Anupam Gupta, Viswanath Nagarajan, and Sahil Singla. Adaptivity gaps for stochastic probing: submodular and xos functions. In *Proceedings of the Twenty-Eighth Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '17, page 1688–1702, USA, 2017. Society for Industrial and Applied Mathematics.
- [GV06] Michel Goemans and Jan Vondrák. Stochastic covering and adaptivity. In *Latin American symposium on theoretical informatics*, pages 532–543. Springer, 2006.
- [IS20] Sharat Ibrahimpur and Chaitanya Swamy. Approximation algorithms for stochastic minimum-norm combinatorial optimization. In *61st IEEE Annual Symposium on Foundations of Computer Science*, pages 966–977, 2020.
- [JLLS20] Haotian Jiang, Jian Li, Daogao Liu, and Sahil Singla. Algorithms and adaptivity gaps for stochastic k-tsp. In *11th Innovations in Theoretical Computer Science Conference (ITCS 2020)*, volume 151, page 45. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2020.
- [KMS23] Thomas Kesselheim, Marco Molinaro, and Sahil Singla. Online and bandit algorithms beyond ℓ_p norms. In *Proceedings of the 2023 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, pages 1566–1593, 2023.
- [KMS24] Thomas Kesselheim, Marco Molinaro, and Sahil Singla. Supermodular approximation of norms and applications. In *Proceedings of the 56th Annual ACM Symposium on Theory of Computing*, STOC 2024, page 1841–1852, New York, NY, USA, 2024. Association for Computing Machinery.
- [LPRY08] Zhen Liu, Srinivasan Parthasarathy, Anand Ranganathan, and Hao Yang. Near-optimal algorithms for shared filter evaluation in data stream systems. In *Proceedings of the 2008 ACM SIGMOD International Conference on Management of Data*, SIGMOD '08, page 133–146, New York, NY, USA, 2008. Association for Computing Machinery.
- [Mes06] Julián Mestre. Greedy in approximation algorithms. In *Proceedings of the 14th Conference on Annual European Symposium - Volume 14*, ESA'06, page 528–539, Berlin, Heidelberg, 2006. Springer-Verlag.
- [MU05] Michael Mitzenmacher and Eli Upfal. *Probability and Computing: Randomized Algorithms and Probabilistic Analysis*. Cambridge University Press, New York, NY, USA, 2005.
- [PRS23] Kalen Patton, Matteo Russo, and Sahil Singla. Submodular Norms with Applications To Online Facility Location and Stochastic Probing. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2023)*, volume 275 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 23:1–23:22, 2023.

- [RSU05] Alvin E. Roth, Tayfun Sönmez, and M. Utku Ünver. Pairwise kidney exchange. *Journal of Economic Theory*, 125(2):151–188, 2005.
- [Sin18] Sahil Singla. The price of information in combinatorial optimization. In *Proceedings of the Twenty-Ninth Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '18, page 2523–2532, USA, 2018. Society for Industrial and Applied Mathematics.
- [You12] W. H. Young. On classes of summable functions and their fourier series. *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, 87(594):225–229, 1912.