The phase coherence of light from extragalactic sources - direct evidence against first order Planck scale fluctuations in time and space

Richard Lieu and Lloyd W. Hillman Department of Physics, University of Alabama, Huntsville, AL 35899, U.S.A.

Abstract

We present a method of directly testing whether time continues to have its usual meaning on scales of $\leq t_P = (\hbar G/c^5)^{\frac{1}{2}} \approx 5.4 \times 10^{-44}$ s, the Planck time. According to quantum gravity, the time t of an event cannot be determined more accurately than a standard deviation of the form $\sigma_t/t = a_o(t_P/t)^{\alpha}$. where a_o and α are positive constants ~ 1; likewise distances are subject to an ultimate uncertainty $c\sigma_t$, where c is the speed of light. As a consequence, the period and wavelength of light cannot be specified precisely; rather, they are independently subject to the same intrinsic limitations in our knowledge of time and space, so that even the most monochromatic plane wave must in reality be a superposition of waves with varying ω and **k**, each having a different phase velocity ω/k . For the entire accessible range of the electromagnetic spectrum this effect is extremely small, but can cumulatively lead to a complete loss of phase information if the emitted radiation propagated a sufficiently large distance. Since, at optical frequencies, the phase coherence of light from a distant point source is a necessary condition for the presence of diffraction patterns when the source is viewed through a telescope, such observations offer by far the most sensitive and uncontroversial test. We show that the HST detection of Airy rings from the active galaxy PKS1413+135, located at a distance of 1.2 Gpc, secures the exclusion of all first order ($\alpha = 1$) quantum gravity fluctuations with an amplitude $a_o > 0.003$. The same result may be used to deduce that the speed of light in vacuo is exact to a few parts in 10^{32} .

1. INTRODUCTION - THE PLANCK SCALE

It is widely believed that time ceases to be well-defined at intervals $\leq t_P$, where quantum fluctuations in the vacuum metric tensor renders General

Relativity an inadequate theory. Both t_P and its corresponding distance scale $l_P = ct_P$, the Planck length, play a vital role in the majority of theoretical models (including superstrings) that constitute innumerable papers attempting to explain how the universe was born, and how it evolved during infancy (see e.g., Silk 2000 and references therein). Given this background, we desperately lack experimental data that reveal even the slightest anomaly in the behavior of time and space at such small scales. Although the recent efforts in utilizing gravitational wave interferometry and the observation of ultra-high energy (UHE) quanta carry potential (Amelino-Camelia 2001, Ng et al 2001, Lieu 2002), they are still some way from delivering a verdict, because the conclusions are invariably subject to interpretational issues. Here we wish to describe how an entirely different yet well established technique has hither been overlooked: not only would it enable direct tests for Planck scale fluctuations (and revealing the detailed properties of any such effects). but also the measurements performed to date could already be used to eliminate prominent theories.

Owing to the variety of proposed models we begin by describing the common feature that define the phenomenon being searched: if a time t is so small that $t \to t_P$ even the best clock ever made will only be able to determine it with an uncertainty $\delta t \ge t$. To express this mathematically we may write the intrinsic standard deviation of time as $\sigma_t/t = f(t_P/t)$, where $f \ll 1$ for $t \gg t_P$ and $f \ge 1$ for $t \le t_P$. Over the range $t \gg t_P$ the (hitherto unknown) function f can be expanded as follows:

$$f(x) = x^{\alpha}(a_o + a_1x + a_2x^2 + ...) \approx a_o x^{\alpha} \text{ for } x \ll 1,$$
(1)

where $x = t_P/t$, and both a_o , α are positive constants (~ 1 for all reasonable scenarios). Since for the rest of this paper we shall be concerned only with times $t \gg t_P$ we may take an approximate form of Eq. (1) as:

$$\frac{\sigma_t}{t} \approx a_o \left(\frac{t_P}{t}\right)^{\alpha} \tag{2}$$

Our appreciation of how Eq. (2) may affect measurements of angular frequencies and wavevectors (ω, \mathbf{k}) arises from the realization that if a quantity $\omega > \omega_P = 2\pi/t_P$ can be determined accurately such a calibration will lead to a 'superclock' that keeps time to within $\delta t < t_P$. Thus as $\omega \to \omega_P$, ω should fluctuate randomly such that $\delta \omega / \omega \to 1$. Indeed for the case of $\sigma_t \approx t_P$ (i.e. Eq. (2) with $\alpha = 1$) the following Eq. was shown by Lieu (2002) to be an immediate consequence:

$$\frac{\sigma_{\omega}}{\omega} \approx a_o \frac{\omega}{\omega_P}, \text{ or } \frac{\sigma_E}{E} \approx a_o \frac{E}{E_P},$$
(3)

where $E = \hbar \omega$ and $E_P = \hbar \omega_P = h/t_P \approx 8.1 \times 10^{28}$ eV. Further, for any value of α it can be proved (see Ng & van Dam 2000) that Eq. (2) leads to:

$$\frac{\sigma_{\omega}}{\omega} \approx a_o \left(\frac{\omega}{\omega_P}\right)^{\alpha}, \text{ or } \frac{\sigma_E}{E} \approx a_o \left(\frac{E}{E_P}\right)^{\alpha}.$$
 (4)

The same reasoning also applies to the intrinsic uncertainty in data on the wavevector \mathbf{k} (note also that for measurements directly taken by an observer $\delta \omega$ and $\delta \mathbf{k}$, like δt and $\delta \mathbf{r}$, are uncorrelated errors), for if any component of \mathbf{k} could be known to high accuracy even in the limit of large p we would be able to surpass the Planck length limitation in distance determination for that direction. Thus a similar equation may then be formulated as:

$$\frac{\sigma_k}{k} \approx a_o \left(\frac{\omega}{\omega_P}\right)^{\alpha} \tag{5}$$

where k is the magnitude of **k** and the right side is identical to the previous equation because $\omega = k$ for photons. Note indeed that Eqs. (4) and (5) hold good for ultra-relativistic particles as well.

About the value of α , the straightforward choice is $\alpha = 1$, which by Eq. (2) implies $\sigma_t \sim t_P$, i.e. the most precise clock has uncertainty $\sim t_P$. Indeed, $\alpha = 1$ is just the first order term in a power series expansion of quantum loop gravity. However, the quantum nature of time at scales $\leq t_P$ may be manifested in other (more contrived) ways. In particular, for random walk models of space-time, where each step has size t_P , $\alpha = 1/2$ (Amelino-Camelia 2000). On the other hand, it was shown (Ng 2002) that as a consequence of the holographic principle (which states that the maximum degrees of freedom allowed within a region of space is given by the volume of the region in Planck units, see Wheeler 1982, Bekenstein 1973, Hawking 1975, 't Hooft 1993, Susskind 1995) σ_t takes the form $\sigma_t/t \sim (t_P/t)^{\frac{2}{3}}$, leading to $\alpha = 2/3$ in Eqs. (3) and (4). Such an undertaking also has the desirable property (Ng 2002) that it readily implies a finite lifetime τ for black holes, viz., $\tau \sim G^2 m^3/\hbar c^4$, in agreement with the earlier calculations of Hawking.

Although the choice of α is not unique, the fact that it appears as an exponent means different values can lead to wildly varying predictions. Specifically, even by taking $E = 10^{20}$ eV (i.e. the highest energy particles known, where Planck scale effects are still only $\sim (E/E_P)^{\alpha} \approx 10^{-9\alpha}$ in significance), an increment of α by 0.5 would demand a detection sensitivity 4.5 orders of magnitude higher. The situation gets much worse as E becomes lower. Thus if an experiment fails to offer confirmation at a given α , one can always raise the value of α , and the search may never end. Fortunately, however, it turns out that all of the three scenarios $\alpha = 1/2, 2/3$, and 1 may be clinched by the rapid advances in observational astronomy.

2. THE PROPAGATION OF LIGHT IN FREE SPACE

How do Eqs. (4) and (5) modify our perception of the radiation dispersion relation? By writing the relation as:

$$\omega^2 - k^2 = 0 \tag{6}$$

the answer becomes clear - one simply needs to calculate the uncertainty in $\omega^2 - k^2$ due to the intrinsic fluctuations in the measurements of ω and **k**, viz. $\delta(\omega^2 - k^2) = 2\omega\delta\omega - 2k\delta k$, bearing in mind that $\delta\omega$ and δk are independent variations, as already discussed. This allows us to obtain the standard deviation

$$\sigma_{\omega^2 - k^2} = 2\sqrt{2}\omega^2 a_o \left(\frac{\omega}{\omega_P}\right)^\alpha \tag{7}$$

Thus, typically Eq. (6) will be replaced by:

$$\omega^2 - k^2 \approx \pm 2\sqrt{2}\omega^2 a_o \left(\frac{\omega}{\omega_P}\right)^{\alpha} \tag{8}$$

In the case of a positive fluctuation on the right hand term of Eq. (6) by unit σ , the phase and group velocities of propagation will read, for $E/E_P \ll 1$, as:

$$v_p = \frac{\omega}{k} \approx 1 + \sqrt{2}a_o \left(\frac{\omega}{\omega_P}\right)^{\alpha}; \quad v_g = \frac{d\omega}{dk} \approx 1 + \sqrt{2}(1+\alpha)a_o \left(\frac{\omega}{\omega_P}\right)^{\alpha}$$
(9)

The results differ from that of a particle - here $\omega^2 - k^2$ is a function of ω and not a constant, so that both v_p and v_g are > 1, i.e. greater than the speed of light v = 1. On the other hand, if the right side of Eq. (6) fluctuates negatively the two wave velocities will read like:

$$v_p = \frac{\omega}{k} \approx 1 - \sqrt{2}a_o \left(\frac{\omega}{\omega_P}\right)^{\alpha}; \quad v_g = \frac{d\omega}{dk} \approx 1 - \sqrt{2}(1+\alpha)a_o \left(\frac{\omega}{\omega_P}\right)^{\alpha}$$
(10)

and will both be < 1.

Is it possible to force a re-interpretation of Eq. (8) in another (more conventional) way, viz., for a particular off-shell mode $\omega^2 - k^2$ typically assumes a *constant* value different from zero by about the unit σ of Eq. (7)? The point, however, is that even in this (highly artificial) approach, the quantities $v_p = \omega/k$ and $v_g = d\omega/dk = k/\omega$ will still disagree with each other randomly by an amount $\sim (\omega/\omega_P)^{\alpha}$, so that the chief outcome of Eqs. (9) and (10) is robust.

3. STELLAR INTERFEROMETRY AS AN ACCURATE TEST

But is such an effect observable? Although an obvious approach is to employ the highest energy radiation, so as to maximize ω/ω_P , such photons are difficult to detect. More familiar types of radiation, e.g. optical light, have much smaller values of ω/ω_P , yet the advantage is that we can measure their properties with great accuracy. Specifically we consider the phase behavior of 1 eV light received from a celestial optical source located at a distance Laway. During the propagation time $\Delta t = L/v_g$, the phase has advanced from its initial value ϕ (which we assume to be well-defined) by an amount:

$$\Delta \phi = 2\pi \frac{v_p \Delta t}{\lambda} = 2\pi \frac{v_p L}{v_q \lambda}$$

According to Eqs. (9) and (10), $\Delta \phi$ should then randomly fluctuate in the following manner:

$$\Delta \phi = 2\pi \frac{L}{\lambda} \left[1 \pm \sqrt{2} \alpha a_o \left(\frac{\omega}{\omega_P} \right)^{\alpha} \right] \tag{11}$$

In the limit when

$$\sqrt{2\alpha}a_o \left(\frac{\omega}{\omega_P}\right)^{\alpha} \frac{L}{\lambda} \ge 1; \text{ or } \frac{\sqrt{2\alpha}a_o}{h} E^{1+\alpha} E_P^{-\alpha} L \ge 1$$
 (12)

the phase of the wave will have appreciable probability of assuming any value between 0 and 2π upon arrival, irrespective of how sharp the initial phase at the source may be. Since a_o and α are free parameters, Eqs. (11) and (12) are a common consequence of many quantum gravity models - both equations can be derived in a variety of ways - though the approach presented here may be taken as representative. From the preceding paragraph, a way towards testing the behavior of time to the limit has become apparent. In stellar interferometry (see e.g. Baldwin & Haniff 2002 for a review) light waves from an astronomical source are incident upon two reflectors (within a terrestrial telescope) and are subsequently coverged to form Young's interference fringes. By Eq. (11), however, we see that if the time ceases to be exact at the Planck scale the phase of light from a sufficiently distant source will appear random - when L is large enough to satisfy Eq. (12) the fringes will disappear. In fact, the value of L at which Eq. (12) holds may readily be calculated for the case of $\alpha = 2/3$ and $\alpha = 1$, with the results:

$$L_{\alpha=\frac{2}{3}} \ge 2.47 \times 10^{15} a_o^{-1} (E/1 \text{ eV})^{-\frac{5}{3}} \text{ cm}; \ L_{\alpha=1} \ge 7.07 \times 10^{24} a_o^{-1} (E/1 \text{ eV})^{-2} \text{ cm}.$$
(13)

For $a_o = 1$ and E = 1 eV these distances correspond respectively to 165 AU (or 8×10^{-4} pc) and 2.3 Mpc.

4. EXAMINING THE TEST IN MORE DETAIL

Since the subject of our search is no small affair we provide here an alternative (and closer) view of the proposed experiment.

In a classical approach to the 'untilted' configuration of Young's interferometry the phase of a plane wave (from a distant source) at the position \mathbf{r} of the double slit system may be written as $\omega t - \mathbf{k} \cdot \mathbf{r}$ where t is the arrival time of the wavefront. The electric fields \mathcal{E}_1 and \mathcal{E}_2 of the waves at some point P behind the slits where they subsequently meet may then assume the form:

$$\mathcal{E}_{1} = \epsilon_{1} |\mathcal{E}_{1}| e^{i(\omega t - \mathbf{k} \cdot \mathbf{r} + \phi_{1})} ; \quad \mathcal{E}_{2} = \epsilon_{2} |\mathcal{E}_{2}| e^{i(\omega t - \mathbf{k} \cdot \mathbf{r} + \phi_{2})}, \quad (14)$$

where $|\mathcal{E}|$ denotes the modulus of the (complex) magnitude of vector \mathcal{E} , ϵ_1 and ϵ_2 are unit vectors, and ϕ_1 , ϕ_2 are the advances in the wave phase during the transits between each of the two slits and P. The intensity of light at P is proportional to $|\mathcal{E}_1 + \mathcal{E}_2|^2$, which contains the term essential to the formation of fringes, viz. $2|\mathcal{E}_1||\mathcal{E}_2|\cos\phi$, where $\phi = \phi_1 - \phi_2$ and ϵ_1 , ϵ_2 are taken to be parallel (as is commonly the case).

If, however, there exist intrinsic and independent uncertainties in one's knowledge of the period and wavelength of light on scales t_P and ct_P respectively, the most monochromatic plane wave will have to be a superposition of many waves, each having slightly varying ω and \mathbf{k} , and the phase velocity $v_p = \omega/k$ will fluctuate according to Eqs. (9) and (10). For optical frequencies the only measurable effect is the phase separation between these waves

after travelling a large distance at different speeds, i.e. Eq. (14) will be replaced by:

$$\mathcal{E}_{1} = \epsilon_{1} |\mathcal{E}_{1}| \sum_{j} a_{j} e^{i(\omega_{j}t - \mathbf{k}_{j} \cdot \mathbf{r} + \phi_{1} + \theta_{j})} \approx \epsilon_{1} |\mathcal{E}_{1}| e^{i(\omega t - \mathbf{k} \cdot \mathbf{r} + \phi_{1})} \sum_{j} a_{j} e^{i\theta_{j}},$$

and:

$$\mathcal{E}_{\mathbf{2}} = \epsilon_{\mathbf{2}} |\mathcal{E}_{2}| \sum_{l} a_{l} e^{i(\omega_{l}t - \mathbf{k}_{1} \cdot \mathbf{r} + \phi_{2} + \theta_{l})} \approx \epsilon_{\mathbf{2}} |\mathcal{E}_{2}| e^{i(\omega t - \mathbf{k} \cdot \mathbf{r} + \phi_{2})} \sum_{l} a_{l} e^{i\theta_{l}},$$

where $\{a_i\}$ are real coefficients (not to be confused with the a_o of Eq. (1)) normalized such that a_i^2 equals the occurrence probability of the *i*th wave, which of course is governed by how far θ_i differs from its zero mean value when compared with the standard deviation in Eq. (11) (viz. $2\sqrt{2\pi\omega/\omega_P}$ when $\alpha = 1$). Note that this time the intensity at P depends on many 'cross' terms, each of the form $2a_ja_l|\mathcal{E}_1||\mathcal{E}_2|cos\phi$, where now $\phi = (\phi_1 - \phi_2) + (\theta_j - \theta_l)$. If the propagation length L is large enough to satisfy Eq. (12) θ_j and θ_l , hence $\theta_j - \theta_l$, will spread over one phase cycle, so that the original term $2|\mathcal{E}_1||\mathcal{E}_2|cos(\phi_1 - \phi_2)$ will no longer be characteristic of the point P. This is the mathematical demonstration of why no appreciable fringe contrast across the detector can be expected. Obviously, the argument can readily be generalized to conclude that if Eq. (12) is fulfilled interference effects from *multiple slits* (or a single large slit as limiting case) will also disappear.

5. THE DIFFRACTION OF LIGHT FROM EXTRAGALACTIC POINT SOURCES - ABSENCE OF ANOMALOUS BEHAVIOR IN TIME AND SPACE AT THE PLANCK SCALE

Let us now consider the observations to date. The Young's type of interference effects were clearly seen at $\lambda = 2.2 \ \mu m$ ($E \approx 0.56 \ eV$) light from a source at 1.012 kpc distance, viz. the star S Ser, using the Infra-red Optical Telescope Array, which enabled a radius determination of the star (van Belle, Thompson, & Creech-Eakman 2002). When comparing with Eq. (13) we see that this result can already be used to completely exclude the $\alpha = 2/3$ model, because for such a value of α and for all reasonable values of a_o , $\Delta \phi$ carries uncertainties $\gg 2\pi$, and the light waves would not have interfered. It is also evident from Eq. (13), however, that no statement about $\alpha = 1$ can be made with the S Ser findings.

Within the context of the previous section's development, it turns out that the well recognized presence of diffraction pattern in the image of extragalactic point sources when they are viewed through the finite aperture of a telescope provides even more stringent constraints on α . Such patterns have been observed from sources located at distances much larger than 1 kpc, implying, as before, that phase coherence of light is maintained at the aperture entrance despite the contrary prediction of quantum gravity. In particular, to clinch the first order prediction $\alpha = 1$ we note that Airy rings (circular diffraction) were clearly visible at both the zeroth and first maxima in an observation of the active galaxy PKS1413+135 (L = 1.216 Gpc) by the HST at 1.6 μm wavelength (Perlman et al 2002). Referring back to Eq. (13), this means exclusion of all $\alpha = 1$ quantum gravity fluctuations that occur at an amplitude $a_o \geq 3.14 \times 10^{-3}$ (moreover the speed of light does not fluctuate fractionally by more than λ/L , or several parts in 10^{32}). To facilitate those who wish to explore the implications in full we offer the following inequality, derivable directly from Eq. (12), which the reader can use to readily find for any value of α the range of a_o still permitted by the PKS1413 result:

$$a_o < 3.14 \times 10^{-3} \frac{(10^{29})^{\alpha - 1}}{\alpha} \tag{15}$$

That Eq. (15) is highly sensitive to α has already been discussed. Two consequences immediately emerge from Eq. (15): (a) for any of the $\alpha < 1$ models to survive, they must involve ridiculously small values of a_o , (b) $\alpha > 1$ remains essentially unconstrained.

Further investigation of sources that lie beyond PKS1413 will scrutinize the $\alpha \leq 1$ scenarios more tightly than what is already a very stringent current limit. More sophisticated ways of pursuing stellar interferometry are necessary to test the cases of $\alpha > 1$, or systematic effects where the dispersion relation of Eq. (6) is modified by definite rather than randomly varying terms.

Nevertheless, the obvious test bed for quantum gravity has indeed been provided by this Gpc distance source; the outcome is negative. No doubt one anticipates interesting propositions on how time and space may have contrived to leave behind not a trace of their quanta. Thus, from Michelson-Morley to extragalactic interferometry there remains no direct experimental evidence of any sort that compels us to abandon the structureless, etherless space-time advocated by Einstein. These points, together with in-depth discussions on how Planck scale phenomenology affects the appearance of the extragalactic sky, will be the subject matter of a paper by Ragazzoni and colleagues.

The authors are grateful to Gerard van Belle, Roberto Ragazzoni, Jonathan Mittaz, Sir Ian Axford, and Lord James McKenzie of the Hebrides and Outer Isles for discussions. They also wish to thank an anonymous referee for help-ful comments.

References

Amelino-Camelia, G., 2000, Towards quantum gravity, Proc. of the XXXV International

Winter School on Theor. Phys., Polanica, Poland, Ed. Jerzy Kowalski-Glikman.

Lecture Notes in Phys., 541, 1, Berlin: Springer-Verlag.

Amelino-Camelia, G., 2001, Nature, 410, 1065.

Baldwin, J.E., & Haniff, C.A., 2002, Phil. Trans. A360, 969.

Bekenstein, J.D., 1973, Phys. Rev. D., 7, 2333.

Hawking, S., 1975, Comm. Math. Phys., 43, 199.

t'Hooft, G., 1993, in *Salamfestschrift*, p. 284, Ed. A Ali et al (World Scientific,

Singapore).

Lieu, R., 2002, ApJ, 568, L67.

Ng, Y. -J., & van Dam, H., 2000, Found. Phys., 30, 795.

Ng, Y. -J., Lee, D. -S., Oh, M.C., & van Dam, H., 2001, Phys. Lett, B507, 236.

Ng, Y. -J., 2002, Int. J. Mod. Phys., D., in press (to appear in Dec. special issue), gr-qc/0201022.

Perlman, E.S., Stocke, J.T., Carilli, C.L., Sugiho, M., Tashiro, M.,

Madejski, G., Wang, Q.D., Conway, J., 2002, AJ, 124, 2401.

Richichi, A., Bloecker, T., Foy, R., Fraix-Burnet, D., Lopez, B., Malbet, F., Stee, P.,

von der Luehe, O.,

Weigelt, G., 2000, Proc. SPIE, 4006, 80.

Silk, J., 2001, The Big Bang, 3rd ed., W.H. Freeman & Co.

Susskind, L., 1995, J. Math. Phys (N.Y.), 36, 6377.

van Belle, G.T., Thompson, R.R., Creech-Eakman, M.J., 2002, AJ, 124, 1706.

Wheeler, J., 1982, Int. J. Theor. Phys., 21, 557.