

CO Molecular Gas in Infrared Luminous Galaxies

Lihong Yao and E. R. Seaquist

Department of Astronomy and Astrophysics, University of Toronto, Toronto, Ontario M5S 3H8, Canada

yao@astro.utoronto.ca, seaquist@astro.utoronto.ca

Nario Kuno

Nobeyama Radio Observatory, Minamimaki, Minamisaku, Nagano, 384-1305, Japan

kuno@nro.nao.ac.jp

and

Loretta Dunne

Department of Physics and Astronomy, University of Wales, Cardiff, CF24 3YB, UK

Loretta.Dunne@astro.cf.ac.uk

ABSTRACT

We present the first statistical survey of the properties of the $^{12}\text{CO}(1-0)$ and $^{12}\text{CO}(3-2)$ line emission from the nuclei of a nearly complete subsample of 60 infrared (IR) luminous galaxies selected from SCUBA Local Universe Galaxy Survey (SLUGS). This subsample is flux limited at $S_{60\mu\text{m}} \geq 5.24$ Jy with far-IR (FIR) luminosities mostly at $L_{\text{FIR}} > 10^{10} L_{\odot}$. We compare the emission line strengths of $^{12}\text{CO}(1-0)$ and $(3-2)$ transitions at a common resolution of $15''$. The measured $^{12}\text{CO}(3-2)$ to $(1-0)$ line intensity ratios r_{31} vary from 0.22 to 1.72 with a mean value of 0.66 for the sources observed, indicating a large spread of the degree of excitation of CO in the sample. These CO data, together with a wide range of data at different wavelengths obtained from the literature, allow us to study the relationship between the CO excitation conditions and the physical properties of gas/dust and star formation in the central regions of galaxies. Our analysis shows that there is a non-linear relation between CO and FIR luminosities, such that their ratio $L_{\text{CO}}/L_{\text{FIR}}$ decreases linearly with increasing L_{FIR} . This behavior was found to be consistent with the Schmidt Law relating star formation rate to molecular gas content, with an index $N = 1.4 \pm 0.3$. We also find a possible

dependence of the degree of CO gas excitation on the efficiency of star forming activity. Using the large velocity gradient (LVG) approximation to model the observed data, we investigate the CO-to-H₂ conversion factor X for the SLUGS sample. The results show that the mean value of X for the SLUGS sample is lower by a factor of 10 compared to the conventional value derived for the Galaxy, if we assume the abundance of CO relative to H₂, $Z_{CO} = 10^{-4}$. For a subset of 12 galaxies with H I maps, we derive a mean total face-on surface density of H₂+H I of about 42 M_⊙ pc⁻² within about 2 kpc of the nucleus. This value is intermediate between that in galaxies like our own and those with strong star formation.

Subject headings: radio lines: galaxies — galaxies: ISM — galaxies: starburst — ISM: evolution — ISM: molecules — surveys

1. Introduction

Knowledge of the properties and evolution of the gas and dust content in IR luminous galaxies is essential for understanding the cause and temporal evolution of starburst activity. In particular, studies of such galaxies in the nearby universe are essential as a step in understanding the role of the starburst phenomenon in the cosmic evolution of galaxies. Therefore, there is a need to investigate large statistical samples of IR luminous galaxies using a multitude of different types of data, including CO, H I, and continuum in the sub-millimeter (sub-mm), FIR and radio, in order to constrain theories of how the interstellar medium (ISM) evolves (e.g. Eales & Edmunds 1996, 1997). A significant step was achieved by Dunne et al. (2000) for the nearby universe by investigating the sub-mm properties of the dust in a complete sample of 104 galaxies. The SLUGS survey is based on the Revised Bright Galaxy Sample of *IRAS* galaxies (Soifer et al. 1987) within $-10^\circ \leq \delta \leq +50^\circ$ and with $cz \geq 1900$ km s⁻¹ and a flux limit of $S_{60\mu m} \geq 5.24$ Jy. By virtue of the selection, the SLUGS galaxies have the FIR luminosity $L_{FIR} \geq 10^{10}$ L_⊙, and hence have greater than the average star formation rate. A study of molecular gas of a complete flux-limited subsample is important to complement the study of the dust.

Low lying CO rotational line transitions at mm and sub-mm wavelengths are often used as tracers of molecular hydrogen. It is valid for molecular clouds in the Galaxy and also for external galaxies. The $J = 3$ level is at 33 K with a critical density of 1.5×10^4 cm⁻³, compared to the $J = 1$ and 2 levels of ¹²CO are lying respectively at 5.5 and 17 K above the ground level, and with respective critical densities 7.4×10^2 and 4.9×10^3 cm⁻³ for a kinetic temperature $T_{kin} = 30$ K in the optically thin limit (Jansen 1995). Therefore,

the $^{12}\text{CO}(3-2)$ transition is a better tracer for the warmer and/or denser molecular gas in star forming regions. Furthermore, the ratio of $^{12}\text{CO}(3-2)$ to $(1-0)$ line emission provides a more sensitive measure of the gas temperature and density than the ratio of $^{12}\text{CO}(2-1)$ to $(1-0)$ lines. The first detection of the $^{12}\text{CO}(3-2)$ line in the nearby starburst galaxy was that in IC 342 reported by Ho (1987). Many observations of the $^{12}\text{CO}(1-0)$ and $^{12}\text{CO}(2-1)$ lines have since been made in local galaxies (e.g. Sanders et al. 1991; Braine & Combes 1992; Braine et al. 1993; Young et al. 1995; Chini et al. 1996, and many others). Most observations of the $^{12}\text{CO}(3-2)$ line cover the central region of nearby objects (e.g. Devereux et al. 1994; Mauersberger et al. 1999; Dumke et al. 2001), where the physical conditions of molecular gas may be different from those prevailing in molecular clouds in the disk of the Galaxy. Extended maps of $^{12}\text{CO}(3-2)$ line for three nearby normal galaxies were reported by Wielebinski et al. (1999).

The weaknesses of most of the previous studies on CO molecular gas are the small numbers of objects or the lack of uniform selection criteria, whereas their strength is the production of maps with high linear resolution. In this paper, we address the aforementioned weaknesses by presenting the largest statistical survey of $^{12}\text{CO}(1-0)$ and $^{12}\text{CO}(3-2)$ point measurements from the central regions in a complete subsample of 60 IR luminous sources from SLUGS (Dunne et al. 2000). The angular resolutions of the $^{12}\text{CO}(1-0)$ observations at the Nobeyama Radio Observatory (NRO) and the $^{12}\text{CO}(3-2)$ observations at the James Clerk Maxwell Telescope (JCMT) are nearly identical, so that no assumptions need be made about differing beam dilution factors. The beamsize ($\sim 15''$) is also nearly identical to that of the SLUGS survey by Dunne et al. (2000). Together with the $850\ \mu\text{m}$ fluxes, plus existing data on H I (Thomas et al. 2002), radio continuum (Condon et al. 1990), and far-IR (Dunne et al. 2000, NED¹), we are able to search for a relationship between the degree of excitation of the CO in this sample and the star forming properties.

It is now suspected that the $^{12}\text{CO}(3-2)$ line emission makes up a substantial fraction of the SCUBA flux at $850\ \mu\text{m}$ for starburst galaxies (e.g. Papadopoulos & Allen 2000; Zhu et al. 2003). Our $^{12}\text{CO}(3-2)$ observations provide an important database for correcting the SCUBA sub-mm data in the SLUGS sample, thus permitting a revised characterization of the dust, which will be presented in a separate paper.

Finally, our investigations on the gas and dust properties in starburst regions provide an opportunity to investigate the controversial parameter X , the ratio of molecular hydrogen mass $M(\text{H}_2)$ and the $^{12}\text{CO}(1-0)$ luminosity L_{CO} , based on the LVG method (Goldreich &

¹The NASA/IPAC Extragalactic Database (NED) is operated by the Jet propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

Kwan 1974). We obtain an estimate for this parameter which depends on Z_{CO} , the abundance of CO relative to H_2 . We also use our estimate of this parameter to investigate the molecular gas content of our SLUGS subsample.

We assume $q_0 = 0.5$ and $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ throughout for consistency with previous work on these galaxies (e.g. Dunne et al. 2000).

2. Observations

2.1. $^{12}\text{CO}(1-0)$ Observations at NRO

Two observing runs measuring the $^{12}\text{CO}(1-0)$ line at 115.27 GHz for 46 SLUGS galaxies were made with NRO 45m telescope between 2001 April 12 - 19 and 2001 November 28 - December 1, using the SIS receiver S100 (single side band or SSB). The wide band acousto-optical spectrometer (AOS) was used as the back end which provided a total usable bandwidth of 250 MHz and a frequency resolution of 250 KHz. At 115.27 GHz, these correspond to a total velocity coverage of 650 km s^{-1} and velocity resolution of 0.65 km s^{-1} . The NRO 45m telescope beamsize (FWHM) at this frequency is $14.6''$. The standard source Mars was used for the determination of the main beam efficiency. The measurement yields $\eta_{mb} = 0.34 \pm 0.04$ after averaging the results from the first and second runs. The average uncertainty in the calibration of spectral line is $\sim 10\%$.

Position switching in azimuth was employed with about 20 seconds each for both ON and OFF positions, 10 sec to move from ON to OFF or from OFF back to ON, so that about one minute was required for each ON-OFF sequence or scan. Since about 30 scans were made for each source, the total integration time per pointing is about 30 minutes. Telescope pointing was checked every half hour by observing strong SiO maser lines from nearby evolved stars with receiver S40 at 43 GHz. The average pointing error was $\sim 3.0''$ (r.m.s.) for the first observing run and $\sim 2.6''$ (r.m.s.) for the second run at low wind speed. The weather conditions were average during the NRO observing runs and the system temperature for good weather conditions was $T_{sys} \sim 535 \text{ K}$. However, half of the observing time was lost for each run due to poor weather conditions. The total number of sources observed at this transition is 51, out of a total sample of 60.

2.2. $^{12}\text{CO}(3-2)$ Observations at JCMT

Observations of the $^{12}\text{CO}(3-2)$ line at 345.80 GHz for the selected 60 SLUGS galaxies were obtained with the 15m JCMT on 2001 April 23 - 28 and 2002 June 25. We used the dual-channel receiver B3 in single-sideband mode, and the Digital Autocorrelation Spectrometer (DAS) spectrometer with a bandwidth of 920 MHz ($\sim 800 \text{ km s}^{-1}$) and a frequency resolution of 750 KHz ($\sim 0.65 \text{ km s}^{-1}$). Focus and pointing were monitored frequently by observing bright continuum sources and planets. The average pointing error (r.m.s.) was found to be $\sim 3.2''$. The observations were performed by using beam-switching at a frequency of 1 Hz with a beam throw of $180''$ in azimuth. The JCMT beamsize (FWHM) at this frequency is $14.4''$. The main-beam efficiency was determined by averaging four measurements based on observations of Mars, yielding a value of $\eta_{mb} = 0.50 \pm 0.05$. The spectral line calibration was checked each night by measuring $^{12}\text{CO}(3-2)$ line emission from IRC+10216. The mean calibration error is $\sim 10\%$. The weather conditions for these observations were good, and the average system temperature was $T_{sys} \sim 600 \text{ K}$.

3. Results

3.1. The Spectra

The reduction of the $^{12}\text{CO}(1-0)$ data was performed using the NRO NEWSTAR package, and the $^{12}\text{CO} \text{ J}=3-2$ data were reduced using the JCMT SPECX package. The $^{12}\text{CO}(1-0)$ line spectra for 51 galaxies together with the spectra of $^{12}\text{CO}(3-2)$ line emission for 59 galaxies are presented in Fig. 1. A linear baseline was subtracted from each spectrum for both data sets, and the spectra were smoothed to 10.2 km s^{-1} and 11.2 km s^{-1} for the NRO and the JCMT data respectively. We converted the measured antenna temperature scale T_a^* of both spectra to the main-beam temperature T_{mb} scale by using the relation $T_{mb} = T_a^*/\eta_{mb}$. The main-beam efficiencies of NRO and JCMT telescopes are given in § 2. The JCMT and NRO line profile shapes mostly agree with each other. The ones that show discrepancies may in some instances be due to pointing error.

3.2. CO and FIR Luminosities

The velocity-integrated intensities of $^{12}\text{CO}(1-0)$ line emission I_{10} and $^{12}\text{CO}(3-2)$ line emission I_{32} corresponding to the common telescope beamsizes of $\sim 15''$ are shown in Table 1. The uncertainties for the integrated line intensities include components due to noise, the un-

certainty in the beam efficiencies and the calibration of spectral line, combined quadratically. The luminosities associated with the beamsize of $\sim 15''$, in units $\text{K km s}^{-1} \text{ Mpc}^2 \Omega_b$, were calculated using equations 1(a) and 1(b) below and are presented in Table 2. Both equations 1(a) and 1(b) were derived by transforming the relevant parameters from the rest frame to the observer frame.

$$L_{12\text{CO}(1-0)} = \left[\frac{D_L^2}{(1+z)^3} \right] I_{10} \Omega_b, \quad (1a)$$

$$L_{12\text{CO}(3-2)} = \left[\frac{D_L^2}{(1+z)^3} \right] I_{32} \Omega_b \quad (1b)$$

where D_L is the luminosity distance of a galaxy in Mpc, $D_L = D (1+z)$, and D is the proper distance of the galaxy (see Table 1), the integrated line intensities I_{10} and I_{32} have units of K km s^{-1} , and Ω_b is the beam solid angle. The numerical value of Ω_b is constant for all observations and has been set to unity for simplicity. For conversion to absolute luminosities, the value of $\Omega_b = 1.1 \times 10^{-3}$ steradians.

Fig. 2 shows a comparison between the CO luminosities L_{CO} and FIR luminosities L_{FIR} for two different FIR color regimes. The quantity L_{FIR} here corresponds to the volume marked by a $15''$ beam obtained by applying a scale factor to the total FIR luminosity (from Dunne et al. 2000). This factor is derived from the original SCUBA images which have a resolution of $15''$ and from radio continuum images (Condon et al. 1990) convolved to $15''$. It measures the ratio of flux within the beam to the total flux. It is less than unity if the source is resolved by the telescope beam and equal to unity for an unresolved source. It is assumed here that the FIR brightness distribution is similar to that for the sub-mm/cm and radio continuum. A linear fit to each transition in the log-log plane yields

$$\log_{10} L_{12\text{CO}(1-0)} = (-0.9 \pm 0.7) + (0.58 \pm 0.07) \log_{10} L_{\text{FIR}}, \quad (2a)$$

$$\log_{10} L_{12\text{CO}(3-2)} = (-2.4 \pm 0.5) + (0.70 \pm 0.04) \log_{10} L_{\text{FIR}} \quad (2b)$$

The r.m.s. dispersion for the fits is about a factor of 3 and 2 for $^{12}\text{CO}(1-0)$ and $^{12}\text{CO}(3-2)$ respectively. The scatter is much larger than the errors measured for the telescopes indicated by the error bars in the plots. Variations in the CO-to- H_2 conversion factor X may be responsible for part of the scatter. The larger scatter seen in the $^{12}\text{CO}(1-0)$ plot may be due to the larger baseline ripple measured at this transition.

It is evident that the CO luminosities at both transitions increase non-linearly with increasing L_{FIR} , such that the ratio L_{CO}/L_{FIR} decreases with increasing L_{FIR} . Fig. 3 shows a plot of the ratio of L_{CO}/L_{FIR} versus the projected beamsize on the galaxies, to test whether this effect is introduced by partially resolving the nearby galaxies. No relationship is evident, indicating that the total luminosity of the galaxy is the primary factor. The implications of this result will be discussed in § 4.1.

3.3. The $^{12}\text{CO}(3-2)/(1-0)$ Line Intensity Ratio

For the 48 galaxies with data at both transitions, we calculate the intensity ratios of the two transitions using the equation,

$$r_{31} = I_{32}/I_{10} \quad (3)$$

The results along with the asymmetric uncertainties are given in Table 1. The asymmetric errors of r_{31} are derived analytically, assuming the error distributions of I_{10} and I_{32} are Gaussian functions. The asymmetric confidence intervals in Table 1 represent an enclosed probability of 68%.

The values of r_{31} range from 0.22 (for NGC 5665) to 1.72 (for NGC 5937). There are seven sources with $r_{31} > 1$, namely UGC 5376, IR 1222-06, UGC 8739, NGC 5433, NGC 5713, NGC 5937, and MCG +01-42, signifying very high excitation coupled with low optical depth. However, UGC 5376, IR 1222-06, and MCG +01-42 have high uncertainties in r_{31} , leaving only four whose r_{31} values are significantly greater than 1.0. In our sample, thirteen galaxies have r_{31} ratios less than 0.4, the average value of r_{31} derived from the giant molecular clouds in the Galaxy (Sanders et al. 1993).

The mean value of r_{31} for SLUGS sample is 0.66, close to that of 0.63 measured for 28 galaxies with strong CO emission published by Mauersberger et al. (1999). The quantity L_{FIR} ranges from 10^9 to $10^{12} L_{\odot}$ in the Mauersberger sample, compared to the range 10^{10} - $10^{12} L_{\odot}$ in SLUGS sample. Fig. 4 shows a comparison between the distributions of r_{31} for the SLUGS sample and those measured by Mauersberger et al. (1999). The probability of a significant difference between these two distributions is $\sim 68\%$ using a *Kolmogorov-Smirnov* test. This implies that these two distributions are not significantly different. The galaxies observed by Mauersberger et al. (1999) are located between 0.8 and 73 Mpc, much closer, and on average less luminous than our objects in the range 25 - 277 Mpc. Thus there is no evidence that r_{31} depends on the distance of the galaxies over the range 0.8 - 277 Mpc, because of the different luminosity ranges represented by these two samples.

A search was conducted for relationships between r_{31} and a number of parameters relating primarily to star formation. The existence of such correlations would be expected to shed light on the factors affecting star formation, or the effects of star formation on gas properties. These are considered below:

3.3.1. Projected Beamsize

Radio continuum maps (Condon et al. 1990), the H I (Thomas et al. 2002), and SCUBA 850 μ m maps (Thomas et al. 2002) show that much of the emission in the SLUGS objects is extended with respect to the beamsize of $\sim 15''$, especially for the nearer objects. On average we detect about 45% of all the emission from these galaxies assuming the CO is distributed in a similar way to the 850 μ m and radio continuum emission. Therefore, the CO measurements may reflect the physical conditions in the center regions for the galaxies at lower cz . Maps of $^{12}\text{CO}(3-2)$ line emission in 12 nearby galaxies by Dumke et al. (2001) have shown that the $^{12}\text{CO}(3-2)/(1-0)$ line intensity ratio decreases outward from the centers of many galaxies. In principle, for a given beamsize, the average line ratios obtained from the single pixel measurement should therefore decrease with increasing distance, since more emission from lesser excited CO lines arising from the disk regions is detected by the telescope beam. Fig. 5 shows a plot of r_{31} versus projected beam size. Similar to the relation between the ratio of $L_{\text{CO}}/L_{\text{FIR}}$ and the linear size sampled (see Fig. 3), no correlation between these quantities is evident. The lack of such a correlation may be due to sample selection effect, however, since the more distant galaxies may have a larger fraction of highly excited CO molecular gas.

3.3.2. Dust and Gas

Large amounts of FIR emission are generated by the dust heated by young massive stars in starburst systems, in which case the FIR luminosity is a good measurement of the star formation rate (SFR) (Kennicutt 1998a). Fig. 6 shows the ^{12}CO line ratio versus dust temperature T_{dust} and FIR luminosity L_{FIR} within a $15''$ beam. The dust temperatures are from Dunne et al. (2000) and the FIR luminosities are scaled to a $15''$ beam using sub-mm and radio maps as described in § 3.2. No significant correlation is evident in these plots. Other SFR indicators are radio continuum luminosity L_{radio} and the optical color index ($B - V$). Plots for these are not shown, but no obvious correlation was found with the parameter r_{31} . In Fig. 7, we examine the dependence of r_{31} on dust mass M_{dust} and molecular hydrogen mass $M(\text{H}_2)$. The dust masses were obtained from Dunne et al. (2000)

and scaled to a $15''$ beam as described earlier. The molecular gas masses for the SLUGS sample are derived from our CO luminosities by applying the conversion factor $X = 2.7 \times 10^{19} \text{ cm}^{-2} [\text{K km s}^{-1}]^{-1}$ derived in § 4.3.1. The result is $M(\text{H}_2) = 1.1 \times 10^3 D_L^2 / (1+z) S_{CO} M_\odot$ by scaling the result of Kenney & Young (1989), where D_L is the luminosity distance of a galaxy in Mpc, and S_{CO} is the $^{12}\text{CO}(1-0)$ flux measured within a $15''$ beam, $S_{CO} = 2.1 I_{10} \text{ Jy km s}^{-1}$. The values of $M(\text{H}_2)$ are also presented in Table 2. Again, no relationship is evident.

3.3.3. Surface Brightnesses and Color Indices

The average surface brightness for all wavebands is defined here as the ratio of the total flux to the square of the optical diameter. Thus it measures the surface brightness averaged over the optical diameter, and provides a measure of the concentration of star forming activity. Fig. 8 shows a plot of r_{31} versus the surface brightness for mid-IR, far-IR, sub-mm, and H I emission. We find two possible correlations, one is between r_{31} and FIR surface brightness, and another one is between the r_{31} and SCUBA 850 μm surface brightness. Fig. 9 shows the line ratio versus different color indices or ratios: $\left[\frac{S_{60\mu\text{m}}}{S_{100\mu\text{m}}} \right]$ and $\left[\frac{S_{HI}}{S_{850\mu\text{m}}} \right]$. No significant correlation is found between these ratios.

3.3.4. Star Formation Efficiency

The ratio $L_{FIR}/M(\text{H}_2)$ is often used as an indicator of the star formation efficiency (*SFE*) which gives the star formation rate per unit mass of molecular gas. In Table 2, we present the results of *SFE* calculations for our SLUGS sample. Here the mass of molecular gas has been derived in the same manner as that in § 3.3.2. These values are an order of magnitude higher than what would be obtained if the conventional value of X is used. Fig. 10 shows a significant correlation between r_{31} and the *SFE* within the telescope beam. The sample shown in Fig. 10 is divided into two ranges by gas mass centered at $M(\text{H}_2) = 10^8 M_\odot$, and also by dust FIR luminosity centered at $L_{FIR} = 10^{10} L_\odot$ which are indicated by four different symbols in the plot. The linear correlation coefficient is obtained from the data with $SFE \leq 200 L_\odot/M_\odot$. The coefficient is 0.74 at a significance level of 1.4×10^{-7} (i.e. probability that r_{31} and *SFE* are unrelated). However, the correlation is diminished at $SFE > 200 L_\odot/M_\odot$, where the ratios spread between 0.5 and 1.72. The significance of the dependence of r_{31} on *SFE* will be discussed in § 4.2.

4. Discussion

4.1. The Relation Between CO and FIR Luminosities

As noted in § 3.2, the non-linear variation of CO with FIR luminosity implies a decrease in the ratio L_{CO}/L_{FIR} with increasing L_{FIR} (see Fig. 2). For $^{12}\text{CO}(1-0)$ this ratio decreases by more than an order of magnitude over the range $L_{FIR} = 5 \times 10^9 - 10^{12} L_{\odot}$. Part of this variation might be attributable to the effect of beamsize if the ratio is dependent on the fraction of the galaxy resolved by the beam. However, as noted in § 3.2, Fig. 3 shows that the beam effects do not seem to influence the L_{CO}/L_{FIR} ratio, and that for any plausible gas excitation gradient (i.e. hotter gas confined in the center) such effects would show in the r_{31} ratio as well, but Fig. 5 shows no evidence that more distant objects have smaller values of r_{31} (see § 3.3.1). In addition, this effect would be expected to produce an increase rather than a decrease in L_{CO}/L_{FIR} with increasing L_{FIR} because of the strong sensitivity of L_{FIR} to dust temperature. Therefore this ratio depends intrinsically on the total FIR luminosity. It should be noted that our results imply that for high z galaxies where L_{FIR} is exceedingly high, the corresponding values of L_{CO} will be seriously overestimated if a linear relation between L_{CO} and L_{FIR} were assumed.

A possible cause of the variation in the L_{CO}/L_{FIR} ratio is an increase in L_{FIR} per unit mass of dust with increasing L_{FIR} . To investigate this point further, we first plot in Fig. 11 L_{CO} versus M_{dust} to determine this relationship is linear. Linear fits to these relations in the log-log plane yield

$$\log_{10} L_{^{12}\text{CO}(1-0)} = (-0.4 \pm 0.6) + (0.77 \pm 0.09) \log_{10} M_{dust}, \quad (4a)$$

$$\log_{10} L_{^{12}\text{CO}(3-2)} = (-1.8 \pm 0.5) + (0.95 \pm 0.07) \log_{10} M_{dust} \quad (4b)$$

We see indeed that the slopes are closer to unity than with $\log_{10} L_{CO}$ versus $\log_{10} L_{FIR}$ (Equation (2a) and (2b)), though there is still a significant departures from a linear relation for $L_{^{12}\text{CO}(1-0)}$. Secondly, we note that Fig. 2 shows a trend that the L_{CO}/L_{FIR} ratio decreases with increasing dust temperature which is indicated by the two different FIR color ranges. This implies that the galaxies of the SLUGS sample which have higher luminosity is because they have a higher temperature in addition to simply more dust and gas, which would alone yield a linear relation between L_{CO} and L_{FIR} . The correlation found between $L_{60\mu m}$ and dust temperature by Dunne et al. (2000) also support this argument. This effect is consistent with the systematically higher SFE 's deduced for such objects, with the warmer IRAS colors, and with the saturation of the r_{31} ratio at high SFE 's (see Fig. 10).

As noted in § 3.3.2, L_{FIR} is widely used as an indicator of the SFR , and $M(H_2)$ is often computed from the CO flux. Thus, the non-linear relation between L_{CO} and L_{FIR} may reflect the behavior of star formation law for the nearby IR luminous galaxies. The star formation law can be represented as a power law that can be expressed in terms of the star formation rate per unit area Σ_{SFR} and the gas density Σ_{gas} , i.e. $\Sigma_{SFR} = A\Sigma_{gas}^N$, first introduced by Schmidt (1959). Previous studies have shown that the activity of disk-averaged star formation is correlated with the mean total molecular gas content (Kennicutt 1989; Boselli 1994; Boselli et al. 1995), and this correlation is found to be stronger with H_2 gas in starburst regions (Kennicutt 1998b). Fig. 12 shows a possible correlation between the SFR per unit area and the H_2 gas density Σ_{H_2} measured within a $15''$ beam for our SLUGS sample. Both Σ_{SFR} and Σ_{H_2} are corrected for the inclination of galaxies in the SLUGS sample. The correction for the galaxy inclination is necessary because the Schmidt Law involves Σ_{SFR} and Σ_{H_2} when viewed normal to the disk. The dispersion in the fit is about a factor of 3. The slope based on a linear fit in the log-log plane yields 1.4 ± 0.3 . This value is consistent with the Schmidt law applied to the circumnuclear starbursts (Kennicutt 1998b), though the uncertainty in the slope is too high to obtain a definitive value for the index. A possible reason for this is that our result refers to different regions in different galaxies, depending on the projected beamsize, whereas the work by Kennicutt and others consistently refer to disk-averaged parameters. The Schmidt Law involves $H\ I+H_2$, but ignoring $H\ I$ gas, which averages about 30% of the total gas mass measured within a $15''$ beam (see § 4.3.2), is most likely not the cause of the scatter seen in Fig. 12, since a study by Wong & Blitz (2002) with a high angular resolution shows no correlation between $H\ I$ and star formation rate.

4.2. Correlation between r_{31} and Star Formation Parameters

We examined in § 3.3 the excitation of CO and its relation with the properties of gas/dust and star formation in the central starburst regions in SLUGS sample. There are no significant correlations between r_{31} and the distance of galaxies, star formation rate, dust temperature and mass, H_2 gas mass, the color indices, and the luminosities of IR and radio continuum. The lack of correlation of r_{31} with properties relating to total star formation reflects a range of localized conditions in the molecular clouds. The possible dependence of r_{31} on FIR/sub-mm surface brightness (see Fig. 8) and SFE (see Fig. 10) suggests that the higher degree of CO excitation is related to a higher concentration and efficiency of star forming activity. Such conditions would arise in an intense starburst where the surface density of such activity is high, consuming most of the gas in the region. The saturation effect of r_{31} seen at higher SFE which is probably due to thermalization of the CO levels, reflecting denser and warmer

gas in regions of high star formation efficiency. This interpretation is consistent also with the non-linear relation between L_{CO} and L_{FIR} (see Fig. 2), and with higher color temperatures associated with the most IR luminous and most efficient star forming galaxies. It is not clear whether these correlations reflect a high excitation ratio as the underlying cause or the effect of star forming activity.

The regime where $r_{31} > 1.0$ requires optically thin CO lines from extremely warm and/or dense gas. The relatively small number of galaxies with high values of r_{31} might be associated with the highly excited gas from a short-lived gas phase in the starburst systems. The lower line ratios may be associated with gas that is cooler and less denser.

4.3. The Molecular Gas Content

The investigation of the gas and dust content depends critically on a reliable estimate of the controversial parameter X . Studies have shown that this parameter varies from galaxy to galaxy (Booth & Aalto 1998; Boselli et al. 2002), and it is thought to be higher in metal-poor galaxies and lower in starburst galaxies than in Galactic molecular clouds, where X is about $2.8 \times 10^{20} \text{ cm}^{-2} [\text{K km s}^{-1}]^{-1}$ (Bloemen et al. 1986; Strong et al. 1988). Thus in starburst galaxies, application of the standard factor can produce a significant overestimate of molecular hydrogen mass (Solomon et al. 1997; Downes & Solomon 1998; Zhu et al. 2003).

4.3.1. The CO-to-H₂ Conversion Factor X from An LVG Method

In this section, we examine the molecular gas content and properties using large velocity gradient (LVG) models. The LVG method is capable in principle of yielding the column density of CO, and hence of H₂, if an abundance ratio $Z_{CO} = [\text{CO}/\text{H}_2]$ is assumed. Hence, this method should yield an estimate of the X factor independent of other considerations such as gas-to-dust ratio, which is discussed in the next section. We assume that the ISM comprises molecular clouds characterized by a single set of average physical properties. It is well understood that models incorporating more data for starburst galaxies require at least two components - a prevailing optically thin warm diffuse component, and an optically thick cool dense component with a smaller filling factor (e.g. Aalto et al. 1995). However, comparisons between single and double component models yield approximately the same column density for H₂, which is weighted strongly toward the prevailing diffuse component (e.g. Zhu et al. 2003).

The LVG approach can be formulated into an equation yielding the X factor from the

LVG parameters given by

$$X = \frac{n(\text{H}_2)\Lambda}{Z_{\text{CO}}T_{\text{rad}}} \quad (5)$$

where $n(\text{H}_2)$ is the H_2 gas density, T_{rad} is the radiation temperature for the $^{12}\text{CO}(1-0)$ line transition, $\Lambda = \frac{Z_{\text{CO}}}{(dv/dr)}$, and (dv/dr) is the velocity gradient associated with the individual clouds (see Appendix A). There are not enough data to uniquely determine the parameters in this equation, which usually require multiple line excitation ratios, measured brightness temperatures, and isotope intensity ratios to fully constrain the conditions in the molecular gas. In our case, it is therefore necessary to make some reasonable assumptions to constrain the LVG solutions.

The highest degree of uncertainty concerns the parameter Z_{CO} which is unknown and needs to be addressed. Values widely quoted for starburst galaxies are in the range 10^{-5} - 10^{-4} (e.g. Booth & Aalto 1998; Mao et al. 2000). Unfortunately, there are few if any reliable determinations for starburst galaxies. Within the Milky Way, measurements of Z_{CO} and chemical models for dark and star forming clouds yield values within the range 5×10^{-5} - 2.7×10^{-4} (Blake et al. 1987; Farquhar et al. 1997; Hartquist et al. 1998; van Dishoeck 1998; van Dishoeck & Blake 1998). It seems unlikely therefore that $Z_{\text{CO}} < 10^{-5}$ for starburst galaxies unless their metallicity is unusually low, which would occur only in low luminosity systems. For the SLUGS sample, therefore, we adopt $Z_{\text{CO}} = 10^{-4}$, bearing in mind the dependence of our computed values for X on this parameter.

We adopt dust temperatures for each galaxy from Dunne et al. (2000), assuming T_{dust} to be representative of gas kinetic temperature T_{kin} . Fig. 13 shows a series of plots of r_{31} versus T_{dust} for our subsample combined with theoretical curves based on LVG models. The theoretical curves represent r_{31} versus T_{kin} for a variety of assumed densities $n(\text{H}_2)$ based on a series of LVG models using $Z_{\text{CO}} = 10^{-4}$, and values of $\Lambda = 10^{-7}$, 10^{-6} , 10^{-5} , and 10^{-4} ($\text{km s}^{-1} \text{ pc}^{-1}$) $^{-1}$. In each case it is assumed that $T_{\text{kin}} = T_{\text{dust}}$. Note the convergence of the curves at higher densities reflecting the saturation of the line ratios due to thermalization of the optically thick CO transitions. Note also that the ratios $r_{31} > 1.0$ can not be readily fit by the models with $\Lambda \geq 10^{-6}$ ($\text{km s}^{-1} \text{ pc}^{-1}$) $^{-1}$, a point to which we return later.

Thus for each galaxy in one of the data plots, it is possible to derive a molecular gas density and radiation temperature T_{rad} which is dependent upon Λ . Using Equation (5), we then compute a range of X parameter for the IR luminous galaxies. For each Λ , only points that fall below the upper limits of r_{31} due to saturation are used for computing the distribution of X . For example, for $\Lambda = 10^{-5}$ ($\text{km s}^{-1} \text{ pc}^{-1}$) $^{-1}$, thermalization occurs at $r_{31} = 0.86$, corresponding to $T_{\text{dust}} = 55$ K, $n(\text{H}_2) = 2.4 \times 10^3 \text{ cm}^{-3}$, and $T_{\text{rad}} = 27$ K. The resulting

range of X is $1 - 3 \times 10^{19} \text{ cm}^{-2} [\text{K km s}^{-1}]^{-1}$. For points falling above the saturation curves ($r_{31} > 0.86$) for $\Lambda = 10^{-5} (\text{km s}^{-1} \text{ pc}^{-1})^{-1}$ (see Fig. 13), we obtain X values in a range $2.4 \times 10^{19} - 6.7 \times 10^{19} \text{ cm}^{-2} [\text{K km s}^{-1}]^{-1}$ from the plots corresponding to $\Lambda = 10^{-6}$ and $10^{-7} (\text{km s}^{-1} \text{ pc}^{-1})^{-1}$. However, the latter value of Λ is virtually unacceptable, since it corresponds to an impossibly high velocity gradient or impossibly low CO abundance. Therefore, T_{kin} would have to be higher than the measured T_{dust} for those points to lie below the saturation curves for $\Lambda \leq 10^{-6} (\text{km s}^{-1} \text{ pc}^{-1})^{-1}$. Fig. 14 shows the resulting distribution of the derived X and its average values for different Λ . It is clear that regardless of Λ , the values of X derived from starburst galaxies are systematically lower than the value $X = 2.8 \times 10^{20} \text{ cm}^{-2} [\text{K km s}^{-1}]^{-1}$ derived from the disk of the Galaxy.

In order to further constrain the range of the X factor and Λ shown in Fig. 14, we compare the CO isotope intensity ratios $R_{10} = I[^{12}\text{CO}(1-0)]/I[^{13}\text{CO}(1-0)]$ predicted by the LVG model with observed values. We adopted the isotope abundance ratio as $\left[\frac{^{12}\text{CO}}{^{13}\text{CO}}\right] = 40 - 75$ (Wilson & Rood 1994). The CO isotope intensity ratios data are taken from Aalto et al. (1995) and Taniguchi et al. (1999) for 34 IR luminous galaxies which have a range of FIR luminosities similar to the SLUGS sample, i.e. $L_{FIR} > 10^{10} L_{\odot}$. Fig. 15 shows a comparison between the distributions of computed and observed isotope ratios for the sample as a whole for each value of Λ , and corresponding value for the mean derived X factor. The plots show that the best agreement with the observed distribution of R_{10} is obtained for $\Lambda = 10^{-5} (\text{km s}^{-1} \text{ pc}^{-1})^{-1}$, corresponding to an average value for $X = 2.7 \times 10^{19} \text{ cm}^{-2} [\text{K km s}^{-1}]^{-1}$. The other plots assuming other values of Λ show marked disagreement with the observed isotope line ratios.

The resulting value for X ($2.7 \times 10^{19} \text{ cm}^{-2} [\text{K km s}^{-1}]^{-1}$) in starburst regions is thus ten times smaller than the conventional value of X for GMC's in the Milky Way. But the result is comparable with $X = 0.5 \times 10^{20} \text{ cm}^{-2} [\text{K km s}^{-1}]^{-1}$ estimated from diffuse clouds in the Galaxy by Polk et al. (1988), and with that found for extreme starbursts in nearby galaxies by Downes & Solomon (1998) and for starburst galaxy M 82 by Mao et al. (2000).

We repeated the LVG model calculations for a higher CO isotope abundance ratio of 75 (Wilson & Rood 1994), and found that the average value of X for $Z_{CO} = 10^{-4}$ becomes $8.2 \times 10^{19} \text{ cm}^{-2} [\text{K km s}^{-1}]^{-1}$. We also repeated the entire analysis by increasing the adopted kinetic temperatures by 50%, and found that the resulting value of X remains essentially the same, since for a given Λ , the lower densities required are largely offset by lower values of the radiation temperature (see Equation (5)). Our LVG analysis also shows that the predicted isotope ratios $^{12}\text{CO}(2-1)/^{13}\text{CO}(2-1)$ are in satisfactory agreement with the observed ratios reported by Taniguchi et al. (1999). The results do not seem to support the conclusion by Taniguchi et al. (1999) that the comparison between CO(2-1) and CO(1-0) isotope intensity

ratios may imply that the unusually large values in starburst systems indicate an isotope abundance anomaly in starburst galaxies. Our results are consistent with effects of excitation and optical depth.

4.3.2. *The $M(\text{H}_2)/M(\text{H I})$ Ratio*

In this section, we briefly consider a small subset of our sample with H I maps available to assess the combined molecular and atomic gas content within the telescope beam. The subset comprises twelve galaxies with an average $L_{FIR} = 4.1 \times 10^{10} L_{\odot}$ (Thomas et al. 2002). The H I masses for our $15''$ beam were estimated from the peak H I column densities maps at $25''$ resolution by assuming a uniform H I brightness distribution within the $25''$ beam. The mean ($\pm$ r.m.s.) projected beamsize for the subset is $3.6 (\pm 1.1)$ kpc.

The overall mean ($\pm$ r.m.s.) values for $M(\text{H}_2)/M(\text{H I})$ ratio and combined $\text{H}_2 + \text{H I}$ face-on surface density are respectively 2.0 ± 1.5 and $42 \pm 18 M_{\odot} \text{ pc}^{-2}$. These quantities must be considered respectively as upper and lower limits however, since the H I masses may be underestimated due to the effects of absorption by H I against the radio continuum in the nuclear regions (Thomas et al. 2002). This is particularly the case for one galaxy, NGC 5900, for which $M(\text{H}_2)/M(\text{H I}) = 21$, and this object was omitted from the mean value quoted above. The mean ratio $M(\text{H}_2)/M(\text{H I})$ is substantially larger than that found for optically selected late type galaxies (0.15) by Boselli et al (2002). This is partly attributable to the fact that their values are global measures, whereas our values refer to the nuclear regions of galaxies selected according to high star formation rates. The $\text{H}_2 + \text{H I}$ surface density is intermediate between the central surface densities for galaxies with lower star forming rates like the Milky Way, which is about $10 M_{\odot} \text{ pc}^{-2}$ (Elmegreen 1989) and those for ultra-luminous IR galaxies, which are up to $10^5 M_{\odot} \text{ pc}^{-2}$ (Aalto et al. 1995).

It should be emphasized that these estimates refer to comparatively warm gas heated by star formation in these systems. The outer disks of galaxies may be hiding substantial amounts of cool H_2 , where the radiative efficiency of CO is low (i.e. X is high). The $M(\text{H}_2)/M_{dust}$ ratio will be discussed in a separate paper (in preparation), where the dust mass of the SLUGS sample (Dunne et al. 2000) will be corrected for the effects of contamination of the SUCBA $850 \mu\text{m}$ filter.

4.3.3. Virial Stability of The Molecular Clouds

Using the virial theorem, the velocity gradient under the virial condition for a spherical cloud with a mean gas density $\langle n \rangle$ is given by,

$$\left(\frac{dv}{dr}\right)_{VIR} = 0.65\alpha^{\frac{1}{2}}\left(\frac{\langle n \rangle}{10^3\text{cm}^{-3}}\right)^{\frac{1}{2}}\text{km s}^{-1}\text{pc}^{-1} \quad (6)$$

where α is in the range 0.5 - 2.5 (Papadopoulos 1998; Bryant & Scoville 1996) depending primarily on the assumed density profile. For a mean gas density of $1.4 \times 10^3 \text{ cm}^{-3}$ for $r_{31} \leq 0.8$ derived from the LVG model for our IR luminous galaxies (see Fig. 13) with $\Lambda = 10^{-5} (\text{km s}^{-1} \text{ pc}^{-1})^{-1}$ and $Z_{CO} = 10^{-4}$, we estimate a virial velocity gradient of $1.1 \text{ km s}^{-1} \text{ pc}^{-1}$ for $\alpha = 2.5$. This virial value is about 9 - 90 times smaller than the velocity gradient of $10 - 100 \text{ km s}^{-1} \text{ pc}^{-1}$ also derived from LVG modeling using $\Lambda = 10^{-5} - 10^{-6} (\text{km s}^{-1} \text{ pc}^{-1})^{-1}$ and $Z_{CO} = 10^{-4}$. The same calculation was made for a lower r_{31} using the source UGC 6436 with $r_{31} = 0.35$, and a corresponding gas density of $7.5 \times 10^2 \text{ cm}^{-3}$. It was found that $\left(\frac{dv}{dr}\right)_{VIR}$ is still about 11 times smaller than the LVG velocity gradient. Since such low values of r_{31} are more sensitive to density than more saturated values near unity, this result provides a strong confirmation of the non-virial expansion velocities. As noted earlier, there are a number of galaxies with high ^{12}CO line ratios ($r_{31} \sim 0.8 - 1.72$) seen in Fig. 13, where Λ may be lower than $10^{-5} (\text{km s}^{-1} \text{ pc}^{-1})^{-1}$. The above estimates imply that the molecular clouds in the starburst regions are not gravitationally bound, confirming the suggestion recently made by Zhu et al. (2003, ApJ in press) for the Antennae galaxies, unless one is willing to accept a significantly lower value for Z_{CO} . The cause could be the destruction by the stellar winds and expanding shells of new-born massive stars. We note that the non-virialized component emanates from the gas phase that dominates the ^{12}CO emission, and not necessarily from the entire cloud mass, in accordance with two-phase gas models (Aalto et al. 1995; Papadopoulos & Seaquist 1999).

5. Conclusions

The main conclusions of this work are:

1. The L_{CO}/L_{FIR} ratios measured from the SLUGS sample decrease with increasing FIR luminosity. The non-linearity between L_{CO}/L_{FIR} and L_{FIR} implies that the more IR luminous galaxies have higher dust temperatures than those at the low luminosity end. This non-linear relation was found to be consistent with that expect for the Schmidt Law for star formation with an index $N = 1.4 \pm 0.3$, though the index is too uncertain to provide a new

and definitive Schmidt relation.

2. The degree of CO excitation measured by the $^{12}\text{CO}(3-2)/(1-0)$ line intensity ratios vary greatly from galaxy to galaxy in the SLUGS sample. There is a trend for the r_{31} ratios to increase with increasing concentration and efficiency of star forming activity. The saturation of r_{31} seen at higher SFE implies that the gas is denser and warmer in regions of high SFE which is consistent with the non-linearity between L_{CO}/L_{FIR} and L_{FIR} .

3. Our ^{12}CO line measurements together with dust and CO isotope data taken from the literature are modeled using the LVG method to estimate that the CO-to- H_2 conversion factor lies between 2.7×10^{19} and $6.6 \times 10^{19} \text{ cm}^{-2} [\text{K km s}^{-1}]^{-1}$ for SLUGS galaxies, which is about 4 - 10 times lower than the conventional X derived from the Galaxy.

4. For a subset of 12 galaxies with available H I maps, the total gas face-on surface density within about 2 kpc of the center averages about $42 \text{ M}_{\odot} \text{ pc}^{-2}$, which is intermediate between that for disk galaxies with low and high star formation rates.

5. Our Virial analysis shows that the molecular clouds in starburst regions are not gravitationally bound unless one is willing to accept a 9 - 90 times lower $[\text{CO}/\text{H}_2]$ abundance ratio. Most of the ^{12}CO line emission originates from the non-virialized warm and diffuse gas clouds.

We thank the staffs of the JCMT and NRO, and Dr. Siow-wang Lee at University of Toronto for their assistance with the observations. We thank Alan Richardson at University of Toronto for his assistance in the analysis of SCUBA 850 μm and cm radio images. We also thank Dr. Helen Thomas at University of Cambridge for providing us the electronic forms of twelve H I maps (from Thomas et al. 2002). L.Y. would like to thank Dr. Phil Arras at Kavli Institute for Theoretical Physics. We thank the referee for many helpful comments. This research was supported by a research grant from the Natural Sciences and Engineering Research Council of Canada, and a Reinhardt Graduate Student Travel Fellowship from the Department of Astronomy and Astrophysics at the University of Toronto.

A. Derivation of The CO-to- H_2 Conversion Factor X From LVG Model

From the LVG model for a spherical molecular cloud, the column density per unit velocity range in the average cloud is given by

$$n(\text{H}_2)\Lambda = \frac{n(\text{H}_2)Z_{CO}}{dv/dr} \quad (\text{A1})$$

where $n(\text{H}_2)$ is the density of H_2 , $\Lambda = \frac{Z_{\text{CO}}}{dv/dr}$, Z_{CO} is the CO abundance ratio $[\text{CO}/\text{H}_2]$, and dv/dr is the velocity gradient within the cloud. The total CO column density per cloud N_{CO} , integrated over velocity is then

$$N_{\text{CO}} = n(\text{H}_2)\Lambda\Delta v \quad (\text{A2})$$

where Δv is the velocity width of an individual cloud.

The column density $\overline{N_{\text{CO}}}$ averaged over the beam area can be written as

$$\overline{N_{\text{CO}}} = n(\text{H}_2)\Lambda\Delta v \frac{N_{\text{cloud}}\sigma}{A_{\text{beam}}} \quad (\text{A3})$$

or,

$$\overline{N_{\text{CO}}} = n(\text{H}_2)\Lambda \left[\frac{N_{\text{cloud}}\sigma}{A_{\text{beam}}} \frac{\Delta v}{\Delta V} \right] \Delta V \quad (\text{A4})$$

where N_{cloud} is the number of clouds in the beam, σ is the cloud cross section area, A_{beam} is the beam area, and ΔV is the line width of all emission detected in the beam.

The quantity inside inside the brackets of Equation (A4) is the brightness dilution factor representing the product of the geometric and velocity filling factors within the beam.

Assuming for simplicity, a rectangular line profile, then

$$\overline{N_{\text{CO}}} = n(\text{H}_2)\Lambda \left[\frac{T_{\text{mb}}}{T_{\text{rad}}} \right] \Delta V \quad (\text{A5})$$

where the radiation temperature T_{rad} is calculated from LVG modeling for $^{12}\text{CO}(1-0)$ or (3-2) line emission, and T_{mb} is the corresponding main-beam temperature.

The quantity $T_{\text{mb}}\Delta V$ can be replaced by $\int T_{\text{mb}}dV$, the line flux in units of brightness. Thus

$$\overline{N_{\text{CO}}} = \frac{n(\text{H}_2)\Lambda}{T_{\text{rad}}} \int T_{\text{mb}}dV \quad (\text{A6})$$

Setting $\overline{N_{\text{CO}}} = Z_{\text{CO}} \overline{N(\text{H}_2)}$, where $\overline{N(\text{H}_2)}$ is the corresponding column density of H_2 , and $X = \frac{\overline{N(\text{H}_2)}}{I_{10}}$, where

$$I_{10} = \int T_{mb} dV \quad (\text{A7})$$

for ^{12}CO (1-0) line emission. We obtain

$$X = \frac{n(H_2)\Lambda}{Z_{CO}T_{rad}}. \quad (\text{A8})$$

REFERENCES

- Aalto, S., Booth, R. S., Black, J. H., & Johansson, L. E. B. 1995, *A&A*, 300, 369
- Blake, G. A., Sutton, E. C., Masson, C. R., & Phillips, T. G. 1987, *ApJ*, 315, 621
- Bloemen, J. B. G. M., et al. 1986, *A&A*, 154, 25
- Booth, R. S. & Aalto, S. 1998, in *The Molecular Astrophysics of Stars and Galaxies*, T. W. Hartquist & D. A. Williams (eds.), Oxford Science Publications, 437
- Boselli, A. 1994, *A&A*, 292, 1
- Boselli, Gavazzi, G., A., Lequeux, J., Buat, V., Casoli, F., Dickey, Z., & Donas, J. 1995, *A&A*, 300, L13
- Boselli, A., Lequeux, J., & Gavazzi, G. 2002, *A&A*, 384, 33
- Braine, M. & Combes, F. 1992, *A&A*, 264, 433
- Braine, M., et al. 1993, *A&AS*, 97, 887
- Bryant, P. M. & Scoville, N. Z. 1996, *ApJ*, 457, 678
- Chini, R., Krügel, E., & Lemke, R. 1996, *A&AS*, 118, 47
- Condon, J. J., Helou, G., Sanders, D. B., & Soifer, B. T. 1990, *ApJS*, 73, 359
- Devereux, N., Taniguchi, Y., Sanders, D. B., Nakai, N., & Young, J. S. 1994, *AJ*, 107, 2006
- Downes, D. & Solomon, P. M. 1998, *ApJ*, 507, 615
- Dumke, M., Nieten, Ch., Thuma, G., Wielebinski, R., & Walsh, W. 2001, *A&A*, 373, 853
- Dunne, L., Eales, S., et al. 2000, *MNRAS*, 315, 115D
- Eales, S. A. & Edmunds, M. G. 1996, *MNRAS*, 280, 1167
- Eales, S. A. & Edmunds, M. G. 1997, *MNRAS*, 286, 732
- Elmegreen, B. G. 1989 *ApJ*, 338, 178
- Farquhar, P. R. A., Millar, T. J., & Herbst, E. 1997, *MNRAS*, 269, 641
- Goldreich, P. & Kwan, J. 1974, *ApJ*, 189, 441

- Hartquist, T. W., Caselli, P., Rawlings, J. M. C., Ruffle, D. P., & Williams, D. A. 1998, in *The Molecular Astrophysics of Stars and Galaxies*, T. W. Hartquist & D. A. Williams (eds.), Oxford Science Publications, 101
- Ho, P. T. P., Turner, J. L., & Martin, R. N. 1987, *ApJ*, 322, L67
- Jansen, D. 1995, PhD Thesis, Leiden Observatory, Netherlands
- Kennicutt, R. C., Jr. 1989, *ApJ*, 344, 685
- Kennicutt, R. C., Jr. 1998, *ARA&A*, 36, 189
- Kennicutt, R. C., Jr. 1998, *ApJ*, 498, 541
- Kenney, J. D. P. & Young, J. S. 1989, *ApJ*, 344, 171
- Mao, R. Q., et al. 2000, *A&A*, 358, 433
- Mauersberger, B., Henkel, C., Walsh, W., & Schulz, A. 1999, *A&A*, 341, 256
- Papadopoulos, P. P. 1998, PhD Thesis, University of Toronto, Canada
- Papadopoulos, P. P. & Seaquist, E. R. 1999 *ApJ*, 514, 95
- Papadopoulos, P. P. & Allen, M. L. 2000 *ApJ*, 537, 631
- Polk, K. S., Knapp, G. R., Stark, A. A., & Soifer, B. T. 1991, *ApJ*, 370, 158
- Sanders, D. B., Scoville, N. Z., & Wilson, R. W. 1988, *ApJ*, 332, 432
- Sanders, D. B. et al. 1993, in *Back to the Galaxy*, edited by F. Verter (Kluwer, Dordrecht), 311
- Schmidt, M., 1959, *ApJ*, 129, 243
- Soifer, B. T., et al. 1987, *ApJ*, 320, 238
- Solomon, P. M., Downes, D., Radford, S. J. E., & Barrett, J. W. 1997, *ApJ*, 478, 144
- Strong, A. W., et al. 1988, *A&A*, 207, 1
- Taniguchi, Y., Ohya, Y., & Sanders, D. B. 1999, *ApJ*, 522, 214
- Thomas, H. C., Dunne, L., et al. 2002, *MNRAS*, 329, 747T
- van Dishoeck, E. F. 1998, *ARA&A*, 36, 53

- van Dishoeck, E. F. & Blake, G. A. 1998, *ARA&A*, 36, 317
- Wielebinski, R., Dumke, M., & Nieten, Ch. 1999, *A&A*, 347, 634
- Wilson, T. L., & Rood, R. T. 1994, *ARA&A*, 32, 191
- Wong, T. & Blitz, L. 2002, *ApJ*, 569, 157
- Young, J. S., et al. 1995, *ApJS*, 98, 219
- Zhu, M., Seaquist, E. R., & Kuno, N. 2003, *ApJ*, in press




Fig. 1.— Spectra of $^{12}\text{CO}(1-0)$ and $^{12}\text{CO}(3-2)$ for individual sources. The velocity scale is Heliocentric. The temperature scale is in units of main beam temperature. The letter p denotes those galaxies that belong to a pair.




Fig. 1.— (continued)




Fig. 1.— (continued)




Fig. 1.— (continued)




Fig. 1.— (continued)




Fig. 2.— Plots of CO luminosity versus FIR luminosity, both associated with the 15'' beam for (a) $^{12}\text{CO}(1-0)$ and (b) $^{12}\text{CO}(3-2)$ emission. The lines represent linear regression fits to the data. The data points have been segregated according to two FIR color regimes in the plot. See text for details.




Fig. 3.— Plots of the ratio of CO-to-FIR luminosity versus the projected beamsize for (a) $^{12}\text{CO}(1-0)$ and (b) $^{12}\text{CO}(3-2)$ emission.




Fig. 4.— Distribution of the ratio r_{31} for IR luminous galaxies, (a) by Mauersberger et al. (1999); (b) this paper.




Fig. 5.— Plot of r_{31} versus the projected beamsize for the SLUGS subsample.




Fig. 6.— Plots of (a) r_{31} versus the dust temperature T_{dust} , (b) r_{31} versus the FIR luminosity L_{FIR} applicable to the 15'' beam.




Fig. 7.— Plots of (a) r_{31} versus dust mass M_{dust} , (b) r_{31} versus molecular gas mass $M(\text{H}_2)$ applicable to the $15''$ beam.




Fig. 8.— Plots showing r_{31} versus the surface brightness averaged over the optical diameter for (a) mid-IR 14.3 μm , (b) far-IR 60 μm , (c) SCUBA 850 μm , and (d) H I 21cm line.




Fig. 9.— Plots of r_{31} versus the color indices or ratios involving (a) far-IR and (b) sub-mm to H I 21cm emission.




Fig. 10.— Plot of r_{31} versus the star formation efficiency $SFE = L_{FIR}/M(\text{H}_2)$ measured within a $15''$ beam. The line represents a linear regression fit to the data with $SFE \leq 200 \text{ L}_\odot/\text{M}_\odot$. The data points have been segregated according to gas mass-class and dust IR luminosity-class to show the relationship between L_{FIR} and $M(\text{H}_2)$ and position in the plot. See text for details.




Fig. 11.— Plots of CO luminosities for the 15'' beam versus dust mass from Dunne et al. (2000) for (a) $^{12}\text{CO}(1-0)$ and (b) $^{12}\text{CO}(3-2)$ emission. The dust masses have not been corrected for contamination of the SCUBA 850 μm filter by $^{12}\text{CO}(3-2)$ line emission.




Fig. 12.— Relation between the *SFR* per unit area and the surface density of H₂ measured within a 15'' beam.




Fig. 13.— Plots of r_{31} versus dust temperature T_{dust} for different densities of molecular hydrogen $\log_{10} n(\text{H}_2)$ based on LVG model computations for four Λ values between 10^{-7} and $10^{-4} (\text{km s}^{-1} \text{pc}^{-1})^{-1}$. Also shown are the observed data, assuming $T_{\text{kin}} = T_{\text{dust}}$. The ^{12}CO abundance Z_{CO} is fixed at 10^{-4} , and the isotope abundance ratio $\left[\frac{^{12}\text{CO}}{^{13}\text{CO}} \right] = 40$.




Fig. 14.— Distribution of the X conversion factor derived from LVG computations and its mean value X_{MEAN} for different values of $\Lambda = \frac{Z_{\text{CO}}}{(dv/dr)}$. The units of X is $\text{cm}^{-2} [\text{K km s}^{-1}]^{-1}$ and Λ is in $(\text{km s}^{-1} \text{ pc}^{-1})^{-1}$.




Fig. 15.— Histograms of the $^{12}\text{CO}(1-0)/^{13}\text{CO}(1-0)$ intensity ratio R_{10} for different values of Λ . Dotted lines: LVG modeling results from this work. Solid lines: data taken from Aalto et al. (1995) and Taniguchi et al. (1999). The $\left[\frac{^{12}\text{CO}}{^{13}\text{CO}}\right]$ abundance ratio is fixed at 40. The values for R_{10} observed correspond to a galaxy sample with $L_{\text{FIR}} \geq 10^{10} L_{\odot}$, similar to the SLUGS subsample.

Table 1. ^{12}CO line parameters of observed 60 IR luminous SLUGS galaxies.

Source ^a	RA ₂₀₀₀ ^b h m s	Dec ₂₀₀₀ ^c ° ' ''	cz ^d (km s ⁻¹)	D ^e (Mpc)	I_{10} ^f (K km s ⁻¹)	$\pm\sigma I_{10}$ ^g	I_{32} ^h (K km s ⁻¹)	$\pm\sigma I_{32}$ ⁱ	r_{31} ^j	$-\sigma_{31}$ ^k	$+\sigma_{31}$ ^l
UGC 5376	10 00 27.1	+03 22 28	2050	27.2	13.0	3.3	15.8	3.3	1.21	-0.45	0.33
NGC 3094	10 01 26.0	+15 46 14	2404	31.9	48.5	10.1	37.5	7.7	0.77	-0.25	0.20
NGC 3110	10 04 01.9	-06 28 29	5034	66.3	50.6	10.7	39.6	8.2	0.78	-0.26	0.21
IR 1017+08	10 19 59.9	+08 13 34	14390	185.2	16.1	4.6	7.4	3.2	0.46	-0.28	0.22
NGC 3221	10 22 20.0	+21 34 10	4110	54.2	33.5	7.1	16.9	3.5	0.51	-0.17	0.13
NGC 3367	10 46 34.8	+13 45 02	3037	40.2	38.0	7.9	12.0	2.5	0.32	-0.11	0.08
IR 1056+24	10 59 18.1	+24 32 34	12921	166.9	38.1	8.0	24.3	5.3	0.64	-0.22	0.17
ARP 148	11 03 53.9	+40 51 00	10350	134.5	8.8	2.9	5.5	1.4	0.62	-0.29	0.21
NGC 3583	11 14 10.8	+48 19 03	2136	28.3	44.0	9.4	18.7	4.0	0.43	-0.14	0.11
MCG +00-29	11 21 10.9	-02 59 13	7646	100.0	...	...	27.5	5.8	...	...	...
UGC 6436	11 25 45.0	+14 40 36	10243	133.2	53.3	11.1	18.4	3.8	0.35	-0.11	0.09
NGC 3994p	11 57 36.8	+32 16 39	3118	41.3	21.2	6.3	14.9	3.6	0.70	-0.31	0.23
NGC 3995p	11 57 44.1	+32 17 40	3254	43.0	23.2	5.1	6.3	1.8	0.27	-0.11	0.09
NGC 4045	12 02 42.3	+01 58 38	1981	26.3	57.3	11.7	14.8	3.1	0.26	-0.08	0.07
IR 1211+03	12 13 46.1	+02 48 40	21885	276.7	...	...	11.0	3.0	...	...	...
NGC 4273	12 19 56.0	+05 20 34	2378	31.5	32.9	6.9	21.5	4.5	0.65	-0.22	0.17
IR 1222-06	12 25 03.9	-06 40 53	7902	103.3	9.1	3.6	11.2	2.5	1.23	-0.63	0.44
NGC 4418	12 26 54.7	-00 52 39	2179	28.9	54.8	11.5	47.4	9.7	0.86	-0.28	0.22
NGC 4433	12 27 38.6	-08 16 45	3000	39.7	108.9	22.7	40.8	8.4	0.37	-0.12	0.10
NGC 4793	12 54 40.7	+28 56 19	2484	32.9	71.5	14.5	26.2	5.3	0.37	-0.12	0.09
NGC 4922	13 01 24.8	+29 18 36	7071	92.6	12.3	5.1	< 2.04	...	...	...	...
NGC 5020	13 12 39.9	+12 35 59	3362	44.5	39.2	8.0	20.2	4.3	0.52	-0.23	0.18
IC 860	13 15 03.5	+24 37 08	3347	44.3	< 5.07	...	< 2.67	...	...	...	...
UGC 8387	13 20 35.3	+34 08 22	7000	91.7	90.6	18.4	63.3	12.9	0.70	-0.21	0.16
NGC 5104	13 21 23.1	+00 20 32	5578	73.4	62.6	12.7	33.6	6.8	0.54	-0.15	0.12
NGC 5256	13 38 17.5	+12 35 58	8353	109.1	...	...	12.3	2.9	...	...	...
NGC 5257p	13 39 53.0	+00 50 22	6757	88.6	33.1	7.1	20.7	4.3	0.63	-0.35	0.28
NGC 5258p	13 39 57.9	+00 49 58	6798	89.1	37.7	8.4	16.1	3.5	0.43	-0.12	0.09
UGC 8739	13 49 14.2	+35 15 23	5032	66.3	44.2	9.0	48.3	9.9	1.09	-0.05	0.05
NGC 5371	13 55 39.0	+40 27 31	2553	33.8	...	...	15.5	3.3	...	...	...
NGC 5394p	13 58 33.6	+37 27 13	3472	45.9	101.5	21.1	36.0	7.3	0.35	-0.63	0.44
NGC 5395p	13 58 37.6	+37 25 32	3491	46.1	30.4	6.6	...	...	...	...	...
NGC 5433	14 02 36.0	+32 30 38	4354	57.4	24.1	7.2	40.6	8.6	1.69	-0.27	0.20
NGC 5426p	14 03 25.1	-06 04 09	2621	34.7	18.2	4.8	5.0	1.7	0.27	-0.31	0.24
NGC 5427p	14 03 25.6	-06 01 42	2678	35.5	12.1	3.1	7.7	2.0	0.64	-0.26	0.19
ZW 247.020	14 19 43.2	+49 14 12	7666	100.3	28.5	6.2	25.9	5.5	0.91	-0.20	0.16
NGC 5600	14 23 49.4	+14 38 21	2319	30.7	11.0	3.0	7.2	1.5	0.65	-0.07	0.06
NGC 5653	14 30 10.4	+31 12 54	3562	47.1	61.1	12.6	38.4	7.7	0.63	-0.61	0.45
NGC 5665	14 32 25.8	+08 04 45	2228	29.5	78.0	16.6	16.8	3.5	0.22	-0.30	0.23
NGC 5676	14 32 46.7	+49 27 29	2114	28.0	...	...	14.7	3.2	...	...	...
NGC 5713	14 40 11.3	-00 17 27	1900	25.2	16.4	4.1	27.2	5.5	1.65	-0.29	0.22
UGC 9618p	14 57 00.5	+24 36 42	9900	128.8	...	...	5.2	1.7	...	...	...
NGC 5792	14 58 22.8	-01 05 27	1924	25.5	35.0	7.4	31.6	6.5	0.90	-0.15	0.11
ZW 049.057	15 13 13.1	+07 13 27	3897	51.5	27.4	6.0	22.8	4.9	0.83	-0.57	0.42
NGC 5900	15 15 05.0	+42 12 32	2511	33.3	74.8	15.4	33.6	6.9	0.45	-0.22	0.16
1 ZW 107	15 18 06.1	+42 44 45	12039	155.8	...	...	14.6	3.3	...	...	...
NGC 5929p	15 26 06.1	+41 40 14	2492	33.0	5.9	3.4	5.1	1.2	0.87	-0.22	0.17
NGC 5930p	15 26 07.9	+41 40 34	2672	35.4	21.5	5.7	11.7	2.6	0.54	-0.60	0.46
IR 1525+36	15 26 59.4	+35 58 37	16602	212.6	...	...	10.3	3.3	...	...	...
NGC 5936	15 30 00.8	+12 59 21	4004	52.9	56.4	11.8	37.8	7.6	0.67	-0.28	0.22
NGC 5937	15 30 46.2	-02 49 45	2807	37.2	18.8	4.3	32.2	6.5	1.72	-0.13	0.10
NGC 5953p	15 34 32.3	+15 11 38	1965	26.1	45.7	9.8	38.7	7.9	0.85	-0.19	0.15
NGC 5954p	15 34 35.0	+15 12 00	1959	26.0	38.6	8.3	14.4	3.2	0.37	-0.10	0.08
ARP 220	15 34 57.1	+23 30 11	5434	71.5	126.3	26.2	74.2	14.9	0.59	-0.11	0.09
IR 1533-05	15 36 11.7	-05 23 52	8186	107.0	...	...	8.9	2.2	...	...	...
NGC 5962	15 36 31.7	+16 36 29	1958	26.0	86.2	17.6	26.9	5.5	0.31	-0.22	0.16
NGC 5990	15 46 16.5	+02 24 56	3839	50.7	86.7	20.0	28.1	5.9	0.32	-0.38	0.29
NGC 6052	16 05 12.9	+20 32 32	4716	62.1	30.6	8.0	17.8	3.7	0.58	-0.12	0.09
MCG +01-42	16 30 56.5	+04 04 59	7342	96.1	26.4	6.3	27.6	5.7	1.05	-0.37	0.28
NGC 6181	16 32 21.0	+19 49 36	2375	31.5	60.0	12.8	21.3	4.4	0.36	-0.12	0.09

^aGalaxy name, the letter p denotes those galaxies that belong to a pair.^bRight ascension J2000 epoch from NED^cDeclination J2000 epoch from NED.^dRecession velocity from NED.^eGalaxy proper distance to the object at the present time.^fThe $^{12}\text{CO}(1-0)$ line intensity $I_{10} = \int T_{mb} dv$ in K km s⁻¹. Upper limits refer to 3 times the standard error due to the noise.^gStandard error of I_{10} .^hThe $^{12}\text{CO}(3-2)$ line intensity $I_{32} = \int T_{mb} dv$ in K km s⁻¹. Upper limits refer to 3 times the standard error due to the noise.ⁱStandard error of I_{32} .^jThe line intensity ratio r_{31} .^kThe negative uncertainty in r_{31} .^lThe positive uncertainty in r_{31} .

Table 2. CO luminosities and masses.

Source ^a	P.B. ^b (kpc)	$\log_{10} L_{12CO(1-0)}^c$ (K km s ⁻¹ Mpc ² Ω_b)	$\log_{10} L_{12CO(3-2)}^d$ (K km s ⁻¹ Mpc ² Ω_b)	$\log_{10} M(\text{H}_2)^e$ ($M_\odot$)	SFE^f ($L_\odot/M_\odot$)
UGC 5376	1.99	3.98	4.06	7.34	177
NGC 3094	2.34	4.69	4.58	8.05	102
NGC 3110	4.90	5.34	5.23	8.71	98
IR 1017+08	14.12	5.72	5.39	9.12	96
NGC 3221	4.00	4.99	4.69	8.36	43
NGC 3367	2.95	4.78	4.28	8.15	32
IR 1056+24	12.66	6.01	5.81	9.40	197
ARP 148	10.12	5.19	4.98	8.57	302
NGC 3583	2.07	4.54	4.17	7.91	68
MCG +00-29	7.46	...	5.43	...	...
UGC 6436	10.02	5.96	5.50	9.35	94
NGC 3994p	3.03	4.55	4.40	7.92	61
NGC 3995p	3.16	4.63	4.07	7.99	59
NGC 4045	1.92	4.59	4.01	7.96	82
IR 1211+03	21.59	...	5.89	...	...
NGC 4273	2.31	4.51	4.33	7.87	84
IR 1222-06	7.71	4.98	5.07	8.36	142
NGC 4418	2.12	4.66	4.59	8.02	461
NGC 4433	2.92	5.23	4.80	8.60	46
NGC 4793	2.41	4.89	4.45	8.25	28
NGC 4922	6.90	5.01	...	8.39	40
NGC 5020	3.27	4.88	4.60	8.25	68
IC 860	3.25	...	...	...	...
UGC 8387	6.83	5.87	5.72	9.25	130
NGC 5104	5.43	5.52	5.25	8.89	96
NGC 5256	8.15	...	5.16	...	...
NGC 5257p	6.59	5.41	5.20	8.78	27
NGC 5258p	6.63	5.47	5.10	8.84	24
UGC 8739	4.90	5.28	5.32	8.65	73
NGC 5371	2.48	...	4.25	...	...
NGC 5394p	3.38	5.32	4.87	8.69	15
NGC 5395p	3.39	4.81	...	8.17	8
NGC 5433	4.24	4.89	5.12	8.26	160
NGC 5426p	2.55	4.34	3.77	7.70	14
NGC 5427p	2.60	4.18	3.98	...	...
ZW 247.020	7.48	5.45	5.40	8.82	169
NGC 5600	2.25	4.01	3.83	7.37	121
NGC 5653	3.46	5.13	4.92	8.49	68
NGC 5665	2.16	4.83	4.16	8.19	11
NGC 5676	2.05	...	4.06	...	...
NGC 5713	1.85	4.02	4.24	7.38	282
UGC 9618p	9.68	...	4.92	...	...
NGC 5792	1.87	4.36	4.31	7.72	127
ZW 049.057	3.79	4.85	4.77	8.22	517
NGC 5900	2.44	4.91	4.57	8.28	53
1 ZW 107	11.79	...	5.53	...	...
NGC 5929p	2.42	3.80	3.74	...	...
NGC 5930p	2.60	4.43	4.16	7.79	99

Table 2—Continued

Source ^a	P.B. ^b (kpc)	$\log_{10} L_{12CO(1-0)}$ ^c (K km s ⁻¹ Mpc ² Ω_b)	$\log_{10} L_{12CO(3-2)}$ ^d (K km s ⁻¹ Mpc ² Ω_b)	$\log_{10} M(\text{H}_2)$ ^e (M _⊙)	SFE ^f (L _⊙ /M _⊙)
IR 1525+36	16.31	...	5.64	...	...
NGC 5936	3.90	5.19	5.02	8.56	82
NGC 5937	2.73	4.41	4.64	7.77	469
NGC 5953p	1.91	4.49	4.42	7.85	58
NGC 5954p	1.90	4.41	3.99	7.78	61
ARP 220	5.29	5.80	5.57	9.17	467
IR 1533-05	7.99	...	5.00	...	...
NGC 5962	1.90	4.76	4.26	8.12	13
NGC 5990	3.73	5.34	4.85	8.71	58
NGC 6052	4.59	5.07	4.83	8.44	102
MCG +01-42	7.16	5.38	5.40	8.75	247
NGC 6181	2.31	4.77	4.32	8.13	26

^aGalaxy name, the letter p denotes those galaxies that belong to a pair.

^bThe projected beamsizes (P.B.) at the source.

^cThe ¹²CO(1-0) luminosity measured within a 15'' beam, $\Omega_b = 1$ beam area.

^dThe ¹²CO(3-2) luminosity measured within a 15'' beam, $\Omega_b = 1$ beam area.

^eMolecular hydrogen gas mass measured within a 15'' beam.

^fStar formation efficiency measured within a 15'' beam.