

Equilibrium Models of Galaxy Clusters with Cooling, Heating and Conduction

M.Brüggen^{1,2}

m.brueggen@iu-bremen.de

ABSTRACT

It is generally argued that most clusters of galaxies host cooling flows in which radiative cooling in the centre causes a slow inflow. However, recent observations by Chandra and XMM conflict with the predicted cooling flow rates. Amongst other mechanisms, heating by a central active galactic nucleus and thermal conduction have been invoked in order to account for the small mass deposition rates. Here, we present a family of hydrostatic models for the intra-cluster medium where radiative losses are exactly balanced by thermal conduction and heating by a central source. We describe the features of this simple model and fit its parameters to the density and temperature profiles of Hydra A.

Subject headings: galaxies: active - galaxies: clusters: cooling flows - X-rays: galaxies

1. Introduction

The X-ray surface brightness of many clusters of galaxies shows a strong central peak which has been interpreted as the signature of a cooling flow (Cowie & Binney 1977, Fabian & Nulsen 1977, Sarazin 1988, Fabian 1994). However, the simple cooling flow model conflicts with a growing number of observations that show that while the temperature is declining in the central region, gas with a temperature below ~ 1 keV is significantly less abundant than predicted. The absence of a cool phase in cooling flows has proven a persistent puzzle. Recently, two main candidates for the heating of the gas in the central regions of clusters have emerged: (i) heating by outflows from active galactic nuclei (AGN) (Tabor & Binney 1993, Churazov et al. 2001a, Binney 2001, Brüggen et al. 2002, Brüggen 2003), and (ii)

¹International University Bremen, Campus Ring 1, 28759 Bremen, Germany

²Max-Planck Institut für Astrophysik, Karl-Schwarzschild-Str 1, 85740 Garching, Germany

transport of heat from the outer regions of the cluster by thermal conduction (Zakamska & Narayan 2003, Ruszkowski & Begelman 2002, Fabian et al. 2001, Voigt et al. 2002, Gruzinov 2002, Friaca 1986, Bertschinger & Meiksin 1986, Meiksin 1988, Bregman & David 1988).

The role of thermal conduction in the ICM has been the subject of a long debate and, owing to the complex physics of MHD turbulence, the value of the effective conductivity remains uncertain. The thermal conductivity of an unmagnetised, fully ionised plasma was calculated by Spitzer (1962). Originally it has been thought that the magnetic field in clusters strongly suppresses the thermal conductivity because the magnetic fields prevent an efficient transport perpendicular to the field lines. Even if the transport can be efficient along the magnetic field lines, the overall isotropic conductivity was thought to be many orders of magnitude less than the Spitzer value. This paradigm has been supported by a number of observations, such as sharp edges at so-called cold fronts, small-scale temperature variations in mergers and sharp boundaries around radio bubbles. It is thought that the existence of these sharp features precludes thermal conduction near the Spitzer value (Markevitch et al. 2000, Vikhlinin, Markevitch & Murray 2001).

Recent theoretical work by Narayan & Medvedev (2001), Malyskin & Kulsrud (2001), Chandran et al. (1999), Chandran & Cowley (1998) and earlier work by Rechester & Rosenbluth (1978) has shown that a turbulent magnetic field is not as efficient in suppressing thermal conduction as previously thought. It is argued that chaotic transverse motions of the tangled magnetic field lines can enhance the cross-field diffusion to an extent that the effective conductivity is of the order of the Spitzer value. Following this work, Zakamska & Narayan (2003) have looked for hydrostatically stable models in which the radiative cooling is exactly balanced by thermal conduction and where the temperatures on the inner and outer boundaries were fixed. For half of the clusters they investigated they found that a thermal conductivity of around 30 % of the Spitzer value yielded good fits to the observed profiles. However, for the other half of the clusters conduction alone appeared unlikely to halt a cooling catastrophe. Consequently, some form of heating is necessary such as mechanical heating by AGN. The work by Zakamska & Narayan (2003) has motivated us to reexamine cluster models that takes into account thermal conduction as well as heating by central radio sources. We include a more generally valid treatment of radiative cooling by utilising a fit to the cooling function that takes into account line cooling. This becomes particularly important in the central regions of cooler clusters.

Radio-loud AGN drive strong outflows in the form of jets that inflate bubbles or lobes. The lobes are filled with hot plasma, and can heat the cluster gas in various ways. The effect of hot bubbles on the ICM consists primarily of heating via pdV work and redistribution of

mass via buoyancy-driven mixing. Hydrodynamic simulations (Brüggen 2003) have shown that a significant fraction ($\sim 10\%$) of the energy residing in radio lobes can be dissipated in the cluster gas.

Clearly, the lifetime of the activity of the central AGN (as brief as $\sim 10^4 - 10^5$ yrs) is short compared to the evolutionary time scale of the cluster gas. Therefore, once the AGN stops supplying energy to the buoyant bubbles, the cluster gas will settle down once again and a full cooling flow may be re-established. The cooling gas flows to the centre of the cluster and may then trigger a further active phase of the AGN. Thus a self-regulating process with cooling periods alternating with brief bursts of AGN activity may be established (Quilis, Bower, & Balogh 2001, Voit & Bryan 2001). The rising bubbles uplift colder material from the vicinity of the AGN and thus disrupt the supply of fresh fuel for the radio jets. This feedback mechanism might automatically regulate the power of the radio jets. It has been found that 71 % of all cD galaxies at the centres of cooling-flow clusters show evidence of radio activity (Burns 1990). This fraction is higher than in non-cooling flow galaxies which again points towards the existence of some form of feedback mechanism where the cooling gas flows to the centre of the cluster and triggers a further active phase of the AGN. The bubbles rise at a speed that is comparable to the sound speed, which in turn is comparable to the dynamical speed. Therefore, the cooling timescale is much longer than the bubble rise timescale, and it is justifiable to treat feedback heating in a time-averaged sense.

In this paper, we will present a simple model that incorporates, both, thermal conduction and heating by bubbles. We compute a class of models that are subject to radiative cooling, thermal conduction, heating by AGN and that are in hydrostatic equilibrium. We will investigate those parameters that yield plausible solutions and will compare our solutions to observations. Finally, we briefly discuss the origin of the energy that is conducted into the cluster.

2. Model

In our model of the cluster, we assume that the dark matter distribution is given by a modified NFW profile (Navarro et al. 1997)

$$\rho_{\text{DM}}(r) = \frac{M_0/2\pi}{(r + r_c)(r + r_s)^2}, \quad (1)$$

where r is the distance from the centre, r_s the scale radius of the NFW profile and r_c a softening radius, within which the density becomes constant and which prevents the temperature from going to zero at $r = 0$. The core radius r_c determines the shape of the potential near

the centre and here we adopt $r_c = r_s/20$ as recommended by Zakamska & Narayan (2003). We can express the characteristic mass M_0 in terms of the commonly used concentration parameter $c = (3M_{\text{vir}}/4\pi 200\rho_{\text{crit}}(z)r_c^3)^{1/3}$, where ρ_{crit} is the critical density of the universe at the redshift of the cluster and M_{vir} is the virial mass:

$$M_0 = 2\pi r_s^2 r_c \rho_{\text{crit}}(z) \left(\frac{200}{3} \right) \frac{c^3}{\ln(1+c) - c/(1+c)} . \quad (2)$$

The conductive heat flux is given by

$$F_c = -\nabla \cdot (\kappa \nabla T), \quad (3)$$

where κ is the thermal conductivity and T temperature. If thermal conduction is due to electrons, the conductivity according to Spitzer (1962) is given by

$$\kappa_{\text{sp}} \approx 9.2 \cdot 10^{-7} T^{5/2} \text{erg s}^{-1} \text{K}^{-1} \text{cm}^{-1}. \quad (4)$$

Here we seek an equilibrium model that is spherically symmetric, static, and time-independent. Moreover, we neglect magnetic fields and the self-gravity of the ICM. In spherical coordinates, the gravitational potential Φ is governed by the dark matter distribution

$$\frac{1}{r^2} \frac{d}{dr} (r^2 \Phi) = 4\pi G \rho_{\text{DM}}(r) = \frac{2GM_0}{(r+r_c)(r+r_s)^2} , \quad (5)$$

where G is the gravitational constant. Following Zakamska & Narayan (2003) the mass-temperature relation of Afshordi & Cen (2002) and the mass-scale relation of Maoz et al. (1997), can be used to determine M_0 and r_s for a given cluster. The equation of hydrostatic equilibrium can now be written as

$$\frac{dp}{dr} = -\rho \frac{d\Phi}{dr} , \quad (6)$$

where p is pressure and ρ gas density. Demanding that radiative cooling and heating are balanced by thermal conduction, we can write

$$\frac{1}{r^2} \frac{d}{dr} (r^2 F_c) = -\rho \mathcal{L} + \mathcal{H} , \quad (7)$$

where $\rho \mathcal{L}$ and $\mathcal{H}$ are the volume cooling and heating functions, respectively, and F_c is the conductive flux, which depends on the temperature gradient as stated above

$$\kappa \frac{dT}{dr} = -F_c, \quad (8)$$

where we write κ as $f\kappa_{\text{sp}}$, f being a suppression factor. We should point out that we omitted a term for the convective flux in equation (7). However, for the levels of heating considered here the ICM is convectively stable and the neglect of convection is a justifiable assumption.

Pressure is related to n_e and T via the ideal gas law

$$p = \frac{\rho kT}{\mu m_u} = \frac{\mu_e}{\mu} n_e kT, \quad (9)$$

where m_u is the atomic mass unit.

For the volume cooling rate $\rho\mathcal{L}$ we use an approximation to the cooling function based on calculations by (Sutherland and Dopita 1993)

$$\rho\mathcal{L} = [C_1(kT)^\alpha + C_2(kT)^\beta + C_3] \frac{\mu_e}{\mu} n_e^2 10^{-22} \text{ erg cm}^3 \text{ s}^{-1}. \quad (10)$$

The units for kT are keV, n_e is the electron number density, and μ and μ_e denote the mean molecular weight per hydrogen atom and per electron, respectively. As in Zakamska & Narayan (2003) we use $\mu = 0.62$ and $\mu_e = 1.18$, corresponding to a fully ionized gas with hydrogen fraction $X = 0.7$ and helium fraction $Y = 0.28$. For an average metallicity $Z = 0.3Z_\odot$ the constants are $\alpha = -1.7$, $\beta = 0.5$, $C_1 = 8.6 \times 10^{-3}$, $C_2 = 5.8 \times 10^{-2}$ and $C_3 = 6.4 \times 10^{-2}$.

The term $\mathcal{H}$ in equation (7) represents the energy input by a central AGN. The energy is deposited in the ICM by the rising bubbles and thus, averaged over time, the heating will be distributed in radius. To quantify this heating, we use a prescription proposed by Begelman (2001) which quantifies the heating of the ICM by the rising bubbles. Assuming that this heating mechanism reaches a quasi-steady state, the details of the bubble motions and geometry should cancel and the energy flux may be written as

$$\dot{e} \propto p_b(r)^{(\gamma_b-1)/\gamma_b}, \quad (11)$$

where $p_b(r)$ is the partial pressure of buoyant gas inside bubbles at radius r and γ_b is the adiabatic index of buoyant gas which we will take to be $4/3$. Assuming that the partial pressure scales with the thermal pressure of the ICM, the volume heating function $\mathcal{H}$ can be expressed as

$$\mathcal{H} \sim -h(r)\nabla \cdot \frac{\dot{e}}{4\pi r^2} = -h(r) \left(\frac{p}{p_0}\right)^{(\gamma_b-1)/\gamma_b} \frac{1}{r} \frac{d \ln p}{d \ln r}, \quad (12)$$

where p_0 is the central pressure and $h(r)$ is the normalization function

$$h(r) = \frac{L}{4\pi r^2} (1 - e^{-r/r_0}) q^{-1}, \quad (13)$$

where L is the luminosity of the AGN. The normalisation factor q is defined by

$$q = \int_{r_{\min}}^{r_{\max}} \left(\frac{p}{p_0}\right)^{(\gamma_b-1)/\gamma_b} \frac{1}{r} \frac{d \ln p}{d \ln r} (1 - e^{-r/r_0}) dr, \quad (14)$$

where r_0 is the inner heating cutoff radius which is determined by the finite size of the central radio source. Here r_0 was taken to be 10 kpc. This heating function mimics a possible feedback mechanism between the AGN and the ICM in the sense that the volume heating function does not depend on the physical conditions at the source alone but on the pressure gradient (Ruszkowski & Begelman 2002). Thus, it resembles thermal conduction, with the difference that the heating rate depends on the gradient of pressure rather than temperature.

Equations (5) - (9) can be combined to yield a set of four first-order differential equations for n_e , $r^2 d\Phi/dr$, T and $r^2 F_c$. We solve these equations as an initial-value problem subject to the initial conditions: $n_e(0) = n_{\text{in}}$, $r^2 d\Phi/dr|_{r=0} = 0$, $T(0) = T_{\text{in}}$ and $r^2 F_c|_{r=0} = 0$, where n_{in} and T_{in} are parameters. Other parameters in this simplified model are M_0 , r_s , f and L . We have investigated the dependence of the resulting density and temperature profiles on some of these parameters, which is discussed in the next session.

3. Results and Discussion

We integrate the system of four ordinary differential equations using a Runge-Kutta method and adopt a characteristic mass of $M_0 = 6.6 \times 10^{14} M_\odot$ and a scale radius of $r_s = 460$ kpc (parameters inferred for the cluster Abell 1795, Zakamska & Narayan 2003). For a cluster with a central temperature of 2 keV and a central density of $n_e = 0.05 \text{ cm}^{-3}$ the resulting density and temperature profiles are shown in Figures 1 and 2. Curves plotted in different styles correspond to different values of the suppression factor f and the luminosity of the central source L . The values assumed for L here correspond to typical energies supplied by the jet to the ICM of $\sim 10^{44} \text{ erg s}^{-1}$ (e.g. Owen, Eilek and Kassim 2000). In Figure 1 one can note that the temperature profile rises less steeply with radius if thermal conduction is

more efficient. In order to maintain a given central temperature, the temperature gradient in the cluster has to be higher if conduction is more suppressed. The higher temperature gradient makes up for the smaller suppression factor so that enough energy is conducted inwards to balance the radiative losses. The highest value for the suppression factor that we have adopted here (0.5) is at the upper end of what is physically plausible. Even higher thermal conductivities seem very unlikely. Moreover, one can see, that the heating term has an effect similar to that of thermal conduction, in that a higher luminosity of the central source flattens out the temperature profile. Because the distributed heating depends on the pressure gradient, a higher luminosity of the central source can afford a smaller pressure gradient, leading to a shallower temperature distribution. Physically speaking, the smaller pressure gradient makes the energy transfer from the bubbles to the ICM less efficient.

For higher values of f and L the electron number density is higher in the centre (< 100 kpc) and drops off faster in the outer regions (see Fig. 2).

Figures 3 and 4 are the corresponding figures for a cluster with a higher central temperature of 4 keV, all other parameters being the same. For this hotter cluster the luminosity of the central source has to be higher in order to produce the same relative effect as in the cooler cluster. In these plots we also included a model with a high luminosity of $L = 3 \times 10^{45}$ erg s $^{-1}$. In this case the temperature distribution is nearly isothermal.

Clearly, by including a physically motivated heating term, there are more degrees of freedom to fit the density and temperature profiles inferred from observations. This can be demonstrated at the example of Hydra A which is a well-studied cooling flow cluster with a FR I radio source (3C218) at its centre. Zakamska & Narayan (2003) have shown that Hydra A cannot be fitted with a conduction model alone unless one allows for unphysically high values for κ . The NFW parameters for Hydra A are $r_s = 370$ kpc and $M_0 = 2.9 \times 10^{14} M_\odot$. The central electron density is $n_e(0) = 0.08$ cm $^{-3}$, the temperature at the centre is $T_{\text{in}} = 3.11$ keV and at 190 kpc is $T(190 \text{ kpc}) = 4.04$ keV. In Figure 5 we show the temperature profile as inferred from CHANDRA observations (David et al. 2001) together with our best fit for $f = 0.5$ ($L = 4.0 \times 10^{45}$ erg s $^{-1}$). As is readily seen, the inclusion of the heating term produces a good fit to the data without having to invoke an abnormally high conductivity.

A question that we have not addressed so far is the question of the origin of the energy that is conducted inwards in order to keep the cluster from collapsing. If we take a cluster model with no heating, i.e. $L = 0$, and a suppression factor of $f = 0.1$ (all other parameters being those of Fig. 1), the asymptotic value of $r^2 F_c$ at large radii is 6.2×10^{44} erg s $^{-1}$. This means that over a period of 5 Gyrs a total energy of $\sim 10^{62}$ erg has to be conducted into the cluster which amounts to ~ 5 % of the total thermal energy of the cluster ($3M_0 k \bar{T} / 2 \mu m_p$). For higher values of f this percentage increases. Ultimately, this energy must be provided

by infalling gas at the accretion shock.

It was pointed out by Loeb (2002) that a significant thermal conductivity will also lead to energy transport out of the cluster because beyond the temperature maximum of the cluster the temperature begins to drop again and the temperature gradient is reversed. Therefore, conduction is also reversed and heat is transported from the cluster to its surrounding envelope in these regions. As estimated by Loeb (2002), heat conduction must be suppressed significantly over the Spitzer value in the outer cluster regions or else the cores of X-ray clusters would have cooled significantly over their life times.

3.1. Stability of the solutions

We have not performed a stability analysis of the model presented here. Instead we refer the reader to relevant works that have been published by others. The thermal stability of stratified atmospheres in the presence of radiative cooling and heat conduction was first analysed by Field (1965) and later generalised by Balbus (1988) and Balbus and Soker (1989) who performed a Lagrangian stability analysis. Recently, the issue of thermal instability in clusters was revisited by Kim & Narayan (2003) who examined the global stability of the models found by Zakamska & Narayan (2003). On the basis of a Lagrangian perturbation analysis they concluded that the growth time of the most unstable global radial mode was $\sim 6 - 9$ times longer than the growth time of the local isobaric modes at the centre if the conductivity was 20 % to 40 % of the Spitzer value. Typical growth times are thus comparable with the time since the last cluster merger. Hence it is argued that the cluster is thermally stable if there is a sufficient amount of thermal conduction. The presence of a heating term as implemented in our model will act to increase the stability of the cluster.

3.2. Summary

We have devised a one-dimensional hydrostatic model for the ICM where radiative losses are balanced by, both, thermal conduction and heating by a central source. By including a physically motivated heating term we have shown that it is possible to fit cluster profiles without having to invoke unphysically high values for the thermal conductivity.

We should concede, however, that our model is very simplified: It assumes that the cluster is in hydrostatic equilibrium and spherically symmetric, it does not allow the gas to condense and drop out of the flow, it ignores magnetic fields and includes a heuristic treatment of heating by AGN. But despite its simplicity, only a few free parameters suffice

to reproduce the observed profiles of X-ray clusters.

I thank Simon White, Eugene Churazov, Bill Forman and Christian Kaiser for helpful discussions.




Fig. 1.— Temperature profiles for a cluster with central temperature $T_{\text{in}} = 2$ keV and central density of $n_e = 0.05 \text{ cm}^{-3}$ with $f = 0.1$, $L = 3 \times 10^{43} \text{ erg s}^{-1}$ (solid) and $f = 0.1$, $L = 3 \times 10^{44} \text{ erg s}^{-1}$ (dotted), $f = 0.5$, $L = 3 \times 10^{43} \text{ erg s}^{-1}$ (dashed) and $f = 0.5$, $L = 3 \times 10^{44} \text{ erg s}^{-1}$ (dash-dot).

REFERENCES

- Afshordi, N., & Cen, R. 2002 ApJ, 564, 669
 Balbus, S.A. 1988, ApJ, 328, 395




Fig. 2.— Electron number density profiles corresponding to the models presented in Fig. 1.




Fig. 3.— Temperature profiles for a cluster with central temperature $T_{\text{in}} = 4$ keV and central density of $n_e = 0.05 \text{ cm}^{-3}$ with $f = 0.1$, $L = 3 \times 10^{43} \text{ erg s}^{-1}$ (solid) and $f = 0.1$, $L = 3 \times 10^{44} \text{ erg s}^{-1}$ (dotted), $f = 0.1$, $L = 3 \times 10^{45} \text{ erg s}^{-1}$ (dashed), $f = 0.5$, $L = 3 \times 10^{43} \text{ erg s}^{-1}$ (dash-dot) and $f = 0.5$, $L = 3 \times 10^{44} \text{ erg s}^{-1}$ (dash-dot-dot).




Fig. 4.— Electron number density profiles corresponding to the models presented in Fig. 3.




Fig. 5.— Gas temperature profile of the Hydra A cluster. The crosses show the results of ACIS data published in David et al. (2001). The line corresponds to the fit with $f = 0.5$ and $L = 4.0 \times 10^{45} \text{ erg s}^{-1}$.

- Balbus, S. A. & Soker, N. 1989, ApJ, 341, 611
- Begelman, M.C. 2001, in *Gas and Galaxy Evolution*, APS Conf. Proc., vol. 240, ed. Hibbard, J.E., Rupen, M.P., and van Gorkom, J.H., p. 363, (astro-ph/0207656)
- Bertschinger, E., & Meiksin, A. 1986, ApJ, 306, L1
- Binney, J. 2001, 'AGN and cooling flows' in *Particles and Fields in Radio Galaxies*, ed. Laing, R. A. , Blundell, K. M., ASP Conference Series, ASP, San Francisco, in press (astro-ph/0103398)
- Bregman, J.N., & David, L.P. 1988, ApJ, 326, 639
- Brüggen, M., Kaiser, C. R., Churazov, E., Ensslin, E. 2002, MNRAS, 331, 545
- Brüggen, M. & Kaiser, C. R. 2002, Nature, 418, 301
- Brüggen, M. 2003, ApJ, submitted
- Burns, J. O. 1990, AJ, 99, 14
- Churazov, E. and Brüggen, M. and Kaiser, C. R. and Böhringer, H. and Forman, W. 2001a, ApJ, 554, 261
- Churazov, E., Sunyaev, R., Forman, W., & Böhringer, H. 2002, MNRAS, 332, 729
- Cowie, L. L., Binney, J. 1977, ApJ, 215, 723
- David, L.P., Nulsen, P.E.J., McNamara, B.R., Forman, W., Jones, C., Ponman, T., Robertson, B., & Wise, M. 2001, ApJ, 557, 546
- Fabian, A. C., Nulsen, P. E. J. 1977, MNRAS, 180, 479
- Fabian, A. C. 1994, ARA&A, 32, 277
- Fabian, A.C., Mushotzky, R.F., Nulsen, P.E.J., & Peterson, J.R. 2001, MNRAS, 321, L20
- Field, G.B. 1965, ApJ, 142, 531
- Friaca, A. C. S. 1986, A&A, 164, 6
- Gruzinov, A. 2002, astro-ph/0203031
- Kim, W.-T., & Narayan, R. 2003, ApJ, submitted (astro-ph/030397)
- Loeb, A. 2002, New Astr., 7, 279

- Malyszhkin, L., & Kulsrud, R. 2001, *ApJ*, 549, 402
- Maoz, D., Rix, H.-W., Gal-Yam, A., & Gould, A. 1997, *ApJ*, 486, 75
- Markevitch, M. et al. 2000, *ApJ*, 541, 542
- Markevitch, M. 2002 (*astro-ph/0205333*)
- Medvedev, M.V., & Narayan, R. 2002, *MNRAS*, submitted (*astro-ph/0107066*)
- Meiksin, A. 1988, *ApJ*, 334, 59
- Narayan, R., & Medvedev, M.V. 2001, *ApJ*, 562, L129
- Navarro, J.F., Frenk, C.S., & White, S.D.M. 1997, *ApJ*, 490, 493
- Owen F.N., Eilek J.A., Kassim N.E., 2000 *ApJ*, submitted, *astro-ph/0006150*
- Quilis, V., Bower, R. G., & Balogh, M. L. 2001, *MNRAS*, 328, 1091
- Rechester, A.B., & Rosenbluth, M.N. 1978, *Phys. Rev. Lett.*, 40, 38
- Rybicki, G.B., & Lightman, A.P. 1979, *Radiative processes in astrophysics* (New York: Wiley)
- Ruszkowski, M., & Begelman, M. C. 2002, *ApJ*, 581, 223
- Sarazin, C. L. 1988, *NATO ASIC Proc. 229: Cooling Flows in Clusters and Galaxies*, Kluwer, Dordrecht, 1
- Spitzer, L. 1962, *Physics of Fully Ionized Gases* (New York: Interscience)
- Sutherland, R.S., & Dopita, M.A. 1993, *ApJS*, 88, 253
- Tabor, G., & Binney, J. 1993, *MNRAS*, 263, 323
- Vikhlinin, A., Markevitch, M., & Murray, S. S. 2001a, *ApJ*, 549, L47
- Vikhlinin, A., Markevitch, M., & Murray, S. S. 2001b, *ApJ*, 551, 160
- Voigt, L.M., Schmidt, R.W., Fabian, A.C., Allen, S.W., & Johnstone R.M., 2002, *MNRAS*, submitted (*astro-ph/0203312*)
- Voit, G. M. & Bryan, G. L. 2001, *Nature*, 414, 425
- Zakamska, N. L., & Narayan, R. 2003, *ApJ*, 582, 162

